

Géométrie vectorielle: Travail #1

Due le 21 mars 2022

Jean-Philippe Miguel-Gagnon

Problem 1

Give an appropriate positive constant c such that $f(n) \leq c \cdot g(n)$ for all $n > 1$.

1. $f(n) = n^2 + n + 1, g(n) = 2n^3$
2. $f(n) = n\sqrt{n} + n^2, g(n) = n^2$
3. $f(n) = n^2 - n + 1, g(n) = n^2/2$

Solution

We solve each solution algebraically to determine a possible constant c .

Part One

$$\begin{aligned} n^2 + n + 1 &= \\ &\leq n^2 + n^2 + n^2 \\ &= 3n^2 \\ &\leq c \cdot 2n^3 \end{aligned}$$

Thus a valid c could be when $c = 2$.

Part Two

$$\begin{aligned} n^2 + n\sqrt{n} &= \\ &= n^2 + n^{3/2} \\ &\leq n^2 + n^{4/2} \\ &= n^2 + n^2 \\ &= 2n^2 \\ &\leq c \cdot n^2 \end{aligned}$$

Thus a valid c is $c = 2$.

Part Three

$$\begin{aligned} n^2 - n + 1 &= \\ &\leq n^2 \\ &\leq c \cdot n^2/2 \end{aligned}$$

Thus a valid c is $c = 2$.