

NouvaDerm

1927nm Thulium Fractional Laser · Pre-Treatment Instructions

PRE-TREATMENT CARE

Please follow these instructions carefully before your NouvaDerm treatment to ensure the best possible results and minimise any risk of adverse reactions.

SKINCARE PREPARATION

- Begin the **Noon Pre-Procedure Program** 10–14 days prior to your appointment.
- Discontinue **retinoids** (tretinoin, retinol) and **AHA/BHA exfoliants** 7 days before treatment.
- Discontinue **vitamin C serums** 3–5 days before treatment.
- Do not use any **self-tanner** or sunless bronzer for 2 weeks prior.
- Arrive with **clean, makeup-free skin** — no creams, serums, or SPF.

SUN & LIFESTYLE

- Avoid direct **sun exposure and tanning beds** for 2–4 weeks prior (strict for Ablative mode; 2 weeks minimum for NOUVAGlo).
- If you have a history of **cold sores (HSV)**, antiviral medication will be prescribed — begin as directed by your physician.
- Avoid **Accutane (isotretinoin)** within 6 months of treatment (12 months for Ablative mode).
- Discontinue **blood thinners and supplements** (fish oil, vitamin E, aspirin unless prescribed) 7 days prior to minimise bruising risk.

DAY OF TREATMENT

- Arrive 30–45 minutes early — topical anaesthetic cream is applied in-clinic and requires time to take effect.
- Avoid caffeine on the day of treatment if you are sensitive.
- Do not drive yourself home following Ablative mode — arrange transport in advance.
- Wear comfortable, loose clothing. Avoid tight collars if treating the neck.

MEDICAL HISTORY — PLEASE INFORM US IF YOU HAVE:

- Active skin infection, open sores, or rash in the treatment area.
- History of keloid or hypertrophic scarring.
- Autoimmune conditions or immunosuppressive medications.
- Pregnancy or breastfeeding.
- Pacemaker or implanted electronic devices.

Questions before your appointment? Contact us at aesthetics@treasuryhealth.ca