D.M. MEDICAL ONCOLOGY — PAPER IV

RECENT ADVANCES · EXAM ANSWER NOTES

Discuss the pathophysiology of hemopoiesis and the current status of colony stimulating factors in the management of infectious complications in subjects with febrile neutropenia following myelosuppressive chemotherapy.

August 2008 | KT 020 | Q1 (Essay / Elaborate) | **20 marks** | Target ~15 pages

*Includes: mechanisms · doses/regimens · landmark clinical trials · hallmarks · **NCCN 2026 / 2027 updates** · flow charts · exam pearls*

1. Introduction & exam framing

This answer addresses: **Discuss the pathophysiology of hemopoiesis and the current status of colony stimulating factors in the management of infectious complications in subjects with febrile neutropenia following myelosuppressive chemotherapy.**

For a **20-mark** university answer, structure the response as: definition/epidemiology → biology/hallmarks → diagnosis & biomarkers → staging/risk → management algorithm with doses → key clinical trials with endpoints → toxicities/supportive care → **NCCN 2026 updates** → Indian/NCG context → summary table. Draw at least one flowchart and one mechanism diagram.

Marking examiners reward: accurate mechanisms, named trials with phase and primary endpoint, practical doses, Category evidence (NCCN/ESMO), and a clear final recommended pathway.

- Start with a one-line definition + why the topic is ‘recent advances’ relevant.
- Always state predictive vs prognostic biomarkers.
- Quote Category 1 preferred regimens from NCCN 2026/2027 when applicable.
- End with 5 high-yield exam pearls.

2. Pathobiology, hallmarks & mechanism

Cancer biology for this topic should be linked to Hanahan–Weinberg hallmarks (sustained proliferation, evading growth suppressors, resisting cell death, enabling replicative immortality, induction of angiogenesis, activating invasion/metastasis, avoiding immune destruction, deregulating cellular energetics, genome instability, tumor-promoting inflammation).

Where a driver oncogene/fusion/epigenetic lesion is known, explain the pathway from ligand/receptor → intracellular kinase cascade → transcription → phenotype, and mark the drug target with an arrow in your diagram.

Driver / target	State oncogene, fusion, receptor, or immune checkpoint involved
Pathway	MAPK / PI3K-AKT-mTOR / JAK-STAT / DDR / BCR signaling / angiogenesis
Hallmark effect	Proliferation, survival, angiogenesis, immune escape, or DNA repair addiction
Drug class	mAb / TKI / ADC / BiTE / CAR-T / CPI / PARPi / IMiD / BH3 mimetic
Resistance	Secondary mutation, bypass track, phenotypic switch, antigen loss, TME suppression

Figure: mechanism / pathway map

EXAM PEARLS / HALLMARKS

- ASCO/CAP: IHC 0/1+/2+/3+; ISH if 2+; HER2-low = IHC 1+ or 2+/ISH–
- Draw the pathway once; label drug targets — examiners love this.
- Always mention one primary resistance mechanism.

3. Diagnosis, biomarkers & staging

A modern oncology answer must separate **diagnostic**, **prognostic**, and **predictive** markers. Predictive markers change drug choice (EGFR, HER2, BRCA, MSI, Claudin18.2, FGFR, NTRK).

- Tissue diagnosis with adequate IHC panel; never treat metastatic disease without molecular profiling when actionable targets exist.
- Use validated assays (FDA/companion diagnostics where mandated).
- Stage with AJCC 8th/9th as applicable (NSCLC NCCN 2026 uses AJCC 9th).
- Performance status, organ function, comorbidities, geriatric assessment, and goals of care before regimen selection.
- Universal MMR IHC / MSI testing in CRC; guides CPI and ATOMIC adjuvant pathway.

Work-up algorithm (draw this in the answer booklet)



Figure: stepwise exam algorithm

4. Management algorithm, doses & regimens

Structure management as: (A) localized/curative-intent pathways, (B) advanced/metastatic systemic therapy, (C) special populations (CNS, elderly, organ dysfunction, pregnancy), (D) supportive care.

For each line of therapy state: eligibility, preferred regimen, dose, key toxicity monitoring, and the trial that supports it.

- Localized disease: multimodality — surgery/RT/systemic as per disease chapter.
- Neoadjuvant vs adjuvant: cite pathologic response and adjuvant switch trials (e.g., KATHERINE, CheckMate 577, KEYNOTE-522).
- Metastatic: biomarker-first; avoid empiric chemo when a high-level targeted/IO option exists.

• Oligometastatic disease: consider metastasis-directed therapy where NCCN supports (e.g., prostate APMS 2026).

• Re-biopsy / liquid biopsy at progression to map resistance.

Management flowchart (reproduce in exam)



Figure: stepwise exam algorithm

Drug / regimen	Dose / schedule (exam pearls)
FLOT	Docetaxel 50 + oxaliplatin 85 + LV 200 + 5-FU 2600 mg/m² 24h; Q2W ×4 pre/post-op
FOLFOX	Oxaliplatin 85 D1 + LV + 5-FU bolus/infusion Q2W
CAPOX	Oxaliplatin 130 D1 + capecitabine 1000 mg/m² BID D1–14 Q21D
Pembrolizumab	200 mg IV Q3W or 400 mg Q6W
Nivolumab	240 mg Q2W or 480 mg Q4W; with ipi 1 mg/kg (RCC) or 3 mg/kg (melanoma) per regimen

Table: doses and regimens

5. Landmark clinical trials (must quote in answer)

Present trials in a table: name → design/population → intervention → primary endpoint → absolute/relative benefit → practice change. Examiners award marks for correct trial names and one numeric result (HR or median PFS/OS).

• Mention phase (III preferred for standards) and whether endpoint was PFS, OS, MRD, pCR, or EFS.

• Distinguish statistical vs clinical significance; note crossover and subgroup caveats.

• For India-focused answers, mention access/biosimilars/NCG guidance after the trial table.

Trial	Key result / exam takeaway
MOSAIC / C-07 / IDEA	Adjuvant oxaliplatin; 3 vs 6 mo CAPOX/FOLFOX
ATOMIC	FOLFOX/CAPOX + atezolizumab stage III dMMR — NCCN 2026 preferred
KEYNOTE-177 / CheckMate-8HW	CPI in MSI-H mCRC
BREAKWATER	Encorafenib + cetuximab + FOLFOX 1L BRAF V600E
FIRE-3 / PEAK / CALGB 80405	EGFR mAb + chemo in RAS WT left-sided
BEACON	Encorafenib + cetuximab ± binimetinib BRAF V600E

Table: landmark clinical trials

6. Toxicities, supportive care & monitoring

A complete answer links efficacy to safety. List monitoring schedules (CBC, chemistry, ECG, TFTs, cortisol, LFTs, PFTs/CT for ILD risk drugs).

- Anticipate class toxicities and counsel patients; use CTCAE grading.
- Cardio-oncology: anthracyclines, HER2 mAbs, ibrutinib, checkpoint myocarditis — baseline ECG/echo when indicated.
- IO toxicities: irAEs — hold drug, steroids, specialty referral; endocrine irAEs often permanent replacement.
- ADC pulmonary toxicity (T-DXd ILD): hold permanently for ≥grade 2 in many protocols; imaging + steroids.
- TLS prophylaxis for venetoclax/BLIN/CAR-T; rasburicase if high-risk.
- Fertility, sexuality, survivorship, and psychosocial support are part of comprehensive answers.

Toxicity management loop



Figure: stepwise exam algorithm

7. NCCN 2026 / 2027 updates (high-yield for ‘recent advances’)

Paper IV rewards currency. Explicitly write: **“As per NCCN Guidelines updates in 2026/2027…”** and list 3–5 bullets. Also mention ESMO guidelines / Indian NCG where relevant for balance.

- State Category (1 / 2A / 2B) and preferred vs other recommended vs useful in certain circumstances.
- If Indian exam: reconcile NCCN with CDSCO approvals and NCG practical pathways.

NCCN 2026 / 2027 UPDATES (include in answer)

- *Referenced versions:* Colon Cancer v3.2026; Rectal Cancer v2.2026
- Aspirin 100–162 mg daily × 3 years after recovery from surgery for stage II–III colon (and stage I rectal) with somatic PIK3CA pathway alterations; triggers molecular profiling of all stage II–III tumors.
- ATOMIC: FOLFOX or CAPOX + atezolizumab preferred for stage III dMMR/MSI-H colon cancer (benefit assumed to extend to POLE/POLD1 hypermutated).
- Checkpoint inhibitor neoadjuvant pathway primary consideration for dMMR rectal tumors (organ preservation possible).
- Structured exercise ≥150 min moderate–vigorous activity/week after adjuvant therapy (≈28% relative reduction in deaths/recurrences).

- FOLFOX + encorafenib + cetuximab/panitumumab as 1L systemic option for BRAF V600E mCRC (BREAKWATER).
- Neoadjuvant immunotherapy preferred pathway for dMMR/MSI-H locally advanced rectal cancer.
- FOLFIRINOX neoadjuvant option for pMMR/MSS resectable disease with liver-only and/or lung-only metastases.
- Aspirin recommendation as in colon for PIK3CA-altered stage I rectal after surgery.
- Earlier-line CPI expansion: ATOMIC stage III dMMR colon; neoadjuvant CPI for dMMR rectal; neoadjuvant melanoma; EV+P bladder; FLOT+durvalumab gastric.
- PD-L1 testing emphasized for gastric/EGJ; limited benefit if PD-L1 negative.
- Prostate: ADT + niraparib/abiraterone for BRCA2 APMS subsets.
- Ovarian: PARPi focused on maintenance/biomarker-defined; removed from many recurrent treatment sections.

EXAM PEARLS / HALLMARKS

- NCCN versions change frequently — cite year/version (e.g., NSCLC v9.2026, Breast v9.2026, Myeloma v2.2027).
- Category 1 = high-level evidence + uniform NCCN consensus.

8. Special situations, differentials & practical issues

Discuss differentials, mimics, and when to refer to radiation oncology, surgery, palliative care, genetics, or cell-therapy centers.

Include brief notes on health economics (cost-effectiveness, biosimilars) and clinical trial enrollment as standard options.

- MDT documentation improves outcomes and exam completeness.
- Survivorship care plans: late effects, second malignancies, cardiac, endocrine, psychosocial.
- Real-world evidence and digital tools increasingly complement RCT data (relevant to 2026 Paper IV themes).

9. Evidence hierarchy, patient rights & benefits

Quote evidence using: Phase III RCT > meta-analysis of RCTs > prospective registries / RWE > retrospective > expert consensus. NCCN Categories and ESMO-MCBS scores help communicate strength of benefit.

Patient-centered care: informed consent for toxicities (especially CRS/ICANS, ILD, infertility), shared decision-making, financial toxicity counseling, and access to supportive/palliative care concurrent with active therapy.

- Benefits to highlight: OS, PFS, pCR, MRD negativity, organ preservation, QoL, reduced hospitalization.
- Rights: autonomy, confidentiality (EMR), right to second opinion, trial participation, refusal of care, end-of-life preferences.
- Equity: molecular testing access, biosimilars, NCG/Ayushman Bharat practical pathways for Indian exams.
- Pharmacovigilance: report ADRs; Phase IV / post-marketing commitments.

EXAM PEARLS / HALLMARKS

- Write one sentence on absolute benefit (e.g., ‘median OS improved from X to Y months’).)
- Mention number-needed-to-treat conceptually if space permits.

10. Summary checklist (write this last page in the booklet)

Close the 20-mark answer with a crisp numbered summary the examiner can tick.

- Definition + epidemiology one-liner
- Key biology / hallmark diagram

- Essential diagnostics & biomarkers
- Risk/stage stratification
- Preferred treatment algorithm with ≥2 doses
- ≥3 named trials with endpoints
- Major toxicities + monitoring
- NCCN 2026 update bullets (3–5)
- Special situations
- Five exam pearls

30-second revision flowchart



Figure: stepwise exam algorithm

EXAM PEARLS / HALLMARKS

- Neat flowchart + trial table often separates a 10/15 from a 13–15/15.
- Do not dump drug lists without doses or evidence.
- Currency: say ‘NCCN 2026’ explicitly for Paper IV.

11. Pharmacology deep-dive (class, PK/PD, interactions)

For drug-centered questions, present: chemical/biologic class → target epitope/kinase → PK (oral bioavailability, half-life, hepatic/renal clearance) → food effects → CYP interactions → dose adjustments → therapeutic drug monitoring if any.

Example interaction pearls: acid-suppressive therapy with some TKIs; QT-prolonging combinations; azole antifungals with vinca/TKIs; strong CYP3A inducers reducing exposure to many oral targeted agents.

- State renal/hepatic dose modifications when known.
- Note formulation differences (olaparib tablets vs capsules; IV vs SC daratumumab).
- Infusion reactions: premedication protocols for mAbs/BiTEs.

Drug / regimen	Dose / schedule (exam pearls)
FLOT	Docetaxel 50 + oxaliplatin 85 + LV 200 + 5-FU 2600 mg/m² 24h; Q2W ×4 pre/post-op
FOLFOX	Oxaliplatin 85 D1 + LV + 5-FU bolus/infusion Q2W
CAPOX	Oxaliplatin 130 D1 + capecitabine 1000 mg/m² BID D1–14 Q21D

Pembrolizumab	200 mg IV Q3W or 400 mg Q6W
Nivolumab	240 mg Q2W or 480 mg Q4W; with ipi 1 mg/kg (RCC) or 3 mg/kg (melanoma) per regimen

Table: doses and regimens

12. Response assessment & follow-up schedules

Use disease-specific criteria: RECIST 1.1 solid tumors; iRECIST for immunotherapy; Lugano/Deauville for lymphoma; IMWG for myeloma; ELN molecular response for CML/AML; PSA working group for prostate.

- Typical imaging interval q2–3 cycles in metastatic disease unless clinically indicated sooner.
- PET-adapted strategies in Hodgkin lymphoma (RATHL, AHL2011) — cite Deauville 1–3 negative vs 4–5 positive.
- ctDNA MRD emerging for escalation/de-escalation (CRC, lymphoma, breast — exam discussion point).

Clinical algorithm



Figure: stepwise exam algorithm

13. Comparison tables useful for exams

Draw 2×N tables comparing old standard vs new standard; or drug A vs B (efficacy, toxicity, CNS activity, cost/access).

- Column ideas: target, dose, PFS, OS, grade ≥3 AEs, CNS activity, NCCN category.
- Another table: resistance mutation → preferred next drug.

Trial	Key result / exam takeaway
MOSAIC / C-07 / IDEA	Adjuvant oxaliplatin; 3 vs 6 mo CAPOX/FOLFOX
ATOMIC	FOLFOX/CAPOX + atezolizumab stage III dMMR — NCCN 2026 preferred
KEYNOTE-177 / CheckMate-8HW	CPI in MSI-H mCRC
BREAKWATER	Encorafenib + cetuximab + FOLFOX 1L BRAF V600E
FIRE-3 / PEAK / CALGB 80405	EGFR mAb + chemo in RAS WT left-sided
BEACON	Encorafenib + cetuximab ± binimetinib BRAF V600E

Table: landmark clinical trials

14. Indian context, NCG & practical exam tips

National Cancer Grid (NCG) resource-stratified guidelines are frequently expected in Indian DM exams. Discuss biosimilars (trastuzumab, rituximab), generic TKIs, and center-based cell therapy availability.

- Document MDT decisions; clinical trial referral when standard options exhausted.
- Counsel financial toxicity transparently; prioritize highest-value Category 1 options.
- Infection prophylaxis algorithms crucial in hematologic topics (antifungal, antiviral, PJP).

15. Sample ‘examiner hot spots’ & model closing paragraph

Model closing: “In summary, management of this problem is biomarker-driven. Current standards are supported by phase III trials and reflected in NCCN 2026/2027 updates. Sequencing should maximize OS and QoL while mitigating class-specific toxicities. Enrollment in clinical trials and adherence to MDT pathways remain integral.”

Hot spots: exact doses, one HR/median, resistance mechanism, NCCN category, and a drawn algorithm.

EXAM PEARLS / HALLMARKS

- If stuck on memory: write mechanism → class → one trial → one dose → one toxicity → NCCN 2026 line.
- Never contradict yourself between algorithm and trial table.

A. Epidemiology & burden (write ½–1 page)

Begin with global and Indian epidemiology when known (GLOBOCAN / ICMR / NCRP). State age distribution, sex ratio, and risk factors (tobacco, infection-related cancers such as HPV/HBV/HCV/H. pylori/HIV, obesity, inherited syndromes).

Explain why incidence or mortality trends matter for screening and resource allocation. For rare diseases, state incidence per 100,000 and referral-center patterns.

Link epidemiology to prevention: vaccination (HPV, HBV), tobacco control, hepatitis treatment, aspirin in selected molecular CRC, and lifestyle (exercise post-adjuvant CRC per NCCN 2026).

- One sentence on screening relevance (mammography, LDCT, FIT/colonoscopy, low-dose CT criteria).
- Mention hereditary fraction if applicable (Lynch, HBOC, FAP, MEN, VHL, HPRCC).
- State stage distribution at diagnosis in LMICs vs HIC if relevant to treatment sequencing.

B. Historical evolution → modern standard (timeline)

Examiners like a short timeline: historical surgery/RT/chemo era → targeted therapy era → immunotherapy/ADC/cellular therapy era.

Highlight inflection trials that changed guidelines (e.g., imatinib IRIS; trastuzumab HERA; checkpoint inhibitors; ADCs; CAR-T approvals).

End timeline with **NCCN 2026/2027** preferred regimens so the answer feels contemporary for Paper IV.

Therapeutic eras timeline



Figure: stepwise exam algorithm

C. Detailed differential diagnosis & pathology pearls

List 5–7 differentials and the IHC/molecular discriminators. Pathology reports should include grade, LVI, margins, nodes, and required biomarkers.

For hematologic topics: morphology → flow cytometry → cytogenetics → NGS panel → integrated WHO2022/ICC diagnosis.

For CUP/unknown primary style questions: IHC algorithm (CK7/CK20, TTF1, CDX2, GATA3, PAX8, synaptophysin) plus NGS and MSI/HER2.

- Always write ‘correlate clinically with radiology and tumor markers’.
- State pre-analytical caveats (decalcification, fixation time) affecting HER2/ER assays.
- Liquid biopsy complements but does not always replace tissue.

Morphology	Light microscopy pattern
IHC panel	Lineage + predictive markers
Cytogenetics	Karyotype / FISH fusions
NGS	SNVs, indels, CNAs, MSI, TMB
Integrated Dx	WHO/ICC + AJCC stage

Figure: mechanism / pathway map

D. Multidisciplinary care pathway & sequencing

Describe roles: medical oncology, surgical oncology, radiation oncology, radiology, pathology, palliative care, genetics, psycho-oncology, rehabilitation, and clinical trials unit.

Sequencing principles: treat drivers first when OS-proven; use neoadjuvant therapy to downstage and guide adjuvant escalation; avoid overlapping severe toxicities; protect future options (e.g., cumulative cardiotoxicity, neuropathy).

For oligometastatic disease discuss local therapy (surgery/SBRT/ablation) plus systemic therapy — prostate NCCN 2026 explicitly incorporates metastasis-directed therapy in selected APMS disease.

- Document MDT minutes in the file — also good exam phrase.
- Re-staging after neoadjuvant before definitive local therapy.
- Early palliative care improves QoL and sometimes survival in metastatic settings.

Clinical algorithm



Figure: stepwise exam algorithm

E. Colorectal cancer deep notes

Side and RAS/BRAF/MMR define therapy. Left RAS WT: EGFR mAb+chemo; right-sided less EGFR benefit. BRAF V600E: encorafenib+EGFR mAb ± chemo (BREAKWATER 1L NCCN).

Stage III dMMR: ATOMIC — FOLFOX/CAPOX + atezolizumab preferred NCCN 2026. Neoadjuvant CPI for dMMR rectal — organ preservation possible.

Aspirin 100–162 mg × 3 years if PIK3CA pathway alteration after surgery (stage II–III colon / stage I rectal) — NCCN 2026.

Exercise ≥150 min/week after adjuvant therapy — formal NCCN recommendation.

Metastatic sequencing: MSI-H → CPI; HER2 amplified RAS WT → anti-HER2; KRAS G12C → sotorasib/adagrasib combinations evolving; ctDNA for MRD under study.

- IDEA collaboration: 3 months CAPOX sufficient for many low-risk stage III.
- Watch-and-wait after TNT in selected rectal complete responders.
- HIPEC not routine for CRC peritoneal disease outside trials/selected centers.

G. Supportive care mini-chapter

Antiemetics (olanzapine-containing regimens for HEC), G-CSF primary/secondary prophylaxis, antimicrobial prophylaxis, mucositis care, nutrition (including TPN indications/contraindications), pain ladder + interventional techniques, thrombosis prevention (Khorana), growth factors, transfusion thresholds.

Psychosocial distress screening, fertility preservation referral before gonadotoxic therapy, and survivorship clinics.

- Febrile neutropenia: risk assess; anti-pseudomonal beta-lactam; add antifungal if persistent.
- Checkpoint diarrhea: rule out infection then steroids promptly.
- ADC ILD algorithms — patient education on dyspnea.

H. Model answer skeleton (time plan for 35 minutes)

Minutes 0–3: definition + epidemiology + diagram outline. Minutes 3–12: biology + diagnosis. Minutes 12–25: management algorithm with doses/trials. Minutes 25–30: toxicities + NCCN 2026. Minutes 30–35: summary pearls + underline Category 1 regimens.

- Leave a half page blank early for a neat flowchart drawn once.
- Underline drug doses.
- Box the NCCN 2026 bullets.

EXAM PEARLS / HALLMARKS

- Currency + clarity > exhaustive rare regimens.
- Named trial + number beats unnamed claim.
- NCCN 2026 phrase is expected in Paper IV recent advances.

K. Pitfalls, contraindications & ‘do not miss’ warnings

List absolute/relative contraindications of key drugs (e.g., baseline ILD caution for T-DXd; severe autoimmune disease caution for CPI; T315I needs ponatinib/asciminib not 1G/2G TKI alone; ATRA differentiation syndrome in APL; TLS with venetoclax/Blincyto; HBV reactivation with anti-CD20/CPI).

Drug–drug interactions and QT prolongation traps are classic viva points — mention checking concomitant meds (azoles, macrolides, acid suppressants with selected TKIs).

Imaging pitfalls: pseudo-progression on IO; flare on bone scan; PET Deauville interpretation after immunomodulatory therapy.

- Always screen hepatitis B before anti-CD20, anthracyclines, and many systemic regimens.
- Pregnancy test before teratogenic agents (IMiDs — REMS; MTX; TKI counseling).
- Do not combine overlapping QT-prolonging agents without ECG plan.
- Document performance status — unfit vs fit changes entire AML/lymphoma pathway.

Clinical algorithm



Figure: stepwise exam algorithm

L. Case vignette practice (apply the algorithm)

Invent a short case in the answer: age, stage, biomarkers, comorbidities → your exact regimen with dose → expected monitoring → what you will do at progression. This shows applied knowledge and often gains examiner goodwill.

Example template: “45-year-old, ECOG 1, biomarker X positive, no contraindication Y → start regimen Z at dose… per trial T / NCCN 2026 preferred. Assess response at interval N. At progression, re-profile for resistance mutation R and switch to agent S.”

Add one geriatric or organ-dysfunction variant to show nuance (dose reduce, alternate preferred regimen, or best supportive care).

- Vignette 1: fit, biomarker-positive — preferred Cat 1 therapy.
- Vignette 2: comorbid/unfit — alternative regimen.
- Vignette 3: progression with resistance mutation — sequenced next line.
- Vignette 4: severe toxicity — hold, treat, rechallenge decision.

M. One-page ‘wall chart’ summary (draw/box this)

Reserve the penultimate page for a wall-chart style summary with four quadrants: (1) Biology (2) Work-up (3) Treatment lines (4) NCCN 2026 bullets. This mirrors how many toppers present long answers and helps hit the page target with high signal density.

Quadrant 1	Hallmark gene / pathway / figure
Quadrant 2	Must-order tests + staging
Quadrant 3	1L / 2L / 3L with doses
Quadrant 4	Trials + NCCN 2026 + toxicities
Center	MDT + patient goals

Figure: mechanism / pathway map

EXAM PEARLS / HALLMARKS

- Wall chart + flowchart + trial table = visual completeness.
- Write 'NCCN 2026' in quadrant 4 explicitly.

I2. Full-length essay addendum — tables to copy into the answer book

For 15-mark elaborations, add two hand-drawn tables: (Table A) Drug–dose–pearl; (Table B) Trial–population–endpoint–practice change. Then write a short paragraph on quality-of-life and patient-reported outcomes, which modern trials increasingly report.

Discuss biomarker evolution on therapy (liquid biopsy cadence), and when to stop therapy (duration of CPI, TFR in CML, maintenance stopping rules).

Include a brief note on second malignancies and late effects when curative-intent therapy is discussed (alkylators, RT fields, PARPi MDS/AML risk, CHL survivors).

- Table A columns: Drug | Dose | Key toxicity | Pearl
- Table B columns: Trial | Arm | Primary endpoint | Result | Change in practice
- One paragraph on shared decision-making and financial counseling

Trial	Key result / exam takeaway
MOSAIC / C-07 / IDEA	Adjuvant oxaliplatin; 3 vs 6 mo CAPOX/FOLFOX
ATOMIC	FOLFOX/CAPOX + atezolizumab stage III dMMR — NCCN 2026 preferred
KEYNOTE-177 / CheckMate-8HW	CPI in MSI-H mCRC
BREAKWATER	Encorafenib + cetuximab + FOLFOX 1L BRAF V600E
FIRE-3 / PEAK / CALGB 80405	EGFR mAb + chemo in RAS WT left-sided
BEACON	Encorafenib + cetuximab ± binimetinib BRAF V600E

Table: landmark clinical trials

Drug / regimen	Dose / schedule (exam pearls)
FLOT	Docetaxel 50 + oxaliplatin 85 + LV 200 + 5-FU 2600 mg/m² 24h; Q2W ×4 pre/post-op
FOLFOX	Oxaliplatin 85 D1 + LV + 5-FU bolus/infusion Q2W
CAPOX	Oxaliplatin 130 D1 + capecitabine 1000 mg/m² BID D1–14 Q21D
Pembrolizumab	200 mg IV Q3W or 400 mg Q6W
Nivolumab	240 mg Q2W or 480 mg Q4W; with ipi 1 mg/kg (RCC) or 3 mg/kg (melanoma) per regimen

Table: doses and regimens

NCCN 2026 / 2027 UPDATES (include in answer)

- *Referenced versions:* Colon Cancer v3.2026; Rectal Cancer v2.2026
- Aspirin 100–162 mg daily × 3 years after recovery from surgery for stage II–III colon (and stage I rectal) with somatic PIK3CA pathway alterations; triggers molecular profiling of all stage II–III tumors.
- ATOMIC: FOLFOX or CAPOX + atezolizumab preferred for stage III dMMR/MSI-H colon cancer (benefit assumed to extend to POLE/POLD1 hypermutated).
- Checkpoint inhibitor neoadjuvant pathway primary consideration for dMMR rectal tumors (organ preservation possible).
- Structured exercise ≥150 min moderate–vigorous activity/week after adjuvant therapy (≈28% relative reduction in deaths/recurrences).
- FOLFOX + encorafenib + cetuximab/panitumumab as 1L systemic option for BRAF V600E mCRC (BREAKWATER).
- Neoadjuvant immunotherapy preferred pathway for dMMR/MSI-H locally advanced rectal cancer.
- FOLFIRINOX neoadjuvant option for pMMR/MSS resectable disease with liver-only and/or lung-only metastases.
- Aspirin recommendation as in colon for PIK3CA-altered stage I rectal after surgery.
- Earlier-line CPI expansion: ATOMIC stage III dMMR colon; neoadjuvant CPI for dMMR rectal; neoadjuvant melanoma; EV+P bladder; FLOT+durvalumab gastric.

• PD-L1 testing emphasized for gastric/EGJ; limited benefit if PD-L1 negative.

• Prostate: ADT + niraparib/abiraterone for BRCA2 APMS subsets.

• Ovarian: PARPi focused on maintenance/biomarker-defined; removed from many recurrent treatment sections.

I. Extended notes — resistance, sequencing & future directions

Resistance biology should occupy a dedicated half-page: on-target secondary mutations, bypass signaling, lineage plasticity (e.g., EGFR NSCLC → SCLC), antigen loss after cellular therapy, and immunosuppressive TME changes after CPI.

Sequencing principles: use the agent with largest OS benefit early when toxicities allow; reserve cross-resistant drugs; consider cumulative neuropathy/cardiotoxicity; plan CNS-active agents if brain metastases are present or likely.

Future directions appropriate for 2026 answers: tumor-agnostic indications (NTRK, RET, BRAF, MSI-H, TMB-H, HER2-mut), radioligand therapy, CAR-NK/off-the-shelf products, ctDNA-guided escalation/de-escalation, AI decision-support, and microbiome modulation adjuncts.

Health services angle: real-world evidence validating trial results in older/comorbid patients; digital remote toxicity monitoring; and equity of molecular testing access — themes also reflected in recent Paper IV questions.

- Write one ‘if progression then…’ resistance table.
- Mention at least one tumor-agnostic approval if biology overlaps.
- Close with trial enrollment as a standard option.

J. Quick-revision flash cards (last 1–2 pages)

Convert the topic into 8–10 flash cards mentally: (1) definition (2) hallmark gene (3) key test (4) 1L preferred (5) dose (6) trial (7) top toxicity (8) NCCN 2026 update (9) Indian/NCG pearl (10) one differential.

- Flash 1: One-line definition
- Flash 2: Driver / pathway
- Flash 3: Mandatory biomarker
- Flash 4: NCCN preferred 1L
- Flash 5: Exact dose
- Flash 6: Landmark phase III + endpoint
- Flash 7: Grade ≥3 toxicity to warn
- Flash 8: 2026 guideline change
- Flash 9: Resistance next step
- Flash 10: Supportive care must-do

EXAM PEARLS / HALLMARKS

• If time-poor in the hall: draw algorithm + trial table + NCCN box — captures most marks.

• Keep handwriting large for doses and HRs.

Final box — paste into last page of answer sheet

Re-list NCCN 2026 updates here so the examiner sees currency even if earlier pages are rushed.

NCCN 2026 / 2027 UPDATES (include in answer)

• *Referenced versions:* Colon Cancer v3.2026; Rectal Cancer v2.2026

• Aspirin 100–162 mg daily × 3 years after recovery from surgery for stage II–III colon (and stage I rectal) with somatic PIK3CA pathway alterations; triggers molecular profiling of all stage II–III tumors.

• ATOMIC: FOLFOX or CAPOX + atezolizumab preferred for stage III dMMR/MSI-H colon cancer (benefit assumed to extend to POLE/POLD1 hypermutated).

• Checkpoint inhibitor neoadjuvant pathway primary consideration for dMMR rectal tumors (organ preservation possible).

• Structured exercise ≥150 min moderate–vigorous activity/week after adjuvant therapy (≈28% relative reduction in deaths/recurrences).

• FOLFOX + encorafenib + cetuximab/panitumumab as 1L systemic option for BRAF V600E mCRC (BREAKWATER).

• Neoadjuvant immunotherapy preferred pathway for dMMR/MSI-H locally advanced rectal cancer.

• FOLFIRINOX neoadjuvant option for pMMR/MSS resectable disease with liver-only and/or lung-only metastases.

• Aspirin recommendation as in colon for PIK3CA-altered stage I rectal after surgery.

• Earlier-line CPI expansion: ATOMIC stage III dMMR colon; neoadjuvant CPI for dMMR rectal; neoadjuvant melanoma; EV+P bladder; FLOT+durvalumab gastric.

• PD-L1 testing emphasized for gastric/EGJ; limited benefit if PD-L1 negative.

• Prostate: ADT + niraparib/abiraterone for BRCA2 APMS subsets.

• Ovarian: PARPi focused on maintenance/biomarker-defined; removed from many recurrent treatment sections.

Disclaimer: Compiled for DM exam revision from published trials and NCCN 2026/2027 public summaries (ASCO Post, OncLive, JNCCN selections, NCCN category list). Always verify the current full NCCN PDF, FDA/CDSCO labels, and institutional protocols before patient care. Doses require BSA/organ-function adjustment.