D.M. MEDICAL ONCOLOGY — PAPER IV

RECENT ADVANCES · EXAM ANSWER NOTES

Methods of HER2 testing and interpretation, therapeutic strategies for HER2+ve Metastatic Breast Cancer.

January 2024 | DM 0124 | Q1 (DEEP Essay / Elaborate (FOCUS 2022–2026)) | **15 marks** | Target ~16 pages

*Includes: mechanisms · doses/regimens · landmark clinical trials · hallmarks · **NCCN 2026 / 2027 updates** · flow charts · exam pearls*

0. How to use this DEEP answer

Sections below are **high-yield, trial-dense content** written for a 15-mark Paper IV elaboration. After the deep core, additional revision modules (algorithms, supportive care, flash cards) extend the note to full booklet length.

EXAM PEARLS / HALLMARKS

- In the exam hall, prioritise the DEEP trial tables + NCCN 2026 boxes first.
- Draw at least one flowchart from the deep sections.

1. Introduction & exam framing

HER2 (ERBB2) testing determines eligibility for anti-HER2 antibodies, tyrosine kinase inhibitors, and antibody–drug conjugates (ADCs). ASCO/CAP guidelines define IHC/ISH interpretation; DESTINY-Breast04/06 expanded therapy into HER2-low / ultralow disease. This essay focuses on testing plus therapeutic strategies for HER2-positive metastatic breast cancer (mBC), with NCCN Breast v9.2026 updates.

Structure: biology → ASCO/CAP methods → interpretation/HER2-low → metastatic sequencing (CLEOPATRA → DB-03 → HER2CLIMB) → doses → toxicities → NCCN 2026 (note imlunestrant is HR+/HER2– ESR1 pathway; still discuss HER2 ADCs) → pearls.

- Clarify HER2-positive (IHC3+ or ISH-amplified) vs HER2-low (IHC1+ or 2+/ISH–).
- Metastatic HER2+ standard: dual blockade + taxane 1L; T-DXd 2L; tucatinib triplet for CNS.
- Flag NCCN 2026 imlunestrant as HR+ ESR1-mutant — not a HER2-directed drug.

EXAM PEARLS / HALLMARKS

- Examiners love a full-page testing algorithm before drugs — do not rush to CLEOPATRA.

2. HER2 pathobiology & drug-target map

HER2 is a ligand-independent receptor tyrosine kinase that heterodimerises (notably with HER3) to activate PI3K–AKT and MAPK cascades → proliferation and survival. Amplification of ERBB2 drives protein overexpression. Trastuzumab binds subdomain IV (ADCC + signalling effects); pertuzumab binds dimerisation domain II; TKIs (lapatinib, neratinib, tucatinib) block the kinase; ADCs deliver payload after internalisation.

T-DXd (trastuzumab deruxtecan): cleavable linker + membrane-permeable topoisomerase-I payload → bystander effect, explaining activity in HER2-low. T-DM1: non-cleavable DM1 payload — less bystander, requires higher HER2 density.

HER2 amp / overexpression	Homodimers & HER2–HER3 → PI3K/MAPK
Trastuzumab	Subdomain IV; ADCC
Pertuzumab	Blocks HER2–HER3 dimerisation
T-DXd	Topo-I ADC; bystander kill; ILD risk
Tucatinib	HER2-selective TKI; CNS penetration
Resistance	PIK3CA, PTEN loss, incomplete blockade, CNS sanctuary

Figure: mechanism / pathway map

3. ASCO/CAP HER2 testing methods

Primary testing: validated IHC on invasive carcinoma. IHC 0 / 1+ = negative (historically); 2+ = equivocal → reflex ISH; 3+ = positive. ISH (dual-probe HER2/CEP17 preferred): report HER2 copy number and HER2/CEP17 ratio per ASCO/CAP cutoffs (classic positive: ratio ≥2.0 with HER2 copies ≥4; additional groups for monosomy/polysomy — quote groups if marks allow).

Preanalytics: cold ischaemia minimised; fixation 6–72 h in 10% neutral buffered formalin; avoid strong decalcification for bone mets when possible; use validated assays and external QA. Repeat testing on metastatic site recommended (discordance ~10%).

- Core biopsy vs excision discordance: retest if unexpected histology/grade or residual disease after NAT.
- Heterogeneous amplification: score invasive component carefully; may need additional blocks.
- NGS ERBB2 mutation ≠ IHC overexpression — neratinib pathways for activating mutations (NCCN).

Drug / regimen	Dose / schedule (exam pearls)
IHC 0	No staining or ≤10% incomplete faint — historically negative; note ultralow debates
IHC 1+	Incomplete faint >10% — HER2-low eligible if met criteria
IHC 2+	Weak–moderate complete membrane — reflex ISH
IHC 3+	Complete intense membrane >10% — HER2-positive
ISH positive	Per ASCO/CAP ratio/copy number groups — treat as HER2+

Table: doses and regimens

4. Interpretation workshop — HER2-low and ultralow

HER2-low = IHC 1+ or IHC 2+ / ISH-negative. DESTINY-Breast04 showed T-DXd superiority over physician's choice chemo in previously treated HER2-low mBC (HR+ and HR– cohorts). DESTINY-Breast06 moved T-DXd earlier in chemo-naive HR+ HER2-low/ultralow after endocrine therapy. Reporting IHC 0 vs ultralow has therapeutic implications — pathology–oncology communication is mandatory.

For classic HER2-positive metastatic disease, dual HER2 blockade strategies still require IHC3+ or ISH amplification — do not treat HER2-low with pertuzumab dual blockade as if HER2+.

ASCO/CAP testing → therapy binning




Figure: stepwise exam algorithm

EXAM PEARLS / HALLMARKS

- Say explicitly: HER2-low is not 'HER2-positive' — different drug algorithms.
- DB-04 mPFS ~10 vs 5 mo range (quote exact cohort numbers if memorised) — ADC expansion story.

5. Staging work-up before metastatic HER2 therapy

Confirm metastatic histology when feasible; restage with CT C/A/P ± bone scan/PET; brain MRI if symptoms or before regimens with CNS implications. Document ER/PR, HER2, prior neoadjuvant/adjuvant anti-HER2 exposure (trastuzumab, pertuzumab, T-DM1), anthracycline history, and baseline LVEF.

De novo vs recurrent metastatic HER2+ may influence residual endocrine options if HR+ (continue endocrine ± anti-HER2 in selected maintenance settings).

- Echo/MUGA baseline and during trastuzumab/pertuzumab/ADC therapy.
- Pregnancy test when relevant — anti-HER2 contraindicated in pregnancy.
- gBRCA rarely changes HER2+ 1L choice but matters for later PARPi if HER2− phenotypes evolve.

6. First-line HER2+ mBC — CLEOPATRA dual blockade

CLEOPATRA: docetaxel + trastuzumab + pertuzumab vs docetaxel + trastuzumab + placebo in 1L HER2+ mBC. Landmark OS improved to ~56–57 months with dual blockade; PFS benefit established new global standard. After chemo completion, continue H+P maintenance until progression/toxicity. Docetaxel typically ≥6 cycles if tolerated; some use paclitaxel backbones.

If HR+/HER2+, endocrine therapy may be combined or sequenced with anti-HER2 maintenance after chemo — individualise. CNS screening as clinically indicated.

Trial	Key result / exam takeaway
CLEOPATRA	H+P+docetaxel vs H+docetaxel: mOS ~57 vs ~40 mo — 1L standard
PERUSE / real-world	Taxane backbone flexibility with dual blockade

PHEREXA / others	Beyond 1L dual blockade lessons — sequencing context

Table: landmark clinical trials

Drug / regimen	Dose / schedule (exam pearls)
Trastuzumab	8 mg/kg IV load → 6 mg/kg Q3W (or SC fixed dose)
Pertuzumab	840 mg IV load → 420 mg Q3W with trastuzumab
Docetaxel	75 mg/m² IV Q3W (CLEOPATRA) × ≥6 cycles if tolerated

Table: doses and regimens

EXAM PEARLS / HALLMARKS

- Memorise OS ~57 months — the CLEOPATRA number every examiner knows.

7. Second-line and beyond — DESTINY-Breast03 (T-DXd)

DESTINY-Breast03 randomised T-DXd vs T-DM1 after trastuzumab ± pertuzumab in HER2+ mBC. mPFS 28.8 vs 6.8 months (stunning separation); OS benefit confirmed on updates. T-DXd is preferred 2L for most patients without contraindications. Monitor interstitial lung disease/pneumonitis — hold permanently for grade ≥2 ILD per label algorithms; counsel on dyspnoea/cough.

T-DM1 (EMILIA) remains an option if T-DXd unavailable/contraindicated or later lines. NCCN sequencing places T-DXd ahead of T-DM1 for eligible 2L patients.

Trial	Key result / exam takeaway
DESTINY-Breast03	T-DXd vs T-DM1: mPFS 28.8 vs 6.8 mo; new 2L standard
EMILIA	T-DM1 vs lapatinib+capecitabine — historical 2L PFS/OS benefit
DESTINY-Breast01/02	Heavily pretreated HER2+ activity; DB02 vs TPC
DESTINY-Breast04/06	HER2-low / ultralow — ADC expansion (mention even in HER2+ essay)

Table: landmark clinical trials

Drug / regimen	Dose / schedule (exam pearls)
T-DXd (breast)	5.4 mg/kg IV Q3W — ILD monitoring mandatory
T-DM1	3.6 mg/kg IV Q3W — thrombocytopenia/LFTs/cardiac

Table: doses and regimens

8. CNS disease — HER2CLIMB and radiotherapy integration

HER2CLIMB: tucatinib + trastuzumab + capecitabine vs placebo + trastuzumab + capecitabine in previously treated HER2+ mBC, including active brain metastases. Improved PFS/OS and intracranial outcomes — preferred systemic option when CNS disease is relevant. Local therapy (SRS/surgery/WBRT) remains essential for symptomatic lesions; coordinate sequencing.

T-DXd also has intracranial activity signals — choose by disease tempo, prior TKI/ADC exposure, and toxicity profile (ILD vs diarrhoea/hand-foot from capecitabine/tucatinib).

Metastatic HER2+ sequencing (exam flowchart)




Figure: stepwise exam algorithm

Trial	Key result / exam takeaway
HER2CLIMB	Tucatinib triplet: PFS/OS benefit; CNS activity practice-changing
NALA	Neratinib + capecitabine vs lapatinib + capecitabine — later-line option
TBCRC / ADC CNS data	T-DXd intracranial responses — evolving sequencing

Table: landmark clinical trials

Drug / regimen	Dose / schedule (exam pearls)
Tucatinib	300 mg PO BID with trastuzumab + capecitabine 1000 mg/m² BID D1–14 Q21D
Neratinib	240 mg PO daily (diarrhoea prophylaxis with loperamide mandatory)
Lapatinib	1250 mg daily with capecitabine in older regimens

Table: doses and regimens

9. HR+/HER2+ metastatic nuances & NCCN 2026 adjacent updates

HR+/HER2+ disease may use endocrine therapy with trastuzumab ± pertuzumab after chemo induction, or in frail patients. NCCN Breast v9.2026 lists abemaciclib + fulvestrant + trastuzumab as Category 2B in selected HR+/HER2+ unresectable/stage IV settings.

Important clarity for exams: imlunestrant is added in NCCN 2026 for ESR1-mutated HR+/HER2– metastatic disease (Cat 2A other recommended) — it is not a HER2-targeted therapy. Mention it under 'NCCN 2026 breast updates' while keeping HER2 ADC pathways (T-DXd, and sacituzumab in other subtypes) in the ADC discussion.

- Neratinib ± trastuzumab or fulvestrant Cat 2A useful in certain circumstances for HER2-activating mutations in stage IV.
- Erdafitinib Cat 2A useful in certain circumstances for FGFR alterations in stage IV — molecular housekeeping.
- Continue anti-HER2 across lines while changing partners — classic teaching.

10. Toxicities & monitoring table

Cardiac dysfunction: risk with trastuzumab/pertuzumab especially after anthracyclines — hold if LVEF declines per criteria; cardio-oncology input. T-DXd: ILD/pneumonitis, nausea, cytopenias, alopecia. T-DM1:

thrombocytopenia, transaminitis, cardiac. Tucatinib/capecitabine: diarrhoea, PPE, hepatotoxicity. Neratinib: early diarrhoea — mandatory prophylaxis.

- Educate on ILD symptoms before cycle 1 of T-DXd; low threshold for CT chest.
- Do not rechallenge T-DXd after symptomatic ILD grade ≥2.
- Antidiarrhoeal protocols for neratinib; dose escalate only if using labelled ramp regimens.

Safety monitoring loop




Figure: stepwise exam algorithm

11. Landmark trial summary table (copy into answer book)

Consolidate the must-quote trials with phase and take-home message.

Trial	Key result / exam takeaway
CLEOPATRA	1L H+P+docetaxel — mOS ~57 mo
EMILIA	T-DM1 beats lapatinib+cape post-trastuzumab
DESTINY-Breast03	T-DXd ■ T-DM1 — mPFS 28.8 vs 6.8 mo
HER2CLIMB	Tucatinib triplet — OS + CNS benefit
NALA	Neratinib+cape vs lapatinib+cape
DESTINY-Breast04	T-DXd in HER2-low — testing implications
DESTINY-Breast06	Earlier T-DXd in HR+ HER2-low/ultralow
KATHERINE (early)	Mention only if asked residual disease adjuvant contrast

Table: landmark clinical trials

12. NCCN Breast Cancer v9.2026 box

Place near the end. Explicitly separate HR+ endocrine updates from HER2-directed standards.

NCCN 2026 / 2027 UPDATES (include in answer)

• *Referenced versions:* Breast Cancer v9.2026

• Metastatic algorithms remain stratified by HR and HER2 status, including HER2-low/ultralow ADC pathways (T-DXd; sacituzumab govitecan in other indicated subtypes).

• Imlunestrant added (Cat 2A other recommended) for ESR1-mutated HR+/HER2– recurrent unresectable/stage IV disease — note: HR+ pathway, not HER2-targeted.

• Abemaciclib + fulvestrant + trastuzumab Cat 2B for selected HR+/HER2+ advanced disease.

• Neratinib ± trastuzumab or fulvestrant Cat 2A useful in certain circumstances for HER2-activating mutations in stage IV.

• Erdafitinib Cat 2A useful in certain circumstances for FGFR1/2/3 fusion/mutation in stage IV.

• NCCN Guidelines Navigator digital tool launched for breast cancer pathways.

EXAM PEARLS / HALLMARKS

• Write one sentence: 'Imlunestrant = ESR1-mutant HR+/HER2–; HER2+ mBC still sequenced CLEOPATRA → T-DXd → tucatinib.'

13. Practical sequencing vignette

Model case: 52-year-old, de novo ER+/HER2+ IHC3+ liver metastases, LVEF 62%. Start docetaxel+H+P (CLEOPATRA). After 6 cycles, maintain H+P ± endocrine. On progression (no ILD risk factors): T-DXd. Later isolated brain progression: add local SRS and consider tucatinib triplet if systemic options fit. Throughout: reassess goals, toxicities, and clinical trial eligibility.

- Oligoprogression: local therapy + continue systemic anti-HER2 when appropriate.
- Performance status decline: drop chemo, keep antibody/TKI if benefiting.
- Indian/NCG context: access to pertuzumab/T-DXd/tucatinib may gate sequence — state alternatives.

14. Summary checklist & exam pearls

Testing page + metastatic algorithm page + trial numbers page + NCCN box = complete answer.

- ASCO/CAP IHC → ISH; report HER2-low separately.
- 1L CLEOPATRA; 2L DB-03 T-DXd; CNS HER2CLIMB.
- Doses: H 8→6 mg/kg; P 840→420; T-DXd 5.4 mg/kg; tucatinib 300 mg BID.
- NCCN v9.2026: imlunestrant is HR+ ESR1; HER2 ADCs still central.

EXAM PEARLS / HALLMARKS

• IHC2+ without ISH is incomplete — always reflex.

• DB-03 28.8 vs 6.8 mo is the ADC number of the decade.

• ILD counselling is part of informed consent for T-DXd.

• HER2CLIMB for brain mets — say 'active CNS included'.

• Do not misclassify imlunestrant as anti-HER2 in the NCCN paragraph.

— Additional exam-length revision modules —

The following modules reinforce algorithms, toxicities, Indian/NCG context, and last-page summary structure.

3. Diagnosis, biomarkers & staging

A modern oncology answer must separate **diagnostic**, **prognostic**, and **predictive** markers. Predictive markers change drug choice (EGFR, HER2, BRCA, MSI, Claudin18.2, FGFR, NTRK).

- Tissue diagnosis with adequate IHC panel; never treat metastatic disease without molecular profiling when actionable targets exist.
- Use validated assays (FDA/companion diagnostics where mandated).
- Stage with AJCC 8th/9th as applicable (NSCLC NCCN 2026 uses AJCC 9th).
- Performance status, organ function, comorbidities, geriatric assessment, and goals of care before regimen selection.
- ER/PR/HER2 mandatory; HER2-low reporting now clinically actionable (DESTINY-Breast04/06).
- gBRCA testing for therapy (olaparib/talazoparib; OlympiA adjuvant).

• ESR1 mutation testing at endocrine resistance — imlunestrant listed in NCCN 2026 updates.

Work-up algorithm (draw this in the answer booklet)




Figure: stepwise exam algorithm

4. Management algorithm, doses & regimens

Structure management as: (A) localized/curative-intent pathways, (B) advanced/metastatic systemic therapy, (C) special populations (CNS, elderly, organ dysfunction, pregnancy), (D) supportive care.

For each line of therapy state: eligibility, preferred regimen, dose, key toxicity monitoring, and the trial that supports it.

• Localized disease: multimodality — surgery/RT/systemic as per disease chapter.

• Neoadjuvant vs adjuvant: cite pathologic response and adjuvant switch trials (e.g., KATHERINE, CheckMate 577, KEYNOTE-522).

• Metastatic: biomarker-first; avoid empiric chemo when a high-level targeted/IO option exists.

• Oligometastatic disease: consider metastasis-directed therapy where NCCN supports (e.g., prostate APMS 2026).

• Re-biopsy / liquid biopsy at progression to map resistance.

• Early HER2+: dual blockade neoadjuvant if tumor >2 cm or node+; adjuvant HP; T-DM1 if residual disease (KATHERINE).

• 1L mBC HER2+: taxane + trastuzumab + pertuzumab (CLEOPATRA).

• 2L: T-DXd preferred (DESTINY-Breast03) over T-DM1.

• HER2+ brain mets: tucatinib triplet (HER2CLIMB); SRS/WBRT as indicated.

Management flowchart (reproduce in exam)




Figure: stepwise exam algorithm

Drug / regimen	Dose / schedule (exam pearls)
Trastuzumab	8 mg/kg → 6 mg/kg IV Q3W (or SC fixed dose)
Pertuzumab	840 mg → 420 mg IV Q3W with trastuzumab
T-DXd	5.4 mg/kg IV Q3W (breast); 6.4 mg/kg gastric (per label) — monitor ILD
T-DM1	3.6 mg/kg IV Q3W
Olaparib	300 mg PO BID (tablets); adjust for toxicities/renal; OlympiA adjuvant 1 year

Table: doses and regimens

5. Landmark clinical trials (must quote in answer)

Present trials in a table: name → design/population → intervention → primary endpoint → absolute/relative benefit → practice change. Examiners award marks for correct trial names and one numeric result (HR or median PFS/OS).

- Mention phase (III preferred for standards) and whether endpoint was PFS, OS, MRD, pCR, or EFS.
- Distinguish statistical vs clinical significance; note crossover and subgroup caveats.
- For India-focused answers, mention access/biosimilars/NCG guidance after the trial table.

Trial	Key result / exam takeaway
HERA / BCIRG-006 / NSABP B-31 / N9831	Adjuvant trastuzumab OS/DFS benefit
CLEOPATRA	Docetaxel + trastuzumab + pertuzumab → OS ~57 mo 1L HER2+ mBC
EMILIA	T-DM1 vs lapatinib+capecitabine — PFS/OS benefit post-trastuzumab
DESTINY-Breast03	T-DXd vs T-DM1 — mPFS 28.8 vs 6.8 mo; new 2L standard
DESTINY-Breast04	T-DXd in HER2-low HR± — practice-changing ADC expansion
DESTINY-Breast06	T-DXd in HER2-ultralow / chemo-naive HR+ — further ADC move earlier
KATHERINE	Adjuvant T-DM1 vs trastuzumab if residual disease post-NAT
APHINITY	Adjuvant pertuzumab + trastuzumab — IDFS benefit especially node+
NeoSphere / TRYPHAENA / KRISTINE	Neoadjuvant dual HER2 blockade ± chemo
NALA / HER2CLIMB	Neratinib; tucatinib+trastuzumab+capecitabine (brain mets)

OlympiA	Adjuvant olaparib in gBRCA HER2– high-risk — iDFS/OS benefit

Table: landmark clinical trials

6. Toxicities, supportive care & monitoring

A complete answer links efficacy to safety. List monitoring schedules (CBC, chemistry, ECG, TFTs, cortisol, LFTs, PFTs/CT for ILD risk drugs).

- Anticipate class toxicities and counsel patients; use CTCAE grading.
- Cardio-oncology: anthracyclines, HER2 mAbs, ibrutinib, checkpoint myocarditis — baseline ECG/echo when indicated.
- IO toxicities: irAEs — hold drug, steroids, specialty referral; endocrine irAEs often permanent replacement.
- ADC pulmonary toxicity (T-DXd ILD): hold permanently for ≥grade 2 in many protocols; imaging + steroids.
- TLS prophylaxis for venetoclax/BLIN/CAR-T; rasburicase if high-risk.
- Fertility, sexuality, survivorship, and psychosocial support are part of comprehensive answers.

Toxicity management loop




Figure: stepwise exam algorithm

7. NCCN 2026 / 2027 updates (high-yield for ‘recent advances’)

Paper IV rewards currency. Explicitly write: **“As per NCCN Guidelines updates in 2026/2027…”** and list 3–5 bullets. Also mention ESMO guidelines / Indian NCG where relevant for balance.

- State Category (1 / 2A / 2B) and preferred vs other recommended vs useful in certain circumstances.
- If Indian exam: reconcile NCCN with CDSCO approvals and NCG practical pathways.

NCCN 2026 / 2027 UPDATES (include in answer)

- *Referenced versions:* Breast Cancer v9.2026
- NCCN Guidelines Navigator launched for breast cancer (digital navigation of guidelines).
- Imlunestrant added (Cat 2A other recommended) for ESR1-mutated HR+/HER2– recurrent unresectable/stage IV disease (1L or subsequent).
- Abemaciclib + fulvestrant + trastuzumab Cat 2B for HR+/HER2+ recurrent unresectable/stage IV.
- Neratinib ± trastuzumab or fulvestrant Cat 2A useful in certain circumstances for HER2-activating mutations in stage IV.
- Erdafitinib Cat 2A useful in certain circumstances for FGFR1/2/3 fusion or mutation in stage IV.

• Metastatic algorithms remain stratified by HR and HER2 status (including HER2-low / ultralow pathways with ADCs such as T-DXd, sacituzumab govitecan).

• Expanding ADC footprint across NCCN 2026: T-DXd (HER2 breast/gastric/NSCLC/urothelial), Dato-DXd (EGFR+ NSCLC), sacituzumab govitecan (breast/urothelial), mirvetuximab (ovarian FRα), enfortumab vedotin (urothelial), belantamab combinations (myeloma), loncastuximab (B-cell).

• Prostate: ADT + niraparib/abiraterone for BRCA2 APMS subsets.

• Ovarian: PARPi focused on maintenance/biomarker-defined; removed from many recurrent treatment sections.

• Breast: continue olaparib/talazoparib pathways per germline BRCA and OlympiA adjuvant context.

EXAM PEARLS / HALLMARKS

• NCCN versions change frequently — cite year/version (e.g., NSCLC v9.2026, Breast v9.2026, Myeloma v2.2027).

• Category 1 = high-level evidence + uniform NCCN consensus.

8. Special situations, differentials & practical issues

Intrinsic subtypes (luminal A/B, HER2-enriched, basal-like) from PAM50 approximate IHC surrogates but are not identical. TNBC biology includes DNA repair defects, AR-positive luminal androgen receptor subset, and immune-rich tumors benefiting from CPI (KEYNOTE-522; KEYNOTE-355 CPS≥10).

• Neoadjuvant pCR is a strong surrogate in HER2+ and TNBC for escalation/de-escalation.

• Antibody-drug conjugates reshaped sequencing: T-DXd before many older anti-HER2 TKIs in metastatic HER2+.

• Pregnancy-associated breast cancer: tailor imaging (no gadolinium/PET routine), surgery, anthracycline after first trimester; anti-HER2 contraindicated in pregnancy.

9. Evidence hierarchy, patient rights & benefits

Quote evidence using: Phase III RCT > meta-analysis of RCTs > prospective registries / RWE > retrospective > expert consensus. NCCN Categories and ESMO-MCBS scores help communicate strength of benefit.

Patient-centered care: informed consent for toxicities (especially CRS/ICANS, ILD, infertility), shared decision-making, financial toxicity counseling, and access to supportive/palliative care concurrent with active therapy.

• Benefits to highlight: OS, PFS, pCR, MRD negativity, organ preservation, QoL, reduced hospitalization.

• Rights: autonomy, confidentiality (EMR), right to second opinion, trial participation, refusal of care, end-of-life preferences.

• Equity: molecular testing access, biosimilars, NCG/Ayushman Bharat practical pathways for Indian exams.

• Pharmacovigilance: report ADRs; Phase IV / post-marketing commitments.

EXAM PEARLS / HALLMARKS

• Write one sentence on absolute benefit (e.g., ‘median OS improved from X to Y months’).)

• Mention number-needed-to-treat conceptually if space permits.

10. Summary checklist (write this last page in the booklet)

Close the 15-mark answer with a crisp numbered summary the examiner can tick.

• Definition + epidemiology one-liner

• Key biology / hallmark diagram

• Essential diagnostics & biomarkers

• Risk/stage stratification

- Preferred treatment algorithm with ≥2 doses
- ≥3 named trials with endpoints
- Major toxicities + monitoring
- NCCN 2026 update bullets (3–5)
- Special situations
- Five exam pearls

30-second revision flowchart




Figure: stepwise exam algorithm

EXAM PEARLS / HALLMARKS

- Neat flowchart + trial table often separates a 10/15 from a 13–15/15.
- Do not dump drug lists without doses or evidence.
- Currency: say 'NCCN 2026' explicitly for Paper IV.

11. Pharmacology deep-dive (class, PK/PD, interactions)

For drug-centered questions, present: chemical/biologic class → target epitope/kinase → PK (oral bioavailability, half-life, hepatic/renal clearance) → food effects → CYP interactions → dose adjustments → therapeutic drug monitoring if any.

Example interaction pearls: acid-suppressive therapy with some TKIs; QT-prolonging combinations; azole antifungals with vinca/TKIs; strong CYP3A inducers reducing exposure to many oral targeted agents.

- State renal/hepatic dose modifications when known.
- Note formulation differences (olaparib tablets vs capsules; IV vs SC daratumumab).
- Infusion reactions: premedication protocols for mAbs/BiTEs.

Drug / regimen	Dose / schedule (exam pearls)
Trastuzumab	8 mg/kg → 6 mg/kg IV Q3W (or SC fixed dose)
Pertuzumab	840 mg → 420 mg IV Q3W with trastuzumab
T-DXd	5.4 mg/kg IV Q3W (breast); 6.4 mg/kg gastric (per label) — monitor ILD
T-DM1	3.6 mg/kg IV Q3W
Olaparib	300 mg PO BID (tablets); adjust for toxicities/renal; OlympiA adjuvant 1 year

Table: doses and regimens

12. Response assessment & follow-up schedules

Use disease-specific criteria: RECIST 1.1 solid tumors; iRECIST for immunotherapy; Lugano/Deauville for lymphoma; IMWG for myeloma; ELN molecular response for CML/AML; PSA working group for prostate.

- Typical imaging interval q2–3 cycles in metastatic disease unless clinically indicated sooner.
- PET-adapted strategies in Hodgkin lymphoma (RATHL, AHL2011) — cite Deauville 1–3 negative vs 4–5 positive.
- ctDNA MRD emerging for escalation/de-escalation (CRC, lymphoma, breast — exam discussion point).

Clinical algorithm




Figure: stepwise exam algorithm

13. Comparison tables useful for exams

Draw 2×N tables comparing old standard vs new standard; or drug A vs B (efficacy, toxicity, CNS activity, cost/access).

- Column ideas: target, dose, PFS, OS, grade ≥3 AEs, CNS activity, NCCN category.
- Another table: resistance mutation → preferred next drug.

Trial	Key result / exam takeaway
HERA / BCIRG-006 / NSABP B-31 / N9831	Adjuvant trastuzumab OS/DFS benefit
CLEOPATRA	Docetaxel + trastuzumab + pertuzumab → OS ~57 mo 1L HER2+ mBC
EMILIA	T-DM1 vs lapatinib+capecitabine — PFS/OS benefit post-trastuzumab
DESTINY-Breast03	T-DXd vs T-DM1 — mPFS 28.8 vs 6.8 mo; new 2L standard
DESTINY-Breast04	T-DXd in HER2-low HR± — practice-changing ADC expansion
DESTINY-Breast06	T-DXd in HER2-ultralow / chemo-naive HR+ — further ADC move earlier

Table: landmark clinical trials

14. Indian context, NCG & practical exam tips

National Cancer Grid (NCG) resource-stratified guidelines are frequently expected in Indian DM exams. Discuss biosimilars (trastuzumab, rituximab), generic TKIs, and center-based cell therapy availability.

- Document MDT decisions; clinical trial referral when standard options exhausted.
- Counsel financial toxicity transparently; prioritize highest-value Category 1 options.
- Infection prophylaxis algorithms crucial in hematologic topics (antifungal, antiviral, PJP).

15. Sample ‘examiner hot spots’ & model closing paragraph

Model closing: “In summary, management of this problem is biomarker-driven. Current standards are supported by phase III trials and reflected in NCCN 2026/2027 updates. Sequencing should maximize OS and QoL while mitigating class-specific toxicities. Enrollment in clinical trials and adherence to MDT pathways remain integral.”

Hot spots: exact doses, one HR/median, resistance mechanism, NCCN category, and a drawn algorithm.

EXAM PEARLS / HALLMARKS

- If stuck on memory: write mechanism → class → one trial → one dose → one toxicity → NCCN 2026 line.
- Never contradict yourself between algorithm and trial table.

A. Epidemiology & burden (write ½–1 page)

Begin with global and Indian epidemiology when known (GLOBOCAN / ICMR / NCRP). State age distribution, sex ratio, and risk factors (tobacco, infection-related cancers such as HPV/HBV/HCV/H. pylori/HIV, obesity, inherited syndromes).

Explain why incidence or mortality trends matter for screening and resource allocation. For rare diseases, state incidence per 100,000 and referral-center patterns.

Link epidemiology to prevention: vaccination (HPV, HBV), tobacco control, hepatitis treatment, aspirin in selected molecular CRC, and lifestyle (exercise post-adjuvant CRC per NCCN 2026).

- One sentence on screening relevance (mammography, LDCT, FIT/colonoscopy, low-dose CT criteria).
- Mention hereditary fraction if applicable (Lynch, HBOC, FAP, MEN, VHL, HPRCC).
- State stage distribution at diagnosis in LMICs vs HIC if relevant to treatment sequencing.

B. Historical evolution → modern standard (timeline)

Examiners like a short timeline: historical surgery/RT/chemo era → targeted therapy era → immunotherapy/ADC/cellular therapy era.

Highlight inflection trials that changed guidelines (e.g., imatinib IRIS; trastuzumab HERA; checkpoint inhibitors; ADCs; CAR-T approvals).

End timeline with **NCCN 2026/2027** preferred regimens so the answer feels contemporary for Paper IV.

Therapeutic eras timeline




Figure: stepwise exam algorithm

C. Detailed differential diagnosis & pathology pearls

List 5–7 differentials and the IHC/molecular discriminators. Pathology reports should include grade, LVI, margins, nodes, and required biomarkers.

For hematologic topics: morphology → flow cytometry → cytogenetics → NGS panel → integrated WHO2022/ICC diagnosis.

For CUP/unknown primary style questions: IHC algorithm (CK7/CK20, TTF1, CDX2, GATA3, PAX8, synaptophysin) plus NGS and MSI/HER2.

- Always write ‘correlate clinically with radiology and tumor markers’.
- State pre-analytical caveats (decalcification, fixation time) affecting HER2/ER assays.
- Liquid biopsy complements but does not always replace tissue.

Morphology	Light microscopy pattern
IHC panel	Lineage + predictive markers
Cytogenetics	Karyotype / FISH fusions
NGS	SNVs, indels, CNAs, MSI, TMB
Integrated Dx	WHO/ICC + AJCC stage

Figure: mechanism / pathway map

D. Multidisciplinary care pathway & sequencing

Describe roles: medical oncology, surgical oncology, radiation oncology, radiology, pathology, palliative care, genetics, psycho-oncology, rehabilitation, and clinical trials unit.

Sequencing principles: treat drivers first when OS-proven; use neoadjuvant therapy to downstage and guide adjuvant escalation; avoid overlapping severe toxicities; protect future options (e.g., cumulative cardiotoxicity, neuropathy).

For oligometastatic disease discuss local therapy (surgery/SBRT/ablation) plus systemic therapy — prostate NCCN 2026 explicitly incorporates metastasis-directed therapy in selected APMS disease.

- Document MDT minutes in the file — also good exam phrase.
- Re-staging after neoadjuvant before definitive local therapy.
- Early palliative care improves QoL and sometimes survival in metastatic settings.

Clinical algorithm




Figure: stepwise exam algorithm

E. Breast oncology deep notes (HER2 / subtypes)

ASCO/CAP HER2 testing: IHC 0, 1+, 2+ (equivocal→ISH), 3+. HER2-low (1+ or 2+/ISH–) and ultralow categories matter after DESTINY-Breast04/06.

Early disease: prefer dual HER2 blockade neoadjuvant for ≥T2 or node-positive; residual disease → adjuvant T-DM1 (KATHERINE). Node-positive adjuvant benefit from pertuzumab (APHINITY).

Metastatic: CLEOPATRA 1L; DESTINY-Breast03 establishes T-DXd over T-DM1 in 2L; HER2CLIMB for brain metastases; continue anti-HER2 beyond progression with changing partners.

Endocrine + anti-HER2 strategies exist for HR+/HER2+; NCCN 2026 adds nuances (e.g., abemaciclib+fulvestrant+trastuzumab Cat 2B in selected HR+/HER2+ advanced disease).

TNBC: KEYNOTE-522 perioperative pembrolizumab; metastatic CPS≥10 pembrolizumab+chemo; sacituzumab govitecan (ASCENT); PARPi if gBRCA; ADCs evolving.

- Cardiotoxicity monitoring with HER2 agents + anthracyclines sequencing.
- Pregnancy: avoid anti-HER2 until after delivery.
- NCCN 2026: imlunestrant for ESR1-mutant HR+/HER2–; neratinib options for HER2-mutant MBC; erdafitinib for FGFR alterations.

Trial	Key result / exam takeaway
CLEOPATRA	H+P+docetaxel OS landmark 1L HER2+ mBC
DESTINY-Breast03	T-DXd ■ T-DM1 2L
KATHERINE	Adjuvant T-DM1 if residual disease
HER2CLIMB	Tucatinib triplet; CNS activity

DESTINY-Breast04/06	HER2-low / ultralow ADC expansion
OlympiA	Adjuvant olaparib gBRCA HER2–

Table: landmark clinical trials

F. HER2 testing interpretation workshop

Spend almost a full page on testing — common 15-mark theme (Jan 2024 paper).

Preanalytics: cold ischemia, fixation 6–72 h in 10% NBF; avoid decalcified bone for HER2 if possible.

Discordance between core and excision: retest if heterogenous, grade/histology unexpected, or residual disease after NAT.

ISH methods: dual-probe HER2/CEP17; ratios and copy number cutoffs per ASCO/CAP updates; polysomy interpretation pitfalls.

Clinical algorithm




Figure: stepwise exam algorithm

G. Supportive care mini-chapter

Antiemetics (olanzapine-containing regimens for HEC), G-CSF primary/secondary prophylaxis, antimicrobial prophylaxis, mucositis care, nutrition (including TPN indications/contraindications), pain ladder + interventional techniques, thrombosis prevention (Khorana), growth factors, transfusion thresholds.

Psychosocial distress screening, fertility preservation referral before gonadotoxic therapy, and survivorship clinics.

- Febrile neutropenia: risk assess; anti-pseudomonal beta-lactam; add antifungal if persistent.
- Checkpoint diarrhea: rule out infection then steroids promptly.
- ADC ILD algorithms — patient education on dyspnea.

H. Model answer skeleton (time plan for 35 minutes)

Minutes 0–3: definition + epidemiology + diagram outline. Minutes 3–12: biology + diagnosis. Minutes 12–25: management algorithm with doses/trials. Minutes 25–30: toxicities + NCCN 2026. Minutes 30–35: summary

pearls + underline Category 1 regimens.

- Leave a half page blank early for a neat flowchart drawn once.
- Underline drug doses.
- Box the NCCN 2026 bullets.

EXAM PEARLS / HALLMARKS

- Currency + clarity > exhaustive rare regimens.
- Named trial + number beats unnamed claim.
- NCCN 2026 phrase is expected in Paper IV recent advances.

K. Pitfalls, contraindications & ‘do not miss’ warnings

List absolute/relative contraindications of key drugs (e.g., baseline ILD caution for T-DXd; severe autoimmune disease caution for CPI; T315I needs ponatinib/asciminib not 1G/2G TKI alone; ATRA differentiation syndrome in APL; TLS with venetoclax/Blincyto; HBV reactivation with anti-CD20/CPI).

Drug–drug interactions and QT prolongation traps are classic viva points — mention checking concomitant meds (azoles, macrolides, acid suppressants with selected TKIs).

Imaging pitfalls: pseudo-progression on IO; flare on bone scan; PET Deauville interpretation after immunomodulatory therapy.

- Always screen hepatitis B before anti-CD20, anthracyclines, and many systemic regimens.
- Pregnancy test before teratogenic agents (IMiDs — REMS; MTX; TKI counseling).
- Do not combine overlapping QT-prolonging agents without ECG plan.
- Document performance status — unfit vs fit changes entire AML/lymphoma pathway.

Clinical algorithm




Figure: stepwise exam algorithm

L. Case vignette practice (apply the algorithm)

Invent a short case in the answer: age, stage, biomarkers, comorbidities → your exact regimen with dose → expected monitoring → what you will do at progression. This shows applied knowledge and often gains examiner goodwill.

Example template: “45-year-old, ECOG 1, biomarker X positive, no contraindication Y → start regimen Z at dose… per trial T / NCCN 2026 preferred. Assess response at interval N. At progression, re-profile for resistance mutation R and switch to agent S.”

Add one geriatric or organ-dysfunction variant to show nuance (dose reduce, alternate preferred regimen, or best supportive care).

- Vignette 1: fit, biomarker-positive — preferred Cat 1 therapy.
- Vignette 2: comorbid/unfit — alternative regimen.
- Vignette 3: progression with resistance mutation — sequenced next line.
- Vignette 4: severe toxicity — hold, treat, rechallenge decision.

M. One-page ‘wall chart’ summary (draw/box this)

Reserve the penultimate page for a wall-chart style summary with four quadrants: (1) Biology (2) Work-up (3) Treatment lines (4) NCCN 2026 bullets. This mirrors how many toppers present long answers and helps hit the page target with high signal density.

Quadrant 1	Hallmark gene / pathway / figure
Quadrant 2	Must-order tests + staging
Quadrant 3	1L / 2L / 3L with doses
Quadrant 4	Trials + NCCN 2026 + toxicities
Center	MDT + patient goals

Figure: mechanism / pathway map

EXAM PEARLS / HALLMARKS

- Wall chart + flowchart + trial table = visual completeness.
- Write ‘NCCN 2026’ in quadrant 4 explicitly.

I2. Full-length essay addendum — tables to copy into the answer book

For 15-mark elaborations, add two hand-drawn tables: (Table A) Drug–dose–pearl; (Table B) Trial–population–endpoint–practice change. Then write a short paragraph on quality-of-life and patient-reported outcomes, which modern trials increasingly report.

Discuss biomarker evolution on therapy (liquid biopsy cadence), and when to stop therapy (duration of CPI, TFR in CML, maintenance stopping rules).

Include a brief note on second malignancies and late effects when curative-intent therapy is discussed (alkylators, RT fields, PARPi MDS/AML risk, CHL survivors).

- Table A columns: Drug | Dose | Key toxicity | Pearl
- Table B columns: Trial | Arm | Primary endpoint | Result | Change in practice
- One paragraph on shared decision-making and financial counseling

Trial	Key result / exam takeaway
HERA / BCIRG-006 / NSABP B-31 / N9831	Adjuvant trastuzumab OS/DFS benefit
CLEOPATRA	Docetaxel + trastuzumab + pertuzumab → OS ~57 mo 1L HER2+ mBC
EMILIA	T-DM1 vs lapatinib+capecitabine — PFS/OS benefit post-trastuzumab

DESTINY-Breast03	T-DXd vs T-DM1 — mPFS 28.8 vs 6.8 mo; new 2L standard
DESTINY-Breast04	T-DXd in HER2-low HR± — practice-changing ADC expansion
DESTINY-Breast06	T-DXd in HER2-ultralow / chemo-naive HR+ — further ADC move earlier
KATHERINE	Adjuvant T-DM1 vs trastuzumab if residual disease post-NAT
APHINITY	Adjuvant pertuzumab + trastuzumab — IDFS benefit especially node+
NeoSphere / TRYPHAENA / KRISTINE	Neoadjuvant dual HER2 blockade ± chemo
NALA / HER2CLIMB	Neratinib; tucatinib+trastuzumab+capecitabine (brain mets)
OlympiA	Adjuvant olaparib in gBRCA HER2– high-risk — iDFS/OS benefit

Table: landmark clinical trials

Drug / regimen	Dose / schedule (exam pearls)
Trastuzumab	8 mg/kg → 6 mg/kg IV Q3W (or SC fixed dose)
Pertuzumab	840 mg → 420 mg IV Q3W with trastuzumab
T-DXd	5.4 mg/kg IV Q3W (breast); 6.4 mg/kg gastric (per label) — monitor ILD
T-DM1	3.6 mg/kg IV Q3W
Olaparib	300 mg PO BID (tablets); adjust for toxicities/renal; OlympiA adjuvant 1 year

Table: doses and regimens

NCCN 2026 / 2027 UPDATES (include in answer)

• *Referenced versions:* Breast Cancer v9.2026

• NCCN Guidelines Navigator launched for breast cancer (digital navigation of guidelines).

• Imlunestrant added (Cat 2A other recommended) for ESR1-mutated HR+/HER2– recurrent unresectable/stage IV disease (1L or subsequent).

• Abemaciclib + fulvestrant + trastuzumab Cat 2B for HR+/HER2+ recurrent unresectable/stage IV.

• Neratinib ± trastuzumab or fulvestrant Cat 2A useful in certain circumstances for HER2-activating mutations in stage IV.

• Erdafitinib Cat 2A useful in certain circumstances for FGFR1/2/3 fusion or mutation in stage IV.

• Metastatic algorithms remain stratified by HR and HER2 status (including HER2-low / ultralow pathways with ADCs such as T-DXd, sacituzumab govitecan).

• Expanding ADC footprint across NCCN 2026: T-DXd (HER2 breast/gastric/NSCLC/urothelial), Dato-DXd (EGFR+ NSCLC), sacituzumab govitecan (breast/urothelial), mirvetuximab (ovarian FRα), enfortumab vedotin (urothelial), belantamab combinations (myeloma), loncastuximab (B-cell).

• Prostate: ADT + niraparib/abiraterone for BRCA2 APMS subsets.

• Ovarian: PARPi focused on maintenance/biomarker-defined; removed from many recurrent treatment sections.

• Breast: continue olaparib/talazoparib pathways per germline BRCA and OlympiA adjuvant context.

I. Extended notes — resistance, sequencing & future directions

Resistance biology should occupy a dedicated half-page: on-target secondary mutations, bypass signaling, lineage plasticity (e.g., EGFR NSCLC → SCLC), antigen loss after cellular therapy, and immunosuppressive TME changes after CPI.

Sequencing principles: use the agent with largest OS benefit early when toxicities allow; reserve cross-resistant drugs; consider cumulative neuropathy/cardiotoxicity; plan CNS-active agents if brain metastases are present or likely.

Future directions appropriate for 2026 answers: tumor-agnostic indications (NTRK, RET, BRAF, MSI-H, TMB-H, HER2-mut), radioligand therapy, CAR-NK/off-the-shelf products, ctDNA-guided escalation/de-escalation, AI decision-support, and microbiome modulation adjuncts.

Health services angle: real-world evidence validating trial results in older/comorbid patients; digital remote toxicity monitoring; and equity of molecular testing access — themes also reflected in recent Paper IV questions.

- Write one ‘if progression then…’ resistance table.
- Mention at least one tumor-agnostic approval if biology overlaps.
- Close with trial enrollment as a standard option.

J. Quick-revision flash cards (last 1–2 pages)

Convert the topic into 8–10 flash cards mentally: (1) definition (2) hallmark gene (3) key test (4) 1L preferred (5) dose (6) trial (7) top toxicity (8) NCCN 2026 update (9) Indian/NCG pearl (10) one differential.

- Flash 1: One-line definition
- Flash 2: Driver / pathway
- Flash 3: Mandatory biomarker
- Flash 4: NCCN preferred 1L
- Flash 5: Exact dose
- Flash 6: Landmark phase III + endpoint
- Flash 7: Grade ≥3 toxicity to warn
- Flash 8: 2026 guideline change
- Flash 9: Resistance next step
- Flash 10: Supportive care must-do

EXAM PEARLS / HALLMARKS

- If time-poor in the hall: draw algorithm + trial table + NCCN box — captures most marks.
- Keep handwriting large for doses and HRs.

Final box — paste into last page of answer sheet

Re-list NCCN 2026 updates here so the examiner sees currency even if earlier pages are rushed.

NCCN 2026 / 2027 UPDATES (include in answer)

- *Referenced versions:* Breast Cancer v9.2026
- NCCN Guidelines Navigator launched for breast cancer (digital navigation of guidelines).
- Imlunestrant added (Cat 2A other recommended) for ESR1-mutated HR+/HER2– recurrent unresectable/stage IV disease (1L or subsequent).
- Abemaciclib + fulvestrant + trastuzumab Cat 2B for HR+/HER2+ recurrent unresectable/stage IV.
- Neratinib ± trastuzumab or fulvestrant Cat 2A useful in certain circumstances for HER2-activating mutations in stage IV.
- Erdafitinib Cat 2A useful in certain circumstances for FGFR1/2/3 fusion or mutation in stage IV.
- Metastatic algorithms remain stratified by HR and HER2 status (including HER2-low / ultralow pathways with ADCs such as T-DXd, sacituzumab govitecan).
- Expanding ADC footprint across NCCN 2026: T-DXd (HER2 breast/gastric/NSCLC/urothelial), Dato-DXd (EGFR+ NSCLC), sacituzumab govitecan (breast/urothelial), mirvetuximab (ovarian FRα), enfortumab vedotin (urothelial), belantamab combinations (myeloma), loncastuximab (B-cell).
- Prostate: ADT + niraparib/abiraterone for BRCA2 APMS subsets.
- Ovarian: PARPi focused on maintenance/biomarker-defined; removed from many recurrent treatment sections.
- Breast: continue olaparib/talazoparib pathways per germline BRCA and OlympiA adjuvant context.

Final NCCN 2026 box

Copy these bullets onto the last page of your answer booklet for Paper IV currency marks.

NCCN 2026 / 2027 UPDATES (include in answer)

• *Referenced versions:* Breast Cancer v9.2026

• NCCN Guidelines Navigator launched for breast cancer (digital navigation of guidelines).

• Imlunestrant added (Cat 2A other recommended) for ESR1-mutated HR+/HER2– recurrent unresectable/stage IV disease (1L or subsequent).

• Abemaciclib + fulvestrant + trastuzumab Cat 2B for HR+/HER2+ recurrent unresectable/stage IV.

• Neratinib ± trastuzumab or fulvestrant Cat 2A useful in certain circumstances for HER2-activating mutations in stage IV.

• Erdafitinib Cat 2A useful in certain circumstances for FGFR1/2/3 fusion or mutation in stage IV.

• Metastatic algorithms remain stratified by HR and HER2 status (including HER2-low / ultralow pathways with ADCs such as T-DXd, sacituzumab govitecan).

• Expanding ADC footprint across NCCN 2026: T-DXd (HER2 breast/gastric/NSCLC/urothelial), Dato-DXd (EGFR+ NSCLC), sacituzumab govitecan (breast/urothelial), mirvetuximab (ovarian FRα), enfortumab vedotin (urothelial), belantamab combinations (myeloma), loncastuximab (B-cell).

• Prostate: ADT + niraparib/abiraterone for BRCA2 APMS subsets.

• Ovarian: PARPi focused on maintenance/biomarker-defined; removed from many recurrent treatment sections.

• Breast: continue olaparib/talazoparib pathways per germline BRCA and OlympiA adjuvant context.

Disclaimer: Compiled for DM exam revision from published trials and NCCN 2026/2027 public summaries (ASCO Post, OncLive, JNCCN selections, NCCN category list). Always verify the current full NCCN PDF, FDA/CDSCO labels, and institutional protocols before patient care. Doses require BSA/organ-function adjustment.