D.M. MEDICAL ONCOLOGY — PAPER IV

RECENT ADVANCES · EXAM ANSWER NOTES

Discuss the role of genes involved in mismatch repair (MMR) mechanisms in various cancers. Discuss testing modalities of MMR genes and their interpretation. Describe the mechanism of MMR as a biomarker for checkpoint inhibitors.

February 2023 | DM 0223 | Q1 (DEEP Essay / Elaborate (FOCUS 2022–2026)) | **15 marks** | Target ~16 pages

*Includes: mechanisms · doses/regimens · landmark clinical trials · hallmarks · **NCCN 2026 / 2027 updates** · flow charts · exam pearls*

0. How to use this DEEP answer

Sections below are **high-yield, trial-dense content** written for a 15-mark Paper IV elaboration. After the deep core, additional revision modules (algorithms, supportive care, flash cards) extend the note to full booklet length.

EXAM PEARLS / HALLMARKS

- In the exam hall, prioritise the DEEP trial tables + NCCN 2026 boxes first.
- Draw at least one flowchart from the deep sections.

1. Introduction & exam framing

Mismatch repair (MMR) is a post-replication DNA repair system that corrects base–base mismatches and insertion–deletion loops arising mainly in microsatellite sequences. Loss of MMR → microsatellite instability–high (MSI-H) / deficient MMR (dMMR) phenotype → hypermutation, high neoantigen load, and exceptional sensitivity to PD-1/PD-L1 checkpoint inhibitors (CPIs) across tumour types (tissue-agnostic approvals).

For a 15-mark answer structure: (1) genes and pathway, (2) cancers and Lynch syndrome, (3) testing modalities with interpretation pitfalls, (4) mechanistic link to CPI response, (5) landmark trials with numbers, (6) NCCN 2026 CRC pathways (ATOMIC; neoadjuvant rectal CPI), (7) exam pearls and a drawn algorithm.

- State predictive vs prognostic: dMMR/MSI-H is both prognostic (better stage-for-stage CRC prognosis historically) and strongly predictive of CPI benefit.
- Always contrast germline Lynch vs sporadic (often MLH1 promoter hypermethylation ± BRAF V600E).
- Quote KEYNOTE-177, CheckMate-8HW, KEYNOTE-158, and ATOMIC with primary endpoints.

EXAM PEARLS / HALLMARKS

- Open with a one-line definition of MMR → MSI-H → neoantigens → CPI — examiners mark this triad.
- Draw IHC 4-protein panel + reflex pathway once; it scores heavily.

2. MMR genes and molecular pathway

Core MutSα heterodimer = MSH2–MSH6 (recognises single-base mismatches and 1–2 bp loops). MutSβ = MSH2–MSH3 (larger insertion–deletion loops). MutLα = MLH1–PMS2 (endonuclease that recruits excision/resynthesis machinery). EPCAM deletions silence MSH2 by transcriptional read-through and are a recognised Lynch mechanism without MSH2 coding mutation.

Biallelic inactivation (germline + second hit, or two somatic hits, or epigenetic silencing) abolishes MMR → MSI. Constitutional mismatch repair deficiency (CMMRD) = biallelic germline MMR mutations → childhood cancers (haematologic, brain, GI).

- Gene → protein pairs to memorise: MLH1↔PMS2 (co-loss common); MSH2↔MSH6.
- Isolated MSH6 or PMS2 loss can be subtle on IHC — do not dismiss without PCR/NGS if clinical suspicion high.
- POLE/POLD1 exonuclease mutations cause ultramutation (MSS often) but also CPI-responsive — mention as differential hypermutators.

Replication error	Polymerase slip at microsatellites → mismatch / indel loop
Recognition	MutSα (MSH2–MSH6) or MutSβ (MSH2–MSH3) binds lesion
Incision	MutLα (MLH1–PMS2) endonuclease nick
Excision / resynthesis	EXOI, DNA pol δ, ligase restore correct sequence
Failure (dMMR)	Frameshift peptides → neoantigens; MSI-H genomic signature

Figure: mechanism / pathway map

3. Role of MMR genes across cancers

Lynch syndrome (LS) spectrum: colorectal (~lifetime risk 40–80% depending on gene), endometrial (especially MSH6/MLH1), gastric, ovarian, urothelial (upper tract classic for MSH2), small bowel, sebaceous tumours (Muir–Torre), brain (Turcot). Gene-specific penetrance matters: MSH6/PMS2 often later onset / endometrial-skewed; MLH1/MSH2 higher CRC risk.

Sporadic MSI-H CRC (~12–15% of CRC): typically older patients, right-sided, BRAF V600E+, MLH1 promoter CpG island hypermethylation — not Lynch. Gastric, endometrial, and small subsets of pancreatic, biliary, prostate, and sarcoma also harbour dMMR/MSI-H. Pan-tumour prevalence ~2–4% but enrichment in CRC/endometrial/gastric.

- Stage II CRC: MSI-H historically associated with good prognosis and limited fluoropyrimidine-alone benefit — classic exam line.
- Metastatic MSI-H CRC: CPI preferred over chemo first-line (KEYNOTE-177 / CheckMate-8HW).
- dMMR rectal cancer: neoadjuvant CPI can enable organ preservation (NCCN 2026 primary consideration).

Trial	Key result / exam takeaway
KEYNOTE-177	1L pembrolizumab vs chemo MSI-H mCRC: mPFS 16.5 vs 8.2 mo; durable OS; chemo crossover
CheckMate-8HW	Nivo+ipi vs chemo (and vs nivo) MSI-H/dMMR mCRC — dual CPI PFS superiority
KEYNOTE-158	Pembro tissue-agnostic MSI-H/dMMR solid tumours — ORR ~34%; basis of agnostic label
ATOMIC (NCCN 2026)	Stage III dMMR colon: FOLFOX/CAPOX + atezolizumab preferred adjuvant pathway

Table: landmark clinical trials

4. Lynch vs sporadic — clinical and molecular distinction

Universal tumour testing (IHC ± MSI) is standard for CRC and endometrial cancer in most guidelines. If MLH1/PMS2 loss: reflex BRAF V600E and/or MLH1 methylation. BRAF V600E+ or methylated MLH1 → sporadic pathway (usually stop germline testing unless strong pedigree). Intact BRAF and unmethylated MLH1 with IHC loss → refer genetics for germline MMR panel (± EPCAM).

Amsterdam/Bethesda criteria are historical triage tools; universal testing catches more LS. Cascade testing of relatives after pathogenic germline variant is the prevention payoff.

Lynch triage after tumour MMR testing (draw this)




Figure: stepwise exam algorithm

EXAM PEARLS / HALLMARKS

- Never call 'Lynch' on IHC alone — IHC shows phenotype; germline confirms syndrome.
- EPCAM deletion → MSH2 protein loss on IHC; include EPCAM in LS panels.

5. Testing modalities — IHC for MMR proteins

Four-antibody IHC (MLH1, MSH2, MSH6, PMS2) on formalin-fixed paraffin-embedded tumour is the workhorse. Loss of nuclear staining in tumour with retained internal control (stroma/lymphocytes) defines dMMR. Paired loss patterns: MLH1+PMS2; MSH2+MSH6; isolated MSH6 or PMS2 possible.

Preanalytics: adequate fixation (typically 6–72 h, 10% NBF), avoid over-decalcification, report heterogeneity. Neoadjuvant therapy can weaken staining — interpret cautiously; correlate MSI.

- IHC = protein expression surrogate; does not directly prove MSI but correlates strongly.
- Indeterminate / focal loss → reflex MSI PCR or NGS MSI / MMR sequencing.
- Post-treatment rectal specimens: prefer pretreatment biopsy for MMR if possible.

MLH1 loss	Usually ± PMS2 loss; check methylation/BRAF
MSH2 loss	Usually ± MSH6 loss; think LS / EPCAM
Isolated MSH6	LS possible; MSI may be low/subtle
Isolated PMS2	LS possible; confirm with genetics
All four retained	pMMR phenotype — still consider POLE if ultramutated

Figure: mechanism / pathway map

6. Testing modalities — MSI PCR and NGS

PCR MSI assays compare tumour vs normal at mononucleotide/dinucleotide loci (e.g., Bethesda/NCI panel or pentaplex mononucleotide panels). MSI-H typically ≥2/5 (or ≥30%) unstable markers; MSI-L = 1 marker; MSS = stable. Mononucleotide panels are more specific.

NGS-based MSI / tumour mutational burden (TMB) from large panels estimates MSI from indel burden at homopolymers and reports MMR gene mutations. Advantages: simultaneous RAS/BRAF/POLE and theranostic genes. Pitfalls: low tumour purity, poor coverage of microsatellites, and MSH6-mutant tumours sometimes MSI-L/MSS on limited panels.

- Concordance IHC ↔ PCR is high (~90–95%); discordances need both modalities + genetics discussion.
- Liquid biopsy MSI is emerging but tissue remains gold standard for treatment decisions when available.
- For CPI eligibility, FDA labels accept IHC dMMR or PCR/NGS MSI-H per companion diagnostics.

Drug / regimen	Dose / schedule (exam pearls)
IHC MMR panel	MLH1 / MSH2 / MSH6 / PMS2 nuclear stain; internal control mandatory
PCR MSI	Tumour vs normal; MSI-H if ≥2/5 (panel-dependent)
NGS MSI/TMB	Panel-based; report MMR variants + MSI score + TMB mut/Mb
Germline LS panel	MLH1, MSH2, MSH6, PMS2, EPCAM ± complementary assays for deletions

Table: doses and regimens

7. Interpretation workshop — pitfalls and reporting

Common pitfalls: (1) scoring cytoplasmic stain as retained; (2) missing weak MSH6 after chemotherapy/radiotherapy; (3) calling LS without methylation/BRAF workup; (4) treating MSI-L as MSI-H; (5) ignoring clonal heterogeneity after neoadjuvant therapy.

Report template should state method, antibodies/panel, result (dMMR vs pMMR or MSI-H/L/MSS), and recommended reflex tests. For exams, write a mini-algorithm and one sentence on genetic counselling triggers.

MMR testing interpretation loop




Figure: stepwise exam algorithm

EXAM PEARLS / HALLMARKS

- Exam trap: BRAF V600E in CRC supports sporadic MLH1 methylation — not LS.
- Exam trap: IHC loss without MSI-H still may qualify as dMMR for CPI if assay validated.

8. Mechanism — why dMMR/MSI-H predicts CPI benefit

Frameshift mutations in coding microsatellites generate mutant neopeptides presented on MHC-I/II. The inflamed TME of MSI-H tumours shows high CD8+ infiltration, upregulated PD-1/PD-L1, and interferon-γ signatures — priming for PD-1 blockade. CPIs release exhausted T cells to recognise these neoantigens → durable responses even in chemo-refractory disease.

Resistance still occurs: antigen presentation defects (B2M, HLA LOH), JAK/STAT IFN-pathway loss, alternative checkpoints, and immunosuppressive myeloid programs. MSS tumours rarely respond to PD-1 alone — hence combination and biomarker research.

- Tissue-agnostic principle: biology > histology when MSI-H/dMMR.
- TMB-high overlaps but is not identical to MSI-H — do not conflate casually.
- CTLA-4 + PD-1 (nivo+ipi) deepens responses in MSI-H mCRC (CheckMate-8HW).

dMMR	Unrepaired indels at microsatellites
Neoantigens	Frameshift peptides → MHC presentation
T-cell priming	High TILs; IFN-γ inflamed phenotype
Checkpoint up	PD-1/PD-L1 adaptive resistance
CPI unlock	Anti–PD-1/PD-L1 ± CTLA-4 → durable control

Figure: mechanism / pathway map

9. Landmark CPI trials (quote numbers)

Memorise phase, population, arms, and primary endpoint for four to six trials. Examiners expect KEYNOTE-177 medians and the tissue-agnostic KEYNOTE-158 ORR. Add ATOMIC for 2026 adjuvant CRC relevance and CheckMate-8HW for dual CPI.

Trial	Key result / exam takeaway
KEYNOTE-016/164	Early signal: pembro in dMMR CRC — high ORR; launched MSI-H CPI era
KEYNOTE-177 (Ph3)	1L pembro vs chemo MSI-H mCRC: PFS 16.5 vs 8.2 mo; 5-y OS ~50%+ with crossover
CheckMate-142	Nivo ± ipi in prior-treated MSI-H mCRC — durable ORR; ipi adds depth
CheckMate-8HW (Ph3)	1L nivo+ipi vs chemo MSI-H/dMMR mCRC — superior PFS; dual CPI standard discussion
KEYNOTE-158	Pembro MSI-H non-CRC: ORR ~34%; DoR durable → tissue-agnostic approval
ATOMIC / Alliance	Adjuvant atezo + FOLFOX/CAPOX stage III dMMR colon — NCCN 2026 preferred
Neoadjuvant dMMR rectal series	Dostarlimab / other CPI — clinical CR and organ preservation possible

Table: landmark clinical trials

EXAM PEARLS / HALLMARKS

- Write one HR or median per major trial — vague 'improved PFS' scores less.
- Mention chemo crossover in KEYNOTE-177 when discussing OS nuance.

10. Therapeutic strategies & doses when MMR is the biomarker

Metastatic MSI-H/dMMR solid tumours: anti–PD-1 monotherapy (pembrolizumab; nivolumab) or nivo+ipi in CRC per trial/label. Continue until progression/toxicity per protocol; pseudo-progression uncommon but known — confirm with clinical status ± repeat imaging.

Localised disease shifts (NCCN 2026): stage III dMMR colon — chemo + atezolizumab pathway; dMMR locally advanced rectal — neoadjuvant CPI as primary consideration aiming for organ preservation. POLE/POLD1 ultramutated tumours: CPI benefit often extrapolated.

MMR biomarker → therapy algorithm




Figure: stepwise exam algorithm

Drug / regimen	Dose / schedule (exam pearls)
Pembrolizumab	200 mg IV Q3W or 400 mg IV Q6W (CRC and tissue-agnostic MSI-H/dMMR)
Nivolumab	240 mg Q2W or 480 mg Q4W; with ipilimumab per CheckMate CRC regimens
Ipilimumab (CRC dual CPI)	1 mg/kg with nivolumab per CheckMate-8HW/142 schedules — monitor irAE
Atezolizumab (ATOMIC context)	With FOLFOX or CAPOX adjuvant stage III dMMR — per protocol/NCCN
Dostarlimab (rectal dMMR)	Neoadjuvant PD-1 in highly selected organ-preservation pathways

Table: doses and regimens

11. Toxicities, monitoring & supportive care

Immune-related AEs (irAEs): dermatitis, colitis, hepatitis, endocrinopathies (thyroid, hypophysitis, adrenal), pneumonitis, nephritis, rarer myocarditis/neurotoxicity. Dual CPI increases grade ≥3 irAE rates — counsel and monitor closely.

Management principles: hold CPI for ≥G2; steroids first-line for most; escalate to infliximab/vedolizumab (colitis), MMF (hepatitis), etc.; permanently discontinue for severe myocarditis/neurotoxicity. Baseline TSH, LFTs, creatinine; educate patients on red flags.

- Endocrine irAEs often need lifelong replacement even after CPI stop — do not 'taper off' thyroxine casually.
- Distinguish infection vs immune colitis before infliximab.
- Vaccinations and infection workup before intensifying immunosuppression.




Figure: stepwise exam algorithm

12. NCCN 2026 updates relevant to MMR / MSI

Weave NCCN Colon v3.2026 and Rectal v2.2026 bullets into the 'recent advances' paragraph. ATOMIC adjuvant atezolizumab for stage III dMMR and neoadjuvant CPI for dMMR rectal are the highest-yield 2026 lines. Also note aspirin for PIK3CA-altered resected CRC (orthogonal molecular point often paired in the same molecular workup discussion).

NCCN 2026 / 2027 UPDATES (include in answer)

- *Referenced versions:* Colon Cancer v3.2026; Rectal Cancer v2.2026; NCCN immunotherapy / disease-specific MSI-H pathways
- ATOMIC: FOLFOX or CAPOX + atezolizumab preferred for stage III dMMR/MSI-H colon (benefit assumed to extend to POLE/POLD1 hypermutated in panel discussion).
- Checkpoint inhibitor neoadjuvant pathway is a primary consideration for dMMR rectal tumours — organ preservation possible.
- Universal MMR/MSI testing informs both hereditary triage and CPI eligibility.
- Metastatic MSI-H CRC: CPI-based 1L preferred; dual CPI (nivo+ipi) per evidence hierarchy.
- Aspirin 100–162 mg daily × 3 years after surgery for stage II–III colon (and stage I rectal) with somatic PIK3CA pathway alterations — triggers broad molecular profiling.
- Structured exercise ≥150 min/week after adjuvant CRC therapy — cite as supportive NCCN 2026 item.

EXAM PEARLS / HALLMARKS

- Say 'NCCN Colon v3.2026 — ATOMIC preferred for stage III dMMR' verbatim in the closing box.

13. Special situations & differentials

CMMRD, Muir–Torre, and Turcot variants expand the syndromic discussion. Immunotherapy in LS-related non-CRC tumours follows agnostic principles. After CPI failure in MSI-H mCRC, options include chemo ± targeted therapy by RAS/BRAF/HER2, trials, and local therapies for oligoprogression while sometimes continuing CPI off-label/trial — acknowledge evidence gaps.

Differentiate MSI-H from TMB-H MSS (e.g., some melanomas, NSCLC) — both may respond to CPI but mechanisms and labels differ. Polymerase proofreading deficiency (POLE) is the classic ultramutated MSS CPI-responsive mimic.

- Pregnancy and CPI: generally avoid; individualise only in life-threatening disease.
- Autoimmune disease: weigh flare risk; often still treat MSI-H metastatic disease with caution.
- Transplant recipients: CPI may precipitate rejection — multidisciplinary decision.

14. Summary checklist & exam pearls

Close the answer with a boxed pathway: genes → tests → Lynch triage → CPI mechanism → key trials → NCCN 2026. Underline that MMR is a pan-tumour predictive biomarker with hereditary implications beyond the index cancer.

- Genes: MLH1, MSH2, MSH6, PMS2 (+ EPCAM).
- Tests: IHC ± MSI PCR/NGS; methylation/BRAF for MLH1-pattern.
- Mechanism: neoantigens + inflamed TME → PD-1 benefit.
- Trials: KEYNOTE-177, 8HW, 158, ATOMIC.
- NCCN 2026: adjuvant atezo stage III dMMR; neoadjuvant CPI dMMR rectal.

35-minute answer time plan




Figure: stepwise exam algorithm

EXAM PEARLS / HALLMARKS

- IHC loss patterns predict which gene to sequence — write the pairs.
- Sporadic MSI-H CRC = MLH1 methylation ± BRAF; Lynch ≠ MSI alone.
- KEYNOTE-177 mPFS 16.5 vs 8.2 mo is the number to memorise.
- ATOMIC + rectal neoadjuvant CPI are the 2026 'recent advances' clinchers.
- Always offer genetic counselling language when LS is possible.

— Additional exam-length revision modules —

The following modules reinforce algorithms, toxicities, Indian/NCG context, and last-page summary structure.

3. Diagnosis, biomarkers & staging

A modern oncology answer must separate **diagnostic**, **prognostic**, and **predictive** markers. Predictive markers change drug choice (EGFR, HER2, BRCA, MSI, Claudin18.2, FGFR, NTRK).

• Tissue diagnosis with adequate IHC panel; never treat metastatic disease without molecular profiling when actionable targets exist.

• Use validated assays (FDA/companion diagnostics where mandated).

• Stage with AJCC 8th/9th as applicable (NSCLC NCCN 2026 uses AJCC 9th).

• Performance status, organ function, comorbidities, geriatric assessment, and goals of care before regimen selection.

• Universal MMR IHC / MSI testing in CRC; guides CPI and ATOMIC adjuvant pathway.

Work-up algorithm (draw this in the answer booklet)




Figure: stepwise exam algorithm

4. Management algorithm, doses & regimens

Structure management as: (A) localized/curative-intent pathways, (B) advanced/metastatic systemic therapy, (C) special populations (CNS, elderly, organ dysfunction, pregnancy), (D) supportive care.

For each line of therapy state: eligibility, preferred regimen, dose, key toxicity monitoring, and the trial that supports it.

• Localized disease: multimodality — surgery/RT/systemic as per disease chapter.

• Neoadjuvant vs adjuvant: cite pathologic response and adjuvant switch trials (e.g., KATHERINE, CheckMate 577, KEYNOTE-522).

• Metastatic: biomarker-first; avoid empiric chemo when a high-level targeted/IO option exists.

• Oligometastatic disease: consider metastasis-directed therapy where NCCN supports (e.g., prostate APMS 2026).

• Re-biopsy / liquid biopsy at progression to map resistance.

Management flowchart (reproduce in exam)




Figure: stepwise exam algorithm

Drug / regimen	Dose / schedule (exam pearls)
Pembrolizumab	200 mg IV Q3W or 400 mg Q6W
Nivolumab	240 mg Q2W or 480 mg Q4W; with ipi 1 mg/kg (RCC) or 3 mg/kg (melanoma) per regimen
Olaparib	300 mg PO BID (tablets); adjust for toxicities/renal; OlympiA adjuvant 1 year
Trastuzumab	8 mg/kg → 6 mg/kg IV Q3W (or SC fixed dose)

Table: doses and regimens

5. Landmark clinical trials (must quote in answer)

Present trials in a table: name → design/population → intervention → primary endpoint → absolute/relative benefit → practice change. Examiners award marks for correct trial names and one numeric result (HR or median PFS/OS).

- Mention phase (III preferred for standards) and whether endpoint was PFS, OS, MRD, pCR, or EFS.
- Distinguish statistical vs clinical significance; note crossover and subgroup caveats.
- For India-focused answers, mention access/biosimilars/NCG guidance after the trial table.

Trial	Key result / exam takeaway
MOSAIC / C-07 / IDEA	Adjuvant oxaliplatin; 3 vs 6 mo CAPOX/FOLFOX
ATOMIC	FOLFOX/CAPOX + atezolizumab stage III dMMR — NCCN 2026 preferred
KEYNOTE-177 / CheckMate-8HW	CPI in MSI-H mCRC
BREAKWATER	Encorafenib + cetuximab + FOLFOX 1L BRAF V600E
FIRE-3 / PEAK / CALGB 80405	EGFR mAb + chemo in RAS WT left-sided
BEACON	Encorafenib + cetuximab ± binimetinib BRAF V600E

Table: landmark clinical trials

6. Toxicities, supportive care & monitoring

A complete answer links efficacy to safety. List monitoring schedules (CBC, chemistry, ECG, TFTs, cortisol, LFTs, PFTs/CT for ILD risk drugs).

- Anticipate class toxicities and counsel patients; use CTCAE grading.
- Cardio-oncology: anthracyclines, HER2 mAbs, ibrutinib, checkpoint myocarditis — baseline ECG/echo when indicated.
- IO toxicities: irAEs — hold drug, steroids, specialty referral; endocrine irAEs often permanent replacement.
- ADC pulmonary toxicity (T-DXd ILD): hold permanently for ≥grade 2 in many protocols; imaging + steroids.
- TLS prophylaxis for venetoclax/BLIN/CAR-T; rasburicase if high-risk.
- Fertility, sexuality, survivorship, and psychosocial support are part of comprehensive answers.
- Algorithm: grade 1 → continue with close watch; grade 2 → hold + steroids; grade 3–4 → discontinue permanently for many toxicities + high-dose steroids ± infliximab/vedolizumab/MMF.

Toxicity management loop




Figure: stepwise exam algorithm

7. NCCN 2026 / 2027 updates (high-yield for ‘recent advances’)

Paper IV rewards currency. Explicitly write: **“As per NCCN Guidelines updates in 2026/2027…”** and list 3–5 bullets. Also mention ESMO guidelines / Indian NCG where relevant for balance.

- State Category (1 / 2A / 2B) and preferred vs other recommended vs useful in certain circumstances.
- If Indian exam: reconcile NCCN with CDSCO approvals and NCG practical pathways.

NCCN 2026 / 2027 UPDATES (include in answer)

- *Referenced versions:* Colon Cancer v3.2026; Rectal Cancer v2.2026
- Aspirin 100–162 mg daily × 3 years after recovery from surgery for stage II–III colon (and stage I rectal) with somatic PIK3CA pathway alterations; triggers molecular profiling of all stage II–III tumors.
- ATOMIC: FOLFOX or CAPOX + atezolizumab preferred for stage III dMMR/MSI-H colon cancer (benefit assumed to extend to POLE/POLD1 hypermutated).
- Checkpoint inhibitor neoadjuvant pathway primary consideration for dMMR rectal tumors (organ preservation possible).
- Structured exercise ≥150 min moderate–vigorous activity/week after adjuvant therapy (≈28% relative reduction in deaths/recurrences).
- FOLFOX + encorafenib + cetuximab/panitumumab as 1L systemic option for BRAF V600E mCRC (BREAKWATER).
- Neoadjuvant immunotherapy preferred pathway for dMMR/MSI-H locally advanced rectal cancer.

- FOLFIRINOX neoadjuvant option for pMMR/MSS resectable disease with liver-only and/or lung-only metastases.
- Aspirin recommendation as in colon for PIK3CA-altered stage I rectal after surgery.
- Earlier-line CPI expansion: ATOMIC stage III dMMR colon; neoadjuvant CPI for dMMR rectal; neoadjuvant melanoma; EV+P bladder; FLOT+durvalumab gastric.
- PD-L1 testing emphasized for gastric/EGJ; limited benefit if PD-L1 negative.
- Prostate: ADT + niraparib/abiraterone for BRCA2 APMS subsets.
- Ovarian: PARPi focused on maintenance/biomarker-defined; removed from many recurrent treatment sections.

EXAM PEARLS / HALLMARKS

- NCCN versions change frequently — cite year/version (e.g., NSCLC v9.2026, Breast v9.2026, Myeloma v2.2027).
- Category 1 = high-level evidence + uniform NCCN consensus.

8. Special situations, differentials & practical issues

Discuss differentials, mimics, and when to refer to radiation oncology, surgery, palliative care, genetics, or cell-therapy centers.

Include brief notes on health economics (cost-effectiveness, biosimilars) and clinical trial enrollment as standard options.

- MDT documentation improves outcomes and exam completeness.
- Survivorship care plans: late effects, second malignancies, cardiac, endocrine, psychosocial.
- Real-world evidence and digital tools increasingly complement RCT data (relevant to 2026 Paper IV themes).

9. Evidence hierarchy, patient rights & benefits

Quote evidence using: Phase III RCT > meta-analysis of RCTs > prospective registries / RWE > retrospective > expert consensus. NCCN Categories and ESMO-MCBS scores help communicate strength of benefit.

Patient-centered care: informed consent for toxicities (especially CRS/ICANS, ILD, infertility), shared decision-making, financial toxicity counseling, and access to supportive/palliative care concurrent with active therapy.

- Benefits to highlight: OS, PFS, pCR, MRD negativity, organ preservation, QoL, reduced hospitalization.
- Rights: autonomy, confidentiality (EMR), right to second opinion, trial participation, refusal of care, end-of-life preferences.
- Equity: molecular testing access, biosimilars, NCG/Ayushman Bharat practical pathways for Indian exams.
- Pharmacovigilance: report ADRs; Phase IV / post-marketing commitments.

EXAM PEARLS / HALLMARKS

- Write one sentence on absolute benefit (e.g., ‘median OS improved from X to Y months’).)
- Mention number-needed-to-treat conceptually if space permits.

10. Summary checklist (write this last page in the booklet)

Close the 15-mark answer with a crisp numbered summary the examiner can tick.

- Definition + epidemiology one-liner
- Key biology / hallmark diagram
- Essential diagnostics & biomarkers
- Risk/stage stratification

- Preferred treatment algorithm with ≥2 doses
- ≥3 named trials with endpoints
- Major toxicities + monitoring
- NCCN 2026 update bullets (3–5)
- Special situations
- Five exam pearls

30-second revision flowchart




Figure: stepwise exam algorithm

EXAM PEARLS / HALLMARKS

- Neat flowchart + trial table often separates a 10/15 from a 13–15/15.
- Do not dump drug lists without doses or evidence.
- Currency: say 'NCCN 2026' explicitly for Paper IV.

11. Pharmacology deep-dive (class, PK/PD, interactions)

For drug-centered questions, present: chemical/biologic class → target epitope/kinase → PK (oral bioavailability, half-life, hepatic/renal clearance) → food effects → CYP interactions → dose adjustments → therapeutic drug monitoring if any.

Example interaction pearls: acid-suppressive therapy with some TKIs; QT-prolonging combinations; azole antifungals with vinca/TKIs; strong CYP3A inducers reducing exposure to many oral targeted agents.

- State renal/hepatic dose modifications when known.
- Note formulation differences (olaparib tablets vs capsules; IV vs SC daratumumab).
- Infusion reactions: premedication protocols for mAbs/BiTEs.

Drug / regimen	Dose / schedule (exam pearls)
Pembrolizumab	200 mg IV Q3W or 400 mg Q6W
Nivolumab	240 mg Q2W or 480 mg Q4W; with ipi 1 mg/kg (RCC) or 3 mg/kg (melanoma) per regimen
Olaparib	300 mg PO BID (tablets); adjust for toxicities/renal; OlympiA adjuvant 1 year
Trastuzumab	8 mg/kg → 6 mg/kg IV Q3W (or SC fixed dose)

Table: doses and regimens

12. Response assessment & follow-up schedules

Use disease-specific criteria: RECIST 1.1 solid tumors; iRECIST for immunotherapy; Lugano/Deauville for lymphoma; IMWG for myeloma; ELN molecular response for CML/AML; PSA working group for prostate.

- Typical imaging interval q2–3 cycles in metastatic disease unless clinically indicated sooner.
- PET-adapted strategies in Hodgkin lymphoma (RATHL, AHL2011) — cite Deauville 1–3 negative vs 4–5 positive.
- ctDNA MRD emerging for escalation/de-escalation (CRC, lymphoma, breast — exam discussion point).

Clinical algorithm




Figure: stepwise exam algorithm

13. Comparison tables useful for exams

Draw 2×N tables comparing old standard vs new standard; or drug A vs B (efficacy, toxicity, CNS activity, cost/access).

- Column ideas: target, dose, PFS, OS, grade ≥3 AEs, CNS activity, NCCN category.
- Another table: resistance mutation → preferred next drug.

Trial	Key result / exam takeaway
MOSAIC / C-07 / IDEA	Adjuvant oxaliplatin; 3 vs 6 mo CAPOX/FOLFOX
ATOMIC	FOLFOX/CAPOX + atezolizumab stage III dMMR — NCCN 2026 preferred
KEYNOTE-177 / CheckMate-8HW	CPI in MSI-H mCRC
BREAKWATER	Encorafenib + cetuximab + FOLFOX 1L BRAF V600E
FIRE-3 / PEAK / CALGB 80405	EGFR mAb + chemo in RAS WT left-sided
BEACON	Encorafenib + cetuximab ± binimetinib BRAF V600E

Table: landmark clinical trials

14. Indian context, NCG & practical exam tips

National Cancer Grid (NCG) resource-stratified guidelines are frequently expected in Indian DM exams. Discuss biosimilars (trastuzumab, rituximab), generic TKIs, and center-based cell therapy availability.

- Document MDT decisions; clinical trial referral when standard options exhausted.
- Counsel financial toxicity transparently; prioritize highest-value Category 1 options.
- Infection prophylaxis algorithms crucial in hematologic topics (antifungal, antiviral, PJP).

15. Sample ‘examiner hot spots’ & model closing paragraph

Model closing: “In summary, management of this problem is biomarker-driven. Current standards are supported by phase III trials and reflected in NCCN 2026/2027 updates. Sequencing should maximize OS and QoL while mitigating class-specific toxicities. Enrollment in clinical trials and adherence to MDT pathways remain integral.”

Hot spots: exact doses, one HR/median, resistance mechanism, NCCN category, and a drawn algorithm.

EXAM PEARLS / HALLMARKS

- If stuck on memory: write mechanism → class → one trial → one dose → one toxicity → NCCN 2026 line.
- Never contradict yourself between algorithm and trial table.

A. Epidemiology & burden (write ½–1 page)

Begin with global and Indian epidemiology when known (GLOBOCAN / ICMR / NCRP). State age distribution, sex ratio, and risk factors (tobacco, infection-related cancers such as HPV/HBV/HCV/H. pylori/HIV, obesity, inherited syndromes).

Explain why incidence or mortality trends matter for screening and resource allocation. For rare diseases, state incidence per 100,000 and referral-center patterns.

Link epidemiology to prevention: vaccination (HPV, HBV), tobacco control, hepatitis treatment, aspirin in selected molecular CRC, and lifestyle (exercise post-adjuvant CRC per NCCN 2026).

- One sentence on screening relevance (mammography, LDCT, FIT/colonoscopy, low-dose CT criteria).
- Mention hereditary fraction if applicable (Lynch, HBOC, FAP, MEN, VHL, HPRCC).
- State stage distribution at diagnosis in LMICs vs HIC if relevant to treatment sequencing.

B. Historical evolution → modern standard (timeline)

Examiners like a short timeline: historical surgery/RT/chemo era → targeted therapy era → immunotherapy/ADC/cellular therapy era.

Highlight inflection trials that changed guidelines (e.g., imatinib IRIS; trastuzumab HERA; checkpoint inhibitors; ADCs; CAR-T approvals).

End timeline with **NCCN 2026/2027** preferred regimens so the answer feels contemporary for Paper IV.

Therapeutic eras timeline




Figure: stepwise exam algorithm

C. Detailed differential diagnosis & pathology pearls

List 5–7 differentials and the IHC/molecular discriminators. Pathology reports should include grade, LVI, margins, nodes, and required biomarkers.

For hematologic topics: morphology → flow cytometry → cytogenetics → NGS panel → integrated WHO2022/ICC diagnosis.

For CUP/unknown primary style questions: IHC algorithm (CK7/CK20, TTF1, CDX2, GATA3, PAX8, synaptophysin) plus NGS and MSI/HER2.

- Always write ‘correlate clinically with radiology and tumor markers’.
- State pre-analytical caveats (decalcification, fixation time) affecting HER2/ER assays.
- Liquid biopsy complements but does not always replace tissue.

Morphology	Light microscopy pattern
IHC panel	Lineage + predictive markers
Cytogenetics	Karyotype / FISH fusions
NGS	SNVs, indels, CNAs, MSI, TMB
Integrated Dx	WHO/ICC + AJCC stage

Figure: mechanism / pathway map

D. Multidisciplinary care pathway & sequencing

Describe roles: medical oncology, surgical oncology, radiation oncology, radiology, pathology, palliative care, genetics, psycho-oncology, rehabilitation, and clinical trials unit.

Sequencing principles: treat drivers first when OS-proven; use neoadjuvant therapy to downstage and guide adjuvant escalation; avoid overlapping severe toxicities; protect future options (e.g., cumulative cardiotoxicity, neuropathy).

For oligometastatic disease discuss local therapy (surgery/SBRT/ablation) plus systemic therapy — prostate NCCN 2026 explicitly incorporates metastasis-directed therapy in selected APMS disease.

- Document MDT minutes in the file — also good exam phrase.
- Re-staging after neoadjuvant before definitive local therapy.
- Early palliative care improves QoL and sometimes survival in metastatic settings.

Clinical algorithm




Figure: stepwise exam algorithm

E. Colorectal cancer deep notes

Side and RAS/BRAF/MMR define therapy. Left RAS WT: EGFR mAb+chemo; right-sided less EGFR benefit. BRAF V600E: encorafenib+EGFR mAb ± chemo (BREAKWATER 1L NCCN).

Stage III dMMR: ATOMIC — FOLFOX/CAPOX + atezolizumab preferred NCCN 2026. Neoadjuvant CPI for dMMR rectal — organ preservation possible.

Aspirin 100–162 mg × 3 years if PIK3CA pathway alteration after surgery (stage II–III colon / stage I rectal) — NCCN 2026.

Exercise ≥150 min/week after adjuvant therapy — formal NCCN recommendation.

Metastatic sequencing: MSI-H → CPI; HER2 amplified RAS WT → anti-HER2; KRAS G12C → sotorasib/adagrasib combinations evolving; ctDNA for MRD under study.

- IDEA collaboration: 3 months CAPOX sufficient for many low-risk stage III.
- Watch-and-wait after TNT in selected rectal complete responders.
- HIPEC not routine for CRC peritoneal disease outside trials/selected centers.

E. Immunotherapy deep notes

CTLA-4 acts priming phase in lymphoid organs; PD-1/PD-L1 in peripheral effector phase. LAG-3 (relatlimab) additional checkpoint.

Biomarkers: PD-L1, MSI/dMMR, TMB — imperfect. TME, gut microbiome modulation is an exam discussion point (2026 short note theme).

Neoadjuvant IO preferred for resectable nodal melanoma (NCCN 2026). Combinations with VEGF in RCC; with chemo in lung/gastric; with ADC in bladder (EV+P).

- irAE timelines: skin/GI early; endocrine anytime; late neuropathies/myocarditis rare but deadly.
- Rechallenge decisions individualized after grade 2 resolution; avoid after grade 4 myocarditis/SJS etc.

G. Supportive care mini-chapter

Antiemetics (olanzapine-containing regimens for HEC), G-CSF primary/secondary prophylaxis, antimicrobial prophylaxis, mucositis care, nutrition (including TPN indications/contraindications), pain ladder + interventional techniques, thrombosis prevention (Khorana), growth factors, transfusion thresholds.

Psychosocial distress screening, fertility preservation referral before gonadotoxic therapy, and survivorship clinics.

- Febrile neutropenia: risk assess; anti-pseudomonal beta-lactam; add antifungal if persistent.
- Checkpoint diarrhea: rule out infection then steroids promptly.
- ADC ILD algorithms — patient education on dyspnea.

H. Model answer skeleton (time plan for 35 minutes)

Minutes 0–3: definition + epidemiology + diagram outline. Minutes 3–12: biology + diagnosis. Minutes 12–25: management algorithm with doses/trials. Minutes 25–30: toxicities + NCCN 2026. Minutes 30–35: summary pearls + underline Category 1 regimens.

- Leave a half page blank early for a neat flowchart drawn once.
- Underline drug doses.
- Box the NCCN 2026 bullets.

EXAM PEARLS / HALLMARKS

- Currency + clarity > exhaustive rare regimens.
- Named trial + number beats unnamed claim.
- NCCN 2026 phrase is expected in Paper IV recent advances.

K. Pitfalls, contraindications & ‘do not miss’ warnings

List absolute/relative contraindications of key drugs (e.g., baseline ILD caution for T-DXd; severe autoimmune disease caution for CPI; T315I needs ponatinib/asciminib not 1G/2G TKI alone; ATRA differentiation syndrome in APL; TLS with venetoclax/Blincyto; HBV reactivation with anti-CD20/CPI).

Drug–drug interactions and QT prolongation traps are classic viva points — mention checking concomitant meds (azoles, macrolides, acid suppressants with selected TKIs).

Imaging pitfalls: pseudo-progression on IO; flare on bone scan; PET Deauville interpretation after immunomodulatory therapy.

- Always screen hepatitis B before anti-CD20, anthracyclines, and many systemic regimens.
- Pregnancy test before teratogenic agents (IMiDs — REMS; MTX; TKI counseling).
- Do not combine overlapping QT-prolonging agents without ECG plan.
- Document performance status — unfit vs fit changes entire AML/lymphoma pathway.

Clinical algorithm




Figure: stepwise exam algorithm

L. Case vignette practice (apply the algorithm)

Invent a short case in the answer: age, stage, biomarkers, comorbidities → your exact regimen with dose → expected monitoring → what you will do at progression. This shows applied knowledge and often gains examiner goodwill.

Example template: “45-year-old, ECOG 1, biomarker X positive, no contraindication Y → start regimen Z at dose… per trial T / NCCN 2026 preferred. Assess response at interval N. At progression, re-profile for resistance mutation R and switch to agent S.”

Add one geriatric or organ-dysfunction variant to show nuance (dose reduce, alternate preferred regimen, or best supportive care).

- Vignette 1: fit, biomarker-positive — preferred Cat 1 therapy.
- Vignette 2: comorbid/unfit — alternative regimen.
- Vignette 3: progression with resistance mutation — sequenced next line.
- Vignette 4: severe toxicity — hold, treat, rechallenge decision.

M. One-page ‘wall chart’ summary (draw/box this)

Reserve the penultimate page for a wall-chart style summary with four quadrants: (1) Biology (2) Work-up (3) Treatment lines (4) NCCN 2026 bullets. This mirrors how many toppers present long answers and helps hit the page target with high signal density.

Quadrant 1	Hallmark gene / pathway / figure
Quadrant 2	Must-order tests + staging
Quadrant 3	1L / 2L / 3L with doses
Quadrant 4	Trials + NCCN 2026 + toxicities
Center	MDT + patient goals

Figure: mechanism / pathway map

EXAM PEARLS / HALLMARKS

- Wall chart + flowchart + trial table = visual completeness.
- Write 'NCCN 2026' in quadrant 4 explicitly.

I2. Full-length essay addendum — tables to copy into the answer book

For 15-mark elaborations, add two hand-drawn tables: (Table A) Drug–dose–pearl; (Table B) Trial–population–endpoint–practice change. Then write a short paragraph on quality-of-life and patient-reported outcomes, which modern trials increasingly report.

Discuss biomarker evolution on therapy (liquid biopsy cadence), and when to stop therapy (duration of CPI, TFR in CML, maintenance stopping rules).

Include a brief note on second malignancies and late effects when curative-intent therapy is discussed (alkylators, RT fields, PARPi MDS/AML risk, CHL survivors).

- Table A columns: Drug | Dose | Key toxicity | Pearl
- Table B columns: Trial | Arm | Primary endpoint | Result | Change in practice
- One paragraph on shared decision-making and financial counseling

Trial	Key result / exam takeaway
MOSAIC / C-07 / IDEA	Adjuvant oxaliplatin; 3 vs 6 mo CAPOX/FOLFOX
ATOMIC	FOLFOX/CAPOX + atezolizumab stage III dMMR — NCCN 2026 preferred
KEYNOTE-177 / CheckMate-8HW	CPI in MSI-H mCRC
BREAKWATER	Encorafenib + cetuximab + FOLFOX 1L BRAF V600E
FIRE-3 / PEAK / CALGB 80405	EGFR mAb + chemo in RAS WT left-sided
BEACON	Encorafenib + cetuximab ± binimetinib BRAF V600E

Table: landmark clinical trials

Drug / regimen	Dose / schedule (exam pearls)
Pembrolizumab	200 mg IV Q3W or 400 mg Q6W
Nivolumab	240 mg Q2W or 480 mg Q4W; with ipi 1 mg/kg (RCC) or 3 mg/kg (melanoma) per regimen
Olaparib	300 mg PO BID (tablets); adjust for toxicities/renal; OlympiA adjuvant 1 year
Trastuzumab	8 mg/kg → 6 mg/kg IV Q3W (or SC fixed dose)

Table: doses and regimens

NCCN 2026 / 2027 UPDATES (include in answer)

- *Referenced versions:* Colon Cancer v3.2026; Rectal Cancer v2.2026
- Aspirin 100–162 mg daily × 3 years after recovery from surgery for stage II–III colon (and stage I rectal) with somatic PIK3CA pathway alterations; triggers molecular profiling of all stage II–III tumors.
- ATOMIC: FOLFOX or CAPOX + atezolizumab preferred for stage III dMMR/MSI-H colon cancer (benefit assumed to extend to POLE/POLD1 hypermutated).
- Checkpoint inhibitor neoadjuvant pathway primary consideration for dMMR rectal tumors (organ preservation possible).
- Structured exercise ≥150 min moderate–vigorous activity/week after adjuvant therapy (≈28% relative reduction in deaths/recurrences).
- FOLFOX + encorafenib + cetuximab/panitumumab as 1L systemic option for BRAF V600E mCRC (BREAKWATER).
- Neoadjuvant immunotherapy preferred pathway for dMMR/MSI-H locally advanced rectal cancer.
- FOLFIRINOX neoadjuvant option for pMMR/MSS resectable disease with liver-only and/or lung-only metastases.
- Aspirin recommendation as in colon for PIK3CA-altered stage I rectal after surgery.
- Earlier-line CPI expansion: ATOMIC stage III dMMR colon; neoadjuvant CPI for dMMR rectal; neoadjuvant melanoma; EV+P bladder; FLOT+durvalumab gastric.

- PD-L1 testing emphasized for gastric/EGJ; limited benefit if PD-L1 negative.
- Prostate: ADT + niraparib/abiraterone for BRCA2 APMS subsets.
- Ovarian: PARPi focused on maintenance/biomarker-defined; removed from many recurrent treatment sections.

I. Extended notes — resistance, sequencing & future directions

Resistance biology should occupy a dedicated half-page: on-target secondary mutations, bypass signaling, lineage plasticity (e.g., EGFR NSCLC → SCLC), antigen loss after cellular therapy, and immunosuppressive TME changes after CPI.

Sequencing principles: use the agent with largest OS benefit early when toxicities allow; reserve cross-resistant drugs; consider cumulative neuropathy/cardiotoxicity; plan CNS-active agents if brain metastases are present or likely.

Future directions appropriate for 2026 answers: tumor-agnostic indications (NTRK, RET, BRAF, MSI-H, TMB-H, HER2-mut), radioligand therapy, CAR-NK/off-the-shelf products, ctDNA-guided escalation/de-escalation, AI decision-support, and microbiome modulation adjuncts.

Health services angle: real-world evidence validating trial results in older/comorbid patients; digital remote toxicity monitoring; and equity of molecular testing access — themes also reflected in recent Paper IV questions.

- Write one ‘if progression then…’ resistance table.
- Mention at least one tumor-agnostic approval if biology overlaps.
- Close with trial enrollment as a standard option.

J. Quick-revision flash cards (last 1–2 pages)

Convert the topic into 8–10 flash cards mentally: (1) definition (2) hallmark gene (3) key test (4) 1L preferred (5) dose (6) trial (7) top toxicity (8) NCCN 2026 update (9) Indian/NCG pearl (10) one differential.

- Flash 1: One-line definition
- Flash 2: Driver / pathway
- Flash 3: Mandatory biomarker
- Flash 4: NCCN preferred 1L
- Flash 5: Exact dose
- Flash 6: Landmark phase III + endpoint
- Flash 7: Grade ≥3 toxicity to warn
- Flash 8: 2026 guideline change
- Flash 9: Resistance next step
- Flash 10: Supportive care must-do

EXAM PEARLS / HALLMARKS

- If time-poor in the hall: draw algorithm + trial table + NCCN box — captures most marks.
- Keep handwriting large for doses and HRs.

Final box — paste into last page of answer sheet

Re-list NCCN 2026 updates here so the examiner sees currency even if earlier pages are rushed.

NCCN 2026 / 2027 UPDATES (include in answer)

- *Referenced versions:* Colon Cancer v3.2026; Rectal Cancer v2.2026
- Aspirin 100–162 mg daily × 3 years after recovery from surgery for stage II–III colon (and stage I rectal) with somatic PIK3CA pathway alterations; triggers molecular profiling of all stage II–III tumors.
- ATOMIC: FOLFOX or CAPOX + atezolizumab preferred for stage III dMMR/MSI-H colon cancer (benefit assumed to extend to POLE/POLD1 hypermutated).
- Checkpoint inhibitor neoadjuvant pathway primary consideration for dMMR rectal tumors (organ preservation possible).
- Structured exercise ≥150 min moderate–vigorous activity/week after adjuvant therapy (≈28% relative reduction in deaths/recurrences).
- FOLFOX + encorafenib + cetuximab/panitumumab as 1L systemic option for BRAF V600E mCRC (BREAKWATER).
- Neoadjuvant immunotherapy preferred pathway for dMMR/MSI-H locally advanced rectal cancer.
- FOLFIRINOX neoadjuvant option for pMMR/MSS resectable disease with liver-only and/or lung-only metastases.
- Aspirin recommendation as in colon for PIK3CA-altered stage I rectal after surgery.
- Earlier-line CPI expansion: ATOMIC stage III dMMR colon; neoadjuvant CPI for dMMR rectal; neoadjuvant melanoma; EV+P bladder; FLOT+durvalumab gastric.
- PD-L1 testing emphasized for gastric/EGJ; limited benefit if PD-L1 negative.
- Prostate: ADT + niraparib/abiraterone for BRCA2 APMS subsets.
- Ovarian: PARPi focused on maintenance/biomarker-defined; removed from many recurrent treatment sections.

Final NCCN 2026 box

Copy these bullets onto the last page of your answer booklet for Paper IV currency marks.

NCCN 2026 / 2027 UPDATES (include in answer)

- *Referenced versions:* Colon Cancer v3.2026; Rectal Cancer v2.2026
- Aspirin 100–162 mg daily × 3 years after recovery from surgery for stage II–III colon (and stage I rectal) with somatic PIK3CA pathway alterations; triggers molecular profiling of all stage II–III tumors.
- ATOMIC: FOLFOX or CAPOX + atezolizumab preferred for stage III dMMR/MSI-H colon cancer (benefit assumed to extend to POLE/POLD1 hypermutated).
- Checkpoint inhibitor neoadjuvant pathway primary consideration for dMMR rectal tumors (organ preservation possible).
- Structured exercise ≥150 min moderate–vigorous activity/week after adjuvant therapy (≈28% relative reduction in deaths/recurrences).
- FOLFOX + encorafenib + cetuximab/panitumumab as 1L systemic option for BRAF V600E mCRC (BREAKWATER).
- Neoadjuvant immunotherapy preferred pathway for dMMR/MSI-H locally advanced rectal cancer.
- FOLFIRINOX neoadjuvant option for pMMR/MSS resectable disease with liver-only and/or lung-only metastases.
- Aspirin recommendation as in colon for PIK3CA-altered stage I rectal after surgery.
- Earlier-line CPI expansion: ATOMIC stage III dMMR colon; neoadjuvant CPI for dMMR rectal; neoadjuvant melanoma; EV+P bladder; FLOT+durvalumab gastric.
- PD-L1 testing emphasized for gastric/EGJ; limited benefit if PD-L1 negative.
- Prostate: ADT + niraparib/abiraterone for BRCA2 APMS subsets.
- Ovarian: PARPi focused on maintenance/biomarker-defined; removed from many recurrent treatment sections.

Disclaimer: Compiled for DM exam revision from published trials and NCCN 2026/2027 public summaries (ASCO Post, OncLive, JNCCN selections, NCCN category list). Always verify the current full NCCN PDF, FDA/CDSCO labels, and institutional protocols before patient care. Doses require BSA/organ-function adjustment.