

Two-Graphs and NSSDs: An Algebraic Approach

Luke Collins Irene Sciriha

Department of Mathematics
University of Malta

S³ Annual Science Conference
2nd March, 2018



Structure of the Talk

- 1 Introduction
 - Definition of a Graph
 - Representing Graphs as Matrices
 - Spectrum and Seidel Switching
 - Defining Two-Graphs
- 2 Regular Two-Graphs
 - The Involution \mathbf{M}
 - Descendant Form of a Regular Two-Graph
 - Results about Descendants of Regular Two-Graphs
- 3 Strongly Regular Graphs
 - Definition
 - Structure of Descendants of Regular Two-Graphs


Definition of a Graph

Definition of a Graph

In mathematics, a *graph* is not one of these:



Definition of a Graph

In mathematics, a *graph* is not one of these:






Definition of a Graph

In mathematics, a *graph* is not one of these:



Definition of a Graph

In mathematics, a *graph* is not one of these:



Definition of a Graph

Definition of a Graph

Definition (Graph)

A **graph** G is a pair (V, E) where V is a non-empty finite set, and E is a set of unordered pairs of the elements of V .

Definition of a Graph

Definition (Graph)

A **graph** G is a pair (V, E) where V is a non-empty finite set, and E is a set of unordered pairs of the elements of V .

The elements of the set V are called *vertices*, and the pairs in E are called *edges*.

Definition of a Graph

Definition (Graph)

A **graph** G is a pair (V, E) where V is a non-empty finite set, and E is a set of unordered pairs of the elements of V .

The elements of the set V are called *vertices*, and the pairs in E are called *edges*.

Example

$V = \{1, 2, 3, 4, 5\}$ and $E = \{\{1, 2\}, \{1, 3\}, \{2, 3\}, \{2, 4\}, \{3, 4\}, \{4, 5\}\}$ define a graph.

Definition of a Graph


Definition (Graph)

A **graph** G is a pair (V, E) where V is a non-empty finite set, and E is a set of unordered pairs of the elements of V .

The elements of the set V are called *vertices*, and the pairs in E are called *edges*.

Example

$V = \{1, 2, 3, 4, 5\}$ and $E = \{\{1, 2\}, \{1, 3\}, \{2, 3\}, \{2, 4\}, \{3, 4\}, \{4, 5\}\}$ define a graph.



Definition of a Graph



Definition (Graph)

A **graph** G is a pair (V, E) where V is a non-empty finite set, and E is a set of unordered pairs of the elements of V .

The elements of the set V are called *vertices*, and the pairs in E are called *edges*.

Example

$V = \{1, 2, 3, 4, 5\}$ and $E = \{\{1, 2\}, \{1, 3\}, \{2, 3\}, \{2, 4\}, \{3, 4\}, \{4, 5\}\}$ define a graph.



Definition of a Graph




Definition (Graph)

A **graph** G is a pair (V, E) where V is a non-empty finite set, and E is a set of unordered pairs of the elements of V .

The elements of the set V are called *vertices*, and the pairs in E are called *edges*.

Example

$V = \{1, 2, 3, 4, 5\}$ and $E = \{\{1, 2\}, \{1, 3\}, \{2, 3\}, \{2, 4\}, \{3, 4\}, \{4, 5\}\}$ define a graph.



Representing Graphs as Matrices

Representing Graphs as Matrices

We usually use the letter n for the number of vertices, that is, $n = |V|$.

Representing Graphs as Matrices

We usually use the letter n for the number of vertices, that is, $n = |V|$.
To encode graphs algebraically, we can use an *adjacency matrix*:

Representing Graphs as Matrices

We usually use the letter n for the number of vertices, that is, $n = |V|$. To encode graphs algebraically, we can use an *adjacency matrix*:

Definition (Adjacency matrix)

The **adjacency matrix** of a graph $G = (V, E)$ is the $n \times n$ matrix (a_{ij}) where

Representing Graphs as Matrices

We usually use the letter n for the number of vertices, that is, $n = |V|$. To encode graphs algebraically, we can use an *adjacency matrix*:

Definition (Adjacency matrix)

The **adjacency matrix** of a graph $G = (V, E)$ is the $n \times n$ matrix (a_{ij}) where

$$a_{ij} = \begin{cases} 1 & \text{if vertex } i \text{ is adjacent to vertex } j, \text{ i.e. } \{v_i, v_j\} \in E \\ 0 & \text{otherwise.} \end{cases}$$


Adjacency Matrix

Example

Adjacency Matrix

Example


The graph from the previous example has the following adjacency matrix.



Adjacency Matrix

Example


The graph from the previous example has the following adjacency matrix.



Adjacency Matrix

Example

The graph from the previous example has the following adjacency matrix.




$$\begin{array}{c}
 \begin{matrix} & 1 & 2 & 3 & 4 & 5 \\
 \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \end{matrix} & \begin{pmatrix} 0 & 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 1 & 0 \\ 1 & 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 \end{pmatrix}
 \end{matrix}
 \end{array}$$

Adjacency Matrix

Example

The graph from the previous example has the following adjacency matrix.




$$\begin{array}{c}
 \begin{matrix} & 1 & 2 & 3 & 4 & 5 \end{matrix} \\
 \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \end{matrix} \begin{pmatrix} 0 & 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 1 & 0 \\ 1 & 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 \end{pmatrix}
 \end{array}$$

Note that in general,

Adjacency Matrix

Example

The graph from the previous example has the following adjacency matrix.



$$\begin{array}{c}
 \begin{matrix} & 1 & 2 & 3 & 4 & 5 \end{matrix} \\
 \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \end{matrix} \begin{pmatrix}
 0 & 1 & 1 & 0 & 0 \\
 1 & 0 & 1 & 1 & 0 \\
 1 & 1 & 0 & 1 & 0 \\
 0 & 1 & 1 & 0 & 1 \\
 0 & 0 & 0 & 1 & 0
 \end{pmatrix}
 \end{array}$$


Note that in general,

- The adjacency matrix is symmetric

Adjacency Matrix

Example

The graph from the previous example has the following adjacency matrix.



$$\begin{array}{c}
 \begin{matrix} & 1 & 2 & 3 & 4 & 5 \end{matrix} \\
 \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \end{matrix} \begin{pmatrix}
 0 & 1 & 1 & 0 & 0 \\
 1 & 0 & 1 & 1 & 0 \\
 1 & 1 & 0 & 1 & 0 \\
 0 & 1 & 1 & 0 & 1 \\
 0 & 0 & 0 & 1 & 0
 \end{pmatrix}
 \end{array}$$


Note that in general,

- The adjacency matrix is symmetric
- Each 1 represents an edge, and each 0 represents a non-edge

Adjacency Matrix

Example

The graph from the previous example has the following adjacency matrix.



$$\begin{array}{c}
 \begin{matrix} & 1 & 2 & 3 & 4 & 5 \end{matrix} \\
 \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \end{matrix} \begin{pmatrix} 0 & 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 1 & 0 \\ 1 & 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 \end{pmatrix}
 \end{array}$$

Note that in general,

- The adjacency matrix is symmetric
- Each 1 represents an edge, and each 0 represents a non-edge
- Each entry on the diagonal is 0, since we consider simple graphs

Seidel Matrix

Seidel Matrix

Another way of encoding graphs is the *Seidel matrix*:

Seidel Matrix

Another way of encoding graphs is the *Seidel matrix*:

Definition (Seidel matrix)

The **Seidel matrix** of a graph $G = (V, E)$ is the $n \times n$ matrix (s_{ij}) where

Seidel Matrix

Another way of encoding graphs is the *Seidel matrix*:

Definition (Seidel matrix)

The **Seidel matrix** of a graph $G = (V, E)$ is the $n \times n$ matrix (s_{ij}) where

$$s_{ij} = \begin{cases} 0 & \text{if } i = j \\ -1 & \text{if vertex } i \text{ and vertex } j \text{ are adjacent} \\ 1 & \text{otherwise.} \end{cases}$$

Seidel Matrix

Another way of encoding graphs is the *Seidel matrix*:

Definition (Seidel matrix)

The **Seidel matrix** of a graph $G = (V, E)$ is the $n \times n$ matrix (s_{ij}) where

$$s_{ij} = \begin{cases} 0 & \text{if } i = j \\ -1 & \text{if vertex } i \text{ and vertex } j \text{ are adjacent} \\ 1 & \text{otherwise.} \end{cases}$$

Eventually we will work with a variant of the Seidel matrix, which we will introduce later.


Seidel Matrix

Example

Seidel Matrix

Example


The graph from the previous example has the following Seidel matrix.



Seidel Matrix

Example


The graph from the previous example has the following Seidel matrix.



Seidel Matrix

Example

The graph from the previous example has the following Seidel matrix.




$$\begin{array}{c}
 \begin{array}{ccccc}
 & 1 & 2 & 3 & 4 & 5 \\
 \begin{array}{c} 1 \\ 2 \\ 3 \\ 4 \\ 5 \end{array} & \begin{pmatrix} 0 & -1 & -1 & 1 & 1 \\ -1 & 0 & -1 & -1 & 1 \\ -1 & -1 & 0 & -1 & 1 \\ 1 & -1 & -1 & 0 & -1 \\ 1 & 1 & 1 & -1 & 0 \end{pmatrix}
 \end{array}
 \end{array}$$

Seidel Matrix

Example

The graph from the previous example has the following Seidel matrix.



$$\begin{matrix} & \begin{matrix} 1 & 2 & 3 & 4 & 5 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \end{matrix} & \begin{pmatrix} 0 & -1 & -1 & 1 & 1 \\ -1 & 0 & -1 & -1 & 1 \\ -1 & -1 & 0 & -1 & 1 \\ 1 & -1 & -1 & 0 & -1 \\ 1 & 1 & 1 & -1 & 0 \end{pmatrix} \end{matrix}$$

Note that in general, if \mathbf{A} and \mathbf{S} are the adjacency and Seidel matrices of a graph G respectively, then

$$\mathbf{S} = \mathbf{J} - \mathbf{I} - 2\mathbf{A}$$

where \mathbf{J} is the matrix consisting entirely of 1's and \mathbf{I} is the identity matrix.

Spectrum of a Graph

Spectrum of a Graph

The distinct eigenvalues $\mu_1, \mu_2, \dots, \mu_s$ of a given matrix \mathbf{X} together with their multiplicities m_1, m_2, \dots, m_s form the **spectrum** of \mathbf{X} , denoted $\mu_1^{(m_1)} \mu_2^{(m_2)} \dots \mu_s^{(m_s)}$.

Spectrum of a Graph

The distinct eigenvalues $\mu_1, \mu_2, \dots, \mu_s$ of a given matrix \mathbf{X} together with their multiplicities m_1, m_2, \dots, m_s form the **spectrum** of \mathbf{X} , denoted $\mu_1^{(m_1)} \mu_2^{(m_2)} \dots \mu_s^{(m_s)}$.

Definition (Spectra)

Spectrum of a Graph

The distinct eigenvalues $\mu_1, \mu_2, \dots, \mu_s$ of a given matrix \mathbf{X} together with their multiplicities m_1, m_2, \dots, m_s form the **spectrum** of \mathbf{X} , denoted $\mu_1^{(m_1)} \mu_2^{(m_2)} \dots \mu_s^{(m_s)}$.

Definition (Spectra)

- 1 The **spectrum** of a graph G is the spectrum of its adjacency matrix

Spectrum of a Graph

The distinct eigenvalues $\mu_1, \mu_2, \dots, \mu_s$ of a given matrix \mathbf{X} together with their multiplicities m_1, m_2, \dots, m_s form the **spectrum** of \mathbf{X} , denoted $\mu_1^{(m_1)} \mu_2^{(m_2)} \dots \mu_s^{(m_s)}$.

Definition (Spectra)

- ① The **spectrum** of a graph G is the spectrum of its adjacency matrix
- ② The **Seidel spectrum** of a graph G is the spectrum of its Seidel matrix

Seidel Switching


Seidel Switching

Given a graph $G = (V, E)$ and a subset of the vertices $U \subseteq V$, the operation of *Seidel switching* with respect to U **exchanges all edges and non-edges** between U and $V \setminus U$ to obtain the graph $SS(U)$.

Seidel Switching

Given a graph $G = (V, E)$ and a subset of the vertices $U \subseteq V$, the operation of *Seidel switching* with respect to U **exchanges all edges and non-edges** between U and $V \setminus U$ to obtain the graph $SS(U)$.


Example



Seidel Switching

Given a graph $G = (V, E)$ and a subset of the vertices $U \subseteq V$, the operation of *Seidel switching* with respect to U **exchanges all edges and non-edges** between U and $V \setminus U$ to obtain the graph $SS(U)$.


Example



Seidel Switching

Given a graph $G = (V, E)$ and a subset of the vertices $U \subseteq V$, the operation of *Seidel switching* with respect to U **exchanges all edges and non-edges** between U and $V \setminus U$ to obtain the graph $SS(U)$.


Example



Seidel Switching

Given a graph $G = (V, E)$ and a subset of the vertices $U \subseteq V$, the operation of *Seidel switching* with respect to U **exchanges all edges and non-edges** between U and $V \setminus U$ to obtain the graph $SS(U)$.


Example



Seidel Switching

Given a graph $G = (V, E)$ and a subset of the vertices $U \subseteq V$, the operation of *Seidel switching* with respect to U **exchanges all edges and non-edges** between U and $V \setminus U$ to obtain the graph $SS(U)$.


Example



Seidel Switching

Given a graph $G = (V, E)$ and a subset of the vertices $U \subseteq V$, the operation of *Seidel switching* with respect to U **exchanges all edges and non-edges** between U and $V \setminus U$ to obtain the graph $SS(U)$.

Example



What Seidel Switching does to the Seidel Matrix


What Seidel Switching does to the Seidel Matrix

We can assume that the vertices of the set $U \subseteq V$ are labelled first (otherwise simply relabel the vertices).


What Seidel Switching does to the Seidel Matrix

We can assume that the vertices of the set $U \subseteq V$ are labelled first (otherwise simply relabel the vertices). In our example, we had the following:

G




$SS(U)$, where $U = \{1, 3\}$




What Seidel Switching does to the Seidel Matrix

We can assume that the vertices of the set $U \subseteq V$ are labelled first (otherwise simply relabel the vertices). In our example, we had the following:

G



$SS(U)$, where $U = \{1, 3\}$




$$\begin{array}{c}
 \begin{array}{cc} 1 & 3 \end{array} \quad \begin{array}{cc} 2 & 4 \end{array} \quad 5 \\
 \begin{array}{c} 1 \\ 3 \\ 2 \\ 4 \\ 5 \end{array} \left(\begin{array}{cc|cc}
 0 & -1 & -1 & 1 & 1 \\
 -1 & 0 & -1 & -1 & 1 \\
 \hline
 -1 & -1 & 0 & -1 & 1 \\
 1 & -1 & -1 & 0 & -1 \\
 1 & 1 & 1 & -1 & 0
 \end{array} \right)
 \end{array}$$


What Seidel Switching does to the Seidel Matrix

We can assume that the vertices of the set $U \subseteq V$ are labelled first (otherwise simply relabel the vertices). In our example, we had the following:

G



$SS(U)$, where $U = \{1, 3\}$



$$\begin{array}{c}
 \begin{array}{cc} 1 & 3 \end{array} \quad \begin{array}{cc} 2 & 4 \end{array} \quad 5 \\
 \begin{array}{c} 1 \\ 3 \\ 2 \\ 4 \\ 5 \end{array} \left(\begin{array}{cc|cc} 0 & -1 & -1 & 1 & 1 \\ -1 & 0 & -1 & -1 & 1 \\ \hline -1 & -1 & 0 & -1 & 1 \\ 1 & -1 & -1 & 0 & -1 \\ 1 & 1 & 1 & -1 & 0 \end{array} \right)
 \end{array}$$

$$\begin{array}{c}
 \begin{array}{cc} 1 & 3 \end{array} \quad \begin{array}{cc} 2 & 4 \end{array} \quad 5 \\
 \begin{array}{c} 1 \\ 3 \\ 2 \\ 4 \\ 5 \end{array} \left(\begin{array}{cc|cc} 0 & -1 & 1 & -1 & -1 \\ -1 & 0 & 1 & 1 & -1 \\ \hline 1 & 1 & 0 & -1 & 1 \\ -1 & 1 & -1 & 0 & -1 \\ -1 & -1 & 1 & -1 & 0 \end{array} \right)
 \end{array}$$

What Seidel Switching does to the Seidel Matrix

What Seidel Switching does to the Seidel Matrix

In general, if \mathbf{S} and $\mathbf{S}_{SS(U)}$ are the Seidel matrices of G and $SS(U)$, then

$$\mathbf{S} = \left(\begin{array}{c|c} \mathbf{S}_U & \mathbf{R} \\ \hline \mathbf{R}^\top & \mathbf{S}_{V \setminus U} \end{array} \right) \iff$$

What Seidel Switching does to the Seidel Matrix

In general, if \mathbf{S} and $\mathbf{S}_{\text{SS}(U)}$ are the Seidel matrices of G and $\text{SS}(U)$, then

$$\mathbf{S} = \left(\begin{array}{c|c} \mathbf{S}_U & \mathbf{R} \\ \hline \mathbf{R}^\top & \mathbf{S}_{V \setminus U} \end{array} \right) \iff \mathbf{S}_{\text{SS}(U)} = \left(\begin{array}{c|c} \mathbf{S}_U & -\mathbf{R} \\ \hline -\mathbf{R}^\top & \mathbf{S}_{V \setminus U} \end{array} \right).$$

What Seidel Switching does to the Seidel Matrix

In general, if \mathbf{S} and $\mathbf{S}_{\text{SS}(U)}$ are the Seidel matrices of G and $\text{SS}(U)$, then

$$\mathbf{S} = \left(\begin{array}{c|c} \mathbf{S}_U & \mathbf{R} \\ \hline \mathbf{R}^\top & \mathbf{S}_{V \setminus U} \end{array} \right) \iff \mathbf{S}_{\text{SS}(U)} = \left(\begin{array}{c|c} \mathbf{S}_U & -\mathbf{R} \\ \hline -\mathbf{R}^\top & \mathbf{S}_{V \setminus U} \end{array} \right).$$

In other words, $\mathbf{S}_{\text{SS}(U)} = \mathbf{D}^{-1} \mathbf{S} \mathbf{D}$, where $\mathbf{D}^{-1} = \mathbf{D}$ is the diagonal matrix with $d_{ii} = +1$ if $i \in U$ and $d_{ii} = -1$ otherwise.

What Seidel Switching does to the Seidel Matrix

In general, if \mathbf{S} and $\mathbf{S}_{SS(U)}$ are the Seidel matrices of G and $SS(U)$, then

$$\mathbf{S} = \left(\begin{array}{c|c} \mathbf{S}_U & \mathbf{R} \\ \hline \mathbf{R}^\top & \mathbf{S}_{V \setminus U} \end{array} \right) \iff \mathbf{S}_{SS(U)} = \left(\begin{array}{c|c} \mathbf{S}_U & -\mathbf{R} \\ \hline -\mathbf{R}^\top & \mathbf{S}_{V \setminus U} \end{array} \right).$$

In other words, $\mathbf{S}_{SS(U)} = \mathbf{D}^{-1} \mathbf{S} \mathbf{D}$, where $\mathbf{D}^{-1} = \mathbf{D}$ is the diagonal matrix with $d_{ii} = +1$ if $i \in U$ and $d_{ii} = -1$ otherwise.

It follows that \mathbf{S} and $\mathbf{S}_{SS(U)}$ are similar, and therefore G and $SS(U)$ have the same Seidel spectrum.

Two-Graphs

Two-Graphs

The operation of Seidel switching defines an equivalence relation on the set of all graphs on n vertices.

Two-Graphs

The operation of Seidel switching defines an equivalence relation on the set of all graphs on n vertices.

Definition (Two-graph)

A **two-graph** or **switching class** is an equivalence class of the Seidel switching equivalence relation.

Two-Graphs

The operation of Seidel switching defines an equivalence relation on the set of all graphs on n vertices.

Definition (Two-graph)

A **two-graph** or **switching class** is an equivalence class of the Seidel switching equivalence relation.

- A two-graph on n vertices consists of all the n -vertex graphs with the same Seidel spectrum.

Two-Graphs

The operation of Seidel switching defines an equivalence relation on the set of all graphs on n vertices.

Definition (Two-graph)

A **two-graph** or **switching class** is an equivalence class of the Seidel switching equivalence relation.

- A two-graph on n vertices consists of all the n -vertex graphs with the same Seidel spectrum.
- The term 'two-graph' originally arose in a combinatorial context, and actually refers to a couple (V, Δ) where $\Delta \subseteq \binom{V}{3}$ is a collection of triples $\{v_1, v_2, v_3\}$ with the property that any 4-subset of V contains an even number of triples of Δ . This is known to be equivalent to our definition.

Regular Two-Graphs

Regular Two-Graphs

Definition (Regular two-graph)

A two-graph is said to be **regular** if the Seidel matrix of any representative has precisely two distinct eigenvalues.

Regular Two-Graphs

Definition (Regular two-graph)

A two-graph is said to be **regular** if the Seidel matrix of any representative has precisely two distinct eigenvalues.

- This is a valid definition because the Seidel spectrum of any member of a two-graph is the same.

Regular Two-Graphs

Definition (Regular two-graph)

A two-graph is said to be **regular** if the Seidel matrix of any representative has precisely two distinct eigenvalues.

- This is a valid definition because the Seidel spectrum of any member of a two-graph is the same.
- Reverting to the combinatorial definition of 'two-graph', (V, Δ) is said to be regular if every pair of vertices lies in the same number of triples of Δ . This is known to be equivalent to our definition.

The Involution M

The Involution \mathbf{M}

Suppose \mathbf{M} is a symmetric matrix which is involutory, that is, $\mathbf{M}^2 = \mathbf{I}$.
Then

The Involution \mathbf{M}

Suppose \mathbf{M} is a symmetric matrix which is involutory, that is, $\mathbf{M}^2 = \mathbf{I}$.
Then

- By spectral decomposition, \mathbf{M} has eigenvalues 1 and -1 .

The Involution \mathbf{M}

Suppose \mathbf{M} is a symmetric matrix which is involutory, that is, $\mathbf{M}^2 = \mathbf{I}$. Then

- By spectral decomposition, \mathbf{M} has eigenvalues 1 and -1 .
- If \mathbf{M} is written as

$$\mathbf{M} = \left(\begin{array}{c|c} \mathbf{B} & \mathbf{v} \\ \hline \mathbf{v}^\top & -\lambda \end{array} \right),$$

then $\mathbf{B}\mathbf{v} = \lambda\mathbf{v}$ and $|\lambda| < 1$.

The Involution \mathbf{M}

Suppose \mathbf{M} is a symmetric matrix which is involutory, that is, $\mathbf{M}^2 = \mathbf{I}$. Then

- By spectral decomposition, \mathbf{M} has eigenvalues 1 and -1 .
- If \mathbf{M} is written as

$$\mathbf{M} = \left(\begin{array}{c|c} \mathbf{B} & \mathbf{v} \\ \hline \mathbf{v}^\top & -\lambda \end{array} \right),$$

then $\mathbf{B}\mathbf{v} = \lambda\mathbf{v}$ and $|\lambda| < 1$.

Furthermore, if the spectrum of \mathbf{M} is $1^{(n-k)}(-1)^{(k)}$, then it follows by Cauchy's interlacing inequalities that the spectrum of \mathbf{B} is

$$1^{(n-k-1)}(-1)^{(k-1)}\lambda(1).$$

$$\mathbf{M} = \alpha \mathbf{S} - \lambda \mathbf{I}$$

$$\mathbf{M} = \alpha \mathbf{S} - \lambda \mathbf{I}$$

If \mathbf{S} is the Seidel matrix of a regular two-graph on n vertices with eigenvalues μ_1, μ_2 , then

$$\mathbf{M} = \alpha \mathbf{S} - \lambda \mathbf{I}$$

where

$$\alpha = \frac{2}{\mu_1 - \mu_2} \quad \text{and} \quad \lambda = \frac{\mu_1 + \mu_2}{\mu_1 - \mu_2}$$

is an involution.

$$\mathbf{M} = \alpha \mathbf{S} - \lambda \mathbf{I}$$

If \mathbf{S} is the Seidel matrix of a regular two-graph on n vertices with eigenvalues μ_1, μ_2 , then

$$\mathbf{M} = \alpha \mathbf{S} - \lambda \mathbf{I}$$

where

$$\alpha = \frac{2}{\mu_1 - \mu_2} \quad \text{and} \quad \lambda = \frac{\mu_1 + \mu_2}{\mu_1 - \mu_2}$$

is an involution.

- This matrix still gives us an encoding of the graph.

$$\mathbf{M} = \alpha \mathbf{S} - \lambda \mathbf{I}$$

If \mathbf{S} is the Seidel matrix of a regular two-graph on n vertices with eigenvalues μ_1, μ_2 , then

$$\mathbf{M} = \alpha \mathbf{S} - \lambda \mathbf{I}$$

where

$$\alpha = \frac{2}{\mu_1 - \mu_2} \quad \text{and} \quad \lambda = \frac{\mu_1 + \mu_2}{\mu_1 - \mu_2}$$

is an involution.


- This matrix still gives us an encoding of the graph.
- $\mu_1 \mu_2 = 1 - n$.

$$\mathbf{M} = \alpha \mathbf{S} - \lambda \mathbf{I}$$

Example ($K_{2,4}$)


$$\mathbf{M} = \alpha \mathbf{S} - \lambda \mathbf{I}$$

Example ($K_{2,4}$)



$$\mathbf{M} = \alpha \mathbf{S} - \lambda \mathbf{I}$$


Example ($K_{2,4}$)



Seidel spectrum: $(-1)^5(5)^1$

$$\mathbf{M} = \alpha \mathbf{S} - \lambda \mathbf{I}$$

Example ($K_{2,4}$)




Seidel spectrum: $(-1)^5(5)^1$

$$\alpha = \frac{2}{\mu_1 - \mu_2} = -\frac{1}{3}$$

$$\lambda = \frac{\mu_1 + \mu_2}{\mu_1 - \mu_2} = -\frac{2}{3}$$

$$\mathbf{M} = \alpha \mathbf{S} - \lambda \mathbf{I}$$

Example ($K_{2,4}$)Seidel spectrum: $(-1)^5(5)^1$


$$\alpha = \frac{2}{\mu_1 - \mu_2} = -\frac{1}{3}$$

$$\lambda = \frac{\mu_1 + \mu_2}{\mu_1 - \mu_2} = -\frac{2}{3}$$

$$\mathbf{S} = \begin{pmatrix} 0 & -1 & 1 & 1 & -1 & 1 \\ -1 & 0 & -1 & -1 & 1 & -1 \\ 1 & -1 & 0 & 1 & -1 & 1 \\ 1 & -1 & 1 & 0 & -1 & 1 \\ -1 & 1 & -1 & -1 & 0 & -1 \\ 1 & -1 & 1 & 1 & -1 & 0 \end{pmatrix}$$

$$\mathbf{M} = \alpha \mathbf{S} - \lambda \mathbf{I}$$

Example ($K_{2,4}$)



Seidel spectrum: $(-1)^5(5)^1$

$$\alpha = \frac{2}{\mu_1 - \mu_2} = -\frac{1}{3}$$

$$\lambda = \frac{\mu_1 + \mu_2}{\mu_1 - \mu_2} = -\frac{2}{3}$$

$$\mathbf{M} = \alpha \mathbf{S} - \lambda \mathbf{I} = \begin{pmatrix} 2/3 & 1/3 & -1/3 & -1/3 & 1/3 & -1/3 \\ 1/3 & 2/3 & 1/3 & 1/3 & -1/3 & 1/3 \\ -1/3 & 1/3 & 2/3 & -1/3 & 1/3 & -1/3 \\ -1/3 & 1/3 & -1/3 & 2/3 & 1/3 & -1/3 \\ 1/3 & -1/3 & 1/3 & 1/3 & 2/3 & 1/3 \\ -1/3 & 1/3 & -1/3 & -1/3 & 1/3 & 2/3 \end{pmatrix}$$

Descendant Form

Descendant Form

Every two-graph on n vertices has a class representative of the form $D \dot{\cup} K_1$ where D is a graph on $n - 1$ vertices.

Descendant Form

Every two-graph on n vertices has a class representative of the form $D \dot{\cup} K_1$ where D is a graph on $n - 1$ vertices.

Definition (Descendant)

Any two-graph representative of the form $D \dot{\cup} K_1$ is said to be in *descendant form*, and the component D is said to be a *descendant* of the two-graph.

Descendant Form

Every two-graph on n vertices has a class representative of the form $D \dot{\cup} K_1$ where D is a graph on $n - 1$ vertices.

Definition (Descendant)

Any two-graph representative of the form $D \dot{\cup} K_1$ is said to be in *descendant form*, and the component D is said to be a *descendant* of the two-graph.

Obtaining a Descendant Form

Consider a representative (V, E) which is not in descendant form.

Descendant Form

Every two-graph on n vertices has a class representative of the form $D \dot{\cup} K_1$ where D is a graph on $n - 1$ vertices.

Definition (Descendant)

Any two-graph representative of the form $D \dot{\cup} K_1$ is said to be in *descendant form*, and the component D is said to be a *descendant* of the two-graph.

Obtaining a Descendant Form

Consider a representative (V, E) which is not in descendant form.

- 1 Pick any vertex $v \in V$.

Descendant Form

Every two-graph on n vertices has a class representative of the form $D \dot{\cup} K_1$ where D is a graph on $n - 1$ vertices.

Definition (Descendant)

Any two-graph representative of the form $D \dot{\cup} K_1$ is said to be in *descendant form*, and the component D is said to be a *descendant* of the two-graph.

Obtaining a Descendant Form

Consider a representative (V, E) which is not in descendant form.

- 1 Pick any vertex $v \in V$.
- 2 Let U be the set of all neighbours of v .

Descendant Form

Every two-graph on n vertices has a class representative of the form $D \dot{\cup} K_1$ where D is a graph on $n - 1$ vertices.

Definition (Descendant)

Any two-graph representative of the form $D \dot{\cup} K_1$ is said to be in *descendant form*, and the component D is said to be a *descendant* of the two-graph.

Obtaining a Descendant Form

Consider a representative (V, E) which is not in descendant form.

- 1 Pick any vertex $v \in V$.
- 2 Let U be the set of all neighbours of v .
- 3 Then the vertex v is isolated in $SS(U)$.

Descendant Form

Example (The Petersen Graph)

Descendant Form


Example (The Petersen Graph)

The famous Petersen graph is contained in a regular two-graph.

Descendant Form

Example (The Petersen Graph)


The famous Petersen graph is contained in a regular two-graph.



Descendant Form

Example (The Petersen Graph)

The famous Petersen graph is contained in a regular two-graph.




Let us isolate vertex 8.

Descendant Form

Example (The Petersen Graph)

The famous Petersen graph is contained in a regular two-graph.




Let us isolate vertex 8. Its set of neighbours is $U = \{7, 3, 9\}$.

Descendant Form

Example (The Petersen Graph)

The famous Petersen graph is contained in a regular two-graph.




Let us isolate vertex 8. Its set of neighbours is $U = \{7, 3, 9\}$. Now we focus on the edges between U and $V \setminus U$.

Descendant Form

Example (The Petersen Graph)

The famous Petersen graph is contained in a regular two-graph.




Let us isolate vertex 8. Its set of neighbours is $U = \{7, 3, 9\}$. Now we focus on the edges between U and $V \setminus U$. And the non-edges.

Descendant Form

Example (The Petersen Graph)

The famous Petersen graph is contained in a regular two-graph.




Let us isolate vertex 8. Its set of neighbours is $U = \{7, 3, 9\}$. Now we focus on the edges between U and $V \setminus U$. And the non-edges. Switch edges and non-edges.

Descendant Form

Example (The Petersen Graph)

The famous Petersen graph is contained in a regular two-graph.




Let us isolate vertex 8. Its set of neighbours is $U = \{7, 3, 9\}$. Now we focus on the edges between U and $V \setminus U$. And the non-edges. Switch edges and non-edges. Obtain $SS(U)$.

Descendant Form

Example (The Petersen Graph)

The famous Petersen graph is contained in a regular two-graph.



Let us isolate vertex 8. Its set of neighbours is $U = \{7, 3, 9\}$. Now we focus on the edges between U and $V \setminus U$. And the non-edges. Switch edges and non-edges. Obtain $SS(U)$. Move vertices around to look nicer.

Results about Descendants of Regular Two-Graphs

Results about Descendants of Regular Two-Graphs

Using the fact that $\mathbf{M}^2 = \mathbf{I}$, we easily obtain the following known results for descendants of regular two-graphs.

Results about Descendants of Regular Two-Graphs

Using the fact that $\mathbf{M}^2 = \mathbf{I}$, we easily obtain the following known results for descendants of regular two-graphs.

- 1 D is a ρ -regular subgraph, each vertex having degree

$$\rho = \frac{n}{2} - \frac{\lambda}{\alpha} - 1.$$

Results about Descendants of Regular Two-Graphs

Using the fact that $\mathbf{M}^2 = \mathbf{I}$, we easily obtain the following known results for descendants of regular two-graphs.

- 1 D is a ρ -regular subgraph, each vertex having degree

$$\rho = \frac{n}{2} - \frac{\lambda}{\alpha} - 1.$$

- 2 Substituting for α and λ , we also get that n and $\mu_1 + \mu_2$ have the same parity (even/odd).

Results and Descendants of a Regular Two-Graphs

Results and Descendants of a Regular Two-Graphs

We prove the first result, that D is ρ -regular with $\rho = \frac{n}{2} - \frac{\lambda}{\alpha} - 1$.

Results and Descendants of a Regular Two-Graphs

We prove the first result, that D is ρ -regular with $\rho = \frac{n}{2} - \frac{\lambda}{\alpha} - 1$.

Proof.

Let the Seidel eigenvalues of G be μ_1 and μ_2 , where G is in descendant form. Using the values of α and λ , the first and last rows of the involution \mathbf{M} are of the form

$$\begin{array}{l} \text{Row 1} \\ \text{Row } n \end{array} \begin{pmatrix} -\lambda & \pm\alpha & \pm\alpha & \cdots & \pm\alpha & \alpha \\ & & \vdots & & & \\ \alpha & \alpha & \alpha & \cdots & \alpha & -\lambda \end{pmatrix}$$

where we are assuming that the last row corresponds to the isolated vertex. The number of $-\alpha$'s in row 1 is the degree of vertex 1. Since $\mathbf{M}^2 = \mathbf{I}$, the inner product $\langle \text{Row 1}, \text{Row } n \rangle = 0$.



Results and Descendants of a Regular Two-Graphs

We prove the first result, that D is ρ -regular with $\rho = \frac{n}{2} - \frac{\lambda}{\alpha} - 1$.

Proof.

$$\begin{array}{l} \text{Row 1} \\ \text{Row } n \end{array} \begin{pmatrix} -\lambda & \pm\alpha & \pm\alpha & \cdots & \pm\alpha & \alpha \\ & & \vdots & & & \\ \alpha & \alpha & \alpha & \cdots & \alpha & -\lambda \end{pmatrix}$$

$$\langle \text{Row 1}, \text{Row } n \rangle = 0 \implies -\alpha\lambda - (n-2)\alpha^2 - 2\rho_1\alpha - \alpha\lambda = 0$$

where ρ_1 denotes the degree of vertex 1.

Note that ρ_1 is independent of the vertex label 1, since

$$\langle \text{Row 1}, \text{Row } i \rangle = 0$$

for all $1 \leq i \leq n-1$. Thus D is ρ -regular. □

Strongly Regular Graphs

Strongly Regular Graphs

Recall: A graph is called *regular* if all the vertices are of the same degree.

Strongly Regular Graphs

Recall: A graph is called *regular* if all the vertices are of the same degree.

Definition (Strongly regular graph)

Strongly Regular Graphs

Recall: A graph is called *regular* if all the vertices are of the same degree.

Definition (Strongly regular graph)

A graph G is said to be a strongly regular graph or an $\text{srg}(n, \rho, e, f)$ if:

Strongly Regular Graphs

Recall: A graph is called *regular* if all the vertices are of the same degree.

Definition (Strongly regular graph)

A graph G is said to be a strongly regular graph or an $\text{srg}(n, \rho, e, f)$ if:

- 1 it has n vertices,

Strongly Regular Graphs

Recall: A graph is called *regular* if all the vertices are of the same degree.

Definition (Strongly regular graph)

A graph G is said to be a strongly regular graph or an $\text{srg}(n, \rho, e, f)$ if:

- 1 it has n vertices,
- 2 each vertex has degree ρ ,

Strongly Regular Graphs

Recall: A graph is called *regular* if all the vertices are of the same degree.

Definition (Strongly regular graph)

A graph G is said to be a strongly regular graph or an $\text{srg}(n, \rho, e, f)$ if:

- 1 it has n vertices,
- 2 each vertex has degree ρ ,
- 3 every two adjacent vertices have e common neighbours, and

Strongly Regular Graphs

Recall: A graph is called *regular* if all the vertices are of the same degree.

Definition (Strongly regular graph)

A graph G is said to be a strongly regular graph or an $\text{srg}(n, \rho, e, f)$ if:

- 1 it has n vertices,
- 2 each vertex has degree ρ ,
- 3 every two adjacent vertices have e common neighbours, and
- 4 every two non-adjacent vertices have f common neighbours.


Strongly Regular Graphs

Example (Descendant of Petersen Graph)

Strongly Regular Graphs

Example (Descendant of Petersen Graph)

The descendant from the last example is an $\text{srg}(9, 4, 1, 2)$.



Structure of Descendants of Regular Two-Graphs

Structure of Descendants of Regular Two-Graphs

Consider a descendant form $D \dot{\cup} K_1$ of a regular two-graph, and for any two adjacent vertices, let \tilde{e} denote the number of common neighbours and let \tilde{e}^\sim denote the number of common non-neighbours.

Structure of Descendants of Regular Two-Graphs

Consider a descendant form $D \dot{\cup} K_1$ of a regular two-graph, and for any two adjacent vertices, let \tilde{e} denote the number of common neighbours and let \tilde{e} denote the number of common non-neighbours.

Similarly, for any two non-adjacent vertices, let \tilde{f} denote the number of common neighbours and let \tilde{f} denote the number of common non-neighbours.

Structure of Descendants of Regular Two-Graphs

Consider a descendant form $D \dot{\cup} K_1$ of a regular two-graph, and for any two adjacent vertices, let \tilde{e} denote the number of common neighbours and let $\tilde{\bar{e}}$ denote the number of common non-neighbours.

Similarly, for any two non-adjacent vertices, let \tilde{f} denote the number of common neighbours and let $\tilde{\bar{f}}$ denote the number of common non-neighbours.

By considering the rows of **M** we obtain the following formulæ:

- $\tilde{e} + \tilde{\bar{e}} = \frac{1}{2}(n-2) - \frac{\lambda}{\alpha}$
 $\tilde{e} - \tilde{\bar{e}} = 2\rho - n$
- $\tilde{f} + \tilde{\bar{f}} = \frac{1}{2}(n-2) + \frac{\lambda}{\alpha}$
 $\tilde{f} - \tilde{\bar{f}} = 2\rho - (n-2)$

Structure of Descendants of Regular Two-Graphs

Consider a descendant form $D \dot{\cup} K_1$ of a regular two-graph, and for any two adjacent vertices, let \tilde{e} denote the number of common neighbours and let $\tilde{\bar{e}}$ denote the number of common non-neighbours.

Similarly, for any two non-adjacent vertices, let \tilde{f} denote the number of common neighbours and let $\tilde{\bar{f}}$ denote the number of common non-neighbours.

By considering the rows of **M** we obtain the following formulæ:

- $\tilde{e} + \tilde{\bar{e}} = \frac{1}{2}(n-2) - \frac{\lambda}{\alpha}$
 $\tilde{e} - \tilde{\bar{e}} = 2\rho - n$
- $\tilde{f} + \tilde{\bar{f}} = \frac{1}{2}(n-2) + \frac{\lambda}{\alpha}$
 $\tilde{f} - \tilde{\bar{f}} = 2\rho - (n-2)$

From these it follows that \tilde{e} , $\tilde{\bar{e}}$, \tilde{f} and $\tilde{\bar{f}}$ are invariant for any pair of adjacent/non-adjacent vertices.

Structure of Descendants of Regular Two-Graphs

Structure of Descendants of Regular Two-Graphs

From the formulæ obtained previously, we get the following results.

Structure of Descendants of Regular Two-Graphs

From the formulæ obtained previously, we get the following results.

- Given a descendant form $D \dot{\cup} K_1$ of a regular two-graph on n vertices, then D is an $\text{srg}(n-1, \rho, e, f)$ where $e = \tilde{e}$ and $f = \tilde{f} = \frac{\rho}{2}$.

Structure of Descendants of Regular Two-Graphs

From the formulæ obtained previously, we get the following results.

- Given a descendant form $D \dot{\cup} K_1$ of a regular two-graph on n vertices, then D is an $\text{srg}(n-1, \rho, e, f)$ where $e = \tilde{e}$ and $f = \tilde{f} = \frac{\rho}{2}$.
- n must be even.

Structure of Descendants of Regular Two-Graphs

From the formulæ obtained previously, we get the following results.

- Given a descendant form $D \dot{\cup} K_1$ of a regular two-graph on n vertices, then D is an $\text{srg}(n-1, \rho, e, f)$ where $e = \tilde{e}$ and $f = \tilde{f} = \frac{\rho}{2}$.
- n must be even.
- $\frac{\lambda}{\alpha}$ is an integer.

Structure of Descendants of Regular Two-Graphs

From the formulæ obtained previously, we get the following results.

- Given a descendant form $D \dot{\cup} K_1$ of a regular two-graph on n vertices, then D is an $\text{srg}(n-1, \rho, e, f)$ where $e = \tilde{e}$ and $f = \tilde{f} = \frac{\rho}{2}$.
- n must be even.
- $\frac{\lambda}{\alpha}$ is an integer.

THE END