

Geometric Origin of CP Phases from Hyperbolic Holonomy

Marvin L. Gentry

the date of receipt and acceptance should be inserted later

Abstract We show that CP-violating phases arise naturally as holonomy invariants associated with flat $U(1)$ connections on compact hyperbolic 3-manifolds. If $M = \mathbb{H}^3/\Gamma$ admits an orientation-reversing isometry Θ , then Θ induces complex conjugation on $U(1)$ character representations of $\pi_1(M)$. Any closed geodesic whose holonomy is not ± 1 therefore provides a rephasing-invariant CP-odd observable. For flat connections these phases are topological invariants and are stable under all metric deformations preserving topological type. We formulate the argument entirely in representation-theoretic language and illustrate the mechanism using the Meyerhoff manifold (OrientableClosedCensus[1], vol = 0.9814, $H_1 = \mathbb{Z}/5$).

The geometric construction developed here provides the theoretical foundation for the companion results in [10,11], where the CKM and PMNS mixing matrices are derived explicitly from the K - and N -factors of the Iwasawa decomposition of compact hyperbolic 3-manifold holonomy [10,11].

1 Hyperbolic Setup

Let Γ be a torsion-free discrete group acting properly discontinuously and cocompactly on $\mathbb{H}^3$ by orientation-preserving isometries. Let

$$M = \mathbb{H}^3/\Gamma$$

be the resulting compact oriented hyperbolic 3-manifold.

Closed geodesics correspond to conjugacy classes $[\gamma] \subset \Gamma$ with γ hyperbolic.

Definition 1 (Flat $U(1)$ connection) A flat $U(1)$ connection on M is equivalently a character

$$\rho : \pi_1(M) \longrightarrow U(1).$$

The holonomy of a closed geodesic represented by γ is

$$W(\gamma) := \rho([\gamma]).$$

Because $U(1)$ is abelian, $W(\gamma)$ depends only on the free homotopy class.

2 Orientation Reversal and CP

Let $\Theta : M \rightarrow M$ be an orientation-reversing isometry. It induces an automorphism

$$\Theta_* : \pi_1(M) \rightarrow \pi_1(M).$$

Definition 2 (CP action on characters) Given a character ρ , define its CP-conjugate character by

$$\rho^\Theta(\gamma) = \overline{\rho(\Theta_*^{-1}\gamma)}.$$

Remark 1 The inverse Θ_*^{-1} appears because Θ induces a *pullback* of representations: if Θ_* is an automorphism of $\pi_1(M)$, the natural action on characters is by precomposition with Θ_*^{-1} , ensuring ρ^Θ is again a group homomorphism.

Lemma 1 (Holonomy conjugation) *Let ρ be a flat $U(1)$ character. For any $\gamma \in \pi_1(M)$,*

$$W_{\rho^\Theta}(\gamma) = \overline{W_\rho(\Theta_*^{-1}\gamma)}.$$

Proof By definition,

$$W_{\rho^\Theta}(\gamma) = \rho^\Theta(\gamma) = \overline{\rho(\Theta_*^{-1}\gamma)} = \overline{W_\rho(\Theta_*^{-1}\gamma)}.$$

Theorem 1 (Geometric CP violation) *Suppose:*

1. M admits an orientation-reversing isometry Θ ,
2. ρ is a flat $U(1)$ character,
3. there exists a hyperbolic element γ whose conjugacy class is fixed by Θ_* (i.e., $\Theta_*[\gamma] = [\gamma]$) and such that $W_\rho(\gamma) \neq \pm 1$.

Then $\arg W_\rho(\gamma)$ is a rephasing-invariant CP-odd observable.

Proof Because the conjugacy class of γ is fixed, we may choose a representative so there exists $\delta \in \pi_1(M)$ with $\Theta_*^{-1}(\gamma) = \delta\gamma\delta^{-1}$. Since $U(1)$ characters are conjugation-invariant, $W_\rho(\Theta_*^{-1}\gamma) = W_\rho(\gamma)$ exactly. Applying the CP action,

$$W_\rho(\gamma) \xrightarrow{\text{CP}} W_{\rho^\Theta}(\gamma) = \overline{W_\rho(\Theta_*^{-1}\gamma)} = \overline{W_\rho(\gamma)}.$$

Hence $\arg W_\rho(\gamma) \mapsto -\arg W_\rho(\gamma)$. If $W_\rho(\gamma) \neq \pm 1$, the phase is nonzero and flips sign under CP, so it is a genuine CP-odd observable. Rephasing invariance follows from Proposition 3.

3 Rephasing Invariance

Proposition 1 *Holonomy phases are invariant under all $U(1)$ gauge transformations.*

Proof Gauge transformations act by conjugation on ρ , but since $U(1)$ is abelian, conjugation is trivial.

Remark 2 The term *rephasing invariance* here refers to $U(1)$ gauge transformations of the flat connection, which are trivial for an abelian structure group. This is distinct from the sector-local rephasings of the companion papers [10, 11], which remove unphysical phases from fermion field definitions. Both notions confirm that $\arg W_\rho(\gamma)$ is invariant and carries physical information.

4 Topological Stability

Theorem 2 *Let ρ be a flat character. Then $W_\rho(\gamma)$ depends only on the homotopy class of γ . It is invariant under:*

1. metric deformations preserving the topological type of M ,
2. continuous deformations of ρ within its connected component in the character variety.

Proof Flatness implies holonomy depends only on the homotopy class. Metric deformations do not change $\pi_1(M)$, so the holonomy of each class remains unchanged. Continuous variations of ρ in character space vary the holonomy continuously; if they stay within the same component, the holonomy can change continuously but remains a topological invariant in the sense that its value is determined by the representation. For the purpose of CP violation, it suffices that the phase is topologically quantised (e.g., a root of unity) and does not drift under small metric perturbations.

Remark 3 (Discrete character variety) When $H_1(M; \mathbb{Z})$ is purely torsion—as for the Meyerhoff manifold with $H_1 = \mathbb{Z}/5$ —the character variety $\text{Hom}(\pi_1(M), U(1))$ consists of finitely many isolated points. For $\mathbb{Z}/5$ there are exactly five, corresponding to the fifth roots of unity. Each connected component is a single point, so no continuous deformation of ρ is possible; the holonomy phase is *topologically quantized* and cannot drift.

5 Example: The Meyerhoff Manifold

The Meyerhoff manifold $M = \text{OrientableClosedCensus}[1]$ (SnapPy [3], vol = 0.9814) is proven arithmetic with invariant trace field $\mathbb{Q}(\sqrt{-283})$ [4]. Its first homology group satisfies

$$H_1(M; \mathbb{Z}) \cong \mathbb{Z}/5.$$

The character variety consists of five isolated points; choose ρ sending the generator to $e^{2\pi i/5}$ and let γ represent the generator of $\mathbb{Z}/5$. Complex conjugation in $\mathbb{Q}(\sqrt{-283})$ induces an orientation-reversing isometry acting by inversion $k \mapsto -k$ on $H_1(M; \mathbb{Z}) = \mathbb{Z}/5$, mapping χ to its complex conjugate [5, 4]. The conjugacy class of γ is fixed, so Theorem 2.3 applies. Therefore

$$W_\rho(\gamma) = e^{2\pi i/5},$$

which is $\neq \pm 1$ and provides a concrete geometric CP-odd phase.

6 Conclusion

We have shown that CP-violating phases can arise from the geometry of compact hyperbolic 3-manifolds. The holonomy of a flat $U(1)$ character around a closed geodesic, when the manifold admits an orientation-reversing isometry that fixes the geodesic's conjugacy class, yields a rephasing-invariant observable that flips sign under CP. For flat connections these phases are topological invariants, stable under metric deformations. The Meyerhoff manifold provides an explicit example realising a fifth root of unity as a CP-odd phase. We note that the twist angles $\phi(\gamma) = \text{Im} \log \lambda(\gamma)$ in the A -factor of the Iwasawa decomposition [10, 11] are precisely the arguments of the flat $U(1)$ holonomies studied here; the $\mathbb{Z}/5$ torsion quantizes these phases, providing the geometric origin of CP violation in that framework. Embedding this mechanism into a complete flavor model remains open.

Data Availability

This manuscript contains no research data.

Conflict of Interest

The author declares no conflicts of interest.

References

1. G. C. Branco, L. Lavoura, and J. P. Silva, *CP Violation*, Oxford University Press, 1999.
2. M. R. Bridson and A. Haefliger, *Metric Spaces of Non-Positive Curvature*, Springer, 1999.
3. M. Culler, N. Dunfield, M. Goerner, and J. R. Weeks, "SnapPy, a computer program for studying the geometry and topology of 3-manifolds," <http://snappy.computop.org>.
4. T. Chinburg, E. Friedman, K. N. Jones, and A. W. Reid, "The arithmetic hyperbolic 3-manifold of smallest volume," *Ann. Scuola Norm. Sup. Pisa* **30**, 1–40 (2001).
5. C. Maclachlan and A. W. Reid, *The Arithmetic of Hyperbolic 3-Manifolds*, Springer, New York (2003).
6. C. Jarlskog, "A basis independent formulation of the connection between quark mass matrices, CP violation and experiment," *Z. Phys. C* **29** (1985), 491.
7. J. G. Ratcliffe, *Foundations of Hyperbolic Manifolds*, Springer, 2006.
8. W. P. Thurston, *Three-Dimensional Geometry and Topology*, Princeton University Press, 1997.
9. J. R. Weeks, "Computation of hyperbolic structures in knot theory," in *Handbook of Knot Theory*, Elsevier, 2005.
10. M. L. Gentry, CKM quark mixing from geodesic axes of hyperbolic 3-manifold holonomy, *Phys. Rev. D* (2026), submitted March 2026, temporary ID EPJC-26-03-349.
11. M. L. Gentry, Lepton mixing from Borel structure of hyperbolic holonomy, *Phys. Rev. D* (2026), submitted March 2026, temporary ID EPJC-26-03-351.