ESSAY · RESEARCH METHODS

Why Most Published Research Findings Are False

John P. A. Ioannidis

University of Ioannina School of Medicine, Ioannina, Greece
Tufts University School of Medicine, Boston, Massachusetts, United States

Originally published in PLoS Medicine 2(8): e124, 30 August 2005 · doi:10.1371/journal.pmed.0020124 · © 2005 John P. A. Ioannidis, Creative Commons Attribution License

SUMMARY

There is increasing concern that most current published research findings are false. The probability that a research claim is true may depend on study power and bias, the number of other studies on the same question, and, importantly, the ratio of true to no relationships among the relationships probed in each scientific field. In this framework, a research finding is less likely to be true when the studies conducted in a field are smaller; when effect sizes are smaller; when there is a greater number and lesser preselection of tested relationships; where there is greater flexibility in designs, definitions, outcomes, and analytical modes; when there is greater financial and other interest and prejudice; and when more teams are involved in a scientific field in chase of statistical significance. Simulations show that for most study designs and settings, it is more likely for a research claim to be false than true. Moreover, for many current scientific fields, claimed research findings may often be simply accurate measures of the prevailing bias. In this essay, I discuss the implications of these problems for the conduct and interpretation of research.

Published research findings are sometimes refuted by subsequent evidence, with ensuing confusion and disappointment. Refutation and controversy is seen across the range of research designs, from clinical trials and traditional epidemiological studies [1–3] to the most modern molecular research [4, 5]. There is increasing concern that in modern research, false findings may be the majority or even the vast majority of published research claims [6–8]. However, this should not be surprising. It can be proven that most claimed research findings are false. Here I will examine the key factors that influence this problem and some corollaries thereof.

Competing interests. The author has declared that no competing interests exist. **Abbreviation.** PPV, positive predictive value.

Modeling the Framework for False Positive Findings

Several methodologists have pointed out [9–11] that the high rate of nonreplication (lack of confirmation) of research discoveries is a consequence of the convenient, yet ill-founded strategy of claiming conclusive research findings solely on the basis of a single study assessed by formal statistical significance, typically for a *p*-value less than 0.05.

As has been shown previously, the probability that a research finding is indeed true depends on the prior probability of it being true (before doing the study), the statistical power of the study, and the level of statistical significance [10, 11]. Consider a 2 × 2 table in which research

findings are compared against the gold standard of true relationships in a scientific field. In a research field both true and false hypotheses can be made about the presence of relationships. Let R be the ratio of the number of "true relationships" to "no relationships" among those tested in the field. R is characteristic of the field and can vary a lot depending on whether the field targets highly likely relationships or searches for only one or a few true relationships among thousands and millions of hypotheses that may be postulated. Let us also consider, for computational simplicity, circumscribed fields where either there is only one true relationship (among many that can be hypothesized) or the power is similar to find any of the several existing true relationships. The pre-study probability of a relationship being true is $R/(R+1)$. The probability of a study finding a true relationship reflects the power $1-\beta$ (one minus the Type II error rate). The probability of claiming a relationship when none truly exists reflects the Type I error rate, α. Assuming that c relationships are being probed in the field, the expected values of the 2 × 2 table are given in Table 1. After a research finding has been claimed based on achieving formal statistical significance, the post-study probability that it is true is the positive predictive value, PPV. The PPV is also the complementary probability of what Wacholder et al. have called the false positive report probability [10]. According to the 2 × 2 table, one gets

$$\mathrm{PPV} = \frac{(1-\beta)R}{R-\beta R+\alpha}. \qquad (1)$$

A research finding is thus more likely true than false if $(1-\beta)R > \alpha$. Since usually the vast majority of investigators depend on $\alpha = 0.05$, this means that a research finding is more likely true than false if $(1-\beta)R > 0.05$.

What is less well appreciated is that bias and the extent of repeated independent testing by different teams of investigators around the globe may further distort this picture and may lead to even smaller probabilities of the research findings being indeed true. We will try to model these two factors in the context of similar 2 × 2 tables.

Bias

First, let us define bias as the combination of various design, data, analysis, and presentation factors that tend to produce research findings when they should not be produced. Let u be the proportion of probed analyses that would not have been "research findings," but nevertheless end up presented and reported as such, because of bias. Bias should not be confused with chance variability that causes some findings to be false by chance even though the study design, data, analysis, and presentation are perfect. Bias can entail manipulation in the analysis or reporting of findings. Selective or distorted reporting is a typical form of such bias. We may assume that u does not depend on whether a true relationship exists or not. This is not an unreasonable assumption, since typically it is impossible to know which relationships are indeed true. In the presence of bias (Table 2), one gets

$$\mathrm{PPV} = \frac{[1-\beta]R + u\beta R}{R+\alpha-\beta R+u-u\alpha+u\beta R}, \qquad (2)$$

and PPV decreases with increasing u, unless $1-\beta \leq \alpha$, i.e., $1-\beta \leq 0.05$ for most situations. Thus, with increasing bias, the chances that a research finding is true diminish considerably. This is shown for different levels of power and for different pre-study odds in Figure 1.

Testing by Several Independent Teams

Several independent teams may be addressing the same sets of research questions. As research efforts are globalized, it is practically the rule that several research teams, often dozens of them, may probe the same or similar questions. The probability that at least one study, among several done on the same question, claims a statistically significant research finding is easy to estimate. For n independent studies of equal power, the 2 × 2 table is shown in Table 3:

$$\mathrm{PPV} = \frac{R(1-\beta^n)}{R+1-[1-\alpha]^n - R\beta^n} \qquad (3)$$

(not considering bias). With increasing number of independent studies, PPV tends to decrease, unless $1-\beta < \alpha$,

Table 1. Research Findings and True Relationships

Research finding	True relationship		
	Yes	No	Total
Yes	$c(1-\beta)R/(R+1)$	$c\alpha/(R+1)$	$c(R+\alpha-\beta R)/(R+1)$
No	$c\beta R/(R+1)$	$c(1-\alpha)/(R+1)$	$c(1-\alpha+\beta R)/(R+1)$
Total	$cR/(R+1)$	$c/(R+1)$	c

Table 2. Research Findings and True Relationships in the Presence of Bias

Research finding	True relationship		
	Yes	No	Total
Yes	$(c[1-\beta]R+uc\beta R)/(R+1)$	$c\alpha+uc(1-\alpha)/(R+1)$	$c(R+\alpha-\beta R+u-u\alpha+u\beta R)/(R+1)$
No	$(1-u)c\beta R/(R+1)$	$(1-u)c(1-\alpha)/(R+1)$	$c(1-u)(1-\alpha+\beta R)/(R+1)$
Total	$cR/(R+1)$	$c/(R+1)$	c




Figure 1. PPV (Probability That a Research Finding Is True) as a Function of the Pre-Study Odds for Various Levels of Bias, u

Panels correspond to power of 0.80, 0.50, and 0.20. Drawn in code from Eq. (2) for the values of u in the 2005 legend, with no bias ($u = 0$) dashed; the curves printed in 2005 match $u = 0$, 0.05, 0.20 and 0.80.

Table 3. Research Findings and True Relationships in the Presence of Multiple Studies

Research finding	True relationship		
	Yes	No	Total
Yes	$cR(1-\beta^n)/(R+1)$	$c(1-[1-\alpha]^n)/(R+1)$	$c(R+1-[1-\alpha]^n-R\beta^n)/(R+1)$
No	$cR\beta^n/(R+1)$	$c(1-\alpha)^n/(R+1)$	$c([1-\alpha]^n+R\beta^n)/(R+1)$
Total	$cR/(R+1)$	$c/(R+1)$	c

i.e., typically $1-\beta < 0.05$. This is shown for different levels of power and for different pre-study odds in Figure 2. For n studies of different power, the term β^n is replaced by the product of the terms β_i for $i = 1$ to n, but inferences are similar.

Corollaries

A practical example is shown in Box 1. Based on the above considerations, one may deduce several interesting corollaries about the probability that a research finding is indeed true.

Corollary 1 **The smaller the studies conducted in a scientific field, the less likely the research findings are to be true.** Small sample size means smaller power and, for all functions above, the PPV for a true research finding decreases as power decreases towards $1-\beta = 0.05$. Thus, other factors being equal, research findings are more likely true in scientific fields that undertake large studies, such as randomized controlled trials in cardiology (several thousand subjects randomized) [12] than in scientific fields with small studies, such as most research of molecular predictors (sample sizes 100-fold smaller) [13].

Corollary 2 **The smaller the effect sizes in a scientific field, the less likely the research findings are to be true.** Power is also related to the effect size. Thus research findings are more likely true in scientific fields with large effects, such as the impact of smoking on cancer or cardiovascular disease (relative risks 3–20), than in




Figure 2. PPV (Probability That a Research Finding Is True) as a Function of the Pre-Study Odds for Various Numbers of Conducted Studies, n

Panels correspond to power of 0.80, 0.50, and 0.20. Drawn in code from Eq. (3).

scientific fields where postulated effects are small, such as genetic risk factors for multigenetic diseases (relative risks 1.1–1.5) [7]. Modern epidemiology is increasingly obliged to target smaller effect sizes [14]. Consequently, the proportion of true research findings is expected to decrease.

Corollary 3 **The greater the number and the lesser the selection of tested relationships in a scientific field, the less likely the research findings are to be true.** As shown above, the post-study probability that a finding is true (PPV) depends a lot on the pre-study odds (R). Thus, research findings are more likely true in confirmatory designs, such as large phase III randomized controlled trials, or meta-analyses thereof, than in hypothesis-generating experiments. Fields considered highly informative and creative given the wealth of the assembled and tested information, such as microarrays and other high-throughput discovery-oriented research [4, 8, 15], should have extremely low PPV.

Corollary 4 **The greater the flexibility in designs, definitions, outcomes, and analytical modes in a scientific field, the less likely the research findings are to be true.** Flexibility increases the potential for transforming what would be "negative" results into "positive" results, i.e., bias, u. For several research designs, e.g., randomized controlled trials [16–18] or meta-analyses [19, 20], there have been efforts to standardize their conduct and reporting. Adherence to common standards is likely to increase the proportion of true findings. Regardless, even in the most stringent research designs, bias seems to be a major problem. For example, there is strong evidence that selective outcome reporting, with manipulation of the outcomes and analyses reported, is a common problem even for randomized trials [21]. Simply abolishing selective publication would not make this problem go away.

Corollary 5 **The greater the financial and other interests and prejudices in a scientific field, the less**

Box 1. An Example: Science at Low Pre-Study Odds

Let us assume that a team of investigators performs a whole genome association study to test whether any of 100,000 gene polymorphisms are associated with susceptibility to schizophrenia. Based on what we know about the extent of heritability of the disease, it is reasonable to expect that probably around ten gene polymorphisms among those tested would be truly associated with schizophrenia, with relatively similar odds ratios around 1.3 for the ten or so polymorphisms and with a fairly similar power to identify any of them. Then $R = 10/100{,}000 = 10^{-4}$, and the pre-study probability for any polymorphism to be associated with schizophrenia is also $R/(R+1) = 10^{-4}$. Let us also suppose that the study has 60% power to find an association with an odds ratio of 1.3 at $\alpha = 0.05$. Then it can be estimated that if a statistically significant association is found with the p-value barely crossing the 0.05 threshold, the post-study probability that this is true increases about 12-fold compared with the pre-study probability, but it is still only 12×10^{-4}.

Now let us suppose that the investigators manipulate their design, analyses, and reporting so as to make more relationships cross the $p = 0.05$ threshold even though this would not have been crossed with a perfectly adhered to design and analysis and with perfect comprehensive reporting of the results, strictly according to the original study plan. In the presence of bias with $u = 0.10$, the post-study probability that a research finding is true is only 4.4×10^{-4}. Furthermore, even in the absence of any bias, when ten independent research teams perform similar experiments around the world, if one of them finds a formally statistically significant association, the probability that the research finding is true is only 1.5×10^{-4}, hardly any higher than the probability we had before any of this extensive research was undertaken!

likely the research findings are to be true. Conflicts of interest and prejudice may increase bias, u. Conflicts of interest are very common in biomedical research [22], and typically they are inadequately and sparsely reported [22, 23]. Prejudice may not necessarily have financial roots. Scientists in a given field may be prejudiced purely because of their belief in a scientific theory or commitment to their own findings. Prestigious investigators may suppress via the peer review process the appearance and dissemination of findings that refute their findings, thus condemning their field to perpetuate false dogma. Empirical evidence on expert opinion shows that it is extremely unreliable [24].

Corollary 6 **The hotter a scientific field (with more scientific teams involved), the less likely the research findings are to be true.** This seemingly paradoxical corollary follows because, as stated above, the PPV of isolated findings decreases when many teams of investigators are involved in the same field. This may explain why we occasionally see major excitement followed rapidly by severe disappointments in fields that draw wide attention. The term Proteus phenomenon has been coined to describe this phenomenon of rapidly alternating extreme research claims and extremely opposite refutations [25]. Empirical evidence suggests that this sequence of extreme opposites is very common in molecular genetics [25].

Most Research Findings Are False for Most Research Designs and for Most Fields

In the described framework, a PPV exceeding 50% is quite difficult to get. Table 4 provides the results of simulations using the formulas developed for the influence of power, ratio of true to non-true relationships, and bias, for various types of situations that may be characteristic of specific study designs and settings. A finding from a well-conducted, adequately powered randomized controlled trial starting with a 50% pre-study chance that the intervention is effective is eventually true about 85% of the time. A fairly similar performance is expected of a confirmatory meta-analysis of good-quality randomized trials: potential bias probably increases, but power and pre-test chances are higher compared to a single randomized trial. Conversely, a meta-analytic finding from inconclusive studies where pooling is used to "correct" the low power of single studies, is probably false if $R \leq 1{:}3$. Research findings from underpowered, early-phase clinical trials would be true about one in four times, or even less frequently if bias is present. Epidemiological studies of an exploratory nature perform even worse, especially when underpowered, but even well-powered epidemiological studies may have only a one in five chance being true, if $R = 1{:}10$. Finally, in discovery-oriented research with massive testing, where tested relationships exceed true ones 1,000-fold (e.g., 30,000 genes tested, of which 30 may be the true culprits) [26, 27], PPV for each claimed relationship is extremely low, even with considerable standardization of laboratory and statistical methods, outcomes, and reporting thereof to minimize bias.

Claimed Research Findings May Often Be Simply Accurate Measures of the Prevailing Bias

As shown, the majority of modern biomedical research is operating in areas with very low pre- and post-study probability for true findings. Let us suppose that in a research field there are no true findings at all to be discovered. In such a "null field," one would ideally expect all observed effect sizes to vary by chance around the null in the absence of bias. The extent that observed findings deviate from what is expected by chance alone would be simply a pure measure of the prevailing bias.

Table 4. PPV of Research Findings for Various Combinations of Power ($1 - \beta$), Ratio of True to Not-True Relationships (R), and Bias (u)

$1 - \beta$	R	u	Practical example	PPV
0.80	1:1	0.10	Adequately powered RCT with little bias and 1:1 pre-study odds	0.85
0.95	2:1	0.30	Confirmatory meta-analysis of good-quality RCTs	0.85
0.80	1:3	0.40	Meta-analysis of small inconclusive studies	0.41
0.20	1:5	0.20	Underpowered, but well-performed phase I/II RCT	0.23
0.20	1:5	0.80	Underpowered, poorly performed phase I/II RCT	0.17
0.80	1:10	0.30	Adequately powered exploratory epidemiological study	0.20
0.20	1:10	0.30	Underpowered exploratory epidemiological study	0.12
0.20	1:1,000	0.80	Discovery-oriented exploratory research with massive testing	0.0010
0.20	1:1,000	0.20	As in previous example, but with more limited bias (more standardized)	0.0015

The estimated PPVs (positive predictive values) are derived assuming $\alpha = 0.05$ for a single study; here they are computed from Eq. (2).
RCT, randomized controlled trial.

For example, let us suppose that no nutrients or dietary patterns are actually important determinants for the risk of developing a specific tumor. Let us also suppose that the scientific literature has examined 60 nutrients and claims all of them to be related to the risk of developing this tumor with relative risks in the range of 1.2 to 1.4 for the comparison of the upper to lower intake tertiles. Then the claimed effect sizes are simply measuring nothing else but the net bias that has been involved in the generation of this scientific literature. Claimed effect sizes are in fact the most accurate estimates of the net bias. It even follows that between "null fields," the fields that claim stronger effects (often with accompanying claims of medical or public health importance) are simply those that have sustained the worst biases.

For fields with very low PPV, the few true relationships would not distort this overall picture much. Even if a few relationships are true, the shape of the distribution of the observed effects would still yield a clear measure of the biases involved in the field. This concept totally reverses the way we view scientific results. Traditionally, investigators have viewed large and highly significant effects with excitement, as signs of important discoveries. Too large and too highly significant effects may actually be more likely to be signs of large bias in most fields of modern research. They should lead investigators to careful critical thinking about what might have gone wrong with their data, analyses, and results.

How Can We Improve the Situation?

Is it unavoidable that most research findings are false, or can we improve the situation? A major problem is that it is impossible to know with 100% certainty what the truth is in any research question. In this regard, the pure "gold" standard is unattainable. However, there are several approaches to improve the post-study probability.

Better powered evidence, e.g., large studies or low-bias meta-analyses, may help, as it comes closer to the unknown "gold" standard. However, large studies may still have biases and these should be acknowledged and avoided. Moreover, one should be cautious that extremely large studies may be more likely to find a formally statistical significant difference for a trivial effect that is not really meaningfully different from the null [28–30].

Second, most research questions are addressed by many teams, and it is misleading to emphasize the statistically significant findings of any single team. What matters is the totality of the evidence. Diminishing bias through enhanced research standards and curtailing of prejudices may also help. However, this may require a change in scientific mentality that might be difficult to achieve. In some research designs, efforts may also be more successful with upfront registration of studies, e.g., randomized trials [31]. Registration would pose a challenge for hypothesis-generating research. Some kind of registration or networking of data collections or investigators within fields may be more feasible than registration of each and every hypothesis-generating experiment. Regardless, even if we do not see a great deal of progress with registration of studies in other fields, the principles of developing and adhering to a protocol could be more widely borrowed from randomized controlled trials.

Finally, instead of chasing statistical significance, we should improve our understanding of the range of R values—the pre-study odds—where research efforts operate [10]. Before running an experiment, investigators should consider what they believe the chances are that they are testing a true rather than a non-true relationship. Speculated high R values may sometimes then be ascertained. As described above, whenever ethically acceptable, large studies with minimal bias should be performed on research findings that are considered relatively established, to see how often they are indeed confirmed. I suspect several established "classics" will fail the test [32].

Nevertheless, most new discoveries will continue to stem from hypothesis-generating research with low or very low pre-study odds. We should then acknowledge that statistical significance testing in the report of a single study gives only a partial picture, without knowing how much testing has been done outside the report and in the relevant field at large. Despite a large statistical literature for multiple testing corrections [33], usually it is impossible to decipher how much data dredging by the reporting authors or other research teams has preceded a reported research finding. Even if determining this were feasible, this would not inform us about the pre-study odds. Thus, it is unavoidable that one should make approximate assumptions on how many relationships are expected to be true among those probed across the relevant research fields and research designs. Even though these assumptions would be considerably subjective, they would still be very useful in interpreting research claims and putting them in context.

ARTICLE INFORMATION

This edition. Abridged from J. P. A. Ioannidis, "Why Most Published Research Findings Are False", *PLoS Medicine* 2(8): e124 (2005), doi:10.1371/journal.pmed.0020124, under the Creative Commons Attribution License. Some sentences and paragraphs are left out, the formulas are numbered, the tables and figures are redrawn and one typing slip is corrected. Set in Gelasio and Sofia Sans Semi Condensed (SIL OFL); formulas by MathJax.

REFERENCES

[1] Ioannidis JP, Haidich AB, Lau J. Any casualties in the clash of randomised and observational evidence? BMJ 2001;322:879–80.

[2] Lawlor DA, Davey Smith G, Kundu D, Bruckdorfer KR, Ebrahim S. Those confounded vitamins: what can we learn from the differences between observational versus randomised trial evidence? Lancet 2004;363:1724–7.

[3] Vandenbroucke JP. When are observational studies as credible as randomised trials? Lancet 2004;363:1728–31.

[4] Michiels S, Koscielny S, Hill C. Prediction of cancer outcome with microarrays: a multiple random validation strategy. Lancet 2005;365:488–92.

[5] Ioannidis JPA, Ntzani EE, Trikalinos TA, Contopoulos-Ioannidis DG. Replication validity of genetic association studies. Nat Genet 2001;29:306–9.

[6] Colhoun HM, McKeigue PM, Davey Smith G. Problems of reporting genetic associations with complex outcomes. Lancet 2003;361:865–72.

[7] Ioannidis JP. Genetic associations: false or true? Trends Mol Med 2003;9:135–8.

[8] Ioannidis JPA. Microarrays and molecular research: noise discovery? Lancet 2005;365:454–5.

[9] Sterne JA, Davey Smith G. Sifting the evidence—what's wrong with significance tests? BMJ 2001;322:226–31.

[10] Wacholder S, Chanock S, Garcia-Closas M, El Ghormli L, Rothman N. Assessing the probability that a positive report is false: an approach for molecular epidemiology studies. J Natl Cancer Inst 2004;96:434–42.

[11] Risch NJ. Searching for genetic determinants in the new millennium. Nature 2000;405:847–56.

[12] Yusuf S, Collins R, Peto R. Why do we need some large, simple randomized trials? Stat Med 1984;3:409–22.

[13] Altman DG, Royston P. What do we mean by validating a prognostic model? Stat Med 2000;19:453–73.

[14] Taubes G. Epidemiology faces its limits. Science 1995;269:164–9.

[15] Golub TR, Slonim DK, Tamayo P, Huard C, Gaasenbeek M, et al. Molecular classification of cancer: class discovery and class prediction by gene expression monitoring. Science 1999;286:531–7.

[16] Moher D, Schulz KF, Altman DG. The CONSORT statement: revised recommendations for improving the quality of reports of parallel-group randomised trials. Lancet 2001;357:1191–4.

[17] Ioannidis JP, Evans SJ, Gøtzsche PC, O'Neill RT, Altman DG, et al. Better reporting of harms in randomized trials: an extension of the CONSORT statement. Ann Intern Med 2004;141:781–8.

[18] International Conference on Harmonisation E9 Expert Working Group. ICH Harmonised Tripartite Guideline. Statistical principles for clinical trials. Stat Med 1999;18:1905–42.

[19] Moher D, Cook DJ, Eastwood S, Olkin I, Rennie D, et al. Improving the quality of reports of meta-analyses of randomised controlled trials: the QUOROM statement. Quality of Reporting of Meta-analyses. Lancet 1999;354:1896–900.

[20] Stroup DF, Berlin JA, Morton SC, Olkin I, Williamson GD, et al. Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. JAMA 2000;283:2008–12.

[21] Chan AW, Hróbjartsson A, Haahr MT, Gøtzsche PC, Altman DG. Empirical evidence for selective reporting of outcomes in randomized trials: comparison of protocols to published articles. JAMA 2004;291:2457–65.

[22] Krimsky S, Rothenberg LS, Stott P, Kyle G. Scientific journals and their authors' financial interests: a pilot study. Psychother Psychosom 1998;67:194–201.

[23] Papanikolaou GN, Baltogianni MS, Contopoulos-Ioannidis DG, Haidich AB, Giannakakis IA, et al. Reporting of conflicts of interest in guidelines of preventive and therapeutic interventions. BMC Med Res Methodol 2001;1:3.

[24] Antman EM, Lau J, Kupelnick B, Mosteller F, Chalmers TC. A comparison of results of meta-analyses of randomized control trials and recommendations of clinical experts. Treatments for myocardial infarction. JAMA 1992;268:240–8.

[25] Ioannidis JP, Trikalinos TA. Early extreme contradictory estimates may appear in published research: the Proteus phenomenon in molecular genetics research and randomized trials. J Clin Epidemiol 2005;58:543–9.

[26] Ntzani EE, Ioannidis JP. Predictive ability of DNA microarrays for cancer outcomes and correlates: an empirical assessment. Lancet 2003;362:1439–44.

[27] Ransohoff DF. Rules of evidence for cancer molecular-marker discovery and validation. Nat Rev Cancer 2004;4:309–14.

[28] Lindley DV. A statistical paradox. Biometrika 1957;44:187–92.

[29] Bartlett MS. A comment on D. V. Lindley's statistical paradox. Biometrika 1957;44:533–4.

[30] Senn SJ. Two cheers for P-values? J Epidemiol Biostat 2001;6:193–204.

[31] De Angelis C, Drazen JM, Frizelle FA, Haug C, Hoey J, et al. Clinical trial registration: a statement from the International Committee of Medical Journal Editors. N Engl J Med 2004;351:1250–1.

[32] Ioannidis JPA. Contradicted and initially stronger effects in highly cited clinical research. JAMA 2005;294:218–28.

[33] Hsueh HM, Chen JJ, Kodell RL. Comparison of methods for estimating the number of true null hypotheses in multiplicity testing. J Biopharm Stat 2003;13:675–89.