ICPSR | INTER-UNIVERSITY CONSORTIUM FOR POLITICAL AND SOCIAL RESEARCH

ICPSR 36800

Research on Pathways to Desistance [Maricopa County, AZ and Philadelphia County, PA]: Subject Measures - Scales, 2000-2010

Edward P. Mulvey
University of Pittsburgh

Codebook for Substance Abuse

Inter-university Consortium for
Political and Social Research
P.O. Box 1248
Ann Arbor, Michigan 48106
www.icpsr.umich.edu

Terms of Use

The terms of use for this study can be found at:
http://www.icpsr.umich.edu/icpsrweb/ICPSR/studies/36800/terms

Information about Copyrighted Content

Some instruments administered as part of this study may contain in whole or substantially in part contents from copyrighted instruments. Reproductions of the instruments are provided as documentation for the analysis of the data associated with this collection. Restrictions on "fair use" apply to all copyrighted content. More information about the reproduction of copyrighted works by educators and librarians is available from the United States Copyright Office.

NOTICE
WARNING CONCERNING COPYRIGHT RESTRICTIONS

The copyright law of the United States (Title 17, United States Code) governs the making of photocopies or other reproductions of copyrighted material. Under certain conditions specified in the law, libraries and archives are authorized to furnish a photocopy or other reproduction. One of these specified conditions is that the photocopy or reproduction is not to be "used for any purpose other than private study, scholarship, or research." If a user makes a request for, or later uses, a photocopy or reproduction for purposes in excess of "fair use," that user may be liable for copyright infringement.

ICPSR 36800

Research on Pathways to Desistance [Maricopa County, AZ and Philadelphia County, PA]: Subject Measures - Scales, 2000-2010

Variable Description and Frequencies

Note: Frequencies displayed for the variables are not weighted. They are purely descriptive and may not be representative of the study population. Please review any sampling or weighting information available with the study.

Summary statistics (minimum, maximum, mean, median, and standard deviation) may not be available for every variable in the codebook. Conversely, a listing of frequencies in table format may not be present for every variable in the codebook either. However, all variables in the dataset are present and display sufficient information about each variable. These decisions are made intentionally and are at the discretion of the archive producing this codebook.

Substance Abuse

CASEID: S0 IntInfo: Subject ID

Based upon 1,354 valid cases out of 1,354 total cases.

Location: 1-5 (width: 5; decimal: 0)
Variable Type: numeric

S0SUBUSE: S0 Subuse: Whats most EVER used alcohol

Value	Label	Unweighted Frequency	%
1	Not at all/not used in lifetime	260	19.2 %
2	1-2 times	217	16.0 %
3	Less than 1x per month	128	9.5 %
4	Once per month	91	6.7 %
5	2-3 times per month	133	9.8 %
6	Once per week	115	8.5 %
7	2-3 times per week	168	12.4 %
8	4-5 times per week	107	7.9 %
9	Everyday	131	9.7 %
	Missing Data		
-200	Data missing	4	0.3 %
	Total	**1,354**	**100%**

Based upon 1,350 valid cases out of 1,354 total cases.

Location: 6-9 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -99 , -98

S0SUBUS2: S0 Subuse: Past 6 months how often used alcohol

Value	Label	Unweighted Frequency	%
1	Not at all	633	46.8 %
2	1-2 times	207	15.3 %
3	Less than 1x per month	111	8.2 %
4	Once per month	51	3.8 %
5	2-3 times per month	80	5.9 %
6	Once per week	73	5.4 %
7	2-3 times per week	109	8.1 %
8	4-5 times per week	47	3.5 %
9	Everyday	38	2.8 %
	Missing Data		
-200	Data missing	4	0.3 %

Value	Label	Unweighted Frequency	%
-99	Dont Know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,349 valid cases out of 1,354 total cases.

Location: 10-13 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -101 , -99 , -98

S0SUBUS3: S0 Subuse: How many drinks do you usually have

Based upon 1,348 valid cases out of 1,354 total cases.

- Mean: 3.76
- Median: 1.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 65.00
- Standard Deviation: 6.13

Location: 14-17 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -102 , -101 , -99 , -98

S0SUBUS4: S0 Subuse: Past 6 months how often had 5 or more drinks at time

Value	Label	Unweighted Frequency	%
1	Not at all	846	62.5 %
2	1-2 times	149	11.0 %
3	Less than 1x per month	74	5.5 %
4	Once per month	36	2.7 %
5	2-3 times per month	51	3.8 %
6	Once per week	54	4.0 %
7	2-3 times per week	74	5.5 %
8	4-5 times per week	38	2.8 %
9	Everyday	27	2.0 %
	Missing Data		
-200	Data missing	4	0.3 %
-99	Dont Know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,349 valid cases out of 1,354 total cases.

Location: 18-21 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -102 , -101 , -99 , -98

S0SUBUS5: S0 Subuse: Past 6 months num times been drunk on alcohol

Value	Label	Unweighted Frequency	%
1	Not at all	806	59.5 %
2	1-2 times	188	13.9 %
3	Less than 1x per month	114	8.4 %
4	Once per month	27	2.0 %
5	2-3 times per month	42	3.1 %
6	Once per week	48	3.5 %
7	2-3 times per week	70	5.2 %
8	4-5 times per week	39	2.9 %
9	Everyday	16	1.2 %
	Missing Data		
-200	Data missing	4	0.3 %
	Total	**1,354**	**100%**

Based upon 1,350 valid cases out of 1,354 total cases.

Location: 22-25 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -102 , -101 , -99 , -98

S0SUBUS6: S0 Subuse: Whats most EVER used marijuana

Value	Label	Unweighted Frequency	%
1	Not at all/not used in lifetime	202	14.9 %
2	1-2 times	131	9.7 %
3	Less than 1x per month	34	2.5 %
4	Once per month	31	2.3 %
5	2-3 times per month	48	3.5 %
6	Once per week	38	2.8 %
7	2-3 times per week	86	6.4 %
8	4-5 times per week	111	8.2 %
9	Everyday	669	49.4 %
	Missing Data		
-200	Data missing	4	0.3 %
	Total	**1,354**	**100%**

Based upon 1,350 valid cases out of 1,354 total cases.

Location: 26-29 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -99 , -98

S0SUBUS7: S0 Subuse: Past 6 months num times used marijuana

Value	Label	Unweighted Frequency	%
1	Not at all	581	42.9 %
2	1-2 times	111	8.2 %
3	Less than 1x per month	60	4.4 %
4	Once per month	25	1.8 %
5	2-3 times per month	53	3.9 %
6	Once per week	32	2.4 %
7	2-3 times per week	83	6.1 %
8	4-5 times per week	109	8.1 %
9	Everyday	295	21.8 %
	Missing Data		
-200	Data missing	4	0.3 %
-99	Dont Know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,349 valid cases out of 1,354 total cases.

Location: 30-33 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -103 , -99 , -98

S0SUBUS8: S0 Subuse: Most EVER used sedatives to get high

Value	Label	Unweighted Frequency	%
1	Not at all/not used in lifetime	1064	78.6 %
2	1-2 times	92	6.8 %
3	Less than 1x per month	27	2.0 %
4	Once per month	32	2.4 %
5	2-3 times per month	29	2.1 %
6	Once per week	17	1.3 %
7	2-3 times per week	43	3.2 %
8	4-5 times per week	19	1.4 %
9	Everyday	26	1.9 %
	Missing Data		
-200	Data missing	4	0.3 %
-99	Dont Know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,349 valid cases out of 1,354 total cases.

Location: 34-37 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -99 , -98

S0SUBUS9: S0 Subuse: Past 6 months num times used sedatives

Value	Label	Unweighted Frequency	%
1	Not at all	1194	88.2 %
2	1-2 times	58	4.3 %
3	Less than 1x per month	17	1.3 %
4	Once per month	16	1.2 %
5	2-3 times per month	17	1.3 %
6	Once per week	8	0.6 %
7	2-3 times per week	22	1.6 %
8	4-5 times per week	8	0.6 %
9	Everyday	10	0.7 %
	Missing Data		
-200	Data missing	4	0.3 %
	Total	**1,354**	**100%**

Based upon 1,350 valid cases out of 1,354 total cases.

Location: 38-41 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -103 , -99 , -98

S0SUBU10: S0 Subuse: Most EVER used stimulants to get high

Value	Label	Unweighted Frequency	%
1	Not at all/not used in lifetime	1144	84.5 %
2	1-2 times	53	3.9 %
3	Less than 1x per month	14	1.0 %
4	Once per month	8	0.6 %
5	2-3 times per month	9	0.7 %
6	Once per week	13	1.0 %
7	2-3 times per week	23	1.7 %
8	4-5 times per week	19	1.4 %
9	Everyday	67	4.9 %
	Missing Data		
-200	Data missing	4	0.3 %
	Total	**1,354**	**100%**

Based upon 1,350 valid cases out of 1,354 total cases.

Location: 42-45 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -99 , -98

S0SUBU11: S0 Subuse: Past 6 months num times used stimulants

Value	Label	Unweighted Frequency	%
1	Not at all	1228	90.7 %
2	1-2 times	23	1.7 %
3	Less than 1x per month	16	1.2 %
4	Once per month	4	0.3 %
5	2-3 times per month	8	0.6 %
6	Once per week	9	0.7 %
7	2-3 times per week	16	1.2 %
8	4-5 times per week	22	1.6 %
9	Everyday	24	1.8 %
	Missing Data		
-200	Data missing	4	0.3 %
	Total	**1,354**	**100%**

Based upon 1,350 valid cases out of 1,354 total cases.

Location: 46-49 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -103 , -99 , -98

S0SUBU12: S0 Subuse: Most EVER used cocaine

Value	Label	Unweighted Frequency	%
1	Not at all/not used in lifetime	1035	76.4 %
2	1-2 times	110	8.1 %
3	Less than 1x per month	21	1.6 %
4	Once per month	21	1.6 %
5	2-3 times per month	35	2.6 %
6	Once per week	25	1.8 %
7	2-3 times per week	42	3.1 %
8	4-5 times per week	22	1.6 %
9	Everyday	39	2.9 %
	Missing Data		
-200	Data missing	4	0.3 %
	Total	**1,354**	**100%**

Based upon 1,350 valid cases out of 1,354 total cases.

Location: 50-53 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -99 , -98

S0SUBU13: S0 Subuse: Past 6 months num times used cocaine

Value	Label	Unweighted Frequency	%
1	Not at all	1183	87.4 %
2	1-2 times	50	3.7 %
3	Less than 1x per month	29	2.1 %
4	Once per month	9	0.7 %
5	2-3 times per month	15	1.1 %
6	Once per week	19	1.4 %
7	2-3 times per week	22	1.6 %
8	4-5 times per week	10	0.7 %
9	Everyday	13	1.0 %
	Missing Data		
-200	Data missing	4	0.3 %
	Total	**1,354**	**100%**

Based upon 1,350 valid cases out of 1,354 total cases.

Location: 54-57 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -103 , -99 , -98

S0SUBU14: S0 Subuse: Most EVER used Opiates

Value	Label	Unweighted Frequency	%
1	Not at all/not used in lifetime	1248	92.2 %
2	1-2 times	42	3.1 %
3	Less than 1x per month	15	1.1 %
4	Once per month	9	0.7 %
5	2-3 times per month	10	0.7 %
6	Once per week	5	0.4 %
7	2-3 times per week	7	0.5 %
8	4-5 times per week	8	0.6 %
9	Everyday	6	0.4 %
	Missing Data		
-200	Data missing	4	0.3 %
	Total	**1,354**	**100%**

Based upon 1,350 valid cases out of 1,354 total cases.

Location: 58-61 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -99 , -98

S0SUBU15: S0 Subuse: Past 6 months num times used opiates

Value	Label	Unweighted Frequency	%
1	Not at all	1300	96.0 %
2	1-2 times	21	1.6 %
3	Less than 1x per month	7	0.5 %
4	Once per month	6	0.4 %
5	2-3 times per month	4	0.3 %
6	Once per week	2	0.1 %
7	2-3 times per week	5	0.4 %
8	4-5 times per week	5	0.4 %
9	Everyday	0	0.0 %
	Missing Data		
-200	Data missing	4	0.3 %
	Total	**1,354**	**100%**

Based upon 1,350 valid cases out of 1,354 total cases.

Location: 62-65 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -103 , -99 , -98

S0SUBU16: S0 Subuse: Most EVER used ecstasy to get high

Value	Label	Unweighted Frequency	%
1	Not at all/not used in lifetime	1130	83.5 %
2	1-2 times	99	7.3 %
3	Less than 1x per month	28	2.1 %
4	Once per month	16	1.2 %
5	2-3 times per month	22	1.6 %
6	Once per week	15	1.1 %
7	2-3 times per week	24	1.8 %
8	4-5 times per week	6	0.4 %
9	Everyday	10	0.7 %
	Missing Data		
-200	Data missing	4	0.3 %
	Total	**1,354**	**100%**

Based upon 1,350 valid cases out of 1,354 total cases.

Location: 66-69 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -99 , -98

S0SUBU17: S0 Subuse: Past 6 months num times used ecstasy

Value	Label	Unweighted Frequency	%
1	Not at all	1240	91.6 %
2	1-2 times	57	4.2 %
3	Less than 1x per month	12	0.9 %
4	Once per month	11	0.8 %
5	2-3 times per month	13	1.0 %
6	Once per week	4	0.3 %
7	2-3 times per week	11	0.8 %
8	4-5 times per week	1	0.1 %
9	Everyday	1	0.1 %
	Missing Data		
-200	Data missing	4	0.3 %
	Total	**1,354**	**100%**

Based upon 1,350 valid cases out of 1,354 total cases.

Location: 70-73 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -103 , -99 , -98

S0SUBU18: S0 Subuse: Most EVER used hallucinogens

Value	Label	Unweighted Frequency	%
1	Not at all/not used in lifetime	983	72.6 %
2	1-2 times	142	10.5 %
3	Less than 1x per month	39	2.9 %
4	Once per month	14	1.0 %
5	2-3 times per month	38	2.8 %
6	Once per week	31	2.3 %
7	2-3 times per week	37	2.7 %
8	4-5 times per week	22	1.6 %
9	Everyday	44	3.2 %
	Missing Data		
-200	Data missing	4	0.3 %
	Total	**1,354**	**100%**

Based upon 1,350 valid cases out of 1,354 total cases.

Location: 74-77 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -99 , -98

S0SUBU19: S0 Subuse: Past 6 months num times used hallucinogens

Value	Label	Unweighted Frequency	%
1	Not at all	1193	88.1 %
2	1-2 times	57	4.2 %
3	Less than 1x per month	22	1.6 %
4	Once per month	11	0.8 %
5	2-3 times per month	17	1.3 %
6	Once per week	11	0.8 %
7	2-3 times per week	12	0.9 %
8	4-5 times per week	15	1.1 %
9	Everyday	12	0.9 %
	Missing Data		
-200	Data missing	4	0.3 %
	Total	**1,354**	**100%**

Based upon 1,350 valid cases out of 1,354 total cases.

Location: 78-81 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -103 , -99 , -98

S0SUBU20: S0 Subuse: Most EVER used inhalants to get high

Value	Label	Unweighted Frequency	%
1	Not at all/not used in lifetime	1170	86.4 %
2	1-2 times	89	6.6 %
3	Less than 1x per month	19	1.4 %
4	Once per month	9	0.7 %
5	2-3 times per month	17	1.3 %
6	Once per week	5	0.4 %
7	2-3 times per week	14	1.0 %
8	4-5 times per week	7	0.5 %
9	Everyday	20	1.5 %
	Missing Data		
-200	Data missing	4	0.3 %
	Total	**1,354**	**100%**

Based upon 1,350 valid cases out of 1,354 total cases.

Location: 82-85 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -99 , -98

S0SUBU21: S0 Subuse: Past 6 months num times used inhalants

Value	Label	Unweighted Frequency	%
1	Not at all	1297	95.8 %
2	1-2 times	22	1.6 %
3	Less than 1x per month	4	0.3 %
4	Once per month	3	0.2 %
5	2-3 times per month	8	0.6 %
6	Once per week	4	0.3 %
7	2-3 times per week	4	0.3 %
8	4-5 times per week	6	0.4 %
9	Everyday	2	0.1 %
	Missing Data		
-200	Data missing	4	0.3 %
	Total	**1,354**	**100%**

Based upon 1,350 valid cases out of 1,354 total cases.

Location: 86-89 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -103 , -99 , -98

S0SUBU22: S0 Subuse: Most EVER used amyl nitrate odorizers

Value	Label	Unweighted Frequency	%
1	Not at all/not used in lifetime	1297	95.8 %
2	1-2 times	29	2.1 %
3	Less than 1x per month	4	0.3 %
4	Once per month	3	0.2 %
5	2-3 times per month	4	0.3 %
6	Once per week	2	0.1 %
7	2-3 times per week	2	0.1 %
8	4-5 times per week	3	0.2 %
9	Everyday	6	0.4 %
	Missing Data		
-200	Data missing	4	0.3 %
	Total	**1,354**	**100%**

Based upon 1,350 valid cases out of 1,354 total cases.

Location: 90-93 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -99 , -98

S0SUBU23: S0 Subuse: Past 6 months num times used amyl nitrate

Value	Label	Unweighted Frequency	%
1	Not at all	1328	98.1 %
2	1-2 times	7	0.5 %
3	Less than 1x per month	2	0.1 %
4	Once per month	3	0.2 %
5	2-3 times per month	2	0.1 %
6	Once per week	3	0.2 %
7	2-3 times per week	1	0.1 %
8	4-5 times per week	2	0.1 %
9	Everyday	2	0.1 %
	Missing Data		
-200	Data missing	4	0.3 %
	Total	**1,354**	**100%**

Based upon 1,350 valid cases out of 1,354 total cases.

Location: 94-97 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -103 , -99 , -98

S0SUBU24: S0 Subuse: Most EVER used any OTHER drugs

Value	Label	Unweighted Frequency	%
1	Not at all/not used in lifetime	1335	98.6 %
2	1-2 times	6	0.4 %
3	Less than 1x per month	3	0.2 %
4	Once per month	0	0.0 %
5	2-3 times per month	1	0.1 %
6	Once per week	1	0.1 %
7	2-3 times per week	1	0.1 %
8	4-5 times per week	2	0.1 %
9	Everyday	1	0.1 %
	Missing Data		
-200	Data missing	4	0.3 %
	Total	**1,354**	**100%**

Based upon 1,350 valid cases out of 1,354 total cases.

Location: 98-101 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -99 , -98

S0SUBU35: S0 Subuse: Most EVER used prescribed medications (own or someone elses)

Value	Label	Unweighted Frequency	%
1	Not at all/not used in lifetime	0	0.0 %
2	1-2 times	1	0.1 %
3	Less than 1x per month	0	0.0 %
4	Once per month	1	0.1 %
5	2-3 times per month	0	0.0 %
6	Once per week	0	0.0 %
7	2-3 times per week	2	0.1 %
8	4-5 times per week	0	0.0 %
9	Everyday	0	0.0 %
	Missing Data		
-800	Data unavailable	1346	99.4 %
-200	Data missing	4	0.3 %
	Total	**1,354**	**100%**

Based upon 4 valid cases out of 1,354 total cases.

Location: 102-105 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -200

S0SUBU25: S0 Subuse: Past 6 months used any other things get high

Value	Label	Unweighted Frequency	%
1	Yes	6	0.4 %
5	No	1344	99.3 %
	Missing Data		
-200	Data missing	4	0.3 %
	Total	**1,354**	**100%**

Based upon 1,350 valid cases out of 1,354 total cases.

Location: 106-109 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -103 , -9 , -8

S0SUBU36: S0 Subuse: Past 6 months num times used prescribed medications (own or someone elses)

Value	Label	Unweighted Frequency	%
1	Not at all/not used in lifetime	1	0.1 %
2	1-2 times	0	0.0 %
3	Less than 1x per month	0	0.0 %
4	Once per month	0	0.0 %
5	2-3 times per month	0	0.0 %

Value	Label	Unweighted Frequency	%
6	Once per week	0	0.0 %
7	2-3 times per week	1	0.1 %
8	4-5 times per week	1	0.1 %
9	Everyday	0	0.0 %
	Missing Data		
-800	Data unavailable	1347	99.5 %
-200	Data missing	4	0.3 %
	Total	**1,354**	**100%**

Based upon 3 valid cases out of 1,354 total cases.

Location: 110-113 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -200

S0SUBU26: S0 Subuse: Past 6 months num times used other to get high

Value	Label	Unweighted Frequency	%
1	Not at all	1344	99.3 %
2	1-2 times	1	0.1 %
3	Less than 1x per month	1	0.1 %
4	Once per month	0	0.0 %
5	2-3 times per month	0	0.0 %
6	Once per week	1	0.1 %
7	2-3 times per week	2	0.1 %
8	4-5 times per week	0	0.0 %
9	Everyday	1	0.1 %
	Missing Data		
-200	Data missing	4	0.3 %
	Total	**1,354**	**100%**

Based upon 1,350 valid cases out of 1,354 total cases.

Location: 114-117 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -104 , -103 , -99 , -98

S0SUBU27: S0 Subuse: Used any drugs past 24 hours

Value	Label	Unweighted Frequency	%
1	Yes	59	4.4 %
5	No	1291	95.3 %
	Missing Data		
-200	Data missing	4	0.3 %

Value	Label	Unweighted Frequency	%
	Total	**1,354**	**100%**

Based upon 1,350 valid cases out of 1,354 total cases.

Location: 118-121 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -9 , -8

S0SUBU01: S0 Subuse: Drug 1 used in past 24 hours

Value	Label	Unweighted Frequency	%
11	Marijuana/hashish	51	3.8 %
12	Sedatives/tranquilizers (valium,xanax, etc.)	3	0.2 %
13	Stimulants/amphetamines (diet pills, methamphetamine, etc.)	2	0.1 %
14	Cocaine	1	0.1 %
15	Opiates	0	0.0 %
16	Ecstasy	0	0.0 %
17	Hallucinogens (acid, LSD, etc.)	1	0.1 %
18	Inhalants (glue, paint, etc.)	0	0.0 %
19	Amyl nitrate/poppers/odorizers	0	0.0 %
96	Other (specify)	0	0.0 %
	Missing Data		
-200	Data missing	4	0.3 %
-106	Hasnt used < 11 drugs in p24hr - question skipped	1	0.1 %
-105	Hasnt used other drug past 24hr - question skipped	1291	95.3 %
	Total	**1,354**	**100%**

Based upon 58 valid cases out of 1,354 total cases.

Location: 122-125 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -106 , -105 , -99 , -98

S0SUBU02: S0 Subuse: Drug 2 used in past 24 hours

Value	Label	Unweighted Frequency	%
11	Marijuana/hashish	1	0.1 %
12	Sedatives/tranquilizers (valium,xanax, etc.)	1	0.1 %
13	Stimulants/amphetamines (diet pills, methamphetamine, etc.)	2	0.1 %
14	Cocaine	0	0.0 %
15	Opiates	0	0.0 %
16	Ecstasy	0	0.0 %
17	Hallucinogens (acid, LSD, etc.)	0	0.0 %
18	Inhalants (glue, paint, etc.)	0	0.0 %

Value	Label	Unweighted Frequency	%
19	Amyl nitrate/poppers/odorizers	0	0.0 %
96	Other (specify)	0	0.0 %
	Missing Data		
-200	Data missing	4	0.3 %
-106	Hasnt used < 11 drugs in p24hr - question skipped	55	4.1 %
-105	Hasnt used other drug past 24hr - question skipped	1291	95.3 %
	Total	**1,354**	**100%**

Based upon 4 valid cases out of 1,354 total cases.

Location: 126-129 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -106 , -105 , -99 , -98

S0SUBU03: S0 Subuse: Drug 3 used in past 24 hours

Value	Label	Unweighted Frequency	%
11	Marijuana/hashish	0	0.0 %
12	Sedatives/tranquilizers (valium,xanax, etc.)	0	0.0 %
13	Stimulants/amphetamines (diet pills, methamphetamine, etc.)	0	0.0 %
14	Cocaine	0	0.0 %
15	Opiates	1	0.1 %
16	Ecstasy	0	0.0 %
17	Hallucinogens (acid, LSD, etc.)	0	0.0 %
18	Inhalants (glue, paint, etc.)	0	0.0 %
19	Amyl nitrate/poppers/odorizers	0	0.0 %
96	Other (specify)	0	0.0 %
	Missing Data		
-200	Data missing	4	0.3 %
-106	Hasnt used < 11 drugs in p24hr - question skipped	58	4.3 %
-105	Hasnt used other drug past 24hr - question skipped	1291	95.3 %
	Total	**1,354**	**100%**

Based upon 1 valid cases out of 1,354 total cases.

Location: 130-133 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -106 , -105 , -99 , -98

S0SUBU04: S0 Subuse: Drug 4 used in past 24 hours

Value	Label	Unweighted Frequency	%
11	Marijuana/hashish	0	0.0 %
12	Sedatives/tranquilizers (valium,xanax, etc.)	0	0.0 %

Value	Label	Unweighted Frequency	%
13	Stimulants/amphetamines (diet pills, methamphetamine, etc.)	0	0.0 %
14	Cocaine	0	0.0 %
15	Opiates	0	0.0 %
16	Ecstasy	0	0.0 %
17	Hallucinogens (acid, LSD, etc.)	0	0.0 %
18	Inhalants (glue, paint, etc.)	0	0.0 %
19	Amyl nitrate/poppers/odorizers	0	0.0 %
96	Other (specify)	0	0.0 %
	Missing Data		
-200	Data missing	4	0.3 %
-106	Hasnt used < 11 drugs in p24hr - question skipped	59	4.4 %
-105	Hasnt used other drug past 24hr - question skipped	1291	95.3 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 134-137 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -106 , -105 , -99 , -98

S0SUBU05: S0 Subuse: Drug 5 used in past 24 hours

Value	Label	Unweighted Frequency	%
11	Marijuana/hashish	0	0.0 %
12	Sedatives/tranquilizers (valium,xanax, etc.)	0	0.0 %
13	Stimulants/amphetamines (diet pills, methamphetamine, etc.)	0	0.0 %
14	Cocaine	0	0.0 %
15	Opiates	0	0.0 %
16	Ecstasy	0	0.0 %
17	Hallucinogens (acid, LSD, etc.)	0	0.0 %
18	Inhalants (glue, paint, etc.)	0	0.0 %
19	Amyl nitrate/poppers/odorizers	0	0.0 %
96	Other (specify)	0	0.0 %
	Missing Data		
-200	Data missing	4	0.3 %
-106	Hasnt used < 11 drugs in p24hr - question skipped	59	4.4 %
-105	Hasnt used other drug past 24hr - question skipped	1291	95.3 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 138-141 (width: 4; decimal: 0)
Variable Type: numeric

(Range of) Missing Values: -200 , -106 , -105 , -99 , -98

S0SUBU06: S0 Subuse: Drug 6 used in past 24 hours

Value	Label	Unweighted Frequency	%
11	Marijuana/hashish	0	0.0 %
12	Sedatives/tranquilizers (valium,xanax, etc.)	0	0.0 %
13	Stimulants/amphetamines (diet pills, methamphetamine, etc.)	0	0.0 %
14	Cocaine	0	0.0 %
15	Opiates	0	0.0 %
16	Ecstasy	0	0.0 %
17	Hallucinogens (acid, LSD, etc.)	0	0.0 %
18	Inhalants (glue, paint, etc.)	0	0.0 %
19	Amyl nitrate/poppers/odorizers	0	0.0 %
96	Other (specify)	0	0.0 %
	Missing Data		
-200	Data missing	4	0.3 %
-106	Hasnt used < 11 drugs in p24hr - question skipped	59	4.4 %
-105	Hasnt used other drug past 24hr - question skipped	1291	95.3 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 142-145 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -106 , -105 , -99 , -98

S0SUBU07: S0 Subuse: Drug 7 used in past 24 hours

Value	Label	Unweighted Frequency	%
11	Marijuana/hashish	0	0.0 %
12	Sedatives/tranquilizers (valium,xanax, etc.)	0	0.0 %
13	Stimulants/amphetamines (diet pills, methamphetamine, etc.)	0	0.0 %
14	Cocaine	0	0.0 %
15	Opiates	0	0.0 %
16	Ecstasy	0	0.0 %
17	Hallucinogens (acid, LSD, etc.)	0	0.0 %
18	Inhalants (glue, paint, etc.)	0	0.0 %
19	Amyl nitrate/poppers/odorizers	0	0.0 %
96	Other (specify)	0	0.0 %
	Missing Data		
-200	Data missing	4	0.3 %
-106	Hasnt used < 11 drugs in p24hr - question skipped	59	4.4 %

Value	Label	Unweighted Frequency	%
-105	Hasnt used other drug past 24hr - question skipped	1291	95.3 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 146-149 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -106 , -105 , -99 , -98

S0SUBU08: S0 Subuse: Drug 8 used in past 24 hours

Value	Label	Unweighted Frequency	%
11	Marijuana/hashish	0	0.0 %
12	Sedatives/tranquilizers (valium,xanax, etc.)	0	0.0 %
13	Stimulants/amphetamines (diet pills, methamphetamine, etc.)	0	0.0 %
14	Cocaine	0	0.0 %
15	Opiates	0	0.0 %
16	Ecstasy	0	0.0 %
17	Hallucinogens (acid, LSD, etc.)	0	0.0 %
18	Inhalants (glue, paint, etc.)	0	0.0 %
19	Amyl nitrate/poppers/odorizers	0	0.0 %
96	Other (specify)	0	0.0 %
	Missing Data		
-200	Data missing	4	0.3 %
-106	Hasnt used < 11 drugs in p24hr - question skipped	59	4.4 %
-105	Hasnt used other drug past 24hr - question skipped	1291	95.3 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 150-153 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -106 , -105 , -99 , -98

S0SUBU09: S0 Subuse: Drug 9 used in past 24 hours

Value	Label	Unweighted Frequency	%
11	Marijuana/hashish	0	0.0 %
12	Sedatives/tranquilizers (valium,xanax, etc.)	0	0.0 %
13	Stimulants/amphetamines (diet pills, methamphetamine, etc.)	0	0.0 %
14	Cocaine	0	0.0 %
15	Opiates	0	0.0 %
16	Ecstasy	0	0.0 %
17	Hallucinogens (acid, LSD, etc.)	0	0.0 %

Value	Label	Unweighted Frequency	%
18	Inhalants (glue, paint, etc.)	0	0.0 %
19	Amyl nitrate/poppers/odorizers	0	0.0 %
96	Other (specify)	0	0.0 %
	Missing Data		
-200	Data missing	4	0.3 %
-106	Hasnt used < 11 drugs in p24hr - question skipped	59	4.4 %
-105	Hasnt used other drug past 24hr - question skipped	1291	95.3 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 154-157 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -106 , -105 , -99 , -98

S0SUBU28: S0 Subuse: Drug 10 used in past 24 hours

Value	Label	Unweighted Frequency	%
11	Marijuana/hashish	0	0.0 %
12	Sedatives/tranquilizers (valium,xanax, etc.)	0	0.0 %
13	Stimulants/amphetamines (diet pills, methamphetamine, etc.)	0	0.0 %
14	Cocaine	0	0.0 %
15	Opiates	0	0.0 %
16	Ecstasy	0	0.0 %
17	Hallucinogens (acid, LSD, etc.)	0	0.0 %
18	Inhalants (glue, paint, etc.)	0	0.0 %
19	Amyl nitrate/poppers/odorizers	0	0.0 %
96	Other (specify)	0	0.0 %
	Missing Data		
-200	Data missing	4	0.3 %
-106	Hasnt used < 11 drugs in p24hr - question skipped	59	4.4 %
-105	Hasnt used other drug past 24hr - question skipped	1291	95.3 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 158-161 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -106 , -105 , -99 , -98

S0SUBU29: S0 Subuse: Drug 11 used in past 24 hours

Value	Label	Unweighted Frequency	%
11	Marijuana/hashish	0	0.0 %

Value	Label	Unweighted Frequency	%
12	Sedatives/tranquilizers (valium,xanax, etc.)	0	0.0 %
13	Stimulants/amphetamines (diet pills, methamphetamine, etc.)	0	0.0 %
14	Cocaine	0	0.0 %
15	Opiates	0	0.0 %
16	Ecstasy	0	0.0 %
17	Hallucinogens (acid, LSD, etc.)	0	0.0 %
18	Inhalants (glue, paint, etc.)	0	0.0 %
19	Amyl nitrate/poppers/odorizers	0	0.0 %
96	Other (specify)	0	0.0 %
	Missing Data		
-200	Data missing	4	0.3 %
-106	Hasnt used < 11 drugs in p24hr - question skipped	59	4.4 %
-105	Hasnt used other drug past 24hr - question skipped	1291	95.3 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 162-165 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -106 , -105 , -99 , -98

S0SUBU37: S0 Subuse: Past 24 hours used prescribed medications (own or someone elses)

Value	Label	Unweighted Frequency	%
1	Yes	2	0.1 %
5	No	0	0.0 %
	Missing Data		
-800	Data unavailable	1352	99.9 %
	Total	**1,354**	**100%**

Based upon 2 valid cases out of 1,354 total cases.

Location: 166-169 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800

S0SUBU30: S0 Subuse: Other drug used past 24 hours (specify)

Based upon 1,354 valid cases out of 1,354 total cases.

Location: 170-170 (width: 1; decimal: 0)
Variable Type: character

S0SUBU31: S0 SubUse: EVER smoked cigarettes

Value	Label	Unweighted Frequency	%
1	Yes	981	72.5 %
5	No	370	27.3 %
	Missing Data		
-200	Data missing	3	0.2 %
	Total	**1,354**	**100%**

Based upon 1,351 valid cases out of 1,354 total cases.

Location: 171-174 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -9 , -8

S0SUBU32: S0 SubUse: When smoking the most how often

Value	Label	Unweighted Frequency	%
0	Never smoked in lifetime	370	27.3 %
1	1-2 times	143	10.6 %
2	Once per month	57	4.2 %
3	2-3 times/week	108	8.0 %
4	4-5 times/week	55	4.1 %
5	Every day	618	45.6 %
	Missing Data		
-200	Data missing	3	0.2 %
	Total	**1,354**	**100%**

Based upon 1,351 valid cases out of 1,354 total cases.

Location: 175-178 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -108 , -9 , -8

S0SUBU33: S0 SubUse: Past 6 months how often smoke cigarettes

Value	Label	Unweighted Frequency	%
1	Not at all	629	46.5 %
2	1-2 times	64	4.7 %
3	3-5 times	47	3.5 %
4	1x per month	18	1.3 %
5	2-3x per month	32	2.4 %
6	1x per week	22	1.6 %
7	2-3x per week	63	4.7 %
8	4-5x per week	65	4.8 %
9	Everyday	411	30.4 %
	Missing Data		

Value	Label	Unweighted Frequency	%
-200	Data missing	3	0.2 %
	Total	**1,354**	**100%**

Based upon 1,351 valid cases out of 1,354 total cases.

Location: 179-182 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -108 , -99 , -98

S0SUBU34: S0 SubUse: Past 6 months how many cigarettes in a day

Value	Label	Unweighted Frequency	%
1	0 cigarettes	640	47.3 %
2	1-4 cigarettes	304	22.5 %
3	5-9 cigarettes	142	10.5 %
4	10-15 cigarettes	104	7.7 %
5	16-20 cigarettes	105	7.8 %
6	More than a pack a day	56	4.1 %
	Missing Data		
-200	Data missing	3	0.2 %
	Total	**1,354**	**100%**

Based upon 1,351 valid cases out of 1,354 total cases.

Location: 183-186 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -109 , -108 , -9 , -8

S0SUSYMA: S0 SubUse: Lifetime family complaints because of alc/drug use

Value	Label	Unweighted Frequency	%
1	Yes	613	45.3 %
5	No	641	47.3 %
	Missing Data		
-200	Data missing	4	0.3 %
-111	Skipped: Has never used drugs or alcohol	95	7.0 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,254 valid cases out of 1,354 total cases.

Location: 187-190 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -111 , -9 , -8

S0SUSYMB: S0 SubUse: Past 6 months family complaints because of alc/drug use

Value	Label	Unweighted Frequency	%
1	Yes	348	25.7 %
5	No	188	13.9 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	567	41.9 %
-111	Never used drugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	151	11.2 %
	Total	**1,354**	**100%**

Based upon 536 valid cases out of 1,354 total cases.

Location: 191-194 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSYMC: S0 SubUse: Was it because of alcohol drugs both

Value	Label	Unweighted Frequency	%
1	Alcohol	80	5.9 %
2	Drugs	324	23.9 %
3	Both	209	15.4 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	490	36.2 %
-111	Never used drugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	151	11.2 %
	Total	**1,354**	**100%**

Based upon 613 valid cases out of 1,354 total cases.

Location: 195-198 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSYM2: S0 SubUse: Lifetime friends complain because of alc/drug use

Value	Label	Unweighted Frequency	%
1	Yes	258	19.1 %
5	No	996	73.6 %
	Missing Data		
-200	Data missing	4	0.3 %
-111	Skipped: Has never used drugs or alcohol	95	7.0 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,254 valid cases out of 1,354 total cases.

Location: 199-202 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -111 , -9 , -8

S0SUSYM3: S0 SubUse: Past 6 months friends complain because of alc/drug use

Value	Label	Unweighted Frequency	%
1	Yes	158	11.7 %
5	No	70	5.2 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	808	59.7 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	218	16.1 %
	Total	**1,354**	**100%**

Based upon 228 valid cases out of 1,354 total cases.

Location: 203-206 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSYM4: S0 SubUse: Was it because of alcohol drugs or both

Value	Label	Unweighted Frequency	%
1	Alcohol	49	3.6 %
2	Drugs	124	9.2 %
3	Both	85	6.3 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	778	57.5 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	218	16.1 %
	Total	**1,354**	**100%**

Based upon 258 valid cases out of 1,354 total cases.

Location: 207-210 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSYM5: S0 SubUse: Lifetime trouble school/work because of alc/drug use

Value	Label	Unweighted Frequency	%
1	Yes	301	22.2 %

Value	Label	Unweighted Frequency	%
5	No	954	70.5 %
	Missing Data		
-200	Data missing	4	0.3 %
-111	Skipped: Has never used drugs or alcohol	95	7.0 %
	Total	**1,354**	**100%**

Based upon 1,255 valid cases out of 1,354 total cases.

Location: 211-214 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -111 , -9 , -8

S0SUSYM6: S0 SubUse: Past 6 months trouble school/work because of alc/drug use

Value	Label	Unweighted Frequency	%
1	Yes	96	7.1 %
5	No	170	12.6 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	774	57.2 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	214	15.8 %
	Total	**1,354**	**100%**

Based upon 266 valid cases out of 1,354 total cases.

Location: 215-218 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSYM7: S0 SubUse: Was it because of alcohol drugs or both

Value	Label	Unweighted Frequency	%
1	Alcohol	26	1.9 %
2	Drugs	217	16.0 %
3	Both	58	4.3 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	739	54.6 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	214	15.8 %
	Total	**1,354**	**100%**

Based upon 301 valid cases out of 1,354 total cases.

Location: 219-222 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSYM8: S0 SubUse: Lifetime missed school/work because of alc/drug use

Value	Label	Unweighted Frequency	%
1	Yes	375	27.7 %
5	No	880	65.0 %
	Missing Data		
-200	Data missing	4	0.3 %
-111	Skipped: Has never used drugs or alcohol	95	7.0 %
	Total	**1,354**	**100%**

Based upon 1,255 valid cases out of 1,354 total cases.

Location: 223-226 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -111 , -9 , -8

S0SUSYM9: S0 SubUse: Past 6 months miss school/work because of alc/drug use

Value	Label	Unweighted Frequency	%
1	Yes	119	8.8 %
5	No	211	15.6 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	725	53.5 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	198	14.6 %
-9	Dont Know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 330 valid cases out of 1,354 total cases.

Location: 227-230 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY10: S0 SubUse: Was it because of alcohol drugs or both

Value	Label	Unweighted Frequency	%
1	Alcohol	76	5.6 %
2	Drugs	180	13.3 %
3	Both	117	8.6 %
	Missing Data		

Value	Label	Unweighted Frequency	%
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	681	50.3 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	198	14.6 %
-9	Dont Know	2	0.1 %
	Total	**1,354**	**100%**

Based upon 373 valid cases out of 1,354 total cases.

Location: 231-234 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY11: S0 SubUse: Lifetime arrested for alc/drug use

Value	Label	Unweighted Frequency	%
1	Yes	353	26.1 %
5	No	901	66.5 %
	Missing Data		
-200	Data missing	5	0.4 %
-111	Skipped: Has never used drugs or alcohol	95	7.0 %
	Total	**1,354**	**100%**

Based upon 1,254 valid cases out of 1,354 total cases.

Location: 235-238 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -111 , -9 , -8

S0SUSY12: S0 SubUse: Past 6 months arrested for alc/drug use

Value	Label	Unweighted Frequency	%
1	Yes	180	13.3 %
5	No	125	9.2 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	746	55.1 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	203	15.0 %
	Total	**1,354**	**100%**

Based upon 305 valid cases out of 1,354 total cases.

Location: 239-242 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY13: S0 SubUse: Was it because of alcohol drugs or both

Value	Label	Unweighted Frequency	%
1	Alcohol	66	4.9 %
2	Drugs	209	15.4 %
3	Both	77	5.7 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	698	51.6 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	203	15.0 %
-9	Dont Know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 352 valid cases out of 1,354 total cases.

Location: 243-246 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY14: S0 SubUse: Lifetime argue family because of alc/drug use

Value	Label	Unweighted Frequency	%
1	Yes	424	31.3 %
5	No	830	61.3 %
	Missing Data		
-200	Data missing	5	0.4 %
-111	Skipped: Has never used drugs or alcohol	95	7.0 %
	Total	**1,354**	**100%**

Based upon 1,254 valid cases out of 1,354 total cases.

Location: 247-250 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -111 , -9 , -8

S0SUSY15: S0 SubUse: Past 6 months argue w family because of alc/drug use

Value	Label	Unweighted Frequency	%
1	Yes	215	15.9 %
5	No	158	11.7 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	701	51.8 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %

Value	Label	Unweighted Frequency	%
-110	No complaints from family about use - question skipped	180	13.3 %
	Total	**1,354**	**100%**

Based upon 373 valid cases out of 1,354 total cases.

Location: 251-254 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY16: S0 SubUse: Was it because of alcohol drugs or both

Value	Label	Unweighted Frequency	%
1	Alcohol	71	5.2 %
2	Drugs	209	15.4 %
3	Both	144	10.6 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	650	48.0 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	180	13.3 %
	Total	**1,354**	**100%**

Based upon 424 valid cases out of 1,354 total cases.

Location: 255-258 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY17: S0 SubUse: Lifetime continue to use w known problems

Value	Label	Unweighted Frequency	%
1	Yes	239	17.7 %
5	No	1015	75.0 %
	Missing Data		
-200	Data missing	5	0.4 %
-111	Skipped: Has never used drugs or alcohol	95	7.0 %
	Total	**1,354**	**100%**

Based upon 1,254 valid cases out of 1,354 total cases.

Location: 259-262 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -111 , -9 , -8

S0SUSY18: S0 SubUse: Past 6 months continue use w known problems

Value	Label	Unweighted Frequency	%
1	Yes	130	9.6 %
5	No	83	6.1 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	804	59.4 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	237	17.5 %
	Total	**1,354**	**100%**

Based upon 213 valid cases out of 1,354 total cases.

Location: 263-266 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY19: S0 SubUse: Was it because of alcohol drugs or both

Value	Label	Unweighted Frequency	%
1	Alcohol	31	2.3 %
2	Drugs	117	8.6 %
3	Both	91	6.7 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	778	57.5 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	237	17.5 %
	Total	**1,354**	**100%**

Based upon 239 valid cases out of 1,354 total cases.

Location: 267-270 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY20: S0 SubUse: Lifetime cant remember party because of alc/drug use

Value	Label	Unweighted Frequency	%
1	Yes	493	36.4 %
5	No	761	56.2 %
	Missing Data		
-200	Data missing	5	0.4 %
-111	Skipped: Has never used drugs or alcohol	95	7.0 %
	Total	**1,354**	**100%**

Based upon 1,254 valid cases out of 1,354 total cases.

Location: 271-274 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -111 , -9 , -8

S0SUSY21: S0 SubUse: Past 6 months cant remember party because of alc/drug use

Value	Label	Unweighted Frequency	%
1	Yes	237	17.5 %
5	No	205	15.1 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	637	47.0 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	175	12.9 %
	Total	**1,354**	**100%**

Based upon 442 valid cases out of 1,354 total cases.

Location: 275-278 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY22: S0 SubUse: Was it because of alcohol drugs or both

Value	Label	Unweighted Frequency	%
1	Alcohol	175	12.9 %
2	Drugs	132	9.7 %
3	Both	186	13.7 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	586	43.3 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	175	12.9 %
	Total	**1,354**	**100%**

Based upon 493 valid cases out of 1,354 total cases.

Location: 279-282 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY23: S0 SubUse: Lifetime used in situations could get hurt

Value	Label	Unweighted Frequency	%
1	Yes	464	34.3 %

Value	Label	Unweighted Frequency	%
5	No	790	58.3 %
	Missing Data		
-200	Data missing	5	0.4 %
-111	Skipped: Has never used drugs or alcohol	95	7.0 %
	Total	**1,354**	**100%**

Based upon 1,254 valid cases out of 1,354 total cases.

Location: 283-286 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -111 , -9 , -8

S0SUSY24: S0 SubUse: Past 6 months used when could get hurt

Value	Label	Unweighted Frequency	%
1	Yes	221	16.3 %
5	No	182	13.4 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	669	49.4 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	182	13.4 %
	Total	**1,354**	**100%**

Based upon 403 valid cases out of 1,354 total cases.

Location: 287-290 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY25: S0 SubUse: Was it because of alcohol drugs or both

Value	Label	Unweighted Frequency	%
1	Alcohol	97	7.2 %
2	Drugs	181	13.4 %
3	Both	186	13.7 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	608	44.9 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	182	13.4 %
	Total	**1,354**	**100%**

Based upon 464 valid cases out of 1,354 total cases.

Location: 291-294 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY26: S0 SubUse: Lifetime had accident because of alc/drug use

Value	Label	Unweighted Frequency	%
1	Yes	152	11.2 %
5	No	1102	81.4 %
	Missing Data		
-200	Data missing	5	0.4 %
-111	Skipped: Has never used drugs or alcohol	95	7.0 %
	Total	**1,354**	**100%**

Based upon 1,254 valid cases out of 1,354 total cases.

Location: 295-298 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -111 , -9 , -8

S0SUSY27: S0 SubUse: Past 6 months accident bc drug alcohol use

Value	Label	Unweighted Frequency	%
1	Yes	57	4.2 %
5	No	79	5.8 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	875	64.6 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	243	17.9 %
	Total	**1,354**	**100%**

Based upon 136 valid cases out of 1,354 total cases.

Location: 299-302 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY28: S0 SubUse: Was it because of alcohol drugs or both

Value	Label	Unweighted Frequency	%
1	Alcohol	51	3.8 %
2	Drugs	46	3.4 %
3	Both	55	4.1 %
	Missing Data		
-200	Data missing	5	0.4 %

Value	Label	Unweighted Frequency	%
-113	Individual item not endorsed, followup skipped	859	63.4 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	243	17.9 %
	Total	**1,354**	**100%**

Based upon 152 valid cases out of 1,354 total cases.

Location: 303-306 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY29: S0 SubUse: Lifetime alc/drugs interfere w work school

Value	Label	Unweighted Frequency	%
1	Yes	384	28.4 %
5	No	870	64.3 %
	Missing Data		
-200	Data missing	5	0.4 %
-111	Skipped: Has never used drugs or alcohol	95	7.0 %
	Total	**1,354**	**100%**

Based upon 1,254 valid cases out of 1,354 total cases.

Location: 307-310 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -111 , -9 , -8

S0SUSY30: S0 SubUse: Past 6 months alc/drugs interfere w work school

Value	Label	Unweighted Frequency	%
1	Yes	181	13.4 %
5	No	153	11.3 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	728	53.8 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	192	14.2 %
	Total	**1,354**	**100%**

Based upon 334 valid cases out of 1,354 total cases.

Location: 311-314 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY31: S0 SubUse: Was it because of alcohol drugs or both

Value	Label	Unweighted Frequency	%
1	Alcohol	38	2.8 %
2	Drugs	217	16.0 %
3	Both	129	9.5 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	678	50.1 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	192	14.2 %
	Total	**1,354**	**100%**

Based upon 384 valid cases out of 1,354 total cases.

Location: 315-318 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY32: S0 SubUse: Lifetime trouble w police because of alc/drug use

Value	Label	Unweighted Frequency	%
1	Yes	311	23.0 %
5	No	943	69.6 %
	Missing Data		
-200	Data missing	5	0.4 %
-111	Skipped: Has never used drugs or alcohol	95	7.0 %
	Total	**1,354**	**100%**

Based upon 1,254 valid cases out of 1,354 total cases.

Location: 319-322 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -111 , -9 , -8

S0SUSY33: S0 SubUse: Past 6 months trouble w police

Value	Label	Unweighted Frequency	%
1	Yes	123	9.1 %
5	No	149	11.0 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	773	57.1 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	209	15.4 %
	Total	**1,354**	**100%**

Based upon 272 valid cases out of 1,354 total cases.

Location: 323-326 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY34: S0 SubUse: Was it because of alcohol drugs or both

Value	Label	Unweighted Frequency	%
1	Alcohol	72	5.3 %
2	Drugs	157	11.6 %
3	Both	82	6.1 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	734	54.2 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	209	15.4 %
	Total	**1,354**	**100%**

Based upon 311 valid cases out of 1,354 total cases.

Location: 327-330 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY35: S0 SubUse: Lifetime had sex for money to buy drugs

Value	Label	Unweighted Frequency	%
1	Yes	12	0.9 %
5	No	1240	91.6 %
	Missing Data		
-200	Data missing	5	0.4 %
-111	Skipped: Has never used drugs or alcohol	95	7.0 %
-9	Dont know	2	0.1 %
	Total	**1,354**	**100%**

Based upon 1,252 valid cases out of 1,354 total cases.

Location: 331-334 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -111 , -9 , -8

S0SUSY36: S0 SubUse: Past 6 months had sex for money for drug

Value	Label	Unweighted Frequency	%
1	Yes	5	0.4 %
5	No	6	0.4 %

Value	Label	Unweighted Frequency	%
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	967	71.4 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	276	20.4 %
	Total	**1,354**	**100%**

Based upon 11 valid cases out of 1,354 total cases.

Location: 335-338 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY37: S0 SubUse: Was it because of alcohol drugs or both

Value	Label	Unweighted Frequency	%
1	Alcohol	1	0.1 %
2	Drugs	10	0.7 %
3	Both	1	0.1 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	966	71.3 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	276	20.4 %
	Total	**1,354**	**100%**

Based upon 12 valid cases out of 1,354 total cases.

Location: 339-342 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY38: S0 SubUse: Lifetime sold drugs to buy drugs

Value	Label	Unweighted Frequency	%
1	Yes	361	26.7 %
5	No	893	66.0 %
	Missing Data		
-200	Data missing	5	0.4 %
-111	Skipped: Has never used drugs or alcohol	95	7.0 %
	Total	**1,354**	**100%**

Based upon 1,254 valid cases out of 1,354 total cases.

Location: 343-346 (width: 4; decimal: 0)
Variable Type: numeric

(Range of) Missing Values: -200 , -111 , -9 , -8

S0SUSY39: S0 SubUse: Past 6 months sold drugs to buy drugs

Value	Label	Unweighted Frequency	%
1	Yes	183	13.5 %
5	No	145	10.7 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	729	53.8 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	197	14.5 %
	Total	**1,354**	**100%**

Based upon 328 valid cases out of 1,354 total cases.

Location: 347-350 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY40: S0 SubUse: Was it because of alcohol drugs or both

Value	Label	Unweighted Frequency	%
1	Alcohol	4	0.3 %
2	Drugs	303	22.4 %
3	Both	54	4.0 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	696	51.4 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	197	14.5 %
	Total	**1,354**	**100%**

Based upon 361 valid cases out of 1,354 total cases.

Location: 351-354 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY41: S0 SubUse: Lifetime robbed/stole money for drugs

Value	Label	Unweighted Frequency	%
1	Yes	260	19.2 %
5	No	994	73.4 %
	Missing Data		
-200	Data missing	5	0.4 %

Value	Label	Unweighted Frequency	%
-111	Skipped: Has never used drugs or alcohol	95	7.0 %
	Total	**1,354**	**100%**

Based upon 1,254 valid cases out of 1,354 total cases.

Location: 355-358 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -111 , -9 , -8

S0SUSY42: S0 SubUse: Past 6 months robbed/stole money for drugs

Value	Label	Unweighted Frequency	%
1	Yes	118	8.7 %
5	No	119	8.8 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	794	58.6 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	223	16.5 %
	Total	**1,354**	**100%**

Based upon 237 valid cases out of 1,354 total cases.

Location: 359-362 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY43: S0 SubUse: Was it because of alcohol drugs or both

Value	Label	Unweighted Frequency	%
1	Alcohol	13	1.0 %
2	Drugs	184	13.6 %
3	Both	62	4.6 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	771	56.9 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	223	16.5 %
-9	Dont Know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 259 valid cases out of 1,354 total cases.

Location: 363-366 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY44: S0 SubUse: Lifetime fought bc of drug/alc use

Value	Label	Unweighted Frequency	%
1	Yes	337	24.9 %
5	No	917	67.7 %
	Missing Data		
-200	Data missing	5	0.4 %
-111	Skipped: Has never used drugs or alcohol	95	7.0 %
	Total	**1,354**	**100%**

Based upon 1,254 valid cases out of 1,354 total cases.

Location: 367-370 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -111 , -9 , -8

S0SUSY45: S0 SubUse: Past 6 months fight bc of drug/alc use

Value	Label	Unweighted Frequency	%
1	Yes	140	10.3 %
5	No	157	11.6 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	749	55.3 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	208	15.4 %
	Total	**1,354**	**100%**

Based upon 297 valid cases out of 1,354 total cases.

Location: 371-374 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY46: S0 SubUse: Was it because of alcohol drugs or both

Value	Label	Unweighted Frequency	%
1	Alcohol	126	9.3 %
2	Drugs	115	8.5 %
3	Both	96	7.1 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	709	52.4 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	208	15.4 %

Value	Label	Unweighted Frequency	%
	Total	**1,354**	**100%**

Based upon 337 valid cases out of 1,354 total cases.

Location: 375-378 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY47: S0 SubUse: Lifetime destroy property bc of drug/alc

Value	Label	Unweighted Frequency	%
1	Yes	220	16.2 %
5	No	1034	76.4 %
	Missing Data		
-200	Data missing	5	0.4 %
-111	Skipped: Has never used drugs or alcohol	95	7.0 %
	Total	**1,354**	**100%**

Based upon 1,254 valid cases out of 1,354 total cases.

Location: 379-382 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -111 , -9 , -8

S0SUSY48: S0 SubUse: Past 6 months destroy property

Value	Label	Unweighted Frequency	%
1	Yes	94	6.9 %
5	No	98	7.2 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	832	61.4 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	229	16.9 %
-9	Dont Know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 192 valid cases out of 1,354 total cases.

Location: 383-386 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY49: S0 SubUse: Was it because of alcohol drugs or both

Value	Label	Unweighted Frequency	%
1	Alcohol	71	5.2 %
2	Drugs	76	5.6 %
3	Both	72	5.3 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	805	59.5 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	229	16.9 %
-8	Refuse	1	0.1 %
	Total	**1,354**	**100%**

Based upon 219 valid cases out of 1,354 total cases.

Location: 387-390 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY50: S0 SubUse: Lifetime urge to use could not stop self

Value	Label	Unweighted Frequency	%
1	Yes	221	16.3 %
5	No	1033	76.3 %
	Missing Data		
-200	Data missing	5	0.4 %
-111	Skipped: Has never used drugs or alcohol	95	7.0 %
	Total	**1,354**	**100%**

Based upon 1,254 valid cases out of 1,354 total cases.

Location: 391-394 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -111 , -9 , -8

S0SUSY51: S0 SubUse: Past 6 months urge to use cant stop self

Value	Label	Unweighted Frequency	%
1	Yes	120	8.9 %
5	No	73	5.4 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	830	61.3 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	231	17.1 %
	Total	**1,354**	**100%**

Based upon 193 valid cases out of 1,354 total cases.

Location: 395-398 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY52: S0 SubUse: Was it because of alcohol drugs or both

Value	Label	Unweighted Frequency	%
1	Alcohol	23	1.7 %
2	Drugs	138	10.2 %
3	Both	60	4.4 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	802	59.2 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	231	17.1 %
	Total	**1,354**	**100%**

Based upon 221 valid cases out of 1,354 total cases.

Location: 399-402 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY53: S0 SubUse: Lifetime want to use cant think anything

Value	Label	Unweighted Frequency	%
1	Yes	182	13.4 %
5	No	1072	79.2 %
	Missing Data		
-200	Data missing	5	0.4 %
-111	Skipped: Has never used drugs or alcohol	95	7.0 %
	Total	**1,354**	**100%**

Based upon 1,254 valid cases out of 1,354 total cases.

Location: 403-406 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -111 , -9 , -8

S0SUSY54: S0 SubUse: Past 6 months want to use cant think

Value	Label	Unweighted Frequency	%
1	Yes	109	8.1 %
5	No	60	4.4 %
	Missing Data		

Value	Label	Unweighted Frequency	%
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	842	62.2 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	243	17.9 %
	Total	**1,354**	**100%**

Based upon 169 valid cases out of 1,354 total cases.

Location: 407-410 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY55: S0 SubUse: Was it because of alcohol drugs or both

Value	Label	Unweighted Frequency	%
1	Alcohol	14	1.0 %
2	Drugs	118	8.7 %
3	Both	50	3.7 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	829	61.2 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	243	17.9 %
	Total	**1,354**	**100%**

Based upon 182 valid cases out of 1,354 total cases.

Location: 411-414 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY56: S0 SubUse: Cut down on DnA have symptoms

Value	Label	Unweighted Frequency	%
1	Yes	193	14.3 %
5	No	1061	78.4 %
	Missing Data		
-200	Data missing	5	0.4 %
-111	Skipped: Has never used drugs or alcohol	95	7.0 %
	Total	**1,354**	**100%**

Based upon 1,254 valid cases out of 1,354 total cases.

Location: 415-418 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -111 , -9 , -8

S0SUSY57: S0 SubUse: Past 6 months cut down have symptoms

Value	Label	Unweighted Frequency	%
1	Yes	116	8.6 %
5	No	59	4.4 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	839	62.0 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	240	17.7 %
	Total	**1,354**	**100%**

Based upon 175 valid cases out of 1,354 total cases.

Location: 419-422 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY58: S0 SubUse: Was it because of alcohol drugs or both

Value	Label	Unweighted Frequency	%
1	Alcohol	27	2.0 %
2	Drugs	125	9.2 %
3	Both	41	3.0 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	821	60.6 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	240	17.7 %
	Total	**1,354**	**100%**

Based upon 193 valid cases out of 1,354 total cases.

Location: 423-426 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY59: S0 SubUse: Lifetime ever use to avoid symptoms

Value	Label	Unweighted Frequency	%
1	Yes	109	8.1 %
5	No	84	6.2 %
	Missing Data		
-200	Data missing	5	0.4 %
-115	Individual items not endorsed, followup skipped	1061	78.4 %

Value	Label	Unweighted Frequency	%
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
	Total	**1,354**	**100%**

Based upon 193 valid cases out of 1,354 total cases.

Location: 427-430 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -115 , -113 , -111 , -9 , -8

S0SUSY60: S0 SubUse: Past 6 months use to avoid symptoms

Value	Label	Unweighted Frequency	%
1	Yes	77	5.7 %
5	No	24	1.8 %
	Missing Data		
-200	Data missing	5	0.4 %
-115	Individual items not endorsed, followup skipped	1061	78.4 %
-113	Individual item not endorsed, followup skipped	79	5.8 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	13	1.0 %
	Total	**1,354**	**100%**

Based upon 101 valid cases out of 1,354 total cases.

Location: 431-434 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -115 , -113 , -111 , -110 , -9 , -8

S0SUSY61: S0 SubUse: Was it because of alcohol drugs or both

Value	Label	Unweighted Frequency	%
1	Alcohol	11	0.8 %
2	Drugs	68	5.0 %
3	Both	30	2.2 %
	Missing Data		
-200	Data missing	5	0.4 %
-115	Individual items not endorsed, followup skipped	1061	78.4 %
-113	Individual item not endorsed, followup skipped	71	5.2 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	13	1.0 %
	Total	**1,354**	**100%**

Based upon 109 valid cases out of 1,354 total cases.

Location: 435-438 (width: 4; decimal: 0)
Variable Type: numeric

(Range of) Missing Values: -200 , -115 , -113 , -111 , -110 , -9 , -8

S0SUSY62: S0 SubUse: Lifetime cut down family act to use

Value	Label	Unweighted Frequency	%
1	Yes	307	22.7 %
5	No	947	69.9 %
	Missing Data		
-200	Data missing	5	0.4 %
-111	Skipped: Has never used drugs or alcohol	95	7.0 %
	Total	**1,354**	**100%**

Based upon 1,254 valid cases out of 1,354 total cases.

Location: 439-442 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -111 , -9 , -8

S0SUSY63: S0 SubUse: Past 6 months cut down family act

Value	Label	Unweighted Frequency	%
1	Yes	155	11.4 %
5	No	117	8.6 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	759	56.1 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	223	16.5 %
	Total	**1,354**	**100%**

Based upon 272 valid cases out of 1,354 total cases.

Location: 443-446 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY64: S0 SubUse: Was it because of alcohol drugs or both

Value	Label	Unweighted Frequency	%
1	Alcohol	26	1.9 %
2	Drugs	187	13.8 %
3	Both	94	6.9 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	724	53.5 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %

Value	Label	Unweighted Frequency	%
-110	No complaints from family about use - question skipped	223	16.5 %
	Total	**1,354**	**100%**

Based upon 307 valid cases out of 1,354 total cases.

Location: 447-450 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY65: S0 SubUse: Lifetime used when promised wouldnt

Value	Label	Unweighted Frequency	%
1	Yes	469	34.6 %
5	No	785	58.0 %
	Missing Data		
-200	Data missing	5	0.4 %
-111	Skipped: Has never used drugs or alcohol	95	7.0 %
	Total	**1,354**	**100%**

Based upon 1,254 valid cases out of 1,354 total cases.

Location: 451-454 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -111 , -9 , -8

S0SUSY66: S0 SubUse: Past 6 months used when promised wouldnt

Value	Label	Unweighted Frequency	%
1	Yes	249	18.4 %
5	No	163	12.0 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	656	48.4 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	186	13.7 %
	Total	**1,354**	**100%**

Based upon 412 valid cases out of 1,354 total cases.

Location: 455-458 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY67: S0 SubUse: Was it because of alcohol drugs or both

Value	Label	Unweighted Frequency	%
1	Alcohol	58	4.3 %
2	Drugs	272	20.1 %
3	Both	139	10.3 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	599	44.2 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	186	13.7 %
	Total	**1,354**	**100%**

Based upon 469 valid cases out of 1,354 total cases.

Location: 459-462 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY68: S0 SubUse: Lifetime used more days than intended

Value	Label	Unweighted Frequency	%
1	Yes	424	31.3 %
5	No	830	61.3 %
	Missing Data		
-200	Data missing	5	0.4 %
-111	Skipped: Has never used drugs or alcohol	95	7.0 %
	Total	**1,354**	**100%**

Based upon 1,254 valid cases out of 1,354 total cases.

Location: 463-466 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -111 , -9 , -8

S0SUSY69: S0 SubUse: Past 6 months used more days than intended

Value	Label	Unweighted Frequency	%
1	Yes	244	18.0 %
5	No	136	10.0 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	688	50.8 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	186	13.7 %
	Total	**1,354**	**100%**

Based upon 380 valid cases out of 1,354 total cases.

Location: 467-470 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY70: S0 SubUse: Was it because of alcohol drugs or both

Value	Label	Unweighted Frequency	%
1	Alcohol	58	4.3 %
2	Drugs	244	18.0 %
3	Both	122	9.0 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	644	47.6 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	186	13.7 %
	Total	**1,354**	**100%**

Based upon 424 valid cases out of 1,354 total cases.

Location: 471-474 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY71: S0 SubUse: Lifetime tried to cut down but couldnt

Value	Label	Unweighted Frequency	%
1	Yes	304	22.5 %
5	No	950	70.2 %
	Missing Data		
-200	Data missing	5	0.4 %
-111	Skipped: Has never used drugs or alcohol	95	7.0 %
	Total	**1,354**	**100%**

Based upon 1,254 valid cases out of 1,354 total cases.

Location: 475-478 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -111 , -9 , -8

S0SUSY72: S0 SubUse: Past 6 months tried to cut down but couldnt

Value	Label	Unweighted Frequency	%
1	Yes	174	12.9 %
5	No	100	7.4 %
	Missing Data		

Value	Label	Unweighted Frequency	%
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	764	56.4 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	216	16.0 %
	Total	**1,354**	**100%**

Based upon 274 valid cases out of 1,354 total cases.

Location: 479-482 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY73: S0 SubUse: Was it because of alcohol drugs or both

Value	Label	Unweighted Frequency	%
1	Alcohol	23	1.7 %
2	Drugs	209	15.4 %
3	Both	72	5.3 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	734	54.2 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	216	16.0 %
	Total	**1,354**	**100%**

Based upon 304 valid cases out of 1,354 total cases.

Location: 483-486 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY74: S0 SubUse: Lifetime needed more to get same effect

Value	Label	Unweighted Frequency	%
1	Yes	414	30.6 %
5	No	839	62.0 %
	Missing Data		
-200	Data missing	5	0.4 %
-111	Skipped: Has never used drugs or alcohol	95	7.0 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,253 valid cases out of 1,354 total cases.

Location: 487-490 (width: 4; decimal: 0)
Variable Type: numeric

(Range of) Missing Values: -200 , -111 , -9 , -8

S0SUSY75: S0 SubUse: Past 6 months needed more to get high

Value	Label	Unweighted Frequency	%
1	Yes	214	15.8 %
5	No	157	11.6 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	684	50.5 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	199	14.7 %
	Total	**1,354**	**100%**

Based upon 371 valid cases out of 1,354 total cases.

Location: 491-494 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY76: S0 SubUse: Was it because of alcohol drugs or both

Value	Label	Unweighted Frequency	%
1	Alcohol	55	4.1 %
2	Drugs	248	18.3 %
3	Both	111	8.2 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	641	47.3 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	199	14.7 %
	Total	**1,354**	**100%**

Based upon 414 valid cases out of 1,354 total cases.

Location: 495-498 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY77: S0 SubUse: Lifetime felt needed or depend on DnA

Value	Label	Unweighted Frequency	%
1	Yes	183	13.5 %
5	No	1070	79.0 %
	Missing Data		
-200	Data missing	5	0.4 %

Value	Label	Unweighted Frequency	%
-111	Skipped: Has never used drugs or alcohol	95	7.0 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,253 valid cases out of 1,354 total cases.

Location: 499-502 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -111 , -9 , -8

S0SUSY78: S0 SubUse: Past 6 months felt needed or depend on DnA

Value	Label	Unweighted Frequency	%
1	Yes	110	8.1 %
5	No	57	4.2 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	842	62.2 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	245	18.1 %
	Total	**1,354**	**100%**

Based upon 167 valid cases out of 1,354 total cases.

Location: 503-506 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY79: S0 SubUse: Was it because of alcohol drugs or both

Value	Label	Unweighted Frequency	%
1	Alcohol	13	1.0 %
2	Drugs	133	9.8 %
3	Both	37	2.7 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	826	61.0 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	245	18.1 %
	Total	**1,354**	**100%**

Based upon 183 valid cases out of 1,354 total cases.

Location: 507-510 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY80: S0 SubUse: Lifetime talked to dr or clergy about DnA

Value	Label	Unweighted Frequency	%
1	Yes	185	13.7 %
5	No	1069	79.0 %
	Missing Data		
-200	Data missing	5	0.4 %
-111	Skipped: Has never used drugs or alcohol	95	7.0 %
	Total	**1,354**	**100%**

Based upon 1,254 valid cases out of 1,354 total cases.

Location: 511-514 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -111 , -9 , -8

S0SUSY81: S0 SubUse: Past 6 months talked to dr or clergy

Value	Label	Unweighted Frequency	%
1	Yes	100	7.4 %
5	No	60	4.4 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	856	63.2 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	238	17.6 %
	Total	**1,354**	**100%**

Based upon 160 valid cases out of 1,354 total cases.

Location: 515-518 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY82: S0 SubUse: Was it because of alcohol drugs or both

Value	Label	Unweighted Frequency	%
1	Alcohol	16	1.2 %
2	Drugs	103	7.6 %
3	Both	66	4.9 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	831	61.4 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	238	17.6 %

Value	Label	Unweighted Frequency	%
	Total	**1,354**	**100%**

Based upon 185 valid cases out of 1,354 total cases.

Location: 519-522 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY83: S0 SubUse: Lifetime attend AA or other self help

Value	Label	Unweighted Frequency	%
1	Yes	217	16.0 %
5	No	1037	76.6 %
	Missing Data		
-200	Data missing	5	0.4 %
-111	Skipped: Has never used drugs or alcohol	95	7.0 %
	Total	**1,354**	**100%**

Based upon 1,254 valid cases out of 1,354 total cases.

Location: 523-526 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -111 , -9 , -8

S0SUSY84: S0 SubUse: Past 6 months attend AA or self help

Value	Label	Unweighted Frequency	%
1	Yes	114	8.4 %
5	No	78	5.8 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	839	62.0 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	223	16.5 %
	Total	**1,354**	**100%**

Based upon 192 valid cases out of 1,354 total cases.

Location: 527-530 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY85: S0 SubUse: Was it because of alcohol drugs or both

Value	Label	Unweighted Frequency	%
1	Alcohol	37	2.7 %

Value	Label	Unweighted Frequency	%
2	Drugs	104	7.7 %
3	Both	76	5.6 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	814	60.1 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	223	16.5 %
	Total	**1,354**	**100%**

Based upon 217 valid cases out of 1,354 total cases.

Location: 531-534 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY86: S0 SubUse: Lifetime got DnA treatment from therapist

Value	Label	Unweighted Frequency	%
1	Yes	107	7.9 %
5	No	1147	84.7 %
	Missing Data		
-200	Data missing	5	0.4 %
-111	Skipped: Has never used drugs or alcohol	95	7.0 %
	Total	**1,354**	**100%**

Based upon 1,254 valid cases out of 1,354 total cases.

Location: 535-538 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -111 , -9 , -8

S0SUSY87: S0 SubUse: Past 6 months DnA treatment from therapist

Value	Label	Unweighted Frequency	%
1	Yes	59	4.4 %
5	No	39	2.9 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	906	66.9 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	250	18.5 %
	Total	**1,354**	**100%**

Based upon 98 valid cases out of 1,354 total cases.

Location: 539-542 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY88: S0 SubUse: Was it because of alcohol drugs or both

Value	Label	Unweighted Frequency	%
1	Alcohol	6	0.4 %
2	Drugs	61	4.5 %
3	Both	40	3.0 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	897	66.2 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	250	18.5 %
	Total	**1,354**	**100%**

Based upon 107 valid cases out of 1,354 total cases.

Location: 543-546 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY89: S0 SubUse: Lifetime been admitted to hosp for DnA

Value	Label	Unweighted Frequency	%
1	Yes	73	5.4 %
5	No	1181	87.2 %
	Missing Data		
-200	Data missing	5	0.4 %
-111	Skipped: Has never used drugs or alcohol	95	7.0 %
	Total	**1,354**	**100%**

Based upon 1,254 valid cases out of 1,354 total cases.

Location: 547-550 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -111 , -9 , -8

S0SUSY90: S0 SubUse: Past 6 months admitted to hosp for DnA

Value	Label	Unweighted Frequency	%
1	Yes	24	1.8 %
5	No	40	3.0 %
	Missing Data		
-200	Data missing	5	0.4 %

Value	Label	Unweighted Frequency	%
-113	Individual item not endorsed, followup skipped	929	68.6 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	261	19.3 %
	Total	**1,354**	**100%**

Based upon 64 valid cases out of 1,354 total cases.

Location: 551-554 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0SUSY91: S0 SubUse: Was it because of alcohol drugs or both

Value	Label	Unweighted Frequency	%
1	Alcohol	24	1.8 %
2	Drugs	35	2.6 %
3	Both	14	1.0 %
	Missing Data		
-200	Data missing	5	0.4 %
-113	Individual item not endorsed, followup skipped	920	67.9 %
-111	Never used rugs or alcohol - questions skipped	95	7.0 %
-110	No complaints from family about use - question skipped	261	19.3 %
	Total	**1,354**	**100%**

Based upon 73 valid cases out of 1,354 total cases.

Location: 555-558 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -113 , -111 , -110 , -9 , -8

S0PSUBA: S0 SubUse: Characterize mom drinking growing up

Value	Label	Unweighted Frequency	%
1	Abstainer (did not drink)	644	47.6 %
2	Occasional light drinker	373	27.5 %
3	Social drinker	146	10.8 %
4	Occasional heavy drinker	59	4.4 %
5	Frequent heavy drinker	35	2.6 %
6	Daily heavy drinker	50	3.7 %
	Missing Data		
-200	Data missing	3	0.2 %
-9	Dont Know	44	3.2 %
	Total	**1,354**	**100%**

Based upon 1,307 valid cases out of 1,354 total cases.

Location: 559-562 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -120 , -9 , -8

S0PSUBB: S0 SubUse: Relatives complain about mom drinking

Value	Label	Unweighted Frequency	%
1	Yes	155	11.4 %
5	No	516	38.1 %
	Missing Data		
-200	Data missing	3	0.2 %
-113	Individual item not endorsed, follow-up skipped	644	47.6 %
-9	Dont Know	36	2.7 %
	Total	**1,354**	**100%**

Based upon 671 valid cases out of 1,354 total cases.

Location: 563-566 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -120 , -113 , -9 , -8

S0PSUBC: S0 SubUse: Has mom ever attended A A bc drinking

Value	Label	Unweighted Frequency	%
1	Yes	101	7.5 %
5	No	547	40.4 %
	Missing Data		
-200	Data missing	3	0.2 %
-113	Individual item not endorsed, follow-up skipped	644	47.6 %
-9	Dont Know	59	4.4 %
	Total	**1,354**	**100%**

Based upon 648 valid cases out of 1,354 total cases.

Location: 567-570 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -120 , -113 , -9 , -8

S0PSUBD: S0 SubUse: Mom drinking create problems w relatives

Value	Label	Unweighted Frequency	%
1	Yes	135	10.0 %
5	No	533	39.4 %
	Missing Data		
-200	Data missing	3	0.2 %

Value	Label	Unweighted Frequency	%
-113	Individual item not endorsed, follow-up skipped	644	47.6 %
-9	Dont Know	39	2.9 %
	Total	**1,354**	**100%**

Based upon 668 valid cases out of 1,354 total cases.

Location: 571-574 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -120 , -113 , -9 , -8

S0PSUBE: S0 SubUse: Mom in trouble at work bc of drinking

Value	Label	Unweighted Frequency	%
1	Yes	22	1.6 %
5	No	635	46.9 %
	Missing Data		
-200	Data missing	3	0.2 %
-113	Individual item not endorsed, follow-up skipped	644	47.6 %
-9	Dont Know	50	3.7 %
	Total	**1,354**	**100%**

Based upon 657 valid cases out of 1,354 total cases.

Location: 575-578 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -120 , -113 , -9 , -8

S0PSUBF: S0 SubUse: Mom go to anyone for help for drinking

Value	Label	Unweighted Frequency	%
1	Yes	97	7.2 %
5	No	555	41.0 %
	Missing Data		
-200	Data missing	3	0.2 %
-113	Individual item not endorsed, follow-up skipped	644	47.6 %
-9	Dont Know	55	4.1 %
	Total	**1,354**	**100%**

Based upon 652 valid cases out of 1,354 total cases.

Location: 579-582 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -120 , -113 , -9 , -8

S0PSUBG: S0 SubUse: Mom been in a hosp because of drinking

Value	Label	Unweighted Frequency	%
1	Yes	41	3.0 %
5	No	612	45.2 %
	Missing Data		
-200	Data missing	3	0.2 %
-113	Individual item not endorsed, follow-up skipped	644	47.6 %
-9	Dont Know	54	4.0 %
	Total	**1,354**	**100%**

Based upon 653 valid cases out of 1,354 total cases.

Location: 583-586 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -120 , -113 , -9 , -8

S0PSUBH: S0 SubUse: Mom ever arrested for drunken driving

Value	Label	Unweighted Frequency	%
1	Yes	47	3.5 %
5	No	606	44.8 %
	Missing Data		
-200	Data missing	3	0.2 %
-113	Individual item not endorsed, follow-up skipped	644	47.6 %
-9	Dont Know	54	4.0 %
	Total	**1,354**	**100%**

Based upon 653 valid cases out of 1,354 total cases.

Location: 587-590 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -120 , -113 , -9 , -8

S0PSUBO: S0 SubUse: Mom ever neglect obligations bc drinking

Value	Label	Unweighted Frequency	%
1	Yes	84	6.2 %
5	No	579	42.8 %
	Missing Data		
-200	Data missing	3	0.2 %
-113	Individual item not endorsed, follow-up skipped	644	47.6 %
-9	Dont Know	44	3.2 %
	Total	**1,354**	**100%**

Based upon 663 valid cases out of 1,354 total cases.

Location: 591-594 (width: 4; decimal: 0)
Variable Type: numeric

(Range of) Missing Values: -200 , -120 , -113 , -9 , -8

S0PSUBI: S0 SubUse: Mom ever arrested for drunken behavior

Value	Label	Unweighted Frequency	%
1	Yes	51	3.8 %
5	No	594	43.9 %
	Missing Data		
-200	Data missing	3	0.2 %
-113	Individual item not endorsed, follow-up skipped	644	47.6 %
-9	Dont Know	62	4.6 %
	Total	**1,354**	**100%**

Based upon 645 valid cases out of 1,354 total cases.

Location: 595-598 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -120 , -113 , -9 , -8

S0PSUBJ: S0 SubUse: Do you think mom is an alcoholic now

Value	Label	Unweighted Frequency	%
1	Yes	38	2.8 %
3	Not sure	41	3.0 %
5	No	1244	91.9 %
	Missing Data		
-200	Data missing	3	0.2 %
-9	Dont Know	28	2.1 %
	Total	**1,354**	**100%**

Based upon 1,323 valid cases out of 1,354 total cases.

Location: 599-602 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -120 , -9 , -8

S0PSUBK: S0 SubUse: Think mom was an alcoholic in past

Value	Label	Unweighted Frequency	%
1	Yes	173	12.8 %
3	Not sure	54	4.0 %
5	No	1100	81.2 %
	Missing Data		
-200	Data missing	3	0.2 %
-9	Dont Know	24	1.8 %
	Total	**1,354**	**100%**

Based upon 1,327 valid cases out of 1,354 total cases.

Location: 603-606 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -121 , -120 , -9 , -8

S0PSUBL: S0 SubUse: Does did mom use street drugs

Value	Label	Unweighted Frequency	%
1	Yes, in the past year	123	9.1 %
2	Yes, but not in the past year	252	18.6 %
5	No, never	924	68.2 %
	Missing Data		
-200	Data missing	3	0.2 %
-9	Dont Know	51	3.8 %
-8	Refuse	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,299 valid cases out of 1,354 total cases.

Location: 607-610 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -9 , -8

S0PSUBM: S0 SubUse: Think mom has drug problem now

Value	Label	Unweighted Frequency	%
1	Yes	53	3.9 %
3	Not sure	11	0.8 %
5	No	1235	91.2 %
	Missing Data		
-200	Data missing	3	0.2 %
-114	Not Applicable	52	3.8 %
	Total	**1,354**	**100%**

Based upon 1,299 valid cases out of 1,354 total cases.

Location: 611-614 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -122 , -114 , -9 , -8

S0PSUBN: S0 SubUse: Think mom had drug problem in past

Value	Label	Unweighted Frequency	%
1	Yes	232	17.1 %
3	Not sure	19	1.4 %
5	No	1048	77.4 %

Value	Label	Unweighted Frequency	%
	Missing Data		
-200	Data missing	3	0.2 %
-114	Not Applicable	52	3.8 %
	Total	**1,354**	**100%**

Based upon 1,299 valid cases out of 1,354 total cases.

Location: 615-618 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -123 , -122 , -114 , -9 , -8

S0PSUBA2: S0 SubUse: Characterize dads drinking growing up

Value	Label	Unweighted Frequency	%
1	Abstainer (did not drink)	311	23.0 %
2	Occasional light drinker	210	15.5 %
3	Social drinker	173	12.8 %
4	Occasional heavy drinker	108	8.0 %
5	Frequent heavy drinker	108	8.0 %
6	Daily heavy drinker	189	14.0 %
	Missing Data		
-200	Data missing	3	0.2 %
-9	Dont Know	250	18.5 %
-8	Refuse	2	0.1 %
	Total	**1,354**	**100%**

Based upon 1,099 valid cases out of 1,354 total cases.

Location: 619-622 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -121 , -9 , -8

S0PSUBB2: S0 SubUse: Relatives complain about dads drinking

Value	Label	Unweighted Frequency	%
1	Yes	342	25.3 %
5	No	488	36.0 %
	Missing Data		
-200	Data missing	3	0.2 %
-116	Individual item not endorsed, follow-up skipped	311	23.0 %
-9	Dont Know	209	15.4 %
-8	Refuse	1	0.1 %
	Total	**1,354**	**100%**

Based upon 830 valid cases out of 1,354 total cases.

Location: 623-626 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -124 , -116 , -9 , -8

S0PSUBC2: S0 SubUse: Dad ever attend A A because of drinking

Value	Label	Unweighted Frequency	%
1	Yes	127	9.4 %
5	No	624	46.1 %
	Missing Data		
-200	Data missing	3	0.2 %
-116	Individual item not endorsed, follow-up skipped	311	23.0 %
-9	Dont Know	288	21.3 %
-8	Refuse	1	0.1 %
	Total	**1,354**	**100%**

Based upon 751 valid cases out of 1,354 total cases.

Location: 627-630 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -124 , -116 , -9 , -8

S0PSUBD2: S0 SubUse: Dads drinking create problems relatives

Value	Label	Unweighted Frequency	%
1	Yes	333	24.6 %
5	No	467	34.5 %
	Missing Data		
-200	Data missing	3	0.2 %
-116	Individual item not endorsed, follow-up skipped	311	23.0 %
-9	Dont Know	239	17.7 %
-8	Refuse	1	0.1 %
	Total	**1,354**	**100%**

Based upon 800 valid cases out of 1,354 total cases.

Location: 631-634 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -124 , -116 , -9 , -8

S0PSUBE2: S0 SubUse: Dad ever in trouble at work for drinking

Value	Label	Unweighted Frequency	%
1	Yes	102	7.5 %
5	No	645	47.6 %

Value	Label	Unweighted Frequency	%
	Missing Data		
-200	Data missing	3	0.2 %
-116	Individual item not endorsed, follow-up skipped	311	23.0 %
-9	Dont Know	292	21.6 %
-8	Refuse	1	0.1 %
	Total	**1,354**	**100%**

Based upon 747 valid cases out of 1,354 total cases.

Location: 635-638 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -124 , -116 , -9 , -8

S0PSUBF2: S0 SubUse: Dad go to anyone for help about drinking

Value	Label	Unweighted Frequency	%
1	Yes	98	7.2 %
5	No	649	47.9 %
	Missing Data		
-200	Data missing	3	0.2 %
-116	Individual item not endorsed, follow-up skipped	311	23.0 %
-9	Dont Know	292	21.6 %
-8	Refuse	1	0.1 %
	Total	**1,354**	**100%**

Based upon 747 valid cases out of 1,354 total cases.

Location: 639-642 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -124 , -116 , -9 , -8

S0PSUBG2: S0 SubUse: Dad been in hosp because of drinking

Value	Label	Unweighted Frequency	%
1	Yes	84	6.2 %
5	No	675	49.9 %
	Missing Data		
-200	Data missing	3	0.2 %
-116	Individual item not endorsed, follow-up skipped	311	23.0 %
-9	Dont Know	280	20.7 %
-8	Refuse	1	0.1 %
	Total	**1,354**	**100%**

Based upon 759 valid cases out of 1,354 total cases.

Location: 643-646 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -124 , -116 , -9 , -8

S0PSUBH2: S0 SubUse: Dad ever arrested for drunken driving

Value	Label	Unweighted Frequency	%
1	Yes	199	14.7 %
5	No	553	40.8 %
	Missing Data		
-200	Data missing	3	0.2 %
-116	Individual item not endorsed, follow-up skipped	311	23.0 %
-9	Dont Know	287	21.2 %
-8	Refuse	1	0.1 %
	Total	**1,354**	**100%**

Based upon 752 valid cases out of 1,354 total cases.

Location: 647-650 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -124 , -116 , -9 , -8

S0PSUBO2: S0 SubUse: Dad ever neglect obligations bc drinking

Value	Label	Unweighted Frequency	%
1	Yes	179	13.2 %
5	No	599	44.2 %
	Missing Data		
-200	Data missing	3	0.2 %
-116	Individual item not endorsed, follow-up skipped	311	23.0 %
-9	Dont Know	261	19.3 %
-8	Refuse	1	0.1 %
	Total	**1,354**	**100%**

Based upon 778 valid cases out of 1,354 total cases.

Location: 651-654 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -124 , -116 , -9 , -8

S0PSUBI2: S0 SubUse: Dad ever arrested for drunken behavior

Value	Label	Unweighted Frequency	%
1	Yes	208	15.4 %
5	No	541	40.0 %
	Missing Data		

Value	Label	Unweighted Frequency	%
-200	Data missing	3	0.2 %
-116	Individual item not endorsed, follow-up skipped	311	23.0 %
-9	Dont Know	290	21.4 %
-8	Refuse	1	0.1 %
	Total	**1,354**	**100%**

Based upon 749 valid cases out of 1,354 total cases.

Location: 655-658 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -124 , -116 , -9 , -8

S0PSUBJ2: S0 SubUse: Do you think dad is an alcoholic now

Value	Label	Unweighted Frequency	%
1	Yes	164	12.1 %
3	Not sure	113	8.3 %
5	No	562	41.5 %
	Missing Data		
-200	Data missing	3	0.2 %
-116	Individual item not endorsed, follow-up skipped	311	23.0 %
-9	Dont Know	200	14.8 %
-8	Refuse	1	0.1 %
	Total	**1,354**	**100%**

Based upon 839 valid cases out of 1,354 total cases.

Location: 659-662 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -124 , -116 , -9 , -8

S0PSUBK2: S0 SubUse: Think dad was an alcoholic in past

Value	Label	Unweighted Frequency	%
1	Yes	423	31.2 %
3	Not sure	92	6.8 %
5	No	344	25.4 %
	Missing Data		
-200	Data missing	3	0.2 %
-116	Individual item not endorsed, follow-up skipped	311	23.0 %
-9	Dont Know	180	13.3 %
-8	Refuse	1	0.1 %
	Total	**1,354**	**100%**

Based upon 859 valid cases out of 1,354 total cases.

Location: 663-666 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -125 , -124 , -116 , -9 , -8

S0PSUBL2: S0 SubUse: Does or did dad use street drugs

Value	Label	Unweighted Frequency	%
1	Yes, in the past year	205	15.1 %
2	Yes, but not in the past year	309	22.8 %
5	No, never	574	42.4 %
	Missing Data		
-200	Data missing	3	0.2 %
-9	Dont Know	261	19.3 %
-8	Refuse	2	0.1 %
	Total	**1,354**	**100%**

Based upon 1,088 valid cases out of 1,354 total cases.

Location: 667-670 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -9 , -8

S0PSUBM2: S0 SubUse: Think dad has a drug problem now

Value	Label	Unweighted Frequency	%
1	Yes	103	7.6 %
3	Not sure	25	1.8 %
5	No	958	70.8 %
	Missing Data		
-200	Data missing	3	0.2 %
-117	Not Applicable	263	19.4 %
-9	Dont Know	2	0.1 %
	Total	**1,354**	**100%**

Based upon 1,086 valid cases out of 1,354 total cases.

Location: 671-674 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -126 , -117 , -9 , -8

S0PSUBN2: S0 SubUse: Think dad had a drug problem in past

Value	Label	Unweighted Frequency	%
1	Yes	346	25.6 %
3	Not sure	38	2.8 %

Value	Label	Unweighted Frequency	%
5	No	702	51.8 %
	Missing Data		
-200	Data missing	3	0.2 %
-117	Not Applicable	263	19.4 %
-9	Dont Know	2	0.1 %
	Total	**1,354**	**100%**

Based upon 1,086 valid cases out of 1,354 total cases.

Location: 675-678 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200 , -127 , -126 , -117 , -9 , -8

S0QFBRMO: S0 SubUse: Quant/freq used beer/wine/liquor past 6 mo

Based upon 1,347 valid cases out of 1,354 total cases.

- Mean: 15.42
- Median: 1.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 315.00
- Standard Deviation: 31.42

Location: 679-682 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -300 , -200 , -100 , -9 , -8

S0EVRUSE: S0 SubUse: Number drugs used in lifetime

Based upon 1,350 valid cases out of 1,354 total cases.

- Mean: 2.14
- Median: 1.00
- Mode: 1.00
- Minimum: 0.00
- Maximum: 9.00
- Standard Deviation: 1.98

Location: 683-686 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -300 , -200 , -100 , -9 , -8

S06MOUSE: S0 SubUse: Number drugs used past 6 mo

Based upon 1,350 valid cases out of 1,354 total cases.

- Mean: 1.19
- Median: 1.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 9.00

- Standard Deviation: 1.52

Location: 687-690 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -300 , -200 , -100 , -9 , -8

S0QFSMK6: S0 SubUse: Quant/Freq used cigarettes past 6 mo

Based upon 1,351 valid cases out of 1,354 total cases.

- Mean: 32.09
- Median: 2.50
- Mode: 0.00
- Minimum: 0.00
- Maximum: 160.00
- Standard Deviation: 49.42

Location: 691-696 (width: 6; decimal: 1)
Variable Type: numeric
(Range of) Missing Values: -300.0 , -200.0 , -100.0 , -9.0 , -8.0

S0ALSC2: S0 SubUse: Lifetime Alcohol ConsequencesL

Based upon 1,351 valid cases out of 1,354 total cases.

- Mean: 2.04
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 15.00
- Standard Deviation: 3.13

Location: 697-700 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200

S0A6SC2: S0 SubUse: Past 6 mo Alcohol consequences

Based upon 1,351 valid cases out of 1,354 total cases.

- Mean: 0.94
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 15.00
- Standard Deviation: 2.10

Location: 701-704 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200

S0DLSC2: S0 SubUse: Lifetime Drug Consequences

Based upon 1,351 valid cases out of 1,354 total cases.

- Mean: 3.33
- Median: 2.00

- Mode: 0.00
- Minimum: 0.00
- Maximum: 16.00
- Standard Deviation: 3.93

Location: 705-708 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200

S0D6SC2: S0 SubUse: Past 6 mo Drug consequences

Based upon 1,351 valid cases out of 1,354 total cases.

- Mean: 1.57
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 15.00
- Standard Deviation: 2.81

Location: 709-712 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200

S0ALDP2: S0 SubUse: Lifetime alcohol dependence symptoms

Based upon 1,351 valid cases out of 1,354 total cases.

- Mean: 0.79
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 10.00
- Standard Deviation: 1.71

Location: 713-716 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200

S0A6DP2: S0 SubUse: Past 6 months alcohol dependence symtpoms

Based upon 1,351 valid cases out of 1,354 total cases.

- Mean: 0.43
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 10.00
- Standard Deviation: 1.35

Location: 717-720 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200

S0DLDP2: S0 SubUse: Lifetime drugs dependence symptoms

Based upon 1,351 valid cases out of 1,354 total cases.

- Mean: 1.85
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 10.00
- Standard Deviation: 2.62

Location: 721-724 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200

S0D6DP2: S0 SubUse: Past 6 months drugs dependence symptoms

Based upon 1,351 valid cases out of 1,354 total cases.

- Mean: 1.03
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 10.00
- Standard Deviation: 2.16

Location: 725-728 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200

S0TOTSCL: S0 SubUse: Lifetime total susbtance use consequences

Based upon 1,351 valid cases out of 1,354 total cases.

- Mean: 4.11
- Median: 3.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 16.00
- Standard Deviation: 4.30

Location: 729-732 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200

S0TOTSC6: S0 SubUse: Past 6 mos total substance use consequences

Based upon 1,351 valid cases out of 1,354 total cases.

- Mean: 1.93
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 15.00
- Standard Deviation: 3.06

Location: 733-736 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200

S0TOTDSL: S0 SubUse: Lifetime total substance use dependency symptoms

Based upon 1,351 valid cases out of 1,354 total cases.

- Mean: 2.08
- Median: 1.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 10.00
- Standard Deviation: 2.71

Location: 737-740 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200

S0TOTDS6: S0 SubUse: Past 6 mos total susbtance use dependency symptoms

Based upon 1,351 valid cases out of 1,354 total cases.

- Mean: 1.16
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 10.00
- Standard Deviation: 2.26

Location: 741-744 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200

S0ALTRTY: S0 SubUse: Lifetime alcohol treatment

Value	Label	Unweighted Frequency	%
0	No	1195	88.3 %
1	Yes	156	11.5 %
	Missing Data		
-200	Data missing	3	0.2 %
	Total	**1,354**	**100%**

Based upon 1,351 valid cases out of 1,354 total cases.

Location: 745-748 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200

S0ALTRTM: S0 SubUse: Past 6 month alcohol treatment

Value	Label	Unweighted Frequency	%
0	No	1276	94.2 %
1	Yes	75	5.5 %
	Missing Data		
-200	Data missing	3	0.2 %
	Total	**1,354**	**100%**

Based upon 1,351 valid cases out of 1,354 total cases.

Location: 749-752 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200

S0DGTRTY: S0 SubUse: Lifetime drug treatment

Value	Label	Unweighted Frequency	%
0	No	1115	82.3 %
1	Yes	236	17.4 %
	Missing Data		
-200	Data missing	3	0.2 %
	Total	**1,354**	**100%**

Based upon 1,351 valid cases out of 1,354 total cases.

Location: 753-756 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200

S0DGTRTM: S0 SubUse: Past 6 month drug treatment

Value	Label	Unweighted Frequency	%
0	No	1225	90.5 %
1	Yes	126	9.3 %
	Missing Data		
-200	Data missing	3	0.2 %
	Total	**1,354**	**100%**

Based upon 1,351 valid cases out of 1,354 total cases.

Location: 757-760 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200

S0TRTY: S0 SubUse: Lifetime Alcohol or Drug Treatment

Value	Label	Unweighted Frequency	%
0	No	1066	78.7 %
1	Yes	285	21.0 %
	Missing Data		
-200	Data missing	3	0.2 %
	Total	**1,354**	**100%**

Based upon 1,351 valid cases out of 1,354 total cases.

Location: 761-764 (width: 4; decimal: 0)
Variable Type: numeric

(Range of) Missing Values: -200

S0TRTM: S0 SubUse: Past 6 Month Alcohol or Drug Treatment

Value	Label	Unweighted Frequency	%
0	No	1198	88.5 %
1	Yes	153	11.3 %
	Missing Data		
-200	Data missing	3	0.2 %
	Total	**1,354**	**100%**

Based upon 1,351 valid cases out of 1,354 total cases.

Location: 765-768 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -200

S0FALSC: S0 SubUse: Alcohol Social Consequences - Father

Based upon 1,154 valid cases out of 1,354 total cases.

- Mean: 1.45
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 9.00
- Standard Deviation: 2.19

Location: 769-772 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -300 , -200 , -100 , -9 , -8

S0MALSC: S0 SubUse: Alcohol Social Consequences - Mother

Based upon 1,321 valid cases out of 1,354 total cases.

- Mean: 0.55
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 9.00
- Standard Deviation: 1.44

Location: 773-776 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -300 , -200 , -100 , -9 , -8

S1SUBUSE1: S1 SubUse: How often had alcohol to drink in recall period

Value	Label	Unweighted Frequency	%
1	Not at all	775	57.2 %
2	1-2 times	160	11.8 %

Value	Label	Unweighted Frequency	%
3	3-5 times	90	6.6 %
4	1X per month	31	2.3 %
5	2-3X per month	63	4.7 %
6	1X per week	54	4.0 %
7	2-3X per week	50	3.7 %
8	4-5X per week	22	1.6 %
9	Every day	16	1.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
	Total	**1,354**	**100%**

Based upon 1,261 valid cases out of 1,354 total cases.

Location: 777-780 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S1SUBUSE2: S1 SubUse: Num alcoholic drinks usually consumed

Based upon 1,260 valid cases out of 1,354 total cases.

- Mean: 2.30
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 40.00
- Standard Deviation: 4.52

Location: 781-784 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -101 , -99 , -98

S1SUBUSE3: S1 SubUse: How often had 5+ drinks at one time in recall period

Value	Label	Unweighted Frequency	%
1	Not at all	947	69.9 %
2	1-2 times	91	6.7 %
3	3-5 times	65	4.8 %
4	1X per month	31	2.3 %
5	2-3X per month	29	2.1 %
6	1X per week	32	2.4 %
7	2-3X per week	37	2.7 %
8	4-5X per week	17	1.3 %
9	Every day	11	0.8 %
	Missing Data		

Value	Label	Unweighted Frequency	%
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-99	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,260 valid cases out of 1,354 total cases.

Location: 785-788 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S1SUBUSE4: S1 SubUse: Num times been drunk on alcohol during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	922	68.1 %
2	1-2 times	135	10.0 %
3	3-5 times	72	5.3 %
4	1X per month	24	1.8 %
5	2-3X per month	34	2.5 %
6	1X per week	27	2.0 %
7	2-3X per week	27	2.0 %
8	4-5X per week	15	1.1 %
9	Every day	5	0.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
	Total	**1,354**	**100%**

Based upon 1,261 valid cases out of 1,354 total cases.

Location: 789-792 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S1SUBUSE6: S1 SubUse: Num times used marijuana/hashish during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	848	62.6 %
2	1-2 times	100	7.4 %
3	3-5 times	68	5.0 %
4	1X per month	21	1.6 %
5	2-3X per month	32	2.4 %
6	1X per week	18	1.3 %
7	2-3X per week	39	2.9 %

Value	Label	Unweighted Frequency	%
8	4-5X per week	45	3.3 %
9	Every day	90	6.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
	Total	**1,354**	**100%**

Based upon 1,261 valid cases out of 1,354 total cases.

Location: 793-796 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S1SUBUSE8: S1 SubUse: Num times used sedative/tranquilizers during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1177	86.9 %
2	1-2 times	27	2.0 %
3	3-5 times	16	1.2 %
4	1X per month	6	0.4 %
5	2-3X per month	12	0.9 %
6	1X per week	6	0.4 %
7	2-3X per week	8	0.6 %
8	4-5X per week	4	0.3 %
9	Every day	4	0.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-99	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,260 valid cases out of 1,354 total cases.

Location: 797-800 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S1SUBUSE10: S1 SubUse: Num times used stimulants/amphetamines during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1189	87.8 %
2	1-2 times	14	1.0 %
3	3-5 times	15	1.1 %
4	1X per month	4	0.3 %

Value	Label	Unweighted Frequency	%
5	2-3X per month	11	0.8 %
6	1X per week	4	0.3 %
7	2-3X per week	11	0.8 %
8	4-5X per week	5	0.4 %
9	Every day	8	0.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
	Total	**1,354**	**100%**

Based upon 1,261 valid cases out of 1,354 total cases.

Location: 801-804 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S1SUBUSE12: S1 SubUse: Num times used cocaine during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1180	87.1 %
2	1-2 times	34	2.5 %
3	3-5 times	13	1.0 %
4	1X per month	4	0.3 %
5	2-3X per month	12	0.9 %
6	1X per week	4	0.3 %
7	2-3X per week	7	0.5 %
8	4-5X per week	5	0.4 %
9	Every day	2	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
	Total	**1,354**	**100%**

Based upon 1,261 valid cases out of 1,354 total cases.

Location: 805-808 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S1SUBUSE14: S1 SubUse: Num times used opiates during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1236	91.3 %
2	1-2 times	11	0.8 %

Value	Label	Unweighted Frequency	%
3	3-5 times	2	0.1 %
4	1X per month	4	0.3 %
5	2-3X per month	1	0.1 %
6	1X per week	0	0.0 %
7	2-3X per week	3	0.2 %
8	4-5X per week	3	0.2 %
9	Every day	1	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
	Total	**1,354**	**100%**

Based upon 1,261 valid cases out of 1,354 total cases.

Location: 809-812 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S1SUBUSE16: S1 SubUse: Num times used ecstasy during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1225	90.5 %
2	1-2 times	23	1.7 %
3	3-5 times	5	0.4 %
4	1X per month	2	0.1 %
5	2-3X per month	3	0.2 %
6	1X per week	1	0.1 %
7	2-3X per week	0	0.0 %
8	4-5X per week	2	0.1 %
9	Every day	0	0.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
	Total	**1,354**	**100%**

Based upon 1,261 valid cases out of 1,354 total cases.

Location: 813-816 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S1SUBUSE18: S1 SubUse: Num times used hallucinogens during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1205	89.0 %
2	1-2 times	29	2.1 %
3	3-5 times	10	0.7 %
4	1X per month	4	0.3 %
5	2-3X per month	5	0.4 %
6	1X per week	2	0.1 %
7	2-3X per week	1	0.1 %
8	4-5X per week	2	0.1 %
9	Every day	3	0.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
	Total	**1,354**	**100%**

Based upon 1,261 valid cases out of 1,354 total cases.

Location: 817-820 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S1SUBUSE20: S1 SubUse: Num times used inhalants during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1232	91.0 %
2	1-2 times	11	0.8 %
3	3-5 times	14	1.0 %
4	1X per month	0	0.0 %
5	2-3X per month	2	0.1 %
6	1X per week	1	0.1 %
7	2-3X per week	0	0.0 %
8	4-5X per week	0	0.0 %
9	Every day	1	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
	Total	**1,354**	**100%**

Based upon 1,261 valid cases out of 1,354 total cases.

Location: 821-824 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S1SUBUSE22: S1 SubUse: Num times used amyl nitrate/odorizers/rush during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1252	92.5 %
2	1-2 times	4	0.3 %
3	3-5 times	4	0.3 %
4	1X per month	0	0.0 %
5	2-3X per month	1	0.1 %
6	1X per week	0	0.0 %
7	2-3X per week	0	0.0 %
8	4-5X per week	0	0.0 %
9	Every day	0	0.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
	Total	**1,354**	**100%**

Based upon 1,261 valid cases out of 1,354 total cases.

Location: 825-828 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S1SUBUSE24: S1 SubUse: Used other drugs during recall period [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	4	0.3 %
5	No	1257	92.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
	Total	**1,354**	**100%**

Based upon 1,261 valid cases out of 1,354 total cases.

Location: 829-832 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE26: S1 SubUse: Num times used other drugs during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1257	92.8 %
2	1-2 times	3	0.2 %
3	3-5 times	0	0.0 %

Value	Label	Unweighted Frequency	%
4	1X per month	0	0.0 %
5	2-3X per month	0	0.0 %
6	1X per week	0	0.0 %
7	2-3X per week	1	0.1 %
8	4-5X per week	0	0.0 %
9	Every day	0	0.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
	Total	**1,354**	**100%**

Based upon 1,261 valid cases out of 1,354 total cases.

Location: 833-836 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S1SUBUSEDY: S1 SubUse: Used any drugs past 24hrs [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	61	4.5 %
5	No	1200	88.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
	Total	**1,354**	**100%**

Based upon 1,261 valid cases out of 1,354 total cases.

Location: 837-840 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_DRUG24MAR: S1 SubUse_Drug24: Used marijuana/hashish past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	54	4.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	89	6.6 %
-104	Skip-in for SUBUSEDY 1	11	0.8 %
-103	Skip-in for SUBUSEDY <> 1	1200	88.6 %
	Total	**1,354**	**100%**

Based upon 54 valid cases out of 1,354 total cases.

Location: 841-844 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_DRUG24TRANQ: S1 SubUse_Drug24: Used sedatives/tranquilizers past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	4	0.3 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	89	6.6 %
-104	Skip-in for SUBUSEDY 1	61	4.5 %
-103	Skip-in for SUBUSEDY <> 1	1200	88.6 %
	Total	**1,354**	**100%**

Based upon 4 valid cases out of 1,354 total cases.

Location: 845-848 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_DRUG24AMPH: S1 SubUse_Drug24: Used stimulants/amphetamines past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	4	0.3 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	89	6.6 %
-104	Skip-in for SUBUSEDY 1	61	4.5 %
-103	Skip-in for SUBUSEDY <> 1	1200	88.6 %
	Total	**1,354**	**100%**

Based upon 4 valid cases out of 1,354 total cases.

Location: 849-852 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_DRUG24COC: S1 SubUse_Drug24: Used Cocaine past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	2	0.1 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	89	6.6 %

Value	Label	Unweighted Frequency	%
-104	Skip-in for SUBUSEDY 1	63	4.7 %
-103	Skip-in for SUBUSEDY <> 1	1200	88.6 %
	Total	**1,354**	**100%**

Based upon 2 valid cases out of 1,354 total cases.

Location: 853-856 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_DRUG24OPI: S1 SubUse_Drug24: Used Opiates past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	89	6.6 %
-104	Skip-in for SUBUSEDY 1	65	4.8 %
-103	Skip-in for SUBUSEDY <> 1	1200	88.6 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 857-860 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_DRUG24ECST: S1 SubUse_Drug24: Used Ecstacy past 24 hours

Value	Label	Unweighted Frequency	%
1	Yes	1	0.1 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	89	6.6 %
-104	Skip-in for SUBUSEDY 1	64	4.7 %
-103	Skip-in for SUBUSEDY <> 1	1200	88.6 %
	Total	**1,354**	**100%**

Based upon 1 valid cases out of 1,354 total cases.

Location: 861-864 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_DRUG24HALL: S1 SubUse_Drug24: Used Hallucinogens past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	1	0.1 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	89	6.6 %
-104	Skip-in for SUBUSEDY 1	64	4.7 %
-103	Skip-in for SUBUSEDY <> 1	1200	88.6 %
	Total	**1,354**	**100%**

Based upon 1 valid cases out of 1,354 total cases.

Location: 865-868 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_DRUG24INH: S1 SubUse_Drug24: Used Inhalants past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	89	6.6 %
-104	Skip-in for SUBUSEDY 1	65	4.8 %
-103	Skip-in for SUBUSEDY <> 1	1200	88.6 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 869-872 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_DRUG24AMYL: S1 SubUse_Drug24: Used Amyl nitrate/poppers/odorizers 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	89	6.6 %
-104	Skip-in for SUBUSEDY 1	65	4.8 %
-103	Skip-in for SUBUSEDY <> 1	1200	88.6 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 873-876 (width: 4; decimal: 0)
Variable Type: numeric

(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_DRUG24OTHER: S1 SubUse_Drug24: Used Other (specify) past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	1	0.1 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	89	6.6 %
-104	Skip-in for SUBUSEDY 1	64	4.7 %
-103	Skip-in for SUBUSEDY <> 1	1200	88.6 %
	Total	**1,354**	**100%**

Based upon 1 valid cases out of 1,354 total cases.

Location: 877-880 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_DRUG24NONE: S1 SubUse_Drug24: Used NA (No drugs) past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	89	6.6 %
-104	Skip-in for SUBUSEDY 1	65	4.8 %
-103	Skip-in for SUBUSEDY <> 1	1200	88.6 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 881-884 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S1SUBUSE29: S1 SubUse: Specify other drug used past 24hrs

Based upon 1,354 valid cases out of 1,354 total cases.

Location: 885-894 (width: 10; decimal: 0)
Variable Type: character

S1SUBUSE30: S1 SubUse: Num times smoked cigarettes during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	694	51.3 %
2	1-2 times	66	4.9 %

Value	Label	Unweighted Frequency	%
3	3-5 times	54	4.0 %
4	1X per month	25	1.8 %
5	2-3X per month	28	2.1 %
6	1X per week	17	1.3 %
7	2-3X per week	54	4.0 %
8	4-5X per week	40	3.0 %
9	Every day	282	20.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-99	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,260 valid cases out of 1,354 total cases.

Location: 895-898 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S1SUBUSE31: S1 SubUse: Num cigarettes smoke per day during recall period

Value	Label	Unweighted Frequency	%
1	0 cigarettes	705	52.1 %
2	1-4 cigarettes	271	20.0 %
3	5-9 cigarettes	108	8.0 %
4	10-15 cigarettes	78	5.8 %
5	16-20 cigarettes	61	4.5 %
6	More than a pack a day cigarettes	38	2.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
	Total	**1,354**	**100%**

Based upon 1,261 valid cases out of 1,354 total cases.

Location: 899-902 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -105 , -9 , -8

S1SUBUSE_SUBSYM01_SUSYMB: S1 SubUse: Family ever complain because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	166	12.3 %

Value	Label	Unweighted Frequency	%
5	No	447	33.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-106	Skip-in for alchstat/drugstat	648	47.9 %
	Total	**1,354**	**100%**

Based upon 613 valid cases out of 1,354 total cases.

Location: 903-906 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM01_SUSYMC: S1 SubUse: Specify cause of family complaint [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	33	2.4 %
2	Drugs	84	6.2 %
3	Both	49	3.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	12	0.9 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-107	Follow-up skipped because gate <> 1	1087	80.3 %
	Total	**1,354**	**100%**

Based upon 166 valid cases out of 1,354 total cases.

Location: 907-910 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM02_SUSYMB: S1 SubUse: Friend ever complain because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	55	4.1 %
5	No	558	41.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-106	Skip-in for alchstat/drugstat	648	47.9 %
	Total	**1,354**	**100%**

Based upon 613 valid cases out of 1,354 total cases.

Location: 911-914 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM02_SUSYMC: S1 SubUse: Specify cause of friend complain [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	13	1.0 %
2	Drugs	25	1.8 %
3	Both	17	1.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	12	0.9 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-107	Follow-up skipped because gate <> 1	1198	88.5 %
	Total	**1,354**	**100%**

Based upon 55 valid cases out of 1,354 total cases.

Location: 915-918 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM03_SUSYMB: S1 SubUse: Ever trouble at work because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	40	3.0 %
5	No	573	42.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-106	Skip-in for alchstat/drugstat	648	47.9 %
	Total	**1,354**	**100%**

Based upon 613 valid cases out of 1,354 total cases.

Location: 919-922 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM03_SUSYMC: S1 SubUse: Specify cause of trouble at work [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	2	0.1 %

Value	Label	Unweighted Frequency	%
2	Drugs	28	2.1 %
3	Both	10	0.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	12	0.9 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-107	Follow-up skipped because gate <> 1	1213	89.6 %
	Total	**1,354**	**100%**

Based upon 40 valid cases out of 1,354 total cases.

Location: 923-926 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM04_SUSYMB: S1 SubUse: Ever missed work/school because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	71	5.2 %
5	No	542	40.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-106	Skip-in for alchstat/drugstat	648	47.9 %
	Total	**1,354**	**100%**

Based upon 613 valid cases out of 1,354 total cases.

Location: 927-930 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM04_SUSYMC: S1 SubUse: Specify cause of missed work/school [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	22	1.6 %
2	Drugs	30	2.2 %
3	Both	19	1.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	12	0.9 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-107	Follow-up skipped because gate <> 1	1182	87.3 %
	Total	**1,354**	**100%**

Based upon 71 valid cases out of 1,354 total cases.

Location: 931-934 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM05_SUSYMB: S1 SubUse: Ever arrested because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	59	4.4 %
5	No	554	40.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-106	Skip-in for alchstat/drugstat	648	47.9 %
	Total	**1,354**	**100%**

Based upon 613 valid cases out of 1,354 total cases.

Location: 935-938 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM05_SUSYMC: S1 SubUse: Specify cause of arrest [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	12	0.9 %
2	Drugs	36	2.7 %
3	Both	11	0.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	12	0.9 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-107	Follow-up skipped because gate <> 1	1194	88.2 %
	Total	**1,354**	**100%**

Based upon 59 valid cases out of 1,354 total cases.

Location: 939-942 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM06_SUSYMB: S1 SubUse: Ever argue family/friends because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	118	8.7 %
5	No	495	36.6 %

Value	Label	Unweighted Frequency	%
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-106	Skip-in for alchstat/drugstat	648	47.9 %
	Total	**1,354**	**100%**

Based upon 613 valid cases out of 1,354 total cases.

Location: 943-946 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM06_SUSYMC: S1 SubUse: Specify cause of argument w/ family/friends [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	30	2.2 %
2	Drugs	59	4.4 %
3	Both	29	2.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	12	0.9 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-107	Follow-up skipped because gate <> 1	1135	83.8 %
	Total	**1,354**	**100%**

Based upon 118 valid cases out of 1,354 total cases.

Location: 947-950 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM07_SUSYMB: S1 SubUse: Ever cont. drug/alcohol use in spite cause health problems in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	84	6.2 %
5	No	529	39.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-106	Skip-in for alchstat/drugstat	648	47.9 %
	Total	**1,354**	**100%**

Based upon 613 valid cases out of 1,354 total cases.

Location: 951-954 (width: 4; decimal: 0)

Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM07_SUSYMC: S1 SubUse: Specify continued using [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	11	0.8 %
2	Drugs	50	3.7 %
3	Both	23	1.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	11	0.8 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-107	Follow-up skipped because gate <> 1	1170	86.4 %
	Total	**1,354**	**100%**

Based upon 84 valid cases out of 1,354 total cases.

Location: 955-958 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM08_SUSYMB: S1 SubUse: Ever not remember prior night because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	152	11.2 %
5	No	461	34.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-106	Skip-in for alchstat/drugstat	648	47.9 %
	Total	**1,354**	**100%**

Based upon 613 valid cases out of 1,354 total cases.

Location: 959-962 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM08_SUSYMC: S1 SubUse: Specify cause of not remember night [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	71	5.2 %
2	Drugs	33	2.4 %
3	Both	48	3.5 %
	Missing Data		

Value	Label	Unweighted Frequency	%
-202	IntCompStat = 2 partial interview, data missing	12	0.9 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-107	Follow-up skipped because gate <> 1	1101	81.3 %
	Total	**1,354**	**100%**

Based upon 152 valid cases out of 1,354 total cases.

Location: 963-966 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM09_SUSYMB: S1 SubUse: Ever used in situations could get hurt in recall period [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	149	11.0 %
5	No	464	34.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-106	Skip-in for alchstat/drugstat	648	47.9 %
	Total	**1,354**	**100%**

Based upon 613 valid cases out of 1,354 total cases.

Location: 967-970 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM09_SUSYMC: S1 SubUse: Specify used drugs in hurtful situations [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	49	3.6 %
2	Drugs	45	3.3 %
3	Both	55	4.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	12	0.9 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-107	Follow-up skipped because gate <> 1	1104	81.5 %
	Total	**1,354**	**100%**

Based upon 149 valid cases out of 1,354 total cases.

Location: 971-974 (width: 4; decimal: 0)
Variable Type: numeric

(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM10_SUSYMB: S1 SubUse: Ever had accident/injury because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	41	3.0 %
5	No	572	42.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-106	Skip-in for alchstat/drugstat	648	47.9 %
	Total	**1,354**	**100%**

Based upon 613 valid cases out of 1,354 total cases.

Location: 975-978 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM10_SUSYMC: S1 SubUse: Specify cause of accident/injury [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	20	1.5 %
2	Drugs	8	0.6 %
3	Both	13	1.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	12	0.9 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-107	Follow-up skipped because gate <> 1	1212	89.5 %
	Total	**1,354**	**100%**

Based upon 41 valid cases out of 1,354 total cases.

Location: 979-982 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM11_SUSYMB: S1 SubUse: Alcohol/drug use interfere work/responsibility school/home/job in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	78	5.8 %
5	No	535	39.5 %
	Missing Data		

Value	Label	Unweighted Frequency	%
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-106	Skip-in for alchstat/drugstat	648	47.9 %
	Total	**1,354**	**100%**

Based upon 613 valid cases out of 1,354 total cases.

Location: 983-986 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM11_SUSYMC: S1 SubUse: Specify cause of interference [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	12	0.9 %
2	Drugs	45	3.3 %
3	Both	21	1.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	12	0.9 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-107	Follow-up skipped because gate <> 1	1175	86.8 %
	Total	**1,354**	**100%**

Based upon 78 valid cases out of 1,354 total cases.

Location: 987-990 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM12_SUSYMB: S1 SubUse: Ever trouble police no arrest because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	68	5.0 %
5	No	545	40.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-106	Skip-in for alchstat/drugstat	648	47.9 %
	Total	**1,354**	**100%**

Based upon 613 valid cases out of 1,354 total cases.

Location: 991-994 (width: 4; decimal: 0)
Variable Type: numeric

(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM12_SUSYMC: S1 SubUse: Specify cause trouble police no arrest [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	28	2.1 %
2	Drugs	29	2.1 %
3	Both	11	0.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	12	0.9 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-107	Follow-up skipped because gate <> 1	1185	87.5 %
	Total	**1,354**	**100%**

Based upon 68 valid cases out of 1,354 total cases.

Location: 995-998 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM13_SUSYMB: S1 SubUse: Sold/exchanged sex for drugs in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	2	0.1 %
5	No	611	45.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-106	Skip-in for alchstat/drugstat	648	47.9 %
	Total	**1,354**	**100%**

Based upon 613 valid cases out of 1,354 total cases.

Location: 999-1002 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM13_SUSYMC: S1 SubUse: Specify cause sold/exchanged sex [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	0	0.0 %
2	Drugs	2	0.1 %
3	Both	0	0.0 %
	Missing Data		

Value	Label	Unweighted Frequency	%
-202	IntCompStat = 2 partial interview, data missing	12	0.9 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-107	Follow-up skipped because gate <> 1	1251	92.4 %
	Total	**1,354**	**100%**

Based upon 2 valid cases out of 1,354 total cases.

Location: 1003-1006 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM14_SUSYMB: S1 SubUse: Sold drugs for money buy drugs in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	92	6.8 %
5	No	521	38.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-106	Skip-in for alchstat/drugstat	648	47.9 %
	Total	**1,354**	**100%**

Based upon 613 valid cases out of 1,354 total cases.

Location: 1007-1010 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM14_SUSYMC: S1 SubUse: Specify cause sold drugs for money to buy [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	1	0.1 %
2	Drugs	79	5.8 %
3	Both	12	0.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	12	0.9 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-107	Follow-up skipped because gate <> 1	1161	85.7 %
	Total	**1,354**	**100%**

Based upon 92 valid cases out of 1,354 total cases.

Location: 1011-1014 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM15_SUSYMB: S1 SubUse: Stole for money buy drugs in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	51	3.8 %
5	No	562	41.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-106	Skip-in for alchstat/drugstat	648	47.9 %
	Total	**1,354**	**100%**

Based upon 613 valid cases out of 1,354 total cases.

Location: 1015-1018 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM15_SUSYMC: S1 SubUse: Specify cause stole for money to buy [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	3	0.2 %
2	Drugs	39	2.9 %
3	Both	9	0.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	12	0.9 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-107	Follow-up skipped because gate <> 1	1202	88.8 %
	Total	**1,354**	**100%**

Based upon 51 valid cases out of 1,354 total cases.

Location: 1019-1022 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM16_SUSYMB: S1 SubUse: Ever had physical fight because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	74	5.5 %
5	No	539	39.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %

Value	Label	Unweighted Frequency	%
-106	Skip-in for alchstat/drugstat	648	47.9 %
	Total	**1,354**	**100%**

Based upon 613 valid cases out of 1,354 total cases.

Location: 1023-1026 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM16_SUSYMC: S1 SubUse: Specify cause of fight [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	32	2.4 %
2	Drugs	26	1.9 %
3	Both	16	1.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	12	0.9 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-107	Follow-up skipped because gate <> 1	1179	87.1 %
	Total	**1,354**	**100%**

Based upon 74 valid cases out of 1,354 total cases.

Location: 1027-1030 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM17_SUSYMB: S1 SubUse: Ever destroy property because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	42	3.1 %
5	No	571	42.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-106	Skip-in for alchstat/drugstat	648	47.9 %
	Total	**1,354**	**100%**

Based upon 613 valid cases out of 1,354 total cases.

Location: 1031-1034 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM17_SUSYMC: S1 SubUse: Specify cause of destroy property [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	17	1.3 %
2	Drugs	11	0.8 %
3	Both	14	1.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	12	0.9 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-107	Follow-up skipped because gate <> 1	1211	89.4 %
	Total	**1,354**	**100%**

Based upon 42 valid cases out of 1,354 total cases.

Location: 1035-1038 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM18_SUSYMB: S1 SubUse: Ever felt strong urge use drug/alcohol no self stop in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	75	5.5 %
5	No	538	39.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-106	Skip-in for alchstat/drugstat	648	47.9 %
	Total	**1,354**	**100%**

Based upon 613 valid cases out of 1,354 total cases.

Location: 1039-1042 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM18_SUSYMC: S1 SubUse: Specify the cause of strong urge [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	10	0.7 %
2	Drugs	54	4.0 %
3	Both	11	0.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	12	0.9 %

Value	Label	Unweighted Frequency	%
-201	IntCompStat = 4 missed interview	89	6.6 %
-107	Follow-up skipped because gate <> 1	1178	87.0 %
	Total	**1,354**	**100%**

Based upon 75 valid cases out of 1,354 total cases.

Location: 1043-1046 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM19_SUSYMB: S1 SubUse: Ever want drug/alcohol think of nothing else in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	64	4.7 %
5	No	549	40.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-106	Skip-in for alchstat/drugstat	648	47.9 %
	Total	**1,354**	**100%**

Based upon 613 valid cases out of 1,354 total cases.

Location: 1047-1050 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM19_SUSYMC: S1 SubUse: Specify cause of want [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	14	1.0 %
2	Drugs	41	3.0 %
3	Both	9	0.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	12	0.9 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-107	Follow-up skipped because gate <> 1	1189	87.8 %
	Total	**1,354**	**100%**

Based upon 64 valid cases out of 1,354 total cases.

Location: 1051-1054 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM20_SUSYMB: S1 SubUse: Ever experience withdrawal from drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	53	3.9 %
5	No	560	41.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-106	Skip-in for alchstat/drugstat	648	47.9 %
	Total	**1,354**	**100%**

Based upon 613 valid cases out of 1,354 total cases.

Location: 1055-1058 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM20_SUSYMC: S1 SubUse: Specify cause of withdraw [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	9	0.7 %
2	Drugs	40	3.0 %
3	Both	4	0.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	12	0.9 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-107	Follow-up skipped because gate <> 1	1200	88.6 %
	Total	**1,354**	**100%**

Based upon 53 valid cases out of 1,354 total cases.

Location: 1059-1062 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM21_SUSYMB: S1 SubUse: Take drug/alcohol to keep from having withdraw in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	31	2.3 %
5	No	22	1.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %
-201	IntCompStat = 4 missed interview	89	6.6 %

Value	Label	Unweighted Frequency	%
-108	Skip-in for SUBSYM20=5	556	41.1 %
-106	Skip-in for alchstat/drugstat	648	47.9 %
	Total	**1,354**	**100%**

Based upon 53 valid cases out of 1,354 total cases.

Location: 1063-1066 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM21_SUSYMC: S1 SubUse: Specify reason for consumption [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	4	0.3 %
2	Drugs	22	1.6 %
3	Both	5	0.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	12	0.9 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-107	Follow-up skipped because gate <> 1	1222	90.3 %
	Total	**1,354**	**100%**

Based upon 31 valid cases out of 1,354 total cases.

Location: 1067-1070 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM22_SUSYMB: S1 SubUse: Give up usual activities/interests to use drug/alcohol in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	57	4.2 %
5	No	556	41.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-106	Skip-in for alchstat/drugstat	648	47.9 %
	Total	**1,354**	**100%**

Based upon 613 valid cases out of 1,354 total cases.

Location: 1071-1074 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM22_SUSYMC: S1 SubUse: Specify cause of giving up activities/interests [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	9	0.7 %
2	Drugs	30	2.2 %
3	Both	18	1.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	12	0.9 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-107	Follow-up skipped because gate <> 1	1196	88.3 %
	Total	**1,354**	**100%**

Based upon 57 valid cases out of 1,354 total cases.

Location: 1075-1078 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM23_SUSYMB: S1 SubUse: Ever used in rp when promised wouldn't [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	156	11.5 %
5	No	457	33.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-106	Skip-in for alchstat/drugstat	648	47.9 %
	Total	**1,354**	**100%**

Based upon 613 valid cases out of 1,354 total cases.

Location: 1079-1082 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM23_SUSYMC: S1 SubUse: Specify what use you started [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	31	2.3 %
2	Drugs	82	6.1 %
3	Both	43	3.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	12	0.9 %

Value	Label	Unweighted Frequency	%
-201	IntCompStat = 4 missed interview	89	6.6 %
-107	Follow-up skipped because gate <> 1	1097	81.0 %
	Total	**1,354**	**100%**

Based upon 156 valid cases out of 1,354 total cases.

Location: 1083-1086 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM24_SUSYMB: S1 SubUse: Ever use drug/alcohol more freq than intended in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	121	8.9 %
5	No	492	36.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-106	Skip-in for alchstat/drugstat	648	47.9 %
	Total	**1,354**	**100%**

Based upon 613 valid cases out of 1,354 total cases.

Location: 1087-1090 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM24_SUSYMC: S1 SubUse: Specify what you used more frequently [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	24	1.8 %
2	Drugs	68	5.0 %
3	Both	29	2.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	12	0.9 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-107	Follow-up skipped because gate <> 1	1132	83.6 %
	Total	**1,354**	**100%**

Based upon 121 valid cases out of 1,354 total cases.

Location: 1091-1094 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM25_SUSYMB: S1 SubUse: Ever tried cut down drug/alcohol use but couldnt in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	88	6.5 %
5	No	525	38.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-106	Skip-in for alchstat/drugstat	648	47.9 %
	Total	**1,354**	**100%**

Based upon 613 valid cases out of 1,354 total cases.

Location: 1095-1098 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM25_SUSYMC: S1 SubUse: Specify what substance use you tried to curtail [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	11	0.8 %
2	Drugs	62	4.6 %
3	Both	15	1.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	12	0.9 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-107	Follow-up skipped because gate <> 1	1165	86.0 %
	Total	**1,354**	**100%**

Based upon 88 valid cases out of 1,354 total cases.

Location: 1099-1102 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM26_SUSYMB: S1 SubUse: Ever need more drug/alcohol get same effect in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	132	9.7 %
5	No	481	35.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %

Value	Label	Unweighted Frequency	%
-106	Skip-in for alchstat/drugstat	648	47.9 %
	Total	**1,354**	**100%**

Based upon 613 valid cases out of 1,354 total cases.

Location: 1103-1106 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM26_SUSYMC: S1 SubUse: Specify substance need more for same effect [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	27	2.0 %
2	Drugs	71	5.2 %
3	Both	34	2.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	12	0.9 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-107	Follow-up skipped because gate <> 1	1121	82.8 %
	Total	**1,354**	**100%**

Based upon 132 valid cases out of 1,354 total cases.

Location: 1107-1110 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM27_SUSYMB: S1 SubUse: Ever used drug/alcohol enough to depend in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	53	3.9 %
5	No	560	41.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-106	Skip-in for alchstat/drugstat	648	47.9 %
	Total	**1,354**	**100%**

Based upon 613 valid cases out of 1,354 total cases.

Location: 1111-1114 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM27_SUSYMC: S1 SubUse: Specify dependency [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	4	0.3 %
2	Drugs	38	2.8 %
3	Both	11	0.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	12	0.9 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-107	Follow-up skipped because gate <> 1	1200	88.6 %
	Total	**1,354**	**100%**

Based upon 53 valid cases out of 1,354 total cases.

Location: 1115-1118 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM28_SUSYMB: S1 SubUse: Ever talked doctor/clergy/guidance counselor drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	67	4.9 %
5	No	546	40.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-106	Skip-in for alchstat/drugstat	648	47.9 %
	Total	**1,354**	**100%**

Based upon 613 valid cases out of 1,354 total cases.

Location: 1119-1122 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM28_SUSYMC: S1 SubUse: Specify talked doctor/clergy/guidance counselor [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	11	0.8 %
2	Drugs	38	2.8 %
3	Both	18	1.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	12	0.9 %

Value	Label	Unweighted Frequency	%
-201	IntCompStat = 4 missed interview	89	6.6 %
-107	Follow-up skipped because gate <> 1	1186	87.6 %
	Total	**1,354**	**100%**

Based upon 67 valid cases out of 1,354 total cases.

Location: 1123-1126 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM29_SUSYMB: S1 SubUse: Ever attended self-help group for drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	98	7.2 %
5	No	515	38.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-106	Skip-in for alchstat/drugstat	648	47.9 %
	Total	**1,354**	**100%**

Based upon 613 valid cases out of 1,354 total cases.

Location: 1127-1130 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM29_SUSYMC: S1 SubUse: Specify reason for self-help [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	21	1.6 %
2	Drugs	52	3.8 %
3	Both	25	1.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	12	0.9 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-107	Follow-up skipped because gate <> 1	1155	85.3 %
	Total	**1,354**	**100%**

Based upon 98 valid cases out of 1,354 total cases.

Location: 1131-1134 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM30_SUSYMB: S1 SubUse: Ever treated by therapist because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	61	4.5 %
5	No	552	40.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-106	Skip-in for alchstat/drugstat	648	47.9 %
	Total	**1,354**	**100%**

Based upon 613 valid cases out of 1,354 total cases.

Location: 1135-1138 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM30_SUSYMC: S1 SubUse: Specify reason for therapist visit [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	9	0.7 %
2	Drugs	30	2.2 %
3	Both	22	1.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	12	0.9 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-107	Follow-up skipped because gate <> 1	1192	88.0 %
	Total	**1,354**	**100%**

Based upon 61 valid cases out of 1,354 total cases.

Location: 1139-1142 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM31_SUSYMB: S1 SubUse: Ever Admitted to hospital because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	7	0.5 %
5	No	606	44.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %

Value	Label	Unweighted Frequency	%
-106	Skip-in for alchstat/drugstat	648	47.9 %
	Total	**1,354**	**100%**

Based upon 613 valid cases out of 1,354 total cases.

Location: 1143-1146 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE_SUBSYM31_SUSYMC: S1 SubUse: Specify reason admitted to hospital [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	3	0.2 %
2	Drugs	3	0.2 %
3	Both	1	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	12	0.9 %
-201	IntCompStat = 4 missed interview	89	6.6 %
-107	Follow-up skipped because gate <> 1	1246	92.0 %
	Total	**1,354**	**100%**

Based upon 7 valid cases out of 1,354 total cases.

Location: 1147-1150 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S1SUBUSE28: S1 SubUse: Num times abused prescribed medications (own or someone elses) during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	0	0.0 %
2	1-2 times	0	0.0 %
3	3-5 times	1	0.1 %
4	1X per month	0	0.0 %
5	2-3X per month	0	0.0 %
6	1X per week	0	0.0 %
7	2-3X per week	1	0.1 %
8	4-5X per week	0	0.0 %
9	Every day	0	0.0 %
	Missing Data		
-800	Question not asked for this subject	1259	93.0 %
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %

Value	Label	Unweighted Frequency	%
-201	IntCompStat = 4 missed interview	89	6.6 %
	Total	**1,354**	**100%**

Based upon 2 valid cases out of 1,354 total cases.

Location: 1151-1154 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S1QFBRMO: S1 SubUse: Quant/freq used beer/wine/liquor - recall period

Based upon 1,260 valid cases out of 1,354 total cases.

- Mean: 8.69
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 240.00
- Standard Deviation: 22.17

Location: 1155-1158 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -300 , -202 , -201 , -100 , -9 , -8

S16MOUSE: S1 SubUse: Number drugs used - recall period

Based upon 1,261 valid cases out of 1,354 total cases.

- Mean: 0.64
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 9.00
- Standard Deviation: 1.20

Location: 1159-1162 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -300 , -202 , -201 , -100 , -9 , -8

S1QFSMK6: S1 SubUse: Quant/freq used cigarettes - recall period

Based upon 1,260 valid cases out of 1,354 total cases.

- Mean: 22.23
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 160.00
- Standard Deviation: 41.25

Location: 1163-1168 (width: 6; decimal: 1)
Variable Type: numeric
(Range of) Missing Values: -300.0 , -202.0 , -201.0 , -100.0 , -9.0 , -8.0

S1A6SC2: S1 SubUse: Recall period alcohol consequences

Based upon 1,261 valid cases out of 1,354 total cases.

- Mean: 0.57
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 14.00
- Standard Deviation: 1.60

Location: 1169-1172 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S1D6SC2: S1 SubUse: Recall period drug consequences

Based upon 1,261 valid cases out of 1,354 total cases.

- Mean: 0.78
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 13.00
- Standard Deviation: 2.04

Location: 1173-1176 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S1A6DP2: S1 SubUse: Recall period alcohol dependence symptoms

Based upon 1,261 valid cases out of 1,354 total cases.

- Mean: 0.26
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 10.00
- Standard Deviation: 0.98

Location: 1177-1180 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S1D6DP2: S1 SubUse: Recall period drug dependence symptoms

Based upon 1,261 valid cases out of 1,354 total cases.

- Mean: 0.54
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 10.00
- Standard Deviation: 1.59

Location: 1181-1184 (width: 4; decimal: 0)

Variable Type: numeric
(Range of) Missing Values: -202 , -201

S1TOTSC6: S1 SubUse: Recall period total substance use consequences

Based upon 1,261 valid cases out of 1,354 total cases.

- Mean: 1.06
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 14.00
- Standard Deviation: 2.36

Location: 1185-1188 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S1TOTDS6: S1 SubUse: Recall period total substance use dependency symp

Based upon 1,261 valid cases out of 1,354 total cases.

- Mean: 0.66
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 10.00
- Standard Deviation: 1.72

Location: 1189-1192 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S1ALTRTM: S1 SubUse: Recall period alcohol treatment

Value	Label	Unweighted Frequency	%
0	No	1197	88.4 %
1	Yes	64	4.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
	Total	**1,354**	**100%**

Based upon 1,261 valid cases out of 1,354 total cases.

Location: 1193-1196 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S1DGTRTM: S1 SubUse: Recall period drug treatment

Value	Label	Unweighted Frequency	%
0	No	1163	85.9 %

Value	Label	Unweighted Frequency	%
1	Yes	98	7.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
	Total	**1,354**	**100%**

Based upon 1,261 valid cases out of 1,354 total cases.

Location: 1197-1200 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S1TRTM: S1 SubUse: Recall period alcohol or drug treatment

Value	Label	Unweighted Frequency	%
0	No	1138	84.0 %
1	Yes	123	9.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	89	6.6 %
	Total	**1,354**	**100%**

Based upon 1,261 valid cases out of 1,354 total cases.

Location: 1201-1204 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S2SUBUSE1: S2 SubUse: How often had alcohol to drink in recall period

Value	Label	Unweighted Frequency	%
1	Not at all	737	54.4 %
2	1-2 times	157	11.6 %
3	3-5 times	97	7.2 %
4	1X per month	47	3.5 %
5	2-3X per month	76	5.6 %
6	1X per week	47	3.5 %
7	2-3X per week	67	4.9 %
8	4-5X per week	23	1.7 %
9	Every day	9	0.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
	Total	**1,354**	**100%**

Based upon 1,260 valid cases out of 1,354 total cases.

Location: 1205-1208 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S2SUBUSE2: S2 SubUse: Num alcoholic drinks usually consumed

Based upon 1,255 valid cases out of 1,354 total cases.

- Mean: 2.46
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 60.00
- Standard Deviation: 4.67

Location: 1209-1212 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -101 , -99 , -98

S2SUBUSE3: S2 SubUse: How often had 5+ drinks at one time in recall period

Value	Label	Unweighted Frequency	%
1	Not at all	928	68.5 %
2	1-2 times	111	8.2 %
3	3-5 times	46	3.4 %
4	1X per month	30	2.2 %
5	2-3X per month	44	3.2 %
6	1X per week	35	2.6 %
7	2-3X per week	39	2.9 %
8	4-5X per week	21	1.6 %
9	Every day	6	0.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
	Total	**1,354**	**100%**

Based upon 1,260 valid cases out of 1,354 total cases.

Location: 1213-1216 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S2SUBUSE4: S2 SubUse: Num times been drunk on alcohol during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	901	66.5 %
2	1-2 times	133	9.8 %

Value	Label	Unweighted Frequency	%
3	3-5 times	87	6.4 %
4	1X per month	30	2.2 %
5	2-3X per month	38	2.8 %
6	1X per week	26	1.9 %
7	2-3X per week	32	2.4 %
8	4-5X per week	11	0.8 %
9	Every day	2	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
	Total	**1,354**	**100%**

Based upon 1,260 valid cases out of 1,354 total cases.

Location: 1217-1220 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S2SUBUSE6: S2 SubUse: Num times used marijuana/hashish during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	848	62.6 %
2	1-2 times	86	6.4 %
3	3-5 times	49	3.6 %
4	1X per month	26	1.9 %
5	2-3X per month	37	2.7 %
6	1X per week	26	1.9 %
7	2-3X per week	49	3.6 %
8	4-5X per week	37	2.7 %
9	Every day	100	7.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-99	Dont know	2	0.1 %
	Total	**1,354**	**100%**

Based upon 1,258 valid cases out of 1,354 total cases.

Location: 1221-1224 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S2SUBUSE8: S2 SubUse: Num times used sedative/tranquilizers during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1193	88.1 %
2	1-2 times	23	1.7 %
3	3-5 times	10	0.7 %
4	1X per month	9	0.7 %
5	2-3X per month	6	0.4 %
6	1X per week	5	0.4 %
7	2-3X per week	9	0.7 %
8	4-5X per week	2	0.1 %
9	Every day	3	0.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
	Total	**1,354**	**100%**

Based upon 1,260 valid cases out of 1,354 total cases.

Location: 1225-1228 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S2SUBUSE10: S2 SubUse: Num times used stimulants/amphetamines during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1189	87.8 %
2	1-2 times	19	1.4 %
3	3-5 times	9	0.7 %
4	1X per month	6	0.4 %
5	2-3X per month	8	0.6 %
6	1X per week	1	0.1 %
7	2-3X per week	8	0.6 %
8	4-5X per week	11	0.8 %
9	Every day	9	0.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
	Total	**1,354**	**100%**

Based upon 1,260 valid cases out of 1,354 total cases.

Location: 1229-1232 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S2SUBUSE12: S2 SubUse: Num times used cocaine during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1202	88.8 %
2	1-2 times	23	1.7 %
3	3-5 times	7	0.5 %
4	1X per month	4	0.3 %
5	2-3X per month	6	0.4 %
6	1X per week	5	0.4 %
7	2-3X per week	7	0.5 %
8	4-5X per week	2	0.1 %
9	Every day	4	0.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
	Total	**1,354**	**100%**

Based upon 1,260 valid cases out of 1,354 total cases.

Location: 1233-1236 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S2SUBUSE14: S2 SubUse: Num times used opiates during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1238	91.4 %
2	1-2 times	7	0.5 %
3	3-5 times	4	0.3 %
4	1X per month	1	0.1 %
5	2-3X per month	2	0.1 %
6	1X per week	0	0.0 %
7	2-3X per week	3	0.2 %
8	4-5X per week	3	0.2 %
9	Every day	2	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
	Total	**1,354**	**100%**

Based upon 1,260 valid cases out of 1,354 total cases.

Location: 1237-1240 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S2SUBUSE16: S2 SubUse: Num times used ecstasy during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1228	90.7 %
2	1-2 times	15	1.1 %
3	3-5 times	6	0.4 %
4	1X per month	5	0.4 %
5	2-3X per month	3	0.2 %
6	1X per week	0	0.0 %
7	2-3X per week	0	0.0 %
8	4-5X per week	1	0.1 %
9	Every day	2	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
	Total	**1,354**	**100%**

Based upon 1,260 valid cases out of 1,354 total cases.

Location: 1241-1244 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S2SUBUSE18: S2 SubUse: Num times used hallucinogens during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1196	88.3 %
2	1-2 times	32	2.4 %
3	3-5 times	15	1.1 %
4	1X per month	4	0.3 %
5	2-3X per month	7	0.5 %
6	1X per week	2	0.1 %
7	2-3X per week	1	0.1 %
8	4-5X per week	1	0.1 %
9	Every day	2	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
	Total	**1,354**	**100%**

Based upon 1,260 valid cases out of 1,354 total cases.

Location: 1245-1248 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S2SUBUSE20: S2 SubUse: Num times used inhalants during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1239	91.5 %
2	1-2 times	8	0.6 %
3	3-5 times	4	0.3 %
4	1X per month	2	0.1 %
5	2-3X per month	4	0.3 %
6	1X per week	0	0.0 %
7	2-3X per week	1	0.1 %
8	4-5X per week	0	0.0 %
9	Every day	2	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
	Total	**1,354**	**100%**

Based upon 1,260 valid cases out of 1,354 total cases.

Location: 1249-1252 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S2SUBUSE22: S2 SubUse: Num times used amyl nitrate/odorizers/rush during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1249	92.2 %
2	1-2 times	3	0.2 %
3	3-5 times	1	0.1 %
4	1X per month	1	0.1 %
5	2-3X per month	4	0.3 %
6	1X per week	0	0.0 %
7	2-3X per week	0	0.0 %
8	4-5X per week	0	0.0 %
9	Every day	2	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
	Total	**1,354**	**100%**

Based upon 1,260 valid cases out of 1,354 total cases.

Location: 1253-1256 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S2SUBUSE24: S2 SubUse: Used other drugs during recall period [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	4	0.3 %
5	No	1256	92.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
	Total	**1,354**	**100%**

Based upon 1,260 valid cases out of 1,354 total cases.

Location: 1257-1260 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE26: S2 SubUse: Num times used other drugs during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1256	92.8 %
2	1-2 times	1	0.1 %
3	3-5 times	2	0.1 %
4	1X per month	0	0.0 %
5	2-3X per month	1	0.1 %
6	1X per week	0	0.0 %
7	2-3X per week	0	0.0 %
8	4-5X per week	0	0.0 %
9	Every day	0	0.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
	Total	**1,354**	**100%**

Based upon 1,260 valid cases out of 1,354 total cases.

Location: 1261-1264 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S2SUBUSEDY: S2 SubUse: Used any drugs past 24hrs [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	84	6.2 %
5	No	1176	86.9 %
	Missing Data		

Value	Label	Unweighted Frequency	%
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
	Total	**1,354**	**100%**

Based upon 1,260 valid cases out of 1,354 total cases.

Location: 1265-1268 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_DRUG24MAR: S2 SubUse_Drug24: Used marijuana/hashish past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	78	5.8 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	92	6.8 %
-104	Skip-in for SUBUSEDY 1	8	0.6 %
-103	Skip-in for SUBUSEDY <> 1	1176	86.9 %
	Total	**1,354**	**100%**

Based upon 78 valid cases out of 1,354 total cases.

Location: 1269-1272 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_DRUG24TRANQ: S2 SubUse_Drug24: Used sedatives/tranquilizers past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	1	0.1 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	92	6.8 %
-104	Skip-in for SUBUSEDY 1	85	6.3 %
-103	Skip-in for SUBUSEDY <> 1	1176	86.9 %
	Total	**1,354**	**100%**

Based upon 1 valid cases out of 1,354 total cases.

Location: 1273-1276 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_DRUG24AMPH: S2 SubUse_Drug24: Used stimulants/amphetamines past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	4	0.3 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	92	6.8 %
-104	Skip-in for SUBUSEDY 1	82	6.1 %
-103	Skip-in for SUBUSEDY <> 1	1176	86.9 %
	Total	**1,354**	**100%**

Based upon 4 valid cases out of 1,354 total cases.

Location: 1277-1280 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_DRUG24COC: S2 SubUse_Drug24: Used Cocaine past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	3	0.2 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	92	6.8 %
-104	Skip-in for SUBUSEDY 1	83	6.1 %
-103	Skip-in for SUBUSEDY <> 1	1176	86.9 %
	Total	**1,354**	**100%**

Based upon 3 valid cases out of 1,354 total cases.

Location: 1281-1284 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_DRUG24OPI: S2 SubUse_Drug24: Used Opiates past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	1	0.1 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	92	6.8 %
-104	Skip-in for SUBUSEDY 1	85	6.3 %
-103	Skip-in for SUBUSEDY <> 1	1176	86.9 %
	Total	**1,354**	**100%**

Based upon 1 valid cases out of 1,354 total cases.

Location: 1285-1288 (width: 4; decimal: 0)
Variable Type: numeric

(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_DRUG24ECST: S2 SubUse_Drug24: Used Ecstacy past 24 hours

Value	Label	Unweighted Frequency	%
1	Yes	1	0.1 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	92	6.8 %
-104	Skip-in for SUBUSEDY 1	85	6.3 %
-103	Skip-in for SUBUSEDY <> 1	1176	86.9 %
	Total	**1,354**	**100%**

Based upon 1 valid cases out of 1,354 total cases.

Location: 1289-1292 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_DRUG24HALL: S2 SubUse_Drug24: Used Hallucinogens past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	2	0.1 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	92	6.8 %
-104	Skip-in for SUBUSEDY 1	84	6.2 %
-103	Skip-in for SUBUSEDY <> 1	1176	86.9 %
	Total	**1,354**	**100%**

Based upon 2 valid cases out of 1,354 total cases.

Location: 1293-1296 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_DRUG24INH: S2 SubUse_Drug24: Used Inhalants past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	1	0.1 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	92	6.8 %
-104	Skip-in for SUBUSEDY 1	85	6.3 %
-103	Skip-in for SUBUSEDY <> 1	1176	86.9 %
	Total	**1,354**	**100%**

Based upon 1 valid cases out of 1,354 total cases.

Location: 1297-1300 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_DRUG24AMYL: S2 SubUse_Drug24: Used Amyl nitrate/poppers/odorizers 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	92	6.8 %
-104	Skip-in for SUBUSEDY 1	86	6.4 %
-103	Skip-in for SUBUSEDY <> 1	1176	86.9 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 1301-1304 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_DRUG24OTHER: S2 SubUse_Drug24: Used Other (specify) past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	1	0.1 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	92	6.8 %
-104	Skip-in for SUBUSEDY 1	85	6.3 %
-103	Skip-in for SUBUSEDY <> 1	1176	86.9 %
	Total	**1,354**	**100%**

Based upon 1 valid cases out of 1,354 total cases.

Location: 1305-1308 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_DRUG24NONE: S2 SubUse_Drug24: Used NA (No drugs) past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	92	6.8 %

Value	Label	Unweighted Frequency	%
-104	Skip-in for SUBUSEDY 1	86	6.4 %
-103	Skip-in for SUBUSEDY <> 1	1176	86.9 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 1309-1312 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S2SUBUSE29: S2 SubUse: Specify other drug used past 24hrs

Based upon 1,354 valid cases out of 1,354 total cases.

Location: 1313-1320 (width: 8; decimal: 0)
Variable Type: character

S2SUBUSE30: S2 SubUse: Num times smoked cigarettes during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	650	48.0 %
2	1-2 times	56	4.1 %
3	3-5 times	45	3.3 %
4	1X per month	19	1.4 %
5	2-3X per month	36	2.7 %
6	1X per week	19	1.4 %
7	2-3X per week	46	3.4 %
8	4-5X per week	44	3.2 %
9	Every day	342	25.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-99	Dont know	3	0.2 %
	Total	**1,354**	**100%**

Based upon 1,257 valid cases out of 1,354 total cases.

Location: 1321-1324 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S2SUBUSE31: S2 SubUse: Num cigarettes smoke per day during recall period

Value	Label	Unweighted Frequency	%
1	0 cigarettes	668	49.3 %
2	1-4 cigarettes	261	19.3 %

Value	Label	Unweighted Frequency	%
3	5-9 cigarettes	127	9.4 %
4	10-15 cigarettes	98	7.2 %
5	16-20 cigarettes	76	5.6 %
6	More than a pack a day cigarettes	28	2.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-9	Dont know	2	0.1 %
	Total	**1,354**	**100%**

Based upon 1,258 valid cases out of 1,354 total cases.

Location: 1325-1328 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -105 , -9 , -8

S2SUBUSE_SUBSYM01_SUSYMB: S2 SubUse: Family ever complain because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	164	12.1 %
5	No	462	34.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-106	Skip-in for alchstat/drugstat	634	46.8 %
	Total	**1,354**	**100%**

Based upon 626 valid cases out of 1,354 total cases.

Location: 1329-1332 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM01_SUSYMC: S2 SubUse: Specify cause of family complaint [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	36	2.7 %
2	Drugs	80	5.9 %
3	Both	48	3.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %
-201	IntCompStat = 4 missed interview	92	6.8 %

Value	Label	Unweighted Frequency	%
-107	Follow-up skipped because gate <> 1	1090	80.5 %
	Total	**1,354**	**100%**

Based upon 164 valid cases out of 1,354 total cases.

Location: 1333-1336 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM02_SUSYMB: S2 SubUse: Friend ever complain because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	73	5.4 %
5	No	553	40.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-106	Skip-in for alchstat/drugstat	634	46.8 %
	Total	**1,354**	**100%**

Based upon 626 valid cases out of 1,354 total cases.

Location: 1337-1340 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM02_SUSYMC: S2 SubUse: Specify cause of friend complain [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	21	1.6 %
2	Drugs	35	2.6 %
3	Both	17	1.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-107	Follow-up skipped because gate <> 1	1181	87.2 %
	Total	**1,354**	**100%**

Based upon 73 valid cases out of 1,354 total cases.

Location: 1341-1344 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM03_SUSYMB: S2 SubUse: Ever trouble at work because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	32	2.4 %
5	No	594	43.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-106	Skip-in for alchstat/drugstat	634	46.8 %
	Total	**1,354**	**100%**

Based upon 626 valid cases out of 1,354 total cases.

Location: 1345-1348 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM03_SUSYMC: S2 SubUse: Specify cause of trouble at work [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	6	0.4 %
2	Drugs	24	1.8 %
3	Both	2	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-107	Follow-up skipped because gate <> 1	1222	90.3 %
	Total	**1,354**	**100%**

Based upon 32 valid cases out of 1,354 total cases.

Location: 1349-1352 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM04_SUSYMB: S2 SubUse: Ever missed work/school because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	60	4.4 %
5	No	566	41.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %

Value	Label	Unweighted Frequency	%
-106	Skip-in for alchstat/drugstat	634	46.8 %
	Total	**1,354**	**100%**

Based upon 626 valid cases out of 1,354 total cases.

Location: 1353-1356 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM04_SUSYMC: S2 SubUse: Specify cause of missed work/school [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	20	1.5 %
2	Drugs	26	1.9 %
3	Both	14	1.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-107	Follow-up skipped because gate <> 1	1194	88.2 %
	Total	**1,354**	**100%**

Based upon 60 valid cases out of 1,354 total cases.

Location: 1357-1360 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM05_SUSYMB: S2 SubUse: Ever arrested because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	47	3.5 %
5	No	579	42.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-106	Skip-in for alchstat/drugstat	634	46.8 %
	Total	**1,354**	**100%**

Based upon 626 valid cases out of 1,354 total cases.

Location: 1361-1364 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM05_SUSYMC: S2 SubUse: Specify cause of arrest [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	14	1.0 %
2	Drugs	28	2.1 %
3	Both	5	0.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	9	0.7 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-107	Follow-up skipped because gate <> 1	1206	89.1 %
	Total	**1,354**	**100%**

Based upon 47 valid cases out of 1,354 total cases.

Location: 1365-1368 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM06_SUSYMB: S2 SubUse: Ever argue family/friends because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	114	8.4 %
5	No	512	37.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-106	Skip-in for alchstat/drugstat	634	46.8 %
	Total	**1,354**	**100%**

Based upon 626 valid cases out of 1,354 total cases.

Location: 1369-1372 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM06_SUSYMC: S2 SubUse: Specify cause of argument w/ family/friends [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	35	2.6 %
2	Drugs	52	3.8 %
3	Both	27	2.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	7	0.5 %

Value	Label	Unweighted Frequency	%
-201	IntCompStat = 4 missed interview	92	6.8 %
-107	Follow-up skipped because gate <> 1	1141	84.3 %
	Total	**1,354**	**100%**

Based upon 114 valid cases out of 1,354 total cases.

Location: 1373-1376 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM07_SUSYMB: S2 SubUse: Ever cont. drug/alcohol use in spite cause health problems in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	88	6.5 %
5	No	538	39.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-106	Skip-in for alchstat/drugstat	634	46.8 %
	Total	**1,354**	**100%**

Based upon 626 valid cases out of 1,354 total cases.

Location: 1377-1380 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM07_SUSYMC: S2 SubUse: Specify continued using [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	23	1.7 %
2	Drugs	40	3.0 %
3	Both	25	1.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-107	Follow-up skipped because gate <> 1	1166	86.1 %
	Total	**1,354**	**100%**

Based upon 88 valid cases out of 1,354 total cases.

Location: 1381-1384 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM08_SUSYMB: S2 SubUse: Ever not remember prior night because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	136	10.0 %
5	No	490	36.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-106	Skip-in for alchstat/drugstat	634	46.8 %
	Total	**1,354**	**100%**

Based upon 626 valid cases out of 1,354 total cases.

Location: 1385-1388 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM08_SUSYMC: S2 SubUse: Specify cause of not remember night [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	67	4.9 %
2	Drugs	23	1.7 %
3	Both	45	3.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	9	0.7 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-107	Follow-up skipped because gate <> 1	1117	82.5 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 135 valid cases out of 1,354 total cases.

Location: 1389-1392 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM09_SUSYMB: S2 SubUse: Ever used in situations could get hurt in recall period [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	150	11.1 %
5	No	476	35.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %

Value	Label	Unweighted Frequency	%
-201	IntCompStat = 4 missed interview	92	6.8 %
-106	Skip-in for alchstat/drugstat	634	46.8 %
	Total	**1,354**	**100%**

Based upon 626 valid cases out of 1,354 total cases.

Location: 1393-1396 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM09_SUSYMC: S2 SubUse: Specify used drugs in hurtful situations [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	51	3.8 %
2	Drugs	50	3.7 %
3	Both	49	3.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-107	Follow-up skipped because gate <> 1	1104	81.5 %
	Total	**1,354**	**100%**

Based upon 150 valid cases out of 1,354 total cases.

Location: 1397-1400 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM10_SUSYMB: S2 SubUse: Ever had accident/injury because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	42	3.1 %
5	No	584	43.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-106	Skip-in for alchstat/drugstat	634	46.8 %
	Total	**1,354**	**100%**

Based upon 626 valid cases out of 1,354 total cases.

Location: 1401-1404 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM10_SUSYMC: S2 SubUse: Specify cause of accident/injury [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	16	1.2 %
2	Drugs	21	1.6 %
3	Both	5	0.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	9	0.7 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-107	Follow-up skipped because gate <> 1	1211	89.4 %
	Total	**1,354**	**100%**

Based upon 42 valid cases out of 1,354 total cases.

Location: 1405-1408 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM11_SUSYMB: S2 SubUse: Alcohol/drug use interfere work/responsibility school/home/job in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	64	4.7 %
5	No	562	41.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-106	Skip-in for alchstat/drugstat	634	46.8 %
	Total	**1,354**	**100%**

Based upon 626 valid cases out of 1,354 total cases.

Location: 1409-1412 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM11_SUSYMC: S2 SubUse: Specify cause of interference [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	11	0.8 %
2	Drugs	37	2.7 %
3	Both	16	1.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %

Value	Label	Unweighted Frequency	%
-201	IntCompStat = 4 missed interview	92	6.8 %
-107	Follow-up skipped because gate <> 1	1190	87.9 %
	Total	**1,354**	**100%**

Based upon 64 valid cases out of 1,354 total cases.

Location: 1413-1416 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM12_SUSYMB: S2 SubUse: Ever trouble police no arrest because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	43	3.2 %
5	No	583	43.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-106	Skip-in for alchstat/drugstat	634	46.8 %
	Total	**1,354**	**100%**

Based upon 626 valid cases out of 1,354 total cases.

Location: 1417-1420 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM12_SUSYMC: S2 SubUse: Specify cause trouble police no arrest [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	16	1.2 %
2	Drugs	22	1.6 %
3	Both	5	0.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-107	Follow-up skipped because gate <> 1	1211	89.4 %
	Total	**1,354**	**100%**

Based upon 43 valid cases out of 1,354 total cases.

Location: 1421-1424 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM13_SUSYMB: S2 SubUse: Sold/exchanged sex for drugs in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	1	0.1 %
5	No	625	46.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-106	Skip-in for alchstat/drugstat	634	46.8 %
	Total	**1,354**	**100%**

Based upon 626 valid cases out of 1,354 total cases.

Location: 1425-1428 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM13_SUSYMC: S2 SubUse: Specify cause sold/exchanged sex [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	0	0.0 %
2	Drugs	1	0.1 %
3	Both	0	0.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	9	0.7 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-107	Follow-up skipped because gate <> 1	1252	92.5 %
	Total	**1,354**	**100%**

Based upon 1 valid cases out of 1,354 total cases.

Location: 1429-1432 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM14_SUSYMB: S2 SubUse: Sold drugs for money buy drugs in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	71	5.2 %
5	No	555	41.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-106	Skip-in for alchstat/drugstat	634	46.8 %

Value	Label	Unweighted Frequency	%
	Total	**1,354**	**100%**

Based upon 626 valid cases out of 1,354 total cases.

Location: 1433-1436 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM14_SUSYMC: S2 SubUse: Specify cause sold drugs for money to buy [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	1	0.1 %
2	Drugs	61	4.5 %
3	Both	9	0.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-107	Follow-up skipped because gate <> 1	1183	87.4 %
	Total	**1,354**	**100%**

Based upon 71 valid cases out of 1,354 total cases.

Location: 1437-1440 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM15_SUSYMB: S2 SubUse: Stole for money buy drugs in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	39	2.9 %
5	No	587	43.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-106	Skip-in for alchstat/drugstat	634	46.8 %
	Total	**1,354**	**100%**

Based upon 626 valid cases out of 1,354 total cases.

Location: 1441-1444 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM15_SUSYMC: S2 SubUse: Specify cause stole for money to buy [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	5	0.4 %
2	Drugs	31	2.3 %
3	Both	3	0.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	9	0.7 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-107	Follow-up skipped because gate <> 1	1214	89.7 %
	Total	**1,354**	**100%**

Based upon 39 valid cases out of 1,354 total cases.

Location: 1445-1448 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM16_SUSYMB: S2 SubUse: Ever had physical fight because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	78	5.8 %
5	No	548	40.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-106	Skip-in for alchstat/drugstat	634	46.8 %
	Total	**1,354**	**100%**

Based upon 626 valid cases out of 1,354 total cases.

Location: 1449-1452 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM16_SUSYMC: S2 SubUse: Specify cause of fight [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	35	2.6 %
2	Drugs	23	1.7 %
3	Both	20	1.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	7	0.5 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-107	Follow-up skipped because gate <> 1	1177	86.9 %
	Total	**1,354**	**100%**

Based upon 78 valid cases out of 1,354 total cases.

Location: 1453-1456 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM17_SUSYMB: S2 SubUse: Ever destroy property because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	39	2.9 %
5	No	587	43.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-106	Skip-in for alchstat/drugstat	634	46.8 %
	Total	**1,354**	**100%**

Based upon 626 valid cases out of 1,354 total cases.

Location: 1457-1460 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM17_SUSYMC: S2 SubUse: Specify cause of destroy property [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	23	1.7 %
2	Drugs	9	0.7 %
3	Both	7	0.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	9	0.7 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-107	Follow-up skipped because gate <> 1	1214	89.7 %
	Total	**1,354**	**100%**

Based upon 39 valid cases out of 1,354 total cases.

Location: 1461-1464 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM18_SUSYMB: S2 SubUse: Ever felt strong urge use drug/alcohol no self stop in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	56	4.1 %

Value	Label	Unweighted Frequency	%
5	No	570	42.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-106	Skip-in for alchstat/drugstat	634	46.8 %
	Total	**1,354**	**100%**

Based upon 626 valid cases out of 1,354 total cases.

Location: 1465-1468 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM18_SUSYMC: S2 SubUse: Specify the cause of strong urge [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	12	0.9 %
2	Drugs	34	2.5 %
3	Both	10	0.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-107	Follow-up skipped because gate <> 1	1198	88.5 %
	Total	**1,354**	**100%**

Based upon 56 valid cases out of 1,354 total cases.

Location: 1469-1472 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM19_SUSYMB: S2 SubUse: Ever want drug/alcohol think of nothing else in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	57	4.2 %
5	No	569	42.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-106	Skip-in for alchstat/drugstat	634	46.8 %
	Total	**1,354**	**100%**

Based upon 626 valid cases out of 1,354 total cases.

Location: 1473-1476 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM19_SUSYMC: S2 SubUse: Specify cause of want [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	12	0.9 %
2	Drugs	36	2.7 %
3	Both	9	0.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-107	Follow-up skipped because gate <> 1	1197	88.4 %
	Total	**1,354**	**100%**

Based upon 57 valid cases out of 1,354 total cases.

Location: 1477-1480 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM20_SUSYMB: S2 SubUse: Ever experience withdrawal from drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	46	3.4 %
5	No	580	42.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-106	Skip-in for alchstat/drugstat	634	46.8 %
	Total	**1,354**	**100%**

Based upon 626 valid cases out of 1,354 total cases.

Location: 1481-1484 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM20_SUSYMC: S2 SubUse: Specify cause of withdraw [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	7	0.5 %
2	Drugs	32	2.4 %

Value	Label	Unweighted Frequency	%
3	Both	7	0.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	7	0.5 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-107	Follow-up skipped because gate <> 1	1209	89.3 %
	Total	**1,354**	**100%**

Based upon 46 valid cases out of 1,354 total cases.

Location: 1485-1488 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM21_SUSYMB: S2 SubUse: Take drug/alcohol to keep from having withdraw in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	26	1.9 %
5	No	20	1.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-108	Skip-in for SUBSYM20=5	577	42.6 %
-106	Skip-in for alchstat/drugstat	635	46.9 %
	Total	**1,354**	**100%**

Based upon 46 valid cases out of 1,354 total cases.

Location: 1489-1492 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM21_SUSYMC: S2 SubUse: Specify reason for consumption [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	5	0.4 %
2	Drugs	19	1.4 %
3	Both	2	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-107	Follow-up skipped because gate <> 1	1228	90.7 %
	Total	**1,354**	**100%**

Based upon 26 valid cases out of 1,354 total cases.

Location: 1493-1496 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM22_SUSYMB: S2 SubUse: Give up usual activities/interests to use drug/alcohol in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	52	3.8 %
5	No	574	42.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-106	Skip-in for alchstat/drugstat	634	46.8 %
	Total	**1,354**	**100%**

Based upon 626 valid cases out of 1,354 total cases.

Location: 1497-1500 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM22_SUSYMC: S2 SubUse: Specify cause of giving up activities/interests [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	9	0.7 %
2	Drugs	32	2.4 %
3	Both	11	0.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-107	Follow-up skipped because gate <> 1	1202	88.8 %
	Total	**1,354**	**100%**

Based upon 52 valid cases out of 1,354 total cases.

Location: 1501-1504 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM23_SUSYMB: S2 SubUse: Ever used in rp when promised wouldn't [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	129	9.5 %
5	No	497	36.7 %

Value	Label	Unweighted Frequency	%
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-106	Skip-in for alchstat/drugstat	634	46.8 %
	Total	**1,354**	**100%**

Based upon 626 valid cases out of 1,354 total cases.

Location: 1505-1508 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM23_SUSYMC: S2 SubUse: Specify what use you started [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	24	1.8 %
2	Drugs	67	4.9 %
3	Both	38	2.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-107	Follow-up skipped because gate <> 1	1125	83.1 %
	Total	**1,354**	**100%**

Based upon 129 valid cases out of 1,354 total cases.

Location: 1509-1512 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM24_SUSYMB: S2 SubUse: Ever use drug/alcohol more freq than intended in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	116	8.6 %
5	No	510	37.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-106	Skip-in for alchstat/drugstat	634	46.8 %
	Total	**1,354**	**100%**

Based upon 626 valid cases out of 1,354 total cases.

Location: 1513-1516 (width: 4; decimal: 0)

Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM24_SUSYMC: S2 SubUse: Specify what you used more frequently [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	29	2.1 %
2	Drugs	59	4.4 %
3	Both	28	2.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	9	0.7 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-107	Follow-up skipped because gate <> 1	1137	84.0 %
	Total	**1,354**	**100%**

Based upon 116 valid cases out of 1,354 total cases.

Location: 1517-1520 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM25_SUSYMB: S2 SubUse: Ever tried cut down drug/alcohol use but couldnt in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	81	6.0 %
5	No	545	40.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-106	Skip-in for alchstat/drugstat	634	46.8 %
	Total	**1,354**	**100%**

Based upon 626 valid cases out of 1,354 total cases.

Location: 1521-1524 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM25_SUSYMC: S2 SubUse: Specify what substance use you tried to curtail [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	11	0.8 %
2	Drugs	55	4.1 %
3	Both	15	1.1 %

Value	Label	Unweighted Frequency	%
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-107	Follow-up skipped because gate <> 1	1173	86.6 %
	Total	**1,354**	**100%**

Based upon 81 valid cases out of 1,354 total cases.

Location: 1525-1528 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM26_SUSYMB: S2 SubUse: Ever need more drug/alcohol get same effect in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	130	9.6 %
5	No	496	36.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-106	Skip-in for alchstat/drugstat	634	46.8 %
	Total	**1,354**	**100%**

Based upon 626 valid cases out of 1,354 total cases.

Location: 1529-1532 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM26_SUSYMC: S2 SubUse: Specify substance need more for same effect [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	29	2.1 %
2	Drugs	71	5.2 %
3	Both	30	2.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	7	0.5 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-107	Follow-up skipped because gate <> 1	1125	83.1 %
	Total	**1,354**	**100%**

Based upon 130 valid cases out of 1,354 total cases.

Location: 1533-1536 (width: 4; decimal: 0)

Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM27_SUSYMB: S2 SubUse: Ever used drug/alcohol enough to depend in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	48	3.5 %
5	No	578	42.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-106	Skip-in for alchstat/drugstat	634	46.8 %
	Total	**1,354**	**100%**

Based upon 626 valid cases out of 1,354 total cases.

Location: 1537-1540 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM27_SUSYMC: S2 SubUse: Specify dependency [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	7	0.5 %
2	Drugs	36	2.7 %
3	Both	5	0.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	7	0.5 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-107	Follow-up skipped because gate <> 1	1207	89.1 %
	Total	**1,354**	**100%**

Based upon 48 valid cases out of 1,354 total cases.

Location: 1541-1544 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM28_SUSYMB: S2 SubUse: Ever talked doctor/clergy/guidance counselor drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	40	3.0 %
5	No	586	43.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %

Value	Label	Unweighted Frequency	%
-201	IntCompStat = 4 missed interview	92	6.8 %
-106	Skip-in for alchstat/drugstat	634	46.8 %
	Total	**1,354**	**100%**

Based upon 626 valid cases out of 1,354 total cases.

Location: 1545-1548 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM28_SUSYMC: S2 SubUse: Specify talked doctor/clergy/guidance counselor [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	9	0.7 %
2	Drugs	21	1.6 %
3	Both	10	0.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	9	0.7 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-107	Follow-up skipped because gate <> 1	1213	89.6 %
	Total	**1,354**	**100%**

Based upon 40 valid cases out of 1,354 total cases.

Location: 1549-1552 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM29_SUSYMB: S2 SubUse: Ever attended self-help group for drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	66	4.9 %
5	No	560	41.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-106	Skip-in for alchstat/drugstat	634	46.8 %
	Total	**1,354**	**100%**

Based upon 626 valid cases out of 1,354 total cases.

Location: 1553-1556 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM29_SUSYMC: S2 SubUse: Specify reason for self-help [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	16	1.2 %
2	Drugs	31	2.3 %
3	Both	19	1.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-107	Follow-up skipped because gate <> 1	1188	87.7 %
	Total	**1,354**	**100%**

Based upon 66 valid cases out of 1,354 total cases.

Location: 1557-1560 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM30_SUSYMB: S2 SubUse: Ever treated by therapist because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	28	2.1 %
5	No	598	44.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-106	Skip-in for alchstat/drugstat	634	46.8 %
	Total	**1,354**	**100%**

Based upon 626 valid cases out of 1,354 total cases.

Location: 1561-1564 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM30_SUSYMC: S2 SubUse: Specify reason for therapist visit [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	4	0.3 %
2	Drugs	17	1.3 %
3	Both	7	0.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	9	0.7 %

Value	Label	Unweighted Frequency	%
-201	IntCompStat = 4 missed interview	92	6.8 %
-107	Follow-up skipped because gate <> 1	1225	90.5 %
	Total	**1,354**	**100%**

Based upon 28 valid cases out of 1,354 total cases.

Location: 1565-1568 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM31_SUSYMB: S2 SubUse: Ever Admitted to hospital because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	11	0.8 %
5	No	615	45.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-106	Skip-in for alchstat/drugstat	634	46.8 %
	Total	**1,354**	**100%**

Based upon 626 valid cases out of 1,354 total cases.

Location: 1569-1572 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE_SUBSYM31_SUSYMC: S2 SubUse: Specify reason admitted to hospital [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	5	0.4 %
2	Drugs	6	0.4 %
3	Both	0	0.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	9	0.7 %
-201	IntCompStat = 4 missed interview	92	6.8 %
-107	Follow-up skipped because gate <> 1	1242	91.7 %
	Total	**1,354**	**100%**

Based upon 11 valid cases out of 1,354 total cases.

Location: 1573-1576 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S2SUBUSE28: S2 SubUse: Num times abused prescribed medications (own or someone elses) during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	0	0.0 %
2	1-2 times	3	0.2 %
3	3-5 times	1	0.1 %
4	1X per month	0	0.0 %
5	2-3X per month	0	0.0 %
6	1X per week	0	0.0 %
7	2-3X per week	0	0.0 %
8	4-5X per week	0	0.0 %
9	Every day	0	0.0 %
	Missing Data		
-800	Question not asked for this subject	1256	92.8 %
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
	Total	**1,354**	**100%**

Based upon 4 valid cases out of 1,354 total cases.

Location: 1577-1580 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S2QFBRMO: S2 SubUse: Quant/freq used beer/wine/liquor - recall period

Based upon 1,255 valid cases out of 1,354 total cases.

- Mean: 9.56
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 420.00
- Standard Deviation: 25.15

Location: 1581-1584 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -300 , -202 , -201 , -100 , -9 , -8

S26MOUSE: S2 SubUse: Number drugs used - recall period

Based upon 1,260 valid cases out of 1,354 total cases.

- Mean: 0.60
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 9.00
- Standard Deviation: 1.14

Location: 1585-1588 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -300 , -202 , -201 , -100 , -9 , -8

S2QFSMK6: S2 SubUse: Quant/freq used cigarettes - recall period

Based upon 1,256 valid cases out of 1,354 total cases.

- Mean: 25.68
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 160.00
- Standard Deviation: 43.32

Location: 1589-1594 (width: 6; decimal: 1)
Variable Type: numeric
(Range of) Missing Values: -300.0 , -202.0 , -201.0 , -100.0 , -9.0 , -8.0

S2A6SC2: S2 SubUse: Recall period alcohol consequences

Based upon 1,260 valid cases out of 1,354 total cases.

- Mean: 0.54
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 10.00
- Standard Deviation: 1.45

Location: 1595-1598 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S2D6SC2: S2 SubUse: Recall period drug consequences

Based upon 1,260 valid cases out of 1,354 total cases.

- Mean: 0.68
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 15.00
- Standard Deviation: 1.91

Location: 1599-1602 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S2A6DP2: S2 SubUse: Recall period alcohol dependence symptoms

Based upon 1,260 valid cases out of 1,354 total cases.

- Mean: 0.24
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00

- Maximum: 10.00
- Standard Deviation: 0.93

Location: 1603-1606 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S2D6DP2: S2 SubUse: Recall period drug dependence symptoms

Based upon 1,260 valid cases out of 1,354 total cases.

- Mean: 0.47
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 10.00
- Standard Deviation: 1.51

Location: 1607-1610 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S2TOTSC6: S2 SubUse: Recall period total substance use consequences

Based upon 1,260 valid cases out of 1,354 total cases.

- Mean: 0.98
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 15.00
- Standard Deviation: 2.22

Location: 1611-1614 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S2TOTDS6: S2 SubUse: Recall period total substance use dependency symp

Based upon 1,260 valid cases out of 1,354 total cases.

- Mean: 0.59
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 10.00
- Standard Deviation: 1.63

Location: 1615-1618 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S2ALTRTM: S2 SubUse: Recall period alcohol treatment

Value	Label	Unweighted Frequency	%
0	No	1216	89.8 %

Value	Label	Unweighted Frequency	%
1	Yes	44	3.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
	Total	**1,354**	**100%**

Based upon 1,260 valid cases out of 1,354 total cases.

Location: 1619-1622 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S2DGTRTM: S2 SubUse: Recall period drug treatment

Value	Label	Unweighted Frequency	%
0	No	1194	88.2 %
1	Yes	66	4.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
	Total	**1,354**	**100%**

Based upon 1,260 valid cases out of 1,354 total cases.

Location: 1623-1626 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S2TRTM: S2 SubUse: Recall period alcohol or drug treatment

Value	Label	Unweighted Frequency	%
0	No	1175	86.8 %
1	Yes	85	6.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	92	6.8 %
	Total	**1,354**	**100%**

Based upon 1,260 valid cases out of 1,354 total cases.

Location: 1627-1630 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S3SUBUSE1: S3 SubUse: How often had alcohol to drink in recall period

Value	Label	Unweighted Frequency	%
1	Not at all	672	49.6 %
2	1-2 times	164	12.1 %
3	3-5 times	88	6.5 %
4	1X per month	37	2.7 %
5	2-3X per month	99	7.3 %
6	1X per week	60	4.4 %
7	2-3X per week	73	5.4 %
8	4-5X per week	24	1.8 %
9	Every day	11	0.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
	Total	**1,354**	**100%**

Based upon 1,228 valid cases out of 1,354 total cases.

Location: 1631-1634 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S3SUBUSE2: S3 SubUse: Num alcoholic drinks usually consumed

Based upon 1,225 valid cases out of 1,354 total cases.

- Mean: 2.70
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 64.00
- Standard Deviation: 5.21

Location: 1635-1638 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -101 , -99 , -98

S3SUBUSE3: S3 SubUse: How often had 5+ drinks at one time in recall period

Value	Label	Unweighted Frequency	%
1	Not at all	886	65.4 %
2	1-2 times	87	6.4 %
3	3-5 times	62	4.6 %
4	1X per month	27	2.0 %
5	2-3X per month	55	4.1 %
6	1X per week	36	2.7 %
7	2-3X per week	54	4.0 %
8	4-5X per week	15	1.1 %

Value	Label	Unweighted Frequency	%
9	Every day	6	0.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
	Total	**1,354**	**100%**

Based upon 1,228 valid cases out of 1,354 total cases.

Location: 1639-1642 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S3SUBUSE4: S3 SubUse: Num times been drunk on alcohol during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	864	63.8 %
2	1-2 times	124	9.2 %
3	3-5 times	84	6.2 %
4	1X per month	25	1.8 %
5	2-3X per month	38	2.8 %
6	1X per week	44	3.2 %
7	2-3X per week	39	2.9 %
8	4-5X per week	9	0.7 %
9	Every day	1	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
	Total	**1,354**	**100%**

Based upon 1,228 valid cases out of 1,354 total cases.

Location: 1643-1646 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S3SUBUSE6: S3 SubUse: Num times used marijuana/hashish during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	809	59.7 %
2	1-2 times	75	5.5 %
3	3-5 times	50	3.7 %
4	1X per month	20	1.5 %
5	2-3X per month	43	3.2 %
6	1X per week	21	1.6 %

Value	Label	Unweighted Frequency	%
7	2-3X per week	46	3.4 %
8	4-5X per week	45	3.3 %
9	Every day	119	8.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
	Total	**1,354**	**100%**

Based upon 1,228 valid cases out of 1,354 total cases.

Location: 1647-1650 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S3SUBUSE8: S3 SubUse: Num times used sedative/tranquilizers during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1165	86.0 %
2	1-2 times	20	1.5 %
3	3-5 times	12	0.9 %
4	1X per month	4	0.3 %
5	2-3X per month	11	0.8 %
6	1X per week	7	0.5 %
7	2-3X per week	2	0.1 %
8	4-5X per week	5	0.4 %
9	Every day	2	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
	Total	**1,354**	**100%**

Based upon 1,228 valid cases out of 1,354 total cases.

Location: 1651-1654 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S3SUBUSE10: S3 SubUse: Num times used stimulants/amphetamines during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1165	86.0 %
2	1-2 times	16	1.2 %
3	3-5 times	4	0.3 %
4	1X per month	4	0.3 %

Value	Label	Unweighted Frequency	%
5	2-3X per month	9	0.7 %
6	1X per week	5	0.4 %
7	2-3X per week	11	0.8 %
8	4-5X per week	10	0.7 %
9	Every day	4	0.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
	Total	**1,354**	**100%**

Based upon 1,228 valid cases out of 1,354 total cases.

Location: 1655-1658 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S3SUBUSE12: S3 SubUse: Num times used cocaine during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1184	87.4 %
2	1-2 times	22	1.6 %
3	3-5 times	6	0.4 %
4	1X per month	7	0.5 %
5	2-3X per month	3	0.2 %
6	1X per week	0	0.0 %
7	2-3X per week	3	0.2 %
8	4-5X per week	2	0.1 %
9	Every day	1	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
	Total	**1,354**	**100%**

Based upon 1,228 valid cases out of 1,354 total cases.

Location: 1659-1662 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S3SUBUSE14: S3 SubUse: Num times used opiates during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1204	88.9 %
2	1-2 times	6	0.4 %

Value	Label	Unweighted Frequency	%
3	3-5 times	5	0.4 %
4	1X per month	3	0.2 %
5	2-3X per month	4	0.3 %
6	1X per week	0	0.0 %
7	2-3X per week	1	0.1 %
8	4-5X per week	3	0.2 %
9	Every day	2	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
	Total	**1,354**	**100%**

Based upon 1,228 valid cases out of 1,354 total cases.

Location: 1663-1666 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S3SUBUSE16: S3 SubUse: Num times used ecstasy during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1204	88.9 %
2	1-2 times	9	0.7 %
3	3-5 times	7	0.5 %
4	1X per month	1	0.1 %
5	2-3X per month	4	0.3 %
6	1X per week	1	0.1 %
7	2-3X per week	0	0.0 %
8	4-5X per week	1	0.1 %
9	Every day	1	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
	Total	**1,354**	**100%**

Based upon 1,228 valid cases out of 1,354 total cases.

Location: 1667-1670 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S3SUBUSE18: S3 SubUse: Num times used hallucinogens during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1174	86.7 %
2	1-2 times	23	1.7 %
3	3-5 times	11	0.8 %
4	1X per month	4	0.3 %
5	2-3X per month	4	0.3 %
6	1X per week	2	0.1 %
7	2-3X per week	3	0.2 %
8	4-5X per week	4	0.3 %
9	Every day	3	0.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
	Total	**1,354**	**100%**

Based upon 1,228 valid cases out of 1,354 total cases.

Location: 1671-1674 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S3SUBUSE20: S3 SubUse: Num times used inhalants during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1212	89.5 %
2	1-2 times	9	0.7 %
3	3-5 times	2	0.1 %
4	1X per month	0	0.0 %
5	2-3X per month	4	0.3 %
6	1X per week	0	0.0 %
7	2-3X per week	0	0.0 %
8	4-5X per week	0	0.0 %
9	Every day	1	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
	Total	**1,354**	**100%**

Based upon 1,228 valid cases out of 1,354 total cases.

Location: 1675-1678 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S3SUBUSE22: S3 SubUse: Num times used amyl nitrate/odorizers/rush during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1222	90.3 %
2	1-2 times	2	0.1 %
3	3-5 times	0	0.0 %
4	1X per month	0	0.0 %
5	2-3X per month	3	0.2 %
6	1X per week	0	0.0 %
7	2-3X per week	0	0.0 %
8	4-5X per week	0	0.0 %
9	Every day	1	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
	Total	**1,354**	**100%**

Based upon 1,228 valid cases out of 1,354 total cases.

Location: 1679-1682 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S3SUBUSE24: S3 SubUse: Used other drugs during recall period [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	2	0.1 %
5	No	1226	90.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
	Total	**1,354**	**100%**

Based upon 1,228 valid cases out of 1,354 total cases.

Location: 1683-1686 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE26: S3 SubUse: Num times used other drugs during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1226	90.5 %
2	1-2 times	1	0.1 %
3	3-5 times	1	0.1 %

Value	Label	Unweighted Frequency	%
4	1X per month	0	0.0 %
5	2-3X per month	0	0.0 %
6	1X per week	0	0.0 %
7	2-3X per week	0	0.0 %
8	4-5X per week	0	0.0 %
9	Every day	0	0.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
	Total	**1,354**	**100%**

Based upon 1,228 valid cases out of 1,354 total cases.

Location: 1687-1690 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S3SUBUSEDY: S3 SubUse: Used any drugs past 24hrs [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	104	7.7 %
5	No	1123	82.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,227 valid cases out of 1,354 total cases.

Location: 1691-1694 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_DRUG24MAR: S3 SubUse_Drug24: Used marijuana/hashish past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	97	7.2 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	125	9.2 %
-104	Skip-in for SUBUSEDY 1	8	0.6 %
-103	Skip-in for SUBUSEDY <> 1	1124	83.0 %
	Total	**1,354**	**100%**

Based upon 97 valid cases out of 1,354 total cases.

Location: 1695-1698 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_DRUG24TRANQ: S3 SubUse_Drug24: Used sedatives/tranquilizers past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	2	0.1 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	125	9.2 %
-104	Skip-in for SUBUSEDY 1	103	7.6 %
-103	Skip-in for SUBUSEDY <> 1	1124	83.0 %
	Total	**1,354**	**100%**

Based upon 2 valid cases out of 1,354 total cases.

Location: 1699-1702 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_DRUG24AMPH: S3 SubUse_Drug24: Used stimulants/amphetamines past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	8	0.6 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	125	9.2 %
-104	Skip-in for SUBUSEDY 1	97	7.2 %
-103	Skip-in for SUBUSEDY <> 1	1124	83.0 %
	Total	**1,354**	**100%**

Based upon 8 valid cases out of 1,354 total cases.

Location: 1703-1706 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_DRUG24COC: S3 SubUse_Drug24: Used Cocaine past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	1	0.1 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	125	9.2 %

Value	Label	Unweighted Frequency	%
-104	Skip-in for SUBUSEDY 1	104	7.7 %
-103	Skip-in for SUBUSEDY <> 1	1124	83.0 %
	Total	**1,354**	**100%**

Based upon 1 valid cases out of 1,354 total cases.

Location: 1707-1710 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_DRUG24OPI: S3 SubUse_Drug24: Used Opiates past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	1	0.1 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	125	9.2 %
-104	Skip-in for SUBUSEDY 1	104	7.7 %
-103	Skip-in for SUBUSEDY <> 1	1124	83.0 %
	Total	**1,354**	**100%**

Based upon 1 valid cases out of 1,354 total cases.

Location: 1711-1714 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_DRUG24ECST: S3 SubUse_Drug24: Used Ecstacy past 24 hours

Value	Label	Unweighted Frequency	%
1	Yes	1	0.1 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	125	9.2 %
-104	Skip-in for SUBUSEDY 1	104	7.7 %
-103	Skip-in for SUBUSEDY <> 1	1124	83.0 %
	Total	**1,354**	**100%**

Based upon 1 valid cases out of 1,354 total cases.

Location: 1715-1718 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_DRUG24HALL: S3 SubUse_Drug24: Used Hallucinogens past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	1	0.1 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	125	9.2 %
-104	Skip-in for SUBUSEDY 1	104	7.7 %
-103	Skip-in for SUBUSEDY <> 1	1124	83.0 %
	Total	**1,354**	**100%**

Based upon 1 valid cases out of 1,354 total cases.

Location: 1719-1722 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_DRUG24INH: S3 SubUse_Drug24: Used Inhalants past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	125	9.2 %
-104	Skip-in for SUBUSEDY 1	105	7.8 %
-103	Skip-in for SUBUSEDY <> 1	1124	83.0 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 1723-1726 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_DRUG24AMYL: S3 SubUse_Drug24: Used Amyl nitrate/poppers/odorizers 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	1	0.1 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	125	9.2 %
-104	Skip-in for SUBUSEDY 1	104	7.7 %
-103	Skip-in for SUBUSEDY <> 1	1124	83.0 %
	Total	**1,354**	**100%**

Based upon 1 valid cases out of 1,354 total cases.

Location: 1727-1730 (width: 4; decimal: 0)
Variable Type: numeric

(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_DRUG24OTHER: S3 SubUse_Drug24: Used Other (specify) past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	1	0.1 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	125	9.2 %
-104	Skip-in for SUBUSEDY 1	104	7.7 %
-103	Skip-in for SUBUSEDY <> 1	1124	83.0 %
	Total	**1,354**	**100%**

Based upon 1 valid cases out of 1,354 total cases.

Location: 1731-1734 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_DRUG24NONE: S3 SubUse_Drug24: Used NA (No drugs) past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	125	9.2 %
-104	Skip-in for SUBUSEDY 1	105	7.8 %
-103	Skip-in for SUBUSEDY <> 1	1124	83.0 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 1735-1738 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S3SUBUSE29: S3 SubUse: Specify other drug used past 24hrs

Based upon 1,354 valid cases out of 1,354 total cases.

Location: 1739-1759 (width: 21; decimal: 0)
Variable Type: character

S3SUBUSE30: S3 SubUse: Num times smoked cigarettes during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	621	45.9 %
2	1-2 times	51	3.8 %

Value	Label	Unweighted Frequency	%
3	3-5 times	36	2.7 %
4	1X per month	17	1.3 %
5	2-3X per month	20	1.5 %
6	1X per week	20	1.5 %
7	2-3X per week	53	3.9 %
8	4-5X per week	28	2.1 %
9	Every day	381	28.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-99	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,227 valid cases out of 1,354 total cases.

Location: 1760-1763 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S3SUBUSE31: S3 SubUse: Num cigarettes smoke per day during recall period

Value	Label	Unweighted Frequency	%
1	0 cigarettes	638	47.1 %
2	1-4 cigarettes	266	19.6 %
3	5-9 cigarettes	109	8.1 %
4	10-15 cigarettes	97	7.2 %
5	16-20 cigarettes	80	5.9 %
6	More than a pack a day cigarettes	38	2.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
	Total	**1,354**	**100%**

Based upon 1,228 valid cases out of 1,354 total cases.

Location: 1764-1767 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -105 , -9 , -8

S3SUBUSE_SUBSYM01_SUSYMB: S3 SubUse: Family ever complain because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	147	10.9 %

Value	Label	Unweighted Frequency	%
5	No	517	38.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-106	Skip-in for alchstat/drugstat	564	41.7 %
	Total	**1,354**	**100%**

Based upon 664 valid cases out of 1,354 total cases.

Location: 1768-1771 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM01_SUSYMC: S3 SubUse: Specify cause of family complaint [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	42	3.1 %
2	Drugs	65	4.8 %
3	Both	40	3.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	13	1.0 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-107	Follow-up skipped because gate <> 1	1069	79.0 %
	Total	**1,354**	**100%**

Based upon 147 valid cases out of 1,354 total cases.

Location: 1772-1775 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM02_SUSYMB: S3 SubUse: Friend ever complain because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	63	4.7 %
5	No	601	44.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-106	Skip-in for alchstat/drugstat	564	41.7 %
	Total	**1,354**	**100%**

Based upon 664 valid cases out of 1,354 total cases.

Location: 1776-1779 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM02_SUSYMC: S3 SubUse: Specify cause of friend complain [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	21	1.6 %
2	Drugs	29	2.1 %
3	Both	13	1.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	15	1.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-107	Follow-up skipped because gate <> 1	1151	85.0 %
	Total	**1,354**	**100%**

Based upon 63 valid cases out of 1,354 total cases.

Location: 1780-1783 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM03_SUSYMB: S3 SubUse: Ever trouble at work because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	22	1.6 %
5	No	641	47.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-106	Skip-in for alchstat/drugstat	564	41.7 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 663 valid cases out of 1,354 total cases.

Location: 1784-1787 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM03_SUSYMC: S3 SubUse: Specify cause of trouble at work [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	3	0.2 %
2	Drugs	17	1.3 %
3	Both	2	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	16	1.2 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-107	Follow-up skipped because gate <> 1	1191	88.0 %
	Total	**1,354**	**100%**

Based upon 22 valid cases out of 1,354 total cases.

Location: 1788-1791 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM04_SUSYMB: S3 SubUse: Ever missed work/school because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	39	2.9 %
5	No	624	46.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-106	Skip-in for alchstat/drugstat	564	41.7 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 663 valid cases out of 1,354 total cases.

Location: 1792-1795 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM04_SUSYMC: S3 SubUse: Specify cause of missed work/school [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	10	0.7 %
2	Drugs	20	1.5 %
3	Both	9	0.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	16	1.2 %
-201	IntCompStat = 4 missed interview	125	9.2 %

Value	Label	Unweighted Frequency	%
-107	Follow-up skipped because gate <> 1	1174	86.7 %
	Total	**1,354**	**100%**

Based upon 39 valid cases out of 1,354 total cases.

Location: 1796-1799 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM05_SUSYMB: S3 SubUse: Ever arrested because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	44	3.2 %
5	No	620	45.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-106	Skip-in for alchstat/drugstat	564	41.7 %
	Total	**1,354**	**100%**

Based upon 664 valid cases out of 1,354 total cases.

Location: 1800-1803 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM05_SUSYMC: S3 SubUse: Specify cause of arrest [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	12	0.9 %
2	Drugs	25	1.8 %
3	Both	7	0.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	13	1.0 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-107	Follow-up skipped because gate <> 1	1172	86.6 %
	Total	**1,354**	**100%**

Based upon 44 valid cases out of 1,354 total cases.

Location: 1804-1807 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM06_SUSYMB: S3 SubUse: Ever argue family/friends because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	94	6.9 %
5	No	570	42.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-106	Skip-in for alchstat/drugstat	564	41.7 %
	Total	**1,354**	**100%**

Based upon 664 valid cases out of 1,354 total cases.

Location: 1808-1811 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM06_SUSYMC: S3 SubUse: Specify cause of argument w/ family/friends [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	32	2.4 %
2	Drugs	46	3.4 %
3	Both	16	1.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	14	1.0 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-107	Follow-up skipped because gate <> 1	1121	82.8 %
	Total	**1,354**	**100%**

Based upon 94 valid cases out of 1,354 total cases.

Location: 1812-1815 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM07_SUSYMB: S3 SubUse: Ever cont. drug/alcohol use in spite cause health problems in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	77	5.7 %
5	No	587	43.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %

Value	Label	Unweighted Frequency	%
-106	Skip-in for alchstat/drugstat	564	41.7 %
	Total	**1,354**	**100%**

Based upon 664 valid cases out of 1,354 total cases.

Location: 1816-1819 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM07_SUSYMC: S3 SubUse: Specify continued using [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	17	1.3 %
2	Drugs	38	2.8 %
3	Both	22	1.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	16	1.2 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-107	Follow-up skipped because gate <> 1	1136	83.9 %
	Total	**1,354**	**100%**

Based upon 77 valid cases out of 1,354 total cases.

Location: 1820-1823 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM08_SUSYMB: S3 SubUse: Ever not remember prior night because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	142	10.5 %
5	No	522	38.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-106	Skip-in for alchstat/drugstat	564	41.7 %
	Total	**1,354**	**100%**

Based upon 664 valid cases out of 1,354 total cases.

Location: 1824-1827 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM08_SUSYMC: S3 SubUse: Specify cause of not remember night [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	77	5.7 %
2	Drugs	24	1.8 %
3	Both	41	3.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	13	1.0 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-107	Follow-up skipped because gate <> 1	1074	79.3 %
	Total	**1,354**	**100%**

Based upon 142 valid cases out of 1,354 total cases.

Location: 1828-1831 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM09_SUSYMB: S3 SubUse: Ever used in situations could get hurt in recall period [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	147	10.9 %
5	No	517	38.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-106	Skip-in for alchstat/drugstat	564	41.7 %
	Total	**1,354**	**100%**

Based upon 664 valid cases out of 1,354 total cases.

Location: 1832-1835 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM09_SUSYMC: S3 SubUse: Specify used drugs in hurtful situations [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	62	4.6 %
2	Drugs	41	3.0 %
3	Both	44	3.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	12	0.9 %

Value	Label	Unweighted Frequency	%
-201	IntCompStat = 4 missed interview	125	9.2 %
-107	Follow-up skipped because gate <> 1	1070	79.0 %
	Total	**1,354**	**100%**

Based upon 147 valid cases out of 1,354 total cases.

Location: 1836-1839 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM10_SUSYMB: S3 SubUse: Ever had accident/injury because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	43	3.2 %
5	No	621	45.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-106	Skip-in for alchstat/drugstat	564	41.7 %
	Total	**1,354**	**100%**

Based upon 664 valid cases out of 1,354 total cases.

Location: 1840-1843 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM10_SUSYMC: S3 SubUse: Specify cause of accident/injury [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	20	1.5 %
2	Drugs	12	0.9 %
3	Both	11	0.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	16	1.2 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-107	Follow-up skipped because gate <> 1	1170	86.4 %
	Total	**1,354**	**100%**

Based upon 43 valid cases out of 1,354 total cases.

Location: 1844-1847 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM11_SUSYMB: S3 SubUse: Alcohol/drug use interfere work/responsibility school/home/job in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	61	4.5 %
5	No	603	44.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-106	Skip-in for alchstat/drugstat	564	41.7 %
	Total	**1,354**	**100%**

Based upon 664 valid cases out of 1,354 total cases.

Location: 1848-1851 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM11_SUSYMC: S3 SubUse: Specify cause of interference [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	17	1.3 %
2	Drugs	36	2.7 %
3	Both	8	0.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	14	1.0 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-107	Follow-up skipped because gate <> 1	1154	85.2 %
	Total	**1,354**	**100%**

Based upon 61 valid cases out of 1,354 total cases.

Location: 1852-1855 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM12_SUSYMB: S3 SubUse: Ever trouble police no arrest because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	55	4.1 %
5	No	609	45.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %

Value	Label	Unweighted Frequency	%
-106	Skip-in for alchstat/drugstat	564	41.7 %
	Total	**1,354**	**100%**

Based upon 664 valid cases out of 1,354 total cases.

Location: 1856-1859 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM12_SUSYMC: S3 SubUse: Specify cause trouble police no arrest [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	26	1.9 %
2	Drugs	22	1.6 %
3	Both	7	0.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	13	1.0 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-107	Follow-up skipped because gate <> 1	1161	85.7 %
	Total	**1,354**	**100%**

Based upon 55 valid cases out of 1,354 total cases.

Location: 1860-1863 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM13_SUSYMB: S3 SubUse: Sold/exchanged sex for drugs in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	1	0.1 %
5	No	663	49.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-106	Skip-in for alchstat/drugstat	564	41.7 %
	Total	**1,354**	**100%**

Based upon 664 valid cases out of 1,354 total cases.

Location: 1864-1867 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM13_SUSYMC: S3 SubUse: Specify cause sold/exchanged sex [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	0	0.0 %
2	Drugs	1	0.1 %
3	Both	0	0.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	16	1.2 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-107	Follow-up skipped because gate <> 1	1212	89.5 %
	Total	**1,354**	**100%**

Based upon 1 valid cases out of 1,354 total cases.

Location: 1868-1871 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM14_SUSYMB: S3 SubUse: Sold drugs for money buy drugs in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	64	4.7 %
5	No	600	44.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-106	Skip-in for alchstat/drugstat	564	41.7 %
	Total	**1,354**	**100%**

Based upon 664 valid cases out of 1,354 total cases.

Location: 1872-1875 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM14_SUSYMC: S3 SubUse: Specify cause sold drugs for money to buy [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	1	0.1 %
2	Drugs	59	4.4 %
3	Both	4	0.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	13	1.0 %

Value	Label	Unweighted Frequency	%
-201	IntCompStat = 4 missed interview	125	9.2 %
-107	Follow-up skipped because gate <> 1	1152	85.1 %
	Total	**1,354**	**100%**

Based upon 64 valid cases out of 1,354 total cases.

Location: 1876-1879 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM15_SUSYMB: S3 SubUse: Stole for money buy drugs in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	36	2.7 %
5	No	628	46.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-106	Skip-in for alchstat/drugstat	564	41.7 %
	Total	**1,354**	**100%**

Based upon 664 valid cases out of 1,354 total cases.

Location: 1880-1883 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM15_SUSYMC: S3 SubUse: Specify cause stole for money to buy [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	1	0.1 %
2	Drugs	31	2.3 %
3	Both	4	0.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	16	1.2 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-107	Follow-up skipped because gate <> 1	1177	86.9 %
	Total	**1,354**	**100%**

Based upon 36 valid cases out of 1,354 total cases.

Location: 1884-1887 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM16_SUSYMB: S3 SubUse: Ever had physical fight because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	65	4.8 %
5	No	599	44.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-106	Skip-in for alchstat/drugstat	564	41.7 %
	Total	**1,354**	**100%**

Based upon 664 valid cases out of 1,354 total cases.

Location: 1888-1891 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM16_SUSYMC: S3 SubUse: Specify cause of fight [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	30	2.2 %
2	Drugs	20	1.5 %
3	Both	15	1.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	15	1.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-107	Follow-up skipped because gate <> 1	1149	84.9 %
	Total	**1,354**	**100%**

Based upon 65 valid cases out of 1,354 total cases.

Location: 1892-1895 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM17_SUSYMB: S3 SubUse: Ever destroy property because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	32	2.4 %
5	No	632	46.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %

Value	Label	Unweighted Frequency	%
-106	Skip-in for alchstat/drugstat	564	41.7 %
	Total	**1,354**	**100%**

Based upon 664 valid cases out of 1,354 total cases.

Location: 1896-1899 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM17_SUSYMC: S3 SubUse: Specify cause of destroy property [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	14	1.0 %
2	Drugs	9	0.7 %
3	Both	9	0.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	16	1.2 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-107	Follow-up skipped because gate <> 1	1181	87.2 %
	Total	**1,354**	**100%**

Based upon 32 valid cases out of 1,354 total cases.

Location: 1900-1903 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM18_SUSYMB: S3 SubUse: Ever felt strong urge use drug/alcohol no self stop in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	59	4.4 %
5	No	605	44.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-106	Skip-in for alchstat/drugstat	564	41.7 %
	Total	**1,354**	**100%**

Based upon 664 valid cases out of 1,354 total cases.

Location: 1904-1907 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM18_SUSYMC: S3 SubUse: Specify the cause of strong urge [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	11	0.8 %
2	Drugs	35	2.6 %
3	Both	13	1.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	15	1.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-107	Follow-up skipped because gate <> 1	1155	85.3 %
	Total	**1,354**	**100%**

Based upon 59 valid cases out of 1,354 total cases.

Location: 1908-1911 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM19_SUSYMB: S3 SubUse: Ever want drug/alcohol think of nothing else in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	53	3.9 %
5	No	611	45.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-106	Skip-in for alchstat/drugstat	564	41.7 %
	Total	**1,354**	**100%**

Based upon 664 valid cases out of 1,354 total cases.

Location: 1912-1915 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM19_SUSYMC: S3 SubUse: Specify cause of want [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	9	0.7 %
2	Drugs	37	2.7 %
3	Both	7	0.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	15	1.1 %

Value	Label	Unweighted Frequency	%
-201	IntCompStat = 4 missed interview	125	9.2 %
-107	Follow-up skipped because gate <> 1	1161	85.7 %
	Total	**1,354**	**100%**

Based upon 53 valid cases out of 1,354 total cases.

Location: 1916-1919 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM20_SUSYMB: S3 SubUse: Ever experience withdrawal from drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	43	3.2 %
5	No	621	45.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-106	Skip-in for alchstat/drugstat	564	41.7 %
	Total	**1,354**	**100%**

Based upon 664 valid cases out of 1,354 total cases.

Location: 1920-1923 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM20_SUSYMC: S3 SubUse: Specify cause of withdraw [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	4	0.3 %
2	Drugs	35	2.6 %
3	Both	4	0.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	14	1.0 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-107	Follow-up skipped because gate <> 1	1172	86.6 %
	Total	**1,354**	**100%**

Based upon 43 valid cases out of 1,354 total cases.

Location: 1924-1927 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM21_SUSYMB: S3 SubUse: Take drug/alcohol to keep from having withdraw in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	24	1.8 %
5	No	19	1.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	9	0.7 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-108	Skip-in for SUBSYM20=5	613	45.3 %
-106	Skip-in for alchstat/drugstat	564	41.7 %
	Total	**1,354**	**100%**

Based upon 43 valid cases out of 1,354 total cases.

Location: 1928-1931 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM21_SUSYMC: S3 SubUse: Specify reason for consumption [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	1	0.1 %
2	Drugs	20	1.5 %
3	Both	3	0.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	14	1.0 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-107	Follow-up skipped because gate <> 1	1191	88.0 %
	Total	**1,354**	**100%**

Based upon 24 valid cases out of 1,354 total cases.

Location: 1932-1935 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM22_SUSYMB: S3 SubUse: Give up usual activities/interests to use drug/alcohol in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	46	3.4 %
5	No	618	45.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %

Value	Label	Unweighted Frequency	%
-201	IntCompStat = 4 missed interview	125	9.2 %
-106	Skip-in for alchstat/drugstat	564	41.7 %
	Total	**1,354**	**100%**

Based upon 664 valid cases out of 1,354 total cases.

Location: 1936-1939 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM22_SUSYMC: S3 SubUse: Specify cause of giving up activities/interests [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	5	0.4 %
2	Drugs	29	2.1 %
3	Both	12	0.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	14	1.0 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-107	Follow-up skipped because gate <> 1	1169	86.3 %
	Total	**1,354**	**100%**

Based upon 46 valid cases out of 1,354 total cases.

Location: 1940-1943 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM23_SUSYMB: S3 SubUse: Ever used in rp when promised wouldn't [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	122	9.0 %
5	No	542	40.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-106	Skip-in for alchstat/drugstat	564	41.7 %
	Total	**1,354**	**100%**

Based upon 664 valid cases out of 1,354 total cases.

Location: 1944-1947 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM23_SUSYMC: S3 SubUse: Specify what use you started [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	28	2.1 %
2	Drugs	68	5.0 %
3	Both	26	1.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	11	0.8 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-107	Follow-up skipped because gate <> 1	1096	80.9 %
	Total	**1,354**	**100%**

Based upon 122 valid cases out of 1,354 total cases.

Location: 1948-1951 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM24_SUSYMB: S3 SubUse: Ever use drug/alcohol more freq than intended in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	109	8.1 %
5	No	555	41.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-106	Skip-in for alchstat/drugstat	564	41.7 %
	Total	**1,354**	**100%**

Based upon 664 valid cases out of 1,354 total cases.

Location: 1952-1955 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM24_SUSYMC: S3 SubUse: Specify what you used more frequently [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	23	1.7 %
2	Drugs	64	4.7 %
3	Both	22	1.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	15	1.1 %

Value	Label	Unweighted Frequency	%
-201	IntCompStat = 4 missed interview	125	9.2 %
-107	Follow-up skipped because gate <> 1	1105	81.6 %
	Total	**1,354**	**100%**

Based upon 109 valid cases out of 1,354 total cases.

Location: 1956-1959 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM25_SUSYMB: S3 SubUse: Ever tried cut down drug/alcohol use but couldnt in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	79	5.8 %
5	No	585	43.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-106	Skip-in for alchstat/drugstat	564	41.7 %
	Total	**1,354**	**100%**

Based upon 664 valid cases out of 1,354 total cases.

Location: 1960-1963 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM25_SUSYMC: S3 SubUse: Specify what substance use you tried to curtail [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	15	1.1 %
2	Drugs	50	3.7 %
3	Both	14	1.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	12	0.9 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-107	Follow-up skipped because gate <> 1	1138	84.0 %
	Total	**1,354**	**100%**

Based upon 79 valid cases out of 1,354 total cases.

Location: 1964-1967 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM26_SUSYMB: S3 SubUse: Ever need more drug/alcohol get same effect in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	122	9.0 %
5	No	542	40.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-106	Skip-in for alchstat/drugstat	564	41.7 %
	Total	**1,354**	**100%**

Based upon 664 valid cases out of 1,354 total cases.

Location: 1968-1971 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM26_SUSYMC: S3 SubUse: Specify substance need more for same effect [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	34	2.5 %
2	Drugs	71	5.2 %
3	Both	17	1.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	13	1.0 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-107	Follow-up skipped because gate <> 1	1094	80.8 %
	Total	**1,354**	**100%**

Based upon 122 valid cases out of 1,354 total cases.

Location: 1972-1975 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM27_SUSYMB: S3 SubUse: Ever used drug/alcohol enough to depend in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	52	3.8 %
5	No	612	45.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %

Value	Label	Unweighted Frequency	%
-106	Skip-in for alchstat/drugstat	564	41.7 %
	Total	**1,354**	**100%**

Based upon 664 valid cases out of 1,354 total cases.

Location: 1976-1979 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM27_SUSYMC: S3 SubUse: Specify dependency [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	8	0.6 %
2	Drugs	37	2.7 %
3	Both	7	0.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	15	1.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-107	Follow-up skipped because gate <> 1	1162	85.8 %
	Total	**1,354**	**100%**

Based upon 52 valid cases out of 1,354 total cases.

Location: 1980-1983 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM28_SUSYMB: S3 SubUse: Ever talked doctor/clergy/guidance counselor drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	40	3.0 %
5	No	624	46.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-106	Skip-in for alchstat/drugstat	564	41.7 %
	Total	**1,354**	**100%**

Based upon 664 valid cases out of 1,354 total cases.

Location: 1984-1987 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM28_SUSYMC: S3 SubUse: Specify talked doctor/clergy/guidance counselor [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	8	0.6 %
2	Drugs	14	1.0 %
3	Both	18	1.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	14	1.0 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-107	Follow-up skipped because gate <> 1	1175	86.8 %
	Total	**1,354**	**100%**

Based upon 40 valid cases out of 1,354 total cases.

Location: 1988-1991 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM29_SUSYMB: S3 SubUse: Ever attended self-help group for drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	55	4.1 %
5	No	609	45.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-106	Skip-in for alchstat/drugstat	564	41.7 %
	Total	**1,354**	**100%**

Based upon 664 valid cases out of 1,354 total cases.

Location: 1992-1995 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM29_SUSYMC: S3 SubUse: Specify reason for self-help [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	11	0.8 %
2	Drugs	24	1.8 %
3	Both	20	1.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	13	1.0 %

Value	Label	Unweighted Frequency	%
-201	IntCompStat = 4 missed interview	125	9.2 %
-107	Follow-up skipped because gate <> 1	1161	85.7 %
	Total	**1,354**	**100%**

Based upon 55 valid cases out of 1,354 total cases.

Location: 1996-1999 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM30_SUSYMB: S3 SubUse: Ever treated by therapist because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	25	1.8 %
5	No	639	47.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-106	Skip-in for alchstat/drugstat	564	41.7 %
	Total	**1,354**	**100%**

Based upon 664 valid cases out of 1,354 total cases.

Location: 2000-2003 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM30_SUSYMC: S3 SubUse: Specify reason for therapist visit [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	2	0.1 %
2	Drugs	16	1.2 %
3	Both	7	0.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	16	1.2 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-107	Follow-up skipped because gate <> 1	1188	87.7 %
	Total	**1,354**	**100%**

Based upon 25 valid cases out of 1,354 total cases.

Location: 2004-2007 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM31_SUSYMB: S3 SubUse: Ever Admitted to hospital because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	6	0.4 %
5	No	658	48.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-106	Skip-in for alchstat/drugstat	564	41.7 %
	Total	**1,354**	**100%**

Based upon 664 valid cases out of 1,354 total cases.

Location: 2008-2011 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE_SUBSYM31_SUSYMC: S3 SubUse: Specify reason admitted to hospital [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	1	0.1 %
2	Drugs	4	0.3 %
3	Both	1	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	16	1.2 %
-201	IntCompStat = 4 missed interview	125	9.2 %
-107	Follow-up skipped because gate <> 1	1207	89.1 %
	Total	**1,354**	**100%**

Based upon 6 valid cases out of 1,354 total cases.

Location: 2012-2015 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S3SUBUSE28: S3 SubUse: Num times abused prescribed medications (own or someone elses) during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	0	0.0 %
2	1-2 times	0	0.0 %
3	3-5 times	0	0.0 %
4	1X per month	0	0.0 %
5	2-3X per month	1	0.1 %

Value	Label	Unweighted Frequency	%
6	1X per week	0	0.0 %
7	2-3X per week	0	0.0 %
8	4-5X per week	0	0.0 %
9	Every day	1	0.1 %
	Missing Data		
-800	Question not asked for this subject	1226	90.5 %
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
	Total	**1,354**	**100%**

Based upon 2 valid cases out of 1,354 total cases.

Location: 2016-2019 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S3QFBRMO: S3 SubUse: Quant/freq used beer/wine/liquor - recall period

Based upon 1,225 valid cases out of 1,354 total cases.

- Mean: 10.65
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 448.00
- Standard Deviation: 28.66

Location: 2020-2023 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -300 , -202 , -201 , -100 , -9 , -8

S36MOUSE: S3 SubUse: Number drugs used - recall period

Based upon 1,228 valid cases out of 1,354 total cases.

- Mean: 0.58
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 9.00
- Standard Deviation: 1.07

Location: 2024-2027 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -300 , -202 , -201 , -100 , -9 , -8

S3QFSMK6: S3 SubUse: Quant/freq used cigarettes - recall period

Based upon 1,227 valid cases out of 1,354 total cases.

- Mean: 28.48

- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 160.00
- Standard Deviation: 46.40

Location: 2028-2033 (width: 6; decimal: 1)
Variable Type: numeric
(Range of) Missing Values: -300.0 , -202.0 , -201.0 , -100.0 , -9.0 , -8.0

S3A6SC2: S3 SubUse: Recall period alcohol consequences

Based upon 1,228 valid cases out of 1,354 total cases.

- Mean: 0.52
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 11.00
- Standard Deviation: 1.49

Location: 2034-2037 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S3D6SC2: S3 SubUse: Recall period drug consequences

Based upon 1,228 valid cases out of 1,354 total cases.

- Mean: 0.61
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 14.00
- Standard Deviation: 1.83

Location: 2038-2041 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S3A6DP2: S3 SubUse: Recall period alcohol dependence symptoms

Based upon 1,228 valid cases out of 1,354 total cases.

- Mean: 0.21
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 10.00
- Standard Deviation: 0.92

Location: 2042-2045 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S3D6DP2: S3 SubUse: Recall period drug dependence symptoms

Based upon 1,228 valid cases out of 1,354 total cases.

- Mean: 0.46
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 10.00
- Standard Deviation: 1.56

Location: 2046-2049 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S3TOTSC6: S3 SubUse: Recall period total substance use consequences

Based upon 1,228 valid cases out of 1,354 total cases.

- Mean: 0.92
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 14.00
- Standard Deviation: 2.16

Location: 2050-2053 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S3TOTDS6: S3 SubUse: Recall period total substance use dependency symp

Based upon 1,228 valid cases out of 1,354 total cases.

- Mean: 0.58
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 10.00
- Standard Deviation: 1.66

Location: 2054-2057 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S3ALTRTM: S3 SubUse: Recall period alcohol treatment

Value	Label	Unweighted Frequency	%
0	No	1193	88.1 %
1	Yes	35	2.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
	Total	**1,354**	**100%**

Based upon 1,228 valid cases out of 1,354 total cases.

Location: 2058-2061 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S3DGTRTM: S3 SubUse: Recall period drug treatment

Value	Label	Unweighted Frequency	%
0	No	1176	86.9 %
1	Yes	52	3.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
	Total	**1,354**	**100%**

Based upon 1,228 valid cases out of 1,354 total cases.

Location: 2062-2065 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S3TRTM: S3 SubUse: Recall period alcohol or drug treatment

Value	Label	Unweighted Frequency	%
0	No	1164	86.0 %
1	Yes	64	4.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	125	9.2 %
	Total	**1,354**	**100%**

Based upon 1,228 valid cases out of 1,354 total cases.

Location: 2066-2069 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S4SUBUSE1: S4 SubUse: How often had alcohol to drink in recall period

Value	Label	Unweighted Frequency	%
1	Not at all	650	48.0 %
2	1-2 times	146	10.8 %
3	3-5 times	112	8.3 %
4	1X per month	50	3.7 %
5	2-3X per month	90	6.6 %
6	1X per week	58	4.3 %

Value	Label	Unweighted Frequency	%
7	2-3X per week	73	5.4 %
8	4-5X per week	26	1.9 %
9	Every day	24	1.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
	Total	**1,354**	**100%**

Based upon 1,229 valid cases out of 1,354 total cases.

Location: 2070-2073 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S4SUBUSE2: S4 SubUse: Num alcoholic drinks usually consumed

Based upon 1,226 valid cases out of 1,354 total cases.

- Mean: 2.64
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 66.00
- Standard Deviation: 4.98

Location: 2074-2077 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -101 , -99 , -98

S4SUBUSE3: S4 SubUse: How often had 5+ drinks at one time in recall period

Value	Label	Unweighted Frequency	%
1	Not at all	864	63.8 %
2	1-2 times	99	7.3 %
3	3-5 times	60	4.4 %
4	1X per month	38	2.8 %
5	2-3X per month	49	3.6 %
6	1X per week	40	3.0 %
7	2-3X per week	45	3.3 %
8	4-5X per week	18	1.3 %
9	Every day	14	1.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-99	Dont know	2	0.1 %
	Total	**1,354**	**100%**

Based upon 1,227 valid cases out of 1,354 total cases.

Location: 2078-2081 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S4SUBUSE4: S4 SubUse: Num times been drunk on alcohol during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	824	60.9 %
2	1-2 times	158	11.7 %
3	3-5 times	85	6.3 %
4	1X per month	35	2.6 %
5	2-3X per month	37	2.7 %
6	1X per week	35	2.6 %
7	2-3X per week	30	2.2 %
8	4-5X per week	18	1.3 %
9	Every day	7	0.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
	Total	**1,354**	**100%**

Based upon 1,229 valid cases out of 1,354 total cases.

Location: 2082-2085 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S4SUBUSE6: S4 SubUse: Num times used marijuana/hashish during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	807	59.6 %
2	1-2 times	70	5.2 %
3	3-5 times	36	2.7 %
4	1X per month	22	1.6 %
5	2-3X per month	38	2.8 %
6	1X per week	27	2.0 %
7	2-3X per week	49	3.6 %
8	4-5X per week	54	4.0 %
9	Every day	125	9.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-99	Dont know	1	0.1 %

Value	Label	Unweighted Frequency	%
	Total	**1,354**	**100%**

Based upon 1,228 valid cases out of 1,354 total cases.

Location: 2086-2089 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S4SUBUSE8: S4 SubUse: Num times used sedative/tranquilizers during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1173	86.6 %
2	1-2 times	25	1.8 %
3	3-5 times	6	0.4 %
4	1X per month	6	0.4 %
5	2-3X per month	7	0.5 %
6	1X per week	2	0.1 %
7	2-3X per week	4	0.3 %
8	4-5X per week	3	0.2 %
9	Every day	3	0.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
	Total	**1,354**	**100%**

Based upon 1,229 valid cases out of 1,354 total cases.

Location: 2090-2093 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S4SUBUSE10: S4 SubUse: Num times used stimulants/amphetamines during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1155	85.3 %
2	1-2 times	10	0.7 %
3	3-5 times	8	0.6 %
4	1X per month	6	0.4 %
5	2-3X per month	12	0.9 %
6	1X per week	5	0.4 %
7	2-3X per week	11	0.8 %
8	4-5X per week	5	0.4 %
9	Every day	17	1.3 %
	Missing Data		

Value	Label	Unweighted Frequency	%
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
	Total	**1,354**	**100%**

Based upon 1,229 valid cases out of 1,354 total cases.

Location: 2094-2097 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S4SUBUSE12: S4 SubUse: Num times used cocaine during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1179	87.1 %
2	1-2 times	20	1.5 %
3	3-5 times	10	0.7 %
4	1X per month	7	0.5 %
5	2-3X per month	7	0.5 %
6	1X per week	3	0.2 %
7	2-3X per week	2	0.1 %
8	4-5X per week	1	0.1 %
9	Every day	0	0.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
	Total	**1,354**	**100%**

Based upon 1,229 valid cases out of 1,354 total cases.

Location: 2098-2101 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S4SUBUSE14: S4 SubUse: Num times used opiates during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1207	89.1 %
2	1-2 times	7	0.5 %
3	3-5 times	1	0.1 %
4	1X per month	2	0.1 %
5	2-3X per month	7	0.5 %
6	1X per week	0	0.0 %
7	2-3X per week	2	0.1 %
8	4-5X per week	2	0.1 %

Value	Label	Unweighted Frequency	%
9	Every day	1	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
	Total	**1,354**	**100%**

Based upon 1,229 valid cases out of 1,354 total cases.

Location: 2102-2105 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S4SUBUSE16: S4 SubUse: Num times used ecstasy during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1211	89.4 %
2	1-2 times	5	0.4 %
3	3-5 times	3	0.2 %
4	1X per month	0	0.0 %
5	2-3X per month	5	0.4 %
6	1X per week	4	0.3 %
7	2-3X per week	1	0.1 %
8	4-5X per week	0	0.0 %
9	Every day	0	0.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
	Total	**1,354**	**100%**

Based upon 1,229 valid cases out of 1,354 total cases.

Location: 2106-2109 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S4SUBUSE18: S4 SubUse: Num times used hallucinogens during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1179	87.1 %
2	1-2 times	22	1.6 %
3	3-5 times	9	0.7 %
4	1X per month	2	0.1 %
5	2-3X per month	7	0.5 %
6	1X per week	4	0.3 %

Value	Label	Unweighted Frequency	%
7	2-3X per week	3	0.2 %
8	4-5X per week	1	0.1 %
9	Every day	2	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
	Total	**1,354**	**100%**

Based upon 1,229 valid cases out of 1,354 total cases.

Location: 2110-2113 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S4SUBUSE20: S4 SubUse: Num times used inhalants during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1210	89.4 %
2	1-2 times	7	0.5 %
3	3-5 times	3	0.2 %
4	1X per month	0	0.0 %
5	2-3X per month	7	0.5 %
6	1X per week	0	0.0 %
7	2-3X per week	0	0.0 %
8	4-5X per week	1	0.1 %
9	Every day	1	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
	Total	**1,354**	**100%**

Based upon 1,229 valid cases out of 1,354 total cases.

Location: 2114-2117 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S4SUBUSE22: S4 SubUse: Num times used amyl nitrate/odorizers/rush during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1217	89.9 %
2	1-2 times	5	0.4 %
3	3-5 times	0	0.0 %
4	1X per month	0	0.0 %

Value	Label	Unweighted Frequency	%
5	2-3X per month	7	0.5 %
6	1X per week	0	0.0 %
7	2-3X per week	0	0.0 %
8	4-5X per week	0	0.0 %
9	Every day	0	0.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
	Total	**1,354**	**100%**

Based upon 1,229 valid cases out of 1,354 total cases.

Location: 2118-2121 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S4SUBUSE24: S4 SubUse: Used other drugs during recall period [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	1229	90.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
	Total	**1,354**	**100%**

Based upon 1,229 valid cases out of 1,354 total cases.

Location: 2122-2125 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE26: S4 SubUse: Num times used other drugs during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1229	90.8 %
2	1-2 times	0	0.0 %
3	3-5 times	0	0.0 %
4	1X per month	0	0.0 %
5	2-3X per month	0	0.0 %
6	1X per week	0	0.0 %
7	2-3X per week	0	0.0 %
8	4-5X per week	0	0.0 %
9	Every day	0	0.0 %

Value	Label	Unweighted Frequency	%
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
	Total	**1,354**	**100%**

Based upon 1,229 valid cases out of 1,354 total cases.

Location: 2126-2129 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S4SUBUSEDY: S4 SubUse: Used any drugs past 24hrs [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	112	8.3 %
5	No	1117	82.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
	Total	**1,354**	**100%**

Based upon 1,229 valid cases out of 1,354 total cases.

Location: 2130-2133 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_DRUG24MAR: S4 SubUse_Drug24: Used marijuana/hashish past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	108	8.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	123	9.1 %
-104	Skip-in for SUBUSEDY 1	6	0.4 %
-103	Skip-in for SUBUSEDY <> 1	1117	82.5 %
	Total	**1,354**	**100%**

Based upon 108 valid cases out of 1,354 total cases.

Location: 2134-2137 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_DRUG24TRANQ: S4 SubUse_Drug24: Used sedatives/tranquilizers past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	1	0.1 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	123	9.1 %
-104	Skip-in for SUBUSEDY 1	113	8.3 %
-103	Skip-in for SUBUSEDY <> 1	1117	82.5 %
	Total	**1,354**	**100%**

Based upon 1 valid cases out of 1,354 total cases.

Location: 2138-2141 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_DRUG24AMPH: S4 SubUse_Drug24: Used stimulants/amphetamines past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	6	0.4 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	123	9.1 %
-104	Skip-in for SUBUSEDY 1	108	8.0 %
-103	Skip-in for SUBUSEDY <> 1	1117	82.5 %
	Total	**1,354**	**100%**

Based upon 6 valid cases out of 1,354 total cases.

Location: 2142-2145 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_DRUG24COC: S4 SubUse_Drug24: Used Cocaine past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	1	0.1 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	123	9.1 %
-104	Skip-in for SUBUSEDY 1	113	8.3 %
-103	Skip-in for SUBUSEDY <> 1	1117	82.5 %
	Total	**1,354**	**100%**

Based upon 1 valid cases out of 1,354 total cases.

Location: 2146-2149 (width: 4; decimal: 0)
Variable Type: numeric

(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_DRUG24OPI: S4 SubUse_Drug24: Used Opiates past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	123	9.1 %
-104	Skip-in for SUBUSEDY 1	114	8.4 %
-103	Skip-in for SUBUSEDY <> 1	1117	82.5 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 2150-2153 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_DRUG24ECST: S4 SubUse_Drug24: Used Ecstacy past 24 hours

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	123	9.1 %
-104	Skip-in for SUBUSEDY 1	114	8.4 %
-103	Skip-in for SUBUSEDY <> 1	1117	82.5 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 2154-2157 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_DRUG24HALL: S4 SubUse_Drug24: Used Hallucinogens past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	123	9.1 %
-104	Skip-in for SUBUSEDY 1	114	8.4 %
-103	Skip-in for SUBUSEDY <> 1	1117	82.5 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 2158-2161 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_DRUG24INH: S4 SubUse_Drug24: Used Inhalants past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	123	9.1 %
-104	Skip-in for SUBUSEDY 1	114	8.4 %
-103	Skip-in for SUBUSEDY <> 1	1117	82.5 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 2162-2165 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_DRUG24AMYL: S4 SubUse_Drug24: Used Amyl nitrate/poppers/odorizers 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	123	9.1 %
-104	Skip-in for SUBUSEDY 1	114	8.4 %
-103	Skip-in for SUBUSEDY <> 1	1117	82.5 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 2166-2169 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_DRUG24OTHER: S4 SubUse_Drug24: Used Other (specify) past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	123	9.1 %

Value	Label	Unweighted Frequency	%
-104	Skip-in for SUBUSEDY 1	114	8.4 %
-103	Skip-in for SUBUSEDY <> 1	1117	82.5 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 2170-2173 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_DRUG24NONE: S4 SubUse_Drug24: Used NA (No drugs) past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	123	9.1 %
-104	Skip-in for SUBUSEDY 1	114	8.4 %
-103	Skip-in for SUBUSEDY <> 1	1117	82.5 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 2174-2177 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S4SUBUSE29: S4 SubUse: Specify other drug used past 24hrs

Based upon 1,354 valid cases out of 1,354 total cases.

Location: 2178-2178 (width: 1; decimal: 0)
Variable Type: character

S4SUBUSE30: S4 SubUse: Num times smoked cigarettes during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	603	44.5 %
2	1-2 times	27	2.0 %
3	3-5 times	22	1.6 %
4	1X per month	9	0.7 %
5	2-3X per month	23	1.7 %
6	1X per week	19	1.4 %
7	2-3X per week	42	3.1 %
8	4-5X per week	35	2.6 %
9	Every day	448	33.1 %

Value	Label	Unweighted Frequency	%
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-99	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,228 valid cases out of 1,354 total cases.

Location: 2179-2182 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S4SUBUSE31: S4 SubUse: Num cigarettes smoke per day during recall period

Value	Label	Unweighted Frequency	%
1	0 cigarettes	617	45.6 %
2	1-4 cigarettes	229	16.9 %
3	5-9 cigarettes	163	12.0 %
4	10-15 cigarettes	107	7.9 %
5	16-20 cigarettes	84	6.2 %
6	More than a pack a day cigarettes	29	2.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
	Total	**1,354**	**100%**

Based upon 1,229 valid cases out of 1,354 total cases.

Location: 2183-2186 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -105 , -9 , -8

S4SUBUSE_SUBSYM01_SUSYMB: S4 SubUse: Family ever complain because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	161	11.9 %
5	No	515	38.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-106	Skip-in for alchstat/drugstat	553	40.8 %
	Total	**1,354**	**100%**

Based upon 676 valid cases out of 1,354 total cases.

Location: 2187-2190 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM01_SUSYMC: S4 SubUse: Specify cause of family complaint [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	39	2.9 %
2	Drugs	73	5.4 %
3	Both	49	3.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	6	0.4 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-107	Follow-up skipped because gate <> 1	1064	78.6 %
	Total	**1,354**	**100%**

Based upon 161 valid cases out of 1,354 total cases.

Location: 2191-2194 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM02_SUSYMB: S4 SubUse: Friend ever complain because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	55	4.1 %
5	No	621	45.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-106	Skip-in for alchstat/drugstat	553	40.8 %
	Total	**1,354**	**100%**

Based upon 676 valid cases out of 1,354 total cases.

Location: 2195-2198 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM02_SUSYMC: S4 SubUse: Specify cause of friend complain [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	14	1.0 %

Value	Label	Unweighted Frequency	%
2	Drugs	26	1.9 %
3	Both	15	1.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-107	Follow-up skipped because gate <> 1	1168	86.3 %
	Total	**1,354**	**100%**

Based upon 55 valid cases out of 1,354 total cases.

Location: 2199-2202 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM03_SUSYMB: S4 SubUse: Ever trouble at work because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	28	2.1 %
5	No	648	47.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-106	Skip-in for alchstat/drugstat	553	40.8 %
	Total	**1,354**	**100%**

Based upon 676 valid cases out of 1,354 total cases.

Location: 2203-2206 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM03_SUSYMC: S4 SubUse: Specify cause of trouble at work [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	6	0.4 %
2	Drugs	15	1.1 %
3	Both	7	0.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	7	0.5 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-107	Follow-up skipped because gate <> 1	1196	88.3 %
	Total	**1,354**	**100%**

Based upon 28 valid cases out of 1,354 total cases.

Location: 2207-2210 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM04_SUSYMB: S4 SubUse: Ever missed work/school because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	44	3.2 %
5	No	632	46.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-106	Skip-in for alchstat/drugstat	553	40.8 %
	Total	**1,354**	**100%**

Based upon 676 valid cases out of 1,354 total cases.

Location: 2211-2214 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM04_SUSYMC: S4 SubUse: Specify cause of missed work/school [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	17	1.3 %
2	Drugs	15	1.1 %
3	Both	12	0.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	7	0.5 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-107	Follow-up skipped because gate <> 1	1180	87.1 %
	Total	**1,354**	**100%**

Based upon 44 valid cases out of 1,354 total cases.

Location: 2215-2218 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM05_SUSYMB: S4 SubUse: Ever arrested because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	72	5.3 %

Value	Label	Unweighted Frequency	%
5	No	604	44.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-106	Skip-in for alchstat/drugstat	553	40.8 %
	Total	**1,354**	**100%**

Based upon 676 valid cases out of 1,354 total cases.

Location: 2219-2222 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM05_SUSYMC: S4 SubUse: Specify cause of arrest [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	20	1.5 %
2	Drugs	41	3.0 %
3	Both	11	0.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	7	0.5 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-107	Follow-up skipped because gate <> 1	1152	85.1 %
	Total	**1,354**	**100%**

Based upon 72 valid cases out of 1,354 total cases.

Location: 2223-2226 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM06_SUSYMB: S4 SubUse: Ever argue family/friends because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	102	7.5 %
5	No	574	42.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-106	Skip-in for alchstat/drugstat	553	40.8 %
	Total	**1,354**	**100%**

Based upon 676 valid cases out of 1,354 total cases.

Location: 2227-2230 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM06_SUSYMC: S4 SubUse: Specify cause of argument w/ family/friends [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	23	1.7 %
2	Drugs	45	3.3 %
3	Both	34	2.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	7	0.5 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-107	Follow-up skipped because gate <> 1	1122	82.9 %
	Total	**1,354**	**100%**

Based upon 102 valid cases out of 1,354 total cases.

Location: 2231-2234 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM07_SUSYMB: S4 SubUse: Ever cont. drug/alcohol use in spite cause health problems in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	80	5.9 %
5	No	596	44.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-106	Skip-in for alchstat/drugstat	553	40.8 %
	Total	**1,354**	**100%**

Based upon 676 valid cases out of 1,354 total cases.

Location: 2235-2238 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM07_SUSYMC: S4 SubUse: Specify continued using [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	27	2.0 %
2	Drugs	27	2.0 %
3	Both	26	1.9 %

Value	Label	Unweighted Frequency	%
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-107	Follow-up skipped because gate <> 1	1143	84.4 %
	Total	**1,354**	**100%**

Based upon 80 valid cases out of 1,354 total cases.

Location: 2239-2242 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM08_SUSYMB: S4 SubUse: Ever not remember prior night because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	140	10.3 %
5	No	536	39.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-106	Skip-in for alchstat/drugstat	553	40.8 %
	Total	**1,354**	**100%**

Based upon 676 valid cases out of 1,354 total cases.

Location: 2243-2246 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM08_SUSYMC: S4 SubUse: Specify cause of not remember night [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	70	5.2 %
2	Drugs	20	1.5 %
3	Both	50	3.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	7	0.5 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-107	Follow-up skipped because gate <> 1	1084	80.1 %
	Total	**1,354**	**100%**

Based upon 140 valid cases out of 1,354 total cases.

Location: 2247-2250 (width: 4; decimal: 0)

Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM09_SUSYMB: S4 SubUse: Ever used in situations could get hurt in recall period [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	152	11.2 %
5	No	524	38.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-106	Skip-in for alchstat/drugstat	553	40.8 %
	Total	**1,354**	**100%**

Based upon 676 valid cases out of 1,354 total cases.

Location: 2251-2254 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM09_SUSYMC: S4 SubUse: Specify used drugs in hurtful situations [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	59	4.4 %
2	Drugs	37	2.7 %
3	Both	56	4.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	6	0.4 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-107	Follow-up skipped because gate <> 1	1073	79.2 %
	Total	**1,354**	**100%**

Based upon 152 valid cases out of 1,354 total cases.

Location: 2255-2258 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM10_SUSYMB: S4 SubUse: Ever had accident/injury because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	43	3.2 %
5	No	633	46.8 %
	Missing Data		

Value	Label	Unweighted Frequency	%
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-106	Skip-in for alchstat/drugstat	553	40.8 %
	Total	**1,354**	**100%**

Based upon 676 valid cases out of 1,354 total cases.

Location: 2259-2262 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM10_SUSYMC: S4 SubUse: Specify cause of accident/injury [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	27	2.0 %
2	Drugs	7	0.5 %
3	Both	9	0.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-107	Follow-up skipped because gate <> 1	1180	87.1 %
	Total	**1,354**	**100%**

Based upon 43 valid cases out of 1,354 total cases.

Location: 2263-2266 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM11_SUSYMB: S4 SubUse: Alcohol/drug use interfere work/responsibility school/home/job in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	77	5.7 %
5	No	599	44.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-106	Skip-in for alchstat/drugstat	553	40.8 %
	Total	**1,354**	**100%**

Based upon 676 valid cases out of 1,354 total cases.

Location: 2267-2270 (width: 4; decimal: 0)
Variable Type: numeric

(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM11_SUSYMC: S4 SubUse: Specify cause of interference [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	22	1.6 %
2	Drugs	31	2.3 %
3	Both	24	1.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	7	0.5 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-107	Follow-up skipped because gate <> 1	1147	84.7 %
	Total	**1,354**	**100%**

Based upon 77 valid cases out of 1,354 total cases.

Location: 2271-2274 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM12_SUSYMB: S4 SubUse: Ever trouble police no arrest because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	63	4.7 %
5	No	613	45.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-106	Skip-in for alchstat/drugstat	553	40.8 %
	Total	**1,354**	**100%**

Based upon 676 valid cases out of 1,354 total cases.

Location: 2275-2278 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM12_SUSYMC: S4 SubUse: Specify cause trouble police no arrest [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	27	2.0 %
2	Drugs	24	1.8 %
3	Both	12	0.9 %

Value	Label	Unweighted Frequency	%
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	7	0.5 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-107	Follow-up skipped because gate <> 1	1161	85.7 %
	Total	**1,354**	**100%**

Based upon 63 valid cases out of 1,354 total cases.

Location: 2279-2282 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM13_SUSYMB: S4 SubUse: Sold/exchanged sex for drugs in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	3	0.2 %
5	No	673	49.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-106	Skip-in for alchstat/drugstat	553	40.8 %
	Total	**1,354**	**100%**

Based upon 676 valid cases out of 1,354 total cases.

Location: 2283-2286 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM13_SUSYMC: S4 SubUse: Specify cause sold/exchanged sex [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	1	0.1 %
2	Drugs	2	0.1 %
3	Both	0	0.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-107	Follow-up skipped because gate <> 1	1220	90.1 %
	Total	**1,354**	**100%**

Based upon 3 valid cases out of 1,354 total cases.

Location: 2287-2290 (width: 4; decimal: 0)
Variable Type: numeric

(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM14_SUSYMB: S4 SubUse: Sold drugs for money buy drugs in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	75	5.5 %
5	No	601	44.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-106	Skip-in for alchstat/drugstat	553	40.8 %
	Total	**1,354**	**100%**

Based upon 676 valid cases out of 1,354 total cases.

Location: 2291-2294 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM14_SUSYMC: S4 SubUse: Specify cause sold drugs for money to buy [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	2	0.1 %
2	Drugs	63	4.7 %
3	Both	10	0.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-107	Follow-up skipped because gate <> 1	1148	84.8 %
	Total	**1,354**	**100%**

Based upon 75 valid cases out of 1,354 total cases.

Location: 2295-2298 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM15_SUSYMB: S4 SubUse: Stole for money buy drugs in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	32	2.4 %
5	No	644	47.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %

Value	Label	Unweighted Frequency	%
-106	Skip-in for alchstat/drugstat	553	40.8 %
	Total	**1,354**	**100%**

Based upon 676 valid cases out of 1,354 total cases.

Location: 2299-2302 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM15_SUSYMC: S4 SubUse: Specify cause stole for money to buy [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	1	0.1 %
2	Drugs	25	1.8 %
3	Both	6	0.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	7	0.5 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-107	Follow-up skipped because gate <> 1	1192	88.0 %
	Total	**1,354**	**100%**

Based upon 32 valid cases out of 1,354 total cases.

Location: 2303-2306 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM16_SUSYMB: S4 SubUse: Ever had physical fight because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	60	4.4 %
5	No	616	45.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-106	Skip-in for alchstat/drugstat	553	40.8 %
	Total	**1,354**	**100%**

Based upon 676 valid cases out of 1,354 total cases.

Location: 2307-2310 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM16_SUSYMC: S4 SubUse: Specify cause of fight [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	31	2.3 %
2	Drugs	15	1.1 %
3	Both	14	1.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	6	0.4 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-107	Follow-up skipped because gate <> 1	1165	86.0 %
	Total	**1,354**	**100%**

Based upon 60 valid cases out of 1,354 total cases.

Location: 2311-2314 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM17_SUSYMB: S4 SubUse: Ever destroy property because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	41	3.0 %
5	No	635	46.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-106	Skip-in for alchstat/drugstat	553	40.8 %
	Total	**1,354**	**100%**

Based upon 676 valid cases out of 1,354 total cases.

Location: 2315-2318 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM17_SUSYMC: S4 SubUse: Specify cause of destroy property [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	13	1.0 %
2	Drugs	19	1.4 %
3	Both	9	0.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %

Value	Label	Unweighted Frequency	%
-201	IntCompStat = 4 missed interview	123	9.1 %
-107	Follow-up skipped because gate <> 1	1182	87.3 %
	Total	**1,354**	**100%**

Based upon 41 valid cases out of 1,354 total cases.

Location: 2319-2322 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM18_SUSYMB: S4 SubUse: Ever felt strong urge use drug/alcohol no self stop in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	66	4.9 %
5	No	610	45.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-106	Skip-in for alchstat/drugstat	553	40.8 %
	Total	**1,354**	**100%**

Based upon 676 valid cases out of 1,354 total cases.

Location: 2323-2326 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM18_SUSYMC: S4 SubUse: Specify the cause of strong urge [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	12	0.9 %
2	Drugs	37	2.7 %
3	Both	17	1.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	7	0.5 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-107	Follow-up skipped because gate <> 1	1158	85.5 %
	Total	**1,354**	**100%**

Based upon 66 valid cases out of 1,354 total cases.

Location: 2327-2330 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM19_SUSYMB: S4 SubUse: Ever want drug/alcohol think of nothing else in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	61	4.5 %
5	No	615	45.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-106	Skip-in for alchstat/drugstat	553	40.8 %
	Total	**1,354**	**100%**

Based upon 676 valid cases out of 1,354 total cases.

Location: 2331-2334 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM19_SUSYMC: S4 SubUse: Specify cause of want [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	9	0.7 %
2	Drugs	40	3.0 %
3	Both	12	0.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	7	0.5 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-107	Follow-up skipped because gate <> 1	1163	85.9 %
	Total	**1,354**	**100%**

Based upon 61 valid cases out of 1,354 total cases.

Location: 2335-2338 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM20_SUSYMB: S4 SubUse: Ever experience withdrawal from drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	48	3.5 %
5	No	628	46.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %

Value	Label	Unweighted Frequency	%
-106	Skip-in for alchstat/drugstat	553	40.8 %
	Total	**1,354**	**100%**

Based upon 676 valid cases out of 1,354 total cases.

Location: 2339-2342 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM20_SUSYMC: S4 SubUse: Specify cause of withdraw [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	10	0.7 %
2	Drugs	31	2.3 %
3	Both	7	0.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-107	Follow-up skipped because gate <> 1	1175	86.8 %
	Total	**1,354**	**100%**

Based upon 48 valid cases out of 1,354 total cases.

Location: 2343-2346 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM21_SUSYMB: S4 SubUse: Take drug/alcohol to keep from having withdraw in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	26	1.9 %
5	No	22	1.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	6	0.4 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-108	Skip-in for SUBSYM20=5	621	45.9 %
-106	Skip-in for alchstat/drugstat	556	41.1 %
	Total	**1,354**	**100%**

Based upon 48 valid cases out of 1,354 total cases.

Location: 2347-2350 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM21_SUSYMC: S4 SubUse: Specify reason for consumption [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	6	0.4 %
2	Drugs	19	1.4 %
3	Both	1	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-107	Follow-up skipped because gate <> 1	1197	88.4 %
	Total	**1,354**	**100%**

Based upon 26 valid cases out of 1,354 total cases.

Location: 2351-2354 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM22_SUSYMB: S4 SubUse: Give up usual activities/interests to use drug/alcohol in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	58	4.3 %
5	No	618	45.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-106	Skip-in for alchstat/drugstat	553	40.8 %
	Total	**1,354**	**100%**

Based upon 676 valid cases out of 1,354 total cases.

Location: 2355-2358 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM22_SUSYMC: S4 SubUse: Specify cause of giving up activities/interests [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	9	0.7 %
2	Drugs	32	2.4 %
3	Both	17	1.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %

Value	Label	Unweighted Frequency	%
-201	IntCompStat = 4 missed interview	123	9.1 %
-107	Follow-up skipped because gate <> 1	1165	86.0 %
	Total	**1,354**	**100%**

Based upon 58 valid cases out of 1,354 total cases.

Location: 2359-2362 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM23_SUSYMB: S4 SubUse: Ever used in rp when promised wouldn't [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	118	8.7 %
5	No	558	41.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-106	Skip-in for alchstat/drugstat	553	40.8 %
	Total	**1,354**	**100%**

Based upon 676 valid cases out of 1,354 total cases.

Location: 2363-2366 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM23_SUSYMC: S4 SubUse: Specify what use you started [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	28	2.1 %
2	Drugs	52	3.8 %
3	Both	38	2.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	7	0.5 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-107	Follow-up skipped because gate <> 1	1106	81.7 %
	Total	**1,354**	**100%**

Based upon 118 valid cases out of 1,354 total cases.

Location: 2367-2370 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM24_SUSYMB: S4 SubUse: Ever use drug/alcohol more freq than intended in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	122	9.0 %
5	No	554	40.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-106	Skip-in for alchstat/drugstat	553	40.8 %
	Total	**1,354**	**100%**

Based upon 676 valid cases out of 1,354 total cases.

Location: 2371-2374 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM24_SUSYMC: S4 SubUse: Specify what you used more frequently [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	28	2.1 %
2	Drugs	63	4.7 %
3	Both	31	2.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	7	0.5 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-107	Follow-up skipped because gate <> 1	1102	81.4 %
	Total	**1,354**	**100%**

Based upon 122 valid cases out of 1,354 total cases.

Location: 2375-2378 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM25_SUSYMB: S4 SubUse: Ever tried cut down drug/alcohol use but couldnt in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	82	6.1 %
5	No	594	43.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %

Value	Label	Unweighted Frequency	%
-106	Skip-in for alchstat/drugstat	553	40.8 %
	Total	**1,354**	**100%**

Based upon 676 valid cases out of 1,354 total cases.

Location: 2379-2382 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM25_SUSYMC: S4 SubUse: Specify what substance use you tried to curtail [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	12	0.9 %
2	Drugs	52	3.8 %
3	Both	18	1.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	7	0.5 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-107	Follow-up skipped because gate <> 1	1142	84.3 %
	Total	**1,354**	**100%**

Based upon 82 valid cases out of 1,354 total cases.

Location: 2383-2386 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM26_SUSYMB: S4 SubUse: Ever need more drug/alcohol get same effect in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	125	9.2 %
5	No	551	40.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-106	Skip-in for alchstat/drugstat	553	40.8 %
	Total	**1,354**	**100%**

Based upon 676 valid cases out of 1,354 total cases.

Location: 2387-2390 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM26_SUSYMC: S4 SubUse: Specify substance need more for same effect [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	28	2.1 %
2	Drugs	65	4.8 %
3	Both	32	2.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	7	0.5 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-107	Follow-up skipped because gate <> 1	1099	81.2 %
	Total	**1,354**	**100%**

Based upon 125 valid cases out of 1,354 total cases.

Location: 2391-2394 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM27_SUSYMB: S4 SubUse: Ever used drug/alcohol enough to depend in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	55	4.1 %
5	No	621	45.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-106	Skip-in for alchstat/drugstat	553	40.8 %
	Total	**1,354**	**100%**

Based upon 676 valid cases out of 1,354 total cases.

Location: 2395-2398 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM27_SUSYMC: S4 SubUse: Specify dependency [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	7	0.5 %
2	Drugs	41	3.0 %
3	Both	7	0.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %

Value	Label	Unweighted Frequency	%
-201	IntCompStat = 4 missed interview	123	9.1 %
-107	Follow-up skipped because gate <> 1	1168	86.3 %
	Total	**1,354**	**100%**

Based upon 55 valid cases out of 1,354 total cases.

Location: 2399-2402 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM28_SUSYMB: S4 SubUse: Ever talked doctor/clergy/guidance counselor drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	27	2.0 %
5	No	649	47.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-106	Skip-in for alchstat/drugstat	553	40.8 %
	Total	**1,354**	**100%**

Based upon 676 valid cases out of 1,354 total cases.

Location: 2403-2406 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM28_SUSYMC: S4 SubUse: Specify talked doctor/clergy/guidance counselor [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	6	0.4 %
2	Drugs	12	0.9 %
3	Both	9	0.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	7	0.5 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-107	Follow-up skipped because gate <> 1	1197	88.4 %
	Total	**1,354**	**100%**

Based upon 27 valid cases out of 1,354 total cases.

Location: 2407-2410 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM29_SUSYMB: S4 SubUse: Ever attended self-help group for drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	40	3.0 %
5	No	636	47.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-106	Skip-in for alchstat/drugstat	553	40.8 %
	Total	**1,354**	**100%**

Based upon 676 valid cases out of 1,354 total cases.

Location: 2411-2414 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM29_SUSYMC: S4 SubUse: Specify reason for self-help [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	11	0.8 %
2	Drugs	18	1.3 %
3	Both	11	0.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	7	0.5 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-107	Follow-up skipped because gate <> 1	1184	87.4 %
	Total	**1,354**	**100%**

Based upon 40 valid cases out of 1,354 total cases.

Location: 2415-2418 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM30_SUSYMB: S4 SubUse: Ever treated by therapist because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	26	1.9 %
5	No	650	48.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %

Value	Label	Unweighted Frequency	%
-106	Skip-in for alchstat/drugstat	553	40.8 %
	Total	**1,354**	**100%**

Based upon 676 valid cases out of 1,354 total cases.

Location: 2419-2422 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM30_SUSYMC: S4 SubUse: Specify reason for therapist visit [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	5	0.4 %
2	Drugs	7	0.5 %
3	Both	14	1.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	6	0.4 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-107	Follow-up skipped because gate <> 1	1199	88.6 %
	Total	**1,354**	**100%**

Based upon 26 valid cases out of 1,354 total cases.

Location: 2423-2426 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM31_SUSYMB: S4 SubUse: Ever Admitted to hospital because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	10	0.7 %
5	No	666	49.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-106	Skip-in for alchstat/drugstat	553	40.8 %
	Total	**1,354**	**100%**

Based upon 676 valid cases out of 1,354 total cases.

Location: 2427-2430 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE_SUBSYM31_SUSYMC: S4 SubUse: Specify reason admitted to hospital [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	4	0.3 %
2	Drugs	4	0.3 %
3	Both	2	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	7	0.5 %
-201	IntCompStat = 4 missed interview	123	9.1 %
-107	Follow-up skipped because gate <> 1	1214	89.7 %
	Total	**1,354**	**100%**

Based upon 10 valid cases out of 1,354 total cases.

Location: 2431-2434 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S4SUBUSE28: S4 SubUse: Num times abused prescribed medications (own or someone elses) during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	0	0.0 %
2	1-2 times	2	0.1 %
3	3-5 times	0	0.0 %
4	1X per month	0	0.0 %
5	2-3X per month	0	0.0 %
6	1X per week	0	0.0 %
7	2-3X per week	1	0.1 %
8	4-5X per week	0	0.0 %
9	Every day	0	0.0 %
	Missing Data		
-800	Question not asked for this subject	1226	90.5 %
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
	Total	**1,354**	**100%**

Based upon 3 valid cases out of 1,354 total cases.

Location: 2435-2438 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S4QFBRMO: S4 SubUse: Quant/freq used beer/wine/liquor - recall period

Based upon 1,226 valid cases out of 1,354 total cases.

- Mean: 10.74
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 480.00
- Standard Deviation: 27.32

Location: 2439-2442 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -300 , -202 , -201 , -100 , -9 , -8

S46MOUSE: S4 SubUse: Number drugs used - recall period

Based upon 1,229 valid cases out of 1,354 total cases.

- Mean: 0.59
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 9.00
- Standard Deviation: 1.07

Location: 2443-2446 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -300 , -202 , -201 , -100 , -9 , -8

S4QFSMK6: S4 SubUse: Quant/freq used cigarettes - recall period

Based upon 1,229 valid cases out of 1,354 total cases.

- Mean: 31.36
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 160.00
- Standard Deviation: 46.71

Location: 2447-2452 (width: 6; decimal: 1)
Variable Type: numeric
(Range of) Missing Values: -300.0 , -202.0 , -201.0 , -100.0 , -9.0 , -8.0

S4A6SC2: S4 SubUse: Recall period alcohol consequences

Based upon 1,229 valid cases out of 1,354 total cases.

- Mean: 0.60
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 17.00
- Standard Deviation: 1.67

Location: 2453-2456 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S4D6SC2: S4 SubUse: Recall period drug consequences

Based upon 1,229 valid cases out of 1,354 total cases.

- Mean: 0.67
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 15.00
- Standard Deviation: 1.91

Location: 2457-2460 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S4A6DP2: S4 SubUse: Recall period alcohol dependence symptoms

Based upon 1,229 valid cases out of 1,354 total cases.

- Mean: 0.27
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 10.00
- Standard Deviation: 1.01

Location: 2461-2464 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S4D6DP2: S4 SubUse: Recall period drug dependence symptoms

Based upon 1,229 valid cases out of 1,354 total cases.

- Mean: 0.50
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 10.00
- Standard Deviation: 1.53

Location: 2465-2468 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S4TOTSC6: S4 SubUse: Recall period total substance use consequences

Based upon 1,229 valid cases out of 1,354 total cases.

- Mean: 1.00
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 17.00
- Standard Deviation: 2.31

Location: 2469-2472 (width: 4; decimal: 0)

Variable Type: numeric
(Range of) Missing Values: -202 , -201

S4TOTDS6: S4 SubUse: Recall period total substance use dependency symp

Based upon 1,229 valid cases out of 1,354 total cases.

- Mean: 0.62
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 10.00
- Standard Deviation: 1.69

Location: 2473-2476 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S4ALTRTM: S4 SubUse: Recall period alcohol treatment

Value	Label	Unweighted Frequency	%
0	No	1193	88.1 %
1	Yes	36	2.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
	Total	**1,354**	**100%**

Based upon 1,229 valid cases out of 1,354 total cases.

Location: 2477-2480 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S4DGTRTM: S4 SubUse: Recall period drug treatment

Value	Label	Unweighted Frequency	%
0	No	1184	87.4 %
1	Yes	45	3.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
	Total	**1,354**	**100%**

Based upon 1,229 valid cases out of 1,354 total cases.

Location: 2481-2484 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S4TRTM: S4 SubUse: Recall period alcohol or drug treatment

Value	Label	Unweighted Frequency	%
0	No	1170	86.4 %
1	Yes	59	4.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	123	9.1 %
	Total	**1,354**	**100%**

Based upon 1,229 valid cases out of 1,354 total cases.

Location: 2485-2488 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S5SUBUSE1: S5 SubUse: How often had alcohol to drink in recall period

Value	Label	Unweighted Frequency	%
1	Not at all	658	48.6 %
2	1-2 times	150	11.1 %
3	3-5 times	89	6.6 %
4	1X per month	49	3.6 %
5	2-3X per month	93	6.9 %
6	1X per week	67	4.9 %
7	2-3X per week	90	6.6 %
8	4-5X per week	24	1.8 %
9	Every day	13	1.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
	Total	**1,354**	**100%**

Based upon 1,233 valid cases out of 1,354 total cases.

Location: 2489-2492 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S5SUBUSE2: S5 SubUse: Num alcoholic drinks usually consumed

Based upon 1,231 valid cases out of 1,354 total cases.

- Mean: 2.38
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00

- Maximum: 30.00
- Standard Deviation: 4.00

Location: 2493-2496 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -101 , -99 , -98

S5SUBUSE3: S5 SubUse: How often had 5+ drinks at one time in recall period

Value	Label	Unweighted Frequency	%
1	Not at all	887	65.5 %
2	1-2 times	92	6.8 %
3	3-5 times	54	4.0 %
4	1X per month	36	2.7 %
5	2-3X per month	42	3.1 %
6	1X per week	43	3.2 %
7	2-3X per week	54	4.0 %
8	4-5X per week	15	1.1 %
9	Every day	10	0.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
	Total	**1,354**	**100%**

Based upon 1,233 valid cases out of 1,354 total cases.

Location: 2497-2500 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S5SUBUSE4: S5 SubUse: Num times been drunk on alcohol during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	848	62.6 %
2	1-2 times	153	11.3 %
3	3-5 times	73	5.4 %
4	1X per month	31	2.3 %
5	2-3X per month	42	3.1 %
6	1X per week	34	2.5 %
7	2-3X per week	38	2.8 %
8	4-5X per week	9	0.7 %
9	Every day	4	0.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %

Value	Label	Unweighted Frequency	%
-99	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,232 valid cases out of 1,354 total cases.

Location: 2501-2504 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S5SUBUSE6: S5 SubUse: Num times used marijuana/hashish during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	811	59.9 %
2	1-2 times	79	5.8 %
3	3-5 times	43	3.2 %
4	1X per month	19	1.4 %
5	2-3X per month	24	1.8 %
6	1X per week	21	1.6 %
7	2-3X per week	57	4.2 %
8	4-5X per week	43	3.2 %
9	Every day	135	10.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-99	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,232 valid cases out of 1,354 total cases.

Location: 2505-2508 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S5SUBUSE8: S5 SubUse: Num times used sedative/tranquilizers during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1188	87.7 %
2	1-2 times	13	1.0 %
3	3-5 times	11	0.8 %
4	1X per month	2	0.1 %
5	2-3X per month	7	0.5 %
6	1X per week	7	0.5 %
7	2-3X per week	4	0.3 %
8	4-5X per week	0	0.0 %

Value	Label	Unweighted Frequency	%
9	Every day	1	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
	Total	**1,354**	**100%**

Based upon 1,233 valid cases out of 1,354 total cases.

Location: 2509-2512 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S5SUBUSE10: S5 SubUse: Num times used stimulants/amphetamines during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1171	86.5 %
2	1-2 times	7	0.5 %
3	3-5 times	8	0.6 %
4	1X per month	5	0.4 %
5	2-3X per month	9	0.7 %
6	1X per week	6	0.4 %
7	2-3X per week	7	0.5 %
8	4-5X per week	7	0.5 %
9	Every day	13	1.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
	Total	**1,354**	**100%**

Based upon 1,233 valid cases out of 1,354 total cases.

Location: 2513-2516 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S5SUBUSE12: S5 SubUse: Num times used cocaine during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1189	87.8 %
2	1-2 times	16	1.2 %
3	3-5 times	12	0.9 %
4	1X per month	2	0.1 %
5	2-3X per month	4	0.3 %
6	1X per week	4	0.3 %

Value	Label	Unweighted Frequency	%
7	2-3X per week	3	0.2 %
8	4-5X per week	1	0.1 %
9	Every day	2	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
	Total	**1,354**	**100%**

Based upon 1,233 valid cases out of 1,354 total cases.

Location: 2517-2520 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S5SUBUSE14: S5 SubUse: Num times used opiates during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1207	89.1 %
2	1-2 times	6	0.4 %
3	3-5 times	2	0.1 %
4	1X per month	2	0.1 %
5	2-3X per month	6	0.4 %
6	1X per week	4	0.3 %
7	2-3X per week	2	0.1 %
8	4-5X per week	0	0.0 %
9	Every day	4	0.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
	Total	**1,354**	**100%**

Based upon 1,233 valid cases out of 1,354 total cases.

Location: 2521-2524 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S5SUBUSE16: S5 SubUse: Num times used ecstasy during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1210	89.4 %
2	1-2 times	13	1.0 %
3	3-5 times	8	0.6 %
4	1X per month	0	0.0 %

Value	Label	Unweighted Frequency	%
5	2-3X per month	1	0.1 %
6	1X per week	0	0.0 %
7	2-3X per week	0	0.0 %
8	4-5X per week	0	0.0 %
9	Every day	1	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
	Total	**1,354**	**100%**

Based upon 1,233 valid cases out of 1,354 total cases.

Location: 2525-2528 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S5SUBUSE18: S5 SubUse: Num times used hallucinogens during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1192	88.0 %
2	1-2 times	20	1.5 %
3	3-5 times	8	0.6 %
4	1X per month	3	0.2 %
5	2-3X per month	3	0.2 %
6	1X per week	0	0.0 %
7	2-3X per week	2	0.1 %
8	4-5X per week	2	0.1 %
9	Every day	3	0.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
	Total	**1,354**	**100%**

Based upon 1,233 valid cases out of 1,354 total cases.

Location: 2529-2532 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S5SUBUSE20: S5 SubUse: Num times used inhalants during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1222	90.3 %
2	1-2 times	6	0.4 %

Value	Label	Unweighted Frequency	%
3	3-5 times	1	0.1 %
4	1X per month	0	0.0 %
5	2-3X per month	2	0.1 %
6	1X per week	0	0.0 %
7	2-3X per week	0	0.0 %
8	4-5X per week	0	0.0 %
9	Every day	2	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
	Total	**1,354**	**100%**

Based upon 1,233 valid cases out of 1,354 total cases.

Location: 2533-2536 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S5SUBUSE22: S5 SubUse: Num times used amyl nitrate/odorizers/rush during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1229	90.8 %
2	1-2 times	1	0.1 %
3	3-5 times	0	0.0 %
4	1X per month	0	0.0 %
5	2-3X per month	2	0.1 %
6	1X per week	0	0.0 %
7	2-3X per week	0	0.0 %
8	4-5X per week	0	0.0 %
9	Every day	1	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
	Total	**1,354**	**100%**

Based upon 1,233 valid cases out of 1,354 total cases.

Location: 2537-2540 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S5SUBUSE24: S5 SubUse: Used other drugs during recall period [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	1	0.1 %
5	No	1232	91.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
	Total	**1,354**	**100%**

Based upon 1,233 valid cases out of 1,354 total cases.

Location: 2541-2544 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE26: S5 SubUse: Num times used other drugs during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1232	91.0 %
2	1-2 times	1	0.1 %
3	3-5 times	0	0.0 %
4	1X per month	0	0.0 %
5	2-3X per month	0	0.0 %
6	1X per week	0	0.0 %
7	2-3X per week	0	0.0 %
8	4-5X per week	0	0.0 %
9	Every day	0	0.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
	Total	**1,354**	**100%**

Based upon 1,233 valid cases out of 1,354 total cases.

Location: 2545-2548 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S5SUBUSEDY: S5 SubUse: Used any drugs past 24hrs [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	125	9.2 %
5	No	1108	81.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %

Value	Label	Unweighted Frequency	%
	Total	**1,354**	**100%**

Based upon 1,233 valid cases out of 1,354 total cases.

Location: 2549-2552 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_DRUG24MAR: S5 SubUse_Drug24: Used marijuana/hashish past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	116	8.6 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	120	8.9 %
-104	Skip-in for SUBUSEDY 1	10	0.7 %
-103	Skip-in for SUBUSEDY <> 1	1108	81.8 %
	Total	**1,354**	**100%**

Based upon 116 valid cases out of 1,354 total cases.

Location: 2553-2556 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_DRUG24TRANQ: S5 SubUse_Drug24: Used sedatives/tranquilizers past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	2	0.1 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	120	8.9 %
-104	Skip-in for SUBUSEDY 1	124	9.2 %
-103	Skip-in for SUBUSEDY <> 1	1108	81.8 %
	Total	**1,354**	**100%**

Based upon 2 valid cases out of 1,354 total cases.

Location: 2557-2560 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_DRUG24AMPH: S5 SubUse_Drug24: Used stimulants/amphetamines past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	12	0.9 %

Value	Label	Unweighted Frequency	%
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	120	8.9 %
-104	Skip-in for SUBUSEDY 1	114	8.4 %
-103	Skip-in for SUBUSEDY <> 1	1108	81.8 %
	Total	**1,354**	**100%**

Based upon 12 valid cases out of 1,354 total cases.

Location: 2561-2564 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_DRUG24COC: S5 SubUse_Drug24: Used Cocaine past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	120	8.9 %
-104	Skip-in for SUBUSEDY 1	126	9.3 %
-103	Skip-in for SUBUSEDY <> 1	1108	81.8 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 2565-2568 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_DRUG24OPI: S5 SubUse_Drug24: Used Opiates past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	1	0.1 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	120	8.9 %
-104	Skip-in for SUBUSEDY 1	125	9.2 %
-103	Skip-in for SUBUSEDY <> 1	1108	81.8 %
	Total	**1,354**	**100%**

Based upon 1 valid cases out of 1,354 total cases.

Location: 2569-2572 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_DRUG24ECST: S5 SubUse_Drug24: Used Ecstacy past 24 hours

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	120	8.9 %
-104	Skip-in for SUBUSEDY 1	126	9.3 %
-103	Skip-in for SUBUSEDY <> 1	1108	81.8 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 2573-2576 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_DRUG24HALL: S5 SubUse_Drug24: Used Hallucinogens past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	1	0.1 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	120	8.9 %
-104	Skip-in for SUBUSEDY 1	125	9.2 %
-103	Skip-in for SUBUSEDY <> 1	1108	81.8 %
	Total	**1,354**	**100%**

Based upon 1 valid cases out of 1,354 total cases.

Location: 2577-2580 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_DRUG24INH: S5 SubUse_Drug24: Used Inhalants past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	120	8.9 %
-104	Skip-in for SUBUSEDY 1	126	9.3 %
-103	Skip-in for SUBUSEDY <> 1	1108	81.8 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 2581-2584 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_DRUG24AMYL: S5 SubUse_Drug24: Used Amyl nitrate/poppers/odorizers 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	120	8.9 %
-104	Skip-in for SUBUSEDY 1	126	9.3 %
-103	Skip-in for SUBUSEDY <> 1	1108	81.8 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 2585-2588 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_DRUG24OTHER: S5 SubUse_Drug24: Used Other (specify) past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	1	0.1 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	120	8.9 %
-104	Skip-in for SUBUSEDY 1	125	9.2 %
-103	Skip-in for SUBUSEDY <> 1	1108	81.8 %
	Total	**1,354**	**100%**

Based upon 1 valid cases out of 1,354 total cases.

Location: 2589-2592 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_DRUG24NONE: S5 SubUse_Drug24: Used NA (No drugs) past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	120	8.9 %

Value	Label	Unweighted Frequency	%
-104	Skip-in for SUBUSEDY 1	126	9.3 %
-103	Skip-in for SUBUSEDY <> 1	1108	81.8 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 2593-2596 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S5SUBUSE29: S5 SubUse: Specify other drug used past 24hrs

Based upon 1,354 valid cases out of 1,354 total cases.

Location: 2597-2617 (width: 21; decimal: 0)
Variable Type: character

S5SUBUSE30: S5 SubUse: Num times smoked cigarettes during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	572	42.2 %
2	1-2 times	35	2.6 %
3	3-5 times	16	1.2 %
4	1X per month	11	0.8 %
5	2-3X per month	8	0.6 %
6	1X per week	25	1.8 %
7	2-3X per week	40	3.0 %
8	4-5X per week	34	2.5 %
9	Every day	492	36.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
	Total	**1,354**	**100%**

Based upon 1,233 valid cases out of 1,354 total cases.

Location: 2618-2621 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S5SUBUSE31: S5 SubUse: Num cigarettes smoke per day during recall period

Value	Label	Unweighted Frequency	%
1	0 cigarettes	592	43.7 %
2	1-4 cigarettes	251	18.5 %
3	5-9 cigarettes	157	11.6 %

Value	Label	Unweighted Frequency	%
4	10-15 cigarettes	127	9.4 %
5	16-20 cigarettes	79	5.8 %
6	More than a pack a day cigarettes	27	2.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
	Total	**1,354**	**100%**

Based upon 1,233 valid cases out of 1,354 total cases.

Location: 2622-2625 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -105 , -9 , -8

S5SUBUSE_SUBSYM01_SUSYMB: S5 SubUse: Family ever complain because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	146	10.8 %
5	No	532	39.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-106	Skip-in for alchstat/drugstat	555	41.0 %
	Total	**1,354**	**100%**

Based upon 678 valid cases out of 1,354 total cases.

Location: 2626-2629 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM01_SUSYMC: S5 SubUse: Specify cause of family complaint [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	33	2.4 %
2	Drugs	74	5.5 %
3	Both	39	2.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	7	0.5 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-107	Follow-up skipped because gate <> 1	1081	79.8 %
	Total	**1,354**	**100%**

Based upon 146 valid cases out of 1,354 total cases.

Location: 2630-2633 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM02_SUSYMB: S5 SubUse: Friend ever complain because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	65	4.8 %
5	No	613	45.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-106	Skip-in for alchstat/drugstat	555	41.0 %
	Total	**1,354**	**100%**

Based upon 678 valid cases out of 1,354 total cases.

Location: 2634-2637 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM02_SUSYMC: S5 SubUse: Specify cause of friend complain [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	17	1.3 %
2	Drugs	35	2.6 %
3	Both	13	1.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	7	0.5 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-107	Follow-up skipped because gate <> 1	1162	85.8 %
	Total	**1,354**	**100%**

Based upon 65 valid cases out of 1,354 total cases.

Location: 2638-2641 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM03_SUSYMB: S5 SubUse: Ever trouble at work because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	30	2.2 %

Value	Label	Unweighted Frequency	%
5	No	648	47.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-106	Skip-in for alchstat/drugstat	555	41.0 %
	Total	**1,354**	**100%**

Based upon 678 valid cases out of 1,354 total cases.

Location: 2642-2645 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM03_SUSYMC: S5 SubUse: Specify cause of trouble at work [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	11	0.8 %
2	Drugs	17	1.3 %
3	Both	2	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-107	Follow-up skipped because gate <> 1	1196	88.3 %
	Total	**1,354**	**100%**

Based upon 30 valid cases out of 1,354 total cases.

Location: 2646-2649 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM04_SUSYMB: S5 SubUse: Ever missed work/school because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	36	2.7 %
5	No	642	47.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-106	Skip-in for alchstat/drugstat	555	41.0 %
	Total	**1,354**	**100%**

Based upon 678 valid cases out of 1,354 total cases.

Location: 2650-2653 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM04_SUSYMC: S5 SubUse: Specify cause of missed work/school [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	18	1.3 %
2	Drugs	11	0.8 %
3	Both	7	0.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-107	Follow-up skipped because gate <> 1	1190	87.9 %
	Total	**1,354**	**100%**

Based upon 36 valid cases out of 1,354 total cases.

Location: 2654-2657 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM05_SUSYMB: S5 SubUse: Ever arrested because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	59	4.4 %
5	No	618	45.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-106	Skip-in for alchstat/drugstat	555	41.0 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 677 valid cases out of 1,354 total cases.

Location: 2658-2661 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM05_SUSYMC: S5 SubUse: Specify cause of arrest [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	18	1.3 %

Value	Label	Unweighted Frequency	%
2	Drugs	31	2.3 %
3	Both	10	0.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	7	0.5 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-107	Follow-up skipped because gate <> 1	1168	86.3 %
	Total	**1,354**	**100%**

Based upon 59 valid cases out of 1,354 total cases.

Location: 2662-2665 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM06_SUSYMB: S5 SubUse: Ever argue family/friends because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	94	6.9 %
5	No	584	43.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-106	Skip-in for alchstat/drugstat	555	41.0 %
	Total	**1,354**	**100%**

Based upon 678 valid cases out of 1,354 total cases.

Location: 2666-2669 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM06_SUSYMC: S5 SubUse: Specify cause of argument w/ family/friends [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	24	1.8 %
2	Drugs	51	3.8 %
3	Both	19	1.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	7	0.5 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-107	Follow-up skipped because gate <> 1	1133	83.7 %
	Total	**1,354**	**100%**

Based upon 94 valid cases out of 1,354 total cases.

Location: 2670-2673 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM07_SUSYMB: S5 SubUse: Ever cont. drug/alcohol use in spite cause health problems in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	71	5.2 %
5	No	607	44.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-106	Skip-in for alchstat/drugstat	555	41.0 %
	Total	**1,354**	**100%**

Based upon 678 valid cases out of 1,354 total cases.

Location: 2674-2677 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM07_SUSYMC: S5 SubUse: Specify continued using [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	16	1.2 %
2	Drugs	33	2.4 %
3	Both	22	1.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	7	0.5 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-107	Follow-up skipped because gate <> 1	1156	85.4 %
	Total	**1,354**	**100%**

Based upon 71 valid cases out of 1,354 total cases.

Location: 2678-2681 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM08_SUSYMB: S5 SubUse: Ever not remember prior night because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	116	8.6 %
5	No	562	41.5 %

Value	Label	Unweighted Frequency	%
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-106	Skip-in for alchstat/drugstat	555	41.0 %
	Total	**1,354**	**100%**

Based upon 678 valid cases out of 1,354 total cases.

Location: 2682-2685 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM08_SUSYMC: S5 SubUse: Specify cause of not remember night [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	59	4.4 %
2	Drugs	21	1.6 %
3	Both	36	2.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	7	0.5 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-107	Follow-up skipped because gate <> 1	1111	82.1 %
	Total	**1,354**	**100%**

Based upon 116 valid cases out of 1,354 total cases.

Location: 2686-2689 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM09_SUSYMB: S5 SubUse: Ever used in situations could get hurt in recall period [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	115	8.5 %
5	No	563	41.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-106	Skip-in for alchstat/drugstat	555	41.0 %
	Total	**1,354**	**100%**

Based upon 678 valid cases out of 1,354 total cases.

Location: 2690-2693 (width: 4; decimal: 0)

Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM09_SUSYMC: S5 SubUse: Specify used drugs in hurtful situations [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	36	2.7 %
2	Drugs	38	2.8 %
3	Both	41	3.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	7	0.5 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-107	Follow-up skipped because gate <> 1	1112	82.1 %
	Total	**1,354**	**100%**

Based upon 115 valid cases out of 1,354 total cases.

Location: 2694-2697 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM10_SUSYMB: S5 SubUse: Ever had accident/injury because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	30	2.2 %
5	No	648	47.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-106	Skip-in for alchstat/drugstat	555	41.0 %
	Total	**1,354**	**100%**

Based upon 678 valid cases out of 1,354 total cases.

Location: 2698-2701 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM10_SUSYMC: S5 SubUse: Specify cause of accident/injury [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	16	1.2 %
2	Drugs	9	0.7 %
3	Both	5	0.4 %

Value	Label	Unweighted Frequency	%
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-107	Follow-up skipped because gate <> 1	1196	88.3 %
	Total	**1,354**	**100%**

Based upon 30 valid cases out of 1,354 total cases.

Location: 2702-2705 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM11_SUSYMB: S5 SubUse: Alcohol/drug use interfere work/responsibility school/home/job in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	62	4.6 %
5	No	616	45.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-106	Skip-in for alchstat/drugstat	555	41.0 %
	Total	**1,354**	**100%**

Based upon 678 valid cases out of 1,354 total cases.

Location: 2706-2709 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM11_SUSYMC: S5 SubUse: Specify cause of interference [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	19	1.4 %
2	Drugs	30	2.2 %
3	Both	13	1.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	7	0.5 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-107	Follow-up skipped because gate <> 1	1165	86.0 %
	Total	**1,354**	**100%**

Based upon 62 valid cases out of 1,354 total cases.

Location: 2710-2713 (width: 4; decimal: 0)

Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM12_SUSYMB: S5 SubUse: Ever trouble police no arrest because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	40	3.0 %
5	No	637	47.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-106	Skip-in for alchstat/drugstat	555	41.0 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 677 valid cases out of 1,354 total cases.

Location: 2714-2717 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM12_SUSYMC: S5 SubUse: Specify cause trouble police no arrest [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	16	1.2 %
2	Drugs	21	1.6 %
3	Both	3	0.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-107	Follow-up skipped because gate <> 1	1186	87.6 %
	Total	**1,354**	**100%**

Based upon 40 valid cases out of 1,354 total cases.

Location: 2718-2721 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM13_SUSYMB: S5 SubUse: Sold/exchanged sex for drugs in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	1	0.1 %
5	No	677	50.0 %
	Missing Data		

Value	Label	Unweighted Frequency	%
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-106	Skip-in for alchstat/drugstat	555	41.0 %
	Total	**1,354**	**100%**

Based upon 678 valid cases out of 1,354 total cases.

Location: 2722-2725 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM13_SUSYMC: S5 SubUse: Specify cause sold/exchanged sex [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	0	0.0 %
2	Drugs	1	0.1 %
3	Both	0	0.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-107	Follow-up skipped because gate <> 1	1225	90.5 %
	Total	**1,354**	**100%**

Based upon 1 valid cases out of 1,354 total cases.

Location: 2726-2729 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM14_SUSYMB: S5 SubUse: Sold drugs for money buy drugs in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	59	4.4 %
5	No	619	45.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-106	Skip-in for alchstat/drugstat	555	41.0 %
	Total	**1,354**	**100%**

Based upon 678 valid cases out of 1,354 total cases.

Location: 2730-2733 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM14_SUSYMC: S5 SubUse: Specify cause sold drugs for money to buy [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	0	0.0 %
2	Drugs	50	3.7 %
3	Both	9	0.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-107	Follow-up skipped because gate <> 1	1167	86.2 %
	Total	**1,354**	**100%**

Based upon 59 valid cases out of 1,354 total cases.

Location: 2734-2737 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM15_SUSYMB: S5 SubUse: Stole for money buy drugs in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	25	1.8 %
5	No	653	48.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-106	Skip-in for alchstat/drugstat	555	41.0 %
	Total	**1,354**	**100%**

Based upon 678 valid cases out of 1,354 total cases.

Location: 2738-2741 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM15_SUSYMC: S5 SubUse: Specify cause stole for money to buy [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	0	0.0 %
2	Drugs	19	1.4 %
3	Both	6	0.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %

Value	Label	Unweighted Frequency	%
-201	IntCompStat = 4 missed interview	120	8.9 %
-107	Follow-up skipped because gate <> 1	1201	88.7 %
	Total	**1,354**	**100%**

Based upon 25 valid cases out of 1,354 total cases.

Location: 2742-2745 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM16_SUSYMB: S5 SubUse: Ever had physical fight because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	53	3.9 %
5	No	624	46.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-106	Skip-in for alchstat/drugstat	555	41.0 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 677 valid cases out of 1,354 total cases.

Location: 2746-2749 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM16_SUSYMC: S5 SubUse: Specify cause of fight [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	31	2.3 %
2	Drugs	12	0.9 %
3	Both	10	0.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	7	0.5 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-107	Follow-up skipped because gate <> 1	1174	86.7 %
	Total	**1,354**	**100%**

Based upon 53 valid cases out of 1,354 total cases.

Location: 2750-2753 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM17_SUSYMB: S5 SubUse: Ever destroy property because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	26	1.9 %
5	No	651	48.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-106	Skip-in for alchstat/drugstat	555	41.0 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 677 valid cases out of 1,354 total cases.

Location: 2754-2757 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM17_SUSYMC: S5 SubUse: Specify cause of destroy property [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	11	0.8 %
2	Drugs	9	0.7 %
3	Both	6	0.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-107	Follow-up skipped because gate <> 1	1200	88.6 %
	Total	**1,354**	**100%**

Based upon 26 valid cases out of 1,354 total cases.

Location: 2758-2761 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM18_SUSYMB: S5 SubUse: Ever felt strong urge use drug/alcohol no self stop in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	62	4.6 %
5	No	616	45.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %

Value	Label	Unweighted Frequency	%
-201	IntCompStat = 4 missed interview	120	8.9 %
-106	Skip-in for alchstat/drugstat	555	41.0 %
	Total	**1,354**	**100%**

Based upon 678 valid cases out of 1,354 total cases.

Location: 2762-2765 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM18_SUSYMC: S5 SubUse: Specify the cause of strong urge [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	13	1.0 %
2	Drugs	36	2.7 %
3	Both	13	1.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	7	0.5 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-107	Follow-up skipped because gate <> 1	1165	86.0 %
	Total	**1,354**	**100%**

Based upon 62 valid cases out of 1,354 total cases.

Location: 2766-2769 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM19_SUSYMB: S5 SubUse: Ever want drug/alcohol think of nothing else in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	53	3.9 %
5	No	625	46.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-106	Skip-in for alchstat/drugstat	555	41.0 %
	Total	**1,354**	**100%**

Based upon 678 valid cases out of 1,354 total cases.

Location: 2770-2773 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM19_SUSYMC: S5 SubUse: Specify cause of want [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	12	0.9 %
2	Drugs	27	2.0 %
3	Both	14	1.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	7	0.5 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-107	Follow-up skipped because gate <> 1	1174	86.7 %
	Total	**1,354**	**100%**

Based upon 53 valid cases out of 1,354 total cases.

Location: 2774-2777 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM20_SUSYMB: S5 SubUse: Ever experience withdrawal from drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	40	3.0 %
5	No	637	47.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-106	Skip-in for alchstat/drugstat	555	41.0 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 677 valid cases out of 1,354 total cases.

Location: 2778-2781 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM20_SUSYMC: S5 SubUse: Specify cause of withdraw [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	3	0.2 %
2	Drugs	34	2.5 %
3	Both	3	0.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %

Value	Label	Unweighted Frequency	%
-201	IntCompStat = 4 missed interview	120	8.9 %
-107	Follow-up skipped because gate <> 1	1186	87.6 %
	Total	**1,354**	**100%**

Based upon 40 valid cases out of 1,354 total cases.

Location: 2782-2785 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM21_SUSYMB: S5 SubUse: Take drug/alcohol to keep from having withdraw in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	24	1.8 %
5	No	16	1.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-108	Skip-in for SUBSYM20=5	632	46.7 %
-106	Skip-in for alchstat/drugstat	557	41.1 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 40 valid cases out of 1,354 total cases.

Location: 2786-2789 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM21_SUSYMC: S5 SubUse: Specify reason for consumption [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	2	0.1 %
2	Drugs	18	1.3 %
3	Both	4	0.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-107	Follow-up skipped because gate <> 1	1202	88.8 %
	Total	**1,354**	**100%**

Based upon 24 valid cases out of 1,354 total cases.

Location: 2790-2793 (width: 4; decimal: 0)

Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM22_SUSYMB: S5 SubUse: Give up usual activities/interests to use drug/alcohol in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	52	3.8 %
5	No	626	46.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-106	Skip-in for alchstat/drugstat	555	41.0 %
	Total	**1,354**	**100%**

Based upon 678 valid cases out of 1,354 total cases.

Location: 2794-2797 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM22_SUSYMC: S5 SubUse: Specify cause of giving up activities/interests [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	8	0.6 %
2	Drugs	36	2.7 %
3	Both	8	0.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-107	Follow-up skipped because gate <> 1	1174	86.7 %
	Total	**1,354**	**100%**

Based upon 52 valid cases out of 1,354 total cases.

Location: 2798-2801 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM23_SUSYMB: S5 SubUse: Ever used in rp when promised wouldn't [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	110	8.1 %
5	No	568	41.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %

Value	Label	Unweighted Frequency	%
-201	IntCompStat = 4 missed interview	120	8.9 %
-106	Skip-in for alchstat/drugstat	555	41.0 %
	Total	**1,354**	**100%**

Based upon 678 valid cases out of 1,354 total cases.

Location: 2802-2805 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM23_SUSYMC: S5 SubUse: Specify what use you started [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	21	1.6 %
2	Drugs	63	4.7 %
3	Both	26	1.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	7	0.5 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-107	Follow-up skipped because gate <> 1	1117	82.5 %
	Total	**1,354**	**100%**

Based upon 110 valid cases out of 1,354 total cases.

Location: 2806-2809 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM24_SUSYMB: S5 SubUse: Ever use drug/alcohol more freq than intended in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	100	7.4 %
5	No	578	42.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-106	Skip-in for alchstat/drugstat	555	41.0 %
	Total	**1,354**	**100%**

Based upon 678 valid cases out of 1,354 total cases.

Location: 2810-2813 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM24_SUSYMC: S5 SubUse: Specify what you used more frequently [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	19	1.4 %
2	Drugs	58	4.3 %
3	Both	23	1.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	7	0.5 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-107	Follow-up skipped because gate <> 1	1127	83.2 %
	Total	**1,354**	**100%**

Based upon 100 valid cases out of 1,354 total cases.

Location: 2814-2817 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM25_SUSYMB: S5 SubUse: Ever tried cut down drug/alcohol use but couldnt in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	69	5.1 %
5	No	609	45.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-106	Skip-in for alchstat/drugstat	555	41.0 %
	Total	**1,354**	**100%**

Based upon 678 valid cases out of 1,354 total cases.

Location: 2818-2821 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM25_SUSYMC: S5 SubUse: Specify what substance use you tried to curtail [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	9	0.7 %
2	Drugs	51	3.8 %
3	Both	9	0.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	7	0.5 %

Value	Label	Unweighted Frequency	%
-201	IntCompStat = 4 missed interview	120	8.9 %
-107	Follow-up skipped because gate <> 1	1158	85.5 %
	Total	**1,354**	**100%**

Based upon 69 valid cases out of 1,354 total cases.

Location: 2822-2825 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM26_SUSYMB: S5 SubUse: Ever need more drug/alcohol get same effect in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	122	9.0 %
5	No	556	41.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-106	Skip-in for alchstat/drugstat	555	41.0 %
	Total	**1,354**	**100%**

Based upon 678 valid cases out of 1,354 total cases.

Location: 2826-2829 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM26_SUSYMC: S5 SubUse: Specify substance need more for same effect [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	26	1.9 %
2	Drugs	70	5.2 %
3	Both	26	1.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	7	0.5 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-107	Follow-up skipped because gate <> 1	1105	81.6 %
	Total	**1,354**	**100%**

Based upon 122 valid cases out of 1,354 total cases.

Location: 2830-2833 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM27_SUSYMB: S5 SubUse: Ever used drug/alcohol enough to depend in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	60	4.4 %
5	No	618	45.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-106	Skip-in for alchstat/drugstat	555	41.0 %
	Total	**1,354**	**100%**

Based upon 678 valid cases out of 1,354 total cases.

Location: 2834-2837 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM27_SUSYMC: S5 SubUse: Specify dependency [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	6	0.4 %
2	Drugs	42	3.1 %
3	Both	12	0.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	7	0.5 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-107	Follow-up skipped because gate <> 1	1167	86.2 %
	Total	**1,354**	**100%**

Based upon 60 valid cases out of 1,354 total cases.

Location: 2838-2841 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM28_SUSYMB: S5 SubUse: Ever talked doctor/clergy/guidance counselor drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	22	1.6 %
5	No	656	48.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %

Value	Label	Unweighted Frequency	%
-106	Skip-in for alchstat/drugstat	555	41.0 %
	Total	**1,354**	**100%**

Based upon 678 valid cases out of 1,354 total cases.

Location: 2842-2845 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM28_SUSYMC: S5 SubUse: Specify talked doctor/clergy/guidance counselor [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	2	0.1 %
2	Drugs	9	0.7 %
3	Both	11	0.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	7	0.5 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-107	Follow-up skipped because gate <> 1	1205	89.0 %
	Total	**1,354**	**100%**

Based upon 22 valid cases out of 1,354 total cases.

Location: 2846-2849 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM29_SUSYMB: S5 SubUse: Ever attended self-help group for drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	44	3.2 %
5	No	634	46.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-106	Skip-in for alchstat/drugstat	555	41.0 %
	Total	**1,354**	**100%**

Based upon 678 valid cases out of 1,354 total cases.

Location: 2850-2853 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM29_SUSYMC: S5 SubUse: Specify reason for self-help [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	11	0.8 %
2	Drugs	17	1.3 %
3	Both	16	1.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	6	0.4 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-107	Follow-up skipped because gate <> 1	1184	87.4 %
	Total	**1,354**	**100%**

Based upon 44 valid cases out of 1,354 total cases.

Location: 2854-2857 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM30_SUSYMB: S5 SubUse: Ever treated by therapist because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	15	1.1 %
5	No	663	49.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-106	Skip-in for alchstat/drugstat	555	41.0 %
	Total	**1,354**	**100%**

Based upon 678 valid cases out of 1,354 total cases.

Location: 2858-2861 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM30_SUSYMC: S5 SubUse: Specify reason for therapist visit [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	0	0.0 %
2	Drugs	8	0.6 %
3	Both	7	0.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %

Value	Label	Unweighted Frequency	%
-201	IntCompStat = 4 missed interview	120	8.9 %
-107	Follow-up skipped because gate <> 1	1211	89.4 %
	Total	**1,354**	**100%**

Based upon 15 valid cases out of 1,354 total cases.

Location: 2862-2865 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM31_SUSYMB: S5 SubUse: Ever Admitted to hospital because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	5	0.4 %
5	No	673	49.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-106	Skip-in for alchstat/drugstat	555	41.0 %
	Total	**1,354**	**100%**

Based upon 678 valid cases out of 1,354 total cases.

Location: 2866-2869 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE_SUBSYM31_SUSYMC: S5 SubUse: Specify reason admitted to hospital [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	1	0.1 %
2	Drugs	0	0.0 %
3	Both	4	0.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	8	0.6 %
-201	IntCompStat = 4 missed interview	120	8.9 %
-107	Follow-up skipped because gate <> 1	1221	90.2 %
	Total	**1,354**	**100%**

Based upon 5 valid cases out of 1,354 total cases.

Location: 2870-2873 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S5SUBUSE28: S5 SubUse: Num times abused prescribed medications (own or someone elses) during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	0	0.0 %
2	1-2 times	1	0.1 %
3	3-5 times	0	0.0 %
4	1X per month	0	0.0 %
5	2-3X per month	0	0.0 %
6	1X per week	1	0.1 %
7	2-3X per week	0	0.0 %
8	4-5X per week	0	0.0 %
9	Every day	1	0.1 %
	Missing Data		
-800	Question not asked for this subject	1230	90.8 %
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
	Total	**1,354**	**100%**

Based upon 3 valid cases out of 1,354 total cases.

Location: 2874-2877 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S5QFBRMO: S5 SubUse: Quant/freq used beer/wine/liquor - recall period

Based upon 1,231 valid cases out of 1,354 total cases.

- Mean: 9.80
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 210.00
- Standard Deviation: 21.62

Location: 2878-2881 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -300 , -202 , -201 , -100 , -9 , -8

S56MOUSE: S5 SubUse: Number drugs used - recall period

Based upon 1,233 valid cases out of 1,354 total cases.

- Mean: 0.55
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 9.00
- Standard Deviation: 1.01

Location: 2882-2885 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -300 , -202 , -201 , -100 , -9 , -8

S5QFSMK6: S5 SubUse: Quant/freq used cigarettes - recall period

Based upon 1,233 valid cases out of 1,354 total cases.

- Mean: 32.86
- Median: 5.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 160.00
- Standard Deviation: 47.10

Location: 2886-2891 (width: 6; decimal: 1)
Variable Type: numeric
(Range of) Missing Values: -300.0 , -202.0 , -201.0 , -100.0 , -9.0 , -8.0

S5A6SC2: S5 SubUse: Recall period alcohol consequences

Based upon 1,233 valid cases out of 1,354 total cases.

- Mean: 0.46
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 12.00
- Standard Deviation: 1.46

Location: 2892-2895 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S5D6SC2: S5 SubUse: Recall period drug consequences

Based upon 1,233 valid cases out of 1,354 total cases.

- Mean: 0.57
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 14.00
- Standard Deviation: 1.74

Location: 2896-2899 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S5A6DP2: S5 SubUse: Recall period alcohol dependence symptoms

Based upon 1,233 valid cases out of 1,354 total cases.

- Mean: 0.21
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00

- Maximum: 10.00
- Standard Deviation: 0.93

Location: 2900-2903 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S5D6DP2: S5 SubUse: Recall period drug dependence symptoms

Based upon 1,233 valid cases out of 1,354 total cases.

- Mean: 0.46
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 10.00
- Standard Deviation: 1.58

Location: 2904-2907 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S5TOTSC6: S5 SubUse: Recall period total substance use consequences

Based upon 1,233 valid cases out of 1,354 total cases.

- Mean: 0.83
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 14.00
- Standard Deviation: 2.08

Location: 2908-2911 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S5TOTDS6: S5 SubUse: Recall period total substance use dependency symp

Based upon 1,233 valid cases out of 1,354 total cases.

- Mean: 0.56
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 10.00
- Standard Deviation: 1.68

Location: 2912-2915 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S5ALTRTM: S5 SubUse: Recall period alcohol treatment

Value	Label	Unweighted Frequency	%
0	No	1202	88.8 %

Value	Label	Unweighted Frequency	%
1	Yes	31	2.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
	Total	**1,354**	**100%**

Based upon 1,233 valid cases out of 1,354 total cases.

Location: 2916-2919 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S5DGTRTM: S5 SubUse: Recall period drug treatment

Value	Label	Unweighted Frequency	%
0	No	1198	88.5 %
1	Yes	35	2.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
	Total	**1,354**	**100%**

Based upon 1,233 valid cases out of 1,354 total cases.

Location: 2920-2923 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S5TRTM: S5 SubUse: Recall period alcohol or drug treatment

Value	Label	Unweighted Frequency	%
0	No	1186	87.6 %
1	Yes	47	3.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	120	8.9 %
	Total	**1,354**	**100%**

Based upon 1,233 valid cases out of 1,354 total cases.

Location: 2924-2927 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S6SUBUSE1: S6 SubUse: How often had alcohol to drink in recall period

Value	Label	Unweighted Frequency	%
1	Not at all	629	46.5 %
2	1-2 times	131	9.7 %
3	3-5 times	86	6.4 %
4	1X per month	56	4.1 %
5	2-3X per month	96	7.1 %
6	1X per week	91	6.7 %
7	2-3X per week	99	7.3 %
8	4-5X per week	26	1.9 %
9	Every day	17	1.3 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,231 valid cases out of 1,354 total cases.

Location: 2928-2931 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S6SUBUSE2: S6 SubUse: Num alcoholic drinks usually consumed

Based upon 1,228 valid cases out of 1,354 total cases.

- Mean: 2.68
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 65.00
- Standard Deviation: 4.87

Location: 2932-2935 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -101 , -99 , -98

S6SUBUSE3: S6 SubUse: How often had 5+ drinks at one time in recall period

Value	Label	Unweighted Frequency	%
1	Not at all	858	63.4 %
2	1-2 times	103	7.6 %
3	3-5 times	37	2.7 %
4	1X per month	39	2.9 %
5	2-3X per month	52	3.8 %
6	1X per week	57	4.2 %
7	2-3X per week	53	3.9 %
8	4-5X per week	18	1.3 %

Value	Label	Unweighted Frequency	%
9	Every day	13	1.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-99	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,230 valid cases out of 1,354 total cases.

Location: 2936-2939 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S6SUBUSE4: S6 SubUse: Num times been drunk on alcohol during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	825	60.9 %
2	1-2 times	147	10.9 %
3	3-5 times	66	4.9 %
4	1X per month	43	3.2 %
5	2-3X per month	44	3.2 %
6	1X per week	51	3.8 %
7	2-3X per week	40	3.0 %
8	4-5X per week	9	0.7 %
9	Every day	6	0.4 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,231 valid cases out of 1,354 total cases.

Location: 2940-2943 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S6SUBUSE6: S6 SubUse: Num times used marijuana/hashish during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	794	58.6 %
2	1-2 times	76	5.6 %
3	3-5 times	53	3.9 %
4	1X per month	22	1.6 %
5	2-3X per month	41	3.0 %

Value	Label	Unweighted Frequency	%
6	1X per week	22	1.6 %
7	2-3X per week	56	4.1 %
8	4-5X per week	31	2.3 %
9	Every day	135	10.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-99	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,230 valid cases out of 1,354 total cases.

Location: 2944-2947 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S6SUBUSE8: S6 SubUse: Num times used sedative/tranquilizers during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1179	87.1 %
2	1-2 times	18	1.3 %
3	3-5 times	6	0.4 %
4	1X per month	9	0.7 %
5	2-3X per month	6	0.4 %
6	1X per week	5	0.4 %
7	2-3X per week	6	0.4 %
8	4-5X per week	1	0.1 %
9	Every day	1	0.1 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,231 valid cases out of 1,354 total cases.

Location: 2948-2951 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S6SUBUSE10: S6 SubUse: Num times used stimulants/amphetamines during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1167	86.2 %
2	1-2 times	5	0.4 %

Value	Label	Unweighted Frequency	%
3	3-5 times	11	0.8 %
4	1X per month	4	0.3 %
5	2-3X per month	5	0.4 %
6	1X per week	8	0.6 %
7	2-3X per week	10	0.7 %
8	4-5X per week	5	0.4 %
9	Every day	16	1.2 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,231 valid cases out of 1,354 total cases.

Location: 2952-2955 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S6SUBUSE12: S6 SubUse: Num times used cocaine during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1182	87.3 %
2	1-2 times	16	1.2 %
3	3-5 times	15	1.1 %
4	1X per month	4	0.3 %
5	2-3X per month	6	0.4 %
6	1X per week	3	0.2 %
7	2-3X per week	2	0.1 %
8	4-5X per week	2	0.1 %
9	Every day	1	0.1 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,231 valid cases out of 1,354 total cases.

Location: 2956-2959 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S6SUBUSE14: S6 SubUse: Num times used opiates during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1205	89.0 %
2	1-2 times	12	0.9 %
3	3-5 times	3	0.2 %
4	1X per month	1	0.1 %
5	2-3X per month	3	0.2 %
6	1X per week	3	0.2 %
7	2-3X per week	1	0.1 %
8	4-5X per week	0	0.0 %
9	Every day	2	0.1 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-99	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,230 valid cases out of 1,354 total cases.

Location: 2960-2963 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S6SUBUSE16: S6 SubUse: Num times used ecstasy during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1211	89.4 %
2	1-2 times	12	0.9 %
3	3-5 times	1	0.1 %
4	1X per month	1	0.1 %
5	2-3X per month	3	0.2 %
6	1X per week	0	0.0 %
7	2-3X per week	0	0.0 %
8	4-5X per week	2	0.1 %
9	Every day	1	0.1 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,231 valid cases out of 1,354 total cases.

Location: 2964-2967 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S6SUBUSE18: S6 SubUse: Num times used hallucinogens during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1196	88.3 %
2	1-2 times	19	1.4 %
3	3-5 times	6	0.4 %
4	1X per month	0	0.0 %
5	2-3X per month	3	0.2 %
6	1X per week	1	0.1 %
7	2-3X per week	2	0.1 %
8	4-5X per week	1	0.1 %
9	Every day	3	0.2 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,231 valid cases out of 1,354 total cases.

Location: 2968-2971 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S6SUBUSE20: S6 SubUse: Num times used inhalants during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1224	90.4 %
2	1-2 times	1	0.1 %
3	3-5 times	1	0.1 %
4	1X per month	1	0.1 %
5	2-3X per month	2	0.1 %
6	1X per week	0	0.0 %
7	2-3X per week	0	0.0 %
8	4-5X per week	0	0.0 %
9	Every day	2	0.1 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,231 valid cases out of 1,354 total cases.

Location: 2972-2975 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S6SUBUSE22: S6 SubUse: Num times used amyl nitrate/odorizers/rush during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1226	90.5 %
2	1-2 times	2	0.1 %
3	3-5 times	0	0.0 %
4	1X per month	0	0.0 %
5	2-3X per month	1	0.1 %
6	1X per week	0	0.0 %
7	2-3X per week	0	0.0 %
8	4-5X per week	0	0.0 %
9	Every day	2	0.1 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,231 valid cases out of 1,354 total cases.

Location: 2976-2979 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S6SUBUSE24: S6 SubUse: Used other drugs during recall period [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	1	0.1 %
5	No	1230	90.8 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,231 valid cases out of 1,354 total cases.

Location: 2980-2983 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE26: S6 SubUse: Num times used other drugs during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1230	90.8 %
2	1-2 times	0	0.0 %
3	3-5 times	0	0.0 %

Value	Label	Unweighted Frequency	%
4	1X per month	0	0.0 %
5	2-3X per month	1	0.1 %
6	1X per week	0	0.0 %
7	2-3X per week	0	0.0 %
8	4-5X per week	0	0.0 %
9	Every day	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,231 valid cases out of 1,354 total cases.

Location: 2984-2987 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S6SUBUSEDY: S6 SubUse: Used any drugs past 24hrs [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	129	9.5 %
5	No	1102	81.4 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,231 valid cases out of 1,354 total cases.

Location: 2988-2991 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_DRUG24MAR: S6 SubUse_Drug24: Used marijuana/hashish past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	120	8.9 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-104	Skip-in for SUBUSEDY 1	9	0.7 %
-103	Skip-in for SUBUSEDY <> 1	1102	81.4 %
	Total	**1,354**	**100%**

Based upon 120 valid cases out of 1,354 total cases.

Location: 2992-2995 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_DRUG24TRANQ: S6 SubUse_Drug24: Used sedatives/tranquilizers past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	1	0.1 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-104	Skip-in for SUBUSEDY 1	128	9.5 %
-103	Skip-in for SUBUSEDY <> 1	1102	81.4 %
	Total	**1,354**	**100%**

Based upon 1 valid cases out of 1,354 total cases.

Location: 2996-2999 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_DRUG24AMPH: S6 SubUse_Drug24: Used stimulants/amphetamines past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	7	0.5 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-104	Skip-in for SUBUSEDY 1	122	9.0 %
-103	Skip-in for SUBUSEDY <> 1	1102	81.4 %
	Total	**1,354**	**100%**

Based upon 7 valid cases out of 1,354 total cases.

Location: 3000-3003 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_DRUG24COC: S6 SubUse_Drug24: Used Cocaine past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	1	0.1 %
5	No	0	0.0 %

Value	Label	Unweighted Frequency	%
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-104	Skip-in for SUBUSEDY 1	128	9.5 %
-103	Skip-in for SUBUSEDY <> 1	1102	81.4 %
	Total	**1,354**	**100%**

Based upon 1 valid cases out of 1,354 total cases.

Location: 3004-3007 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_DRUG24OPI: S6 SubUse_Drug24: Used Opiates past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	1	0.1 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-104	Skip-in for SUBUSEDY 1	128	9.5 %
-103	Skip-in for SUBUSEDY <> 1	1102	81.4 %
	Total	**1,354**	**100%**

Based upon 1 valid cases out of 1,354 total cases.

Location: 3008-3011 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_DRUG24ECST: S6 SubUse_Drug24: Used Ecstacy past 24 hours

Value	Label	Unweighted Frequency	%
1	Yes	1	0.1 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-104	Skip-in for SUBUSEDY 1	128	9.5 %
-103	Skip-in for SUBUSEDY <> 1	1102	81.4 %
	Total	**1,354**	**100%**

Based upon 1 valid cases out of 1,354 total cases.

Location: 3012-3015 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_DRUG24HALL: S6 SubUse_Drug24: Used Hallucinogens past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	1	0.1 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-104	Skip-in for SUBUSEDY 1	128	9.5 %
-103	Skip-in for SUBUSEDY <> 1	1102	81.4 %
	Total	**1,354**	**100%**

Based upon 1 valid cases out of 1,354 total cases.

Location: 3016-3019 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_DRUG24INH: S6 SubUse_Drug24: Used Inhalants past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-104	Skip-in for SUBUSEDY 1	129	9.5 %
-103	Skip-in for SUBUSEDY <> 1	1102	81.4 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 3020-3023 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_DRUG24AMYL: S6 SubUse_Drug24: Used Amyl nitrate/poppers/odorizers 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		

Value	Label	Unweighted Frequency	%
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-104	Skip-in for SUBUSEDY 1	129	9.5 %
-103	Skip-in for SUBUSEDY <> 1	1102	81.4 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 3024-3027 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_DRUG24OTHER: S6 SubUse_Drug24: Used Other (specify) past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-104	Skip-in for SUBUSEDY 1	129	9.5 %
-103	Skip-in for SUBUSEDY <> 1	1102	81.4 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 3028-3031 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_DRUG24NONE: S6 SubUse_Drug24: Used NA (No drugs) past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-104	Skip-in for SUBUSEDY 1	129	9.5 %
-103	Skip-in for SUBUSEDY <> 1	1102	81.4 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 3032-3035 (width: 4; decimal: 0)
Variable Type: numeric

(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S6SUBUSE29: S6 SubUse: Specify other drug used past 24hrs

Based upon 1,354 valid cases out of 1,354 total cases.

Location: 3036-3036 (width: 1; decimal: 0)
Variable Type: character

S6SUBUSE30: S6 SubUse: Num times smoked cigarettes during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	567	41.9 %
2	1-2 times	24	1.8 %
3	3-5 times	17	1.3 %
4	1X per month	9	0.7 %
5	2-3X per month	14	1.0 %
6	1X per week	19	1.4 %
7	2-3X per week	50	3.7 %
8	4-5X per week	22	1.6 %
9	Every day	509	37.6 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,231 valid cases out of 1,354 total cases.

Location: 3037-3040 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S6SUBUSE31: S6 SubUse: Num cigarettes smoke per day during recall period

Value	Label	Unweighted Frequency	%
1	0 cigarettes	590	43.6 %
2	1-4 cigarettes	245	18.1 %
3	5-9 cigarettes	155	11.4 %
4	10-15 cigarettes	126	9.3 %
5	16-20 cigarettes	88	6.5 %
6	More than a pack a day cigarettes	26	1.9 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,230 valid cases out of 1,354 total cases.

Location: 3041-3044 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -105 , -9 , -8

S6SUBUSE_SUBSYM01_SUSYMB: S6 SubUse: Family ever complain because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	142	10.5 %
5	No	570	42.1 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-106	Skip-in for alchstat/drugstat	518	38.3 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 712 valid cases out of 1,354 total cases.

Location: 3045-3048 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM01_SUSYMC: S6 SubUse: Specify cause of family complaint [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	36	2.7 %
2	Drugs	67	4.9 %
3	Both	39	2.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-107	Follow-up skipped because gate <> 1	1086	80.2 %
	Total	**1,354**	**100%**

Based upon 142 valid cases out of 1,354 total cases.

Location: 3049-3052 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM02_SUSYMB: S6 SubUse: Friend ever complain because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	49	3.6 %
5	No	664	49.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-106	Skip-in for alchstat/drugstat	518	38.3 %
	Total	**1,354**	**100%**

Based upon 713 valid cases out of 1,354 total cases.

Location: 3053-3056 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM02_SUSYMC: S6 SubUse: Specify cause of friend complain [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	13	1.0 %
2	Drugs	26	1.9 %
3	Both	10	0.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-107	Follow-up skipped because gate <> 1	1179	87.1 %
	Total	**1,354**	**100%**

Based upon 49 valid cases out of 1,354 total cases.

Location: 3057-3060 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM03_SUSYMB: S6 SubUse: Ever trouble at work because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	23	1.7 %
5	No	690	51.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-106	Skip-in for alchstat/drugstat	518	38.3 %

Value	Label	Unweighted Frequency	%
	Total	**1,354**	**100%**

Based upon 713 valid cases out of 1,354 total cases.

Location: 3061-3064 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM03_SUSYMC: S6 SubUse: Specify cause of trouble at work [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	5	0.4 %
2	Drugs	15	1.1 %
3	Both	3	0.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-107	Follow-up skipped because gate <> 1	1205	89.0 %
	Total	**1,354**	**100%**

Based upon 23 valid cases out of 1,354 total cases.

Location: 3065-3068 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM04_SUSYMB: S6 SubUse: Ever missed work/school because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	43	3.2 %
5	No	670	49.5 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-106	Skip-in for alchstat/drugstat	518	38.3 %
	Total	**1,354**	**100%**

Based upon 713 valid cases out of 1,354 total cases.

Location: 3069-3072 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM04_SUSYMC: S6 SubUse: Specify cause of missed work/school [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	19	1.4 %
2	Drugs	16	1.2 %
3	Both	8	0.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-107	Follow-up skipped because gate <> 1	1185	87.5 %
	Total	**1,354**	**100%**

Based upon 43 valid cases out of 1,354 total cases.

Location: 3073-3076 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM05_SUSYMB: S6 SubUse: Ever arrested because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	69	5.1 %
5	No	644	47.6 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-106	Skip-in for alchstat/drugstat	518	38.3 %
	Total	**1,354**	**100%**

Based upon 713 valid cases out of 1,354 total cases.

Location: 3077-3080 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM05_SUSYMC: S6 SubUse: Specify cause of arrest [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	23	1.7 %
2	Drugs	37	2.7 %
3	Both	9	0.7 %
	Missing Data		

Value	Label	Unweighted Frequency	%
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-107	Follow-up skipped because gate <> 1	1159	85.6 %
	Total	**1,354**	**100%**

Based upon 69 valid cases out of 1,354 total cases.

Location: 3081-3084 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM06_SUSYMB: S6 SubUse: Ever argue family/friends because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	90	6.6 %
5	No	623	46.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-106	Skip-in for alchstat/drugstat	518	38.3 %
	Total	**1,354**	**100%**

Based upon 713 valid cases out of 1,354 total cases.

Location: 3085-3088 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM06_SUSYMC: S6 SubUse: Specify cause of argument w/ family/friends [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	26	1.9 %
2	Drugs	43	3.2 %
3	Both	21	1.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-107	Follow-up skipped because gate <> 1	1138	84.0 %
	Total	**1,354**	**100%**

Based upon 90 valid cases out of 1,354 total cases.

Location: 3089-3092 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM07_SUSYMB: S6 SubUse: Ever cont. drug/alcohol use in spite cause health problems in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	87	6.4 %
5	No	626	46.2 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-106	Skip-in for alchstat/drugstat	518	38.3 %
	Total	**1,354**	**100%**

Based upon 713 valid cases out of 1,354 total cases.

Location: 3093-3096 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM07_SUSYMC: S6 SubUse: Specify continued using [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	25	1.8 %
2	Drugs	40	3.0 %
3	Both	22	1.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-107	Follow-up skipped because gate <> 1	1141	84.3 %
	Total	**1,354**	**100%**

Based upon 87 valid cases out of 1,354 total cases.

Location: 3097-3100 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM08_SUSYMB: S6 SubUse: Ever not remember prior night because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	126	9.3 %

Value	Label	Unweighted Frequency	%
5	No	587	43.4 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-106	Skip-in for alchstat/drugstat	518	38.3 %
	Total	**1,354**	**100%**

Based upon 713 valid cases out of 1,354 total cases.

Location: 3101-3104 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM08_SUSYMC: S6 SubUse: Specify cause of not remember night [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	73	5.4 %
2	Drugs	19	1.4 %
3	Both	34	2.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-107	Follow-up skipped because gate <> 1	1102	81.4 %
	Total	**1,354**	**100%**

Based upon 126 valid cases out of 1,354 total cases.

Location: 3105-3108 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM09_SUSYMB: S6 SubUse: Ever used in situations could get hurt in recall period [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	127	9.4 %
5	No	586	43.3 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-106	Skip-in for alchstat/drugstat	518	38.3 %
	Total	**1,354**	**100%**

Based upon 713 valid cases out of 1,354 total cases.

Location: 3109-3112 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM09_SUSYMC: S6 SubUse: Specify used drugs in hurtful situations [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	52	3.8 %
2	Drugs	36	2.7 %
3	Both	39	2.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-107	Follow-up skipped because gate <> 1	1101	81.3 %
	Total	**1,354**	**100%**

Based upon 127 valid cases out of 1,354 total cases.

Location: 3113-3116 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM10_SUSYMB: S6 SubUse: Ever had accident/injury because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	42	3.1 %
5	No	671	49.6 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-106	Skip-in for alchstat/drugstat	518	38.3 %
	Total	**1,354**	**100%**

Based upon 713 valid cases out of 1,354 total cases.

Location: 3117-3120 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM10_SUSYMC: S6 SubUse: Specify cause of accident/injury [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	20	1.5 %
2	Drugs	14	1.0 %
3	Both	8	0.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-107	Follow-up skipped because gate <> 1	1186	87.6 %
	Total	**1,354**	**100%**

Based upon 42 valid cases out of 1,354 total cases.

Location: 3121-3124 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM11_SUSYMB: S6 SubUse: Alcohol/drug use interfere work/responsibility school/home/job in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	56	4.1 %
5	No	657	48.5 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-106	Skip-in for alchstat/drugstat	518	38.3 %
	Total	**1,354**	**100%**

Based upon 713 valid cases out of 1,354 total cases.

Location: 3125-3128 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM11_SUSYMC: S6 SubUse: Specify cause of interference [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	18	1.3 %
2	Drugs	26	1.9 %
3	Both	12	0.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	122	9.0 %

Value	Label	Unweighted Frequency	%
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-107	Follow-up skipped because gate <> 1	1172	86.6 %
	Total	**1,354**	**100%**

Based upon 56 valid cases out of 1,354 total cases.

Location: 3129-3132 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM12_SUSYMB: S6 SubUse: Ever trouble police no arrest because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	41	3.0 %
5	No	672	49.6 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-106	Skip-in for alchstat/drugstat	518	38.3 %
	Total	**1,354**	**100%**

Based upon 713 valid cases out of 1,354 total cases.

Location: 3133-3136 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM12_SUSYMC: S6 SubUse: Specify cause trouble police no arrest [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	15	1.1 %
2	Drugs	22	1.6 %
3	Both	4	0.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-107	Follow-up skipped because gate <> 1	1187	87.7 %
	Total	**1,354**	**100%**

Based upon 41 valid cases out of 1,354 total cases.

Location: 3137-3140 (width: 4; decimal: 0)
Variable Type: numeric

(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM13_SUSYMB: S6 SubUse: Sold/exchanged sex for drugs in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	4	0.3 %
5	No	709	52.4 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-106	Skip-in for alchstat/drugstat	518	38.3 %
	Total	**1,354**	**100%**

Based upon 713 valid cases out of 1,354 total cases.

Location: 3141-3144 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM13_SUSYMC: S6 SubUse: Specify cause sold/exchanged sex [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	1	0.1 %
2	Drugs	3	0.2 %
3	Both	0	0.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-107	Follow-up skipped because gate <> 1	1224	90.4 %
	Total	**1,354**	**100%**

Based upon 4 valid cases out of 1,354 total cases.

Location: 3145-3148 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM14_SUSYMB: S6 SubUse: Sold drugs for money buy drugs in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	56	4.1 %
5	No	657	48.5 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %

Value	Label	Unweighted Frequency	%
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-106	Skip-in for alchstat/drugstat	518	38.3 %
	Total	**1,354**	**100%**

Based upon 713 valid cases out of 1,354 total cases.

Location: 3149-3152 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM14_SUSYMC: S6 SubUse: Specify cause sold drugs for money to buy [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	3	0.2 %
2	Drugs	43	3.2 %
3	Both	10	0.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-107	Follow-up skipped because gate <> 1	1172	86.6 %
	Total	**1,354**	**100%**

Based upon 56 valid cases out of 1,354 total cases.

Location: 3153-3156 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM15_SUSYMB: S6 SubUse: Stole for money buy drugs in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	19	1.4 %
5	No	694	51.3 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-106	Skip-in for alchstat/drugstat	518	38.3 %
	Total	**1,354**	**100%**

Based upon 713 valid cases out of 1,354 total cases.

Location: 3157-3160 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM15_SUSYMC: S6 SubUse: Specify cause stole for money to buy [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	1	0.1 %
2	Drugs	14	1.0 %
3	Both	4	0.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-107	Follow-up skipped because gate <> 1	1209	89.3 %
	Total	**1,354**	**100%**

Based upon 19 valid cases out of 1,354 total cases.

Location: 3161-3164 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM16_SUSYMB: S6 SubUse: Ever had physical fight because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	43	3.2 %
5	No	669	49.4 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-106	Skip-in for alchstat/drugstat	518	38.3 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 712 valid cases out of 1,354 total cases.

Location: 3165-3168 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM16_SUSYMC: S6 SubUse: Specify cause of fight [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	20	1.5 %
2	Drugs	14	1.0 %
3	Both	9	0.7 %

Value	Label	Unweighted Frequency	%
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-107	Follow-up skipped because gate <> 1	1185	87.5 %
	Total	**1,354**	**100%**

Based upon 43 valid cases out of 1,354 total cases.

Location: 3169-3172 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM17_SUSYMB: S6 SubUse: Ever destroy property because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	31	2.3 %
5	No	682	50.4 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-106	Skip-in for alchstat/drugstat	518	38.3 %
	Total	**1,354**	**100%**

Based upon 713 valid cases out of 1,354 total cases.

Location: 3173-3176 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM17_SUSYMC: S6 SubUse: Specify cause of destroy property [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	17	1.3 %
2	Drugs	9	0.7 %
3	Both	5	0.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-107	Follow-up skipped because gate <> 1	1197	88.4 %
	Total	**1,354**	**100%**

Based upon 31 valid cases out of 1,354 total cases.

Location: 3177-3180 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM18_SUSYMB: S6 SubUse: Ever felt strong urge use drug/alcohol no self stop in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	44	3.2 %
5	No	669	49.4 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-106	Skip-in for alchstat/drugstat	518	38.3 %
	Total	**1,354**	**100%**

Based upon 713 valid cases out of 1,354 total cases.

Location: 3181-3184 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM18_SUSYMC: S6 SubUse: Specify the cause of strong urge [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	13	1.0 %
2	Drugs	24	1.8 %
3	Both	7	0.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-107	Follow-up skipped because gate <> 1	1184	87.4 %
	Total	**1,354**	**100%**

Based upon 44 valid cases out of 1,354 total cases.

Location: 3185-3188 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM19_SUSYMB: S6 SubUse: Ever want drug/alcohol think of nothing else in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	38	2.8 %
5	No	675	49.9 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-106	Skip-in for alchstat/drugstat	518	38.3 %
	Total	**1,354**	**100%**

Based upon 713 valid cases out of 1,354 total cases.

Location: 3189-3192 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM19_SUSYMC: S6 SubUse: Specify cause of want [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	10	0.7 %
2	Drugs	20	1.5 %
3	Both	8	0.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-107	Follow-up skipped because gate <> 1	1190	87.9 %
	Total	**1,354**	**100%**

Based upon 38 valid cases out of 1,354 total cases.

Location: 3193-3196 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM20_SUSYMB: S6 SubUse: Ever experience withdrawal from drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	37	2.7 %
5	No	676	49.9 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-106	Skip-in for alchstat/drugstat	518	38.3 %
	Total	**1,354**	**100%**

Based upon 713 valid cases out of 1,354 total cases.

Location: 3197-3200 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM20_SUSYMC: S6 SubUse: Specify cause of withdraw [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	6	0.4 %
2	Drugs	24	1.8 %
3	Both	7	0.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-107	Follow-up skipped because gate <> 1	1191	88.0 %
	Total	**1,354**	**100%**

Based upon 37 valid cases out of 1,354 total cases.

Location: 3201-3204 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM21_SUSYMB: S6 SubUse: Take drug/alcohol to keep from having withdraw in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	18	1.3 %
5	No	19	1.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-108	Skip-in for SUBSYM20=5	673	49.7 %
-106	Skip-in for alchstat/drugstat	520	38.4 %
	Total	**1,354**	**100%**

Based upon 37 valid cases out of 1,354 total cases.

Location: 3205-3208 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM21_SUSYMC: S6 SubUse: Specify reason for consumption [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	3	0.2 %
2	Drugs	13	1.0 %
3	Both	2	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-107	Follow-up skipped because gate <> 1	1210	89.4 %
	Total	**1,354**	**100%**

Based upon 18 valid cases out of 1,354 total cases.

Location: 3209-3212 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM22_SUSYMB: S6 SubUse: Give up usual activities/interests to use drug/alcohol in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	51	3.8 %
5	No	661	48.8 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-106	Skip-in for alchstat/drugstat	518	38.3 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 712 valid cases out of 1,354 total cases.

Location: 3213-3216 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM22_SUSYMC: S6 SubUse: Specify cause of giving up activities/interests [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	9	0.7 %
2	Drugs	30	2.2 %
3	Both	12	0.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %

Value	Label	Unweighted Frequency	%
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-107	Follow-up skipped because gate <> 1	1177	86.9 %
	Total	**1,354**	**100%**

Based upon 51 valid cases out of 1,354 total cases.

Location: 3217-3220 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM23_SUSYMB: S6 SubUse: Ever used in rp when promised wouldn't [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	119	8.8 %
5	No	594	43.9 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-106	Skip-in for alchstat/drugstat	518	38.3 %
	Total	**1,354**	**100%**

Based upon 713 valid cases out of 1,354 total cases.

Location: 3221-3224 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM23_SUSYMC: S6 SubUse: Specify what use you started [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	34	2.5 %
2	Drugs	57	4.2 %
3	Both	28	2.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-107	Follow-up skipped because gate <> 1	1109	81.9 %
	Total	**1,354**	**100%**

Based upon 119 valid cases out of 1,354 total cases.

Location: 3225-3228 (width: 4; decimal: 0)
Variable Type: numeric

(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM24_SUSYMB: S6 SubUse: Ever use drug/alcohol more freq than intended in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	91	6.7 %
5	No	622	45.9 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-106	Skip-in for alchstat/drugstat	518	38.3 %
	Total	**1,354**	**100%**

Based upon 713 valid cases out of 1,354 total cases.

Location: 3229-3232 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM24_SUSYMC: S6 SubUse: Specify what you used more frequently [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	25	1.8 %
2	Drugs	53	3.9 %
3	Both	13	1.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-107	Follow-up skipped because gate <> 1	1137	84.0 %
	Total	**1,354**	**100%**

Based upon 91 valid cases out of 1,354 total cases.

Location: 3233-3236 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM25_SUSYMB: S6 SubUse: Ever tried cut down drug/alcohol use but couldnt in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	73	5.4 %
5	No	640	47.3 %

Value	Label	Unweighted Frequency	%
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-106	Skip-in for alchstat/drugstat	518	38.3 %
	Total	**1,354**	**100%**

Based upon 713 valid cases out of 1,354 total cases.

Location: 3237-3240 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM25_SUSYMC: S6 SubUse: Specify what substance use you tried to curtail [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	16	1.2 %
2	Drugs	48	3.5 %
3	Both	9	0.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-107	Follow-up skipped because gate <> 1	1155	85.3 %
	Total	**1,354**	**100%**

Based upon 73 valid cases out of 1,354 total cases.

Location: 3241-3244 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM26_SUSYMB: S6 SubUse: Ever need more drug/alcohol get same effect in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	117	8.6 %
5	No	596	44.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-106	Skip-in for alchstat/drugstat	518	38.3 %
	Total	**1,354**	**100%**

Based upon 713 valid cases out of 1,354 total cases.

Location: 3245-3248 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM26_SUSYMC: S6 SubUse: Specify substance need more for same effect [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	29	2.1 %
2	Drugs	61	4.5 %
3	Both	27	2.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-107	Follow-up skipped because gate <> 1	1111	82.1 %
	Total	**1,354**	**100%**

Based upon 117 valid cases out of 1,354 total cases.

Location: 3249-3252 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM27_SUSYMB: S6 SubUse: Ever used drug/alcohol enough to depend in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	48	3.5 %
5	No	665	49.1 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-106	Skip-in for alchstat/drugstat	518	38.3 %
	Total	**1,354**	**100%**

Based upon 713 valid cases out of 1,354 total cases.

Location: 3253-3256 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM27_SUSYMC: S6 SubUse: Specify dependency [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	6	0.4 %

Value	Label	Unweighted Frequency	%
2	Drugs	34	2.5 %
3	Both	8	0.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-107	Follow-up skipped because gate <> 1	1180	87.1 %
	Total	**1,354**	**100%**

Based upon 48 valid cases out of 1,354 total cases.

Location: 3257-3260 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM28_SUSYMB: S6 SubUse: Ever talked doctor/clergy/guidance counselor drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	24	1.8 %
5	No	689	50.9 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-106	Skip-in for alchstat/drugstat	518	38.3 %
	Total	**1,354**	**100%**

Based upon 713 valid cases out of 1,354 total cases.

Location: 3261-3264 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM28_SUSYMC: S6 SubUse: Specify talked doctor/clergy/guidance counselor [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	3	0.2 %
2	Drugs	15	1.1 %
3	Both	6	0.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %

Value	Label	Unweighted Frequency	%
-107	Follow-up skipped because gate <> 1	1204	88.9 %
	Total	**1,354**	**100%**

Based upon 24 valid cases out of 1,354 total cases.

Location: 3265-3268 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM29_SUSYMB: S6 SubUse: Ever attended self-help group for drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	37	2.7 %
5	No	675	49.9 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-106	Skip-in for alchstat/drugstat	518	38.3 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 712 valid cases out of 1,354 total cases.

Location: 3269-3272 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM29_SUSYMC: S6 SubUse: Specify reason for self-help [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	10	0.7 %
2	Drugs	12	0.9 %
3	Both	15	1.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-107	Follow-up skipped because gate <> 1	1191	88.0 %
	Total	**1,354**	**100%**

Based upon 37 valid cases out of 1,354 total cases.

Location: 3273-3276 (width: 4; decimal: 0)
Variable Type: numeric

(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM30_SUSYMB: S6 SubUse: Ever treated by therapist because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	20	1.5 %
5	No	693	51.2 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-106	Skip-in for alchstat/drugstat	518	38.3 %
	Total	**1,354**	**100%**

Based upon 713 valid cases out of 1,354 total cases.

Location: 3277-3280 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM30_SUSYMC: S6 SubUse: Specify reason for therapist visit [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	3	0.2 %
2	Drugs	10	0.7 %
3	Both	7	0.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-107	Follow-up skipped because gate <> 1	1208	89.2 %
	Total	**1,354**	**100%**

Based upon 20 valid cases out of 1,354 total cases.

Location: 3281-3284 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM31_SUSYMB: S6 SubUse: Ever Admitted to hospital because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	9	0.7 %
5	No	704	52.0 %

Value	Label	Unweighted Frequency	%
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-106	Skip-in for alchstat/drugstat	518	38.3 %
	Total	**1,354**	**100%**

Based upon 713 valid cases out of 1,354 total cases.

Location: 3285-3288 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE_SUBSYM31_SUSYMC: S6 SubUse: Specify reason admitted to hospital [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	4	0.3 %
2	Drugs	4	0.3 %
3	Both	1	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
-107	Follow-up skipped because gate <> 1	1219	90.0 %
	Total	**1,354**	**100%**

Based upon 9 valid cases out of 1,354 total cases.

Location: 3289-3292 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S6SUBUSE28: S6 SubUse: Num times abused prescribed medications (own or someone elses) during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	0	0.0 %
2	1-2 times	3	0.2 %
3	3-5 times	0	0.0 %
4	1X per month	0	0.0 %
5	2-3X per month	0	0.0 %
6	1X per week	0	0.0 %
7	2-3X per week	0	0.0 %
8	4-5X per week	0	0.0 %

Value	Label	Unweighted Frequency	%
9	Every day	1	0.1 %
	Missing Data		
-800	Question not asked for this subject	1227	90.6 %
-201	IntCompStat = 4 missed interview	122	9.0 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	1	0.1 %
	Total	**1,354**	**100%**

Based upon 4 valid cases out of 1,354 total cases.

Location: 3293-3296 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S6QFBRMO: S6 SubUse: Quant/freq used beer/wine/liquor - recall period

Based upon 1,228 valid cases out of 1,354 total cases.

- Mean: 11.40
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 400.00
- Standard Deviation: 25.59

Location: 3297-3300 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -300 , -202 , -201 , -100 , -9 , -8

S66MOUSE: S6 SubUse: Number drugs used - recall period

Based upon 1,231 valid cases out of 1,354 total cases.

- Mean: 0.56
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 9.00
- Standard Deviation: 0.99

Location: 3301-3304 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -300 , -202 , -201 , -100 , -9 , -8

S6QFSMK6: S6 SubUse: Quant/freq used cigarettes - recall period

Based upon 1,230 valid cases out of 1,354 total cases.

- Mean: 33.66
- Median: 7.50
- Mode: 0.00
- Minimum: 0.00
- Maximum: 160.00

- Standard Deviation: 47.81

Location: 3305-3310 (width: 6; decimal: 1)
Variable Type: numeric
(Range of) Missing Values: -300.0 , -202.0 , -201.0 , -100.0 , -9.0 , -8.0

S6A6SC2: S6 SubUse: Recall period alcohol consequences

Based upon 1,232 valid cases out of 1,354 total cases.

- Mean: 0.49
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 17.00
- Standard Deviation: 1.49

Location: 3311-3314 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S6D6SC2: S6 SubUse: Recall period drug consequences

Based upon 1,232 valid cases out of 1,354 total cases.

- Mean: 0.55
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 13.00
- Standard Deviation: 1.67

Location: 3315-3318 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S6A6DP2: S6 SubUse: Recall period alcohol dependence symptoms

Based upon 1,232 valid cases out of 1,354 total cases.

- Mean: 0.22
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 10.00
- Standard Deviation: 0.89

Location: 3319-3322 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S6D6DP2: S6 SubUse: Recall period drug dependence symptoms

Based upon 1,232 valid cases out of 1,354 total cases.

- Mean: 0.39
- Median: 0.00

- Mode: 0.00
- Minimum: 0.00
- Maximum: 10.00
- Standard Deviation: 1.30

Location: 3323-3326 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S6TOTSC6: S6 SubUse: Recall period total substance use consequences

Based upon 1,232 valid cases out of 1,354 total cases.

- Mean: 0.85
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 17.00
- Standard Deviation: 2.05

Location: 3327-3330 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S6TOTDS6: S6 SubUse: Recall period total substance use dependency symp

Based upon 1,232 valid cases out of 1,354 total cases.

- Mean: 0.52
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 10.00
- Standard Deviation: 1.46

Location: 3331-3334 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S6ALTRTM: S6 SubUse: Recall period alcohol treatment

Value	Label	Unweighted Frequency	%
0	No	1201	88.7 %
1	Yes	31	2.3 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
	Total	**1,354**	**100%**

Based upon 1,232 valid cases out of 1,354 total cases.

Location: 3335-3338 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S6DGTRTM: S6 SubUse: Recall period drug treatment

Value	Label	Unweighted Frequency	%
0	No	1195	88.3 %
1	Yes	37	2.7 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
	Total	**1,354**	**100%**

Based upon 1,232 valid cases out of 1,354 total cases.

Location: 3339-3342 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S6TRTM: S6 SubUse: Recall period alcohol or drug treatment

Value	Label	Unweighted Frequency	%
0	No	1183	87.4 %
1	Yes	49	3.6 %
	Missing Data		
-201	IntCompStat = 4 missed interview	122	9.0 %
	Total	**1,354**	**100%**

Based upon 1,232 valid cases out of 1,354 total cases.

Location: 3343-3346 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S7SUBUSE1: S7 SubUse: How often had alcohol to drink in recall period

Value	Label	Unweighted Frequency	%
1	Not at all	508	37.5 %
2	1-5 times	188	13.9 %
3	6-11 times	84	6.2 %
4	1X per month	61	4.5 %
5	2-3X per month	129	9.5 %
6	1X per week	83	6.1 %
7	2-3X per week	119	8.8 %
8	4-5X per week	21	1.6 %
9	Every day	17	1.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %

Value	Label	Unweighted Frequency	%
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
	Total	**1,354**	**100%**

Based upon 1,210 valid cases out of 1,354 total cases.

Location: 3347-3350 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S7SUBUSE2: S7 SubUse: Num alcoholic drinks usually consumed

Based upon 1,209 valid cases out of 1,354 total cases.

- Mean: 3.31
- Median: 2.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 67.00
- Standard Deviation: 5.07

Location: 3351-3354 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -101 , -99 , -98

S7SUBUSE3: S7 SubUse: How often had 5+ drinks at one time in recall period

Value	Label	Unweighted Frequency	%
1	Not at all	737	54.4 %
2	1-5 times	166	12.3 %
3	6-11 times	50	3.7 %
4	1X per month	35	2.6 %
5	2-3X per month	80	5.9 %
6	1X per week	53	3.9 %
7	2-3X per week	64	4.7 %
8	4-5X per week	14	1.0 %
9	Every day	11	0.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
	Total	**1,354**	**100%**

Based upon 1,210 valid cases out of 1,354 total cases.

Location: 3355-3358 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S7SUBUSE4: S7 SubUse: Num times been drunk on alcohol during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	702	51.8 %
2	1-5 times	244	18.0 %
3	6-11 times	50	3.7 %
4	1X per month	40	3.0 %
5	2-3X per month	63	4.7 %
6	1X per week	48	3.5 %
7	2-3X per week	44	3.2 %
8	4-5X per week	13	1.0 %
9	Every day	6	0.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
	Total	**1,354**	**100%**

Based upon 1,210 valid cases out of 1,354 total cases.

Location: 3359-3362 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S7SUBUSE6: S7 SubUse: Num times used marijuana/hashish during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	719	53.1 %
2	1-5 times	111	8.2 %
3	6-11 times	30	2.2 %
4	1X per month	28	2.1 %
5	2-3X per month	34	2.5 %
6	1X per week	25	1.8 %
7	2-3X per week	58	4.3 %
8	4-5X per week	46	3.4 %
9	Every day	159	11.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
	Total	**1,354**	**100%**

Based upon 1,210 valid cases out of 1,354 total cases.

Location: 3363-3366 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S7SUBUSE8: S7 SubUse: Num times used sedative/tranquilizers during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1144	84.5 %
2	1-5 times	33	2.4 %
3	6-11 times	5	0.4 %
4	1X per month	2	0.1 %
5	2-3X per month	5	0.4 %
6	1X per week	3	0.2 %
7	2-3X per week	11	0.8 %
8	4-5X per week	2	0.1 %
9	Every day	5	0.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
	Total	**1,354**	**100%**

Based upon 1,210 valid cases out of 1,354 total cases.

Location: 3367-3370 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S7SUBUSE10: S7 SubUse: Num times used stimulants/amphetamines during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1128	83.3 %
2	1-5 times	17	1.3 %
3	6-11 times	12	0.9 %
4	1X per month	7	0.5 %
5	2-3X per month	6	0.4 %
6	1X per week	6	0.4 %
7	2-3X per week	4	0.3 %
8	4-5X per week	7	0.5 %
9	Every day	22	1.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %

Value	Label	Unweighted Frequency	%
-99	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,209 valid cases out of 1,354 total cases.

Location: 3371-3374 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S7SUBUSE12: S7 SubUse: Num times used cocaine during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1148	84.8 %
2	1-5 times	37	2.7 %
3	6-11 times	4	0.3 %
4	1X per month	4	0.3 %
5	2-3X per month	5	0.4 %
6	1X per week	1	0.1 %
7	2-3X per week	7	0.5 %
8	4-5X per week	1	0.1 %
9	Every day	3	0.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
	Total	**1,354**	**100%**

Based upon 1,210 valid cases out of 1,354 total cases.

Location: 3375-3378 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S7SUBUSE14: S7 SubUse: Num times used opiates during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1172	86.6 %
2	1-5 times	16	1.2 %
3	6-11 times	4	0.3 %
4	1X per month	2	0.1 %
5	2-3X per month	5	0.4 %
6	1X per week	2	0.1 %
7	2-3X per week	4	0.3 %
8	4-5X per week	2	0.1 %

Value	Label	Unweighted Frequency	%
9	Every day	2	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-99	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,209 valid cases out of 1,354 total cases.

Location: 3379-3382 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S7SUBUSE16: S7 SubUse: Num times used ecstasy during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1167	86.2 %
2	1-5 times	31	2.3 %
3	6-11 times	1	0.1 %
4	1X per month	2	0.1 %
5	2-3X per month	2	0.1 %
6	1X per week	2	0.1 %
7	2-3X per week	3	0.2 %
8	4-5X per week	1	0.1 %
9	Every day	1	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
	Total	**1,354**	**100%**

Based upon 1,210 valid cases out of 1,354 total cases.

Location: 3383-3386 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S7SUBUSE18: S7 SubUse: Num times used hallucinogens during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1170	86.4 %
2	1-5 times	26	1.9 %
3	6-11 times	3	0.2 %

Value	Label	Unweighted Frequency	%
4	1X per month	1	0.1 %
5	2-3X per month	1	0.1 %
6	1X per week	1	0.1 %
7	2-3X per week	5	0.4 %
8	4-5X per week	0	0.0 %
9	Every day	3	0.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
	Total	**1,354**	**100%**

Based upon 1,210 valid cases out of 1,354 total cases.

Location: 3387-3390 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S7SUBUSE20: S7 SubUse: Num times used inhalants during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1204	88.9 %
2	1-5 times	2	0.1 %
3	6-11 times	1	0.1 %
4	1X per month	0	0.0 %
5	2-3X per month	1	0.1 %
6	1X per week	0	0.0 %
7	2-3X per week	1	0.1 %
8	4-5X per week	0	0.0 %
9	Every day	1	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
	Total	**1,354**	**100%**

Based upon 1,210 valid cases out of 1,354 total cases.

Location: 3391-3394 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S7SUBUSE22: S7 SubUse: Num times used amyl nitrate/odorizers/rush during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1206	89.1 %
2	1-5 times	1	0.1 %
3	6-11 times	0	0.0 %
4	1X per month	0	0.0 %
5	2-3X per month	2	0.1 %
6	1X per week	0	0.0 %
7	2-3X per week	0	0.0 %
8	4-5X per week	0	0.0 %
9	Every day	1	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
	Total	**1,354**	**100%**

Based upon 1,210 valid cases out of 1,354 total cases.

Location: 3395-3398 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S7SUBUSE24: S7 SubUse: Used other drugs during recall period [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	1	0.1 %
5	No	1209	89.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
	Total	**1,354**	**100%**

Based upon 1,210 valid cases out of 1,354 total cases.

Location: 3399-3402 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE26: S7 SubUse: Num times used other drugs during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1209	89.3 %
2	1-5 times	0	0.0 %
3	6-11 times	0	0.0 %

Value	Label	Unweighted Frequency	%
4	1X per month	0	0.0 %
5	2-3X per month	0	0.0 %
6	1X per week	0	0.0 %
7	2-3X per week	1	0.1 %
8	4-5X per week	0	0.0 %
9	Every day	0	0.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
	Total	**1,354**	**100%**

Based upon 1,210 valid cases out of 1,354 total cases.

Location: 3403-3406 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S7SUBUSEDY: S7 SubUse: Used any drugs past 24hrs [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	118	8.7 %
5	No	1092	80.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
	Total	**1,354**	**100%**

Based upon 1,210 valid cases out of 1,354 total cases.

Location: 3407-3410 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_DRUG24MAR: S7 SubUse_Drug24: Used marijuana/hashish past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	113	8.3 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-104	Skip-in for SUBUSEDY 1	6	0.4 %

Value	Label	Unweighted Frequency	%
-103	Skip-in for SUBUSEDY <> 1	1092	80.6 %
	Total	**1,354**	**100%**

Based upon 113 valid cases out of 1,354 total cases.

Location: 3411-3414 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_DRUG24TRANQ: S7 SubUse_Drug24: Used sedatives/tranquilizers past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-104	Skip-in for SUBUSEDY 1	119	8.8 %
-103	Skip-in for SUBUSEDY <> 1	1092	80.6 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 3415-3418 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_DRUG24AMPH: S7 SubUse_Drug24: Used stimulants/amphetamines past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	3	0.2 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-104	Skip-in for SUBUSEDY 1	116	8.6 %
-103	Skip-in for SUBUSEDY <> 1	1092	80.6 %
	Total	**1,354**	**100%**

Based upon 3 valid cases out of 1,354 total cases.

Location: 3419-3422 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_DRUG24COC: S7 SubUse_Drug24: Used Cocaine past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	2	0.1 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-104	Skip-in for SUBUSEDY 1	117	8.6 %
-103	Skip-in for SUBUSEDY <> 1	1092	80.6 %
	Total	**1,354**	**100%**

Based upon 2 valid cases out of 1,354 total cases.

Location: 3423-3426 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_DRUG24OPI: S7 SubUse_Drug24: Used Opiates past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	3	0.2 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-104	Skip-in for SUBUSEDY 1	116	8.6 %
-103	Skip-in for SUBUSEDY <> 1	1092	80.6 %
	Total	**1,354**	**100%**

Based upon 3 valid cases out of 1,354 total cases.

Location: 3427-3430 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_DRUG24ECST: S7 SubUse_Drug24: Used Ecstacy past 24 hours

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-104	Skip-in for SUBUSEDY 1	119	8.8 %
-103	Skip-in for SUBUSEDY <> 1	1092	80.6 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 3431-3434 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_DRUG24HALL: S7 SubUse_Drug24: Used Hallucinogens past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-104	Skip-in for SUBUSEDY 1	119	8.8 %
-103	Skip-in for SUBUSEDY <> 1	1092	80.6 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 3435-3438 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_DRUG24INH: S7 SubUse_Drug24: Used Inhalants past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-104	Skip-in for SUBUSEDY 1	119	8.8 %
-103	Skip-in for SUBUSEDY <> 1	1092	80.6 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 3439-3442 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_DRUG24AMYL: S7 SubUse_Drug24: Used Amyl nitrate/poppers/odorizers 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %

Value	Label	Unweighted Frequency	%
	Missing Data		
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-104	Skip-in for SUBUSEDY 1	119	8.8 %
-103	Skip-in for SUBUSEDY <> 1	1092	80.6 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 3443-3446 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_DRUG24OTHER: S7 SubUse_Drug24: Used Other (specify) past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-104	Skip-in for SUBUSEDY 1	119	8.8 %
-103	Skip-in for SUBUSEDY <> 1	1092	80.6 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 3447-3450 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_DRUG24NONE: S7 SubUse_Drug24: Used NA (No drugs) past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-104	Skip-in for SUBUSEDY 1	119	8.8 %
-103	Skip-in for SUBUSEDY <> 1	1092	80.6 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 3451-3454 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S7SUBUSE29: S7 SubUse: Specify other drug used past 24hrs

Based upon 1,354 valid cases out of 1,354 total cases.

Location: 3455-3455 (width: 1; decimal: 0)
Variable Type: character

S7SUBUSE30: S7 SubUse: Num times smoked cigarettes during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	502	37.1 %
2	1-5 times	26	1.9 %
3	6-11 times	15	1.1 %
4	1X per month	8	0.6 %
5	2-3X per month	18	1.3 %
6	1X per week	19	1.4 %
7	2-3X per week	43	3.2 %
8	4-5X per week	33	2.4 %
9	Every day	545	40.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-99	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,209 valid cases out of 1,354 total cases.

Location: 3456-3459 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S7SUBUSE31: S7 SubUse: Num cigarettes smoke per day during recall period

Value	Label	Unweighted Frequency	%
1	0 cigarettes	531	39.2 %
2	1-4 cigarettes	231	17.1 %
3	5-9 cigarettes	181	13.4 %
4	10-15 cigarettes	135	10.0 %
5	16-20 cigarettes	106	7.8 %
6	More than a pack a day cigarettes	25	1.8 %
	Missing Data		

Value	Label	Unweighted Frequency	%
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,209 valid cases out of 1,354 total cases.

Location: 3460-3463 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -105 , -9 , -8

S7SUBUSE_SUBSYM01_SUSYMB: S7 SubUse: Family ever complain because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	189	14.0 %
5	No	600	44.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-106	Skip-in for alchstat/drugstat	421	31.1 %
	Total	**1,354**	**100%**

Based upon 789 valid cases out of 1,354 total cases.

Location: 3464-3467 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM01_SUSYMC: S7 SubUse: Specify cause of family complaint [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	57	4.2 %
2	Drugs	84	6.2 %
3	Both	48	3.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-107	Follow-up skipped because gate <> 1	1021	75.4 %
	Total	**1,354**	**100%**

Based upon 189 valid cases out of 1,354 total cases.

Location: 3468-3471 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM02_SUSYMB: S7 SubUse: Friend ever complain because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	79	5.8 %
5	No	710	52.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-106	Skip-in for alchstat/drugstat	421	31.1 %
	Total	**1,354**	**100%**

Based upon 789 valid cases out of 1,354 total cases.

Location: 3472-3475 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM02_SUSYMC: S7 SubUse: Specify cause of friend complain [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	26	1.9 %
2	Drugs	36	2.7 %
3	Both	17	1.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-107	Follow-up skipped because gate <> 1	1131	83.5 %
	Total	**1,354**	**100%**

Based upon 79 valid cases out of 1,354 total cases.

Location: 3476-3479 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM03_SUSYMB: S7 SubUse: Ever trouble at work because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	33	2.4 %
5	No	756	55.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-106	Skip-in for alchstat/drugstat	421	31.1 %
	Total	**1,354**	**100%**

Based upon 789 valid cases out of 1,354 total cases.

Location: 3480-3483 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM03_SUSYMC: S7 SubUse: Specify cause of trouble at work [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	19	1.4 %
2	Drugs	9	0.7 %
3	Both	5	0.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-107	Follow-up skipped because gate <> 1	1177	86.9 %
	Total	**1,354**	**100%**

Based upon 33 valid cases out of 1,354 total cases.

Location: 3484-3487 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM04_SUSYMB: S7 SubUse: Ever missed work/school because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	51	3.8 %
5	No	738	54.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %

Value	Label	Unweighted Frequency	%
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-106	Skip-in for alchstat/drugstat	421	31.1 %
	Total	**1,354**	**100%**

Based upon 789 valid cases out of 1,354 total cases.

Location: 3488-3491 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM04_SUSYMC: S7 SubUse: Specify cause of missed work/school [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	29	2.1 %
2	Drugs	13	1.0 %
3	Both	9	0.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-107	Follow-up skipped because gate <> 1	1159	85.6 %
	Total	**1,354**	**100%**

Based upon 51 valid cases out of 1,354 total cases.

Location: 3492-3495 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM05_SUSYMB: S7 SubUse: Ever arrested because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	105	7.8 %
5	No	684	50.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-106	Skip-in for alchstat/drugstat	421	31.1 %
	Total	**1,354**	**100%**

Based upon 789 valid cases out of 1,354 total cases.

Location: 3496-3499 (width: 4; decimal: 0)

Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM05_SUSYMC: S7 SubUse: Specify cause of arrest [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	35	2.6 %
2	Drugs	59	4.4 %
3	Both	11	0.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-107	Follow-up skipped because gate <> 1	1105	81.6 %
	Total	**1,354**	**100%**

Based upon 105 valid cases out of 1,354 total cases.

Location: 3500-3503 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM06_SUSYMB: S7 SubUse: Ever argue family/friends because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	120	8.9 %
5	No	669	49.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-106	Skip-in for alchstat/drugstat	421	31.1 %
	Total	**1,354**	**100%**

Based upon 789 valid cases out of 1,354 total cases.

Location: 3504-3507 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM06_SUSYMC: S7 SubUse: Specify cause of argument w/ family/friends [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	38	2.8 %
2	Drugs	58	4.3 %

Value	Label	Unweighted Frequency	%
3	Both	24	1.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-107	Follow-up skipped because gate <> 1	1090	80.5 %
	Total	**1,354**	**100%**

Based upon 120 valid cases out of 1,354 total cases.

Location: 3508-3511 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM07_SUSYMB: S7 SubUse: Ever cont. drug/alcohol use in spite cause health problems in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	97	7.2 %
5	No	692	51.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-106	Skip-in for alchstat/drugstat	421	31.1 %
	Total	**1,354**	**100%**

Based upon 789 valid cases out of 1,354 total cases.

Location: 3512-3515 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM07_SUSYMC: S7 SubUse: Specify continued using [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	25	1.8 %
2	Drugs	42	3.1 %
3	Both	30	2.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-107	Follow-up skipped because gate <> 1	1113	82.2 %

Value	Label	Unweighted Frequency	%
	Total	**1,354**	**100%**

Based upon 97 valid cases out of 1,354 total cases.

Location: 3516-3519 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM08_SUSYMB: S7 SubUse: Ever not remember prior night because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	149	11.0 %
5	No	640	47.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-106	Skip-in for alchstat/drugstat	421	31.1 %
	Total	**1,354**	**100%**

Based upon 789 valid cases out of 1,354 total cases.

Location: 3520-3523 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM08_SUSYMC: S7 SubUse: Specify cause of not remember night [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	88	6.5 %
2	Drugs	22	1.6 %
3	Both	39	2.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-107	Follow-up skipped because gate <> 1	1061	78.4 %
	Total	**1,354**	**100%**

Based upon 149 valid cases out of 1,354 total cases.

Location: 3524-3527 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM09_SUSYMB: S7 SubUse: Ever used in situations could get hurt in recall period [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	150	11.1 %
5	No	639	47.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-106	Skip-in for alchstat/drugstat	421	31.1 %
	Total	**1,354**	**100%**

Based upon 789 valid cases out of 1,354 total cases.

Location: 3528-3531 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM09_SUSYMC: S7 SubUse: Specify used drugs in hurtful situations [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	51	3.8 %
2	Drugs	44	3.2 %
3	Both	55	4.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-107	Follow-up skipped because gate <> 1	1060	78.3 %
	Total	**1,354**	**100%**

Based upon 150 valid cases out of 1,354 total cases.

Location: 3532-3535 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM10_SUSYMB: S7 SubUse: Ever had accident/injury because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	36	2.7 %
5	No	753	55.6 %
	Missing Data		

Value	Label	Unweighted Frequency	%
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-106	Skip-in for alchstat/drugstat	421	31.1 %
	Total	**1,354**	**100%**

Based upon 789 valid cases out of 1,354 total cases.

Location: 3536-3539 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM10_SUSYMC: S7 SubUse: Specify cause of accident/injury [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	21	1.6 %
2	Drugs	8	0.6 %
3	Both	7	0.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-107	Follow-up skipped because gate <> 1	1174	86.7 %
	Total	**1,354**	**100%**

Based upon 36 valid cases out of 1,354 total cases.

Location: 3540-3543 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM11_SUSYMB: S7 SubUse: Alcohol/drug use interfere work/responsibility school/home/job in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	73	5.4 %
5	No	716	52.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-106	Skip-in for alchstat/drugstat	421	31.1 %
	Total	**1,354**	**100%**

Based upon 789 valid cases out of 1,354 total cases.

Location: 3544-3547 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM11_SUSYMC: S7 SubUse: Specify cause of interference [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	21	1.6 %
2	Drugs	34	2.5 %
3	Both	18	1.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-107	Follow-up skipped because gate <> 1	1137	84.0 %
	Total	**1,354**	**100%**

Based upon 73 valid cases out of 1,354 total cases.

Location: 3548-3551 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM12_SUSYMB: S7 SubUse: Ever trouble police no arrest because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	60	4.4 %
5	No	729	53.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-106	Skip-in for alchstat/drugstat	421	31.1 %
	Total	**1,354**	**100%**

Based upon 789 valid cases out of 1,354 total cases.

Location: 3552-3555 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM12_SUSYMC: S7 SubUse: Specify cause trouble police no arrest [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	24	1.8 %
2	Drugs	26	1.9 %
3	Both	10	0.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-107	Follow-up skipped because gate <> 1	1150	84.9 %
	Total	**1,354**	**100%**

Based upon 60 valid cases out of 1,354 total cases.

Location: 3556-3559 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM13_SUSYMB: S7 SubUse: Sold/exchanged sex for drugs in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	4	0.3 %
5	No	785	58.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-106	Skip-in for alchstat/drugstat	421	31.1 %
	Total	**1,354**	**100%**

Based upon 789 valid cases out of 1,354 total cases.

Location: 3560-3563 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM13_SUSYMC: S7 SubUse: Specify cause sold/exchanged sex [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	1	0.1 %
2	Drugs	2	0.1 %
3	Both	1	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %

Value	Label	Unweighted Frequency	%
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-107	Follow-up skipped because gate <> 1	1206	89.1 %
	Total	**1,354**	**100%**

Based upon 4 valid cases out of 1,354 total cases.

Location: 3564-3567 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM14_SUSYMB: S7 SubUse: Sold drugs for money buy drugs in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	70	5.2 %
5	No	718	53.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-106	Skip-in for alchstat/drugstat	421	31.1 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 788 valid cases out of 1,354 total cases.

Location: 3568-3571 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM14_SUSYMC: S7 SubUse: Specify cause sold drugs for money to buy [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	1	0.1 %
2	Drugs	62	4.6 %
3	Both	7	0.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-107	Follow-up skipped because gate <> 1	1140	84.2 %
	Total	**1,354**	**100%**

Based upon 70 valid cases out of 1,354 total cases.

Location: 3572-3575 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM15_SUSYMB: S7 SubUse: Stole for money buy drugs in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	33	2.4 %
5	No	755	55.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-106	Skip-in for alchstat/drugstat	421	31.1 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 788 valid cases out of 1,354 total cases.

Location: 3576-3579 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM15_SUSYMC: S7 SubUse: Specify cause stole for money to buy [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	0	0.0 %
2	Drugs	30	2.2 %
3	Both	3	0.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-107	Follow-up skipped because gate <> 1	1177	86.9 %
	Total	**1,354**	**100%**

Based upon 33 valid cases out of 1,354 total cases.

Location: 3580-3583 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM16_SUSYMB: S7 SubUse: Ever had physical fight because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	76	5.6 %
5	No	713	52.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-106	Skip-in for alchstat/drugstat	421	31.1 %
	Total	**1,354**	**100%**

Based upon 789 valid cases out of 1,354 total cases.

Location: 3584-3587 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM16_SUSYMC: S7 SubUse: Specify cause of fight [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	41	3.0 %
2	Drugs	16	1.2 %
3	Both	19	1.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-107	Follow-up skipped because gate <> 1	1134	83.8 %
	Total	**1,354**	**100%**

Based upon 76 valid cases out of 1,354 total cases.

Location: 3588-3591 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM17_SUSYMB: S7 SubUse: Ever destroy property because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	38	2.8 %
5	No	751	55.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %

Value	Label	Unweighted Frequency	%
-106	Skip-in for alchstat/drugstat	421	31.1 %
	Total	**1,354**	**100%**

Based upon 789 valid cases out of 1,354 total cases.

Location: 3592-3595 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM17_SUSYMC: S7 SubUse: Specify cause of destroy property [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	16	1.2 %
2	Drugs	14	1.0 %
3	Both	8	0.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-107	Follow-up skipped because gate <> 1	1172	86.6 %
	Total	**1,354**	**100%**

Based upon 38 valid cases out of 1,354 total cases.

Location: 3596-3599 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM18_SUSYMB: S7 SubUse: Ever felt strong urge use drug/alcohol no self stop in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	73	5.4 %
5	No	716	52.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-106	Skip-in for alchstat/drugstat	421	31.1 %
	Total	**1,354**	**100%**

Based upon 789 valid cases out of 1,354 total cases.

Location: 3600-3603 (width: 4; decimal: 0)
Variable Type: numeric

(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM18_SUSYMC: S7 SubUse: Specify the cause of strong urge [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	16	1.2 %
2	Drugs	46	3.4 %
3	Both	11	0.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-107	Follow-up skipped because gate <> 1	1137	84.0 %
	Total	**1,354**	**100%**

Based upon 73 valid cases out of 1,354 total cases.

Location: 3604-3607 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM19_SUSYMB: S7 SubUse: Ever want drug/alcohol think of nothing else in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	61	4.5 %
5	No	728	53.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-106	Skip-in for alchstat/drugstat	421	31.1 %
	Total	**1,354**	**100%**

Based upon 789 valid cases out of 1,354 total cases.

Location: 3608-3611 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM19_SUSYMC: S7 SubUse: Specify cause of want [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	18	1.3 %
2	Drugs	32	2.4 %

Value	Label	Unweighted Frequency	%
3	Both	11	0.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-107	Follow-up skipped because gate <> 1	1149	84.9 %
	Total	**1,354**	**100%**

Based upon 61 valid cases out of 1,354 total cases.

Location: 3612-3615 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM20_SUSYMB: S7 SubUse: Ever experience withdrawal from drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	56	4.1 %
5	No	733	54.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-106	Skip-in for alchstat/drugstat	421	31.1 %
	Total	**1,354**	**100%**

Based upon 789 valid cases out of 1,354 total cases.

Location: 3616-3619 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM20_SUSYMC: S7 SubUse: Specify cause of withdraw [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	6	0.4 %
2	Drugs	39	2.9 %
3	Both	11	0.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-107	Follow-up skipped because gate <> 1	1154	85.2 %

Value	Label	Unweighted Frequency	%
	Total	**1,354**	**100%**

Based upon 56 valid cases out of 1,354 total cases.

Location: 3620-3623 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM21_SUSYMB: S7 SubUse: Take drug/alcohol to keep from having withdraw in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	31	2.3 %
5	No	25	1.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-108	Skip-in for SUBSYM20=5	733	54.1 %
-106	Skip-in for alchstat/drugstat	421	31.1 %
	Total	**1,354**	**100%**

Based upon 56 valid cases out of 1,354 total cases.

Location: 3624-3627 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM21_SUSYMC: S7 SubUse: Specify reason for consumption [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	3	0.2 %
2	Drugs	24	1.8 %
3	Both	4	0.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-107	Follow-up skipped because gate <> 1	1179	87.1 %
	Total	**1,354**	**100%**

Based upon 31 valid cases out of 1,354 total cases.

Location: 3628-3631 (width: 4; decimal: 0)
Variable Type: numeric

(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM22_SUSYMB: S7 SubUse: Give up usual activities/interests to use drug/alcohol in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	62	4.6 %
5	No	727	53.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-106	Skip-in for alchstat/drugstat	421	31.1 %
	Total	**1,354**	**100%**

Based upon 789 valid cases out of 1,354 total cases.

Location: 3632-3635 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM22_SUSYMC: S7 SubUse: Specify cause of giving up activities/interests [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	16	1.2 %
2	Drugs	34	2.5 %
3	Both	12	0.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-107	Follow-up skipped because gate <> 1	1148	84.8 %
	Total	**1,354**	**100%**

Based upon 62 valid cases out of 1,354 total cases.

Location: 3636-3639 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM23_SUSYMB: S7 SubUse: Ever used in rp when promised wouldn't [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	132	9.7 %
5	No	657	48.5 %

Value	Label	Unweighted Frequency	%
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-106	Skip-in for alchstat/drugstat	421	31.1 %
	Total	**1,354**	**100%**

Based upon 789 valid cases out of 1,354 total cases.

Location: 3640-3643 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM23_SUSYMC: S7 SubUse: Specify what use you started [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	36	2.7 %
2	Drugs	73	5.4 %
3	Both	23	1.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-107	Follow-up skipped because gate <> 1	1078	79.6 %
	Total	**1,354**	**100%**

Based upon 132 valid cases out of 1,354 total cases.

Location: 3644-3647 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM24_SUSYMB: S7 SubUse: Ever use drug/alcohol more freq than intended in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	134	9.9 %
5	No	655	48.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-106	Skip-in for alchstat/drugstat	421	31.1 %

Value	Label	Unweighted Frequency	%
	Total	**1,354**	**100%**

Based upon 789 valid cases out of 1,354 total cases.

Location: 3648-3651 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM24_SUSYMC: S7 SubUse: Specify what you used more frequently [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	36	2.7 %
2	Drugs	73	5.4 %
3	Both	25	1.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-107	Follow-up skipped because gate <> 1	1076	79.5 %
	Total	**1,354**	**100%**

Based upon 134 valid cases out of 1,354 total cases.

Location: 3652-3655 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM25_SUSYMB: S7 SubUse: Ever tried cut down drug/alcohol use but couldnt in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	100	7.4 %
5	No	689	50.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-106	Skip-in for alchstat/drugstat	421	31.1 %
	Total	**1,354**	**100%**

Based upon 789 valid cases out of 1,354 total cases.

Location: 3656-3659 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM25_SUSYMC: S7 SubUse: Specify what substance use you tried to curtail [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	16	1.2 %
2	Drugs	69	5.1 %
3	Both	15	1.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-107	Follow-up skipped because gate <> 1	1110	82.0 %
	Total	**1,354**	**100%**

Based upon 100 valid cases out of 1,354 total cases.

Location: 3660-3663 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM26_SUSYMB: S7 SubUse: Ever need more drug/alcohol get same effect in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	144	10.6 %
5	No	645	47.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-106	Skip-in for alchstat/drugstat	421	31.1 %
	Total	**1,354**	**100%**

Based upon 789 valid cases out of 1,354 total cases.

Location: 3664-3667 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM26_SUSYMC: S7 SubUse: Specify substance need more for same effect [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	43	3.2 %
2	Drugs	74	5.5 %
3	Both	27	2.0 %

Value	Label	Unweighted Frequency	%
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-107	Follow-up skipped because gate <> 1	1066	78.7 %
	Total	**1,354**	**100%**

Based upon 144 valid cases out of 1,354 total cases.

Location: 3668-3671 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM27_SUSYMB: S7 SubUse: Ever used drug/alcohol enough to depend in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	65	4.8 %
5	No	724	53.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-106	Skip-in for alchstat/drugstat	421	31.1 %
	Total	**1,354**	**100%**

Based upon 789 valid cases out of 1,354 total cases.

Location: 3672-3675 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM27_SUSYMC: S7 SubUse: Specify dependency [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	9	0.7 %
2	Drugs	43	3.2 %
3	Both	13	1.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-107	Follow-up skipped because gate <> 1	1145	84.6 %
	Total	**1,354**	**100%**

Based upon 65 valid cases out of 1,354 total cases.

Location: 3676-3679 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM28_SUSYMB: S7 SubUse: Ever talked doctor/clergy/guidance counselor drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	28	2.1 %
5	No	761	56.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-106	Skip-in for alchstat/drugstat	421	31.1 %
	Total	**1,354**	**100%**

Based upon 789 valid cases out of 1,354 total cases.

Location: 3680-3683 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM28_SUSYMC: S7 SubUse: Specify talked doctor/clergy/guidance counselor [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	11	0.8 %
2	Drugs	10	0.7 %
3	Both	7	0.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-107	Follow-up skipped because gate <> 1	1182	87.3 %
	Total	**1,354**	**100%**

Based upon 28 valid cases out of 1,354 total cases.

Location: 3684-3687 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM29_SUSYMB: S7 SubUse: Ever attended self-help group for drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	69	5.1 %
5	No	720	53.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-106	Skip-in for alchstat/drugstat	421	31.1 %
	Total	**1,354**	**100%**

Based upon 789 valid cases out of 1,354 total cases.

Location: 3688-3691 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM29_SUSYMC: S7 SubUse: Specify reason for self-help [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	26	1.9 %
2	Drugs	22	1.6 %
3	Both	20	1.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-107	Follow-up skipped because gate <> 1	1141	84.3 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 68 valid cases out of 1,354 total cases.

Location: 3692-3695 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM30_SUSYMB: S7 SubUse: Ever treated by therapist because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	20	1.5 %
5	No	769	56.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %

Value	Label	Unweighted Frequency	%
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-106	Skip-in for alchstat/drugstat	421	31.1 %
	Total	**1,354**	**100%**

Based upon 789 valid cases out of 1,354 total cases.

Location: 3696-3699 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM30_SUSYMC: S7 SubUse: Specify reason for therapist visit [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	3	0.2 %
2	Drugs	10	0.7 %
3	Both	7	0.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-107	Follow-up skipped because gate <> 1	1190	87.9 %
	Total	**1,354**	**100%**

Based upon 20 valid cases out of 1,354 total cases.

Location: 3700-3703 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM31_SUSYMB: S7 SubUse: Ever Admitted to hospital because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	8	0.6 %
5	No	781	57.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-106	Skip-in for alchstat/drugstat	421	31.1 %
	Total	**1,354**	**100%**

Based upon 789 valid cases out of 1,354 total cases.

Location: 3704-3707 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE_SUBSYM31_SUSYMC: S7 SubUse: Specify reason admitted to hospital [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	4	0.3 %
2	Drugs	3	0.2 %
3	Both	1	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
-107	Follow-up skipped because gate <> 1	1202	88.8 %
	Total	**1,354**	**100%**

Based upon 8 valid cases out of 1,354 total cases.

Location: 3708-3711 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S7SUBUSE28: S7 SubUse: Num times abused prescribed medications (own or someone elses) during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	0	0.0 %
2	1-5 times	0	0.0 %
3	6-11 times	1	0.1 %
4	1X per month	0	0.0 %
5	2-3X per month	0	0.0 %
6	1X per week	0	0.0 %
7	2-3X per week	0	0.0 %
8	4-5X per week	0	0.0 %
9	Every day	0	0.0 %
	Missing Data		
-800	Question not asked for this subject	1209	89.3 %
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	4	0.3 %
	Total	**1,354**	**100%**

Based upon 1 valid cases out of 1,354 total cases.

Location: 3712-3715 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S7QFBRMO: S7 SubUse: Quant/freq used beer/wine/liquor - recall period

Based upon 1,209 valid cases out of 1,354 total cases.

- Mean: 12.64
- Median: 3.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 402.00
- Standard Deviation: 24.59

Location: 3716-3719 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -300 , -202 , -201 , -100 , -9 , -8

S76MOUSE: S7 SubUse: Number drugs used - recall period

Based upon 1,210 valid cases out of 1,354 total cases.

- Mean: 0.69
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 9.00
- Standard Deviation: 1.09

Location: 3720-3723 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -300 , -202 , -201 , -100 , -9 , -8

S7QFSMK6: S7 SubUse: Quant/freq used cigarettes - recall period

Based upon 1,208 valid cases out of 1,354 total cases.

- Mean: 37.77
- Median: 15.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 160.00
- Standard Deviation: 49.32

Location: 3724-3729 (width: 6; decimal: 1)
Variable Type: numeric
(Range of) Missing Values: -300.0 , -202.0 , -201.0 , -100.0 , -9.0 , -8.0

S7A6SC2: S7 SubUse: Recall period alcohol consequences

Based upon 1,214 valid cases out of 1,354 total cases.

- Mean: 0.66
- Median: 0.00

- Mode: 0.00
- Minimum: 0.00
- Maximum: 13.00
- Standard Deviation: 1.74

Location: 3730-3733 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S7D6SC2: S7 SubUse: Recall period drug consequences

Based upon 1,214 valid cases out of 1,354 total cases.

- Mean: 0.72
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 13.00
- Standard Deviation: 1.89

Location: 3734-3737 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S7A6DP2: S7 SubUse: Recall period alcohol dependence symptoms

Based upon 1,214 valid cases out of 1,354 total cases.

- Mean: 0.29
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 10.00
- Standard Deviation: 1.08

Location: 3738-3741 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S7D6DP2: S7 SubUse: Recall period drug dependence symptoms

Based upon 1,214 valid cases out of 1,354 total cases.

- Mean: 0.54
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 10.00
- Standard Deviation: 1.64

Location: 3742-3745 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S7TOTSC6: S7 SubUse: Recall period total substance use consequences

Based upon 1,214 valid cases out of 1,354 total cases.

- Mean: 1.12
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 13.00
- Standard Deviation: 2.36

Location: 3746-3749 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S7TOTDS6: S7 SubUse: Recall period total substance use dependency symp

Based upon 1,214 valid cases out of 1,354 total cases.

- Mean: 0.71
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 10.00
- Standard Deviation: 1.81

Location: 3750-3753 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S7ALTRTM: S7 SubUse: Recall period alcohol treatment

Value	Label	Unweighted Frequency	%
0	No	1163	85.9 %
1	Yes	51	3.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
	Total	**1,354**	**100%**

Based upon 1,214 valid cases out of 1,354 total cases.

Location: 3754-3757 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S7DGTRTM: S7 SubUse: Recall period drug treatment

Value	Label	Unweighted Frequency	%
0	No	1169	86.3 %
1	Yes	45	3.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %

Value	Label	Unweighted Frequency	%
	Total	**1,354**	**100%**

Based upon 1,214 valid cases out of 1,354 total cases.

Location: 3758-3761 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S7TRTM: S7 SubUse: Recall period alcohol or drug treatment

Value	Label	Unweighted Frequency	%
0	No	1138	84.0 %
1	Yes	76	5.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	1	0.1 %
-201	IntCompStat = 4 missed interview	139	10.3 %
	Total	**1,354**	**100%**

Based upon 1,214 valid cases out of 1,354 total cases.

Location: 3762-3765 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S8SUBUSE1: S8 SubUse: How often had alcohol to drink in recall period

Value	Label	Unweighted Frequency	%
1	Not at all	506	37.4 %
2	1-5 times	160	11.8 %
3	6-11 times	63	4.7 %
4	1X per month	71	5.2 %
5	2-3X per month	129	9.5 %
6	1X per week	87	6.4 %
7	2-3X per week	133	9.8 %
8	4-5X per week	28	2.1 %
9	Every day	23	1.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
	Total	**1,354**	**100%**

Based upon 1,200 valid cases out of 1,354 total cases.

Location: 3766-3769 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S8SUBUSE2: S8 SubUse: Num alcoholic drinks usually consumed

Based upon 1,200 valid cases out of 1,354 total cases.

- Mean: 3.40
- Median: 2.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 60.00
- Standard Deviation: 5.17

Location: 3770-3773 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -101 , -99 , -98

S8SUBUSE3: S8 SubUse: How often had 5+ drinks at one time in recall period

Value	Label	Unweighted Frequency	%
1	Not at all	727	53.7 %
2	1-5 times	147	10.9 %
3	6-11 times	50	3.7 %
4	1X per month	51	3.8 %
5	2-3X per month	66	4.9 %
6	1X per week	57	4.2 %
7	2-3X per week	82	6.1 %
8	4-5X per week	10	0.7 %
9	Every day	10	0.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
	Total	**1,354**	**100%**

Based upon 1,200 valid cases out of 1,354 total cases.

Location: 3774-3777 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S8SUBUSE4: S8 SubUse: Num times been drunk on alcohol during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	691	51.0 %

Value	Label	Unweighted Frequency	%
2	1-5 times	226	16.7 %
3	6-11 times	60	4.4 %
4	1X per month	47	3.5 %
5	2-3X per month	53	3.9 %
6	1X per week	55	4.1 %
7	2-3X per week	57	4.2 %
8	4-5X per week	5	0.4 %
9	Every day	5	0.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-99	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,199 valid cases out of 1,354 total cases.

Location: 3778-3781 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S8SUBUSE6: S8 SubUse: Num times used marijuana/hashish during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	757	55.9 %
2	1-5 times	96	7.1 %
3	6-11 times	32	2.4 %
4	1X per month	20	1.5 %
5	2-3X per month	31	2.3 %
6	1X per week	24	1.8 %
7	2-3X per week	51	3.8 %
8	4-5X per week	44	3.2 %
9	Every day	145	10.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
	Total	**1,354**	**100%**

Based upon 1,200 valid cases out of 1,354 total cases.

Location: 3782-3785 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S8SUBUSE8: S8 SubUse: Num times used sedative/tranquilizers during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1140	84.2 %
2	1-5 times	18	1.3 %
3	6-11 times	8	0.6 %
4	1X per month	8	0.6 %
5	2-3X per month	6	0.4 %
6	1X per week	4	0.3 %
7	2-3X per week	11	0.8 %
8	4-5X per week	1	0.1 %
9	Every day	4	0.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
	Total	**1,354**	**100%**

Based upon 1,200 valid cases out of 1,354 total cases.

Location: 3786-3789 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S8SUBUSE10: S8 SubUse: Num times used stimulants/amphetamines during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1132	83.6 %
2	1-5 times	15	1.1 %
3	6-11 times	8	0.6 %
4	1X per month	8	0.6 %
5	2-3X per month	9	0.7 %
6	1X per week	4	0.3 %
7	2-3X per week	4	0.3 %
8	4-5X per week	9	0.7 %
9	Every day	11	0.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %

Value	Label	Unweighted Frequency	%
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
	Total	**1,354**	**100%**

Based upon 1,200 valid cases out of 1,354 total cases.

Location: 3790-3793 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S8SUBUSE12: S8 SubUse: Num times used cocaine during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1142	84.3 %
2	1-5 times	26	1.9 %
3	6-11 times	5	0.4 %
4	1X per month	8	0.6 %
5	2-3X per month	8	0.6 %
6	1X per week	6	0.4 %
7	2-3X per week	2	0.1 %
8	4-5X per week	1	0.1 %
9	Every day	2	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
	Total	**1,354**	**100%**

Based upon 1,200 valid cases out of 1,354 total cases.

Location: 3794-3797 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S8SUBUSE14: S8 SubUse: Num times used opiates during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1159	85.6 %
2	1-5 times	17	1.3 %
3	6-11 times	3	0.2 %
4	1X per month	5	0.4 %
5	2-3X per month	4	0.3 %
6	1X per week	5	0.4 %

Value	Label	Unweighted Frequency	%
7	2-3X per week	4	0.3 %
8	4-5X per week	0	0.0 %
9	Every day	3	0.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
	Total	**1,354**	**100%**

Based upon 1,200 valid cases out of 1,354 total cases.

Location: 3798-3801 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S8SUBUSE16: S8 SubUse: Num times used ecstasy during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1158	85.5 %
2	1-5 times	27	2.0 %
3	6-11 times	4	0.3 %
4	1X per month	2	0.1 %
5	2-3X per month	6	0.4 %
6	1X per week	1	0.1 %
7	2-3X per week	0	0.0 %
8	4-5X per week	0	0.0 %
9	Every day	2	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
	Total	**1,354**	**100%**

Based upon 1,200 valid cases out of 1,354 total cases.

Location: 3802-3805 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S8SUBUSE18: S8 SubUse: Num times used hallucinogens during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1166	86.1 %
2	1-5 times	18	1.3 %
3	6-11 times	3	0.2 %
4	1X per month	3	0.2 %
5	2-3X per month	5	0.4 %
6	1X per week	2	0.1 %
7	2-3X per week	1	0.1 %
8	4-5X per week	1	0.1 %
9	Every day	1	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
	Total	**1,354**	**100%**

Based upon 1,200 valid cases out of 1,354 total cases.

Location: 3806-3809 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S8SUBUSE20: S8 SubUse: Num times used inhalants during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1189	87.8 %
2	1-5 times	8	0.6 %
3	6-11 times	0	0.0 %
4	1X per month	0	0.0 %
5	2-3X per month	2	0.1 %
6	1X per week	0	0.0 %
7	2-3X per week	0	0.0 %
8	4-5X per week	0	0.0 %
9	Every day	1	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
	Total	**1,354**	**100%**

Based upon 1,200 valid cases out of 1,354 total cases.

Location: 3810-3813 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S8SUBUSE22: S8 SubUse: Num times used amyl nitrate/odorizers/rush during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1195	88.3 %
2	1-5 times	1	0.1 %
3	6-11 times	0	0.0 %
4	1X per month	0	0.0 %
5	2-3X per month	3	0.2 %
6	1X per week	0	0.0 %
7	2-3X per week	0	0.0 %
8	4-5X per week	0	0.0 %
9	Every day	1	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
	Total	**1,354**	**100%**

Based upon 1,200 valid cases out of 1,354 total cases.

Location: 3814-3817 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S8SUBUSE24: S8 SubUse: Used other drugs during recall period [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	1	0.1 %
5	No	1199	88.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
	Total	**1,354**	**100%**

Based upon 1,200 valid cases out of 1,354 total cases.

Location: 3818-3821 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE26: S8 SubUse: Num times used other drugs during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1199	88.6 %
2	1-5 times	1	0.1 %
3	6-11 times	0	0.0 %
4	1X per month	0	0.0 %
5	2-3X per month	0	0.0 %
6	1X per week	0	0.0 %
7	2-3X per week	0	0.0 %
8	4-5X per week	0	0.0 %
9	Every day	0	0.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
	Total	**1,354**	**100%**

Based upon 1,200 valid cases out of 1,354 total cases.

Location: 3822-3825 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S8SUBUSEDY: S8 SubUse: Used any drugs past 24hrs [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	127	9.4 %
5	No	1073	79.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
	Total	**1,354**	**100%**

Based upon 1,200 valid cases out of 1,354 total cases.

Location: 3826-3829 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_DRUG24MAR: S8 SubUse_Drug24: Used marijuana/hashish past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	118	8.7 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-104	Skip-in for SUBUSEDY 1	12	0.9 %
-103	Skip-in for SUBUSEDY <> 1	1073	79.2 %
	Total	**1,354**	**100%**

Based upon 118 valid cases out of 1,354 total cases.

Location: 3830-3833 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_DRUG24TRANQ: S8 SubUse_Drug24: Used sedatives/tranquilizers past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	2	0.1 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-104	Skip-in for SUBUSEDY 1	128	9.5 %
-103	Skip-in for SUBUSEDY <> 1	1073	79.2 %
	Total	**1,354**	**100%**

Based upon 2 valid cases out of 1,354 total cases.

Location: 3834-3837 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_DRUG24AMPH: S8 SubUse_Drug24: Used stimulants/amphetamines past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	5	0.4 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %

Value	Label	Unweighted Frequency	%
-104	Skip-in for SUBUSEDY 1	125	9.2 %
-103	Skip-in for SUBUSEDY <> 1	1073	79.2 %
	Total	**1,354**	**100%**

Based upon 5 valid cases out of 1,354 total cases.

Location: 3838-3841 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_DRUG24COC: S8 SubUse_Drug24: Used Cocaine past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	6	0.4 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-104	Skip-in for SUBUSEDY 1	124	9.2 %
-103	Skip-in for SUBUSEDY <> 1	1073	79.2 %
	Total	**1,354**	**100%**

Based upon 6 valid cases out of 1,354 total cases.

Location: 3842-3845 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_DRUG24OPI: S8 SubUse_Drug24: Used Opiates past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	3	0.2 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-104	Skip-in for SUBUSEDY 1	127	9.4 %
-103	Skip-in for SUBUSEDY <> 1	1073	79.2 %
	Total	**1,354**	**100%**

Based upon 3 valid cases out of 1,354 total cases.

Location: 3846-3849 (width: 4; decimal: 0)
Variable Type: numeric

(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_DRUG24ECST: S8 SubUse_Drug24: Used Ecstacy past 24 hours

Value	Label	Unweighted Frequency	%
1	Yes	2	0.1 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-104	Skip-in for SUBUSEDY 1	128	9.5 %
-103	Skip-in for SUBUSEDY <> 1	1073	79.2 %
	Total	**1,354**	**100%**

Based upon 2 valid cases out of 1,354 total cases.

Location: 3850-3853 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_DRUG24HALL: S8 SubUse_Drug24: Used Hallucinogens past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-104	Skip-in for SUBUSEDY 1	130	9.6 %
-103	Skip-in for SUBUSEDY <> 1	1073	79.2 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 3854-3857 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_DRUG24INH: S8 SubUse_Drug24: Used Inhalants past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		

Value	Label	Unweighted Frequency	%
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-104	Skip-in for SUBUSEDY 1	130	9.6 %
-103	Skip-in for SUBUSEDY <> 1	1073	79.2 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 3858-3861 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_DRUG24AMYL: S8 SubUse_Drug24: Used Amyl nitrate/poppers/odorizers 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-104	Skip-in for SUBUSEDY 1	130	9.6 %
-103	Skip-in for SUBUSEDY <> 1	1073	79.2 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 3862-3865 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_DRUG24OTHER: S8 SubUse_Drug24: Used Other (specify) past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-104	Skip-in for SUBUSEDY 1	130	9.6 %
-103	Skip-in for SUBUSEDY <> 1	1073	79.2 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 3866-3869 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_DRUG24NONE: S8 SubUse_Drug24: Used NA (No drugs) past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-104	Skip-in for SUBUSEDY 1	130	9.6 %
-103	Skip-in for SUBUSEDY <> 1	1073	79.2 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 3870-3873 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S8SUBUSE29: S8 SubUse: Specify other drug used past 24hrs

Based upon 1,354 valid cases out of 1,354 total cases.

Location: 3874-3874 (width: 1; decimal: 0)
Variable Type: character

S8SUBUSE30: S8 SubUse: Num times smoked cigarettes during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	524	38.7 %
2	1-5 times	20	1.5 %
3	6-11 times	4	0.3 %
4	1X per month	5	0.4 %
5	2-3X per month	12	0.9 %
6	1X per week	11	0.8 %
7	2-3X per week	44	3.2 %
8	4-5X per week	23	1.7 %
9	Every day	556	41.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %

Value	Label	Unweighted Frequency	%
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-99	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,199 valid cases out of 1,354 total cases.

Location: 3875-3878 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S8SUBUSE31: S8 SubUse: Num cigarettes smoke per day during recall period

Value	Label	Unweighted Frequency	%
1	0 cigarettes	542	40.0 %
2	1-4 cigarettes	193	14.3 %
3	5-9 cigarettes	167	12.3 %
4	10-15 cigarettes	158	11.7 %
5	16-20 cigarettes	116	8.6 %
6	More than a pack a day cigarettes	22	1.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-9	Dont know	2	0.1 %
	Total	**1,354**	**100%**

Based upon 1,198 valid cases out of 1,354 total cases.

Location: 3879-3882 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -105 , -9 , -8

S8SUBUSE_SUBSYM01_SUSYMB: S8 SubUse: Family ever complain because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	172	12.7 %
5	No	616	45.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %

Value	Label	Unweighted Frequency	%
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-106	Skip-in for alchstat/drugstat	412	30.4 %
	Total	**1,354**	**100%**

Based upon 788 valid cases out of 1,354 total cases.

Location: 3883-3886 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM01_SUSYMC: S8 SubUse: Specify cause of family complaint [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	53	3.9 %
2	Drugs	72	5.3 %
3	Both	47	3.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-107	Follow-up skipped because gate <> 1	1027	75.8 %
	Total	**1,354**	**100%**

Based upon 172 valid cases out of 1,354 total cases.

Location: 3887-3890 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM02_SUSYMB: S8 SubUse: Friend ever complain because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	64	4.7 %
5	No	724	53.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-106	Skip-in for alchstat/drugstat	412	30.4 %
	Total	**1,354**	**100%**

Based upon 788 valid cases out of 1,354 total cases.

Location: 3891-3894 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM02_SUSYMC: S8 SubUse: Specify cause of friend complain [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	18	1.3 %
2	Drugs	35	2.6 %
3	Both	11	0.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-107	Follow-up skipped because gate <> 1	1135	83.8 %
	Total	**1,354**	**100%**

Based upon 64 valid cases out of 1,354 total cases.

Location: 3895-3898 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM03_SUSYMB: S8 SubUse: Ever trouble at work because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	21	1.6 %
5	No	767	56.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-106	Skip-in for alchstat/drugstat	412	30.4 %
	Total	**1,354**	**100%**

Based upon 788 valid cases out of 1,354 total cases.

Location: 3899-3902 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM03_SUSYMC: S8 SubUse: Specify cause of trouble at work [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	8	0.6 %
2	Drugs	13	1.0 %
3	Both	0	0.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-107	Follow-up skipped because gate <> 1	1178	87.0 %
	Total	**1,354**	**100%**

Based upon 21 valid cases out of 1,354 total cases.

Location: 3903-3906 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM04_SUSYMB: S8 SubUse: Ever missed work/school because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	56	4.1 %
5	No	732	54.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-106	Skip-in for alchstat/drugstat	412	30.4 %
	Total	**1,354**	**100%**

Based upon 788 valid cases out of 1,354 total cases.

Location: 3907-3910 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM04_SUSYMC: S8 SubUse: Specify cause of missed work/school [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	26	1.9 %

Value	Label	Unweighted Frequency	%
2	Drugs	18	1.3 %
3	Both	12	0.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-107	Follow-up skipped because gate <> 1	1143	84.4 %
	Total	**1,354**	**100%**

Based upon 56 valid cases out of 1,354 total cases.

Location: 3911-3914 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM05_SUSYMB: S8 SubUse: Ever arrested because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	79	5.8 %
5	No	709	52.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-106	Skip-in for alchstat/drugstat	412	30.4 %
	Total	**1,354**	**100%**

Based upon 788 valid cases out of 1,354 total cases.

Location: 3915-3918 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM05_SUSYMC: S8 SubUse: Specify cause of arrest [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	38	2.8 %
2	Drugs	33	2.4 %
3	Both	8	0.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %

Value	Label	Unweighted Frequency	%
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-107	Follow-up skipped because gate <> 1	1120	82.7 %
	Total	**1,354**	**100%**

Based upon 79 valid cases out of 1,354 total cases.

Location: 3919-3922 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM06_SUSYMB: S8 SubUse: Ever argue family/friends because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	114	8.4 %
5	No	674	49.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-106	Skip-in for alchstat/drugstat	412	30.4 %
	Total	**1,354**	**100%**

Based upon 788 valid cases out of 1,354 total cases.

Location: 3923-3926 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM06_SUSYMC: S8 SubUse: Specify cause of argument w/ family/friends [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	43	3.2 %
2	Drugs	47	3.5 %
3	Both	24	1.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %

Value	Label	Unweighted Frequency	%
-107	Follow-up skipped because gate <> 1	1085	80.1 %
	Total	**1,354**	**100%**

Based upon 114 valid cases out of 1,354 total cases.

Location: 3927-3930 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM07_SUSYMB: S8 SubUse: Ever cont. drug/alcohol use in spite cause health problems in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	78	5.8 %
5	No	710	52.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-106	Skip-in for alchstat/drugstat	412	30.4 %
	Total	**1,354**	**100%**

Based upon 788 valid cases out of 1,354 total cases.

Location: 3931-3934 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM07_SUSYMC: S8 SubUse: Specify continued using [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	23	1.7 %
2	Drugs	32	2.4 %
3	Both	23	1.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-107	Follow-up skipped because gate <> 1	1121	82.8 %
	Total	**1,354**	**100%**

Based upon 78 valid cases out of 1,354 total cases.

Location: 3935-3938 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM08_SUSYMB: S8 SubUse: Ever not remember prior night because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	157	11.6 %
5	No	631	46.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-106	Skip-in for alchstat/drugstat	412	30.4 %
	Total	**1,354**	**100%**

Based upon 788 valid cases out of 1,354 total cases.

Location: 3939-3942 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM08_SUSYMC: S8 SubUse: Specify cause of not remember night [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	92	6.8 %
2	Drugs	21	1.6 %
3	Both	44	3.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-107	Follow-up skipped because gate <> 1	1042	77.0 %
	Total	**1,354**	**100%**

Based upon 157 valid cases out of 1,354 total cases.

Location: 3943-3946 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM09_SUSYMB: S8 SubUse: Ever used in situations could get hurt in recall period [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	151	11.2 %
5	No	637	47.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-106	Skip-in for alchstat/drugstat	412	30.4 %
	Total	**1,354**	**100%**

Based upon 788 valid cases out of 1,354 total cases.

Location: 3947-3950 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM09_SUSYMC: S8 SubUse: Specify used drugs in hurtful situations [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	61	4.5 %
2	Drugs	31	2.3 %
3	Both	59	4.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-107	Follow-up skipped because gate <> 1	1048	77.4 %
	Total	**1,354**	**100%**

Based upon 151 valid cases out of 1,354 total cases.

Location: 3951-3954 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM10_SUSYMB: S8 SubUse: Ever had accident/injury because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	37	2.7 %
5	No	751	55.5 %
	Missing Data		

Value	Label	Unweighted Frequency	%
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-106	Skip-in for alchstat/drugstat	412	30.4 %
	Total	**1,354**	**100%**

Based upon 788 valid cases out of 1,354 total cases.

Location: 3955-3958 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM10_SUSYMC: S8 SubUse: Specify cause of accident/injury [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	18	1.3 %
2	Drugs	11	0.8 %
3	Both	8	0.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-107	Follow-up skipped because gate <> 1	1162	85.8 %
	Total	**1,354**	**100%**

Based upon 37 valid cases out of 1,354 total cases.

Location: 3959-3962 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM11_SUSYMB: S8 SubUse: Alcohol/drug use interfere work/responsibility school/home/job in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	56	4.1 %
5	No	732	54.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %

Value	Label	Unweighted Frequency	%
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-106	Skip-in for alchstat/drugstat	412	30.4 %
	Total	**1,354**	**100%**

Based upon 788 valid cases out of 1,354 total cases.

Location: 3963-3966 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM11_SUSYMC: S8 SubUse: Specify cause of interference [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	18	1.3 %
2	Drugs	21	1.6 %
3	Both	17	1.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-107	Follow-up skipped because gate <> 1	1143	84.4 %
	Total	**1,354**	**100%**

Based upon 56 valid cases out of 1,354 total cases.

Location: 3967-3970 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM12_SUSYMB: S8 SubUse: Ever trouble police no arrest because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	55	4.1 %
5	No	733	54.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-106	Skip-in for alchstat/drugstat	412	30.4 %
	Total	**1,354**	**100%**

Based upon 788 valid cases out of 1,354 total cases.

Location: 3971-3974 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM12_SUSYMC: S8 SubUse: Specify cause trouble police no arrest [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	22	1.6 %
2	Drugs	26	1.9 %
3	Both	7	0.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-107	Follow-up skipped because gate <> 1	1144	84.5 %
	Total	**1,354**	**100%**

Based upon 55 valid cases out of 1,354 total cases.

Location: 3975-3978 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM13_SUSYMB: S8 SubUse: Sold/exchanged sex for drugs in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	3	0.2 %
5	No	785	58.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-106	Skip-in for alchstat/drugstat	412	30.4 %
	Total	**1,354**	**100%**

Based upon 788 valid cases out of 1,354 total cases.

Location: 3979-3982 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM13_SUSYMC: S8 SubUse: Specify cause sold/exchanged sex [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	1	0.1 %
2	Drugs	2	0.1 %
3	Both	0	0.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-107	Follow-up skipped because gate <> 1	1196	88.3 %
	Total	**1,354**	**100%**

Based upon 3 valid cases out of 1,354 total cases.

Location: 3983-3986 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM14_SUSYMB: S8 SubUse: Sold drugs for money buy drugs in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	67	4.9 %
5	No	721	53.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-106	Skip-in for alchstat/drugstat	412	30.4 %
	Total	**1,354**	**100%**

Based upon 788 valid cases out of 1,354 total cases.

Location: 3987-3990 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM14_SUSYMC: S8 SubUse: Specify cause sold drugs for money to buy [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	1	0.1 %

Value	Label	Unweighted Frequency	%
2	Drugs	60	4.4 %
3	Both	6	0.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-107	Follow-up skipped because gate <> 1	1132	83.6 %
	Total	**1,354**	**100%**

Based upon 67 valid cases out of 1,354 total cases.

Location: 3991-3994 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM15_SUSYMB: S8 SubUse: Stole for money buy drugs in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	29	2.1 %
5	No	759	56.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-106	Skip-in for alchstat/drugstat	412	30.4 %
	Total	**1,354**	**100%**

Based upon 788 valid cases out of 1,354 total cases.

Location: 3995-3998 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM15_SUSYMC: S8 SubUse: Specify cause stole for money to buy [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	0	0.0 %
2	Drugs	25	1.8 %
3	Both	4	0.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %

Value	Label	Unweighted Frequency	%
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-107	Follow-up skipped because gate <> 1	1170	86.4 %
	Total	**1,354**	**100%**

Based upon 29 valid cases out of 1,354 total cases.

Location: 3999-4002 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM16_SUSYMB: S8 SubUse: Ever had physical fight because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	73	5.4 %
5	No	715	52.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-106	Skip-in for alchstat/drugstat	412	30.4 %
	Total	**1,354**	**100%**

Based upon 788 valid cases out of 1,354 total cases.

Location: 4003-4006 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM16_SUSYMC: S8 SubUse: Specify cause of fight [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	37	2.7 %
2	Drugs	20	1.5 %
3	Both	16	1.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-107	Follow-up skipped because gate <> 1	1126	83.2 %

Value	Label	Unweighted Frequency	%
	Total	**1,354**	**100%**

Based upon 73 valid cases out of 1,354 total cases.

Location: 4007-4010 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM17_SUSYMB: S8 SubUse: Ever destroy property because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	33	2.4 %
5	No	755	55.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-106	Skip-in for alchstat/drugstat	412	30.4 %
	Total	**1,354**	**100%**

Based upon 788 valid cases out of 1,354 total cases.

Location: 4011-4014 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM17_SUSYMC: S8 SubUse: Specify cause of destroy property [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	13	1.0 %
2	Drugs	15	1.1 %
3	Both	5	0.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-107	Follow-up skipped because gate <> 1	1166	86.1 %
	Total	**1,354**	**100%**

Based upon 33 valid cases out of 1,354 total cases.

Location: 4015-4018 (width: 4; decimal: 0)

Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM18_SUSYMB: S8 SubUse: Ever felt strong urge use drug/alcohol no self stop in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	51	3.8 %
5	No	737	54.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-106	Skip-in for alchstat/drugstat	412	30.4 %
	Total	**1,354**	**100%**

Based upon 788 valid cases out of 1,354 total cases.

Location: 4019-4022 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM18_SUSYMC: S8 SubUse: Specify the cause of strong urge [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	9	0.7 %
2	Drugs	29	2.1 %
3	Both	13	1.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-107	Follow-up skipped because gate <> 1	1148	84.8 %
	Total	**1,354**	**100%**

Based upon 51 valid cases out of 1,354 total cases.

Location: 4023-4026 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM19_SUSYMB: S8 SubUse: Ever want drug/alcohol think of nothing else in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	58	4.3 %
5	No	730	53.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-106	Skip-in for alchstat/drugstat	412	30.4 %
	Total	**1,354**	**100%**

Based upon 788 valid cases out of 1,354 total cases.

Location: 4027-4030 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM19_SUSYMC: S8 SubUse: Specify cause of want [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	9	0.7 %
2	Drugs	36	2.7 %
3	Both	13	1.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-107	Follow-up skipped because gate <> 1	1141	84.3 %
	Total	**1,354**	**100%**

Based upon 58 valid cases out of 1,354 total cases.

Location: 4031-4034 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM20_SUSYMB: S8 SubUse: Ever experience withdrawal from drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	47	3.5 %
5	No	741	54.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %

Value	Label	Unweighted Frequency	%
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-106	Skip-in for alchstat/drugstat	412	30.4 %
	Total	**1,354**	**100%**

Based upon 788 valid cases out of 1,354 total cases.

Location: 4035-4038 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM20_SUSYMC: S8 SubUse: Specify cause of withdraw [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	7	0.5 %
2	Drugs	34	2.5 %
3	Both	6	0.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-107	Follow-up skipped because gate <> 1	1152	85.1 %
	Total	**1,354**	**100%**

Based upon 47 valid cases out of 1,354 total cases.

Location: 4039-4042 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM21_SUSYMB: S8 SubUse: Take drug/alcohol to keep from having withdraw in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	25	1.8 %
5	No	22	1.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-108	Skip-in for SUBSYM20=5	741	54.7 %

Value	Label	Unweighted Frequency	%
-106	Skip-in for alchstat/drugstat	412	30.4 %
	Total	**1,354**	**100%**

Based upon 47 valid cases out of 1,354 total cases.

Location: 4043-4046 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM21_SUSYMC: S8 SubUse: Specify reason for consumption [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	3	0.2 %
2	Drugs	18	1.3 %
3	Both	4	0.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-107	Follow-up skipped because gate <> 1	1174	86.7 %
	Total	**1,354**	**100%**

Based upon 25 valid cases out of 1,354 total cases.

Location: 4047-4050 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM22_SUSYMB: S8 SubUse: Give up usual activities/interests to use drug/alcohol in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	39	2.9 %
5	No	749	55.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-106	Skip-in for alchstat/drugstat	412	30.4 %
	Total	**1,354**	**100%**

Based upon 788 valid cases out of 1,354 total cases.

Location: 4051-4054 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM22_SUSYMC: S8 SubUse: Specify cause of giving up activities/interests [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	5	0.4 %
2	Drugs	23	1.7 %
3	Both	11	0.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-107	Follow-up skipped because gate <> 1	1160	85.7 %
	Total	**1,354**	**100%**

Based upon 39 valid cases out of 1,354 total cases.

Location: 4055-4058 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM23_SUSYMB: S8 SubUse: Ever used in rp when promised wouldn't [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	120	8.9 %
5	No	668	49.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-106	Skip-in for alchstat/drugstat	412	30.4 %
	Total	**1,354**	**100%**

Based upon 788 valid cases out of 1,354 total cases.

Location: 4059-4062 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM23_SUSYMC: S8 SubUse: Specify what use you started [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	31	2.3 %
2	Drugs	60	4.4 %
3	Both	29	2.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-107	Follow-up skipped because gate <> 1	1079	79.7 %
	Total	**1,354**	**100%**

Based upon 120 valid cases out of 1,354 total cases.

Location: 4063-4066 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM24_SUSYMB: S8 SubUse: Ever use drug/alcohol more freq than intended in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	114	8.4 %
5	No	674	49.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-106	Skip-in for alchstat/drugstat	412	30.4 %
	Total	**1,354**	**100%**

Based upon 788 valid cases out of 1,354 total cases.

Location: 4067-4070 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM24_SUSYMC: S8 SubUse: Specify what you used more frequently [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	30	2.2 %

Value	Label	Unweighted Frequency	%
2	Drugs	55	4.1 %
3	Both	29	2.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-107	Follow-up skipped because gate <> 1	1085	80.1 %
	Total	**1,354**	**100%**

Based upon 114 valid cases out of 1,354 total cases.

Location: 4071-4074 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM25_SUSYMB: S8 SubUse: Ever tried cut down drug/alcohol use but couldnt in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	85	6.3 %
5	No	703	51.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-106	Skip-in for alchstat/drugstat	412	30.4 %
	Total	**1,354**	**100%**

Based upon 788 valid cases out of 1,354 total cases.

Location: 4075-4078 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM25_SUSYMC: S8 SubUse: Specify what substance use you tried to curtail [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	14	1.0 %
2	Drugs	56	4.1 %
3	Both	15	1.1 %
	Missing Data		

Value	Label	Unweighted Frequency	%
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-107	Follow-up skipped because gate <> 1	1114	82.3 %
	Total	**1,354**	**100%**

Based upon 85 valid cases out of 1,354 total cases.

Location: 4079-4082 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM26_SUSYMB: S8 SubUse: Ever need more drug/alcohol get same effect in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	122	9.0 %
5	No	666	49.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-106	Skip-in for alchstat/drugstat	412	30.4 %
	Total	**1,354**	**100%**

Based upon 788 valid cases out of 1,354 total cases.

Location: 4083-4086 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM26_SUSYMC: S8 SubUse: Specify substance need more for same effect [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	28	2.1 %
2	Drugs	58	4.3 %
3	Both	36	2.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %

Value	Label	Unweighted Frequency	%
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-107	Follow-up skipped because gate <> 1	1077	79.5 %
	Total	**1,354**	**100%**

Based upon 122 valid cases out of 1,354 total cases.

Location: 4087-4090 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM27_SUSYMB: S8 SubUse: Ever used drug/alcohol enough to depend in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	54	4.0 %
5	No	734	54.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-106	Skip-in for alchstat/drugstat	412	30.4 %
	Total	**1,354**	**100%**

Based upon 788 valid cases out of 1,354 total cases.

Location: 4091-4094 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM27_SUSYMC: S8 SubUse: Specify dependency [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	7	0.5 %
2	Drugs	39	2.9 %
3	Both	8	0.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-107	Follow-up skipped because gate <> 1	1145	84.6 %
	Total	**1,354**	**100%**

Based upon 54 valid cases out of 1,354 total cases.

Location: 4095-4098 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM28_SUSYMB: S8 SubUse: Ever talked doctor/clergy/guidance counselor drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	29	2.1 %
5	No	759	56.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-106	Skip-in for alchstat/drugstat	412	30.4 %
	Total	**1,354**	**100%**

Based upon 788 valid cases out of 1,354 total cases.

Location: 4099-4102 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM28_SUSYMC: S8 SubUse: Specify talked doctor/clergy/guidance counselor [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	7	0.5 %
2	Drugs	14	1.0 %
3	Both	8	0.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-107	Follow-up skipped because gate <> 1	1170	86.4 %
	Total	**1,354**	**100%**

Based upon 29 valid cases out of 1,354 total cases.

Location: 4103-4106 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM29_SUSYMB: S8 SubUse: Ever attended self-help group for drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	59	4.4 %
5	No	729	53.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-106	Skip-in for alchstat/drugstat	412	30.4 %
	Total	**1,354**	**100%**

Based upon 788 valid cases out of 1,354 total cases.

Location: 4107-4110 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM29_SUSYMC: S8 SubUse: Specify reason for self-help [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	22	1.6 %
2	Drugs	20	1.5 %
3	Both	17	1.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-107	Follow-up skipped because gate <> 1	1140	84.2 %
	Total	**1,354**	**100%**

Based upon 59 valid cases out of 1,354 total cases.

Location: 4111-4114 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM30_SUSYMB: S8 SubUse: Ever treated by therapist because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	30	2.2 %

Value	Label	Unweighted Frequency	%
5	No	758	56.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-106	Skip-in for alchstat/drugstat	412	30.4 %
	Total	**1,354**	**100%**

Based upon 788 valid cases out of 1,354 total cases.

Location: 4115-4118 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM30_SUSYMC: S8 SubUse: Specify reason for therapist visit [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	7	0.5 %
2	Drugs	14	1.0 %
3	Both	9	0.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-107	Follow-up skipped because gate <> 1	1169	86.3 %
	Total	**1,354**	**100%**

Based upon 30 valid cases out of 1,354 total cases.

Location: 4119-4122 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM31_SUSYMB: S8 SubUse: Ever Admitted to hospital because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	15	1.1 %
5	No	773	57.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %

Value	Label	Unweighted Frequency	%
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-106	Skip-in for alchstat/drugstat	412	30.4 %
	Total	**1,354**	**100%**

Based upon 788 valid cases out of 1,354 total cases.

Location: 4123-4126 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE_SUBSYM31_SUSYMC: S8 SubUse: Specify reason admitted to hospital [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	2	0.1 %
2	Drugs	8	0.6 %
3	Both	5	0.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
-107	Follow-up skipped because gate <> 1	1184	87.4 %
	Total	**1,354**	**100%**

Based upon 15 valid cases out of 1,354 total cases.

Location: 4127-4130 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S8SUBUSE28: S8 SubUse: Num times abused prescribed medications (own or someone elses) during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	0	0.0 %
2	1-5 times	0	0.0 %
3	6-11 times	0	0.0 %
4	1X per month	0	0.0 %
5	2-3X per month	0	0.0 %
6	1X per week	0	0.0 %
7	2-3X per week	0	0.0 %

Value	Label	Unweighted Frequency	%
8	4-5X per week	0	0.0 %
9	Every day	0	0.0 %
	Missing Data		
-800	Question not asked for this subject	1200	88.6 %
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
-110	Skipped: Interview was conducted in federal facility	1	0.1 %
-109	Skipped: Interviewer incorrectly indicated subject was in federal facility	3	0.2 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 4131-4134 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S8QFBRMO: S8 SubUse: Quant/freq used beer/wine/liquor - recall period

Based upon 1,200 valid cases out of 1,354 total cases.

- Mean: 13.82
- Median: 3.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 300.00
- Standard Deviation: 24.24

Location: 4135-4138 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -300 , -202 , -201 , -100 , -9 , -8

S86MOUSE: S8 SubUse: Number drugs used - recall period

Based upon 1,200 valid cases out of 1,354 total cases.

- Mean: 0.64
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 9.00
- Standard Deviation: 1.11

Location: 4139-4142 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -300 , -202 , -201 , -100 , -9 , -8

S8QFSMK6: S8 SubUse: Quant/freq used cigarettes - recall period

Based upon 1,198 valid cases out of 1,354 total cases.

- Mean: 39.92

- Median: 15.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 160.00
- Standard Deviation: 50.84

Location: 4143-4148 (width: 6; decimal: 1)
Variable Type: numeric
(Range of) Missing Values: -300.0 , -202.0 , -201.0 , -100.0 , -9.0 , -8.0

S8A6SC2: S8 SubUse: Recall period alcohol consequences

Based upon 1,204 valid cases out of 1,354 total cases.

- Mean: 0.63
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 11.00
- Standard Deviation: 1.61

Location: 4149-4152 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S8D6SC2: S8 SubUse: Recall period drug consequences

Based upon 1,204 valid cases out of 1,354 total cases.

- Mean: 0.64
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 16.00
- Standard Deviation: 1.89

Location: 4153-4156 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S8A6DP2: S8 SubUse: Recall period alcohol dependence symptoms

Based upon 1,204 valid cases out of 1,354 total cases.

- Mean: 0.25
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 9.00
- Standard Deviation: 0.96

Location: 4157-4160 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S8D6DP2: S8 SubUse: Recall period drug dependence symptoms

Based upon 1,204 valid cases out of 1,354 total cases.

- Mean: 0.48
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 10.00
- Standard Deviation: 1.55

Location: 4161-4164 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S8TOTSC6: S8 SubUse: Recall period total substance use consequences

Based upon 1,204 valid cases out of 1,354 total cases.

- Mean: 1.03
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 16.00
- Standard Deviation: 2.26

Location: 4165-4168 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S8TOTDS6: S8 SubUse: Recall period total substance use dependency symp

Based upon 1,204 valid cases out of 1,354 total cases.

- Mean: 0.59
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 10.00
- Standard Deviation: 1.66

Location: 4169-4172 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S8ALTRTM: S8 SubUse: Recall period alcohol treatment

Value	Label	Unweighted Frequency	%
0	No	1156	85.4 %
1	Yes	48	3.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
	Total	**1,354**	**100%**

Based upon 1,204 valid cases out of 1,354 total cases.

Location: 4173-4176 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S8DGTRTM: S8 SubUse: Recall period drug treatment

Value	Label	Unweighted Frequency	%
0	No	1156	85.4 %
1	Yes	48	3.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
	Total	**1,354**	**100%**

Based upon 1,204 valid cases out of 1,354 total cases.

Location: 4177-4180 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S8TRTM: S8 SubUse: Recall period alcohol or drug treatment

Value	Label	Unweighted Frequency	%
0	No	1131	83.5 %
1	Yes	73	5.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	147	10.9 %
	Total	**1,354**	**100%**

Based upon 1,204 valid cases out of 1,354 total cases.

Location: 4181-4184 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S9SUBUSE1: S9 SubUse: How often had alcohol to drink in recall period

Value	Label	Unweighted Frequency	%
1	Not at all	452	33.4 %
2	1-5 times	154	11.4 %
3	6-11 times	62	4.6 %
4	1X per month	68	5.0 %
5	2-3X per month	129	9.5 %
6	1X per week	94	6.9 %

Value	Label	Unweighted Frequency	%
7	2-3X per week	152	11.2 %
8	4-5X per week	33	2.4 %
9	Every day	25	1.8 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-99	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,169 valid cases out of 1,354 total cases.

Location: 4185-4188 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S9SUBUSE2: S9 SubUse: Num alcoholic drinks usually consumed

Based upon 1,167 valid cases out of 1,354 total cases.

- Mean: 3.57
- Median: 2.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 48.00
- Standard Deviation: 5.20

Location: 4189-4192 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -101 , -99 , -98

S9SUBUSE3: S9 SubUse: How often had 5+ drinks at one time in recall period

Value	Label	Unweighted Frequency	%
1	Not at all	676	49.9 %
2	1-5 times	152	11.2 %
3	6-11 times	41	3.0 %
4	1X per month	45	3.3 %
5	2-3X per month	66	4.9 %
6	1X per week	63	4.7 %
7	2-3X per week	88	6.5 %
8	4-5X per week	26	1.9 %
9	Every day	12	0.9 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-99	Dont know	1	0.1 %

Value	Label	Unweighted Frequency	%
	Total	**1,354**	**100%**

Based upon 1,169 valid cases out of 1,354 total cases.

Location: 4193-4196 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S9SUBUSE4: S9 SubUse: Num times been drunk on alcohol during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	636	47.0 %
2	1-5 times	235	17.4 %
3	6-11 times	48	3.5 %
4	1X per month	54	4.0 %
5	2-3X per month	67	4.9 %
6	1X per week	49	3.6 %
7	2-3X per week	56	4.1 %
8	4-5X per week	15	1.1 %
9	Every day	9	0.7 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-99	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,169 valid cases out of 1,354 total cases.

Location: 4197-4200 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S9SUBUSE6: S9 SubUse: Num times used marijuana/hashish during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	730	53.9 %
2	1-5 times	91	6.7 %
3	6-11 times	22	1.6 %
4	1X per month	25	1.8 %
5	2-3X per month	34	2.5 %
6	1X per week	25	1.8 %
7	2-3X per week	57	4.2 %
8	4-5X per week	45	3.3 %
9	Every day	140	10.3 %

Value	Label	Unweighted Frequency	%
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-99	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,169 valid cases out of 1,354 total cases.

Location: 4201-4204 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S9SUBUSE8: S9 SubUse: Num times used sedative/tranquilizers during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1107	81.8 %
2	1-5 times	25	1.8 %
3	6-11 times	5	0.4 %
4	1X per month	7	0.5 %
5	2-3X per month	11	0.8 %
6	1X per week	2	0.1 %
7	2-3X per week	5	0.4 %
8	4-5X per week	3	0.2 %
9	Every day	4	0.3 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-99	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,169 valid cases out of 1,354 total cases.

Location: 4205-4208 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S9SUBUSE10: S9 SubUse: Num times used stimulants/amphetamines during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1120	82.7 %
2	1-5 times	14	1.0 %
3	6-11 times	1	0.1 %
4	1X per month	5	0.4 %
5	2-3X per month	9	0.7 %

Value	Label	Unweighted Frequency	%
6	1X per week	6	0.4 %
7	2-3X per week	6	0.4 %
8	4-5X per week	2	0.1 %
9	Every day	6	0.4 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-99	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,169 valid cases out of 1,354 total cases.

Location: 4209-4212 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S9SUBUSE12: S9 SubUse: Num times used cocaine during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1123	82.9 %
2	1-5 times	22	1.6 %
3	6-11 times	6	0.4 %
4	1X per month	1	0.1 %
5	2-3X per month	7	0.5 %
6	1X per week	2	0.1 %
7	2-3X per week	3	0.2 %
8	4-5X per week	2	0.1 %
9	Every day	3	0.2 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-99	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,169 valid cases out of 1,354 total cases.

Location: 4213-4216 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S9SUBUSE14: S9 SubUse: Num times used opiates during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1125	83.1 %

Value	Label	Unweighted Frequency	%
2	1-5 times	22	1.6 %
3	6-11 times	2	0.1 %
4	1X per month	2	0.1 %
5	2-3X per month	5	0.4 %
6	1X per week	2	0.1 %
7	2-3X per week	9	0.7 %
8	4-5X per week	1	0.1 %
9	Every day	1	0.1 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-99	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,169 valid cases out of 1,354 total cases.

Location: 4217-4220 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S9SUBUSE16: S9 SubUse: Num times used ecstasy during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1138	84.0 %
2	1-5 times	21	1.6 %
3	6-11 times	1	0.1 %
4	1X per month	2	0.1 %
5	2-3X per month	4	0.3 %
6	1X per week	2	0.1 %
7	2-3X per week	0	0.0 %
8	4-5X per week	0	0.0 %
9	Every day	1	0.1 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-99	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,169 valid cases out of 1,354 total cases.

Location: 4221-4224 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S9SUBUSE18: S9 SubUse: Num times used hallucinogens during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1135	83.8 %
2	1-5 times	21	1.6 %
3	6-11 times	2	0.1 %
4	1X per month	5	0.4 %
5	2-3X per month	3	0.2 %
6	1X per week	0	0.0 %
7	2-3X per week	1	0.1 %
8	4-5X per week	0	0.0 %
9	Every day	2	0.1 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-99	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,169 valid cases out of 1,354 total cases.

Location: 4225-4228 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S9SUBUSE20: S9 SubUse: Num times used inhalants during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1165	86.0 %
2	1-5 times	2	0.1 %
3	6-11 times	1	0.1 %
4	1X per month	0	0.0 %
5	2-3X per month	1	0.1 %
6	1X per week	0	0.0 %
7	2-3X per week	0	0.0 %
8	4-5X per week	0	0.0 %
9	Every day	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-99	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,169 valid cases out of 1,354 total cases.

Location: 4229-4232 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S9SUBUSE22: S9 SubUse: Num times used amyl nitrate/odorizers/rush during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1166	86.1 %
2	1-5 times	2	0.1 %
3	6-11 times	0	0.0 %
4	1X per month	0	0.0 %
5	2-3X per month	1	0.1 %
6	1X per week	0	0.0 %
7	2-3X per week	0	0.0 %
8	4-5X per week	0	0.0 %
9	Every day	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-99	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,169 valid cases out of 1,354 total cases.

Location: 4233-4236 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S9SUBUSE24: S9 SubUse: Used other drugs during recall period [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	2	0.1 %
5	No	1167	86.2 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,169 valid cases out of 1,354 total cases.

Location: 4237-4240 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE26: S9 SubUse: Num times used other drugs during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1167	86.2 %
2	1-5 times	2	0.1 %
3	6-11 times	0	0.0 %
4	1X per month	0	0.0 %
5	2-3X per month	0	0.0 %
6	1X per week	0	0.0 %
7	2-3X per week	0	0.0 %
8	4-5X per week	0	0.0 %
9	Every day	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-102	Skipped: Has not used any other drugs in recall period	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,169 valid cases out of 1,354 total cases.

Location: 4241-4244 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S9SUBUSEDY: S9 SubUse: Used any drugs past 24hrs [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	99	7.3 %
5	No	1070	79.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 1,169 valid cases out of 1,354 total cases.

Location: 4245-4248 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_DRUG24MAR: S9 SubUse_Drug24: Used marijuana/hashish past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	91	6.7 %
5	No	0	0.0 %
	Missing Data		

Value	Label	Unweighted Frequency	%
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-104	Skip-in for SUBUSEDY 1	8	0.6 %
-103	Skip-in for SUBUSEDY <> 1	1071	79.1 %
	Total	**1,354**	**100%**

Based upon 91 valid cases out of 1,354 total cases.

Location: 4249-4252 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_DRUG24TRANQ: S9 SubUse_Drug24: Used sedatives/tranquilizers past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	3	0.2 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-104	Skip-in for SUBUSEDY 1	96	7.1 %
-103	Skip-in for SUBUSEDY <> 1	1071	79.1 %
	Total	**1,354**	**100%**

Based upon 3 valid cases out of 1,354 total cases.

Location: 4253-4256 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_DRUG24AMPH: S9 SubUse_Drug24: Used stimulants/amphetamines past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	2	0.1 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-104	Skip-in for SUBUSEDY 1	97	7.2 %
-103	Skip-in for SUBUSEDY <> 1	1071	79.1 %
	Total	**1,354**	**100%**

Based upon 2 valid cases out of 1,354 total cases.

Location: 4257-4260 (width: 4; decimal: 0)
Variable Type: numeric

(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_DRUG24COC: S9 SubUse_Drug24: Used Cocaine past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-104	Skip-in for SUBUSEDY 1	99	7.3 %
-103	Skip-in for SUBUSEDY <> 1	1071	79.1 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 4261-4264 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_DRUG24OPI: S9 SubUse_Drug24: Used Opiates past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	5	0.4 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-104	Skip-in for SUBUSEDY 1	94	6.9 %
-103	Skip-in for SUBUSEDY <> 1	1071	79.1 %
	Total	**1,354**	**100%**

Based upon 5 valid cases out of 1,354 total cases.

Location: 4265-4268 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_DRUG24ECST: S9 SubUse_Drug24: Used Ecstacy past 24 hours

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %

Value	Label	Unweighted Frequency	%
-104	Skip-in for SUBUSEDY 1	99	7.3 %
-103	Skip-in for SUBUSEDY <> 1	1071	79.1 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 4269-4272 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_DRUG24HALL: S9 SubUse_Drug24: Used Hallucinogens past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	1	0.1 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-104	Skip-in for SUBUSEDY 1	98	7.2 %
-103	Skip-in for SUBUSEDY <> 1	1071	79.1 %
	Total	**1,354**	**100%**

Based upon 1 valid cases out of 1,354 total cases.

Location: 4273-4276 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_DRUG24INH: S9 SubUse_Drug24: Used Inhalants past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-104	Skip-in for SUBUSEDY 1	99	7.3 %
-103	Skip-in for SUBUSEDY <> 1	1071	79.1 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 4277-4280 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_DRUG24AMYL: S9 SubUse_Drug24: Used Amyl nitrate/poppers/odorizers 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-104	Skip-in for SUBUSEDY 1	99	7.3 %
-103	Skip-in for SUBUSEDY <> 1	1071	79.1 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 4281-4284 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_DRUG24OTHER: S9 SubUse_Drug24: Used Other (specify) past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-104	Skip-in for SUBUSEDY 1	99	7.3 %
-103	Skip-in for SUBUSEDY <> 1	1071	79.1 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 4285-4288 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_DRUG24NONE: S9 SubUse_Drug24: Used NA (No drugs) past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-104	Skip-in for SUBUSEDY 1	99	7.3 %

Value	Label	Unweighted Frequency	%
-103	Skip-in for SUBUSEDY <> 1	1071	79.1 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 4289-4292 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

S9SUBUSE29: S9 SubUse: Specify other drug used past 24hrs

Based upon 1,354 valid cases out of 1,354 total cases.

Location: 4293-4293 (width: 1; decimal: 0)
Variable Type: character

S9SUBUSE30: S9 SubUse: Num times smoked cigarettes during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	514	38.0 %
2	1-5 times	7	0.5 %
3	6-11 times	7	0.5 %
4	1X per month	10	0.7 %
5	2-3X per month	15	1.1 %
6	1X per week	17	1.3 %
7	2-3X per week	38	2.8 %
8	4-5X per week	23	1.7 %
9	Every day	537	39.7 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-99	Dont know	2	0.1 %
	Total	**1,354**	**100%**

Based upon 1,168 valid cases out of 1,354 total cases.

Location: 4294-4297 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S9SUBUSE31: S9 SubUse: Num cigarettes smoke per day during recall period

Value	Label	Unweighted Frequency	%
1	0 cigarettes	529	39.1 %
2	1-4 cigarettes	195	14.4 %
3	5-9 cigarettes	154	11.4 %

Value	Label	Unweighted Frequency	%
4	10-15 cigarettes	149	11.0 %
5	16-20 cigarettes	106	7.8 %
6	More than a pack a day cigarettes	35	2.6 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-9	Dont know	2	0.1 %
	Total	**1,354**	**100%**

Based upon 1,168 valid cases out of 1,354 total cases.

Location: 4298-4301 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -105 , -9 , -8

S9SUBUSE_SUBSYM01_SUSYMB: S9 SubUse: Family ever complain because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	149	11.0 %
5	No	641	47.3 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-106	Skip-in for alchstat/drugstat	379	28.0 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 790 valid cases out of 1,354 total cases.

Location: 4302-4305 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM01_SUSYMC: S9 SubUse: Specify cause of family complaint [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	49	3.6 %
2	Drugs	61	4.5 %
3	Both	39	2.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	175	12.9 %

Value	Label	Unweighted Frequency	%
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-107	Follow-up skipped because gate <> 1	1017	75.1 %
	Total	**1,354**	**100%**

Based upon 149 valid cases out of 1,354 total cases.

Location: 4306-4309 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM02_SUSYMB: S9 SubUse: Friend ever complain because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	66	4.9 %
5	No	724	53.5 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-106	Skip-in for alchstat/drugstat	379	28.0 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 790 valid cases out of 1,354 total cases.

Location: 4310-4313 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM02_SUSYMC: S9 SubUse: Specify cause of friend complain [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	27	2.0 %
2	Drugs	31	2.3 %
3	Both	8	0.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-107	Follow-up skipped because gate <> 1	1100	81.2 %
	Total	**1,354**	**100%**

Based upon 66 valid cases out of 1,354 total cases.

Location: 4314-4317 (width: 4; decimal: 0)

Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM03_SUSYMB: S9 SubUse: Ever trouble at work because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	17	1.3 %
5	No	773	57.1 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-106	Skip-in for alchstat/drugstat	379	28.0 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 790 valid cases out of 1,354 total cases.

Location: 4318-4321 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM03_SUSYMC: S9 SubUse: Specify cause of trouble at work [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	7	0.5 %
2	Drugs	8	0.6 %
3	Both	2	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-107	Follow-up skipped because gate <> 1	1149	84.9 %
	Total	**1,354**	**100%**

Based upon 17 valid cases out of 1,354 total cases.

Location: 4322-4325 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM04_SUSYMB: S9 SubUse: Ever missed work/school because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	45	3.3 %

Value	Label	Unweighted Frequency	%
5	No	745	55.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-106	Skip-in for alchstat/drugstat	379	28.0 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 790 valid cases out of 1,354 total cases.

Location: 4326-4329 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM04_SUSYMC: S9 SubUse: Specify cause of missed work/school [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	29	2.1 %
2	Drugs	9	0.7 %
3	Both	7	0.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-107	Follow-up skipped because gate <> 1	1121	82.8 %
	Total	**1,354**	**100%**

Based upon 45 valid cases out of 1,354 total cases.

Location: 4330-4333 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM05_SUSYMB: S9 SubUse: Ever arrested because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	56	4.1 %
5	No	734	54.2 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-106	Skip-in for alchstat/drugstat	379	28.0 %

Value	Label	Unweighted Frequency	%
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 790 valid cases out of 1,354 total cases.

Location: 4334-4337 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM05_SUSYMC: S9 SubUse: Specify cause of arrest [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	26	1.9 %
2	Drugs	19	1.4 %
3	Both	11	0.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-107	Follow-up skipped because gate <> 1	1110	82.0 %
	Total	**1,354**	**100%**

Based upon 56 valid cases out of 1,354 total cases.

Location: 4338-4341 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM06_SUSYMB: S9 SubUse: Ever argue family/friends because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	102	7.5 %
5	No	688	50.8 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-106	Skip-in for alchstat/drugstat	379	28.0 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 790 valid cases out of 1,354 total cases.

Location: 4342-4345 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM06_SUSYMC: S9 SubUse: Specify cause of argument w/ family/friends [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	44	3.2 %
2	Drugs	34	2.5 %
3	Both	24	1.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-107	Follow-up skipped because gate <> 1	1064	78.6 %
	Total	**1,354**	**100%**

Based upon 102 valid cases out of 1,354 total cases.

Location: 4346-4349 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM07_SUSYMB: S9 SubUse: Ever cont. drug/alcohol use in spite cause health problems in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	77	5.7 %
5	No	712	52.6 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-106	Skip-in for alchstat/drugstat	379	28.0 %
-9	Dont know	2	0.1 %
	Total	**1,354**	**100%**

Based upon 789 valid cases out of 1,354 total cases.

Location: 4350-4353 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM07_SUSYMC: S9 SubUse: Specify continued using [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	33	2.4 %
2	Drugs	25	1.8 %
3	Both	19	1.4 %

Value	Label	Unweighted Frequency	%
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-107	Follow-up skipped because gate <> 1	1089	80.4 %
	Total	**1,354**	**100%**

Based upon 77 valid cases out of 1,354 total cases.

Location: 4354-4357 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM08_SUSYMB: S9 SubUse: Ever not remember prior night because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	149	11.0 %
5	No	641	47.3 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-106	Skip-in for alchstat/drugstat	379	28.0 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 790 valid cases out of 1,354 total cases.

Location: 4358-4361 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM08_SUSYMC: S9 SubUse: Specify cause of not remember night [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	93	6.9 %
2	Drugs	15	1.1 %
3	Both	41	3.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-107	Follow-up skipped because gate <> 1	1017	75.1 %

Value	Label	Unweighted Frequency	%
	Total	**1,354**	**100%**

Based upon 149 valid cases out of 1,354 total cases.

Location: 4362-4365 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM09_SUSYMB: S9 SubUse: Ever used in situations could get hurt in recall period [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	149	11.0 %
5	No	641	47.3 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-106	Skip-in for alchstat/drugstat	379	28.0 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 790 valid cases out of 1,354 total cases.

Location: 4366-4369 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM09_SUSYMC: S9 SubUse: Specify used drugs in hurtful situations [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	72	5.3 %
2	Drugs	32	2.4 %
3	Both	45	3.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-107	Follow-up skipped because gate <> 1	1017	75.1 %
	Total	**1,354**	**100%**

Based upon 149 valid cases out of 1,354 total cases.

Location: 4370-4373 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM10_SUSYMB: S9 SubUse: Ever had accident/injury because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	22	1.6 %
5	No	768	56.7 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-106	Skip-in for alchstat/drugstat	379	28.0 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 790 valid cases out of 1,354 total cases.

Location: 4374-4377 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM10_SUSYMC: S9 SubUse: Specify cause of accident/injury [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	12	0.9 %
2	Drugs	6	0.4 %
3	Both	4	0.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-107	Follow-up skipped because gate <> 1	1144	84.5 %
	Total	**1,354**	**100%**

Based upon 22 valid cases out of 1,354 total cases.

Location: 4378-4381 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM11_SUSYMB: S9 SubUse: Alcohol/drug use interfere work/responsibility school/home/job in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	59	4.4 %
5	No	731	54.0 %
	Missing Data		

Value	Label	Unweighted Frequency	%
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-106	Skip-in for alchstat/drugstat	379	28.0 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 790 valid cases out of 1,354 total cases.

Location: 4382-4385 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM11_SUSYMC: S9 SubUse: Specify cause of interference [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	17	1.3 %
2	Drugs	25	1.8 %
3	Both	17	1.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-107	Follow-up skipped because gate <> 1	1107	81.8 %
	Total	**1,354**	**100%**

Based upon 59 valid cases out of 1,354 total cases.

Location: 4386-4389 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM12_SUSYMB: S9 SubUse: Ever trouble police no arrest because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	27	2.0 %
5	No	763	56.4 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-106	Skip-in for alchstat/drugstat	379	28.0 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 790 valid cases out of 1,354 total cases.

Location: 4390-4393 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM12_SUSYMC: S9 SubUse: Specify cause trouble police no arrest [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	15	1.1 %
2	Drugs	8	0.6 %
3	Both	4	0.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-107	Follow-up skipped because gate <> 1	1139	84.1 %
	Total	**1,354**	**100%**

Based upon 27 valid cases out of 1,354 total cases.

Location: 4394-4397 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM13_SUSYMB: S9 SubUse: Sold/exchanged sex for drugs in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	4	0.3 %
5	No	786	58.1 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-106	Skip-in for alchstat/drugstat	379	28.0 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 790 valid cases out of 1,354 total cases.

Location: 4398-4401 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM13_SUSYMC: S9 SubUse: Specify cause sold/exchanged sex [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	1	0.1 %
2	Drugs	2	0.1 %
3	Both	1	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-107	Follow-up skipped because gate <> 1	1162	85.8 %
	Total	**1,354**	**100%**

Based upon 4 valid cases out of 1,354 total cases.

Location: 4402-4405 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM14_SUSYMB: S9 SubUse: Sold drugs for money buy drugs in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	51	3.8 %
5	No	739	54.6 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-106	Skip-in for alchstat/drugstat	379	28.0 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 790 valid cases out of 1,354 total cases.

Location: 4406-4409 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM14_SUSYMC: S9 SubUse: Specify cause sold drugs for money to buy [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	1	0.1 %
2	Drugs	45	3.3 %
3	Both	5	0.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	175	12.9 %

Value	Label	Unweighted Frequency	%
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-107	Follow-up skipped because gate <> 1	1115	82.3 %
	Total	**1,354**	**100%**

Based upon 51 valid cases out of 1,354 total cases.

Location: 4410-4413 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM15_SUSYMB: S9 SubUse: Stole for money buy drugs in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	16	1.2 %
5	No	774	57.2 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-106	Skip-in for alchstat/drugstat	379	28.0 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 790 valid cases out of 1,354 total cases.

Location: 4414-4417 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM15_SUSYMC: S9 SubUse: Specify cause stole for money to buy [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	3	0.2 %
2	Drugs	13	1.0 %
3	Both	0	0.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-107	Follow-up skipped because gate <> 1	1150	84.9 %
	Total	**1,354**	**100%**

Based upon 16 valid cases out of 1,354 total cases.

Location: 4418-4421 (width: 4; decimal: 0)
Variable Type: numeric

(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM16_SUSYMB: S9 SubUse: Ever had physical fight because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	60	4.4 %
5	No	730	53.9 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-106	Skip-in for alchstat/drugstat	379	28.0 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 790 valid cases out of 1,354 total cases.

Location: 4422-4425 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM16_SUSYMC: S9 SubUse: Specify cause of fight [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	41	3.0 %
2	Drugs	11	0.8 %
3	Both	8	0.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-107	Follow-up skipped because gate <> 1	1106	81.7 %
	Total	**1,354**	**100%**

Based upon 60 valid cases out of 1,354 total cases.

Location: 4426-4429 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM17_SUSYMB: S9 SubUse: Ever destroy property because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	33	2.4 %
5	No	757	55.9 %

Value	Label	Unweighted Frequency	%
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-106	Skip-in for alchstat/drugstat	379	28.0 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 790 valid cases out of 1,354 total cases.

Location: 4430-4433 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM17_SUSYMC: S9 SubUse: Specify cause of destroy property [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	18	1.3 %
2	Drugs	9	0.7 %
3	Both	6	0.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-107	Follow-up skipped because gate <> 1	1133	83.7 %
	Total	**1,354**	**100%**

Based upon 33 valid cases out of 1,354 total cases.

Location: 4434-4437 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM18_SUSYMB: S9 SubUse: Ever felt strong urge use drug/alcohol no self stop in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	53	3.9 %
5	No	737	54.4 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-106	Skip-in for alchstat/drugstat	379	28.0 %
-9	Dont know	1	0.1 %

Value	Label	Unweighted Frequency	%
	Total	**1,354**	**100%**

Based upon 790 valid cases out of 1,354 total cases.

Location: 4438-4441 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM18_SUSYMC: S9 SubUse: Specify the cause of strong urge [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	12	0.9 %
2	Drugs	26	1.9 %
3	Both	15	1.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-107	Follow-up skipped because gate <> 1	1113	82.2 %
	Total	**1,354**	**100%**

Based upon 53 valid cases out of 1,354 total cases.

Location: 4442-4445 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM19_SUSYMB: S9 SubUse: Ever want drug/alcohol think of nothing else in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	59	4.4 %
5	No	731	54.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-106	Skip-in for alchstat/drugstat	379	28.0 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 790 valid cases out of 1,354 total cases.

Location: 4446-4449 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM19_SUSYMC: S9 SubUse: Specify cause of want [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	18	1.3 %
2	Drugs	25	1.8 %
3	Both	16	1.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-107	Follow-up skipped because gate <> 1	1107	81.8 %
	Total	**1,354**	**100%**

Based upon 59 valid cases out of 1,354 total cases.

Location: 4450-4453 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM20_SUSYMB: S9 SubUse: Ever experience withdrawal from drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	36	2.7 %
5	No	754	55.7 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-106	Skip-in for alchstat/drugstat	379	28.0 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 790 valid cases out of 1,354 total cases.

Location: 4454-4457 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM20_SUSYMC: S9 SubUse: Specify cause of withdraw [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	10	0.7 %
2	Drugs	23	1.7 %
3	Both	3	0.2 %
	Missing Data		

Value	Label	Unweighted Frequency	%
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-107	Follow-up skipped because gate <> 1	1130	83.5 %
	Total	**1,354**	**100%**

Based upon 36 valid cases out of 1,354 total cases.

Location: 4458-4461 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM21_SUSYMB: S9 SubUse: Take drug/alcohol to keep from having withdraw in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	17	1.3 %
5	No	19	1.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	2	0.1 %
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-108	Skip-in for SUBSYM20=5	751	55.5 %
-106	Skip-in for alchstat/drugstat	380	28.1 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 36 valid cases out of 1,354 total cases.

Location: 4462-4465 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM21_SUSYMC: S9 SubUse: Specify reason for consumption [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	5	0.4 %
2	Drugs	9	0.7 %
3	Both	3	0.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %

Value	Label	Unweighted Frequency	%
-107	Follow-up skipped because gate <> 1	1149	84.9 %
	Total	**1,354**	**100%**

Based upon 17 valid cases out of 1,354 total cases.

Location: 4466-4469 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM22_SUSYMB: S9 SubUse: Give up usual activities/interests to use drug/alcohol in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	49	3.6 %
5	No	741	54.7 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-106	Skip-in for alchstat/drugstat	379	28.0 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 790 valid cases out of 1,354 total cases.

Location: 4470-4473 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM22_SUSYMC: S9 SubUse: Specify cause of giving up activities/interests [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	11	0.8 %
2	Drugs	25	1.8 %
3	Both	13	1.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-107	Follow-up skipped because gate <> 1	1117	82.5 %
	Total	**1,354**	**100%**

Based upon 49 valid cases out of 1,354 total cases.

Location: 4474-4477 (width: 4; decimal: 0)
Variable Type: numeric

(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM23_SUSYMB: S9 SubUse: Ever used in rp when promised wouldn't [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	109	8.1 %
5	No	681	50.3 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-106	Skip-in for alchstat/drugstat	379	28.0 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 790 valid cases out of 1,354 total cases.

Location: 4478-4481 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM23_SUSYMC: S9 SubUse: Specify what use you started [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	31	2.3 %
2	Drugs	53	3.9 %
3	Both	25	1.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-107	Follow-up skipped because gate <> 1	1057	78.1 %
	Total	**1,354**	**100%**

Based upon 109 valid cases out of 1,354 total cases.

Location: 4482-4485 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM24_SUSYMB: S9 SubUse: Ever use drug/alcohol more freq than intended in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	95	7.0 %
5	No	695	51.3 %

Value	Label	Unweighted Frequency	%
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-106	Skip-in for alchstat/drugstat	379	28.0 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 790 valid cases out of 1,354 total cases.

Location: 4486-4489 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM24_SUSYMC: S9 SubUse: Specify what you used more frequently [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	31	2.3 %
2	Drugs	42	3.1 %
3	Both	22	1.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-107	Follow-up skipped because gate <> 1	1071	79.1 %
	Total	**1,354**	**100%**

Based upon 95 valid cases out of 1,354 total cases.

Location: 4490-4493 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM25_SUSYMB: S9 SubUse: Ever tried cut down drug/alcohol use but couldnt in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	72	5.3 %
5	No	718	53.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-106	Skip-in for alchstat/drugstat	379	28.0 %
-9	Dont know	1	0.1 %

Value	Label	Unweighted Frequency	%
	Total	**1,354**	**100%**

Based upon 790 valid cases out of 1,354 total cases.

Location: 4494-4497 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM25_SUSYMC: S9 SubUse: Specify what substance use you tried to curtail [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	12	0.9 %
2	Drugs	49	3.6 %
3	Both	11	0.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-107	Follow-up skipped because gate <> 1	1094	80.8 %
	Total	**1,354**	**100%**

Based upon 72 valid cases out of 1,354 total cases.

Location: 4498-4501 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM26_SUSYMB: S9 SubUse: Ever need more drug/alcohol get same effect in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	113	8.3 %
5	No	677	50.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-106	Skip-in for alchstat/drugstat	379	28.0 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 790 valid cases out of 1,354 total cases.

Location: 4502-4505 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM26_SUSYMC: S9 SubUse: Specify substance need more for same effect [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	36	2.7 %
2	Drugs	54	4.0 %
3	Both	23	1.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-107	Follow-up skipped because gate <> 1	1053	77.8 %
	Total	**1,354**	**100%**

Based upon 113 valid cases out of 1,354 total cases.

Location: 4506-4509 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM27_SUSYMB: S9 SubUse: Ever used drug/alcohol enough to depend in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	55	4.1 %
5	No	735	54.3 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-106	Skip-in for alchstat/drugstat	379	28.0 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 790 valid cases out of 1,354 total cases.

Location: 4510-4513 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM27_SUSYMC: S9 SubUse: Specify dependency [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	13	1.0 %
2	Drugs	34	2.5 %
3	Both	8	0.6 %

Value	Label	Unweighted Frequency	%
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-107	Follow-up skipped because gate <> 1	1111	82.1 %
	Total	**1,354**	**100%**

Based upon 55 valid cases out of 1,354 total cases.

Location: 4514-4517 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM28_SUSYMB: S9 SubUse: Ever talked doctor/clergy/guidance counselor drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	20	1.5 %
5	No	770	56.9 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-106	Skip-in for alchstat/drugstat	379	28.0 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 790 valid cases out of 1,354 total cases.

Location: 4518-4521 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM28_SUSYMC: S9 SubUse: Specify talked doctor/clergy/guidance counselor [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	5	0.4 %
2	Drugs	9	0.7 %
3	Both	6	0.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-107	Follow-up skipped because gate <> 1	1146	84.6 %

Value	Label	Unweighted Frequency	%
	Total	**1,354**	**100%**

Based upon 20 valid cases out of 1,354 total cases.

Location: 4522-4525 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM29_SUSYMB: S9 SubUse: Ever attended self-help group for drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	62	4.6 %
5	No	728	53.8 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-106	Skip-in for alchstat/drugstat	379	28.0 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 790 valid cases out of 1,354 total cases.

Location: 4526-4529 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM29_SUSYMC: S9 SubUse: Specify reason for self-help [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	25	1.8 %
2	Drugs	14	1.0 %
3	Both	23	1.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-107	Follow-up skipped because gate <> 1	1104	81.5 %
	Total	**1,354**	**100%**

Based upon 62 valid cases out of 1,354 total cases.

Location: 4530-4533 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM30_SUSYMB: S9 SubUse: Ever treated by therapist because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	21	1.6 %
5	No	769	56.8 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-106	Skip-in for alchstat/drugstat	379	28.0 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 790 valid cases out of 1,354 total cases.

Location: 4534-4537 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM30_SUSYMC: S9 SubUse: Specify reason for therapist visit [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	6	0.4 %
2	Drugs	8	0.6 %
3	Both	7	0.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-107	Follow-up skipped because gate <> 1	1145	84.6 %
	Total	**1,354**	**100%**

Based upon 21 valid cases out of 1,354 total cases.

Location: 4538-4541 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM31_SUSYMB: S9 SubUse: Ever Admitted to hospital because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	11	0.8 %
5	No	779	57.5 %
	Missing Data		

Value	Label	Unweighted Frequency	%
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-106	Skip-in for alchstat/drugstat	379	28.0 %
-9	Dont know	1	0.1 %
	Total	**1,354**	**100%**

Based upon 790 valid cases out of 1,354 total cases.

Location: 4542-4545 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE_SUBSYM31_SUSYMC: S9 SubUse: Specify reason admitted to hospital [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	5	0.4 %
2	Drugs	3	0.2 %
3	Both	3	0.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	4	0.3 %
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
-107	Follow-up skipped because gate <> 1	1155	85.3 %
	Total	**1,354**	**100%**

Based upon 11 valid cases out of 1,354 total cases.

Location: 4546-4549 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

S9SUBUSE28: S9 SubUse: Num times abused prescribed medications (own or someone elses) during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	0	0.0 %
2	1-5 times	0	0.0 %
3	6-11 times	0	0.0 %
4	1X per month	0	0.0 %
5	2-3X per month	0	0.0 %
6	1X per week	0	0.0 %
7	2-3X per week	0	0.0 %
8	4-5X per week	0	0.0 %

Value	Label	Unweighted Frequency	%
9	Every day	0	0.0 %
	Missing Data		
-800	Question not asked for this subject	1170	86.4 %
-201	IntCompStat = 4 missed interview	175	12.9 %
-110	Skipped: Interview was conducted in federal facility	9	0.7 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 4550-4553 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

S9QFBRMO: S9 SubUse: Quant/freq used beer/wine/liquor - recall period

Based upon 1,167 valid cases out of 1,354 total cases.

- Mean: 15.79
- Median: 4.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 336.00
- Standard Deviation: 28.66

Location: 4554-4557 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -300 , -202 , -201 , -100 , -9 , -8

S96MOUSE: S9 SubUse: Number drugs used - recall period

Based upon 1,169 valid cases out of 1,354 total cases.

- Mean: 0.61
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 9.00
- Standard Deviation: 0.99

Location: 4558-4561 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -300 , -202 , -201 , -100 , -9 , -8

S9QFSMK6: S9 SubUse: Quant/freq used cigarettes - recall period

Based upon 1,168 valid cases out of 1,354 total cases.

- Mean: 40.11
- Median: 15.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 160.00

- Standard Deviation: 51.64

Location: 4562-4567 (width: 6; decimal: 1)
Variable Type: numeric
(Range of) Missing Values: -300.0 , -202.0 , -201.0 , -100.0 , -9.0 , -8.0

S9A6SC2: S9 SubUse: Recall period alcohol consequences

Based upon 1,179 valid cases out of 1,354 total cases.

- Mean: 0.62
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 17.00
- Standard Deviation: 1.63

Location: 4568-4571 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S9D6SC2: S9 SubUse: Recall period drug consequences

Based upon 1,179 valid cases out of 1,354 total cases.

- Mean: 0.50
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 11.00
- Standard Deviation: 1.50

Location: 4572-4575 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S9A6DP2: S9 SubUse: Recall period alcohol dependence symptoms

Based upon 1,179 valid cases out of 1,354 total cases.

- Mean: 0.27
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 10.00
- Standard Deviation: 1.10

Location: 4576-4579 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S9D6DP2: S9 SubUse: Recall period drug dependence symptoms

Based upon 1,179 valid cases out of 1,354 total cases.

- Mean: 0.41
- Median: 0.00

- Mode: 0.00
- Minimum: 0.00
- Maximum: 10.00
- Standard Deviation: 1.39

Location: 4580-4583 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S9TOTSC6: S9 SubUse: Recall period total substance use consequences

Based upon 1,179 valid cases out of 1,354 total cases.

- Mean: 0.92
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 17.00
- Standard Deviation: 2.01

Location: 4584-4587 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S9TOTDS6: S9 SubUse: Recall period total substance use dependency symp

Based upon 1,179 valid cases out of 1,354 total cases.

- Mean: 0.56
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 10.00
- Standard Deviation: 1.62

Location: 4588-4591 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S9ALTRTM: S9 SubUse: Recall period alcohol treatment

Value	Label	Unweighted Frequency	%
0	No	1122	82.9 %
1	Yes	57	4.2 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
	Total	**1,354**	**100%**

Based upon 1,179 valid cases out of 1,354 total cases.

Location: 4592-4595 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S9DGTRTM: S9 SubUse: Recall period drug treatment

Value	Label	Unweighted Frequency	%
0	No	1129	83.4 %
1	Yes	50	3.7 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
	Total	**1,354**	**100%**

Based upon 1,179 valid cases out of 1,354 total cases.

Location: 4596-4599 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

S9TRTM: S9 SubUse: Recall period alcohol or drug treatment

Value	Label	Unweighted Frequency	%
0	No	1102	81.4 %
1	Yes	77	5.7 %
	Missing Data		
-201	IntCompStat = 4 missed interview	175	12.9 %
	Total	**1,354**	**100%**

Based upon 1,179 valid cases out of 1,354 total cases.

Location: 4600-4603 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

SASUBUSE1: SA SubUse: How often had alcohol to drink in recall period

Value	Label	Unweighted Frequency	%
1	Not at all	426	31.5 %
2	1-5 times	153	11.3 %
3	6-11 times	61	4.5 %
4	1X per month	65	4.8 %
5	2-3X per month	119	8.8 %
6	1X per week	72	5.3 %
7	2-3X per week	157	11.6 %
8	4-5X per week	40	3.0 %
9	Every day	31	2.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %

Value	Label	Unweighted Frequency	%
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
	Total	**1,354**	**100%**

Based upon 1,124 valid cases out of 1,354 total cases.

Location: 4604-4607 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

SASUBUSE2: SA SubUse: Num alcoholic drinks usually consumed

Based upon 1,124 valid cases out of 1,354 total cases.

- Mean: 3.72
- Median: 2.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 60.00
- Standard Deviation: 5.30

Location: 4608-4611 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -101 , -99 , -98

SASUBUSE3: SA SubUse: How often had 5+ drinks at one time in recall period

Value	Label	Unweighted Frequency	%
1	Not at all	637	47.0 %
2	1-5 times	147	10.9 %
3	6-11 times	29	2.1 %
4	1X per month	48	3.5 %
5	2-3X per month	57	4.2 %
6	1X per week	51	3.8 %
7	2-3X per week	108	8.0 %
8	4-5X per week	25	1.8 %
9	Every day	22	1.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
	Total	**1,354**	**100%**

Based upon 1,124 valid cases out of 1,354 total cases.

Location: 4612-4615 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

SASUBUSE4: SA SubUse: Num times been drunk on alcohol during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	617	45.6 %
2	1-5 times	234	17.3 %
3	6-11 times	45	3.3 %
4	1X per month	41	3.0 %
5	2-3X per month	54	4.0 %
6	1X per week	44	3.2 %
7	2-3X per week	61	4.5 %
8	4-5X per week	18	1.3 %
9	Every day	10	0.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
	Total	**1,354**	**100%**

Based upon 1,124 valid cases out of 1,354 total cases.

Location: 4616-4619 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

SASUBUSE6: SA SubUse: Num times used marijuana/hashish during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	739	54.6 %
2	1-5 times	82	6.1 %
3	6-11 times	27	2.0 %
4	1X per month	24	1.8 %
5	2-3X per month	41	3.0 %
6	1X per week	20	1.5 %
7	2-3X per week	37	2.7 %
8	4-5X per week	37	2.7 %
9	Every day	117	8.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
	Total	**1,354**	**100%**

Based upon 1,124 valid cases out of 1,354 total cases.

Location: 4620-4623 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

SASUBUSE8: SA SubUse: Num times used sedative/tranquilizers during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1069	79.0 %
2	1-5 times	17	1.3 %
3	6-11 times	5	0.4 %
4	1X per month	4	0.3 %
5	2-3X per month	8	0.6 %
6	1X per week	2	0.1 %
7	2-3X per week	6	0.4 %
8	4-5X per week	3	0.2 %
9	Every day	10	0.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
	Total	**1,354**	**100%**

Based upon 1,124 valid cases out of 1,354 total cases.

Location: 4624-4627 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

SASUBUSE10: SA SubUse: Num times used stimulants/amphetamines during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1086	80.2 %
2	1-5 times	11	0.8 %
3	6-11 times	5	0.4 %
4	1X per month	3	0.2 %
5	2-3X per month	5	0.4 %
6	1X per week	3	0.2 %
7	2-3X per week	1	0.1 %
8	4-5X per week	7	0.5 %
9	Every day	3	0.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %

Value	Label	Unweighted Frequency	%
	Total	**1,354**	**100%**

Based upon 1,124 valid cases out of 1,354 total cases.

Location: 4628-4631 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

SASUBUSE12: SA SubUse: Num times used cocaine during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1075	79.4 %
2	1-5 times	24	1.8 %
3	6-11 times	7	0.5 %
4	1X per month	4	0.3 %
5	2-3X per month	5	0.4 %
6	1X per week	3	0.2 %
7	2-3X per week	2	0.1 %
8	4-5X per week	2	0.1 %
9	Every day	2	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
	Total	**1,354**	**100%**

Based upon 1,124 valid cases out of 1,354 total cases.

Location: 4632-4635 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

SASUBUSE14: SA SubUse: Num times used opiates during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1073	79.2 %
2	1-5 times	17	1.3 %
3	6-11 times	1	0.1 %
4	1X per month	4	0.3 %
5	2-3X per month	9	0.7 %
6	1X per week	4	0.3 %
7	2-3X per week	5	0.4 %
8	4-5X per week	4	0.3 %
9	Every day	7	0.5 %

Value	Label	Unweighted Frequency	%
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
	Total	**1,354**	**100%**

Based upon 1,124 valid cases out of 1,354 total cases.

Location: 4636-4639 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

SASUBUSE16: SA SubUse: Num times used ecstasy during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1088	80.4 %
2	1-5 times	19	1.4 %
3	6-11 times	4	0.3 %
4	1X per month	2	0.1 %
5	2-3X per month	7	0.5 %
6	1X per week	0	0.0 %
7	2-3X per week	0	0.0 %
8	4-5X per week	2	0.1 %
9	Every day	2	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
	Total	**1,354**	**100%**

Based upon 1,124 valid cases out of 1,354 total cases.

Location: 4640-4643 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

SASUBUSE18: SA SubUse: Num times used hallucinogens during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1089	80.4 %
2	1-5 times	17	1.3 %
3	6-11 times	0	0.0 %
4	1X per month	4	0.3 %
5	2-3X per month	4	0.3 %

Value	Label	Unweighted Frequency	%
6	1X per week	4	0.3 %
7	2-3X per week	1	0.1 %
8	4-5X per week	2	0.1 %
9	Every day	3	0.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
	Total	**1,354**	**100%**

Based upon 1,124 valid cases out of 1,354 total cases.

Location: 4644-4647 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

SASUBUSE20: SA SubUse: Num times used inhalants during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1116	82.4 %
2	1-5 times	1	0.1 %
3	6-11 times	2	0.1 %
4	1X per month	0	0.0 %
5	2-3X per month	4	0.3 %
6	1X per week	0	0.0 %
7	2-3X per week	0	0.0 %
8	4-5X per week	0	0.0 %
9	Every day	1	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
	Total	**1,354**	**100%**

Based upon 1,124 valid cases out of 1,354 total cases.

Location: 4648-4651 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

SASUBUSE22: SA SubUse: Num times used amyl nitrate/odorizers/rush during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1116	82.4 %

Value	Label	Unweighted Frequency	%
2	1-5 times	2	0.1 %
3	6-11 times	0	0.0 %
4	1X per month	0	0.0 %
5	2-3X per month	5	0.4 %
6	1X per week	0	0.0 %
7	2-3X per week	0	0.0 %
8	4-5X per week	0	0.0 %
9	Every day	1	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
	Total	**1,354**	**100%**

Based upon 1,124 valid cases out of 1,354 total cases.

Location: 4652-4655 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

SASUBUSE24: SA SubUse: Used other drugs during recall period [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	1	0.1 %
5	No	1123	82.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
	Total	**1,354**	**100%**

Based upon 1,124 valid cases out of 1,354 total cases.

Location: 4656-4659 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE26: SA SubUse: Num times used other drugs during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	1123	82.9 %
2	1-5 times	0	0.0 %
3	6-11 times	1	0.1 %
4	1X per month	0	0.0 %

Value	Label	Unweighted Frequency	%
5	2-3X per month	0	0.0 %
6	1X per week	0	0.0 %
7	2-3X per week	0	0.0 %
8	4-5X per week	0	0.0 %
9	Every day	0	0.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
	Total	**1,354**	**100%**

Based upon 1,124 valid cases out of 1,354 total cases.

Location: 4660-4663 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

SASUBUSEDY: SA SubUse: Used any drugs past 24hrs [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	91	6.7 %
5	No	1033	76.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
	Total	**1,354**	**100%**

Based upon 1,124 valid cases out of 1,354 total cases.

Location: 4664-4667 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_DRUG24MAR: SA SubUse_Drug24: Used marijuana/hashish past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	90	6.6 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-104	Skip-in for SUBUSEDY 1	4	0.3 %
-103	Skip-in for SUBUSEDY <> 1	1033	76.3 %

Value	Label	Unweighted Frequency	%
	Total	**1,354**	**100%**

Based upon 90 valid cases out of 1,354 total cases.

Location: 4668-4671 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

SASUBUSE_DRUG24TRANQ: SA SubUse_Drug24: Used sedatives/tranquilizers past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	2	0.1 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-104	Skip-in for SUBUSEDY 1	92	6.8 %
-103	Skip-in for SUBUSEDY <> 1	1033	76.3 %
	Total	**1,354**	**100%**

Based upon 2 valid cases out of 1,354 total cases.

Location: 4672-4675 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

SASUBUSE_DRUG24AMPH: SA SubUse_Drug24: Used stimulants/amphetamines past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	1	0.1 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-104	Skip-in for SUBUSEDY 1	93	6.9 %
-103	Skip-in for SUBUSEDY <> 1	1033	76.3 %
	Total	**1,354**	**100%**

Based upon 1 valid cases out of 1,354 total cases.

Location: 4676-4679 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

SASUBUSE_DRUG24COC: SA SubUse_Drug24: Used Cocaine past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	1	0.1 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-104	Skip-in for SUBUSEDY 1	93	6.9 %
-103	Skip-in for SUBUSEDY <> 1	1033	76.3 %
	Total	**1,354**	**100%**

Based upon 1 valid cases out of 1,354 total cases.

Location: 4680-4683 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

SASUBUSE_DRUG24OPI: SA SubUse_Drug24: Used Opiates past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	2	0.1 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-104	Skip-in for SUBUSEDY 1	92	6.8 %
-103	Skip-in for SUBUSEDY <> 1	1033	76.3 %
	Total	**1,354**	**100%**

Based upon 2 valid cases out of 1,354 total cases.

Location: 4684-4687 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

SASUBUSE_DRUG24ECST: SA SubUse_Drug24: Used Ecstacy past 24 hours

Value	Label	Unweighted Frequency	%
1	Yes	1	0.1 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-104	Skip-in for SUBUSEDY 1	93	6.9 %
-103	Skip-in for SUBUSEDY <> 1	1033	76.3 %
	Total	**1,354**	**100%**

Based upon 1 valid cases out of 1,354 total cases.

Location: 4688-4691 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

SASUBUSE_DRUG24HALL: SA SubUse_Drug24: Used Hallucinogens past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-104	Skip-in for SUBUSEDY 1	94	6.9 %
-103	Skip-in for SUBUSEDY <> 1	1033	76.3 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 4692-4695 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

SASUBUSE_DRUG24INH: SA SubUse_Drug24: Used Inhalants past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-104	Skip-in for SUBUSEDY 1	94	6.9 %
-103	Skip-in for SUBUSEDY <> 1	1033	76.3 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 4696-4699 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

SASUBUSE_DRUG24AMYL: SA SubUse_Drug24: Used Amyl nitrate/poppers/odorizers 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %

Value	Label	Unweighted Frequency	%
	Missing Data		
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-104	Skip-in for SUBUSEDY 1	94	6.9 %
-103	Skip-in for SUBUSEDY <> 1	1033	76.3 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 4700-4703 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

SASUBUSE_DRUG24OTHER: SA SubUse_Drug24: Used Other (specify) past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-104	Skip-in for SUBUSEDY 1	94	6.9 %
-103	Skip-in for SUBUSEDY <> 1	1033	76.3 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 4704-4707 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

SASUBUSE_DRUG24NONE: SA SubUse_Drug24: Used NA (No drugs) past 24hrs

Value	Label	Unweighted Frequency	%
1	Yes	0	0.0 %
5	No	0	0.0 %
	Missing Data		
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-104	Skip-in for SUBUSEDY 1	94	6.9 %
-103	Skip-in for SUBUSEDY <> 1	1033	76.3 %
	Total	**1,354**	**100%**

Based upon 0 valid cases out of 1,354 total cases.

Location: 4708-4711 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -106 , -104 , -103 , -9 , -8

SASUBUSE29: SA SubUse: Specify other drug used past 24hrs

Based upon 1,354 valid cases out of 1,354 total cases.

Location: 4712-4712 (width: 1; decimal: 0)
Variable Type: character

SASUBUSE30: SA SubUse: Num times smoked cigarettes during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	489	36.1 %
2	1-5 times	14	1.0 %
3	6-11 times	10	0.7 %
4	1X per month	6	0.4 %
5	2-3X per month	20	1.5 %
6	1X per week	13	1.0 %
7	2-3X per week	29	2.1 %
8	4-5X per week	24	1.8 %
9	Every day	519	38.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
	Total	**1,354**	**100%**

Based upon 1,124 valid cases out of 1,354 total cases.

Location: 4713-4716 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

SASUBUSE31: SA SubUse: Num cigarettes smoke per day during recall period

Value	Label	Unweighted Frequency	%
1	0 cigarettes	514	38.0 %
2	1-4 cigarettes	199	14.7 %
3	5-9 cigarettes	165	12.2 %
4	10-15 cigarettes	127	9.4 %
5	16-20 cigarettes	87	6.4 %
6	More than a pack a day cigarettes	32	2.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %

Value	Label	Unweighted Frequency	%
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
	Total	**1,354**	**100%**

Based upon 1,124 valid cases out of 1,354 total cases.

Location: 4717-4720 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -105 , -9 , -8

SASUBUSE_SUBSYM01_SUSYMB: SA SubUse: Family ever complain because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	123	9.1 %
5	No	628	46.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-106	Skip-in for alchstat/drugstat	373	27.5 %
	Total	**1,354**	**100%**

Based upon 751 valid cases out of 1,354 total cases.

Location: 4721-4724 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM01_SUSYMC: SA SubUse: Specify cause of family complaint [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	55	4.1 %
2	Drugs	42	3.1 %
3	Both	26	1.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-107	Follow-up skipped because gate <> 1	1001	73.9 %
	Total	**1,354**	**100%**

Based upon 123 valid cases out of 1,354 total cases.

Location: 4725-4728 (width: 4; decimal: 0)

Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM02_SUSYMB: SA SubUse: Friend ever complain because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	51	3.8 %
5	No	700	51.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-106	Skip-in for alchstat/drugstat	373	27.5 %
	Total	**1,354**	**100%**

Based upon 751 valid cases out of 1,354 total cases.

Location: 4729-4732 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM02_SUSYMC: SA SubUse: Specify cause of friend complain [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	17	1.3 %
2	Drugs	22	1.6 %
3	Both	12	0.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-107	Follow-up skipped because gate <> 1	1073	79.2 %
	Total	**1,354**	**100%**

Based upon 51 valid cases out of 1,354 total cases.

Location: 4733-4736 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM03_SUSYMB: SA SubUse: Ever trouble at work because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	17	1.3 %

Value	Label	Unweighted Frequency	%
5	No	734	54.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-106	Skip-in for alchstat/drugstat	373	27.5 %
	Total	**1,354**	**100%**

Based upon 751 valid cases out of 1,354 total cases.

Location: 4737-4740 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM03_SUSYMC: SA SubUse: Specify cause of trouble at work [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	10	0.7 %
2	Drugs	6	0.4 %
3	Both	1	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-107	Follow-up skipped because gate <> 1	1107	81.8 %
	Total	**1,354**	**100%**

Based upon 17 valid cases out of 1,354 total cases.

Location: 4741-4744 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM04_SUSYMB: SA SubUse: Ever missed work/school because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	35	2.6 %
5	No	716	52.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %

Value	Label	Unweighted Frequency	%
-106	Skip-in for alchstat/drugstat	373	27.5 %
	Total	**1,354**	**100%**

Based upon 751 valid cases out of 1,354 total cases.

Location: 4745-4748 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM04_SUSYMC: SA SubUse: Specify cause of missed work/school [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	23	1.7 %
2	Drugs	7	0.5 %
3	Both	5	0.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-107	Follow-up skipped because gate <> 1	1089	80.4 %
	Total	**1,354**	**100%**

Based upon 35 valid cases out of 1,354 total cases.

Location: 4749-4752 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM05_SUSYMB: SA SubUse: Ever arrested because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	61	4.5 %
5	No	690	51.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-106	Skip-in for alchstat/drugstat	373	27.5 %
	Total	**1,354**	**100%**

Based upon 751 valid cases out of 1,354 total cases.

Location: 4753-4756 (width: 4; decimal: 0)
Variable Type: numeric

(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM05_SUSYMC: SA SubUse: Specify cause of arrest [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	22	1.6 %
2	Drugs	31	2.3 %
3	Both	8	0.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-107	Follow-up skipped because gate <> 1	1063	78.5 %
	Total	**1,354**	**100%**

Based upon 61 valid cases out of 1,354 total cases.

Location: 4757-4760 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM06_SUSYMB: SA SubUse: Ever argue family/friends because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	92	6.8 %
5	No	659	48.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-106	Skip-in for alchstat/drugstat	373	27.5 %
	Total	**1,354**	**100%**

Based upon 751 valid cases out of 1,354 total cases.

Location: 4761-4764 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM06_SUSYMC: SA SubUse: Specify cause of argument w/ family/friends [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	42	3.1 %
2	Drugs	26	1.9 %

Value	Label	Unweighted Frequency	%
3	Both	24	1.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-107	Follow-up skipped because gate <> 1	1032	76.2 %
	Total	**1,354**	**100%**

Based upon 92 valid cases out of 1,354 total cases.

Location: 4765-4768 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM07_SUSYMB: SA SubUse: Ever cont. drug/alcohol use in spite cause health problems in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	73	5.4 %
5	No	678	50.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-106	Skip-in for alchstat/drugstat	373	27.5 %
	Total	**1,354**	**100%**

Based upon 751 valid cases out of 1,354 total cases.

Location: 4769-4772 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM07_SUSYMC: SA SubUse: Specify continued using [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	26	1.9 %
2	Drugs	29	2.1 %
3	Both	18	1.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-107	Follow-up skipped because gate <> 1	1051	77.6 %

Value	Label	Unweighted Frequency	%
	Total	**1,354**	**100%**

Based upon 73 valid cases out of 1,354 total cases.

Location: 4773-4776 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM08_SUSYMB: SA SubUse: Ever not remember prior night because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	133	9.8 %
5	No	618	45.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-106	Skip-in for alchstat/drugstat	373	27.5 %
	Total	**1,354**	**100%**

Based upon 751 valid cases out of 1,354 total cases.

Location: 4777-4780 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM08_SUSYMC: SA SubUse: Specify cause of not remember night [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	86	6.4 %
2	Drugs	15	1.1 %
3	Both	32	2.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-107	Follow-up skipped because gate <> 1	991	73.2 %
	Total	**1,354**	**100%**

Based upon 133 valid cases out of 1,354 total cases.

Location: 4781-4784 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM09_SUSYMB: SA SubUse: Ever used in situations could get hurt in recall period [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	126	9.3 %
5	No	625	46.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-106	Skip-in for alchstat/drugstat	373	27.5 %
	Total	**1,354**	**100%**

Based upon 751 valid cases out of 1,354 total cases.

Location: 4785-4788 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM09_SUSYMC: SA SubUse: Specify used drugs in hurtful situations [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	60	4.4 %
2	Drugs	26	1.9 %
3	Both	40	3.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-107	Follow-up skipped because gate <> 1	998	73.7 %
	Total	**1,354**	**100%**

Based upon 126 valid cases out of 1,354 total cases.

Location: 4789-4792 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM10_SUSYMB: SA SubUse: Ever had accident/injury because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	36	2.7 %
5	No	715	52.8 %
	Missing Data		

Value	Label	Unweighted Frequency	%
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-106	Skip-in for alchstat/drugstat	373	27.5 %
	Total	**1,354**	**100%**

Based upon 751 valid cases out of 1,354 total cases.

Location: 4793-4796 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM10_SUSYMC: SA SubUse: Specify cause of accident/injury [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	25	1.8 %
2	Drugs	4	0.3 %
3	Both	7	0.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-107	Follow-up skipped because gate <> 1	1088	80.4 %
	Total	**1,354**	**100%**

Based upon 36 valid cases out of 1,354 total cases.

Location: 4797-4800 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM11_SUSYMB: SA SubUse: Alcohol/drug use interfere work/responsibility school/home/job in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	49	3.6 %
5	No	702	51.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-106	Skip-in for alchstat/drugstat	373	27.5 %
	Total	**1,354**	**100%**

Based upon 751 valid cases out of 1,354 total cases.

Location: 4801-4804 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM11_SUSYMC: SA SubUse: Specify cause of interference [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	19	1.4 %
2	Drugs	21	1.6 %
3	Both	9	0.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-107	Follow-up skipped because gate <> 1	1075	79.4 %
	Total	**1,354**	**100%**

Based upon 49 valid cases out of 1,354 total cases.

Location: 4805-4808 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM12_SUSYMB: SA SubUse: Ever trouble police no arrest because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	28	2.1 %
5	No	723	53.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-106	Skip-in for alchstat/drugstat	373	27.5 %
	Total	**1,354**	**100%**

Based upon 751 valid cases out of 1,354 total cases.

Location: 4809-4812 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM12_SUSYMC: SA SubUse: Specify cause trouble police no arrest [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	14	1.0 %
2	Drugs	6	0.4 %
3	Both	8	0.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-107	Follow-up skipped because gate <> 1	1096	80.9 %
	Total	**1,354**	**100%**

Based upon 28 valid cases out of 1,354 total cases.

Location: 4813-4816 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM13_SUSYMB: SA SubUse: Sold/exchanged sex for drugs in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	4	0.3 %
5	No	747	55.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-106	Skip-in for alchstat/drugstat	373	27.5 %
	Total	**1,354**	**100%**

Based upon 751 valid cases out of 1,354 total cases.

Location: 4817-4820 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM13_SUSYMC: SA SubUse: Specify cause sold/exchanged sex [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	1	0.1 %
2	Drugs	3	0.2 %
3	Both	0	0.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %

Value	Label	Unweighted Frequency	%
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-107	Follow-up skipped because gate <> 1	1120	82.7 %
	Total	**1,354**	**100%**

Based upon 4 valid cases out of 1,354 total cases.

Location: 4821-4824 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM14_SUSYMB: SA SubUse: Sold drugs for money buy drugs in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	39	2.9 %
5	No	712	52.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-106	Skip-in for alchstat/drugstat	373	27.5 %
	Total	**1,354**	**100%**

Based upon 751 valid cases out of 1,354 total cases.

Location: 4825-4828 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM14_SUSYMC: SA SubUse: Specify cause sold drugs for money to buy [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	0	0.0 %
2	Drugs	36	2.7 %
3	Both	3	0.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-107	Follow-up skipped because gate <> 1	1085	80.1 %
	Total	**1,354**	**100%**

Based upon 39 valid cases out of 1,354 total cases.

Location: 4829-4832 (width: 4; decimal: 0)
Variable Type: numeric

(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM15_SUSYMB: SA SubUse: Stole for money buy drugs in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	11	0.8 %
5	No	740	54.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-106	Skip-in for alchstat/drugstat	373	27.5 %
	Total	**1,354**	**100%**

Based upon 751 valid cases out of 1,354 total cases.

Location: 4833-4836 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM15_SUSYMC: SA SubUse: Specify cause stole for money to buy [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	0	0.0 %
2	Drugs	11	0.8 %
3	Both	0	0.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-107	Follow-up skipped because gate <> 1	1113	82.2 %
	Total	**1,354**	**100%**

Based upon 11 valid cases out of 1,354 total cases.

Location: 4837-4840 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM16_SUSYMB: SA SubUse: Ever had physical fight because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	46	3.4 %
5	No	705	52.1 %

Value	Label	Unweighted Frequency	%
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-106	Skip-in for alchstat/drugstat	373	27.5 %
	Total	**1,354**	**100%**

Based upon 751 valid cases out of 1,354 total cases.

Location: 4841-4844 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM16_SUSYMC: SA SubUse: Specify cause of fight [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	37	2.7 %
2	Drugs	4	0.3 %
3	Both	5	0.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-107	Follow-up skipped because gate <> 1	1078	79.6 %
	Total	**1,354**	**100%**

Based upon 46 valid cases out of 1,354 total cases.

Location: 4845-4848 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM17_SUSYMB: SA SubUse: Ever destroy property because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	25	1.8 %
5	No	726	53.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-106	Skip-in for alchstat/drugstat	373	27.5 %
	Total	**1,354**	**100%**

Based upon 751 valid cases out of 1,354 total cases.

Location: 4849-4852 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM17_SUSYMC: SA SubUse: Specify cause of destroy property [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	16	1.2 %
2	Drugs	6	0.4 %
3	Both	3	0.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-107	Follow-up skipped because gate <> 1	1099	81.2 %
	Total	**1,354**	**100%**

Based upon 25 valid cases out of 1,354 total cases.

Location: 4853-4856 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM18_SUSYMB: SA SubUse: Ever felt strong urge use drug/alcohol no self stop in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	52	3.8 %
5	No	699	51.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-106	Skip-in for alchstat/drugstat	373	27.5 %
	Total	**1,354**	**100%**

Based upon 751 valid cases out of 1,354 total cases.

Location: 4857-4860 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM18_SUSYMC: SA SubUse: Specify the cause of strong urge [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	18	1.3 %
2	Drugs	25	1.8 %
3	Both	9	0.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-107	Follow-up skipped because gate <> 1	1072	79.2 %
	Total	**1,354**	**100%**

Based upon 52 valid cases out of 1,354 total cases.

Location: 4861-4864 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM19_SUSYMB: SA SubUse: Ever want drug/alcohol think of nothing else in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	49	3.6 %
5	No	702	51.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-106	Skip-in for alchstat/drugstat	373	27.5 %
	Total	**1,354**	**100%**

Based upon 751 valid cases out of 1,354 total cases.

Location: 4865-4868 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM19_SUSYMC: SA SubUse: Specify cause of want [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	13	1.0 %
2	Drugs	25	1.8 %
3	Both	11	0.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %

Value	Label	Unweighted Frequency	%
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-107	Follow-up skipped because gate <> 1	1075	79.4 %
	Total	**1,354**	**100%**

Based upon 49 valid cases out of 1,354 total cases.

Location: 4869-4872 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM20_SUSYMB: SA SubUse: Ever experience withdrawal from drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	43	3.2 %
5	No	708	52.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-106	Skip-in for alchstat/drugstat	373	27.5 %
	Total	**1,354**	**100%**

Based upon 751 valid cases out of 1,354 total cases.

Location: 4873-4876 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM20_SUSYMC: SA SubUse: Specify cause of withdraw [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	13	1.0 %
2	Drugs	24	1.8 %
3	Both	6	0.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-107	Follow-up skipped because gate <> 1	1081	79.8 %
	Total	**1,354**	**100%**

Based upon 43 valid cases out of 1,354 total cases.

Location: 4877-4880 (width: 4; decimal: 0)
Variable Type: numeric

(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM21_SUSYMB: SA SubUse: Take drug/alcohol to keep from having withdraw in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	33	2.4 %
5	No	10	0.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-108	Skip-in for SUBSYM20=5	708	52.3 %
-106	Skip-in for alchstat/drugstat	373	27.5 %
	Total	**1,354**	**100%**

Based upon 43 valid cases out of 1,354 total cases.

Location: 4881-4884 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM21_SUSYMC: SA SubUse: Specify reason for consumption [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	10	0.7 %
2	Drugs	18	1.3 %
3	Both	5	0.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-107	Follow-up skipped because gate <> 1	1091	80.6 %
	Total	**1,354**	**100%**

Based upon 33 valid cases out of 1,354 total cases.

Location: 4885-4888 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM22_SUSYMB: SA SubUse: Give up usual activities/interests to use drug/alcohol in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	42	3.1 %
5	No	709	52.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-106	Skip-in for alchstat/drugstat	373	27.5 %
	Total	**1,354**	**100%**

Based upon 751 valid cases out of 1,354 total cases.

Location: 4889-4892 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM22_SUSYMC: SA SubUse: Specify cause of giving up activities/interests [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	14	1.0 %
2	Drugs	18	1.3 %
3	Both	10	0.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-107	Follow-up skipped because gate <> 1	1082	79.9 %
	Total	**1,354**	**100%**

Based upon 42 valid cases out of 1,354 total cases.

Location: 4893-4896 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM23_SUSYMB: SA SubUse: Ever used in rp when promised wouldn't [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	95	7.0 %
5	No	656	48.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %

Value	Label	Unweighted Frequency	%
-106	Skip-in for alchstat/drugstat	373	27.5 %
	Total	**1,354**	**100%**

Based upon 751 valid cases out of 1,354 total cases.

Location: 4897-4900 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM23_SUSYMC: SA SubUse: Specify what use you started [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	37	2.7 %
2	Drugs	36	2.7 %
3	Both	22	1.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-107	Follow-up skipped because gate <> 1	1029	76.0 %
	Total	**1,354**	**100%**

Based upon 95 valid cases out of 1,354 total cases.

Location: 4901-4904 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM24_SUSYMB: SA SubUse: Ever use drug/alcohol more freq than intended in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	103	7.6 %
5	No	648	47.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-106	Skip-in for alchstat/drugstat	373	27.5 %
	Total	**1,354**	**100%**

Based upon 751 valid cases out of 1,354 total cases.

Location: 4905-4908 (width: 4; decimal: 0)
Variable Type: numeric

(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM24_SUSYMC: SA SubUse: Specify what you used more frequently [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	42	3.1 %
2	Drugs	37	2.7 %
3	Both	24	1.8 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-107	Follow-up skipped because gate <> 1	1021	75.4 %
	Total	**1,354**	**100%**

Based upon 103 valid cases out of 1,354 total cases.

Location: 4909-4912 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM25_SUSYMB: SA SubUse: Ever tried cut down drug/alcohol use but couldnt in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	67	4.9 %
5	No	684	50.5 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-106	Skip-in for alchstat/drugstat	373	27.5 %
	Total	**1,354**	**100%**

Based upon 751 valid cases out of 1,354 total cases.

Location: 4913-4916 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM25_SUSYMC: SA SubUse: Specify what substance use you tried to curtail [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	20	1.5 %

Value	Label	Unweighted Frequency	%
2	Drugs	34	2.5 %
3	Both	13	1.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-107	Follow-up skipped because gate <> 1	1057	78.1 %
	Total	**1,354**	**100%**

Based upon 67 valid cases out of 1,354 total cases.

Location: 4917-4920 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM26_SUSYMB: SA SubUse: Ever need more drug/alcohol get same effect in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	109	8.1 %
5	No	642	47.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-106	Skip-in for alchstat/drugstat	373	27.5 %
	Total	**1,354**	**100%**

Based upon 751 valid cases out of 1,354 total cases.

Location: 4921-4924 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM26_SUSYMC: SA SubUse: Specify substance need more for same effect [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	44	3.2 %
2	Drugs	47	3.5 %
3	Both	18	1.3 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %

Value	Label	Unweighted Frequency	%
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-107	Follow-up skipped because gate <> 1	1015	75.0 %
	Total	**1,354**	**100%**

Based upon 109 valid cases out of 1,354 total cases.

Location: 4925-4928 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM27_SUSYMB: SA SubUse: Ever used drug/alcohol enough to depend in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	52	3.8 %
5	No	699	51.6 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-106	Skip-in for alchstat/drugstat	373	27.5 %
	Total	**1,354**	**100%**

Based upon 751 valid cases out of 1,354 total cases.

Location: 4929-4932 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM27_SUSYMC: SA SubUse: Specify dependency [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	15	1.1 %
2	Drugs	31	2.3 %
3	Both	6	0.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-107	Follow-up skipped because gate <> 1	1072	79.2 %
	Total	**1,354**	**100%**

Based upon 52 valid cases out of 1,354 total cases.

Location: 4933-4936 (width: 4; decimal: 0)
Variable Type: numeric

(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM28_SUSYMB: SA SubUse: Ever talked doctor/clergy/guidance counselor drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	35	2.6 %
5	No	716	52.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-106	Skip-in for alchstat/drugstat	373	27.5 %
	Total	**1,354**	**100%**

Based upon 751 valid cases out of 1,354 total cases.

Location: 4937-4940 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM28_SUSYMC: SA SubUse: Specify talked doctor/clergy/guidance counselor [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	9	0.7 %
2	Drugs	17	1.3 %
3	Both	9	0.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-107	Follow-up skipped because gate <> 1	1089	80.4 %
	Total	**1,354**	**100%**

Based upon 35 valid cases out of 1,354 total cases.

Location: 4941-4944 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM29_SUSYMB: SA SubUse: Ever attended self-help group for drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	55	4.1 %

Value	Label	Unweighted Frequency	%
5	No	696	51.4 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-106	Skip-in for alchstat/drugstat	373	27.5 %
	Total	**1,354**	**100%**

Based upon 751 valid cases out of 1,354 total cases.

Location: 4945-4948 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM29_SUSYMC: SA SubUse: Specify reason for self-help [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	20	1.5 %
2	Drugs	21	1.6 %
3	Both	14	1.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-107	Follow-up skipped because gate <> 1	1069	79.0 %
	Total	**1,354**	**100%**

Based upon 55 valid cases out of 1,354 total cases.

Location: 4949-4952 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM30_SUSYMB: SA SubUse: Ever treated by therapist because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	18	1.3 %
5	No	733	54.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %

Value	Label	Unweighted Frequency	%
-106	Skip-in for alchstat/drugstat	373	27.5 %
	Total	**1,354**	**100%**

Based upon 751 valid cases out of 1,354 total cases.

Location: 4953-4956 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM30_SUSYMC: SA SubUse: Specify reason for therapist visit [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	0	0.0 %
2	Drugs	9	0.7 %
3	Both	9	0.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-107	Follow-up skipped because gate <> 1	1106	81.7 %
	Total	**1,354**	**100%**

Based upon 18 valid cases out of 1,354 total cases.

Location: 4957-4960 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM31_SUSYMB: SA SubUse: Ever Admitted to hospital because drug/alcohol use in rp [Y/N]

Value	Label	Unweighted Frequency	%
1	Yes	7	0.5 %
5	No	744	54.9 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-106	Skip-in for alchstat/drugstat	373	27.5 %
	Total	**1,354**	**100%**

Based upon 751 valid cases out of 1,354 total cases.

Location: 4961-4964 (width: 4; decimal: 0)
Variable Type: numeric

(Range of) Missing Values: -202 , -201 , -110 , -109 , -108 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE_SUBSYM31_SUSYMC: SA SubUse: Specify reason admitted to hospital [drug/alcohol/both]

Value	Label	Unweighted Frequency	%
1	Alcohol	2	0.1 %
2	Drugs	3	0.2 %
3	Both	2	0.1 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
-107	Follow-up skipped because gate <> 1	1117	82.5 %
	Total	**1,354**	**100%**

Based upon 7 valid cases out of 1,354 total cases.

Location: 4965-4968 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201 , -110 , -109 , -107 , -106 , -104 , -103 , -9 , -8

SASUBUSE28: SA SubUse: Num times abused prescribed medications (own or someone elses) during recall period

Value	Label	Unweighted Frequency	%
1	Not at all	0	0.0 %
2	1-5 times	0	0.0 %
3	6-11 times	1	0.1 %
4	1X per month	0	0.0 %
5	2-3X per month	0	0.0 %
6	1X per week	0	0.0 %
7	2-3X per week	0	0.0 %
8	4-5X per week	0	0.0 %
9	Every day	0	0.0 %
	Missing Data		
-800	Question not asked for this subject	1123	82.9 %
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
-110	Skipped: Interview was conducted in federal facility	7	0.5 %
	Total	**1,354**	**100%**

Based upon 1 valid cases out of 1,354 total cases.

Location: 4969-4972 (width: 4; decimal: 0)
Variable Type: numeric

(Range of) Missing Values: -800 , -202 , -201 , -110 , -109 , -102 , -101 , -99 , -98

SAQFBRMO: SA SubUse: Quant/freq used beer/wine/liquor - recall period

Based upon 1,124 valid cases out of 1,354 total cases.

- Mean: 17.37
- Median: 4.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 480.00
- Standard Deviation: 32.85

Location: 4973-4976 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -300 , -202 , -201 , -100 , -9 , -8

SA6MOUSE: SA SubUse: Number drugs used - recall period

Based upon 1,124 valid cases out of 1,354 total cases.

- Mean: 0.59
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 9.00
- Standard Deviation: 1.08

Location: 4977-4980 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -300 , -202 , -201 , -100 , -9 , -8

SAQFSMK6: SA SubUse: Quant/freq used cigarettes - recall period

Based upon 1,124 valid cases out of 1,354 total cases.

- Mean: 37.70
- Median: 15.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 160.00
- Standard Deviation: 49.80

Location: 4981-4986 (width: 6; decimal: 1)
Variable Type: numeric
(Range of) Missing Values: -300.0 , -202.0 , -201.0 , -100.0 , -9.0 , -8.0

SAA6SC2: SA SubUse: Recall period alcohol consequences

Based upon 1,131 valid cases out of 1,354 total cases.

- Mean: 0.58
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 14.00
- Standard Deviation: 1.60

Location: 4987-4990 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

SAD6SC2: SA SubUse: Recall period drug consequences

Based upon 1,131 valid cases out of 1,354 total cases.

- Mean: 0.44
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 13.00
- Standard Deviation: 1.42

Location: 4991-4994 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

SAA6DP2: SA SubUse: Recall period alcohol dependence symptoms

Based upon 1,131 valid cases out of 1,354 total cases.

- Mean: 0.31
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 10.00
- Standard Deviation: 1.23

Location: 4995-4998 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

SAD6DP2: SA SubUse: Recall period drug dependence symptoms

Based upon 1,131 valid cases out of 1,354 total cases.

- Mean: 0.37
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 10.00
- Standard Deviation: 1.41

Location: 4999-5002 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

SATOTSC6: SA SubUse: Recall period total substance use consequences

Based upon 1,131 valid cases out of 1,354 total cases.

- Mean: 0.84
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00

- Maximum: 15.00
- Standard Deviation: 1.97

Location: 5003-5006 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

SATOTDS6: SA SubUse: Recall period total substance use dependency symp

Based upon 1,131 valid cases out of 1,354 total cases.

- Mean: 0.57
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 10.00
- Standard Deviation: 1.70

Location: 5007-5010 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

SAALTRTM: SA SubUse: Recall period alcohol treatment

Value	Label	Unweighted Frequency	%
0	No	1090	80.5 %
1	Yes	41	3.0 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
	Total	**1,354**	**100%**

Based upon 1,131 valid cases out of 1,354 total cases.

Location: 5011-5014 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201

SADGTRTM: SA SubUse: Recall period drug treatment

Value	Label	Unweighted Frequency	%
0	No	1087	80.3 %
1	Yes	44	3.2 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
	Total	**1,354**	**100%**

Based upon 1,131 valid cases out of 1,354 total cases.

Location: 5015-5018 (width: 4; decimal: 0)

Variable Type: numeric
(Range of) Missing Values: -202 , -201

SATRTM: SA SubUse: Recall period alcohol or drug treatment

Value	Label	Unweighted Frequency	%
0	No	1067	78.8 %
1	Yes	64	4.7 %
	Missing Data		
-202	IntCompStat = 2 partial interview, data missing	3	0.2 %
-201	IntCompStat = 4 missed interview	220	16.2 %
	Total	**1,354**	**100%**

Based upon 1,131 valid cases out of 1,354 total cases.

Location: 5019-5022 (width: 4; decimal: 0)
Variable Type: numeric
(Range of) Missing Values: -202 , -201