

# Chi-Square Tests

R Handout

*Derek H. Ogle*

---

## First Commands

```
> library(NCStats)
```

## Chi-Square Test

### From Summarized Observed Table

When [Chinook Salmon](#) (*Oncorhynchus tshawytscha*) were first introduced to Lake Superior there was concern that they would compete with native [Lake Trout](#) (*Salvelinus namaycush*) for [Cisco](#) (*Coregonus artedii*). Preliminarily, fisheries biologists classified the dominant food items (Cisco, [Smelt](#) (*Osmerus mordax*) (another type of fish), or [Mysis](#) (an invertebrate)) in the diets of 50 Lake Trout and 40 Chinook Salmon. They found that 32, 10, and 8 Lake Trout diets were dominated by Cisco, Smelt, and *Mysis*, respectively. Similarly, 18, 18, and 4 Chinook Salmon diets were dominated by Cisco, Smelt, and *Mysis*, respectively. Test (at the 10% level) if the distribution of dominant food items differs at the 5% level?

```
> freq <- c(32,10,8,18,18,4)
> ( obs <- matrix(freq,nrow=2,byrow=TRUE) )
```

```
      [,1] [,2] [,3]
[1,]    32    10     8
[2,]    18    18     4
```

```
> rownames(obs) <- c("Lake Trout","Chinook Salmon")
> colnames(obs) <- c("Cisco","Smelt","Mysis")
> obs
```

```
      Cisco Smelt Mysis
Lake Trout    32    10     8
Chinook Salmon 18    18     4
```

```
> chi1 <- chisq.test(obs,correct=FALSE)
> chi1$expected
```

```
      Cisco  Smelt  Mysis
Lake Trout 27.77778 15.55556 6.666667
Chinook Salmon 22.22222 12.44444 5.333333
```

```
> chi1$expected >= 5
```

	Cisco	Smelt	Mysis
Lake Trout	TRUE	TRUE	TRUE
Chinook Salmon	TRUE	TRUE	TRUE

```
> chi1
```

Pearson's Chi-squared test with obs  
X-squared = 6.5083, df = 2, p-value = 0.03861

```
> chi1$residuals
```

	Cisco	Smelt	Mysis
Lake Trout	0.8011103	-1.408590	0.5163978
Chinook Salmon	-0.8956686	1.574852	-0.5773503

```
> percTable(obs,margin=1,digits=1)
```

	Cisco	Smelt	Mysis	Sum
Lake Trout	64	20	16	100
Chinook Salmon	45	45	10	100

```
> ( obs2 <- obs[,-2] )
```

	Cisco	Mysis
Lake Trout	32	8
Chinook Salmon	18	4

```
> ( chi2 <- chisq.test(obs2,correct=FALSE) )
```

Warning in chisq.test(obs2, correct = FALSE): Chi-squared approximation may be incorrect

Pearson's Chi-squared test with obs2  
X-squared = 0.0301, df = 1, p-value = 0.8624

## From Raw Data

The [General Social Survey](#) (GSS) is a nationwide survey that has been administered since 1972 to gather data on contemporary American society in an attempt to monitor and explain trends in attitudes, behaviors, and attributes of American society. One part of that survey asked respondents to state their opinion on how true the following statement was – “*All radioactivity is made by humans.*” Respondents were also categorized by their highest educational degree. The results from this portion of the GSS is in [SciTest1.csv](#). Use these data to determine, at the 5% level, if the distribution of responses to this statement differs among levels of education.

```
> setwd("C:/aaaWork/Web/GitHub/NCMTH107/modules/Chi_Square")
> ST1 <- read.csv("SciTest1.csv")
> str(ST1)
```

```
'data.frame': 2549 obs. of 2 variables:
 $ degree : Factor w/ 5 levels "bach","grad",...: 5 5 5 5 5 5 5 5 5 ...
 $ scitest: Factor w/ 4 levels "def.not","def.true",...: 2 2 2 2 2 2 2 2 2 ...
```

```
> levels(ST1$degree)
```

```
[1] "bach" "grad" "hs" "jc" "lt.hs"
```

```
> ST1$fdegree <- factor(ST1$degree,levels=c("lt.hs","hs","jc","bach","grad"))
> levels(ST1$scitest)
```

```
[1] "def.not" "def.true" "prob.not" "prob.true"
```

```
> ST1$fscitest <- factor(ST1$scitest,levels=c("def.not","prob.not","prob.true","def.true"))
```

```
> ( freq.tbl <- xtabs(~fdegree+fscitest,data=ST1) )
```

	fscitest			
fdegree	def.not	prob.not	prob.true	def.true
lt.hs	52	112	155	70
hs	366	451	437	114
jc	60	44	36	9
bach	214	135	78	12
grad	123	57	18	6

```
> ST1.chi <- chisq.test(freq.tbl,correct=FALSE)
> ST1.chi$expected
```

	fscitest			
fdegree	def.not	prob.not	prob.true	def.true
lt.hs	124.37623	121.93448	110.48882	32.20047
hs	437.39506	428.80816	388.55708	113.23970
jc	47.64025	46.70498	42.32091	12.33386
bach	140.36289	137.60730	124.69047	36.33935
grad	65.22558	63.94508	57.94272	16.88662

```
> ST1.chi$expected >= 5
```

	fscitest			
fdegree	def.not	prob.not	prob.true	def.true
lt.hs	TRUE	TRUE	TRUE	TRUE
hs	TRUE	TRUE	TRUE	TRUE
jc	TRUE	TRUE	TRUE	TRUE
bach	TRUE	TRUE	TRUE	TRUE
grad	TRUE	TRUE	TRUE	TRUE

```
> all(chi1$expected >= 5)
```

```
[1] TRUE
```

```
> ST1.chi
```

```
Pearson's Chi-squared test with freq.tbl  
X-squared = 288.2331, df = 12, p-value < 2.2e-16
```

```
> ST1.chi$residuals
```

```
      fscitest  
fdegree  def.not  prob.not  prob.true  def.true  
lt.hs   -6.4897392 -0.8996675  4.2345762  6.6612431  
hs      -3.4137460  1.0716718  2.4575530  0.0714471  
jc       1.7906992 -0.3958064 -0.9716327 -0.9492868  
bach     6.2154218 -0.2222643 -4.1813025 -4.0375730  
grad     7.1536326 -0.8685073 -5.2473280 -2.6492428
```

```
> percTable(freq.tbl,margin=1,digits=1)
```

```
      fscitest  
fdegree  def.not  prob.not  prob.true  def.true  Sum  
lt.hs    13.4    28.8    39.8    18.0 100.0  
hs        26.8    33.0    31.9     8.3 100.0  
jc        40.3    29.5    24.2     6.0 100.0  
bach      48.7    30.8    17.8     2.7 100.0  
grad      60.3    27.9     8.8     2.9 99.9
```