

Univariate EDA

R Handout

Derek H. Ogle

Background

Karagas et al. (1996) conducted a pilot study to assess the utility of arsenic concentrations in the toenail as an indicator of ingestion of arsenic-containing water. They interviewed 21 individuals whose household drinking water supply was provided by a private (unregulated) well, including 10 individuals who lived in areas of New Hampshire where elevated water levels of arsenic had been reported previously. Each participant also provided a sample of water and toenail clippings.

The data are recorded in Arsenic.csv. Descriptions of the variables are below.

- **age**: Age (yrs) of person
 - **sex**: Sex of person
 - **usedrink**: How much (fraction of time) the well is used for drinking – A=" $< \frac{1}{4}$ ", B=" $\approx \frac{1}{4}$ ", C=" $\approx \frac{1}{2}$ ", D=" $\approx \frac{3}{4}$ ", E=" $> \frac{3}{4}$ "
 - **usecook**: How much (fraction of time) the well is used for cooking – A=" $< \frac{1}{4}$ ", B=" $\approx \frac{1}{4}$ ", C=" $\approx \frac{1}{2}$ ", D=" $\approx \frac{3}{4}$ ", E=" $> \frac{3}{4}$ "
 - **arswater**: Arsenic in water (ppm)
 - **arsnails**: Arsenic in toenails (ppm)
-

Getting the Data

```
> library(NCStats)
```

```
> setwd("C:/stats/")
```

```
> ars <- read.csv("Arsenic.csv")
```

```
> str(ars)
```

```
'data.frame':  21 obs. of  6 variables:
 $ age      : int  44 45 44 66 37 45 47 38 41 49 ...
 $ sex      : Factor w/ 2 levels "F","M": 1 1 2 1 2 1 2 1 1 1 ...
 $ usedrink: Factor w/ 5 levels "A","B","C","D",...: 5 4 5 3 2 5 5 4 3 4 ...
 $ usecook  : Factor w/ 2 levels "B","E": 2 2 2 2 2 2 2 2 1 2 ...
 $ arswater: num  0.00087 0.00021 0 0.00115 0 0 0.00013 0.00069 0.00039 0 ...
 $ arsnails: num  0.119 0.118 0.099 0.118 0.277 0.358 0.08 0.158 0.31 0.105 ...
```

```
> headtail(ars)
```

	age	sex	usedrink	usecook	arswater	arsnails
1	44	F	E	E	0.00087	0.119
2	45	F	D	E	0.00021	0.118
3	44	M	E	E	0.00000	0.099
19	42	M	E	E	0.01650	0.275
20	62	M	E	E	0.00012	0.135
21	36	M	E	E	0.00410	0.175

Univariate EDA – Categorical

```
> ( tbl.drink <- xtabs(~usedrink,data=ars) )
```

```
usedrink  
  A  B  C  D  E  
1  1  2  3 14
```

```
> percTable(tbl.drink,digits=1)
```

```
usedrink  
  A  B  C  D  E  
4.8 4.8 9.5 14.3 66.7
```

```
> barplot(tbl.drink,xlab="Rating of Use for Drinking",ylab="Frequency",col="gray90")
```

