

Two-Way ANOVA

R Handout

Derek H. Ogle

Initialization

```
> library(NCStats)
> library(multcomp)      # glht()
```

Bacteria Example

Background

What is the optimal temperature (27,35,43°C) and concentration (0.6,0.8,1.0,1.2,1.4% by weight) of the nutrient, tryptone, for culturing the *Staphylococcus aureus* bacterium. Each treatment was repeated twice. The number of bacteria was recorded in millions CFU/mL (CFU=Colony Forming Units).

```
> setwd("C:/aaaWork/Web/GitHub/NCMTH207/modules/Anova-2Way")
> bact <- read.csv("Bacteria.csv")
> str(bact)
```

```
'data.frame':  30 obs. of  3 variables:
 $ temp : int  27 27 27 27 27 35 35 35 35 35 ...
 $ conc  : num  0.6 0.8 1 1.2 1.4 0.6 0.8 1 1.2 1.4 ...
 $ cells: int  55 120 186 260 151 82 166 179 223 178 ...
```

```
> bact$ftemp <- factor(bact$temp)
> bact$fconc <- factor(bact$conc)
> str(bact)
```

```
'data.frame':  30 obs. of  5 variables:
 $ temp : int  27 27 27 27 27 35 35 35 35 35 ...
 $ conc  : num  0.6 0.8 1 1.2 1.4 0.6 0.8 1 1.2 1.4 ...
 $ cells: int  55 120 186 260 151 82 166 179 223 178 ...
 $ ftemp: Factor w/ 3 levels "27","35","43": 1 1 1 1 1 2 2 2 2 2 ...
 $ fconc: Factor w/ 5 levels "0.6","0.8","1",...: 1 2 3 4 5 1 2 3 4 5 ...
```

Initial Summaries

```
> sumTable(cells~ftemp*fconc,data=bact,FUN=length)
```

```
      0.6 0.8 1 1.2 1.4
27    2   2 2   2   2
35    2   2 2   2   2
43    2   2 2   2   2
```

```
> sumTable(cells~ftemp*fconc,data=bact,FUN=mean,digits=0)
```

```
      0.6 0.8   1 1.2 1.4
27 102 106 160 267 131
35   88 161 170 230 198
43  134 166 136 208 164
```

```
> sumTable(cells~ftemp*fconc,data=bact,FUN=sd,digits=1)
```

```
      0.6  0.8    1  1.2  1.4
27 67.2 20.5 37.5  9.9 28.3
35  8.5  7.1 13.4  9.2 29.0
43 26.9 28.3  0.7 27.6 27.6
```

Model Fitting and Summary



```
> lm1 <- lm(cells~ftemp*fconc,data=bact)
> anova(lm1)
```

Analysis of Variance Table



Response: cells

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
ftemp	2	1313	656.4	0.8557	0.44473
fconc	4	51596	12899.1	16.8154	2.041e-05
ftemp:fconc	8	14703	1837.8	2.3958	0.06886
Residuals	15	11507	767.1		

```
> fitPlot(lm1, main="") # left
> fitPlot(lm1,interval=FALSE,change.order=TRUE,xlab="Concentration (%)",
          ylab="Mean Number of Cells",legend="topleft", main="") # right
```



```
> fitPlot(lm1, which="ftemp", ylim=c(60, 270), xlab="Temperature (C)",
  ylab="Mean Number of Cells", main="") # left
> fitPlot(lm1, which="fconc", ylim=c(60, 270), xlab="Concentration (%)",
  ylab="Mean Number of Cells", main="") # right
```



Multiple Comparisons

```
> bact.mc1 <- glht(lm1, mcp(fconc="Tukey"))
```

Warning in mcp2matrix(model, linfct = linfct): covariate interactions found -- default contrast might be inappropriate

```
> summary(bact.mc1)
```

Simultaneous Tests for General Linear Hypotheses

Multiple Comparisons of Means: Tukey Contrasts


Fit: `lm(formula = cells ~ ftemp * fconc, data = bact)`

Linear Hypotheses:

	Estimate	Std. Error	t value	Pr(> t)
0.8 - 0.6 == 0	3.0	27.7	0.108	0.99996
1 - 0.6 == 0	57.0	27.7	2.058	0.28718
1.2 - 0.6 == 0	164.5	27.7	5.939	< 0.001
1.4 - 0.6 == 0	28.5	27.7	1.029	0.83818
1 - 0.8 == 0	54.0	27.7	1.950	0.33498
1.2 - 0.8 == 0	161.5	27.7	5.831	< 0.001
1.4 - 0.8 == 0	25.5	27.7	0.921	0.88452
1.2 - 1 == 0	107.5	27.7	3.881	0.01103
1.4 - 1 == 0	-28.5	27.7	-1.029	0.83819
1.4 - 1.2 == 0	-136.0	27.7	-4.910	0.00148

(Adjusted p values reported -- single-step method)

```
> fitPlot(lm1,which="fconc",xlab="Concentration (%)",ylab="Mean Number of Cells",main="")  
> addSigLetters(lm1,which="fconc",lets=c("a","a","a","b","a"),pos=c(2,2,4,2,4))
```



Soil Phosphorous Example

Background

Soil phosphorous is important for the invasion of native vegetation by exotic weeds. Clements (1983) studied the soil phosphorous in the Sydney region (Australia) to determine how soil phosphorous varied with topographical location and soil type. Bushland sites were chosen in Brisbane Waters National Park, Ku-ring-gai Chase National Park and Royal National Park. These areas were relatively unaffected by suburban development, were free from immediate roadside or track effects, and had not been burned for at least two years. Shale-derived and sandstone-derived soils in four topographic locations were examined with three 250 m² quadrats in each of the eight combinations of soil type and topography. Cores of soil of 75 mm depth and 25 mm diameter, free from surface litter, were collected from each of five randomly selected points in each quadrat. The five soil samples were pooled and the total soil phosphorous (ppm) was determined for each pooled sample. Determine the effect of soil type and topography on total soil phosphorous level.

```
> sp <- read.csv("SoilPhosphorous.csv")
> str(sp)
```


```
'data.frame':  24 obs. of  3 variables:
 $ soil: Factor w/ 2 levels "sandstone","shale": 2 2 2 2 2 2 2 2 2 2 ...
 $ topo: Factor w/ 4 levels "hilltop","north",...: 4 4 4 2 2 2 3 3 3 1 ...
 $ phos: int  98 172 185 78 77 100 117 54 96 83 ...
```

Analysis

```
> lm1 <- lm(phos~soil*topo,data=sp)
> levenesTest(lm1)
```

```
Levene's Test for Homogeneity of Variance (center = median)
      Df F value Pr(>F)
group  7  0.3741 0.9043
      16
```

```
> residPlot(lm1)
```



```
> adTest(lm1$residuals)
```

Anderson-Darling normality test with x
A = 0.2126, p-value = 0.8351

```
> outlierTest(lm1)
```

No Studentized residuals with Bonferonni $p < 0.05$
Largest |rstudent|:
rstudent unadjusted p-value Bonferonni p
1 -2.824098 0.012821 0.30769

```
> anova(lm1)
```

Analysis of Variance Table

Response: phos

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
soil	1	17876.0	17876.0	22.9818	0.0001988
topo	3	9693.8	3231.3	4.1542	0.0235128
soil:topo	3	11390.8	3796.9	4.8814	0.0134826
Residuals	16	12445.3	777.8		

```
> sp$comb <- sp$soil:sp$topo
> view(sp)
```

	soil	topo	phos	comb
2	shale	valley	172	shale:valley
3	shale	valley	185	shale:valley
4	shale	north	78	shale:north
6	shale	north	100	shale:north
19	sandstone	south	28	sandstone:south
21	sandstone	south	72	sandstone:south

```
> lm1a <- lm(phos~comb,data=sp)
> anova(lm1a)
```

Analysis of Variance Table

Response: phos

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
comb	7	38961	5565.8	7.1555	0.0005729
Residuals	16	12445	777.8		

```
> spint.mc <- glht(lm1a, mcp(comb="Tukey"))
```

```
> summary(spint.mc)
```

Simultaneous Tests for General Linear Hypotheses

Multiple Comparisons of Means: Tukey Contrasts

Fit: lm(formula = phos ~ comb, data = sp)

Linear Hypotheses:

	Estimate	Std. Error	t value	Pr(> t)
sandstone:north - sandstone:hilltop == 0	1.667	22.772	0.073	1.00000
sandstone:south - sandstone:hilltop == 0	19.333	22.772	0.849	0.98683
sandstone:valley - sandstone:hilltop == 0	-4.000	22.772	-0.176	1.00000
shale:hilltop - sandstone:hilltop == 0	4.667	22.772	0.205	1.00000
shale:north - sandstone:hilltop == 0	53.333	22.772	2.342	0.33025
shale:south - sandstone:hilltop == 0	57.333	22.772	2.518	0.25548
shale:valley - sandstone:hilltop == 0	120.000	22.772	5.270	0.00157
sandstone:south - sandstone:north == 0	17.667	22.772	0.776	0.99218
sandstone:valley - sandstone:north == 0	-5.667	22.772	-0.249	1.00000
shale:hilltop - sandstone:north == 0	3.000	22.772	0.132	1.00000
shale:north - sandstone:north == 0	51.667	22.772	2.269	0.36503
shale:south - sandstone:north == 0	55.667	22.772	2.445	0.28476
shale:valley - sandstone:north == 0	118.333	22.772	5.196	0.00180
sandstone:valley - sandstone:south == 0	-23.333	22.772	-1.025	0.96344
shale:hilltop - sandstone:south == 0	-14.667	22.772	-0.644	0.99747
shale:north - sandstone:south == 0	34.000	22.772	1.493	0.80062
shale:south - sandstone:south == 0	38.000	22.772	1.669	0.70549
shale:valley - sandstone:south == 0	100.667	22.772	4.421	0.00797
shale:hilltop - sandstone:valley == 0	8.667	22.772	0.381	0.99992
shale:north - sandstone:valley == 0	57.333	22.772	2.518	0.25527
shale:south - sandstone:valley == 0	61.333	22.772	2.693	0.19367
shale:valley - sandstone:valley == 0	124.000	22.772	5.445	0.00101
shale:north - shale:hilltop == 0	48.667	22.772	2.137	0.43361
shale:south - shale:hilltop == 0	52.667	22.772	2.313	0.34411
shale:valley - shale:hilltop == 0	115.333	22.772	5.065	0.00221
shale:south - shale:north == 0	4.000	22.772	0.176	1.00000
shale:valley - shale:north == 0	66.667	22.772	2.928	0.13149
shale:valley - shale:south == 0	62.667	22.772	2.752	0.17653

(Adjusted p values reported -- single-step method)

```
> glhtSig(spint.mc)
```

```
[1] "shale:valley - sandstone:hilltop" "shale:valley - sandstone:north"
[3] "shale:valley - sandstone:south"   "shale:valley - sandstone:valley"
[5] "shale:valley - shale:hilltop"
```

```
> fitPlot(lm1,change.order=TRUE,interval=FALSE,main="",ylim=c(20,160),
          ylab="Mean Phosphorous Level",xlab="Topographic Location",legend="topleft")
> addSigLetters(lm1,change.order=TRUE,lets=c("a","a","a","ab","a","ab","a","b"),
               pos=c(1,3,1,3,1,1,3,1))
```

