

Growth of Pronghorn Antelopes

1. The indicator variable is named $TAME$ where $TAME = 1$ if the pronghorn is in the “tame” group and $TAME = 0$ if it is in the “diet-curtailed” group.
2. The full model is $\mu_{WEIGHT} = \alpha + \beta TIME + \delta_1 TAME + \gamma_1 TAME * TIME$
3. The submodels for both groups are shown in the table below

Group	$Tame=$	Submodel ($\mu_{WEIGHT} =$)
Diet-curtailed	0	$= \alpha + \beta TIME$
Tame	1	$= (\alpha + \delta_1) + (\beta + \gamma_1) TIME$

4. Interpretations of the coefficients are listed below.
 - α is the intercept of diet-curtailed (reference) group
 - β is the slope of diet-curtailed (reference) group
 - δ_1 is the difference in intercept of tame and diet-curtailed groups (i.e., tame-dietcurtailed)
 - γ_1 is the difference in slopes of tame and diet-curtailed groups (i.e., tame-dietcurtailed)
5. The models for the “parallel lines test” are shown below

$$H_O : \mu_{WEIGHT|\dots} = \alpha + \beta TIME + \delta_1 TAME$$

$$H_A : \mu_{WEIGHT|\dots} = \alpha + \beta TIME + \delta_1 TAME + \gamma_1 TAME * TIME$$

6. The models for the “equal-intercepts test” (assuming parallele lines) are shown below

$$H_O : \mu_{WEIGHT|\dots} = \alpha + \beta TIME$$

$$H_A : \mu_{WEIGHT|\dots} = \alpha + \beta TIME + \delta_1 TAME$$

Food Intake for Rainbow Trout

1. The three required indicator variables are shown below

- $FOUR = 1$ if in the “four-day starved” group, $FOUR = 0$ otherwise
- $EIGHT = 1$ if in the “eight-day starved” group, $EIGHT = 0$ otherwise
- $STEEN = 1$ if in the “sixteen-day starved” group, $STEEN = 0$ otherwise

2. The full model is

$$\mu_{stomvol} = \alpha + \beta intake + \delta_1 FOUR + \delta_2 EIGHT + \delta_3 STEEN \\ + \gamma_1 FOUR * intake + \gamma_2 EIGHT * intake + \gamma_3 STEEN * intake$$

3. The sub-models for all four groups are shown in the table below

Group	$FOUR=$	$EIGHT=$	$STEEN=$	Submodel ($\mu_{stomvol} =$)
1-day starved	0	0	0	$= \alpha + \beta intake$
4-day starved	1	0	0	$= (\alpha + \delta_1) + (\beta + \gamma_1) intake$
8-day starved	0	1	0	$= (\alpha + \delta_2) + (\beta + \gamma_2) intake$
16-day starved	0	0	1	$= (\alpha + \delta_3) + (\beta + \gamma_3) intake$

4. Interpretations of the coefficients are listed below.

- α is the intercept of the 1-day starved (reference) group
- β is the slope of the 1-day starved (reference) group
- δ_1 is the difference in the intercepts of the 4-day and 1-day starved groups
- γ_1 is the difference in the slopes of the 4-day and 1-day starved groups
- δ_2 is the difference in the intercepts of the 8-day and 1-day starved groups
- γ_2 is the difference in the slopes of the 8-day and 1-day starved groups
- δ_3 is the difference in the intercepts of the 16-day and 1-day starved groups
- γ_3 is the difference in the slopes of the 16-day and 1-day starved groups

5. Models for the “parallel lines test” are shown below

$$H_O : \mu_{stomvol} = \alpha + \beta intake + \delta_1 FOUR + \delta_2 EIGHT + \delta_3 STEEN \\ H_A : \mu_{stomvol} = \alpha + \beta intake + \delta_1 FOUR + \delta_2 EIGHT + \delta_3 STEEN \\ + \gamma_1 FOUR * intake + \gamma_2 EIGHT * intake + \gamma_3 STEEN * intake$$

6. Models for the “equal-intercepts test” (assuming parallel lines) are shown below

$$H_O : \mu_{stomvol} = \alpha + \beta intake \\ H_A : \mu_{stomvol} = \alpha + \beta intake + \delta_1 FOUR + \delta_2 EIGHT + \delta_3 STEEN$$