

One-Way ANOVA

R Handout

Derek H. Ogle

Initialization

```
> library(NCStats)
> library(multcomp)      # glht()
```

Raspberry Example

Background

A researcher is interested in the effect of irrigation on fruit production by raspberry plants. The researcher has determined that he will examine the effects of 100 ml (a maintenance amount), 200, 400, and 800 ml of water per pot. The researcher had 16 identical planting pots available and much more than that number of raspberry plant seedlings. A square table for growing the plants in a greenhouse was also available. He had enough time to let the plants mature to the point of producing fruit (i.e. berries) or not. At the end of this period, the total weight (g) of mature berries was recorded.

```
> setwd("C:/aaaWork/Web/GitHub/NCMTH207/modules/Anova-1Way")
> rasp <- read.csv("Raspberry.csv")
> str(rasp)
```

```
'data.frame':   16 obs. of  2 variables:
 $ water : int   100 100 100 100 200 200 200 200 400 400 ...
 $ weight: num   8.1 10.9 11.1 13.9 12.2 11.5 11.4 6.8 6.5 5.5 ...
```

```
> rasp$water <- factor(rasp$water)
> str(rasp)
```

```
'data.frame':   16 obs. of  2 variables:
 $ water : Factor w/ 4 levels "100","200","400",...: 1 1 1 1 2 2 2 2 3 3 ...
 $ weight: num   8.1 10.9 11.1 13.9 12.2 11.5 11.4 6.8 6.5 5.5 ...
```

Fitting the Linear Model

```
> lm1 <- lm(weight~water,data=rasp)
> anova(lm1)
```

Analysis of Variance Table

Response: weight

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
water	3	115.043	38.348	10.793	0.001004
Residuals	12	42.635	3.553		

```
> summary(lm1)
```

Coefficients:


	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	11.0000	0.9425	11.672	6.58e-08
water200	-0.5250	1.3328	-0.394	0.70057
water400	-5.6250	1.3328	-4.220	0.00119
water800	-5.6000	1.3328	-4.202	0.00123

Residual standard error: 1.885 on 12 degrees of freedom

Multiple R-squared: 0.7296, Adjusted R-squared: 0.662

F-statistic: 10.79 on 3 and 12 DF, p-value: 0.001004

```
> fitPlot(lm1,xlab="Water Treatment (ml)",ylab="Weight (g)",main="")
```



Multiple Comparison Tests

```
> rasp.mc <- glht(lm1, mcp(water = "Tukey"))  
> summary(rasp.mc)
```

Simultaneous Tests for General Linear Hypotheses

Multiple Comparisons of Means: Tukey Contrasts

```
Fit: lm(formula = weight ~ water, data = rasp)
```

Linear Hypotheses:

	Estimate	Std. Error	t value	Pr(> t)
200 - 100 == 0	-0.525	1.333	-0.394	0.97834
400 - 100 == 0	-5.625	1.333	-4.220	0.00537
800 - 100 == 0	-5.600	1.333	-4.202	0.00589
400 - 200 == 0	-5.100	1.333	-3.826	0.01104
800 - 200 == 0	-5.075	1.333	-3.808	0.01133
800 - 400 == 0	0.025	1.333	0.019	1.00000

(Adjusted p values reported -- single-step method)

```
> confint(rasp.mc)
```

Simultaneous Confidence Intervals

Multiple Comparisons of Means: Tukey Contrasts

```
Fit: lm(formula = weight ~ water, data = rasp)
```


Quantile = 2.9683

95% family-wise confidence level

Linear Hypotheses:

	Estimate	lwr	upr
200 - 100 == 0	-0.5250	-4.4813	3.4313
400 - 100 == 0	-5.6250	-9.5813	-1.6687
800 - 100 == 0	-5.6000	-9.5563	-1.6437
400 - 200 == 0	-5.1000	-9.0563	-1.1437
800 - 200 == 0	-5.0750	-9.0313	-1.1187
800 - 400 == 0	0.0250	-3.9313	3.9813

```
> fitPlot(lm1,xlab="Water Treatment (ml)",ylab="Weight (g)",main="")
> addSigLetters(lm1,lets=c("a","a","b","b"),pos=c(2,4,2,4))
```




Checking the Assumptions

```
> levenesTest(lm1)
```

```
Levene's Test for Homogeneity of Variance (center = median)
      Df F value Pr(>F)
group  3  0.3256 0.8069
      12
```

```
> residPlot(lm1)
```



```
> adTest(lm1$residuals)
```

```
Anderson-Darling normality test with x  
A = 0.4308, p-value = 0.2688
```

```
> outlierTest(lm1)
```

```
No Studentized residuals with Bonferonni p < 0.05  
Largest |rstudent|:  
      rstudent unadjusted p-value Bonferonni p  
8 -2.836044      0.016196      0.25914
```

Benthic Infaunal Example

Background

Australian researchers were interested in the effect of effluent releases on benthic organisms in the release area. To examine the effect, the researchers recorded the total abundance of benthic organisms at 8 haphazardly-selected sublocations at each of 8 control locations (thought to have not been impacted by the effluent release) and 1 potentially impacted location. The results are recorded in **BenthicInfaunal.csv**. Use these data to determine if the mean abundance of benthic organisms differs between the locations (and, especially, if the impacted location differs from any of the control locations).

```
> ben <- read.csv("BenthicInfaunal.csv")  
> ben$site <- factor(ben$site)  
> str(ben)
```


```
'data.frame':  72 obs. of  2 variables:  
 $ site      : Factor w/ 9 levels "1","2","3","4",...: 1 1 1 1 1 1 1 1 2 2 ...  
 $ abundance: num  14.4 20.4 21.2 17.6 29 ...
```

Assumption Checking with Possible Transformations

```
> lm2 <- lm(abundance~site,data=ben)  
> levenesTest(lm2)
```

```
Levene's Test for Homogeneity of Variance (center = median)  
      Df F value    Pr(>F)  
group  8  3.2452 0.003726  
      63
```

```
> residPlot(lm2)
```



```
> adTest(lm2$residuals)
```

Anderson-Darling normality test with x
A = 1.6389, p-value = 0.0002996

```
> outlierTest(lm2)
```


	rstudent	unadjusted p-value	Bonferonni p
20	6.624666	9.5554e-09	6.8799e-07

```
> ## transChooser(lm2) # interactive, result not shown
> ben$logab <- log(ben$abundance)
> lm3 <- lm(logab~site,data=ben)
> levenesTest(lm3)
```

Levene's Test for Homogeneity of Variance (center = median)

	Df	F value	Pr(>F)
group	8	1.5339	0.1636
	63		

```
> residPlot(lm3)
```



```
> adTest(lm3$residuals)
```

Anderson-Darling normality test with x
A = 0.3323, p-value = 0.5062

```
> outlierTest(lm3)
```

No Studentized residuals with Bonferonni $p < 0.05$
Largest |rstudent|:

	rstudent	unadjusted p-value	Bonferonni p
20	2.928889	0.004754	0.34229

Model Summarization

```
> anova(lm3)
```

Analysis of Variance Table

Response: logab

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
site	8	8.6683	1.08353	29.066	< 2.2e-16
Residuals	63	2.3485	0.03728		

```
> ben.mc <- glht(lm3, mcp(site = "Dunnett"))
> summary(ben.mc)
```

Simultaneous Tests for General Linear Hypotheses

Multiple Comparisons of Means: Dunnett Contrasts


```
Fit: lm(formula = logab ~ site, data = ben)
```

Linear Hypotheses:

	Estimate	Std. Error	t value	Pr(> t)
2 - 1 == 0	-0.218435	0.096537	-2.263	0.14575
3 - 1 == 0	0.703189	0.096537	7.284	< 0.001
4 - 1 == 0	0.453836	0.096537	4.701	< 0.001
5 - 1 == 0	-0.414859	0.096537	-4.297	< 0.001
6 - 1 == 0	-0.004238	0.096537	-0.044	1.00000
7 - 1 == 0	0.140280	0.096537	1.453	0.57978
8 - 1 == 0	-0.371867	0.096537	-3.852	0.00199
9 - 1 == 0	0.168668	0.096537	1.747	0.37998

(Adjusted p values reported -- single-step method)

```
> fitPlot(lm3,ylab="Log Abundance",xlab="Site",main="")
> addSigLetters(lm3,lets=c("", "", "***", "***", "***", "", "", "***", ""),pos=c(2,4,2,4,2,2,4,2,4))
```



```
> confint(ben.mc)
```

Simultaneous Confidence Intervals

Multiple Comparisons of Means: Dunnett Contrasts

```
Fit: lm(formula = logab ~ site, data = ben)
```

Quantile = 2.7282

95% family-wise confidence level

Linear Hypotheses:

	Estimate	lwr	upr
2 - 1 == 0	-0.218435	-0.481812	0.044943
3 - 1 == 0	0.703189	0.439811	0.966566
4 - 1 == 0	0.453836	0.190459	0.717214
5 - 1 == 0	-0.414859	-0.678237	-0.151482
6 - 1 == 0	-0.004238	-0.267615	0.259140
7 - 1 == 0	0.140280	-0.123097	0.403658
8 - 1 == 0	-0.371867	-0.635245	-0.108490
9 - 1 == 0	0.168668	-0.094709	0.432045

```
> exp(confint(ben.mc)$confint)
```

	Estimate	lwr	upr
2 - 1	0.8037761	0.6176580	1.0459769
3 - 1	2.0201841	1.5524011	2.6289235
4 - 1	1.5743404	1.2097946	2.0487343
5 - 1	0.6604332	0.5075068	0.8594406
6 - 1	0.9957713	0.7651959	1.2958258
7 - 1	1.1505965	0.8841705	1.4973042
8 - 1	0.6894457	0.5298014	0.8971955
9 - 1	1.1837272	0.9096297	1.5404181

attr(,"conf.level")
[1] 0.95
attr(,"calpha")
[1] 2.728333