von Bertalanffy Growth Function - Nunavut Exercise

1.	Load the PG008_orig	ginal.xlsx file	into a	data.frame	object	and	$\operatorname{restrict}$	the	data	to	only	those	fish	captured	in
	$freshwater \ {\rm in} \ 2010.$	Use these data:	for the	following qu	uestions										

- a. Examine the plot of length versus age. Do the data look linear or curved, is there an obvious asymptote, are young fish well represented, how variable are lengths within ages?
- b. Fit the typical parameterization of the VBGF to these data. Construct a fitted-line plot (i.e., superimpose the fitted VBGF onto the length versus age plot) and a residual plot. Comment on model fit.
- c. Compute the correlation between parameter values. Comment
- d. Compute the parameter estimates. Carefully interpret the value of each parameter. Comment on how realistic you think each estimate is.
- e. Construct 95% likelihood profile and bootstrap confidence intervals. How do the relative widths of the confidence intervals compare (between methods).
- f. Predict the mean length, with 95% confidence interval, for an age-18 Charr. Comment on the width of this confidence interval?
- 2. [Time Permitting] Repeat the previous question but using either the original, Gallucci and Quinn, or Mooij parameterizations of the VBGF. [Note that you can see the equations for these VBGFs with, for example, growthFunShow("vonBertalanffy",param="GallucciQuinn",plot=TRUE). You can declare a function for these VBGFs by using, for example, vb <- vbFuns("GallucciQuinn").]
 - a. How does the fit of this model (and estimates of the common parameters) compare with the results from the typical VBGF fit in the previous question?
- 3. [Time Permitting] Repeat the first question but using either the Gompertz or logistic growth functions. [Note that you can declare a logistic growth function by using, for example, lgf <- logisticFuns(msg=TRUE).]
 - a. How does the fit of this growth function compare with the results from the typical VBGF fit in the first question?