



Loops and the Fundamental Group

Dror Atariah
Freie Universität Berlin

What is Seminar, 21/11/2008

Motivation

Loops

- Relations Between Loops
- Loops Concatenation



Group Structure on Loops

Examples

- Subset of plane
- The Circle
- The Torus


- ▶ We want to study topological spaces

- ▶ We want to study topological spaces
- ▶ What is the difference between the following two spaces?




Let's have some loops

First let us have a look into the first example:



Let's have some loops


First let us have a look into the first example:



Each loop can be continuously contracted into a point.


And what about loops here?

Next, lets consider the second example:



And what about loops here?


Next, lets consider the second example:



Here, the *red* loop cannot be contracted to a point.

And what about loops here?

Next, lets consider the second example:




Here, the *red* loop cannot be contracted to a point.
Let's introduce some definitions!

Definition (Loop)

Given a topological space X and the unit interval $I \subset \mathbb{R}$, a loop is a continuous map

$$\lambda : I \rightarrow X$$

such that $\lambda(0) = \lambda(1)$.



- Later on, we will consider topological spaces with a base point (X, x_0) , where X is the topological space, and x_0 is the base point.

- ▶ Later on, we will consider topological spaces with a base point (X, x_0) , where X is the topological space, and x_0 is the base point.
- ▶ For any loop λ we have $\lambda(0) = \lambda(1) = x_0$

Motivation

Loops

Relations Between Loops

Loops Concatenation

Group Structure on Loops

Examples


Subset of plane


The Circle

The Torus

How To Distinguish Between Loops?

Consider the following picture





Homotopy


Definition (Homotopy of loops)

Two loops $\lambda, \mu : I \rightarrow (X, x_0)$ are called *homotopic* with base point held fixed, or for short:

$$\lambda \simeq \mu \quad \text{rel}(0, 1)$$

if there exists a continuous map $F : I \times I \rightarrow (X, x_0)$ such that the following holds:

1. $F(s, 0) = \lambda(s), \quad \forall s \in I$
2. $F(s, 1) = \mu(s), \quad \forall s \in I$
3. $F(0, t) = F(1, t) = x_0, \quad \forall t \in I$



Homotopy



Definition (Homotopy of loops)

Two loops $\lambda, \mu : I \rightarrow (X, x_0)$ are called *homotopic* with base point held fixed, or for short:

$$\lambda \simeq \mu \quad \text{rel}(0, 1)$$

if there exists a continuous map $F : I \times I \rightarrow (X, x_0)$ such that the following holds:

1. $F(s, 0) = \lambda(s), \quad \forall s \in I$
2. $F(s, 1) = \mu(s), \quad \forall s \in I$
3. $F(0, t) = F(1, t) = x_0, \quad \forall t \in I$



Homotopy

Definition (Homotopy of loops)



Two loops $\lambda, \mu : I \rightarrow (X, x_0)$ are called *homotopic* with base point held fixed, or for short:

$$\lambda \simeq \mu \quad \text{rel}(0, 1)$$

if there exists a continuous map $F : I \times I \rightarrow (X, x_0)$ such that the following holds:


1. $F(s, 0) = \lambda(s), \quad \forall s \in I$
2. $F(s, 1) = \mu(s), \quad \forall s \in I$
3. $F(0, t) = F(1, t) = x_0, \quad \forall t \in I$

The map F is called the *homotopy* between λ and μ .




Example of Homotopy

Let's have a look at the following figure:



Example of Homotopy

Let's have a look at the following figure:



- The red loops are homotopic, so are the green loops.

- 11

Motivation

Loops

Relations Between Loops

Loops Concatenation

Group Structure on Loops

Examples

Subset of plane


The Circle

The Torus

Definition

Given two loops $\lambda, \mu : I \rightarrow X$ with a base point x_0 , we define the concatenation of them as follows:

$$\lambda * \mu(t) = \begin{cases} \lambda(2t) & 0 \leq t \leq \frac{1}{2} \\ \mu(2t - 1) & \frac{1}{2} \leq t \leq 1 \end{cases}$$



We first traverse along the **red** loop and then along the **green** one.

The homotopy relation between loops is an *equivalence relation*, that is:

The homotopy relation between loops is an *equivalence relation*, that is:

- ▶ $\lambda \simeq \lambda \quad \text{rel}(0, 1)$
- ▶ $\lambda \simeq \mu \quad \text{rel}(0, 1) \Rightarrow \mu \simeq \lambda \quad \text{rel}(0, 1)$
- ▶ $\lambda \simeq \mu \quad \text{rel}(0, 1) \text{ and } \mu \simeq \tau \quad \text{rel}(0, 1) \Rightarrow \lambda \simeq \tau \quad \text{rel}(0, 1)$

The homotopy relation between loops is an *equivalence relation*, that is:

- ▶ $\lambda \simeq \lambda \quad \text{rel}(0, 1)$
- ▶ $\lambda \simeq \mu \quad \text{rel}(0, 1) \Rightarrow \mu \simeq \lambda \quad \text{rel}(0, 1)$
- ▶ $\lambda \simeq \mu \quad \text{rel}(0, 1) \text{ and } \mu \simeq \tau \quad \text{rel}(0, 1) \Rightarrow \lambda \simeq \tau \quad \text{rel}(0, 1)$

Equivalence Classes

So we can consider the set of equivalence classes of loops $[\lambda]$ over a topological space X with base point x_0 .

In order to have a group structure on the set of equivalence classes, we have to define a group operation.

In order to have a group structure on the set of equivalence classes, we have to define a group operation.

Concatenation

We recall the concatenation $*$

In order to have a group structure on the set of equivalence classes, we have to define a group operation.

Concatenation

We recall the concatenation $*$

Definition (Group Operation)

Given two loop classes $[\lambda]$ and $[\mu]$ we define:

1. $[\lambda] * [\mu] := [\lambda * \mu]$
2. The inverse of $[\lambda]$ is given by $[\lambda^{-1}]$ that is $[\lambda]^{-1} = [\lambda^{-1}]$, where $\lambda^{-1}(t) = \lambda(1 - t)$.

Definition

Given a topological space X with a base point x_0 , the *fundamental group*, $\pi_1(X, x_0)$, is the set of equivalence classes of loops along with the operation $*$

FU Berlin, Loops and Groups, Nov. 2008

Two More Definitions

Before we go on, for the sake of completeness!

Definition (The Constant Loop)

The constant loop in a pointed topological space (X, x_0) is the the map:

$$\xi : I \rightarrow x_0$$

that is $\xi(t) = x_0$ for all $t \in I$.

Two More Definitions

Before we go on, for the sake of completeness!

Definition (The Constant Loop)

The constant loop in a pointed topological space (X, x_0) is the the map:

$$\xi : I \rightarrow x_0$$

that is $\xi(t) = x_0$ for all $t \in I$.

Definition (Null-Homotopic)

A loop λ is called *null-homotopic* if it is homotopic to the constant loop.

Two More Definitions

Before we go on, for the sake of completeness!

Definition (The Constant Loop)

The constant loop in a pointed topological space (X, x_0) is the the map:

$$\xi : I \rightarrow x_0$$

that is $\xi(t) = x_0$ for all $t \in I$.

Definition (Null-Homotopic)

A loop λ is called *null-homotopic* if it is homotopic to the constant loop.

The Group's Unit Element

Note that the class of null-homotopic loops is the unit element of the fundamental group.

-

Loops

Loops Concatenation


Examples


The Circle

20


Set With No Holes

Consider the following pointed space





- Here all loops are null-homotopic, i.e. all are homotopic to the constant loop




- ▶ Here all loops are null-homotopic, i.e. all are homotopic to the constant loop
- ▶ This means that the fundamental group is trivial


-


Two topological spaces X and Y are called homotopically equivalent, or of the same homotopy type, if there exists two maps $f : X \rightarrow Y$ and $g : Y \rightarrow X$ such that $f \circ g \simeq id_X$ and $g \circ f \simeq id_Y$.

Set With a Hole

Now Consider the following space








Here we have two types of loops:

- ▶ Those homotopically equivalent to the constant loop
- ▶ Those which enclose the hole



- ▶ Those homotopically equivalent to the constant loop
- ▶ Those which enclose the hole

What is the fundamental group in this case?

Motivation

Loops

Relations Between Loops

Loops Concatenation


Group Structure on Loops

Examples

Subset of plane

The Circle

The Torus



The Fundamental Group of the Circle

Theorem

We have for any point $x_0 \in S^1$

$$\pi_1(S^1, x_0) = \mathbb{Z}$$

The Fundamental Group of the Circle

Theorem

We have for any point $x_0 \in S^1$

$$\pi_1(S^1, x_0) = \mathbb{Z}$$

Proof Outline.

- ▶ Up to rotations, all loops in S^1 are characterized by the number of times they wind around the origin
- ▶ A negative integer i is isomorphic to a loop winding i times clockwise
- ▶ Concatenation of loops in $\pi_1(S^1, x_0)$ is equivalent to addition of integers in \mathbb{Z}



Motivation

Loops

Relations Between Loops

Loops Concatenation

Group Structure on Loops

Examples

Subset of plane

The Circle

The Torus

The Fundamental Group of The Torus


- ▶ The torus can be given as a product of two circles: $T^2 = S^1 \times S^1$



The Fundamental Group of The Torus

- ▶ The torus can be given as a product of two circles: $T^2 = S^1 \times S^1$
- ▶ Thus, $\pi_1(T^2) = \pi_1(S^1 \times S^1) \cong \pi_1(S^1) \times \pi_1(S^1) \cong \mathbb{Z} \times \mathbb{Z}$

The Fundamental Group of The Torus

- ▶ The torus can be given as a product of two circles: $T^2 = S^1 \times S^1$
- ▶ Thus, $\pi_1(T^2) = \pi_1(S^1 \times S^1) \cong \pi_1(S^1) \times \pi_1(S^1) \cong \mathbb{Z} \times \mathbb{Z}$
- ▶ The pair (i, j) of integers corresponds to a loop winding i times around the first circle and j times around the other one

- 



Thank you!

atariah@mi.fu-berlin.de