

Almost periodic Functions and the Scott Analysis of Linear Orderings

David Gonzalez

UC Berkeley

July 2024

TU Wein

Outline

- ▶ The definitions and history of Scott analysis
- ▶ Classification for linear orderings
- ▶ A recent construction using almost periodic functions

$\mathcal{L}_{\omega_1, \omega}$ formulas and their complexity

- ▶ $\mathcal{L}_{\omega_1, \omega}$ is infinitary logic; it extends first order logic by allowing countable conjunctions and disjunctions.

$\mathcal{L}_{\omega_1, \omega}$ formulas and their complexity

- ▶ $\mathcal{L}_{\omega_1, \omega}$ is infinitary logic; it extends first order logic by allowing countable conjunctions and disjunctions.
- ▶ $\varphi \in \mathcal{L}_{\omega_1, \omega}$ is in $\Sigma_0^{\text{in}} = \Pi_0^{\text{in}}$ if it is quantifier free and has no infinitary disjunctions or conjunctions.
- ▶ For $\alpha \in \omega_1$, φ is $\Sigma_\alpha^{\text{in}}$ if $\varphi = \bigvee_i \exists \bar{x} \psi_i(\bar{x})$ for $\psi_i \in \Pi_\beta^{\text{in}}$ with $\beta < \alpha$.
- ▶ For $\alpha \in \omega_1$, φ is Π_α^{in} if $\varphi = \bigwedge_i \forall \bar{x} \psi_i(\bar{x})$ for $\psi_i \in \Sigma_\beta^{\text{in}}$ with $\beta < \alpha$.

$\mathcal{L}_{\omega_1, \omega}$ formulas and their complexity

- ▶ $\mathcal{L}_{\omega_1, \omega}$ is infinitary logic; it extends first order logic by allowing countable conjunctions and disjunctions.
- ▶ $\varphi \in \mathcal{L}_{\omega_1, \omega}$ is in $\Sigma_0^{\text{in}} = \Pi_0^{\text{in}}$ if it is quantifier free and has no infinitary disjunctions or conjunctions.
- ▶ For $\alpha \in \omega_1$, φ is $\Sigma_\alpha^{\text{in}}$ if $\varphi = \bigvee_i \exists \bar{x} \psi_i(\bar{x})$ for $\psi_i \in \Pi_\beta^{\text{in}}$ with $\beta < \alpha$.
- ▶ For $\alpha \in \omega_1$, φ is Π_α^{in} if $\varphi = \bigwedge_i \forall \bar{x} \psi_i(\bar{x})$ for $\psi_i \in \Sigma_\beta^{\text{in}}$ with $\beta < \alpha$.
- ▶ For $\alpha \in \omega_1$, φ is $d\text{-}\Sigma_\alpha^{\text{in}}$ if $\varphi = \psi \wedge \chi$ for $\psi \in \Sigma_\alpha^{\text{in}}$ and $\chi \in \Pi_\alpha^{\text{in}}$

$\mathcal{L}_{\omega_1, \omega}$ formulas and their complexity

- ▶ For two models M, N we say $M \leq_\alpha N$ if $\Pi_\alpha^{\text{in}} - \text{Th}(M) \subseteq \Pi_\alpha^{\text{in}} - \text{Th}(N)$.
- ▶ Note that $M \geq_\alpha N$ if and only if $\Sigma_\alpha^{\text{in}} - \text{Th}(M) \subseteq \Sigma_\alpha^{\text{in}} - \text{Th}(N)$.
- ▶ We put $M \equiv_\alpha N$ if both of the above hold.

Scott Complexity

Theorem: [Scott] For every countable structure M there is a sentence $\varphi \in \mathcal{L}_{\omega_1, \omega}$ such that $N \cong M \iff N \models \varphi$.

Scott Complexity

Theorem: [Scott] For every countable structure M there is a sentence $\varphi \in \mathcal{L}_{\omega_1, \omega}$ such that $N \cong M \iff N \models \varphi$.


Definition: A φ as in the theorem statement is called a *Scott sentence*.

Scott Complexity

Theorem: [Scott] For every countable structure M there is a sentence $\varphi \in \mathcal{L}_{\omega_1, \omega}$ such that $N \cong M \iff N \models \varphi$.

Definition: A φ as in the theorem statement is called a *Scott sentence*.

Definition: The *Scott complexity* (SC) of M is the least among $\{\Sigma_{\alpha}^{\text{in}}, \Pi_{\alpha}^{\text{in}}, d\text{-}\Sigma_{\alpha}^{\text{in}}\}_{\alpha \in \omega_1}$ such that M has a Scott sentence of said complexity.



Robustness and Scott Rank

Definition: [Montalbán] The parametrized *Scott rank* (pSR) of M is the least $\alpha \in \omega_1$ such that M has a $\Sigma_{\alpha+2}^{in}$ Scott sentence.

Robustness and Scott Rank

Definition: [Montalbán] The parametrized *Scott rank* (pSR) of M is the least $\alpha \in \omega_1$ such that M has a $\Sigma_{\alpha+2}^{in}$ Scott sentence.

Definition: [Montalbán] The unparametrized *Scott rank* (uSR) of M is the least $\alpha \in \omega_1$ such that M has a $\Pi_{\alpha+1}^{in}$ Scott sentence.

Robustness and Scott Rank

Definition: [Montalbán] The parametrized *Scott rank* (pSR) of M is the least $\alpha \in \omega_1$ such that M has a $\Sigma_{\alpha+2}^{in}$ Scott sentence.

Definition: [Montalbán] The unparametrized *Scott rank* (uSR) of M is the least $\alpha \in \omega_1$ such that M has a $\Pi_{\alpha+1}^{in}$ Scott sentence.

Theorem: [Montalbán] The (un)parameterized Scott rank of M is the $\alpha \in \omega_1$ such that M the automorphism orbits of all tuples in M are definable in a Σ_{α}^{in} way with(out) parameters. There are also *many* other equivalent statements.

Scott analysis

For $T \in \mathcal{L}_{\omega_1, \omega}$ let

$$I_{SC}(T, \Gamma) = |\{M : M \models T \wedge SC(M) = \Gamma\}|,$$

$$I_u(T, \alpha) = |\{M : M \models T \wedge uSR(M) = \alpha\}|,$$

$$I_p(T, \alpha) = |\{M : M \models T \wedge pSR(M) = \alpha\}|.$$

Scott analysis

For $T \in \mathcal{L}_{\omega_1, \omega}$ let

$$I_{SC}(T, \Gamma) = |\{M : M \models T \wedge SC(M) = \Gamma\}|,$$

$$I_u(T, \alpha) = |\{M : M \models T \wedge uSR(M) = \alpha\}|,$$

$$I_p(T, \alpha) = |\{M : M \models T \wedge pSR(M) = \alpha\}|.$$

Scott analysis generally refers to any inquiry into the behavior of the above functions.

Linear orderings

Between G. and Rosseger 2023 and G., Harrison-Trainor and Ho 2024 we fully characterized the I_{SC} function for linear orderings.

Linear orderings

Between G. and Rosseger 2023 and G., Harrison-Trainor and Ho 2024 we fully characterized the I_{SC} function for linear orderings.

1. $I_{SC}(LO, \Pi_n) = 1$ if $n \leq 2$
2. $I_{SC}(LO, \Sigma_n) = 0$ if $n \leq 3$

Linear orderings

Between G. and Rosseger 2023 and G., Harrison-Trainor and Ho 2024 we fully characterized the I_{SC} function for linear orderings.

1. $I_{SC}(LO, \Pi_n) = 1$ if $n \leq 2$
2. $I_{SC}(LO, \Sigma_n) = 0$ if $n \leq 3$
3. $I_{SC}(LO, d\text{-}\Sigma_n) = \aleph_0$ if $n \leq 4$
4. $I_{SC}(LO, \Sigma_4) = \aleph_0$
5. $I_{SC}(LO, \Pi_3) = \aleph_0$

Linear orderings

Between G. and Rosseger 2023 and G., Harrison-Trainor and Ho 2024 we fully characterized the I_{SC} function for linear orderings.

1. $I_{SC}(LO, \Pi_n) = 1$ if $n \leq 2$
2. $I_{SC}(LO, \Sigma_n) = 0$ if $n \leq 3$
3. $I_{SC}(LO, d\text{-}\Sigma_n) = \aleph_0$ if $n \leq 4$
4. $I_{SC}(LO, \Sigma_4) = \aleph_0$
5. $I_{SC}(LO, \Pi_3) = \aleph_0$
6. $I_{SC}(LO, \Gamma) = 2^{\aleph_0}$ otherwise

The Relationship of the Concepts

Alvir, Greenberg, Harrison-Trainor and Turetsky (AGHTT) showed that Scott sentence complexity is related to Montalbán's Scott ranks.

SC	pSR	uSR	complexity of parameters
$\Sigma_{\alpha+2}^{\text{in}}$	α	$\alpha + 2$	$\Pi_{\alpha+1}^{\text{in}}$
$d\text{-}\Sigma_{\alpha+1}^{\text{in}}$	α	$\alpha + 1$	Π_{α}^{in}
$\Pi_{\alpha+1}^{\text{in}}$	α	α	none
α limit			
$\Sigma_{\alpha+1}^{\text{in}}$	α	$\alpha + 1$	Π_{α}^{in}
Π_{α}^{in}	α	α	none

For limit α , $\Sigma_{\alpha}^{\text{in}}$ and $d\text{-}\Sigma_{\alpha}^{\text{in}}$ are not possible.

The Relationship of the Concepts

Alvir, Greenberg, Harrison-Trainor and Turetsky (AGHTT) showed that Scott sentence complexity is related to Montalbán's Scott ranks.

SC	pSR	uSR	complexity of parameters
$\Sigma_{\alpha+2}^{\text{in}}$	α	$\alpha + 2$	$\Pi_{\alpha+1}^{\text{in}}$
$d\text{-}\Sigma_{\alpha+1}^{\text{in}}$	α	$\alpha + 1$	Π_{α}^{in}
$\Pi_{\alpha+1}^{\text{in}}$	α	α	none
α limit			
$\Sigma_{\alpha+1}^{\text{in}}$	α	$\alpha + 1$	Π_{α}^{in}
Π_{α}^{in}	α	α	none

For limit α , $\Sigma_{\alpha}^{\text{in}}$ and $d\text{-}\Sigma_{\alpha}^{\text{in}}$ are not possible.

Notice the limit case is left ambiguous in their work.

Limit levels: History of the Mystery

- ▶ In 1983, A. Miller gave examples of all possible Scott complexities except for $\Sigma_{\lambda+1}$.

Limit levels: History of the Mystery

- ▶ In 1983, A. Miller gave examples of all possible Scott complexities except for $\Sigma_{\lambda+1}$.
- ▶ 38 years later AGHTT gave an example of Scott complexity $\Sigma_{\lambda+1}$.

Limit levels: History of the Mystery

- ▶ In 1983, A. Miller gave examples of all possible Scott complexities except for $\Sigma_{\lambda+1}$.
- ▶ 38 years later AGHTT gave an example of Scott complexity $\Sigma_{\lambda+1}$.
- ▶ The example is very pretty, but quite complex.

Limit levels: History of the Mystery II

- ▶ The results of AGHTT did not give robust criteria for limit levels.

Limit levels: History of the Mystery II

- ▶ The results of AGHTT did not give robust criteria for limit levels.
- ▶ G. and Rossegger developed the following two years later.

Definition: A λ -unstable sequence in \mathcal{A} is a fundamental sequence $\delta_i \rightarrow \lambda$ along with tuple $a_i \in \mathcal{A}$ with $a_i \equiv_{\delta_i} a_{i+1}$ and $a_i \not\equiv_{\delta_{i+1}} a_{i+1}$.

Limit levels: History of the Mystery II

- ▶ The results of AGHTT did not give robust criteria for limit levels.
- ▶ G. and Rossegger developed the following two years later.

Definition: A λ -unstable sequence in \mathcal{A} is a fundamental sequence $\delta_i \rightarrow \lambda$ along with tuple $a_i \in \mathcal{A}$ with $a_i \equiv_{\delta_i} a_{i+1}$ and $a_i \not\equiv_{\delta_{i+1}} a_{i+1}$.

Theorem: \mathcal{A} with $\text{uSR}(\mathcal{A}) = \text{pSR}(\mathcal{A}) = \lambda$ has $\text{SC}(\mathcal{A}) = \Pi_{\lambda+1}$ if and only if it has some λ -unstable sequence.

Limit levels: History of the Mystery II

- ▶ The results of AGHTT did not give robust criteria for limit levels.
- ▶ G. and Rossegger developed the following two years later.

Definition: A λ -unstable sequence in \mathcal{A} is a fundamental sequence $\delta_i \rightarrow \lambda$ along with tuple $a_i \in \mathcal{A}$ with $a_i \equiv_{\delta_i} a_{i+1}$ and $a_i \not\equiv_{\delta_{i+1}} a_{i+1}$.

Theorem: \mathcal{A} with $\text{uSR}(\mathcal{A}) = \text{pSR}(\mathcal{A}) = \lambda$ has $\text{SC}(\mathcal{A}) = \Pi_{\lambda+1}$ if and only if it has some λ -unstable sequence.

Theorem: \mathcal{A} with $\text{uSR}(\mathcal{A}) = \lambda$ and $\text{pSR}(\mathcal{A}) = \lambda + 1$ has $\text{SC}(\mathcal{A}) = \Sigma_{\lambda+1}$ if and only if it has no λ -unstable sequence over some parameters.

Limit levels: History of the Mystery III

- ▶ G. and Rossegger showed there is no Linear ordering of Scott complexity Σ_3 and gave examples of all other possibilities... except for $\Sigma_{\lambda+1}$ which was left open.

Limit levels: History of the Mystery III

- ▶ G. and Rossegger showed there is no Linear ordering of Scott complexity Σ_3 and gave examples of all other possibilities... except for $\Sigma_{\lambda+1}$ which was left open.
- ▶ G., Harrison-Trainor and Ho showed there is a linear ordering of Scott complexity $\Sigma_{\lambda+1}$.

Limit levels: History of the Mystery III

- ▶ G. and Rossegger showed there is no Linear ordering of Scott complexity Σ_3 and gave examples of all other possibilities... except for $\Sigma_{\lambda+1}$ which was left open.
- ▶ G., Harrison-Trainor and Ho showed there is a linear ordering of Scott complexity $\Sigma_{\lambda+1}$.
- ▶ We will now focus on this construction and its surprising life.

The Construction: almost periodic Functions

Definition: A function $f : \mathbb{Z} \rightarrow \mathbb{N}$ is *almost periodic* if for all n $f_n := \min(f, n)$ is periodic but f itself is not periodic.

The Construction: almost periodic Functions

Definition: A function $f : \mathbb{Z} \rightarrow \mathbb{N}$ is *almost periodic* if for all n $f_n := \min(f, n)$ is periodic but f itself is not periodic.

Example: Let $\sigma_0 = 0$ and $\sigma_{i+1} = \sigma_i \frown (i+1) \frown \sigma_i$. Limits of these finite strings produce almost periodic function.

$\dots, 4, 0, 1, 0, 2, 0, 1, 0, 3, 0, 1, 0, 2, 0, 1, 0, 5, 0, 1, 0, 2, 0, 1, 0, \dots$

The Construction: almost periodic Functions

Definition: A function $f : \mathbb{Z} \rightarrow \mathbb{N}$ is *almost periodic* if for all n $f_n := \min(f, n)$ is periodic but f itself is not periodic.

Example: Let $\sigma_0 = 0$ and $\sigma_{i+1} = \sigma_i \frown (i+1) \frown \sigma_i$. Limits of these finite strings produce almost periodic function.

$\dots, 4, 0, 1, 0, 2, 0, 1, 0, 3, 0, 1, 0, 2, 0, 1, 0, 5, 0, 1, 0, 2, 0, 1, 0, \dots$

The Construction: almost periodic Functions

Definition: A function $f : \mathbb{Z} \rightarrow \mathbb{N}$ is *almost periodic* if for all n $f_n := \min(f, n)$ is periodic but f itself is not periodic.

Example: Let $\sigma_0 = 0$ and $\sigma_{i+1} = \sigma_i \frown (i+1) \frown \sigma_i$. Limits of these finite strings produce almost periodic function.

$\dots, 4, 0, 1, 0, 2, 0, 1, 0, 3, 0, 1, 0, 2, 0, 1, 0, 5, 0, 1, 0, 2, 0, 1, 0, \dots$

The Construction: almost periodic Functions

Definition: A function $f : \mathbb{Z} \rightarrow \mathbb{N}$ is *almost periodic* if for all n $f_n := \min(f, n)$ is periodic but f itself is not periodic.

Example: Let $\sigma_0 = 0$ and $\sigma_{i+1} = \sigma_i \frown (i+1) \frown \sigma_i$. Limits of these finite strings produce almost periodic function.

$\dots, 4, 0, 1, 0, 2, 0, 1, 0, 3, 0, 1, 0, 2, 0, 1, 0, 5, 0, 1, 0, 2, 0, 1, 0, \dots$

The Construction: almost periodic Functions

Definition: A function $f : \mathbb{Z} \rightarrow \mathbb{N}$ is *almost periodic* if for all n $f_n := \min(f, n)$ is periodic but f itself is not periodic.

Example: Let $\sigma_0 = 0$ and $\sigma_{i+1} = \sigma_i \frown (i+1) \frown \sigma_i$. Limits of these finite strings produce almost periodic function.

$\dots, 4, 0, 1, 0, 2, 0, 1, 0, 3, 0, 1, 0, 2, 0, 1, 0, 5, 0, 1, 0, 2, 0, 1, 0, \dots$

The Construction: almost periodic Functions

Definition: A function $f : \mathbb{Z} \rightarrow \mathbb{N}$ is *almost periodic* if for all n $f_n := \min(f, n)$ is periodic but f itself is not periodic.

Example: Let $\sigma_0 = 0$ and $\sigma_{i+1} = \sigma_i \frown (i+1) \frown \sigma_i$. Limits of these finite strings produce almost periodic function.

$\cdots, 4, 0, 1, 0, 2, 0, 1, 0, 3, 0, 1, 0, 2, 0, 1, 0, 5, 0, 1, 0, 2, 0, 1, 0, \cdots$

The Construction: almost periodic Functions

Definition: A function $f : \mathbb{Z} \rightarrow \mathbb{N}$ is *almost periodic* if for all n $f_n := \min(f, n)$ is periodic but f itself is not periodic.

Example: Let $\sigma_0 = 0$ and $\sigma_{i+1} = \sigma_i \frown (i+1) \frown \sigma_i$. Limits of these finite strings produce almost periodic function.

$\dots, 4, 0, 1, 0, 2, 0, 1, 0, 3, 0, 1, 0, 2, 0, 1, 0, 5, 0, 1, 0, 2, 0, 1, 0, \dots$

The Construction: λ -mixable pairs

Definition

An ordered pair $(\{L_i\}_{i \in \omega}, K)$ of a sequence of linear orderings $\{L_i\}_{i \in \omega}$ and a single linear ordering K is called a λ -mixable pair if the following properties hold for some non-zero fundamental sequence $\delta_i \rightarrow \lambda$:

1. $\text{uSR}(K) < \lambda$
2. $\text{uSR}(L_i) < \lambda$

The Construction: λ -mixable pairs

Definition

An ordered pair $(\{L_i\}_{i \in \omega}, K)$ of a sequence of linear orderings $\{L_i\}_{i \in \omega}$ and a single linear ordering K is called a λ -mixable pair if the following properties hold for some non-zero fundamental sequence $\delta_i \rightarrow \lambda$:

1. $\text{uSR}(K) < \lambda$
2. $\text{uSR}(L_i) < \lambda$
3. $L_i \equiv_{\delta_i} L_{i+1}$
4. $L_i \not\equiv_{\delta_{i+1}} L_{i+1}$

The Construction: λ -mixable pairs

Definition

An ordered pair $(\{L_i\}_{i \in \omega}, K)$ of a sequence of linear orderings $\{L_i\}_{i \in \omega}$ and a single linear ordering K is called a λ -mixable pair if the following properties hold for some non-zero fundamental sequence $\delta_i \rightarrow \lambda$:

1. $\text{uSR}(K) < \lambda$
2. $\text{uSR}(L_i) < \lambda$
3. $L_i \equiv_{\delta_i} L_{i+1}$
4. $L_i \not\equiv_{\delta_{i+1}} L_{i+1}$
5. any finite alternating sum
 $1 + L_{a_0} + 1 + K + 1 + L_{a_1} + 1 + K + \cdots + 1 + L_{a_n}$ has intervals isomorphic to K only within the written K blocks (or as the entire written K block).

The Construction: λ -mixable pair examples

► $L_i = \omega^{\delta_i}$ and $K = \zeta$

The Construction: λ -mixable pair examples

- ▶ $L_i = \omega^{\delta_i}$ and $K = \zeta$
- ▶ $L_i = \zeta^{\delta_i}$ and $K = \omega$

The Construction: λ -mixable pair examples

- ▶ $L_i = \omega^{\delta_i}$ and $K = \zeta$
- ▶ $L_i = \zeta^{\delta_i}$ and $K = \omega$
- ▶ $L_i = \sum_{n < i} (n + \zeta^{\delta_n}) + \sum_{n \geq i} (n + \zeta^{\delta_i})$ and $K = \omega$.

The Construction

Given any λ -mixable pair $(\{L_i\}_{i \in \omega}, K)$ and almost periodic function $f : \mathbb{Z} \rightarrow \mathbb{N}$,

$$L_f = \sum_{n \in \mathbb{Z}} (1 + L_{f(n)} + 1 + K)$$

has Scott complexity $\Sigma_{\lambda+1}$.

The Construction

Given any λ -mixable pair $(\{L_i\}_{i \in \omega}, K)$ and almost periodic function $f : \mathbb{Z} \rightarrow \mathbb{N}$,

$$L_f = \sum_{n \in \mathbb{Z}} (1 + L_{f(n)} + 1 + K)$$

has Scott complexity $\Sigma_{\lambda+1}$.

Example:

$$\cdots + \omega + \zeta + \omega^2 + \zeta + \omega + \zeta + \omega^3 + \zeta + \omega + \zeta + \omega^2 + \zeta + \omega + \cdots$$

Going Further: The Structure of almost periodic Functions

Definition: $f \in E_{\mathbb{Z}} g$ if there is an $n \in \mathbb{Z}$ such that for all $m \in \mathbb{Z}$
 $f(m) = g(m + n)$.

Going Further: The Structure of almost periodic Functions

Definition: $f \in \mathbb{Z}g$ if there is an $n \in \mathbb{Z}$ such that for all $m \in \mathbb{Z}$
 $f(m) = g(m + n)$.

Definition: $f \in_{fin} g$ if for all $n \in \mathbb{Z}$ $f_n \in \mathbb{Z}g_n$.

Going Further: The Structure of almost periodic Functions

Definition: $fE_{\mathbb{Z}}g$ if there is an $n \in \mathbb{Z}$ such that for all $m \in \mathbb{Z}$ $f(m) = g(m + n)$.

Definition: $fE_{fin}g$ if for all n $f_nE_{\mathbb{Z}}g_n$.

Translation: $fE_{\mathbb{Z}}g$ if and only if $L_f \cong L_g$. $fE_{fin}g$ if and only if $L_f \equiv_{\lambda} L_g$.

Going Further: Similar and Simple

There were no known previous examples of a pair \mathcal{A}, \mathcal{B} with $SC(\mathcal{A}) = SC(\mathcal{B}) = \Sigma_{\alpha+1}$ with $\mathcal{A} \equiv_{\alpha} \mathcal{B}$ (even away from the limit).

Going Further: Similar and Simple

There were no known previous examples of a pair \mathcal{A}, \mathcal{B} with $SC(\mathcal{A}) = SC(\mathcal{B}) = \Sigma_{\alpha+1}$ with $\mathcal{A} \equiv_{\alpha} \mathcal{B}$ (even away from the limit).

We found continuum many \equiv_{λ} -equivalent linear orderings all with Scott complexity $\Sigma_{\alpha+1}$.

Going Further: Similar and Simple

There were no known previous examples of a pair \mathcal{A}, \mathcal{B} with $SC(\mathcal{A}) = SC(\mathcal{B}) = \Sigma_{\alpha+1}$ with $\mathcal{A} \equiv_{\alpha} \mathcal{B}$ (even away from the limit).

We found continuum many \equiv_{λ} -equivalent linear orderings all with Scott complexity $\Sigma_{\alpha+1}$.

Combinatorial solution: There are continuum many E_{fin} classes in a single $E_{\mathbb{Z}}$ class. In fact, $E_0 \leq_B E_{\mathbb{Z}}|_{[b]_{E_{fin}}}$.

Thank you!

Going Further: Scott Skipping

- ▶ In 2013 Montalbán asked if there could be a Π_α sentence φ where every $M \models \varphi$ has $\text{uSR}(M) > \alpha$.

Going Further: Scott Skipping

- ▶ In 2013 Montalbán asked if there could be a Π_α sentence φ where every $M \models \varphi$ has $\text{uSR}(M) > \alpha$.
- ▶ In his thesis, Matthew Harrison-Trainer carefully produced a Π_2 formula with only models of Scott rank α for any α .

Going Further: Scott Skipping

- ▶ In 2013 Montalbán asked if there could be a Π_α sentence φ where every $M \models \varphi$ has $\text{uSR}(M) > \alpha$.
- ▶ In his thesis, Matthew Harrison-Trainer carefully produced a Π_2 formula with only models of Scott rank α for any α .
- ▶ This was the only known example

Theorem: There is a Scott complexity $\Sigma_{\lambda+1}$ linear ordering L such that if $M \equiv_\lambda L$ then $\text{SC}(M) = \Sigma_{\lambda+1}$.

Going Further: Scott Skipping

Naive Desire: Find a L_f such that all $M \equiv_\lambda L_f$ are isomorphic to L_g for a semi-periodic g .

Going Further: Scott Skipping

Naive Desire: Find a L_f such that all $M \equiv_\lambda L_f$ are isomorphic to L_g for a semi-periodic g .

Close! Need to extend semi-periodic functions to output infinite values $f : \mathbb{Z} \rightarrow \mathbb{N} \cup \{\infty\}$.

Going Further: Scott Skipping

Naive Desire: Find a L_f such that all $M \equiv_\lambda L_f$ are isomorphic to L_g for a semi-periodic g .

Close! Need to extend semi-periodic functions to output infinite values $f : \mathbb{Z} \rightarrow \mathbb{N} \cup \{\infty\}$.

Example: The 2-adic valuation - it is infinite at 0.

Going Further: Scott Skipping

Naive Desire: Find a L_f such that all $M \equiv_\lambda L_f$ are isomorphic to L_g for a semi-periodic g .

Close! Need to extend semi-periodic functions to output infinite values $f : \mathbb{Z} \rightarrow \mathbb{N} \cup \{\infty\}$.

Example: The 2-adic valuation - it is infinite at 0.

Everything works out using a λ -mixable pair that has a unique (i.e. Scott complexity Π_λ) limit structure like

$$L_i = \sum_{n < i} (n + \zeta^{\delta_n}) + \sum_{n \geq i} (n + \zeta^{\delta_i}).$$

A Dual Surprise

Theorem: Any structure \mathcal{A} is \equiv_λ to at most one Scott complexity $\Pi_{\lambda+1}$ structure. Every \equiv_λ -class contains a structure that is no Scott complexity $\Pi_{\lambda+1}$.

A Dual Surprise

Theorem: Any structure \mathcal{A} is \equiv_λ to at most one Scott complexity $\Pi_{\lambda+1}$ structure. Every \equiv_λ -class contains a structure that is no Scott complexity $\Pi_{\lambda+1}$.

This means there is no analogue for Harrison-Trainor's general Scott skipping result at the level of Scott complexity.

A Dual Surprise

Theorem: Any structure \mathcal{A} is \equiv_λ to at most one Scott complexity $\Pi_{\lambda+1}$ structure. Every \equiv_λ -class contains a structure that is no Scott complexity $\Pi_{\lambda+1}$.

This means there is no analogue for Harrison-Trainor's general Scott skipping result at the level of Scott complexity.

A new paradigm: $\Sigma_{\lambda+1}$ was considered rare for sociological reasons, $\Pi_{\lambda+1}$ is actually rare even if it has very simple examples like ω^λ .