

Scott ranks of models of arithmetic

Mateusz Łełyk

Faculty of Philosophy, University of Warsaw

15.07.2024



The first-order arithmetic

The language of arithmetic \mathcal{L}_{PA} , officially contains

The first-order arithmetic

The language of arithmetic \mathcal{L}_{PA} , officially contains

- constants 0, 1,

The first-order arithmetic

The language of arithmetic \mathcal{L}_{PA} , officially contains

- constants $0, 1$,
- function symbols $+, \times$,

The first-order arithmetic

The language of arithmetic \mathcal{L}_{PA} , officially contains

- constants $0, 1$,
- function symbols $+, \times$,
- relation $<$.

The first-order arithmetic

The language of arithmetic \mathcal{L}_{PA} , officially contains

- constants $0, 1$,
- function symbols $+, \times$,
- relation $<$.

Definition

The system PA^- contains the axioms of non-negative parts of discretely-ordered rings.

The first-order arithmetic

The language of arithmetic \mathcal{L}_{PA} , officially contains

- constants 0, 1,
- function symbols $+$, \times ,
- relation $<$.

Definition

The system PA^- contains the axioms of non-negative parts of discretely-ordered rings. PA extends PA^- with the induction scheme, to wit, the collection of sentences

$$\phi(0) \wedge \forall x(\phi(x) \rightarrow \phi(x+1)) \longrightarrow \forall x\phi(x),$$

where ϕ might contain free variables other than x .

The first-order arithmetic

The language of arithmetic \mathcal{L}_{PA} , officially contains

- constants 0, 1,
- function symbols $+$, \times ,
- relation $<$.

Definition

The system PA^- contains the axioms of non-negative parts of discretely-ordered rings. PA extends PA^- with the induction scheme, to wit, the collection of sentences

$$\phi(0) \wedge \forall x(\phi(x) \rightarrow \phi(x+1)) \longrightarrow \forall x\phi(x),$$

where ϕ might contain free variables other than x .

Remark

PA is biinterpretable with $\text{ZF} \setminus \{\text{InfTy}\} + \neg\text{InfTy} + \text{TC}$.

Some salient classes of models of PA

Suppose $\mathcal{M} \models \text{PA}$

- Prime models. Let $K(\mathcal{M})$ be the submodel of \mathcal{M} generated by $\text{Def}(\mathcal{M})$.

Some salient classes of models of PA

Suppose $\mathcal{M} \models \text{PA}$

- Prime models. Let $K(\mathcal{M})$ be the submodel of \mathcal{M} generated by $\text{Def}(\mathcal{M})$. Then $K(\mathcal{M}) \preceq \mathcal{M}$.

Some salient classes of models of PA

Suppose $\mathcal{M} \models \text{PA}$

- Prime models. Let $K(\mathcal{M})$ be the submodel of \mathcal{M} generated by $\text{Def}(\mathcal{M})$. Then $K(\mathcal{M}) \preceq \mathcal{M}$. In fact, $K(\mathcal{M})$ is the prime model of $\text{Th}(\mathcal{M})$.

Some salient classes of models of PA

Suppose $\mathcal{M} \models \text{PA}$

- Prime models. Let $K(\mathcal{M})$ be the submodel of \mathcal{M} generated by $\text{Def}(\mathcal{M})$. Then $K(\mathcal{M}) \preceq \mathcal{M}$. In fact, $K(\mathcal{M})$ is the prime model of $\text{Th}(\mathcal{M})$.
- Finitely generated models. Let $K(\mathcal{M}, a)$ be the submodel of \mathcal{M} generated by $\text{Def}(\mathcal{M}, a)$. Then $a \in K(\mathcal{M}, a) \preceq \mathcal{M}$.

Some salient classes of models of PA

Suppose $\mathcal{M} \models \text{PA}$

- Prime models. Let $K(\mathcal{M})$ be the submodel of \mathcal{M} generated by $\text{Def}(\mathcal{M})$. Then $K(\mathcal{M}) \preceq \mathcal{M}$. In fact, $K(\mathcal{M})$ is the prime model of $\text{Th}(\mathcal{M})$.
- Finitely generated models. Let $K(\mathcal{M}, a)$ be the submodel of \mathcal{M} generated by $\text{Def}(\mathcal{M}, a)$. Then $a \in K(\mathcal{M}, a) \preceq \mathcal{M}$.
- Recursively saturated models. \mathcal{M} is recursively saturated if every recursive set of formulae with finitely many parameters which is consistent with $\text{Th}((\mathcal{M}, a)_{a \in M})$ is realized in \mathcal{M} .

Some salient classes of models of PA

Suppose $\mathcal{M} \models \text{PA}$

- Prime models. Let $K(\mathcal{M})$ be the submodel of \mathcal{M} generated by $\text{Def}(\mathcal{M})$. Then $K(\mathcal{M}) \preceq \mathcal{M}$. In fact, $K(\mathcal{M})$ is the prime model of $\text{Th}(\mathcal{M})$.
- Finitely generated models. Let $K(\mathcal{M}, a)$ be the submodel of \mathcal{M} generated by $\text{Def}(\mathcal{M}, a)$. Then $a \in K(\mathcal{M}, a) \preceq \mathcal{M}$.
- Recursively saturated models. \mathcal{M} is recursively saturated if every recursive set of formulae with finitely many parameters which is consistent with $\text{Th}((\mathcal{M}, a)_{a \in M})$ is realized in \mathcal{M} .

Remark

Prime models and countable recursively saturated models of PA are homogeneous.

The structural complexity of models of PA

Theorem (Montalbán and Rossegger, 2023)

- ① *For every $\alpha \geq \omega$ there is a model of PA of Scott rank exactly α .*
- ② *Assume that $\mathcal{M} \models \text{PA}$.*
 - *If \mathcal{M} is standard, then $\text{SR}(\mathcal{M}) = 1$. If \mathcal{M} is nonstandard, then $\text{SR}(\mathcal{M}) \geq \omega$.*
 - *If $\mathcal{M} = K(\mathcal{M})$ is nonstandard, then $\text{SR}(\mathcal{M}) = \omega$.*
 - *If \mathcal{M} is homogeneous, then $\text{SR}(\mathcal{M}) \leq \omega + 1$.*

The structural complexity of models of PA

Theorem (Montalbán and Rossegger, 2023)

- ① *For every $\alpha \geq \omega$ there is a model of PA of Scott rank exactly α .*
- ② *Assume that $\mathcal{M} \models \text{PA}$.*
 - *If \mathcal{M} is standard, then $\text{SR}(\mathcal{M}) = 1$. If \mathcal{M} is nonstandard, then $\text{SR}(\mathcal{M}) \geq \omega$.*
 - *If $\mathcal{M} = K(\mathcal{M})$ is nonstandard, then $\text{SR}(\mathcal{M}) = \omega$.*
 - *If \mathcal{M} is homogeneous, then $\text{SR}(\mathcal{M}) \leq \omega + 1$.*

Question

What are the models of PA of rank ω ?

Theorem (Ł., Szlufik)

Assume $\mathcal{M} \models \text{PA}$ and $a \in M$ is undefinable. Then the automorphism orbit of a is not $\Sigma_\omega^{\text{inf}}$ definable.

Key ingredients: Σ_n -recursive saturation.

Let $\mathcal{M} \models \text{PA}$.

Key ingredients: Σ_n -recursive saturation.

Let $\mathcal{M} \models \text{PA}$. We say that a type $p(x, a)$ *is coded in* \mathcal{M}

Key ingredients: Σ_n -recursive saturation.

Let $\mathcal{M} \models \text{PA}$. We say that a type $p(x, a)$ *is coded in* \mathcal{M} if there is a $c \in \mathcal{M}$ such that for every formula $\phi(x, y)$

$$\phi(x, a) \in p(x, a) \iff \mathcal{M} \models \ulcorner \phi(x, y) \urcorner \in c.$$

Key ingredients: Σ_n -recursive saturation.

Let $\mathcal{M} \models \text{PA}$. We say that a type $p(x, a)$ is *coded in \mathcal{M}* if there is a $c \in \mathcal{M}$ such that for every formula $\phi(x, y)$

$$\phi(x, a) \in p(x, a) \iff \mathcal{M} \models \ulcorner \phi(x, y) \urcorner \in c.$$

Lemma

Let $p(x, a)$ be a Σ_n -type over \mathcal{M} . Then

- ① If $p(x, a)$ is coded in \mathcal{M} , then $p(x, a)$ is realized in \mathcal{M} .
- ② If for some b , $p(x, a) = \{\phi(x, a) \in \Sigma_n : \mathcal{M} \models \phi(b, a)\}$, then it is coded.

Key ingredients: Σ_n -recursive saturation.

Let $\mathcal{M} \models \text{PA}$. We say that a type $p(x, a)$ is coded in \mathcal{M} if there is a $c \in \mathcal{M}$ such that for every formula $\phi(x, y)$

$$\phi(x, a) \in p(x, a) \iff \mathcal{M} \models \ulcorner \phi(x, y) \urcorner \in c.$$

Lemma

Let $p(x, a)$ be a Σ_n -type over \mathcal{M} . Then

- ① If $p(x, a)$ is coded in \mathcal{M} , then $p(x, a)$ is realized in \mathcal{M} .
- ② If for some b , $p(x, a) = \{\phi(x, a) \in \Sigma_n : \mathcal{M} \models \phi(b, a)\}$, then it is coded.

Proof (1). Assume that c codes $p(x, a)$ and consider a formula $\psi(y)$

$$\exists x \forall \phi < y (\phi \in c \rightarrow \text{Sat}_n(\phi, \langle x, a \rangle)).$$

Key ingredients: Σ_n -recursive saturation.

Let $\mathcal{M} \models \text{PA}$. We say that a type $p(x, a)$ is coded in \mathcal{M} if there is a $c \in \mathcal{M}$ such that for every formula $\phi(x, y)$

$$\phi(x, a) \in p(x, a) \iff \mathcal{M} \models \ulcorner \phi(x, y) \urcorner \in c.$$

Lemma

Let $p(x, a)$ be a Σ_n -type over \mathcal{M} . Then

- ① If $p(x, a)$ is coded in \mathcal{M} , then $p(x, a)$ is realized in \mathcal{M} .
- ② If for some b , $p(x, a) = \{\phi(x, a) \in \Sigma_n : \mathcal{M} \models \phi(b, a)\}$, then it is coded.

Proof (2). Assume d realizes $p(x, a)$ and consider the type $p'(x, a, d)$

$$\{\phi(d, a) \equiv \ulcorner \phi(v, w) \urcorner \in x : \phi \in \Sigma_n\}$$

Key ingredients: Σ_n -recursive saturation.

Lemma

Let $p(x, a)$ be a Σ_n -type over \mathcal{M} . Then

- ① If $p(x, a)$ is coded in \mathcal{M} , then $p(x, a)$ is realized in \mathcal{M} .
- ② If $p(x, a)$ is Σ_n -complete and realized in \mathcal{M} , then it is coded.

Key ingredients: Σ_n -recursive saturation.

Lemma

Let $p(x, a)$ be a Σ_n -type over \mathcal{M} . Then

- ① If $p(x, a)$ is coded in \mathcal{M} , then $p(x, a)$ is realized in \mathcal{M} .
- ② If $p(x, a)$ is Σ_n -complete and realized in \mathcal{M} , then it is coded.

Lemma

Suppose $\mathcal{M} \models \text{PA}$, $a, b \in M$. Then, for every n , the following are equivalent:

- Σ_n^{inf} -types of a, b are the same.
- Σ_n -types of a, b are the same.

Key ingredients: The Ehrenfeucht Lemma

Lemma (Ehrenfeucht)

If $\mathcal{M} \models \text{PA}$, $a \neq b \in M$ and b is definable from a , then the types of a and b are different.

Key ingredients: The Ehrenfeucht Lemma

Lemma (Ehrenfeucht)

If $\mathcal{M} \models \text{PA}$, $a \neq b \in M$ and b is definable from a , then the types of a and b are different.

Proof sketch (due to Fuchs-Gitman-Hamkins): assume $F(a) = b$, where F is an \mathcal{M} -definable total function.

Key ingredients: The Ehrenfeucht Lemma

Lemma (Ehrenfeucht)

If $\mathcal{M} \models \text{PA}$, $a \neq b \in M$ and b is definable from a , then the types of a and b are different.

Proof sketch (due to Fuchs-Gitman-Hamkins): assume $F(a) = b$, where F is an \mathcal{M} -definable total function. Assume $a < b$ (the other case being symmetric).

Key ingredients: The Ehrenfeucht Lemma

Lemma (Ehrenfeucht)

If $\mathcal{M} \models \text{PA}$, $a \neq b \in M$ and b is definable from a , then the types of a and b are different.

Proof sketch (due to Fuchs-Gitman-Hamkins): assume $F(a) = b$, where F is an \mathcal{M} -definable total function. Assume $a < b$ (the other case being symmetric). Put

$$xRy \iff F(y) = x \wedge y < x.$$

Key ingredients: The Ehrenfeucht Lemma

Lemma (Ehrenfeucht)

If $\mathcal{M} \models \text{PA}$, $a \neq b \in M$ and b is definable from a , then the types of a and b are different.

Proof sketch (due to Fuchs-Gitman-Hamkins): assume $F(a) = b$, where F is an \mathcal{M} -definable total function. Assume $a < b$ (the other case being symmetric). Put

$$xRy \iff F(y) = x \wedge y < x.$$

Then (\mathcal{M}, R) is a forest and we have bRa .

Key ingredients: The Ehrenfeucht Lemma

Lemma (Ehrenfeucht)

If $\mathcal{M} \models \text{PA}$, $a \neq b \in M$ and b is definable from a , then the types of a and b are different.

Proof sketch (due to Fuchs-Gitman-Hamkins): assume $F(a) = b$, where F is an \mathcal{M} -definable total function. Assume $a < b$ (the other case being symmetric). Put

$$xRy \iff F(y) = x \wedge y < x.$$

Then (\mathcal{M}, R) is a forest and we have bRa . Now a, b differ on the following formula

"The shortest path between me and the least element of my connected component is of even length".

Proof sketch

Assume that $a \in M$ is undefinable but the automorphism orbit of a is Σ_{ω}^{inf} -definable.

Proof sketch

Assume that $a \in M$ is undefinable but the automorphism orbit of a is Σ_ω^{inf} -definable. It follows that for some $n \in \omega$, the orbit of a is Σ_n^{inf} -definable.

Assume that $a \in M$ is undefinable but the automorphism orbit of a is Σ_ω^{inf} -definable. It follows that for some $n \in \omega$, the orbit of a is Σ_n^{inf} -definable. By the lemmas it is enough to find $b \in K(\mathcal{M}, a)$ which has the same Σ_n type as a .

Assume that $a \in M$ is undefinable but the automorphism orbit of a is Σ_ω^{inf} -definable. It follows that for some $n \in \omega$, the orbit of a is Σ_n^{inf} -definable. By the lemmas it is enough to find $b \in K(\mathcal{M}, a)$ which has the same Σ_n type as a . By MOPA folklore $K(\mathcal{M}, a) \prec \mathcal{M}$.

Assume that $a \in M$ is undefinable but the automorphism orbit of a is Σ_ω^{inf} -definable. It follows that for some $n \in \omega$, the orbit of a is Σ_n^{inf} -definable. By the lemmas it is enough to find $b \in K(\mathcal{M}, a)$ which has the same Σ_n type as a . By MOPA folklore $K(\mathcal{M}, a) \prec \mathcal{M}$. By induction, the Σ_n -type of a is coded (in $K(\mathcal{M}, a)$) and so is the type

$$p(x) := \{\phi(x) \wedge x \neq a : K(\mathcal{M}, a) \models \phi(a), \phi \in \Sigma_n\}.$$

Proof sketch

Assume that $a \in M$ is undefinable but the automorphism orbit of a is Σ_ω^{inf} -definable. It follows that for some $n \in \omega$, the orbit of a is Σ_n^{inf} -definable. By the lemmas it is enough to find $b \in K(\mathcal{M}, a)$ which has the same Σ_n type as a . By MOPA folklore $K(\mathcal{M}, a) \prec \mathcal{M}$. By induction, the Σ_n -type of a is coded (in $K(\mathcal{M}, a)$) and so is the type

$$p(x) := \{\phi(x) \wedge x \neq a : K(\mathcal{M}, a) \models \phi(a), \phi \in \Sigma_n\}.$$

By induction and the undefinability of a , it follows that $p(x)$ is realized in $K(\mathcal{M}, a)$ and we are done.

Definition (Kalociński)

Theory T has the simplest model property, SMP, if up to an iso, T has exactly one model of the least rank.

Definition (Kalociński)

Theory T has the simplest model property, SMP, if up to an iso, T has exactly one model of the least rank.

Corollary

Every completion of PA has SMP.

Consequences

Definition (Kalociński)

Theory T has the simplest model property, SMP, if up to an iso, T has exactly one model of the least rank.

Corollary

Every completion of PA has SMP.

Corollary

If $\mathcal{M} \models \text{Th}(\mathbb{N})$ is nonstandard, then $\text{SR}(\mathcal{M}) > \omega$.

Consequences

Definition (Kalociński)

Theory T has the simplest model property, SMP, if up to an iso, T has exactly one model of the least rank.

Corollary

Every completion of PA has SMP.

Corollary

If $\mathcal{M} \models \text{Th}(\mathbb{N})$ is nonstandard, then $\text{SR}(\mathcal{M}) > \omega$.

Corollary

If \mathcal{M} is recursively saturated, then $\text{SR}(\mathcal{M}) = \omega + 1$.

Proposition

If \mathcal{M} is finitely generated by an undefinable element, then $\text{SR}(\mathcal{M}) = \omega + 1$.

Proposition

If \mathcal{M} is finitely generated by an undefinable element, then $\text{SR}(\mathcal{M}) = \omega + 1$.

Proposition

The models of PA with the parametrized Scott rank ω are exactly the finitely generated models of PA.

Fragments of PA

Recall that

- the quantifier $\forall x\phi$ ($\exists x\phi$) is *bounded* iff ϕ is of the form $x < t \rightarrow \psi$ ($x < t \wedge \psi$) and t does not mention x .
- ϕ is Δ_0 if all the quantifiers in ϕ are bounded.

Fragments of PA

Recall that

- the quantifier $\forall x\phi$ ($\exists x\phi$) is *bounded* iff ϕ is of the form $x < t \rightarrow \psi$ ($x < t \wedge \psi$) and t does not mention x .
- ϕ is Δ_0 if all the quantifiers in ϕ are bounded.

$$\Sigma_0 = \Pi_0 = \Delta_0$$

$$\Sigma_{n+1} = \exists x \Pi_n$$

$$\Pi_{n+1} = \forall x \Sigma_n$$

Fragments of PA

Recall that

- the quantifier $\forall x\phi$ ($\exists x\phi$) is *bounded* iff ϕ is of the form $x < t \rightarrow \psi$ ($x < t \wedge \psi$) and t does not mention x .
- ϕ is Δ_0 if all the quantifiers in ϕ are bounded.

$$\Sigma_0 = \Pi_0 = \Delta_0$$

$$\Sigma_{n+1} = \exists x \Pi_n$$

$$\Pi_{n+1} = \forall x \Sigma_n$$

For a class of formulae Γ , $I\Gamma$ denotes the extension of PA^- with induction restricted to formulae from Γ .

Fragments of PA

Recall that

- the quantifier $\forall x\phi$ ($\exists x\phi$) is *bounded* iff ϕ is of the form $x < t \rightarrow \psi$ ($x < t \wedge \psi$) and t does not mention x .
- ϕ is Δ_0 if all the quantifiers in ϕ are bounded.

$$\Sigma_0 = \Pi_0 = \Delta_0$$

$$\Sigma_{n+1} = \exists x \Pi_n$$

$$\Pi_{n+1} = \forall x \Sigma_n$$

For a class of formulae Γ , $I\Gamma$ denotes the extension of PA^- with induction restricted to formulae from Γ .

Remark

$I\Sigma_1$ proves MRDP theorem, so actually over $I\Sigma_1$ Σ_n -definable sets are \exists_n -definable.

Proposition

If $\mathcal{M} \models \text{I}\Sigma_n$ and \mathcal{M} contains a nonstandard definable element, then $\text{SR}(\mathcal{M}) > n$.

Proposition

If $\mathcal{M} \models \text{I}\Sigma_n$ and \mathcal{M} contains a nonstandard definable element, then $\text{SR}(\mathcal{M}) > n$.

Corollary

Suppose $\mathcal{M} \models \text{I}\Sigma_1$ is nonstandard. Then $\text{SR}(\mathcal{M}) > 1$.

Proposition

If $\mathcal{M} \models \text{I}\Sigma_n$ and \mathcal{M} contains a nonstandard definable element, then $\text{SR}(\mathcal{M}) > n$.

Corollary

Suppose $\mathcal{M} \models \text{I}\Sigma_1$ is nonstandard. Then $\text{SR}(\mathcal{M}) > 1$.

Corollary

$\text{I}\Sigma_1$ has SMP.

Proposition

If $\mathcal{M} \models I\Sigma_n$ and \mathcal{M} contains a nonstandard definable element, then $\text{SR}(\mathcal{M}) > n$.

Corollary

Suppose $\mathcal{M} \models I\Sigma_1$ is nonstandard. Then $\text{SR}(\mathcal{M}) > 1$.

Corollary

$I\Sigma_1$ has SMP. But there are nonstandard models of $\Pi_2\text{-Cons}(I\Sigma_1)$ of rank 1.

Lemma (Essentially Ehrenfeucht)

Suppose that $\mathcal{M} \models \text{IS}_{n+1}$ and $a \neq b \in M$ are such that b is Σ_n definable from a . Then Σ_{n+2} -types of a, b are different.

Lemma (Essentially Ehrenfeucht)

Suppose that $\mathcal{M} \models I\Sigma_{n+1}$ and $a \neq b \in M$ are such that b is Σ_n definable from a . Then Σ_{n+2} -types of a, b are different.

Proposition

Suppose $\mathcal{M} \models I\Sigma_{n+1}$ is nonstandard. Then $\text{SR}(\mathcal{M}) \geq n$.