



# Exploiting Imprecise Information Sources in Sequential Decision Making Problems under Uncertainty.

Ph.D defense **N.Drougard**

doctoral school: EDSYS,

institution: ISAE-SUPAERO,

laboratory: ONERA-The French Aerospace Lab.



retour sur innovation


- 1 Context
- 2 An hybrid POMDP
- 3 Benefiting from factorized structures
- 4 Conclusion/Perspectives

- 1 Context
- 2 An hybrid POMDP
- 3 Benefiting from factorized structures
- 4 Conclusion/Perspectives

# Context

## Partially Observable Markov Decision Process (POMDP)


**POMDP:** model for sequential decision making under uncertainty



# Context

## Partially Observable Markov Decision Process (POMDP)

**POMDP:** model for sequential decision making under uncertainty




$s \in \mathcal{S}$ : **system states;**

# Context

## Partially Observable Markov Decision Process (POMDP)


**POMDP:** model for sequential decision making under uncertainty



# Context

## Partially Observable Markov Decision Process (POMDP)



**POMDP:** model for sequential decision making under uncertainty



# Context

## Partially Observable Markov Decision Process (POMDP)

**POMDP:** model for sequential decision making under uncertainty







# Context

## Partially Observable Markov Decision Process (POMDP)



**POMDP:** model for sequential decision making under uncertainty



# Context

## Partially Observable Markov Decision Process (POMDP)

**POMDP:** model for sequential decision making under uncertainty



## Partially Observable Markov Decision Process (POMDP)

$o \in \mathcal{O}$ : observations;

$a \in \mathcal{A}$ : agent's actions;

***b*: belief state.**



# Context

belief state, strategy, criterion.

**POMDP:**  $\langle \mathcal{S}, \mathcal{A}, \mathcal{O}, T, O, r, \gamma \rangle$ ,

- **transition** function  $T(s, a, s') = \mathbf{p}(s' \mid s, a)$ ;
- **observation** function  $O(s', a, o') = \mathbf{p}(o' \mid s', a)$ .

# Context

belief state, strategy, criterion.

**POMDP:**  $\langle \mathcal{S}, \mathcal{A}, \mathcal{O}, T, O, r, \gamma \rangle$ ,

- **transition** function  $T(s, a, s') = \mathbf{p}(s' \mid s, a)$ ;
- **observation** function  $O(s', a, o') = \mathbf{p}(o' \mid s', a)$ .

**belief state:**  $b_t(s) = \mathbb{P}(s_t = s \mid a_0, o_1, \dots, a_{t-1}, o_t)$

# Context

belief state, strategy, criterion.

**POMDP:**  $\langle \mathcal{S}, \mathcal{A}, \mathcal{O}, T, O, r, \gamma \rangle$ ,

- **transition** function  $T(s, a, s') = \mathbf{p}(s' | s, a)$ ;
- **observation** function  $O(s', a, o') = \mathbf{p}(o' | s', a)$ .

**belief state:**  $b_t(s) = \mathbb{P}(s_t = s | a_0, o_1, \dots, a_{t-1}, o_t)$

**probabilistic belief update**

$$b_{t+1}(s') \propto \mathbf{p}(o' | s', a) \cdot \sum_{s \in \mathcal{S}} \mathbf{p}(s' | s, a) \cdot b_t(s)$$

# Context

belief state, strategy, criterion.

**POMDP:**  $\langle \mathcal{S}, \mathcal{A}, \mathcal{O}, T, O, r, \gamma \rangle$ ,

- **transition** function  $T(s, a, s') = \mathbf{p}(s' | s, a)$ ;
- **observation** function  $O(s', a, o') = \mathbf{p}(o' | s', a)$ .

**belief state:**  $b_t(s) = \mathbb{P}(s_t = s | a_0, o_1, \dots, a_{t-1}, o_t)$

## probabilistic belief update

$$b_{t+1}(s') \propto \mathbf{p}(o' | s', a) \cdot \sum_{s \in \mathcal{S}} \mathbf{p}(s' | s, a) \cdot b_t(s)$$

**action choices:** strategy  $\delta(b_t) = a_t \in \mathcal{A}$

$$\text{maximizing } \mathbb{E}_{s_0 \sim b_0} \left[ \sum_{t=0}^{+\infty} \gamma^t \cdot r(s_t, \delta(b_t)) \right], \quad 0 < \gamma < 1.$$

# Flaws of the POMDP model

## POMDPs in practice

- optimal strategy computation  $\geq$  **PSPACE**;
- probabilities are **imprecisely known** in practice;
- agent's **ignorance** not taken into account.



# Why model ignorance?

knowledge is not always encouraged with POMDPs

- initial belief deterministic  $s_0 = s_A$ .


$s_A$

$$b_0(s_A) = 1$$

# Why model ignorance?

knowledge is not always encouraged with POMDPs


- initial belief deterministic  $s_0 = s_A$ .



# Why model ignorance?

knowledge is not always encouraged with POMDPs


- initial belief deterministic  $s_0 = s_A$ .



# Why model ignorance?

knowledge is not always encouraged with POMDPs

- initial belief deterministic  $s_0 = s_A$ .




- $\{s_B, s_C, s_D\} \xrightarrow{\text{deterministic}} s_E, r(s_E) = 0.$

# Why model ignorance?

knowledge is not always encouraged with POMDPs

- initial belief deterministic  $s_0 = s_A$ .



- $\{s_B, s_C, s_D\} \xrightarrow{\text{deterministic}} s_E, r(s_E) = 0$ .

$$\mathbb{E}_{s_0 \sim b_0} \left[ \sum_{t=0}^{+\infty} \gamma^t \cdot r(s_t) \mid a_0 = \tilde{a} \text{ or } a' \right] = r(s_0) + 5\gamma.$$

**the safe action is not preferred.**

# Qualitative Possibility Theory

an hybrid model with possibilistic belief states

## Qualitative Possibility Theory

- **simplification/imprecision** taken into account,  
**BUT frequentist information lost**;
- **ignorance** modeling;
- possibilistic belief states already studied:  $\pi$ -POMDP  
(Sabbadin UAI98, Drougard UAI13, AAAI14).

# Qualitative Possibility Theory

an hybrid model with possibilistic belief states

## Qualitative Possibility Theory

- **simplification/imprecision** taken into account,  
**BUT frequentist information lost**;
  - **ignorance** modeling;
  - possibilistic belief states already studied:  $\pi$ -POMDP (Sabbadin UAI98, Drougard UAI13, AAAI14).
- 
- **POMDP with possibilistic belief states**
    - heuristic for solving POMDPs;
    - results in a standard MDP.

# Qualitative Possibility Theory

an hybrid model with possibilistic belief states

## Qualitative Possibility Theory

- **simplification/imprecision** taken into account,  
**BUT frequentist information lost**;
  - **ignorance** modeling;
  - possibilistic belief states already studied:  $\pi$ -POMDP  
(Sabbadin UAI98, Drougard UAI13, AAAI14).
- 
- **POMDP with possibilistic belief states**
    - heuristic for solving POMDPs;
    - results in a standard MDP.
- 
- **defined distributions  $\pi$ :**  
 $\mathbb{P} \rightarrow \pi$  transformations: pignistic, specific, ...



# Qualitative Possibility Theory

## presentation

$1 = l_1 > l_2 > \dots > l_{\#\mathcal{L}} = 0$  form the **finite scale**  $\mathcal{L}$ .

events  $e \subset \Omega$  (universe)

**sorted** using possibility **degrees**  $\pi(e) \in \mathcal{L}$ ,

$\neq$

**quantified** with **frequencies**  $\mathbf{p}(e) \in [0, 1]$  (probabilities).

# Qualitative Possibility Theory

## presentation

$1 = l_1 > l_2 > \dots > l_{\#\mathcal{L}} = 0$  form the **finite scale**  $\mathcal{L}$ .

events  $e \subset \Omega$  (universe)

**sorted** using possibility **degrees**  $\pi(e) \in \mathcal{L}$ ,

$\neq$

**quantified** with **frequencies**  $p(e) \in [0, 1]$  (probabilities).

$e_1 \neq e_2$ , 2 events  $\subset \Omega$

■  $\pi(e_1) < \pi(e_2) \Leftrightarrow$  “ $e_1$  is less plausible than  $e_2$ ”;

# Qualitative Possibility Theory

presentation

$1 = l_1 > l_2 > \dots > l_{\#\mathcal{L}} = 0$  form the **finite scale**  $\mathcal{L}$ .

events  $e \subset \Omega$  (universe)

**sorted** using possibility **degrees**  $\pi(e) \in \mathcal{L}$ ,

$\neq$

**quantified** with **frequencies**  $\mathbf{p}(e) \in [0, 1]$  (probabilities).

$e_1 \neq e_2$ , 2 events  $\subset \Omega$

■  $\pi(e_1) < \pi(e_2) \Leftrightarrow$  “ $e_1$  is less plausible than  $e_2$ ”;

**Probability ( $\mathbb{P}$ )** / **Possibility ( $\Pi$ ):**

$e_1$ or $e_2$	$\mathbf{p}(e_1) + \mathbf{p}(e_2 \cap \overline{e_1})$	$\max \{ \pi(e_1), \pi(e_2) \}$
$e_1$ and $e_2$	$\mathbf{p}(e_1) \cdot \mathbf{p}(e_2 \mid e_1)$	$\min \{ \pi(e_1), \pi(e_2 \mid e_1) \}$

- 1 Context
- 2 An hybrid POMDP
- 3 Benefiting from factorized structures
- 4 Conclusion/Perspectives

# A possibilistic belief state

belief space discretization

$$\Pi_S = \left\{ \text{possibility distributions} \right\}: \# \Pi_S < +\infty$$

→ **belief space discretization.**

# A possibilistic belief state

belief space discretization

$$\Pi_S = \left\{ \text{possibility distributions} \right\}: \# \Pi_S < +\infty$$

→ **belief space discretization.**

$$b_t^\pi(s) = \pi(s_t = s \mid a_0, o_1, \dots, a_{t-1}, o_t)$$

# A possibilistic belief state

belief space discretization

$$\Pi_S = \left\{ \text{possibility distributions} \right\}: \# \Pi_S < +\infty$$

→ **belief space discretization.**

$$b_t^\pi(s) = \pi(s_t = s \mid a_0, o_1, \dots, a_{t-1}, o_t)$$

update – **possibilistic** belief state

$$b_{t+1}^\pi(s') = \begin{cases} 1 & \text{if } \pi(o', s' \mid b_t^\pi, a) = \pi(o' \mid b_t^\pi, a) \\ \pi(o', s' \mid b_t^\pi, a) & \text{otherwise.} \end{cases}$$

denoted by  $b_{t+1}^\pi(s') \propto^\pi \pi(o', s' \mid b_t^\pi, a)$

# A possibilistic belief state

belief space discretization

$$\Pi_S = \left\{ \text{possibility distributions} \right\}: \# \Pi_S < +\infty$$

→ **belief space discretization.**

$$b_t^\pi(s) = \pi(s_t = s \mid a_0, o_1, \dots, a_{t-1}, o_t)$$

update – **possibilistic** belief state

$$b_{t+1}^\pi(s') = \begin{cases} 1 & \text{if } \pi(o', s' \mid b_t^\pi, a) = \pi(o' \mid b_t^\pi, a) \\ \pi(o', s' \mid b_t^\pi, a) & \text{otherwise.} \end{cases}$$

denoted by  $b_{t+1}^\pi(s') \propto^\pi \pi(o', s' \mid b_t^\pi, a)$

- $\pi(o', s' \mid b_t^\pi, a) = \max_{s \in \mathcal{S}} \min \left\{ \pi(o' \mid s', a), \pi(s' \mid s, a), b_t^\pi(s) \right\};$
- $\pi(o' \mid s', a) = \max_{s' \in \mathcal{S}} \pi(o', s' \mid b_t^\pi, a).$



# A possibilistic belief state

belief space discretization

$$\Pi_S = \left\{ \text{possibility distributions} \right\}: \# \Pi_S < +\infty$$

→ **belief space discretization.**

$$b_t^\pi(s) = \pi(s_t = s \mid a_0, o_1, \dots, a_{t-1}, o_t)$$

update – **possibilistic** belief state

$$b_{t+1}^\pi(s') = \begin{cases} 1 & \text{if } \pi(o', s' \mid b_t^\pi, a) = \pi(o' \mid b_t^\pi, a) \\ \pi(o', s' \mid b_t^\pi, a) & \text{otherwise.} \end{cases}$$

denoted by  $b_{t+1}^\pi(s') \propto^\pi \pi(o', s' \mid b_t^\pi, a)$

- the update **only depends on  $o'$  and  $a$ .**

# Pignistic transformation and transitions

## Pignistic transformation

numbering of the  $n = \#\mathcal{S}$  system states:

$$1 = b^\pi(s_1) \geq \dots \geq b^\pi(s_n) \geq b^\pi(s_{n+1}) = 0.$$

**pignistic transformation** –  $P : \Pi_{\mathcal{S}} \rightarrow \mathbb{P}_{\mathcal{S}}$

$$\overline{b^\pi}(s_i) = \sum_{j=i}^{\#\mathcal{S}} \frac{b^\pi(s_j) - b^\pi(s_{j+1})}{j}.$$

- probability distribution  $\overline{b^\pi} =$  **gravity center** of the represented probabilistic distributions;
- **Laplace principle**: ignorance  $\rightarrow$  uniform probability.

# Pignistic transformation

Examples of pignistic transformations (red) of possibility distributions (blue)

# Pignistic transformation and transitions

## Transition function of epistemic states

Approximation of the probabilities over the observations:

- $\mathbf{p}(o' \mid s, a) = \sum_{s' \in \mathcal{S}} O(s', a, o') \cdot T(s, a, s')$ ;
- $\mathbf{p}(o' \mid b^\pi, a) := \sum_{s \in \mathcal{S}} \mathbf{p}(o' \mid s, a) \cdot \overline{b^\pi}(s).$

$$\Rightarrow \mathbf{p}\left((b^\pi)' \mid b^\pi, a\right) = \sum_{\substack{o' \text{ t.q.} \\ u(b^\pi, a, o') = (b^\pi)'}} \mathbf{p}(o' \mid b^\pi, a).$$

notation: if  $a \in \mathcal{A}$  selected,  $o' \in \mathcal{O}$  received,

$$b_{t+1}^\pi = u(o', a, b_t^\pi) = \text{update of } b_t^\pi.$$

# Choquet integral and rewards

pessimistic evaluation of the rewards – necessity measure

imprecision of  $b^\pi$  = agent ignorance + discretization:  
**pessimistic reward** about these imprecisions.

# Choquet integral and rewards

pessimistic evaluation of the rewards – necessity measure

imprecision of  $b^\pi$  = agent ignorance + discretization:  
**pessimistic reward** about these imprecisions.

Dual measure of  $\Pi : 2^{\mathcal{S}} \rightarrow \mathcal{L}$

necessity  $\mathcal{N}$  such that  $\forall A \subseteq \mathcal{S}, \mathcal{N}(A) = 1 - \Pi(\overline{A})$ .

# Choquet integral and rewards

pessimistic evaluation of the rewards – necessity measure

imprecision of  $b^\pi$  = agent ignorance + discretization:

**pessimistic reward** about these imprecisions.

Dual measure of  $\Pi : 2^{\mathcal{S}} \rightarrow \mathcal{L}$

necessity  $\mathcal{N}$  such that  $\forall A \subseteq \mathcal{S}, \mathcal{N}(A) = 1 - \Pi(\bar{A})$ .

$r_1 > r_2 > \dots > r_{k+1} = 0$  represents elements of  $\{r(s, a) | s \in \mathcal{S}\}$ .

Choquet integral of  $r$  with respect to  $\mathcal{N}$

$$Ch(r, \mathcal{N}) = \sum_{i=1}^k (r_i - r_{i+1}) \cdot \mathcal{N}(\{r(s) \geq r_i\}) \quad (1)$$

(2)

# Choquet integral and rewards

pessimistic evaluation of the rewards – necessity measure

imprecision of  $b^\pi$  = agent ignorance + discretization:

**pessimistic reward** about these imprecisions.

Dual measure of  $\Pi : 2^{\mathcal{S}} \rightarrow \mathcal{L}$

necessity  $\mathcal{N}$  such that  $\forall A \subseteq \mathcal{S}, \mathcal{N}(A) = 1 - \Pi(\bar{A})$ .

$r_1 > r_2 > \dots > r_{k+1} = 0$  represents elements of  $\{r(s, a) | s \in \mathcal{S}\}$ .

Choquet integral of  $r$  with respect to  $\mathcal{N}$

$$Ch(r, \mathcal{N}) = \sum_{i=1}^k (r_i - r_{i+1}) \cdot \mathcal{N}(\{r(s) \geq r_i\}) \quad (1)$$


$$= \sum_{i=1}^{\#\mathcal{L}-1} (l_i - l_{i+1}) \cdot \min_{\substack{s \in \mathcal{S} \text{ s.t.} \\ b^\pi(s) \geq l_i}} r(s). \quad (2)$$

notation  $\mathcal{L} = \{l_1 = 1, l_2, l_3, \dots, 0\}$ .




# Choquet integral and rewards

back to the example about ignorance



# Choquet integral and rewards

back to the example about ignorance




- $Ch(r, N_{b_1^\pi} \mid a_0 = \tilde{a}) = r(s_D, \tilde{a}) = 5,$
- $Ch(r, N_{b_1^\pi} \mid a_0 = a') = \min_{s \in \mathcal{S}} r(s, a') = 0.$

the safe action is preferred! **dispersion reduced**

# Choquet integral and rewards

back to the example about ignorance



- $Ch(r, N_{b_1^\pi} \mid a_0 = \tilde{a}) = r(s_D, \tilde{a}) = 5,$
- $Ch(r, N_{b_1^\pi} \mid a_0 = a') = \min_{s \in \mathcal{S}} r(s, a') = 0.$

the safe action is preferred! **dispersion reduced**

if  $\mathcal{N}_{b_1^\pi}$  replaced by  $b_1 \Rightarrow Ch(r, b_1) = \mathbb{E}_{s \sim b_1} [r(s, a)]$ .

# resulting MDP

## translation summary

input: a POMDP  $\langle \mathcal{S}, \mathcal{A}, \mathcal{O}, T, O, r, \gamma \rangle$ ;

output: the MDP  $\langle \tilde{\mathcal{S}}, \mathcal{A}, \tilde{T}, \tilde{r}, \gamma \rangle$ :

# resulting MDP

## translation summary

input: a POMDP  $\langle \mathcal{S}, \mathcal{A}, \mathcal{O}, T, O, r, \gamma \rangle$ ;

output: the MDP  $\langle \tilde{\mathcal{S}}, \mathcal{A}, \tilde{T}, \tilde{r}, \gamma \rangle$ :

- **state space**  $\tilde{\mathcal{S}} = \Pi_{\mathcal{S}}$ ,  
the set of the possibility distributions over  $\mathcal{S}$ ;

# resulting MDP

## translation summary

input: a POMDP  $\langle \mathcal{S}, \mathcal{A}, \mathcal{O}, T, O, r, \gamma \rangle$ ;

output: the MDP  $\langle \tilde{\mathcal{S}}, \mathcal{A}, \tilde{T}, \tilde{r}, \gamma \rangle$ :

- **state space**  $\tilde{\mathcal{S}} = \Pi_{\mathcal{S}}$ ,  
the set of the possibility distributions over  $\mathcal{S}$ ;
- $\forall b^\pi, (b^\pi)'$  possibilistic belief states  $\in \Pi_{\mathcal{S}}, \forall a \in \mathcal{A}$ ,  
**transitions**  $\tilde{T}(b^\pi, a, (b^\pi)') = \mathbf{p}((b^\pi)' | b^\pi, a)$ ;

# resulting MDP

## translation summary

input: a POMDP  $\langle \mathcal{S}, \mathcal{A}, \mathcal{O}, T, O, r, \gamma \rangle$ ;

output: the MDP  $\langle \tilde{\mathcal{S}}, \mathcal{A}, \tilde{T}, \tilde{r}, \gamma \rangle$ :

- **state space**  $\tilde{\mathcal{S}} = \Pi_{\mathcal{S}}$ ,  
the set of the possibility distributions over  $\mathcal{S}$ ;
- $\forall b^\pi, (b^\pi)'$  possibilistic belief states  $\in \Pi_{\mathcal{S}}, \forall a \in \mathcal{A}$ ,  
**transitions**  $\tilde{T}(b^\pi, a, (b^\pi)') = \mathbf{p}((b^\pi)' | b^\pi, a)$ ;
- **reward**  $\tilde{r}(a, b^\pi) = Ch(r(a, \cdot), \mathcal{N}_{b^\pi})$ ,  
 $\mathcal{N}_{b^\pi}$  necessity measure computed from  $b^\pi$ .

# resulting MDP

## translation summary

input: a POMDP  $\langle \mathcal{S}, \mathcal{A}, \mathcal{O}, T, O, r, \gamma \rangle$ ;

output: the MDP  $\langle \tilde{\mathcal{S}}, \mathcal{A}, \tilde{T}, \tilde{r}, \gamma \rangle$ :

- **state space**  $\tilde{\mathcal{S}} = \Pi_{\mathcal{S}}$ ,  
the set of the possibility distributions over  $\mathcal{S}$ ;
- $\forall b^\pi, (b^\pi)'$  possibilistic belief states  $\in \Pi_{\mathcal{S}}, \forall a \in \mathcal{A}$ ,  
**transitions**  $\tilde{T}(b^\pi, a, (b^\pi)') = \mathbf{p}((b^\pi)' | b^\pi, a)$ ;
- **reward**  $\tilde{r}(a, b^\pi) = Ch(r(a, \cdot), \mathcal{N}_{b^\pi})$ ,  
 $\mathcal{N}_{b^\pi}$  necessity measure computed from  $b^\pi$ .

$$\text{criterion: } \mathbb{E}_{(b_t^\pi) \sim \tilde{T}} \left[ \sum_{t=0}^{+\infty} \gamma^t \cdot \tilde{r}(b_t^\pi, d_t) \right].$$



# hybrid POMDP and $\pi$ -POMDP

differences with possibilistic models

	hybrid POMDP	$\pi$ -POMDP
transitions	probabilities	qualitative possibility
rewards	quantitative $\in \mathbb{R}$	qualitative $\in \mathcal{L}$
situation	-some imprecisions -large POMDP	few quantitative
issues	$\pi$ definition	commensurability
in practice	MDP	$\pi$ -MDP

# hybrid POMDP and $\pi$ -POMDP

differences with possibilistic models

	hybrid POMDP	$\pi$ -POMDP
transitions	probabilities	qualitative possibility
rewards	quantitative $\in \mathbb{R}$	qualitative $\in \mathcal{L}$
situation	-some imprecisions -large POMDP	few quantitative
issues	$\pi$ definition	commensurability
in practice	MDP	$\pi$ -MDP

## hybrid model:

- only belief states are possibilistic:
  - agent knowledge = **possibility** distribution;
- probabilistic dynamics:
  - **approximated** (prob.) transition between epistemic states.

- 1 Context
- 2 An hybrid POMDP
- 3 Benefiting from factorized structures
- 4 Conclusion/Perspectives

# factorized POMDP

## definition

- $\mathcal{S}$  described by  $\mathbb{S} = \{s_1, \dots, s_m\}$ :  $\mathcal{S} = s_1 \times \dots \times s_m$ .  
Notation:  $\mathbb{S}' = \{s'_1, \dots, s'_m\}$ ;

# factorized POMDP

## definition

- $\mathcal{S}$  described by  $\mathbb{S} = \{s_1, \dots, s_m\}$ :  $\mathcal{S} = s_1 \times \dots \times s_m$ .  
Notation:  $\mathbb{S}' = \{s'_1, \dots, s'_m\}$ ;
- **transition** function of  $s'_j$ ,  
 $T_j^a(\mathbb{S}, s'_j) = \mathbf{p}(s'_j \mid \mathbb{S}, a), \forall j \in \{1, \dots, m\}$  et  $\forall a \in \mathcal{A}$ ;

- $\mathcal{S}$  described by  $\mathbb{S} = \{s_1, \dots, s_m\}$ :  $\mathcal{S} = s_1 \times \dots \times s_m$ .  
Notation:  $\mathbb{S}' = \{s'_1, \dots, s'_m\}$ ;
- **transition** function of  $s'_j$ ,  
 $T_j^a(\mathbb{S}, s'_j) = \mathbf{p}(s'_j \mid \mathbb{S}, a), \forall j \in \{1, \dots, m\}$  et  $\forall a \in \mathcal{A}$ ;
- $\mathcal{O}$  described by  $\mathbb{O} = \{o_1, \dots, o_n\}$ :  $\mathcal{O} = o_1 \times \dots \times o_n$ ;

- $\mathcal{S}$  described by  $\mathbb{S} = \{s_1, \dots, s_m\}$ :  $\mathcal{S} = s_1 \times \dots \times s_m$ .  
Notation:  $\mathbb{S}' = \{s'_1, \dots, s'_m\}$ ;
- **transition** function of  $s'_j$ ,  
 $T_j^a(\mathbb{S}, s'_j) = \mathbf{p}(s'_j \mid \mathbb{S}, a)$ ,  $\forall j \in \{1, \dots, m\}$  et  $\forall a \in \mathcal{A}$ ;
- $\mathcal{O}$  described by  $\mathbb{O} = \{o_1, \dots, o_n\}$ :  $\mathcal{O} = o_1 \times \dots \times o_n$ ;
- **observation** function of  $o'_i$ ,  
 $O_i^a(\mathbb{S}', o'_i) = \mathbf{p}(o'_i \mid \mathbb{S}', a)$ ,  $\forall i \in \{1, \dots, n\}$  et  $\forall a \in \mathcal{A}$ .

# factorized POMDP

## definition

- $\mathcal{S}$  described by  $\mathbb{S} = \{s_1, \dots, s_m\}$ :  $\mathcal{S} = s_1 \times \dots \times s_m$ .  
Notation:  $\mathbb{S}' = \{s'_1, \dots, s'_m\}$ ;
- **transition** function of  $s'_j$ ,  
 $T_j^a(\mathbb{S}, s'_j) = \mathbf{p}(s'_j \mid \mathbb{S}, a)$ ,  $\forall j \in \{1, \dots, m\}$  et  $\forall a \in \mathcal{A}$ ;
- $\mathcal{O}$  described by  $\mathbb{O} = \{o_1, \dots, o_n\}$ :  $\mathcal{O} = o_1 \times \dots \times o_n$ ;
- **observation** function of  $o'_i$ ,  
 $O_i^a(\mathbb{S}', o'_i) = \mathbf{p}(o'_i \mid \mathbb{S}', a)$ ,  $\forall i \in \{1, \dots, n\}$  et  $\forall a \in \mathcal{A}$ .

## independences:

- $\rightarrow \forall s'_i, s'_j \in \mathbb{S}', \quad s'_i \perp\!\!\!\perp s'_j \mid \{\mathbb{S}, a \in \mathcal{A}\},$
- $\rightarrow \forall o'_i, o'_j \in \mathbb{O}', \quad o'_i \perp\!\!\!\perp o'_j \mid \{\mathbb{S}', a \in \mathcal{A}\}.$



# Notations

some variables does not interact with each other

variables about the **current** system state,

$s_1$

$\vdots$

$s_{j_1}$

$\vdots$

$s_{j_2}$

$\vdots$

$s_{j_k}$

$\vdots$

$s_m$

variable  $s'_j$  about  
the **next** state.

$s'_j$

# Notations

some variables does not interact with each other


variables about the **current** system state,

$$s_k \rightarrow s'_j$$



$\exists a \in \mathcal{A}$ , such that

$T_j^a(\mathbb{S}, s'_j)$  depends on  $s_k$ .




# Notations

some variables does not interact with each other

variables about the **current** system state,

$s_k \rightarrow s'_j$   
 $\Updownarrow$   
 $\exists a \in \mathcal{A}$ , such that  
 $T_j^a(\mathbb{S}, s'_j)$  depends on  $s_k$ .




the parents of  $s'_j$  are:

$$\begin{aligned}\mathcal{P}(s'_j) &= \{s \in \mathbb{S} \mid s \rightarrow s'_j\} \\ &= \{s_{j_1}, s_{j_2}, \dots, s_{j_k}\}.\end{aligned}$$

# Notations

concerning observation variables

next state



# Notations

concerning observation variables

$$s'_j \rightarrow o'_i$$




$\exists a \in \mathcal{A}$ , such that

$$O_i^a(S', o'_i)$$

depends on  $s'_j$ .

next state



# Notations

concerning observation variables

$$s'_j \rightarrow o'_i$$




$\exists a \in \mathcal{A}$ , such that

$$O_i^a(\mathbb{S}', o'_i)$$

depends on  $s'_j$ .

next state



# Notations


concerning observation variables

$$s'_j \rightarrow o'_i$$




$\exists a \in \mathcal{A}$ , such that

$O_i^a(S', o'_i)$   
depends on  $s'_j$ .



**next state**



# Notations


concerning observation variables

$$s'_j \rightarrow o'_i$$



$\exists a \in \mathcal{A}$ , such that

$O_i^a(S', o'_i)$   
depends on  $s'_j$ .





# Notations

concerning observation variables


$$s'_j \rightarrow o'_i$$



$\exists a \in \mathcal{A}$ , such that


$$O_i^a(S', o'_i)$$

depends on  $s'_j$ .



# Notations

concerning observation variables



# Rewritings of parameters

## PROBABILISTIC parameters

- $T_j^a(\mathbb{S}, s'_j) = T_j^a(\mathcal{P}(s'_j), s'_j);$
- $O_i^a(\mathbb{S}', o'_i) = O_i^a(\mathcal{P}(o'_i), o'_i).$

# Rewritings of parameters

## PROBABILISTIC parameters

- $T_j^a(\mathbb{S}, s'_j) = T_j^a(\mathcal{P}(s'_j), s'_j);$
- $O_i^a(\mathbb{S}', o'_i) = O_i^a(\mathcal{P}(o'_i), o'_i).$

consequences on the joint distribution

$$\begin{aligned}\mathbf{p}(o'_i, \mathcal{P}(o'_i) \mid \mathbb{S}, a) &= O_i^a(\mathcal{P}(o'_i), o'_i) \cdot \prod_{s'_j \in \mathcal{P}(o'_i)} T_j^a(\mathcal{P}(s'_j), s'_j) \\ &= \mathbf{p}(o'_i, \mathcal{P}(o'_i) \mid \mathcal{Q}(o'_i), a).\end{aligned}$$

# Rewritings of parameters

## PROBABILISTIC parameters

- $T_j^a(\mathbb{S}, s'_j) = T_j^a(\mathcal{P}(s'_j), s'_j);$
- $O_i^a(\mathbb{S}', o'_i) = O_i^a(\mathcal{P}(o'_i), o'_i).$

consequences on the joint distribution

$$\begin{aligned}\mathbf{p}(o'_i, \mathcal{P}(o'_i) \mid \mathbb{S}, a) &= O_i^a(\mathcal{P}(o'_i), o'_i) \cdot \prod_{s'_j \in \mathcal{P}(o'_i)} T_j^a(\mathcal{P}(s'_j), s'_j) \\ &= \mathbf{p}(o'_i, \mathcal{P}(o'_i) \mid \mathcal{Q}(o'_i), a).\end{aligned}$$

observation probabilities

epistemic state

$$b^\pi(\mathbb{S}) \xrightarrow{\text{marginalization}} b^\pi(\mathcal{Q}(o'_i)) \xrightarrow{\text{pignistic transformation}} \overline{b}^\pi(\mathcal{Q}(o'_i))$$

$$\mathbf{p}(o'_i \mid b^\pi, a) = \sum_{2^{\mathcal{P}(o'_i)}, 2^{\mathcal{Q}(o'_i)}} \mathbf{p}(o'_i, \mathcal{P}(o'_i) \mid \mathcal{Q}(o'_i), a) \cdot \overline{b}^\pi(\mathcal{Q}(o'_i))$$

# Parameters rewritings

## POSSIBILISTIC parameters

- $\pi(s'_j \mid \mathbb{S}, a) = \pi(s'_j \mid \mathcal{P}(s'_j), a);$
- $\pi(o'_i \mid \mathbb{S}', a) = \pi(o'_i \mid \mathcal{P}(o'_i), a).$

- $\pi(s'_j \mid \mathbb{S}, a) = \pi(s'_j \mid \mathcal{P}(s'_j), a);$
- $\pi(o'_i \mid \mathbb{S}', a) = \pi(o'_i \mid \mathcal{P}(o'_i), a).$

### marginal possibilistic belief states

$$\forall o'_i \in \mathbb{O},$$
$$b_{t+1}^\pi(\mathcal{P}(o'_i)) \propto^\pi \pi(o'_i, \mathcal{P}(o'_i) \mid a_0, o_1, \dots, a_{t-1}, o_t)$$

- $\pi(s'_j \mid \mathbb{S}, a) = \pi(s'_j \mid \mathcal{P}(s'_j), a);$
- $\pi(o'_i \mid \mathbb{S}', a) = \pi(o'_i \mid \mathcal{P}(o'_i), a).$

### marginal possibilistic belief states

$$\begin{aligned} \forall o'_i \in \mathbb{O}, \\ b_{t+1}^\pi(\mathcal{P}(o'_i)) &\propto^\pi \pi(o'_i, \mathcal{P}(o'_i) \mid a_0, o_1, \dots, a_{t-1}, o_t) \\ &= \max_{2^{\mathcal{Q}(o'_i)}} \min \left\{ \pi(o'_i, \mathcal{P}(o'_i) \mid \mathcal{Q}(o'_i), a), b_t^\pi(\mathcal{Q}(o'_i)) \right\} \end{aligned}$$



- $\pi(s'_j \mid \mathbb{S}, a) = \pi(s'_j \mid \mathcal{P}(s'_j), a);$
- $\pi(o'_i \mid \mathbb{S}', a) = \pi(o'_i \mid \mathcal{P}(o'_i), a).$

### marginal possibilistic belief states

$\forall o'_i \in \mathbb{O},$

$$\begin{aligned} b_{t+1}^\pi(\mathcal{P}(o'_i)) &\propto^\pi \pi(o'_i, \mathcal{P}(o'_i) \mid a_0, o_1, \dots, a_{t-1}, o_t) \\ &= \max_{2^{\mathcal{Q}(o'_i)}} \min \left\{ \pi(o'_i, \mathcal{P}(o'_i) \mid \mathcal{Q}(o'_i), a), b_t^\pi(\mathcal{Q}(o'_i)) \right\} \\ &\quad \text{denoted by } \pi(o'_i, \mathcal{P}(o'_i) \mid b_t^\pi, a). \end{aligned}$$

# Variable classification

3 classes of state variables

variable: visible  $s_v \in \mathbb{S}_v$

$s'_v$

---

inferred hidden  $s_h \in \mathbb{S}_h$

$s'_h$

---


fully hidden  $s_f \in \mathbb{S}_f$

$s'_f$

# Variable classification

3 classes of state variables

variable: visible  $s_v \in \mathbb{S}_v$



---

inferred hidden  $s_h \in \mathbb{S}_h$



---


fully hidden  $s_f \in \mathbb{S}_f$



# Variable classification

3 classes of state variables

variable: **visible**  $s_v \in \mathbb{S}_v$



$$\mathbf{p}(s'_v | b_t^\pi, a) = \sum_{2^{\mathcal{P}(s'_v)}} T^a(\mathcal{P}(s'_v), s'_v) \cdot \overline{b}_t^\pi(\mathcal{P}(s'_v)).$$

---

**inferred hidden**  $s_h \in \mathbb{S}_h$



---

**fully hidden**  $s_f \in \mathbb{S}_f$




# Variable classification

3 classes of state variables

variable: **visible**  $s_v \in \mathbb{S}_v$

$\Leftrightarrow$  deterministic belief variable.



$$\mathbf{p}(s'_v | b_t^\pi, a) = \sum_{2^{\mathcal{P}(s'_v)}} T^a(\mathcal{P}(s'_v), s'_v) \cdot \overline{b}_t^\pi(\mathcal{P}(s'_v)).$$

---

**inferred hidden**  $s_h \in \mathbb{S}_h$



---

**fully hidden**  $s_f \in \mathbb{S}_f$




# Variable classification

3 classes of state variables

variable: **visible**  $s_v \in \mathbb{S}_v$


$\Leftrightarrow$  deterministic belief variable.

$$\mathbf{p}(s'_v | b_t^\pi, a) = \sum_{2^{\mathcal{P}(s'_v)}} T^a(\mathcal{P}(s'_v), s'_v) \cdot \overline{b}_t^\pi(\mathcal{P}(s'_v)).$$



---

**inferred hidden**  $s_h \in \mathbb{S}_h$



---

**fully hidden**  $s_f \in \mathbb{S}_f$




# Variable classification

3 classes of state variables

variable: **visible**  $s_v \in \mathbb{S}_v$


$\Leftrightarrow$  deterministic belief variable.

$$\mathbf{p}(s'_v | b_t^\pi, a) = \sum_{2^{\mathcal{P}(s'_v)}} T^a(\mathcal{P}(s'_v), s'_v) \cdot \overline{b}_t^\pi(\mathcal{P}(s'_v)).$$



inferred hidden  $s_h \in \mathbb{S}_h$

$$b_{t+1}^\pi(\mathcal{P}(o'_i)) = b_{t+1}^\pi(s_h, s_h^a, s_h^b, s_h^c)$$



fully hidden  $s_f \in \mathbb{S}_f$




# Variable classification

3 classes of state variables

variable: **visible**  $s_v \in \mathbb{S}_v$

$\Leftrightarrow$  deterministic belief variable.


$$\mathbf{p}(s'_v \mid b_t^\pi, a) = \sum_{2^{\mathcal{P}(s'_v)}} T^a(\mathcal{P}(s'_v), s'_v) \cdot \overline{b}_t^\pi(\mathcal{P}(s'_v)).$$



**inferred hidden**  $s_h \in \mathbb{S}_h$

$$b_{t+1}^\pi(\mathcal{P}(o'_i)) = b_{t+1}^\pi(s_h, s_h^a, s_h^b, s_h^c)$$

$$\propto^\pi \pi(o'_i, \mathcal{P}(o'_i) \mid b_t^\pi, a).$$



**fully hidden**  $s_f \in \mathbb{S}_f$






# Variable classification

3 classes of state variables

variable: **visible**  $s_v \in \mathbb{S}_v$

$\Leftrightarrow$  deterministic belief variable.


$$\mathbf{p}(s'_v | b_t^\pi, a) = \sum_{2^{\mathcal{P}(s'_v)}} T^a(\mathcal{P}(s'_v), s'_v) \cdot \overline{b}_t^\pi(\mathcal{P}(s'_v)).$$



**inferred hidden**  $s_h \in \mathbb{S}_h$

$$b_{t+1}^\pi(\mathcal{P}(o'_i)) = b_{t+1}^\pi(s_h, s_h^a, s_h^b, s_h^c)$$

$$\propto^\pi \pi(o'_i, \mathcal{P}(o'_i) | b_t^\pi, a).$$



$\triangle$   $\mathcal{P}(o'_i)$  may contain visible variables.

**fully hidden**  $s_f \in \mathbb{S}_f$




# Variable classification

3 classes of state variables

variable: **visible**  $s_v \in \mathbb{S}_v$

$\Leftrightarrow$  deterministic belief variable.


$$\mathbf{p}(s'_v | b_t^\pi, a) = \sum_{2^{\mathcal{P}(s'_v)}} T^a(\mathcal{P}(s'_v), s'_v) \cdot \overline{b}_t^\pi(\mathcal{P}(s'_v)).$$



**inferred hidden**  $s_h \in \mathbb{S}_h$

$$b_{t+1}^\pi(\mathcal{P}(o'_i)) = b_{t+1}^\pi(s_h, s_h^a, s_h^b, s_h^c)$$

$$\propto^\pi \pi(o'_i, \mathcal{P}(o'_i) | b_t^\pi, a).$$



$\triangle!$   $\mathcal{P}(o'_i)$  may contain visible variables.

**fully hidden**  $s_f \in \mathbb{S}_f$




# Variable classification

3 classes of state variables

variable: **visible**  $s_v \in \mathbb{S}_v$

$\Leftrightarrow$  deterministic belief variable.


$$\mathbf{p}(s'_v | b_t^\pi, a) = \sum_{2^{\mathcal{P}(s'_v)}} T^a(\mathcal{P}(s'_v), s'_v) \cdot \overline{b}_t^\pi(\mathcal{P}(s'_v)).$$



**inferred hidden**  $s_h \in \mathbb{S}_h$

$$b_{t+1}^\pi(\mathcal{P}(o'_i)) = b_{t+1}^\pi(s_h, s_h^a, s_h^b, s_h^c)$$

$$\propto^\pi \pi(o'_i, \mathcal{P}(o'_i) | b_t^\pi, a).$$



$\triangle!$   $\mathcal{P}(o'_i)$  may contain visible variables.

**fully hidden**  $s_f \in \mathbb{S}_f$



$$b_{t+1}^\pi(s'_f) = \max_{2^{\mathcal{P}(s'_f)}} \min \left\{ \pi(s'_f | \mathcal{P}(s'_f), a), b_t^\pi(\mathcal{P}(s'_f)) \right\}.$$


# Variable classification

3 classes of state variables

variable: **visible**  $s_v \in \mathbb{S}_v$

$\Leftrightarrow$  deterministic belief variable.

$$\mathbf{p}(s'_v \mid b_t^\pi, a) = \sum_{2^{\mathcal{P}(s'_v)}} T^a(\mathcal{P}(s'_v), s'_v) \cdot \overline{b}_t^\pi(\mathcal{P}(s'_v)).$$




**inferred hidden**  $s_h \in \mathbb{S}_h$

$$b_{t+1}^\pi(\mathcal{P}(o'_i)) = b_{t+1}^\pi(s_h, s_h^a, s_h^b, s_h^c)$$

$$\propto^\pi \pi(o'_i, \mathcal{P}(o'_i) \mid b_t^\pi, a).$$

$\triangle!$   $\mathcal{P}(o'_i)$  may contain visible variables.



**fully hidden**  $s_f \in \mathbb{S}_f$

$\rightarrow$  observations don't  
inform belief state on  $s'_f$ .

$$b_{t+1}^\pi(s'_f) = \max_{2^{\mathcal{P}(s'_f)}} \min \left\{ \pi(s'_f \mid \mathcal{P}(s'_f), a), b_t^\pi(\mathcal{P}(s'_f)) \right\}.$$



# Possibilistic belief variables

global belief state

$$\mathbb{O}_h = \mathbb{O} \setminus \mathbb{S}_v.$$

bound over the global belief state

$$b_{t+1}^\pi(\mathbb{S}') = \pi(\mathbb{S}' \mid a_0, o_1, \dots, a_t, o_{t+1})$$

$$\leq \beta_{t+1}(\mathbb{S}')$$

$$= \min \left\{ \min_{s'_j \in \mathbb{S}_v} \left[ \mathbb{1}_{\{s'_j = o'_j\}} \right], \min_{s'_j \in \mathbb{S}_f} \left[ b_{t+1}^\pi(s'_j) \right], \min_{o'_i \in \mathbb{O}_h} \left[ b_{t+1}^\pi(\mathcal{P}(o'_i)) \right] \right\}$$

# Possibilistic belief variables

global belief state

$$\mathbb{O}_h = \mathbb{O} \setminus \mathbb{S}_v.$$

bound over the global belief state

$$b_{t+1}^\pi(\mathbb{S}') = \pi(\mathbb{S}' \mid a_0, o_1, \dots, a_t, o_{t+1})$$

$$\leq \beta_{t+1}(\mathbb{S}')$$

$$= \min \left\{ \min_{s'_j \in \mathbb{S}_v} \left[ \mathbb{1}_{\{s'_j = o'_j\}} \right], \min_{s'_j \in \mathbb{S}_f} \left[ b_{t+1}^\pi(s'_j) \right], \min_{o'_i \in \mathbb{O}_h} \left[ b_{t+1}^\pi(\mathcal{P}(o'_i)) \right] \right\}$$

- $\beta_t$  = **less informative** version of the belief state:  
 $b_t^\pi \leq \beta_t$ ;
- computed using **marginal belief states**  $\leftrightarrow$  **factorization**.

# Variables de croyance

different according to the class of the variable

$$\lambda = \#\mathcal{L}$$

# Variables de croyance

different according to the class of the variable

$$\lambda = \#\mathcal{L}$$

- $\forall s'_v \in \mathbb{S}_v$ , 1 variable  $\beta'_v$  is enough.



# Variables de croyance

different according to the class of the variable

$$\lambda = \#\mathcal{L}$$

■  $\forall s'_v \in \mathbb{S}_v$ , 1 variable  $\beta'_v$  is enough.

■  $p_i = \#\mathcal{P}(o'_i)$ .

$\forall o_i \in \mathbb{O} \setminus \mathbb{S}_v$ ,  $\lambda^{2^{p_i}} - (\lambda - 1)^{2^{p_i}}$  belief states,

$\Rightarrow \lceil \log_2(\lambda^{2^{p_i}} - (\lambda - 1)^{2^{p_i}}) \rceil$  boolean variables  $\beta'_h$ .

# Variables de croyance

different according to the class of the variable

$$\lambda = \#\mathcal{L}$$

■  $\forall s'_v \in \mathbb{S}_v$ , 1 variable  $\beta'_v$  is enough.

■  $p_i = \#\mathcal{P}(o'_i)$ .

$\forall o_i \in \mathbb{O} \setminus \mathbb{S}_v$ ,  $\lambda^{2^{p_i}} - (\lambda - 1)^{2^{p_i}}$  belief states,

$\Rightarrow \lceil \log_2(\lambda^{2^{p_i}} - (\lambda - 1)^{2^{p_i}}) \rceil$  boolean variables  $\beta'_h$ .

■  $\forall s'_f \in \mathbb{S}_f$ ,  $\lambda^2 - (\lambda - 1)^2 = 2\lambda - 1$  belief states,

$\Rightarrow \lceil \log_2(2\lambda - 1) \rceil$  boolean variables  $\beta'_f$ .

# resulting MDP in practice

trick: “flipflop” variable

boolean variable “*flipflop*”  $f$  changes state at each time step  
→ defines 2 phases:

- 1 *observation generation*,
- 2 *belief update* (deterministic knowing the observation).

MDP variables:

$\tilde{S} =$


**beliefs:**  $\beta = \beta_v^1 \times \dots \times \beta_v^{m_v} \times \beta_h^1 \times \dots \times \beta_h^{m_h} \times \beta_f^1 \times \dots \times \beta_f^{m_f}$

$\times$

**visible variables** :  $v = f \times s_v^1 \times \dots \times s_v^{m_v} \times o_1 \times \dots \times o_k.$

# resulting MDP in practice

## final structured MDP



$\tilde{\mathcal{S}} =$


**beliefs:**  $\beta = \beta_v^1 \times \dots \times \beta_v^{m_v} \times \beta_h^1 \times \dots \times \beta_h^{m_h} \times \beta_f^1 \times \dots \times \beta_f^{m_f}$

$\times$

**visible variables** :  $v = f \times s_v^1 \times \dots \times s_v^{m_v} \times o_1 \times \dots \times o_k.$

# resulting MDP in practice

## final structured MDP



# factorized model's variables:  $\#\mathbb{O} + \#\mathbb{S}_v +$


$$+ \sum_{i=1}^{\#\mathbb{O}_h} \left[ \log_2 (\lambda^{2^{p_i}} - (\lambda - 1)^{2^{p_i}}) \right] + \#\mathbb{S}_f \cdot \left[ \log_2 (2\lambda - 1) \right]$$

$\ll$  # initial hybrid model's variables:

$$\left[ \log_2 (\lambda^{2^{\#\mathbb{S}}} - (\lambda - 1)^{2^{\#\mathbb{S}}}) \right]$$

# resulting MDP in practice

## final structured MDP



# factorized model's variables:

$$\leq \#\mathbb{O} + \#S_v + \sum_{i=1}^{\#\mathbb{O}_h} \log_2(\lambda) \cdot 2^{p_i} + \#S_f \cdot (1 + \log_2(\lambda))$$

$\ll$  # initial hybrid model's variables:  
 $\geq \log_2(\lambda) \cdot (2^{\#\mathbb{S}} - 1).$

- 1 Context
- 2 An hybrid POMDP
- 3 Benefiting from factorized structures
- 4 Conclusion/Perspectives

POMDP  $\xrightarrow{\text{translation}}$  MDP with finite state space

- transition probabilities on the **possibilistic belief states**;



POMDP  $\xrightarrow{\text{translation}}$  MDP with finite state space

- transition probabilities on the **possibilistic belief states**;
- **pessimistic** evaluation of the **rewards**  
(Choquet integral);

POMDP  $\xrightarrow{\text{translation}}$  MDP with finite state space

- transition probabilities on the **possibilistic belief states**;
- **pessimistic** evaluation of the **rewards**  
(Choquet integral);
- factorized POMDP  $\xrightarrow{\text{translation}}$  **factorized MPD**.

POMDP  $\xrightarrow{\text{translation}}$  MDP with finite state space

- transition probabilities on the **possibilistic belief states**;
- **pessimistic** evaluation of the **rewards** (Choquet integral);
- factorized POMDP  $\xrightarrow{\text{translation}}$  **factorized MPD**.

**perspectives:**

- IPPC problems (factorized POMDPs);

POMDP  $\xrightarrow{\text{translation}}$  MDP with finite state space

- transition probabilities on the **possibilistic belief states**;
- **pessimistic** evaluation of the **rewards** (Choquet integral);
- factorized POMDP  $\xrightarrow{\text{translation}}$  **factorized MPD**.

## perspectives:

- IPPC problems (factorized POMDPs);
- tests of this approach:
  - 1 **simplification**:  $\pi$  distributions definition ( $\pi$ -normalization, pignistic transformation, most specific, ... );
  - 2 **imprecision**: robust in practice?

Thank you!