Module 4
Lecture Notes

Competencies 97 to 132. One section per competency, each headed with its number and what you should be able to do, with worked exam questions at the end of every topic.

Solano Community College
Dr. Sharilyn Rennie

Contents

Each topic and every competency inside it. Use the competency number to find the section, and the competency packet to see the exact wording you are held to.

The respiratory system	**97 to 104**	**4**
97. Upper and lower tracts, conducting and respiratory zones		
98. Nose, nasal cavity, and pharynx		
99. Larynx		
100. Trachea and bronchial tree		
101. Lungs and pleurae		
102. Respiratory zone and alveolar structure		
103. Thoracic wall and diaphragm		
104. Respiratory disorders by structure affected		
The lymphatic system	**105 to 108**	**18**
105. Lymph pathway		
106. Vessel and node structure		
107. Primary and secondary organs		
108. Lymphatic disorders		
The digestive system, part I: through the esophagus	**109 to 121**	**29**
109. Four-layer gut wall		
110. Enteric nerve plexuses		
111. Peritoneum and folds		
112. Retroperitoneal organs		
113. Mouth, pharynx, esophagus		
114. Major canal sphincters		
120. Teeth and tongue		
121. Salivary glands		
The digestive system, part II: stomach, pancreas, liver, and gallbladder	**115 to 125**	**40**
115. Stomach regions and features		
122. Liver gross and lobule anatomy		
123. Gallbladder and bile pathway		
124. Pancreas structure		
125. Clinical anatomy of accessory disorders		
The digestive system, part III: small intestine, large intestine, and rectum	**116 to 119**	**49**
116. Small intestine and surface area		
117. Large intestine features		
118. Trace the pathway of food		
119. Clinical anatomy of canal disorders		
Blood flow of the abdomen and lower limb	**126 to 127**	**57**
126. Abdominal aorta and portal system		
127. Lower limb vessels and nerves		

Endocrine anatomy	**128 to 132**	**65**
128. Major endocrine glands and locations		
129. Pituitary gross anatomy and lobes		
130. Thyroid and parathyroid anatomy		
131. Adrenal gland zones		
132. Pancreas, gonads, and organs with endocrine tissue		

BIO 004 Human Anatomy · Module 4 notes · Dr. Sharilyn Rennie

The respiratory system

The written notes for the respiratory lectures. Module 4 competencies 97 to 104, with eight worked questions at the end.

Before the competencies: what the system does

In your course notes: Module 4 packet, Chapter 1 introduction (page 4).

Dr. Rennie's video for this part: Respiratory system concept video, chapter Respiratory system overview, and Ventilation and gas exchange.

Respiration is four linked jobs. Two of them belong to this system, one belongs to the cardiovascular system, and the last one happens out in the tissues. The respiratory system and the cardiovascular system are so tightly linked that if either fails, oxygen never reaches the cells.

1. Pulmonary ventilation (breathing): air moves into the lungs (inspiration) and out of the lungs (expiration), like a bellows.
2. External respiration: oxygen crosses from the alveoli into the blood and carbon dioxide crosses from the blood into the alveoli, at the respiratory membrane.
3. Transport of gases: the cardiovascular system carries oxygen from the lungs to the tissues and carbon dioxide back. The heart and vessels are the delivery truck; the blood is the cargo.
4. Internal respiration: oxygen moves from the systemic capillaries into the tissue cells and carbon dioxide moves out of them.

- Cellular respiration is something else: the mitochondria using oxygen and making carbon dioxide to produce ATP. It is a job of every cell, not of the respiratory system.
- Moving air also gives you smell and speech.
- The structures in order, nose to alveolus: nose, nasal cavity, paranasal sinuses, pharynx, larynx, trachea, bronchi and their branches, lungs, alveoli. (Nice Nurses Practice Thorough Bronchial Lung Assessments.)
- The respiratory muscles, the diaphragm and its helpers, create the pressure changes that move the air. They are competency 103.

Competency 97. Upper and lower tracts, conducting and respiratory zones

In your course notes: Module 4 packet, Chapter 1, competency 97 (page 4).

Dr. Rennie's video for this part: Respiratory system concept video, chapter Respiratory system overview.

You should be able to: Distinguish the upper and lower respiratory tracts and the conducting and respiratory zones, and assign the major airway structures to each.

The respiratory system moves air into and out of the body and brings that air close to the blood. Air follows one continuous path from the nose to the air sacs of the lungs. Anatomists group that path two ways, and the two groupings use different dividing lines, so you have to learn both.

One breath, step by step

1. Air enters at the nares (nostrils) and passes the nasal vestibule, where coarse hairs catch large particles.
2. It moves through the nasal cavity, tumbling past the conchae, and is warmed and moistened.
3. It leaves through the choanae into the nasopharynx (air only), then the oropharynx and laryngopharynx (shared with food).
4. It turns forward into the larynx and passes the glottis, between the vocal folds.
5. It runs down the trachea to the carina, where the trachea forks.

6. It enters a main bronchus (one per lung), then a lobar bronchus (one per lobe), then a segmental bronchus (one per bronchopulmonary segment).
7. It passes through bronchioles, which have no cartilage, to a terminal bronchiole, the last part of the conducting zone.
8. It crosses into a respiratory bronchiole, the first airway of the respiratory zone, then an alveolar duct, an alveolar sac, and an alveolus, where gases cross into the blood.

Upper respiratory tract
The nose, nasal cavity, paranasal sinuses, and pharynx. These sit in the head and neck.

Lower respiratory tract
The larynx, trachea, bronchi and all their branches, and the lungs. The larynx is the first structure of the lower tract, so the line between the tracts falls where the pharynx meets the larynx. Some clinical sources put the larynx in the upper tract; these notes follow the anatomy textbook split.

Conducting zone
Every passage that only carries air, from the nose down to the terminal bronchioles. Its walls are too thick for gases to cross. It warms, moistens, and filters the air on the way through.

Respiratory zone
Where gas exchange happens: the respiratory bronchioles, alveolar ducts, alveolar sacs, and alveoli. The line between the zones is deep inside the lung, where a terminal bronchiole gives way to a respiratory bronchiole.

Structure	Tract	Zone
Nose and nasal cavity	Upper	Conducting
Paranasal sinuses	Upper	Air spaces off the nasal cavity, no gas exchange
Pharynx	Upper	Conducting
Larynx	Lower	Conducting
Trachea	Lower	Conducting
Main, lobar, and segmental bronchi	Lower	Conducting
Bronchioles and terminal bronchioles	Lower	Conducting
Respiratory bronchioles	Lower	Respiratory
Alveolar ducts and sacs	Lower	Respiratory
Alveoli	Lower	Respiratory

- Tract tells you where a structure is. Zone tells you what it does.
- Because the lines fall in different places, one structure can be lower tract and conducting zone at the same time. The trachea is the usual example.
- Every upper tract structure is conducting. The lower tract holds the rest of the conducting zone plus the whole respiratory zone.
- The conducting zone ends at the terminal bronchioles, not at the bronchi and not where the lung begins. Gas exchange itself is physiology; this course covers the structures.

Competency 98. Nose, nasal cavity, and pharynx

In your course notes: Module 4 packet, Chapter 1, competency 98 (page 6).

Dr. Rennie's video for this part: Respiratory system concept video, chapter Respiratory system overview.

You should be able to: Identify the parts of the nasal cavity, paranasal sinuses, and the three regions of the pharynx.

The nose is the main entrance for air. It warms, moistens, and filters the air, it houses the receptors for smell, and it acts as a resonating chamber for the voice. Bone forms the upper part of the external nose (nasal bones, frontal bone, maxillae) and hyaline cartilage forms the lower part, which is why the tip bends.

The external nose

Feature	What it is
Root	Where the nose meets the forehead, between the eyebrows (the glabella is the flat area between the brows)
Bridge	The upper, bony part
Dorsum nasi	The anterior margin, from the root to the apex
Apex	The tip
Alae	The flared sides of the nostrils
Nares	The nostrils, the external openings

- Framework: nasal bones and the frontal bone above, the maxillae at the sides, and plates of hyaline cartilage below (septal and alar cartilages and the lateral processes of the septal cartilage).
- The nose does five things: it is an airway, it warms and moistens the air, it filters and cleans it, it resonates for speech, and it houses the smell receptors.

Parts of the nasal cavity

Nasal septum
The midline wall that splits the cavity into right and left halves. Septal cartilage in front; the perpendicular plate of the ethmoid above the vomer behind.

Nasal vestibule
The area just inside each nostril, lined with skin, sebaceous glands, and coarse hairs that trap large particles.

Nasal conchae
Three scroll-shaped shelves on each lateral wall: superior, middle, and inferior. The superior and middle conchae are parts of the ethmoid bone. The inferior concha is a separate bone. They make the air tumble so more of it touches the warm, moist lining.

Nasal meatuses
The passages under each concha: superior, middle, and inferior meatus. A meatus is the space, not the shelf.

Hard palate
The bony front part of the floor, separating the nasal cavity from the mouth.

Soft palate
The muscular back part of the floor. It lifts to close off the nasopharynx during swallowing.

Choanae
The two back openings where the nasal cavity becomes the nasopharynx.

Two linings

Mucosa	Epithelium	Job
Olfactory mucosa	Olfactory epithelium, a small patch in the roof	Houses the smell receptors
Respiratory mucosa	Pseudostratified ciliated columnar with goblet cells	Secretes mucus that traps dust and microbes; the cilia sweep it back to the pharynx to be swallowed

- The vestibule is lined with stratified squamous epithelium, because it takes the mechanical wear at the entrance.
- The seromucous glands secrete lysozyme, an antibacterial enzyme, and the epithelial cells secrete defensins. The high water content of the mucus humidifies the air.

- A dense capillary bed close to the surface warms incoming air. It is also why nosebleeds (epistaxis) are common.
- Nerve endings in the lining trigger the sneeze reflex when dust or pollen irritates them.
- Cold air slows the cilia, so mucus stops moving back toward the throat and drips out the front. That is why your nose runs in cold weather. On the way out, water vapor condenses on the cooler lining, so the conchae reclaim heat and moisture during exhalation.

The paranasal sinuses

Air-filled spaces inside four skull bones around the nasal cavity, each named for its bone and lined with the same mucosa as the nasal cavity. Each drains through a small opening into the nasal cavity. They lighten the skull, help warm and moisten air, and add resonance to the voice.

Sinus	Bone	Where it sits	Drains into
Frontal sinus	Frontal	In the forehead, above the orbits	Middle meatus
Ethmoidal air cells	Ethmoid	Many small spaces between the nasal cavity and the orbits	Middle meatus (front and middle cells), superior meatus (back cells)
Sphenoidal sinus	Sphenoid	Deep in the skull, behind the nasal cavity	The sphenoethmoidal recess, above the superior concha
Maxillary sinus	Maxilla	In the cheek, below each orbit; the largest	Middle meatus

The pharynx, top to bottom

The pharynx (throat) is a funnel-shaped muscular tube from the base of the skull down to the level of the cricoid cartilage, where it continues as the esophagus. It connects the nasal cavity and mouth to the larynx and esophagus.

Region	Location	Carries	Epithelium and structures
Nasopharynx	Behind the nasal cavity, above the soft palate	Air only	Pseudostratified ciliated columnar. Pharyngeal tonsil (adenoid) on the back wall, auditory tube openings on the side walls
Oropharynx	Behind the oral cavity, soft palate to epiglottis	Air and food	Stratified squamous, which resists friction. Palatine tonsils in the side walls, lingual tonsil on the back of the tongue
Laryngopharynx	Behind the larynx, epiglottis to esophagus	Air and food	Stratified squamous. Where air turns forward into the larynx and food goes back into the esophagus

- When you swallow, the soft palate and uvula lift and seal the nasopharynx, so food does not enter the nose.
- The epithelium changes with the job. Where only air passes, the lining is ciliated and moves mucus. Where food also passes, it is stratified squamous and stands up to rubbing.

Competency 99. Larynx

In your course notes: Module 4 packet, Chapter 1, competency 99 (page 8).

Dr. Rennie's video for this part: Respiratory system concept video, chapter Larynx and voice passage.

You should be able to: Identify the cartilages and folds of the larynx, including the thyroid and cricoid cartilages, epiglottis, arytenoids, vocal cords, and glottis.

The larynx (voice box) connects the laryngopharynx to the trachea. It sits in the front of the neck at about C3 to C6 and hangs from the hyoid bone, which is above it and not part of it. It has three jobs: hold the airway open, keep food and drink out of the lower airway, and produce the voice.

The nine cartilages

Cartilage	Number	Cartilage type	What it is
Thyroid cartilage	Single	Hyaline	The large shield. Its front ridge is the laryngeal prominence (Adam's apple). Open at the back
Cricoid cartilage	Single	Hyaline	The signet ring just below the thyroid cartilage, narrow in front and wide behind. The only complete ring of cartilage in the airway. The trachea hangs from its lower edge
Epiglottis	Single	Elastic	The spoon-shaped flap attached inside the thyroid cartilage. It tips down over the laryngeal inlet during a swallow
Arytenoid cartilages	Paired	Mostly hyaline	Sit on top of the back of the cricoid. They anchor the vocal folds and pivot to move them, so they are the most important paired cartilages
Corniculate cartilages	Paired	Elastic	Small cartilages on top of the arytenoids
Cuneiform cartilages	Paired	Elastic	Small cartilages in the fold running from each arytenoid up to the epiglottis

- The cricothyroid ligament fills the gap between the thyroid and cricoid cartilages in front. It lies just under the skin, below the vocal folds, with no cartilage over it, which is why it is the landmark for an emergency airway when the larynx above is blocked.
- The epiglottis is elastic because it bends over the inlet with every swallow and springs back. The thyroid, cricoid, and arytenoid cartilages are hyaline (only the tips of the arytenoids are elastic).

The folds and the glottis

Vocal folds

The true vocal cords, the lower pair. Each contains a vocal ligament, an elastic band running from an arytenoid cartilage behind to the thyroid cartilage in front. Very few blood vessels, so they look pearly white. They vibrate as air passes between them, and that vibration is sound.

Vestibular folds

The false vocal cords, the upper pair. They help close the airway during swallowing and play no part in sound.

Glottis

The vocal folds together with the opening between them (the rima glottidis), the narrowest point air passes in the larynx. Books draw the edges a little differently: some use the name for the opening alone, and OpenStax includes the vestibular folds.

Laryngeal inlet

The opening into the larynx from the laryngopharynx, which the epiglottis covers during a swallow.

- When the arytenoids pivot, the vocal folds swing apart to widen the glottis for breathing or come together to close it.
- The lining changes at the vocal folds. Above them, where food can rub, stratified squamous. Below them, pseudostratified ciliated columnar, with cilia sweeping mucus up toward the pharynx.
- The vocal folds make the sound, but the pharynx, oral cavity, and sinuses amplify it like the body of an instrument, and the lips, tongue, and soft palate shape it into words.
- The larynx also works as a sphincter. When you strain, lift something heavy, or empty your bowels, the vocal folds close and hold air in the chest (the Valsalva maneuver), which stiffens the trunk and raises abdominal pressure.
- The laryngeal prominence grows larger in males at puberty under testosterone, which is why the voice drops.

Competency 100. Trachea and bronchial tree

In your course notes: Module 4 packet, Chapter 1, competency 100 (page 10).

Dr. Rennie's video for this part: Respiratory system concept video, chapter Trachea and breathing tubes.

You should be able to: Trace air through the trachea and bronchial tree in order from the trachea and carina down to the terminal bronchioles.

Below the larynx the airway becomes a branching tree. The tubes get narrower and more numerous at every step, and every one of them down to the terminal bronchioles belongs to the conducting zone.

From the trachea to a terminal bronchiole, in order

1. Trachea: the windpipe, from the lower edge of the cricoid cartilage into the mediastinum, about 10 to 12 cm long. Sixteen to twenty C-shaped rings of hyaline cartilage hold it open. The open side of each C faces back toward the esophagus, and the trachealis muscle bridges the gap, so the esophagus can bulge forward when you swallow.
2. Carina: the internal ridge formed by the lowest tracheal cartilage at the fork, about at the sternal angle (T4 to T5). Its lining is very sensitive; touching it sets off a strong cough.
3. Main (primary) bronchi: one enters each lung at the hilum. The right main bronchus is wider, shorter, and more vertical than the left, so an inhaled object usually lands in the right lung.
4. Lobar (secondary) bronchi: one to each lobe, three on the right and two on the left.
5. Segmental (tertiary) bronchi: one to each bronchopulmonary segment.
6. Bronchioles: tubes less than 1 mm across, with smooth muscle and no cartilage.
7. Terminal bronchioles: the last and smallest conducting tubes. The next tubes, the respiratory bronchioles, begin the respiratory zone.

The four layers of the tracheal wall

Layer	What is in it
Mucosa	Pseudostratified ciliated columnar epithelium with goblet cells on a lamina propria rich in elastic fibers. The cilia move mucus up toward the pharynx
Submucosa	Connective tissue with seromucous glands that produce sheets of mucus
Cartilaginous layer	The 16 to 20 C-shaped rings of hyaline cartilage, open at the back, bridged by the trachealis muscle. The last cartilage forms the carina
Adventitia	The outer connective tissue that ties the trachea to its neighbors

- The elastic fibers let the trachea stretch and recoil with each breath.
- When the trachealis contracts it narrows the trachea, which speeds the air during a cough and helps blast secretions out.

How the wall changes down the tree

Level	Cartilage	Epithelium	Smooth muscle
Trachea	C-shaped hyaline rings	Pseudostratified ciliated columnar with goblet cells	Trachealis across the back of each ring
Main bronchi	C-shaped rings	Pseudostratified ciliated columnar with goblet cells	A thin layer
Lobar and segmental bronchi	Irregular plates	Pseudostratified ciliated columnar, fewer goblet cells in smaller branches	A more complete layer
Bronchioles	None	Simple columnar to simple cuboidal	A complete layer, much of the wall
Terminal bronchioles	None	Simple cuboidal	A complete layer

- Going down the tree, cartilage shrinks and smooth muscle takes over. Rings, then plates, then none.
- Cartilage is already gone at the bronchioles, not just at the terminal bronchioles.

- Without cartilage to hold them open, bronchioles narrow a great deal when their smooth muscle contracts. That is the structural reason asthma is so dangerous.

Competency 101. Lungs and pleurae

In your course notes: Module 4 packet, Chapter 1, competency 101 (page 11).

Dr. Rennie's video for this part: Respiratory system concept video, chapter Lungs position and diaphragm, and Lung sections and blood flow.

You should be able to: Identify the lobes, fissures, and surfaces of each lung and the parietal and visceral pleurae with the pleural cavity.

The two lungs fill most of the thoracic cavity, one on each side of the mediastinum (the central compartment holding the heart, great vessels, trachea, and esophagus), with the diaphragm beneath them. Each lung is cone-shaped, soft, and spongy. The two are not the same size: the heart tilts left and takes room, so the left lung is smaller; the liver pushes the right dome of the diaphragm up, so the right lung is shorter and wider.

Surfaces and landmarks

Apex
The narrow top of the lung, rising just above the clavicle into the root of the neck.

Base
The broad, curved bottom surface resting on the diaphragm.

Costal surface
The large outer surface against the ribs.

Mediastinal surface
The medial surface facing the mediastinum. The hilum is on it.

Hilum
The indented area on the mediastinal surface where the bronchus, blood vessels, nerves, and lymphatic vessels enter and leave.

Root of the lung
The bundle passing through the hilum: main bronchus, pulmonary artery, pulmonary veins, bronchial vessels, nerves, and lymphatics.

Cardiac notch
The indentation in the front edge of the left lung that makes room for the heart.

Lingula
The tongue-shaped part of the left superior lobe just below the cardiac notch. It sits where the middle lobe sits on the right.

- In the root, the pulmonary artery carries oxygen-poor blood into the lung and the pulmonary veins carry oxygen-rich blood back to the heart.
- Telling right from left at the hilum on the cadaver: in the left hilum the pulmonary artery is the highest structure; in the right hilum a bronchus sits at or above the artery. In both, the pulmonary veins are lowest.

Lobes and fissures

Lung	Lobes	Fissures	Notes
Right lung	Three: superior, middle, inferior	Oblique fissure (separates the inferior lobe) and horizontal fissure (separates superior from middle)	Larger; shorter and wider because the liver raises the right diaphragm. Ten bronchopulmonary segments
Left lung	Two: superior, inferior	Oblique fissure only	Smaller, with the cardiac notch and the lingula. About 8 to 10 segments

- Each lobe is divided into bronchopulmonary segments, each served by its own segmental bronchus, artery, and vein. Because each segment is its own unit, a surgeon can remove a diseased segment without damaging the segments around it, like taking one section out of an orange.
- Right has three, left has two. The left gave up its middle lobe to make room for the heart, and the lingula is what is left in its place.

Two blood supplies to the lung

Circulation	Pressure and volume	Job
Pulmonary circulation (pulmonary arteries and veins)	Low pressure, high volume	Gas exchange at the alveoli
Bronchial circulation (bronchial arteries off the aorta)	High pressure, low volume	Oxygen-rich systemic blood to feed the lung tissue itself

The lungs are supplied by parasympathetic and sympathetic motor fibers and by visceral sensory fibers. The sympathetic fibers widen the airways; the parasympathetic fibers narrow them.

The pleurae

Parietal pleura
: The serous membrane lining the inside of the thoracic wall, the top of the diaphragm, and the sides of the mediastinum.

Visceral pleura
: The serous membrane clinging to the lung surface and dipping into the fissures. The two layers are one continuous sheet that turns back on itself at the hilum.

Pleural cavity
: The very thin space between the two layers, holding only a film of pleural fluid. There is a right and a left pleural cavity, kept apart by the mediastinum.

Costodiaphragmatic recess
: The low pocket where the parietal pleura turns from the ribs onto the diaphragm. The lowest part of the pleural cavity, where extra fluid collects.

- The fluid does two jobs: it lets the lung glide against the chest wall, and it holds the two layers together so the lung follows the chest wall as it expands.
- Because the two pleural cavities are separate, air or fluid in one does not collapse the other lung.

Competency 102. Respiratory zone and alveolar structure

In your course notes: Module 4 packet, Chapter 1, competency 102 (page 13).

Dr. Rennie's video for this part: Respiratory system concept video, chapter Alveoli and gas exchange.

You should be able to: Describe the structure of an alveolus and identify the type I and type II alveolar cells, alveolar macrophages, respiratory membrane, and pulmonary capillaries.

Past the terminal bronchioles the airway walls thin into tiny air sacs. The respiratory zone is the bulk of the lung tissue, and at low power it looks lacy because it is mostly air spaces separated by thin walls.

Respiratory bronchioles
: The first airways of the respiratory zone, with a few alveoli budding from their walls.

Alveolar ducts
: Passages whose walls are lined entirely with alveoli.

Alveolar sacs
: Clusters of alveoli opening into a shared space, like a bunch of grapes.

Alveoli
: The single tiny air sacs where gas exchange happens. The two lungs hold hundreds of millions of them.

Alveolar pores
: Small openings connecting neighboring alveoli.

The alveolar wall and the respiratory membrane

Structure	What it is	Job
Type I alveolar cells	Very thin, flat simple squamous cells covering nearly all of the alveolar surface	Gases cross the wall through them
Type II alveolar cells	Rounder, cuboidal cells scattered among the type I cells	Secrete surfactant, which lowers surface tension and keeps the alveoli from collapsing; also secrete antimicrobial proteins
Alveolar macrophages	The dust cells, moving freely over the inner surface of the alveoli	Engulf dust, microbes, and debris that got past the conducting zone
Pulmonary capillaries	A dense net wrapping each alveolus, fed by branches of the pulmonary arteries and drained by the pulmonary veins	Bring blood within a membrane's thickness of the air
Respiratory membrane	The barrier between alveolar air and blood where a capillary presses on an alveolus	Three layers, air to blood: the type I alveolar cell, the fused basement membranes of the alveolus and capillary, and the capillary endothelial cell

- Type I is thin, for exchange. Type II makes surfactant. Macrophages clean up.
- Elastic fibers surround each alveolus and let the lung stretch on the way in and recoil on the way out.
- The membrane is extremely thin and the alveoli are extremely many, so the lungs have a huge surface for gas exchange. That surface is fragile, which is the story of emphysema.

Competency 103. Thoracic wall and diaphragm

In your course notes: Module 4 packet, Chapter 1, competency 103 (page 14).

Dr. Rennie's video for this part: Respiratory system concept video, chapter Breathing muscles and final tips.

You should be able to: Identify the diaphragm, the intercostal muscles, and the three structures passing through the diaphragm at T8, T10, and T12.

The lungs sit inside the thoracic wall: ribs and costal cartilages, the sternum in front, the thoracic vertebrae behind, muscle between the ribs, and the diaphragm closing the bottom. Breathing works by changing the size of this cage. The lungs follow the wall because the pleural fluid holds them to it.

The diaphragm

- A dome-shaped sheet of skeletal muscle forming the floor of the thoracic cavity and separating it from the abdominal cavity. Its fibers attach around the lower rib cage, the sternum, and the lumbar vertebrae and pull toward the central tendon, a flat sheet at the top of the dome.
- The right dome sits higher because the liver is under it.
- When it contracts it flattens and moves down, making the thoracic cavity taller. It is the main muscle of breathing.
- Supplied by the right and left phrenic nerves, from spinal nerves C3, C4, and C5. On the cadaver each phrenic nerve runs down the side of the pericardium, in front of the root of the lung. C3, 4, and 5 keep the diaphragm alive.

The intercostal muscles and the accessory muscles

Muscle	Fiber direction or shape	Role in breathing	Nerve
Diaphragm	Dome, fibers rising to the central tendon	Main muscle of inspiration; contracts, flattens, moves down	Phrenic nerves (C3 to C5)
External intercostals	Downward and forward, like hands sliding into front pockets	Lift the ribs in inspiration, quiet and forced	Intercostal nerves
Internal intercostals	At right angles to the externals, downward and backward	Pull the ribs down in forced expiration (a hard cough, blowing out candles)	Intercostal nerves

- Quiet expiration needs no muscle effort; the lungs and chest wall recoil on their own.
- Harder inspiration recruits the scalenes, sternocleidomastoid, and pectoralis minor. Forced expiration uses the internal intercostals and the abdominal muscles.

Three structures through the diaphragm

Vertebral level	Opening	Main structure	Also passes through
T8	Caval opening, in the central tendon	Inferior vena cava	
T10	Esophageal hiatus, in the muscle	Esophagus	Vagus nerves
T12	Aortic hiatus, behind the diaphragm between the crura	Aorta	Thoracic duct

- I-8, E-10, A-12. Inferior vena cava at T8, Esophagus at T10, Aorta at T12. These are vertebral levels, listed top to bottom. (I Eat At 8, 10, 12 o'clock.)
- Above the diaphragm the vessel is the thoracic aorta; below it, the abdominal aorta. The crura are the muscular legs that anchor the diaphragm to the lumbar vertebrae.

Vessels you will find in the upper thorax (lab)

Structure	Artery or vein	Where it is
Brachiocephalic trunk	Artery	First branch of the aortic arch, right side only; splits into the right subclavian and right common carotid
Common carotid arteries	Artery	Up the neck to the head; the right from the brachiocephalic trunk, the left straight off the arch
Subclavian arteries	Artery	Out to the upper limbs; the right from the brachiocephalic trunk, the left straight off the arch
Internal jugular veins	Vein	Drain the head, down the neck beside the common carotids
Subclavian veins	Vein	Drain the upper limbs
Brachiocephalic veins	Vein	Each forms where an internal jugular joins a subclavian vein; the two join to form the superior vena cava
Pulmonary arteries	Artery	From the heart into each lung at the hilum
Pulmonary veins	Vein	From each lung back to the heart

For each one, say the side and whether it is an artery or a vein before you name it.

Competency 104. Respiratory disorders by structure affected

In your course notes: Module 4 packet, Chapter 1, competency 104 (page 16).

You should be able to: Name common respiratory disorders and state which airway, alveolar, or pleural structure each one affects.

Most respiratory disorders do one of three things. Some narrow or block the airways. Some damage or fill the alveoli, so the exchange surface is lost. Some affect the pleurae and pleural cavity, so the lung cannot stay against the chest wall. Learn the structure first, then what happens to it.

Disorder	Structure affected	What changes in that structure
Rhinitis	Nasal mucosa	The lining is inflamed, swollen, and makes extra mucus (allergies or infection)
Sinusitis	Paranasal sinuses	The sinus lining is inflamed, often when the openings into the nasal cavity are blocked
Laryngitis	Larynx, vocal folds	The vocal folds are inflamed and swollen, cannot vibrate normally, and the voice turns hoarse
Asthma	Bronchi and bronchioles	The lining is inflamed, smooth muscle tightens, extra mucus collects, and the airways narrow. Bronchioles have no cartilage to hold them open
Chronic bronchitis	Bronchi	The lining stays inflamed and makes too much mucus, which blocks the airways
Cystic fibrosis	Bronchi and bronchioles	An inherited disorder; the mucus is thick and sticky, the cilia cannot clear it, and it clogs the airways
Emphysema	Alveoli	Alveolar walls break down and many small alveoli merge into fewer, larger spaces. The exchange surface shrinks and the lung loses elastic recoil. The change is permanent
COPD	Airways and alveoli	Chronic obstructive pulmonary disease, a long-term blockage of airflow; mainly emphysema and chronic bronchitis together
Pneumonia	Alveoli	An infection inflames the alveoli and fills them with fluid. The walls are not destroyed
Tuberculosis	Lung tissue	A bacterial infection that forms small nodules (tubercles) and scar tissue
Infant respiratory distress syndrome	Alveoli (type II cells)	A premature newborn's type II cells do not yet make enough surfactant, so the alveoli collapse with each breath
Lung cancer	Lung tissue	A malignant tumor, usually starting in the epithelium lining the airways or alveoli; strongly linked to smoking
Pleurisy	Pleurae	The pleural layers are inflamed and rough, so they rub with each breath and cause sharp pain
Pleural effusion	Pleural cavity	Extra fluid collects in the pleural cavity and presses on the lung; in an upright person it pools in the costodiaphragmatic recess
Pneumothorax	Pleural cavity	Air enters the pleural cavity, the layers separate, and the lung on that side collapses

What this means for patients

- Asthma narrows the bronchioles so much because they have no cartilage to hold them open.
- Emphysema and pneumonia both take away working alveoli, but only emphysema destroys the walls. Pneumonia fills the sacs; emphysema erases them.

- A pneumothorax collapses one lung, not both, because each lung has its own sealed pleural cavity. The same thin, sealed space that lets the lung glide is what makes it vulnerable.
- Because the right main bronchus is wider, shorter, and more vertical, an inhaled object usually ends up in the right lung.
- The cricothyroid ligament sits just under the skin below the vocal folds with no cartilage over it, so it is the landmark for an emergency airway when the larynx above is blocked.

Worked questions

Eight questions in the style of the lecture exam, one for each competency. Read the answer and the reasons each wrong option is wrong. The wrong options are where the learning is.

QUESTION 1 · COMPETENCY 97

Which structure is the last part of the conducting zone?

A. Segmental bronchus

B. Terminal bronchiole

C. Respiratory bronchiole

D. Alveolar duct

Answer: B. The conducting zone runs from the nose down to the terminal bronchioles. They are the last tubes that only carry air.

Why the others are wrong

- **A** A segmental bronchus is in the conducting zone, but the tree keeps branching past it into bronchioles, so it is not the last part.
- **C** The respiratory bronchiole is the first airway of the respiratory zone. It has alveoli in its wall, so gas exchange starts there.
- **D** Alveolar ducts are lined entirely with alveoli and sit deep in the respiratory zone.

QUESTION 2 · COMPETENCY 98

Which region of the pharynx is lined with pseudostratified ciliated columnar epithelium?

A. Nasopharynx

B. Oropharynx

C. Laryngopharynx

D. All three regions

Answer: A. The nasopharynx carries only air, so it keeps the ciliated, mucus-moving lining of the nasal cavity.

Why the others are wrong

- **B** The oropharynx carries food as well as air, so it is lined with stratified squamous epithelium that resists rubbing.
- **C** The laryngopharynx also carries food and is stratified squamous for the same reason.
- **D** Only the nasopharynx is ciliated. The lining changes where food starts to pass.

QUESTION 3 · COMPETENCY 99

Which laryngeal cartilage is the only complete ring of cartilage in the airway?

A. Thyroid cartilage

B. Epiglottis

C. Cricoid cartilage

D. Arytenoid cartilage

Answer: C. The cricoid is shaped like a signet ring, narrow in front and wide behind, and it goes all the way around. The trachea hangs from its lower edge.

Why the others are wrong

- **A** The thyroid cartilage is the biggest, but it is a shield that is open at the back, not a ring.
- **B** The epiglottis is a single elastic flap, not a ring.
- **D** The arytenoids are small paired cartilages sitting on the back of the cricoid. They anchor the vocal folds.

QUESTION 4 · COMPETENCY 100

An inhaled peanut is most likely to lodge in the right lung because the right main bronchus is

A. narrower, longer, and more horizontal than the left

B. wider, shorter, and more vertical than the left

C. lined with more cilia than the left

D. the only main bronchus with C-shaped cartilage

Answer: B. The right main bronchus is wider, shorter, and more vertical, so an object falling down the trachea tends to drop straight into it.

Why the others are wrong

- **A** That describes the left main bronchus, which angles more sharply to get around the heart.
- **C** Both main bronchi have the same ciliated lining. Cilia move mucus; they do not steer objects.
- **D** Both main bronchi have C-shaped cartilage rings. The difference is the angle and width, not the cartilage.

QUESTION 5 · COMPETENCY 101

A student labels a horizontal fissure on the left lung. What is wrong with the drawing?

A. Nothing; both lungs have a horizontal fissure

B. The left lung has only an oblique fissure

C. The horizontal fissure belongs on the base, not the front

D. Fissures are only visible on the right lung

Answer: B. The left lung has two lobes separated by one oblique fissure. The horizontal fissure separates the superior and middle lobes of the right lung, and the left lung has no middle lobe.

Why the others are wrong

- **A** Only the right lung has three lobes and therefore two fissures.
- **C** The horizontal fissure runs across the front and side of the right lung at about the level of the fourth rib; it is not on the base.
- **D** The left lung has a clearly visible oblique fissure. It is the horizontal fissure it lacks.

QUESTION 6 · COMPETENCY 102

Which cell secretes surfactant?

A. Type I alveolar cell

B. Type II alveolar cell

C. Alveolar macrophage

D. Capillary endothelial cell

Answer: B. Type II alveolar cells are the cuboidal cells scattered in the alveolar wall. They make surfactant, which lowers surface tension so the alveoli do not collapse.

Why the others are wrong

- **A** Type I cells are the thin, flat cells that make up most of the wall. Gases cross through them, but they do not make surfactant.
- **C** Alveolar macrophages are the dust cells. They engulf debris; they do not secrete surfactant.
- **D** The capillary endothelium is the blood-side layer of the respiratory membrane. It belongs to the vessel, not the alveolus.

QUESTION 7 · COMPETENCY 103

The esophagus passes through the diaphragm at the level of which vertebra?

A. T8

B. T10

C. T12

D. L1

Answer: B. The esophageal hiatus is at T10. I-8, E-10, A-12: the inferior vena cava at T8, the esophagus at T10, the aorta at T12.

Why the others are wrong

- **A** T8 is the caval opening in the central tendon, for the inferior vena cava.
- **C** T12 is the aortic hiatus, behind the diaphragm between the crura, for the aorta and thoracic duct.
- **D** Nothing on this list passes the diaphragm at L1. The celiac trunk and superior mesenteric artery branch off the aorta near that level, below the diaphragm.

QUESTION 8 · COMPETENCY 104

Which disorder is air in the pleural cavity with collapse of the lung on that side?

A. Pleural effusion

B. Pneumonia

C. Pneumothorax

D. Emphysema

Answer: C. Pneumothorax is air between the parietal and visceral pleurae. The seal breaks, the lung pulls away from the chest wall, and that lung collapses. Only one side is affected because each lung has its own pleural cavity.

Why the others are wrong

- **A** Pleural effusion is fluid in the pleural cavity, not air. It pools in the costodiaphragmatic recess and presses on the lung.
- **B** Pneumonia is an infection inside the alveoli, which fill with fluid. The pleural cavity is not involved.
- **D** Emphysema destroys alveolar walls inside the lung. It is not air in the pleural space.

BIO 004 Human Anatomy · Module 4 notes · Dr. Sharilyn Rennie

The lymphatic system

The written notes for the lymphatic system lecture: the one-way lymph pathway, the structure of lymphatic vessels and lymph nodes, the primary and secondary lymphoid organs, and the common lymphatic disorders. Module 4 competencies 105 to 108, with eight worked questions.

Competency 105. Lymph pathway

In your course notes: Module 4 packet, Chapter 2, competency 105 (page 18).

Dr. Rennie's video for this part: Lymphatic system concept video.

You should be able to: Define lymph and trace its one-way route from lymphatic capillaries through collecting vessels, nodes, trunks, and ducts back to the subclavian veins.

Blood capillaries leak. All day long, fluid filters out of them into the spaces between cells, and that fluid is called interstitial fluid. The blood capillaries take most of it back, but some fluid and a small amount of plasma protein stay behind in the tissues. The lymphatic system collects this leftover fluid, about 3 liters a day, and returns it to the blood. That keeps fluid from building up in the tissues and keeps blood volume steady.

Once interstitial fluid enters a lymphatic vessel, it is called lymph. Lymph is clear and watery. It carries water, plasma proteins, and whatever else it picks up in the tissues, such as cell debris, bacteria, and sometimes cancer cells. Lymph coming from the small intestine after a fatty meal looks milky white. This fat-rich lymph is called chyle.

The fluids and the system

Lymphatic system
: A network of vessels, tissues, and organs that drains tissue fluid and houses the body's defenses.

Interstitial fluid
: The fluid in the spaces between cells, filtered out of the blood capillaries. In the tissue spaces it is still interstitial fluid, not lymph.

Lymph
: Interstitial fluid once it has entered a lymphatic vessel. It becomes lymph the moment it enters a lymphatic capillary. Clear and watery.

Chyle
: Milky, fat-rich lymph coming from the lacteals of the small intestine.

Three functions
: Returns excess interstitial fluid to the blood, absorbs dietary fats from the small intestine, and houses the immune defenses.

The one-way route, in order

The lymphatic system is a one-way system. It does not form a loop the way blood vessels do. Lymph starts in the tissues and flows only toward the heart. As it travels, small vessels join into larger and larger vessels, the way small streams join to form a river. On the way, the lymph passes through lymph nodes, which filter it before it reaches the blood.

1. Lymphatic capillaries: blind-ended tubes in the tissues take up excess interstitial fluid. The fluid is now lymph.
2. Lymphatic collecting vessels: larger vessels with valves carry the lymph onward and keep it moving one way.
3. Lymph nodes: nodes spaced along the collecting vessels filter the lymph. Lymph usually passes through several nodes in a row.
4. Lymphatic trunks: collecting vessels merge into large trunks. Each trunk drains one major region of the body.
5. Lymphatic ducts: the trunks empty into one of two ducts, the right lymphatic duct or the larger thoracic duct.
6. Subclavian veins: each duct empties into the blood where the subclavian vein meets the internal jugular vein on its own side. At this point lymph becomes part of the blood plasma again.

- Worked example from the note sheet, starting at the left foot: a lymphatic capillary in the foot, a collecting vessel with valves running up the leg, an inguinal node in the groin, the left lumbar trunk running up the back wall of the abdomen, the cisterna chyli, the thoracic duct up through the thorax, and finally the junction of the left subclavian and left internal jugular veins. Every arrow points toward the heart and none point back.

The nine trunks

There are nine lymphatic trunks. Each one is named for the region it drains. Four of them come in left and right pairs, and one, the intestinal trunk, is single.

Trunk	Region drained	Empties into
Lumbar (left and right)	Lower limbs, pelvis, and abdominal wall	Thoracic duct, at the cisterna chyli
Intestinal (single)	Digestive organs of the abdomen	Thoracic duct, at the cisterna chyli
Bronchomediastinal (left and right)	Thoracic organs and the thoracic wall	Left: thoracic duct. Right: right lymphatic duct
Subclavian (left and right)	Upper limbs	Left: thoracic duct. Right: right lymphatic duct
Jugular (left and right)	Head and neck	Left: thoracic duct. Right: right lymphatic duct

The two ducts

The right lymphatic duct is short and small. It drains the right upper limb and the right side of the head, neck, and thorax. The thoracic duct is the largest lymphatic vessel in the body. It drains everything else, which is about three quarters of the body. The thoracic duct begins in the abdomen as a sac called the cisterna chyli, which sits in front of the upper lumbar vertebrae. From there it passes up through the diaphragm with the aorta and climbs through the thorax beside the vertebral column.

Both ducts end in the base of the neck. Each one empties into the junction of the subclavian vein and the internal jugular vein on its own side. The right lymphatic duct empties on the right side. The thoracic duct empties on the left side.

Feature	Right lymphatic duct	Thoracic duct
Size	Short and small	Largest lymphatic vessel
Drains	Right upper limb and the right side of the head, neck, and thorax	Everything else, about three quarters of the body
Trunks it receives	Right jugular, right subclavian, right bronchomediastinal	Both lumbar, intestinal, left jugular, left subclavian, left bronchomediastinal
Starts at	The right side of the base of the neck	The cisterna chyli, a sac in the abdomen
Empties into	Junction of the right subclavian and right internal jugular veins	Junction of the left subclavian and left internal jugular veins

- Memory tip: split the body into four quarters. The right upper quarter (right arm, right side of the head, neck, and chest) drains into the right lymphatic duct. The other three quarters drain into the thoracic duct. The right leg is thoracic duct territory, not right lymphatic duct territory.

How lymph moves without a pump

The lymphatic system has no heart to push the lymph. Lymph is moved by forces from the structures around the vessels, and the valves keep it from sliding backward.

- Skeletal muscle pump: contracting muscles squeeze the vessels that run between them.
- Respiratory pump: pressure changes in the thorax with each breath pull lymph upward.
- Smooth muscle in the walls of the larger lymphatic vessels contracts in rhythm.
- Pulsing of nearby arteries presses on the vessels.

- Valves in the collecting vessels keep all of this movement going one way, toward the heart.

Competency 106. Vessel and node structure

In your course notes: Module 4 packet, Chapter 2, competency 106 (page 20).

Dr. Rennie's video for this part: Lymphatic system concept video.

You should be able to: Describe the structure of lymphatic capillaries, valved collecting vessels, and a lymph node including capsule, cortex, medulla, hilum, and afferent and efferent vessels.

Lymphatic capillaries

Lymphatic capillaries are the smallest lymphatic vessels. They lie in the tissues, woven between the blood capillaries. Each one is a thin tube that is closed at one end, so it is called blind-ended. Its wall is a single layer of endothelial cells.

The endothelial cells of a lymphatic capillary overlap at their edges. These loose, overlapping endothelial cells act as one-way flap valves. When fluid builds up in the tissue, the flaps push open and fluid flows in. When pressure inside the capillary rises, the flaps press shut, so the fluid cannot flow back out. Fine collagen strands called anchoring filaments tie the endothelial cells to the surrounding tissue. When the tissue swells, the filaments pull on the cells and hold the capillary open instead of letting it collapse.

Because the gaps between the cells are large, lymphatic capillaries are more permeable than blood capillaries. Proteins, cell debris, bacteria, and even cancer cells can get in. Lymphatic capillaries are found in nearly every tissue. They are absent from the central nervous system (the brain and spinal cord), bone and red bone marrow, and avascular tissues such as cartilage, the epidermis, and the cornea.

Feature	Lymphatic capillary	Blood capillary
Ends	Blind (closed) at one end	Open at both ends, part of a loop
Size	Wider	Narrower
Wall	Overlapping endothelial cells act as one-way flap valves	Endothelial cells joined more tightly
Held open by	Anchoring filaments tied to the surrounding tissue	Pressure of the blood inside
What gets in	Fluid, proteins, cells, pathogens, and even cancer cells	Mostly small molecules
Flow	One way, away from the tissues	Through, from the artery side to the vein side

- In the small intestine, each villus has a special lymphatic capillary in its center called a lacteal. Lacteals absorb dietary fats, which are packaged in particles too large to enter blood capillaries. The fat turns the lymph milky, and this lymph (chyle) drains to the cisterna chyli and up the thoracic duct.

Collecting vessels and their valves

Lymphatic capillaries join to form lymphatic collecting vessels. These are built like small veins. Their walls have the same three layers as a vein, but the layers are thinner. Collecting vessels have more internal valves than veins do. These lymphatic valves are one-way flaps, so lymph can move forward but cannot flow backward. When a collecting vessel is full, the stretches between valves bulge, which gives the vessel a beaded look.

- Superficial collecting vessels run in the subcutaneous layer of the skin alongside the superficial veins. Deep collecting vessels run alongside the deep arteries.
- Along their length, the collecting vessels pass through lymph nodes.

The lymph node

A lymph node is a small, bean-shaped organ that filters lymph. Hundreds of nodes lie along the collecting vessels. They are clustered in the neck (cervical nodes), the armpit (axillary nodes), and the groin (inguinal nodes), and there are many more deep in the thorax and abdomen.

The outside of a node is a capsule of dense connective tissue. Strands of connective tissue called trabeculae

extend inward from the capsule and divide the node into compartments. Inside, a mesh of reticular fibers holds the lymphocytes and macrophages in place.

The node has two regions. The outer region is the cortex. The outer cortex holds rounded clusters of lymphocytes called lymphatic follicles, which are mostly B cells. Many follicles have a pale germinal center where B cells multiply. The deep cortex is mostly T cells. The inner region is the medulla. It contains medullary cords, strands of lymphocytes and plasma cells, separated by medullary sinuses, open channels lined with macrophages.

One side of the node is indented. This is the hilum. Blood vessels enter and leave the node at the hilum, and lymph leaves there too. Afferent lymphatic vessels carry lymph into the node. There are several of them, and they enter at many points on the rounded (convex) surface through the capsule. Efferent lymphatic vessels carry filtered lymph out of the node, and they leave only at the hilum. Because there are more afferent vessels than efferent vessels, lymph backs up and moves slowly through the node. This gives the macrophages and lymphocytes time to work.

Structure	What it is
Capsule	Dense connective tissue covering; trabeculae extend inward from it
Afferent lymphatic vessels	Several vessels that carry lymph into the node through the capsule, on the convex surface
Subcapsular sinus	The space just under the capsule where lymph first enters
Cortex	Outer region; the outer cortex has lymphatic follicles of B cells with germinal centers, and the deep cortex is mostly T cells
Medulla	Inner region of medullary cords and medullary sinuses lined with macrophages
Hilum	The indented side of the node, where blood vessels enter and leave and efferent vessels leave
Efferent lymphatic vessels	Fewer vessels that carry filtered lymph out of the node at the hilum

The path of lymph through a node

1. Afferent lymphatic vessels: lymph enters through the capsule on the convex surface.
2. Subcapsular sinus: lymph spreads out in the space just under the capsule.
3. Cortical sinuses: lymph seeps through the cortex, past the follicles.
4. Medullary sinuses: macrophages lining these channels engulf bacteria, debris, and cancer cells.
5. Efferent lymphatic vessels: filtered lymph leaves at the hilum and moves on to the next node or a trunk.

What keeps lymph moving one way at each step

Structure	Wall or covering	What keeps lymph moving one way	Where lymph goes next
Lymphatic capillary	One layer of overlapping endothelial cells, held open by anchoring filaments	The overlapping cells act as flap valves	A collecting vessel
Collecting vessel	Built like a vein, with three thin layers	Many valves, more than veins have	A lymph node
Lymph node	A capsule of dense connective tissue	Lymph comes in only through the afferent vessels and leaves only through the efferent vessels	Out through the efferent vessels at the hilum, on toward a trunk

- Memory tip: afferent vessels arrive, efferent vessels exit. The same two words mean "in" and "out" in other systems too, such as the arterioles of the kidney and the neurons of the nervous system.

- Afferent vessels enter all around the capsule, but efferent vessels leave only at the hilum. In a drawing, more should come in than go out.

Competency 107. Primary and secondary organs

In your course notes: Module 4 packet, Chapter 2, competency 107 (page 22).

Dr. Rennie's video for this part: Lymphatic system concept video.

You should be able to: Sort red bone marrow, thymus, lymph nodes, spleen, tonsils, and MALT into primary and secondary lymphoid organs with location and role.

The main cells of the lymphatic system are lymphocytes, a type of white blood cell. The two kinds you need here are B cells and T cells. All lymphocytes are made in the red bone marrow. Before they can fight anything, they have to finish maturing. B cells mature in the red bone marrow. T cell precursors leave the marrow and travel in the blood to the thymus, where they mature.

This gives a simple way to sort the lymphoid (lymphatic) organs. Primary lymphoid organs are where lymphocytes are made and mature. There are only two: the red bone marrow and the thymus. Secondary lymphoid organs are where mature lymphocytes gather and meet pathogens. These are the lymph nodes, the spleen, the tonsils, and MALT. Mature lymphocytes leave the marrow and thymus and travel out to the secondary organs. Each secondary organ sits where it can catch pathogens from one source: the lymph nodes check lymph, the spleen checks blood, and the tonsils and MALT guard the linings of open tracts.

Organ	Group	Location	Role
Red bone marrow	Primary	Spongy bone of the axial skeleton (such as the sternum and vertebrae), the limb girdles (such as the hip bones), and the proximal ends of the femur and humerus	Where all lymphocytes form, and where B cells mature
Thymus	Primary	Mediastinum, above the heart and behind the sternum	Where T cells mature
Lymph nodes	Secondary	Along the collecting vessels, clustered in the cervical, axillary, and inguinal regions (neck, armpits, and groin)	Filter the lymph
Spleen	Secondary	Upper left abdomen, to the left of and behind the stomach, under the lower left ribs	Filters the blood
Tonsils	Secondary	A ring of lymphatic tissue in the walls of the pharynx and at the base of the tongue, around the entrance to the throat	Guard against inhaled and swallowed pathogens
MALT	Secondary	Mucous membranes of the digestive, respiratory, urinary, and reproductive tracts	Guards the mucosal surfaces

- Memory tip: only two organs are primary, the red bone marrow and the thymus, because those are the only places lymphocytes are made and matured. B cells mature in the bone marrow and T cells mature in the thymus. Everything else on the list is secondary.
- The spleen is the largest lymphatic organ, but it is secondary. Size does not make an organ primary; making and maturing lymphocytes does.

The thymus

The thymus sits in the mediastinum, above the heart and just behind the sternum. It is large in infants and children. After puberty it shrinks, and in older adults it is mostly fat. Its outer cortex is packed with immature T cells. Its inner medulla has fewer, more mature T cells and small round structures called thymic corpuscles. The thymus does not filter lymph. Unlike a lymph node, it has no afferent lymphatic vessels and no follicles.

The spleen

The spleen is the largest lymphatic organ. It sits in the upper left abdomen against the diaphragm, to the left of and behind the stomach, and it is protected by the lower left ribs. The splenic artery and splenic vein pass through its hilum. The spleen filters the blood. It removes worn-out red blood cells and platelets and responds to

pathogens carried in the blood. Its tissue comes in two kinds.

Structure	What it is
Spleen	The largest lymphatic organ, in the upper left abdomen under the lower left ribs
White pulp	Small islands of lymphocytes clustered around central arteries; carries out the spleen's immune functions
Red pulp	Most of the organ; venous sinuses and splenic cords full of red blood cells and macrophages that filter the blood and remove worn-out erythrocytes

The tonsils and MALT

MALT stands for mucosa-associated lymphatic tissue. It is lymphatic tissue in the mucous membranes that line the open tracts of the body, where pathogens can get in.

The largest and best organized pieces of MALT are the tonsils, the Peyer patches, and the appendix.

The tonsils form a ring of lymphatic tissue around the entrance to the pharynx, where they meet inhaled and swallowed material. Their surface has deep pits called crypts that trap bacteria and debris.

Structure	Location	Notes
Pharyngeal tonsil	Posterior wall of the nasopharynx	Single; also called the adenoid
Palatine tonsils	Lateral walls of the oropharynx, between the palatoglossal and palatopharyngeal arches	Paired and the largest; the ones usually infected and removed
Lingual tonsils	Base (posterior third) of the tongue, facing the oropharynx	Guard swallowed material
Tubal tonsils	Around the openings of the auditory tubes in the nasopharynx	Paired
Peyer patches	Wall of the ileum, the last part of the small intestine	Clusters of lymphatic follicles that watch the intestinal contents
Appendix	Off the cecum of the large intestine	Wall packed with lymphatic follicles

Competency 108. Lymphatic disorders

In your course notes: Module 4 packet, Chapter 2, competency 108 (page 24).

You should be able to: Name edema, lymphedema, lymphadenopathy, lymphoma, splenomegaly, and tonsillitis and identify the structure each one affects.

Lymphatic disorders fall into two groups. In the first group, drainage fails and fluid collects in the tissues, which shows up as swelling. In the second group, the lymphatic tissue itself changes. A node, the spleen, or the tonsils become enlarged, inflamed, or cancerous.

Edema is swelling caused by excess fluid in the interstitial space. It happens when fluid leaks out of the blood capillaries faster than the lymphatic vessels can carry it away. Lymphedema is a specific kind of edema. It is long-term swelling caused by blocked, damaged, or missing lymphatic vessels. It can follow surgery that removes lymph nodes, because the vessels that passed through those nodes no longer drain the region. The difference between the two is where the problem sits: edema is a problem in the tissue space, while lymphedema is a problem in the lymphatic vessels themselves.

Lymphadenopathy means enlarged lymph nodes. Nodes often swell when they are fighting an infection nearby, and they can also enlarge when cancer cells lodge in them. Lymphoma is a cancer of the lymphocytes in lymphatic tissue. It usually starts in lymph nodes and comes in two main groups, Hodgkin lymphoma and non-Hodgkin lymphoma. Lymphadenopathy is any enlarged node, for any reason; lymphoma is a cancer of the lymphocytes. Splenomegaly is an enlarged spleen. It can

happen with an infection such as mononucleosis or with a blood disorder such as leukemia. Tonsillitis is inflammation of the tonsils, usually the palatine tonsils, and it is common in childhood.

Disorder	Structure affected	What changes
Edema	The interstitial space	Excess interstitial fluid builds up and the tissue swells, because fluid leaks out faster than the lymphatics can drain it
Lymphedema	The lymphatic vessels	Blocked, damaged, or missing vessels cause long-term swelling of the region they drain, sometimes after lymph node removal
Lymphadenopathy	The lymph nodes	The nodes enlarge, often while fighting an infection, and sometimes because cancer cells have lodged in them
Lymphoma	Lymphatic tissue, usually the lymph nodes	Lymphocytes become cancerous and the tissue enlarges; Hodgkin or non-Hodgkin lymphoma
Splenomegaly	The spleen	The spleen enlarges, for example in mononucleosis or leukemia, and becomes easier to rupture
Tonsillitis	The tonsils, usually the palatine tonsils	The tonsils become infected, red, and swollen

What this means for patients

- Swollen glands are lymphadenopathy. Lymph nodes enlarge as they trap and fight pathogens, which is why a clinician feels for tender nodes in the neck, armpit, and groin during an exam.
- Because lymph passes through nodes in order, the lymphatic system also maps how some cancers spread. The first node a region drains into is the sentinel node, and checking it tells a surgeon whether a cancer has begun to travel. Removing many nodes can block drainage and leave lasting lymphedema in the limb.
- The spleen sits just under the lower left ribs. Its soft tissue tears easily, and it holds a large volume of blood, so a blow to that area can rupture the spleen and cause heavy internal bleeding. An enlarged spleen, as in mononucleosis, ruptures even more easily, which is why contact sports are dangerous during mononucleosis.

Worked questions

Eight questions in the style of the lecture exam, two for each competency. Read the answer and the reasons each wrong option is wrong. The wrong options are where the learning is.

QUESTION 1 · COMPETENCY 105

The thoracic duct empties lymph into the blood at which point?

A. The junction of the right subclavian and right internal jugular veins

B. The cisterna chyli

C. The superior vena cava

D. The junction of the left subclavian and left internal jugular veins

Answer: D. Both ducts empty where a subclavian vein meets an internal jugular vein, each on its own side. The thoracic duct ends on the left, so it empties into the junction of the left subclavian and left internal jugular veins.

Why the others are wrong

- **A** That is where the right lymphatic duct empties. The thoracic duct ends on the left side of the base of the neck.
- **B** The cisterna chyli is where the thoracic duct begins, a sac in the abdomen in front of the upper lumbar vertebrae. Lymph enters the duct there; it does not enter the blood there.
- **C** The ducts end at the vein junctions in the base of the neck. Blood from those junctions flows on into the superior vena cava, but the ducts do not open into it directly.

QUESTION 2 · COMPETENCY 105

Lymph from the right leg returns to the blood through which vessel?

A. The right lymphatic duct

B. The thoracic duct

C. The right subclavian trunk

D. The right jugular trunk

Answer: B. The right lymphatic duct drains only the right upper quarter of the body: the right arm and the right side of the head, neck, and thorax. Every other region, including both legs, drains through the lumbar trunks into the cisterna chyli and up the thoracic duct.

Why the others are wrong

- **A** The right lymphatic duct is short and small and takes only the right jugular, right subclavian, and right bronchomediastinal trunks. The right leg is not in its territory.
- **C** The subclavian trunks drain the upper limbs. The right leg drains into the right lumbar trunk.
- **D** The jugular trunks drain the head and neck, not a limb.

QUESTION 3 · COMPETENCY 106

Which vessels carry lymph out of a lymph node, and where do they leave?

A. Efferent vessels, leaving only at the hilum

B. Afferent vessels, leaving only at the hilum

C. Efferent vessels, leaving through the capsule all over the convex surface

D. Afferent vessels, leaving through the capsule all over the convex surface

Answer: A. Efferent vessels exit. Filtered lymph leaves a node through a few efferent vessels at the hilum, the indented side where the blood vessels also enter and leave. Because several afferent vessels bring lymph in and fewer efferent vessels take it out, lymph slows down inside the node.

Why the others are wrong

- **B** Afferent vessels arrive. They carry lymph into the node, and they do not use the hilum.
- **C** Lymph enters through the capsule on the convex surface; it does not leave there. Efferent vessels are found only at the hilum.
- **D** Afferent vessels do enter through the capsule on the convex surface, but they carry lymph in, not out.

QUESTION 4 · COMPETENCY 106

What holds a lymphatic capillary open when the surrounding tissue swells with fluid?

A. The pressure of the lymph inside it

B. A thick layer of smooth muscle in its wall

C. Anchoring filaments tied to the surrounding tissue

D. Valves in its lumen

Answer: C. Anchoring filaments are fine collagen strands that tie the endothelial cells of the lymphatic capillary to the tissue around it. When the tissue swells, the filaments pull on the cells and hold the capillary open, so fluid can flow in instead of the tube collapsing.

Why the others are wrong

- **A** Internal pressure holds a blood capillary open. In a lymphatic capillary, rising pressure inside presses the overlapping flaps shut; it is not what keeps the tube open.
- **B** A lymphatic capillary wall is a single layer of endothelial cells with no muscle. Smooth muscle appears in the walls of the larger lymphatic vessels.
- **D** Lymphatic capillaries have no internal valves. Their one-way flow comes from the overlapping endothelial cells acting as flap valves. Internal valves are a feature of the collecting vessels.

QUESTION 5 · COMPETENCY 107

Which two organs are the primary lymphoid organs?

A. Spleen and thymus

B. Red bone marrow and thymus

C. Red bone marrow and spleen

D. Lymph nodes and thymus

Answer: B. Primary lymphoid organs are where lymphocytes are made and mature. All lymphocytes form in the red bone marrow, B cells mature there, and T cells mature in the thymus. Those are the only two places that happens.

Why the others are wrong

- **A** The thymus is primary, but the spleen is secondary. It filters blood and is where mature lymphocytes meet pathogens; it does not make or mature them. Being the largest lymphatic organ does not make it primary.
- **C** Red bone marrow is primary, but the spleen is secondary for the same reason.
- **D** The thymus is primary, but lymph nodes are secondary. They filter lymph and hold mature lymphocytes.

QUESTION 6 · COMPETENCY 107

Which lymphoid organ filters the blood, removing worn-out red blood cells and platelets?

A. Thymus

B. Lymph node

C. Palatine tonsil

D. Spleen

Answer: D. The spleen filters blood. Its red pulp is full of venous sinuses, splenic cords, and macrophages that remove worn-out erythrocytes and platelets, and its white pulp responds to pathogens carried in the blood. It sits in the upper left abdomen under the lower left ribs.

Why the others are wrong

- **A** The thymus is where T cells mature. It does not filter blood or lymph, and it has no afferent lymphatic vessels and no follicles.
- **B** A lymph node filters lymph, not blood. Each secondary organ checks one source, and nodes check the lymph that passes through them.
- **C** The palatine tonsils are part of the ring of lymphatic tissue around the entrance to the pharynx. They guard against inhaled and swallowed pathogens; they do not filter blood.

QUESTION 7 · COMPETENCY 108

A patient has long-term swelling of one arm after surgery that removed the axillary lymph nodes. Which disorder is this, and which structure is affected?

A. Edema; the interstitial space

B. Lymphadenopathy; the lymph nodes

C. Lymphedema; the lymphatic vessels

D. Lymphoma; lymphatic tissue

Answer: C. Lymphedema is long-term swelling caused by blocked, damaged, or missing lymphatic vessels. Removing the axillary nodes cuts the vessels that passed through them, so the arm no longer drains and the swelling lasts.

Why the others are wrong

- **A** Edema is swelling in the interstitial space because fluid leaks out faster than healthy lymphatics can drain it. Here the problem is the lymphatic vessels themselves, which makes it lymphedema, a specific kind of edema.
- **B** Lymphadenopathy is enlarged lymph nodes. In this patient the nodes have been removed, not enlarged.
- **D** Lymphoma is a cancer of the lymphocytes in lymphatic tissue. Nothing here is cancerous; the swelling is a drainage problem.

QUESTION 8 · COMPETENCY 108

A student with mononucleosis is told to avoid contact sports for several weeks. Which disorder makes this necessary, and which organ is at risk?

A. Splenomegaly; the spleen could rupture

B. Tonsillitis; the tonsils could rupture

C. Lymphadenopathy; the neck nodes could rupture

D. Lymphoma; the lymph nodes could rupture

Answer: A. Mononucleosis can enlarge the spleen, which is splenomegaly. The spleen sits just under the lower left ribs, its soft tissue tears easily, and it holds a large volume of blood, so a blow to that area can rupture it and cause heavy internal bleeding. An enlarged spleen ruptures even more easily.

Why the others are wrong

- **B** Tonsillitis is inflammation of the tonsils, usually the palatine tonsils. It can be part of an illness, but it is not the reason to avoid contact sports.
- **C** Lymphadenopathy is enlarged lymph nodes, which do occur in infections, but the structure at risk from a blow is the spleen, not the nodes.
- **D** Lymphoma is a cancer of the lymphocytes. Mononucleosis is an infection, not a cancer.

BIO 004 Human Anatomy · Module 4 notes · Dr. Sharilyn Rennie

The digestive system, part I: through the esophagus

The written notes for the first digestive lecture: the tube and its wall, the enteric plexuses, the peritoneum, the mouth, salivary glands, tongue, pharynx, esophagus, and the sphincters. Module 4 competencies 109 to 114, 120, and 121, with nine worked questions.

Before the competencies: the two groups of organs and the six jobs

In your course notes: Module 4 packet, Chapter 3 introduction (page 25).

Dr. Rennie's video for this part: Digestive system concept video, chapter Digestive organ overview, and GI tract core processes.

The digestive system is two groups of organs. The gastrointestinal (GI) tract, also called the alimentary canal, is one continuous tube from the mouth to the anus that runs through the thoracic and abdominal cavities: mouth, most of the pharynx, esophagus, stomach, small intestine, and large intestine. The accessory digestive organs sit beside the tube: the teeth and tongue, which touch food directly, and the salivary glands, liver, gallbladder, and pancreas, which make or store secretions and send them into the tube through ducts.

- In a living person the tract is about 5 to 7 m (16.5 to 23 feet) long, because its muscle is in a steady state of partial contraction (tonus). After death the muscle relaxes and the tube measures about 7 to 9 m (23 to 29.5 feet).

The six basic processes of the GI tract

1. Ingestion: taking food and liquid into the mouth.
2. Secretion: water, acid, buffers, and enzymes released into the lumen.
3. Mixing and propulsion: the smooth muscle of the wall contracts and relaxes, mixing food with secretions and pushing it forward.
4. Digestion: breaking food into smaller molecules. Mechanical digestion is the cutting and grinding of the teeth and the churning of the stomach and small intestine. Chemical digestion is enzymes splitting carbohydrates, lipids, proteins, and nucleic acids by hydrolysis. Vitamins, ions, water, and cholesterol can be absorbed without chemical digestion.
5. Absorption: digested molecules cross the epithelium from the lumen into the blood or lymph.
6. Defecation: eliminating what cannot be used (feces: wastes, indigestible material, bacteria, and cells shed from the lining).

Competency 109. Four-layer gut wall

In your course notes: Module 4 packet, Chapter 3, competency 109 (page 25).

Dr. Rennie's video for this part: Digestive system concept video, chapter GI wall layers and mucosa, and Epithelial types and functions.

You should be able to: Name the four layers of the alimentary canal wall from lumen outward and state the dominant tissue of each.

From the esophagus to the anal canal the wall of the alimentary canal has the same four layers. From the lumen outward: mucosa, submucosa, muscularis, serosa (or adventitia). Every region changes the details to suit its job, but the order never changes.

The mucosa, the inner lining, has three parts

1. Epithelium, in direct contact with the food. Nonkeratinized stratified squamous epithelium in the mouth, pharynx, esophagus, and anal canal, where the job is protection from rubbing. Simple columnar epithelium in the stomach and intestines, where the job is secretion and absorption. Tight junctions seal the cells together so nothing leaks between them, and the cells turn over every 5 to 7 days. Exocrine cells secrete mucus and fluid into the lumen; enteroendocrine cells secrete hormones into the blood.

2. Lamina propria: areolar connective tissue carrying the blood and lymphatic vessels that pick up absorbed nutrients. It supports the epithelium and holds mucosa-associated lymphatic tissue (MALT), the lymphatic nodules that guard the tract. MALT is heaviest in the tonsils, the small intestine, the appendix, and the large intestine.
3. Muscularis mucosae: a thin layer of smooth muscle. In the stomach and small intestine it throws the lining into small folds, which adds surface area and keeps every cell exposed to the contents.

The other three layers

Layer	Main tissue	What it holds
Submucosa	Areolar connective tissue (OpenStax calls it dense); binds the mucosa to the muscularis	Larger blood and lymphatic vessels that receive absorbed food molecules; the submucosal plexus; glands in some regions
Muscularis	Smooth muscle in most of the tube, in two sheets: an inner circular layer and an outer longitudinal layer, with the myenteric plexus between them	Skeletal muscle instead in the mouth, pharynx, and the upper and middle esophagus (voluntary swallowing) and in the external anal sphincter (voluntary defecation)
Serosa	A serous membrane: simple squamous epithelium (mesothelium) over areolar connective tissue. It is the visceral peritoneum and suspends the tract in the abdominal cavity	Where the tube is not in the peritoneal cavity, an adventitia of areolar connective tissue replaces it: most of the esophagus, and the back of retroperitoneal organs

- Read the wall from the inside out: mucosa, submucosa, muscularis, serosa.
- The muscularis mucosae is a thin layer inside the mucosa. It is not the muscularis. Keep them in separate rows of your table.
- Two places break the muscle pattern. The stomach has a third, oblique layer (competency 115). The large intestine gathers its outer longitudinal layer into three bands, the teniae coli (competency 117).

Competency 110. Enteric nerve plexuses

In your course notes: Module 4 packet, Chapter 3, competency 110 (page 26).

Dr. Rennie's video for this part: Digestive system concept video, chapter Muscle layers and gut nerves, and Autonomic control and plexuses.

You should be able to: Locate the submucosal and myenteric plexuses in the gut wall and match each to the wall layer it sits within.

The tract has its own nervous system built into its wall, the enteric nervous system, sometimes called the brain of the gut. About 100 million neurons run from the esophagus to the anus, arranged in two plexuses. Each sits in a different layer, and the layer tells you the job.

Plexus	Wall layer	Motor neurons supply	Controls
Myenteric plexus (Auerbach)	Muscularis, between the inner circular and outer longitudinal layers	The two smooth muscle layers	Motility: how often and how hard the muscularis contracts
Submucosal plexus (Meissner)	Submucosa, just deep to the mucosa	The secretory cells of the mucosa and the vessels to it	Secretion by the GI organs, and blood flow to the lining

- Each plexus holds motor neurons, interneurons, and sensory neurons. Interneurons connect the two plexuses. Sensory neurons reach the mucosa, where chemoreceptors respond to chemicals in the food and mechanoreceptors respond to stretch when the wall fills.
- The name tells you the layer. Sub-mucosal is in the submucosa. Myenteric comes from myo, muscle, so it sits in the muscularis.
- Secretion goes with the submucosal plexus; movement goes with the myenteric plexus.

The autonomic nervous system adjusts the enteric system

- Parasympathetic fibers reach the tract through the vagus nerve (cranial nerve X) for everything down to about the first half of the large intestine, and through pelvic splanchnic nerves from the sacral cord for the last half. Preganglionic fibers synapse on neurons in the myenteric and submucosal plexuses; some reach the smooth muscle and glands directly. Parasympathetic activity increases secretion and motility.
- Sympathetic fibers come from the thoracic and upper lumbar cord. Postganglionic fibers synapse on plexus neurons and inhibit them. Sympathetic activity decreases secretion and motility, which is why anger, fear, and anxiety slow digestion.
- GI reflexes: receptors in the wall sense stretch or chemicals, signal neurons in the enteric system, the CNS, or the ANS, and those neurons turn GI glands and smooth muscle up or down.

Competency 111. Peritoneum and folds

In your course notes: Module 4 packet, Chapter 3, competency 111 (page 27).

Dr. Rennie's video for this part: Digestive system concept video, chapter Peritoneum and abdominal folds.

You should be able to: Identify the parietal and visceral peritoneum, the peritoneal cavity, and the five folds greater omentum, lesser omentum, falciform ligament, mesentery, and mesocolon and state what each fold connects.

The peritoneum is the largest serous membrane in the body: a layer of simple squamous epithelium (mesothelium) on areolar connective tissue. Like the pleura, it has two layers with a film of serous fluid between them so the organs slide without friction.

Parietal peritoneum
Lines the walls of the abdominopelvic cavity.

Visceral peritoneum
Covers the organs. On a canal organ it is the serosa.

Peritoneal cavity
The thin space between the two layers, holding a little lubricating peritoneal fluid.

Ascites
In liver disease, pancreatic disease, and some cancers, the peritoneal cavity fills with liters of fluid that often has to be drained.

Unlike the pleura, the peritoneum has large folds that weave between the organs. A fold is a double layer of peritoneum. The folds hold organs to each other and to the walls, store fat between their layers, and carry the blood vessels, lymphatic vessels, and nerves to the organs. Five are named in this course. For each one, learn its two ends.

Fold	What it connects	Other features
Greater omentum	Hangs from the greater curvature of the stomach and the first part of the duodenum, drapes down in front of the small intestine, then folds back up and attaches to the transverse colon	The largest fold. Folds on itself into four layers. A fatty apron that grows with weight gain and holds many lymph nodes that fight infections of the tract
Falciform ligament	Attaches the liver to the anterior abdominal wall and the diaphragm	The liver is the only organ attached to the anterior wall. Its free lower edge holds the ligamentum teres
Lesser omentum	From the lesser curvature of the stomach and the first part of the duodenum up to the liver	Its free edge carries the hepatic portal vein, the hepatic artery, and the common bile duct, with some lymph nodes
Mesentery	A fan from the posterior abdominal wall to the jejunum and ileum	Double-layered, full of fat, with the vessels, lymphatics, and lymph nodes of the small intestine between its layers
Mesocolon	Two separate folds from the posterior wall: the transverse mesocolon to the transverse colon and the sigmoid mesocolon to the sigmoid colon	Carries the vessels to those parts of the colon. With the mesentery, it holds the intestines loosely so they can move as they mix and push food

- Greater goes with the greater curvature and hangs low. Lesser goes with the lesser curvature and runs up to the liver. Meso means middle: the mesentery and mesocolon hold the intestines out from the back wall like a hammock.
- The falciform ligament reaches the front wall; the mesentery and mesocolon reach the back wall.
- On the cadaver, the greater omentum is the first thing you see when the abdomen is opened. Lift it toward the head to find the transverse colon on its back surface and the small intestine below.

What this means for patients

- Peritonitis is acute inflammation of the peritoneum. Microbes get in through a surgical or accidental wound, a ruptured appendix, or a perforated intestine, and the infection can be life-threatening.
- Inflamed peritoneal surfaces rubbing together is also peritonitis, painful but not usually dangerous. People on peritoneal dialysis are at risk.

Competency 112. Retroperitoneal organs

In your course notes: Module 4 packet, Chapter 3, competency 112 (page 28).

Dr. Rennie's video for this part: Digestive system concept video, chapter Peritoneum and abdominal folds.

You should be able to: Distinguish retroperitoneal organs from intraperitoneal ones and name the kidneys, pancreas, duodenum, and parts of the colon as retroperitoneal.

Intraperitoneal organ

Wrapped almost completely in visceral peritoneum and hanging into the cavity on a fold, so it can move: the stomach, jejunum, ileum, transverse colon, sigmoid colon, liver, and spleen.

Retroperitoneal organ

Lies on the posterior abdominal wall in the retroperitoneum, the region behind the parietal peritoneum, with peritoneum on its front surface only. It is not in the peritoneal cavity, has no mesentery, and is held firmly in place. Its back surface is covered by an adventitia.

Organ or part	Position	How you can tell
Stomach	Intraperitoneal	Hangs between the lesser and greater omenta
Duodenum	Retroperitoneal (except its first part)	Fixed to the back wall in a C around the head of the pancreas
Jejunum and ileum	Intraperitoneal	Hang on the mesentery
Ascending colon	Retroperitoneal	Fixed to the right posterior wall
Transverse colon	Intraperitoneal	Hangs on the transverse mesocolon
Descending colon	Retroperitoneal	Fixed to the left posterior wall
Sigmoid colon	Intraperitoneal	Hangs on the sigmoid mesocolon
Pancreas	Retroperitoneal (except the tip of the tail)	Lies across the back wall behind the stomach
Kidneys	Retroperitoneal	Against the back wall on either side of the spine

- The colon alternates: ascending fixed, transverse hanging, descending fixed, sigmoid hanging. The two parts with a mesocolon are the two intraperitoneal parts.
- SADPUCKER for the long list: Suprarenal (adrenal) glands, Aorta and inferior vena cava, Duodenum (except the first part), Pancreas (except the tail), Ureters, Colon (ascending and descending), Kidneys, Esophagus (the abdominal part), Rectum.

Competency 113. Mouth, pharynx, esophagus

In your course notes: Module 4 packet, Chapter 3, competency 113 (page 29).

Dr. Rennie's video for this part: Digestive system concept video, chapter Mouth and pharynx anatomy.

You should be able to: Identify the oral cavity boundaries, hard and soft palate, uvula, fauces, tonsils, the three pharyngeal regions, and the esophagus with its hiatus.

The mouth (oral cavity, buccal cavity)

Boundary	What forms it
Anterior	The lips (labia): fleshy folds of skin outside and nonkeratinized stratified squamous mucosa inside, with the orbicularis oris between. The labial frenulum, a midline fold, ties each lip to its gum
Lateral	The cheeks: skin outside, mucosa inside, the buccinator muscle between. Lips and cheeks keep food between the teeth and help with speech
Superior (roof)	The hard palate in front and the soft palate behind
Inferior (floor)	The tongue and the muscles under it (mylohyoid)
Posterior	The fauces, the opening into the oropharynx

Oral vestibule
: The space between the lips and cheeks on the outside and the gums and teeth on the inside.

Oral cavity proper
: The space from the gums and teeth back to the fauces.

Hard palate
: The bony front of the roof, formed by the palatine processes of the maxillae and the horizontal plates of the palatine bones under mucosa. It walls the mouth off from the nasal cavity, so you can chew and breathe at the same time, and gives the tongue a firm surface to press food against.

Soft palate
: The muscular, arch-shaped back of the roof, between the oropharynx and nasopharynx. Its back border opens into the oropharynx through the fauces.

Uvula
: The finger-shaped muscular tag hanging from the back edge of the soft palate. During a swallow the soft palate and uvula lift and seal the nasopharynx, so food and liquid do not enter the nose.

Palatoglossal arch
: The front arch, running from the soft palate down to the base of the tongue.

Palatopharyngeal arch
: The back arch, running from the soft palate to the side of the pharynx.

Palatine tonsils
: Between the two arches on each side. These are the tonsils you see in an open mouth.

Lingual tonsil
: On the back (root) of the tongue.

The pharynx

A funnel-shaped tube of skeletal muscle lined with mucous membrane, from the choanae to the esophagus behind and the larynx in front. Three regions: the nasopharynx (air only), the oropharynx, and the laryngopharynx (air and food). The pharyngeal tonsil sits in the nasopharynx. A swallowed bolus goes mouth, oropharynx, laryngopharynx, esophagus, stomach, pushed by the pharyngeal constrictors. Breathing stops for the moment the bolus passes.

The esophagus

- A collapsible muscular tube about 25 cm (10 inches) long. It starts at the lower end of the laryngopharynx at the level of the cricoid cartilage, runs through the mediastinum behind the trachea and in front of the vertebral column, pierces the diaphragm, and joins the stomach at the cardia.
- The esophageal hiatus is the opening in the diaphragm it passes through, at about T10.
- It secretes mucus and carries food to the stomach. No digestion and no absorption happen here.

Layer	In the esophagus
Mucosa	Nonkeratinized stratified squamous epithelium (wear and tear), lamina propria, muscularis mucosae. Mucous glands near the stomach
Submucosa	Areolar connective tissue, vessels, mucous glands
Muscularis	Upper third skeletal muscle, middle third mixed, lower third smooth muscle. Two sphincters: the upper esophageal sphincter (skeletal) and the lower esophageal (cardiac) sphincter (smooth), near the heart
Adventitia	Areolar connective tissue with no mesothelium, blending into the mediastinum to anchor the esophagus. Only the short abdominal segment has a serosa

What this means for patients

- Hiatal hernia: part of the stomach pushes up through the esophageal hiatus into the thorax. It weakens the support the diaphragm gives the lower esophageal sphincter and often comes with reflux (competency 119).

Competency 114. Major canal sphincters

In your course notes: Module 4 packet, Chapter 3, competency 114 (page 31).

You should be able to: Locate the upper and lower esophageal, pyloric, ileocecal, and internal and external anal sphincters and state which regions each one separates.

A sphincter is a ring of muscle around the tube that can squeeze it shut. Sphincters sit at borders between regions. They control when contents move on and most keep contents from flowing backward. Learn each one by the two regions on either side of it.

Sphincter	Where	Separates	Muscle
Upper esophageal sphincter	Top of the esophagus, at the cricoid cartilage	Laryngopharynx from esophagus	Skeletal; stays closed between swallows so air does not enter the esophagus
Lower esophageal sphincter (also called the cardiac sphincter or gastroesophageal sphincter)	Bottom of the esophagus, just under the diaphragm	Esophagus from stomach (cardia)	Smooth; the diaphragm around the hiatus helps hold it shut, keeping acid out of the esophagus
Pyloric sphincter	Around the pylorus at the stomach exit	Stomach from duodenum	Smooth, a thickening of the circular layer; meters small amounts of chyme into the duodenum
Ileocecal sphincter	End of the ileum where it opens into the cecum	Small intestine (ileum) from large intestine (cecum)	Smooth; with folds of mucosa it forms the ileocecal valve, which keeps colon contents out of the ileum
Internal anal sphincter	Around the anal canal, inside the external sphincter	Anal canal from the outside	Smooth, involuntary
External anal sphincter	Around the anal canal, outside the internal sphincter	Anal canal from the outside	Skeletal, voluntary

- Only the two ends of the canal have skeletal sphincters: the upper esophageal at the top and the external anal at the bottom. Everything in between is smooth muscle.
- The sphincter of Oddi (hepatopancreatic sphincter) is on your lab list with the small intestine, but it is not a sphincter of the canal. It controls bile and pancreatic juice entering the duodenum at the major duodenal papilla (competencies 123 and 124).

Competency 121. Salivary glands

In your course notes: Module 4 packet, Chapter 4, competency 121 (page 40).

Dr. Rennie's video for this part: Digestive system concept video, chapter Mouth and pharynx anatomy.

You should be able to: Locate the parotid, submandibular, and sublingual glands and state where the duct of each opens.

Saliva keeps the mouth and pharynx moist, cleans the mouth and teeth, and during a meal lubricates and dissolves food and starts the chemical breakdown of starch. Small glands in the lining (labial, buccal, palatal, and lingual glands) keep the mouth wet all the time. Most saliva comes from three pairs of major glands outside the lining, each sending its saliva in through a duct.

Gland	Location	Where the duct opens	Saliva
Parotid gland	In front of and below the ear, between the skin and the masseter muscle; the largest	The parotid duct crosses the masseter, pierces the buccinator, and opens into the vestibule opposite the upper second molar	Watery (serous), with salivary amylase
Submandibular gland	Floor of the mouth, along the inner surface of the mandible	The submandibular duct runs forward under the mucosa of the floor and opens just lateral to the lingual frenulum	Mostly serous with some mucus, a little thicker
Sublingual gland	Under the tongue in the floor of the mouth, above the submandibular gland; the smallest	Many small sublingual ducts opening along the floor of the mouth	Mostly mucus, thick, very little amylase

- Each name tells you the spot. Parotid, beside the ear. Submandibular, below the mandible. Sublingual, below the tongue.
- Only the parotid duct opens in the cheek. The other two open under the tongue.
- Saliva is about 99.5 percent water. The rest is ions (sodium, potassium, chloride, bicarbonate, phosphate), dissolved gases, and organic substances: urea, uric acid, mucus, IgA, lysozyme, and salivary amylase, which splits starch into maltose, maltotriose, and alpha-dextrins.

What this means for patients

- Mumps is a paramyxovirus infection that targets the parotid glands. The glands swell on one or both sides in front of the ramus of the mandible, with fever, malaise, and throat pain on swallowing or with sour foods. In about 30 percent of males past puberty the testes become inflamed; sterility is rare unless both sides are involved. A vaccine prevents it.

Competency 120. Teeth and tongue

In your course notes: Module 4 packet, Chapter 4, competency 120 (page 38).

Dr. Rennie's video for this part: Digestive system concept video, chapter Mouth and pharynx anatomy.

You should be able to: Identify the teeth and tongue with the lingual frenulum, papillae, intrinsic and extrinsic muscles, and lingual glands.

The tongue

The tongue is skeletal muscle covered with mucous membrane. With its muscles it forms the floor of the oral cavity. A median septum divides it into right and left halves along its whole length, and it attaches below to the hyoid bone, the styloid process of the temporal bone, and the mandible.

Muscle group	Origin and insertion	Action
Extrinsic (genioglossus, hyoglossus, styloglossus, palatoglossus)	Start on bone or the soft palate outside the tongue, end in the connective tissue of the tongue	Move the whole tongue: side to side, in and out, up and down, to position food for chewing, shape the bolus, and push it back to swallow. They also hold the tongue in place
Intrinsic (superior and inferior longitudinal, transverse, vertical)	Start and end inside the tongue	Change the shape and size of the tongue for speech and swallowing: curl, flatten, narrow, shorten

- The hypoglossal nerve (cranial nerve XII) supplies all the tongue muscles except palatoglossus, which the vagus supplies.
- Extrinsic starts outside the tongue and moves it. Intrinsic sits within it and changes its shape.

Lingual frenulum

The midline fold of mucosa under the tongue that ties it to the floor of the mouth and limits how far back it can move. A short or rigid frenulum is ankyloglossia (tongue-tie), which impairs speech and can be corrected surgically.

Lingual papillae

Projections of the lamina propria covered with stratified squamous epithelium on the top and sides of the tongue. Some carry taste buds; all give the tongue grip on food.

Lingual glands

Small glands in the lamina propria of the tongue. They release mucus and a watery fluid with lingual lipase, which starts the breakdown of dietary triglycerides into fatty acids and diglycerides.

Papillae	Where and what they look like	Taste buds
Filiform	Thin, pointed, the most numerous, over most of the top surface	No. They give the tongue friction
Fungiform	Mushroom-shaped, scattered among the filiform, most on the tip and sides	Yes
Vallate (circumvallate)	Large, round, about 8 to 12 in a V-shaped row near the back, each in a trench	Yes
Foliate	Leaf-like folds on the sides toward the back	Yes

The teeth

- Each tooth sits in a socket (an alveolus) in the maxilla or mandible, held by a periodontal ligament. The joint is a gomphosis, a fibrous joint.
- Children have 20 deciduous teeth. Adults have 32 permanent teeth: 8 incisors (cut), 4 canines (tear), 8 premolars (crush), and 12 molars (grind), including the four third molars (wisdom teeth).

Crown, neck, root

The crown is above the gum (gingiva), the neck is at the gum line, and the root sits in the socket.

Enamel

Covers the crown. The hardest substance in the body.

Dentin

Forms most of the tooth, under the enamel.

Pulp cavity and root canal

The center of the tooth, holding the pulp (vessels, nerves, connective tissue). It continues down each root as the root canal.

Cementum

Covers the root and anchors it to the periodontal ligament.

Worked questions

Nine questions in the style of the lecture exam. Read the answer and the reasons each wrong option is wrong. The wrong options are where the learning is.

QUESTION 1 · COMPETENCY 109

Which layer of the gut wall contains the large blood vessels, lymphatic vessels, and the submucosal plexus?

A. Mucosa

B. Submucosa

C. Muscularis

D. Serosa

Answer: B. The submucosa is the connective tissue layer between the mucosa and the muscularis. It carries the larger vessels that receive absorbed nutrients and holds the submucosal plexus.

Why the others are wrong

- **A** The mucosa has small capillaries and lymphatic tissue in its lamina propria, but the large vessels and the plexus are one layer out.
- **C** The muscularis is the two smooth muscle layers. The plexus inside it is the myenteric plexus, not the submucosal.
- **D** The serosa is the thin outer serous membrane, the visceral peritoneum. It is not where the vessels run.

QUESTION 2 · COMPETENCY 109

Which region of the canal is lined with simple columnar epithelium?

A. Esophagus

B. Oropharynx

C. Stomach

D. Anal canal

Answer: C. The stomach and both intestines secrete and absorb, so they are lined with simple columnar epithelium.

Why the others are wrong

- **A** The esophagus carries rough food past, so its lining is stratified squamous for protection.
- **B** The oropharynx carries food and air and is stratified squamous.
- **D** The anal canal takes friction at the exit of the tube and is stratified squamous near the anus.

QUESTION 3 · COMPETENCY 110

The myenteric plexus is found

A. in the lamina propria of the mucosa

B. in the submucosa, just under the mucosa

C. between the inner circular and outer longitudinal layers of the muscularis

D. between the muscularis and the serosa

Answer: C. Myenteric means in the muscle. It sits between the two smooth muscle layers it controls, and it runs motility.

Why the others are wrong

- **A** The lamina propria holds vessels and MALT, not a plexus.
- **B** That is the submucosal plexus, which runs secretion and blood flow to the lining.
- **D** Nothing named in this course sits between the muscularis and the serosa. Both plexuses are inside the wall.

QUESTION 4 · COMPETENCY 111

Which peritoneal fold attaches the liver to the anterior abdominal wall and the diaphragm?

A. Lesser omentum

B. Greater omentum

C. Falciform ligament

D. Mesentery

Answer: C. The falciform ligament is the sickle-shaped fold from the front of the liver to the anterior wall and diaphragm. The liver is the only organ attached to the anterior wall.

Why the others are wrong

- **A** The lesser omentum connects the stomach and the first part of the duodenum to the liver. It does not reach the body wall.
- **B** The greater omentum hangs from the greater curvature of the stomach and attaches to the transverse colon.
- **D** The mesentery holds the jejunum and ileum to the posterior wall.

QUESTION 5 · COMPETENCY 112

Which of these is retroperitoneal?

A. Transverse colon

B. Jejunum

C. Stomach

D. Descending colon

Answer: D. The descending colon is fixed to the left posterior wall behind the parietal peritoneum. Ascending and descending colon are the two retroperitoneal parts.

Why the others are wrong

- **A** The transverse colon hangs on the transverse mesocolon, so it is intraperitoneal.
- **B** The jejunum hangs on the mesentery, wrapped in visceral peritoneum.
- **C** The stomach hangs between the lesser and greater omenta inside the peritoneal cavity.

QUESTION 6 · COMPETENCY 113

During a swallow, which structure lifts to seal off the nasopharynx?

A. The epiglottis

B. The soft palate and uvula

C. The hard palate

D. The palatine tonsils

Answer: B. The soft palate is muscle, so it can move. It and the uvula rise and press against the back wall of the pharynx, closing the nasopharynx so food does not enter the nose.

Why the others are wrong

- **A** The epiglottis tips down over the laryngeal inlet to keep food out of the airway below, not the nose above.
- **C** The hard palate is bone and cannot move.
- **D** The palatine tonsils are lymphatic tissue between the arches. They guard the throat; they do not seal anything.

QUESTION 7 · COMPETENCY 114

Which sphincter is skeletal muscle under voluntary control?

A. Lower esophageal sphincter

B. Pyloric sphincter

C. Internal anal sphincter

D. External anal sphincter

Answer: D. The external anal sphincter is skeletal muscle, the one you control. Only the two ends of the canal, the upper esophageal and the external anal sphincters, are skeletal.

Why the others are wrong

- **A** The lower esophageal sphincter is smooth muscle helped by the diaphragm.
- **B** The pyloric sphincter is a thickening of the circular smooth muscle layer.
- **C** The internal anal sphincter is smooth muscle and involuntary. It sits inside the external one.

QUESTION 8 · COMPETENCY 121

The parotid duct opens

A. beside the lingual frenulum

B. along the floor of the mouth under the tongue

C. inside the cheek opposite the upper second molar

D. at the base of the tongue

Answer: C. The parotid duct runs forward across the masseter, pierces the buccinator, and opens into the vestibule opposite the upper second molar.

Why the others are wrong

- **A** That is the submandibular duct.
- **B** That is where the many small sublingual ducts open.
- **D** The base of the tongue holds the lingual tonsil, not a salivary duct opening.

QUESTION 9 · COMPETENCY 120

Which papillae have no taste buds?

A. Fungiform

B. Vallate

C. Foliate

D. Filiform

Answer: D. Filiform papillae are the thin, pointed, most numerous papillae. They give the tongue friction and carry no taste buds.

Why the others are wrong

- **A** Fungiform papillae on the tip and sides carry taste buds.
- **B** Vallate papillae, the big ones in the V at the back, carry taste buds.
- **C** Foliate papillae on the sides carry taste buds.

BIO 004 Human Anatomy · Module 4 notes · Dr. Sharilyn Rennie

The digestive system, part II: stomach, pancreas, liver, and gallbladder

The written notes for the second digestive lecture: the stomach, the pancreas, the liver and its lobule, the gallbladder and the path of bile, and the accessory organ disorders. Module 4 competencies 115 and 122 to 125, with nine worked questions.

Competency 115. Stomach regions and features

In your course notes: Module 4 packet, Chapter 3, competency 115 (page 32).

Dr. Rennie's video for this part: Digestive system concept video, chapter Stomach, pancreas, liver, gallbladder.

You should be able to: Identify the cardia, fundus, body, pyloric part, curvatures, rugae, and the three-layered muscularis of the stomach.

The stomach is a J-shaped widening of the canal just under the diaphragm, connecting the esophagus above to the duodenum below. It is a mixing chamber and a holding tank: it stores the meal, churns it with gastric juice until the semisolid bolus is a liquid called chyme, begins the digestion of starch, triglycerides, and proteins, absorbs a few substances, and releases chyme into the duodenum a little at a time. Empty, it is about the size of a large sausage. Full, it stretches to hold 2 to 3 liters.

The four regions

Region	Where it is
Cardia (cardiac region)	The small region around the opening from the esophagus
Fundus	The rounded dome that bulges above and to the left of the cardia, up under the diaphragm
Body	The large central part below the fundus
Pyloric part (pyloric region)	The funnel that connects the body to the duodenum, in three parts in order: the pyloric antrum (wide, joins the body), the pyloric canal (narrow), and the pylorus (the outlet, surrounded by the pyloric sphincter)

Lesser curvature
: The short, concave border on the right (medial) side, facing the liver. The lesser omentum attaches here and runs up to the liver.

Greater curvature
: The long, convex border on the left (lateral) side. The greater omentum hangs from here and drapes over the intestines.

Rugae
: Large folds of mucosa and submucosa on the inside of an empty stomach. They flatten as the stomach fills, which is how it stretches.

Pyloric sphincter
: The ring of smooth muscle around the pylorus that meters chyme into the duodenum. A thickening of the circular layer.

- Draw the fundus up beside the esophagus, not at the bottom of the J. Food collects there, but it does not have to pass through it to move on.
- Our notes call the curvatures medial and lateral (OpenStax); your slides call them superior and inferior. Both describe the same two borders.

The wall of the stomach

Layer	In the stomach
Mucosa	Simple columnar epithelium of surface mucous cells. The surface is dotted with gastric pits, narrow channels that gastric glands open into. Lamina propria of areolar connective tissue; muscularis mucosae of smooth muscle
Submucosa	Areolar connective tissue
Muscularis	Three layers of smooth muscle instead of two: inner oblique (the extra layer, mostly in the body), middle circular (thickens into the pyloric sphincter), outer longitudinal. The oblique fibers let the stomach twist and squeeze as well as push
Serosa	Visceral peritoneum. At the lesser curvature it continues up to the liver as the lesser omentum; at the greater curvature it continues down as the greater omentum

The cells of the gastric glands

Cell	Secretes
Mucous neck cells	Mucus
Chief cells	Pepsinogen (the inactive form of the protein-digesting enzyme pepsin) and gastric lipase
Parietal cells	Hydrochloric acid, and intrinsic factor, which is needed to absorb vitamin B12
G cells (enteroendocrine), mostly in the pyloric antrum	Gastrin, a hormone that stimulates gastric activity

Together the exocrine secretions are gastric juice.

What this means for patients

- Pylorospasm: in some infants the pyloric sphincter fails to relax, food cannot leave the stomach, the stomach overfills, and the baby vomits to relieve the pressure. Treated with drugs that relax the sphincter.
- Pyloric stenosis: a narrowing of the pyloric sphincter in infants. The hallmark is projectile vomiting. It needs surgical correction.
- Vomiting (emesis) is the forceful expulsion of the contents of the stomach, and sometimes the duodenum, through the mouth. Irritation or stretch of the stomach, unpleasant sights, general anesthesia, dizziness, and drugs such as morphine and digitalis all signal the vomiting center in the medulla (the area postrema). The diaphragm and abdominal muscles squeeze the stomach and the contents come up through relaxed esophageal sphincters. Prolonged vomiting in infants and the elderly is dangerous: loss of acidic gastric juice shifts blood pH toward alkalosis, the person dehydrates, and acid damages the esophagus and teeth.

Competency 124. Pancreas structure

In your course notes: Module 4 packet, Chapter 4, competency 124 (page 44).

Dr. Rennie's video for this part: Digestive system concept video, chapter Stomach, pancreas, liver, gallbladder.

You should be able to: Identify the head, body, and tail of the pancreas with its main and accessory ducts, major duodenal papilla, sphincter of Oddi, acini, and islets.

Most chemical digestion happens in the small intestine, and the pancreas, liver, and gallbladder supply the tools. The pancreas is a long, soft gland lying across the posterior abdominal wall behind the greater curvature of the stomach. It is retroperitoneal. It is an exocrine gland (pancreatic juice down ducts into the duodenum) and an endocrine gland (hormones into the blood).

Structure	What it is
Head	The wide end, in the C-shaped curve of the duodenum, right of the midline
Body	The middle part, crossing the midline behind the stomach, in front of the aorta and the vertebral column
Tail	The narrow left end, near the spleen
Main pancreatic duct (duct of Wirsung)	The larger duct. Runs the whole length of the gland from tail to head, collecting pancreatic juice. In the wall of the duodenum it joins the common bile duct at the hepatopancreatic ampulla (ampulla of Vater)
Accessory pancreatic duct (duct of Santorini)	The smaller duct. Drains part of the head and opens on its own into the duodenum at the minor duodenal papilla, a little above the major papilla
Hepatopancreatic ampulla	The short widened chamber where the common bile duct and main pancreatic duct meet
Major duodenal papilla	The bump in the duodenal lining where the ampulla opens, in the descending part of the duodenum below the pyloric sphincter
Sphincter of Oddi (hepatopancreatic sphincter)	The ring of smooth muscle around the ampulla that controls bile and pancreatic juice flowing into the duodenum
Acini	Small clusters of gland cells around tiny ducts: the exocrine part, about 99 percent of the gland. They make pancreatic juice, 1200 to 1500 mL a day of clear fluid: water, salts, sodium bicarbonate, and digestive enzymes
Pancreatic islets (islets of Langerhans)	Pale round patches scattered among the acini: the endocrine part, about 1 percent. They release glucagon, insulin, somatostatin, and pancreatic polypeptide into the blood

- Match the size words. The main duct opens at the major papilla. The accessory duct opens at the minor papilla.
- Only the main duct joins the common bile duct. The accessory duct empties on its own.

Competency 122. Liver gross and lobule anatomy

In your course notes: Module 4 packet, Chapter 4, competency 122 (page 40).

Dr. Rennie's video for this part: Digestive system concept video, chapter Stomach, pancreas, liver, gallbladder.

You should be able to: Identify the four liver lobes, falciform ligament, and ligamentum teres, and name the hepatocytes, sinusoids, canaliculi, and portal triad of the lobule.

The liver is the largest gland in the body. It sits under the diaphragm, filling most of the right hypochondriac region and part of the epigastric region, protected by the lower ribs. Visceral peritoneum covers most of it, and under that a capsule of dense irregular connective tissue covers all of it.

Lobes and ligaments

Structure	Where and what
Right lobe	The largest lobe, right of the falciform ligament
Left lobe	Left of the falciform ligament, reaching over the stomach
Quadrate lobe	A small square lobe on the inferior surface, between the gallbladder and the ligamentum teres
Caudate lobe	A small lobe on the posterior surface beside the inferior vena cava
Porta hepatis	The doorway between the quadrate and caudate lobes. The hepatic portal vein and hepatic artery proper enter here; the right and left hepatic ducts leave here
Falciform ligament	The sickle-shaped fold of peritoneum that suspends the liver from the diaphragm and the anterior abdominal wall and marks the line between the right and left lobes on the front
Ligamentum teres (round ligament)	The fibrous cord in the free edge of the falciform ligament, running from the umbilicus to the liver. It is what remains of the fetal umbilical vein
Coronary ligaments	Narrow extensions of parietal peritoneum that hang the liver from the diaphragm
Lesser omentum	The fold from the lesser curvature of the stomach and the start of the duodenum to the visceral surface of the liver. Its free edge carries the portal vein, hepatic artery proper, and common bile duct

- From the front you see only two lobes. The quadrate and caudate lobes are underneath and behind.
- The ligamentum teres comes from the umbilical vein, which carried oxygen-rich blood from the placenta. Vein, not artery.

The liver lobule

Liver tissue is thousands of small six-sided units. Hepatocytes are the working cells: specialized epithelial cells with 5 to 12 sides making up about 80 percent of the liver's volume. They are arranged in plates one cell thick, the hepatic laminae, that radiate out from a central vein like spokes. Between the plates run the sinusoids.

Structure	Where it is	What it does
Central vein	The center of the lobule	Collects blood from the sinusoids
Hepatocytes	In plates (hepatic laminae) radiating from the central vein	The working cells: they make bile and do the liver's metabolic work
Hepatic sinusoids	Between the plates	Wide, leaky, endothelium-lined capillaries carrying mixed blood from the edge of the lobule toward the central vein. Stellate macrophages (Kupffer cells) line them and destroy worn-out blood cells, bacteria, and foreign matter arriving from the gut
Bile canaliculi	Grooves between neighboring hepatocytes inside each plate	Collect the bile the hepatocytes make and carry it outward to the edge of the lobule
Portal triad	At each corner of the lobule	A branch of the hepatic portal vein, a branch of the hepatic artery, and a bile duct. The two vessels bring blood in; the duct carries bile out

- Blood flows in toward the central vein. Bile flows out toward the portal triad. Opposite directions.
- Bile is a yellow-brown, sometimes olive-green liquid. It is both an excretory product and a digestive secretion.

Blood through the liver, in order

1. The hepatic artery proper (oxygen-rich blood from the aorta) and the hepatic portal vein (oxygen-poor blood carrying newly absorbed nutrients, drugs, microbes, and toxins from the gut and spleen) enter at the porta hepatis.

2. Branches of both run to the portal triads at the corners of every lobule.
3. The two streams mix in the sinusoids and flow past the hepatocytes toward the center.
4. The central vein collects the blood.
5. Central veins join into hepatic veins, which leave the back of the liver.
6. The hepatic veins empty into the inferior vena cava just under the diaphragm, and the blood returns to the right atrium.

Because so much blood from the gut passes through the liver, it is a very common site for cancers of the digestive tract to spread.

What the liver does

- Makes bile and the bile salts in it, which emulsify fats in the small intestine so they can be absorbed.
- Carbohydrate metabolism: keeps blood glucose steady and stores extra glucose as glycogen.
- Lipid metabolism and protein metabolism, including removing the amino group from amino acids (deamination) and sending the ammonia out as urea in the urine.
- Processes drugs and hormones, including alcohol and antibiotics.
- Excretes bilirubin, the pigment left over from breaking down old red blood cells.
- Stores glycogen and vitamins A, B12, D, E, and K.
- Phagocytosis by the Kupffer cells, and the final activation of vitamin D.

Competency 123. Gallbladder and bile pathway

In your course notes: Module 4 packet, Chapter 4, competency 123 (page 42).

Dr. Rennie's video for this part: Digestive system concept video, chapter Stomach, pancreas, liver, gallbladder.

You should be able to: Identify the gallbladder regions and trace bile from the canaliculi through the hepatic, cystic, and common bile ducts to the hepatopancreatic ampulla and duodenum.

The gallbladder is a pear-shaped muscular sac about 8 to 10 cm long, sitting in a shallow hollow on the visceral surface of the right lobe of the liver and hanging a little past its lower edge. It stores bile between meals and concentrates it by removing water. When fatty food reaches the duodenum, the smooth muscle in its wall contracts and squeezes bile out.

Structure	What it is
Fundus	The wide, rounded end that sticks out past the lower border of the liver
Body	The main middle part
Neck	The narrow end that points toward the porta hepatis and continues as the cystic duct
Right and left hepatic ducts	Drain bile from the two sides of the liver and leave at the porta hepatis
Common hepatic duct	Formed where the right and left hepatic ducts join
Cystic duct	Runs from the neck of the gallbladder and joins the common hepatic duct. Carries bile in both directions
Common bile duct	Formed where the common hepatic duct and the cystic duct join. Runs down in the edge of the lesser omentum, behind the first part of the duodenum, and through the head of the pancreas
Hepatopancreatic ampulla, sphincter of Oddi, major duodenal papilla	Where the common bile duct meets the main pancreatic duct and empties into the descending duodenum (competency 124)

A drop of bile, from the liver to the duodenum

1. Hepatocytes make the bile.
2. Bile canaliculi between the hepatocytes collect it and carry it to the edge of the lobule.

3. Bile ductules and the bile ducts of the portal triads carry it out of the lobules and join into larger ducts.
4. Right and left hepatic ducts leave the liver at the porta hepatis.
5. They join to form the common hepatic duct.
6. Side trip: when the sphincter of Oddi is closed, bile backs up and turns up the cystic duct into the gallbladder for storage. When the gallbladder contracts, bile comes back down the same cystic duct.
7. Common hepatic duct plus cystic duct form the common bile duct.
8. At the hepatopancreatic ampulla the common bile duct joins the main pancreatic duct.
9. The sphincter of Oddi opens and bile enters the duodenum through the major duodenal papilla, where it mixes with food.

- The duct names tell you where each duct comes from. Hepatic means liver. Cystic means bladder, this bladder. The common bile duct is where both come together.
- The cystic duct joins the common hepatic duct. It does not run straight to the duodenum. The gallbladder is a side trip, not a step every drop takes.

Competency 125. Clinical anatomy of accessory disorders

In your course notes: Module 4 packet, Chapter 4, competency 125 (page 45).

You should be able to: Match gallstones, cholecystitis, hepatitis, cirrhosis, jaundice, pancreatitis, and mumps to the accessory organ each one affects and explain how a stone at the ampulla can block bile and pancreatic flow at once.

Disorder	Organ affected	What changes
Gallstones	Gallbladder and bile ducts	Hard deposits form from bile in the gallbladder. A stone can move into the cystic duct or the common bile duct and block it
Cholecystitis	Gallbladder	The wall is inflamed and swollen, most often because a stone blocks the cystic duct and traps bile inside
Hepatitis	Liver	The liver is inflamed and hepatocytes are damaged, most often by a virus, also by alcohol or some drugs
Cirrhosis	Liver	After long-term damage, hepatocytes are replaced by bands of scar tissue and lumps of regrowing cells. The lobules lose their pattern and blood has trouble getting through
Jaundice	Liver or bile ducts, seen in the skin and eyes	Bilirubin builds up in the blood and stains the skin and the whites of the eyes yellow, because the liver cannot clear it into bile, a duct is blocked, or red cells break down too fast
Pancreatitis	Pancreas	The pancreas is inflamed and swollen. Digestive enzymes become active inside the gland and digest its own tissue. Gallstones and alcohol are common causes
Mumps	Parotid glands	A viral infection swells the parotids, so the face puffs out in front of and below the ears

Shared ducts, shared problems

- The common bile duct and the main pancreatic duct meet at the hepatopancreatic ampulla and share one exit into the duodenum. A gallstone that travels down the common bile duct and lodges in the ampulla plugs that shared exit.
- Bile backs up into the liver and the blood: jaundice.
- Pancreatic juice backs up into the pancreas, where its enzymes damage the gland: pancreatitis.
- One stone, two organs. That is one way gallstones lead to pancreatitis.
- Mumps swells the parotid gland, not the pancreas. Cholecystitis is the gallbladder; hepatitis and cirrhosis are the liver.

Worked questions

Nine questions in the style of the lecture exam. Read the answer and the reasons each wrong option is wrong. The wrong options are where the learning is.

QUESTION 1 · COMPETENCY 115

The dome-shaped region of the stomach that bulges above and to the left of the esophageal opening is the

A. cardia

B. fundus

C. body

D. pyloric antrum

Answer: B. The fundus is the rounded dome up under the diaphragm, above and left of the cardia. Food can collect there without passing through it.

Why the others are wrong

- **A** The cardia is the small region right around the opening from the esophagus.
- **C** The body is the large central part below the fundus.
- **D** The pyloric antrum is the wide first part of the funnel at the exit of the stomach.

QUESTION 2 · COMPETENCY 115

Which muscle layer is found in the stomach but not in the rest of the canal?

A. Inner oblique

B. Middle circular

C. Outer longitudinal

D. Muscularis mucosae

Answer: A. The stomach has three layers of smooth muscle. The inner oblique layer is the extra one; its diagonal fibers let the stomach twist and squeeze food.

Why the others are wrong

- **B** The circular layer is in the whole tube. In the stomach it is the middle layer and thickens at the pylorus.
- **C** The longitudinal layer is in the whole tube as the outer layer.
- **D** The muscularis mucosae is the thin muscle layer inside the mucosa of the whole tube, not a layer of the muscularis.

QUESTION 3 · COMPETENCY 115

Which gastric gland cell secretes hydrochloric acid and intrinsic factor?

A. Chief cell

B. Mucous neck cell

C. Parietal cell

D. G cell

Answer: C. Parietal cells make hydrochloric acid and intrinsic factor, the protein needed to absorb vitamin B12.

Why the others are wrong

- **A** Chief cells secrete pepsinogen and gastric lipase.
- **B** Mucous neck cells secrete mucus.
- **D** G cells are enteroendocrine cells that release the hormone gastrin into the blood.

QUESTION 4 · COMPETENCY 124

Which pancreatic duct joins the common bile duct at the hepatopancreatic ampulla?

A. The accessory duct (duct of Santorini)

B. The main pancreatic duct (duct of Wirsung)

C. Both ducts

D. Neither; pancreatic juice enters at the minor papilla only

Answer: B. The main pancreatic duct runs the whole length of the gland and joins the common bile duct at the ampulla, which opens at the major duodenal papilla.

Why the others are wrong

- **A** The accessory duct drains part of the head and opens on its own at the minor duodenal papilla.
- **C** Only the main duct joins the bile duct. The accessory duct empties separately.
- **D** Most pancreatic juice enters through the ampulla at the major papilla. Only the accessory duct uses the minor papilla.

QUESTION 5 · COMPETENCY 124

On a slide of the pancreas, the pale round patches among the darker clusters are

A. acini, the exocrine part

B. pancreatic islets, the endocrine part

C. bile canaliculi

D. hepatic sinusoids

Answer: B. The islets of Langerhans are the pale round patches, about 1 percent of the gland, releasing hormones such as insulin and glucagon into the blood.

Why the others are wrong

- **A** The acini are the darker clusters around them, about 99 percent of the gland, making pancreatic juice.
- **C** Bile canaliculi are in the liver, between hepatocytes.
- **D** Sinusoids are the leaky capillaries of the liver lobule, not pancreatic tissue.

QUESTION 6 · COMPETENCY 122

The ligamentum teres is the remnant of which fetal vessel?

A. Umbilical artery

B. Umbilical vein

C. Ductus arteriosus

D. Ductus venosus

Answer: B. The ligamentum teres (round ligament) runs from the umbilicus to the liver in the free edge of the falciform ligament. Before birth it was the umbilical vein carrying oxygen-rich blood from the placenta.

Why the others are wrong

- **A** The umbilical arteries become the medial umbilical ligaments on the inside of the anterior abdominal wall.
- **C** The ductus arteriosus was the fetal shunt between the pulmonary trunk and the aorta; it becomes the ligamentum arteriosum.
- **D** The ductus venosus was the fetal shunt through the liver; it becomes the ligamentum venosum, not the round ligament.

QUESTION 7 · COMPETENCY 122

In a liver lobule, blood flows

A. from the central vein outward to the portal triads

B. from the portal triads through the sinusoids to the central vein

C. from the bile canaliculi into the sinusoids

D. from the hepatic veins into the portal triads

Answer: B. The two vessels of each portal triad feed the sinusoids, and the blood flows past the hepatocytes toward the central vein in the middle. Bile goes the opposite way, out toward the triads.

Why the others are wrong

- **A** That is the direction of bile, not blood.
- **C** Bile canaliculi carry bile and do not connect to the sinusoids.
- **D** Hepatic veins carry blood out of the liver to the inferior vena cava. They are downstream of the central veins, not upstream of the triads.

QUESTION 8 · COMPETENCY 123

The common bile duct is formed by the union of the

A. right and left hepatic ducts

B. common hepatic duct and the cystic duct

C. cystic duct and the main pancreatic duct

D. common hepatic duct and the accessory pancreatic duct

Answer: B. The right and left hepatic ducts first form the common hepatic duct. The cystic duct from the gallbladder joins it, and from that point the duct is the common bile duct.

Why the others are wrong

- **A** The right and left hepatic ducts form the common hepatic duct, one step earlier.
- **C** The main pancreatic duct joins the common bile duct further down, at the hepatopancreatic ampulla. The cystic duct never meets it directly.
- **D** The accessory pancreatic duct opens on its own at the minor papilla and joins nothing.

QUESTION 9 · COMPETENCY 125

A gallstone lodges in the hepatopancreatic ampulla. Which two problems can follow?

A. Jaundice and pancreatitis

B. Mumps and hepatitis

C. Cholecystitis and cirrhosis

D. Hepatitis and pancreatitis

Answer: A. The ampulla is the shared exit for the common bile duct and the main pancreatic duct. Blocked, bile backs up into the liver and blood (jaundice) and pancreatic enzymes back up into the pancreas (pancreatitis).

Why the others are wrong

- **B** Mumps is a viral infection of the parotid glands; it has nothing to do with a stone. Hepatitis is inflammation of the liver by a virus, alcohol, or drugs.
- **C** Cholecystitis usually follows a stone in the cystic duct, which traps bile in the gallbladder, not a stone at the ampulla. Cirrhosis is the scarring left by long-term liver damage.
- **D** Pancreatitis is right, but hepatitis is inflammation of the liver by a virus or toxin, not a blocked duct. The blocked duct produces jaundice.

BIO 004 Human Anatomy · Module 4 notes · Dr. Sharilyn Rennie

The digestive system, part III: small intestine, large intestine, and rectum

The written notes for the third digestive lecture: the small intestine and its three levels of folding, the large intestine, the rectum and anal canal, the whole path of food, and the canal disorders. Module 4 competencies 116 to 119, with nine worked questions.

Competency 116. Small intestine and surface area

In your course notes: Module 4 packet, Chapter 3, competency 116 (page 33).

Dr. Rennie's video for this part: Digestive system concept video, chapter Small and large intestines.

You should be able to: Identify the duodenum, jejunum, and ileum and name the circular folds, villi, lacteal, and microvilli as features that expand absorptive surface area.

The small intestine runs from the pyloric sphincter to the ileocecal valve. It is the longest part of the canal, about 3 m (10 feet) in a living person, and fills most of the middle and lower abdomen. Most chemical digestion and nearly all absorption of nutrients happen here.

Region	Length	Location and features
Duodenum	About 25 cm (10 inches); the shortest	A C-shaped curve from the pyloric sphincter around the head of the pancreas. Mostly retroperitoneal. Bile and pancreatic juice enter through the major duodenal papilla on its inner wall
Jejunum	About 1 m (3 feet)	The middle region, mostly in the upper left abdomen, hanging on the mesentery. The name means empty, which is how it is found after death
Ileum	About 2 m (6 feet); the longest	The last region, mostly in the lower right abdomen, hanging on the mesentery. Ends at the ileocecal valve. Its wall holds Peyer patches

- There is no sharp border between the jejunum and the ileum; one becomes the other along the mesentery.
- The ileum is the longest part, not the jejunum. Your slides give the duodenum as 12 inches; OpenStax and Tortora give about 25 cm (10 inches). Either is accepted.

The wall of the small intestine

Layer	In the small intestine
Mucosa	Simple columnar epithelium with several cell types: absorptive cells (enzymes on their surface, microvilli for absorption), goblet cells (mucus), and in the intestinal glands (crypts of Lieberkuhn) Paneth cells (lysozyme, a bactericidal enzyme, and phagocytosis) and enteroendocrine cells: S cells (secretin), CCK cells (cholecystokinin), and K cells (glucose-dependent insulinotropic peptide). Lamina propria of areolar connective tissue with MALT: solitary lymphatic nodules, most numerous in the distal ileum, and aggregated lymphatic follicles (Peyer patches) in the ileum. Muscularis mucosae of smooth muscle
Submucosa	In the duodenum, duodenal (Brunner) glands that secrete alkaline mucus to neutralize the acid in chyme. Lymphatic tissue sometimes extends down from the lamina propria
Muscularis	Two layers: a thicker inner circular layer and a thinner outer longitudinal layer
Serosa	Visceral peritoneum completely surrounds the small intestine, except the retroperitoneal part of the duodenum

Three levels of folding, largest to smallest

Feature	What it is	Seen with
Circular folds (plicae circulares)	Permanent ridges of mucosa and submucosa running around the inside of the tube, from the proximal duodenum to the middle of the ileum. Some go all the way around, some part way. Unlike rugae they do not flatten when the tube fills. They increase the surface and make chyme spiral instead of running straight through	The naked eye
Villi	Finger-like projections of the mucosa, 20 to 40 per square millimeter, that give the lining a velvety look. Each has a core of lamina propria with an arteriole, a venule, a capillary network, and a lacteal	A light microscope
Microvilli	Tiny projections of the apical membrane of each absorptive cell, each a bundle of 20 to 30 actin filaments, about 200 million per square millimeter. Together they form the brush border, which also carries digestive enzymes	An electron microscope (a fuzzy edge on a light slide)

- Fold, finger, fuzz. Circular folds are folds of the wall. Villi are fingers on the folds. Microvilli are the fuzz on each cell of each finger.
- The lacteal is the lymphatic capillary in the center of each villus. Most absorbed fats are too large to enter a blood capillary, so they enter the lacteal and ride the lymph before reaching the blood. Sugars and amino acids enter the blood capillaries.

Two hormones from the small intestine

Hormone	Released by	Trigger	Main effects
Cholecystokinin (CCK)	CCK cells of the intestinal glands	Chyme with amino acids and fatty acids	Pancreatic juice rich in enzymes; contraction of the gallbladder, squeezing bile down the cystic and common bile ducts; relaxation of the sphincter of Oddi; slower gastric emptying; a feeling of fullness (satiety) through the hypothalamus; growth of the pancreas; enhances secretin
Secretin	S cells of the intestinal glands	Acidic chyme	Pancreatic juice rich in bicarbonate, which buffers the acid; less gastric acid and gastric juice; growth of the pancreas; enhances CCK

Together they keep the duodenum from being overloaded with more chyme than it can handle. Other gut hormones: motilin, substance P, and bombesin stimulate motility; vasoactive intestinal peptide (VIP) stimulates secretion of ions and water and inhibits gastric acid; gastrin-releasing peptide stimulates gastrin release; somatostatin inhibits it.

Competency 117. Large intestine features

In your course notes: Module 4 packet, Chapter 3, competency 117 (page 34).

Dr. Rennie's video for this part: Digestive system concept video, chapter Small and large intestines.

You should be able to: Identify the cecum, appendix, four parts of the colon, rectum, and anal canal along with the teniae coli, haustra, and omental appendices.

The large intestine is the last part of the canal, from the ileocecal valve to the anus, about 1.5 m (5 feet) long and about 6.5 cm (2.5 inches) across, the same in the living person and the cadaver. It frames the small intestine on three sides. It absorbs water, ions, and some vitamins, forms and stores feces, and empties them. Bacteria living in it break down leftover proteins and make some B vitamins and vitamin K. Haustral churning, peristalsis, and mass peristalsis drive the contents toward the rectum.

The regions, in order

Region	Where it is
Cecum	A blind pouch in the lower right abdomen hanging below the ileocecal valve. The first part of the large intestine
Vermiform appendix	A narrow, worm-shaped tube attached to the cecum, full of lymphatic tissue. Its own small mesentery, the mesoappendix, ties it to the ileum. Food never travels through it
Ascending colon	Up the right side to the liver. Retroperitoneal
Hepatic (right colic) flexure	The bend under the liver where the ascending colon turns left
Transverse colon	Across the upper abdomen from right to left, below the stomach. Hangs on the transverse mesocolon
Splenic (left colic) flexure	The bend near the spleen where the transverse colon turns down
Descending colon	Down the left side. Retroperitoneal
Sigmoid colon	The S-shaped part beginning near the left iliac crest, curving medially into the pelvis to join the rectum. Hangs on the sigmoid mesocolon
Rectum	The straight part in the pelvis, in front of the sacrum and coccyx, from the rectosigmoid junction to the rectoanal junction. Stores feces temporarily; its lower widened part, the ampulla, holds them until defecation. Three transverse rectal folds, one on the right and two on the left
Anal canal	The last 2 to 3 cm, passing through the pelvic floor to open at the anus. Its mucosa is in longitudinal folds, the anal columns, with arteries and veins. Lined with stratified squamous epithelium near the anus. Surrounded by the internal anal sphincter (smooth, involuntary) and the external anal sphincter (skeletal, voluntary), which keep the anus closed except during defecation

- The colon has four parts: ascending, transverse, descending, sigmoid. The two flexures are bends between them, not separate parts.
- Trace it as a frame: up the right side, a bend at the liver, across the top, a bend at the spleen, down the left side, then an S into the pelvis.
- Rectum and anal canal lengths differ between sources, so you are asked where each one sits, not how long it is.

The wall of the large intestine

Layer	In the large intestine
Mucosa	Simple columnar epithelium with absorptive cells (water absorption) and many goblet cells (mucus to lubricate the passage), both in intestinal glands that run the full thickness of the mucosa. No circular folds and no villi, only microvilli on the absorptive cells, so far less absorption than the small intestine. Lamina propria with solitary lymph nodules
Submucosa	Areolar connective tissue
Muscularis	The outer longitudinal layer is gathered into three thick bands, the teniae coli, with little or no longitudinal muscle between them. The inner circular layer is a complete sheet
Serosa	Visceral peritoneum, with the omental appendices hanging from it

Three features found only on the large intestine

Teniae coli

Three flat ribbons of smooth muscle running the length of the cecum and colon: the outer longitudinal layer gathered into bands instead of a full sheet. They spread back into a complete layer at the rectum, so the rectum has no haustra. The three meet at the base of the appendix, which is one way to find it.

Haustra

The row of pouches that gives the colon its puckered look. The teniae are shorter than the wall and keep a tonic contraction, so the wall bunches up between them.

Omental (epiploic) appendices

Small fat-filled pouches of visceral peritoneum hanging from the surface of the colon, mostly along the teniae.

These three are how you tell large from small intestine at a glance on a model or the cadaver. Count the teniae: three, not two or four.

Blood and nerves of the rectum

- Arteries: the superior rectal artery (from the inferior mesenteric artery) supplies the upper rectum; the right and left middle rectal arteries supply the lower rectum.
- Veins: a venous plexus in the wall drained by the rectal veins.
- Nerves: sympathetic fibers from the lumbar splanchnic nerves through the hypogastric plexus; parasympathetic and visceral sensory fibers through the pelvic splanchnic nerves.

Competency 118. Trace the pathway of food

In your course notes: Module 4 packet, Chapter 3, competency 118 (page 35).

You should be able to: Trace a bite of food through every named segment of the alimentary canal in order from mouth to anus.

A bite of food moves through the alimentary canal in one direction only. Every segment below is one the food passes through, in order, with the sphincters at the borders where they sit.

1. Oral cavity: food comes in through the lips, is chewed by the teeth, mixed with saliva, and shaped into a bolus by the tongue.
2. Fauces: the tongue pushes the bolus back through the opening into the throat.
3. Oropharynx: the soft palate and uvula rise to seal the nasopharynx above.
4. Laryngopharynx: the epiglottis covers the larynx, so the bolus passes behind it toward the esophagus.
5. Upper esophageal sphincter relaxes.
6. Esophagus: waves of muscle contraction carry the bolus down through the thorax and the esophageal hiatus.
7. Lower esophageal sphincter relaxes.
8. Cardia of the stomach, then the body, where the food is churned with gastric juice into chyme.
9. Pyloric antrum, pyloric canal, pylorus: the funnel out of the stomach.
10. Pyloric sphincter lets small amounts of chyme through.
11. Duodenum: bile and pancreatic juice enter at the major duodenal papilla.
12. Jejunum: digestion continues and most nutrients are absorbed.
13. Ileum: absorption continues.
14. Ileocecal sphincter (valve) lets the remaining material into the large intestine.
15. Cecum, then ascending colon, hepatic flexure, transverse colon, splenic flexure, descending colon, sigmoid colon.
16. Rectum: feces are stored.
17. Anal canal, through the internal and external anal sphincters, to the anus.

- Three named structures are off the path. The nasopharynx carries only air and is sealed during a swallow. The fundus bulges above the cardia, so food can collect there without passing through it. The appendix is a blind tube off the cecum.
- The cecum comes right after the ileum. Do not jump from the ileum to the ascending colon.

- Group the path as the six organs first (mouth, pharynx, esophagus, stomach, small intestine, large intestine), then fill in the named parts inside each one and put a sphincter at each border.
- Most absorption happens in the small intestine. Water is absorbed and feces form in the large intestine.

Competency 119. Clinical anatomy of canal disorders

In your course notes: Module 4 packet, Chapter 3, competency 119 (page 36).

You should be able to: Match GERD, hiatal hernia, peptic ulcer, appendicitis, diverticulosis, IBD, hemorrhoids, and colorectal cancer to the canal structure each one affects.

Each disorder starts in a specific structure. If you know where that structure sits and what layers it has, you can predict where the patient feels it and how serious it can become.

Disorder	Structure affected	What changes
Gastroesophageal reflux disease (GERD)	Lower esophageal sphincter and lower esophagus	The sphincter does not close well, so acidic stomach contents flow back up. The stratified squamous lining of the esophagus has no protection against acid and becomes inflamed. Felt as heartburn, a burning behind the sternum, because the esophagus lies in the thorax just behind it
Hiatal hernia	Esophageal hiatus and upper stomach	Part of the stomach pushes up through the hiatus into the thorax. The diaphragm loses its grip on the lower esophageal sphincter, so reflux often follows
Peptic ulcer	Mucosa of the stomach or duodenum	An open sore where the mucosa has worn away. Common causes: infection with Helicobacter pylori and regular use of NSAID pain relievers. It deepens layer by layer: through the submucosa, where it can open a vessel and bleed, through the muscularis, and through the serosa, where it perforates and the contents spill into the peritoneal cavity
Appendicitis	Vermiform appendix	The lumen blocks, the appendix swells and inflames, and pain lands in the lower right abdomen where the cecum sits. If it bursts, its contents spill into the peritoneal cavity: peritonitis
Diverticulosis	Wall of the colon, most often the sigmoid	Small pouches (diverticula) of mucosa and submucosa bulge out through weak spots in the muscularis. When the pouches inflame, it is diverticulitis
Inflammatory bowel disease (IBD)	Wall of the small and large intestine	Long-term inflammation of the wall. Crohn disease can affect any part from mouth to anus, most often the ileum and colon, through every layer. Ulcerative colitis affects only the colon and rectum, mostly the mucosa
Hemorrhoids	Veins in the wall of the anal canal and lower rectum	The veins swell and stretch into soft bulges. External hemorrhoids are thrombosed veins of the external plexus, often with pregnancy or constipation, and painful because they have somatic innervation. Internal hemorrhoids are dilated veins of the internal rectal plexus above the dentate line, visceral innervation, so painless unless they ulcerate
Colorectal cancer	Mucosa of the colon or rectum	Lining cells grow out of control, usually starting as a polyp, then growing through the deeper layers of the wall

What this means for patients

- Where a problem sits predicts the symptom. A weak lower esophageal sphincter burns behind the sternum. An inflamed appendix hurts in the lower right.
- The four-layer wall explains how a problem gets worse. Ulcers, diverticula, IBD, and cancer all become more serious as they reach deeper layers, and a perforation through the serosa turns a gut problem into peritonitis.

- GERD and hiatal hernia are two different things that often travel together: one is a weak sphincter, the other is stomach through the diaphragm.

Worked questions

Nine questions in the style of the lecture exam. Read the answer and the reasons each wrong option is wrong. The wrong options are where the learning is.

QUESTION 1 · COMPETENCY 116

Which region of the small intestine is the longest?

A. Duodenum

B. Jejunum

C. Ileum

D. They are all about the same length

Answer: C. The ileum is about 2 m (6 feet), the jejunum about 1 m (3 feet), and the duodenum about 25 cm (10 inches). The ileum is the last and longest region.

Why the others are wrong

- **A** The duodenum is the shortest, a C around the head of the pancreas.
- **B** The jejunum is the middle region and about half the length of the ileum.
- **D** The three differ a lot; the duodenum is a fraction of the length of the other two.

QUESTION 2 · COMPETENCY 116

The lacteal in the center of a villus is

A. a blood capillary that absorbs sugars

B. a lymphatic capillary that absorbs most fats

C. a duct that carries bile into the villus

D. a nerve of the submucosal plexus

Answer: B. The lacteal is a lymphatic capillary. Absorbed fats are too large for blood capillaries, so they enter the lacteal and travel in lymph before reaching the blood.

Why the others are wrong

- **A** The villus does have a blood capillary network, and sugars and amino acids enter it, but that is not the lacteal.
- **C** Bile enters the duodenum at the major duodenal papilla and stays in the lumen. No duct runs into a villus.
- **D** The submucosal plexus sits in the submucosa, below the mucosa. The lacteal is a vessel, not a nerve.

QUESTION 3 · COMPETENCY 116

Put the three surface-expanding features of the small intestine in order from largest to smallest.

A. Villi, circular folds, microvilli

B. Microvilli, villi, circular folds

C. Circular folds, villi, microvilli

D. Circular folds, microvilli, villi

Answer: C. Circular folds are ridges of the whole wall you can see with your eyes. Villi are fingers of mucosa on the folds, seen with a light microscope. Microvilli are the fuzz on one cell, seen with an electron microscope. Fold, finger, fuzz.

Why the others are wrong

- **A** Circular folds are larger than villi; villi sit on the folds.
- **B** This is smallest to largest, the reverse order.
- **D** Microvilli are the smallest, on the surface of a single absorptive cell on a villus.

QUESTION 4 · COMPETENCY 117

Which two parts of the colon hang on a mesocolon?

A. Ascending and descending

B. Ascending and transverse

C. Transverse and sigmoid

D. Descending and sigmoid

Answer: C. The transverse colon hangs on the transverse mesocolon and the sigmoid colon on the sigmoid mesocolon. Those are the two intraperitoneal parts.

Why the others are wrong

- **A** Ascending and descending are retroperitoneal, fixed to the back wall.
- **B** The transverse colon is right, but the ascending colon is retroperitoneal.
- **D** The sigmoid colon is right, but the descending colon is retroperitoneal.

QUESTION 5 · COMPETENCY 117

The teniae coli are

A. three bands of the outer longitudinal muscle layer

B. three bands of the inner circular muscle layer

C. fat-filled pouches of peritoneum on the colon

D. the pouches that give the colon its puckered look

Answer: A. The outer longitudinal layer of the muscularis is gathered into three flat ribbons instead of a complete sheet. Because they are shorter than the wall, the wall bunches into haustra between them.

Why the others are wrong

- **B** The inner circular layer stays a complete sheet in the large intestine.
- **C** Those are the omental appendices, which hang along the teniae.
- **D** Those are the haustra, the pouches the teniae create.

QUESTION 6 · COMPETENCY 117

Which sphincter of the anal canal is smooth muscle and not under voluntary control?

A. External anal sphincter

B. Internal anal sphincter

C. Ileocecal sphincter

D. Pyloric sphincter

Answer: B. The internal anal sphincter is a thickening of the circular smooth muscle of the anal canal. It is involuntary. The external sphincter outside it is skeletal and voluntary.

Why the others are wrong

- **A** The external anal sphincter is skeletal muscle, the one you control.
- **C** The ileocecal sphincter is smooth and involuntary, but it sits between the ileum and the cecum, not at the anal canal.
- **D** The pyloric sphincter is at the exit of the stomach.

QUESTION 7 · COMPETENCY 118

A bite of food leaves the ileum. Which segment does it enter next?

A. Ascending colon

B. Cecum

C. Transverse colon

D. Rectum

Answer: B. The ileum opens through the ileocecal valve into the cecum, the pouch at the start of the large intestine. The ascending colon comes after the cecum.

Why the others are wrong

- **A** The ascending colon rises from the cecum; the food reaches it one step later.
- **C** The transverse colon is two steps and one flexure further on.
- **D** The rectum is near the end of the path, after the sigmoid colon.

QUESTION 8 · COMPETENCY 119

A patient has a burning pain behind the sternum after meals because stomach acid is flowing back into the esophagus. Which structure is failing?

A. Pyloric sphincter

B. Upper esophageal sphincter

C. Lower esophageal sphincter

D. Ileocecal valve

Answer: C. This is GERD. The lower esophageal sphincter does not close well, acid comes up onto the stratified squamous lining of the esophagus, and the esophagus sits in the thorax just behind the sternum, so that is where it burns.

Why the others are wrong

- **A** The pyloric sphincter controls flow out of the stomach into the duodenum. It has nothing to do with backflow into the esophagus.
- **B** The upper esophageal sphincter sits at the top of the esophagus. Acid would have to pass the lower sphincter first.
- **D** The ileocecal valve is between the small and large intestine, far from the sternum.

QUESTION 9 · COMPETENCY 119

A peptic ulcer starts in which layer and erodes in which direction?

A. Serosa, inward toward the lumen

B. Muscularis, in both directions

C. Mucosa, outward toward the serosa

D. Submucosa, inward toward the lumen

Answer: C. An ulcer begins on the inside, where acid and H. pylori meet the mucosa. It deepens through the submucosa (where it can bleed), the muscularis, and finally the serosa, where it perforates into the peritoneal cavity.

Why the others are wrong

- **A** The serosa is the outside of the tube; the acid is on the inside. Ulcers do not start from the peritoneal cavity.
- **B** The muscularis is the third layer in; an ulcer reaches it only after eating through the mucosa and submucosa.
- **D** The submucosa is the second layer. The ulcer gets there from the mucosa and continues outward, not inward.

BIO 004 Human Anatomy · Module 4 notes · Dr. Sharilyn Rennie

Blood flow of the abdomen and lower limb

The written notes for the blood flow lecture: the abdominal aorta and its branches, the inferior vena cava and its tributaries, the hepatic portal system, and the vessels and nerves of the lower limb. Module 4 competencies 126 and 127, with nine worked questions.

Competency 126. The abdominal aorta and its branches

In your course notes: Module 4 packet, Chapter 5, competency 126 (page 46).

Dr. Rennie's video for this part: No concept video for this chapter. Use the flow charts in the Module 4 lab packet and the cadaver.

You should be able to: Identify the major branches of the abdominal aorta and the vessels of the hepatic portal system on the cadaver.

The abdominal aorta is the aorta below the diaphragm. It enters the abdomen through the aortic hiatus at about T12, runs down the front of the lumbar vertebrae just left of the midline with the inferior vena cava on its right, and ends at about L4 by dividing into the right and left common iliac arteries. Its branches come in two groups: unpaired branches leave the front and feed the digestive organs; paired branches leave the sides and feed the kidneys, the gonads, and other structures that come in twos.

Branch	About where it leaves	What it supplies
Celiac trunk (unpaired)	T12, just below the diaphragm	Short; splits almost at once into the left gastric, splenic, and common hepatic arteries
Left gastric artery	From the celiac trunk	The stomach along the lesser curvature and the lower esophagus
Splenic artery	From the celiac trunk	Runs left along the top of the pancreas to the spleen, with branches to the pancreas and stomach
Common hepatic artery	From the celiac trunk	Runs right; gives the hepatic artery proper to the liver and gallbladder, plus branches to the stomach, duodenum, and pancreas
Superior mesenteric artery (unpaired)	L1	Runs in the mesentery to most of the small intestine, part of the pancreas, the cecum, appendix, ascending colon, and most of the transverse colon
Renal arteries (paired)	L1 to L2	The kidneys
Gonadal arteries (paired)	L2	The testes (testicular arteries) or the ovaries (ovarian arteries)
Inferior mesenteric artery (unpaired)	L3	The end of the transverse colon, the descending colon, the sigmoid colon, and the upper rectum
Right and left common iliac arteries	L4, where the aorta ends	The pelvis and lower limbs, through the internal and external iliac arteries

- The three gut branches go in the same order as the gut. Celiac feeds the top (stomach, liver, spleen, pancreas, start of the duodenum). Superior mesenteric feeds the middle (small intestine and the colon through most of the transverse). Inferior mesenteric feeds the end (the rest of the colon and the upper rectum).
- The three unpaired branches are the celiac trunk, superior mesenteric, and inferior mesenteric. Renal and gonadal arteries are paired.
- The hepatic artery proper rides up in the free edge of the lesser omentum to the porta hepatis.

Arteries of the pelvis

- Each common iliac artery divides in front of the sacroiliac joint. The internal iliac artery dives into the pelvis and supplies the bladder, rectum, reproductive organs, pelvic walls, gluteal muscles, and external genitalia. The external iliac artery runs along the psoas major, passes under the inguinal ligament, and becomes the femoral artery.
- The medial umbilical ligament is a fibrous cord on the inside of the anterior abdominal wall running from the pelvis toward the umbilicus. It is what remains of the fetal umbilical artery, a branch of the internal iliac. The part nearest the internal iliac stays open and supplies the bladder.

Competency 126. The inferior vena cava and the veins of the abdomen

In your course notes: Module 4 packet, Chapter 5, competency 126 (page 48).

The inferior vena cava (IVC) forms at about L5 where the right and left common iliac veins join. It runs up the right side of the aorta, passes through the central tendon of the diaphragm at about T8, and empties into the right atrium.

Vein	Drains	Empties into
External iliac vein	The lower limb; it is the femoral vein above the inguinal ligament	Common iliac vein
Internal iliac vein	The pelvis	Common iliac vein
Right and left common iliac veins	The pelvis and lower limb	Join to form the inferior vena cava
Renal veins	The kidneys. The left renal vein is longer because it crosses in front of the aorta to reach the IVC	Inferior vena cava
Right gonadal vein	The right testis or ovary	Inferior vena cava
Left gonadal vein	The left testis or ovary	Left renal vein
Hepatic veins	The liver, after the sinusoids and central veins	Inferior vena cava, just below the diaphragm
Inferior vena cava	Most of the body below the diaphragm	Right atrium

- The gonadal veins are not symmetric. Right into the IVC, left into the left renal vein.
- The stomach, intestines, spleen, and pancreas are not on this list. Their blood goes to the liver first, through the hepatic portal system.

Competency 126. The hepatic portal system

In your course notes: Module 4 packet, Chapter 5, competency 126 (page 48), and Chapter 4, competency 122 (page 41).

A portal system is a route in which blood passes through two capillary beds in a row before it returns to the heart. In the hepatic portal system the first bed is in the walls of the stomach, intestines, spleen, and pancreas, and the second bed is the sinusoids of the liver. The hepatic portal vein connects them. Everything absorbed from the gut, nutrients and whatever came with them, is processed by the liver before it reaches the rest of the body. It is also why cancers of the digestive tract so often spread to the liver first.

Vessel	Drains	Empties into
Superior mesenteric vein	The small intestine, cecum, ascending colon, and most of the transverse colon; runs beside the superior mesenteric artery	Joins the splenic vein to form the hepatic portal vein
Inferior mesenteric vein	The descending colon, sigmoid colon, and upper rectum	Usually the splenic vein
Splenic vein	The spleen, pancreas, and part of the stomach; runs along the back of the pancreas	Joins the superior mesenteric vein to form the hepatic portal vein
Hepatic portal vein	All of the above. Forms behind the neck of the pancreas, runs up in the free edge of the lesser omentum, and enters the liver at the porta hepatis	The hepatic sinusoids
Hepatic veins	The liver, from the central veins	Inferior vena cava

A glucose molecule, from a villus to the heart

1. Capillaries of an intestinal villus in the jejunum. The glucose enters the blood here: the first capillary bed.
2. Small veins of the intestinal wall and mesentery.
3. Superior mesenteric vein.
4. Hepatic portal vein, formed where the superior mesenteric and splenic veins join.
5. Porta hepatis, then a portal vein branch in a portal triad.
6. Hepatic sinusoids, past the hepatocytes: the second capillary bed.
7. Central vein of the lobule.
8. Hepatic veins, out of the back of the liver.
9. Inferior vena cava, through the diaphragm at T8.
10. Right atrium.

- The hepatic portal vein carries blood into the liver. The hepatic veins carry it out. Portal in, hepatic out.
- Two things enter the porta hepatis (the portal vein and the hepatic artery proper) and the hepatic ducts leave it. The hepatic veins do not use the porta hepatis; they leave the posterior surface straight into the vena cava.
- Portal vein blood is nutrient-rich and oxygen-poor. The hepatic artery brings the oxygen. They mix in the sinusoids and flow in toward the central vein, while bile flows out toward the triads.

Competency 127. Arteries and veins of the lower limb

In your course notes: Module 4 packet, Chapter 5, competency 127 (page 49).

You should be able to: Identify the femoral and lower-limb blood vessels and the major nerves of the abdomen and lower limb on the cadaver.

In the lower limb, arteries, veins, and nerves run together in bundles between the muscles you already know. Use the muscles as landmarks. When you name a vessel, say right or left and artery or vein.

One artery, renamed at each landmark

1. External iliac artery: along the pelvic brim from the common iliac.
2. Femoral artery: the same vessel after it passes under the inguinal ligament. Its pulse is felt about halfway between the anterior superior iliac spine and the pubic symphysis. It runs through the femoral triangle, gives off the deep femoral artery to most of the thigh muscles, then runs down under the sartorius.
3. Popliteal artery: the same vessel after it passes through the adductor hiatus in adductor magnus to reach the back of the knee.
4. Anterior and posterior tibial arteries: the popliteal divides just below the knee. The anterior tibial goes forward into the front of the leg and becomes the dorsalis pedis artery on the top of the foot. The posterior tibial, the larger branch, runs down the back of the leg deep to the gastrocnemius and soleus beside the tibial nerve, gives off the fibular artery to the lateral leg, passes behind the medial malleolus (second pulse point), and divides into the plantar arteries in the sole.

Femoral triangle
: The space just below the inguinal ligament, bounded by the inguinal ligament above, the sartorius laterally, and the adductor longus medially. From lateral to medial: femoral nerve, femoral artery, femoral vein, empty space, lymphatics (NAVEL). The vein is medial to the artery.

Adductor hiatus
: The opening in adductor magnus through which the femoral artery and vein pass to the back of the knee, where their names change to popliteal.

Veins of the lower limb

Vessel	Deep or superficial	Course and landmarks
Posterior and anterior tibial veins, popliteal vein, femoral vein	Deep	Run beside the arteries and share their names. The tibial veins join into the popliteal vein, which becomes the femoral vein up the thigh (medial to the artery in the triangle), which becomes the external iliac vein above the inguinal ligament. The deep veins carry most of the blood out of the limb
Great saphenous vein	Superficial	The longest vein in the body. Starts on the medial side of the top of the foot, passes in front of the medial malleolus, runs up the medial leg and thigh in the fat under the skin, and empties into the femoral vein just below the inguinal ligament. Surgeons use it as a graft in coronary bypass surgery
Small saphenous vein	Superficial	Up the back of the calf into the popliteal vein

- The name changes at three landmarks on the way to the foot: the diaphragm (thoracic to abdominal aorta), the inguinal ligament (external iliac to femoral), and the adductor hiatus (femoral to popliteal).
- The great saphenous vein is superficial and medial. The femoral vein is the deep one.

Competency 127. Nerves of the abdomen and lower limb

In your course notes: Module 4 packet, Chapter 5, competency 127 (page 50).

Only the femoral nerve is on your Module 4 structure list, and the plexus notes also name the obturator nerve. The rest are here so you can recognize what you see on the posterior abdominal wall and in the limb. You identify nerves by location; you are not asked which nerve supplies each muscle.

On the posterior abdominal wall

The lumbar plexus forms inside the psoas major from the anterior rami of L1 to L4. Its branches come out of the psoas and cross the posterior wall, which is where you see them on the cadaver.

Nerve	Spinal nerves	Where you see it	What it supplies
Iliohypogastric	L1	Crossing the posterior wall toward the iliac crest	Lower abdominal wall muscles, skin above the pubis
Ilioinguinal	L1	Just below the iliohypogastric, then through the inguinal canal	Lower abdominal wall, skin of the upper medial thigh and external genitalia
Genitofemoral	L1 to L2	Down the front of the psoas major	Skin of the external genitalia and a patch of upper anterior thigh
Lateral femoral cutaneous	L2 to L3	Across the iliacus toward the anterior superior iliac spine	Skin of the lateral thigh
Femoral	L2 to L4	Out of the lateral side of the psoas, in the groove between psoas major and iliacus	Anterior thigh muscles and skin
Obturator	L2 to L4	Out of the medial side of the psoas, to the obturator foramen	Adductor muscles and skin of the medial thigh

- Most of the anterolateral abdominal wall is supplied by the lower intercostal nerves (T7 to T11) and the subcostal nerve (T12); the lumbar plexus supplies only the lowest strip above the pubis and the groin.
- The anterior and posterior vagal trunks (vagus, cranial nerve X) enter the abdomen on the esophagus through the esophageal hiatus and supply the stomach and other abdominal organs.

In the lower limb

Nerve	Plexus	Course	Muscles supplied
Femoral nerve	Lumbar plexus (L2 to L4)	Under the inguinal ligament, lateral to the femoral artery, into the femoral triangle	Anterior thigh: quadriceps femoris and sartorius; a branch continues to the skin of the medial leg
Sciatic nerve	Sacral plexus (L4 to S3)	Out the greater sciatic foramen, deep to gluteus maximus, down the back of the thigh deep to the hamstrings; the largest nerve in the body	The hamstrings; splits above the knee into the tibial and common fibular nerves
Tibial nerve	Sacral, from the sciatic	Straight down through the back of the knee and the posterior leg deep to gastrocnemius and soleus, beside the posterior tibial artery, behind the medial malleolus into the sole	Posterior leg muscles and the sole of the foot
Common fibular nerve	Sacral, from the sciatic	Along the biceps femoris tendon to the lateral knee, around the neck of the fibula just under the skin, where it splits into superficial and deep branches	Superficial branch: lateral leg (fibularis longus and brevis). Deep branch: anterior leg (tibialis anterior, extensor hallucis longus, extensor digitorum longus, fibularis tertius)

- The tibial and common fibular nerves come off the sciatic nerve, not straight off the plexus.
- Because the common fibular nerve lies against the neck of the fibula just under the skin, a blow or a tight cast there can injure it and weaken the muscles that lift the foot: foot drop.

Worked questions

Nine questions in the style of the lecture exam. Read the answer and the reasons each wrong option is wrong. The wrong options are where the learning is.

QUESTION 1 · COMPETENCY 126

Which of these is an unpaired branch of the abdominal aorta?

A. Renal artery

B. Gonadal artery

C. Superior mesenteric artery

D. Common iliac artery

Answer: C. The superior mesenteric artery leaves the front of the aorta at about L1 as a single vessel and feeds the middle of the gut. The three unpaired branches are the celiac trunk, the superior mesenteric, and the inferior mesenteric.

Why the others are wrong

- **A** The renal arteries are paired, one to each kidney, leaving the sides of the aorta.
- **B** The gonadal arteries are paired, one to each testis or ovary.
- **D** The common iliac arteries are the paired terminal branches where the aorta ends at L4.

QUESTION 2 · COMPETENCY 126

The celiac trunk splits into which three arteries?

A. Left gastric, splenic, common hepatic

B. Superior mesenteric, renal, gonadal

C. Left gastric, right gastric, inferior mesenteric

D. Splenic, renal, common hepatic

Answer: A. The celiac trunk is short and splits almost at once into the left gastric artery (stomach and lower esophagus), the splenic artery (spleen, pancreas, stomach), and the common hepatic artery (liver, gallbladder, stomach, duodenum, pancreas).

Why the others are wrong

- **B** Those are separate branches of the aorta itself, below the celiac trunk.
- **C** The inferior mesenteric artery is its own unpaired branch at L3, not part of the celiac trunk.
- **D** The renal arteries come straight off the sides of the aorta, not from the celiac trunk.

QUESTION 3 · COMPETENCY 126

The left gonadal vein empties into the

A. inferior vena cava

B. left renal vein

C. left common iliac vein

D. hepatic portal vein

Answer: B. The gonadal veins are not symmetric. The right one drains straight into the inferior vena cava; the left one drains into the left renal vein, which then crosses in front of the aorta to the IVC.

Why the others are wrong

- **A** That is the right gonadal vein.
- **C** The common iliac veins collect from the pelvis and lower limb, below the level of the gonadal veins.
- **D** The hepatic portal vein drains the digestive organs and spleen, not the gonads.

QUESTION 4 · COMPETENCY 126

What makes the hepatic portal system a portal system?

A. It carries oxygen-rich blood to the liver

B. Its blood passes through two capillary beds in a row before returning to the heart

C. It drains directly into the right atrium

D. It is the only vein that carries blood away from the liver

Answer: B. A portal system is two capillary beds in series. The first is in the wall of the gut, spleen, and pancreas; the second is the hepatic sinusoids. The hepatic portal vein connects them.

Why the others are wrong

- **A** Portal vein blood is oxygen-poor and nutrient-rich. The hepatic artery brings the oxygen.
- **C** The portal vein empties into the liver sinusoids. The hepatic veins and the inferior vena cava carry the blood on to the right atrium.
- **D** The portal vein carries blood into the liver. The hepatic veins carry it away.

QUESTION 5 · COMPETENCY 126

Which vessel carries blood out of the liver?

A. Hepatic portal vein

B. Hepatic artery proper

C. Hepatic veins

D. Splenic vein

Answer: C. The central veins of the lobules join into the hepatic veins, which leave the back of the liver and empty into the inferior vena cava just under the diaphragm.

Why the others are wrong

- **A** The hepatic portal vein brings nutrient-rich blood from the gut into the liver at the porta hepatis.
- **B** The hepatic artery proper brings oxygen-rich blood into the liver.
- **D** The splenic vein drains the spleen, pancreas, and part of the stomach and joins the superior mesenteric vein to form the portal vein, upstream of the liver.

QUESTION 6 · COMPETENCY 127

The femoral artery becomes the popliteal artery when it

A. passes under the inguinal ligament

B. gives off the deep femoral artery

C. passes through the adductor hiatus to the back of the knee

D. divides below the knee

Answer: C. The femoral artery runs down under the sartorius and through the adductor hiatus, the opening in adductor magnus, to reach the popliteal fossa. From there it is the popliteal artery.

Why the others are wrong

- **A** At the inguinal ligament the external iliac artery becomes the femoral artery, one name change earlier.
- **B** The deep femoral branch leaves in the femoral triangle, but the femoral artery keeps its name and continues down the thigh.
- **D** Below the knee the popliteal artery divides into the anterior and posterior tibial arteries.

QUESTION 7 · COMPETENCY 127

In the femoral triangle, from lateral to medial, the order is

A. vein, artery, nerve

B. artery, vein, nerve

C. nerve, artery, vein

D. nerve, vein, artery

Answer: C. NAVEL: nerve, artery, vein, empty space, lymphatics, from lateral to medial. The femoral vein lies on the medial side of the artery.

Why the others are wrong

- **A** This is the reverse order; the nerve is the most lateral structure.
- **B** The nerve is lateral to the artery, not medial to the vein.
- **D** The artery lies between the nerve and the vein.

QUESTION 8 · COMPETENCY 127

Which vein is superficial and runs up the medial side of the leg and thigh, passing in front of the medial malleolus?

A. Femoral vein

B. Popliteal vein

C. Great saphenous vein

D. Posterior tibial vein

Answer: C. The great saphenous vein is the longest vein in the body. It starts on the medial top of the foot, passes in front of the medial malleolus, runs up the medial leg and thigh in the fat under the skin, and drains into the femoral vein.

Why the others are wrong

- **A** The femoral vein is deep, beside the femoral artery.
- **B** The popliteal vein is deep, behind the knee.
- **D** The posterior tibial vein is deep, running with the posterior tibial artery behind the medial malleolus, not in front of it.

QUESTION 9 · COMPETENCY 127

A tight cast around the neck of the fibula weakens the muscles that lift the foot. Which nerve is injured?

A. Tibial nerve

B. Common fibular nerve

C. Femoral nerve

D. Obturator nerve

Answer: B. The common fibular nerve wraps around the neck of the fibula just under the skin, and its deep branch supplies the anterior leg muscles that dorsiflex the foot. Pressure there causes foot drop.

Why the others are wrong

- **A** The tibial nerve runs deep in the back of the leg and supplies the posterior leg muscles, which point the foot down.
- **C** The femoral nerve supplies the anterior thigh (quadriceps and sartorius) and never reaches the fibula.
- **D** The obturator nerve supplies the adductors of the medial thigh.

BIO 004 Human Anatomy · Module 4 notes · Dr. Sharilyn Rennie

Endocrine anatomy

The written notes for the endocrine lecture: the major glands and where each sits, the pituitary and its two lobes, the thyroid and parathyroid glands, the adrenal zones, and the pancreas, gonads, and other organs with endocrine tissue. Module 4 competencies 128 to 132, with nine worked questions.

Before the competencies: what an endocrine gland is

In your course notes: Module 4 packet, Chapter 6 introduction (page 52).

Dr. Rennie's video for this part: Endocrine system concept video, chapter Endocrine system overview and purpose.

The endocrine system is a scattered set of glands and tissues that release hormones, chemical messengers that travel in the blood. Unlike most organ systems, most of its glands are not physically connected to each other. They sit in the head, neck, thorax, abdomen, pelvis, and scrotum, and the blood is what links them.

An endocrine gland is a ductless gland. Its cells release their product into the surrounding interstitial fluid, and from there it enters the blood. That is why endocrine glands have such a rich supply of capillaries. An exocrine gland is different: it has a duct that carries its product onto a surface or into a cavity, the way the salivary glands empty into the mouth. Some organs, such as the pancreas, have both kinds of tissue.

Endocrine system
: The glands and tissues that release hormones into the blood to signal distant cells.

Hormone
: A chemical messenger released by an endocrine cell that travels in the blood to target cells.

Endocrine gland
: A ductless gland that releases its product into the surrounding fluid and the blood.

Exocrine gland
: A gland that releases its product through a duct onto a surface or into a cavity. Not part of the endocrine system.

Target cell
: A cell that carries the receptor for a given hormone and so responds to it.

- Ductless and into the blood means endocrine. Through a duct and onto a surface means exocrine.
- Because hormones travel in the blood, an endocrine gland can be far from the cells it controls.

Competency 128. Major endocrine glands and locations

In your course notes: Module 4 packet, Chapter 6, competency 128 (page 52).

Dr. Rennie's video for this part: Endocrine system concept video, chapter Major glands and their roles.

You should be able to: Name the major endocrine glands and locate each one in the body from the hypothalamus to the gonads.

Learn the major glands from the top of the body down. Three are in the brain or hang from it: the hypothalamus, the pituitary gland, and the pineal gland. Two are in the neck: the thyroid gland and the parathyroid glands behind it. The thymus is in the thorax. The adrenal glands and the pancreas are in the abdomen, and the endocrine tissue of the pancreas is the pancreatic islets. The gonads are the ovaries in the female pelvis and the testes in the male scrotum.

Where each gland sits

Gland	Body region	Location
Hypothalamus	Brain	The lowest part of the diencephalon, below the thalamus. It forms the floor and lower walls of the third ventricle. It links the nervous and endocrine systems and controls the pituitary
Pituitary gland	Brain	A pea-sized gland in the sella turcica of the sphenoid bone, hanging below the hypothalamus by the infundibulum
Pineal gland	Brain	A small, cone-shaped gland in the epithalamus, projecting from the posterior roof of the third ventricle
Thyroid gland	Neck	A butterfly-shaped gland in the anterior neck just below the larynx. Its two lobes lie beside the trachea and its isthmus crosses in front of it. The largest purely endocrine gland
Parathyroid glands	Neck	Usually four tiny glands on the posterior surface of the thyroid lobes, two on each side
Thymus	Thorax	In the mediastinum, behind the sternum and above the heart. Large in childhood, shrinks after puberty
Adrenal glands	Abdomen	Also called the suprarenal glands. One on the superior pole of each kidney, on the posterior abdominal wall behind the peritoneum
Pancreatic islets	Abdomen	Small clusters of endocrine cells scattered through the pancreas, which lies behind the stomach
Ovaries	Pelvis	In the pelvic cavity, one on each side of the uterus
Testes	Scrotum	In the scrotum, outside the abdominopelvic cavity

- Memory tip: learn the glands from top to bottom by body region. Brain: hypothalamus, pituitary, pineal. Neck: thyroid and parathyroids. Chest: thymus. Abdomen: adrenals and pancreas. Pelvis: ovaries, or testes in the scrotum.
- The pineal gland sits at the back of the third ventricle and the pituitary hangs below the hypothalamus, in front and below. They are not in the same spot.
- Four of the glands come in pairs or more: the parathyroids (usually four, not two), the adrenals, the ovaries, and the testes. The pancreatic islets are many small clusters inside one pancreas.
- The thyroid is the largest gland in the body that is purely endocrine. The thymus is large in childhood and shrinks after puberty.

What this means for patients

- The pituitary gland sits in the bony sella turcica, so a growing pituitary tumor usually expands upward out of the sella. Directly above it is the optic chiasm, and pressure there can cause a loss of peripheral vision. That vision change can be the first sign that a gland the size of a pea has enlarged.
- The thyroid gland lies just under the skin of the anterior neck, so an enlarged thyroid, a goiter, is visible and can be felt. The same superficial position lets a clinician palpate the gland and makes thyroid surgery accessible, though the tiny parathyroid glands on its back surface must be carefully spared.
- Each adrenal gland rests on the superior pole of a kidney. Imaging ordered for the kidney also captures the adrenal, which is why an adrenal mass is sometimes found by accident.

Competency 129. Pituitary gross anatomy and lobes

In your course notes: Module 4 packet, Chapter 6, competency 129 (page 54).

Dr. Rennie's video for this part: Endocrine system concept video, chapter Hypothalamus and pituitary control.

You should be able to: Describe the pituitary in the sella turcica, its infundibulum link to the hypothalamus, and compare the anterior and posterior lobes by tissue type.

The pituitary gland, also called the hypophysis, is about the size of a pea. It sits in the sella turcica, a saddle-shaped pocket in the body of the sphenoid bone in the middle of the base of the skull. The bone forms the floor and the front and back walls of this pocket and protects the gland. The optic chiasm lies just above and in front of it.

The pituitary hangs from the hypothalamus by a slender stalk called the infundibulum. The infundibulum is the physical link between the brain and the gland. Nerve fibers from the hypothalamus run down through it, and small blood vessels run down along it.

Hypophysis
: Another name for the pituitary gland.

Sella turcica
: The saddle-shaped pocket of the sphenoid bone that cradles and protects the pituitary.

Infundibulum
: The slender stalk that connects the pituitary to the hypothalamus above it.

Hypothalamo-hypophyseal tract
: The bundle of nerve fibers that runs from the hypothalamus down through the infundibulum to the posterior lobe.

Hypophyseal portal system
: The blood vessels that carry hypothalamic signals directly to the anterior lobe.

The two lobes

The pituitary has two lobes that develop from two different tissues in the embryo, so it is really two glands joined together. The anterior lobe, or adenohypophysis, grows up from the roof of the embryonic mouth. It is made of glandular epithelial tissue. Its cells are arranged in cords and clusters with wide capillaries running between them. The anterior lobe is the larger of the two.

The posterior lobe, or neurohypophysis, grows down from the floor of the brain. It is made of nervous tissue and is an extension of the hypothalamus itself. It contains the axons of neurons whose cell bodies sit in the hypothalamus, along with supporting glial cells called pituicytes. The posterior lobe makes no hormones of its own. It stores and releases what the hypothalamic neurons make and send down their axons.

Because the tissues differ, each lobe connects to the hypothalamus in a different way. The posterior lobe connects by nerve fibers, the hypothalamo-hypophyseal tract, which runs down through the infundibulum. The anterior lobe has no nerve link to the hypothalamus. It connects through blood vessels, the hypophyseal portal system, which has three parts in order:

1. Primary capillary plexus: a capillary bed at the base of the hypothalamus, where the infundibulum begins, picks up hypothalamic signals.
2. Hypophyseal portal veins: small veins carry the blood down along the infundibulum.
3. Secondary capillary plexus: a second capillary bed in the anterior lobe delivers the signals to the glandular cells.

Feature	Anterior lobe	Posterior lobe
Other name	Adenohypophysis	Neurohypophysis
Tissue type	Glandular epithelial tissue	Nervous tissue, an extension of the hypothalamus
Develops from	An upgrowth of the roof of the embryonic mouth	A downgrowth of the floor of the brain
Link to the hypothalamus	Blood vessels, the hypophyseal portal system	Nerve fibers, the hypothalamo-hypophyseal tract, through the infundibulum
On a slide	Densely packed gland cells, many stained pink or blue, in cords with capillaries between them	Pale, fibrous-looking tissue of axons and scattered pituicyte nuclei

- Memory tip: "neuro" means nervous tissue and "adeno" means gland. The posterior lobe is nervous tissue, so it connects by nerve fibers. The anterior lobe is glandular epithelium, so it connects by blood vessels.
- The tissue type predicts the link. Nervous tissue links by nerve fibers; glandular tissue links by blood vessels. The nerve fibers go to the posterior lobe and the blood vessels go to the anterior lobe, never the other way around.

- A pituitary tumor grows up and out of the sella turcica, because bone closes the floor and walls, and the first thing it presses on is the optic chiasm.

Competency 130. Thyroid and parathyroid anatomy

In your course notes: Module 4 packet, Chapter 6, competency 130 (page 55).

Dr. Rennie's video for this part: Endocrine system concept video, chapter Thyroid and calcium regulation.

You should be able to: Identify the thyroid lobes, isthmus, follicles, colloid, follicular and parafollicular cells, and the parathyroid chief and oxyphil cells.

The thyroid gland, gross anatomy

The thyroid gland is a butterfly-shaped gland in the anterior neck, just below the thyroid cartilage of the larynx. It has a right lobe and a left lobe. Each lobe lies against the side of the larynx and trachea. The two lobes are joined across the midline by a narrow bridge of thyroid tissue called the isthmus, which crosses in front of the upper rings of the trachea, just below the cricoid cartilage. A thin connective tissue capsule covers the gland. The thyroid is the largest gland in the body that is purely endocrine, and it has a very rich blood supply.

The thyroid gland, on the slide

Under the microscope, the thyroid is made of thousands of tiny hollow spheres called thyroid follicles. Each follicle is a ring of cells around a center filled with colloid. Colloid is a thick, protein-rich fluid. It stains pink and fills the center of each follicle.

The wall of each follicle is a single layer of follicular cells. They form a simple cuboidal epithelium that sits around the colloid. Follicular cells are the main hormone-making cells of the thyroid. Between the follicles, or tucked into the outer edge of a follicle wall, are the parafollicular cells, also called C cells. They are larger and paler than follicular cells, and they do not touch the colloid. They occur alone or in small groups.

Structure	What it is	How to find it
Right and left lobes	The two lateral parts of the gland	On each side of the larynx and trachea, below the thyroid cartilage
Isthmus	The bridge of thyroid tissue that connects the two lobes	Crossing the front of the upper trachea in the midline
Thyroid follicles	Microscopic hollow spheres that make up the gland	Rings of cells packed together across the slide
Colloid	The protein-rich fluid stored in the center of each follicle	Pink material filling each ring
Follicular cells	The simple cuboidal cells forming the wall of a follicle	The single layer of cells lining each ring of colloid
Parafollicular (C) cells	Endocrine cells between the follicles	Larger, pale cells between follicles or at the outer edge of a follicle wall, not touching the colloid

- The lobes and isthmus are things you identify on the model or the cadaver. Follicles, colloid, and C cells are too small to see by eye, so those are slide structures only.
- Colloid fills the inside of the follicle. The C cells sit between follicles or at the outer edge of the wall, and they never touch the colloid.

The parathyroid glands

The parathyroid glands are small, round glands, each about the size of a grain of rice. Most people have four, a superior and an inferior gland on the posterior surface of each thyroid lobe. Each one has its own thin capsule that separates it from the thyroid tissue. The number and exact position vary from person to person.

Parathyroid tissue has no follicles and no colloid. Its cells are packed in cords and clusters with capillaries between them. Two cell types are seen on a slide. Chief

cells (also called principal cells) are the small, numerous cells that make up most of the gland. They are the hormone-making cells of the parathyroid. Oxyphil cells are larger and much fewer. Their cytoplasm is packed with mitochondria, so it stains dark pink. Oxyphil cells become more common with age. Their function is not fully understood.

Structure	What it is	How to find it
Parathyroid glands	Usually four small, round glands on the posterior surface of the thyroid lobes	Two on the back of each lobe, each in its own capsule
Chief cells	The main hormone-making cells of the parathyroid glands	Small, numerous cells with round nuclei and pale cytoplasm, making up most of the tissue
Oxyphil cells	Larger, fewer cells whose function is not fully understood	Larger cells with dark pink, granular cytoplasm, alone or in small groups among the chief cells

- Memory tip: follicles and colloid mean thyroid. Packed cords of cells with no colloid mean parathyroid. On a parathyroid slide, the small, plentiful cells are chief cells and the big pink ones are oxyphil cells.

What this means for patients

- The parathyroid glands are tiny and embedded on the back of the thyroid, so during thyroid surgery they can be damaged or removed along with it. A surgeon has to find and spare them.
- Because the thyroid sits just under the skin of the front of the neck, an enlarged thyroid (a goiter) can be seen and felt.

Competency 131. Adrenal gland zones

In your course notes: Module 4 packet, Chapter 6, competency 131 (page 56).

Dr. Rennie's video for this part: Endocrine system concept video, chapter Major glands and their roles.

You should be able to: Identify the adrenal cortex zones in order and the adrenal medulla with its chromaffin cells.

An adrenal gland, also called a suprarenal gland, sits on the superior pole of each kidney. Both adrenal glands lie on the posterior abdominal wall behind the peritoneum, enclosed with the kidney in fat and the renal fascia. The right gland is shaped like a pyramid. The left gland is shaped like a crescent. Each gland is covered by a connective tissue capsule.

Each adrenal gland is two glands in one. The outer adrenal cortex makes up most of the gland. It is glandular epithelial tissue that develops from mesoderm. The inner adrenal medulla is a small core in the center. It is modified nervous tissue that develops from the neural crest, the same embryonic tissue that forms the sympathetic ganglia. On a slide the cortex looks like layered bands and the medulla looks like a different tissue in the middle.

Capsule
: The connective tissue covering of the gland.

Adrenal cortex
: The outer region and the bulk of the gland. Glandular epithelial tissue arranged in three zones.

Adrenal medulla
: The inner core. Modified nervous tissue that belongs to the sympathetic nervous system.

The three zones of the cortex, from the capsule inward

The cortex is built in three zones. Each zone is named for how its cells are arranged. Follow them in order, from the capsule inward to the medulla.

1. Zona glomerulosa: the thin outer zone, just under the capsule. Its cells form small rounded clusters and arches. "Glomerulus" means a little ball.
2. Zona fasciculata: the thick middle zone, the widest of the three. Its cells line up in long, straight columns running toward the center. "Fascicle" means a bundle. The cells look pale and foamy because they hold many lipid droplets.

3. Zona reticularis: the thin inner zone, against the medulla. Its cells form a branching network. "Reticulum" means a network. These cells stain darker than the fasciculata.

The adrenal medulla and chromaffin cells

The medulla is made of chromaffin cells. These are large, round to many-sided cells arranged in clusters around blood vessels. Chromaffin cells are modified sympathetic neurons. They have no axons or dendrites, and they release their product straight into the blood. They are supplied by preganglionic sympathetic nerve fibers, so the medulla works like a sympathetic ganglion. Their name comes from the brown color they turn when stained with chromium salts.

Layer	Part of	Position and thickness	Tissue type	How the cells are arranged
Capsule	Covering	Outermost, thin	Dense connective tissue	A connective tissue sheet
Zona glomerulosa	Cortex	Outer, thin	Glandular epithelial tissue	Rounded clusters and arches
Zona fasciculata	Cortex	Middle, the thickest of the three	Glandular epithelial tissue	Long, straight columns of pale, foamy cells
Zona reticularis	Cortex	Inner, against the medulla, thinner than the fasciculata	Glandular epithelial tissue	A branching network of darker cells
Adrenal medulla	Medulla	The core	Modified nervous tissue	Clusters of large chromaffin cells around blood vessels

- Memory tip: GFR, from the outside in. Glomerulosa, Fasciculata, Reticularis. Then the medulla, full of chromaffin cells, in the center.
- The middle zone is the thickest. If your drawing has the glomerulosa as the widest band, the order or the widths are wrong.
- The three cortex zones are all glandular tissue. The medulla is the only part that is nervous tissue.

What this means for patients

- The adrenal gland rests on the top of the kidney, so imaging ordered for the kidney usually shows the adrenal as well. That is how an adrenal mass is sometimes found by accident.

Competency 132. Pancreas, gonads, and organs with endocrine tissue

In your course notes: Module 4 packet, Chapter 6, competency 132 (page 58).

Dr. Rennie's video for this part: Endocrine system concept video, chapter Pancreas hormones and blood sugar.

You should be able to: Locate the pancreatic islets and gonads and name other organs such as the heart, kidneys, and skin that contain endocrine tissue.

The pancreas and its islets

The pancreas is a long, soft gland in the upper abdomen. It lies behind the stomach, against the posterior abdominal wall and mostly behind the peritoneum. Its head sits in the C-shaped curve of the duodenum. Its body crosses behind the stomach, and its tail reaches toward the spleen on the left.

The pancreas is a mixed gland, with both exocrine and endocrine tissue. Most of it is exocrine tissue. This tissue is made of acini, clusters of cells that make digestive enzymes and send them through ducts to the duodenum. Scattered among the acini are about 1 to 2 million small islands of endocrine cells called the pancreatic islets, or islets of Langerhans. The islets have no ducts. They release their hormones into the many capillaries that run through them. On a slide, an islet is a round, pale-staining patch of cells surrounded by darker-staining acini.

The gonads

The gonads are mixed organs too. They make gametes, and they also contain endocrine cells. The ovaries are the paired female gonads. They sit in the pelvic cavity, one on each side of the uterus, against the lateral pelvic wall. Their endocrine tissue is the cells of the developing ovarian follicles and the corpus luteum that forms after

ovulation. The testes are the paired male gonads. They sit in the scrotum, outside the abdominopelvic cavity. Their endocrine tissue is the interstitial cells (Leydig cells), which lie in the spaces between the seminiferous tubules.

Structure	Location	Endocrine part
Pancreas	Upper abdomen behind the stomach; head in the curve of the duodenum, tail toward the spleen	Both exocrine and endocrine; a mixed gland
Pancreatic islets	Scattered throughout the pancreas	The endocrine cell clusters (islets of Langerhans)
Acini	Most of the pancreas	None. The exocrine clusters that make digestive enzymes
Ovaries	Pelvic cavity, one on each side of the uterus	Cells of the ovarian follicles and the corpus luteum
Testes	Scrotum	Interstitial (Leydig) cells between the seminiferous tubules

- The islets are the endocrine part of the pancreas and the acini are the exocrine part. Islets are pale, acini are dark, and the islets are the small patches.
- These notes call three organs mixed organs: the pancreas, the ovaries, and the testes. An organ is mixed when it does an endocrine job alongside another role, such as making digestive enzymes or making gametes.

Other organs with endocrine tissue

Many organs whose main job is something else still contain endocrine cells. These organs are not counted as endocrine glands, and they are not called mixed organs either, but their endocrine cells still release hormones into the blood. The heart has hormone-making cells in the muscle of the atrial walls. The kidneys have endocrine cells in the renal cortex. The skin takes part too: cells of the epidermis begin making a hormone when the skin is exposed to sunlight. The stomach and small intestine, the liver, adipose tissue, and the placenta during pregnancy also contain endocrine cells.

Organ	Where it sits	Where the endocrine cells are
Hypothalamus	Brain, floor and walls of the third ventricle	Neurosecretory cells. A neuroendocrine organ that controls the pituitary gland
Heart	Chest, in the mediastinum	Cardiac muscle cells in the walls of the atria
Kidneys	Abdomen, against the posterior wall	Endocrine cells in the renal cortex, including the juxtaglomerular cells in the walls of the afferent arterioles
Skin	Body surface	Keratinocytes of the epidermis, which begin making a hormone when exposed to sunlight
Stomach and small intestine	Abdomen	Enteroendocrine cells scattered in the mucosa
Liver	Upper right abdomen	Hepatocytes, the main liver cells
Adipose tissue	Under the skin and around organs	Adipocytes (fat cells)
Placenta	In the uterus during pregnancy	Placental cells. A temporary organ, because it exists only during pregnancy

- The heart, kidneys, and skin contain endocrine tissue, but they are not endocrine glands and these notes do not count them as mixed organs. Only the pancreas, ovaries, and testes get that label.
- The placenta is the one temporary organ on the list. It forms during pregnancy and is gone afterward.

Worked questions

Nine questions in the style of the lecture exam, at least one for each competency. Read the answer and the reasons each wrong option is wrong. The wrong options are where the learning is.

QUESTION 1 · COMPETENCY 128

Which endocrine gland sits in the sella turcica of the sphenoid bone?

A. Pineal gland

B. Hypothalamus

C. Pituitary gland

D. Thyroid gland

Answer: C. The pituitary is the pea-sized gland that sits in the sella turcica, the saddle-shaped pocket of the sphenoid bone, hanging below the hypothalamus by the infundibulum.

Why the others are wrong

- **A** The pineal gland is in the epithalamus, projecting from the posterior roof of the third ventricle. It is in the brain, not in a pocket of bone.
- **B** The hypothalamus is the floor and lower walls of the third ventricle, above the pituitary. It is brain tissue, not a gland in the sella.
- **D** The thyroid is in the anterior neck, below the larynx, with its lobes beside the trachea.

QUESTION 2 · COMPETENCY 128

Which gland is the largest purely endocrine gland in the body?

A. Thyroid gland

B. Pancreas

C. Thymus

D. Adrenal gland

Answer: A. The thyroid, the butterfly-shaped gland in the front of the neck, is the largest gland that is entirely endocrine.

Why the others are wrong

- **B** The pancreas is larger, but it is a mixed gland. Most of it is exocrine acini, and only the scattered islets are endocrine.
- **C** The thymus in the mediastinum is large in childhood but shrinks after puberty. It is not the largest purely endocrine gland.
- **D** Each adrenal gland is a small gland capping the top of a kidney, much smaller than the thyroid.

QUESTION 3 · COMPETENCY 129

The posterior lobe of the pituitary (neurohypophysis) is made of which tissue?

A. Glandular epithelial tissue

B. Dense connective tissue

C. Simple cuboidal epithelium

D. Nervous tissue

Answer: D. The posterior lobe grows down from the floor of the brain and is an extension of the hypothalamus. It holds the axons of hypothalamic neurons and supporting pituicytes. "Neuro" means nervous tissue.

Why the others are wrong

- **A** Glandular epithelial tissue is the anterior lobe, the adenohypophysis, which grows up from the roof of the embryonic mouth.
- **B** Dense connective tissue forms capsules such as the one around the adrenal gland, not a pituitary lobe.
- **C** Simple cuboidal epithelium is the follicular cell layer of the thyroid follicle, not pituitary tissue.

QUESTION 4 · COMPETENCY 129

How does the anterior lobe of the pituitary receive its signals from the hypothalamus?

A. Through the hypothalamo-hypophyseal tract of nerve fibers

B. Through the blood vessels of the hypophyseal portal system

C. Through the axons of pituicytes

D. Through the optic chiasm

Answer: B. The anterior lobe is glandular epithelium with no nerve link to the hypothalamus. Hypothalamic signals enter the primary capillary plexus, travel down the hypophyseal portal veins along the infundibulum, and reach the gland cells at the secondary capillary plexus.

Why the others are wrong

- **A** The hypothalamo-hypophyseal tract is the nerve fiber bundle to the posterior lobe, the nervous-tissue lobe.
- **C** Pituicytes are the supporting glial cells of the posterior lobe. They do not carry signals to the anterior lobe.
- **D** The optic chiasm lies just above and in front of the pituitary. A pituitary tumor presses on it, but it carries no signals to the gland.

QUESTION 5 · COMPETENCY 130

On a thyroid slide, where is the colloid?

A. Filling the center of each follicle

B. Between the follicles

C. In the cytoplasm of the parafollicular cells

D. In the connective tissue capsule

Answer: A. Each thyroid follicle is a ring of follicular cells around a center filled with colloid, the thick, protein-rich fluid that stains pink.

Why the others are wrong

- **B** Between the follicles are the parafollicular (C) cells and capillaries, not colloid. The C cells never touch the colloid.
- **C** Parafollicular cells are larger, paler cells that sit outside the follicle wall. Colloid is stored inside the follicle, not inside any cell.
- **D** The capsule is a thin connective tissue covering on the outside of the gland.

QUESTION 6 · COMPETENCY 130

Which cells make up most of a parathyroid gland and are its hormone-making cells?

A. Follicular cells

B. Oxyphil cells

C. Chief cells

D. Parafollicular cells

Answer: C. Chief cells (principal cells) are the small, numerous cells with round nuclei and pale cytoplasm that make up most of the parathyroid tissue, and they are the gland's hormone-making cells.

Why the others are wrong

- **A** Follicular cells form the wall of a thyroid follicle. They are thyroid cells, not parathyroid cells.
- **B** Oxyphil cells are the larger, fewer parathyroid cells with dark pink cytoplasm. They become more common with age and their function is not fully understood.
- **D** Parafollicular (C) cells are thyroid cells sitting between the follicles.

QUESTION 7 · COMPETENCY 131

Which list gives the zones of the adrenal cortex in order from the capsule inward?

A. Zona reticularis, zona fasciculata, zona glomerulosa

B. Zona glomerulosa, zona fasciculata, zona reticularis

C. Zona fasciculata, zona glomerulosa, zona reticularis

D. Zona glomerulosa, zona reticularis, zona fasciculata

Answer: B. GFR from the outside in: the thin glomerulosa under the capsule, the thick fasciculata in the middle, and the thin reticularis against the medulla.

Why the others are wrong

- **A** That is the order from the medulla outward, backwards from the capsule.
- **C** The glomerulosa is the outer zone, just under the capsule. The fasciculata is the middle zone, not the outer one.
- **D** The reticularis is the inner zone against the medulla. The fasciculata sits between the glomerulosa and the reticularis.

QUESTION 8 · COMPETENCY 131

Chromaffin cells are found in which part of the adrenal gland?

A. Zona fasciculata

B. Capsule

C. Zona glomerulosa

D. Adrenal medulla

Answer: D. The medulla, the core of the gland, is made of chromaffin cells. They are modified sympathetic neurons with no axons or dendrites, clustered around blood vessels, and they release their product straight into the blood.

Why the others are wrong

- **A** The zona fasciculata is the thick middle cortex zone, made of pale, foamy gland cells in straight columns.
- **B** The capsule is dense connective tissue covering the outside of the gland.
- **C** The zona glomerulosa is the thin outer cortex zone, made of gland cells in rounded clusters and arches.

QUESTION 9 · COMPETENCY 132

Which organ contains endocrine tissue but is not counted as an endocrine gland or a mixed organ in these notes?

A. Pancreas

B. Heart

C. Testis

D. Thyroid gland

Answer: B. The heart's main job is pumping blood, but cardiac muscle cells in the walls of the atria also release a hormone. It contains endocrine tissue without being an endocrine gland, and these notes do not call it a mixed organ.

Why the others are wrong

- **A** The pancreas is one of the three mixed organs in these notes: exocrine acini plus endocrine islets.
- **C** The testis is a mixed organ too. It makes gametes and has endocrine interstitial (Leydig) cells between the seminiferous tubules.
- **D** The thyroid is a purely endocrine gland, the largest one in the body.