# nebula3d

Manual

3D Reciprocal-Space Diffuse Neutron Scattering Analysis
Powder Ring Removal · Bragg Punch & Fill · 3D-ΔPDF Transform

Version 0.4.0

| | |
|---|---|
| **Language** | Python ≥ 3.10 |
| **License** | AGPL-3.0-or-later |
| **Dependencies** | numpy, scipy, h5py, matplotlib |
| **Instrument** | Mantid NeXus (MDHistoWorkspace) |
| **Reference data** | Mantid NeXus HKL volumes |

**Tsung-Han Yang**
© 2026 Tsung-Han Yang. All rights reserved.

# Table of Contents

**1. Introduction** 4
- 1.1 Scope of This Manual 4
- 1.2 What nebula3d Does Not Do 4

**2. Background Theory** 5
- 2.1 Neutron Scattering Fundamentals 5
- 2.2 The 3D Difference Pair Distribution Function 5
- 2.3 Coordinate Systems and Conventions 5
- 2.4 Powder Rings: Physical Origin 6

**3. Package Architecture** 7
- 3.1 Overview 7
- 3.2 The HKLVolume Data Structure 7
- 3.3 File Formats 7

**4. Algorithms** 9
- 4.1 Stage 1: Powder Ring Removal 9
- 4.2 Stage 2: Bragg Peak Punching 10
- 4.3 Stage 3: Bragg-Hole Backfill 11
- 4.4 Stage 4: 3D-ΔPDF Transform 13

**5. Installation and Setup** 15
- 5.1 Requirements 15
- 5.2 Install from Source 15
- 5.3 Recommended Runtime Environment 15
- 5.4 Verifying the Installation 15

**6. End-to-End Workflow** 16
- 6.1 One-Command Pipeline 16
- 6.2 Stage-by-Stage Commands 16
- 6.3 Multi-Volume Workflow 17
- 6.4 Python API 17

**7. Configuration Reference** 19
- 7.1 Ring Removal Parameters 19
- 7.2 Bragg Punch Parameters 19
- 7.3 Backfill Parameters 20
- 7.4 3D-ΔPDF Transform Parameters 20

**8. Visualization and Interactive Exploration** 21
- 8.1 Visualization API Overview 21
- 8.2 plot_slice() 21
- 8.3 plot_radial_profile() and plot_azimuthal_map() 21
- 8.4 plot_overview() 21
- 8.5 Interactive Viewers 22

**9. Worked Examples** 23
- 9.1 Quick Inspection of a New Dataset 23
- 9.2 Locating and Profiling a Powder Ring 23

9.3 Checking Bragg Punch Quality 23
9.4 Computing and Displaying the 3D-ΔPDF 24
**10. Testing** 25
10.1 Running the Test Suite 25
10.2 Key Regression Tests 25
**11. Known Limitations and Troubleshooting** 26
11.1 Near-Origin Spike 26
11.2 Axis Cross in ΔPDF 26
11.3 Residual Ring After Subtraction 26
11.4 Search Mode Punching Diffuse Structure 26
11.5 Rank-1 SVD Failure in Ring Model 26
11.6 Performance Notes 26
**12. Glossary** 28
**13. References** 29

# 1. Introduction

**Where the rest is documented.** This manual covers the Python package. The browser console (in-browser and native) and the NEBULA Pilot agent are documented in docs/web.md and docs/pilot.md, and the pipeline's current stages in docs/pipeline.md, at github.com/drthyang/nebula3d/tree/main/docs.

**nebula3d** is a Python 3.10+ toolkit for the complete analysis of three-dimensional (3D) diffuse neutron scattering volumes. It implements a production-ready four-stage pipeline that transforms raw single-crystal Mantid NeXus output into real-space 3D difference pair distribution functions (3D-ΔPDF):

**Stage 1 — Powder-ring removal.** Subtracts azimuthally smooth ring contamination from the cryostat, sample holder, and sample environment without masking real diffuse signal.

**Stage 2 — Bragg punch.** Identifies and masks sharp Bragg and satellite peaks using lattice-aware integer-node detection, data-driven shaping, and an hkl-agnostic search mode for off-integer satellites.

**Stage 3 — Backfill.** Replaces masked voxels with physically reasonable estimates derived from the surrounding radial background or crystal symmetry equivalents.

**Stage 4 — 3D-ΔPDF transform.** Converts the cleaned reciprocal-space volume to a real-space pair-correlation map via a correctly centred 3D discrete Fourier transform with apodization and smooth-background subtraction.

The package is designed around a single core data structure (HKLVolume) that carries the 3D intensity array, per-voxel standard deviations, a validity mask, HKL coordinate axes, and the crystal orientation (UB) matrix. All pipeline stages operate on this type, making it straightforward to inspect intermediate results, apply individual stages, or extend the pipeline with custom algorithms.

Visualization is provided through a thin, scriptable layer of Matplotlib primitives: 2D plane slices, radial profiles, azimuthal ring-texture maps, a 2×2 overview diagnostic, and three interactive orthogonal-cut real-space viewers for quality assurance at every stage.

## 1.1 Scope of This Manual

This manual covers the background physics and mathematics of 3D diffuse neutron scattering analysis, the detailed algorithms and their theoretical basis, a step-by-step workflow guide, the full Python API, configuration parameters, generic worked examples for single-volume and multi-volume workflows, interactive visualization, known limitations, and a complete reference list.

## 1.2 What nebula3d Does Not Do

The package does not perform raw detector reduction, normalization to an absolute cross-section, or Reverse Monte Carlo (RMC) structural refinement. It assumes the input is a Mantid background-subtracted MDHistoWorkspace volume (*_cc_sub_bkg.nxs). For the reduction steps that precede this, use Mantid directly.

# 2. Background Theory

## 2.1 Neutron Scattering Fundamentals

In a neutron scattering experiment a monochromatic beam of neutrons with wavevector $\mathbf{k}_i$ is scattered by the sample. Neutrons with final wavevector $\mathbf{k}_f$ are detected as a function of the momentum transfer

$$\mathbf{Q} = \mathbf{k}_f - \mathbf{k}_i \quad (|\mathbf{Q}| = Q = 4\pi \sin\theta / \lambda)$$

For an elastic measurement ($|\mathbf{k}_i| = |\mathbf{k}_f|$) on a single crystal, the scattered intensity $I(\mathbf{Q})$ is recorded over a 3D grid in reciprocal space indexed by the fractional Miller indices ($h$, $k$, $l$).

The total measured intensity separates into two components:

$$I_{total}(\mathbf{Q}) = I_{Bragg}(\mathbf{Q}) + I_{diffuse}(\mathbf{Q})$$

The **Bragg contribution** $I_{Bragg}$ consists of sharp peaks at integer lattice nodes (and satellite peaks for modulated structures). It encodes the average periodic structure. The **diffuse contribution** $I_{diffuse}$ is the remaining broad, continuous signal that encodes deviations from the average: atomic displacements, occupancy disorder, short-range magnetic correlations, and phonon/magnon scattering.

## 2.2 The 3D Difference Pair Distribution Function

The three-dimensional difference pair distribution function (3D-ΔPDF) is defined as the Fourier transform of the diffuse scattering intensity:

$$\Delta\rho(\mathbf{r}) = FT[\, I_{diffuse}(\mathbf{Q}) \,] = FT[\, I_{total}(\mathbf{Q}) - I_{Bragg}(\mathbf{Q}) \,]$$

In real space, $\Delta\rho(\mathbf{r})$ is a three-dimensional map of interatomic correlation deviations. A positive value at vector $\mathbf{r}$ means there are *more* interatomic pairs separated by $\mathbf{r}$ than the average structure predicts; a negative value means *fewer*. Peaks in $\Delta\rho(\mathbf{r})$ at specific real-space positions directly identify the geometry of correlated displacements or ordered nano-domains.

The key practical relation connecting measurement to analysis is:

$$\Delta\rho(\mathbf{r}) \approx FT[\, I_{total}(\mathbf{Q}) - I_{Bragg}(\mathbf{Q}) \,] = FT[\, I_{cleaned}(\mathbf{Q}) \,]$$

where $I_{cleaned}(\mathbf{Q})$ is the diffuse volume after ring removal, Bragg punching, and backfill. The accuracy of $\Delta\rho$ therefore depends directly on the quality of each preprocessing stage.

> **Key Reference**
> Weber & Simonov, Z. Kristallogr. 227, 238–247 (2012), establish the 3D-ΔPDF formalism; Kobas, Weber & Steurer, Phys. Rev. B 71, 224205 (2005), introduced the punch-and-fill reduction; Simonov, Weber & Steurer, J. Appl. Cryst. 47, 2011–2018 (2014), quantify its experimental uncertainties.

## 2.3 Coordinate Systems and Conventions

Two coordinate systems are used throughout the package:

- **Fractional HKL.** Reciprocal-lattice coordinates ($h$, $k$, $l$) measured in units of the reciprocal basis vectors **a***, **b***, **c***. Integer values correspond to Bragg positions; fractional values index the continuous diffuse volume.
- **Cartesian Q.** Physical momentum transfer in units of Å■¹, related to HKL by the UB matrix: **Q** = UB · [$h$, $k$, $l$]■. The magnitude $|\mathbf{Q}| = 2\pi/d$ (physics convention, as opposed to the crystallographic $1/d$ used in Mantid).

The **UB matrix** combines the orientation matrix U (aligning the crystal axes to the laboratory frame) and the reciprocal-metric matrix B (encoding the lattice parameters). Mantid files store the crystallographic UB (no 2π factor); nebula3d reads this and multiplies by 2π to convert to the physics convention.

$$\mathbf{Q} = 2\pi \cdot \mathrm{UB}_{\mathrm{cryst}} \cdot [h, k, l] \quad (\text{Å}^{-1})$$

## 2.4 Powder Rings: Physical Origin

Polycrystalline material in the beam path — the sample cryostat, aluminium heat shields, sample holder, and sample capsule walls — produces *powder diffraction rings*: bands of elevated intensity at fixed $|\mathbf{Q}|$ values corresponding to the *d*-spacings of the polycrystalline material. For aluminium (FCC, $a$ = 4.046 Å), the strongest ring is Al(111) at $|\mathbf{Q}|$ = 2.69 Å$^{-1}$.

Rings are not isotropic. Detector solid-angle coverage, absorption path length, and normalization artifacts modulate the ring intensity with azimuthal direction $\phi$. A useful physical model for the ring contribution at a voxel with scattering vector magnitude $|\mathbf{Q}|$ and azimuthal angle $\phi$ is:

$$I_{\mathrm{ring}}(Q, \phi) = T(\phi) \times \Sigma_i A_i \cdot G(|Q| - q_i, \sigma_i)$$

where $G(|\mathbf{Q}|-q_i, \sigma_i)$ is a Gaussian radial profile for ring $i$, $A_i$ is the per-ring amplitude, and $T(\phi)$ is the shared azimuthal texture function. The measured signal is:

$$I_{\mathrm{measured}}(Q, \phi) = I_{\mathrm{diffuse}}(Q) + I_{\mathrm{ring}}(Q, \phi)$$

Since the diffuse signal of interest does *not* share the ring's radial peak structure or azimuthal texture, it is in principle separable from the ring contribution.

# 3. Package Architecture

## 3.1 Overview

The package is installed from source under src/nebula3d/ and follows a strict separation between the core data structure, I/O, the four algorithmic stages, and visualization:

| Module | Responsibility |
|---|---|
| nebula3d/core.py | HKLVolume dataclass — the universal data carrier |
| nebula3d/io/ | Mantid NeXus reader, HDF5 load/save, ASCII HKL I/O |
| nebula3d/preprocessing/ | Powder-ring models, radial background, backfill |
| nebula3d/analysis/ | Bragg punch/backfill, 3D-ΔPDF FFT |
| nebula3d/visualization/ | Slice plots, radial profiles, interactive viewers |
| nebula3d/utils/ | UB matrix, d-spacing, Q↔HKL conversion utilities |
| examples/ | Pipeline scripts and interactive viewer entry points |

## 3.2 The HKLVolume Data Structure

Every stage of the pipeline operates on an HKLVolume dataclass defined in src/nebula3d/core.py:

```
@dataclass
class HKLVolume:
data : NDArray[np.float64] # shape (nh, nk, nl) intensity
sigma : NDArray[np.float64] # shape (nh, nk, nl) standard deviation
mask : NDArray[np.bool_] # True = valid (not masked)
h_axis : NDArray # 1D fractional-HKL H coordinates
k_axis : NDArray # 1D fractional-HKL K coordinates
l_axis : NDArray # 1D fractional-HKL L coordinates
ub_matrix : NDArray[np.float64]# (3, 3) physics-convention UB (Angstrom^-1)
instrument: str # provenance label
```

Key methods provided by HKLVolume:

| Method | Description |
|---|---|
| `hkl_grid()` | Returns (H, K, L) meshgrids of shape (nh, nk, nl) |
| `q_magnitude()` | Returns \|Q\| in Å■¹ for every voxel |
| `masked_data()` | Returns data with masked voxels set to NaN |
| `apply_mask(m)` | Updates the mask in place |

## 3.3 File Formats

### *Input: Mantid NeXus*

Raw input is a Mantid MDHistoWorkspace saved as a NeXus file (*_cc_sub_bkg.nxs). The reader (nebula3d/io/mantid_nxs.py) extracts the signal, variance, mask, bin-edge arrays, and the UB matrix from the standard Mantid HDF5 hierarchy. The Mantid crystallographic UB is rescaled by 2π on read.

### *Native HDF5*

All intermediate and output files are stored as HDF5 (.h5). Load with vol = nebula3d.load(path); save with nebula3d.save(vol, path). The ΔPDF output additionally stores the transform configuration and direct-lattice constants as HDF5 attributes for use by the interactive viewers.

# 4. Algorithms

## 4.1 Stage 1: Powder Ring Removal

### *4.1.1 Design Principle*

Ring removal is strictly *subtractive*. The algorithm estimates only the azimuthally smooth ring intensity and subtracts it. It never masks or replaces voxels solely because they exhibit radial excess, because radial excess can be genuine anisotropic diffuse scattering.

### *4.1.2 Non-Parametric Patch-Based Method (Production)*

The production algorithm is implemented in nebula3d/preprocessing/radial_background.py as PatchedRadialRingModel. It processes the volume one H-slice (0kl plane) at a time and builds a non-parametric radial ring model in azimuthal patches:

**1. Azimuthal patches.** The reference plane (default hk0) defines the azimuthal angle $\phi$ = atan2(k_Q, h_Q). The full range $\phi \in [0, 2\pi)$ is divided into N overlapping patches, each weighted by a Hann window to ensure smooth blending across patch boundaries.

**2. Robust radial profile.** Within each azimuthal patch, voxels are binned by |Q| and a trimmed-mean profile is computed. The trimmed mean rejects high-tail outliers (Bragg peaks) and low-tail outliers (detector gaps), yielding a clean estimate of the combined diffuse + ring signal in each bin.

**3. SNIP baseline estimation.** A smooth diffuse baseline is estimated from the robust profile using a morphological opening: successive minimum and maximum filters with a kernel wider than the expected ring width. Light polynomial smoothing is applied. This is equivalent to the Sensitive Nonlinear Iterative Peak (SNIP) clipping algorithm common in nuclear spectroscopy.

**4. Ring component.** The ring component for each patch is: ring(|Q|) = max(0, prof – base). Taking the positive part ensures no diffuse signal is subtracted when the profile lies below the baseline (noise fluctuations).

**5. Subtraction.** Each voxel receives a Hann-weighted blend of the ring estimates from its two nearest azimuthal patches, interpolated to the voxel's exact |Q|. Cross-H confirmed ring shells and amplitude ceilings from adjacent H planes are propagated to prevent integer-H Bragg artifacts from masquerading as powder rings in nearby planes.

### *4.1.3 Factored Gaussian/SVD Model (Legacy)*

The older algorithm (PatchedRingModel) remains in the package and is useful background. It fits a rank-1 factored model to the amplitude matrix:

$$A[i, P] \approx A_i \times T[P]$$

where index *i* runs over rings and *P* over azimuthal patches. A rank-1 SVD extracts per-ring amplitudes $A_i$ and patch textures T[P]. A Fourier series $T(\phi)$ is then fitted to give a smooth, periodic texture function. The quality of the rank-1 approximation is monitored via rank1_variance; values ≥ 0.90 confirm the shared-texture assumption.

### 4.1.4 Diagnostic Metrics

| Diagnostic | Interpretation |
|---|---|
| `rank1_variance` | Fraction of amplitude-matrix variance explained by rank-1 SVD. ≥ 0.90 confirms shared T(ϕ). Lower values suggest per-ring texture fitting is needed. |
| Per-patch counts | Sparse patches receive lower weight; uniform coverage is ideal. |
| Raw vs baseline profiles | Plot with plot_radial_profile() to visually verify ring isolation. |

## 4.2 Stage 2: Bragg Peak Punching

Bragg punching is implemented in nebula3d/analysis/bragg.py by the BraggRemover class. The recommended production mode is mode="both", which combines a lattice-aware integer-node pass followed by an hkl-agnostic search pass.

### 4.2.1 Integer-Node Path

The integer-node path exploits the known lattice geometry:

- **Enumerate.** All integer (h, k, l) nodes within the HKL volume extent are listed.
- **Detect.** A local HKL window of half-width detect_window_hkl is inspected around each node. By default (integer_detect="significance") the node is retained when the excess at the window's brightest voxel over its own background, the median of a shell between 1× and 2× the resolution ellipsoid, reaches min_significance standard errors, corrected for picking the brightest of the window's voxels. Nothing in this test depends on the intensity units, so X-ray rates and neutron counts are judged alike, and a broad diffuse maximum at a node, no higher than its own surroundings, is left alone. With integer_max_shell_fraction (pipeline 0.15) a node is also rejected when it is resolvably broader than a Bragg peak: its 1×–2× shell still holds more than that fraction of its core's excess over a 3×–4× shell. A punch sized for a peak would take the core of such a maximum and leave its skirt. integer_detect="floors" keeps the earlier test: the peak exceeds min_intensity and the local median by min_prominence, in data units.
- **Recentre.** The punch centre is moved to the measured local maximum within the detection window.
- **Fit shape (optional).** If integer_optimize_shape=True, the covariance of the peak core (the connected voxels above a threshold fraction of the peak height) is computed in Q and corrected for the cut, giving a tilted ellipsoid along the peak's own principal axes. It always contains the resolution ellipsoid of the punch frame and is capped at max_radius_scale times it; peaks too weak to measure get the resolution ellipsoid.
- **Guard (H-slab).** The parameter integer_h_guard_hkl clips each integer-node punch to a slab of half-width $H_{guard}$ centred on the integer-H plane. This prevents strong Bragg holes at H=0 from extending into the diffuse planes at H=±1/3 or H=±2/3.

### 4.2.2 Weak Bragg at Integer Nodes

With the floors test (integer_detect="floors"), weak peaks at integer nodes can fall below the absolute min_intensity and min_prominence floors but still be sharp local outliers. The parameter integer_local_prominence_n_mad catches these by requiring the prominence to exceed a multiple of the local Median Absolute Deviation (MAD) within the detection window. Because this criterion is *locked to integer lattice nodes* (never applied at fractional positions), it cannot punch the q=1/3 diffuse planes. An additional small absolute floor integer_local_min_prominence avoids false positives in very flat regions.

### 4.2.3 Search Path (HKL-Agnostic)

The search path detects off-integer satellites and any sharp feature missed by the integer pass. At each |**Q**| shell, a robust background level is estimated as:

```
bg_threshold = median(I_shell) + n_mad × MAD(I_shell)
```

Local maxima above this threshold and the floors search_min_intensity and search_min_prominence are retained as punch centres. The pipeline gives the floors in units of the diffuse scatter (search_floor_unit="scatter": the median over |**Q**| shells of each shell's 1.4826·MAD), so they follow the data's intensity scale. Because the search does not know the crystal lattice, structured diffuse planes must be explicitly protected:

- **Explicit exclusion.** A list of H-plane centres (search_exclude_h_centers) defines fixed exclusion slabs of half-width search_exclude_h_half_width.
- **Periodic exclusion (preferred).** search_exclude_h_fractions defines fractional parts modulo 1. For example, fractions [0.3333, 0.6667] protect every H = n ± 1/3 and n ± 2/3 plane across the full H range, covering higher-order satellites at H = ±4/3, ±5/3, … that a fixed-centre list would miss.

### 4.2.4 Punch Shape

Each identified peak is punched as an anisotropic ellipsoid in HKL space:

```
(h − h_0)^2/r_h^2 + (k − k_0)^2/r_k^2 + (l − l_0)^2/r_l^2 ≤ 1
```

where ($h_0$, $k_0$, $l_0$) is the fitted punch centre and ($r_h$, $r_k$, $r_l$) are the HKL semi-axes. A guard margin is added to all radii. Optionally, the radii are scaled by the cube root of the peak intensity (relative to a reference), capped at max_radius_scale.

### 4.2.5 Direct Beam

The direct beam at **Q** = 0 is much broader than ordinary Bragg peaks and is treated separately. It is punched after all other peaks, using independent radii specified by incident_beam_ellipsoid_radii_hkl. The backfill of the direct-beam hole uses a specially shifted radial shell just outside the punch boundary, preventing the over-subtraction halo from contaminating the fill.

## 4.3 Stage 3: Bragg-Hole Backfill

After punching, masked voxels must be replaced with physically reasonable estimates before the Fourier transform. The dedicated wrapper is backfill_bragg() in nebula3d/analysis/bragg_fill.py.

Each hole is filled from the diffuse intensity *around* it (the 3D-ΔPDF punch-and-fill convention). Every punch sits on a reciprocal-lattice node, so any bias of the fill at the nodes repeats on the lattice and Fourier-transforms into spurious ΔPDF features at the lattice vectors, where the real correlations are.

### 4.3.1 Local Fill (Default)

method="local" fills each connected punched component from a dilated-shell median of nearby valid voxels, falling back to the global median if too few neighbours are available. It is robust to Bragg tails leaking past the punch edge; the fill is flat, leaving a small step at the hole edge.

### 4.3.2 Laplace Fill

method="laplace" solves the discrete Laplace equation (6-neighbour stencil) in every hole, so the fill continues the surrounding diffuse smoothly with no edge step. All holes form one sparse block-diagonal system solved by Jacobi-preconditioned conjugate gradients. The Dirichlet boundary sits laplace_gap (default 1) voxels outside the punch so leaked Bragg tails do not pull the fill up; the band is filled too, so the fill meets the kept data at its boundary with no step.

### 4.3.3 Q-Shell Fill (Comparison Only)

method="q_shell" fills each hole with the median of all valid voxels in its $|\mathbf{Q}|$ bin (step q_shell_step). It ignores the diffuse structure at the lattice nodes (correlations at lattice-vector separations peak or dip exactly there), so it is biased at every node and leaves lattice-vector artefacts in the $\Delta$PDF. Kept for comparison only.

### 4.3.3 Removed: generic image inpainting

Total-variation (Chambolle–Pock), Laue-symmetry copying, RBF and biharmonic inpainting were removed. They carry no model of diffuse scattering: TV assumes a piecewise-constant image and leaves staircase artefacts, and every symmetry copy of a punched Bragg node is itself punched. Coverage gaps are best filled by symmetrising the data before it reaches the pipeline.

## 4.4 Stage 4: 3D-ΔPDF Transform

### 4.4.1 Correct FFT Recipe

The cleaned volume stores **Q**=0 at the *array centre* (index n//2), but NumPy's fftn expects the origin at index [0,0,0]. A correct, centred Fourier transform of the real, centrosymmetric I(**Q**) must therefore be:

$$\Delta\rho = \texttt{fftshift( fftn( ifftshift( I_windowed ) ) ).real}$$

Applied step by step in nebula3d/analysis/delta_pdf.py:

- **Step 1: Fill masked voxels.** Replace NaN (masked) voxels with 0. The backfilled volume should already be NaN-free; this is a safety guard.
- **Step 2: Optional Q-space crop.** Symmetrically crop to |H| ≤ h_max, |K| ≤ k_max, |L| ≤ l_max, keeping the array centred on Q=0.
- **Step 3: Subtract smooth background.** Compute I_bg = GaussianBlur(I, σ) with σ ≈ 1.5 r.l.u. and subtract: I_new = I – I_bg. This removes the broad separable diffuse envelope that would otherwise produce an axis cross in real space (see Section 4.4.3).
- **Step 4: Apodize.** Multiply by a separable window W(H,K,L) = w(H)×w(K)×w(L). The Hann window w(x) = cos²(πx/(2x_max)) (default) suppresses termination ripples from the finite |Q| range. Alternatives: Gaussian with width gaussian_sigma × Q_max, or no window.
- **Step 5: Remove DC.** Subtract the mean *after* windowing: I_w –= mean(I_w). This ensures Σ I = 0 exactly, zeroing the r=0 self-correlation spike. (Subtracting before windowing leaves a nonzero windowed sum, creating a spurious large peak at r=0.)
- **Step 6: Zero-pad symmetrically.** Pad to the next fast FFT length (5-smooth) in each dimension, keeping Q=0 on the new centre. One-sided padding shifts the origin and breaks the ifftshift below. Zero-padding provides sinc-interpolation of the real-space grid; it does not increase intrinsic resolution, which is set by the Q_max and apodization window.
- **Step 7: Transform.** ifftshift → fftn → fftshift: move Q=0 to the corner, transform, then recentre r=0.
- **Step 8: Take real part.** The transform of centrosymmetric data I(Q)=I(–Q) is real; the imaginary part is numerical noise and a useful diagnostic.

### 4.4.2 Real-Space Axes

The real-space coordinate along axis *a* is computed from the FFT frequency grid and the UB matrix:

$$\Delta\texttt{h} = \texttt{(h_max – h_min) / (n_h – 1)}$$

$$\texttt{freq_h = fftshift( fftfreq(n_h_padded, d=}\Delta\texttt{h) ) [cycles per HKL unit]}$$

$$\texttt{x_a = freq_h} \times 2\pi \texttt{ / |UB[:,0]| [Å]}$$

The factor 2π/|**a***| converts cycles per HKL unit to Å in direct space, where **a*** = UB[:,0] is the first column of the UB matrix (the reciprocal basis vector along H).

### 4.4.3 The Axis-Cross Artifact and Its Removal

A common artifact in 3D-ΔPDF maps is a bright cross along the $y_K$=0 and $z_L$=0 axes in the $y_K$–$z_L$ plane. Diagnosis showed:

- The cross is present on planes with *no* Bragg peaks (e.g. H=1/3).
- The input has 0% masked voxels along the axis lines.
- Replacing the exact K=0 or L=0 input lines with neighbour averages has no effect.

The root cause is a broad, slowly-varying *diffuse envelope* centred near K=L=0. This envelope is approximately separable ($I \approx f(K) + g(L)$), and the Fourier transform of a separable function concentrates on the two principal axes. The Hann window, itself a centred separable hump, multiplies this effect. Subtraction of the scalar mean only removes the DC term; it leaves the envelope shape untouched.

The fix is smooth-background subtraction before windowing. Using a Gaussian blur with $\sigma \approx 1.5$ r.l.u. (per-axis control available):

$$\mathbf{I_new = I - GaussianBlur(I, \sigma)}$$

This removes the separable envelope while preserving the oscillatory modulation (the genuine diffuse structure). As with all ΔPDF background subtractions, genuine very-long-period / low-*r* correlations at the same length scale as the background cannot be cleanly separated and are also attenuated.

> **Parameter recommendation**
> Use SUBTRACT_BG="0,1.5,1.5" to apply a per-axis Gaussian blur with σ_H=0 (slice-wise background, preserves H-layering), σ_K=1.5, σ_L=1.5 r.l.u. Tune the blur widths for the length scale of the smooth background in your volume.

### *4.4.4 Near-Origin Spike*

A strong feature at r < ~3 Å remains in all 3D-ΔPDF maps. It originates from residual high-|Q| Bragg leakage, discontinuities at the punch-hole boundaries, and the direct-beam punch. Colour scales are therefore set from the p99 of |Δρ| at r > 3 Å so this near-origin spike does not dominate the display.

### *4.4.5 The Centring Bug (Fixed 2026-06-05)*

Earlier code computed fftshift(fftn(data)) without ifftshift and used one-sided zero-padding. With Q=0 at the array centre, the missing ifftshift introduces a linear phase ramp $e^{-i\pi k} = (-1)^k$ across the output. Taking the real part then *flips the sign of real-space features by pixel parity*, so each correlation peak splits into mixed positive/negative lobes. This has been fixed in all current code; the regression test test_delta_pdf_centring_positive_peak guards against reintroduction.

# 5. Installation and Setup

## 5.1 Requirements

| Package | Version | Purpose |
|---|---|---|
| Python | ≥ 3.10 | Core language |
| numpy | ≥ 1.24 | Array operations, FFT |
| scipy | ≥ 1.10 | Interpolation, Gaussian filters |
| h5py | ≥ 3.8 | HDF5 file I/O |
| matplotlib | ≥ 3.7 | Visualization |

## 5.2 Install from Source

```
git clone https://github.com/user/nebula3d
cd nebula3d
pip install -e ".[dev]"
```

The [dev] extra installs pytest, ruff, and mypy for the test suite and linters.

## 5.3 Recommended Runtime Environment

The recommended environment uses the sci-general conda environment with all dependencies pre-installed. Set the following shell variables before running any script:

```
export PY=/opt/homebrew/Caskroom/miniforge/base/envs/sci-general/bin/python
export PYTHONPATH=src
export MPLCONFIGDIR=/private/tmp/nebula3d-mpl
```

MPLCONFIGDIR keeps Matplotlib cache files outside the repository. If your Python 3.10+ environment already has the dependencies active, replace $PY with python3.

## 5.4 Verifying the Installation

```
PYTHONPATH=src python -c "import nebula3d; print(nebula3d.__version__)"
PYTHONPATH=src python -m pytest -o addopts='' tests/
```

# 6. End-to-End Workflow

## 6.1 One-Command Pipeline

The script examples/run_pipeline.py orchestrates all four stages automatically. It detects which output files already exist and skips those stages, enabling fast incremental reprocessing:

```
PYTHONPATH=src MPLCONFIGDIR=/tmp/mpl \
python examples/run_pipeline.py
```

Key environment overrides:

| Variable | Effect |
|---|---|
| DATA_FILE | Path to input .nxs file |
| NO_VIEWER=1 | Skip GUI stages; stop after writing _delta_pdf.h5 |
| FORCE=1 | Recompute every stage even if output exists |
| FORCE_FROM=rings\|punch\|backfill\|pdf | Recompute from the named stage onward |
| SLICE_AXIS=H\|K\|L | Direction along which ring removal iterates planes (default H) |

## 6.2 Stage-by-Stage Commands

### *Stage 1: Ring Removal*

```
PYTHONPATH=src MPLCONFIGDIR=/tmp/mpl RING_PRESET=cc_on \
python examples/remove_rings_3d.py
```

Output: data/processed/*_ringremoved.h5

Use RING_PRESET=cc_off for more aggressive subtraction on noisy data.

### *Stage 2: Bragg Punch*

```
PYTHONPATH=src MPLCONFIGDIR=/tmp/mpl \
PUNCH_PRESET=cc_on MODE=both MIN_I=0.8 MIN_PROM=0.8 \
INTEGER_FIT_POSITION=1 INTEGER_FIT_SHAPE=1 \
INTEGER_H_GUARD=0.12 \
SEARCH_EXCLUDE_H_FRACTIONS=0.3333,0.6667 \
SEARCH_EXCLUDE_H_WIDTH=0.08 \
python examples/punch_bragg_3d.py
```

Output: data/processed/*_braggpunched.h5

### *Stage 3: Backfill*

```
PYTHONPATH=src METHOD=local \
python examples/backfill_bragg_3d.py
```

Output: data/processed/*_braggpunched_backfilled.h5

### *Stage 4: 3D-ΔPDF*

```
PYTHONPATH=src MPLCONFIGDIR=/tmp/mpl \
SUBTRACT_BG="0,1.5,1.5" CROP_H=4 CROP_K=8 CROP_L=15 \
APODIZE=gaussian GAUSSIAN_SIGMA=0.4 \
python examples/delta_pdf.py
```

Outputs: examples/_delta_pdf.h5 and PNG central-cut images.

## 6.3 Multi-Volume Workflow

Run the pipeline once per input volume by overriding DATA_FILE:

```
# condition A
NO_VIEWER=1 \
DATA_FILE="data/raw/condition_a_cc_sub_bkg.nxs" \
python examples/run_pipeline.py

# condition B
NO_VIEWER=1 \
DATA_FILE="data/raw/condition_b_cc_sub_bkg.nxs" \
python examples/run_pipeline.py

# condition C
NO_VIEWER=1 \
DATA_FILE="data/raw/condition_c_cc_sub_bkg.nxs" \
python examples/run_pipeline.py
```

To generate persistent per-condition ΔPDF files in data/processed/ use OUT_FILE and PROC_FILE:

```
PROC_FILE="data/processed/condition_a_backfilled.h5" \
OUT_FILE="data/processed/condition_a_delta_pdf.h5" \
SUBTRACT_BG="0,1.5,1.5" CROP_H=4 CROP_K=8 CROP_L=15 \
APODIZE=gaussian GAUSSIAN_SIGMA=0.4 \
python examples/delta_pdf.py
```

## 6.4 Python API

All pipeline stages can be called programmatically:

```
import nebula3d
from nebula3d.analysis import BraggRemover, backfill_bragg, compute_delta_pdf

# Load ring-removed volume
vol = nebula3d.load("data/processed/sample_ringremoved.h5")

# Bragg punch
remover = BraggRemover(
mode="both",
punch_spherical_radii=(0.097, 0.072, 0.115), # Å■¹
integer_detect="significance", # no intensity floor at the nodes
min_significance=5.0,
integer_max_shell_fraction=0.15, # no broader than a Bragg peak
integer_optimize_position=True,
integer_optimize_shape=True,
integer_h_guard_hkl=0.12,
search_n_mad=4.0,
search_floor_unit="scatter", # floors × the diffuse scatter
search_min_intensity=27.0,
search_min_prominence=27.0,
search_exclude_h_fractions=(1/3, 2/3),
search_exclude_h_half_width=0.08,
incident_beam_ellipsoid_radii_hkl=(0.15, 0.50, 1.00),
)
punched = remover.apply(vol)

# Backfill
filled = backfill_bragg(punched) # method="local"

# 3D-DeltaPDF
dpdf = compute_delta_pdf(
filled,
apodization="gaussian",
gaussian_sigma=0.4,
crop_hkl=(4, 8, 15),
subtract_smooth_bg=(0, 1.5, 1.5),
)
print(f"DeltaPDF shape: {dpdf.data.shape}")
print(f"Real-space range: x +/-{dpdf.x_axis.max():.1f} A")
```

# 7. Configuration Reference

## 7.1 Ring Removal Parameters

| Parameter | Type | Default | Description |
|---|---|---|---|
| RING_PRESET | str | cc_on | Preset: cc_on (conservative) or cc_off (aggressive) |
| Q_STEP | float | 0.02 | Radial bin width for profile computation (Å■¹) |
| N_FOURIER | int | 8 | Number of Fourier terms for T(ϕ) fit |
| PROFILE_METHOD | str | trimmed_mean | Bin statistic: trimmed_mean or median |
| TEXTURE_Q_SMOOTH | float | 0.1 | Smoothing σ for texture along \|Q\| (Å■¹) |
| SLICE_AXIS | str | H | Axis along which to iterate slices: H, K, or L |

## 7.2 Bragg Punch Parameters

| Parameter | Type | Default | Description |
|---|---|---|---|
| MODE | str | both | Detection: integer, search (auto), or both |
| PUNCH_PRESET | str | cc_on | Preset for punch radii and thresholds |
| MIN_I | float | 0.8 | Minimum intensity threshold for Bragg detection |
| MIN_PROM | float | 0.8 | Minimum local-median prominence |
| INTEGER_FIT_POSITION | 0/1 | 1 | Optimize punch centre to measured maximum |
| INTEGER_FIT_SHAPE | 0/1 | 1 | Fit anisotropic punch radii from data |
| INTEGER_H_GUARD | float | 0.12 | H-slab half-width for integer-node punches |
| SEARCH_EXCLUDE_H_FRACTIONS | str | 0.3333,0.6667 | Periodic H exclusions (fractional parts mod 1) |
| SEARCH_EXCLUDE_H_WIDTH | float | 0.08 | Half-width of search-exclusion slabs |
| INCIDENT_ELLIPSOID_R_HKL | str | 0.15,0.50,1.00 | Direct-beam punch ellipsoid semi-axes (HKL) |

## 7.3 Backfill Parameters

| Parameter | Type | Default | Description |
|---|---|---|---|
| `METHOD` | str | `local` | Fill method: local, laplace, q_shell (comparison) |
| `LAPLACE_GAP` | int | `1` | Voxels outside the punch where the laplace fill takes its boundary |
| `Q_SHELL_STEP` | float | `0.05` | Bin width for q-shell fill (Å■¹) |
| `Q_SHELL_MIN_COUNT` | int | `10` | Min valid voxels per bin; fall back to local if fewer |

## 7.4 3D-ΔPDF Transform Parameters

| Parameter | Type | Default | Description |
|---|---|---|---|
| `APODIZE` | str | `gaussian` | Window function: hann, gaussian, or none |
| `GAUSSIAN_SIGMA` | float | `0.4` | Gaussian window width as fraction of Q_max |
| `CROP_H / CROP_K / CROP_L` | float | `4 / 8 / 15` | Symmetric Q-crop limits in HKL |
| `SUBTRACT_BG` | str | `0,1.5,1.5` | Per-axis Gaussian blur σ for background subtraction (H,K,L r.l.u.) |
| `REAL_SPACE_ANGSTROM` | 0/1 | `1` | Output real-space axes in Å (1) or HKL units (0) |
| `OUT_FILE` | str | `examples/_delta_pdf.h5` | Output HDF5 path for the ΔPDF volume |

# 8. Visualization and Interactive Exploration

## 8.1 Visualization API Overview

All visualization functions live in nebula3d.visualization and follow a primitive-first design: each function accepts an HKLVolume, draws into a caller-supplied Matplotlib Axes or Figure, and returns it. This makes the same calls work in one-shot scripts, IPython sessions, Jupyter notebooks, and the interactive viewer scripts.

```
from nebula3d.visualization import (
extract_slice, plot_slice,
plot_radial_profile, plot_azimuthal_map,
plot_overview, SliceData,
)
```

## 8.2 plot_slice()

Two-dimensional intensity slice through an HKLVolume. The plane argument is read as (horizontal, vertical) and selects the two displayed axes; the remaining axis is cut at value:

| Plane | x-axis | y-axis | Fixed (cut by value) |
|---|---|---|---|
| <tt>'kl'</tt> or <tt>'0kl'</tt> | K | L | H |
| <tt>'hl'</tt> or <tt>'h0l'</tt> | H | L | K |
| <tt>'hk'</tt> or <tt>'hk0'</tt> | H | K | L |

```
# Log-scale background ring view
plot_slice(bkg, "kl", value=0.0, log_scale=True)

# Exact fractional plane with manual colour limits
plot_slice(data, "hk", value=0.3333, interp=True, vmin=0.0, vmax=0.4)
```

**Off-grid interpolation.** Pass interp=True to linearly interpolate between the two bracketing planes so the exact value is honoured. The interpolation is NaN-aware: where only one bracketing plane is valid, that value is used.

## 8.3 plot_radial_profile() and plot_azimuthal_map()

```
# Radial profile with Al(111) ring marker
plot_radial_profile(data, mark_q=[2.69])

# Azimuthal texture of the Al(111) ring
plot_azimuthal_map(data, q_center=2.69)
```

plot_radial_profile bins voxels by |**Q**| and plots the mean or median intensity. plot_azimuthal_map bins by azimuthal angle $\phi$ within a thin |**Q**| shell, showing the ring texture $T(\phi)$.

## 8.4 plot_overview()

```
fig = plot_overview(data, log_scale=True)
```

A 2×2 diagnostic figure: KL (H=0), HL (K=0), HK (L=0) slices and a radial profile. The first look at any new volume.

## 8.5 Interactive Viewers

### *Cleanup QA Viewer*

Four-panel view (raw → ring-removed → Bragg-punched → backfilled) with H/K/L plane selector and cut-position slider. Use to verify that integer-H Bragg peaks are cleanly removed and fractional-H diffuse planes are preserved:

```
PYTHONPATH=src MPLCONFIGDIR=/tmp/mpl \
PUNCH_PRESET=cc_on MODE=both MIN_I=0.8 MIN_PROM=0.8 \
INTEGER_FIT_POSITION=1 INTEGER_FIT_SHAPE=1 INTEGER_H_GUARD=0.12 \
SEARCH_EXCLUDE_H_FRACTIONS=0.3333,0.6667 SEARCH_EXCLUDE_H_WIDTH=0.08 \
H_VALUE=0.3333 \
python examples/explore_slice.py
```

### *3D-ΔPDF Orthoslice Viewer (Recommended)*

All three orthogonal real-space planes simultaneously ($x_H$–$y_K$, $x_H$–$z_L$, $y_K$–$z_L$) with independent cut sliders, a contrast multiplier, and a unit-cell gridline toggle:

```
PYTHONPATH=src MPLCONFIGDIR=/tmp/mpl \
PDF_FILE=data/processed/condition_a_delta_pdf.h5 RMAX=50 \
python examples/explore_delta_pdf_ortho.py
```

### *Multi-Volume Comparison Grid*

A 3×3 grid (rows = three related volumes, columns = three orthogonal cuts) with shared cut sliders and a single global colour scale so intensities are directly comparable:

```
PYTHONPATH=src MPLCONFIGDIR=/tmp/mpl \
PDF_FILES=data/processed/condition_a_delta_pdf.h5,data/processed/condition_b_del
ta_pdf.h5,data/processed/condition_c_delta_pdf.h5 \
PDF_LABELS="condition A,condition B,condition C" \
python examples/explore_delta_pdf_multi.py
```

# 9. Worked Examples

## 9.1 Quick Inspection of a New Dataset

Load a raw volume and display the 2×2 diagnostic overview:

```
import nebula3d
from nebula3d.visualization import plot_overview

vol = nebula3d.load("path/to/sample.nxs") # or .h5
fig = plot_overview(vol, log_scale=True)
fig.savefig("overview.png", dpi=120)
```

## 9.2 Locating and Profiling a Powder Ring

```
from nebula3d.visualization import plot_radial_profile, plot_azimuthal_map
import matplotlib.pyplot as plt

# Step 1: identify ring positions from radial profile
fig, ax = plt.subplots()
plot_radial_profile(vol, ax=ax, mark_q=[2.69, 4.39, 5.07])
plt.title("Radial profile: Al ring positions")

# Step 2: check azimuthal texture T(phi) at Al(111)
plot_azimuthal_map(vol, q_center=2.69, q_width=0.05)
plt.title("Al(111) ring texture")
```

## 9.3 Checking Bragg Punch Quality

Compare an integer-H plane (Bragg present) to a fractional-H diffuse plane:

```
import nebula3d
from nebula3d.visualization import plot_slice
import matplotlib.pyplot as plt

punched = nebula3d.load("data/processed/..._braggpunched.h5")

fig, axes = plt.subplots(1, 2, figsize=(12, 5))

# Integer-H: check all Bragg holes are punched
plot_slice(punched, "kl", value=0.0, ax=axes[0],
log_scale=True, title="H=0 (integer): Bragg punched")

# Fractional-H: check diffuse is preserved
plot_slice(punched, "kl", value=0.3333, interp=True, ax=axes[1],
log_scale=True, title="H=1/3 (fractional): diffuse intact")

fig.tight_layout()
fig.savefig("bragg_check.png", dpi=110)
```

## 9.4 Computing and Displaying the 3D-ΔPDF

```
import nebula3d
from nebula3d.analysis import compute_delta_pdf
import matplotlib.pyplot as plt

filled = nebula3d.load("data/processed/..._backfilled.h5")

dpdf = compute_delta_pdf(
filled,
apodization="gaussian",
gaussian_sigma=0.4,
crop_hkl=(4, 8, 15),
subtract_smooth_bg=(0, 1.5, 1.5),
)
nebula3d.save(dpdf, "output_delta_pdf.h5")

# Plot the central hk0 slice
hk0 = dpdf.slice_hk0() # h-k plane at l=0
vmax = abs(hk0.data).max()

fig, ax = plt.subplots(figsize=(7, 6))
im = ax.imshow(hk0.data.T, origin="lower",
extent=[hk0.x_axis[0], hk0.x_axis[-1],
hk0.y_axis[0], hk0.y_axis[-1]],
cmap="RdBu_r", vmin=-vmax, vmax=vmax)
ax.set_xlabel(r"$x_H$ (Å)")
ax.set_ylabel(r"$y_K$ (Å)")
fig.colorbar(im, ax=ax, label="Δρ (arb.)")
fig.savefig("delta_pdf_hk0.png", dpi=150)
```

# 10. Testing

## 10.1 Running the Test Suite

```
# Full suite with coverage
PYTHONPATH=src python -m pytest -o addopts='' tests/

# Lint (ruff)
python -m ruff check src/ tests/

# Type check (mypy)
python -m mypy src/nebula3d --ignore-missing-imports
```

Note: The -o addopts='' flag is required because the project uses conda rather than a virtualenv; without it pytest may pick up unexpected plugins.

## 10.2 Key Regression Tests

| Test | What it verifies |
|---|---|
| `test_delta_pdf_centring_positive_peak` | Guards the FFT centring fix: verifies a positive cosine input produces a positive real-space peak (not a sign-flipped lobe). |
| Ring subtraction (synthetic) | Verifies that a known Gaussian ring on a flat background is cleanly subtracted without affecting the flat region. |
| Bragg punch / search modes | Checks that integer-node and search-mode masks are consistent with expected punch radii and guard conditions. |

# 11. Known Limitations and Troubleshooting

## 11.1 Near-Origin Spike

**Symptom.** A very strong feature at r < ~3 Å in all ΔPDF maps that dominates the colour scale.

**Cause.** Residual high-|Q| Bragg leakage, discontinuities at punch-hole boundaries, and the direct-beam punch.

**Mitigation.** Set colour scales from the p99 of |Δρ| at r > 3 Å (this is the default in all viewer scripts). For quantitative analysis of short-range correlations, tapered punch boundaries and softer high-|Q| windows are planned for a future release.

## 11.2 Axis Cross in ΔPDF

**Symptom.** A bright cross along $y_K$=0 and $z_L$=0 axes in the real-space map.

**Cause.** Residual separable diffuse envelope (see Section 4.4.3).

**Fix.** Use SUBTRACT_BG="0,1.5,1.5" (or the Python API parameter subtract_smooth_bg=(0, 1.5, 1.5)).

## 11.3 Residual Ring After Subtraction

**Symptom.** Faint ring pattern still visible after ring removal.

**Causes and mitigations.**

- Gaussian width σ_i too narrow: widen via Q_STEP and re-run detection.
- Texture mismatch (rank1_variance < 0.90): check diagnostic and consider per-ring T_i(ϕ) fitting (planned feature).
- Ring preset mismatch: try RING_PRESET=cc_off for noisier or more strongly ringed data.

## 11.4 Search Mode Punching Diffuse Structure

**Symptom.** Structured diffuse on fractional-H planes is partially masked.

**Fix.** Add the fractional H values to the periodic exclusion list: SEARCH_EXCLUDE_H_FRACTIONS=0.3333,0.6667. Use the periodic form rather than an explicit centre list to catch higher-order satellites automatically.

## 11.5 Rank-1 SVD Failure in Ring Model

**Symptom.** rank1_variance < 0.90 after ring fitting.

**Meaning.** Different rings have significantly different azimuthal textures; the shared T(ϕ) model is insufficient.

**Current workaround.** Use the non-parametric production path (PatchedRadialRingModel), which handles per-patch textures without assuming a shared T(ϕ). Per-ring T_i(ϕ) fitting is a planned improvement for the factored model.

## 11.6 Performance Notes

Benchmarks are for a 401×401×301 volume (~48 M voxels) on a typical laptop. Ring removal is the bottleneck; it is embarrassingly parallel per H-slice and a future release will add optional concurrent.futures.ProcessPoolExecutor parallelisation.

| Operation | Typical time | Notes |
|---|---|---|
| Ring removal (per-slice) | ~2–5 min | Sequential per-H-slice; embarrassingly parallel |
| Bragg punch (detection + mask) | ~1–2 min | Scales with n_peaks |
| Backfill (q-shell) | ~30 sec | Linear in n_voxels |
| 3D-ΔPDF (FFT + windowing) | ~10 sec | Dominated by FFT of zero-padded array |

# 12. Glossary

**Apodization.** A window function applied to I(Q) before FFT to suppress termination ripples from the finite Q range. Common choices: Hann, Gaussian.

**Backfill.** Replacement of masked (punched) voxels with estimated background intensities derived from surrounding valid voxels.

**Biharmonic.** A fill based on iterative solution of $\nabla^4 u = 0$; produces very smooth interpolations.

**Bragg peaks.** Sharp peaks at integer (and satellite) reciprocal-lattice nodes; encode the average periodic structure.

**Centrosymmetric.** Having inversion symmetry $I(Q) = I(-Q)$; ensures the Fourier transform is real-valued.

**Diffuse scattering.** Broad, continuous intensity encoding structural disorder, short-range order, and dynamic correlations.

**Fractional HKL.** Reciprocal-lattice coordinates (h,k,l) in units of a*, b*, c*; integers are Bragg positions.

**Hann window.** A cosine-squared taper $w(x) = \cos^2(\pi x / 2x_{max})$; smoothly suppresses the signal to zero at the Q boundary.

**HKLVolume.** The central data structure: a 3D array plus axes, mask, σ, UB matrix, and instrument label.

**Laue class.** Crystal point-group symmetry class (e.g. mmm, m3m, 4/mmm) determining equivalent Q positions.

**MAD.** Median Absolute Deviation; a robust measure of spread: MAD = median(|x – median(x)|).

**Powder ring.** Azimuthally smooth band of intensity at fixed |Q| from polycrystalline material in the beam.

**Punch.** Mask and remove a Bragg peak by setting a region of voxels invalid.

**Q.** Cartesian momentum transfer vector in $Å^{-1}$; $Q = UB \cdot [h,k,l]$; $|Q| = 2\pi/d$ (physics convention).

**SNIP.** Sensitive Nonlinear Iterative Peak clipping; a morphological method for baseline estimation in 1D profiles.

**UB matrix.** Orientation (U) × metric (B) matrix; $Q = UB \cdot [h,k,l]$.

**3D-ΔPDF.** Three-dimensional difference pair distribution function; the Fourier transform of diffuse scattering, mapping to real-space pair correlations.

# 13. References


[1] T. Weber and A. Simonov, “The three-dimensional pair distribution function analysis of disordered single crystals: basic concepts,” *Z. Kristallogr.* **227**, 238–247 (2012). DOI: 10.1524/zkri.2012.1504. **Establishes the 3D-ΔPDF formalism.**

[2] A. Simonov, T. Weber, and W. Steurer, “Experimental uncertainties of three-dimensional pair distribution function investigations exemplified on the diffuse scattering from a tris-*tert*-butyl-1,3,5-benzene tricarboxamide single crystal,” *J. Appl. Cryst.* **47**, 2011–2018 (2014). DOI: 10.1107/S1600576714023668. **Uncertainties of the 3D-ΔPDF reduction.**

[3] A. Chambolle and T. Pock, “A First-Order Primal-Dual Algorithm for Convex Problems with Applications to Imaging,” *J. Math. Imaging Vision* **40**, 120–145 (2011). DOI: 10.1007/s10851-010-0251-1. **Primal-dual TV inpainting (formerly an option; removed).**

[4] M. Bertalmio, G. Sapiro, V. Caselles, and C. Ballester, “Image inpainting,” *Proc. ACM SIGGRAPH*, 417–424 (2000). DOI: 10.1145/344779.344972. **Origin of PDE-based diffusion inpainting.**

[5] M. Bertero and P. Boccacci, *Introduction to Inverse Problems in Imaging*, IOP Publishing, Bristol, UK (1998). ISBN: 978-0750304351. **General theory of regularised inverse problems.**

[6] Mantid Project, “Mantid — Manipulation and Analysis Toolkit for Instrument Data,” *Mantid* (2013–present). www.mantidproject.org. **Data reduction and MDHistoWorkspace I/O.**

[7] C. G. Ryan, E. Clayton, W. L. Griffin, S. H. Sie, and D. R. Cousens, “SNIP, a statistics-sensitive background treatment for the quantitative analysis of PIXE spectra in geoscience applications,” *Nucl. Instrum. Methods Phys. Res. B* **34**, 396–402 (1988). DOI: 10.1016/0168-583X(88)90063-8. **SNIP baseline algorithm used in radial profile processing.**

[8] M. Kobas, T. Weber, and W. Steurer, “Structural disorder in the decagonal Al–Co–Ni. I. Patterson analysis of diffuse x-ray scattering data,” *Phys. Rev. B* **71**, 224205 (2005). DOI: 10.1103/PhysRevB.71.224205. **Origin of the punch-and-fill reduction.**

[9] A. Simonov, T. Weber, and W. Steurer, “*Yell*: a computer program for diffuse scattering analysis via three-dimensional delta pair distribution function refinement,” *J. Appl. Cryst.* **47**, 1146–1152 (2014). DOI: 10.1107/S1600576714008668. **3D-ΔPDF model refinement.**