

# Fields and Galois Theory

David Rufino

Apr 2020

## Contents

<b>1</b>	<b>Fields and Galois Theory</b>	<b>1</b>
1.1	Field extensions . . . . .	1
1.2	Polynomials over a field . . . . .	2
1.2.1	Separable Polynomials . . . . .	3
1.3	Algebraic Extensions . . . . .	4
1.4	Algebraic Closure . . . . .	5
1.5	Separability . . . . .	6
1.6	Splitting Fields . . . . .	7
1.7	Perfect Fields . . . . .	8
1.8	Normal Extensions . . . . .	8
1.9	Galois Correspondence . . . . .	10

## 1 Fields and Galois Theory

Some concise notes on fields and galois theory for algebraic extensions  $K/k$ . Mainly follows LangAlgebra72, with a couple of changes. The definition of separable extension is the more obvious and general one. The proof of key lemma  $[K : K^H] \leq \#H$  uses independence argument rather than Primitive Element Theorem. This is taken from JMilne and Garling.

### 1.1 Field extensions

A field extension  $K/k$  is a pair  $(K, i_{kK})$  where  $i_{kK} : k \rightarrow K$  is an injective ring homomorphism (embedding). Note every field is an extension over its prime subfield ( $k = \mathbb{Q}$  or  $\mathbb{F}_p$ ) by inclusion, so we lose no generality by always considering field extensions. Then a morphism  $\sigma : K/k \rightarrow L/k$  of field extensions is simply a ring homomorphism which commutes i.e.  $\sigma \circ i_{kK} = i_{kL}$ . We may also consider a “tower” of extensions

$$K_n / \dots / K_1 / K_0 = k$$

with embeddings  $i_{K_i K_{i+1}} : K_i \rightarrow K_{i+1}$ , with the picture that these usually correspond to inclusions. Typically if we have a family of morphisms

$$\sigma_i : K_i \rightarrow M$$

they would commute with these embeddings. In particular we may abuse notation by defining  $\sigma_i|_{K_j} = \sigma_i \circ i_{K_{i-1} K_i} \circ \dots \circ i_{K_j K_{j+1}}$ .

Note every field morphism is automatically injective, and is an isomorphism onto its image. We may sometimes omit the reference to the base field  $k$ , particularly when describing morphisms.

Observe every extension  $K/k$  may be viewed as a  $k$ -vector space with  $\lambda \cdot \alpha = i_{kK}(\lambda)\alpha$ .

#### Definition 1.1

A field extension  $K/k$  is said to be finite if it has finite dimension as a  $k$ -vector space. In this case denote its dimension by  $[K : k]$

Field homomorphisms are automatically injective

#### Proposition 1.1

Let  $\sigma : K \rightarrow L$  be a field homomorphism then  $\sigma$  is injective. Therefore we may refer to it as an embedding. We write the set of such maps compatible with  $k$ -structure as

$$\text{Mor}_k((K, i_{kK}), (L, i_{kL})) = \text{Mor}_k(K, L)$$

Note we will mostly omit the dependence on the maps  $i_{kK}$  and  $i_{kL}$ .

*Proof.* Recall that for a ring homomorphism  $\sigma$  is injective iff  $\ker(\sigma) = (0)$ . Suppose that  $x \neq 0$ , then  $1 = f(1) = f(xx^{-1}) = f(x)f(x)^{-1}$ , hence  $f(x) \neq 0$ . Therefore  $\sigma$  is injective.

Alternatively since a field has no non-trivial ideals, so  $\ker(\sigma) = (0)$  or  $K$ . □

In many circumstances they are also surjective, by using counting arguments.

**Proposition 1.2**

Let  $K/k$  be a finite field extension, then every  $k$ -embedding  $\sigma : K \rightarrow K$  is automatically surjective and therefore an isomorphism.

$$\text{Mor}_k(K, K) = \text{Aut}(K/k)$$

*Proof.* This follows from results on finite vector spaces. □

## 1.2 Polynomials over a field

**Proposition 1.3**

The ring  $k[X]$  is a principal ideal domain, that is every ideal has the form

$$\mathfrak{a} = (f)$$

Moreover we may choose  $f$  to be monic, and in this case it is unique.

**Corollary 1.4 (UFD)**

$k[X]$  is a unique factorization domain (UFD). More precisely every non-zero polynomial may be decomposed uniquely as follows

$$f = c(f) \prod_{i \in I} p_i^{v_{p_i}(f)}$$

where the  $p_i$  run over all distinct monic irreducible polynomials,  $0 \neq c(f) \in k$  is the leading coefficient of  $f$  and  $v_{p_i}(f)$  are non-negative integers with only finitely many non-zero so that the product is well-defined.

*Proof.* The fact that  $k[X]$  is a UFD follows from standard results about PIDs. Note that the units of  $k[X]$  are precisely the constant polynomials, so  $f, g$  are associates iff  $f = \lambda g$ , and every irreducible polynomial is associated to precisely one monic irreducible polynomial. Therefore we're able to determine a unique decomposition as above. □

**Remark 1.2**

The polynomials  $(X - \alpha)$  are irreducible.

Note by uniqueness  $v_p$  satisfies

$$v_p(fg) = v_p(f) + v_p(g)$$

and furthermore

$$\deg(f) = \sum_i \deg(p_i) v_{p_i}(f)$$

The support of  $f$  is the set of irreducible polynomials for which  $v_p(f)$  is non-zero.

**Lemma 1.5 (Divisibility)**

$f$  divides  $g$  iff  $v_p(f) \leq v_p(g) \ \forall p$  iff  $(g) \subseteq (f)$

**Lemma 1.6 (Roots and Multiplicity)**

For  $f \in k[X]$  a non-constant polynomial and  $\alpha \in k$  we have

$$f(\alpha) = 0 \iff (X - \alpha) \mid f \iff v_{(X-\alpha)}(f) > 0$$

In this case  $r := v_{(X-\alpha)}(f)$  is the multiplicity of the root  $\alpha$ , and observe

$$f(X) = c(X - \alpha)^r g(X)$$

with  $g(\alpha) \neq 0$  (equivalently  $v_{(X-\alpha)}(g) = 0$ ).

*Proof.* The right to left implication is obvious. Conversely by the division algorithm we may write

$$f(X) = f(\alpha) + (X - \alpha)Q(X)$$

Then if  $f(\alpha) = 0$  we clearly have  $v_{(X-\alpha)}(f) > 0$ . Finally we may construct

$$g(X) = \prod_{p \neq (X-\alpha)} p^{v_p(f)}$$

It's clear that for every  $p$  appearing in the product  $p(\alpha) \neq 0$  because otherwise we would have  $(X - \alpha) \mid p$  and by irreducibility  $(X - \alpha) = p$ . Therefore  $g(\alpha) \neq 0$  as required. □

**Definition 1.3**

Let  $\sigma : K \rightarrow L$  be a field morphism and  $f \in K[X]$  such that

$$f(X) = a_0 + a_1X + \dots + a_nX^n$$

then define  $f^\sigma \in L[X]$  by

$$f^\sigma(X) := \sigma(a_0) + \sigma(a_1)X + \dots + \sigma(a_n)X^n$$

Note this has the property that

$$f^\sigma(\sigma(\alpha)) = \sigma(f(\alpha)) \quad \forall \alpha \in K$$

In the case of a field extension  $K/k$  we may abuse notation by writing  $f^{i_{kK}}$  as simply  $f$ .

**Definition 1.4** (Splitting Polynomial)

Let  $K/k$  be a field extension and  $f \in k[X]$ . We say a polynomial  $f$  splits completely in  $K$  if the irreducible decomposition of  $f$  in  $K[X]$  is

$$f(X) = c(f) \prod_{i=1}^n (X - \alpha_i)^{r_i}$$

where  $\alpha_i$  are the distinct roots of  $f(X)$  in  $K$  and  $r_i := v_{(X-\alpha_i)}(f)$  are the multiplicities. Equivalently  $f$  splits in  $K$  if

$$p \in K[X] \text{ irreducible} \wedge \deg(p) > 1 \implies v_p(f^i) = 0 \quad (1)$$

Observe that the number of roots counting multiplicities is  $\deg(f)$

$$\deg(f) = \sum_{i=1}^n v_{(X-\alpha_i)}(f^i)$$

**1.2.1 Separable Polynomials**

NB in this section we abuse notation by identifying  $f \in k[X]$  and  $f^{i_{kK}} \in K[X]$

We are interested characterizing polynomials  $f \in k[X]$  which do not have multiple roots in any extension field  $K/k$ . This may be achieved by considering the formal derivative  $f'(X)$  as follows

**Proposition 1.7** (Criteria for Multiple Roots)

Let  $f(X) \in k[X]$  be a polynomial and either  $\text{char}(k) = 0$  or  $r < \text{char}(k)$ . Then  $\alpha \in K$  is a root of multiplicity  $r$  precisely when

$$f(\alpha) = f^{(1)}(\alpha) = \dots = f^{(r-1)}(\alpha) = 0$$

and  $f^{(r)}(\alpha) \neq 0$ .

Therefore the multiple roots are precisely the common roots of  $f(X)$  and  $f'(X)$  (irrespective of the characteristic).

*Proof.* Note that by Lemma 1.6

$$f^{(1)}(X) = (X - \alpha)^{r-1} [rg(X) + (X - \alpha)g'(X)]$$

with  $g(\alpha) \neq 0$  and  $r$  the multiplicity of the root. If  $r = 1$ , then  $f^{(1)}(\alpha) = g(\alpha) \neq 0$  as required. If  $r > 1$ , then  $f^{(1)}(X)$  has  $\alpha$  as a root of multiplicity  $r - 1$ , so it follows by induction.

The second statement is simply the case  $r = 1$ . □

**Definition 1.5** (Coprime)

We say two elements  $x, y$  of a ring are coprime if  $(x, y) = (1) \iff ax + by = 1$  for some  $a, b$

**Definition 1.6**

A polynomial  $f \in k[X]$  is separable if  $f$  and  $f'$  are coprime.

**Proposition 1.8** (Separable Polynomial)

A separable polynomial  $f \in k[X]$  has no multiple roots in any extension field  $K/k$

*Proof.* Since  $(f, f') = 1$  we have  $af + bf' = 1$ . Clearly  $f$  and  $f'$  have no common roots, and therefore  $f$  has no multiple roots by Proposition 1.7. □

We can provide a partial converse by working in a large enough extension field

**Proposition 1.9** (Separability)

Let  $K/k$  be a field extension and  $f \in k[X]$  a polynomial which splits completely in  $K$ . Then TFAE

1.  $f$  is separable
2.  $f$  has no multiple roots in  $K$
3.  $f$  has  $\deg(f)$  distinct roots in  $K$

*Proof.* Using the formula

$$\deg(f) = \sum_{i=1}^n v_{(X-\alpha_i)}(f)$$

we see easily that  $3 \iff 2$ . The previous Proposition shows that  $1 \implies 2$ .

Conversely suppose  $f$  is not separable, then by Lemma 1.10  $f$  and  $f'$  must have a non-trivial common divisor  $h$ . Using 1.5 and (1) we see that  $h$  splits in  $K$ . Any root of  $h$  is a common root of  $f$  and  $f'$  in  $K$ , which by Proposition 1.7 is a multiple root of  $f$  in  $K$ .  $\square$

We used the following

**Lemma 1.10** (Co-prime elements in a PID)

Let  $A$  be a PID, then  $x, y$  are co-prime if and only if they have no non-trivial common divisors.

*Proof.* First suppose  $(x, y) = 1$ , then  $ax + by = 1$  and any common divisor  $d$  must divide 1 and therefore be invertible.

Conversely suppose  $(x, y) \neq (1)$ , since  $A$  is a PID it must equal  $(d)$  for some non-invertible  $d$ , which by definition is a non-trivial common divisor.  $\square$

We will need the following

**Lemma 1.11**

If  $f \mid g$  and  $g$  is separable then  $f$  is separable

*Proof.* By assumption we have  $g = hf$  and  $ag + bg' = 1$  (see). By the product rule we have  $g' = h'f + hf'$ . Therefore  $1 = (ah)f + b(h'f + hf') = (ah + bh')f + bhf'$ , so  $f$  and  $f'$  are coprime as required.  $\square$

### 1.3 Algebraic Extensions

**Definition 1.7** (Algebraic element and Minimal Polynomial)

If  $K/k$  is a field extension we say that  $\alpha \in K$  is algebraic (over  $k$ ) if it satisfies

$$f^i(\alpha) = 0$$

for some polynomial  $f \in k[X]$ , or equivalently if the evaluation morphism

$$\phi_\alpha : k[X] \rightarrow K$$

has a non-zero kernel. There is a unique monic polynomial  $m_{\alpha,k}(X)$  such that  $(m_{\alpha,k}) = \ker(\phi_\alpha)$ , which we call the minimal polynomial. This is irreducible. It is the unique monic polynomial such that

$$f^i(\alpha) = 0 \iff m_{\alpha,k} \mid f$$

*Proof.* Note that  $k[X]/\ker(\phi_\alpha)$  is isomorphic to  $\text{im}(\phi_\alpha)$ . As the image is an integral domain, being a subring of a field, the kernel must be prime. This means that  $m_{\alpha,k}$  is irreducible.  $\square$

**Definition 1.8** (Algebraic Extension)

A field extension  $K/k$  is algebraic if every element  $\alpha \in K$  is algebraic.

A field extension  $K/k$  is simple if  $K = k(\alpha)$  for  $\alpha$  algebraic.

**Proposition 1.12** (f.g. algebraic  $\implies$  finite)

A simple extension  $K = k(\alpha)$  is finite of dimension  $\deg(m_{\alpha,k})$ .

More generally a finitely generated algebraic extension  $K = k(\alpha_1, \dots, \alpha_n)$  is finite over  $k$ .

*Proof.* The first part follows from an isomorphism with  $k[\alpha] = k[X]/(m_\alpha)$ .

The second part follows by induction and taking simple extensions.  $\square$

This allows us to prove a stronger version of Proposition 1.2

**Proposition 1.13**

Let  $K/k$  be an algebraic extension then any  $k$ -embedding to itself is surjective and therefore an isomorphism. In other words

$$\text{Mor}_k(K, K) = \text{Aut}(K/k)$$

*Proof.*  $\square$

We prove the first lifting theorem

**Proposition 1.14** (Lifting to simple extensions)

Let  $K(\alpha)/K$  be a simple algebraic extension, and  $\sigma : K \rightarrow L$  a field embedding such that  $m_{\alpha,K}^\sigma$  has a root in  $L$ . Then there exists a lifting  $\sigma : K(\alpha) \rightarrow L$ .

More precisely the number of extensions is equal to the number of distinct roots of  $m_{\alpha,K}^\sigma$  in  $L$ .

In particular if  $m_{\alpha,K}^\sigma$  is separable and splits completely in  $L$  then there are precisely  $\deg(m_\alpha) = [K(\alpha) : K]$  such extensions.

*Proof.* Essentially we just need that  $K(\alpha)$  and  $\sigma(K)(\alpha')$  are both isomorphic to the quotient ring  $K[X]/(m_{\alpha,K})$ .  $\square$

**Corollary 1.15**

Let  $K(\alpha)/K$  and  $K(\beta)/K$  be two simple extensions such that

$$m_{\alpha,K} = m_{\beta,K}$$

Then there exists a unique  $K$ -isomorphism

$$\sigma : K(\alpha) \rightarrow K(\beta)$$

such that

$$\sigma(\alpha) = \beta$$

*Proof.* The embedding  $K \rightarrow K(\beta)$  extends by the previous proposition to a  $K$ -embedding  $K(\alpha) \rightarrow K(\beta)$ . The image is a field containing  $\beta$  and therefore clearly surjective.  $\square$

## 1.4 Algebraic Closure

The theory is somewhat easier to develop if one may assume the existence of a maximal algebraic extension, so we develop this first.

**Definition 1.9** (Algebraically Closed)

A field  $M$  is algebraically closed if one of the following equivalent conditions holds

- Every algebraic extension  $M'/M$  is trivial
- Every non-constant polynomial in  $M[X]$  has a root in  $M$
- Every non-constant polynomial in  $M[X]$  splits in  $M$

If  $M$  is an extension field of  $k$ , then we say it is algebraically closed over  $k$ .

**Definition 1.10** (Algebraic Closure)

An algebraic closure  $\bar{k}$  of  $k$  is a field extension  $\bar{k}/k$  which is algebraic and for which  $\bar{k}$  is algebraically closed.

**Proposition 1.16** (Existence of Algebraic Closure)

Given a field  $k$  there exists an algebraic closure  $\bar{k}/k$

**Proposition 1.17** (Generic Lifting Theorem)

Let  $K/k$  be a field extension. Suppose  $L/K$  is an algebraic extension and  $\sigma : K \rightarrow M$  a  $k$ -embedding into an algebraically closed field. Then there exists an extension  $\tilde{\sigma} : L \rightarrow M$ . In other words there is a surjection

$$\text{Mor}_k(L, M) \rightarrow \text{Mor}_k(K, M)$$

$$\sigma \rightarrow \sigma \circ i_{KL}$$

*Proof.* Broadly speaking consider the poset of extensions to subfields of  $L$  ordered under consistency and take a maximal element by Zorn's Lemma. Apply Proposition 1.14 to show that this maximal element must be an extension to  $L$ .

If  $[L : K] < \infty$ , then we may simply proceed by induction on dimension.  $\square$

**Corollary 1.18** (Unique up to isomorphism)

An algebraic closure  $\bar{k}$  of  $k$  is unique up to (non-unique) isomorphism.

**Remark 1.11**

Note if  $K/k$  is an algebraic extension then the Proposition shows that we may consider construct an embedding  $K \rightarrow \bar{k}$  commuting with  $k \rightarrow \bar{k}$ .

In general given a tower of algebraic extensions

$$K_n / \dots / K_1 / K_0 = k$$

we will assume the existence of compatible embeddings  $i_{K_i} : K_i \rightarrow \bar{k}$  such that  $i_{K_{i+1}} \circ i_{K_i, K_{i+1}} = i_{K_i}$ .

## 1.5 Separability

The concept of separable extensions is important, as illustrated by Proposition 1.14.

**Definition 1.12** (Separability)

An algebraic element  $x \in K$  is separable over  $k$  if its minimal polynomial is separable.

**Definition 1.13** (Separable Extension)

An algebraic extension  $K/k$  is separable if every element is separable.

**Remark 1.14**

Lang72 initially only defines separability for finite field extensions by  $[K : k]_s = [K : k]$ . Then defines for algebraic extensions for f.g. (i.e. finite) sub-extensions.

**Proposition 1.19**

Let  $K/F/k$  be a tower of extensions, then  $K/k$  is separable  $\iff K/F, F/k$  separable.

*Proof.* Suppose  $K/k$  is separable. Clearly  $F/k$  is separable. Furthermore  $m_{\alpha, F} | m_{\alpha, k}$  by definition, and so the former is separable by Lemma 1.11.

Conversely TODO □

**Proposition 1.20**

$K = k(\alpha_1, \dots, \alpha_n)$  is separable over  $k$  if and only if each  $\alpha_i$  is separable over  $k$ .

*Proof.* □

**Definition 1.15** (Separability degree)

The cardinality of  $\text{Mor}_k(K, (\bar{k}, i_k))$  is independent of the choice of embedding  $i_k : k \rightarrow \bar{k}$ . We write this as  $[K : k]_s$

*Proof.* Essentially if  $L/k$  and  $M/k$  are isomorphic then so are  $\text{Mor}_k(K, L)$  and  $\text{Mor}_k(K, M)$ , since  $\text{Mor}_k(K, -)$  is a covariant functor. If  $L/k$  and  $M/k$  are two algebraic closures of  $k$ , then we have seen they are isomorphic and the result follows. □

From Proposition 1.14 we've seen that  $[k(\alpha) : k]_s \leq [k(\alpha) : k]$  with equality iff  $\alpha$  is separable iff  $k(\alpha)$  is separable. We may show this more generally

**Proposition 1.21** (Multiplicativity of separability degree)

If  $L/K/k$  is a tower of finite extensions then

$$[L : k]_s = [L : K]_s [K : k]_s.$$

*Proof.* We assume compatible embeddings in  $\bar{k}$ . The restriction map

$$\psi : \text{Mor}_k(L, \bar{k}) \rightarrow \text{Mor}_k(K, \bar{k})$$

is surjective by Proposition 1.17. Consider  $\sigma \in \text{Mor}_k(K, \bar{k})$  then the fibre  $\psi^{-1}(\sigma)$  is equal to  $\text{Mor}_K(L, (\bar{k}, \sigma))$ . As we've noted the cardinality does not depend on the embedding  $\sigma$  so the result follows. □

**Proposition 1.22** (Bounds on  $\text{Mor}_k(K, \bar{k})$ )

Let  $K/k$  be a finite extension then

$$[K : k]_s \leq [K : k]$$

with equality if and only if  $K/k$  is separable.

*Proof.* For a general finite extension we may obtain a tower of simple extensions

$$K = K_n / \dots / K_1 / K_0 = k$$

with  $K_i = K_{i-1}(\alpha_i)$ . We have by Proposition 1.21

$$[K : k]_s = \prod_{i=1}^n [K_i : K_{i-1}]_s$$

and from Proposition 1.14 that  $[K_i : K_{i-1}]_s \leq [K_i : K_{i-1}] = \deg(m_{\alpha_i, K_{i-1}})$ . The inequality follows from multiplicativity of extension and separability degrees.

When  $K/k$  is separable then Proposition 1.14 also shows that  $[K_i : K_{i-1}]_s = [K_i : K_{i-1}]$  and again we have equality by multiplicity of both degrees.

Conversely if  $[K : k]_s = [K : k]$  then we must have equality at each stage, which implies  $\alpha_i$  is separable over  $K_{i-1}$ . The choice of  $\alpha_1$  was arbitrary, so we see that every element of  $K$  is separable over  $k$  in this case, as required. □

**Definition 1.16** (Bounds on  $\text{Aut}(K/k)$ )

Let  $K/k$  be an algebraic extension and  $i_K : K \rightarrow \bar{k}$  a fixed  $k$ -embedding. Then there is a natural injection

$$\begin{aligned} i_K^\# : \text{Aut}(K/k) &\rightarrow \text{Mor}_k(K, \bar{k}) \\ \sigma &\rightarrow i_K \circ \sigma \end{aligned}$$

In particular in the finite case

$$\# \text{Aut}(K/k) \leq [K : k]_s \leq [K : k] < \infty$$

If  $i$  is inclusion, then this we may regard  $\text{Aut}(K/k)$  as a subset of  $\text{Mor}_k(K, \bar{k})$

**Corollary 1.23** (Extension to algebraic closure II)

Let  $K/k$  be an algebraic extension and  $i_K : K \rightarrow \bar{k}$  a  $k$ -embedding. Then every  $\sigma : K \rightarrow \bar{k}$  lifts to  $\sigma : \bar{k} \rightarrow \bar{k}$ . i.e. there is a canonical surjection by restriction

$$\begin{aligned} i_K^\flat : \text{Aut}(\bar{k}/k) &\longrightarrow \text{Mor}_k(K, \bar{k}) \\ \sigma &\longrightarrow \sigma \circ i_K \end{aligned}$$

When  $i$  is inclusion then this is simply the restriction to  $K$ .

*Proof.* Given  $\sigma \in \text{Mor}_k(K, \bar{k})$  there exists an extension  $\tilde{\sigma} \in \text{Mor}_k(\bar{k}, \bar{k})$  by Proposition 1.17 with  $L = M = \bar{k}$ . This is automatically an automorphism as required.  $\square$

**Corollary 1.24** (Conjugate elements)

We say  $\alpha, \beta \in \bar{k}$  are conjugate if they have the same minimal polynomial.

This is the case if and only if there is an element  $\sigma \in \text{Aut}(\bar{k}/k)$  such that  $\sigma(\alpha) = \beta$ .

*Proof.* By Corollary 1.15 there is an isomorphism  $k(\alpha) \rightarrow k(\beta)$ . Corollary 1.23 gives the required automorphism. The converse is easy.  $\square$

As an application of the concept of separability degree we prove

**Proposition 1.25** (Primitive Element Theorem)

Let  $K/k$  be a finite separable extension of  $k$  then  $K = k(\alpha)$  is simple.

*Proof.* We only prove the case  $k$  is infinite. The finite case can be proven separately by showing that the  $K^*$  is cyclic.

Consider the set  $\text{Mor}_k(K, \bar{k}) = \{\sigma_1, \dots, \sigma_n\}$  which by Proposition 1.22 has order  $n = [K : k]$ . By induction we can assume that  $K = k(\alpha, \beta)$ . We claim that there exists  $0 \neq c \in k$  such that  $\sigma_i(\alpha + c\beta)$  are all distinct. In this case we clearly have  $\# \text{Mor}_k(k(\alpha + c\beta), \bar{k}) \geq n$  so by the same result  $[k(\alpha + c\beta) : k] \geq n$  whence  $k(\alpha + c\beta) = K$ .

We have  $\sigma_i(\alpha + c\beta) = \sigma_j(\alpha + c\beta) \iff c(\sigma_i(\beta) - \sigma_j(\beta)) = (\sigma_i(\alpha) - \sigma_j(\alpha))$ . Therefore consider the polynomial

$$f(X) = \prod_{i \neq j} (X(\sigma_i(\beta) - \sigma_j(\beta)) - (\sigma_i(\alpha) - \sigma_j(\alpha)))$$

Then the embeddings are distinct precisely when  $f(c) \neq 0$ . Since  $f(X)$  has at most finitely many roots and  $k$  is infinite, there must exist such a  $c$ .  $\square$

## 1.6 Splitting Fields

**Definition 1.17** (Splitting Field)

A field extension  $K/k$  is a splitting field for  $f \in k[X]$  if  $f$  splits in  $K$  and does not in any proper subfield. Equivalently

$$K = k(\alpha_1, \dots, \alpha_n).$$

where  $\alpha_i$  are the roots of  $f^i$  in  $K$ . Similarly  $K$  is said to be a splitting field for a family of polynomials  $\{f_i\}$  if it splits every polynomial and no proper subfield does so.

A splitting field always exists

**Proposition 1.26** (Existence of Splitting Fields)

Every family of polynomials has a splitting field. Moreover any two such splitting fields are  $k$ -isomorphic.

*Proof.* This easiest way is to assume the existence of an algebraic closure and take the smallest field under which the polynomials split. Alternatively if the family of polynomials is finite we may proceed by induction on the total degree and adjoin roots as simple extensions. This gives a bound on the total degree of the extension.

Uniqueness (up to non-unique isomorphism!) requires an extension theorem and a rigidity theorem...  $\square$

## 1.7 Perfect Fields

For large classes of base fields all algebraic extensions are separable :

**Proposition 1.27** (Perfect field)

Let  $k$  be a field. Then TFAE

- Every irreducible polynomial in  $k[X]$  is separable
- Every algebraic extension  $K/k$  is separable
- $\bar{k}$  is separable

In this case we say  $k$  is perfect.

**Proposition 1.28** (Criteria for perfectness)

$k$  is perfect if and only if one of the following holds

- $k$  has characteristic 0
- $k$  has characteristic  $p$  and every element is a  $p$ -th power

In particular finite fields are perfect.

## 1.8 Normal Extensions

A normal extension is an algebraic extension in which every minimal polynomial splits completely. In some sense it is “closed” because automorphisms over the algebraic closure map the field into itself (since morphisms must permute roots). In what follows it is more convenient to work with an algebraic closure, but strictly speaking that is not necessary (see Garling). First we prove a trivial Lemma.

**Lemma 1.29**

Let  $K/k$ ,  $L/k$  be field extensions and  $\sigma : K \rightarrow L$  a  $k$ -embedding. Suppose that  $f \in k[X]$  is a polynomial, then  $\alpha$  is a root of  $f$  if and only if  $\sigma(\alpha)$  is a root of  $f^\sigma$ .

If  $f$  splits in  $K$ , then it splits in  $L$  and  $i$  induces a bijection between the roots of  $f$  in  $K$  and  $L$  preserving multiplicity.

**Proposition 1.30** (Normal Extension)

Let  $K/k$  be an algebraic extension and a given algebraic closure  $\bar{k}$ , then the following are equivalent

1. For any two  $k$ -embedding of  $\sigma, \tau \in \text{Mor}_k(K, \bar{k})$  we have  $\sigma(K) = \tau(K)$ .
2.  $K$  is the splitting field of some family of polynomials  $f_i \in k[X]$ .
3. Every irreducible polynomial in  $k[X]$  with at least one root in  $K$  splits in  $K$ .

This is called a normal extension.

*Proof.* Clearly  $3 \implies 2$ , for  $K$  is the splitting field of all the minimal polynomials of elements in  $K$ .

We show  $2 \implies 1$ . Define  $T_j = \{\alpha \in K \mid f_j(\alpha) = 0\}$  and  $T'_j = \{\alpha \in \bar{k} \mid f_j(\alpha) = 0\}$ . By hypothesis  $K = k(\cup_j T_j)$ , and we claim that for any such  $\sigma$  we have  $\sigma(K) = k(\cup_j T'_j) =: K' \subset \bar{k}$ . By Lemma 1.29 that any  $\sigma$  induces a bijection between  $T_j$  and  $T'_j$ . Therefore  $\cup_j T'_j \subseteq \sigma(K) \implies K' \subseteq \sigma(K)$ . Furthermore  $\sigma(K) \subseteq k(\sigma(\cup_j T_j)) \subseteq k(\cup_j T'_j) = K'$ .

Finally we show  $1 \implies 3$ . Wlog assume  $k \subset K \subset \bar{k}$ . Suppose  $f(X)$  is an irreducible polynomial with root  $\alpha_1, \dots, \alpha_n \in \bar{k}$  and  $\alpha_1 \in K$ . By Corollary 1.24 there is a morphism  $\phi \in \text{Aut}(\bar{k}/k)$  such that  $\phi(\alpha_1) = \alpha_j$ . Therefore the inclusion map  $i$  and the composite map  $\phi \circ i$  must have the same image,  $K$ . Therefore  $\alpha_j \in K$  as required.  $\square$

We provide another criterion

**Proposition 1.31**

Let  $K/k$  be an algebraic extension and  $i_K : K \rightarrow \bar{k}$  a given  $k$ -embedding then the following are equivalent

- $K/k$  is normal
- The embedding  $i_K^\# : \text{Aut}(K/k) \rightarrow \text{Mor}_k(K, \bar{k})$  (1.16) is a bijection

When  $i_K$  is inclusion it's necessary and sufficient that every  $\sigma \in \text{Mor}_k(K, \bar{k})$  has image  $K$ .

When  $K/k$  is finite it's necessary and sufficient that

$$\# \text{Aut}(K/k) = [K : k]_s$$



*Proof.* If  $K/k$  is normal, then given  $\sigma' \in \text{Mor}_k(K, \bar{k})$  by hypothesis the image of  $i$  and  $\sigma'$  is the same. Therefore we can define  $\sigma(x) = i^{-1}(\sigma'(x))$ .

Conversely any  $\sigma' \in \text{Mor}_k(K, \bar{k})$  can be written as  $i \circ \sigma$  for  $\sigma$  invertible, so must have image  $i(K)$ . As this is true for every such  $\sigma'$  then  $K/k$  must be normal.

Clearly if  $K/k$  is finite, then we've shown in Proposition 1.22 that  $[K : k]_s < \infty$ . So the embedding  $i^\#$  is bijective if and only if  $\# \text{Aut}(K/k) = [K : k]_s$  as required.  $\square$

**Corollary 1.32** (Subfield is Normal)

Let  $K/k$  be a normal extension and  $F \subset K$  a subfield, then  $K/F$  is normal.

*Proof.* We assume the tower  $F/k$  has compatible embeddings into  $\bar{k}$ , then we have

$$\text{Mor}_F(K, \bar{k}) \subset \text{Mor}_k(K, \bar{k})$$

and the result follows from the first normality criterion.  $\square$

**Proposition 1.33** (Automorphisms of Normal Extension)

Let  $K/k$  be a normal extension with compatible embeddings in  $\bar{k}$ . There is a canonical isomorphism of groups

$$\begin{aligned} \text{Aut}(\bar{k}/k) / \text{Aut}(\bar{k}/K) &\rightarrow \text{Aut}(K/k). \\ \sigma &\rightarrow i_K^{-1} \circ \sigma \circ i_K \end{aligned}$$

When  $K \subset \bar{k}$  then this map is simply restriction to  $K$ .

*Proof.* By Proposition 1.30 i)  $(\sigma \circ i_K)(K) = i_K(K)$  so this is well-defined and by Corollary 1.23 it's surjective. It's clear the kernel is  $\text{Aut}(\bar{k}/K)$  so the result follows.  $\square$

In fact we can replace  $\bar{k}$  with a normal overfield and obtain results corresponding to Corollary 1.23 and Proposition 1.31 respectively. Due to the explicit embeddings in  $\bar{k}$  the arguments become slightly awkward.

**Corollary 1.34** (Extension to normal overfield)

Let  $L/k$  be a normal extension such that  $K \subset L$  then there is a canonical surjective monoid morphism

$$\text{Aut}(L/k) \rightarrow \text{Mor}_k(K, L)$$

by restriction. The kernel is  $\text{Aut}(L/K)$

*Proof.* Assume an embedding  $i_L : L \rightarrow \bar{k}$ .

Consider  $\sigma \in \text{Mor}_k(K, L)$ , then by Corollary 1.23 there exists an extension  $\tilde{\sigma} \in \text{Aut}(\bar{k}/k)$  such that  $i_L \circ \sigma = \tilde{\sigma} \circ i_K$ . As in the previous Proposition there exists  $\hat{\sigma} \in \text{Aut}(L/k)$  such that  $\tilde{\sigma} \circ i_L = i_L \circ \hat{\sigma}$ . Therefore  $i_L \circ \sigma = i_L \circ \hat{\sigma}|_K$  which implies  $\sigma = \hat{\sigma}|_K$  as required.  $\square$

**Corollary 1.35** (Automorphisms of Normal subextension)

Let  $L/k$  be a normal extension and  $K \subset L$ , then the following are equivalent

1.  $K/k$  is normal
2.  $\text{Mor}_k(K, L) = \text{Aut}(K/k)$ , i.e. every  $k$ -embedding  $\sigma : K \rightarrow L$  has  $\sigma(K) = K$ .
3. For every  $\sigma \in \text{Aut}(L/k)$  we have  $\sigma(K) = K$

In this case  $\text{Aut}(L/K) \triangleleft \text{Aut}(L/k)$  is normal and we have a canonical group isomorphism

$$\text{Aut}(L/k) / \text{Aut}(L/K) \rightarrow \text{Aut}(K/k)$$

*Proof.* 1  $\implies$  2). Given  $\sigma \in \text{Mor}_k(K, L)$ , then by definition we have  $(i_L \circ \sigma)(K) = (i_L)(K)$ , which means  $\sigma(K) = K$

2  $\implies$  1). Given any  $\sigma \in \text{Mor}_k(K, \bar{k})$ , this extends to  $\tilde{\sigma} \in \text{Mor}_k(L, \bar{k})$ . Since  $L$  is normal  $\tilde{\sigma}(L) = i_L(L)$ , we see  $\hat{\sigma} := i_L^{-1} \circ \tilde{\sigma} \in \text{Aut}(L/k)$ . By hypothesis  $\hat{\sigma}(K) = K$ . Furthermore  $\sigma(K) = \tilde{\sigma}(K) = i_L(\hat{\sigma}(K)) = i_L(K)$ . Since  $\sigma$  is arbitrary then it shows  $K$  is normal.

2  $\implies$  3). This is clear

3  $\implies$  2). Given  $\sigma \in \text{Mor}_k(K, L)$  by the previous Corollary we can extend to  $\text{Aut}(L/k)$  and the result follows easily. By the previous corollary the canonical map given is surjective, which yields the isomorphism.  $\square$

The following is straight-forward

**Corollary 1.36** (Conjugate Elements)

Let  $K/k$  be a normal extension and two elements  $\alpha, \beta \in K$  be two elements with the same minimal polynomial. Then there exists  $\sigma \in \text{Aut}(K/k)$  such that  $\sigma(\alpha) = \beta$ .

*Proof.* By Corollary 1.15 there is an isomorphism  $k(\alpha) \rightarrow k(\beta)$ . Corollary 1.34 gives the required extension.  $\square$

## 1.9 Galois Correspondence

We've seen that for  $K/k$  a finite extension (Definition 1.16)

$$\# \operatorname{Aut}(K/k) \leq [K : k]_s \leq [K : k]$$

The first inequality is an equality if and only if  $K$  is normal (Proposition 1.31), and the second is an equality if and only if  $K$  is separable (Proposition 1.22). We have shown

**Proposition 1.37** (Order of Galois Group)

*A finite extension  $K/k$  is normal and separable if and only if  $\# \operatorname{Aut}(K/k) = [K : k]$*

*A normal and separable extension  $K/k$  is called Galois and its group of automorphisms denoted  $\operatorname{Gal}(K/k)$*

**Remark 1.18**

*If  $k$  is perfect then  $\bar{k}/k$  is Galois.*

The main result of Galois Theory is that in the finite case there is an order-reversing bijection between subgroups and subfields

$$\begin{aligned} \{H \leq \operatorname{Gal}(K/k)\} &\longleftrightarrow \{F \subseteq K\} \\ \phi : H &\longrightarrow K^H := \{x \in K \mid h(x) = x \quad \forall h \in H\} \\ \psi : \operatorname{Gal}(K/F) &\longleftarrow F \end{aligned}$$

Such an order reversing map is usually called an (antitone) Galois connection, as the first such type arose from Galois Theory. Note it is well-defined because of the following proposition.

**Proposition 1.38**

*If  $K/k$  is Galois and  $F \subset K$  then  $K/F$  is Galois.*

*Proof.* This follows from Corollary 1.32 and 1.19 □

We need to show that  $\phi \circ \psi = \operatorname{id}$  and  $\psi \circ \phi = \operatorname{id}$ . The first is marginally easier, and follows purely from the definition of Galois without making any finiteness assumptions.

**Proposition 1.39** (Fixed field of Galois group)

*If  $K/k$  is Galois and  $F \subset K$  then*

$$K^{\operatorname{Gal}(K/F)} = F$$

*or in other words  $\phi \circ \psi = \operatorname{id}$ , and in particular  $\phi$  is injective.*

*Proof.* Clearly  $F \subseteq K^{\operatorname{Gal}(K/F)}$ . Conversely given  $\alpha \in K \setminus F$ , then  $\deg m_{\alpha, F} > 1$ . Since  $\alpha$  is separable it must have another root  $\beta \in K$ . By Corollary 1.36 there is an element  $\sigma \in \operatorname{Gal}(K/F)$  such that  $\sigma(\alpha) = \beta$ . In other words  $\alpha \notin K^{\operatorname{Gal}(K/F)}$ , which shows the reverse inclusion. □

**Proposition 1.40**

*Let  $K/k$  be a field extension and  $H \subseteq \operatorname{Aut}(K/k)$  a finite subgroup then  $K/K^H$  is finite Galois with*

$$H = \operatorname{Gal}(K/K^H)$$

*and order equal to  $[K : K^H]$ . In particular for a finite Galois Extension  $K/k$  we have  $\psi \circ \phi = \operatorname{id}$ .*

*Proof.* Firstly observe that trivially  $H \subseteq \operatorname{Aut}(K/K^H)$ . If we know that  $[K : K^H] < \infty$ , then by Definition 1.16 and Proposition 1.22 we have

$$\#H \leq \# \operatorname{Aut}(K/K^H) \leq [K : K^H]_s \leq [K : K^H]$$

We can prove equality everywhere if we show that  $[K : K^H] \leq \#H$ , which is shown either by Lemma 1.41 or Lemma 1.42. Note equality also shows that  $K/K^H$  is finite Galois by Proposition 1.37. □

We present two approaches to showing the inequality  $[K : K^H] \leq \#H$ . The first uses independence of characters style argument (see Garling, JMilne), and the second which is more straightforward uses the action of  $H$  to show that every element has degree at most  $\#H$  (Artin).

**Lemma 1.41** (Bound degree of fixed field I)

*Let  $K/k$  be a field extension and  $H \subset \operatorname{Aut}(K/k)$  a finite subgroup. Then  $[K : K^H] \leq \#H$*

*Proof.* Let  $H = \{\sigma_1, \dots, \sigma_n\}$  with  $\sigma_1 = \text{id}$  and  $\alpha_1, \dots, \alpha_m$  a  $K^H$ -basis for  $K$ .

Consider the vector space  $K^n$  and the elements  $\hat{\alpha}_j = (\sigma_1(\alpha_j), \dots, \sigma_n(\alpha_j))$  for  $j = 1 \dots m$ . It's enough to show that these are linearly independent over  $K$ , as that shows  $m \leq n$ .

Let  $S(K) := \{v \in K^m \mid \sum_{j=1}^m v_j \hat{\alpha}_j = 0\}$ , we aim to show that  $S(K) = \{0\}$ . If we also consider  $S(K^H)$ , any non-zero elements will be a  $K^H$  linear-dependence for  $\alpha_1, \dots, \alpha_m$  by considering the first component ( $\sigma_1 = \text{id}$ ). Therefore by assumption  $S(K^H) = \{0\}$ . Finally we see it's enough to show that  $S(K) \neq \{0\} \implies S(K^H) \neq \{0\}$ .

First observe that  $K^\star$  and  $H$  both act on  $S(K)$ . The first by scaling and the second component-wise. This is well-defined because  $v \in S(K)$  if and only if

$$\sum_j v_j \sigma(\alpha_j) = 0 \quad \forall \sigma \in H.$$

Apply  $\tau$  to obtain

$$\sum_j \tau(v_j) (\tau \circ \sigma)(\alpha_j) = 0 \quad \forall \sigma \in H$$

and since multiplication by  $\tau$  permutes  $H$  we see  $\tau(v) \in S(K)$  as required.

If there exists  $0 \neq v \in S(K)$ , consider  $v$  with a minimal number of non-zero components. By scaling we can assume  $\lambda v$  has at least one component in  $K^H$ . The vector  $\tau(\lambda v) - \lambda v$  then has at least one fewer non-zero components, so by minimality must be zero. Since  $\tau$  was arbitrary we see  $0 \neq \lambda v \in S(K^H)$  as required.  $\square$

**Lemma 1.42** (Bound degree of fixed field II)

Let  $K/k$  be a field extension and  $H$  a finite subgroup of  $\text{Aut}(K/k)$ . Then  $K/K^H$  is finite separable, and simple, with  $[K : K^H] \leq \#H$

*Proof.* We show that  $K/K^H$  is separable and every element has degree at most  $\#H$ . For any  $\alpha \in K$ , consider the orbit  $H(\alpha) = \{\sigma(\alpha) \mid \sigma \in H\}$ , which is of order at most  $\#H$ . Then the polynomial

$$f(X) = \prod_{\beta \in H(\alpha)} (X - \beta)$$

has  $\alpha$  as a root and is separable by Proposition 1.9. Furthermore  $f^\tau = f$  because  $\tau$  permutes  $H(\alpha)$  (it's injective and hence bijective). Therefore  $f \in K^H[X]$  and  $m_{\alpha, K^H} \mid f$ . We see that  $\alpha$  has degree at most  $\#H$  and is separable by Lemma 1.11.

If  $K/k$  is finite, then a-fortiori  $K/K^H$  is finite, so we may apply the Primitive Element Theorem 1.25 directly to show the result.

More generally let  $K^H(\alpha)$  be a simple subfield of  $K$  of maximal degree. This exists because the degree of  $\alpha$  is bounded above by  $\#H$ . We claim  $K^H(\alpha) = K$ , for if not then  $K^H \subset K^H(\alpha) \subset K^H(\alpha, \beta)$  is a finite separable extension of  $K^H$ , whence it must be simple by the Primitive Element Theorem 1.25, contradicting maximality. Finally the degree of  $[K : K^H]$  is the degree of  $\alpha$ , which we've seen is bounded above by  $\#H$ .  $\square$

Now we may demonstrate straightforward criteria for subfield to be normal

**Proposition 1.43**

Let  $K/k$  be a finite Galois extension and  $F \subset K$  a subfield.

Then  $F/k$  is Galois if and only if  $\text{Gal}(K/F) \triangleleft \text{Gal}(K/k)$  is normal. In this case we have a canonical isomorphism

$$\text{Gal}(K/k) / \text{Gal}(K/F) \rightarrow \text{Gal}(F/k)$$

*Proof.* Recall from Corollary 1.35 we have  $F/k$  is normal iff  $\sigma(F) = F$  for all  $\sigma \in \text{Gal}(K/k)$ . We also observe that

$$\text{Gal}(K/\sigma(F)) = \sigma \text{Gal}(K/F) \sigma^{-1}$$

By the correspondence  $\text{Gal}(K/F) = \text{Gal}(K/F') \iff F = F'$ .

Therefore

$$\begin{aligned} F/k \text{ normal} &\iff \sigma(F) = F \quad \forall \sigma \in \text{Gal}(K/k) \\ &\iff \text{Gal}(K/\sigma(F)) = \text{Gal}(K/F) \quad \forall \sigma \in \text{Gal}(K/k) \\ &\iff \sigma \text{Gal}(K/F) \sigma^{-1} = \text{Gal}(K/F) \quad \forall \sigma \in \text{Gal}(K/k) \\ &\iff \text{Gal}(K/F) \triangleleft \text{Gal}(K/k) \end{aligned}$$

Finally the natural restriction map is surjective by Corollary 1.34 and has the required kernel.  $\square$

## References