4.2 Sheaves

Sheaves are typically used to organise sets of functions which are defined by a local differentiability condition (e.g. holomorphic, smooth, etc.). However in some cases it may be convenient to take an abstract category-theoretic approach as the proofs become more conceptual and naturally lead to further generalisations where the topological space is replaced by an arbitrary category. However for the sheafification construction we take (the standard) concrete approach as this is simpler and nicely coincides with the notion of "étale space".

In general we assume that the value category $\mathcal{C}$ admits arbitrary products and equalisers (and therefore also limits, and in particular terminal objects), as well as filtered colimits / direct limits. In some cases we may also require it to be algebraic.

Definition 4.2.1 (Abstract Presheaf)
*Let X be a topological space, $\mathcal{B}$ a base for the topology and $\mathcal{C}$ a category. A $\mathcal{B}$-**presheaf** on X is a functor*

$$\mathcal{F} : (\mathcal{B}, \supseteq) \longrightarrow \mathcal{C}$$

from the poset category of basic open subsets to $\mathcal{C}$ ordered by reverse inclusion for which

$$\mathcal{F}(\emptyset) = \star \text{ is a terminal object in } \mathcal{C}$$

This means for every pair of open subsets $U, V \in \mathcal{B}$ such that $V \subseteq U$ there exist morphisms $\rho_{UV} : \mathcal{F}(U) \to \mathcal{F}(V)$ which satisfy the following properties

a) $\rho_{UU} = 1_{\mathcal{F}(U)}$
b) $\rho_{VW} \circ \rho_{UV} = \rho_{UW}$ for any basic open subset $W \subset \mathcal{B}$

A morphism of presheaves is precisely a natural transformation. That is $\alpha : \mathcal{F} \to \mathcal{G}$ is a family of morphisms $\alpha_U : \mathcal{F}(U) \to \mathcal{G}(U)$ for which

$$\rho_{UV} \circ \alpha_U = \alpha_V \circ \rho_{UV}$$

This determines a category $\mathbf{PSh}(\mathcal{B}; \mathcal{C})$. We may omit the value category $\mathcal{C}$ if there are no additional hypotheses.

Definition 4.2.2 (Abstract Sheaf)
Let X be a topological space, $\mathcal{B}$ a base for the topology closed under finite intersections and $\mathcal{F}$ a $\mathcal{B}$-presheaf.

For every family of non-empty basic open subsets $\{U_i\}_{i \in I}$ with $U = \bigcup_{i \in I} U_i$, consider the sequence of morphisms

$$\mathcal{F}(U) \xrightarrow{\ e\ } \prod_{i \in I} \mathcal{F}(U_i) \underset{d_1}{\overset{d_0}{\rightrightarrows}} \prod_{(i,j) \in I \times I} \mathcal{F}(U_i \cap U_j)$$

where $e = \left(\rho_{UU_i}\right)_{i \in I}$, $d_0 = \left(\rho_{U_i(U_i \cap U_j)} \circ \pi_i\right)_{(i,j) \in I \times I}$ and $d_1 = \left(\rho_{U_j(U_i \cap U_j)} \circ \pi_j\right)_{(i,j) \in I \times I}$.

*We say that $\mathcal{F}$ is a $\mathcal{B}$-**sheaf** if e is an equaliser for (d_0, d_1) for all such families of open subsets.*

This determines a full subcategory $\mathbf{Sh}(\mathcal{B}; \mathcal{C})$ of $\mathbf{PSh}(\mathcal{B}; \mathcal{C})$.

*When $\mathcal{B}$ is just the collection of all open subsets of X then we say $\mathcal{F}$ is a **sheaf**, and we denote the category by $\mathbf{Sh}(X; \mathcal{C})$.*

It may be convenient to write the sheaf condition as the existence of a limit.

Lemma 4.2.3 (Sheaf Condition as a Limit)
Let $\mathcal{F}$ be a presheaf on X and $U = \bigcup_{i \in I} U_i$ a cover; write $U_{ij} := U_i \cap U_j$ for $(i,j) \in I \times I$, and let $d_0, d_1 : \prod_{i \in I} \mathcal{F}(U_i) \to \prod_{(i,j) \in I \times I} \mathcal{F}(U_{ij})$ be as in (4.2.2).

Let $I^{(1)}$ be the small category with objects $I \sqcup (I \times I)$ and, for each $(i,j) \in I \times I$, morphisms $u_{ij} : i \to (i,j)$ and $v_{ij} : j \to (i,j)$ (together with identities, and no further morphisms), and let $D_{\mathcal{F}} : I^{(1)} \to \mathcal{C}$ be the diagram

$$D_{\mathcal{F}}(i) := \mathcal{F}(U_i), \quad D_{\mathcal{F}}((i,j)) := \mathcal{F}(U_{ij}), \quad D_{\mathcal{F}}(u_{ij}) := \rho_{U_i U_{ij}}, \quad D_{\mathcal{F}}(v_{ij}) := \rho_{U_j U_{ij}}$$

Then, as $\mathcal{C}$ has arbitrary products, the contravariant functors $\mathcal{C}^{op} \to \mathbf{Set}$

$$X \mapsto \left\{ \xi : X \to \prod_{i \in I} \mathcal{F}(U_i) \;\middle|\; d_0 \circ \xi = d_1 \circ \xi \right\}, \quad X \mapsto \{\text{cones over } D_{\mathcal{F}} \text{ with apex } X\}$$

© David Rufino CC BY-NC-ND 4.0

(both via precomposition) are naturally isomorphic, via $\xi \mapsto (\pi_i \circ \xi)_{i\in I}$*, extended to* $(i,j) \in I \times I$ *by* $D_{\mathcal{F}}(u_{ij}) \circ \pi_i \circ \xi$*. Consequently* $D_{\mathcal{F}}$ *has* $\left(\mathcal{F}(U), \{\rho_{UU_i}\}_{i\in I}, \{\rho_{UU_{ij}}\}_{(i,j)\in I\times I}\right)$ *as a limit over* $I^{(1)}$ *if and only if* $\mathcal{F}$ *satisfies the sheaf condition for this cover (4.2.2).*

Proof. Given ξ with $d_0 \circ \xi = d_1 \circ \xi$, set $\xi_i := \pi_i \circ \xi$ and $\xi_{(i,j)} := D_{\mathcal{F}}(u_{ij}) \circ \xi_i$; then also $\xi_{(i,j)} = D_{\mathcal{F}}(v_{ij}) \circ \xi_j$, since both sides equal $\pi_{(i,j)} \circ d_0 \circ \xi = \pi_{(i,j)} \circ d_1 \circ \xi$ by definition of d_0, d_1. So $\left(X, \{\xi_k\}_{k\in I^{(1)}}\right)$ is a cone over $D_{\mathcal{F}}$, the only nontrivial conditions being exactly this equality at u_{ij}, v_{ij}.

Conversely, given a cone $\left(X, \{\xi_k\}_{k\in I^{(1)}}\right)$, bundling $(\xi_i)_{i\in I}$ via the universal property of the product (2.6.81) gives $\xi := (\xi_i)_i : X \to \prod_{i\in I} \mathcal{F}(U_i)$ with $\pi_i \circ \xi = \xi_i$; then for all (i,j), $\pi_{(i,j)} \circ d_0 \circ \xi = D_{\mathcal{F}}(u_{ij}) \circ \xi_i = \xi_{(i,j)} = D_{\mathcal{F}}(v_{ij}) \circ \xi_j = \pi_{(i,j)} \circ d_1 \circ \xi$, using the cone conditions at u_{ij}, v_{ij}, so $d_0 \circ \xi = d_1 \circ \xi$.

These two constructions are mutually inverse, directly from $\pi_i \circ \xi = \xi_i$ and the universal property of the product, and are evidently compatible with precomposition, i.e. natural in X: for $\phi : X' \to X$, both sides simply precompose ξ, resp. every ξ_k, by ϕ. So $D_{\mathcal{F}}$ has a limit over $I^{(1)}$ iff (d_0, d_1) has an equaliser, and the two agree. ∎

Proposition 4.2.4 (Sheaf Conditions for Concrete Categories)
Let X *be a topological space,* $\mathcal{B}$ *a base for the topology closed under finite intersections and* $\mathcal{C}$ *is an algebraic category (2.6.185). Then a* $\mathcal{B}$*-presheaf* $\mathcal{F}$ *is a sheaf if and only if it satisfies the following conditions, for every open cover* $U = \bigcup_{i\in I} U_i$ *with* U_i *non-empty*

a) $\sigma, \tau \in \mathcal{F}(U)$ *such that* $\rho_{UU_i}(\sigma) = \rho_{UU_i}(\tau)$ *for all* $i \in I$ *implies* $\sigma = \tau$
b) $\sigma_i \in \mathcal{F}(U_i)$ *such that* $\rho_{U_i(U_i\cap U_j)}(\sigma_i) = \rho_{U_j(U_i\cap U_j)}(\sigma_j)$ *for all* $(i,j) \in I \times I$ *implies there exists* $\sigma \in \mathcal{F}(U)$ *such that* $\rho_{UU_i}(\sigma) = \sigma_i$.

Proof. Suppose that $\mathcal{F}$ is a sheaf in the abstract sense. By assumption the forgetful functor $\mathcal{C} \to \mathbf{Set}$ preserves limits and so by (2.6.198) e is injective and

$$\mathrm{Im}(e) = \left\{(\sigma_i)_{i\in I} \;\middle|\; \sigma_i|_{U_i\cap U_j} = \sigma_j|_{U_i\cap U_j}\right\}$$

which gives both concrete sheaf conditions.

Conversely this condition ensures the sequence is an equaliser in **Set**, and so by hypothesis is so in $\mathcal{C}$. ∎

4.2.1 Stalks

Definition 4.2.5 (Stalk of a presheaf)
Let X *be a topological space,* $\mathcal{B}$ *a base closed under finite intersections and* $\mathcal{F}$ *a* $\mathcal{B}$*-presheaf.*

Let Z *be an irreducible subset of* X*, then the family of open subsets* $U \in \mathcal{B}$ *meeting* Z *is closed under finite intersections (4.1.60) and therefore forms a directed preorder (2.6.124) ordered by reverse inclusion.*

Therefore we may define the ***stalk*** *at* Z *by the direct limit (2.6.126)*

$$\mathcal{F}_Z := \varinjlim_{U\in\mathcal{B},\, U\cap Z\neq\emptyset} \mathcal{F}(U)$$

This means there is a family of morphisms $\{\rho_{UZ} : \mathcal{F}(U) \to \mathcal{F}_Z\}$ *such that*

a) all $V \in \mathcal{B}$ *with* $V \subset U$ *intersecting* Z *satisfy* $\rho_{VZ} \circ \rho_{UV} = \rho_{UZ}$*,*
b) any collection of morphisms $\alpha_U : \mathcal{F}(U) \to \mathcal{A}$ *satisfying* $\alpha_U = \alpha_V \circ \rho_{UV}$ *factors uniquely through a morphism* $\alpha_Z : \mathcal{F}_Z \to \mathcal{A}$

When $Z = \{x\}$ *we denote this simply by* $\mathcal{F}_x$*.*

Proposition 4.2.6 (Explicit form of Stalk)
Consider the situation (4.2.5) where the value category $\mathcal{C}$ *is algebraic (2.6.185). Then*

$$U(\mathcal{F}_Z) = (U \circ \mathcal{F})_Z$$

and so the stalk $\mathcal{F}_Z$ *may be constructed explicitly as follows*

$$\mathcal{F}_Z := \{(U,\sigma) \mid U \in \mathcal{B}, \sigma \in \mathcal{F}(U)\}/\sim$$
$$(U,\sigma) \sim (V,\tau) \iff \exists W \in \mathcal{B} \text{ s.t. } W \cap Z \neq \emptyset \text{ and } \sigma|_W = \tau|_W$$

with morphisms

© David Rufino CC BY-NC-ND 4.0

$$\begin{aligned}\rho_{UZ} : \mathcal{F}(U) &\to \mathcal{F}_Z \\ \sigma &\to [(U, \sigma)]\end{aligned}$$

In particular every element of the stalk $\sigma_Z \in \mathcal{F}_Z$ is the image of some section $\sigma \in \mathcal{F}(U)$ under ρ_{UZ} for some $U \in \mathcal{B}$.

Proof. By (2.6.190), the forgetful functor $U : \mathcal{C} \to \mathbf{Set}$ creates directed colimits, and so the result follows from (2.6.129). ∎

Proposition 4.2.7 (Functoriality of Stalks)
Let X be a topological space with base $\mathcal{B}$ closed under finite intersections, $\phi : \mathcal{F} \to \mathcal{G}$ a morphism of $\mathcal{B}$-presheaves and $Z \subset X$ an irreducible subset. Then there exists a unique morphism $\phi_Z : \mathcal{F}_Z \to \mathcal{G}_Z$ ensuring the following diagram commute for all open subsets U meeting Z

$$\begin{array}{ccc} \mathcal{F}(U) & \xrightarrow{\phi_U} & \mathcal{G}(U) \\ {\scriptstyle \rho_{UZ}}\downarrow & & \downarrow{\scriptstyle \rho_{UZ}} \\ \mathcal{F}_Z & \xrightarrow{\phi_Z} & \mathcal{G}_Z \end{array}$$

Furthermore $(\psi \circ \phi)_Z = \psi_Z \circ \phi_Z$. In the explicit case (4.2.6) then ϕ_Z is given by

$$[(U, \sigma)] \to [(U, \phi_U(\sigma))]$$

This also means that for any $A \in \mathcal{C}$ the bijection

$$\mathrm{Mor}(\mathcal{F}_Z, A) \longrightarrow \left(\mathcal{F}|_{\mathcal{N}_Z} \downarrow A\right)$$

is natural in $\mathcal{F}$, where $\mathcal{N}_Z$ is the category of open sets intersecting Z ordered by reverse inclusion.

Lemma 4.2.8 (Lifting Stalks)
Suppose we are in the situation of (4.2.6) and $\sigma \in \mathcal{F}(U)$, $\tau \in \mathcal{F}(V)$ are sections such that $Z \cap U \cap V \neq \emptyset$.

a) Then $\sigma_Z = \tau_Z$ if and only if there is a neighbourhood $W \in \mathcal{B}$, $W \subseteq U \cap V$, meeting Z such that $\sigma|_W = \tau|_W$.
b) If $\sigma_x = \tau_x$ for all $x \in U \cap V$, then there is an open cover $U \cap V = \bigcup_i U_i$ such that $\sigma|_{U_i} = \tau|_{U_i}$
c) If in addition $\mathcal{F}$ is a sheaf then $\sigma|_{U\cap V} = \tau|_{U\cap V}$.

Proof. Drafted by Claude Sonnet, reviewed by David Rufino.

By (4.2.6), $\mathcal{F}_Z = \{(W, \rho) \mid W \in \mathcal{B}, \rho \in \mathcal{F}(W)\}/\sim$ with $(W, \rho) \sim (W', \rho')$ iff there is $N \in \mathcal{B}$, $N \subseteq W \cap W'$, $N \cap Z \neq \emptyset$, with $\rho|_N = \rho'|_N$; and $\sigma_Z = [(U, \sigma)]$, $\tau_Z = [(V, \tau)]$. So $\sigma_Z = \tau_Z$ if and only if $(U, \sigma) \sim (V, \tau)$, which is exactly the stated condition, proving a).

For b), suppose $\sigma_x = \tau_x$ for all $x \in U \cap V$. For each such x, a) (applied with $Z := \{x\}$, an irreducible subset meeting $U \cap V$) gives $W_x \in \mathcal{B}$, $W_x \subseteq U \cap V$, $x \in W_x$, with $\sigma|_{W_x} = \tau|_{W_x}$. Then $\{W_x\}_{x\in U\cap V}$ is an open cover of $U \cap V$ with the required property.

For c), as $\mathcal{B}$ is closed under finite intersections and $U, V \in \mathcal{B}$, $U \cap V \in \mathcal{B}$, so $\sigma|_{U\cap V}, \tau|_{U\cap V} \in \mathcal{F}(U \cap V)$ are defined. By b) there is an open cover $U \cap V = \bigcup_i W_i$ with $\sigma|_{W_i} = \tau|_{W_i}$, that is $\rho_{(U\cap V)W_i}(\sigma|_{U\cap V}) = \rho_{(U\cap V)W_i}(\tau|_{U\cap V})$ for all i; as $\mathcal{F}$ is a sheaf, the separation condition (4.2.4) gives $\sigma|_{U\cap V} = \tau|_{U\cap V}$. ∎

Proposition 4.2.9 (Stalks when restricting to a base)
Let $\mathcal{B}_2 \subset \mathcal{B}_1$ be bases for X which generate the same topology. Suppose $\mathcal{F}$ is a $\mathcal{B}_1$-presheaf and $Z \subset X$ is an irreducible subset. Then there is a unique morphism making the following diagram commute for all $U \in \mathcal{B}_2$ intersecting Z

$$\begin{array}{ccc} \mathcal{F}|_{\mathcal{B}_2}(U) & = & \mathcal{F}(U) \\ {\scriptstyle \rho_{UZ}}\downarrow & & \downarrow{\scriptstyle \rho_{UZ}} \\ \left(\mathcal{F}|_{\mathcal{B}_2}\right)_Z & \xrightarrow{\sim} & \mathcal{F}_Z \end{array}$$

and it is an isomorphism.

In the situation of (4.2.6) the map is of the obvious form.

Proof. Let $\mathcal{I}_2$ and $\mathcal{I}_1$ be the diagram of open subsets U of $\mathcal{B}_2$ and $\mathcal{B}_1$ meeting Z, ordered by reverse inclusion. Then $\mathcal{I}_2$ is a full subcategory of $\mathcal{I}_1$ which is cofinal (2.6.127). So the result follows from (2.6.128)

© David Rufino CC BY-NC-ND 4.0

In the explicit case this may be shown more directly. ∎

Lemma 4.2.10 (Pointwise Injectivity and Injectivity on Stalks)
Let $\mathcal{C}$ be algebraic (2.6.185) and $\phi : \mathcal{F} \to \mathcal{G}$ a morphism of presheaves such that ϕ_U is injective for all $U \subset X$. Then ϕ_x is injective for all $x \in X$.

Conversely, if ϕ is a morphism of sheaves and ϕ_x is injective for all $x \in X$, then ϕ_U is injective for all $U \subset X$.

Proof. Drafted by Claude Sonnet, reviewed by David Rufino.

Fix $x \in X$ and suppose $a, b \in \mathcal{F}_x$ with $\phi_x(a) = \phi_x(b)$. By (4.2.6), $a = (\sigma_1)_x$, $b = (\sigma_2)_x$ for some $\sigma_1 \in \mathcal{F}(V_1)$, $\sigma_2 \in \mathcal{F}(V_2)$, $V_1, V_2 \in \mathcal{B}$ containing x; setting $V := V_1 \cap V_2 \in \mathcal{B}$, $\sigma := \sigma_1|_V$, $\tau := \sigma_2|_V$, we have $\sigma, \tau \in \mathcal{F}(V)$ with $\sigma_x = a$, $\tau_x = b$. Then

$$(\phi_V(\sigma))_x = \phi_x(\sigma_x) = \phi_x(a) = \phi_x(b) = \phi_x(\tau_x) = (\phi_V(\tau))_x$$

so by (4.2.8).a) (with both opens taken to be V) there is $W \in \mathcal{B}$, $W \subseteq V$, $x \in W$, with $\phi_V(\sigma)|_W = \phi_V(\tau)|_W$, that is $\phi_W(\sigma|_W) = \phi_W(\tau|_W)$. As ϕ_W is injective, $\sigma|_W = \tau|_W$, so $a = \sigma_x = (\sigma|_W)_x = (\tau|_W)_x = \tau_x = b$.

Conversely, fix $U \subset X$ and $\sigma, \tau \in \mathcal{F}(U)$ with $\phi_U(\sigma) = \phi_U(\tau)$. For every $x \in U$, $\phi_x(\sigma_x) = (\phi_U(\sigma))_x = (\phi_U(\tau))_x = \phi_x(\tau_x)$, so $\sigma_x = \tau_x$ as ϕ_x is injective. As $\mathcal{F}$ is a sheaf, (4.2.8).c) (taking both opens of the lemma to be U) gives $\sigma = \tau$. ∎

Proposition 4.2.11 (Isomorphisms of Presheaves and Sheaves)
Let $\phi : \mathcal{F} \to \mathcal{G}$ be a morphism of presheaves (resp. sheaves). Then the following are equivalent

a) ϕ is an isomorphism
b) ϕ_U is an isomorphism for all $U \subset X$

If $\mathcal{C}$ is algebraic (2.6.185), then a), b) are further equivalent to

c) ϕ_U is bijective for all $U \subset X$

If moreover $\mathcal{F}, \mathcal{G}$ are sheaves, a)–c) are equivalent to

d) $\phi_x : \mathcal{F}_x \to \mathcal{G}_x$ is an isomorphism for all $x \in X$

Proof. Drafted by Claude Sonnet, reviewed by David Rufino.

$b) \Longrightarrow a)$: If each ϕ_U is an isomorphism, define $\psi_U := \phi_U^{-1}$; for $V \subseteq U$, inverting both sides of the naturality square $\phi_V \circ \rho_{UV} = \rho_{UV} \circ \phi_U$ gives $\rho_{UV} \circ \psi_U = \psi_V \circ \rho_{UV}$, so $\psi := \{\psi_U\}$ is a morphism of presheaves (resp. sheaves), and $\psi\phi = 1_{\mathcal{F}}$, $\phi\psi = 1_{\mathcal{G}}$ hold pointwise, hence as morphisms of presheaves. So ϕ is an isomorphism.

$a) \Longrightarrow b)$: If ϕ is an isomorphism with inverse ψ, then $\phi_U\psi_U = 1_{\mathcal{G}(U)}$ and $\psi_U\phi_U = 1_{\mathcal{F}(U)}$ for every U, so ϕ_U is an isomorphism.

$b) \Longleftrightarrow c)$ (when $\mathcal{C}$ is algebraic): an isomorphism is in particular bijective (a functor preserves isomorphisms, and an isomorphism in **Set** is a bijection), while conversely a bijective morphism is an isomorphism in $\mathcal{C}$ as U reflects isomorphisms (2.6.190) apply this pointwise at each U.

$b) \Longrightarrow d)$: with $\psi := \{\psi_U\}$ as in $b) \Longrightarrow a)$ above, ψ is a morphism of sheaves with $\psi\phi = 1_{\mathcal{F}}$, $\phi\psi = 1_{\mathcal{G}}$. By functoriality of stalks (4.2.7), $\psi_x \circ \phi_x = (\psi\phi)_x = 1_{\mathcal{F}_x}$ and $\phi_x \circ \psi_x = (\phi\psi)_x = 1_{\mathcal{G}_x}$, so ϕ_x is an isomorphism.

$d) \Longrightarrow c)$: Suppose ϕ_x is an isomorphism, in particular bijective, for every $x \in X$. By (4.2.10), ϕ_U is injective for every $U \subset X$. For surjectivity: fix $U \subset X$ and $\tau \in \mathcal{G}(U)$. For each $x \in U$, surjectivity of ϕ_x gives $a_x \in \mathcal{F}_x$ with $\phi_x(a_x) = \tau_x$; write $a_x = (\rho_x)_x$ for some $\rho_x \in \mathcal{F}(W_x)$, $W_x \in \mathcal{B}$, $x \in W_x \subseteq U$ (shrinking W_x if necessary). Then $\left(\phi_{W_x}(\rho_x)\right)_x = \phi_x\left((\rho_x)_x\right) = \phi_x(a_x) = \tau_x = \left(\tau|_{W_x}\right)_x$, so by (4.2.8).a) there is $W_{x'} \in \mathcal{B}$, $x \in W_{x'} \subseteq W_x$, with $\phi_{W_x}(\rho_x)|_{W_{x'}} = \tau|_{W_{x'}}$; write $\sigma_x := \rho_x|_{W_{x'}} \in \mathcal{F}(W_{x'})$, so that $\phi_{W_{x'}}(\sigma_x) = \tau|_{W_{x'}}$ exactly.

The $\{W_{x'}\}_{x \in U}$ cover U; for $x, y \in U$, on $W_{x'} \cap W_{y'} \in \mathcal{B}$,

$$\phi_{W_{x'} \cap W_{y'}}\left(\sigma_x|_{W_{x'} \cap W_{y'}}\right) = \phi_{W_{x'}}(\sigma_x)|_{W_{x'} \cap W_{y'}} = \tau|_{W_{x'} \cap W_{y'}} = \phi_{W_{y'}}(\sigma_y)|_{W_{x'} \cap W_{y'}} = \phi_{W_{x'} \cap W_{y'}}\left(\sigma_y|_{W_{x'} \cap W_{y'}}\right)$$

so as $\phi_{W_{x'} \cap W_{y'}}$ is injective, $\sigma_x|_{W_{x'} \cap W_{y'}} = \sigma_y|_{W_{x'} \cap W_{y'}}$. As $\mathcal{F}$ is a sheaf, the gluing condition (4.2.4) gives $\sigma \in \mathcal{F}(U)$ with $\sigma|_{W_{x'}} = \sigma_x$ for all x. Then $\phi_U(\sigma)|_{W_{x'}} = \phi_{W_{x'}}\left(\sigma|_{W_{x'}}\right) = \phi_{W_{x'}}(\sigma_x) = \tau|_{W_{x'}}$ for every $x \in U$, so as $\mathcal{G}$ is a sheaf, the separation condition (4.2.4) gives $\phi_U(\sigma) = \tau$. So ϕ_U is surjective; being also injective, ϕ_U is bijective, that is c). ∎

© David Rufino CC BY-NC-ND 4.0

The following somewhat tedious result shows that morphisms are uniquely determined by morphisms on basic open sets.

Proposition 4.2.12
Let X be a topological space, $\mathcal{B}$ a base for the topology closed under finite intersections. Then the functor

$$(-)|_{\mathcal{B}} : \mathbf{Sh}(X;\mathcal{C}) \to \mathbf{Sh}(\mathcal{B};\mathcal{C})$$

is full and faithful.

Proof. Let $\phi, \psi : \mathcal{F} \to \mathcal{G}$ be morphisms which agree on basic open subsets $U \in \mathcal{B}$. Consider an arbitrary open subset $U \subset X$ then by definition $U = \bigcup_{i\in I} U_i$ for $U_i \in \mathcal{B}$. By assumption there is a commutative diagram

$$\begin{array}{ccc} \mathcal{F}(U) & \xrightarrow{e} & \prod_{i\in I}\mathcal{F}(U_i) \\ {\scriptstyle\psi_U}\downarrow\downarrow{\scriptstyle\phi_U} & & \downarrow{\scriptstyle\prod_{i\in I}\phi|_{U_i} = \prod_{i\in I}\psi|_{U_i}} \\ \mathcal{G}(U) & \xrightarrow{e} & \prod_{i\in I}\mathcal{G}(U_i) \end{array}$$

which shows $e \circ \phi_U = e \circ \psi_U$. But e is a monomorphism (2.6.136) (being part of an equaliser) so $\phi_U = \psi_U$. Therefore the functor is faithful.

Let $\phi : \mathcal{F}|_{\mathcal{B}} \to \mathcal{G}|_{\mathcal{B}}$ be a morphism of $\mathcal{B}$-sheaves. For an open set U choose a basic open cover $U = \bigcup_{i\in I} U_i$ then we have a commutative diagram

$$\begin{array}{ccccc} \mathcal{F}(U) & \xrightarrow{e} & \prod_{i\in I}\mathcal{F}(U_i) & \rightrightarrows & \prod_{(i,j)\in I\times I}\mathcal{F}(U_i\cap U_j) \\ \downarrow{\scriptstyle\tilde{\phi}_U} & & \downarrow{\scriptstyle\prod_{i\in I}\phi_{U_i}} & & \downarrow{\scriptstyle\prod_{(i,j)\in I\times I}\phi_{U_i\cap U_j}} \\ \mathcal{G}(U) & \xrightarrow{e} & \prod_{i\in I}\mathcal{G}(U_i) & \rightrightarrows & \prod_{(i,j)\in I\times I}\mathcal{G}(U_i\cap U_j) \end{array}$$

where for the right hand square we must pick the top arrows or bottom arrows. Chasing the diagram shows that $\left(\prod_{i\in I}\phi_{U_i}\right)\circ e$ is equalised by the bottom two arrows, and so the factorisation exists by the universal property and $\tilde{\phi}_U$ the unique morphism which makes the left square commute, i.e. such that $\phi_{U_i}\circ\rho_{UU_i} = \rho_{UU_i}\circ\tilde{\phi}_U$ for all $i \in I$.

We claim that for any basic open subset $W \subset U$ the following diagram commutes

$$\begin{array}{ccccc} \mathcal{F}(U) & \longrightarrow & \mathcal{F}(W) & \xrightarrow{e} & \prod_{i\in I}\mathcal{F}(W\cap U_i) \\ \downarrow{\scriptstyle\tilde{\phi}_U} & & \downarrow{\scriptstyle\phi_W} & & \downarrow{\scriptstyle\prod_{i\in I}\phi_{W\cap U_i}} \\ \mathcal{G}(U) & \longrightarrow & \mathcal{G}(W) & \xrightarrow{e} & \prod_{i\in I}\mathcal{G}(W\cap U_i) \end{array}$$

and $\tilde{\phi}_U$ is the unique morphism such that this commutes for all $W \in \mathcal{B}$. The right hand square commutes by assumption. As e is a monomorphism it is enough to show that

$$\left(\prod_{i\in I}\phi_{W\cap U_i}\right)\circ e\circ\rho_{UW} = e\circ\phi_W\circ\rho_{UW} = e\circ\rho_{UW}\circ\tilde{\phi}_U$$

First observe we have a commutative diagram

$$\begin{array}{ccc} \mathcal{F}(U) & \xrightarrow{e} & \prod_{i\in I}\mathcal{F}(U_i) \\ {\scriptstyle\rho_{UW}}\downarrow & & \downarrow{\scriptstyle\prod_{i\in I}\rho_{U_i(W\cap U_i)}} \\ \mathcal{F}(W) & \xrightarrow{e} & \prod_{i\in I}\mathcal{F}(W\cap U_i) \end{array}$$

which follows because both directions are equal to the diagonal $\left(\rho_{U(W\cap U_i)}\right)_{i\in I}$. Therefore we may combine the two diagrams to find

© David Rufino CC BY-NC-ND 4.0

$$\begin{aligned}
e \circ \phi_W \circ \rho_{UW} &= \left(\prod_{i\in I} \phi_{W\cap U_i}\right) \circ e \circ \rho_{UW} \\
&= \left(\prod_{i\in I} \phi_{W\cap U_i}\right) \circ \left(\prod_{i\in I} \rho_{U_i(W\cap U_i)}\right) \circ e \\
&= \left(\prod_{i\in I} \phi_{W\cap U_i} \circ \rho_{U_i(W\cap U_i)}\right) \circ e \\
&= \left(\prod_{i\in I} \rho_{U_i(W\cap U_i)} \circ \phi_{U_i}\right) \circ e \\
&= \left(\prod_{i\in I} \rho_{U_i(W\cap U_i)}\right) \circ \left(\prod_{i\in I} \phi_{U_i}\right) \circ e \\
&= \left(\prod_{i\in I} \rho_{U_i(W\cap U_i)}\right) \circ e \circ \tilde{\phi}_U \\
&= e \circ \rho_{UW} \circ \tilde{\phi}_U
\end{aligned}$$

As e is a monomorphism (2.6.136) then $\phi_W \circ \rho_{UW} = \rho_{UW} \circ \tilde{\phi}_U$ as required. Then $\tilde{\phi}_U$ is the unique morphism such that this holds for all $W \in \mathcal{B}$ and $W \subset U$, because it is already uniquely determined by $W = U_i$. This also shows that $\tilde{\phi}_W = \phi_W$.

In particular $\tilde{\phi}_U$ does not depend on the choice of cover $\{U_i\}$ of U: if $\tilde{\phi}'_U$ is constructed from a different basic cover of U, it satisfies the same property $\phi_W \circ \rho_{UW} = \rho_{UW} \circ \tilde{\phi}'_U$ for all $W \in \mathcal{B}$, $W \subset U$; taking W to range over the original cover $\{U_i\}$ shows $\tilde{\phi}'_U$ also satisfies the equalising property defining $\tilde{\phi}_U$, so $\tilde{\phi}'_U = \tilde{\phi}_U$ by the uniqueness established above.

Finally for an arbitrary open subset $V = \bigcup_{i\in I} W_i \subset U$, $W_i \in \mathcal{B}$, we wish to show we have a commutative diagram

$$\begin{array}{ccccc}
\mathcal{F}(U) & \longrightarrow & \mathcal{F}(V) & \xrightarrow{e} & \prod_{i\in I} \mathcal{F}(W_i) \\
\downarrow{\scriptstyle \tilde{\phi}_U} & & \downarrow{\scriptstyle \tilde{\phi}_V} & & \downarrow{\scriptstyle \prod_{i\in I} \phi_{W_i}} \\
\mathcal{G}(U) & \longrightarrow & \mathcal{G}(V) & \xrightarrow{e} & \prod_{i\in I} \mathcal{G}(W_i)
\end{array}$$

We've already shown that the large square and right square commute. This shows that

$$e \circ \tilde{\phi}_V \circ \rho_{UV} = e \circ \rho_{UV} \circ \tilde{\phi}_U$$

and as e is a monomorphism (2.6.136) the left square also commutes.

Therefore $\tilde{\phi}$ is a morphism of sheaves and $\tilde{\phi}|_{\mathcal{B}} = \phi$. Therefore the functor is full.

Alternatively, when $\mathcal{C}$ is algebraic (2.6.185), this can be shown more directly by working with elements via the forgetful functor $U : \mathcal{C} \to \mathbf{Set}$, citing (4.2.4) throughout with the base taken to be all of $\mathcal{O}(X)$ (a valid choice of base, since $\mathcal{F}, \mathcal{G}$ are sheaves on all of X (4.2.2)).

Drafted by Claude Sonnet, reviewed by David Rufino.

Faithful. Let $\phi, \psi : \mathcal{F} \to \mathcal{G}$ agree on $U \in \mathcal{B}$, and fix an arbitrary open $V \subset X$ and $\sigma \in \mathcal{F}(V)$. Choose, for each $x \in V$, a basic open $U_x \in \mathcal{B}$ with $x \in U_x \subseteq V$. Then

$$\rho_{VU_x}(\phi_V(\sigma)) = \phi_{U_x}\left(\rho_{VU_x}(\sigma)\right) = \psi_{U_x}\left(\rho_{VU_x}(\sigma)\right) = \rho_{VU_x}(\psi_V(\sigma))$$

using naturality of ϕ, ψ and that they agree on $U_x \in \mathcal{B}$. As $\{U_x\}_{x\in V}$ is a cover of V, separation for $\mathcal{G}$ gives $\phi_V(\sigma) = \psi_V(\sigma)$. As σ, V were arbitrary, $\phi = \psi$.

Full. Let $\phi : \mathcal{F}|_{\mathcal{B}} \to \mathcal{G}|_{\mathcal{B}}$ be a morphism of $\mathcal{B}$-sheaves; by (4.2.7) this determines stalk maps $\phi_x : \mathcal{F}_x \to \mathcal{G}_x$ for every $x \in X$.

Construction. Fix an open $V \subset X$ and $\sigma \in \mathcal{F}(V)$. Choose, for each $x \in V$, a basic open $U_x \in \mathcal{B}$ with $x \in U_x \subseteq V$, and set $\tau_x := \phi_{U_x}\left(\sigma|_{U_x}\right) \in \mathcal{G}(U_x)$. For $x, y \in V$, on $U_x \cap U_y \in \mathcal{B}$,

$$\tau_x|_{U_x\cap U_y} = \phi_{U_x\cap U_y}\left(\sigma|_{U_x\cap U_y}\right) = \tau_y|_{U_x\cap U_y}$$

© David Rufino CC BY-NC-ND 4.0

using naturality of ϕ and that $\big(\sigma|_{U_x}\big)|_{U_x\cap U_y} = \sigma|_{U_x\cap U_y} = \big(\sigma|_{U_y}\big)|_{U_x\cap U_y}$. So by the gluing condition for $\mathcal{G}$, applied to the cover $\{U_x\}_{x\in V}$ of V, there is $\tau\in\mathcal{G}(V)$ with $\tau|_{U_x}=\tau_x$ for all x; define $\tilde{\phi}_V(\sigma) \coloneqq \tau$. By construction and functoriality of stalks (4.2.7), $\tilde{\phi}_V(\sigma)$ satisfies the property

$$(\star)\qquad \big(\tilde{\phi}_V(\sigma)\big)_x = \big(\tau|_{U_x}\big)_x = (\tau_x)_x = \phi_x\Big(\big(\sigma|_{U_x}\big)_x\Big) = \phi_x(\sigma_x) \qquad \forall x\in V$$

Well-definedness. Suppose $\tau,\tau'\in\mathcal{G}(V)$ both satisfy $(\star)$ for the same σ, that is $\tau_x=\phi_x(\sigma_x)=\tau'_x$ for all $x\in V$. For each $x\in V$, choose $W\in\mathcal{B}$ with $x\in W\subseteq V$; then $(\tau|_W)_x=\tau_x=\tau'_x=(\tau'|_W)_x$, so by (4.2.8).a), applied to the sections $\tau|_W,\tau'|_W\in\mathcal{G}(W)$, there is $W_x\in\mathcal{B}$, $x\in W_x\subseteq W$, with $\tau|_{W_x}=\tau'|_{W_x}$. As $\{W_x\}_{x\in V}$ covers V, separation for $\mathcal{G}$ gives $\tau=\tau'$. So $\tilde{\phi}_V(\sigma)$ does not depend on the choice of cover $\{U_x\}$, and is the unique element of $\mathcal{G}(V)$ satisfying $(\star)$.

$\tilde{\phi}$ restricts to ϕ on $\mathcal{B}$. For $U\in\mathcal{B}$ and $\sigma\in\mathcal{F}(U)$, $\phi_U(\sigma)$ itself satisfies $(\phi_U(\sigma))_x=\phi_x(\sigma_x)$ for $x\in U$ (4.2.7), so by uniqueness $\tilde{\phi}_U(\sigma)=\phi_U(\sigma)$.

$\tilde{\phi}$ is a morphism of presheaves. For $V\subseteq U$ and $\sigma\in\mathcal{F}(U)$, both $\tilde{\phi}_V(\sigma|_V)$ and $\tilde{\phi}_U(\sigma)|_V$ satisfy $(\star)$ for $\sigma|_V$: for $x\in V$, $\big(\tilde{\phi}_V(\sigma|_V)\big)_x=\phi_x\big((\sigma|_V)_x\big)=\phi_x(\sigma_x)=\big(\tilde{\phi}_U(\sigma)\big)_x=\big(\tilde{\phi}_U(\sigma)|_V\big)_x$. By uniqueness, $\tilde{\phi}_V(\sigma|_V)=\tilde{\phi}_U(\sigma)|_V$, so $\tilde{\phi}$ is a morphism of sheaves with $\tilde{\phi}|_{\mathcal{B}}=\phi$. Therefore the functor is full. ∎

4.2.2 Sheafification

Definition 4.2.13 (Sheafification)
*Let X be a topological space, $\mathcal{B}$ a basis of open sets for X closed under finite intersections and $\mathcal{F}$ a $\mathcal{B}$-presheaf (with values in a category $\mathcal{C}$). Then we say that a pair $(\eta_{\mathcal{F}},\mathcal{F}^+)$ is a **sheafification** of $\mathcal{F}$ if*

a) $\mathcal{F}^+$ is a sheaf (with values in a category $\mathcal{C}$)
b) $\eta_{\mathcal{F}}:\mathcal{F}\to\mathcal{F}^+|_{\mathcal{B}}$ is a morphism of $\mathcal{B}$-presheaves
c) for every sheaf $\mathcal{G}$ there is a bijection

$$\begin{aligned}(-)^\sharp : \mathrm{Mor}(\mathcal{F}^+,\mathcal{G}) &\xrightarrow{\sim} \mathrm{Mor}(\mathcal{F},\mathcal{G}|_{\mathcal{B}})\\ \phi &\to \ \phi|_{\mathcal{B}}\circ\eta_{\mathcal{F}}\end{aligned}$$

which is natural in $\mathcal{G}$, with the inverse map denoted by $(-)^\flat$.

In other words $\big(\mathcal{F}^+,(-)^\sharp\big)$ represents the functor $\mathcal{G}\to\mathrm{Mor}(\mathcal{F},\mathcal{G}|_{\mathcal{B}})$.

In this case we denote by $\varepsilon_{\mathcal{G}}:(\mathcal{G}|_{\mathcal{B}})^+\to\mathcal{G}$ the unique morphism which maps to $1_{\mathcal{G}|_{\mathcal{B}}}$ under the bijection.

Proposition 4.2.14 (Sheafification is Functorial)
Using the notation of (4.2.13) suppose there exists a sheafification $(\eta_{\mathcal{F}},\mathcal{F}^+)$ for all pre-sheaves $\mathcal{F}$. Then this can be extended to a functor

$$(-)^+ : \mathbf{PSh}(X;\mathcal{B})\to\mathbf{Sh}(X)$$

uniquely determined by either the condition $(-)^+$ is left-adjoint to $(-)|_{\mathcal{B}}$ or that the following diagram commutes

$$\begin{array}{ccc}\mathcal{F}_1 & \xrightarrow{\ \phi\ } & \mathcal{F}_2\\ {\scriptstyle\eta_{\mathcal{F}_1}}\downarrow & & \downarrow{\scriptstyle\eta_{\mathcal{F}_2}}\\ \mathcal{F}_1^+|_{\mathcal{B}} & \xrightarrow[\phi^+|_{\mathcal{B}}]{\ \ \ \ \ } & \mathcal{F}_2^+|_{\mathcal{B}}\end{array}$$

for all morphisms $\phi:\mathcal{F}_1\to\mathcal{F}_2$. More precisely we have the relationship

$$\phi^+ = \big(\eta_{\mathcal{F}_2}\circ\phi\big)^\flat$$

In particular $(\psi\circ\phi)^+=\psi^+\circ\phi^+$ for all morphisms $\psi:\mathcal{F}_2\to\mathcal{F}_3$ and $\eta_{\mathcal{F}}$ is the unit of the adjunction and $\varepsilon_{\mathcal{G}}$ the counit.

Furthermore $(-)|_{\mathcal{B}}:\mathbf{Sh}(X)\to\mathbf{PSh}(X;\mathcal{B})$ is fully faithful, and ε is a natural isomorphism.

Proof. See (2.6.74).

By (4.2.12), $\mathbf{Sh}(X)\to\mathbf{Sh}(\mathcal{B};\mathcal{C})$ is fully faithful; composing with the (trivially fully faithful) inclusion $\mathbf{Sh}(\mathcal{B};\mathcal{C})\hookrightarrow\mathbf{PSh}(X;\mathcal{B})$ shows $(-)|_{\mathcal{B}}:\mathbf{Sh}(X)\to\mathbf{PSh}(X;\mathcal{B})$ is fully faithful, so by (2.6.72), ε is a natural isomorphism. ∎

© David Rufino CC BY-NC-ND 4.0

Proposition 4.2.15
Using the notation of (4.2.13) suppose there exists a sheafification $(\eta_{\mathcal{F}}, \mathcal{F}^+)$ for $\mathcal{F}$. Then for every $x \in X$ there is a unique morphism making the following diagram commute for all $U \in \mathcal{B}$ containing x

$$\begin{array}{ccc} \mathcal{F}(U) & \xrightarrow{\eta_U} & \mathcal{F}^+(U) \\ \downarrow & & \downarrow \\ \mathcal{F}_x & \dashrightarrow & (\mathcal{F}^+)_x \end{array}$$

Proof. By (4.2.7) and (4.2.9) there is a commutative diagram for all $U \in \mathcal{B}$ and $x \in U$

$$\begin{array}{ccccc} \mathcal{F}(U) & \xrightarrow{\eta_U} & \mathcal{F}^+(U) & = & \mathcal{F}^+(U) \\ \downarrow & & \downarrow & & \downarrow \\ \mathcal{F}_x & \longrightarrow & (\mathcal{F}^+|_{\mathcal{B}})_x & \xrightarrow{\sim} & \mathcal{F}^+_x \end{array}$$

where the bottom left arrow is uniquely determined. This shows that the required map exists and is uniquely determined according to this condition. ∎

Proposition 4.2.16 (Sheafification)
Let $\mathcal{C}$ be an algebraic category (2.6.185) then the sheafification (4.2.13) exists.

More precisely define

$$\mathcal{F}^+(U) := \left\{ s : U \to \coprod_{x \in U} \mathcal{F}_x \;\middle|\; U = \bigcup_{i \in I} U_i, U_i \in \mathcal{B}, \sigma_i \in \mathcal{F}(U_i), s(x) = (\sigma_i)_x \quad \forall x \in U_i \right\}$$

Then this is evidently a sheaf and the adjunction $\phi \to \phi^\sharp$ is the unique presheaf morphism such that

$$\phi^\sharp_U(\sigma)_x = \phi\left((\sigma_x)_{x \in U}\right)_x \quad \forall U \in \mathcal{B}, \sigma \in \mathcal{F}(U), x \in U$$

and the inverse adjunction $\psi^\flat \leftarrow \psi$ is the unique sheaf morphism such that

$$\psi^\flat(s)(x) = \psi_x(s(x)) \quad \forall x \in U$$

(using (4.2.9) to identify $(\mathcal{G}|_{\mathcal{B}})_x$ and $\mathcal{G}_x$). The unit of the adjunction is given by

$$(\eta_{\mathcal{F}})_U(\sigma) = (\sigma_x)_{x \in U}$$

and for a morphism of $\mathcal{B}$-sheaves $\phi : \mathcal{F} \to \mathcal{G}$ the morphism ϕ^+ defined in (4.2.14) is simply

$$\phi^+(s)(x) = \phi_x(s(x)).$$

Furthermore $\eta_{\mathcal{F}}$ is an isomorphism if and only if $\mathcal{F}$ is a $\mathcal{B}$-sheaf.

Proof. We first demonstrate this when $\mathcal{C} = \mathbf{Set}$. The sheaf conditions are easy to verify. There is an obvious unit morphism

$$\begin{aligned} \eta_{\mathcal{F}}(U) : \mathcal{F}(U) &\to \mathcal{F}^+(U) \\ \sigma &\to x \to \sigma_x \end{aligned}$$

which is a natural transformation by (4.2.7). According to (2.6.74) it is enough to ensure the following map is a bijection and natural in $\mathcal{G}$

$$\begin{aligned} \mathrm{Mor}(\mathcal{F}^+, \mathcal{G}) &\to \mathrm{Mor}(\mathcal{F}, \mathcal{G}|_{\mathcal{B}}) \\ \phi &\to \phi|_{\mathcal{B}} \circ \eta_{\mathcal{F}} \end{aligned}$$

Naturality in $\mathcal{G}$ follows formally. Suppose $\phi, \psi : \mathcal{F}^+ \to \mathcal{G}$ are morphisms such that $\phi|_{\mathcal{B}} \circ \eta_{\mathcal{F}} = \psi|_{\mathcal{B}} \circ \eta_{\mathcal{F}}$. For a given $s \in \mathcal{F}^+(U)$ we wish to show that $\phi(s) = \psi(s)$. Suppose that s is determined by sections $\sigma_i \in \mathcal{F}(U_i)$, that is to say $s|_{U_i} = \eta_{\mathcal{F}}(\sigma_i)$. Therefore by hypothesis $\phi(s)|_{U_i} = \phi\left(s|_{U_i}\right) = \psi\left(s|_{U_i}\right) = \psi(s)|_{U_i}$. By the concrete sheaf conditions (4.2.4) on $\mathcal{G}$ we conclude that $\phi(s) = \psi(s)$, and so the given map is injective.

© David Rufino CC BY-NC-ND 4.0

Conversely given $\phi : \mathcal{F} \to \mathcal{G}|_{\mathcal{B}}$, we wish to construct a morphism of sheaves $\phi^\flat : \mathcal{F}^+(U) \to \mathcal{G}(U)$ such that $\phi^{\flat}\big((\sigma_x)_{x\in U}\big) = \phi(\sigma)$ for all $\sigma \in \mathcal{F}(U)$ and $U \in \mathcal{B}$. We claim that there exists a family of maps $\big\{\phi^\flat_U : \mathcal{F}^+ \to \mathcal{G}\big\}$ such that $\phi^\flat_U(s)_x = \phi_x(s(x))$ for all $s \in \mathcal{F}^+(U)$. Such a family is unique by (4.2.8).

To show existence consider a section $s \in \mathcal{F}^+(U)$, then by definition $s(x) = \big((\sigma_i)_x\big)$ for some $\sigma_i \in \mathcal{F}(U_i)$, $U_i \in \mathcal{B}$ and all $x \in U_i$. Define $\tau_i \coloneqq \phi(\sigma_i)$ then by (4.2.7) $(\tau_i)_x = \phi_x\big((\sigma_i)_x\big) = \phi_x(s(x))$. This shows that $(\tau_i)_x = \big(\tau_j\big)_x$ for all $x \in U_i \cap U_j$ so by (4.2.8) $\tau_i|_{U_i\cap U_j} = \tau_j|_{U_i\cap U_j}$. By (4.2.4) there exists $\tau \in \mathcal{G}(U)$ such that $\tau|_{U_i} = \tau_i$. This yields a map $\phi^\flat_U$ satisfying $\phi^\flat_U(s)_x = \tau_x = (\tau_i)_x = \phi(\sigma_i)_x = \phi_x\big((\sigma_i)_x\big) = \phi_x(s(x))$ as required.

It remains to show that $\phi^\flat$ is a morphism of sheaves, that is the following diagram is commutative for all open sets $V \subset U$

$$\begin{array}{ccc}
\mathcal{F}^+(U) & \xrightarrow{\phi^\flat_U} & \mathcal{G}(U) \\
\downarrow{\scriptstyle \rho_{UV}} & & \downarrow{\scriptstyle \rho_{UV}} \\
\mathcal{F}^+(V) & \xrightarrow{\phi^\flat_V} & \mathcal{G}(V) \\
\downarrow & & \downarrow{\scriptstyle \rho_x} \\
\mathcal{F}_x & \xrightarrow{\phi_x} & \mathcal{G}_x
\end{array}$$

The large and bottom squares commute by construction. Therefore for $s \in \mathcal{F}^+(U)$ we have $\big(\phi^\flat_U(s)|_V\big)_x = \phi_x(s|_V\ (x)) = \phi^\flat_V(s|_V)_x$. As this holds for all $x \in V$ we find from (4.2.8) that $\phi^\flat_U(s)|_V = \phi^\flat_V(s|_V)$ as required.

As $(-)^\sharp$ is injective it is sufficient by (2.6.36) to show that it has a right inverse, specifically $\big(\psi^\flat\big)^\sharp = \psi$. However $\psi(\sigma)_x = \psi^\flat\big((\sigma_{x'})_{x'\in U}\big)(x) = \big(\psi^\flat\big)^\sharp(\sigma)_x$ by construction so we are done by (4.2.8).

Evidently when $\eta_{\mathcal{F}}$ is an isomorphism then $\mathcal{F}$ is a $\mathcal{B}$-sheaf. We claim that the converse holds, namely suppose $\mathcal{F}$ is a $\mathcal{B}$-sheaf, then we require to show $(\eta_{\mathcal{F}})_U$ is injective and surjective. It is injective by (4.2.8). Similarly it is surjective for suppose $s \in \mathcal{F}^+(U)$ for $U \in \mathcal{B}$ and $s(x) = (\sigma_i)_x$ for all $x \in U_i$ and $\sigma_i \in \mathcal{F}(U_i)$. Then by the same result we have $(\sigma_i)|_{U_i\cap U_j} = \big(\sigma_j\big)|_{U_i\cap U_j}$, whence by (4.2.4) these lift to a section $\sigma \in \mathcal{F}(U)$ such that $\sigma|_{U_i} = \sigma_i$, and in particular $\sigma_x = (\sigma_i)_x = s(x)$ for all $x \in U_i$. Therefore we conclude the map $(\eta_{\mathcal{F}})_U$ is surjective and therefore an isomorphism in **Set**, which means $\eta_{\mathcal{F}}$ is an isomorphism of sheaves.

We now turn to the case of a general algebraic category $\mathcal{C}$, determined by (Ω, E) (2.6.185) write $U : \mathcal{C} \to \mathbf{Set}$ for the forgetful functor.

The remainder of this proof is drafted by Claude Sonnet 5, reviewed by David Rufino.

By (2.6.191), for $U \in \mathcal{B}$ the product $\prod_{x\in U} \mathcal{F}_x$ exists in $\mathcal{C}$, with each $\omega \in \Omega$ acting coordinatewise.

$\mathcal{F}^+(U)$ is a subalgebra of $\prod_{x\in U} \mathcal{F}_x$. Given $\omega \in \Omega$ and $s^{(1)}, ..., s^{(n_\omega)} \in \mathcal{F}^+(U)$, and $x \in U$: choose for each $r = 1, ..., n_\omega$ a representative $\sigma^{(r)} \in \mathcal{F}(V_r)$, $V_r \in \mathcal{B}$, $x \in V_r$, with $s^{(r)}(y) = \big(\sigma^{(r)}\big)_y$ for all $y \in V_r$. As n_ω is finite and $\mathcal{B}$ is closed under finite intersections, $N \coloneqq \bigcap_r V_r \in \mathcal{B}$, and, by functoriality of stalks (4.2.7), $\tau \coloneqq \omega_{\mathcal{F}(N)}\big(\sigma^{(1)}|_N, ..., \sigma^{(n_\omega)}|_N\big) \in \mathcal{F}(N)$ has $\tau_y = \omega_{\mathcal{F}_y}\big(s^{(1)}(y), ..., s^{(n_\omega)}(y)\big)$ for every $y \in N$. So the germ-function $y \mapsto \omega_{\mathcal{F}_y}\big(s^{(1)}(y), ..., s^{(n_\omega)}(y)\big)$ — which is exactly ω applied coordinatewise to $s^{(1)}, ..., s^{(n_\omega)}$ in $\prod_{x\in U} \mathcal{F}_x$ — is locally represented by τ near every point, hence lies in $\mathcal{F}^+(U)$. So $\mathcal{F}^+(U)$ is closed under every $\omega \in \Omega$, and by (2.6.194) is a subalgebra, in particular an object of $\mathcal{C}$, with $\omega_{\mathcal{F}^+(U)}$ given pointwise as above.

The maps constructed above are morphisms of $\mathcal{C}$. If $h : A \to B$ is a homomorphism and $S \subseteq U(A)$, $T \subseteq U(B)$ are subalgebras with $h(S) \subseteq T$, then $h|_S : S \to T$ is a homomorphism, as ω_S, ω_T are literally restrictions of ω_A, ω_B. We apply this three times.

For the restriction maps: the coordinate projection $\prod_{x\in U} \mathcal{F}_x \to \prod_{x\in V} \mathcal{F}_x$ ($V \subseteq U$ in $\mathcal{B}$) is a homomorphism, carries $\mathcal{F}^+(U)$ into $\mathcal{F}^+(V)$ by definition of restriction of germ-functions, and restricts to ρ_{UV}.

For the unit: each stalk map $\rho_{Ux} : \mathcal{F}(U) \to \mathcal{F}_x$ is a morphism of $\mathcal{C}$ (4.2.5), so $(\rho_{Ux})_{x\in U} : \mathcal{F}(U) \to \prod_{x\in U} \mathcal{F}_x$ is a homomorphism (a map into a product is a homomorphism iff each component is); its image lies in $\mathcal{F}^+(U)$ (represented by σ itself, via the cover $\{U\}$), and it restricts to $(\eta_{\mathcal{F}})_U$.

For $\phi^\flat$: given a morphism $\phi : \mathcal{F} \to \mathcal{G}|_{\mathcal{B}}$ of $\mathcal{C}$-presheaves with $\mathcal{G}$ a $\mathcal{B}$-sheaf, each $\phi_x : \mathcal{F}_x \to \mathcal{G}_x$ is a morphism of $\mathcal{C}$ (4.2.7), so $h \coloneqq \prod_{x\in U} \phi_x : \prod_{x\in U} \mathcal{F}_x \to \prod_{x\in U} \mathcal{G}_x$ is a homomorphism. As above, $(\rho_{Ux})_{x\in U} : \mathcal{G}(U) \to \prod_{x\in U} \mathcal{G}_x$ is an injective (by (4.2.8), as $\mathcal{G}$ is a sheaf) homomorphism, so identifies $\mathcal{G}(U)$ with a subalgebra T of $\prod_{x\in U} \mathcal{G}_x$ — its image, which

© David Rufino CC BY-NC-ND 4.0

is a subalgebra since the image of a homomorphism is always closed under the operations of the target. By the construction of $\phi^\flat_U$ above, $h(\mathcal{F}^+(U)) \subseteq T$: indeed for $s \in \mathcal{F}^+(U)$ we exhibited $\tau \in \mathcal{G}(U)$ with $\tau_x = \phi_x(s(x))$ for all $x \in U$. So h restricts to a homomorphism $\mathcal{F}^+(U) \to T \cong \mathcal{G}(U)$, which is $\phi^\flat_U$.

Given this, every map constructed in the case $\mathcal{C} = \mathbf{Set}$ above is now a morphism of $\mathcal{C}$, and the remaining verifications — injectivity of $\phi \mapsto \phi|_{\mathcal{B}} \circ \eta_{\mathcal{F}}$, existence and commutativity of $\phi^\flat$, and the final isomorphism criterion — proceed entirely by manipulating elements of underlying sets via (4.2.8) and (4.2.4), both already stated for general algebraic categories, so the argument given there applies verbatim; in the final step, bijectivity of $(\eta_{\mathcal{F}})_U$ on underlying sets upgrades to an isomorphism of $\mathcal{C}$ since U reflects isomorphisms (2.6.190). ∎

Corollary 4.2.17
Let $\mathcal{C}$ be an algebraic category (2.6.185) then there is an equivalence of categories

$$\mathbf{Sh}(X;\mathcal{B}) \underset{(-)|_{\mathcal{B}}}{\overset{(-)^+}{\rightleftarrows}} \mathbf{Sh}(X)$$

Proof. By the last statement of (4.2.16), and (2.6.73) the essential image of $(-)|_{\mathcal{B}}$ is the full subcategory of $\mathcal{B}$-sheaves, and the unit and counit natural transformations induce the equivalence of categories. ∎

Lemma 4.2.18 (Sheafification Preserves Stalks)
Let $\mathcal{C}$ be algebraic (2.6.185), $\mathcal{F}$ a presheaf on X and $x \in X$. Then $\eta_{\mathcal{F},x} : \mathcal{F}_x \to (\mathcal{F}^+)_x$ (4.2.7) is an isomorphism.

Proof. Drafted by Claude Sonnet, reviewed by David Rufino.

Surjectivity. Let $\xi \in (\mathcal{F}^+)_x$; by (4.2.6), $\xi = s_x$ for some $s \in \mathcal{F}^+(U)$, $U \ni x$. By (4.2.16), $s(y) = (\sigma_i)_y$ for all $y \in U_i$, for some cover $U = \bigcup_i U_i$ and $\sigma_i \in \mathcal{F}(U_i)$; choosing i with $x \in U_i$, this says $s|_{U_i} = \eta_{U_i}(\sigma_i)$, so

$$\xi = s_x = \left(s|_{U_i}\right)_x = \left(\eta_{U_i}(\sigma_i)\right)_x = \eta_x\left((\sigma_i)_x\right)$$

so $\xi \in \mathrm{Im}(\eta_x)$.

Injectivity. Suppose $\eta_x(a) = \eta_x(b)$ for $a, b \in \mathcal{F}_x$. By (4.2.6), $a = \sigma_x$, $b = \tau_x$ for some $\sigma \in \mathcal{F}(U)$, $\tau \in \mathcal{F}(V)$, $U, V \ni x$; then $\eta_x(a) = (\eta_U(\sigma))_x$ and $\eta_x(b) = (\eta_V(\tau))_x$, so by (4.2.8).a) there is $W \subseteq U \cap V$, $x \in W$, with $\eta_U(\sigma)|_W = \eta_V(\tau)|_W$ in $\mathcal{F}^+(W)$, that is $\eta_W(\sigma|_W) = \eta_W(\tau|_W)$. As elements of $\mathcal{F}^+(W)$ are functions $W \to \coprod_{y \in W} \mathcal{F}_y$ (4.2.16), this equality holds in particular at $y = x$:

$$(\sigma|_W)_x = \eta_W(\sigma|_W)(x) = \eta_W(\tau|_W)(x) = (\tau|_W)_x$$

that is $\sigma_x = \tau_x$, so $a = b$. ∎

4.2.3 Inverse Image Functor

Definition 4.2.19 (Inverse Image Sheaf)
*Let $f : X \to Y$ be a continuous map and $\mathcal{G}$ a presheaf on Y. We define as follows the **inverse image sheaf***

$$(f^{-1}\mathcal{G})(U) := \left\{ s : U \to \coprod_{x \in U} \mathcal{G}_{f(x)} \;\middle|\; \forall x \in U, \exists V \in \mathcal{O}_{f(x)} \text{ and } \sigma \in \mathcal{G}(V) \text{ s.t. } s(y) = \sigma_y \text{ for all } y \in f^{-1}(V) \right\}$$

By definition it is a sheaf. Furthermore we see there is a canonical isomorphism

$$\begin{aligned} (f^{-1}\mathcal{G})_x &\xrightarrow{\sim} \mathcal{G}_{f(x)} \\ (U, s) &\to \quad s(x) \end{aligned}$$

4.2.4 Skyscraper Sheaf

Lemma 4.2.20
Consider a family of objects in $\{A_i\}_{i \in I}$ such that $i \in I_0 \implies A_i$ is terminal. Then there is a canonical isomorphism for which the following diagram commutes for all morphisms f

$$\begin{array}{ccc} B & \xrightarrow{f} & \prod_{i \in I} A_i \\ & \searrow^{(\pi_i \circ f)_{i \in I_0}} & \downarrow \sim \\ & & \prod_{i \in I_0} A_i \end{array}$$

© David Rufino CC BY-NC-ND 4.0

namely that given by $(\pi_i)_{i\in I_0}$.

Proof. Let $\phi = (\pi_i)_{i\in I_0}$ and let $\psi : \prod_{i\in I_0} A_i \to \prod_{i\in I} A_i$ be the unique morphism such that

$$\pi_j \circ \psi = \begin{cases} \pi_j & j \in I_0 \\ !_{\prod_{i\in I_0} A_i} & j \notin I_0 \end{cases}$$

Then $\pi_i \circ \psi \circ \phi = \pi_i \circ \phi = \pi_i$ when $i \in I_0$ and it equals $!_{\prod_{i\in I_0} A_i} \circ \phi = !_{\prod_{i\in I} A_i} = \pi_i$ when $i \notin I_0$, whence $\psi \circ \phi = 1$. The reverse is similar. ∎

Proposition 4.2.21
Let $A \in \mathcal{C}$ be an object, $\star$ a terminal object and $x \in X$. The following assignment

$$i_x(A)(U) := \begin{cases} A & x \in U \\ \star & x \notin U \end{cases}$$

determines a sheaf with restriction maps given by

$$\rho_{UV} := \begin{cases} 1_A & x \in V \\ !_A & x \in U \setminus V \\ 1_\star & x \notin U \end{cases}$$

This is known as the ***skyscraper sheaf*** *at x.*

Proof. Suppose that $U = \bigcup_{i\in I} U_i$ and let $I_0 = \{i \in I \mid x \in U_i\}$ then we may show there is a commutative diagram

$$\begin{array}{ccccc} \mathcal{F}(U) & \xrightarrow{e} & \prod_{i\in I} \mathcal{F}(U_i) & \underset{d_1}{\overset{d_0}{\rightrightarrows}} & \prod_{(i,j)\in I\times I} \mathcal{F}(U_i \cap U_j) \\ & \searrow^{e} & \downarrow \sim & & \downarrow \sim \\ & & \prod_{i\in I_0} \mathcal{F}(U_i) & \underset{d_1}{\overset{d_0}{\rightrightarrows}} & \prod_{(i,j)\in I_0\times I_0} \mathcal{F}(U_i \cap U_j) \end{array}$$

where $e = \left(\rho_{UU_i}\right)$, $d_0 = \left(\rho_{U_i(U_i\cap U_j)} \circ \pi_i\right)$ and $d_1 = \left(\rho_{U_j(U_i\cap U_j)} \circ \pi_j\right)$. Then it's enough to show that the bottom diagram is an equaliser.

For $i, j \in I_0$ we have $\mathcal{F}(U) = \mathcal{F}(U_i) = \mathcal{F}(U_i \cap U_j) = A$.

Suppose $f : B \to \prod_{i\in I_0} \mathcal{F}(U_i)$ such that $d_0 \circ f = d_1 \circ f$. Then

$$\begin{aligned} d_0 \circ f = d_1 \circ f &\implies \pi_{(i,j)} \circ d_0 \circ f = \pi_{(i,j)} \circ d_1 \circ f \quad \forall i, j \in I_0 \\ &\implies 1_A \circ \pi_i \circ f = 1_A \circ \pi_j \circ f \quad \forall i, j \in I_0 \\ &\implies \pi_i \circ f = \pi_j \circ f \quad \forall i, j \in I_0 \end{aligned}$$

Therefore define the morphism $f' : B \to \mathcal{F}(U)$ to be $\pi_j \circ f$ for some $j \in I_0$. Observing $e = (1_A)_{i\in I_0}$ we have $e \circ f' = (f')_{i\in I_0} = \left(\pi_j \circ f\right)_{i\in I_0} = (\pi_i \circ f)_{i\in I_0} = f$. ∎

Proposition 4.2.22 (Stalk functor is left-adjoint to Sky-Scraper Sheaf)
Let X be a topological space, $\mathcal{C}$ a category with a terminal object $\star$ and directed colimits, $A \in \mathcal{C}$ and $\mathcal{F}$ a sheaf on X. Then for all $x \in X$ there is a bijection

$$\begin{aligned} \mathrm{Mor}(\mathcal{F}_x, A) &\xrightarrow{\sim} \mathrm{Mor}(\mathcal{F}, i_x(A)) \\ \phi \to \quad \phi^\sharp_U &:= \begin{cases} \phi \circ \rho_{Ux} & x \in U \\ !_{\mathcal{F}(U)} & x \notin U \end{cases} \end{aligned}$$

which is natural in both $\mathcal{F}$ and A.

Proof. Let $\mathcal{N}_x$ be the category of open subsets U of X containing x ordered by reverse inclusion. Then by definition of the stalk (4.2.7) there is a bijection

$$\begin{aligned} \mathrm{Mor}(\mathcal{F}_x, A) &\xrightarrow{\sim} \left(\mathcal{F}|_{\mathcal{N}_x} \downarrow A\right) \\ \phi &\to (\phi \circ \rho_{Ux})_{U\in\mathcal{N}_x} \end{aligned}$$

© David Rufino CC BY-NC-ND 4.0

which is natural in $\mathcal{F}$ and A. It then remains to verify that there is a natural bijection

$$\left(\mathcal{F}|_{\mathcal{N}_x} \downarrow A\right) \xrightarrow{\sim} \mathrm{Mor}(\mathcal{F}, i_x(A))$$
$$\left(\Phi_U\right)_{U \ni x} \to \begin{cases} \Phi_U & x \in U \\ !_{\mathcal{F}(U)} & x \notin U \end{cases}$$

■

4.2.5 Limit Sheaf

Proposition 4.2.23 (Limits of Presheaves and Sheaves may be constructed pointwise) *Let K be a small category and $F : K \to \mathbf{PSh}(X;\mathcal{C})$ (resp. $\mathbf{Sh}(X;\mathcal{C})$) a diagram, that is a presheaf (resp. sheaf) $F(k)$ for each $k \in K$ together with a morphism $F(\kappa) : F(k) \to F(k')$ for each arrow $\kappa : k \to k'$, functorially. Write $F(-)(U) : K \to \mathcal{C}$ for the diagram $k \mapsto F(k)(U)$, and suppose $\lim_K F(-)(U)$ exists in $\mathcal{C}$ for every $U \subset X$.*

Then the assignment

$$\left(\lim_K F\right)(U) \coloneqq \lim_K F(-)(U)$$

can be made into a presheaf (resp. sheaf) uniquely such that the projections $\lambda_k : \lim_K F \to F(k)$ are morphisms of presheaves (resp. sheaves) for every $k \in K$. Furthermore $\left(\lim_K F, \{\lambda_k\}_{k\in K}\right)$ is a limit for F in the category of presheaves (resp. sheaves).

When $\mathcal{C}$ is an algebraic category (2.6.185) this may be constructed in the obvious way

$$\left(\lim_K F\right)(U) \coloneqq \left\{ (\sigma_k)_{k\in K} \in \prod_{k\in K} F(k)(U) \;\middle|\; F(\kappa)_U(\sigma_k) = \sigma_{k'} \text{ for every arrow } \kappa : k \to k' \text{ in } K \right\}$$

Proof. Drafted by Claude Sonnet, reviewed by David Rufino.

Restriction maps. Fix $V \subseteq U$, and write $\{\lambda_{k,U}\}_{k\in K}$, $\{\lambda_{k,V}\}_{k\in K}$ for the given limit cones for $F(-)(U)$, $F(-)(V)$. For every arrow $\kappa : k \to k'$ of K, using that $F(\kappa) : F(k) \to F(k')$ is a morphism of presheaves and the cone condition $F(\kappa)_U \circ \lambda_{k,U} = \lambda_{k',U}$,

$$F(\kappa)_V \circ \left(\rho_{UV}^{F(k)} \circ \lambda_{k,U}\right) = \rho_{UV}^{F(k')} \circ F(\kappa)_U \circ \lambda_{k,U} = \rho_{UV}^{F(k')} \circ \lambda_{k',U}$$

so $\left\{\rho_{UV}^{F(k)} \circ \lambda_{k,U}\right\}_{k\in K}$ is a cone over $F(-)(V)$ with apex $\lim_K F(-)(U)$. By the universal property of $\lim_K F(-)(V)$, there is a unique morphism $\rho_{UV} : \lim_K F(-)(U) \to \lim_K F(-)(V)$ such that $\lambda_{k,V} \circ \rho_{UV} = \rho_{UV}^{F(k)} \circ \lambda_{k,U}$ for every $k \in K$; this is precisely the condition for each λ_k to be a morphism of presheaves.

Taking $U = V$, both ρ_{UU} and $1_{\lim_K F(-)(U)}$ satisfy $\lambda_{k,U} \circ (-) = \lambda_{k,U}$ for every k, so $\rho_{UU} = 1_{\lim_K F(-)(U)}$ by uniqueness. For $W \subseteq V \subseteq U$, both $\rho_{VW} \circ \rho_{UV}$ and ρ_{UW} satisfy, for every k, $\lambda_{k,W} \circ (-) = \rho_{VW}^{F(k)} \circ \rho_{UV}^{F(k)} \circ \lambda_{k,U} = \rho_{UW}^{F(k)} \circ \lambda_{k,U}$, so $\rho_{VW} \circ \rho_{UV} = \rho_{UW}$ by uniqueness. So $\lim_K F$, with these restriction maps, is a presheaf, and each λ_k is a morphism of presheaves; uniqueness of restriction maps making every λ_k a morphism of presheaves is immediate from the uniqueness clause of the universal property of $\lim_K F(-)(V)$.

Limit in the category of presheaves. Let $\mathcal{K}$ be a presheaf and $\{\mu_k : \mathcal{K} \to F(k)\}_{k\in K}$ a cone over F in $\mathbf{PSh}(X;\mathcal{C})$, that is $F(\kappa) \circ \mu_k = \mu_{k'}$ for every arrow $\kappa : k \to k'$. For each U, $\left\{(\mu_k)_U\right\}_{k\in K}$ is a cone over $F(-)(U)$ with apex $\mathcal{K}(U)$, so the universal property of $\lim_K F(-)(U)$ gives a unique $\mu_U : \mathcal{K}(U) \to \lim_K F(-)(U)$ with $\lambda_{k,U} \circ \mu_U = (\mu_k)_U$ for every k. For $V \subseteq U$, both $\rho_{UV} \circ \mu_U$ and $\mu_V \circ \rho_{UV}^{\mathcal{K}}$ satisfy, for every k, $\lambda_{k,V} \circ (-) = \rho_{UV}^{F(k)} \circ (\mu_k)_U = (\mu_k)_V \circ \rho_{UV}^{\mathcal{K}}$ (naturality of μ_k, and the defining properties of μ_U, μ_V), so $\rho_{UV} \circ \mu_U = \mu_V \circ \rho_{UV}^{\mathcal{K}}$ by uniqueness in the universal property of $\lim_K F(-)(V)$: that is, $\mu \coloneqq \{\mu_U\}$ is a morphism of presheaves, with $\lambda_k \circ \mu = \mu_k$ for every k by construction. If μ' is another morphism of presheaves with $\lambda_k \circ \mu' = \mu_k$ for every k, then μ'_U satisfies $\lambda_{k,U} \circ \mu'_U = (\mu_k)_U$ for every k, so $\mu'_U = \mu_U$ by the pointwise uniqueness above, that is $\mu' = \mu$.

Sheaf case. Suppose $F(k)$ is a sheaf for every $k \in K$; we show $\lim_K F$ is also a sheaf. Fix $U = \bigcup_{i\in I} U_i$, and let $I^{(1)}$, D_- be as in (4.2.3).

Define a bifunctor $H : K \times I^{(1)} \to \mathcal{C}$ by $H(k,-) \coloneqq D_{F(k)}$ for $k \in K$, and $H(\kappa, 1_l) \coloneqq F(\kappa)_{U_l}$ for every arrow $\kappa : k \to k'$ of K and $l \in I^{(1)}$ (writing U_l for U_i, resp. $U_i \cap U_j$, when $l = i$, resp. $l = (i,j)$); this is a bifunctor precisely because each $F(\kappa)$ is a morphism of presheaves, so commutes with the maps $D_{F(k)}(u_{ij}), D_{F(k)}(v_{ij})$ (and likewise for $F(k')$).

© David Rufino CC BY-NC-ND 4.0

Since each $F(k)$ is a sheaf, (4.2.3) gives $\lim_{I^{(1)}} H(k,-) = F(k)(U)$ for every $k \in K$; applying (2.6.99) (with the roles of I, J there taken to be our $K, I^{(1)}$), this determines a functor $L : K \to \mathcal{C}$ (2.6.96) with $L(k) = F(k)(U)$ for every k; as $\lim_{k\in K} L(k) = \lim_K F(-)(U)$ exists by hypothesis, (2.6.99) gives that $\lim_{K\times I^{(1)}} H$ exists, with limit cone $\left(\lim_K F(-)(U), \left\{\eta_{k,l} \circ \lambda_{k,U}\right\}_{(k,l)}\right)$, where $\left(\eta_{k,l}\right)_{l\in I^{(1)}}$ is the limit cone for $F(k)(U)$ over $I^{(1)}$ furnished by (4.2.3) in particular the component at (k,l), for $l \in I^{(1)}$, is $\rho^{F(k)}_{UU_l} \circ \lambda_{k,U}$.

Applying (2.6.99) directly to H (whose symmetric hypothesis, that $\lim_{k\in K} H(k,l)$ exists for every $l \in I^{(1)}$, is exactly $\lim_K F(-)(U_l) = D_{\lim_K F}(l)$ existing by hypothesis) to the cone on $\lim_K F(-)(U)$ constructed above: writing $R(l) \coloneqq \lim_{k\in K} H(k,l) = D_{\lim_K F}(l)$, $\lim_{I^{(1)}} R$ exists, with limit cone component at $l \in I^{(1)}$ the unique morphism $\sigma_l : \lim_K F(-)(U) \to \lim_K F(-)(U_l)$ satisfying $\lambda_{k,U_l} \circ \sigma_l = \rho^{F(k)}_{UU_l} \circ \lambda_{k,U}$ for every $k \in K$. This is exactly the defining property of the restriction map ρ_{UU_l} of $\lim_K F$ established above, so $\sigma_l = \rho_{UU_l}$ by uniqueness.

So $D_{\lim_K F}$ has $\left(\lim_K F(-)(U), \left\{\rho_{UU_l}\right\}_{l\in I^{(1)}}\right)$ as a limit over $I^{(1)}$, and by (4.2.3) this says exactly that $\lim_K F$ satisfies the sheaf condition for this cover, as required.

Alternatively, when $\mathcal{C}$ is algebraic (2.6.185), this can be shown more explicitly, using the concrete construction noted above. By (2.6.85) and (2.6.190) (the forgetful functor $U : \mathcal{C} \to \mathbf{Set}$ creates limits), $\lim_K F(-)(U)$ is exactly $\left\{(\sigma_k)_{k\in K} \in \prod_{k\in K} F(k)(U) \;\middle|\; F(\kappa)_U(\sigma_k) = \sigma_{k'}\ \forall \kappa : k \to k'\right\}$ for every U, with $\lambda_{k,U}$ the coordinate projections, and ρ_{UV} acting coordinatewise (restricting each σ_k).

By (4.2.4) it suffices to verify the two concrete sheaf conditions for $\lim_K F$ directly, using that each $F(k)$ satisfies them.

Separation. Suppose $(\sigma_k)_k, (\sigma'_k)_k \in \left(\lim_K F\right)(U)$ satisfy $\rho_{UU_i}\left((\sigma_k)_k\right) = \rho_{UU_i}\left((\sigma'_k)_k\right)$ for all $i \in I$. Since restriction acts coordinatewise, $\rho^{F(k)}_{UU_i}(\sigma_k) = \rho^{F(k)}_{UU_i}(\sigma'_k)$ for every $k \in K$ and $i \in I$, so $\sigma_k = \sigma'_k$ by separation for $F(k)$; as this holds for every k, $(\sigma_k)_k = (\sigma'_k)_k$.

Gluing. Suppose $\left(\sigma_{k,i}\right)_k \in \left(\lim_K F\right)(U_i)$ for $i \in I$ satisfy $\rho_{U_i(U_i\cap U_j)}\left(\left(\sigma_{k,i}\right)_k\right) = \rho_{U_j(U_i\cap U_j)}\left(\left(\sigma_{k,j}\right)_k\right)$ for all $(i,j) \in I \times I$. Fix $k \in K$; since restriction acts coordinatewise, $\rho^{F(k)}_{U_i(U_i\cap U_j)}(\sigma_{k,i}) = \rho^{F(k)}_{U_j(U_i\cap U_j)}(\sigma_{k,j})$ for all (i,j), so gluing for $F(k)$ gives $\sigma_k \in F(k)(U)$ with $\rho^{F(k)}_{UU_i}(\sigma_k) = \sigma_{k,i}$ for all i. For an arrow $\kappa : k \to k'$ of K, and every $i \in I$,

$$\rho^{F(k')}_{UU_i}(F(\kappa)_U(\sigma_k)) = F(\kappa)_{U_i}\left(\rho^{F(k)}_{UU_i}(\sigma_k)\right) = F(\kappa)_{U_i}(\sigma_{k,i}) = \sigma_{k',i} = \rho^{F(k')}_{UU_i}(\sigma_{k'})$$

using naturality of $F(\kappa)$, the defining property of σ_k, that $\left(\sigma_{k,i}\right)_k \in \left(\lim_K F\right)(U_i)$, and the defining property of $\sigma_{k'}$. As $\{U_i\}$ covers U, separation for $F(k')$ gives $F(\kappa)_U(\sigma_k) = \sigma_{k'}$. As κ was arbitrary, $(\sigma_k)_k \in \left(\lim_K F\right)(U)$, with $\rho_{UU_i}\left((\sigma_k)_k\right) = \left(\sigma_{k,i}\right)_k$ for all i, as required. ∎

4.2.6 Equaliser Sheaf

Proposition 4.2.24 (Equaliser Sheaf may be constructed pointwise)
Let $\alpha, \beta : \mathcal{F} \rightrightarrows \mathcal{G}$ be morphisms of presheaves (resp. sheaves) with a family of pointwise equalisers $\{e_U : \mathrm{eq}(\alpha_U, \beta_U) \to \mathcal{F}(U)\}_{U\subset X}$. Then the assignment

$$\mathrm{eq}(\alpha,\beta)(U) \coloneqq \mathrm{eq}(\alpha_U, \beta_U)$$

can be made into a presheaf (resp. sheaf) uniquely such that

$$e : \mathrm{eq}(\alpha,\beta) \to \mathcal{F}$$

is a morphism of presheaves (resp. sheaves). Further e is the equaliser of (α,β) in the category of presheaves (resp. sheaves).

When $\mathcal{C}$ is an algebraic category this may be constructed in the obvious way

$$\mathrm{eq}(\alpha,\beta)(U) \coloneqq \{\sigma \in \mathcal{F}(U) \mid \alpha_U(\sigma) = \beta_U(\sigma)\}$$

Proof. Drafted by Claude Sonnet, reviewed by David Rufino.

Let $\mathbf{2} = \{\emptyset \rightrightarrows 1\}$ and let $F : \mathbf{2} \to \mathbf{PSh}(X;\mathcal{C})$ (resp. $\mathbf{Sh}(X;\mathcal{C})$) be the diagram $F(\emptyset) \coloneqq \mathcal{F}$, $F(1) \coloneqq \mathcal{G}$, sending the two morphisms $\emptyset \rightrightarrows 1$ to α, β respectively. Then $F(-)(U) : \mathbf{2} \to \mathcal{C}$ is exactly the pair $\alpha_U, \beta_U : \mathcal{F}(U) \rightrightarrows \mathcal{G}(U)$, whose limit is $\mathrm{eq}(\alpha_U, \beta_U)$ (2.6.135), which exists by hypothesis for every U. By (4.2.23), the assignment $U \mapsto \mathrm{eq}(\alpha_U, \beta_U)$ can therefore be made into a presheaf (resp. sheaf) uniquely such that the projection e to $\mathcal{F}$ is a morphism of presheaves

© David Rufino CC BY-NC-ND 4.0

(resp. sheaves), and $\mathrm{eq}(\alpha,\beta) \coloneqq \lim_{\mathbf{2}} F$, with this projection e, is a limit for F in the category of presheaves (resp. sheaves), that is the equaliser of (α,β) there (2.6.135), as required.

Alternatively, when $\mathcal{C}$ is algebraic (2.6.185), this follows directly from the algebraic case of (4.2.23): writing $(\sigma_\emptyset,\sigma_1) \in \mathcal{F}(U)\times\mathcal{G}(U)$ for the coordinates indexed by $\mathbf{2} = \{\emptyset \rightrightarrows 1\}$, the compatibility condition $\alpha_U(\sigma_\emptyset) = \sigma_1 = \beta_U(\sigma_\emptyset)$ pins σ_1 down in terms of $\sigma_\emptyset$, so under the projection $(\sigma_\emptyset,\sigma_1)\mapsto\sigma_\emptyset$ this recovers exactly $\mathrm{eq}(\alpha,\beta)(U) = \{\sigma\in\mathcal{F}(U) \mid \alpha_U(\sigma) = \beta_U(\sigma)\}$ noted above. ∎

Corollary 4.2.25 (Kernel may be constructed pointwise)
Let $\mathcal{C}$ be a pointed category and $\alpha:\mathcal{F}\to\mathcal{G}$ a morphism of presheaves (resp. sheaves). Suppose for all $U\subset X$ we have a kernel $(i_U,\ker(\alpha_U))$ of α_U.

Then we may define $\ker(\alpha)$ as follows

$$\ker(\alpha)(U) \coloneqq \ker(\alpha_U)$$

and this may be made into a presheaf (resp. sheaf) uniquely such that i is a morphism of presheaves (resp. sheaves). Furthermore $(i,\ker(\alpha))$ is a kernel in the category of presheaves (resp. sheaves).

When $\mathcal{C}$ is an algebraic category (2.6.185) this may be constructed in the obvious way namely

$$\ker(\alpha)(U) \coloneqq \{\sigma\in\mathcal{F}(U) \mid \alpha_U(\sigma) = 0\}$$

Proof. Drafted by Claude Sonnet, reviewed by David Rufino.

By (2.6.151), $(i_U,\ker(\alpha_U))$ is a kernel for α_U if and only if i_U is an equaliser for $(\alpha_U,0_U)$, where $0:\mathcal{F}\to\mathcal{G}$ is the zero morphism of presheaves (computed pointwise, since $\mathbf{PSh}(X;\mathcal{C})$, resp. $\mathbf{Sh}(X;\mathcal{C})$, is pointed with zero morphisms computed pointwise). So the hypothesis gives a family of pointwise equalisers $\{i_U:\mathrm{eq}(\alpha_U,0_U)\to\mathcal{F}(U)\}_{U\subset X}$ for $(\alpha,0)$, and applying (4.2.24) to this pair, $\mathrm{eq}(\alpha,0)(U) \coloneqq \mathrm{eq}(\alpha_U,0_U) = \ker(\alpha_U) = \ker(\alpha)(U)$ can be made into a presheaf (resp. sheaf) uniquely such that i is a morphism of presheaves (resp. sheaves), and i is the equaliser of $(\alpha,0)$ in the category of presheaves (resp. sheaves); by (2.6.151) again, this is exactly $(i,\ker(\alpha))$ being a kernel for α.

When $\mathcal{C}$ is algebraic, the algebraic case of (4.2.24), applied to $(\alpha,0)$, gives $\mathrm{eq}(\alpha,0)(U) = \{\sigma\in\mathcal{F}(U) \mid \alpha_U(\sigma) = 0_U(\sigma)\} = \{\sigma\in\mathcal{F}(U) \mid \alpha_U(\sigma) = 0\}$, that is $\ker(\alpha)(U) = \{\sigma\in\mathcal{F}(U) \mid \alpha_U(\sigma) = 0\}$, as stated. ∎

4.2.7 Product Presheaf

Proposition 4.2.26 (Product Presheaf may be constructed pointwise)
Let $\{\mathcal{F}_i\}_{i\in I}$ be a family of presheaves (resp. sheaves) with a family of pointwise products $\left\{\left(\prod_{i\in I}\mathcal{F}_i(U),\{\pi_{i,U}\}_{i\in I}\right)\right\}_{U\subset X}$. Then the assignment

$$\left(\prod_{i\in I}\mathcal{F}_i\right)(U) \coloneqq \prod_{i\in I}\mathcal{F}_i(U)$$

can be made into a presheaf (resp. sheaf) uniquely such that $\pi_i:\prod_{i\in I}\mathcal{F}_i\to\mathcal{F}_i$ is a morphism of presheaves (resp. sheaves) for every $i\in I$. Furthermore $\left(\prod_{i\in I}\mathcal{F}_i,\{\pi_i\}_{i\in I}\right)$ is a product of $\{\mathcal{F}_i\}_{i\in I}$ in the category of presheaves (resp. sheaves).

When $\mathcal{C}$ is algebraic (2.6.185) this may be constructed in the obvious way

$$\left(\prod_{i\in I}\mathcal{F}_i\right)(U) \coloneqq \left\{(\sigma_i)_{i\in I} \;\middle|\; \sigma_i\in\mathcal{F}_i(U)\right\}$$

Proof. Drafted by Claude Sonnet, reviewed by David Rufino.

Regard I as a discrete category and let $F:I\to\mathbf{PSh}(X;\mathcal{C})$ (resp. $\mathbf{Sh}(X;\mathcal{C})$) be the diagram $F(i) \coloneqq \mathcal{F}_i$ (a diagram since I is discrete, so there is nothing further to specify). Then $F(-)(U):I\to\mathcal{C}$ is exactly the discrete family $\{\mathcal{F}_i(U)\}_{i\in I}$, whose limit is $\prod_{i\in I}\mathcal{F}_i(U)$ (2.6.81), which exists by hypothesis for every U. By (4.2.23), the assignment $U\mapsto\prod_{i\in I}\mathcal{F}_i(U)$ can therefore be made into a presheaf (resp. sheaf) uniquely such that each projection π_i is a morphism of presheaves (resp. sheaves), and $\left(\prod_{i\in I}\mathcal{F}_i,\{\pi_i\}_{i\in I}\right)$ is a limit for F in the category of presheaves (resp. sheaves), that is a product of $\{\mathcal{F}_i\}_{i\in I}$ there (2.6.81), as required.

When $\mathcal{C}$ is algebraic, this is exactly the algebraic case of (4.2.23) specialised to the discrete category I, where (there being no arrows) the compatibility condition is vacuous, leaving the unconstrained tuples $(\sigma_i)_{i\in I}\in\prod_{i\in I}\mathcal{F}_i(U)$ noted above. ∎

© David Rufino CC BY-NC-ND 4.0

4.2.8 Kernel Pair Sheaf

Corollary 4.2.27 (Kernel Pair Presheaf may be constructed pointwise)
Let $\mu : \mathcal{H} \to \mathcal{G}$ be a morphism of presheaves (resp. sheaves), and suppose for every $U \subset X$ there is a kernel pair $(Z_U, p_{1,U}, p_{2,U})$ of μ_U.

Then we may define $(\mathcal{H} \times_{\mathcal{G}} \mathcal{H})(U) \coloneqq Z_U$ and this may be made into a presheaf (resp. sheaf) uniquely such that p_1, p_2 are morphisms of presheaves (resp. sheaves). Furthermore $(\mathcal{H} \times_{\mathcal{G}} \mathcal{H}, p_1, p_2)$ is a kernel pair of μ in the category of presheaves (resp. sheaves).

When $\mathcal{C}$ is algebraic (2.6.185) this may be constructed in the obvious way

$$(\mathcal{H} \times_{\mathcal{G}} \mathcal{H})(U) \coloneqq \{(\sigma_1, \sigma_2) \in \mathcal{H}(U) \times \mathcal{H}(U) \mid \mu_U(\sigma_1) = \mu_U(\sigma_2)\}$$

Proof. Drafted by Claude Sonnet, reviewed by David Rufino.

Let $F : \mathbf{\Lambda} \to \mathbf{PSh}(X; \mathcal{C})$ (resp. $\mathbf{Sh}(X; \mathcal{C})$) be the diagram $F(a) \coloneqq \mathcal{H}$, $F(b) \coloneqq \mathcal{H}$, $F(c) \coloneqq \mathcal{G}$, sending both morphisms $a \to c$, $b \to c$ to μ (2.6.145). Then $F(-)(U) : \mathbf{\Lambda} \to \mathcal{C}$ is exactly the cospan $\mu_U : \mathcal{H}(U) \to \mathcal{G}(U) \leftarrow \mathcal{H}(U) : \mu_U$, whose limit is the pullback $(Z_U, p_{1,U}, p_{2,U})$ of (μ_U, μ_U), that is a kernel pair of μ_U, which exists by hypothesis for every U. By (4.2.23), the assignment $U \mapsto Z_U$ can therefore be made into a presheaf (resp. sheaf) uniquely such that the projections p_1, p_2 are morphisms of presheaves (resp. sheaves), and $(\mathcal{H} \times_{\mathcal{G}} \mathcal{H}, p_1, p_2) \coloneqq \left(\lim_{\mathbf{\Lambda}} F, p_1, p_2\right)$ is a limit for F in the category of presheaves (resp. sheaves), that is a pullback of (μ, μ), i.e. a kernel pair of μ, there, as required.

When $\mathcal{C}$ is algebraic, this is the algebraic case of (4.2.23) specialised to $\mathbf{\Lambda}$: writing $(\sigma_a, \sigma_b, \sigma_c)$ for the coordinates, the compatibility condition $\mu_U(\sigma_a) = \sigma_c = \mu_U(\sigma_b)$ pins σ_c down in terms of σ_a, σ_b, so under the projection $(\sigma_a, \sigma_b, \sigma_c) \mapsto (\sigma_a, \sigma_b)$ this recovers exactly $\{(\sigma_1, \sigma_2) \in \mathcal{H}(U) \times \mathcal{H}(U) \mid \mu_U(\sigma_1) = \mu_U(\sigma_2)\}$ noted above. ∎

Proposition 4.2.28 (Monomorphisms of Presheaves and Sheaves)
Let $\phi : \mathcal{F} \to \mathcal{G}$ be a morphism of presheaves (resp. sheaves). Then the following are equivalent

a) ϕ is a monomorphism
b) ϕ_U is monic in $\mathcal{C}$ for all $U \subset X$

If $\mathcal{C}$ is algebraic (2.6.185), then a), b) are further equivalent to

c) ϕ_U is injective for all $U \subset X$

If moreover $\mathcal{F}, \mathcal{G}$ are sheaves, a)–c) are equivalent to

d) ϕ_x is injective for all $x \in X$

Proof. Drafted by Claude Sonnet, reviewed by David Rufino.

b) $\implies$ a): Suppose $\alpha, \beta : \mathcal{H} \to \mathcal{F}$ satisfy $\phi\alpha = \phi\beta$. Composition of morphisms of presheaves is computed pointwise, so $\phi_U \alpha_U = \phi_U \beta_U$ for every U; as ϕ_U is monic, $\alpha_U = \beta_U$ for every U, that is $\alpha = \beta$.

a) $\implies$ b): Since $\mathcal{C}$ has products and equalisers, by (2.6.146) ϕ_U has a kernel pair for every U, so by (4.2.27) these assemble into a kernel pair $(\mathcal{F} \times_{\mathcal{G}} \mathcal{F}, p_1, p_2)$ of ϕ in the category of presheaves (resp. sheaves), with U-components $\left(\mathcal{F}(U) \times_{\mathcal{G}(U)} \mathcal{F}(U), p_{1,U}, p_{2,U}\right)$ a kernel pair of ϕ_U. As ϕ is monic, (2.6.148) gives $p_1 = p_2$; since morphisms of presheaves are equal iff equal pointwise, $p_{1,U} = p_{2,U}$ for every U, so by (2.6.148) again (applied in $\mathcal{C}$), ϕ_U is monic for every U.

b) $\iff$ c) (when $\mathcal{C}$ is algebraic): immediate from (2.6.195) (monic iff injective), applied at each U.

c) $\implies$ d): (4.2.10).

d) $\implies$ c): (4.2.10). ∎

Corollary 4.2.29 (Monomorphism of Sheaves is an Isomorphism iff Surjective on Stalks)
Let $\mathcal{C}$ be algebraic (2.6.185) and $m : \mathcal{F} \to \mathcal{G}$ a monomorphism of sheaves. Then m is an isomorphism if and only if m_x is surjective for every $x \in X$.

Proof. Drafted by Claude Sonnet, reviewed by David Rufino.

By (4.2.28), m_U is monic, hence injective (2.6.195) (as $\mathcal{C}$ is algebraic), for every $U \subset X$, so by (4.2.10), m_x is injective for every $x \in X$.

As $\mathcal{C}$ is algebraic, U reflects isomorphisms (2.6.190), so an injective morphism of $\mathcal{C}$ is an isomorphism if and only if it is surjective; applying this to m_x (already known injective) for each x, m_x is an isomorphism for every $x \in X$ if and only if m_x is surjective for every $x \in X$. The result now follows from (4.2.11) (a) $\iff$ d)). ∎

© David Rufino CC BY-NC-ND 4.0

Definition 4.2.30 (Subpresheaf, Subsheaf)
*Let $\mathcal{C}$ be algebraic (2.6.185) and $\mathcal{F}$ a presheaf on X. A **subpresheaf** of $\mathcal{F}$ is an assignment, to every open $U \subset X$, of a subalgebra $\mathcal{G}(U) \subseteq U(\mathcal{F}(U))$ (2.6.193), such that $\rho^{\mathcal{F}}_{UV}(\mathcal{G}(U)) \subseteq \mathcal{G}(V)$ for every $V \subseteq U$.*

By (2.6.194), each $\mathcal{G}(U)$ carries a unique $\mathcal{C}$-algebra structure making the inclusion $\iota_U : \mathcal{G}(U) \hookrightarrow \mathcal{F}(U)$ a morphism of $\mathcal{C}$; since $\rho^{\mathcal{F}}_{UV}$ is a homomorphism carrying $\mathcal{G}(U)$ into $\mathcal{G}(V)$, it restricts to a morphism $\rho^{\mathcal{G}}_{UV} : \mathcal{G}(U) \to \mathcal{G}(V)$ of $\mathcal{C}$, making $\mathcal{G}$ into a presheaf and $\iota := \{\iota_U\} : \mathcal{G} \to \mathcal{F}$ into a morphism of presheaves.

*We say $\mathcal{G}$ is a **subsheaf** of $\mathcal{F}$ if moreover $\mathcal{G}$, with this presheaf structure, is a sheaf.*

Remark 4.2.31
Each ι_U is injective, hence monic (2.6.195), so ι itself is monic (4.2.28). Conversely, if $\iota : \mathcal{G} \to \mathcal{F}$ is a monomorphism of presheaves (resp. sheaves), then each ι_U is injective (4.2.28) and (2.6.195), and its image $U(\iota_U)(U(\mathcal{G}(U))) \subseteq U(\mathcal{F}(U))$ is a subalgebra (being the image of a homomorphism), so ι identifies $\mathcal{G}$ with a subpresheaf (resp. subsheaf) of $\mathcal{F}$.

Proposition 4.2.32 (Sheafification of a Subpresheaf)
Let $\mathcal{C}$ be algebraic (2.6.185) and $\iota : \mathcal{F} \to \mathcal{G}$ a monomorphism of presheaves with $\mathcal{G}$ a sheaf. Define

$$\mathcal{F}^+(U) := \left\{ \tau \in U(\mathcal{G}(U)) \;\middle|\; U = \bigcup_{i \in I} U_i \text{ for some open cover, with } \tau|_{U_i} \in U\big(\iota_{U_i}\big)(U(\mathcal{F}(U_i))) \text{ for every } i \right\}$$

Then $\mathcal{F}^+$ is a subsheaf of $\mathcal{G}$, and the corestriction $\iota' : \mathcal{F} \to \mathcal{F}^+$ of ι is a sheafification of $\mathcal{F}$ (4.2.13) (justifying, by (2.6.64), the use of the notation $\mathcal{F}^+$ for it, as usual).

Proof. Drafted by Claude Sonnet, reviewed by David Rufino.

$\mathcal{F}^+$ is a subsheaf. For $\omega \in \Omega$ and $\tau_1, ..., \tau_{n_\omega} \in \mathcal{F}^+(U)$, choose covers $U = \bigcup_i U_i^{(r)}$ witnessing $\tau_r \in \mathcal{F}^+(U)$; the common refinement $\left\{U_{i_1}^{(1)} \cap \cdots \cap U_{i_{n_\omega}}^{(n_\omega)}\right\}$ is again a cover of U, and on each piece W of it, $\tau_r|_W = \iota_W(\sigma_r)$ for some $\sigma_r \in \mathcal{F}(W)$ (restricting $\tau_r|_{U_{i_r}^{(r)}} = \iota_{U_{i_r}^{(r)}}(\sigma_r)$ further, as ι is a morphism of presheaves), so

$$\omega_{\mathcal{G}(U)}\big(\tau_1, ..., \tau_{n_\omega}\big)|_W = \omega_{\mathcal{G}(W)}(\tau_1|_W, ...) = \omega_{\mathcal{G}(W)}(\iota_W(\sigma_1), ...) = \iota_W\big(\omega_{\mathcal{F}(W)}(\sigma_1, ...)\big) \in \iota_W(\mathcal{F}(W))$$

As this holds for every W in the cover, $\omega_{\mathcal{G}(U)}\big(\tau_1, ..., \tau_{n_\omega}\big) \in \mathcal{F}^+(U)$, so $\mathcal{F}^+(U)$ is a subalgebra of $\mathcal{G}(U)$ (2.6.193). Restricting a witnessing cover for $\tau \in \mathcal{F}^+(U)$ to $V \subseteq U$ shows $\rho^{\mathcal{G}}_{UV}(\mathcal{F}^+(U)) \subseteq \mathcal{F}^+(V)$, so $\mathcal{F}^+$ is a subpresheaf of $\mathcal{G}$ (4.2.30).

Separation for $\mathcal{F}^+$ is immediate, since $\mathcal{F}^+(U) \subseteq U(\mathcal{G}(U))$ and $\mathcal{G}$ is separated. For gluing, let $\tau_i \in \mathcal{F}^+(U_i)$ for a cover $U = \bigcup_i U_i$, agreeing on overlaps; as $\mathcal{G}$ is a sheaf there is $\tau \in \mathcal{G}(U)$ with $\tau|_{U_i} = \tau_i$. Each τ_i has a witnessing cover $U_i = \bigcup_k U_{i,k}$ with $\tau_i|_{U_{i,k}} \in \iota_{U_{i,k}}\big(\mathcal{F}\big(U_{i,k}\big)\big)$; then $\big\{U_{i,k}\big\}_{i,k}$ is a cover of U with $\tau|_{U_{i,k}} = \tau_i|_{U_{i,k}} \in \iota_{U_{i,k}}\big(\mathcal{F}\big(U_{i,k}\big)\big)$, so $\tau \in \mathcal{F}^+(U)$. So $\mathcal{F}^+$ is a subsheaf of $\mathcal{G}$ (4.2.30). (None of this used the germ-function construction of (4.2.16).)

ι' satisfies the universal property. We show directly that for every sheaf $\mathcal{K}$,

$$(-)^\sharp : \mathrm{Mor}(\mathcal{F}^+, \mathcal{K}) \to \mathrm{Mor}(\mathcal{F}, \mathcal{K}), \quad \phi \mapsto \phi \circ \iota'$$

is a bijection; it is then manifestly natural in $\mathcal{K}$, being given by post-composition with the fixed ι', so this establishes (4.2.13) for $(\iota', \mathcal{F}^+)$.

Construction of $\psi^\flat$. Let $\psi : \mathcal{F} \to \mathcal{K}$ and $\tau \in \mathcal{F}^+(U)$, with witnessing cover $U = \bigcup_i U_i$, $\tau|_{U_i} = \iota_{U_i}(\sigma_i)$ for (unique, as ι is monic) $\sigma_i \in \mathcal{F}(U_i)$. On overlaps, $\iota_{U_i \cap U_j}\big(\sigma_i|_{U_i \cap U_j}\big) = \tau|_{U_i \cap U_j} = \iota_{U_i \cap U_j}\big(\sigma_j|_{U_i \cap U_j}\big)$, so as ι is monic, $\sigma_i|_{U_i \cap U_j} = \sigma_j|_{U_i \cap U_j}$, whence $\psi_{U_i}(\sigma_i)|_{U_i \cap U_j} = \psi_{U_j}\big(\sigma_j\big)|_{U_i \cap U_j}$. As $\mathcal{K}$ is a sheaf there is therefore a unique $\psi^\flat_U(\tau) \in \mathcal{K}(U)$ with $\psi^\flat_U(\tau)|_{U_i} = \psi_{U_i}(\sigma_i)$ for every i.

Independence of the witnessing cover. If $U = \bigcup_j V_j$, $\tau|_{V_j} = \iota_{V_j}\big(\rho_j\big)$ is another witnessing cover, then on $U_i \cap V_j$, $\iota_{U_i \cap V_j}\big(\sigma_i|_{U_i \cap V_j}\big) = \tau|_{U_i \cap V_j} = \iota_{U_i \cap V_j}\big(\rho_j|_{U_i \cap V_j}\big)$, so (as ι monic) $\sigma_i|_{U_i \cap V_j} = \rho_j|_{U_i \cap V_j}$, whence $\psi_{U_i}(\sigma_i)|_{U_i \cap V_j} = \psi_{V_j}\big(\rho_j\big)|_{U_i \cap V_j}$; as $\big\{U_i \cap V_j\big\}_{i,j}$ covers U, the two candidate values for $\psi^\flat_U(\tau)$ agree on a common refinement, so are equal by separation of $\mathcal{K}$.

$\psi^\flat_U$ is a homomorphism. Given $\tau_1, ..., \tau_{n_\omega} \in \mathcal{F}^+(U)$, using the common refinement cover $\{W\}$ from the first paragraph (with local representatives $\sigma_r \in \mathcal{F}(W)$ for τ_r), independence of witnessing cover gives $\psi^\flat_U(\tau_r)|_W = \psi_W(\sigma_r)$; as this same cover witnesses $\omega_{\mathcal{F}^+(U)}\big(\tau_1, ..., \tau_{n_\omega}\big) \in \mathcal{F}^+(U)$ with local representatives $\omega_{\mathcal{F}(W)}(\sigma_1, ...)$ (shown above), independence of witnessing cover also gives $\psi^\flat_U\big(\omega_{\mathcal{F}^+(U)}(\tau_1, ...)\big)|_W = \psi_W\big(\omega_{\mathcal{F}(W)}(\sigma_1, ...)\big) = \omega_{\mathcal{K}(W)}(\psi_W(\sigma_1), ...) = \omega_{\mathcal{K}(U)}\big(\psi^\flat_U(\tau_1), ...\big)|_W$ for every W, so by separation of $\mathcal{K}$, $\psi^\flat_U\big(\omega_{\mathcal{F}^+(U)}\big(\tau_1, ..., \tau_{n_\omega}\big)\big) = \omega_{\mathcal{K}(U)}\big(\psi^\flat_U(\tau_1), ..., \psi^\flat_U\big(\tau_{n_\omega}\big)\big)$.

© David Rufino CC BY-NC-ND 4.0

$\psi^\flat$ is a morphism of presheaves. For $V \subseteq U$ and $\tau \in \mathcal{F}^+(U)$ with witnessing cover $\{U_i\}$, $\{U_i \cap V\}$ witnesses $\tau|_V$, so $\psi^\flat_V(\tau|_V)|_{U_i\cap V} = \psi_{U_i\cap V}\left(\sigma_i|_{U_i\cap V}\right) = \psi_{U_i}(\sigma_i)|_{U_i\cap V} = \psi^\flat_U(\tau)|_{U_i\cap V}$ for every i, so $\psi^\flat_V(\tau|_V) = \psi^\flat_U(\tau)|_V$ by separation of $\mathcal{K}$.

$\psi^{\flat\sharp} = \psi$. For $\sigma \in \mathcal{F}(U)$, $\iota'_U(\sigma) = \iota_U(\sigma) \in \mathcal{F}^+(U)$ is witnessed by the trivial cover $\{U\}$ with representative σ itself, so $\psi^\flat_U(\iota'_U(\sigma)) = \psi_U(\sigma)$, that is $\psi^\flat \circ \iota' = \psi$.

Uniqueness. If $\phi : \mathcal{F}^+ \to \mathcal{K}$ also satisfies $\phi \circ \iota' = \psi$, then for $\tau \in \mathcal{F}^+(U)$ with witnessing cover $\{U_i\}$, $\phi_U(\tau)|_{U_i} = \phi_{U_i}\left(\tau|_{U_i}\right) = \phi_{U_i}\left(\iota'_{U_i}(\sigma_i)\right) = \psi_{U_i}(\sigma_i) = \psi^\flat_U(\tau)|_{U_i}$ for every i, so $\phi_U(\tau) = \psi^\flat_U(\tau)$ by separation of $\mathcal{K}$; as U, τ were arbitrary, $\phi = \psi^\flat$.

So $(-)^\sharp$ is a bijection with inverse $\psi \mapsto \psi^\flat$, for every sheaf $\mathcal{K}$: $(\iota', \mathcal{F}^+)$ is a sheafification of $\mathcal{F}$ (4.2.13). ∎

4.2.9 Coequaliser Sheaf

Proposition 4.2.33 (Coequaliser Presheaf)
Let $\alpha, \beta : \mathcal{F} \rightrightarrows \mathcal{G}$ be morphisms of presheaves with a family of pointwise coequalisers $\{p_U : \mathcal{G}(U) \to \operatorname{coeq}(\alpha_U, \beta_U)\}_{U\subset X}$. Then we may define

$$\operatorname{coeq}(\alpha, \beta)(U) \coloneqq \operatorname{coeq}(\alpha_U, \beta_U)$$

and this may be made into a presheaf uniquely such that p is a morphism of presheaves. Furthermore $(p, \operatorname{coeq}(\alpha, \beta))$ is a coequaliser in the category of presheaves.

In particular if $\mathcal{C}$ has all coequalisers then so does $\mathbf{PSh}(X; \mathcal{C})$.

Proof. Drafted by Claude Sonnet, reviewed by David Rufino.

We require to show that for all open subsets $V \subset U$ there is a unique restriction morphism $\rho_{UV} : \operatorname{coeq}(\alpha_U, \beta_U) \to \operatorname{coeq}(\alpha_V, \beta_V)$ ensuring the following diagram commutes


$$\begin{array}{ccc} \mathcal{G}(U) & \xrightarrow{p_U} & \operatorname{coeq}(\alpha_U, \beta_U) \\ {\scriptstyle \rho_{UV}}\downarrow & & \downarrow{\scriptstyle \rho_{UV}} \\ \mathcal{G}(V) & \xrightarrow[p_V]{} & \operatorname{coeq}(\alpha_V, \beta_V) \end{array}$$

We have that

$$p_V \circ \rho_{UV} \circ \alpha_U = p_V \circ \alpha_V \circ \rho_{UV} = p_V \circ \beta_V \circ \rho_{UV} = p_V \circ \rho_{UV} \circ \beta_U$$

whence by the universal property applied to $\operatorname{coeq}(\alpha_U, \beta_U)$ there exists a unique factorisation of $p_V \circ \rho_{UV}$ through p_U. The uniqueness then shows that this satisfies the conditions to be a presheaf ($\rho_{UU} = 1$ and $\rho_{VW} \circ \rho_{UV} = \rho_{UW}$ both follow from uniqueness of the factorisation of $p_W \circ \rho_{UW}$ through p_U), and p is a morphism of presheaves.

We now show that p is a coequaliser of (α, β) in the category of presheaves. Evidently $p \circ \alpha = p \circ \beta$, since this holds pointwise by hypothesis. Suppose that $p' : \mathcal{G} \to \mathcal{H}$ is a morphism of presheaves with $p' \circ \alpha = p' \circ \beta$, then $p'_U \circ \alpha_U = p'_U \circ \beta_U$ whence there is a unique factorisation h_U such that $h_U \circ p_U = p'_U$.



The large square and left-hand square commute, which we may use to show that

$$h_V \circ \rho_{UV} \circ p_U = h_V \circ p_V \circ \rho_{UV} = p'_V \circ \rho_{UV} = \rho_{UV} \circ p'_U = \rho_{UV} \circ h_U \circ p_U$$

© David Rufino CC BY-NC-ND 4.0

As p_U is a coequaliser it is an epimorphism (2.6.136), whence $h_V \circ \rho_{UV} = \rho_{UV} \circ h_U$, which shows that the right-hand square commutes. This means precisely that h is a morphism of presheaves, and by construction $h \circ p = p'$.

Finally suppose h' is another morphism of presheaves with $h' \circ p = p'$. Then for every U, $h'_U \circ p_U = p'_U = h_U \circ p_U$, so by the uniqueness of the pointwise factorisation $h'_U = h_U$, that is $h' = h$.

For the final statement, if $\mathcal{C}$ has all coequalisers then the family of pointwise coequalisers $\{p_U : \mathcal{G}(U) \to \mathrm{coeq}(\alpha_U, \beta_U)\}_{U \subset X}$ exists for every pair $\alpha, \beta : \mathcal{F} \rightrightarrows \mathcal{G}$ of morphisms of presheaves, so the above applies to every such pair and $\mathbf{PSh}(X;\mathcal{C})$ has all coequalisers. ∎

Proposition 4.2.34
Let $\alpha, \beta : \mathcal{F} \rightrightarrows \mathcal{G}$ be morphisms of sheaves and $p : \mathcal{G} \to \mathrm{coeq}^{\mathrm{psh}}(\alpha, \beta)$ be a coequaliser in $\mathbf{PSh}(X;\mathcal{C})$. Then

$$\eta \circ p : \mathcal{G} \to \mathrm{coeq}(\alpha, \beta) := \left(\mathrm{coeq}^{\mathrm{psh}}(\alpha, \beta)\right)^+$$

is a coequaliser in $\mathbf{Sh}(X;\mathcal{C})$.

In particular if $\mathcal{C}$ has all coequalisers then so does $\mathbf{Sh}(X;\mathcal{C})$.

Proof. Drafted by Claude Sonnet, reviewed by David Rufino.

By (4.2.13) and (4.2.14), $(-)^+$ is left adjoint to the inclusion $\mathbf{Sh}(X;\mathcal{C}) \hookrightarrow \mathbf{PSh}(X;\mathcal{C})$ with unit η, and this inclusion is (trivially) fully faithful, since a morphism of sheaves is by definition just a morphism of the underlying presheaves. Hence this is an instance of (2.6.144), applied to this reflective subcategory with the coequaliser p of (α, β) in $\mathbf{PSh}(X;\mathcal{C})$.

For the final statement, if $\mathcal{C}$ has all coequalisers then by (4.2.33) $\mathbf{PSh}(X;\mathcal{C})$ has all coequalisers, so the hypotheses of this proposition are satisfied for every pair (α, β) of morphisms of sheaves, and $\mathbf{Sh}(X;\mathcal{C})$ has all coequalisers. ∎

Definition 4.2.35 (Cokernel Presheaf, Cokernel Sheaf)
*Let $\mathcal{C}$ be pointed and algebraic (2.6.185) and $\alpha : \mathcal{F} \to \mathcal{G}$ a morphism of presheaves. Define the **cokernel presheaf** of α by*

$$\mathrm{coker}^{\mathrm{psh}}(\alpha) := \mathrm{coeq}^{\mathrm{psh}}(\alpha, 0), \quad q_U(\sigma) := \sigma \bmod \alpha_U(\mathcal{F}(U)) \quad (4.2.33)$$


*with $q : \mathcal{G} \to \mathrm{coker}^{\mathrm{psh}}(\alpha)$ the coequaliser projection of α and the zero morphism in $\mathbf{PSh}(X;\mathcal{C})$. When $\mathcal{F}, \mathcal{G}$ are sheaves, define the **cokernel sheaf** of α by*

$$\mathrm{coker}(\alpha)(U) := \left\{ \tau : U \to \coprod_{x \in U} \mathcal{G}_x / \alpha_x(\mathcal{F}_x) \;\middle|\; U = \bigcup_i U_i \text{ for some open cover, and } \sigma_i \in \mathcal{G}(U_i) \right.$$
$$\left. \text{such that } \tau(x) = \sigma_{i,x} \bmod \alpha_x(\mathcal{F}_x) \text{ for all } x \in U_i \right\}$$

*with restriction maps given by restricting functions, and the **cokernel map** $p : \mathcal{G} \to \mathrm{coker}(\alpha)$ by $p_U(\sigma)(x) := \sigma_x \bmod \alpha_x(\mathcal{F}_x)$.*

Remark 4.2.36
When α is (in particular) a monomorphism, each α_U is injective (4.2.28), identifying $\mathcal{F}$ with a subsheaf of $\mathcal{G}$; in this case we also write $\mathcal{G}/\mathcal{F} := \mathrm{coker}(\alpha)$.

Lemma 4.2.37 (Cokernel is a Sheaf, and Commutes with Stalks)
Let $\mathcal{C}$ be pointed and algebraic, $\alpha : \mathcal{F} \to \mathcal{G}$ a morphism of sheaves, and p as in (4.2.35). Then $\mathrm{coker}(\alpha)$ is a sheaf, $(\mathrm{coker}(\alpha), p)$ is a coequaliser of $(\alpha, 0)$ in $\mathbf{Sh}(X;\mathcal{C})$, and, for $x \in X$ there is a unique morphism $\theta_x : \mathrm{coker}(\alpha)_x \to \mathcal{G}_x/\alpha_x(\mathcal{F}_x)$ making



commute, and θ_x is (moreover) an isomorphism.

Proof. Drafted by Claude Sonnet, reviewed by David Rufino.

© David Rufino CC BY-NC-ND 4.0

An auxiliary sheafification-based model. By (4.2.34), $p^0 \coloneqq \eta \circ q : \mathcal{G} \to \operatorname{coker}^0(\alpha) \coloneqq \left(\operatorname{coker}^{\mathrm{psh}}(\alpha)\right)^+$ is a co-equaliser of $(\alpha, 0)$ in $\mathbf{Sh}(X;\mathcal{C})$, with $q, \operatorname{coker}^{\mathrm{psh}}(\alpha)$ as in (4.2.35) and η the sheafification unit (4.2.16) this auxiliary model is used only within this proof, to relate the definition above to the general existence of coequalisers.

η_x is an isomorphism. (4.2.18).

q_x is surjective with kernel $\alpha_x(\mathcal{F}_x)$. Since $\operatorname{coker}^{\mathrm{psh}}(\alpha)(U) = \mathcal{G}(U)/\alpha_U(\mathcal{F}(U))$ (4.2.33), by definition of stalk $\left(\operatorname{coker}^{\mathrm{psh}}(\alpha)\right)_x = \varinjlim_{U \ni x} \mathcal{G}(U)/\alpha_U(\mathcal{F}(U))$, with $q_x(\sigma_x)$ the class of σ in this direct limit for $\sigma \in \mathcal{G}(U)$, $U \ni x$.

Surjective: every element of the direct limit is represented by the class of some $\sigma \in \mathcal{G}(U)$, $U \ni x$, that is $q_x(\sigma_x)$.

Kernel $\supseteq \alpha_x(\mathcal{F}_x)$: for $\rho \in \mathcal{F}_x$, represented by $\tau \in \mathcal{F}(U)$, $\alpha_x(\rho) = (\alpha_U(\tau))_x$; as $\alpha_U(\tau) \in \alpha_U(\mathcal{F}(U))$, its class in $\mathcal{G}(U)/\alpha_U(\mathcal{F}(U))$ is 0, so $q_x(\alpha_x(\rho)) = 0$.

Kernel $\subseteq \alpha_x(\mathcal{F}_x)$: suppose $\sigma \in \mathcal{G}(U)$, $U \ni x$, with $q_x(\sigma_x) = 0$, that is, the class of σ becomes 0 in the direct limit. By (4.2.6) applied to the presheaf $U \mapsto \mathcal{G}(U)/\alpha_U(\mathcal{F}(U))$, this means there is $W \subseteq U$, $x \in W$, with $\sigma|_W \in \alpha_W(\mathcal{F}(W))$, that is $\sigma|_W = \alpha_W(\tau)$ for some $\tau \in \mathcal{F}(W)$; then $\sigma_x = (\sigma|_W)_x = (\alpha_W(\tau))_x = \alpha_x(\tau_x) \in \alpha_x(\mathcal{F}_x)$.

Existence and uniqueness of ϑ^0_x with $\vartheta^0_x \circ q_x = \pi_x$. Since q_x is surjective with kernel $\alpha_x(\mathcal{F}_x)$, it is (up to the canonical comparison isomorphism) the quotient map $\mathcal{G}_x \to \mathcal{G}_x/\alpha_x(\mathcal{F}_x)$, hence satisfies the corresponding universal property: any morphism out of $\mathcal{G}_x$ vanishing on $\ker(q_x)$ factors uniquely through q_x. As π_x vanishes on $\ker(\pi_x) = \alpha_x(\mathcal{F}_x) = \ker(q_x)$, this gives a unique $\vartheta^0_x : \left(\operatorname{coker}^{\mathrm{psh}}(\alpha)\right)_x \to \mathcal{G}_x/\alpha_x(\mathcal{F}_x)$ with $\vartheta^0_x \circ q_x = \pi_x$; uniqueness alone accounts for this, and says nothing yet about ϑ^0_x being an isomorphism.

ϑ^0_x is an isomorphism. This is a separate check. *Surjective*: given $z \in \mathcal{G}_x/\alpha_x(\mathcal{F}_x)$, as π_x is surjective $z = \pi_x(g)$ for some $g \in \mathcal{G}_x$, and $z = \vartheta^0_x(q_x(g))$. *Injective*: every element of $\left(\operatorname{coker}^{\mathrm{psh}}(\alpha)\right)_x$ is $q_x(g)$ for some g, since q_x is surjective; if $\vartheta^0_x(q_x(g)) = 0$ then $\pi_x(g) = 0$, that is $g \in \ker(\pi_x) = \ker(q_x)$, so $q_x(g) = 0$.

$\vartheta_x \coloneqq \vartheta^0_x \circ \eta_x^{-1}$ is an isomorphism $\operatorname{coker}^0(\alpha)_x \to \mathcal{G}_x/\alpha_x(\mathcal{F}_x)$ with $\vartheta_x \circ p^0_x = \pi_x$. It is an isomorphism, being a composite of the isomorphisms η_x^{-1} and ϑ^0_x. As $p^0_x = \eta_x \circ q_x$,

$$\vartheta_x \circ p^0_x = \vartheta^0_x \circ \eta_x^{-1} \circ \eta_x \circ q_x = \vartheta^0_x \circ q_x = \pi_x$$

Relating $\operatorname{coker}^0(\alpha)$ to $\operatorname{coker}(\alpha)$. By (4.2.16), elements of $\operatorname{coker}^0(\alpha)(U)$ are functions $\tau^0 : U \to \coprod_{x \in U} \operatorname{coker}^0(\alpha)_x$ such that, for some open cover $U = \bigcup_i U_i$ and $\sigma_i \in \mathcal{G}(U_i)$, $\tau^0(x) = p^0_{U_i}(\sigma_i)_x$ for $x \in U_i$. Using the isomorphisms ϑ_x, define

$$\vartheta : \coprod_{x \in X} \operatorname{coker}^0(\alpha)_x \longrightarrow \coprod_{x \in X} \mathcal{G}_x/\alpha_x(\mathcal{F}_x), \quad \vartheta|_{\operatorname{coker}^0(\alpha)_x} \coloneqq \vartheta_x$$

a bijection, being a disjoint union of the bijections ϑ_x. Set $\Psi_U(\tau^0) \coloneqq \vartheta \circ \tau^0$ for $\tau^0 \in \operatorname{coker}^0(\alpha)(U)$.

Ψ_U is a well-defined bijection onto $\operatorname{coker}(\alpha)(U)$. As ϑ is a bijection carrying each fibre $\operatorname{coker}^0(\alpha)_x$ to $\mathcal{G}_x/\alpha_x(\mathcal{F}_x)$, post-composition by ϑ is a bijection from functions $U \to \coprod_{x \in U} \operatorname{coker}^0(\alpha)_x$ to functions $U \to \coprod_{x \in U} \mathcal{G}_x/\alpha_x(\mathcal{F}_x)$, carrying the representing data (U_i, σ_i) for τ^0 to representing data for $\Psi_U(\tau^0)$: $\vartheta\left(p^0_{U_i}(\sigma_i)_x\right) = \vartheta_x\left(p^0_{U_i}(\sigma_i)_x\right) = \sigma_{i,x} \bmod \alpha_x(\mathcal{F}_x)$, so $\Psi_U(\tau^0)(x) = \sigma_{i,x} \bmod \alpha_x(\mathcal{F}_x)$ for $x \in U_i$, that is, exactly membership in $\operatorname{coker}(\alpha)(U)$ (4.2.35). Hence Ψ_U restricts to a bijection $\operatorname{coker}^0(\alpha)(U) \to \operatorname{coker}(\alpha)(U)$.

Ψ is an isomorphism of presheaves, and $\operatorname{coker}(\alpha)$ *is a sheaf.* For $V \subseteq U$ and $\tau^0 \in \operatorname{coker}^0(\alpha)(U)$, $\Psi_V(\tau^0|_V) = \vartheta \circ (\tau^0|_V) = (\vartheta \circ \tau^0)|_V = \Psi_U(\tau^0)|_V$, since restricting a function and then composing with ϑ agrees with composing then restricting; hence Ψ commutes with restriction, and is a natural transformation, bijective at every U — an isomorphism of presheaves $\operatorname{coker}^0(\alpha) \cong \operatorname{coker}(\alpha)$. As $\operatorname{coker}^0(\alpha)$ is a sheaf and the sheaf condition transports across an isomorphism of presheaves, $\operatorname{coker}(\alpha)$ is also a sheaf.

$\Psi \circ p^0 = p$. For $\sigma \in \mathcal{G}(U)$, $x \in U$, $\Psi_U\left(p^0_U(\sigma)\right)(x) = \vartheta_x\left(p^0_U(\sigma)_x\right) = \sigma_x \bmod \alpha_x(\mathcal{F}_x) = p_U(\sigma)(x)$, again using $\vartheta_x \circ p^0_x = \pi_x$.

$(\operatorname{coker}(\alpha), p)$ is a coequaliser of $(\alpha, 0)$. Since $p^0 \circ \alpha = p^0 \circ 0$ (as p^0 coequalises $(\alpha, 0)$) and $p = \Psi \circ p^0$, also $p \circ \alpha = \Psi \circ p^0 \circ \alpha = \Psi \circ p^0 \circ 0 = p \circ 0$, so $(\operatorname{coker}(\alpha), p)$ is a co-cone for $(\alpha, 0)$; and Ψ, an isomorphism of presheaves between sheaves, is also an isomorphism of $\mathbf{Sh}(X;\mathcal{C})$ (this inclusion into $\mathbf{PSh}(X;\mathcal{C})$ being fully faithful (4.2.34)). As $(\operatorname{coker}^0(\alpha), p^0)$ is a coequaliser of $(\alpha, 0)$, that is, a colimit of the corresponding diagram (2.6.141), (2.6.90) shows $(\operatorname{coker}(\alpha), p)$ is also such a colimit, that is, a coequaliser of $(\alpha, 0)$.

Conclusion. Set $\theta_x \coloneqq \vartheta_x \circ \Psi_x^{-1}$, an isomorphism, being a composite of the isomorphisms ϑ_x and Ψ_x^{-1}. Since $p_x = \Psi_x \circ p^0_x$ (4.2.7),

© David Rufino CC BY-NC-ND 4.0

$$\theta_x \circ p_x = \vartheta_x \circ \Psi_x^{-1} \circ \Psi_x \circ p_x^0 = \vartheta_x \circ p_x^0 = \pi_x$$

and θ_x is unique with this property, since $p_x = \Psi_x \circ p_x^0$ is surjective (a composite of the surjections Ψ_x, an isomorphism, and $p_x^0 = \eta_x \circ q_x$), hence epic, so at most one morphism out of $\operatorname{coker}(\alpha)_x$ can satisfy $\theta_x \circ p_x = \pi_x$. ∎

4.2.10 Image Sheaf

Unlike kernels and equalisers (4.2.24), the image is not itself a limit — it is initial in the category of factorisations through a monomorphism (2.6.168) — so the argument via (4.2.3) and (2.6.99) used there does not apply, and the pointwise image presheaf need not satisfy the sheaf condition even when $\mathcal{F}, \mathcal{G}$ are sheaves; instead an image sheaf is obtained by sheafifying the image presheaf (4.2.40). We do however first record the following easier pointwise construction of an image presheaf, whenever $\mathcal{C}$ has regular epi-mono factorisations.

Lemma 4.2.38 (Regular Epimorphisms of Presheaves may be Determined Pointwise)
Let $e : \mathcal{F} \to \mathcal{G}$ be a morphism of presheaves such that e_U is a regular epimorphism in $\mathcal{C}$ for every $U \subset X$. Then e is a regular epimorphism of $\mathbf{PSh}(X;\mathcal{C})$.

Proof. Drafted by Claude Sonnet, reviewed by David Rufino.


Since $\mathcal{C}$ has products and equalisers, by (2.6.146) e_U has a kernel pair for every U, so by (4.2.27) these assemble into a kernel pair $(\mathcal{F} \times_{\mathcal{G}} \mathcal{F}, p_1, p_2)$ of e in $\mathbf{PSh}(X;\mathcal{C})$, with U-components $\left(\mathcal{F}(U) \times_{\mathcal{G}(U)} \mathcal{F}(U), p_{1,U}, p_{2,U}\right)$ a kernel pair of e_U.

Since e_U is regular, (2.6.150) shows e_U is the coequaliser of $\left(p_{1,U}, p_{2,U}\right)$, for every U. So applying (4.2.33) to the pair of morphisms of presheaves $(p_1, p_2) : \mathcal{F} \times_{\mathcal{G}} \mathcal{F} \rightrightarrows \mathcal{F}$, this pointwise family of coequalisers assembles into a presheaf $\operatorname{coeq}(p_1, p_2)$ with restriction maps uniquely determined by making the induced map $q := \{e_U\}$ a morphism of presheaves, and $(q, \operatorname{coeq}(p_1, p_2))$ is a coequaliser of (p_1, p_2) in $\mathbf{PSh}(X;\mathcal{C})$.

The presheaf $\mathcal{G}$ also has U-component $\mathcal{G}(U) = \operatorname{coeq}(p_1, p_2)(U)$, with restriction maps uniquely determined by the same condition, that e is a morphism of presheaves; by this uniqueness, $\mathcal{G} = \operatorname{coeq}(p_1, p_2)$ as presheaves and $e = q$. Hence e is a coequaliser of presheaves, i.e. a regular epimorphism of $\mathbf{PSh}(X;\mathcal{C})$. ∎

Proposition 4.2.39 (Image Presheaf may be constructed pointwise)
If $\mathcal{C}$ has regular epi-mono factorisations for every morphism, e.g. **Set** *(2.6.176) or an Abelian category (2.7.7), then $\mathbf{PSh}(X;\mathcal{C})$ has regular epi-mono factorisations for every morphism.*

More precisely, let $\alpha : \mathcal{F} \to \mathcal{G}$ be a morphism of presheaves with a family of pointwise regular epi-mono factorisations $\{\alpha_U = m_U \circ e_U\}_{U \subset X}$




Then the assignment $\operatorname{Im}(\alpha)(U) := \operatorname{Im}(\alpha_U)$ can be made into a presheaf uniquely such that $m_\alpha : \operatorname{Im}(\alpha) \to \mathcal{G}$, $(m_\alpha)_U := m_U$, and $e_\alpha : \mathcal{F} \to \operatorname{Im}(\alpha)$, $(e_\alpha)_U := e_U$, are morphisms of presheaves, and $m_\alpha \circ e_\alpha = \alpha$; furthermore this is a regular epi-mono factorisation of α in $\mathbf{PSh}(X;\mathcal{C})$, and hence $(\operatorname{Im}(\alpha), e_\alpha, m_\alpha)$ is an image of α there (2.6.175).

If moreover $\mathcal{C}$ is algebraic (2.6.185), the pointwise factorisations may be taken as in (2.6.197), namely $\operatorname{Im}(\alpha_U) := U(\alpha_U)(U(\mathcal{F}(U))) \subseteq U(\mathcal{G}(U))$; then $\operatorname{Im}(\alpha)$ is simply the subpresheaf of $\mathcal{G}$ of sections pointwise in the image of α (4.2.30), with restriction maps inherited from $\mathcal{G}$ and m_α the inclusion.

Proof. Drafted by Claude Sonnet, reviewed by David Rufino.

Restriction maps. Fix $V \subseteq U$. Since α is a morphism of presheaves, $\rho^{\mathcal{G}}_{UV} \circ \alpha_U = \alpha_V \circ \rho^{\mathcal{F}}_{UV}$, so applying (2.6.178) to the regular epi-mono factorisations $\alpha_U = m_U \circ e_U$, $\alpha_V = m_V \circ e_V$ with $a := \rho^{\mathcal{F}}_{UV}$, $b := \rho^{\mathcal{G}}_{UV}$ gives a unique morphism $\rho^{\mathrm{Im}}_{UV} : \operatorname{Im}(\alpha_U) \to \operatorname{Im}(\alpha_V)$ making the following diagram commute

© David Rufino CC BY-NC-ND 4.0



namely

$$\rho^{\mathrm{Im}}_{UV} \circ e_U = e_V \circ \rho^{\mathcal{F}}_{UV} \quad \text{and} \quad m_V \circ \rho^{\mathrm{Im}}_{UV} = \rho^{\mathcal{G}}_{UV} \circ m_U$$

The first identity is precisely the condition for e_α to be a morphism of presheaves, and the second for m_α to be one.

$\mathrm{Im}(\alpha)$ is a presheaf. By the functoriality clause of (2.6.178) (taking $a = 1_{\mathcal{F}(U)}$, $b = 1_{\mathcal{G}(U)}$), $\rho^{\mathrm{Im}}_{UU} = 1_{\mathrm{Im}(\alpha_U)}$; and for $W \subseteq V \subseteq U$, taking $a := \rho^{\mathcal{F}}_{UV}$, $b := \rho^{\mathcal{G}}_{UV}$, $c := \rho^{\mathcal{F}}_{VW}$, $d := \rho^{\mathcal{G}}_{VW}$, the composition law gives $\rho^{\mathrm{Im}}_{VW} \circ \rho^{\mathrm{Im}}_{UV} = \rho^{\mathrm{Im}}_{UW}$ (using $\rho^{\mathcal{F}}_{VW} \circ \rho^{\mathcal{F}}_{UV} = \rho^{\mathcal{F}}_{UW}$, and likewise for $\mathcal{G}$). So $\mathrm{Im}(\alpha)$, with these restriction maps, is a presheaf, and e_α, m_α are morphisms of presheaves. Uniqueness of restriction maps making e_α, m_α into morphisms of presheaves is immediate from the uniqueness clause of (2.6.178), since these two identities are exactly what is required of e_α, m_α.

Finally $(m_\alpha \circ e_\alpha)_U = m_U \circ e_U = \alpha_U$ for every U, so $m_\alpha \circ e_\alpha = \alpha$.

Regularity. By (4.2.38) (using that $\mathcal{C}$ has products and equalisers, and each e_U is regular), e_α is a regular epimorphism of $\mathbf{PSh}(X;\mathcal{C})$. Since m_U is monic in $\mathcal{C}$ for every U, m_α is monic in $\mathbf{PSh}(X;\mathcal{C})$ (4.2.28). So $\alpha = m_\alpha \circ e_\alpha$ is a regular epi-mono factorisation of α in $\mathbf{PSh}(X;\mathcal{C})$ (2.6.174), and hence also an image of α there (2.6.175).

For the first statement, if $\mathcal{C}$ has regular epi-mono factorisations for every morphism then every $\alpha_U : \mathcal{F}(U) \to \mathcal{G}(U)$ has one, giving such a family for every morphism α of presheaves, so the above applies to every such α and $\mathbf{PSh}(X;\mathcal{C})$ has regular epi-mono factorisations for every morphism.

The algebraic case. If $\mathcal{C}$ is algebraic, taking $\mathrm{Im}(\alpha_U) := U(\alpha_U)(U(\mathcal{F}(U)))$ as in (2.6.197), the restriction maps constructed above may be identified with those inherited from $\mathcal{G}$: since α is a morphism of presheaves, for $\sigma \in \mathcal{F}(U)$, $\rho^{\mathcal{G}}_{UV}(\alpha_U(\sigma)) = \alpha_V\big(\rho^{\mathcal{F}}_{UV}(\sigma)\big) \in \mathrm{Im}(\alpha_V)$, so $\rho^{\mathcal{G}}_{UV}(\mathrm{Im}(\alpha_U)) \subseteq \mathrm{Im}(\alpha_V)$; by the uniqueness clause of (2.6.178), this restriction of $\rho^{\mathcal{G}}_{UV}$ is exactly ρ^{Im}_{UV}. So in this case $\mathrm{Im}(\alpha)$ is simply the subpresheaf of $\mathcal{G}$ of sections pointwise in the image of α (4.2.30), with e_α the (pointwise surjective) corestriction of α and m_α the inclusion — without needing (2.6.178) to construct the restriction maps in the first place. ∎

Proposition 4.2.40 (Image Sheaf)
Let $\mathcal{C}$ be algebraic (2.6.185), hence have regular epi-mono factorisations for every morphism (2.6.197). Then $\mathbf{Sh}(X;\mathcal{C})$ has regular epi-mono factorisations for every morphism.

More precisely, let $\alpha : \mathcal{F} \to \mathcal{G}$ be a morphism of sheaves, and define

$$\mathrm{Im}(\alpha)(U) := \left\{ \tau \in U(\mathcal{G}(U)) \;\middle|\; U = \bigcup_{i\in I} U_i \text{ for some open cover, with } \tau|_{U_i} \in U\big(\alpha_{U_i}\big)(U(\mathcal{F}(U_i))) \text{ for every } i \right\}$$

the subsheaf of $\mathcal{G}$ of sections locally in the (pointwise) image of α (4.2.32). Then

$$e : \mathcal{F} \longrightarrow \mathrm{Im}(\alpha), e_U(\sigma) := \alpha_U(\sigma) \quad \text{and} \quad m : \mathrm{Im}(\alpha) \hookrightarrow \mathcal{G} \text{ the inclusion}$$

are morphisms of sheaves with $m \circ e = \alpha$, and this is a regular epi-mono factorisation of α in $\mathbf{Sh}(X;\mathcal{C})$; hence $(\mathrm{Im}(\alpha), e, m)$ is an image of α there (2.6.175).

Proof. Drafted by Claude Sonnet, reviewed by David Rufino.

Setup. Let $\big(\mathrm{Im}^{\mathrm{psh}}(\alpha), e_\alpha, m_\alpha\big)$ be the image of α in $\mathbf{PSh}(X;\mathcal{C})$, so $\alpha = m_\alpha \circ e_\alpha$ is a regular epi-mono factorisation there (4.2.39), with $\mathrm{Im}^{\mathrm{psh}}(\alpha)(U) = \mathrm{Im}(\alpha_U) \subseteq \mathcal{G}(U)$ and $(m_\alpha)_U$ the inclusion. In particular m_α is a monomorphism of presheaves (4.2.28), and "locally in the image of m_α" is the same condition as "locally in the image of α", so applying

© David Rufino CC BY-NC-ND 4.0

(4.2.32) to $m_\alpha\colon \mathrm{Im}(\alpha)$ (as defined above) is a subsheaf of $\mathcal{G}$, and the corestriction $\iota' : \mathrm{Im}^{\mathrm{psh}}(\alpha) \to \mathrm{Im}(\alpha)$ of m_α is a sheafification of $\mathrm{Im}^{\mathrm{psh}}(\alpha)$. Since ι' is a corestriction of the injective map $(m_\alpha)_U$ for every U, ι' is itself monic (4.2.28).

Factorisation. For $\sigma \in \mathcal{F}(U)$, $\iota'_U\big((e_\alpha)_U(\sigma)\big) = (m_\alpha)_U\big((e_\alpha)_U(\sigma)\big) = \alpha_U(\sigma)$, so $e = \iota' \circ e_\alpha$; as $\mathcal{F}$ and $\mathrm{Im}(\alpha)$ are sheaves and ι', e_α are morphisms of presheaves, e is a morphism of sheaves. Writing $j : \mathrm{Im}(\alpha) \hookrightarrow \mathcal{G}$ for the inclusion, $j \circ \iota' = m_\alpha$, so

$$m \circ e = j \circ \iota' \circ e_\alpha = m_\alpha \circ e_\alpha = \alpha$$

Monicity. m is the inclusion of the subsheaf $\mathrm{Im}(\alpha)$ into $\mathcal{G}$, hence monic by the remark following (4.2.30).

Regularity. By (4.2.27), e_α has a kernel pair $\big(\mathcal{F} \times_{\mathrm{Im}^{\mathrm{psh}}(\alpha)} \mathcal{F}, p_1, p_2\big)$ in $\mathbf{PSh}(X;\mathcal{C})$, with U-components $\big\{(\sigma,\tau) \in \mathcal{F}(U)^2 \;\big|\; (e_\alpha)_U(\sigma) = (e_\alpha)_U(\tau)\big\}$; as each $(e_\alpha)_U = e_U$ is regular (4.2.38) exhibits e_α as its coequaliser there.

As ι' is monic, $(e_\alpha)_U(\sigma) = (e_\alpha)_U(\tau)$ if and only if $\iota'_U\big((e_\alpha)_U(\sigma)\big) = \iota'_U\big((e_\alpha)_U(\tau)\big)$, that is $e_U(\sigma) = e_U(\tau)$, so the U-components of the kernel pairs of e_α and of e agree, with the same projections p_1, p_2 into $\mathcal{F}$; by the uniqueness clause of (4.2.27), the two kernel-pair presheaves coincide. Since $\mathcal{F}$ and $\mathrm{Im}(\alpha)$ are both sheaves, applying (4.2.27) to e directly, in its sheaf case, shows this common kernel pair is itself a sheaf, with p_1, p_2 morphisms of sheaves.

It remains to show e is a coequaliser of (p_1, p_2) in $\mathbf{Sh}(X;\mathcal{C})$. Let $q' : \mathcal{F} \to \mathcal{K}$ be a morphism of sheaves with $q' \circ p_1 = q' \circ p_2$. As e_α is a coequaliser of (p_1, p_2) in $\mathbf{PSh}(X;\mathcal{C})$, there is a unique morphism of presheaves $\tilde{q} : \mathrm{Im}^{\mathrm{psh}}(\alpha) \to \mathcal{K}$ with $\tilde{q} \circ e_\alpha = q'$. Since $\mathcal{K}$ is a sheaf, the bijection $(-)^\sharp : \mathrm{Mor}(\mathrm{Im}(\alpha), \mathcal{K}) \to \mathrm{Mor}\big(\mathrm{Im}^{\mathrm{psh}}(\alpha), \mathcal{K}\big)$, $\phi \mapsto \phi \circ \iota'$, from the universal property of ι' (4.2.32) gives a unique $\bar{q} : \mathrm{Im}(\alpha) \to \mathcal{K}$ with $\bar{q} \circ \iota' = \tilde{q}$. Then

$$\bar{q} \circ e = \bar{q} \circ \iota' \circ e_\alpha = \tilde{q} \circ e_\alpha = q'$$

For uniqueness, if $\bar{q}' : \mathrm{Im}(\alpha) \to \mathcal{K}$ also satisfies $\bar{q}' \circ e = q'$, then $(\bar{q}' \circ \iota') \circ e_\alpha = \bar{q}' \circ e = q' = \tilde{q} \circ e_\alpha$, and as e_α is epic (2.6.136), $\bar{q}' \circ \iota' = \tilde{q}$; by uniqueness in the bijection $(-)^\sharp$, $\bar{q}' = \bar{q}$.

So $(p_1, p_2) : \mathcal{F} \times_{\mathrm{Im}^{\mathrm{psh}}(\alpha)} \mathcal{F} \rightrightarrows \mathcal{F}$ are morphisms of $\mathbf{Sh}(X;\mathcal{C})$ with coequaliser e there. In particular e is a regular epimorphism of $\mathbf{Sh}(X;\mathcal{C})$, so $\alpha = m \circ e$ is a regular epi-mono factorisation of α in $\mathbf{Sh}(X;\mathcal{C})$ (2.6.174), and hence $(\mathrm{Im}(\alpha), e, m)$ is an image of α there (2.6.175). ∎

Remark 4.2.41
*As established in the **Setup** above, $\iota' : \mathrm{Im}^{\mathrm{psh}}(\alpha) \to \mathrm{Im}(\alpha)$ is a sheafification of the pointwise image presheaf $\mathrm{Im}^{\mathrm{psh}}(\alpha)$ (4.2.39): this recovers the abstract description of the image sheaf, promised at the start of this subsection, as obtained by sheafifying the image presheaf, from the concrete construction above.*

Corollary 4.2.42 (Stalk Criterion for Surjectivity)
Let $\mathcal{C}$ be algebraic (2.6.185), $\alpha : \mathcal{F} \to \mathcal{G}$ a morphism of sheaves, and $\alpha = m \circ e$ a regular epi-mono factorisation of α in $\mathbf{Sh}(X;\mathcal{C})$ (4.2.40). Then m is an isomorphism if and only if α_x is surjective for every $x \in X$.

Proof. Drafted by Claude Sonnet, reviewed by David Rufino.

The canonical factorisation. Write $\alpha = m_0 \circ e_0$ for the canonical factorisation of (4.2.40), where $e_0 = \eta \circ e_\alpha$, with $\big(\mathrm{Im}^{\mathrm{psh}}(\alpha), e_\alpha, m_\alpha\big)$ the image of α in $\mathbf{PSh}(X;\mathcal{C})$, built from pointwise regular epi-mono factorisations $\{\alpha_U = m_U \circ e_U\}_{U \subset X}$ (4.2.39), and $\eta : \mathrm{Im}^{\mathrm{psh}}(\alpha) \to \mathrm{Im}(\alpha) =: \big(\mathrm{Im}^{\mathrm{psh}}(\alpha)\big)^+$ the corestriction of m_α, a sheafification of $\mathrm{Im}^{\mathrm{psh}}(\alpha)$ (4.2.32), i.e. the unit of the sheafification as defined in (4.2.13).

Fix $x \in X$. As $\mathcal{C}$ is algebraic and e_U is a regular epimorphism of $\mathcal{C}$ for every U, (2.6.196) shows e_U is surjective for every U. Given $\xi \in \mathrm{Im}^{\mathrm{psh}}(\alpha)_x$, by (4.2.6) $\xi = \iota_x$ for some $\iota \in \mathrm{Im}^{\mathrm{psh}}(\alpha)(U) = \mathrm{Im}(\alpha_U)$, $U \ni x$; as e_U is surjective, $\iota = e_U(\sigma)$ for some $\sigma \in \mathcal{F}(U)$, so $\xi = \iota_x = (e_U(\sigma))_x = (e_\alpha)_x(\sigma_x)$. So $(e_\alpha)_x$ is surjective.

By (4.2.18), η_x is an isomorphism, in particular surjective. Hence $(e_0)_x = \eta_x \circ (e_\alpha)_x$, being a composite of surjections, is surjective.

Since $\alpha_x = (m_0)_x \circ (e_0)_x$ (4.2.7) with $(e_0)_x$ surjective, α_x is surjective if and only if $(m_0)_x$ is surjective: if $(m_0)_x$ is surjective this is immediate, as a composite of surjections; conversely if α_x is surjective then $\mathrm{Im}\big((m_0)_x\big) \supseteq \mathrm{Im}(\alpha_x) = \mathcal{G}_x$. So by (4.2.29), m_0 is an isomorphism if and only if α_x is surjective for every $x \in X$.

General regular epi-mono factorisations. Let $\alpha = m \circ e$ be any regular epi-mono factorisation of α in $\mathbf{Sh}(X;\mathcal{C})$. By (2.6.179), applied to this and the canonical factorisation $\alpha = m_0 \circ e_0$, there is an isomorphism ϕ with $m_0 \circ \phi = m$. So m is an isomorphism if and only if m_0 is (composing with the isomorphism ϕ on the right preserves being an isomorphism, in either direction), which by the above holds if and only if α_x is surjective for every $x \in X$. ∎

© David Rufino CC BY-NC-ND 4.0

4.2.11 Exactness

Definition 4.2.43 (Exact Sequence of Sheaves)
*Let $\mathcal{C}$ be pointed and algebraic; then $\mathbf{Sh}(X;\mathcal{C})$ is evidently pointed (zero morphisms computed pointwise, the constant zero presheaf being a zero object), with a regular epi-mono factorisation (4.2.40) for every morphism, so (2.6.201) applies with $\mathbf{Sh}(X;\mathcal{C})$ in place of $\mathcal{C}$: a pair of composable morphisms of sheaves $\mathcal{F} \xrightarrow{\alpha} \mathcal{G} \xrightarrow{\beta} \mathcal{H}$ is **exact at $\mathcal{G}$** if $\mathrm{Im}(\alpha) = \ker(\beta)$, that is $(\mathrm{Im}(\alpha), m)$ is (literally) a kernel of β, for any (equivalently, by the same argument, every) choice of image factorisation $\alpha = m \circ e$ — in particular we may as well take the canonical construction of (4.2.40).*

Proposition 4.2.44 (Concrete Criterion for Exactness of Sheaves)
Let $\mathcal{C}$ be pointed and algebraic and $\mathcal{F} \xrightarrow{\alpha} \mathcal{G} \xrightarrow{\beta} \mathcal{H}$ a pair of composable morphisms of sheaves. Regard $\mathrm{Im}(\alpha)$ (4.2.40) and $\ker(\beta)$ (4.2.25) as the canonical constructions, both literal subsheaves of $\mathcal{G}$ since $\mathcal{C}$ is algebraic. Then the following are equivalent

a) the pair is exact at $\mathcal{G}$ (4.2.43) (for any choice of image factorisation of α)
b) $\mathrm{Im}(\alpha)(U) = \ker(\beta)(U)$, as subsets of $\mathcal{G}(U)$, for every open $U \subset X$
c) $\beta \circ \alpha = 0$, and every section of $\ker(\beta)(U)$ is locally in the image of α, for every open $U \subset X$

Proof. Drafted by Claude Sonnet, reviewed by David Rufino.

Write $\alpha = m \circ e$ for the canonical image factorisation (4.2.40) and $i : \ker(\beta) \hookrightarrow \mathcal{G}$ for the canonical kernel (4.2.25).

a) $\iff$ b) **Reduction to the canonical representatives.** By (4.2.43) (applied with $\mathbf{Sh}(X;\mathcal{C})$ in place of $\mathcal{C}$), exactness at $\mathcal{G}$ does not depend on the choice of image factorisation of α, so it suffices to treat the canonical representative m: the pair is exact at $\mathcal{G}$ if and only if $(\mathrm{Im}(\alpha), m)$ is a kernel of β. Since $(\ker(\beta), i)$ is also a kernel of β, by (2.6.80) this holds if and only if there is an isomorphism $\theta : \mathrm{Im}(\alpha) \to \ker(\beta)$ with $i \circ \theta = m$.

($\Longleftarrow$) If $\mathrm{Im}(\alpha)(U) = \ker(\beta)(U)$ for every U, then $m = i$ as morphisms of sheaves, both being the inclusion of this common subsheaf into $\mathcal{G}$, with the (unique, (2.6.194)) induced structure making it a morphism. So $\theta := 1_{\mathrm{Im}(\alpha)}$ satisfies $i \circ \theta = i = m$.

($\Longrightarrow$) Suppose there is an isomorphism $\theta : \mathrm{Im}(\alpha) \to \ker(\beta)$ with $i \circ \theta = m$. Fix $U \subset X$ and $\sigma \in \mathrm{Im}(\alpha)(U)$; since m_U is the inclusion of $\mathrm{Im}(\alpha)(U)$ into $\mathcal{G}(U)$, $i_U(\theta_U(\sigma)) = m_U(\sigma) = \sigma$, and i_U is also an inclusion, so $\theta_U(\sigma) = \sigma$; hence $\sigma \in \ker(\beta)(U)$, giving $\mathrm{Im}(\alpha)(U) \subseteq \ker(\beta)(U)$. As θ is an isomorphism, θ_U is a bijection onto $\ker(\beta)(U)$, so every element of $\ker(\beta)(U)$ equals $\theta_U(\sigma) = \sigma$ for some $\sigma \in \mathrm{Im}(\alpha)(U)$, giving $\ker(\beta)(U) \subseteq \mathrm{Im}(\alpha)(U)$. So $\mathrm{Im}(\alpha)(U) = \ker(\beta)(U)$ for every U.

b) $\iff$ c) Condition *b)* is, for every U, the conjunction of $\mathrm{Im}(\alpha)(U) \subseteq \ker(\beta)(U)$ and $\ker(\beta)(U) \subseteq \mathrm{Im}(\alpha)(U)$; the latter is precisely "every section of $\ker(\beta)(U)$ is locally in the image of α" (4.2.40), so it remains to show $\mathrm{Im}(\alpha)(U) \subseteq \ker(\beta)(U)$ for every U is equivalent to $\beta \circ \alpha = 0$.

($\Longleftarrow$) Suppose $\beta \circ \alpha = 0$. For $\tau \in \mathrm{Im}(\alpha)(U)$, take a witnessing cover $U = \bigcup_j U_j$ with $\tau|_{U_j} = \alpha_{U_j}(\sigma_j)$ for some $\sigma_j \in \mathcal{F}(U_j)$; then $\beta_U(\tau)|_{U_j} = \beta_{U_j}(\tau|_{U_j}) = \beta_{U_j}(\alpha_{U_j}(\sigma_j)) = (\beta \circ \alpha)_{U_j}(\sigma_j) = 0$ for every j, so by separation of $\mathcal{H}$ (4.2.4), $\beta_U(\tau) = 0$, i.e. $\tau \in \ker(\beta)(U)$.

($\Longrightarrow$) Suppose $\mathrm{Im}(\alpha)(U) \subseteq \ker(\beta)(U)$ for every U. For any U and $\sigma \in \mathcal{F}(U)$, $\alpha_U(\sigma) \in \mathrm{Im}(\alpha)(U)$ (witnessed by the trivial cover $\{U\}$), so $\alpha_U(\sigma) \in \ker(\beta)(U)$, that is $\beta_U(\alpha_U(\sigma)) = 0$. As U, σ were arbitrary, $\beta \circ \alpha = 0$. ∎

Lemma 4.2.45 (Subsheaves are Determined by Stalks)
Let $\mathcal{C}$ be algebraic and $\mathcal{G}_1, \mathcal{G}_2$ be subsheaves of a sheaf $\mathcal{G}$ (4.2.30). Then $\mathcal{G}_1(U) = \mathcal{G}_2(U)$, as subsets of $\mathcal{G}(U)$, for every open $U \subset X$, if and only if $(\mathcal{G}_1)_x = (\mathcal{G}_2)_x$, as subsets of $\mathcal{G}_x$ (via the canonical, injective (4.2.28) stalk maps of the inclusions), for every $x \in X$.

Proof. Drafted by Claude Sonnet, reviewed by David Rufino.

($\Longrightarrow$) Immediate: if $\mathcal{G}_1(U) = \mathcal{G}_2(U)$ for every U, then by (4.2.6) every element of $(\mathcal{G}_1)_x$ is σ_x for some $\sigma \in \mathcal{G}_1(U) = \mathcal{G}_2(U)$, $U \ni x$, hence lies in $(\mathcal{G}_2)_x$, and symmetrically.

($\Longleftarrow$) Suppose $(\mathcal{G}_1)_x = (\mathcal{G}_2)_x$ for every $x \in X$. Fix $U \subset X$ and $\sigma \in \mathcal{G}_1(U) \subseteq \mathcal{G}(U)$; we show $\sigma \in \mathcal{G}_2(U)$ (the reverse inclusion, and hence $\mathcal{G}_1(U) = \mathcal{G}_2(U)$, then follows by symmetry).

For $x \in U$, $\sigma_x \in (\mathcal{G}_1)_x = (\mathcal{G}_2)_x$, so by (4.2.6) $\sigma_x = \tau_x$ for some $\tau \in \mathcal{G}_2(V_x) \subseteq \mathcal{G}(V_x)$, $x \in V_x$; by (4.2.8).a) (applied in $\mathcal{G}$, as σ, τ agree at the stalk x) there is $W_x \subseteq U \cap V_x$, $x \in W_x$, with $\sigma|_{W_x} = \tau|_{W_x}$ in $\mathcal{G}(W_x)$; as $\mathcal{G}_2$ is a subpresheaf, $\tau|_{W_x} \in \mathcal{G}_2(W_x)$, so $\sigma|_{W_x} \in \mathcal{G}_2(W_x)$.

© David Rufino CC BY-NC-ND 4.0

So $\{W_x\}_{x\in U}$ is an open cover of U with $\sigma|_{W_x}\in\mathcal{G}_2(W_x)$ for every x; these restrictions agree pairwise on overlaps (all restricting from the common σ), so as $\mathcal{G}_2$ is a sheaf, gluing gives $\rho\in\mathcal{G}_2(U)$ with $\rho|_{W_x}=\sigma|_{W_x}$ for every x. Regarding ρ as an element of $\mathcal{G}(U)$ (via the inclusion $\mathcal{G}_2\hookrightarrow\mathcal{G}$), ρ and σ agree on the cover $\{W_x\}$ of U, so by separation of $\mathcal{G}$ (4.2.4), $\rho=\sigma$; hence $\sigma\in\mathcal{G}_2(U)$. ∎

Lemma 4.2.46 (Kernels and Stalks Commute)
Let $\mathcal{C}$ be pointed and algebraic, $\beta:\mathcal{G}\to\mathcal{H}$ a morphism of sheaves, and $\ker(\beta)(U)=\{\sigma\in\mathcal{G}(U)\mid\beta_U(\sigma)=0\}$ its kernel (4.2.25), regarded as a subsheaf of $\mathcal{G}$. Then for every $x\in X$, the canonical stalk inclusion identifies $\ker(\beta)_x$ with the subset $\ker(\beta_x):=\{\xi\in\mathcal{G}_x\mid\beta_x(\xi)=0\}$ of $\mathcal{G}_x$.

Proof. Drafted by Claude Sonnet, reviewed by David Rufino.

Write $i:\ker(\beta)\hookrightarrow\mathcal{G}$ for the inclusion. By (2.6.151) and (2.6.136), i is monic, so by (4.2.28).d) (as $\mathcal{C}$ is algebraic and $\ker(\beta),\mathcal{G}$ are sheaves), i_x is injective for every $x\in X$.

Since $\beta\circ i=0$ (2.6.151), stalk-functoriality (4.2.7) gives $\beta_x\circ i_x=(\beta\circ i)_x=0_x=0$ (the stalk of the zero presheaf, valued 0 at every open with identity restrictions, is 0, so the stalk functor sends the zero morphism to the zero morphism). So $i_x(\eta)\in\ker(\beta_x)$ for every $\eta\in\ker(\beta)_x$.

Conversely let $\xi\in\ker(\beta_x)$, i.e. $\xi\in\mathcal{G}_x$ with $\beta_x(\xi)=0$. By (4.2.6), $\xi=\sigma_x$ for some $\sigma\in\mathcal{G}(U)$, $U\in\mathcal{B}$, $x\in U$. Then $(\beta_U(\sigma))_x=\beta_x(\sigma_x)=\beta_x(\xi)=0=0_x$, so by (4.2.8).a) there is $W\in\mathcal{B}$, $W\subseteq U$, $x\in W$, with $\beta_U(\sigma)|_W=0|_W$, that is $\beta_W(\sigma|_W)=0$, so $\sigma|_W\in\ker(\beta)(W)$. Then $i_x\big((\sigma|_W)_x\big)=(i_W(\sigma|_W))_x=(\sigma|_W)_x=\sigma_x=\xi$, so $\xi\in\mathrm{Im}(i_x)$.

So i_x is injective with image exactly $\ker(\beta_x)$, that is, identifies $\ker(\beta)_x$ with this subset of $\mathcal{G}_x$. ∎

Lemma 4.2.47 (Images and Stalks Commute)
Let $\mathcal{C}$ be algebraic, $\alpha:\mathcal{F}\to\mathcal{G}$ a morphism of sheaves, and $\alpha=m\circ e$ its image factorisation (4.2.40). Then for every $x\in X$, the canonical stalk inclusion identifies $\mathrm{Im}(\alpha)_x$ with the subset $\mathrm{Im}(\alpha_x):=\alpha_x(\mathcal{F}_x)$ of $\mathcal{G}_x$.

Proof. Drafted by Claude Sonnet, reviewed by David Rufino.

Retain the notation from the proof of (4.2.42): writing $e=\eta\circ e_\alpha$ with $\big(\mathrm{Im}^{\mathrm{psh}}(\alpha),e_\alpha,m_\alpha\big)$ the image of α in $\mathbf{PSh}(X;\mathcal{C})$ (4.2.39) and η the sheafification unit exhibiting $\mathrm{Im}(\alpha)=\big(\mathrm{Im}^{\mathrm{psh}}(\alpha)\big)^+$, it is shown there that $(e_\alpha)_x$ is surjective (each e_U being regular, hence surjective, (2.6.196)) and η_x is an isomorphism (4.2.18), so $e_x=\eta_x\circ(e_\alpha)_x$ is surjective. Also m_x is injective by (4.2.28).d), as m is a monomorphism of sheaves.

Since $m_x\circ e_x=\alpha_x$ and e_x is surjective onto $\mathrm{Im}(\alpha)_x$,

$$m_x(\mathrm{Im}(\alpha)_x)=m_x(e_x(\mathcal{F}_x))=\alpha_x(\mathcal{F}_x)=\mathrm{Im}(\alpha_x)$$

so, m_x being injective, it identifies $\mathrm{Im}(\alpha)_x$ with the subset $\mathrm{Im}(\alpha_x)$ of $\mathcal{G}_x$. ∎

Proposition 4.2.48 (Stalk Criterion for Exactness)
Let $\mathcal{C}$ be pointed and algebraic and $\mathcal{F}\xrightarrow{\alpha}\mathcal{G}\xrightarrow{\beta}\mathcal{H}$ a pair of composable morphisms of sheaves. Then this pair is exact at $\mathcal{G}$ (4.2.43) if and only if, for every $x\in X$, $\mathrm{Im}(\alpha_x)=\ker(\beta_x)$, as subsets of $\mathcal{G}_x$, that is, the stalk pair $\mathcal{F}_x\xrightarrow{\alpha_x}\mathcal{G}_x\xrightarrow{\beta_x}\mathcal{H}_x$ is exact at $\mathcal{G}_x$ in $\mathcal{C}$ (in the sense of (4.2.43), applied verbatim in $\mathcal{C}$).

Proof. Drafted by Claude Sonnet, reviewed by David Rufino.

By (4.2.44), the pair is exact at $\mathcal{G}$ if and only if $\mathrm{Im}(\alpha)(U)=\ker(\beta)(U)$ for every $U\subset X$ (using the canonical constructions of $\mathrm{Im}(\alpha),\ker(\beta)$). By (4.2.45) applied to these subsheaves of $\mathcal{G}$, this holds if and only if $\mathrm{Im}(\alpha)_x=\ker(\beta)_x$, as subsets of $\mathcal{G}_x$, for every $x\in X$. By (4.2.47) and (4.2.46), this is exactly the condition $\mathrm{Im}(\alpha_x)=\ker(\beta_x)$, as subsets of $\mathcal{G}_x$, for every $x\in X$. ∎

4.2.12 Constant Sheaves

Definition 4.2.49
*Let $A\in\mathcal{C}$ and X a topological space. Define the **constant presheaf** as follows*

$$\mathcal{A}^{\mathrm{pre}}(U):=A$$

with restriction maps the identity.

Proposition 4.2.50
Let X be an irreducible topological space then the constant presheaf $\mathcal{A}^{\mathrm{pre}}(U)=A$ is a sheaf, which we denote by $\underline{A}$.

© David Rufino CC BY-NC-ND 4.0

Proof. Suppose that $U = \bigcup_{i \in I} U_i$ where U_i are non-empty, then as U is irreducible (4.1.70), by definition every $U_i \cap U_j \neq \emptyset$. Therefore for sections on U_i are only compatible if they are equal, and so there is an obvious extension to U. ∎

Proposition 4.2.51 (Quotient of Constant Sheaf)
Let $\mathcal{F}$ be a subsheaf of the constant sheaf $\underline{A}$, define the quotient presheaf as follows

$$(\underline{A}/\mathcal{F})(U) \coloneqq \left\{ \sigma : U \to \coprod_{x \in U} A/\mathcal{F}_x \;\middle|\; \sigma(x) \in A/\mathcal{F}_x \right\}$$

© David Rufino CC BY-NC-ND 4.0