# Algebra, Geometry and Number Theory



David Rufino

# Contents


1 Introduction ... 8
2 Foundations ... 10
2.1 Set Theory ... 11
2.1.1 Relations ... 11
2.1.2 Functions ... 12
2.1.3 Partial Orders ... 12
2.1.4 Lattices ... 13
2.1.5 Distributive Lattice ... 17
2.1.6 Galois Connections ... 18
2.1.7 Axiom of Choice ... 20
2.1.8 Chain Conditions ... 21
2.2 Numbers ... 22
2.2.1 Integers ... 22
2.2.2 Arithmetic ... 23
2.2.3 Prime Factorization ... 25
2.3 Matroids ... 26
2.4 Decomposition in Noetherian and Distributive Lattices ... 29
2.5 Krull Dimension ... 31
2.6 Category Theory ... 35
2.6.1 Categories ... 35
2.6.2 Comma and Slice Categories ... 37
2.6.3 Product Categories and Bifunctors ... 37
2.6.4 Equivalence of categories ... 39
2.6.5 Properties of Morphisms ... 40
2.6.6 Pointed Categories ... 43
2.6.7 Yoneda Lemma ... 44
2.6.8 Representable Functors ... 45
2.6.9 Adjoint Functors ... 48
2.6.10 Limits ... 51
2.6.11 Colimit ... 53
2.6.12 Functorial Limits ... 55
2.6.13 Biproduct ... 59
2.6.14 Biproduct Additivity ... 62
2.6.15 Direct Limit / Directed Colimit ... 64
2.6.16 Inverse Limit ... 65
2.6.17 Equaliser and Coequaliser ... 68
2.6.18 Pullback and Kernel Pairs ... 70
2.6.19 Kernels and Cokernels ... 71
2.6.20 Normal, Regular and Extremal Morphisms ... 72
2.6.21 Images and Coimages ... 75
2.6.22 Epi-Mono Factorisations ... 76
2.6.23 Algebraic Categories ... 79
2.6.24 Exact Sequences ... 86
2.7 Abelian Categories ... 87
2.7.1 Images, Coimages and the First Isomorphism Theorem ... 88
2.7.2 Exact Sequences ... 90
2.7.3 Biproducts ... 91
3 Algebra ... 94
3.1 Magmas and Monoids ... 95
3.2 Groups ... 97
3.2.1 Cyclic Groups ... 99
3.2.2 Group Actions ... 101
3.2.3 Symmetric Group ... 102
3.2.4 Shuffle Permutations ... 103
3.2.5 Totally Ordered Abelian Group ... 104

3.2.6 Sums in an Abelian Group ... 104
3.3 Rings and Modules ... 108
3.3.1 Commutative Rings ... 108
3.3.2 Sums in a Ring ... 110
3.3.3 Modules I ... 110
3.3.4 Operations on Ideals ... 112
3.3.5 Quotient Rings ... 116
3.3.6 Chinese Remainder Theorem ... 118
3.3.7 Irreducible and Reduced rings ... 118
3.3.8 Algebra over a Commutative Ring ... 119
3.3.9 Finite-type Algebras ... 120
3.3.10 Bimodules ... 120
3.3.11 Module Direct Product and Sum ... 122
3.3.12 Free Modules ... 123
3.3.13 Exact Sequences ... 124
3.3.14 Dual Module ... 125
3.3.15 Matrices ... 126
3.3.16 Vector Spaces ... 128
3.3.16.1 Dual Space ... 131
3.3.16.2 Bilinear Pairings ... 132
3.3.17 Inverse Limits ... 134
3.4 Tensor Products ... 136
3.4.1 Commutative Tensor Product ... 136
3.4.2 Bimodule Tensor Product ... 138
3.4.3 Extensions of Scalars ... 140
3.4.4 Tensor Product Commutes with Direct Sum ... 142
3.4.5 Tensor Product Exact Sequences ... 144
3.4.6 Vector Space Tensor Product ... 145
3.4.7 Algebra Tensor Product ... 145
3.5 Multilinear Algebra ... 149
3.5.1 Multilinear Maps and Determinants ... 149
3.5.2 Trace Map ... 154
3.5.3 Block Matrices ... 156
3.5.4 Exterior Product ... 158
3.5.5 Matrix Rank ... 159
3.6 Localisation ... 161
3.6.1 Rings ... 161
3.6.2 Modules ... 163
3.6.3 Ideals ... 164
3.6.4 Change of Rings ... 165
3.6.5 Algebras ... 168
3.6.6 Localisation at an element ... 170
3.6.7 Localisation at a prime ideal ... 171
3.6.8 Finiteness Results ... 173
3.7 Monoid Ring ... 174
3.8 Polynomial Rings in One Variable ... 175
3.9 Laurent Polynomials ... 179
3.10 Formal Power Series and Laurent Series ... 180
3.11 Polynomial Rings in Many Variables ... 186
3.12 $\Delta$-Graded Rings ... 188
3.13 Graded Rings ... 193
3.14 Filtered Rings and Completions ... 195
3.14.1 Hensel's Lemma ... 204
3.15 Chain Conditions ... 206
3.16 Principal Ideal Domains ... 208
3.17 Factorisation ... 210
3.17.1 Polynomials over a Field ... 214

3.17.2 Polynomial Ring is a UFD ... 216
3.18 Cayley-Hamilton Theorem ... 218
3.18.1 The $A[X]$-module associated to an endomorphism ... 219
3.19 Fields and Galois Theory ... 221
3.19.1 Prime Fields ... 221
3.19.2 Separability ... 221
3.19.3 Field Extensions ... 222
3.19.4 Algebraic Extensions ... 225
3.19.5 Galois Theory Summary ... 228
3.19.6 Splitting Fields and Algebraic Closure ... 229
3.19.7 Normal Extensions ... 230
3.19.8 Separability (Algebraic Case) ... 232
3.19.9 Purely Inseparable Extensions and Separable Closure ... 234
3.19.10 Separable closure ... 236
3.19.11 Perfect Fields ... 237
3.19.12 Applications of Separability ... 238
3.19.13 Normal Extensions II ... 238
3.19.14 Finite Fields ... 239
3.19.15 Galois Theory ... 240
3.19.16 Norm and Trace ... 242
3.19.17 Transcendental Field Extensions ... 246
3.19.18 Separating Transcendence Base ... 248
3.20 Local Rings ... 251
3.21 Modules over Local Rings (Nakayama's Lemma) ... 252
3.22 Krull Dimension ... 254
3.23 Lying over, Incomparability, Going Up and Going Down ... 257
3.24 Integral Ring Extensions ... 259
3.24.1 Integral Closure in Finite Extensions ... 264
3.25 Valuation Rings and Places ... 265
3.26 Derivations ... 270
3.27 Hauptidealsatz ... 274
3.28 Regular Local Rings ... 277
3.29 Local Rings of Dimension 0 ... 278
3.30 Fractional Ideals ... 279
3.31 Discrete Valuation Rings ... 281
3.32 Dedekind Domains ... 284
3.32.1 Local Monogenicity ... 289
3.33 Extensions of Power Series Rings ... 290
3.34 Complete Discrete Valuation Rings ... 293
3.35 Affine Algebras ... 303
3.35.1 Reduction ... 303
3.35.2 Normalisation ... 303
3.35.3 Nullstellensatz ... 307
3.35.4 Morphisms of Affine Algebras ... 309
3.35.5 Krull Dimension of Affine Algebras ... 309
3.35.5.1 * Biequidimensionality by Strong Normalisation * ... 312
3.35.6 Derivations of Affine Algebras ... 313
3.35.7 Linearly Disjoint Algebras ... 317
3.35.8 Finiteness of Integral Closure ... 318
3.36 Affine Algebras under Base Change ... 320
3.36.1 Etale Algebras ... 321
3.36.2 Geometrically Reduced Algebras ... 324
3.36.3 Geometrically Irreducible Algebras ... 326
3.36.4 Geometrically Integral Algebras ... 330
3.37 Jacobson Rings ... 334
4 Topology and Sheaves ... 335
4.1 Topological Spaces ... 336

4.1.1 Axioms of Countability ... 337
4.1.2 Closure ... 337
4.1.3 Continuous Maps ... 338
4.1.4 Quasi-Homeomorphism ... 339
4.1.5 Kolmogorov Spaces ... 340
4.1.6 Symmetric Spaces ... 341
4.1.7 Hausdorff Spaces ... 342
4.1.8 Convergent Sequences ... 342
4.1.9 Connected Topological Spaces ... 343
4.1.10 Irreducible Topological Spaces ... 344
4.1.11 Noetherian Topological Spaces ... 346
4.1.12 Krull Dimension ... 346
4.1.13 Product Topology ... 349
4.1.14 Quasi-Compactness ... 350
4.1.15 Connectedness ... 353
4.1.16 Order Topology ... 353
4.2 Sheaves ... 354
4.2.1 Stalks ... 355
4.2.2 Sheafification ... 360
4.2.3 Inverse Image Functor ... 363
4.2.4 Skyscraper Sheaf ... 363
4.2.5 Limit Sheaf ... 365
4.2.6 Equaliser Sheaf ... 366
4.2.7 Product Presheaf ... 367
4.2.8 Kernel Pair Sheaf ... 368
4.2.9 Coequaliser Sheaf ... 370
4.2.10 Image Sheaf ... 373
4.2.11 Exactness ... 376
4.2.12 Constant Sheaves ... 377
4.3 Space with Functions ... 379
4.3.1 Local Rings ... 379
4.3.2 Open Immersions ... 381
4.3.3 Closed Immersions ... 383
4.3.4 Locally Closed Subspace ... 384
4.3.5 Glueing ... 384
5 Analysis ... 386
5.1 Real Numbers ... 387
5.1.1 Sequences in an Ordered Field ... 388
5.1.2 Sequential Completeness ... 390
5.1.3 Uniqueness of Reals ... 391
5.1.4 Existence of Reals ... 391
5.1.5 $n$-th Root ... 391
5.1.6 Limsup and Liminf ... 393
5.2 Complex Numbers ... 395
5.3 Metric Spaces ... 397
5.3.1 Completeness ... 398
5.3.2 Compactness ... 398
5.3.3 Uniform Continuity ... 399
5.4 Normed Vector Spaces ... 400
5.4.1 Continuous Functions ... 401
5.4.2 Product Space ... 401
5.4.3 Convergent Sequences ... 402
5.4.4 Finite-Dimensional Normed Spaces ... 403
5.4.5 Convergent Series ... 404
5.4.6 Function Spaces ... 406
5.4.7 Continuous Linear Maps ... 407
5.5 Differentiable Functions ... 409

5.6 Power Series ..... 410
5.7 Real Analysis ..... 412
5.7.1 Closed Intervals ..... 412
5.7.2 Power Function ..... 412
5.7.3 Countability ..... 412
5.7.4 Bolzano-Weierstrass Theorem ..... 412
5.7.5 Boundedness Theorem ..... 414
5.7.6 Intermediate Value Theorem ..... 414
5.7.7 Mean-Value Theorem ..... 415
5.8 Integration ..... 416
5.8.1 Algebras of Sets ..... 416
5.8.2 Set Functions ..... 417
5.8.3 Integration of Regulated Functions ..... 419
5.8.4 Measure Spaces ..... 421
5.8.5 Borel Measure on $\mathbb{R}$ ..... 425
5.8.6 Product Measure ..... 427
5.8.7 Borel Measure on $\mathbb{R}^n$ ..... 428
5.8.8 Measurable Functions ..... 429
5.8.9 Integration over a Measure ..... 431
5.9 Differential Calculus on Banach Spaces ..... 432
5.9.1 Total Derivatives ..... 432
5.9.2 Partial Derivatives ..... 433
5.9.3 Second Derivative ..... 435
5.9.4 Differentiation under Integral Sign ..... 436
5.9.5 Taylor's Theorem ..... 437
5.9.6 Differential Forms on Banach Spaces ..... 438
5.10 Complex Analysis ..... 440
5.10.1 Cauchy-Riemann Equations ..... 440
5.10.2 Holomorphic Functions ..... 441
5.10.3 Exponential and Trigonometric Functions ..... 441
5.10.4 Polar Coordinates of a Path ..... 444
5.10.5 Path Integrals ..... 445
5.11 Riemann Surfaces ..... 447
5.11.1 Morphisms of Riemann Surfaces ..... 448
5.11.2 Tangent and Cotangent Space ..... 449
5.11.3 Holomorphic 1-forms ..... 450
6 Algebraic Geometry ..... 452
6.1 Affine Varieties ..... 453
6.1.1 Topological Properties ..... 456
6.1.2 Structure Sheaf ..... 458
6.1.3 Regular Morphisms of Affine Varieties ..... 460
6.1.4 Dominant Morphisms ..... 463
6.1.5 Dimension ..... 463
6.1.6 Local Rings ..... 465
6.1.7 Rational Points ..... 467
6.1.8 Generic Points ..... 467
6.1.9 Tangent Space and Non-Singular Points ..... 467
6.1.10 Zeta Function over Finite Fields ..... 471
6.1.11 Base Change ..... 472
6.1.12 Valuation Rings on the Function Field ..... 474
6.1.13 Products of Affine Varieties ..... 474
6.1.14 Affine Curves ..... 477
6.2 Abstract Varieties ..... 478
6.2.1 Topological Properties ..... 478
6.2.2 Local Ring ..... 479
6.2.3 Affine Varieties ..... 481
6.2.4 Lifting Stalk Maps ..... 484

6.2.5 Open Subvarieties ... 486
6.2.6 Closed Subvarieties ... 487
6.2.7 Locally Closed Subvariety ... 488
6.2.8 Rational Points ... 488
6.2.9 Dimension ... 489
6.2.10 Galois Action ... 490
6.2.11 Product Variety ... 491
6.2.12 Separated Varieties ... 492
6.2.13 Rational Maps ... 494
6.2.14 Tangent Space ... 495
6.2.15 Completeness ... 496
6.2.16 Projective Space ... 499
6.2.17 Affine and Finite Morphisms ... 501
6.2.18 Normal Varieties ... 501
6.2.19 Weil Divisors ... 502
6.2.20 Algebraic Function Fields ... 503
6.2.21 Algebraic Curves ... 506
6.3 Function Fields and the Riemann-Roch Theorem ... 511
6.3.1 Relation to the Literature ... 511
6.3.2 Riemann-Roch Theorem ... 512
6.3.3 Meromorphic Differentials in the case $k = \mathbb{C}$ ... 517
6.3.4 Derivations on the Function Field ... 518
6.3.5 Residue Symbol at a Rational Place ... 518
6.3.6 The Algebraic Residue Theorem over an Algebraically Closed Field ... 522
6.3.7 Base Extension of the Constant Field ... 526
6.3.8 Residue Symbol at a General Place ... 529
6.3.9 The Algebraic Residue Theorem ... 531
6.3.10 Differentials are Weil Differentials, for General $k$ ... 531
6.3.11 Tate's Approach to Residues: A Sketch ... 532
6.3.12 Sheaf-Theoretic Interpretation of Repartitions ... 533
6.4 Projective Varieties ... 536
6.4.1 Projective Algebraic Sets ... 536
6.4.2 Affine Charts ... 539
6.4.3 Galois Orbits ... 541
6.4.4 Projective Varieties are Abstract Varieties ... 542
6.4.5 Regular Morphisms ... 543
6.4.6 Local Ring ... 545
6.4.7 Dimension ... 546
6.4.8 Valuative Completeness ... 547
6.5 Arithmetic Varieties ... 549
6.6 Scheme Notes ... 550
6.6.1 Schemes over $\mathbb{Z}$ ... 550
6.6.2 Closed Points vs Maximal Ideals ... 550

## Chapter 1

# Introduction

The main purpose of these notes is to provide a detailed exposition of Abstract Algebra, Galois Theory and Algebraic Varieties over non-algebraically closed fields, with particular interest in the Weil Conjectures.

**A note on AI assistance** This book has been in development since 2020. Since sometime in 2026 I have used Claude Sonnet 5 to assist with editing, checking, and writing exposition. The overall architecture and goals of the book remain the same and are my own. Where a proof itself was drafted by Claude rather than by me, this is disclosed explicitly at the start of that proof. I generally intend to replace these Claude-drafted proofs with my own, more idiomatic (and less pedantic) versions as time allows.

This pulls together many sources (standard or otherwise) in order to create a self-contained presentation with almost no prerequisites. At various points I have had to interpolate between an abstract, conceptual approach and a more concrete, down-to-earth approach. Many choices have been towards a more concrete approach (for example, focus on algebraic varieties rather than schemes, concrete construction of sheafification and a more explicit form of Krull's Hauptidealsatz) where I feel the abstract approach is not particularly more enlightening and can anyway be easily derived from the more concrete approach.

An interesting example is the theory of Algebraic Function Fields of One Variable / Regular Algebraic Curves, where I have interpolated between several classical treatments while departing from all of them in a number of respects – not yet ported to this edition.

On the other hand, various approaches to simpler matters have taken a more conceptual approach (for example, Krull Dimension, Dimension Theory of Field Extensions, Multilinear Algebra, Theory of Riemann Surfaces, extensive use of category-theoretic framing of various constructions such as tensor products).

For the section on Algebraic Geometry I've tried to interpolate a little between a more elementary approach (e.g. [Ser55], [Mil17]) and a scheme-theoretic approach [Stacks] [Gro64]. The idea is we may work over non-algebraically closed fields without the full (tedious) machinery of schemes. In these notes the proofs are technically simpler than the corresponding scheme results but also convey the key ideas used in the fully general case.

Similarly to [Stacks], I have made extensive use of cross-referencing so that logical chains of reasoning can be more easily followed. This naturally means many of the proofs are not as self-contained as might otherwise be the case. On the other hand we do obtain some "economy of scale", for example

- Commutative Algebra is the base for Galois Theory, Algebraic Number Theory and Algebraic Varieties
- Matroid theory may be used for both vector spaces and finitely generated field extensions
- "Krull Dimension" may be abstracted to work for both Rings and Topological Spaces / Varieties
- Dedekind Domain results may be applied for both Algebraic Number Fields and Regular Algebraic Curves
- Theory of Representable and Adjoint Functors may be applied fruitfully to many cases, to for example tensor products and sheaves

I also list a non-exhaustive set of references I found useful.

**Set Theory, Lattices, Category Theory**

- On the Structure of Abstract Algebras [Bir35]
- Lattice Theory [Bir40]
- A Course in Universal Algebra [BS81]
- Algebraic Theories [ARV10]
- Abelian Categories [Fre64]
- Handbook of Categorical Algebra 2: Categories and Structures [Bor94]
- Categories for the Working Mathematician [Lan98]
- Algebra: Volume I [Wae91]

**Algebra**

- Algebra [Lan11]

© David Rufino CC BY-NC-ND 4.0

- Introduction to Commutative Algebra [AM69]
- Commutative Algebra [Mat70]
- Commutative Rings [Kap74]
- Steps in Commutative Algebra [Sha00]
- Commutative Algebra I [ZS75]
- Algebra I: Chapters 1–3 [Bou98a]
- Algebra: Chapters 4–7 [Bou89]
- Algebraic Number Theory [Neu99]
- Local Fields [Ser79]
- Residues of Differentials on Curves [Tat68]
- Stacks Project [Stacks]

**Algebraic Geometry**

- Algebraic Geometry [Har13]
- Algebraic Geometry [Mil17]
- Algebraic Geometry and Arithmetic Curves [Liu06]
- Algebraic Curves [Ful08]
- The Red Book of Varieties and Schemes [Mum99]
- Faisceaux algébriques cohérents [Ser55]
- Éléments de géométrie algébrique IV [Gro64]

**Algebraic Function Fields**

- Introduction to the Theory of Algebraic Functions of One Variable [Che51]
- Groupes algébriques et corps de classes [Ser59]
- Algebraic Function Fields and Codes [Sti09]
- Algebraic Geometry (Riemann surfaces and function fields) [Bum98]
- Introduction to Algebraic and Abelian Functions [Lan82]

**Topology, Analysis and Geometry**

- General Topology [Kel71]
- General Topology [Wil70]
- Commutative Algebra: Chapters 1–7 [Bou98b]
- Some remarks on biequidimensionality of topological spaces and Noetherian schemes [Hei17]
- Differential Calculus [Car71]
- Foundations of Differentiable Manifolds and Lie Groups [War13]
- Manifolds, Sheaves, and Cohomology [Wed16]
- Real and Functional Analysis [Lan12]

© David Rufino CC BY-NC-ND 4.0

## Chapter 2

# Foundations

© David Rufino CC BY-NC-ND 4.0

# 2.1 Set Theory

## 2.1.1 Relations

**Definition 2.1.1 (Binary Relation)**
*A **binary relation** (or just **relation**) $R$ on a pair of sets $(X, Y)$ is subset of the cartesian product $X \times Y$.*

*We write $xRy$ to mean precisely $(x, y) \in R$.*

**Definition 2.1.2 (Converse Relation)**
*Let $R$ be a binary relation on $(X, Y)$ then we the converse relation $R^T$ on $(Y, X)$ given by*

$$yR^T x \iff xRy$$

**Definition 2.1.3 (Domain and Range)**
*Let $R$ be a relation on $(X, Y)$. We define the **domain** of $R$ to be*

$$\text{dom}(R) \coloneqq \{x \in X \mid \exists y \in Y \text{ s.t. } xRy\}$$

*and the **range** of $R$*

$$\text{range}(R) \coloneqq \{y \in X \mid \exists x \in X \text{ s.t. } xRy\}$$

**Definition 2.1.4 (Equivalence Relation)**
*Let $R$ be a binary relation on $(X, X)$. It is said to be*

*a) **reflexive** if $xRx$ for all $x \in X$*
*b) **symmetric** if $xRy \implies yRx$ for all $x, y \in X$*
*c) **transitive** if $xRy \wedge yRz \implies xRz$ for all $x, y, z \in X$*

*A relation which satisfies all these properties is called an **equivalence relation** on $X$. In this case we would write*

$$x \sim y$$

*instead of $xRy$. For an element $x \in X$ denote the **equivalence class** of $x$ by*

$$[x]_R = \{y \mid xRy\}$$

*Note that $R^T = R$.*

**Definition 2.1.5 (Partition)**
*Let $X$ be a set an $\mathcal{F}$ a family of subsets of $X$. It is said to be a **partition** if*

*a) $X = \bigcup_{A \in \mathcal{F}} A$*
*b) $A, B \in \mathcal{F} \implies A = B$ or $A \cap B = \varnothing$*

**Proposition 2.1.6 (Equivalence Classes form a Partition)**
*Let $E$ be an equivalence relation on $X$. The family*

$$\mathcal{F} = \{[x]_E \mid x \in X\}$$

*forms a partition of $X$. Denote by $X/E$ the family of equivalence classes, called the **quotient** of $X$ with respect to $E$.*

*Proof.* It's clear that

$$X = \bigcup_{A \in \mathcal{F}} A$$

because by reflexive-ness $x \in [x]_E$ for all $x \in X$.

We claim that for any $z \in [x]_E$ we have $[z]_E = [x]_E$. Suppose $y \in [x]_E$ then $xRz$ and $xRy$. By symmetry and transitivity we then have $zRy$ which implies $y \in [z]_E$. In other words $[x]_E \subseteq [z]_E$. By symmetry of $R$ we have $x \in [z]_E$, so by the same token $[z]_E \subseteq [x]_E$, which shows they are equal.

Therefore it's clear that $[x]_E \cap [y]_E \neq \varnothing \implies [x]_E = [y]_E$ and thus $\mathcal{F}$ forms a partition. ∎

**Definition 2.1.7 (Composition of Relation)**
*Suppose $R$ is a relation on $(X, Y)$ and $S$ a relation on $(Y, Z)$. We define the composition $S \circ R$ on $(X, Z)$*

$$S \circ R = \{(x, z) \mid \exists y \in Y \text{ s.t. } xRy \text{ and } yRz\}$$

© David Rufino CC BY-NC-ND 4.0

### 2.1.2 Functions

**Definition 2.1.8 (Function)**
*A **function** $f : X \to Y$ consists of a binary relation $\Gamma(f)$ on $(X, Y)$ such that*

- $\mathrm{dom}(f) = X$
- $\Gamma(f)$ *is* ***single-valued*** *that is* $x\Gamma(f)y \wedge x\Gamma(f)y' \implies y = y'$

*Equivalently for all $x \in X$ there exists precisely one $y \in Y$ such that $x\Gamma(f)y$.*

*We write $f(x) = y$ for the unique element $y \in Y$ such that $x\Gamma(f)y$.*

**Proposition 2.1.9 (Equality of Functions)**
*Two functions $f, g : X \to Y$ are equal if and only if $f(x) = g(x)$ for all $x \in X$.*

**Proposition 2.1.10 (Composition of Functions)**
*Let $f : X \to Y$ and $g : Y \to Z$ be functions then the composition $\Gamma(g) \circ \Gamma(f)$ is still a function, which we write $g \circ f$, and*

$$(g \circ f)(x) = g(f(x))$$

*Furthermore composition is **associative** in the sense that*

$$(h \circ g) \circ f = h \circ (g \circ f)$$

**Definition 2.1.11 (Injective, Surjective and Bijective)**
*Let $f : X \to Y$ be a function then we say*

- *$f$ is **injective** if $f(x) = f(x') \implies x = x'$*
- *$f$ is **surjective** if for all $y \in Y$ there exists $x$ such that $f(x) = y$*
- *$f$ is **bijective** if it is both injective and surjective*

**Definition 2.1.12 (Inverse Function)**
*Let $f : X \to Y$ and $g : Y \to X$ be functions. We say*

- *$g$ is a **left inverse** for $f$ if $g \circ f = 1_X$*
- *$g$ is a **right inverse** for $f$ if $f \circ g = 1_Y$*
- *$g$ is a **two-sided inverse** for $f$ if it is both a left and right inverse*

**Proposition 2.1.13**
*Let $f : X \to Y$ be a function then*

- *$f$ is **injective** if and only if it has a **left inverse***
- *$f$ is **surjective** if and only if it has a **right inverse***
- *$f$ is **bijective** if and only if it has a **two-sided inverse***

**Definition 2.1.14 (Idempotent Function)**
*A function $p : X \to X$ is **idempotent** $p \circ p = p$.*

**Lemma 2.1.15 (Idempotent Criterion)**
*Let $p : X \to X$ be a function. Then $\mathrm{Fix}(p) \subseteq \mathrm{Im}(p)$ and these are equal if and only if $p$ is idempotent.*

### 2.1.3 Partial Orders

**Definition 2.1.16 (Preorder)**
*A binary relation $\leq$ on $(X, X)$ is a **preorder** if*

- ***reflexivity** $x \leq x$*
- ***transitivity** $x \leq y$ and $y \leq z \implies x \leq z$*

*We may refer to $(X, \leq)$ as a **preordered set** or **preorder**.*

**Definition 2.1.17 (Poset)**
*A binary relation $\leq$ on $(X, X)$ is a **partial order** if*

- ***reflexivity** $x \leq x$*

© David Rufino CC BY-NC-ND 4.0

- ***antisymmetry*** $x \leq y$ and $y \leq x \implies y = x$
- ***transitivity*** $x \leq y$ and $y \leq z \implies x \leq z$

*We may refer to $(X, \leq)$ as a **partially ordered set** or **poset**. Equivalently, a poset is a preorder which is additionally antisymmetric.*

**Definition 2.1.18 (Dual Poset)**
*Given a poset $(X, \leq)$ denote the set $X$ with the converse relation by $\left(X, \overset{d}{\leq}\right)$. This is the **dual poset** to $(X, \leq)$.*

**Example 2.1.19**
*Let $\mathcal{F}$ be a family of subsets of a fixed set $E$. Then $(\mathcal{F}, \subseteq)$ is a poset ordered under inclusion.*

**Definition 2.1.20 (Top and Bottom)**
*Let $(X, \leq)$ we say $\top$ (resp. $\bot$) is a **top element** (resp. **bottom element**) if it is greater than (resp. less than) every element of $x$. In this case it is unique.*

**Definition 2.1.21 (Monotone/Antitone Function)**
*Let $(X, \leq)$ and $(Y, \leq)$ be posets. A function $f : X \to Y$ is*

- ***monotone / order-preserving*** *if $x \leq y \implies f(x) \leq f(y)$*
- ***antitone / order-reversing*** *if $x \leq y \implies f(y) \leq f(x)$*
- *a **monotone embedding** if $x \leq y \iff f(x) \leq f(y)$*
- *an **order isomorphism** if it is bijective and monotone*
- *a **dual isomorphism** if it is bijective and antitone*

**Proposition 2.1.22**
*Let $f : X \to Y$ be a monotone function. Then it is an embedding if and only if it is injective.*

In what follows the notion of closure and kernel operator will be important.

**Definition 2.1.23 (Closure operator)**
*Let $(X, \leq)$ be a partially ordered set. A function $c : X \to X$ is a **closure operator** if it is*

*a)* ***extensive*** $x \leq c(x)$
*b)* ***monotone*** $x \leq y \implies c(x) \leq c(y)$
*c)* ***idempotent*** $c(c(x)) = c(x)$

**Definition 2.1.24 (Kernel operator)**
*Let $(X, \leq)$ be a partially ordered set. A function $\kappa : X \to X$ is a **kernel operator** if it is*

- ***co-extensive*** $\kappa(x) \leq x$
- ***monotone*** $x \leq y \implies \kappa(x) \leq \kappa(y)$
- ***idempotent*** $\kappa(\kappa(x)) = \kappa(x)$

Note these definitions are "dual" with respect to the ordering on $X$.

### 2.1.4 Lattices

Certain families of subsets of algebraic structures (e.g. ideals, subgroups, normal subgroups, submodules) form a "sublattice" of the power set. Certain operations on, and results about, these subsets share common features regardless of the type of algebraic structure. Therefore we detail some elements of "Lattice Theory" (see Birkhoff) which may clarify the exposition.

**Definition 2.1.25 (Upper and Lower Bounds)**
*Let $(X, \leq)$ be a poset and $S \subseteq X$. Define the set of **upper bounds** for $S$ by*

$$S^{\uparrow} = \{x \in X \mid s \leq x \quad \forall s \in S\}$$

*and the set of **lower bounds** for $S$ by*

$$S^{\downarrow} = \{x \in X \mid x \leq s \quad \forall s \in S\}$$

*Note by convention $\varnothing^{\uparrow} = \varnothing^{\downarrow} = X$. Furthermore*

© David Rufino CC BY-NC-ND 4.0

$$X^{\uparrow} = \begin{cases} \{\top\} & X \text{ has a top element} \\ \varnothing & \text{otherwise} \end{cases}$$

*and*

$$X^{\downarrow} = \begin{cases} \{\bot\} & X \text{ has a bottom element} \\ \varnothing & \text{otherwise} \end{cases}$$

**Lemma 2.1.26 (Upper/Lower bounds are antitone maps)**
*Let $(X, \leq)$ be a poset and $S, T$ subsets of $X$ then*

- ***antitone*** $S \subseteq T \implies T^{\uparrow} \subseteq S^{\uparrow}$ *and* $T^{\downarrow} \subseteq S^{\downarrow}$
- ***unit-counit relations*** $S \subseteq S^{\uparrow\downarrow}$ *and* $T \subseteq T^{\downarrow\uparrow}$
- ***triangular identities*** $S^{\uparrow} = S^{\uparrow\downarrow\uparrow}$ *and* $T^{\downarrow} = T^{\downarrow\uparrow\downarrow}$

*Proof.* We prove only the first triangular identity as the others are straightforward consequences of the definitions. Firstly $S \subseteq S^{\uparrow\downarrow} \implies S^{\uparrow\downarrow\uparrow} \subseteq S^{\uparrow}$ by the antitone property. Given the relation $T \subseteq T^{\downarrow\uparrow}$ substitute $T = S^{\uparrow}$ to get the reverse inclusion. ■

**Lemma 2.1.27**
*Let $(X, \leq)$ be a poset and $S, T$ subsets of $X$. Then the intersections $S \cap S^{\uparrow}$ and $T \cap T^{\downarrow}$ contain at most one element. **When they exist** write the elements as $\top_S$ and $\bot_T$ respectively, and are referred to as the **maximum** and **minimum** elements respectively.*

*Proof.* Given $x, y \in S \cap S^{\uparrow}$ then by definition $x \leq y$ and $y \leq x$. By anti-symmetry we have $x = y$ as required. ■

**Definition 2.1.28 (Supremum and Infimum)**
*Let $(X, \leq)$ be a poset and $S \subseteq X$ a subset. We say a **supremum** of $S$ is the minimal upper bound, i.e. the unique element of*

$$S^{\uparrow} \cap S^{\uparrow\downarrow}$$

*when it exists and write this as $\sup S$. Similarly an **infimum** of $S$ is the maximal lower bound, i.e. the unique element of*

$$S^{\downarrow} \cap S^{\downarrow\uparrow}$$

*when it exists and write this as $\inf X$.*

**Lemma 2.1.29 (Maximum = Supremum)**
*Let $(X, \leq)$ be a poset and $S \subseteq X$ a subset. Then $\top_S$ exists if and only if $\sup S$ exists and is a member of $S$. In this case $\top_S = \sup S$.*

**Lemma 2.1.30**
*Let $(X, \leq)$ be a poset. Then $\{\sup S\}^{\uparrow} = S^{\uparrow}$ and $\{\inf T\}^{\downarrow} = T^{\downarrow}$ when these exist.*

**Lemma 2.1.31 (Sup is monotone and Inf is antitone)**
*Let $(X, \leq)$ be a poset and $S, T$ subsets of $X$. Then $S \subseteq T \implies \sup S \leq \sup T$ and $\inf T \leq \inf S$ when these exist.*

*Proof.* Note $S \subseteq T \implies T^{\uparrow} \subseteq S^{\uparrow}$ so $\sup T \in S^{\uparrow}$. By definition $\sup S \in S^{\uparrow\downarrow}$ therefore $\sup S \leq \sup T$.

Similarly $S \subseteq T \implies T^{\downarrow} \subseteq S^{\downarrow}$. By definition $\inf T \in T^{\downarrow} \implies \inf T \in S^{\downarrow}$. By definition $\inf S \in S^{\downarrow\uparrow}$ therefore $\inf T \leq \inf S$.
■

**Remark 2.1.32**
*Note that $\varnothing^{\uparrow} = X$ and therefore $\sup \varnothing = \bot$ when it exists. Similarly $\inf \varnothing = \top$ when it exists.*

*When $\top$ exists $\sup X = \top$, otherwise it is not defined. Similarly when $\bot$ exists $\inf X = \bot$, otherwise it is not defined.*

**Definition 2.1.33 (Lattice)**
*A poset $(X, \leq)$ is a **lattice** if every pair of elements $\{x, y\}$ admits both a supremum and infimum. In this case we write*

$$a \vee b := \sup\{a, b\}$$

*and*

$$a \wedge b := \inf\{a, b\}$$

© David Rufino CC BY-NC-ND 4.0

*These are called the **join** and **meet** operations. A subset $Y$ is called a **sub-lattice** if*

$$a, b \in Y \implies a \wedge b \in Y \text{ and } a \vee b \in Y.$$

*Similarly it is a **complete lattice** if every subset $S$ admits both a supremum and infimum. This is written*

$$\bigvee S := \sup S$$

*and*

$$\bigwedge S := \inf S$$

*Note a complete lattice has both a top and a bottom element (by considering $\sup \varnothing$ and $\inf \varnothing$), and a lattice admits **finite** joins and meets.*

*Trivially*

$$\bigwedge \{x\} = \bigvee \{x\} = x$$

**Example 2.1.34 (Power Set)**
*For a fixed set $E$ the collection of subsets $\mathcal{P}(E)$ is a complete lattice under the union and intersection operator **with the convention that empty intersection is the whole set and empty union is the empty set***

*In this case $\top = E$ and $\bot = \varnothing$.*

**Proposition 2.1.35 (Principal down-sets are lattices)**
*Let $(X, \leq)$ be a lattice and $x, y \in X$. Then the subsets $\{x\}^\uparrow$, $\{x\}^\downarrow$ and $\{x\}^\uparrow \cap \{y\}^\downarrow$ are sub-lattices.*

Verifying a poset is a lattice is slightly easier than it may first appear.

**Lemma 2.1.36 (Supremum is Infimum of upper bounds)**
*Let $(X, \leq)$ be a poset and $S$ a subset of $X$. Then*

$$\sup S = \inf S^\uparrow$$

*when either exists. Dually*

$$\inf S = \sup S^\downarrow$$

*Proof.* By definition $\sup S$ is the unique element of $S^\uparrow \cap S^{\uparrow\downarrow}$ and $\inf S^\uparrow$ is the unique element of $S^{\uparrow\downarrow} \cap S^{\uparrow\downarrow\uparrow}$. By (2.1.26) $S^{\uparrow\downarrow\uparrow} = S^\uparrow$ so they are equivalent. ∎

**Proposition 2.1.37 (Criteria to be a Complete Lattice)**
*Let $(X, \leq)$ be a poset. Then the following are equivalent*

*a) $X$ is a complete lattice*
*b) $X$ admits arbitrary infimums (and in particular has $\top = \inf \varnothing$)*
*c) $X$ admits arbitrary supremums (and in particular has $\bot = \sup \varnothing$)*

*In this case we have the relationships*

$$\bigvee S = \bigwedge S^\uparrow$$

$$\bigwedge S = \bigvee S^\downarrow$$

*Proof.* a) $\implies$ b,c) is clear.

b,c) $\implies$ a) follows from the previous Lemma. ∎

**Lemma 2.1.38**
*Let $(X, \leq)$ be a poset and $(Y, \leq)$ a sub-poset. Let $S \subseteq Y$ be a subset. Then $\inf_Y S$ exists if and only if $\inf_X S$ exists and belongs to $Y$. In this case they are equal.*

*Proof.* Note in general that $T^{\downarrow,Y} = T^{\downarrow,X} \cap Y$ and $T^{\uparrow,Y} = T^{\uparrow,X} \cap Y$. Therefore

$$S^{\downarrow,Y} \cap S^{\downarrow\uparrow,Y} = S^{\downarrow,X} \cap S^{\downarrow\uparrow,X} \cap Y$$

Recall $\inf S$ is the unique element of $S^\downarrow \cap S^{\downarrow\uparrow}$ if it exists. Then the result follows easily. ∎

© David Rufino CC BY-NC-ND 4.0

**Definition 2.1.39 (Moore Family)**
*Let $(X, \leq)$ be a complete lattice. A sub-poset $(Y, \leq)$ is a **Moore family** over $X$ if it satisfies the following property*

$$S \subseteq Y \implies \bigwedge_X S \in Y$$

*In particular this includes the case $S = \varnothing$ and so $\top \in Y$.*

**Example 2.1.40 (Moore family of sets)**
*Given a fixed set $E$, then $\mathcal{P}(E)$ is a complete lattice ordered under inclusion. Then a family of subsets $\mathcal{F}$ is a Moore family precisely when*

- $E \in \mathcal{F}$
- $A_{i\in I} \in \mathcal{F} \implies \bigcap_{i\in I} A_i \in \mathcal{F}$

**Proposition 2.1.41 (Equivalent Formulations of Complete Sub-lattice)**
*Let $(X, \leq)$ be a complete lattice and $(Y, \leq)$ a sub-poset. Then the following are equivalent*

*a) $(Y, \leq)$ is a Moore family*
*b) $(Y, \leq)$ is a complete lattice*
*c) $Y$ is the image of some closure operator $c : X \to X$*

*In this case the closure operator is given by*

$$c(x) = \bigwedge_X \{y \in Y \mid x \leq y\}$$

*For $S \subseteq Y$*

$$\bigwedge_Y S = \bigwedge_X S$$

$$\bigvee_Y S = c\left(\bigvee_X S\right)$$

*and for $S \subseteq X$ we have*

$$c\left(\bigvee_X S\right) = \bigwedge_X (S^\uparrow \cap Y)$$

*Proof.* a) $\implies$ b) By (2.1.38) $S \subseteq Y \implies \bigwedge_Y S = \bigwedge_X S$. By (2.1.37) then $Y$ is a complete lattice.

b) $\implies$ c) Suppose that $(Y, \leq)$ is a complete lattice then define the function $c : X \to X$ by $c(x) = \bigwedge_X \Gamma_x$ where $\Gamma_x = \{y \in Y \mid x \leq y\}$. We need to show that it is a closure operator. Evidently $x \in \Gamma_x^\downarrow$ and $c(x) \in \Gamma_x^{\downarrow\uparrow}$ by definition of infimum. Therefore $x \leq c(x)$ and $c$ is extensive. Note $x \leq y \implies \Gamma_y \subseteq \Gamma_x$. By (2.1.31) we have $\inf \Gamma_x \leq \inf \Gamma_y$, whence $c(x) \leq c(y)$ and $c$ is monotone.

$Y$ is a complete lattice, so by (2.1.38) we have $c(x) \in Y$ so that $\mathrm{Im}(c) \subseteq Y$. We claim that $x \in Y \implies c(x) = x$. In this case $x \in \Gamma_x$ and $c(x) \in \Gamma_x^\downarrow$ whence $c(x) \leq x$ and therefore $x = c(x)$ as required. Therefore $Y \subseteq \mathrm{Fix}(c) \subseteq \mathrm{Im}(c) \subseteq Y$, whence $Y = \mathrm{Im}(c) = \mathrm{Fix}(c)$ and $c$ is idempotent by (2.1.15). As $c$ is extensive, monotone and idempotent it is by definition a closure operator.

c) $\implies$ a) In order for $Y \coloneqq \mathrm{Im}(c)$ to be a Moore family, we need to show $S \subseteq Y \implies \bigwedge_X S \in Y$. We claim that by properties of $c$ we have

$$S \subseteq Y \implies c\big(S^\downarrow\big) \subseteq S^\downarrow$$

$$T \subseteq X \implies c(T^\uparrow) \subseteq T^\uparrow$$

Therefore $c$ maps the singleton set $S^\downarrow \cap S^{\downarrow\uparrow} = \left\{\bigwedge_X S\right\}$ to itself. In otherwords $\bigwedge_X S \in \mathrm{Fix}(c) = \mathrm{Im}(c) = Y$ as required.

Define $\Gamma_x \coloneqq \{y \in Y \mid x \leq y\}$. We wish to show that $c(x) = \bigwedge_X \Gamma_x$. As $x \leq c(x)$ we have $c(x) \in \Gamma_x$. Furthermore $y \in \Gamma_x \implies c(x) \leq c(y) = y$. So $c(x) \in \Gamma_x^\downarrow$. Therefore $c(x) = \bot_{\Gamma_x} = \bigwedge_X \Gamma_x$ as required.

Finally by (2.1.30) $\left\{\bigvee_X S\right\}^\uparrow = S^\uparrow$ for any $S \subseteq X$. Therefore, as $c(x) = \bigwedge_X \Gamma_x$ we find

© David Rufino CC BY-NC-ND 4.0

$$c\left(\bigvee_X S\right) = \bigwedge_X \left(\left\{\bigvee_X S\right\}^{\uparrow} \cap Y\right) = \bigwedge_X (S^{\uparrow} \cap Y)$$

as required. In particular when $S \subseteq Y$ we find by (2.1.41)

$$\bigvee_Y S = \bigwedge_Y S^{\uparrow, Y} = \bigwedge_X S^{\uparrow, Y} = \bigwedge_X (S^{\uparrow} \cap Y) = c\left(\bigvee_X S\right)$$

■

**Remark 2.1.42**
*For a given complete lattice $(X, \leq)$ we have established a correspondence between*

$$\{\text{closure operators } \ c : X \to X\} \longleftrightarrow \{\text{complete sub-lattices } \ (Y, \leq)\}$$

**Corollary 2.1.43 (Moore family admits a closure operator)**
*Let $E$ be a fixed set and $\mathcal{F}$ a Moore family over $(\mathcal{P}(E), \subseteq)$. Then there exists a surjective closure operator $c : \mathcal{P}(E) \to \mathcal{F}$ given by*

$$c(F) = \bigcap_{F \subseteq E_\alpha \in \mathcal{F}} E_\alpha$$

*Any such closure operator $c : \mathcal{P}(E) \to \mathcal{P}(E)$ gives rise to a Moore family $\mathcal{F} := \mathrm{Im}(c)$.*

**Proposition 2.1.44 (Alternative expression for join)**
*Let $(X, \leq)$ be a complete lattice and $c : X \to X$ a closure operator with image $Y$. Then for any subset $S \subset X$*

$$c\left(\bigvee_X S\right) = c\left(\bigvee_X c(S)\right) = \bigvee_Y c(S) = \bigwedge_X (S^{\uparrow} \cap Y)$$

*i.e. it's the smallest "closed" set containing each element of $S$.*

*Proof.* By (2.1.41) the expression for $c\left(\bigvee_X S\right)$ yields

$$c\left(\bigvee_X c(S)\right) = \bigwedge_X (c(S)^{\uparrow} \cap Y) = \bigwedge_X (S^{\uparrow} \cap Y) = c\left(\bigvee_X S\right)$$

where the middle equality follows because if $y \in Y$ then $c(s) \leq y \iff s \leq y$. Furthermore $c(S) \subseteq Y$ so the expression for $\bigvee_Y$ yields

$$c\left(\bigvee_X c(S)\right) = \bigvee_Y c(S).$$

as required. ■

## 2.1.5 Distributive Lattice

**Proposition 2.1.45**
*Let $(X, \leq)$ be a lattice then the following relations hold*

*a)* $x \wedge (y \vee z) \geq (x \wedge y) \vee (x \wedge z)$
*b)* $x \vee (y \wedge z) \leq (x \vee y) \wedge (x \vee z)$

*Proof.* a) By definition $x \wedge y \leq x$ and $x \wedge y \leq y \leq y \vee z$. Therefore $x \wedge y \leq x \wedge (y \vee z)$. By symmetry in $y$ and $z$ we have $x \wedge z \leq x \wedge (y \vee z)$. Whence $(x \wedge y) \vee (x \wedge z) \leq x \wedge (y \vee z)$ as required.

b) follows by duality ■

**Definition 2.1.46 (Distributive Lattice)**
*We say a lattice $(X, \leq)$ is **distributive** if it satisfies the following relations for all $x, y, z \in X$*

- $x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z)$
- $x \vee (y \wedge z) = (x \vee y) \wedge (x \vee z)$

© David Rufino CC BY-NC-ND 4.0

**Proposition 2.1.47**
*Let $(X, \leq)$ be a lattice then TFAE*

*a)* $(X, \leq)$ *is a distributive lattice*
*b)* $x \wedge (y \vee z) \leq (x \wedge y) \vee (x \wedge z)$
*c)* $x \vee (y \wedge z) \geq (x \vee y) \wedge (x \vee z)$

**Example 2.1.48**
*Any family of subsets closed under finite intersection and union is a distributive lattice.*

### 2.1.6 Galois Connections

**Definition 2.1.49 (Galois Connection)**
*Let $\left(X, \underset{X}{\leq}\right)$ and $\left(Y, \underset{Y}{\leq}\right)$ be posets. A pair of functions $(f_\star, f^\star)$*

$$X \underset{f^\star}{\overset{f_\star}{\rightleftharpoons}} Y$$

*is called an **antitone Galois connection** if it satisfies the **adjoint property***

- $x \underset{X}{\leq} f^\star(y) \iff y \underset{Y}{\leq} f_\star(x) \quad \forall x \in X, y \in Y$

*We say it is a **monotone Galois connection** if instead*

- $x \underset{X}{\leq} f^\star(y) \iff f_\star(x) \underset{Y}{\leq} y \quad \forall x \in X, y \in Y$

*We will assume that if not otherwise specified the connection is antitone.*

**Proposition 2.1.50 (Equivalent Condition for Galois Connection)**
*Let $\left(X, \underset{X}{\leq}\right)$ and $\left(Y, \underset{Y}{\leq}\right)$ be posets. Consider a pair of functions*

$$X \underset{f^\star}{\overset{f_\star}{\rightleftharpoons}} Y$$

*Then this constitutes an **antitone Galois Connection** if and only if*

- $f_\star$ *and* $f^\star$ *are both antitone*
- $x \underset{X}{\leq} f^\star(f_\star(x))$ *and* $y \underset{Y}{\leq} f_\star(f^\star(y))$ *for all* $x \in X, y \in Y$ *(i.e.* $f^\star \circ f_\star$ *and* $f_\star \circ f^\star$ *are extensive)*

*Similarly it constitutes a **monotone Galois Connection** if and only if*

- $f_\star$ *and* $f^\star$ *are both monotone*
- $x \underset{X}{\leq} f^\star(f_\star(x))$ *and* $f_\star(f^\star(y)) \underset{Y}{\leq} y$ *for all* $x \in X, y \in Y$

*Proof.* We consider only the antitone case, as the monotone follows from duality (flip $\underset{Y}{\leq}$).

Suppose that $(f_\star, f^\star)$ satisfies the adjoint property

$$f_\star(x) = f_\star(x) \implies f_\star(x) \underset{Y}{\leq} f_\star(x) \implies x \underset{X}{\leq} f^\star(f_\star(x))$$

$$f^\star(y) = f^\star(y) \implies f^\star(y) \underset{Y}{\leq} f^\star(y) \implies y \underset{Y}{\leq} f_\star(f^\star(y))$$

whence the extensive property follows. Furthermore

$$x \underset{X}{\leq} x' \implies x \underset{X}{\leq} f^\star(f_\star(x')) \implies f_\star(x') \underset{Y}{\leq} f_\star(x)$$

$$y \underset{Y}{\leq} y' \implies y \underset{Y}{\leq} f_\star(f^\star(y')) \implies f^\star(y) \underset{X}{\leq} f^\star(y')$$

which shows that the functions $f_\star$ and $f^\star$ are antitone.

Conversely suppose they satisfy the given conditions. Then by the antitone and extensive properties in turn

$$x \underset{X}{\leq} f^\star(y) \implies f_\star(f^\star(y)) \underset{Y}{\leq} f_\star(x) \implies y \underset{Y}{\leq} f_\star(x)$$

and

© David Rufino CC BY-NC-ND 4.0

$$y \underset{Y}{\leq} f_\star(x) \implies f^\star(f_\star(x)) \underset{X}{\leq} f^\star(y) \implies x \underset{X}{\leq} f^\star(y).$$

which is the adjoint property as required. ∎

**Definition 2.1.51 (Closed sets)**
*Let $(f_\star, f^\star)$ be a Galois connection. Then define the **closed sets** to be*

$$X^\star := f^\star(Y)$$

$$Y^\star := f_\star(X)$$

**Proposition 2.1.52 (Isomorphism on closed sets)**
*Consider an antitone (resp. monotone) Galois connection $X \underset{f^\star}{\overset{f_\star}{\rightleftarrows}} Y$. Then it restricts to a dual isomorphism (resp. order isomorphism) on **closed sets***

$$X^\star \underset{f^\star}{\overset{f_\star}{\rightleftarrows}} Y^\star$$

*Furthermore the following properties hold*

- $f_\star \circ f^\star \circ f_\star = f_\star$
- $f^\star \circ f_\star \circ f^\star = f^\star$
- *In the antitone case $f^\star \circ f_\star$ and $f_\star \circ f^\star$ are closure operators.*
- *In the monotone case $f^\star \circ f_\star$ is a closure operator and $f_\star \circ f^\star$ is a kernel operator.*
- $X^\star = \mathrm{Fix}(f^\star \circ f_\star) = \mathrm{Im}(f^\star \circ f_\star)$
- $Y^\star = \mathrm{Fix}(f_\star \circ f^\star) = \mathrm{Im}(f_\star \circ f^\star)$

*Proof.* We detail the antitone case as the monotone case follows by duality. We first prove so-called triangular identities, for by the extensive property (2.1.50)

$$x \leq f^\star(f_\star(x)) \implies f_\star(f^\star(f_\star(x))) \leq f_\star(x)$$

and by the other extensive property

$$f_\star(x) \leq f_\star(f^\star(f_\star(x)))$$

whence they are equal. The other case is similar.

It's immediate that $f^\star \circ f_\star$ and $f_\star \circ f^\star$ are idempotent, and they are extensive by (2.1.50). And the composition of two antitone functions is monotone so $f^\star \circ f_\star$ and $f_\star \circ f^\star$ are closure operators.

Observe

$$\mathrm{Im}(f^\star \circ f_\star) \subseteq \mathrm{Im}(f^\star) \subseteq \mathrm{Fix}(f^\star \circ f_\star)$$

where the first inclusion is trivial and the second inclusion follows from the second triangular identity. However both sides are equal by (2.1.15) and the expression for $X^\star$ follows. The expression for $Y^\star$ follows similarly.

This shows that the maps are mutual inverses as required. ∎

In certain circumstances we may consider a smaller subset of $X$, by applying a suitable closure operator which is compatible with the Galois correspondence:

**Proposition 2.1.53 (Subordinated Closure Operator)**
*Let $X \underset{f^\star}{\overset{f_\star}{\rightleftarrows}} Y$ be a Galois connection and $c : X \to X$ be a closure operator with image $X_c$. Then*

$$c(x) \leq (f^\star \circ f_\star)(x) \quad \forall x \in X \iff \mathrm{Im}(f^\star) \subseteq \mathrm{Fix}(c) \iff X^\star \subseteq X_c$$

*In this case*

$$f_\star(c(x)) = f_\star(x)$$

*Proof.* Suppose $c(x) \leq (f^\star \circ f_\star)(x)$. Substitute $x = f^\star(y)$ then, because $c$ is extensive,

$$f^\star(y) \leq c(f^\star(y)) \leq (f^\star \circ f_\star \circ f^\star)(y) = f^\star(y).$$

© David Rufino CC BY-NC-ND 4.0

Therefore $c(f^\star(y)) = f^\star(y)$ and $\mathrm{Im}(f^\star) \subseteq \mathrm{Fix}(c)$ as required. Conversely suppose this holds, then by the monotone property of $c$ and extensive property of $f^\star \circ f_\star$

$$x \leq (f^\star \circ f_\star)(x) \implies c(x) \leq c((f^\star \circ f_\star)(x)) = (f^\star \circ f_\star)(x).$$

as required. Finally by the extensive property of $c$

$$x \leq c(x) \leq (f^\star \circ f_\star)(x)$$

and by the antitone/monotone property of $f_\star$ and triangular identity

$$f_\star(x) \leq f_\star(c(x)) \leq f_\star(x)$$

whence $f_\star(c(x)) = f(x)$ as required. ∎

The meaning of the "adjoint" criterion can be explained by the following rather generic situation

**Example 2.1.54 (Canonical example of an antitone Galois connection)**
*Suppose there is a predicate*

$$\psi : X \times Y \to \{0, 1\}$$

*Define a connection*

$$\mathcal{P}(X) \underset{f^\star}{\overset{f_\star}{\rightleftarrows}} \mathcal{P}(Y)$$

*by*

$$f_\star(S) = \{y \in Y \mid \psi(x, y) = 1 \quad \forall x \in S\}$$

$$f^\star(T) = \{x \in X \mid \psi(x, y) = 1 \quad \forall y \in T\}$$

*Then*

$$S \subseteq f^\star(T) \iff \psi(s, t) = 1 \quad \forall s \in S \quad t \in T \iff T \subseteq f_\star(S)$$

**Proposition 2.1.55 (Joins under Galois Correspondence)**
*Let* $\left(X, \underset{X}{\leq}\right)$ *and* $\left(Y, \underset{Y}{\leq}\right)$ *be complete lattices with an antitone Galois connection* $X \underset{f^\star}{\overset{f_\star}{\rightleftarrows}} Y$*. Then for* $S \subseteq X$

$$f_\star\Big(\bigvee S\Big) = \bigwedge f_\star(S)$$

*Similarly for* $T \subseteq Y$ *we have*

$$f^\star\Big(\bigvee T\Big) = \bigwedge f^\star(T)$$

*Proof.* Let $a = \bigvee S$ and $b = \bigwedge f_\star(S)$. Then $s \leq a \implies f_\star(a) \leq f_\star(s)$ for all $s \in S$, which implies $f_\star(a) \leq b$. Similarly $b \leq f_\star(s) \implies s \leq f^\star(b)$ by the adjoint criterion. Therefore $a \leq f^\star(b)$ by definition of join, which implies $b \leq f_\star(a)$ by the adjoint criterion again. Whence $f_\star(a) = b$ as required.

The second statement follows from duality. ∎

## 2.1.7 Axiom of Choice

**Theorem 2.1.56 (Axiom of choice)**
*There are a number of essentially equivalent forrmulations of the axiom of choice*

*a) The Cartesian product of a non-empty family of sets is non-empty*
*b) For any set $X$ of non-empty sets there exists a function $f : X \to \bigcup X$ such that $A \in X \implies f(A) \in A$.*
*c) **Zorn's Lemma** Suppose a partially ordered set $(X, \leq)$ is such that every chain in $X$ has an upper bound in $X$. Then $X$ contains at least one maximal element.*
*d) Every surjective function has a right inverse.*

**Corollary 2.1.57 (Choose representatives)**
*Let $\pi : X \to Y$ be a surjective function and $T \subseteq Y$ a subset. Then there exists a subset $S \subseteq X$ such that $\pi|_S$ is bijective.*

*When $T$ is finite $\#S = \#T$.*

© David Rufino CC BY-NC-ND 4.0

**Definition 2.1.58 (Finite Character)**
*A family of sets $\mathcal{F}$ has **finite character** if it satisfies the following property*

$$A \in \mathcal{F} \iff (B \subseteq A \text{ and finite} \implies B \in \mathcal{F})$$

**Corollary 2.1.59 (Tukey's Lemma)**
*Let $\mathcal{F}$ be a family of sets of finite character. Then it is chain-complete when ordered by inclusion.*

*In particular every set is contained in a maximal set.*

## 2.1.8 Chain Conditions

**Definition 2.1.60 (Totally ordered / chains)**
*A poset $(\mathcal{F}, \leq)$ is **totally ordered** if $x \leq y$ or $y \leq x$ for all $x, y \in \mathcal{F}$.*

**Definition 2.1.61 (Chain)**
*A non-empty subset $C$ of $\mathcal{F}$ is a **chain** if it is totally ordered under $\leq$.*

*The length of the chain is simply $\ell(C) := |C| - 1$.*

*A chain $C$ is*

- ***saturated** if $x \leq z \leq y$ and $x, y \in C \implies z \in C$.*
- ***maximal** if it's not contained properly in another chain.*

**Definition 2.1.62 (Chain-Complete)**
*A poset $(\mathcal{F}, \leq)$ is **chain complete** if every chain $C$ has a supremum in $\mathcal{F}$. It is **co-chain complete** if every chain $C$ has an infimum in $\mathcal{F}$.*

**Proposition 2.1.63 (Noetherian / Artinian Poset)**
*Let $(X, \leq)$ be a poset then the following conditions are equivalent*

*a) Any ascending chain*

$$x_1 \leq x_2 \leq ... \leq x_n \leq ...$$

*eventually stabilizes*

*b) Any non-empty subset $Y \subseteq X$ has a maximal element*

*Such a poset is called **Noetherian**. If it satisfies the dual condition then it is called **Artinian**.*

*Proof.* a) $\implies$ b) If $Y$ has no maximal elements then we may (by axiom of dependent choice) construct a strictly increasing sequence, which by definition does not stabilize.

b) $\implies$ a) Clear. ■

© David Rufino CC BY-NC-ND 4.0

## 2.2 Numbers

Informally we consider the set of integers

$$\mathbb{Z} = \{..., -3, -2, -1, 0, 1, 2, 3, ...\}$$

and the subset of natural numbers

$$\mathbb{N} = \{0, 1, 2, ..., \}$$

Although it's possible to construct the integers painstakingly from a small set of axioms (see ...) we instead for brevity simply state the most commonly used results as axioms.

### 2.2.1 Integers

We suppose the existence of a set $\mathbb{Z}$ with distinguished elements $0 \neq 1$ together with

- A binary operation

  $$+ : \mathbb{Z} \times \mathbb{Z} \to \mathbb{Z}$$

  and an involution

  $$(-) : \mathbb{Z} \to \mathbb{Z}$$

  satisfying

  - $-0 = 0$
  - $-(-x) = x$
  - $x = -x \iff x = 0$
  - $x + 0 = 0 = 0 + x$
  - $x + y = y + x$
  - $(x + y) + z = x + (y + z)$
  - $x + (-x) = 0 = (-x) + x$
  - $-(x + y) = (-x) + (-y)$

- A subset $\mathbb{N}$ such that

  - $0, 1 \in \mathbb{N}$
  - $x, y \in \mathbb{N} \implies x + y \in \mathbb{N}$
  - $x \in \mathbb{Z} \implies (x \in \mathbb{N}) \vee (-x \in \mathbb{N})$
  - $x \in \mathbb{N} \wedge -x \in \mathbb{N} \implies x = 0$

  which also satisfies the principle of induction

- Let $S \subseteq \mathbb{N}$ be a set such that
  - $0 \in S$
  - $x \in S \implies x + 1 \in S$

  then $S = \mathbb{N}$

It's possible to use these to show the existence of multiplication

**Proposition 2.2.1 (Multiplication exists)**
*There exists a binary operation*

$$\cdot : \mathbb{Z} \times \mathbb{Z} \to \mathbb{Z}$$

*such that*

- $x \cdot 0 = 0 = 0 \cdot x$
- $x \cdot 1 = x = 1 \cdot x$
- $xy = yx$
- $(xy)z = x(yz)$

© David Rufino CC BY-NC-ND 4.0

- $x(y+z) = xy + xz$
- $(y+z)x = yx + zx$
- $(-x)(y) = -(xy) = x(-y)$

We may also show the existence a partial ordering

**Proposition 2.2.2 (Order exists)**
*There exists a relation $\leq$ on $\mathbb{Z}$ given by*

$$x \leq y \iff y - x \in \mathbb{N}$$

*which satisfies*

$$x \leq y \vee y \leq x$$

$$x \leq y \wedge y \leq x \implies x = y$$

*Define $x < y$ in the obvious way then it satisfies the usual trichotomy law, namely precisely one of the following holds*

$$x < y, \quad x = y, \quad y < x$$

*and further*

- *$z > 0$ then $x < y \iff xz < yz$*
- *$z < 0$ then $x < y \iff yz < xz$*
- *$y > 1$ and $x > 0$ then $x < xy$*

Finally we can construct an absolute value function

**Proposition 2.2.3**
*There exists an **absolute value** function*

$$|\cdot| : \mathbb{Z} \to \mathbb{N}$$

*such that*

$$|x| = \begin{cases} x & 0 < x \\ 0 & x = 0 \\ -x & x < 0 \end{cases}$$

*It satisfies*

- $|x| = 0 \iff x = 0$
- $|x| = |y| \iff x = \pm y$
- $|xy| = |x||y|$
- $|x + y| \leq |x| + |y|$

In many cases it may be more convenient to use the following form of induction

**Proposition 2.2.4 (Well-Ordering Principle)**
*Let $S \subset \mathbb{N}$ be a non-empty subset. Then it contains a minimal element $d$.*

### 2.2.2 Arithmetic

**Proposition 2.2.5 (Division Algorithm)**
*Let $x, y \in \mathbb{Z}$ be non-zero integers then there exists $q, r$ such that*

$$x = yq + r$$

*and*

$$0 \leq r < |y|$$

*Furthermore $(q, r)$ is the unique such pair.*

*Proof.* Suppose first that $x, y > 0$. Let $S = \{x - yn \mid n \in \mathbb{Z}\} \cap \mathbb{N}$. Then $x \in S$ so it is non-empty. By the Well-Ordering principle it has a minimal element $r$. By assumption

$$x = yq + r$$

© David Rufino CC BY-NC-ND 4.0

for some $q \in \mathbb{Z}$ and $r \geq 0$. Suppose $r \geq y$, then $0 \leq x - y(q+1) < r$ contradicting minimality.

The case $x > 0, y < 0$ is then straightforward, as is the case $x < 0$.

For uniqueness suppose $yq' + r' = yq + r$ then $|y||q - q'| = |r' - r| < |y|$ from which it follows $|q - q'| = 0 \implies q = q' \implies r = r'$. ■

**Corollary 2.2.6 (Ideals are Principal)**
*Let $S \subseteq \mathbb{Z}$ be a non-empty set such that*

$$x, y \in S \implies x \pm y \in S$$

*Then $S = d\mathbb{Z}$ for a unique $d \geq 0$.*

*Proof.* First we claim that $0 \in S$. For if $x \in S$ then $0 = x - x \in S$ by assumption. Furthermore $x \in S \implies -x = 0 - x \in S$.

Consider the set $S' = (S \cap \mathbb{N}) \setminus \{0\}$. If it's empty then $S = \{0\}$ (for $x \in S \implies -x \in S$) and $d = 0$.

Otherwise it has a minimal element $d > 0$ by the well-ordering principle. Then by induction $d\mathbb{Z} \subseteq S$. Conversely suppose $y \in S$ then by the division algorithm $y = qd + r$ with $0 \leq r < d$. By assumption $r = y - qd \in S$ and by minimality must be equal to 0. Therefore $y \in d\mathbb{Z}$ and $d\mathbb{Z} = S$ as required. ■

**Definition 2.2.7 (Divisibility)**
*Let $x, y \in \mathbb{Z}$ be two integers. We say that $x$ divides $y$ if there exists $a$ such that $ax = y$. In this case we write*

$$x \mid y$$

*and*

$$\frac{y}{x}$$

*for the unique integer $a$ such that $ax = y$.*

**Lemma 2.2.8**
*Let $x, y \in \mathbb{Z}$ be two integers then*

$$x \mid y \implies |x| \leq |y|$$

*In particular $x \mid y \wedge y \mid x \implies x = \pm y$.*

**Proposition 2.2.9 (Bezout's Theorem)**
*Let $x, y$ be non-zero integers. Then there exists a unique positive integer $d$ such that*

- *$d$ is a common divisor of $x, y$*
- *For any other common divisor $e$ we have $e \mid d$.*

*Further there exists integers $a, b$ such that $ax + by = d$. We write this as $(x, y)$.*

*Proof.* Let $S = \{ax + by \mid a, b \in \mathbb{Z}\}$. Then by (2.2.6) we have $S = d\mathbb{Z}$ for a unique $d > 0$. As $x, y \in S$ by definition $d$ is a common divisor, and by definition $d = d \cdot 1 = ax + by$ for some integers $a, b$. Suppose $e$ is a common divisor then $d = ax + by = e(ap + bq)$ and $e \mid d$ as required.

Any two such common divisors have $d = \pm d'$ by the previous Lemma. Since they are positive and non-zero we have $d = d'$. ■

**Proposition 2.2.10**
*Let $a, x, y$ be non-zero integers then*

$$|a|(x, y) = (ax, ay)$$

*In particular*

$$\left(\frac{x}{x, y}, \frac{y}{x, y}\right) = 1$$

*Proof.* This follows from the characterization of $(x, y)$ as the minimal positive integer in the set $\{mx + ny\}$. ■

© David Rufino CC BY-NC-ND 4.0

### 2.2.3 Prime Factorization

**Definition 2.2.11**
*Let $x \in \mathbb{Z}$ be a non-zero integer. We say that $x$*

- *is a **unit** if it's equal to $1$ or $-1$.*
- *is **prime** if it's not a unit and $x \mid p$ implies $x = \pm 1$ or $x = \pm p$*
- ***composite** otherwise*

**Lemma 2.2.12**
*Let $p$ be a positive prime and $a$ a non-zero integer. Then precisely one of the following holds*

- $(p, a) = 1$
- $(p, a) = p$ *and* $p \mid a$

*Proof.* Note that $(p, a)$ is positive and divides both $p$ and $a$ so the result follows by definition of prime. ■

**Proposition 2.2.13 (Euclid's Lemma)**
*Suppose $x \mid ab$ then $\frac{x}{x,a} \mid b$.*

*In particular if $p$ is a prime and $p \mid ab$, then $p \mid a$ or $p \mid b$.*

*Proof.* First suppose that $(x, a) = 1$. Then by assumption $zx = ab$ and by Bezout's Theorem $mx + na = 1$ for some integers $m, n$. Multiply by $z$ to find that $abm + na = a(bm + n) = z$. Therefore $a(bm + n)x = ab$ and cancel $a$ to find $x \mid b$ as required.

For the general case define $x' = x/(x, a)$ and $a' = a/(x, a)$. Then by (2.2.10) $(x', a') = 1$. Furthermore it's clear that $x' \mid a'b$ so we have $x' \mid b$ by the special case just proven.

Finally suppose $p \mid ab$. If $(p, a) = 1$ then $p \mid b$ by the first result. By (2.2.12) if this does not hold then $p \mid a$ as required.■

Using these results we may show that there exists a unique factorization into primes, unique up to multiplication by a unit.

© David Rufino CC BY-NC-ND 4.0

## 2.3 Matroids

The theory of bases of vector spaces (the Vector Spaces subsection) and transcendence bases of field extensions (the Transcendental Field Extensions subsection) have some formal similarities, as noted in [Wae91]. Here we use the theory of Matroids to formalise this precisely so that the proofs need not be repeated in each case.

**Definition 2.3.1 (Matroid)**
*Consider a set $X$ together with a closure operator $c : \mathcal{P}(X) \to \mathcal{P}(X)$ ("span" operator) such that $c(\varnothing) = \varnothing$. We say*

- *$S \subseteq X$ is **independent** if $x \in S \implies x \notin c(S \setminus \{x\})$*
- *$\Gamma \subseteq X$ is **spanning** if $c(\Gamma) = X$.*

*Note by definition $X$ is spanning and $\varnothing$ is independent. Moreover all singletons $\{x\}$ are independent.*

*We call the pair $(X, c)$ a **matroid** if it also satisfies the following properties*

- ***Finitary** $x \in c(\Gamma) \implies x \in c(\Gamma')$ for some finite subset $\Gamma'$ of $\Gamma$.*
- ***Exchange Property** For all $x, y \in X$ and $Y \subseteq X$ we have*

$$x \in c(Y \cup \{y\}) \setminus c(Y) \implies y \in c(Y \cup \{x\})$$

*We say $(X, c)$ has **finite rank** if it has a finite spanning set.*

*Finally we say $\mathcal{B} \subseteq X$ is a **basis** if it is both **spanning** and **independent**.*

We begin with some elementary characterizations of independent sets

**Lemma 2.3.2**
*Suppose $S \subseteq X$ is a subset*

*a) $A \subseteq c(S) \implies c(S \cup A) = c(S)$*
*b) $S$ is independent if and only if no proper subset has the same span.*

*Proof.* We prove each in turn

a) By monotonicity

$$c(S) \subseteq c(S \cup A) \subseteq c(c(S) \cup A) = c(c(S)) = c(S)$$

b) Suppose $S$ is independent and $S' \subsetneq S$ is a proper subset such that $c(S') = c(S)$. Choose $x \in S \setminus S'$ then by definition $x \in S \implies x \in c(S) = c(S') \subseteq c(S \setminus \{x\})$ contradicting independence.

Conversely suppose for some $x \in S$ we have $x \in c(S \setminus \{x\})$. Define $S' \coloneqq S \setminus \{x\}$. Then $x \in c(S')$ implies $c(S) = c(S')$ by a). As $S'$ is a proper subset this contradicts the hypothesis.

■

**Lemma 2.3.3**
*Every subset of an independent set is independent. Furthermore the family of independent sets has finite character.*

*Proof.* The first statement is straightforward. Suppose $S$ a dependent set such that every finite subset is independent. Then there exists $x \in S$ such that $x \in c(S \setminus \{x\})$. Then by the finitary property there exists a finite subset $S' \subseteq S \setminus \{x\}$ such that $x \in c(S')$. Therefore by definition $S' \cup \{x\}$ is not independent, a contradiction. ■

The finitary condition ensures that $\mathcal{E}$ is "inductively ordered"

**Corollary 2.3.4**
*Let $\{S_i\}_{i \in I}$ be a chain of independent subsets. Then $S = \bigcup_{i \in I} S_i$ is also independent.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Let $F \subseteq S$ be a finite subset. For each $x \in F$ choose $i_x \in I$ with $x \in S_{i_x}$. As $\{S_i\}_{i \in I}$ is a chain and $F$ is finite, the sets $\left\{S_{i_x}\right\}_{x \in F}$ are totally ordered by inclusion and so admit a maximum $S_{i_0}$, whence $F \subseteq S_{i_0}$. By (2.3.3) $F$ is independent, being a subset of the independent set $S_{i_0}$. As $F \subseteq S$ was an arbitrary finite subset, $S$ is independent by (2.3.3). ■

**Lemma 2.3.5 (Extension Property)**
*Suppose $S$ is an independent set and $x \notin c(S)$, then $S \cup \{x\}$ is independent.*

*Proof.* We require to prove that for all $y \in S$ we have $y \notin c(S \cup \{x\} \setminus \{y\})$. By independence of $S$ we have $y \notin c(S \setminus \{y\})$, so by the Exchange Property $y \in c(S \cup \{x\} \setminus \{y\})$ would imply $x \in c(S)$, contradicting the hypothesis. ■

© David Rufino CC BY-NC-ND 4.0

**Proposition 2.3.6**
*Let $\mathcal{F}$ be a family of independent sets which satisfies the following properties*

- $\varnothing \in \mathcal{F}$
- ***extension property*** $S \in \mathcal{F}$ *and* $x \notin c(S) \implies S \cup \{x\} \in \mathcal{F}$
- $\mathcal{F}$ *has finite character*

*then $\mathcal{F}$ consists of all independent sets.*

*Proof.* It's enough to show that $\mathcal{F}$ contains all finite independent sets, which follows by induction on $\#S$. For given an independent set $S$ and $x \in S$, then by definition $x \notin c(S \setminus \{x\})$. By the induction hypothesis $S \setminus \{x\} \in \mathcal{F}$ whence by the extension property $S \in \mathcal{F}$ as required. ∎

**Proposition 2.3.7 (Basis exists)**
*Let $(X, c)$ be a matroid, $S$ independent and $\Gamma$ a subset such that $S \subseteq \Gamma$. Then there exists an independent set $\mathcal{B}$ such that $S \subseteq \mathcal{B} \subseteq \Gamma$ and $c(\mathcal{B}) = c(\Gamma)$.*

*In particular if $\Gamma$ is spanning then $\mathcal{B}$ is a basis.*

*Proof.* Consider the collection

$$\mathcal{I} = \{T \text{ independent} \mid S \subseteq T \subseteq \Gamma\}$$

By (2.3.4), $\mathcal{I}$ is chain-complete. Therefore it has a maximal element $\mathcal{B}$ by Zorn's Lemma. Suppose $x \in \Gamma \setminus c(\mathcal{B})$ then $\mathcal{B} \cup \{x\}$ is independent by (2.3.5), contradicting maximality. Therefore $\Gamma \subseteq c(\mathcal{B}) \implies c(\Gamma) \subseteq c(\mathcal{B})$. The reverse inequality is clear so that $c(\Gamma) = c(\mathcal{B})$. ∎

**Corollary 2.3.8 (Criteria for bases)**
*Let $(X, c)$ be a matroid. Then the following are equivalent*

*a) $\mathcal{B}$ is a basis*
*b) $\mathcal{B}$ is a minimal spanning set*
*c) $\mathcal{B}$ is maximally independent (possibly in some spanning set $\Gamma$)*

*Proof.* a) $\implies$ b). Let $\mathcal{B}$ be a basis and $\Gamma \subseteq \mathcal{B}$ a spanning set. Then by (2.3.2) b) $\Gamma = \mathcal{B}$.

b) $\implies$ a). Let $\mathcal{B}$ be a minimal spanning set, then by (2.3.7) there exists a subset $\mathcal{B}'$ which is a basis, and in particular spanning. By minimality $\mathcal{B} = \mathcal{B}'$.

c) $\implies$ a). By (2.3.7) there exists a basis $\mathcal{B}'$ containing $\mathcal{B}$. By maximality $\mathcal{B}' = \mathcal{B}$.

a) $\implies$ c). Suppose $\mathcal{B} \subseteq S$ where $S$ is independent. Then $S$ is spanning too, and so by (2.3.2) b) $\mathcal{B} = S$. ∎

**Definition 2.3.9 (Sub-matroid)**
*A subset $Y \subseteq X$ is a **sub-matroid** if $c(Y) = Y$. In this case $S \subseteq Y \implies c(S) \subseteq Y$ and so we have an induced matroid structure $(Y, c)$.*

**Lemma 2.3.10 (Mini Exchange Lemma)**
*Let $S \subseteq \Gamma$ be finite sets and $x \in X \setminus S$ such that*

- *$S$ is independent*
- *$x \in c(\Gamma) \setminus c(S)$*

*Then there exists $y \in \Gamma \setminus S$ such that $c(\Gamma \setminus \{y\} \cup \{x\}) = c(\Gamma)$.*

*Proof.* We may assume without loss of generality that $x \notin \Gamma$.

Consider $\tilde{\Gamma} \subseteq \Gamma$ minimal subject to $S \subseteq \tilde{\Gamma}$ and $x \in c(\tilde{\Gamma})$. If $S = \tilde{\Gamma}$ then $x \in c(S)$ a contradiction. Therefore we may choose $y \in \tilde{\Gamma} \setminus S$. By minimality we have $x \in c(\tilde{\Gamma}) \setminus c(\tilde{\Gamma} \setminus \{y\})$. Therefore by the Exchange Property we have $y \in c(\tilde{\Gamma} \setminus \{y\} \cup \{x\})$. Then by (2.3.2) applied twice

$$c(\Gamma \setminus \{y\} \cup \{x\}) = c(\Gamma \cup \{x\}) = c(\Gamma)$$

as required. ∎

**Proposition 2.3.11 (Exchange Lemma)**
*Let $S$ be an independent set and $\Gamma$ be a finite set such that $S \subseteq c(\Gamma)$. Then there exists a subset $T \subseteq \Gamma$ such that*

- *$\#T = \#S$*

© David Rufino CC BY-NC-ND 4.0

- $c(\Gamma \setminus T \cup S) = c(\Gamma)$.

*In particular* $\#S \leq \#\Gamma$.

*Proof.* By considering the sub-matroid $(c(\Gamma), c)$ we may assume without loss of generality that $X = c(\Gamma)$.

Let $n = \#\Gamma$ and consider the set of pairs

$$\mathcal{F} \coloneqq \{(A, B) \mid A \subseteq S, \quad B \subseteq \Gamma, \quad \#A = \#B, \quad c(\Gamma \setminus B \cup A) = X\}$$

Essentially $\mathcal{F}$ is the set of swaps we may perform from $S$ to $\Gamma$ whilst preserving the span. It is non-empty because $(\varnothing, \varnothing) \in \mathcal{F}$ and we wish to show that $(S, B) \in \mathcal{F}$ for some $B$.

Observe that for all $(A, B) \in \mathcal{F}$ we have $\#B \leq n$ so choose a pair such that $\#B$ is maximal. We wish to show that in this case $A = S$, so we suppose to the contrary that $A \subsetneq S$.

We claim that $B \subsetneq \Gamma$, for $B = \Gamma$ implies by construction $c(A) = X$ which would imply $c(A) = c(S)$, contradicting the criteria for independence of $S$ given by (2.3.2) b).

Define $\Gamma' \coloneqq \Gamma \setminus B \cup A$. Then by assumption $c(\Gamma') = X$ and $A \subseteq \Gamma'$ is independent. Choose $x \in S \setminus A$ then by definition of independence $x \notin c(A)$. By (2.3.10) there exists $y \in \Gamma' \setminus A$ such that

$$c(\Gamma' \setminus \{y\} \cup \{x\}) = c(\Gamma') = X$$

Note by construction that $y \notin B$, so we see that $(A \cup \{x\}, B \cup \{y\}) \in \mathcal{F}$ has greater length, which contradicts maximality. ■

**Corollary 2.3.12 (Bases have the same cardinality)**
*Every base of a finite rank matroid is finite and of the same size. Denote this by* $r(X)$.

*Proof.* There is a finite basis by (2.3.7). Then apply (2.3.11). ■

**Corollary 2.3.13**
*Let* $(X, c)$ *be a finite-rank matroid and* $S \subseteq X$ *an independent subset, then* $\#S \leq r(X)$. *Similarly a spanning subset* $\Gamma$ *satisfies* $\#\Gamma \geq r(X)$.

*Proof.* Apply (2.3.7) and (2.3.12). ■

**Corollary 2.3.14 (Basis Criteria)**
*Let* $\mathcal{B}$ *be a subset of a finite-rank matroid* $(X, c)$. *Then the following are equivalent*

*a)* $\mathcal{B}$ *is a basis*
*b)* $\mathcal{B}$ *is independent and* $\#\mathcal{B} \geq r(X)$
*c)* $\mathcal{B}$ *is spanning and* $\#\mathcal{B} \leq r(X)$

*Proof.* Apply (2.3.7) and (2.3.12). ■

**Proposition 2.3.15**
*Let* $Y \subseteq X$ *be a sub-matroid of a finite-rank matroid. Then* $Y = X \iff r(Y) = r(X)$.

*Proof.* Let $\mathcal{B}$ be a basis for $Y$ then $\#\mathcal{B} = r(Y) = r(X)$ and is a-fortiori independent in $X$. Therefore by (2.3.14) $\mathcal{B}$ is a basis for $X$ and hence $X = c(\mathcal{B}) = Y$. ■

© David Rufino CC BY-NC-ND 4.0

## 2.4 Decomposition in Noetherian and Distributive Lattices

An analogue of irreducible factorization in rings (see the Factorisation section) applies to Noetherian Lattices. Furthermore uniqueness holds when the lattice is distributive. For a canonical reference see [Bir40].

**Definition 2.4.1 (Meet-Prime and Meet-Irreducible)**
*Let $(X, \leq)$ be a lattice and $x \in X$. Then we say that $x$ is*

- ***meet-irreducible*** *if $y \wedge z = x \implies y = x$ or $z = x$*
- ***meet-prime*** *if $y \wedge z \leq x \implies y \leq x$ or $z \leq x$*
- ***join-irreducible*** *if $y \vee z = x \implies y = x$ or $y = z$*
- ***join-prime*** *if $x \leq y \vee z \implies x \leq y$ or $x \leq z$*

The following result is proven in [Bir40] Ch. IX Lemma 4.1.

**Proposition 2.4.2 (Prime = Irreducible)**
*Let $(X, \leq)$ be a lattice. In general meet-prime $\implies$ meet-irreducible and join-prime $\implies$ join-irreducible. If $X$ is a distributive lattice, then the converse holds.*

*In the distributive case we denote by $\mathcal{M}(X)$ and $\mathcal{J}(X)$ the sub-poset of meet-prime and join-prime elements respectively.*

*Proof.* The first statement is straightforward. Conversely suppose $X$ is a distributive lattice and $x$ is meet-irreducible. If $y \wedge z \leq x$ then $x = x \vee (y \wedge z) = (y \vee x) \wedge (z \vee x) \implies x = y \vee x$ or $x = z \vee x$, whence the result follows. ∎

**Proposition 2.4.3**
*Let $(X, \leq)$ be a distributive lattice and $Y$ a sub-lattice, then*

$$\mathcal{M}(Y) = \mathcal{M}(X) \cap Y$$

$$\mathcal{J}(Y) = \mathcal{J}(X) \cap Y$$

*In particular this holds when $Y = \{x\}^{\uparrow}, \{y\}^{\downarrow}, \{x\}^{\uparrow} \cap \{y\}^{\downarrow}$.*

**Proposition 2.4.4**
*Let $(X, \leq)$ be a distributive lattice. If it is chain-complete (resp. co-chain-complete) then so is $\mathcal{J}(X)$ (resp. $\mathcal{M}(X)$).*

*Every join-prime element is bounded above by a maximal join-prime element, and every meet-prime element is bounded below by a minimal meet-prime element*

*Proof.* Let $C$ be a chain of join-prime elements and $x := \bigvee C$. Suppose that $y \vee z \geq x$ then $y \vee z \geq w$ for all $w \in C$. Then $y \geq w$ or $z \geq w$ for all $w \in C$. Let $C_1 := \{w \in C \mid y \geq w\}$. If $C_1 = C$ then we are done as $x \leq y$. Otherwise suppose $w_0 \notin C_1$ then by prime-ness $z \geq w_0$. Clearly $w \leq w_0 \implies w \leq z$. Further $w \geq w_0 \implies w \not\leq y$ (as otherwise $w_0 \leq y$) whence $w \leq z$. Therefore $x \leq z$.

The last statement follows from Zorn's Lemma by considering the sub-lattices $\{x\}^{\uparrow}$ and $\{y\}^{\downarrow}$ which inherit the chain complete properties. ∎

**Definition 2.4.5**
*Let $(X, \leq)$ be a lattice and $Y \subseteq X$ a finite subset. Then we say that $Y$ is*

- ***(meet-)irredundant*** *if no proper subset has the same meet*
- ***incomparable*** *(or an **antichain**) if no two elements are comparable*

**Lemma 2.4.6 (incomparable $\iff$ irredundant)**
*Let $(X, \leq)$ be a lattice and $Y \subseteq X$ then $Y$ meet-irredundant $\implies$ $Y$ incomparable. Conversely if $Y$ is a finite incomparable subset of meet-prime elements then $Y$ is meet-irredundant.*

*Proof.* The first part is straightforward, for if $y_1 \lneq y_2$ are elements of $Y$ then $\bigwedge Y = \bigwedge Y \setminus \{y_2\}$.

Conversely suppose $Y' \subsetneq Y$ is such that $\bigwedge Y' = \bigwedge Y$. Choose $y_2 \in Y \setminus Y'$, then $\bigwedge Y' \leq y_2$, whence by definition of meet-prime (and induction) $y_1 \leq y_2$ for some $y_1 \in Y'$. ∎

The following is [Bir40] Chapter IX Theorem 9

**Proposition 2.4.7 (Decomposition in Noetherian Lattice)**
*Let $(X, \leq)$ be a Noetherian distributive lattice. Then every element $x \in X$ has a unique decomposition*

© David Rufino CC BY-NC-ND 4.0

$$x = x_1 \wedge ... \wedge x_n$$

*where $x_i$ are meet-prime and meet-irredundant (equivalently incomparable). These are precisely the meet-primes minimal over $x$.*

*Dually, if $(X, \leq)$ is an Artinian distributive lattice, then every element $x \in X$ has a unique decomposition*

$$x = x_1 \vee ... \vee x_n$$

*where $x_i$ are join-prime and join-irredundant (equivalently incomparable). These are precisely the join-primes maximal below $x$.*

*Proof.* Let $Y$ be the subset of elements which are not finite meets of meet-prime elements, and suppose it is non-empty. Then by (2.1.63) $Y$ has a maximal element $x_0$. It cannot be meet-prime, and therefore not meet-irreducible (2.4.2), so there must exist elements $y_0, z_0 \in X$ such that $x_0 = y_0 \wedge z_0$ but $x_0 \lneq y_0$ and $x_0 \lneq z_0$. By maximality $y_0, z_0$ are finite meets of prime elements, and therefore so is $x_0$ a contradiction.

Therefore we have a decomposition into distinct primes

$$x = x_1 \wedge ... \wedge x_n$$

Consider the family of subsets of $\{x_1, ..., x_n\}$ which have the same meet. Then there exists a minimal subset which by definition is meet-irredundant and by (2.4.6) incomparable. Suppose there is another such decomposition $x = x'_1 \wedge ... \wedge x'_m$. Then for every $i = 1...n$ we have $x'_{\sigma(i)} \leq x_i$ and for every $j = 1...m$ we have $x_{\tau(j)} \leq x'_j$ whence $x_{\tau(\sigma(i))} \leq x'_{\sigma(i)} \leq x_i$. As the decomposition is incomparable we have $\tau(\sigma(i)) = i$ and $x'_{\sigma(i)} = x_i$. Therefore $\sigma$ is injective and $n \leq m$. By symmetry $m \leq n$ and $\sigma$ is a bijection. In otherwords the decomposition is unique.

Note $x \leq z$ and $z$ meet-prime implies $x_j \leq z$ for some $j$. Therefore if $z$ is a minimal prime then $x_j = z$ and all the meet-primes minimal over $x$ must occur in the decomposition. Similarly if $z \leq x_i$ then by incomparability $x_j = z = x_i$. Therefore each $x_i$ is also minimal. ∎

© David Rufino CC BY-NC-ND 4.0

## 2.5 Krull Dimension

The purpose of this section is to abstract the notions of Krull Dimension in commutative ring theory and topology. A more standard approach (eg EGA IV) would be to develop the topological notion first, and then link to commutative ring case using the prime spectrum. Generally the concept is not well-behaved, so stronger conditions are defined which generally hold in geometric cases. Principle references are (EGA0 IV 14.3, Heinrich).

**Definition 2.5.1 (Finite-Dimensional Poset)**
*Let $(\mathcal{G}, \leq)$ be a poset, we say that it is **finite-dimensional** if*

$$\dim(\mathcal{G}) := \sup\{\ell(C) \mid C \subseteq \mathcal{G} \text{ a chain}\} < \infty$$

*In this case we define*

$$\begin{aligned}\dim(x) &:= \dim\big(\{x\}^{\downarrow}\big)\\ \operatorname{codim}(y) &:= \dim\big(\{y\}^{\uparrow}\big)\\ \operatorname{codim}(y, x) &:= \dim\big(\{x\}^{\downarrow} \cap \{y\}^{\uparrow}\big)\end{aligned}$$

*Note $\mathcal{G}$ is both Noetherian and Artinian, but finite-dimensionality is a stronger condition. Note also that $\{x\}^{\downarrow}, \{y\}^{\uparrow}$ and $\{x\}^{\downarrow} \cap \{y\}^{\uparrow}$ are finite-dimensional posets*

**Definition 2.5.2 (Krull Lattice)**
*Let $(\mathcal{F}, \leq)$ be an **Artinian** distributive lattice. We say it is a **Krull Lattice** if the poset of join-prime elements $\mathcal{J}(\mathcal{F})$ is finite-dimensional and define*

$$\dim(\mathcal{F}) := \dim(\mathcal{J}(\mathcal{F}))$$

*By (2.4.3) we have $\mathcal{H} = \{x\}^{\downarrow}, \{y\}^{\uparrow}, \{x\}^{\downarrow} \cap \{y\}^{\uparrow}$ are Krull Lattices such that*

$$\mathcal{J}(\mathcal{H}) = \mathcal{J}(\mathcal{F}) \cap \mathcal{H}$$

*For $x, y \in \mathcal{F}$ we have a unique decomposition into maximal join-prime elements $x_i, y_j \in \mathcal{J}(\mathcal{F})$ (2.4.7)*

$$x = x_1 \vee \ldots \vee x_n$$

$$y = y_1 \vee \ldots \vee y_m$$

*Note that $y \leq x \iff$ for all $j$ we have $y_j \leq x_i$ for some $i$ and we may define*

$$\begin{aligned}\dim(x) &:= \max_i \dim(x_i) = \dim\big(\{x\}^{\downarrow}\big)\\ \operatorname{codim}(y) &:= \min_j \operatorname{codim}\big(y_j\big)\\ \operatorname{codim}(y, x) &:= \min_j \max_i \big\{\operatorname{codim}\big(y_j, x_i\big) \mid y_j \leq x_i\big\}\\ &= \min_j \operatorname{codim}\big(y_j, x\big)\end{aligned}$$

*note it's required to be careful in definition of co-dimension in order to have a sensible co-dimension formula. Note also that*

$$\dim\big(y; \{x\}^{\downarrow}\big) = \dim(y)$$

**Remark 2.5.3**
*For the topological case, we would define $\mathcal{F}$ to be the closed subsets of $X$ and $\mathcal{J}(\mathcal{F})$ would be the collection of irreducible closed subsets.*

**Proposition 2.5.4 (Extending chains)**
*Let $(\mathcal{G}, \leq)$ be a finite-dimensional poset*

- *Every chain is contained in a saturated chain*
- *Every chain is contained in a maximal chain*
- *Every maximal chain is of the form*

  $$x_0 \lneq x_1 \ldots \lneq x_n$$

  *for $x_0$ minimal and $x_n$ maximal in $\mathcal{G}$.*

© David Rufino CC BY-NC-ND 4.0

**Definition 2.5.5 (Properties)**
*Let $\mathcal{G}$ be a finite-dimensional poset. Then we say it is*

- ***Irreducible*** *if it has a top element*
- ***Equidimensional*** *if every maximal element has the same dimension*
- ***Equicodimensional*** *if every minimal element has the same codimension*
- ***Biequidimensional*** *if every maximal chain has the same length*
- ***Quasi-Biequidimensional*** *if $\{x\}^{\downarrow}$ is biequidimensional for all $x$ (equivalently all maximal $x$)*
- ***Catenary*** *if for every pair $y \leq x$, every saturated chain in $[y,x] := \{y\}^{\uparrow} \cap \{x\}^{\downarrow}$ has the same length, namely* $\operatorname{codim}(y,x)$.

*If $\mathcal{F}$ is a Krull Lattice then we say it inherits these properties from $\mathcal{J}(\mathcal{F})$. Note if $\mathcal{F}$ is irreducible then it also has (the same) top element.*

*Trivially irreducible implies equidimensional. Similarly biequidimensional implies both equidimensional and equicodimensional, but not conversely.*

*Finally* ***quasi-biequidimensional + equidimensional $\iff$ biequidimensional.***

**Proposition 2.5.6 (Simple Properties)**
*Let $\mathcal{G}$ be a finite-dimensional poset then*

*a) If $x$ is maximal then* $\operatorname{codim}(x) = 0$
*b) If $x$ is minimal then* $\dim(x) = 0$
*c)* $\dim(\mathcal{G}) = \sup\{\dim(x) \mid x \text{ maximal}\} = \sup\{\operatorname{codim}(x) \mid x \text{ minimal}\}$
*d) For all $z \leq y \leq x$ we have* $\operatorname{codim}(z,y) + \operatorname{codim}(y,x) \leq \operatorname{codim}(z,x)$

*If $\mathcal{F}$ is a Krull Lattice then*

*e) For all $y \leq x$ we have* $\dim(y) + \operatorname{codim}(y,x) \leq \dim(x)$
*f) For all $y \leq x$ we have* $\operatorname{codim}(y,x) = 0 \iff y_j = x_i$ *some $i,j$*

*Alternatively* $\operatorname{codim}(y,x) > 0$ *if and only if* $\big(y_j \leq x_i \implies y_j \neq x_i\big)$.

*Proof.* e) The case of a finite-dimensional poset is (relatively) clear. In the general case then we have

$$\dim\big(y_j\big) + \operatorname{codim}(y,x) \leq \dim\big(y_j\big) + \operatorname{codim}\big(y_j,x\big) = \max_i\big(\dim\big(y_j\big) + \operatorname{codim}\big(y_j,x_i\big)\big) \leq \max_i \dim(x_i) = \dim(x)$$

and taking max over $j$ yields the result.

f) The case $x,y \in \mathcal{J}(\mathcal{F})$ is clear by (2.5.4). For the general case $\operatorname{codim}(y,x) = 0 \iff \operatorname{codim}\big(y_j,x\big) = 0$ for some $j \iff \big(y_j \leq x_i \implies \operatorname{codim}\big(y_j,x_i\big) = 0\big) \iff y_j = x_i$ for some $i,j$. ∎

**Corollary 2.5.7 (Codimension 1 formula)**
*Let $\mathcal{F}$ be a Krull Lattice with $y \in \mathcal{F}$, $x \in \mathcal{J}(\mathcal{F})$ and $y \leq x$. Then*

$$\dim(y) = \dim(x) - 1 \implies \operatorname{codim}(y,x) = 1$$

*Suppose further that $\mathcal{F}$ is irreducible then*

$$\dim(y) = \dim(\mathcal{F}) - 1 \implies \operatorname{codim}(y) = 1$$

*Proof.* By (2.5.6) e) $\operatorname{codim}(y,x) \leq 1$. If $\operatorname{codim}(y,x) = 0$ then by f) we see that $y_j = x$ for some $j$, whence $y = x$ which contradicts $\dim(y) = \dim(x) - 1$. Therefore $\operatorname{codim}(y,x) = 1$ as required.

If $\mathcal{F}$ is irreducible the result follows with $x = \top$. ∎

**Remark 2.5.8 (Duality)**
*We note that the concepts of* ***dimension*** *(of a poset).* ***biequidimensional*** *and* ***catenary*** *are* self-dual, *in the sense that they are preserved when considering the dual poset* $\left(\mathcal{G}, \overset{d}{\leq}\right)$.

*Similarly the concepts of* ***equidimensional*** *and* ***equicodimensional*** *are dual to each other.*

**Proposition 2.5.9**
*Let $\mathcal{G}$ be a finite-dimensional poset. Then the following are equivalent*

- *$\mathcal{G}$ is catenary*

© David Rufino CC BY-NC-ND 4.0

- *For every triplet $z < y < x$ in $\mathcal{G}$ we have*

$$\operatorname{codim}(z, x) = \operatorname{codim}(z, y) + \operatorname{codim}(y, x)$$

*When $\mathcal{G}$ is irreducible this is equivalent to*

$$\operatorname{codim}(z) = \operatorname{codim}(z, y) + \operatorname{codim}(y)$$

*for all $z < y$.*

*Proof.* Suppose $\mathcal{G}$ is catenary. Choose saturated chains $C_1$ in $[z, y]$ and $C_2$ in $[y, x]$. One may show that $C_1 \cap C_2 = \{y\}$ and $C_1 \cup C_2$ is a saturated chain in $[x, z]$. The result follows by definition of catenary.

Conversely, consider a saturated chain between $x$ and $y$ of length $n$

$$x = x_0 \lneq x_1 \lneq ... \lneq x_n = y$$

Then clearly $\operatorname{codim}(x_i, x_{i+1}) = 1$ for all $i = 0...n-1$. Therefore by repeatedly applying the relation we may show that

$$\operatorname{codim}(x, y) = n$$

and therefore every saturated chain between $x$ and $y$ has the same length.

When $\mathcal{G}$ is irreducible we may deduce the second formula by setting $x = \top$. Conversely we may see that

$$\operatorname{codim}(z, x) = \operatorname{codim}(z) - \operatorname{codim}(x) = \operatorname{codim}(z) - \operatorname{codim}(y) + \operatorname{codim}(y) - \operatorname{codim}(x) = \operatorname{codim}(z, y) + \operatorname{codim}(y, x)$$

■

**Lemma 2.5.10**
*Let $\mathcal{G}$ be a biequidimensional finite-dimensional poset. Then for $x \in \mathcal{G}$ we have*

*a) $\{x\}^{\downarrow}$ and $\{x\}^{\uparrow}$ are biequidimensional*
*b) $\dim(\mathcal{G}) = \dim(x) + \operatorname{codim}(x)$*
*c) If $x$ is maximal then $\dim(x) = \dim(\mathcal{G})$ and in particular $\mathcal{G}$ is equidimensional*
*d) If $x$ is minimal then $\operatorname{codim}(x) = \dim(\mathcal{G})$ and in particular $\mathcal{G}$ is equicodimensional*

*In particular $\mathcal{G}$ is quasi-biequidimensional.*

*Proof.* Consider a fixed maximal chain $C$ of $\{x\}^{\uparrow}$, necessarily containing $x$. Any maximal chain $C'$ of $\{x\}^{\downarrow}$ also contains $x$ and combines with $C$ to yield a maximal chain of $\mathcal{F}$. Whence $\ell(C') + \ell(C) = \dim(\mathcal{F})$ and $\{x\}^{\downarrow}$ is biequidimensional. By duality $\{x\}^{\uparrow}$ is biequidimensional and $\ell(C) = \operatorname{codim}(x)$ from which the formula follows.

If $x$ is maximal then clearly $\operatorname{codim}(x) = 0$, and similarly if $x$ is minimal then $\dim(x) = 0$, so the last two statements follow immediately. ■

**Proposition 2.5.11 (Equivalent Characterizations of Biequidimensionality)**
*Let $\mathcal{G}$ be a finite-dimensional poset. Then the following are equivalent*

*a) $\mathcal{G}$ is quasi-biequidimensional*
*b) $\mathcal{G}$ is catenary and for every maximal $x$ we have $\{x\}^{\downarrow}$ is equicodimensional*
*c) $\mathcal{G}$ satisfies $\dim(x) = \dim(y) + \operatorname{codim}(y, x)$ for $y \leq x$*
*d) $\mathcal{G}$ satisfies c) whenever $\operatorname{codim}(y, x) = 1$*

*Furthermore the following relationship holds*

$$\operatorname{codim}(y) = \operatorname{codim}(y, x) + \operatorname{codim}(x) \quad y \leq x$$

*Proof.* a) $\implies$ c). By assumption $\{x\}^{\downarrow}$ is biequidimensional. Then

$$\dim(x) = \dim\big(\{x\}^{\downarrow}\big) \overset{(2.5.10)}{=} \dim\big(y; \{x\}^{\downarrow}\big) + \operatorname{codim}\big(y; \{x\}^{\downarrow}\big) = \dim(y) + \operatorname{codim}(y, x)$$

For c) $\implies$ b). Suppose $z < y < x$ in $\mathcal{G}$ then by the codimension formula applied twice

$$\operatorname{codim}(z, x) = \dim(x) - \dim(z) = \dim(x) - \dim(y) + \dim(y) - \dim(z) = \operatorname{codim}(y, x) + \operatorname{codim}(z, y)$$

so by (2.5.9) $\mathcal{G}$ is catenary. Let $x$ be a maximal element and $z$ a minimal element of $\{x\}^{\downarrow}$, then by the codimension formula

$$\operatorname{codim}(z, x) = \dim(x) - \dim(z) = \dim(x)$$

© David Rufino CC BY-NC-ND 4.0

whence $\{x\}^{\downarrow}$ is equicodimensional.

b) $\implies$ a). Let $x$ be a maximal element and $C$ a maximal chain in $\{x\}^{\downarrow}$ with minimum element $y$ then, as $\mathcal{G}$ is catenary, we have $\ell(C) = \operatorname{codim}(y, x)$. As $\{x\}^{\downarrow}$ is equicodimensional we have $\operatorname{codim}(y, x) = \dim(x)$ which shows that $\{x\}^{\downarrow}$ is biequidimensional.

Clearly c) $\implies$ d). Conversely for d) $\implies$ a) consider a maximal chain in $\{x\}^{\downarrow}$

$$x_0 \lneq \ldots \lneq x_n = x$$

Then clearly $\operatorname{codim}(x_i, x_{i+1}) = 1$ whence by induction $\ell(C) = \dim(x)$.

The final statement follows from (2.5.9) because $\{x\}^{\downarrow}$ is irreducible and catenary. ∎

**Remark 2.5.12**
*This is a corrected version of EGA IV 14.3.3, as noted by Heinrich.*

**Corollary 2.5.13**
*When $\mathcal{F}$ is a quasi-biequidimensional Krull Lattice then we have the following codimension formulas for all $x \in \mathcal{J}(\mathcal{F})$ and $y \in \mathcal{F}$ such that $y \leq x$*

$$\dim(x) = \dim(y) + \operatorname{codim}(y, x)$$
$$\operatorname{codim}(y) = \operatorname{codim}(y, x) + \operatorname{codim}(x)$$

*Furthermore if $\mathcal{F}$ is biequidimensional then for all $y \in \mathcal{F}$*

$$\dim(\mathcal{F}) = \dim(y) + \operatorname{codim}(y)$$

*Proof.* The first two relations hold when $x, y \in \mathcal{J}(\mathcal{F})$ by (2.5.11). We may generalise this as follows.

For the first relation and fixed $x \in \mathcal{J}(\mathcal{F})$ we see $\dim(y_j)$ is maximal precisely when $\operatorname{codim}(y_j, x)$ is minimal. Therefore it holds for all $y \in \mathcal{F}$.

Similarly for the second relation we see that $\operatorname{codim}(y_j, x)$ is minimal precisely when $\operatorname{codim}(y_j)$ is minimal.

The final statement follows from the first relation by taking the supremum over all maximal elements $x$. ∎

**Corollary 2.5.14**
*Let $\mathcal{G}$ be an irreducible finite-dimensional poset. Then the following are equivalent*

*a) $\mathcal{G}$ is biequidimensional*
*b) $\mathcal{G}$ is quasi-biequidimensional*
*c) $\mathcal{G}$ is catenary and equicodimensional*
*d)* $\dim(x) = \dim(y) + \operatorname{codim}(y, x) \quad \forall y \leq x$
*e) $\mathcal{G}$ satisfies d) when* $\operatorname{codim}(y, x) = 1$

*Proof.* The equivalence follows from (2.5.11) by noting that an irreducible poset has only one maximal element and is in particular equidimensional. ∎

**Proposition 2.5.15**
*Let $\mathcal{F}$ be a quasi-biequidimensional Krull Lattice. Then for all $x \in \mathcal{F}$ we have $\{x\}^{\downarrow}$ is a quasi-biequidimensional Krull Lattice.*

*Proof.* Consider $y \in \mathcal{J}(\{x\}^{\downarrow}) \overset{(2.4.3)}{=} \mathcal{J}(\mathcal{F}) \cap \{x\}^{\downarrow}$. Then by definition $\{y\}^{\downarrow}$ is biequidimensional, and so $\{x\}^{\downarrow}$ is quasi-biequidimensional. ∎

© David Rufino CC BY-NC-ND 4.0

# 2.6 Category Theory

## 2.6.1 Categories

**Definition 2.6.1 (Category)**
*A (locally small)* ***category*** $\mathcal{C}$ *consists of*

- *a class* $\mathrm{ob}(\mathcal{C})$ *of objects*
- *for every pair of objects* $X, Y \in \mathrm{ob}(\mathcal{C})$ *a* set *of* ***morphisms*** $\mathrm{Mor}(X, Y)$
- *for every three objects* $X, Y, Z$ *a* ***law of composition***

$$\begin{aligned} \mathrm{Mor}(X,Y) \times \mathrm{Mor}(Y,Z) &\to \mathrm{Mor}(X,Z) \\ (g,f) &\to g \circ f \end{aligned}$$

*such that the following conditions hold*

- $h \circ (g \circ f) = (h \circ g) \circ f$ *(**associativity**)*
- *There exists* $1_X \in \mathrm{Mor}(X,X)$ *such that* $1_X \circ f = f$ *and* $g \circ 1_X = g$.

**Example 2.6.2**
*The category of sets is* **Set** *with maps in the usual way. Note associativity is automatically satisfied.*

**Example 2.6.3 ($n$-pointed category)**
*Given a category* $\mathcal{C}$ *where objects are sets, we may consider the pointed category* $(\mathcal{C}, \star^n)$ *consisting of pairs* $(A, a)$ *where* $A \in \mathrm{ob}(\mathcal{C})$ *and* $a \in A^n$. *We consider only morphisms* $f : A \to B$ *such that* $f(a_i) = b_i$.

**Definition 2.6.4 (Opposite Category)**
*Given a category* $\mathcal{C}$ *the* ***opposite category*** *is denoted* $\mathcal{C}^{op}$ *and given by*

- *The same class of objects* $\mathrm{ob}(\mathcal{C}^{op}) = \mathrm{ob}(\mathcal{C})$
- *For every pair of objects* $X, Y \in \mathrm{ob}(\mathcal{C})$ *the morphisms are reversed*

$$\mathrm{Mor}^{op}(X,Y) := \mathrm{Mor}(Y,X)$$

- *The law of composition is reversed*

$$\begin{aligned} \mathrm{Mor}^{op}(X,Y) \times \mathrm{Mor}^{op}(Y,Z) &\to \mathrm{Mor}^{op}(X,Z) \\ (g,f) &\to f \circ g \end{aligned}$$

**Example 2.6.5**
*The polynomial ring* $A[X]$ *is an initial object in the category of pointed* $A$*-algebras.*

**Definition 2.6.6 (Isomorphism)**
*A morphism* $f : X \to Y$ *is an* ***isomorphism*** *if there exists* $g : Y \to X$ *such that*

$$g \circ f = 1_X \quad \text{and} \quad f \circ g = 1_Y.$$

**Definition 2.6.7 (Functor)**
*A* ***covariant functor*** $F : \mathcal{C} \to \mathcal{D}$ *consists of a mapping of objects*

$$F : \mathrm{ob}(\mathcal{C}) \to \mathrm{ob}(\mathcal{D})$$

*together with a mapping of morphisms*

$$F(-) : \mathrm{Mor}(X,Y) \to \mathrm{Mor}(F(X), F(Y))$$

*which satisfies*

- $F(1_X) = 1_{F(X)}$
- $F(f \circ g) = F(f) \circ F(g)$

*A* ***contravariant functor*** $\mathcal{C} \to \mathcal{D}$ *is equivalent to a covariant functor on the opposite category* $\mathcal{C}^{op} \to \mathcal{D}$, *namely where arrows are reversed.*

**Definition 2.6.8 (Full and faithful)**
*A functor* $F$ *is said to be*

© David Rufino CC BY-NC-ND 4.0

- ***Faithful*** *if* $F(-)$ *is injective.*
- ***Full*** *if* $F(-)$ *is surjective.*

**Definition 2.6.9 (Concrete Category)**
*A* ***concrete category*** *is a pair* $(\mathcal{C}, U)$ *where* $\mathcal{C}$ *and a "****forgetful functor****"* $U : \mathcal{C} \to \mathbf{Set}$ *which is faithful.*

**Example 2.6.10 (Forgetful Functor)**
*The category of groups (resp. rings, modules, ...) is a concrete category in the obvious way.*

**Definition 2.6.11 (Mor functor)**
*For any objects* $X, Y, Z \in \mathrm{ob}(\mathcal{C})$*, there is a canonical covariant functor*

$$\begin{aligned} \mathrm{Mor}(X, -) : \mathcal{C} &\longrightarrow \mathbf{Set} \\ Y &\longrightarrow \mathrm{Mor}(X, Y) \end{aligned}$$

*which acts on a morphism* $f : Y \to Z$ *by*

$$\begin{aligned} \mathrm{Mor}(X, f) : \mathrm{Mor}(X, Y) &\to \mathrm{Mor}(X, Z) \\ g &\to f \circ g \end{aligned}$$

*It's a functor precisely because composition of functions is associative. Similarly there's a canonical contravariant functor* $\mathrm{Mor}(-, Y)$*.*

**Definition 2.6.12 (Natural Transformation)**
*Let* $F, G : \mathcal{C} \to \mathcal{D}$ *be covariant functors. A* ***natural transformation*** $\eta : F \Rightarrow G$ *consists of a family of morphisms*

$$\eta_Z : F(Z) \to G(Z) \quad Z \in \mathrm{ob}(\mathcal{C})$$

*such that the following diagram commutes for all* $f : Z \to Z'$

$$\begin{array}{ccc} F(Z) & \xrightarrow{\eta_Z} & G(Z) \\ \Big\downarrow{\scriptstyle F(f)} & & \Big\downarrow{\scriptstyle G(f)} \\ F(Z') & \xrightarrow{\eta_{Z'}} & G(Z') \end{array}$$

**Definition 2.6.13 (Natural isomorphism)**
*A natural transformation* $\eta : F \Rightarrow G$ *is a natural isomorphism if* $\eta_Z$ *is an isomorphism for all* $Z \in \mathcal{C}$*.*

**Definition 2.6.14 (Functor Category)**
*Let* $I$ *be a small category and* $\mathcal{D}$ *a category. The* ***functor category*** $\mathcal{D}^I$ *has*

- *objects: functors* $F : I \to \mathcal{D}$*;*
- *morphisms* $F \to G$*: natural transformations* $\eta : F \Rightarrow G$ *(2.6.12);*
- *composition: for* $\eta : F \Rightarrow G$ *and* $\theta : G \Rightarrow H$*,* $(\theta \circ \eta)_Z := \theta_Z \circ \eta_Z$ *for* $Z \in \mathrm{ob}(I)$ *– again a natural transformation, since for* $f : Z \to Z'$ *in* $I$*, using naturality of* $\theta$ *then of* $\eta$*,*

$$H(f) \circ \theta_Z \circ \eta_Z = \theta_{Z'} \circ G(f) \circ \eta_Z = \theta_{Z'} \circ \eta_{Z'} \circ F(f);$$

- *identities:* $1_F$*, with* $(1_F)_Z := 1_{F(Z)}$ *(trivially natural).*

*Associativity and the identity laws hold pointwise at each* $Z \in \mathrm{ob}(I)$*, being inherited from those of* $\mathcal{D}$*; so* $\mathcal{D}^I$ *is indeed a category.*

**Definition 2.6.15**
*We say* $\mathcal{C}' \subset \mathcal{C}$ *is a* ***subcategory*** *if*

- $\mathrm{ob}(\mathcal{C}') \subseteq \mathrm{ob}(\mathcal{C})$
- $\mathrm{Mor}_{\mathcal{C}'}(Z, W) \subseteq \mathrm{Mor}_{\mathcal{C}}(Z, W)$
- *Composition agrees when it is well-defined*

*We say* $\mathcal{C}'$ *is a* ***full subcategory*** *if additionally* $\mathrm{Mor}_{\mathcal{C}'}(Z, W) = \mathrm{Mor}_{\mathcal{C}}(Z, W)$*.*

© David Rufino CC BY-NC-ND 4.0

### 2.6.2 Comma and Slice Categories

**Definition 2.6.16 (Comma Category)**
*Let $F : \mathcal{C} \to \mathcal{E}$ and $G : \mathcal{D} \to \mathcal{E}$ be functors. The **comma category** $(F \downarrow G)$ is the category with*

- *objects: triples $(C, D, f)$ where $C \in \mathcal{C}$, $D \in \mathcal{D}$, and $f : F(C) \to G(D)$ is a morphism in $\mathcal{E}$*
- *morphisms: a morphism $(C, D, f) \to (C', D', f')$ is a pair $(g, h)$ where $g : C \to C'$ is a morphism in $\mathcal{C}$ and $h : D \to D'$ is a morphism in $\mathcal{D}$, such that*

$$G(h) \circ f = f' \circ F(g)$$

*with composition $(g', h') \circ (g, h) \coloneqq (g' \circ g, h' \circ h)$ and identities $1_{(C,D,f)} \coloneqq (1_C, 1_D)$, inherited from $\mathcal{C}$ and $\mathcal{D}$.*

**Definition 2.6.17 (Slice Category)**
*Let $\mathcal{C}$ be a category and $X \in \mathcal{C}$, and let $\underline{X} : \mathbf{1} \to \mathcal{C}$ be the functor with $\underline{X}(\star) = X$. The **slice category** $\mathcal{C}/X$ (also denoted $\mathcal{C} \downarrow X$) is the comma category $(\mathrm{id}_{\mathcal{C}} \downarrow \underline{X})$ (2.6.16).*

*Explicitly its objects are pairs $(Y, f)$ where $Y \in \mathrm{ob}(\mathcal{C})$ and $f : Y \to X$ is a morphism in $\mathcal{C}$, and a morphism $(Y, f) \to (Y', f')$ is a morphism $g : Y \to Y'$ in $\mathcal{C}$ such that $f' \circ g = f$.*

*Dually the **coslice category** $X/\mathcal{C}$ (also denoted $X \downarrow \mathcal{C}$) is the comma category $(\underline{X} \downarrow \mathrm{id}_{\mathcal{C}})$, with objects pairs $(Y, f)$ where $Y \in \mathrm{ob}(\mathcal{C})$ and $f : X \to Y$ is a morphism in $\mathcal{C}$, and a morphism $(Y, f) \to (Y', f')$ is a morphism $g : Y \to Y'$ in $\mathcal{C}$ such that $g \circ f = f'$.*

### 2.6.3 Product Categories and Bifunctors

**Definition 2.6.18 (Product Category)**
*Given two categories $\mathcal{C}$, $\mathcal{D}$ we may construct the product category $\mathcal{C} \times \mathcal{D}$ as follows*

- *The objects are given by pairs*

$$\mathrm{ob}(\mathcal{C} \times \mathcal{D}) \coloneqq \mathrm{ob}(\mathcal{C}) \times \mathrm{ob}(\mathcal{D})$$

- *The morphisms are given by pairs*

$$\mathrm{Mor}((C, C'), (D, D')) \coloneqq \mathrm{Mor}(C, C') \times \mathrm{Mor}(D, D')$$

  *Concretely given $f : C \to C'$ and $g : D \to D'$ write $(f, g) : (C, D) \to (C', D')$*
- *The law of composition is determined by*

$$(f, g) \circ (h, k) \coloneqq (f \circ h, g \circ k)$$

*It's clear that the law of composition is associative and the identity morphism for $(C, D)$ is $(1_C, 1_D)$. Furthermore we observe the following property*

$$(f, 1_D) \circ (1_C, g) = (f, g) = (1_C, g) \circ (f, 1_D) \tag{*}$$

**Definition 2.6.19 (Bifunctor)**
*A functor on a product category, $F : \mathcal{C} \times \mathcal{D} \to \mathcal{E}$ is termed a **bifunctor**. $F$ induces a family of functors*

- *$F(C, -) : \mathcal{D} \to \mathcal{E}$ given by*

$$F(C, g) \coloneqq F(1_C, g)$$

- *$F(-, D) : \mathcal{C} \to \mathcal{E}$ given by*

$$F(f, D) \coloneqq F(f, 1_D)$$

*which satisfy the compatibility conditions*

$$\begin{array}{ccc} F(C, D) & \xrightarrow{F(f, D)} & F(C', D) \\ {\scriptstyle F(C, g)}\downarrow & \searrow^{F(f, g)} & \downarrow{\scriptstyle F(C', g)} \\ F(C, D') & \xrightarrow[F(f, D')]{} & F(C', D') \end{array}$$

© David Rufino CC BY-NC-ND 4.0

*by applying $F$ to (*).*

**Proposition 2.6.20 (Reconstruct Bifunctor)**
*Consider a family of functors $\{F_L(C) : \mathcal{D} \to \mathcal{E}\}_{C \in \mathrm{ob}(\mathcal{C})}$ and $\{F_R(D) : \mathcal{C} \to \mathcal{E}\}_{D \in \mathrm{ob}(\mathcal{D})}$. Suppose the following conditions hold*

- $F_L(C)(D) = F_R(D)(C)$
- $F_L(C')(g) \circ F_R(D)(f) = F_R(D')(f) \circ F_L(C)(g)$

*then these determine a well-defined bifunctor given by*

$$F(f \times g) \coloneqq F_L(C')(g) \circ F_R(D)(f)$$

**Proposition 2.6.21**
*Let $F : \mathcal{C} \times \mathcal{D} \to \mathcal{E}$ be a bifunctor. For $f : C \to C'$ then there is a natural transformation*

$$F(f, -) : F(C, -) \Rightarrow F(C', -)$$

*given by $F(f, -)_D \coloneqq F(f, 1_D)$. Similarly for $g : D \to D'$ then there is a natural transformation*

$$F(-, g) : F(-, D) \Rightarrow F(-, D')$$

*Proof.* The naturality condition is immediate from (*) and the commutative diagram in (2.6.19). ∎

**Proposition 2.6.22 (Criteria for Natural Transformation)**
*Let $F, G : \mathcal{C} \times \mathcal{D} \to \mathcal{E}$ be two bifunctors and suppose we have a family of natural transformations*

$$\eta_C : F(C, -) \Rightarrow G(C, -)$$

*for all $C \in \mathcal{C}$. Then the following are equivalent*

*a) $\eta : F(-, -) \Rightarrow G(-, -)$ is a natural transformation*
*b) The following diagram commutes for all $D \in \mathcal{D}$ and $f : C \to C'$*

$$\begin{array}{ccc} F(C, D) & \xrightarrow{\eta_{C,D}} & G(C, D) \\ {\scriptstyle F(f, 1_D)}\downarrow & & \downarrow{\scriptstyle G(f, 1_D)} \\ F(C', D) & \xrightarrow{\eta_{C',D}} & G(C', D) \end{array}$$

*Proof.* a) $\implies$ b) This diagram is a special case of the naturality condition.

b) $\implies$ a) Suppose $f : C \to C'$ and $g : D \to D'$ then we require to show that

$$G(f, g)\big(\eta_{C,D}(\phi)\big) = \eta_{C',D'}(F(f, g)(\phi))$$

However

$$\begin{aligned} G(f, g)\big(\eta_{C,D}(\phi)\big) &= G(f, 1_{D'})\big(G(1_C, g)\big(\eta_{C,D}(\phi)\big)\big) \\ &= G(f, 1_{D'})\eta_{C,D'}(F(1_C, g)(\phi)) \\ &= \eta_{C',D'}(F(f, 1_{D'})(F(1_C, g)(\phi))) \\ &= \eta_{C',D'}(F(f, g)(\phi)) \end{aligned}$$

where we have used that $\eta_{C,D}$ is natural in $D$ and $C$ individually. ∎

**Example 2.6.23 (Mor is a bifunctor)**
*The canonical example is the following, given any locally small category $\mathcal{C}$, we have a bifunctor*

$$\mathrm{Mor} : \mathcal{C}^{op} \times \mathcal{C} \to \mathbf{Set}$$

*given by the set of morphisms. The action on morphisms is given by*

$$\mathrm{Mor}(f, g)(\phi) = g \circ \phi \circ f$$

*and we verify by associativity of $\mathcal{C}$ that*

$$\mathrm{Mor}((f, g) \circ (h, k)) = \mathrm{Mor}(f, g) \circ \mathrm{Mor}(h, k)$$

© David Rufino CC BY-NC-ND 4.0

*and it's clear that the commutativity condition in (*) is satisfied.*

**Example 2.6.24 (Concrete bifunctors)**
*Let $\mathcal{C}$, $\mathcal{D}$ be concrete categories and consider the following bifunctor*

$$F : \mathcal{D}^{op} \times \mathcal{C} \to \mathbf{Set}$$
$$F(D, C) := \{\phi : D \to C \mid P(\phi)\}$$

*where $P(-)$ is some predicate such that $P(\phi) \implies P(g \circ \phi \circ f)$.*

## 2.6.4 Equivalence of categories

**Definition 2.6.25 (Equivalence of categories)**
*Let $\mathcal{C}$, $\mathcal{D}$ be categories. An **equivalence of categories** consists of a pair of functors (either both covariant or both contravariant)*

$$\mathcal{C} \underset{G}{\overset{F}{\rightleftarrows}} \mathcal{D}$$

*together with natural isomorphisms*

$$\eta : \mathrm{id}_{\mathcal{C}} \Rightarrow GF$$

$$\epsilon : FG \Rightarrow \mathrm{id}_{\mathcal{D}}$$

*We say $F$ is an equivalence of categories if there exists some $G$ satisfying these conditions.*

**Definition 2.6.26**
*We say a functor $F : \mathcal{C} \to \mathcal{D}$ is **essentially surjective** if for all $D \in \mathcal{D}$ there exists $C \in \mathcal{C}$ such that $F(C)$ is isomorphic to $D$.*

**Lemma 2.6.27**
*Let $F : \mathcal{C} \to \mathcal{D}$, $G : \mathcal{D} \to \mathcal{C}$ be functors.*

*If there exists a natural isomorphism $\eta : \mathrm{id}_{\mathcal{C}} \Rightarrow GF$ then $F$ is faithful. Explicitly $F(-)$ has a left-inverse given by*

$$g \to \eta_{C'}^{-1} \circ G(g) \circ \eta_C$$

*Furthermore $GF(\eta_C) = \eta_{GF(C)}$.*

*Proof.* Consider the sequence of maps

$$\mathrm{Mor}(C, C') \overset{F(-)}{\longrightarrow} \mathrm{Mor}(F(C), F(C')) \overset{G(-)}{\longrightarrow} \mathrm{Mor}(GF(C), GF(C')) \overset{\mathrm{Mor}\left(\eta_C, \eta_{C'}^{-1}\right)}{\longrightarrow} \mathrm{Mor}(C, C')$$

Note that the composite of this map is given by

$$f \to \eta_{C'}^{-1} \circ GF(f) \circ \eta_C = \eta_{C'}^{-1} \circ \eta_{C'} \circ f = f$$

in other words $F(-)$ has a left inverse and therefore $F$ is faithful.

Note by naturality we have $GF(\eta_C) \circ \eta_C = \eta_{GF(C)} \circ \eta_C$. Since $\eta_C$ is an isomorphism we may cancel to find $GF(\eta_C) = \eta_{GF(C)}$. ∎

**Proposition 2.6.28 (Equivalence is full and faithful)**
*Let $F : \mathcal{C} \to \mathcal{D}$ be a functor then the following are equivalent*

- *$F$ is full, faithful and essentially surjective (2.6.26)*
- *$F$ is an equivalence of categories (2.6.25)*

*In other words $F(-)$ is bijective and hence has a two-sided inverse. Explicitly it is given by*

$$\mathrm{Mor}(C, C') \longleftrightarrow \mathrm{Mor}(F(C), F(C'))$$
$$f \to F(f)$$
$$\eta_{C'}^{-1} \circ G(g) \circ \eta_C \leftarrow g$$

*Proof.* We prove only the second implies the first. By assumption there is an equivalence with $G$ and by the previous Lemma both $F$ and $G$ are faithful by considering $\eta$ and $\epsilon^{-1}$ in turn. Further the given map is already shown to be a left inverse. We claim it's also a right inverse, for consider

© David Rufino CC BY-NC-ND 4.0

$$g' := F\left(\eta_{C'}^{-1}\right) \circ FG(g) \circ F(\eta_C).$$

We claim that $G(g') = G(g)$. As $G$ is faithful this would imply $g' = g$ and the given map is a right inverse as required. Observe

$$G(g') = GF\left(\eta_{C'}^{-1}\right) \circ GFG(g) \circ GF(\eta_C) = \eta_{GF(C')}^{-1} \circ GFG(g) \circ \eta_{GF(C)} = \eta_{GF(C')}^{-1} \circ \eta_{GF(C')} \circ G(g) = G(g)$$

where we have used the result that $GF(\eta_C) = \eta_{GF(C)}$. Since the maps are mutual inverses we see that $F$ is full and faithful as required.

Given $D \in \mathcal{D}$ then $F(G(D))$ is isomorphic to $D$ via $\epsilon$ so $F$ is essentially surjective. ■

**Proposition 2.6.29 (Duality)**
*Let $(-)^\star : \mathcal{C} \to \mathcal{C}$ be a contravariant functor such that there is a natural isomorphism*

$$\eta : \mathrm{id}_{\mathcal{C}} \Rightarrow (-)^{\star\star}$$

*then $(-)^\star$ is an equivalence of categories and in particular full and faithful.*

*Proof.* Define $\epsilon = \eta^{-1}$ to determine the equivalence of categories. By the previous result then $(-)^\star$ is full and faithful.■

## 2.6.5 Properties of Morphisms

**Definition 2.6.30 (Injective, Surjective and Bijective)**
*Let $(\mathcal{C}, U)$ be a concrete category (2.6.9) and $f : X \to Y$ a morphism. Then we say*

- *$f$ is **injective** if $U(f)$ is injective*
- *$f$ is **surjective** if $U(f)$ is surjective*

**Remark 2.6.31**
*Note if $f$ is both surjective and injective it need not be an isomorphism.*

The concepts of monic/split-monic, epic/split-epic, iso generalize the notion of injective, surjective and bijective to general categories as we shall see.

**Definition 2.6.32 (Monomorphism / Epimorphism)**
*A morphism $f : X \to Y$ is said to be a **monomorphism** (or **monic**) if*

$$f \circ g_1 = f \circ g_2 \Longrightarrow g_1 = g_2$$

*for all $g_1, g_2 : Z \to X$.*

*Dually, $f$ is said to be an **epimorphism** (or **epic**) if*

$$g_1 \circ f = g_2 \circ f \Longrightarrow g_1 = g_2$$

*for all $g_1, g_2 : Y \to Z$.*

**Remark 2.6.33 (Monic/Epic Duality)**
*A morphism $f : X \to Y$ of $\mathcal{C}$ is monic if and only if $f^{op} : Y \to X$ is epic in $\mathcal{C}^{op}$, and dually $f$ is epic if and only if $f^{op}$ is monic in $\mathcal{C}^{op}$ (2.6.4): composition in $\mathcal{C}^{op}$ reverses the order of composition in $\mathcal{C}$, so the defining cancellation property of monic in $\mathcal{C}$ is precisely the defining cancellation property of epic in $\mathcal{C}^{op}$, and vice versa.*

**Definition 2.6.34 (Split-monic / Split-epic)**
*A morphism $f : X \to Y$ is **split-monic** if it has a left inverse, $g : Y \to X$*

$$g \circ f = 1_X$$

*We say $g$ is a **section** of $f$.*

*Dually, $f$ is **split-epic** if it has a right inverse, $g : Y \to X$*

$$f \circ g = 1_Y$$

*We say $g$ is a **retraction** of $f$.*

**Proposition 2.6.35 (Split Monic⇒Monic)**
*For a general category $\mathcal{C}$ we have*

- *split-monic $\Longrightarrow$ monic*

© David Rufino CC BY-NC-ND 4.0

- *split-epic* $\implies$ *epic*

We say an **isomorphism** is a morphism with a two-sided inverse. We can refine the criteria for $f$ to be an isomorphism using the notions just defined

**Proposition 2.6.36 (Isomorphism Criteria)**
*Let $f : X \to Y$ be a morphism. Then the following are equivalent*

*a) $f$ is an **isomorphism***
*b) $f$ is both **split-epic** and **split-monic***
*c) $f$ is **split-epic** and **monic***
*d) $f$ is **split-monic** and **epic***

*In this case a morphism $g$ is a retraction if and only if it is a section. And such a $g$ is unique, so we denote it by $f^{-1}$*

*Proof.* This is mostly formal

a) $\implies$ b) Clear.

b) $\implies$ c,d) This follows from (2.6.35).

c) $\implies$ b) Suppose $g$ is a retraction of $f$, that is $fg = 1_Y$. Then $f(gf) = (fg)f = 1_Y \circ f = f = f \circ 1_X$. As $f$ is monic we conclude that $gf = 1_X$ and $g$ is a section of $f$.

d) $\implies$ b) Analogous

b) $\implies$ a) We've shown that any retraction is a section and vice-versa. Furthermore by monic/epic-ness a retraction or section is unique. ■

**Lemma 2.6.37 (Factors of an Isomorphism)**
*Let $f : X \to Y$ be an isomorphism.*

*a) If $f = m \circ g$ for some monomorphism $m$, then $m$ and $g$ are both isomorphisms.*
*b) If $f = h \circ e$ for some epimorphism $e$, then $h$ and $e$ are both isomorphisms.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

a) Since $f$ is an isomorphism, $m \circ (g \circ f^{-1}) = f \circ f^{-1} = 1_Y$, so $m$ is split-epic. As $m$ is also monic, (2.6.36) gives that $m$ is an isomorphism, and so $g = m^{-1} \circ f$ is also an isomorphism, being a composite of isomorphisms.

b) Dually, since $f$ is an isomorphism, $(f^{-1} \circ h) \circ e = f^{-1} \circ f = 1_X$, so $e$ is split-monic. As $e$ is also epic, (2.6.36) gives that $e$ is an isomorphism, and so $h = f \circ e^{-1}$ is also an isomorphism. ■

**Proposition 2.6.38**
*For the category* **Set** *we have*

- *split-monic* $\iff$ *monic* $\iff$ *injective*
- *split-epic* $\iff$ *epic* $\iff$ *surjective*
- *isomorphism* $\iff$ *bijective*

**Definition 2.6.39 (Balanced Category)**
*A category $\mathcal{C}$ is **balanced** if every morphism which is both monic and epic is an isomorphism.*

**Example 2.6.40**
**Set** *is balanced: by (2.6.38), the monic and epic morphisms are precisely the injective and surjective ones, so a morphism which is both is bijective, hence an isomorphism. The category of groups is balanced for the same reason, a bijective group homomorphism has a group homomorphism as its (set-theoretic) inverse.*

*Not every category is balanced. In* **Top** *the map $[0, 2\pi) \to S^1$, $t \mapsto e^{it}$ is continuous and bijective, hence monic and epic, but its inverse is not continuous, so it is not an isomorphism. Likewise in* **Rng** *the inclusion $\mathbb{Z} \hookrightarrow \mathbb{Q}$ is monic and epic (any two ring homomorphisms out of $\mathbb{Q}$ agreeing on $\mathbb{Z}$ agree everywhere) but is not an isomorphism.*

**Definition 2.6.41 (Preserves/Reflects)**
*Let $\mathcal{P}$ be a property of morphisms and $F : \mathcal{C} \to \mathcal{D}$ be a functor then we say*

- *$F$ **preserves** $\mathcal{P}$ if ($f$ satisfies $\mathcal{P} \implies F(f)$ satisfies $\mathcal{P}$)*
- *$F$ **reflects** $\mathcal{P}$ if ($F(f)$ satisfies $\mathcal{P} \implies f$ satisfies $\mathcal{P}$)*

© David Rufino CC BY-NC-ND 4.0

**Proposition 2.6.42**
*Let $F : \mathcal{C} \to \mathcal{D}$ be a covariant functor then*

- *$F$ preserves split-monic, split-epic and iso morphisms.*

*If in addition $F$ is faithful then*

- *$F$ reflects monic and epic morphisms.*

*and if $F$ is full and faithful then*

- *$F$ reflects split-epic, split-monic and isomorphisms.*

*Similar statements apply when $F$ is contravariant.*

*Proof.* The first statement is easy, for example if $gf = 1_X$ then $F(g) \circ F(f) = 1_{F(X)}$.

Suppose $F$ is faithful, $F(f)$ is monic and $fg_1 = fg_2$. Then $F(f)F(g_1) = F(f)F(g_2) \implies F(g_1) = F(g_2)$ by assumption. As $F$ is faithful $g_1 = g_2$ as required. The other statement is similar.

Suppose $F$ is full and faithful and $F(f)$ is split-monic. Then $hF(f) = 1_{F(X)}$. As $F$ is full $h = F(g)$ and $1_{F(X)} = F(gf)$. As $F$ is faithful then $gf = 1_X$. the other statements are similar. ∎

**Proposition 2.6.43**
*Let $(\mathcal{C}, U)$ be a concrete category (2.6.9) then*

- *$f$ split-monic $\implies$ $f$ injective $\implies$ $f$ monic*
- *$f$ split-epic $\implies$ $f$ surjective $\implies$ $f$ epic*
- *$f$ isomorphism $\implies$ $f$ bijective*

*Proof.* Suppose $f$ is split-monic, then $U(f)$ is split-monic by (2.6.42) and so by (2.6.38) $U(f)$ is injective.

Suppose $U(f)$ injective, then by (2.6.38) $U(f)$ is monic. By (2.6.42) $U$ reflects monics and so $f$ is monic.

The other statements are similar. ∎

We can restate the criteria for split-epic/split-monic

**Proposition 2.6.44**
*Let $f : X \to Y$ be a morphism then*

- *$f$ is split-monic if and only if $\mathrm{Mor}(f, Z)$ is surjective for all $Z \in \mathcal{C}$*
- *$f$ is epic if and only if $\mathrm{Mor}(f, Z)$ is injective for all $Z \in \mathcal{C}$*

*dually*

- *$f$ is split-epic if and only if $\mathrm{Mor}(Z, f)$ is surjective for all $Z \in \mathcal{C}$*
- *$f$ is monic if and only if $\mathrm{Mor}(Z, f)$ is injective for all $Z \in \mathcal{C}$*

*Proof.* $f$ is epic (resp. monic) iff $\mathrm{Mor}(f, Z)$ (resp. $\mathrm{Mor}(Z, f)$) is injective precisely by the definitions.

Suppose $f$ is split-monic, then $gf = 1_X \implies (hg)f = h$ for any $h$. That is $\mathrm{Mor}(f, Z)$ is surjective. Conversely if it's surjective then let $Z = X$ and choose $g$ such that $gf = \mathrm{Mor}(f, X)(g) = 1_X$.

A similar statement follows dually for $f$ split-epic, and $f$ monic. ∎

**Corollary 2.6.45 (Isomorphism Criteria)**
*Let $f : X \to Y$ be a morphism then TFAE*

- *$f$ is an isomorphism*
- *$\mathrm{Mor}(f, Z)$ is bijective for all $Z \in \mathcal{C}$*
- *$\mathrm{Mor}(Z, f)$ is bijective for all $Z \in \mathcal{C}$*

*Proof.* This follows from combining (2.6.44) with (2.6.36). ∎

**Definition 2.6.46 (Conservative)**
*We say a functor $F : \mathcal{C} \to \mathcal{D}$ is **conservative** if it reflects isomorphisms.*

*We say a concrete category $(\mathcal{C}, U)$ is **conservative** if the functor $U : \mathcal{C} \to \mathbf{Set}$ is conservative.*

© David Rufino CC BY-NC-ND 4.0

**Proposition 2.6.47**
*Let $(\mathcal{C}, U)$ be a conservative concrete category which admits directed limits. Then the following are equivalent*

*a) $U$ preserves directed colimits*
*b) $U$ reflects directed colimits*

### 2.6.6 Pointed Categories

**Definition 2.6.48 (Initial object)**
*An **initial object** of a category $\mathcal{C}$ is an object $\varnothing$ such that for all objects $X$*

$$\mathrm{Mor}(\varnothing, X) = \{\eta_{\varnothing}^{X}\}$$

*consists of a single element.*

*In this case an initial object is unique up to unique isomorphism (2.6.49).*

**Proposition 2.6.49 (Criteria to be Initial)**
*Let $\varnothing$ be an initial object and $X \in \mathcal{C}$. Then the following are equivalent*

*a) $X$ is an initial object*
*b) there is an isomorphism $\phi : X \xrightarrow{\sim} \varnothing$*
*c) there exists a morphism $\phi : X \to \varnothing$ and only one endomorphism $1_X : X \to X$*

*In this case the isomorphism is unique.*

*Proof.* a) $\implies$ c) by definition

c) $\implies$ b) By assumption $\eta_{\varnothing}^{X} \circ \phi = 1_X$, and by definition of initial $\phi \circ \eta_{\varnothing}^{X} = 1_{\varnothing}$.

b) $\implies$ a) We have a bijection

$$\begin{aligned} \mathrm{Mor}(\varnothing, Y) &\longleftrightarrow \mathrm{Mor}(X, Y) \\ f &\to f \circ \phi \\ g \circ \phi^{-1} &\leftarrow g \end{aligned}$$

so as $\varnothing$ is initial, $\mathrm{Mor}(\varnothing, Y)$ is a singleton for every $Y$, hence so is $\mathrm{Mor}(X, Y)$, that is $X$ is initial. This closes the cycle a) $\implies$ c) $\implies$ b) $\implies$ a), so all three are equivalent.

Finally, since $X$ is then initial, $\mathrm{Mor}(X, \varnothing)$ is a singleton, so there is at most one morphism $X \to \varnothing$; in particular the isomorphism $\phi$ is unique. ∎

**Definition 2.6.50 (Terminal Object)**
*A **terminal object** of a category $\mathcal{C}$ is an object $\star$ such that for all objects $X \in \mathcal{C}$ we have*

$$\mathrm{Mor}(X, \star) = \{!_X\}$$

*is a singleton.*

*In this case a terminal object is unique up to unique isomorphism (2.6.51).*

**Proposition 2.6.51 (Criteria to be Terminal)**
*Let $\star$ be a terminal object and $X \in \mathcal{C}$. Then the following are equivalent*

*a) $X$ is a terminal object*
*b) there is an isomorphism $\phi : \star \xrightarrow{\sim} X$*
*c) there exists a morphism $\phi : \star \to X$ and only one endomorphism $1_X : X \to X$*

*In this case the isomorphism is unique.*

*Proof.* By (2.6.4), $\mathrm{Mor}_{\mathcal{C}^{\mathrm{op}}}(\varnothing, X) = \mathrm{Mor}_{\mathcal{C}}(X, \varnothing)$ for all $X$, so $\star$ is a terminal object of $\mathcal{C}$ if and only if $\star$ is an initial object of $\mathcal{C}^{\mathrm{op}}$; likewise a morphism $\star \to X$ (resp. isomorphism, resp. endomorphism of $X$) in $\mathcal{C}$ is exactly a morphism $X \to \star$ (resp. isomorphism, resp. endomorphism of $X$) in $\mathcal{C}^{\mathrm{op}}$. So applying (2.6.49) in $\mathcal{C}^{\mathrm{op}}$, with $\varnothing := \star$, gives exactly the stated equivalence of a), b), c), together with the uniqueness of the isomorphism. ∎

**Lemma 2.6.52 (Equivalences of Categories Preserve Terminal Objects)**
*Let $F : \mathcal{C} \to \mathcal{D}$ be an equivalence of categories, with $G$ and natural isomorphisms $\eta : \mathrm{id}_{\mathcal{C}} \Rightarrow GF$, $\epsilon : FG \Rightarrow \mathrm{id}_{\mathcal{D}}$ as in (2.6.25). Then $T \in \mathcal{C}$ is a terminal object of $\mathcal{C}$ if and only if $F(T)$ is a terminal object of $\mathcal{D}$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

© David Rufino CC BY-NC-ND 4.0

Suppose $T$ is terminal. By (2.6.28), $F$ is full and faithful, so for every $C \in \mathcal{C}$ the map $\mathrm{Mor}(C,T) \to \mathrm{Mor}(F(C),F(T))$, $f \mapsto F(f)$, is a bijection; as $T$ is terminal $\mathrm{Mor}(C,T)$ is a singleton, so $\mathrm{Mor}(F(C),F(T))$ is too. Now let $Y \in \mathcal{D}$ be arbitrary and set $C \coloneqq G(Y)$; the isomorphism $\epsilon_Y : F(C) \to Y$ gives a bijection $\mathrm{Mor}(F(C),F(T)) \to \mathrm{Mor}(Y,F(T))$, $g \mapsto g \circ \epsilon_Y^{-1}$, so combining with the above, $\mathrm{Mor}(Y,F(T))$ is a singleton. As $Y \in \mathcal{D}$ was arbitrary, $F(T)$ is terminal.

Conversely, $G$ is itself an equivalence of categories (with $F$, $\epsilon^{-1} : \mathrm{id}_{\mathcal{D}} \Rightarrow FG$ and $\eta^{-1} : GF \Rightarrow \mathrm{id}_{\mathcal{C}}$ playing the roles of $F$, $\eta$, $\epsilon$ in (2.6.25)), so if $F(T)$ is terminal then applying the above (with $F,G,T$ replaced by $G,F,F(T)$) shows $GF(T)$ is terminal; as $\eta_T : T \to GF(T)$ is an isomorphism, $T$ is then terminal too by (2.6.51). ∎

**Definition 2.6.53 (Terminal Category)**
*The **terminal category** $\mathbf{1}$ is the category with a single object $\star$ and a single morphism $1_\star$.*

*For any category $\mathcal{C}$ and object $X \in \mathcal{C}$, we write $\underline{X} : \mathbf{1} \to \mathcal{C}$ for the functor with $\underline{X}(\star) = X$.*

**Definition 2.6.54 (Zero Object / Pointed Category)**
*A **zero object** of a category $\mathcal{C}$ is both an initial and terminal object. We denote this by $\mathbf{0}$ or $\mathbf{0}_{\mathcal{C}}$ when we need to emphasise the category $\mathcal{C}$*

*For two objects $X,Y \in \mathcal{C}$ we denote by $0_{X,Y}$ the composite morphism $X \to 0 \to Y$. This is independent of the choice of zero object.*

*A category with a zero object is **pointed**. A category $\mathcal{C}$ is pointed iff $\mathcal{C}^{op}$ is pointed. Further $\mathbf{0}_{\mathcal{C}} = \mathbf{0}_{\mathcal{C}^{op}}$.*

**Proposition 2.6.55**
*Let $\mathcal{C}$ be a pointed category and $X \in \mathcal{C}$. Then the following are equivalent*

*a) $X$ is an initial object*
*b) $X$ is a terminal object*
*c) $X$ is a zero object*

**Proposition 2.6.56**
*Let $\mathcal{C}$ be a pointed category. Then the morphism $0 \to X$ is a monomorphism and the morphism $Y \to 0$ is an epimorphism.*

**Proposition 2.6.57 (Functors Preserve Zeros)**
*Let $F : \mathcal{C} \to \mathcal{D}$ be a functor of pointed categories for which $F(0_{\mathcal{C}})$ is a zero object of $\mathcal{D}$. Then for all objects $X,Y$ we have*

$$F(0_{X,Y}) = 0_{F(X),F(Y)}$$

## 2.6.7 Yoneda Lemma

The motivation for the following result becomes more clear in the next section. Roughly speaking properties of an object, $X$, are encoded in the functor $\mathrm{Mor}(X,-)$.

**Proposition 2.6.58 (Yoneda Lemma)**
*Suppose $X, X' \in \mathcal{C}$ then there is a bijection between morphisms and natural transformation of functors*

$$\begin{aligned} \mathrm{Mor}(X',X) &\xrightarrow{\sim} \mathrm{Nat}(\mathrm{Mor}(X,-),\mathrm{Mor}(X',-)) \\ f &\longrightarrow \mathrm{Mor}(f,-) : (\phi \to \phi \circ f) \\ \alpha_X(1_X) &\longleftarrow \alpha \end{aligned}$$

*Observe that under this correspondence*

- $f = \mathrm{Mor}(f,X)(1_X)$
- $\mathrm{Mor}(f \circ g,-) = \mathrm{Mor}(g,-) \circ \mathrm{Mor}(f,-)$
- *$f$ is an isomorphism $\iff$ $\mathrm{Mor}(f,-)$ is a natural isomorphism.*

*In the latter case* $\mathrm{Mor}(f,-)^{-1} = \mathrm{Mor}(f^{-1},-)$

In many cases the set of morphisms $\mathrm{Mor}(X,X')$ has an additional structure (typically an abelian group or module). We encode this with the following definition

**Definition 2.6.59 (Enriched Hom Functor)**
*Let $(\mathcal{S},U)$ be a concrete category (2.6.9) for which $U$ reflects isomorphisms. We say a bifunctor*

$$\mathrm{Hom} : \mathcal{C}^{op} \times \mathcal{C} \to \mathcal{S}$$

© David Rufino CC BY-NC-ND 4.0

*is an* ***enriched hom functor*** *if we have*

- $U(\mathrm{Hom}(X,Y)) = \mathrm{Mor}(X,Y)$
- $U(\mathrm{Hom}(f,Y)) = \mathrm{Mor}(f,Y)$
- $U(\mathrm{Hom}(X,g)) = \mathrm{Mor}(X,g)$

*and bijections*

$$\begin{array}{ccc} & \mathrm{Mor}(X',X) & \\ \mathrm{Hom}(f,-) \swarrow \sim & & \sim \searrow \mathrm{Mor}(f,-) \\ \mathrm{Nat}(\mathrm{Hom}(X,-),\mathrm{Hom}(X',-)) & \xrightarrow[\sim]{U} & \mathrm{Nat}(\mathrm{Mor}(X,-),\mathrm{Mor}(X',-)) \end{array}$$

*Write the inverse of* $f \to \mathrm{Hom}(f,-)$ *as* $\mathcal{Y}$*. Then for* $\alpha \in \mathrm{Nat}(\mathrm{Hom}(X,-),\mathrm{Hom}(X',-))$

*a)* $\alpha$ *is a natural isomorphism* $\iff U\alpha$ *is a natural isomorphism* $\iff \mathcal{Y}(\alpha)$ *is an isomorphism.*
*b)* $\mathcal{Y}(\alpha) = U(\alpha_X)(1_X)$

*Note the right hand arrow is always a bijection by Yoneda's Lemma (2.6.58) and the diagram always commutes by assumption.*

*Proof.* Suppose $\alpha$ is a natural isomorphism then $U\alpha^{-1} \circ U\alpha = U\left(1_{\mathrm{Hom}(X,-)}\right) = 1_{\mathrm{Mor}(X,-)}$ and similarly $U\alpha \circ U\alpha^{-1} = 1_{\mathrm{Mor}(X',-)}$, so $U\alpha$ is a natural isomorphism. ∎

Note this trivially includes the usual case $\mathcal{S} = \mathbf{Set}$.

## 2.6.8 Representable Functors

Many algebraic constructions (e.g. tensor product) may be formalised in terms of "representable" functors (2.6.60). The usual definition involves the set-valued functor $\mathrm{Mor}(X,-)$ as defined in (2.6.11).

As a generalisation follows we consider an enriched hom functor (2.6.59) $\mathrm{Hom}(-,-)$ taking values in the concrete category $(\mathcal{S},U)$ where $U$ reflects isomorphisms. This clearly generalises the usual case.

**Definition 2.6.60 (Representable Functor)**
*Let* $F : \mathcal{C} \to \mathcal{S}$ *a covariant functor. We say* $F$ *is* ***representable*** *if there is a pair* $(X,\Phi)$ *where* $X \in \mathcal{C}$ *and* $\Phi$ *is a natural isomorphism*

$$\mathrm{Hom}(X,-) \xrightarrow[\sim]{\Phi} F(-)$$

*We say* $F$ *is* ***represented*** *by the pair* $(X,\Phi)$*. Note this implies* $U \circ F$ *is represented by* $U\Phi$*.*

It is often useful to have another conception of representable functor in terms of universal properties as this is more intuitive for applications.

**Definition 2.6.61 (Universal Element)**
*Let* $F : \mathcal{C} \to \mathcal{S}$ *be a covariant functor, then we say that a pair* $(X,i)$*, where* $X \in \mathcal{C}, i \in (U \circ F)(X)$*, is a* ***universal element*** *for* $F$ *if for all* $Y \in \mathcal{C}$ *there are bijections*

$$\begin{aligned} \mathrm{Mor}(X,Y) &\longrightarrow (U \circ F)(Y) \\ f &\to (U \circ F)(f)(i) \end{aligned}$$

*In this case we may also say* $F$ *is* ***represented*** *by the universal element* $(X,i)$*.*

We show the equivalence of the two notions by some abstract nonsense

**Proposition 2.6.62 (Representable Set-Valued Functor = Universal Element)**
*Let* $F : \mathcal{C} \to \mathbf{Set}$ *be a covariant functor then there is a natural correspondence between representations and universal elements*

$$\begin{aligned} \{\text{representations of } F\} &\longleftrightarrow \{\text{universal elements of } F\} \\ (X,\Phi) &\longrightarrow (X,\Phi(1_X)) \\ (X, f \to F(f)(i)) &\longleftarrow (X,i) \end{aligned}$$

The following may in particular be used to show that representations are unique up to isomorphism.

© David Rufino CC BY-NC-ND 4.0

**Proposition 2.6.63 (Morphisms Between Representations)**
*Suppose $(X, \Phi)$ and $(X', \Phi')$ represent the covariant functors $F, F' : \mathcal{C}^{op} \to \mathcal{S}$ respectively. Then there are bijections*

$$\begin{aligned} \mathrm{Mor}(X, X') &\xrightarrow{\sim} \mathrm{Nat}(\mathrm{Hom}(X', -), \mathrm{Hom}(X, -)) \xrightarrow{\sim} \mathrm{Nat}(F'(-), F(-)) \\ f &\to \mathrm{Hom}(f, -) && \to \hat{f} \\ U(\eta_{X'})(1_{X'}) &\leftarrow \eta \qquad \alpha^\star && \leftarrow \alpha \end{aligned}$$

*where we define the mutual inverses*

$$\begin{aligned} \hat{f} &= \Phi \circ \mathrm{Hom}(f, -) \circ (\Phi')^{-1} \\ \alpha^\star &= U\big(\Phi_{X'}^{-1} \circ \alpha_{X'} \circ \Phi'_{X'}\big)(1_{X'}) \end{aligned}$$

*Note this has the following properties*

- $\widehat{1_X} = \mathrm{id}_F$
- $\widehat{g \circ f} = \hat{f} \circ \hat{g}$ *where* $g : X' \to X''$ *is morphism and* $(X'', \Phi'')$ *represents* $F''$
- $f$ *is an isomorphism* $\iff \hat{f}$ *is a natural isomorphism and in this case* $\widehat{f^{-1}} = \hat{f}^{-1}$.

*Further $\alpha^\star \in \mathrm{Mor}(X, X')$ is the unique morphism such that the following diagram of natural transformations commutes*

$$\begin{array}{ccc} \mathrm{Hom}(X', -) & \xRightarrow{\Phi'} & F'(-) \\ {\scriptstyle \mathrm{Hom}(\alpha^\star, -)}\Vdots & & \Vdots{\scriptstyle \alpha} \\ \mathrm{Hom}(X, -) & \xRightarrow{\Phi} & F(-) \end{array}$$

*and satisfies*

- $\mathrm{id}_F^\star = 1_X$
- $(\beta \circ \alpha)^\star = \alpha^\star \circ \beta^\star$
- $\alpha^\star$ *is an isomorphism* $\iff \alpha$ *is a natural isomorphism*

*Proof.* The first bijection is simply the Yoneda Lemma and the second bijection is obvious. ∎

**Corollary 2.6.64 (Representation is Unique)**
*Let $F : \mathcal{C} \to \mathcal{S}$ be a covariant functor which is represented by pairs $(X, \Phi)$ and $(X', \Phi')$. Then they are isomorphic with two-sided inverses*

$$X \underset{U(\Phi'^{-1})(i)}{\overset{U(\Phi^{-1})(i')}{\rightleftarrows}} X'$$

*where $i \coloneqq (U\Phi)(1_X)$ and $i' \coloneqq (U\Phi')(1_{X'})$.*

*Proof.* We may apply the previous result with $F = F'$ and the identity natural transformation $\mathrm{id}_F$. ∎

In practice we typically have a family of representations, and we want to show the construction is functorial

**Definition 2.6.65 (Representable Bifunctor)**
*Let $F : \mathcal{D}^{op} \times \mathcal{C} \to \mathcal{S}$ be a bifunctor. We say that it is* ***representable*** *by $(G, \Phi)$ where $G : \mathcal{D} \to \mathcal{C}$ is a functor, if $\Phi$ is a natural isomorphism of bifunctors*

$$\Phi : \mathrm{Hom}(G(-), -) \xRightarrow{\sim} F(-, -)$$

The following shows we only need to construct the representation pointwise.

**Proposition 2.6.66**
*Let $F : \mathcal{D}^{op} \times \mathcal{C} \to \mathcal{S}$ be a bifunctor such that $F(D, -)$ is representable for all $D \in \mathcal{D}$, by $(G(D), \Phi_D)$. Then $G$ can be made into a covariant functor uniquely such that $\Phi$ constitutes a natural isomorphism of bifunctors*

$$\Phi_{D,C} : \mathrm{Hom}(G(D), C) \xRightarrow{\sim} F(D, C)$$

*Explicitly for any morphism $h : D \to D'$ we have the identity*

© David Rufino CC BY-NC-ND 4.0

$$G(h) = \Phi^{-1}_{D,G(D')}\Big(\Phi_{D',G(D')}\big(1_{G(D')}\big) \circ h\Big)$$

*Proof.* It is straightforward to verify that for any covariant functor $G$, the mapping $\mathrm{Mor}(G(-),-)$ is a bifunctor (contravariant in the first "slot").

Let $h : D \to D'$ be a morphism then by (2.6.21) $F(h,-) : F(D',-) \to F(D,-)$ is a natural transformation. Therefore we may define

$$G(h) \coloneqq F(h,-)^\star$$

in the notation of (2.6.63); it is the unique morphism making the following diagram commute

$$\begin{array}{ccc} \mathrm{Hom}(G(D'),-) & \xRightarrow{\Phi_{D'}} & F(D',-) \\ \Big\Vert\!\Big\Vert\!\Big\Vert\!\

*Proof.* In order to demonstrate that $\mathrm{Hom}(f_D, 1_C)$ is a natural transformation of bifunctors it's enough by (2.6.21) to show the following diagram commutes for $h : D' \to D$

$$\begin{array}{ccc} \mathrm{Hom}(G'(D), C) & \xrightarrow{-\circ f_D} & \mathrm{Hom}(G(D), C) \\ \downarrow{\scriptstyle -\circ G'(h)} & & \downarrow{\scriptstyle -\circ G(h)} \\ \mathrm{Hom}(G'(D'), C) & \xrightarrow{-\circ f_{D'}} & \mathrm{Hom}(G(D'), C) \end{array}$$

However $f_D \circ G(h) = G'(h) \circ f_{D'}$ by definition of natural transformation, from which it follows immediately. The map is a pointwise bijection (for every $D \in \mathcal{D}$) with the inverse given by the usual Yoneda Lemma.

The first two properties are straightforward. From these it follows that if $f$ is a natural isomorphism then so is $\mathrm{Hom}(f, -)$. Conversely if $\mathrm{Hom}(f, -)$ is a natural isomorphism then it has a two-sided inverse which is of the form $\mathrm{Hom}(g, -)$. It's then straightforward to argue that $g$ is a two-sided inverse of $f$. ∎

### 2.6.9 Adjoint Functors

Some universal constructions may be expressed as an adjoint pair of functors. My view is the most illustrative are the module Tensor-Hom adjunction and the sheafification construction. Using this concept we can simplify the verification of universal properties by appealing to general criteria for adjoint functors as below.

**Definition 2.6.69 (Adjoint Pair)**
*Let $F : \mathcal{C} \to \mathcal{D}$ and $G : \mathcal{D} \to \mathcal{C}$ be functors. We say that $F$ is **left adjoint** to $G$ if there is a bijection*

$$\psi_{C,D} : \mathrm{Mor}(F(C), D) \longrightarrow \mathrm{Mor}(C, G(D))$$

*which is natural in $C$ and $D$ in the following sense. Let $\alpha : C' \to C$, $\beta : D \to D'$, then for all $f : F(C) \to D$ we have*

$$\psi_{C',D'}(\beta \circ f \circ F(\alpha)) = G(\beta) \circ \psi_{C,D}(f) \circ \alpha(1)$$

*or equivalently for all $g : C \to G(D)$*

$$\beta \circ \psi^{-1}_{C,D}(g) \circ F(\alpha) = \psi^{-1}_{C',D'}(G(\beta) \circ g \circ \alpha)(2)$$

*Equivalently $F(-)$ represents the bifunctor $\mathrm{Mor}(-, G(-))$ (2.6.65).*

**Proposition 2.6.70 (Adjoint⇒unit, counit)**
*Let $F : \mathcal{C} \to \mathcal{D}$ and $G : \mathcal{D} \to \mathcal{C}$ be adjoint functors with relationship*

$$\psi_{C,D} : \mathrm{Mor}(F(C), D) \xrightarrow{\sim} \mathrm{Mor}(C, G(D))$$

*Then we have two natural transformations (**unit** and **counit** respectively)*

$$\eta : \mathrm{id}_{\mathcal{C}} \Rightarrow G \circ F$$
$$\epsilon : F \circ G \Rightarrow \mathrm{id}_{\mathcal{D}}$$

*defined by*

$$\eta_C = \psi_{C,F(C)}\left(1_{F(C)}\right)$$
$$\epsilon_D = \psi^{-1}_{G(D),D}\left(1_{G(D)}\right)$$

*Furthermore we may recover the adjoint relationship via*

$$\psi_{C,D}(f) = G(f) \circ \eta_C$$

$$\psi^{-1}_{C,D}(g) = \epsilon_D \circ F(g)$$

*Proof.* We show that the transformations given are natural. Suppose $\alpha : C \to C'$ then

© David Rufino CC BY-NC-ND 4.0

$$\begin{aligned} G(F(\alpha)) \circ \eta_C &= G(F(\alpha)) \circ \psi_{C,F(C)}\left(1_{F(C)}\right) \\ &= \psi_{C,F(C')}\left(F(\alpha) \circ 1_{F(C)}\right) \quad (1) \\ &= \psi_{C,F(C')}\left(1_{F(C')} \circ F(\alpha)\right) \\ &= \psi_{C',F(C')}\left(1_{F(C')}\right) \circ \alpha \quad (1) \\ &= \eta_{C'} \circ \alpha \end{aligned}$$

so $\eta$ is a natural transformation. Furthermore

$$\psi_{C,D}(f) = \psi_{C,D}\left(f \circ 1_{F(C)}\right) = G(f) \circ \psi_{C,F(C)}(1) = G(f) \circ \eta_C$$

as required. Similarly for $\beta : D \to D'$

$$\begin{aligned} \beta \circ \epsilon_D &= \beta \circ \psi^{-1}_{G(D),D}\left(1_{G(D)}\right) \\ &= \psi^{-1}_{G(D),D'}\left(G(\beta) \circ 1_{G(D)}\right) \quad (2) \\ &= \psi^{-1}_{G(D),D'}\left(1_{G(D')} \circ G(\beta)\right) \\ &= \psi^{-1}_{G(D'),D'}\left(1_{G(D')}\right) \circ F(G(\beta)) \quad (2) \\ &= \epsilon_{D'} \circ F(G(\beta)) \end{aligned}$$

■

Given two natural transformations we may recover a corresponding adjoint

**Proposition 2.6.71 (Adjoint from unit and counit)**
*Let $F : \mathcal{C} \to \mathcal{D}$ and $G : \mathcal{D} \to \mathcal{C}$ be functors with two natural transformations*

$$\begin{aligned} \eta &: \mathrm{id}_{\mathcal{C}} \Rightarrow G \circ F \\ \epsilon &: F \circ G \Rightarrow \mathrm{id}_{\mathcal{D}} \end{aligned}$$

*Then the following are equivalent*

*a) $F$ is left adjoint to $G$ with unit and counit $\eta$, $\epsilon$, and*
*b) the so-called **triangular identities** are satisfied*

$$1_{G(D)} : G(D) \overset{\eta_{G(D)}}{\longrightarrow} GFG(D) \overset{G(\epsilon_D)}{\longrightarrow} G(D)$$

$$1_{F(C)} : F(C) \overset{F(\eta_C)}{\longrightarrow} FGF(C) \overset{\epsilon_{F(C)}}{\longrightarrow} F(C)$$

*More precisely the adjunction is given by*

$$\begin{aligned} \mathrm{Mor}(F(C), D) &\underset{\phi}{\overset{\psi}{\rightleftharpoons}} \mathrm{Mor}(C, G(D)) \\ f &\longrightarrow G(f) \circ \eta_C \\ \epsilon_D \circ F(g) &\longleftarrow g \end{aligned}$$

*Proof.* Let $\psi, \phi$ denote the proposed adjunction maps. We will use the triangular identities to show that these are mutually inverse. First observe by naturality of $\eta$ and $\epsilon$ that

$$\begin{aligned} \psi\phi(g) &= G(\epsilon_D \circ F(g)) \circ \eta_C = G(\epsilon_D) \circ \eta_{G(D)} \circ g (3) \\ \phi\psi(f) &= \epsilon_D \circ F(G(f) \circ \eta_C) = f \circ \epsilon_{F(C)} \circ F(\eta_C) \end{aligned}$$

It's then immediate that these are mutually inverse maps if and only if the triangular identities are satisfied (one way is obvious, the other way consider $f = 1_{F(C)}$ and $g = 1_{G(D)}$).

Further one may easily verify that $\psi, \phi$ so-defined are natural in $C$ and $D$

$$\begin{aligned} \psi(\beta \circ f \circ F(\alpha)) &= G(\beta) \circ G(f) \circ GF(\alpha) \circ \eta_C \\ &= G(\beta) \circ G(f) \circ \eta_{C'} \circ \alpha \\ &= G(\beta) \circ \psi(f) \circ \alpha \end{aligned}$$

© David Rufino CC BY-NC-ND 4.0

■

**Proposition 2.6.72 (Criteria for right adjoint to be full and faithful)**
*Let $\mathcal{C} \underset{G}{\overset{F}{\rightleftarrows}} \mathcal{D}$ be adjoint functors with $\eta, \epsilon$ unit and counit transformations. Then*

- *$G$ is faithful if and only if $\epsilon$ is pointwise epic*
- *$G$ is full if and only if $\epsilon$ is pointwise split-monic*
- *$G$ is full and faithful if and only if $\epsilon$ is a pointwise isomorphism*

*Proof.* Consider the composite map

$$\mathrm{Mor}(D', D) \overset{\mathrm{Mor}(\epsilon_{D'}, D)}{\longrightarrow} \mathrm{Mor}(F(G(D')), D) \overset{\psi_{G(D'),D}}{\to} \mathrm{Mor}(G(D'), G(D))$$

which is natural in $D$ and $D'$. Note that image of $\alpha \in \mathrm{Mor}(D', D)$ is

$$\psi_{G(D'),D}(\alpha \circ \epsilon_{D'}) = G(\alpha) \circ \psi_{G(D'),D'}(\epsilon_{D'}) = G(\alpha)$$

so the composite is just $G(-)$. The second map is bijective by the adjoint assumption. Therefore the first map is injective (resp. surjective) if and only if $G$ is faithful (resp. full).

By (2.6.44) $\mathrm{Mor}(\epsilon_{D'}, D)$ is injective (resp. surjective) for all $D, D'$ if and only if $\epsilon_{D'}$ is epic (resp. split-monic) for all $D'$.

Then the first two statements follow easily. The last statement follows from the previous two, combined with (2.6.36).■

**Proposition 2.6.73 (Reflective Sub-Category)**
*Let $\mathcal{C} \underset{G}{\overset{F}{\rightleftarrows}} \mathcal{D}$ be adjoint functors such that $G$ is fully faithful. Then for $C \in \mathcal{C}$ we have $\eta_C$ is an isomorphism iff $C$ is in the essential image of $G$ (i.e. is isomorphic to $G(D)$ for some $D$).*

*In particular let $\mathcal{C}'$ denote the **essential image** of $\mathcal{D}$ under $G$. Then this restricts to an equivalence of categories $\mathcal{C}' \underset{G}{\overset{F}{\rightleftarrows}} \mathcal{D}$*

*Proof.* If $\eta_C : C \to GF(C)$ is an isomorphism define $D \coloneqq F(C)$, and we are done.

Conversely we first show that $\eta_{G(D)}$ is an isomorphism. By the triangular identities (2.6.71) we have

$$G(\epsilon_D) \circ \eta_{G(D)} = 1_{G(D)}$$

so $\eta_{G(D)}$ has a left inverse. Recall by (2.6.72) $\epsilon_D$ is an isomorphism, whence so is $G(\epsilon_D)$ (2.6.42). Therefore we may show directly that it is also a right inverse, and therefore $\eta_{G(D)}$ is an isomorphism.

Now assume that we have an isomorphism $\psi : C \xrightarrow{\sim} G(D)$ for some $D \in \mathcal{D}$. Then by naturality we have a commutative diagram

$$\begin{array}{ccc} C & \overset{\psi}{\underset{\sim}{\longrightarrow}} & G(D) \\ {\scriptstyle \eta_C}\downarrow & & \sim\downarrow{\scriptstyle \eta_{G(D)}} \\ GF(C) & \overset{\sim}{\underset{GF(\psi)}{\longrightarrow}} & GFG(D) \end{array}$$

observing that $GF(\psi)$ is also an isomorphism. Then we conclude $\eta_C$ is an isomorphism as required. ■

The following criteria will be useful

**Proposition 2.6.74 (Adjoints can be constructed pointwise)**
*Let $G : \mathcal{D} \to \mathcal{C}$ be a functor. Suppose that for all $C \in \mathrm{ob}(\mathcal{C})$ there exists $F(C) \in \mathrm{ob}(\mathcal{D})$ such that there is a bijection natural in $D$*

$$\psi_{C,D} : \mathrm{Mor}(F(C), D) \xrightarrow{\sim} \mathrm{Mor}(C, G(D))$$

*Then there exists a unique functor $F$ which ensures this is natural in $C$, more precisely for $g : C \to C'$ this is given by*

© David Rufino CC BY-NC-ND 4.0

$$F(g) = \psi^{-1}_{C,F(C')}(\eta_{C'} \circ g)$$
$$\eta_C \coloneqq \psi_{C,F(C)}\left(1_{F(C)}\right)$$

*This is also the unique functor which ensures $\eta_C$ is a natural transformation.*

*Proof.* The hypothesis is precisely that, for each $C \in \mathcal{C}$, $\left(F(C), \psi_{C,-}\right)$ represents the functor $\mathrm{Mor}(C, G(-)) : \mathcal{D} \to \mathcal{S}$; equivalently, this is the hypothesis of (2.6.66) applied to the (already well-defined, since $G$ is a functor) bifunctor $\mathrm{Mor}(-, G(-)) : \mathcal{C}^{\mathrm{op}} \times \mathcal{D} \to \mathcal{S}$, with the roles of the two categories there swapped so that $F(C)$ plays the role of the representing object varying over $\mathcal{C}$. That proposition's conclusion is precisely that $F$ can be made into a functor uniquely such that $\psi$ is a natural isomorphism of bifunctors, giving the first uniqueness; its explicit formula for $g : C \to C'$ reads

$$F(g) = \psi^{-1}_{C,F(C')}\left(\psi_{C',F(C')}\left(1_{F(C')}\right) \circ g\right) = \psi^{-1}_{C,F(C')}(\eta_{C'} \circ g)$$

using the definition of $\eta_{C'}$ for the second equality.

We have already shown that if $\psi_{C,D}$ is natural in $C$ then $\eta_C$ is a natural transformation (2.6.70). Conversely suppose that $\eta_C$ is a natural transformation. Since we have assumed that $\psi_{C,D}$ is natural in $D$, applying this to $\beta \coloneqq F(g)$ and $f \coloneqq 1_{F(C)}$ gives

$$\begin{aligned}\psi_{C,F(C')}(F(g)) &= \psi_{C,F(C')}\left(F(g) \circ 1_{F(C)}\right)\\ &= G(F(g)) \circ \psi_{C,F(C)}\left(1_{F(C)}\right)\\ &= GF(g) \circ \eta_C\\ &= \eta_{C'} \circ g\end{aligned}$$

using naturality of $\eta$ for the last equality, whence $F(g) = \psi^{-1}_{C,F(C')}(\eta_{C'} \circ g)$. ∎

**Proposition 2.6.75**
*Let $\mathcal{C} \underset{G}{\overset{F}{\rightleftarrows}} \mathcal{D}$ be a pair of adjoint functors. Then $F$ preserves initial objects and $G$ preserves terminal objects.*

*If $\mathcal{C}, \mathcal{D}$ are pointed categories then $F$ and $G$ also preserve zero objects and zero morphisms.*

*Proof.* We only prove that $F$ preserves initial objects, as the second statement follows dually. By hypothesis there is a bijection natural in $D$

$$\mathrm{Mor}(F(\varnothing), D) \xrightarrow{\sim} \mathrm{Mor}(\varnothing, G(D))$$

Therefore if $\varnothing$ is initial, then so is $F(\varnothing)$.

The last statement follows from (2.6.55) and (2.6.57). ∎

## 2.6.10 Limits

**Definition 2.6.76**
*A functor $F : I \to \mathcal{C}$ where $I$ is a small category is referred to as a **diagram***

**Definition 2.6.77 (Cones)**
*Let $F : I \to \mathcal{C}$ be a diagram. We say that the pair $\left(L, \{\lambda_i : L \to F(i)\}_{i \in I}\right)$ is a **cone** over $F$ if*

$$F(\alpha) \circ \lambda_i = \lambda_j \quad \text{for all morphisms} \quad \alpha : i \to j$$

*The cones for $F$ constitute a **cone category** $(\Delta \downarrow F)$ where a morphism $\alpha : \left(L, \{\lambda_i\}_{i \in I}\right) \to \left(L', \{\lambda'_i\}_{i \in I}\right)$ is just a morphism $\alpha : L \to L'$ such that $\lambda'_i \circ \alpha = \lambda_i$ for all $i \in I$*

**Definition 2.6.78 (Small Limit)**
*Let $F : I \to \mathcal{C}$ be a diagram. A cone $\left(L, \{\lambda_i\}_{i \in I}\right)$ over $F$ is a **limit** for $F$ if it is a **terminal object** (2.6.50) in the cone category.*

*Explicitly for every cone $\left(L', \{\lambda'_i\}_{i \in I}\right)$ there is a unique morphism $\alpha : L' \to L$ such that $\lambda_i \circ \alpha = \lambda'_i$.*

*We may denote this as $\lim F$.*

**Proposition 2.6.79**
*Let $F : I \to \mathcal{C}$ be a diagram, $\left(L, \{\lambda_i\}_{i \in I}\right)$ a limit of $F$ and $\left(L', \{\lambda'_i\}_{i \in I}\right)$ a cone. Then the following are equivalent*

*a) $L'$ is a limit for $F$*

© David Rufino CC BY-NC-ND 4.0

*b) there is an isomorphism of cones* $L' \xrightarrow{\sim} L$

*c) there is a morphism of cones* $L' \to L$ *and precisely one (cone) endomorphism of* $L'$.

*In this case the (cone) isomorphism is the unique cone morphism.*

*Proof.* By definition $\left(L, \{\lambda_i\}_{i\in I}\right)$ is a terminal object of the cone category $(\Delta \downarrow F)$, so this is precisely (2.6.51) applied with $\star := \left(L, \{\lambda_i\}_{i\in I}\right)$ and $X := \left(L', \{\lambda'_i\}_{i\in I}\right)$. ∎

**Proposition 2.6.80 (Uniqueness of Limits)**
*Let* $F : I \to \mathcal{C}$ *be a diagram and* $\left(L, \{\lambda_i\}_{i\in I}\right)$ *and* $\left(L', \{\lambda'_i\}_{i\in I}\right)$ *limits of* $F$. *Then there is a unique morphism* $\psi : L \to L'$ *satisfying* $\lambda'_i \circ \psi = \lambda_i$, *which is an isomorphism.*

*Proof.* Since $\left(L', \{\lambda'_i\}_{i\in I}\right)$ is a limit for $F$, (2.6.79) gives a unique isomorphism of cones $\psi : L \to L'$, that is, a unique isomorphism $\psi$ with $\lambda'_i \circ \psi = \lambda_i$ for all $i \in I$. ∎

**Proposition 2.6.81 (Product)**
*Let* $I$ *be a discrete small category and* $\{X_i\}_{i\in I}$ *be a family of objects in* $\mathcal{C}$. *Then there is a unique functor* $F : I \to \mathcal{C}$ *such that* $F(i) = X_i$.

*Consider an object* $X \in \mathcal{C}$ *and family of morphisms* $\pi_i : X \to X_i$, *then the following are equivalent*

*a) The cone* $\left(X, \{\pi_i\}_{i\in I}\right)$ *is a limit for* $F$

*b) For any family of morphisms* $\lambda_i : X' \to X_i$ *there exists a unique* $\lambda : X' \to X$ *such that* $\pi_i \circ \lambda = \lambda_i$

*c) For all objects* $X'$ *the map*

$$\Phi : \mathrm{Mor}(X', X) \longrightarrow \prod_{i\in I} \mathrm{Mor}(X', X_i)$$
$$f \to \pi_i \circ f$$

*is a bijection, i.e.* $(X, \Phi)$ *represents the contravariant functor* $\prod_{i\in I} \mathrm{Mor}(-, X_j)$.

*In this case we say that* $\left(X, \{\pi_i\}_{i\in I}\right)$ *is a* ***product*** *for* $\{X_i\}_{i\in I}$ *and denote this by* $\prod_{i\in I} X_i$.

*For any family of morphisms* $\lambda_i : X' \to X_i$ *we write the induced morphism* $\lambda : X' \to \prod_{i\in I} X_i$ *as* $(\lambda_i)_{i\in I}$.

**Proposition 2.6.82 (Functoriality of Product)**
*Let* $I$ *be a discrete small category and suppose that* $\mathcal{C}$ *has* $I$*-shaped limits. Then we may define a map on objects*

$$\prod_{i\in I}(-) : \mathcal{C}^I \to \mathcal{C}$$

*Recall that a morphism* $\prod_{i\in I} X_i \to \prod_{i\in I} Y_i$ *in* $\mathcal{C}^I$ *is simply a family of morphisms* $(f_i : X_i \to Y_i)_{i\in I}$ *in* $\mathcal{C}$. *Then we define*

$$\prod_{i\in I} f_i := (f_i \circ \pi_i)_{i\in I}$$

*which is the unique morphism satisfying*

$$\pi_i \circ \prod_{i\in I} f_i = f_i \circ \pi_i$$

*This makes* $\prod_{i\in I}(-)$ *into a functor.*

**Proposition 2.6.83 (Products of Monomorphisms)**
*Let* $I$ *be a discrete small category and* $\{f_i : X_i \to Y_i\}_{i\in I}$ *a family of monomorphisms such that* $\prod_{i\in I} X_i$ *and* $\prod_{i\in I} Y_i$ *exist. Then* $\prod_{i\in I} f_i : \prod_{i\in I} X_i \to \prod_{i\in I} Y_i$ *is a monomorphism.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Suppose $a, b : Z \to \prod_{i\in I} X_i$ satisfy $\left(\prod_{i\in I} f_i\right) \circ a = \left(\prod_{i\in I} f_i\right) \circ b$. For each $i \in I$, composing with the projection $\pi_i : \prod_{i\in I} Y_i \to Y_i$ and using $\pi_i \circ \prod_{i\in I} f_i = f_i \circ \pi_i$ gives

$$f_i \circ (\pi_i \circ a) = f_i \circ (\pi_i \circ b)$$

and since $f_i$ is monic, $\pi_i \circ a = \pi_i \circ b$. As this holds for every $i \in I$, by the uniqueness clause of the universal property of the product $\prod_{i\in I} X_i$ (2.6.81) we conclude $a = b$. ∎

© David Rufino CC BY-NC-ND 4.0

**Proposition 2.6.84 (Functoriality of Binary Product)**
*Let $X, X', Y, Y' \in \mathcal{C}$ be such that $X \times Y$ and $X' \times Y'$ exists. For $f : X \to X'$ and $g : Y \to Y'$ there exists a unique morphism*

$$f \times g : X \times Y \to X' \times Y'$$

*such that*

$$\pi_{X'} \circ (f \times g) = f \circ \pi_X$$

*and*

$$\pi_{Y'} \circ (f \times g) = g \circ \pi_Y$$

*If $\mathcal{C}$ has all binary products then this determines a bifunctor*

$$\times : \mathcal{C} \times \mathcal{C} \to \mathcal{C}$$

**Proposition 2.6.85 (Limits exist in Set)**
*Limits exist in* **Set** *more precisely for a diagram $F : I \to \mathbf{Set}$*

$$\lim F = \left\{ (x_i)_{i \in I} \in \prod_{i \in I} F(i) \;\middle|\; F(\alpha)(x_i) = x_j \text{ for all } \alpha : i \to j \right\}$$

**Definition 2.6.86**
*Let $U : \mathcal{C} \to \mathcal{D}$ be a functor and $I$ a small category.*

*We say that $U$ **creates limits** of shape $I$ if for all diagrams $F : I \to \mathcal{C}$ such that $\lim U \circ F$ exists then $\lim F$ exists and is the unique cone such that $U(\lim F) = \lim(U \circ F)$.*

*We say that $U$ **preserves limits** of shape $I$ if whenever $\lim F$ exists in $\mathcal{C}$ we have $\lim U \circ F$ exists in $\mathcal{D}$ and is equal to $U(\lim F)$.*

*If we restrict to the case $I$ is discrete (resp. $I = \mathbf{2}$) then we say that $U$ creates/preserves products (resp. equalisers).*

**Proposition 2.6.87 (Creates⇒Preserves)**
*Let $U : \mathcal{C} \to \mathcal{D}$ which creates limits of shape $I$ and $\mathcal{D}$ has all limits of shape $I$. Then $U$ preserves limits of shape $I$.*

*In particular this holds unconditionally for $\mathcal{D} = \mathbf{Set}$ since this has all limits.*

## 2.6.11 Colimit

**Definition 2.6.88 (Co-cones)**
*Let $F : I \to \mathcal{C}$ be a diagram. We say that the pair $\left(L, \{\lambda_i : F(i) \to L\}_{i \in I}\right)$ is a **co-cone** over $F$ if*

$$\lambda_j \circ F(\alpha) = \lambda_i \quad \text{for all morphisms} \quad \alpha : i \to j$$

*The cones for $F$ constitute a **co-cone category** $(F \downarrow \Delta)$ where a morphism $\alpha : \left(L, \{\lambda_i\}_{i \in I}\right) \to \left(L', \{\lambda'_i\}_{i \in I}\right)$ is just a morphism $\alpha : L \to L'$ such that $\alpha \circ \lambda_i = \lambda'_i$ for all $i \in I$*

**Definition 2.6.89 (Small Colimit)**
*Let $F : I \to \mathcal{C}$ be a diagram. A co-cone $\left(L, \{\lambda_i\}_{i \in I}\right)$ over $F$ is a **colimit** for $F$ if it is an **initial object** (2.6.48) in the co-cone category.*

*Explicitly for every co-cone $\left(L', \{\lambda'_i\}_{i \in I}\right)$ there is a unique morphism $\alpha : L \to L'$ such that $\lambda'_i \circ \alpha = \lambda_i$.*

*We may denote this as $\operatorname{colim} F$.*

**Proposition 2.6.90**
*Let $F : I \to \mathcal{C}$ be a diagram, $\left(L, \{\lambda_i\}_{i \in I}\right)$ a colimit of $F$ and $\left(L', \{\lambda'_i\}_{i \in I}\right)$ a co-cone. Then the following are equivalent*

*a) $L'$ is a colimit for $F$*
*b) there is an isomorphism of co-cones $L' \xrightarrow{\sim} L$*
*c) there is a morphism of co-cones $L' \to L$ and precisely one (co-cone) endomorphism of $L'$.*

*In this case the (co-cone) isomorphism is unique.*

*Proof.* By definition $\left(L, \{\lambda_i\}_{i \in I}\right)$ is an initial object of the co-cone category $(F \downarrow \Delta)$, so this is precisely (2.6.49) applied with $\varnothing := \left(L, \{\lambda_i\}_{i \in I}\right)$ and $X := \left(L', \{\lambda'_i\}_{i \in I}\right)$. ∎

© David Rufino CC BY-NC-ND 4.0

**Proposition 2.6.91 (Uniqueness of Colimits)**
*Let $F : I \to \mathcal{C}$ be a diagram and $\left(L, \{\lambda_i\}_{i\in I}\right)$ and $\left(L', \{\lambda'_i\}_{i\in I}\right)$ colimits of $F$. Then there is a unique morphism $\psi : L \to L'$ satisfying $\psi \circ \lambda_i = \lambda'_i$, which is an isomorphism.*

*Proof.* Since $\left(L', \{\lambda'_i\}_{i\in I}\right)$ is a colimit for $F$, (2.6.90) gives a unique isomorphism of co-cones $\phi : L' \to L$, that is, a unique isomorphism $\phi$ with $\phi \circ \lambda'_i = \lambda_i$. Setting $\psi := \phi^{-1} : L \to L'$ gives the required unique isomorphism with $\psi \circ \lambda_i = \lambda'_i$. ∎

**Proposition 2.6.92 (Limits and Colimits swap under $\mathcal{C}^{\mathrm{op}}$)**
*Let $I$ be a small category and $F : I \to \mathcal{C}$ a diagram. Write $F^{op} : I^{op} \to \mathcal{C}^{op}$ for the diagram with $F^{op}(i) := F(i)$ on objects and $F^{op}(\alpha^{op}) := F(\alpha)^{op}$ for $\alpha^{op} : j \to i$ the morphism of $I^{op}$ corresponding to $\alpha : i \to j$ of $I$. Then, regarding each $\lambda_i : L \to F(i)$ as a morphism $\lambda_i^{op} : F(i) \to L$ of $\mathcal{C}^{op}$, $\left(L, \{\lambda_i\}_{i\in I}\right)$ is a limit for $F$ in $\mathcal{C}$ if and only if $\left(L, \{\lambda_i^{op}\}_{i\in I}\right)$ is a colimit for $F^{op}$ in $\mathcal{C}^{op}$.*

*Dually, $\left(L, \{\lambda_i\}_{i\in I}\right)$ is a colimit for $F$ in $\mathcal{C}$ if and only if $\left(L, \{\lambda_i^{op}\}_{i\in I}\right)$ is a limit for $F^{op}$ in $\mathcal{C}^{op}$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

We only prove the first statement, the second following by applying it to $\mathcal{C}^{\mathrm{op}}$ in place of $\mathcal{C}$, using $(\mathcal{C}^{\mathrm{op}})^{\mathrm{op}} = \mathcal{C}$.

Define $\Sigma : (\Delta \downarrow F) \to (F^{\mathrm{op}} \downarrow \Delta)^{\mathrm{op}}$, $\left(L, \{\lambda_i\}_{i\in I}\right) \mapsto \left(L, \{\lambda_i^{\mathrm{op}}\}_{i\in I}\right)$, acting on a cone morphism $\mu : (L, \lambda) \to (L', \lambda')$ (i.e. $\mu : L \to L'$ with $\lambda'_i \circ \mu = \lambda_i$ for all $i$) by $\Sigma(\mu) := \mu^{\mathrm{op}}$. This is well-defined: upon reversing every arrow (using $\mathrm{Mor}_{\mathcal{C}^{\mathrm{op}}}(X, Y) = \mathrm{Mor}_{\mathcal{C}}(Y, X)$ and that composition in $\mathcal{C}^{\mathrm{op}}$ reverses that of $\mathcal{C}$ (2.6.4)), the cone condition $F(\alpha) \circ \lambda_i = \lambda_j$ for $\alpha : i \to j$ of $I$ becomes exactly the co-cone condition $\lambda_i^{\mathrm{op}} \circ F^{\mathrm{op}}(\alpha^{\mathrm{op}}) = \lambda_j^{\mathrm{op}}$ for $\alpha^{\mathrm{op}} : j \to i$ of $I^{\mathrm{op}}$, so $\{\lambda_i^{\mathrm{op}}\}$ is a co-cone over $F^{\mathrm{op}}$; and the cone morphism condition $\lambda'_i \circ \mu = \lambda_i$ becomes $\mu^{\mathrm{op}} \circ (\lambda'_i)^{\mathrm{op}} = \lambda_i^{\mathrm{op}}$, so $\mu^{\mathrm{op}}$ is a morphism $(L', \lambda'^{\mathrm{op}}) \to (L, \lambda^{\mathrm{op}})$ of $(F^{\mathrm{op}} \downarrow \Delta)$, that is, a morphism $(L, \lambda^{\mathrm{op}}) \to (L', \lambda'^{\mathrm{op}})$ of $(F^{\mathrm{op}} \downarrow \Delta)^{\mathrm{op}}$. Thus $\Sigma$ is well-defined, and is manifestly functorial.

Dually define $\Sigma^{-1} : (F^{\mathrm{op}} \downarrow \Delta)^{\mathrm{op}} \to (\Delta \downarrow F)$, $\left(L, \{\mu_i\}_{i\in I}\right) \mapsto \left(L, \{\mu_i^{\mathrm{op}}\}_{i\in I}\right)$, acting on morphisms by $\nu \mapsto \nu^{\mathrm{op}}$. Since $(-)^{\mathrm{op}}$ is an involution on morphisms, $\Sigma^{-1}\Sigma = 1$ and $\Sigma\Sigma^{-1} = 1$, so $\Sigma$ is an isomorphism of categories, in particular an equivalence of categories (2.6.25).

By (2.6.4), a category $\mathcal{D}$ has $T$ as a terminal object if and only if $\mathcal{D}^{\mathrm{op}}$ has $T$ as an initial object (since $\mathrm{Mor}_{\mathcal{D}^{\mathrm{op}}}(X, T) = \mathrm{Mor}_{\mathcal{D}}(T, X)$ for all $X$). So $(L, \{\lambda_i\})$ is a limit for $F$, i.e. terminal in $(\Delta \downarrow F)$ (2.6.78), iff, by (2.6.52), $\Sigma(L, \{\lambda_i\}) = (L, \{\lambda_i^{\mathrm{op}}\})$ is terminal in $(F^{\mathrm{op}} \downarrow \Delta)^{\mathrm{op}}$, iff (applying the noted equivalence to $\mathcal{D} := (F^{\mathrm{op}} \downarrow \Delta)$) $(L, \{\lambda_i^{\mathrm{op}}\})$ is initial in $(F^{\mathrm{op}} \downarrow \Delta)$, that is, a colimit for $F^{\mathrm{op}}$ (2.6.89). ∎

**Proposition 2.6.93 (Co-product)**
*Let $I$ be a discrete small category and $\{X_i\}_{i\in I}$ be a family of objects in $\mathcal{C}$. Then there is a unique functor $F : I \to \mathcal{C}$ such that $F(i) = X_i$.*

*Consider an object $X \in \mathcal{C}$ and family of morphisms $s_i : X_i \to X$, then the following are equivalent*

*a) The co-cone $\left(X, \{s_i\}_{i\in I}\right)$ is a colimit for $F$*
*b) For any family of morphisms $\lambda_i : X_i \to X'$ there exists a unique $\lambda : X \to X'$ such that $\lambda \circ s_i = \lambda_i$*
*c) For all objects $X'$ the map*

$$\Phi : \mathrm{Mor}(X, X') \longrightarrow \prod_{i\in I} \mathrm{Mor}(X_i, X')$$
$$\lambda \to (\lambda \circ s_i)_{i\in I}$$

*is a bijection, i.e. $(X, \Phi)$ represents the covariant functor $\prod_{i\in I} \mathrm{Mor}(X_i, -)$.*

*In this case we say that $\left(X, \{s_i\}_{i\in I}\right)$ is a **co-product** for $\{X_i\}_{i\in I}$ and denote this by $\coprod_{i\in I} X_i$.*

*For any family of morphisms $\lambda_i : X_i \to X'$ we write the induced morphism $\lambda : \coprod_{i\in I} X_i \to X'$ as $[\lambda_i]_{i\in I}$.*

**Definition 2.6.94**
*Let $U : \mathcal{C} \to \mathcal{D}$ be a functor and $I$ a small category.*

*We say that $U$ **creates co-limits** of shape $I$ if for all diagrams $F : I \to \mathcal{C}$ such that $\lim U \circ F$ exists then $\operatorname{colim} F$ exists and is the unique cone such that $U(\operatorname{colim} F) = \operatorname{colim}(U \circ F)$.*

*We say that $U$ **preserves co-limits** of shape $I$ if whenever $\operatorname{colim} F$ exists in $\mathcal{C}$ we have $\operatorname{colim} U \circ F$ exists in $\mathcal{D}$ and is equal to $U(\operatorname{colim} F)$.*

© David Rufino CC BY-NC-ND 4.0

*If we restrict to the case $I$ is discrete (resp. $I = \mathbf{2}$) then we say that $U$ creates/preserves coproducts (resp. coequalisers).*

**Proposition 2.6.95 (Creates⇒Preserves)**
*Let $U : \mathcal{C} \to \mathcal{D}$ which creates colimits of shape $I$ and $\mathcal{D}$ has all colimits of shape $I$. Then $U$ preserves colimits of shape $I$.*

*In particular this holds unconditionally for $\mathcal{D} = \mathbf{Set}$ since this has all colimits.*

## 2.6.12 Functorial Limits

Having fixed a single diagram at a time so far, we now let the diagram vary. By (2.6.96), taking the limit in one variable of a bifunctor is functorial in the other – so a family of limits indexed by $\mathcal{C}$ assembles into a genuine functor $\mathcal{C} \to \mathcal{E}$. Specialising this to the evaluation bifunctor recovers the more familiar statement, (2.6.97): the limit $\lim_J(-)$ is itself a functor out of (a subcategory of) the functor category $\mathcal{C}^J$, so a natural transformation of diagrams induces a morphism of their limits, compatibly with composition and identities – and in particular carries isomorphisms of diagrams to isomorphisms of limits. We also record that limits are invariant under isomorphism of the indexing category, that a limit over a product of two index categories may be computed either simultaneously or iteratively one variable at a time (2.6.99), with the two iterated orders agreeing whenever both make sense (2.6.100), and dual statements for colimits, together with a comparison morphism relating a colimit of limits to a limit of colimits, which need not be an isomorphism in general.

**Proposition 2.6.96 (Pointwise Limits are Functorial)**
*Let $J$ be a small category and $F : \mathcal{C} \times J \to \mathcal{E}$ a bifunctor.*

*Suppose that for every $X \in \mathcal{C}$ there is a limit $\left(\lim_{j\in J} F(X,j), \{\lambda_{X,j}\}_{j\in J}\right)$ for $F(X,-)$.*

*Then $\lim_{j\in J} F(-,j)$ is functorial in the sense there is a unique morphism making the following diagram commute*

$$\begin{array}{ccc} \lim_{j\in J} F(X,j) & \overset{\lim_{j\in J} F(f,j)}{\dashrightarrow} & \lim_{j\in J} F(X',j) \\ \downarrow{\scriptstyle \lambda_{X,j}} & & \downarrow{\scriptstyle \lambda_{X',j'}} \\ F(X,j) & \xrightarrow{F(f,\beta)} & F(X',j') \end{array}$$

*for all $f : X \to X'$ and $\beta : j \to j'$.*

*Therefore this determines a functor $\lim_{j\in J} F(-,j) : \mathcal{C} \to \mathcal{E}$.*

*Proof.* Recall that $\left(\lim_{j\in J} F(X,j), \{\lambda_{X,j}\}_{j\in J}\right)$ is a terminal object in the category of cones $(\Delta \downarrow F(X,-))$. Suppose $f : X \to X'$ is a morphism. Since $F$ is a bifunctor (2.6.19), for every $\beta : j \to j'$ in $J$

$$F(X',\beta) \circ F\big(f, 1_j\big) \circ \lambda_{X,j} = F(f,\beta) \circ \lambda_{X,j} = F\big(f, 1_{j'}\big) \circ F(X,\beta) \circ \lambda_{X,j} = F\big(f, 1_{j'}\big) \circ \lambda_{X,j'}$$

so $\left(\lim_{j\in J} F(X,j), \{F\big(f,1_j\big) \circ \lambda_{X,j}\}_{j\in J}\right)$ is a cone over $F(X',-)$.

Therefore there is a unique morphism $h : \lim_{j\in J} F(X,j) \to \lim_{j\in J} F(X',j)$ into the terminal object of $(\Delta \downarrow F(X',-))$ with $\lambda_{X',j} \circ h = F\big(f,1_j\big) \circ \lambda_{X,j}$ for every $j \in J$. For general $\beta : j \to j'$, the target cone condition and the bifunctoriality identity above give

$$\lambda_{X',j'} \circ h = F(X',\beta) \circ \lambda_{X',j} \circ h = F(X',\beta) \circ F\big(f,1_j\big) \circ \lambda_{X,j} = F(f,\beta) \circ \lambda_{X,j}$$

as required.

Writing $\lim_{j\in J} F(f,j) \coloneqq h$, the identity and composition laws needed for functoriality of $\lim_{j\in J} F(-,j)$ follow directly from this uniqueness: for $f = 1_X$, $h = 1_{\lim_j F(X,j)}$ has the defining property of $\lim_j F(1_X,j)$, so equals it by uniqueness; and for $f : X \to X'$, $f' : X' \to X''$, the morphism $\lim_j F(f',j) \circ \lim_j F(f,j)$ satisfies the same defining property (compatibility with the $\lambda_{X'',j}$) as $\lim_j F(f'f,j)$, so the two agree by uniqueness (this is the same pattern used for image factorisations, see later). ∎

**Corollary 2.6.97 (Limits are Functorial in the Diagram)**
*Let $J$ be a small category and $\mathcal{C}$ a category, and let $\mathcal{C}^J_{\lim}$ be the full subcategory of the functor category $\mathcal{C}^J$ (2.6.14) spanned by those diagrams $D : J \to \mathcal{C}$ for which $\lim_J D$ exists. Then*

$$\lim_J(-) : \mathcal{C}^J_{\lim} \to \mathcal{C}, \quad D \mapsto \lim_J D$$

© David Rufino CC BY-NC-ND 4.0

*is a functor. Consequently, a natural transformation $\eta : D \Rightarrow D'$ of $\mathcal{C}_{\lim}^{J}$ induces $\lim_J \eta : \lim_J D \to \lim_J D'$, satisfying the composition and identity laws of a functor; and if $\eta$ is a natural isomorphism, $\lim_J \eta$ is an isomorphism, with inverse $\lim_J(\eta^{-1})$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Define the **evaluation bifunctor** $\mathrm{ev} : \mathcal{C}_{\lim}^{J} \times J \to \mathcal{C}$ by $\mathrm{ev}(D, j) := D(j)$ on objects and, for a morphism $(\eta, \beta) : (D, j) \to (D', j')$ of $\mathcal{C}_{\lim}^{J} \times J$ (so $\eta : D \Rightarrow D'$ a natural transformation and $\beta : j \to j'$ in $J$), $\mathrm{ev}(\eta, \beta) := D'(\beta) \circ \eta_j$ $(= \eta_{j'} \circ D(\beta)$, by naturality of $\eta$ at $\beta)$. This preserves identities $(\mathrm{ev}(1_D, 1_j) = D(1_j) \circ (1_D)_j = 1_{D(j)})$ and composition: for a further morphism $(\eta', \beta') : (D', j') \to (D'', j'')$,

$$\mathrm{ev}(\eta', \beta') \circ \mathrm{ev}(\eta, \beta) = D''(\beta') \circ \eta'_{j'} \circ D'(\beta) \circ \eta_j = D''(\beta') \circ D''(\beta) \circ \eta'_j \circ \eta_j = \mathrm{ev}((\eta' \eta, \beta' \beta)),$$

using naturality of $\eta'$ at $\beta$ (i.e. $\eta'_{j'} \circ D'(\beta) = D''(\beta) \circ \eta'_j$) and functoriality of $D''$; so ev is a bifunctor.

By definition of $\mathcal{C}_{\lim}^{J}$, $\lim_{j \in J} \mathrm{ev}(D, j) = \lim_J D$ exists for every $D \in \mathcal{C}_{\lim}^{J}$, so (2.6.96) applies to ev (with $\mathcal{C}_{\lim}^{J}$ in the role of the first factor), giving that $\lim_{j \in J} \mathrm{ev}(-, j) = \lim_J(-) : \mathcal{C}_{\lim}^{J} \to \mathcal{C}$ is a functor, with $\lim_J \eta$ its action on a morphism $\eta$, satisfying the composition and identity laws by functoriality. If $\eta$ is a natural isomorphism, i.e. an isomorphism of $\mathcal{C}_{\lim}^{J}$, then $\lim_J \eta$ is an isomorphism with inverse $\lim_J(\eta^{-1})$ by (2.6.42). ∎

**Proposition 2.6.98 (Limits are Invariant under Isomorphism of the Indexing Category)**
*Let $\sigma : J \to I$ be an isomorphism of categories with inverse $\sigma^{-1}$, and $F : I \to \mathcal{C}$ a diagram. Then $\left(L, \{\lambda_i\}_{i \in I}\right)$ is a limit for $F$ if and only if $\left(L, \left\{\lambda_{\sigma(j)}\right\}_{j \in J}\right)$ is a limit for $F\sigma : J \to \mathcal{C}$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Define $\Sigma : (\Delta \downarrow F) \to (\Delta \downarrow F\sigma)$, $\left(L, \{\lambda_i\}_{i \in I}\right) \mapsto \left(L, \left\{\lambda_{\sigma(j)}\right\}_{j \in J}\right)$, acting as the identity on the underlying morphism of apexes; this is well-defined since $\left\{\lambda_{\sigma(j)}\right\}_{j \in J}$ is a cone over $F\sigma$ whenever $\{\lambda_i\}_{i \in I}$ is a cone over $F$ (the cone condition for $F\sigma$ at a morphism $\beta$ of $J$ is exactly the cone condition for $F$ at $\sigma(\beta)$), and $\Sigma$ is manifestly functorial.

Dually define $\Sigma^{-1} : (\Delta \downarrow F\sigma) \to (\Delta \downarrow F)$, $\left(L, \{\mu_j\}_{j \in J}\right) \mapsto \left(L, \left\{\mu_{\sigma^{-1}(i)}\right\}_{i \in I}\right)$. Since $\sigma^{-1}\sigma = 1_J$ and $\sigma\sigma^{-1} = 1_I$, we have $\Sigma^{-1}\Sigma = 1$ and $\Sigma\Sigma^{-1} = 1$ on the nose, so $\Sigma$ is an isomorphism of categories, in particular an equivalence of categories (2.6.25) (with identity natural isomorphisms).

By (2.6.52), $(\Delta \downarrow F)$ has a terminal object iff $(\Delta \downarrow F\sigma)$ does, and $\Sigma$ carries a terminal object of $(\Delta \downarrow F)$, i.e. a limit of $F$, to a terminal object of $(\Delta \downarrow F\sigma)$, i.e. a limit of $F\sigma$, as required. ∎

**Proposition 2.6.99 (Iterated Limits)**
*Let $I, J$ be small categories and $F : I \times J \to \mathcal{C}$ a bifunctor. Suppose that for every $i \in I$ there is a limit $\left(L(i), \{\lambda_{i,j}\}_{j \in J}\right)$ for $F(i, -)$, so that by (2.6.96) this determines a functor $L : I \to \mathcal{C}$.*

*Then $\lim_{(i,j) \in I \times J} F(i, j)$ exists if and only if $\lim_{i \in I} L(i)$ exists, and in this case*

- *a) given a limit $\left(M, \{\nu_i\}_{i \in I}\right)$ for $L$, the cone $\left(M, \{\lambda_{i,j} \circ \nu_i\}_{(i,j) \in I \times J}\right)$ is a limit for $F$;*
- *b) conversely, given a limit $\left(N, \{\pi_{i,j}\}_{(i,j) \in I \times J}\right)$ for $F$, for each $i \in I$ there is a unique morphism $\sigma_i : N \to L(i)$ such that $\lambda_{i,j} \circ \sigma_i = \pi_{i,j}$ for all $j \in J$, and $\left(N, \{\sigma_i\}_{i \in I}\right)$ is a limit for $L$.*

*Symmetrically, suppose that for every $j \in J$ there is a limit $\left(R(j), \{\mu_{i,j}\}_{i \in I}\right)$ for $F(-, j)$, so that by (2.6.96) this determines a functor $R : J \to \mathcal{C}$. Then $\lim_{(i,j) \in I \times J} F(i, j)$ exists if and only if $\lim_{j \in J} R(j)$ exists, and in this case*

- *c) given a limit $\left(M', \{\xi_j\}_{j \in J}\right)$ for $R$, the cone $\left(M', \{\mu_{i,j} \circ \xi_j\}_{(i,j) \in I \times J}\right)$ is a limit for $F$;*
- *d) conversely, given a limit $\left(N, \{\pi_{i,j}\}_{(i,j) \in I \times J}\right)$ for $F$, for each $j \in J$ there is a unique morphism $\tau_j : N \to R(j)$ such that $\mu_{i,j} \circ \tau_j = \pi_{i,j}$ for all $i \in I$, and $\left(N, \{\tau_j\}_{j \in J}\right)$ is a limit for $R$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

We construct mutually inverse functors $\Xi : (\Delta \downarrow L) \to (\Delta \downarrow F)$ and $\Theta : (\Delta \downarrow F) \to (\Delta \downarrow L)$; as $\Theta \circ \Xi = 1$, $\Xi \circ \Theta = 1$, $\Xi$ is in particular an equivalence of categories (2.6.25) (with identity natural isomorphisms), so by (2.6.52), $(\Delta \downarrow F)$ has a terminal object iff $(\Delta \downarrow L)$ does, and $\Xi$ carries a terminal object of $(\Delta \downarrow L)$, i.e. a limit of $L$, to a terminal object of $(\Delta \downarrow F)$, i.e. a limit of $F$, as required.

**The functor $\Xi$.** Given a cone $\left(X, \{\eta_i\}_{i \in I}\right)$ over $L$, define $\Xi(X, \{\eta_i\}) := \left(X, \{\lambda_{i,j} \circ \eta_i\}_{(i,j) \in I \times J}\right)$. This is a cone over $F$: for $(\alpha, \beta) : (i, j) \to (i', j')$, writing $(\alpha, \beta) = (\alpha, 1_{j'}) \circ (1_i, \beta)$,

© David Rufino CC BY-NC-ND 4.0

$$\begin{aligned} F(\alpha,\beta)\circ\lambda_{i,j}\circ\eta_i &= F\big(\alpha,1_{j'}\big)\circ F(1_i,\beta)\circ\lambda_{i,j}\circ\eta_i \\ &= F\big(\alpha,1_{j'}\big)\circ\lambda_{i,j'}\circ\eta_i \\ &= \lambda_{i',j'}\circ L(\alpha)\circ\eta_i \\ &= \lambda_{i',j'}\circ\eta_{i'} \end{aligned}$$

using in turn that $\lambda_{i,_}$ is a cone over $F(i,-)$, the defining property of $L(\alpha)$ (2.6.96), and that $\eta$ is a cone over $L$. On morphisms $\Xi$ is the identity on the underlying morphism of apexes, which is trivially functorial.

**The functor $\Theta$.** Given a cone $\Big(X,\{\rho_{i,j}\}_{(i,j)\in I\times J}\Big)$ over $F$, for each $i\in I$ the family $\{\rho_{i,j}\}_{j\in J}$ is a cone over $F(i,-)$, so by the universal property of $\Big(L(i),\{\lambda_{i,j}\}_j\Big)$ there is a unique morphism $\theta_i : X\to L(i)$ such that $\lambda_{i,j}\circ\theta_i=\rho_{i,j}$ for all $j\in J$. Define $\Theta\big(X,\{\rho_{i,j}\}\big) := \big(X,\{\theta_i\}_{i\in I}\big)$.

We check $\{\theta_i\}$ is a cone over $L$. Fix $\alpha : i\to i'$ in $I$; then for every $j'\in J$

$$\lambda_{i',j'}\circ L(\alpha)\circ\theta_i = F\big(\alpha,1_{j'}\big)\circ\lambda_{i,j'}\circ\theta_i = F\big(\alpha,1_{j'}\big)\circ\rho_{i,j'} = \rho_{i',j'} = \lambda_{i',j'}\circ\theta_{i'}$$

using the defining property of $L(\alpha)$, the defining property of $\theta_i$, that $\rho$ is a cone over $F$ applied to $\big(\alpha,1_{j'}\big):(i,j')\to(i',j')$, and the defining property of $\theta_{i'}$. As this holds for all $j'\in J$, the universal property of $\Big(L(i'),\{\lambda_{i',j'}\}_{j'}\Big)$ gives $L(\alpha)\circ\theta_i=\theta_{i'}$.

On morphisms $\Theta$ again acts as the identity on apexes: if $\phi:\big(X,\{\rho_{i,j}\}\big)\to\big(X',\{\rho'_{i,j}\}\big)$ is a morphism of cones, then $\lambda_{i,j}\circ\theta'_i\circ\phi=\rho'_{i,j}\circ\phi=\rho_{i,j}=\lambda_{i,j}\circ\theta_i$ for all $j$, so by uniqueness in the universal property of $L(i)$, $\theta'_i\circ\phi=\theta_i$, i.e. $\phi$ is also a morphism of cones $(X,\{\theta_i\})\to(X',\{\theta'_i\})$.

**Mutually inverse.** Given a cone $(X,\{\eta_i\})$ over $L$, applying $\Xi$ then $\Theta$ recovers, for each $i$, the unique morphism $X\to L(i)$ factoring $\big(\lambda_{i,j}\circ\eta_i\big)_j$ through $\{\lambda_{i,j}\}_j$; since $\eta_i$ already satisfies this by definition, uniqueness gives $\Theta(\Xi(X,\{\eta_i\}))=(X,\{\eta_i\})$. Conversely, given a cone $\big(X,\{\rho_{i,j}\}\big)$ over $F$, applying $\Theta$ then $\Xi$ gives $\big(X,\{\lambda_{i,j}\circ\theta_i\}\big)=\big(X,\{\rho_{i,j}\}\big)$ directly from the defining property of $\theta_i$. Both functors act as the identity on the underlying morphism of apexes, so $\Xi$ and $\Theta$ are mutually inverse isomorphisms of categories.

Finally, the symmetric statement follows by applying the above to the bifunctor $F^T : J\times I\to\mathcal{C}$, $F^T(j,i):=F(i,j)$ (swapping the roles of $I,J$, with $R$ playing the role of $L$), and reindexing via (2.6.98) along the swap isomorphism $I\times J\cong J\times I$. ∎

**Corollary 2.6.100 (Comparison of Iterated Limits)**
*Let $I,J$ be small categories and $F:I\times J\to\mathcal{C}$ a bifunctor, and suppose (as in (2.6.99)) that $L(i):=\lim_{j\in J}F(i,j)$ exists for every $i\in I$ and $R(j):=\lim_{i\in I}F(i,j)$ exists for every $j\in J$. If any one of $\lim_{(i,j)\in I\times J}F(i,j)$, $\lim_{i\in I}L(i)$, $\lim_{j\in J}R(j)$ exists, then all three exist, and there is a unique isomorphism $\Phi$ such that the following diagram commutes*

$$\begin{array}{ccccc} \lim_{i\in I}L(i) & & \xrightarrow{\quad\Phi\quad} & & \lim_{j\in J}R(j) \\ {\scriptstyle\nu_{i'}}\downarrow & & & & \downarrow{\scriptstyle\xi_{j'}} \\ L(i') & \xrightarrow[\lambda_{i',j'}]{} & F(i',j') & \xleftarrow[\mu_{i',j'}]{} & R(j') \end{array}$$

*for all $i'\in I$, $j'\in J$, where $\nu_{i'}:\lim_{i\in I}L(i)\to L(i')$ and $\xi_{j'}:\lim_{j\in J}R(j)\to R(j')$ are the respective limit cones.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

By (2.6.99), $\lim_{(i,j)\in I\times J}F(i,j)$ exists if and only if $\lim_{i\in I}L(i)$ exists, and if and only if $\lim_{j\in J}R(j)$ exists; so if any one of the three exists, all three do. In this case, taking $(M,\{\nu_i\}):=(\lim_{i\in I}L(i),\{\nu_i\})$ in part a), the cone $\Big(\lim_{i\in I}L(i),\{\lambda_{i,j}\circ\nu_i\}_{(i,j)\in I\times J}\Big)$ is a limit for $F$; likewise, taking $\big(M',\{\xi_j\}\big):=\big(\lim_{j\in J}R(j),\{\xi_j\}\big)$ in part c), the cone $\Big(\lim_{j\in J}R(j),\{\mu_{i,j}\circ\xi_j\}_{(i,j)\in I\times J}\Big)$ is also a limit for $F$. By (2.6.80) there is therefore a unique isomorphism $\Phi$ as required. ∎

**Proposition 2.6.101 (Pointwise Colimits are Functorial)**
*Let $I$ be a small category and $F:I\times\mathcal{D}\to\mathcal{E}$ a bifunctor.*

*Suppose that for every $X\in\mathcal{D}$ there is a colimit $\Big(\operatorname{colim}_{i\in I}F(i,X),\{\mu_{i,X}\}_{i\in I}\Big)$ for $F(-,X)$*

*Then $\operatorname{colim}_{i\in I}F(i,-)$ is functorial in the sense there is a unique morphism making the following diagram commute*

© David Rufino CC BY-NC-ND 4.0

$$\begin{array}{ccc} F(i,X) & \xrightarrow{F(\alpha,g)} & F(i',X') \\ \downarrow{\scriptstyle \mu_{i,X}} & & \downarrow{\scriptstyle \mu_{i',X'}} \\ \operatorname{colim}_{i\in I} F(i,X) & \xrightarrow{\operatorname{colim}_{i\in I} F(i,g)} & \operatorname{colim}_{i\in I} F(i,X') \end{array}$$

*for all* $g : X \to X'$ *and* $\alpha : i \to i'$.

**Proposition 2.6.102 (Comparison Map between Limits and Colimits)**
*Let* $I, J$ *be small categories and* $F : I \times J \to \mathcal{C}$ *be a bifunctor.*

*Suppose that there exists*

*a) limits* $i \to \lim_{j\in J} F(i,j)$ *(which is automatically functorial in* $i$ *(2.6.96))*
*b) colimits* $j \to \operatorname{colim}_{i\in I} F(i,j)$ *(which us automatically functorial in* $j$ *(2.6.101))*
*c) a colimit* $\operatorname{colim}_{i\in I}\big(\lim_{j\in J} F(i,j)\big)$
*d) a limit* $\lim_{j\in J} \operatorname{colim}_{i\in I} F(i,j)$

*(which holds if* $\mathcal{C}$ *has limits of shape* $J$ *and colimits of shape* $I$*). Then there exists a unique "comparison morphism"*

$$\begin{array}{ccccc} \operatorname{colim}_{i\in I} \lim_{j\in J} F(i,j) & & \xrightarrow{\quad\Phi\quad} & & \lim_{j\in J} \operatorname{colim}_{i\in I} F(i,j) \\ {\scriptstyle \lambda_{i'}}\uparrow & & & & \downarrow{\scriptstyle \mu_{j'}} \\ \lim_{j\in J} F(i',j) & \xrightarrow[\lambda_{i',j'}]{} & F(i',j') & \xrightarrow[\mu_{i',j'}]{} & \operatorname{colim}_{i\in I} F(i,j') \end{array}$$

*making the diagram commute for all* $i' \in I$ *and* $j' \in J$.

*Proof.* The morphisms $\lambda_{i'}$, $\lambda_{i',j'}$, $\mu_{i',j'}$ and $\mu_{j'}$ are the usual cone morphisms arising from the limit and colimit constructions. Suppose $\beta : j' \to j''$ then we have a commutative diagram, using both functoriality of colimit (2.6.101) and definition of cone $\lambda_{i',-}$

$$\begin{array}{ccccc} \lim_{j\in J} F(i',j) & \xrightarrow{\lambda_{i',j'}} & F(i',j') & \xrightarrow{\mu_{i',j'}} & \operatorname{colim}_{i\in I} F(i,j') \\ & \searrow^{\lambda_{i',j''}} & \downarrow{\scriptstyle F(1,\beta)} & & \downarrow{\scriptstyle \operatorname{colim}_{i\in I} F(i,\beta)} \\ & & F(i',j'') & \xrightarrow{\mu_{i',j''}} & \operatorname{colim}_{i\in I} F(i,j'') \end{array}$$

By universal property of the (top-right) limit to $\mu_{i',j'} \circ \lambda_{i',j'}$ we may find for each $i' \in I$ a unique morphism

$$\Phi_{i'} : \lim_{j\in J} F(i',j) \longrightarrow \lim_{j\in J} \operatorname{colim}_{i\in I} F(i,j)$$

satisfying

$$\mu_{j'} \circ \Phi_{i'} = \mu_{i',j'} \circ \lambda_{i',j'} \quad \forall j' \in J$$

Then by functoriality of limit (2.6.96) and definition of the cocone $\mu_{-,j'}$ we have a commutative diagram for all $\alpha : i' \to i''$

$$\begin{array}{ccccc} \lim_{j\in J} F(i',j) & \xrightarrow{\lambda_{i',j'}} & F(i',j') & \xrightarrow{\mu_{i',j'}} & \operatorname{colim}_{i\in I} F(i,j') \\ {\scriptstyle \lim_{j\in J} F(\alpha,j)}\downarrow & & {\scriptstyle F(\alpha,1)}\downarrow & \nearrow_{\mu_{i',j''}} & \\ \lim_{j\in J} F(i'',j) & \xrightarrow{\lambda_{i'',j'}} & F(i',j'') & & \end{array}$$

so that for all $j' \in J$

$$\begin{aligned} \mu_{j'} \circ \Phi_{i''} \circ \lim_{j\in J} F(\alpha,j) &= \mu_{i'',j''} \circ \lambda_{i'',j''} \circ \lim_{j\in J} F(\alpha,j) \\ &= \mu_{i',j'} \circ \lambda_{i',j'} \end{aligned}$$

whence by uniqueness

$$\Phi_{i''} \circ \lim_{j\in J} F(\alpha,j) = \Phi_{i'}$$

© David Rufino CC BY-NC-ND 4.0

Therefore by the definition of the (top-left) co-limit there is a unique morphism

$$\Phi : \operatorname{colim}_{i\in I} \lim_{j\in J} F(i,j) \longrightarrow \lim_{j\in J} \operatorname{colim}_{i\in I} F(i,j)$$

such that we have a commutative diagram




Then if there is another morphism $\Phi'$ which makes the square commute define $\Phi'_{i'} = \Phi' \circ \lambda_{i'}$, we may use uniqueness to deduce that $\Phi'_{i'} = \Phi_{i'}$ and $\Phi = \Phi'$. ■

## 2.6.13 Biproduct

In **AbGrp** and $A$-**Mod** coproducts and products coincide, so we formalise this property.

**Definition 2.6.103 (Finite Biproduct)**
*Let $\mathcal{C}$ be a pointed category, $I$ a* finite *set and $\{X_i\}_{i\in I}$ be a family of objects. We say a triple $\left(\bigoplus_{i\in I} X_i, \{s_i\}_{i\in I}, \{\pi_i\}_{i\in I}\right)$ is a **biproduct** for the family $\{X_i\}_{i\in I}$ if the following holds*

*a) $\left(\bigoplus_{i\in I} X_i, \{\pi_i\}_{i\in I}\right)$ is a product*
*b) $\left(\bigoplus_{i\in I} X_i, \{s_i\}_{i\in I}\right)$ is a coproduct*
*c) The morphisms $\pi_i, s_j$ satisfy the orthogonality conditions*

$$\pi_i \circ s_j = \delta_{ij} := \begin{cases} 1_{X_i} & i = j \\ 0 & i \neq j \end{cases}$$

**Proposition 2.6.104 (Criteria for Existence of Biproduct)**
*Let $\mathcal{C}$ be a pointed category, $I$ a finite set and $\{X_i\}_{i\in I}$ be a family of objects. Suppose both the product $\prod_{i\in I} X_i$ and coproduct $\coprod_{i\in I} X_i$ exists. Then there is a unique comparison morphism*

$$\psi : \coprod_{i\in I} X_i \longrightarrow \prod_{i\in I} X_i$$

*such that*

$$\pi_i \circ \psi \circ s_j = \delta_{ij}$$

*Then the following are equivalent*

*a) $\psi$ is an isomorphism*
*b) $\left(\coprod_{i\in I} X_i, \{s_i\}_{i\in I}, \{\pi_i \circ \psi\}_{i\in I}\right)$ is a biproduct*
*c) $\left(\prod_{i\in I} X_i, \{\psi \circ s_i\}_{i\in I}, \{\pi_i\}_{i\in I}\right)$ is a biproduct*
*d) a biproduct $\bigoplus_{i\in I} X_i$ exists*

*Proof.* We may write $\psi = \left[\left(\delta_{ij}\right)_{i\in I}\right]_{j\in I}$ then $\psi$ is the unique map such that $\psi \circ s_j = \left(\delta_{ij}\right)_{i\in I}$, which is the unique map such that $\pi_i \circ \psi \circ s_j = \delta_{ij}$. Suppose $\psi'$ is another such map then $\psi' \circ s_j = \psi \circ s_j$ whence $\psi' = \psi$.

a) $\implies$ b) We show that $\psi$ is a morphism of cones, and the result follows from (2.6.79).

a) $\implies$ c) We show that $\psi^{-1}$ is a morphism of co-cones, and the result follows from (2.6.90).

b) $\implies$ a) By (2.6.80) $\psi$ is uniquely determined as the isomorphism of limits.

c) $\implies$ a) Similarly.

b) $\implies$ d) immediate

d) $\implies$ a) Let $\pi'_i, s'_j$ be the biproduct maps for $\bigoplus_{i\in I} X_i$. Then by (2.6.80) (2.6.91) there is a diagram

© David Rufino CC BY-NC-ND 4.0




where $\alpha$ is the unique morphism satisfying $\alpha \circ s_i = s_i'$, $\beta$ is the unique morphism satisfying $\pi_i \circ \beta = \pi_i'$, and both $\alpha$ and $\beta$ are isomorphisms. Therefore $\pi_i \circ \beta \circ \alpha \circ s_j = \pi_i' \circ s_j' = \delta_{ij}$. Therefore by uniqueness $\psi = \beta \circ \alpha$ and is an isomorphism as required. ∎

**Proposition 2.6.105 (Binary biproducts are enough)**
*Let $\mathcal{C}$ be a pointed category. Then it has all (finite) biproducts iff it has all binary biproducts (i.e. the case $\#I = 2$).*

*Proof.* We prove that $\bigoplus_{i\in I} X_i$ exists for $I = \{1, ..., n\}$ by induction on $n$, where $n = 1$ is immediate and $n = 2$ is the base case. Consider a biproduct $\left(\bigoplus_{i=1}^n X_i, \{s_i\}, \{\pi_i\}\right)$ which we denote by $Y$, and the binary biproduct $(Y \oplus X_{n+1}, \{f, g\}, \{p, q\})$. Then we claim that

$$Z := (Y \oplus X_{n+1}, \{f \circ s_1, ..., f \circ s_n, g\}, \{\pi_1 \circ p, ..., \pi_n \circ p, q\})$$

is a biproduct. There is a natural isomorphism

$$\begin{aligned}\mathrm{Mor}(Z, -) &\xrightarrow{\sim} \mathrm{Mor}(Y, -) \times \mathrm{Mor}(X_{n+1}, -) \\ &\xrightarrow{\sim} \prod_{i=1}^{n+1} \mathrm{Mor}(X_i, -)\end{aligned}$$

which shows that $Z$ is a product with the given projection maps. Similarly we may show that $Z$ is a coproduct. Finally

$$\begin{aligned}\pi_i \circ p \circ f \circ s_j &= \pi_i \circ 1_Y \circ s_j = \delta_{ij} \\ \pi_i \circ p \circ g &= \pi_i \circ 0 = 0 \\ q \circ f \circ s_j &= 0 \circ s_j = 0\end{aligned}$$

so that the cone and co-cone morphisms satisfy the required orthogonality conditions. ∎

In **AbGrp** and $A$-**Mod**, a homomorphism between direct sums $\bigoplus_{j\in J} X_j \to \bigoplus_{i\in I} Y_i$ is determined by, and can be recovered from, a matrix of homomorphisms $X_j \to Y_i$; we record this categorically.

**Definition 2.6.106 (Matrix of a Morphism)**
*Let $\mathcal{C}$ be a pointed category with biproducts $\bigoplus_{j\in J} X_j$ (injections $s_j$, projections $\pi_j^X$) and $\bigoplus_{i\in I} Y_i$ (injections $s_i^Y$, projections $\pi_i^Y$), and $f : \bigoplus_{j\in J} X_j \to \bigoplus_{i\in I} Y_i$ a morphism. The **matrix** of $f$ is the family $\left(f_{ij}\right)_{i\in I, j\in J}$ of morphisms*

$$f_{ij} := \pi_i^Y \circ f \circ s_j : X_j \to Y_i$$

*When $I = \{1, ..., m\}$ and $J = \{1, ..., n\}$ we display this as an $m \times n$ array $\left(f_{ij}\right)$, as for ordinary matrices.*

**Remark 2.6.107**
*Calling $\left(f_{ij}\right)$ a **matrix** is not yet fully justified: for this notation to behave like ordinary matrix notation, composition should correspond to matrix multiplication, $(f \circ g)_{ik} = \sum_j f_{ij} \circ g_{jk}$, but this formula presupposes an addition on $\mathrm{Mor}(X_k, Y_i)$, which is no part of the data of a pointed category with biproducts. We can nonetheless justify composition whenever it requires no addition, namely when each row of $f$ (or column of $g$) has at most one nonzero entry (2.6.114); addition on hom-sets is only introduced once biproducts are used to construct such an addition (2.6.118), after which the general composition formula above is recovered in full (2.6.122).*

**Proposition 2.6.108 (Matrix Notation for Biproducts)**
*In the setting of (2.6.106), the assignment $f \mapsto \left(f_{ij}\right)_{i\in I, j\in J}$ is a bijection*

$$\mathrm{Mor}\left(\bigoplus_{j\in J} X_j, \bigoplus_{i\in I} Y_i\right) \xrightarrow{\sim} \prod_{i\in I, j\in J} \mathrm{Mor}(X_j, Y_i)$$

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Given a family $\left(g_{ij}\right)_{i\in I, j\in J}$ of morphisms $g_{ij} : X_j \to Y_i$, for each $j \in J$ the family $\left(g_{ij}\right)_{i\in I}$ determines, by the universal property of the product $\bigoplus_{i\in I} Y_i$ (2.6.81), a unique morphism $c_j : X_j \to \bigoplus_{i\in I} Y_i$ with $\pi_i^Y \circ c_j = g_{ij}$ for every $i \in I$. In

© David Rufino CC BY-NC-ND 4.0

turn, the family $(c_j)_{j\in J}$ determines, by the universal property of the coproduct $\bigoplus_{j\in J} X_j$ (2.6.93), a unique morphism $g : \bigoplus_{j\in J} X_j \to \bigoplus_{i\in I} Y_i$ with $g \circ s_j = c_j$ for every $j \in J$. We show $f \mapsto (f_{ij})$ and $(g_{ij}) \mapsto g$ are mutually inverse.

Starting from $f$, form its matrix $(f_{ij})$ and then $c_j, g$ as above. Since $\pi_i^Y \circ (f \circ s_j) = f_{ij}$ for every $i$, $f \circ s_j$ satisfies the defining property of $c_j$, so $c_j = f \circ s_j$ by uniqueness; then $f$ itself satisfies $f \circ s_j = c_j$ for every $j$, so $g = f$ by uniqueness.

Starting from a family $(g_{ij})$, form $c_j, g$ as above and then the matrix of $g$: for every $i, j$, the $(i, j)$ entry of the matrix of $g$ is $\pi_i^Y \circ g \circ s_j = \pi_i^Y \circ c_j = g_{ij}$, using $g \circ s_j = c_j$ and then the defining property of $c_j$. So the matrix of $g$ is again $(g_{ij})$. ∎

**Remark 2.6.109**
*The matrix of $1_{\bigoplus_{i\in I} X_i}$ is the **identity matrix** $(\delta_{ij} 1_{X_i})_{i,j\in I}$, immediate from the orthogonality conditions $\pi_i \circ s_j = \delta_{ij}$ (2.6.103).*

**Remark 2.6.110**
*Any object $X$ is trivially a biproduct of the one-element family $\{X\}$, via $(X, \{1_X\}, \{1_X\})$ (the orthogonality condition reduces to $1_X \circ 1_X = 1_X$). This lets us apply (2.6.108) whenever one side of a morphism is a single object rather than a genuine biproduct, by regarding it as such.*

**Remark 2.6.111 (Injections and Projections as Matrices)**
*For the binary biproduct $X_1 \oplus X_2$, the injections and projections are themselves matrices in the sense of (2.6.106), namely the columns and rows of the identity matrix*

$$s_1 = \begin{pmatrix} 1_{X_1} \\ 0 \end{pmatrix} \quad s_2 = \begin{pmatrix} 0 \\ 1_{X_2} \end{pmatrix} \quad \pi_1 = \begin{pmatrix} 1_{X_1} & 0 \end{pmatrix} \quad \pi_2 = \begin{pmatrix} 0 & 1_{X_2} \end{pmatrix}$$

*immediate from $\pi_i \circ s_j = \delta_{ij}$ (2.6.103). More generally, for $\bigoplus_{i\in I} X_i$, regarding $X_{i_0}$ as the trivial biproduct of $\{X_{i_0}\}$, $s_{i_0}$ is the $i_0$-th column and $\pi_{i_0}$ the $i_0$-th row of the identity matrix $(\delta_{ij} 1_{X_i})_{i,j\in I}$.*

**Lemma 2.6.112 (A Matrix is a Row of Columns and a Column of Rows)**
*Let $M : X_1 \oplus X_2 \to Y_1 \oplus Y_2$ have matrix $\begin{pmatrix} M_{11} & M_{12} \\ M_{21} & M_{22} \end{pmatrix}$. Then*

$$M = \begin{pmatrix} \begin{pmatrix} M_{11} \\ M_{21} \end{pmatrix} & \begin{pmatrix} M_{12} \\ M_{22} \end{pmatrix} \end{pmatrix} = \begin{pmatrix} \begin{pmatrix} M_{11} & M_{12} \end{pmatrix} \\ \begin{pmatrix} M_{21} & M_{22} \end{pmatrix} \end{pmatrix}$$

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Write $c_1 \coloneqq M \circ s_1$, $c_2 \coloneqq M \circ s_2$ for the columns of $M$. The morphism $\begin{pmatrix} c_1 & c_2 \end{pmatrix} : X_1 \oplus X_2 \to Y_1 \oplus Y_2$ is, by definition of the coproduct-induced notation (2.6.93), the unique morphism satisfying $\begin{pmatrix} c_1 & c_2 \end{pmatrix} \circ s_1 = c_1$ and $\begin{pmatrix} c_1 & c_2 \end{pmatrix} \circ s_2 = c_2$. Since $M$ itself satisfies $M \circ s_1 = c_1$ and $M \circ s_2 = c_2$ by definition of $c_1, c_2$, uniqueness gives $M = \begin{pmatrix} c_1 & c_2 \end{pmatrix}$.

Dually, writing $r_1 \coloneqq \pi_1 \circ M$, $r_2 \coloneqq \pi_2 \circ M$ for the rows of $M$, $\begin{pmatrix} r_1 \\ r_2 \end{pmatrix}$ is the unique morphism with $\pi_1 \circ \begin{pmatrix} r_1 \\ r_2 \end{pmatrix} = r_1$ and $\pi_2 \circ \begin{pmatrix} r_1 \\ r_2 \end{pmatrix} = r_2$ (2.6.81), and $M$ satisfies these by definition of $r_1, r_2$, so $M = \begin{pmatrix} r_1 \\ r_2 \end{pmatrix}$. ∎

**Lemma 2.6.113 (Composing a Column or Row on the Outside)**
*This is a general fact about products and coproducts, not requiring biproducts: let $p : Z \to Y_1$, $q : Z \to Y_2$ and $N : W \to Z$. Then*

$$\begin{pmatrix} p \\ q \end{pmatrix} \circ N = \begin{pmatrix} p \circ N \\ q \circ N \end{pmatrix}$$

*Dually, for $a : X_1 \to Z$, $b : X_2 \to Z$ and $N : Z \to W$*

$$N \circ \begin{pmatrix} a & b \end{pmatrix} = \begin{pmatrix} N \circ a & N \circ b \end{pmatrix}$$

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Write $\pi_1, \pi_2$ for the projections of $Y_1 \oplus Y_2$. Since $\pi_1 \circ \begin{pmatrix} p \\ q \end{pmatrix} = p$ and $\pi_2 \circ \begin{pmatrix} p \\ q \end{pmatrix} = q$ (2.6.106), for any $N : W \to Z$, by associativity of composition

$$\pi_1 \circ \left( \begin{pmatrix} p \\ q \end{pmatrix} \circ N \right) = p \circ N \qquad \pi_2 \circ \left( \begin{pmatrix} p \\ q \end{pmatrix} \circ N \right) = q \circ N$$

so $\begin{pmatrix} p \\ q \end{pmatrix} \circ N$ satisfies the defining property of $\begin{pmatrix} p \circ N \\ q \circ N \end{pmatrix}$, and the two are equal by the uniqueness clause of (2.6.81).

© David Rufino CC BY-NC-ND 4.0

Dually, writing $s_1, s_2$ for the injections of $X_1 \oplus X_2$, since $(a\ \ b) \circ s_1 = a$ and $(a\ \ b) \circ s_2 = b$, for any $N : Z \to W$

$$(N \circ (a\ \ b)) \circ s_1 = N \circ a \qquad (N \circ (a\ \ b)) \circ s_2 = N \circ b$$

so $N \circ (a\ \ b)$ satisfies the defining property of $(N \circ a\ \ N \circ b)$, and the two are equal by the uniqueness clause of (2.6.93).■

## 2.6.14 Biproduct Additivity

Following [Fre64] [Lan98], our eventual goal is to show that in a pointed category with (binary) biproducts, each hom-set $\mathrm{Mor}(X, Y)$ carries a canonical addition, making composition bilinear – so that additivity, usually imposed as an axiom, is instead a theorem, derived here purely from the existence of biproducts, via matrix notation.

**Proposition 2.6.114 (Composition Without Cross Terms)**
*Let $\mathcal{C}$ be a pointed category with binary biproducts, hence all finite biproducts (2.6.105), $I, J$ finite sets, $\sigma : I \to J$ a function, and $M : \bigoplus_{j\in J} X_j \to \bigoplus_{i\in I} Y_i$ a morphism whose matrix satisfies $M_{ij} = 0$ for every $i \in I$, $j \neq \sigma(i)$. Then for any $N : \bigoplus_{k\in K} W_k \to \bigoplus_{j\in J} X_j$ with matrix $(N_{jk})$*

$$(M \circ N)_{ik} = M_{i,\sigma(i)} \circ N_{\sigma(i),k} \quad \text{for all } i \in I, k \in K$$

*Dually, if $\tau : I \to J$ and $M' : \bigoplus_{i\in I} X_i \to \bigoplus_{j\in J} Y_j$ has matrix satisfying $M'_{ji} = 0$ for every $i \in I$, $j \neq \tau(i)$, then for any $N : \bigoplus_{j\in J} Y_j \to \bigoplus_{k\in K} Z_k$ with matrix $(N_{kj})$*

$$(N \circ M')_{ki} = N_{k,\tau(i)} \circ M'_{\tau(i),i} \quad \text{for all } k \in K, i \in I$$

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

For $i \in I$ and any $j \in J$, $a_i \circ \pi_{\sigma(i)} \circ s_j = a_i \circ \delta_{\sigma(i),j}$ (using $\pi_{\sigma(i)} \circ s_j = \delta_{\sigma(i),j}$ (2.6.103)), which is $a_i = M_{i,\sigma(i)}$ if $j = \sigma(i)$ and $0 = M_{ij}$ otherwise (by hypothesis on $M$); this matches $\pi_i \circ M \circ s_j = M_{ij}$ for every $j \in J$, so by the uniqueness clause of (2.6.93) (applied to maps out of $\bigoplus_{j\in J} X_j$), $\pi_i \circ M = a_i \circ \pi_{\sigma(i)}$.

Hence, for every $i \in I$, $k \in K$

$$\pi_i \circ M \circ N \circ s_k = a_i \circ \pi_{\sigma(i)} \circ N \circ s_k = a_i \circ N_{\sigma(i),k}$$

giving the matrix of $M \circ N$.

Dually, for $i \in I$ and any $j \in J$, $\pi_j \circ s_{\tau(i)} \circ c_i = \delta_{j,\tau(i)} c_i$, which matches $\pi_j \circ M' \circ s_i = M'_{ji}$ for every $j \in J$ (equal to $c_i$ if $j = \tau(i)$ and 0 otherwise, by hypothesis), so by the uniqueness clause of (2.6.81) (applied to maps into $\bigoplus_{j\in J} Y_j$), $M' \circ s_i = s_{\tau(i)} \circ c_i$; hence, for every $k \in K$, $i \in I$

$$\pi_k \circ N \circ M' \circ s_i = \pi_k \circ N \circ s_{\tau(i)} \circ c_i = N_{k,\tau(i)} \circ c_i$$

giving the matrix of $N \circ M'$. ■

**Example 2.6.115**
*(2.6.114) recovers familiar matrix identities, at various arities, without needing addition, since none of the matrices below have cross terms. For $a : X \to Y$, regarding $X$ (resp. $Y$) as the trivial biproduct of $\{X\}$ (resp. $\{Y\}$) (2.6.108)*

$$(a\ \ 0) = a \circ (1_X\ \ 0) \qquad \begin{pmatrix} a \\ 0 \end{pmatrix} = \begin{pmatrix} 1_Y \\ 0 \end{pmatrix} \circ a$$

*for $a : X_1 \to Y_1$, $b : X_2 \to Y_2$, $p : W \to X_1$ and $q : W \to X_2$*

$$\begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix} \circ \begin{pmatrix} p \\ q \end{pmatrix} = \begin{pmatrix} a \circ p \\ b \circ q \end{pmatrix}$$

*and for $u : X_1 \to Z$*

$$(u\ \ 0) \circ \begin{pmatrix} p \\ q \end{pmatrix} = u \circ p$$

**Definition 2.6.116 (Diagonal and Codiagonal)**
*Let $\mathcal{C}$ be a pointed category with binary biproduct $X \oplus X$, with injections $s_1, s_2$ and projections $\pi_1, \pi_2$. Regarding $X$ as the trivial biproduct of $\{X\}$, (2.6.108) identifies $\mathrm{Mor}(X, X \oplus X)$ with $2 \times 1$ matrices and $\mathrm{Mor}(X \oplus X, X)$ with $1 \times 2$ matrices over $\mathrm{Mor}(X, X)$. The **diagonal** $\Delta_X : X \to X \oplus X$ and **codiagonal** (or **summation map**) $\nabla_X : X \oplus X \to X$ are the morphisms with matrices*

© David Rufino CC BY-NC-ND 4.0

$$\Delta_X = \begin{pmatrix} 1_X \\ 1_X \end{pmatrix} \qquad \nabla_X = \begin{pmatrix} 1_X & 1_X \end{pmatrix}$$

*that is, the unique morphisms with* $\pi_1 \circ \Delta_X = \pi_2 \circ \Delta_X = 1_X$ *and* $\nabla_X \circ s_1 = \nabla_X \circ s_2 = 1_X$ *respectively.*

**Remark 2.6.117**
*In* **AbGrp**, *where* $X \oplus X$ *is the direct sum* $X \times X$ *with* $s_1(x) = (x, 0)$, $s_2(x) = (0, x)$, $\pi_1(x, y) = x$, $\pi_2(x, y) = y$, *the codiagonal is exactly the addition map:* $\nabla_X(x, 0) = x$ *and* $\nabla_X(0, y) = y$ *force, by additivity of the homomorphism* $\nabla_X$, $\nabla_X(x, y) = \nabla_X(x, 0) + \nabla_X(0, y) = x + y$. *Dually the diagonal is* $\Delta_X(x) = (x, x)$.

**Definition 2.6.118 (Sum of Parallel Morphisms)**
*Let* $\mathcal{C}$ *be a pointed category with binary biproducts and* $f, g : X \to Y$. *We define*

$$f + g \coloneqq \begin{pmatrix} f & g \end{pmatrix} \circ \Delta_X = \nabla_Y \circ \begin{pmatrix} f & 0 \\ 0 & g \end{pmatrix} \circ \Delta_X = \nabla_Y \circ \begin{pmatrix} f \\ g \end{pmatrix} : X \to Y$$

*the three expressions agreeing by (2.6.114), since* $\begin{pmatrix} f & 0 \\ 0 & g \end{pmatrix}$ *has no cross terms.*

**Proposition 2.6.119 (Interchange Law)**
*Let* $\mathcal{C}$ *be a pointed category with binary biproducts and* $w, x, y, z : X \to Y$. *Then* $(w + x) + (y + z) = (w + y) + (x + z)$.

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.* Following [Fre64] Theorem 2.38.

Let $M \coloneqq \begin{pmatrix} w & x \\ y & z \end{pmatrix} : X \oplus X \to Y \oplus Y$ (2.6.108). Then

$$M \circ \Delta_X \overset{(2.6.112)}{=} \begin{pmatrix} w & x \\ y & z \end{pmatrix} \circ \Delta_X \overset{(2.6.113)}{=} \begin{pmatrix} w & x \end{pmatrix} \circ \Delta_X \\ \overset{(2.6.118)}{=} \begin{pmatrix} w + x \\ y + z \end{pmatrix}$$

$$\nabla_Y \circ M \overset{(2.6.112)}{=} \nabla_Y \circ \begin{pmatrix} \begin{pmatrix} w \\ y \end{pmatrix} & \begin{pmatrix} x \\ z \end{pmatrix} \end{pmatrix} \overset{(2.6.113)}{=} \begin{pmatrix} \nabla_Y \circ \begin{pmatrix} w \\ y \end{pmatrix} & \nabla_Y \circ \begin{pmatrix} x \\ z \end{pmatrix} \end{pmatrix} \\ \overset{(2.6.118)}{=} \begin{pmatrix} w + y & x + z \end{pmatrix}$$

so, applying $\nabla_Y \circ -$ to the first display and $- \circ \Delta_X$ to the second (each a further instance of (2.6.118))

$$(w + x) + (y + z) = \nabla_Y \circ (M \circ \Delta_X) = (\nabla_Y \circ M) \circ \Delta_X = (w + y) + (x + z)$$

■

**Remark 2.6.120 (Eckmann--Hilton Phenomenon)**
*The Interchange Law is an instance of the classical* ***Eckmann–Hilton argument****: whenever a set carries two unital binary operations* $*, \cdot$ *satisfying the interchange law* $(a * b) \cdot (c * d) = (a \cdot c) * (b \cdot d)$ *and sharing a common unit, the two operations coincide and are commutative and associative. Here the two operations are the row-first and column-first ways of combining the block* $\begin{pmatrix} w & x \\ y & z \end{pmatrix}$ *into a single morphism via* $\Delta_X, \nabla_Y$; *(2.6.118) already records that they coincide (as a single well-defined* $+$*), and (2.6.119) is exactly the interchange law for it. Commutativity and associativity, proved below, are the usual Eckmann–Hilton consequences, recovered by the substitutions* $w = z \coloneqq 0$ *and* $y \coloneqq 0$ *respectively.*

**Proposition 2.6.121 (**$\mathrm{Mor}(X, Y)$ **is a Commutative Monoid, and Composition is Bilinear)**
*Let* $\mathcal{C}$ *be a pointed category with binary biproducts. For objects* $X, Y$, $(\mathrm{Mor}(X, Y), +)$ *is a commutative monoid with identity* $0$. *Moreover, for* $f, g : X \to Y$, $h : Y \to Z$ *and* $k : W \to X$

$$h \circ (f + g) = h \circ f + h \circ g \qquad (f + g) \circ k = f \circ k + g \circ k$$

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

*Identity.* Since $\begin{pmatrix} f & 0 \end{pmatrix}$ and $\begin{pmatrix} 0 & f \end{pmatrix}$ each have a single nonzero entry, (2.6.114) gives

$$f + 0 = \begin{pmatrix} f & 0 \end{pmatrix} \circ \begin{pmatrix} 1_X \\ 1_X \end{pmatrix} = f \circ 1_X = f \qquad 0 + f = \begin{pmatrix} 0 & f \end{pmatrix} \circ \begin{pmatrix} 1_X \\ 1_X \end{pmatrix} = f \circ 1_X = f$$

*Commutativity.* By (2.6.119), setting $w \coloneqq 0$, $x \coloneqq f$, $y \coloneqq g$, $z \coloneqq 0$ gives $(0 + f) + (g + 0) = (0 + g) + (f + 0)$, that is $f + g = g + f$ using the identity just proved.

*Associativity.* For a third morphism $h : X \to Y$, setting $w \coloneqq f$, $x \coloneqq g$, $y \coloneqq 0$, $z \coloneqq h$ in (2.6.119) gives $(f + g) + (0 + h) = (f + 0) + (g + h)$, that is $(f + g) + h = f + (g + h)$.

*Bilinearity.*

© David Rufino CC BY-NC-ND 4.0

$$h \circ (f+g) \overset{(2.6.118)}{=} h \circ (f \;\; g) \circ \Delta_X \overset{(2.6.113)}{=} (h \circ f \;\; h \circ g) \circ \Delta_X \overset{(2.6.118)}{=} h \circ f + h \circ g$$

and dually

$$(f+g) \circ k \overset{(2.6.118)}{=} \nabla_Y \circ \begin{pmatrix} f \\ g \end{pmatrix} \circ k \overset{(2.6.113)}{=} \nabla_Y \circ \begin{pmatrix} f \circ k \\ g \circ k \end{pmatrix} \overset{(2.6.118)}{=} f \circ k + g \circ k$$

■

**Proposition 2.6.122 (Composition is Matrix Multiplication)**
*Let $\mathcal{C}$ be a pointed category with binary biproducts, $I, J, K$ finite sets, $M : \bigoplus_{j\in J} X_j \to \bigoplus_{i\in I} Y_i$ with matrix $(M_{ij})$ and $N : \bigoplus_{k\in K} W_k \to \bigoplus_{j\in J} X_j$ with matrix $(N_{jk})$. Then*

$$(M \circ N)_{ik} = \sum_{j\in J} M_{ij} \circ N_{jk} \quad \text{for all } i \in I, k \in K$$

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

We first show that any column $C : Z \to \bigoplus_{j\in J} X_j$, with entries $C_j \coloneqq \pi_j \circ C$, satisfies $C = \sum_{j\in J} s_j \circ C_j$. Indeed, for each $i \in J$, by (2.6.121) (bilinearity, iterated over the finite sum)

$$\pi_i \circ \sum_{j\in J} s_j \circ C_j = \sum_{j\in J} \pi_i \circ s_j \circ C_j = \sum_{j\in J} \delta_{ij} C_j = C_i$$

using $\pi_i \circ s_j = \delta_{ij}$ (2.6.103) and that every term with $j \neq i$ vanishes (composing with the zero morphism), so this matches the matrix of $C$ entrywise, whence $\sum_{j\in J} s_j \circ C_j = C$ by (2.6.108). Dually, any row $R : \bigoplus_{i\in I} Y_i \to Z$ satisfies $R = \sum_{i\in I} R_i \circ \pi_i$, where $R_i \coloneqq R \circ s_i$.

Now fix $i \in I$, $k \in K$. The $k$-th column of $N$ is $N \circ s_k$, with entries $N_{jk} = \pi_j \circ N \circ s_k$, so by the column decomposition, $N \circ s_k = \sum_{j\in J} s_j \circ N_{jk}$. Hence, using bilinearity to distribute $M \circ -$ and then $\pi_i \circ -$ over the finite sum

$$(M \circ N)_{ik} = \pi_i \circ M \circ N \circ s_k = \pi_i \circ M \circ \sum_{j\in J} s_j \circ N_{jk} = \sum_{j\in J} \pi_i \circ M \circ s_j \circ N_{jk} = \sum_{j\in J} M_{ij} \circ N_{jk}$$

■

## 2.6.15 Direct Limit / Directed Colimit

Here we consider only a certain type of colimit, which is referred to variously as **direct limit** [AM69], **limite inductive** [Gro64] or **filtered colimit** [Lan98]. We also restrict to the case where the indexing category is "thin", which is more properly refered to as a directed colimit.

**Definition 2.6.123 (Directed Preorder)**
*We say a category $I$ is a **directed preorder** if it is small, thin and filtered, that is for any $i, j \in I$*

- *there exists at most one morphism $i \to j$, and*
- *there exists $k \in I$ and morphisms $i \to k$ and $j \to k$*

*If there exists a morphism $i \to j$ then we write $i \prec j$.*

**Example 2.6.124**
*A typical example is any family of subsets of a set $X$ ordered by reversion inclusion which are closed under finite intersection.*

**Definition 2.6.125 (Direct limit / Directed colimit)**
*Let $I$ be a directed preorder and $F : I \to \mathcal{C}$ a diagram.*

*A **cocone** over $F$ is a pair $(L, \{\mu : F(i) \to L\}_{i\in I})$ such that*

$$i \prec j \implies \mu_j \circ F(i \to j) = \mu_i$$

*The cocones over $F$ constitute a category $(F \downarrow \Delta)$ where morphisms $\alpha : (L, \{\mu_i\}) \to (L', \{\mu'_i\})$ are simply morphisms $\alpha : L \to L'$ such that $\alpha \circ \mu_i = \mu'_i$ for all $i \in I$.*

*A cone $\left(L, \{\mu_i\}_{i\in I}\right)$ over $F$ is a **direct limit** for $F$ if it is an **initial object** (2.6.48) in the cocone category $(F \downarrow \Delta)$. Explicitly for every cocone $\left(L', \{\mu'_i\}_{i\in I}\right)$ there is a unique morphism $\alpha : L \to L'$ such that $\alpha \circ \mu_i = \mu'_i$ for all $i \in I$.*

© David Rufino CC BY-NC-ND 4.0

*We denote this by* $\varinjlim_{i\in I} F(i)$.

**Definition 2.6.126 (Cofinal Subpreorder)**
*Let $I$ be a directed preorder. Then a full subcategory $I' \subset I$ is said to be* ***cofinal*** *if*

$$\forall i \in I \quad \exists i' \in I' \text{ s.t. } i \leq i'$$

*In this case $I'$ is also directed.*

**Proposition 2.6.127 (Cofinal Direct Limit)**
*Let $I$ be a directed preorder and $I'$ a full subcategory which is also directed. Suppose $F : I \to \mathcal{C}$ is a diagram then there is a unique morphism of cocones*

$$\varinjlim_{i\in I'} F(i) \longrightarrow \varinjlim_{i\in I} F(i)$$

*provided both these exist.*

*If $I'$ is cofinal in $I$ then this is an isomorphism.*

**Proposition 2.6.128 (Direct Limits of Sets)**
*Let $I$ be a directed preorder and $F : I \to \mathbf{Set}$ a diagram. Consider*

$$L := \{(i, x) \mid i \in I, x \in F(i)\}/\sim$$

*where $(i, x) \sim (j, y)$ if for some $k \in I$ we have $i \prec k, j \prec k$ and $F(i \to k)(x) = F(j \to k)(y)$, and let $\mu_i : F(i) \to L$, $\mu_i(x) := [(i, x)]$.*

*Then $\sim$ is an equivalence relation, and $\left(L, \{\mu_i\}_{i\in I}\right)$ is a direct limit for $F$ (2.6.125), that is $L = \varinjlim_{i\in I} F(i)$. Moreover the $\mu_i$ are jointly surjective, that is every element of $L$ is of the form $\mu_i(x)$ for some $i \in I$, $x \in F(i)$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

**$\sim$ is an equivalence relation.** Reflexivity and symmetry are immediate, using $i \prec i$ (via the identity morphism). For transitivity, suppose $(i, x) \sim (j, y)$ via $k$ (so $i, j \prec k$, $F(i \to k)(x) = F(j \to k)(y)$) and $(j, y) \sim (l, z)$ via $m$ (so $j, l \prec m$, $F(j \to m)(y) = F(l \to m)(z)$). As $I$ is directed there is $n$ with $k, m \prec n$, and applying $F(k \to n)$, resp. $F(m \to n)$, to the two witnessing equations and using functoriality of $F$,

$$F(i \to n)(x) = F(k \to n)(F(i \to k)(x)) = F(k \to n)(F(j \to k)(y)) = F(j \to n)(y)$$

$$F(j \to n)(y) = F(m \to n)(F(j \to m)(y)) = F(m \to n)(F(l \to m)(z)) = F(l \to n)(z)$$

so $F(i \to n)(x) = F(l \to n)(z)$, that is $(i, x) \sim (l, z)$ via $n$.

**Joint surjectivity.** Immediate, as every element of $L$ is by construction the class $[(i, x)] = \mu_i(x)$ of some pair.

**$(L, \{\mu_i\})$ is a cocone.** For $i \prec j$, taking $k := j$ in the equivalence relation (using $F(j \to j) = 1_{F(j)}$) gives $(i, x) \sim (j, F(i \to j)(x))$, that is $\mu_j(F(i \to j)(x)) = \mu_i(x)$, as required by (2.6.125).

**$(L, \{\mu_i\})$ is initial.** Let $(L', \{\mu'_i\})$ be any cocone over $F$. Define $\alpha : L \to L'$ by $\alpha([(i, x)]) := \mu'_i(x)$; this is well-defined, for if $(i, x) \sim (j, y)$ via $k$ then, using the cocone condition for $\mu'$,

$$\mu'_i(x) = \mu'_k(F(i \to k)(x)) = \mu'_k(F(j \to k)(y)) = \mu'_j(y)$$

and by construction $\alpha \circ \mu_i = \mu'_i$ for all $i$. Any morphism of cocones $\beta : L \to L'$ satisfies $\beta(\mu_i(x)) = \mu'_i(x)$ for all $i, x$, so $\beta = \alpha$ by joint surjectivity of the $\mu_i$; hence $\alpha$ is the unique such morphism, and $(L, \{\mu_i\})$ is initial in $(F \downarrow \Delta)$, that is a direct limit for $F$. ∎

## 2.6.16 Inverse Limit

**Definition 2.6.129 (Inverse Limit / Directed Limit)**
*Let $I$ be a directed preorder and $F : I^{op} \to \mathcal{C}$ a diagram – equivalently, an* ***inverse system*** *indexed by $I$: for $i \prec j$ in $I$, a* ***restriction*** *morphism $\rho_{ij} : F(j) \to F(i)$ in $\mathcal{C}$, with $\rho_{ii} = 1_{F(i)}$ and $\rho_{ik} = \rho_{ij} \circ \rho_{jk}$ whenever $i \prec j \prec k$.*

*A limit for $F$ in the sense of (2.6.78) (applied to the small category $I^{op}$) is called an* ***inverse limit*** *for the system, denoted $\varprojlim_{i\in I} F(i)$: explicitly, a cone $\left(L, \{\lambda_i : L \to F(i)\}_{i\in I}\right)$ with $\rho_{ij} \circ \lambda_j = \lambda_i$ for every $i \prec j$, terminal among such cones.*

© David Rufino CC BY-NC-ND 4.0

**Proposition 2.6.130 (Inverse Limits of Sets)**
*Let $I$ be a directed preorder and $F : I^{op} \to \mathbf{Set}$ an inverse system, with restriction maps $\rho_{ij} : F(j) \to F(i)$ $(i \prec j)$. Then*

$$\lim_{\longleftarrow i\in I} F(i) = \left\{ (x_i)_{i\in I} \in \prod_{i\in I} F(i) \;\middle|\; \rho_{ij}(x_j) = x_i \text{ for all } i \prec j \right\},$$

*with projections $\lambda_i\Big((x_j)_j\Big) := x_i$.*

*Proof.* This is (2.6.85) applied to the diagram $F : I^{\mathrm{op}} \to \mathbf{Set}$ ($I^{\mathrm{op}}$ a small category, as $I$ is). ∎

**Proposition 2.6.131 (Cofinal Inverse Limit)**
*Let $I$ be a directed preorder, $I' \subset I$ a full subcategory which is also directed, and $F : I^{op} \to \mathcal{C}$ an inverse system, with restriction maps $\rho_{ij}$ ($i \prec j$ in $I$). Then, provided both exist, there is a unique morphism of cones*

$$\lim_{\longleftarrow i\in I} F(i) \longrightarrow \lim_{\longleftarrow i\in I'} F(i)$$

*(restricting the limit cone over $I$ to the subcategory $I'$).*

*If $I'$ is cofinal in $I$ (2.6.126) then this is an isomorphism.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Write $L' := \lim_{\longleftarrow i\in I'} F(i)$, with projections $\lambda'_i : L' \to F(i)$ $(i \in I')$. The cone $\left(L', \{\lambda'_i\}_{i\in I'}\right)$ over $F|_{I'^{\mathrm{op}}}$ is in particular a cone over all of $F$ restricted along the inclusion $I'^{\mathrm{op}} \subset I^{\mathrm{op}}$, so composing with the (terminal, by definition of $\lim_{\longleftarrow i\in I} F(i)$) limit cone's universal property in the other direction gives nothing yet – rather, the stated morphism is simply the unique one afforded by terminality of $L'$ among cones over $F|_{I'^{\mathrm{op}}}$, applied to the cone $\left(\lim_{\longleftarrow i\in I} F(i), \{\lambda_i\}_{i\in I'}\right)$ (the limit cone over $I$, restricted in its indexing set to $I'$ – still a cone over $F|_{I'^{\mathrm{op}}}$, by the cone condition on $I$ restricted to $i \prec j$ both in $I'$): this gives the unique morphism $\lim_{\longleftarrow i\in I} F(i) \to L'$ of the statement.

Now suppose $I'$ is cofinal. For $i \in I$, choose (by cofinality) $i' \in I'$ with $i \prec i'$, and define $\mu_i := \rho_{i,i'} \circ \lambda'_{i'} : L' \to F(i)$.

**$\mu_i$ is independent of the choice of $i'$.** If $i \prec i'_1, i'_2$ with $i'_1, i'_2 \in I'$, directedness of $I'$ gives $i'_3 \in I'$ with $i'_1, i'_2 \prec i'_3$; then, using the cone condition for $\lambda'$ (within $I'$, at $i'_1 \prec i'_3$) and the restriction composition law of (2.6.129) (at $i \prec i'_1 \prec i'_3$),

$$\rho_{i,i'_1} \circ \lambda'_{i'_1} = \rho_{i,i'_1} \circ \rho_{i'_1,i'_3} \circ \lambda'_{i'_3} = \rho_{i,i'_3} \circ \lambda'_{i'_3},$$

and symmetrically $\rho_{i,i'_2} \circ \lambda'_{i'_2} = \rho_{i,i'_3} \circ \lambda'_{i'_3}$; so both choices agree.

**$\left(L', \{\mu_i\}_{i\in I}\right)$ is a cone over $F$.** Let $i \prec j$ in $I$. Choose $j' \in I'$ with $j \prec j'$; then also $i \prec j'$, so $j'$ may be used to compute both $\mu_i$ and $\mu_j$:

$$\rho_{ij} \circ \mu_j = \rho_{ij} \circ \rho_{j,j'} \circ \lambda'_{j'} = \rho_{i,j'} \circ \lambda'_{j'} = \mu_i,$$

using the restriction composition law at $i \prec j \prec j'$. Moreover for $i \in I'$ itself, taking $i' := i$ (with $\rho_{ii} = 1_{F(i)}$) gives $\mu_i = \lambda'_i$: the cone $\left(L', \{\mu_i\}_{i\in I}\right)$ restricts, on $I'$, to the original cone $\left(L', \{\lambda'_i\}_{i\in I'}\right)$.

**$\left(L', \{\mu_i\}_{i\in I}\right)$ is terminal.** Let $\left(N, \{\nu_i\}_{i\in I}\right)$ be any cone over $F$. Restricting to $I'$ gives a cone over $F|_{I'^{\mathrm{op}}}$, so by terminality of $L'$ there is a unique morphism $\gamma : N \to L'$ with $\lambda'_i \circ \gamma = \nu_i$ for all $i \in I'$. For general $i \in I$, choosing $i' \in I'$ with $i \prec i'$,

$$\mu_i \circ \gamma = \rho_{i,i'} \circ \lambda'_{i'} \circ \gamma = \rho_{i,i'} \circ \nu_{i'} = \nu_i$$

using the cone condition for $\nu$ (over all of $I$) at $i \prec i'$; so $\gamma$ is a morphism of cones $\left(N, \{\nu_i\}_{i\in I}\right) \to \left(L', \{\mu_i\}_{i\in I}\right)$. Any such morphism $\delta$ satisfies, in particular for $i \in I'$ (where $\mu_i = \lambda'_i$), $\lambda'_i \circ \delta = \nu_i$, so $\delta = \gamma$ by the uniqueness already established for $\gamma$. Hence $\left(L', \{\mu_i\}_{i\in I}\right)$ is terminal among cones over $F$ on all of $I$, that is, $\left(L', \{\mu_i\}_{i\in I}\right)$ is $\lim_{\longleftarrow i\in I} F(i)$.

By uniqueness of limits up to unique isomorphism of cones (2.6.78), there is a (unique) isomorphism $\lim_{\longleftarrow i\in I} F(i) \xrightarrow{\sim} L'$ compatible with the projections $\{\mu_i\}_{i\in I}$ (on the $L'$ side) and $\{\lambda_i\}_{i\in I}$ (on the $\lim_{\longleftarrow i\in I} F(i)$ side); restricting this compatibility to $i \in I'$ (where $\mu_i = \lambda'_i$) shows it is compatible with $\{\lambda'_i\}_{i\in I'}$ too, so by the uniqueness clause established in the first paragraph, this isomorphism is exactly the morphism of cones $\lim_{\longleftarrow i\in I} F(i) \to \lim_{\longleftarrow i\in I'} F(i) = L'$ of the statement. ∎

© David Rufino CC BY-NC-ND 4.0

**Corollary 2.6.132 (Functoriality of the Inverse Limit)**
*Let $I$ be a directed preorder, and let $\mathcal{C}^{I^{op}}_{\lim}$ denote the full subcategory of the functor category $\mathcal{C}^{I^{op}}$ (2.6.14) spanned by those inverse systems $F : I^{op} \to \mathcal{C}$ for which $\varprojlim_i F(i)$ exists. Then*

$$\varprojlim_i (-) : \mathcal{C}^{I^{op}}_{\lim} \longrightarrow \mathcal{C}, \quad F \mapsto \varprojlim_i F(i)$$

*is a functor.*

*More precisely: for a natural transformation $\sigma = (\sigma_i : F(i) \to G(i))_{i \in I}$ there is a unique morphism*

$$\varprojlim_i \sigma_i : \varprojlim_i F(i) \longrightarrow \varprojlim_i G(i)$$

*compatible with the projections; and*

*a) for a further natural transformation $\tau = (\tau_i : G(i) \to K(i))_i$ (so $\tau\sigma := (\tau_i \sigma_i)_i$, itself a natural transformation $F \Rightarrow K$),*

$$\varprojlim_i (\tau_i \sigma_i) = \varprojlim_i \tau_i \circ \varprojlim_i \sigma_i;$$

*b) $\varprojlim_i 1_{F(i)} = 1_{\varprojlim_i F(i)}$.*

*Consequently, if every $\sigma_i$ is an isomorphism, so is $\varprojlim_i \sigma_i$ (2.6.42).*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

A natural transformation $\sigma = (\sigma_i)_i$ satisfying $\sigma_i \circ \rho^F_{ij} = \rho^G_{ij} \circ \sigma_j$ for all $i \prec j$ is exactly a morphism $F \Rightarrow G$ of the functor category $\mathcal{C}^{I^{op}}$ (2.6.14) (the naturality square for $\sigma$ at the morphism $j \to i$ of $I^{op}$ corresponding to $i \prec j$), invertible in $\mathcal{C}^{I^{op}}$ exactly when every $\sigma_i$ is an isomorphism (with inverse $(\sigma_i^{-1})_i$). So this is precisely the case $J := I^{op}$ of (2.6.97). ∎

**Corollary 2.6.133 (Inverse Limits Commute with Products)**
*Let $I$ be a directed preorder, $K$ a set, and $(F_k)_{k \in K}$ inverse systems $F_k : I^{op} \to \mathcal{C}$, with limit-cone projections $\lambda_{k,i} : \varprojlim_i F_k(i) \to F_k(i)$, such that $\varprojlim_i F_k(i)$ exists for every $k \in K$; and suppose $\prod_{k \in K} F_k(i)$ exists for every $i \in I$, with product projections $\pi_{k,i} : \prod_{k' \in K} F_{k'}(i) \to F_k(i)$. If $\prod_{k \in K} \varprojlim_i F_k(i)$ exists, with product projections $p_k$, then $\varprojlim_i \left(\prod_{k \in K} F_k(i)\right)$ exists too, with limit-cone projections $\Lambda_i$, and there is a unique morphism*

$$\Phi : \prod_{k \in K} \varprojlim_i F_k(i) \longrightarrow \varprojlim_i \left( \prod_{k \in K} F_k(i) \right)$$

*such that the following commutes for every $k \in K$, $i \in I$*

$$\begin{array}{ccccc}
\prod_{k \in K} \varprojlim_i F_k(i) & & \xrightarrow{\quad\Phi\quad} & & \varprojlim_i \left(\prod_{k \in K} F_k(i)\right) \\
\Big\downarrow{\scriptstyle p_k} & & & & \Big\downarrow{\scriptstyle \Lambda_i} \\
\varprojlim_i F_k(i) & \xrightarrow[\lambda_{k,i}]{} & F_k(i) & \xleftarrow[\pi_{k,i}]{} & \prod_{k \in K} F_k(i)
\end{array}$$

*and $\Phi$ is an isomorphism.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Regard $K$ as a discrete small category and define a bifunctor $H : K \times I^{op} \to \mathcal{C}$ by $H(k,i) := F_k(i)$ (a morphism of $K \times I^{op}$ is a pair $(1_k, \alpha)$ for $\alpha$ a morphism of $I^{op}$, since $K$ is discrete, and $H(1_k, \alpha) := F_k(\alpha)$; this is a bifunctor as each $F_k$ is a functor). By (2.6.81) (limits over a discrete category are products), $\left(\prod_{k \in K} F_k(i), \{\pi_{k,i}\}_k\right)$ is $\lim_{k \in K} H(k,i)$ for every $i \in I^{op}$, and by (2.6.129), $\left(\varprojlim_i F_k(i), \{\lambda_{k,i}\}_i\right)$ is $\lim_{i \in I^{op}} H(k,i)$ for every $k \in K$; both exist by hypothesis, so (writing $L(k) := \varprojlim_i F_k(i)$, $R(i) := \prod_{k \in K} F_k(i)$) (2.6.100) applies to $H$.

As $\left(\prod_{k \in K} \varprojlim_i F_k(i), \{p_k\}_k\right) = \left(\lim_{k \in K} L(k), \{p_k\}_k\right)$ exists by hypothesis, (2.6.100) gives that $\lim_{i \in I^{op}} R(i) = \varprojlim_i \left(\prod_{k \in K} F_k(i)\right)$ also exists, with limit-cone projections $\Lambda_i$ say, together with a unique isomorphism $\Phi$ making the stated diagram commute for every $k, i$ (that proposition's diagram, with $\nu_k := p_k$, $\xi_i := \Lambda_i$, $\lambda_{k,i}$ as here, and $\mu_{k,i} := \pi_{k,i}$). ∎

© David Rufino CC BY-NC-ND 4.0

## 2.6.17 Equaliser and Coequaliser

**Definition 2.6.134 (Equaliser)**
*Consider a commutative diagram in a category $\mathcal{C}$*

$$X \xrightarrow{e} Y \underset{\beta}{\overset{\alpha}{\rightrightarrows}} Z$$

*such that $\alpha \circ e = \beta \circ e$. We say that $(X, e)$ is an **equaliser** for $(\alpha, \beta)$ if for all $e' : X' \to Y$ such that $\alpha \circ e' = \beta \circ e'$ there is a unique factorisation*




*This is equivalent to $(X, e)$ being a limit for the obvious functor $F : \mathbf{2} \to \mathcal{C}$ where $\mathbf{2}$ denotes the category $\cdot \rightrightarrows \cdot$.*

**Proposition 2.6.135**
*Every equaliser is a monomorphism, and every coequaliser is an epimorphism.*

*Proof.* Suppose that $(X, e)$ is an equaliser for $\alpha, \beta : Y \rightrightarrows Z$, and $e \circ f = e \circ g = h$. Then evidentally $\alpha \circ h = \beta \circ h$ so there is a unique morphism $k$ such that $e \circ k = h$, whence $k = f = g$.

The second statement follows by duality. ∎

**Proposition 2.6.136**
*Suppose $(X, e) = \mathrm{eq}(\alpha, \beta)$. Then $\alpha = \beta$ if and only if $e$ is an isomorphism.*

*Proof.* By hypothesis $\alpha \circ e = \beta \circ e$ so if $e$ is an isomorphism $\alpha = \alpha \circ e \circ e^{-1} = \beta \circ e \circ e^{-1} = \beta$ as required.

Conversely if $\alpha = \beta$, then $\alpha \circ 1_Y = \beta \circ 1_Y$ so there is a morphism $i : Y \to X$ such that $e \circ i = 1_Y$. Therefore $e$ is split-epic and monic (2.6.135) and therefore an isomorphism (2.6.36). ∎

**Proposition 2.6.137**
*Let $(X, e)$ be an equaliser for $\alpha, \beta : Y \rightrightarrows Z$. Suppose $\phi : X' \xrightarrow{\sim} X$ be an isomorphism. Then $(X', e \circ \phi)$ is also an equaliser for $(\alpha, \beta)$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

By definition of Equaliser, $(X, e)$ is a limit for the diagram $F : \mathbf{2} \to \mathcal{C}$ determined by $(\alpha, \beta)$, and since $\alpha \circ e = \beta \circ e$, $(X', e \circ \phi)$ is a cone for $F$. As $e \circ \phi = e \circ \phi$, the isomorphism $\phi : X' \to X$ is (tautologically) an isomorphism of cones $(X', e \circ \phi) \to (X, e)$, so by (2.6.79) $(X', e \circ \phi)$ is also a limit for $F$, that is, an equaliser for $(\alpha, \beta)$. ∎

**Proposition 2.6.138**
*Let $(X, e)$ be a coequaliser for $\alpha, \beta : Y \rightrightarrows Z$. Suppose $\phi : X \xrightarrow{\sim} X'$ is an isomorphism. Then $(X', \phi \circ e)$ is also a coequaliser for $(\alpha, \beta)$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

By definition of Coequaliser, $(X, e)$ is a colimit for the diagram $F : \mathbf{2} \to \mathcal{C}$ determined by $(\alpha, \beta)$, and since $e \circ \alpha = e \circ \beta$, $(X', \phi \circ e)$ is a co-cone for $F$. As $\phi \circ e = \phi \circ e$, the isomorphism $\phi : X \to X'$ is (tautologically) an isomorphism of co-cones $(X, e) \to (X', \phi \circ e)$, so by (2.6.90) $(X', \phi \circ e)$ is also a colimit for $F$, that is, a coequaliser for $(\alpha, \beta)$. ∎

**Proposition 2.6.139**
*A diagram in* **Set**

$$X \xrightarrow{i} Y \underset{\beta}{\overset{\alpha}{\rightrightarrows}} Z$$

*is an equaliser if and only if $i$ is injective and*

$$\mathrm{Im}(i) = \{y \in Y \mid \alpha(y) = \beta(y)\}$$

© David Rufino CC BY-NC-ND 4.0

**Definition 2.6.140 (Coequaliser)**
*Consider a commutative diagram in a category $\mathcal{C}$*

$$X \underset{\beta}{\overset{\alpha}{\rightrightarrows}} Y \xrightarrow{\pi} Z$$

*such that $\pi \circ \alpha = \pi \circ \beta$. We say that $\pi$ is a **co-equaliser** for $(\alpha, \beta)$ if for all $\pi' : Y \to Z'$ such that $\pi' \circ \alpha = \pi' \circ \beta$ there is a unique factorisation*

$$\begin{array}{ccccc} X & \underset{\beta}{\overset{\alpha}{\rightrightarrows}} & Y & \xrightarrow{\pi} & Z \\ & & & \searrow^{\pi'} & \downarrow \\ & & & & Z' \end{array}$$

*This is equivalent to $(Z, \pi)$ being a colimit for the obvious functor $F : \mathbf{2} \to \mathcal{C}$ where $\mathbf{2}$ denotes the category $\cdot \rightrightarrows \cdot$.*

**Proposition 2.6.141**
*The following diagram is a coequaliser*

$$X \underset{\beta}{\overset{\alpha}{\rightrightarrows}} Y \xrightarrow{\pi} Z$$

*if and only if the following is an equaliser in* **Set** *for all $W \in \mathcal{C}$*

$$\mathrm{Mor}(Z, W) \xrightarrow{\mathrm{Mor}(\pi, W)} \mathrm{Mor}(Y, W) \underset{\mathrm{Mor}(\beta, W)}{\overset{\mathrm{Mor}(\alpha, W)}{\rightrightarrows}} \mathrm{Mor}(X, W)$$

*Proof.* We see from the definition that $\pi$ is a coequaliser for $(\alpha, \beta)$ if and only if $\mathrm{Mor}(\pi, W)$ is injective and

$$\mathrm{Im}(\mathrm{Mor}(\pi, W)) = \{\pi' : Y \to W \mid \pi' \circ \alpha = \pi' \circ \beta\}$$

which is exactly the criteria (2.6.139) for an equaliser in **Set**. ∎

**Proposition 2.6.142 (Left Adjoints Preserve Coequalisers)**
*Let $\mathcal{C} \underset{G}{\overset{F}{\rightleftarrows}} \mathcal{D}$ a pair of adjoint functors. Then $F$ preserves coequalisers and $G$ preserves equalisers.*

*Proof.* Consider a coequaliser

$$X \underset{\beta}{\overset{\alpha}{\rightrightarrows}} Y \xrightarrow{\pi} Z$$

By the adjoint relationship we have a commutative diagram for all $D \in \mathcal{D}$

$$\begin{array}{ccccc} \mathrm{Mor}(F(Y), D) & \xrightarrow{\mathrm{Mor}(F(\pi), D)} & \mathrm{Mor}(F(X), D) & \underset{\mathrm{Mor}(F(\beta), D)}{\overset{\mathrm{Mor}(F(\alpha), D)}{\rightrightarrows}} & \mathrm{Mor}(F(Z), D) \\ \downarrow \sim & & \downarrow \sim & & \downarrow \sim \\ \mathrm{Mor}(Y, G(D)) & \xrightarrow{\mathrm{Mor}(\pi, G(D))} & \mathrm{Mor}(X, G(D)) & \underset{\mathrm{Mor}(\beta, G(D))}{\overset{\mathrm{Mor}(\alpha, G(D))}{\rightrightarrows}} & \mathrm{Mor}(Y, G(D)) \end{array}$$

By (2.6.141) the bottom row is an equaliser, and so by the isomorphisms so is the top row. Therefore we conclude by the same result that the following is an equaliser as required

$$F(X) \underset{F(\beta)}{\overset{F(\alpha)}{\rightrightarrows}} F(Y) \xrightarrow{F(\pi)} F(Z)$$

∎

**Proposition 2.6.143 (Coequalisers in a Reflective Subcategory)**
*Let $\mathcal{C} \underset{G}{\overset{F}{\rightleftarrows}} \mathcal{D}$ be adjoint functors with $G$ fully faithful and unit $\eta$; by (2.6.73) we may regard $\mathcal{D}$ as a full subcategory*

© David Rufino CC BY-NC-ND 4.0

*of $\mathcal{C}$ via $G$. Let $\alpha, \beta : X \rightrightarrows Y$ be morphisms of $\mathcal{D}$ and suppose $\pi : Y \to Z$ is a coequaliser of $(\alpha, \beta)$ in $\mathcal{C}$. Then $\eta_Z \circ \pi : Y \to F(Z)$ is a coequaliser of $(\alpha, \beta)$ in $\mathcal{D}$.*

*In particular if every pair of morphisms of $\mathcal{D}$ has a coequaliser in $\mathcal{C}$ then $\mathcal{D}$ has all coequalisers.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Write $q \coloneqq \eta_Z \circ \pi$. Since $\pi \circ \alpha = \pi \circ \beta$ we have $q \circ \alpha = \eta_Z \circ \pi \circ \alpha = \eta_Z \circ \pi \circ \beta = q \circ \beta$.

Suppose $q' : Y \to H$ is a morphism of $\mathcal{D}$ with $q' \circ \alpha = q' \circ \beta$. Regarding $q'$ as a morphism of $\mathcal{C}$ (as $\mathcal{D}$ is a subcategory), the universal property of the coequaliser $\pi$ gives a unique morphism $\tilde{q} : Z \to H$ of $\mathcal{C}$ such that $\tilde{q} \circ \pi = q'$.

Since $F$ is left adjoint to $G$ with unit $\eta$ (2.6.71), there is a bijection $\mathrm{Mor}(F(Z), H) \xrightarrow{\sim} \mathrm{Mor}(Z, H)$, $\phi \mapsto \phi \circ \eta_Z$, so there is a unique morphism $\overline{q} : F(Z) \to H$ of $\mathcal{D}$ such that $\overline{q} \circ \eta_Z = \tilde{q}$. Then

$$\overline{q} \circ q = \overline{q} \circ \eta_Z \circ \pi = \tilde{q} \circ \pi = q'$$

so $\overline{q}$ is a factorisation of $q'$ through $q$.

Suppose $\overline{q}'$ is another morphism of $\mathcal{D}$ with $\overline{q}' \circ q = q'$, that is $(\overline{q}' \circ \eta_Z) \circ \pi = q' = \tilde{q} \circ \pi$. As $\pi$ is a coequaliser it is an epimorphism (2.6.135), whence $\overline{q}' \circ \eta_Z = \tilde{q}$, and by the uniqueness of the factorisation above $\overline{q}' = \overline{q}$.

Therefore $q = \eta_Z \circ \pi$ satisfies the universal property of the coequaliser of $(\alpha, \beta)$ in $\mathcal{D}$.

For the final statement, if every pair of morphisms of $\mathcal{D}$ has a coequaliser in $\mathcal{C}$ then the above applies to every such pair, so $\mathcal{D}$ has all coequalisers. ∎

## 2.6.18 Pullback and Kernel Pairs

**Definition 2.6.144 (Pullback)**
*Let $f : X \to Z$, $g : Y \to Z$ be morphisms in $\mathcal{C}$. We say $(W, p_1, p_2)$, with $p_1 : W \to X$ and $p_2 : W \to Y$, is a **pullback** of $(f, g)$ if $f \circ p_1 = g \circ p_2$, and for every $(W', p_1', p_2')$ with $f \circ p_1' = g \circ p_2'$ there is a unique morphism $\phi : W' \to W$ such that $p_1 \circ \phi = p_1'$ and $p_2 \circ \phi = p_2'$*




*This is equivalent to $(W, \{p_1, p_2\})$ being a limit for the evident diagram $F : \mathbf{\Lambda} \to \mathcal{C}$ determined by $(f, g)$, where $\mathbf{\Lambda} \coloneqq \{a \to c \leftarrow b\}$ denotes the category with three objects and, besides identities, two further morphisms $a \to c$, $b \to c$. We write $W =: X \times_Z Y$.*

**Proposition 2.6.145 (Pullback from Products and Equalisers)**
*Let $f : X \to Z$, $g : Y \to Z$ be morphisms such that $X \times Y$ exists with projections $\pi_X, \pi_Y$, and such that $(W, e) \coloneqq \mathrm{eq}(f \circ \pi_X, g \circ \pi_Y : X \times Y \to Z)$ exists. Then $(W, \pi_X \circ e, \pi_Y \circ e)$ is a pullback of $(f, g)$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Write $p_1 \coloneqq \pi_X \circ e$, $p_2 \coloneqq \pi_Y \circ e$. By definition of the equaliser, $f \circ \pi_X \circ e = g \circ \pi_Y \circ e$, that is $f \circ p_1 = g \circ p_2$.

Suppose $(W', p_1', p_2')$ satisfies $f \circ p_1' = g \circ p_2'$. By the universal property of the product there is a unique $h : W' \to X \times Y$ with $\pi_X \circ h = p_1'$, $\pi_Y \circ h = p_2'$. Then $f \circ \pi_X \circ h = f \circ p_1' = g \circ p_2' = g \circ \pi_Y \circ h$, so $h$ equalises $(f \circ \pi_X, g \circ \pi_Y)$, whence by the universal property of the equaliser there is a unique $\phi : W' \to W$ with $e \circ \phi = h$. Then $p_1 \circ \phi = \pi_X \circ e \circ \phi = \pi_X \circ h = p_1'$, and similarly $p_2 \circ \phi = p_2'$.

If $\phi'$ also satisfies $p_1 \circ \phi' = p_1'$, $p_2 \circ \phi' = p_2'$, then $\pi_X \circ (e \circ \phi') = p_1' = \pi_X \circ h$ and $\pi_Y \circ (e \circ \phi') = p_2' = \pi_Y \circ h$, so by uniqueness in the universal property of the product, $e \circ \phi' = h$; as $e$ is monic (2.6.135), $\phi' = \phi$. ∎

**Definition 2.6.146 (Kernel Pair)**
*Let $f : X \to Y$ be a morphism. A **kernel pair** of $f$ is a pullback $(Z, p_1, p_2)$ of $(f, f)$, that is $p_1, p_2 : Z \to X$ with $f \circ p_1 = f \circ p_2$, universal among such pairs*

© David Rufino CC BY-NC-ND 4.0

$$\begin{array}{ccc} Z & \xrightarrow{p_2} & X \\ {\scriptstyle p_1}\downarrow & & \downarrow{\scriptstyle f} \\ X & \xrightarrow[f]{} & Y \end{array}$$

*In* **Set**, *the kernel pair of $f$ is precisely the relation $\{(x,x') \in X \times X \mid f(x) = f(x')\} \subseteq X \times X$, together with its two projections: it records exactly which pairs of elements $f$ identifies. So $f$ is injective if and only if this relation is trivial, i.e. reduces to the diagonal $\{(x,x) \mid x \in X\}$ – this is the picture behind (2.6.147) below, which characterises monomorphisms in general as those morphisms whose kernel pair is trivial.*

**Proposition 2.6.147 (Monomorphisms via Kernel Pairs)**
*Let $\mu : X \to Y$ be a morphism with a kernel pair $(Z, p_1, p_2)$. Then $\mu$ is monic if and only if $p_1 = p_2$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

($\Leftarrow$) Suppose $p_1 = p_2$. If $a, b : W \to X$ satisfy $\mu \circ a = \mu \circ b$, then by the universal property of the kernel pair there is a unique $\psi : W \to Z$ with $p_1 \circ \psi = a$ and $p_2 \circ \psi = b$; since $p_1 = p_2$ this gives $a = p_1 \circ \psi = p_2 \circ \psi = b$. So $\mu$ is monic.

($\Rightarrow$) Suppose $\mu$ is monic. Since $\mu \circ p_1 = \mu \circ p_2$ by definition of the kernel pair (2.6.146), monicity of $\mu$ gives $p_1 = p_2$. ■

**Corollary 2.6.148 (Monomorphisms have Trivial Kernel Pair)**
*Let $\mu : X \to Y$ be a morphism. Then $\mu$ is monic if and only if $(X, 1_X, 1_X)$ is a kernel pair of $\mu$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

($\Leftarrow$) This is the case $Z \coloneqq X$, $p_1 = p_2 \coloneqq 1_X$ of (2.6.147) ($\Leftarrow$).

($\Rightarrow$) Suppose $\mu$ is monic. Given $a, b : W \to X$ with $\mu \circ a = \mu \circ b$, monicity gives $a = b$; then $\phi \coloneqq a$ satisfies $1_X \circ \phi = a$ and $1_X \circ \phi = b$ (as $a = b$), and is the unique such morphism, since any $\phi'$ with $1_X \circ \phi' = a$ must equal $a$. So $(X, 1_X, 1_X)$ satisfies the universal property of a kernel pair of $\mu$ (2.6.146). ■

**Proposition 2.6.149 (Regular Epimorphism is the Coequaliser of its Kernel Pair)**
*Let $e : X \to Y$ be a regular epimorphism with kernel pair $(Z, p_1, p_2)$. Then $e$ is the coequaliser of $(p_1, p_2)$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Since $e$ is regular, by (2.6.158) there are morphisms $\gamma, \delta : W \rightrightarrows X$ such that $e$ is the coequaliser of $(\gamma, \delta)$. Since $e \circ \gamma = e \circ \delta$, the universal property of the kernel pair $(Z, p_1, p_2)$ gives a unique morphism $h : W \to Z$ with $p_1 \circ h = \gamma$ and $p_2 \circ h = \delta$.

By definition of the kernel pair, $e \circ p_1 = e \circ p_2$. Suppose $k : X \to V$ satisfies $k \circ p_1 = k \circ p_2$. Then

$$k \circ \gamma = k \circ p_1 \circ h = k \circ p_2 \circ h = k \circ \delta$$

so, as $e$ is the coequaliser of $(\gamma, \delta)$, there is a unique morphism $l : Y \to V$ such that $l \circ e = k$. Since $e$ is (in particular) an epimorphism (2.6.158), this $l$ is the unique morphism satisfying $l \circ e = k$ among all morphisms $Y \to V$, in particular the unique one factoring $k$ through $e$. So $e$ satisfies the universal property of the coequaliser of $(p_1, p_2)$. ■

## 2.6.19 Kernels and Cokernels

**Proposition 2.6.150 (Kernel)**
*Let $\mathcal{C}$ be a pointed category (2.6.54). Suppose we have a sequence*

$$\ker(\alpha) \xrightarrow{i} X \xrightarrow{\alpha} Y$$

*then the following are equivalent*

*a) $i$ is the equaliser of $(\alpha, 0)$*
*b) $\alpha \circ i = 0$ and $\alpha \circ i' = 0 \Longrightarrow \exists!\, e$ s.t. $i \circ e = i'$*

*In this case we say that $(i, \ker(\alpha))$ is the* ***kernel*** *of $\alpha$*

**Lemma 2.6.151 ($f$ monic $\Longrightarrow \ker(f) = 0$)**
*Let $\mathcal{C}$ be a pointed category and $f : X \to Y$ monic. Then $\ker(f) = 0$.*

*Dually $f$ epic $\Longrightarrow \operatorname{coker}(f) = 0$.*

© David Rufino CC BY-NC-ND 4.0

*Proof.* Evidentally $f \circ 0 = 0_{0,X}$ and for any object $X'$ we have a unique morphism $0_{X',0} : X' \to 0$, so satisfies the uniqueness property. ∎

**Remark 2.6.152**
*The converse does not hold, for we may define a monoid homomorphism $f : \mathbb{N} \to \mathbb{N}$ given by $f(n) := \min(n, 1)$. Then $f$ is a homomorphism, $\ker(f) = \{0\}$ but $f$ is not injective.*

**Proposition 2.6.153 (Cokernel)**
*Let $\mathcal{C}$ be a pointed category (2.6.54). Suppose we have a sequence*

$$X \xrightarrow{\alpha} Y \xrightarrow{\pi} \operatorname{coker}(\alpha)$$

*then the following are equivalent*

*a) $\pi$ is the coequaliser of $(\alpha, 0)$,*
*b) $\pi \circ \alpha = 0$ and $\pi' \circ \alpha = 0 \implies \exists! \, h$ s.t. $h \circ \pi = \pi'$,*

*In this case we say that $(\pi, \operatorname{coker}(\alpha))$ is the **cokernel** of $f$*

**Lemma 2.6.154 (Kernels and Cokernels swap under $\mathcal{C}^{\mathrm{op}}$)**
*Let $\mathcal{C}$ be a pointed category and $\alpha : X \to Y$ a morphism, with kernel $\ker(\alpha) \xrightarrow{i} X$ and cokernel $Y \xrightarrow{\pi} \operatorname{coker}(\alpha)$. Regarding $i$ and $\pi$ as morphisms of $\mathcal{C}^{op}$,*

$$(\ker(\alpha), i) \text{ is a cokernel of } \alpha^{op} \text{ in } \mathcal{C}^{op}, \quad (\operatorname{coker}(\alpha), \pi) \text{ is a kernel of } \alpha^{op} \text{ in } \mathcal{C}^{op}.$$

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

The defining property of $(\ker(\alpha), i)$ (2.6.150) – namely $\alpha \circ i = 0$, and for every $i' : X' \to X$ with $\alpha \circ i' = 0$ there is a unique $e : X' \to \ker(\alpha)$ with $i \circ e = i'$ – is, upon reversing every arrow, exactly the defining property (2.6.153) of $(\ker(\alpha), i)$ as a cokernel of $\alpha^{\mathrm{op}}$ in $\mathcal{C}^{\mathrm{op}}$. The dual statement follows similarly. ∎

**Proposition 2.6.155 (Left Adjoints Preserve Cokernels)**
*Let $\mathcal{C}, \mathcal{D}$ be pointed categories and $\mathcal{C} \underset{G}{\overset{F}{\rightleftarrows}} \mathcal{D}$ a pair of adjoint functors.*

*Then $F$ preserves cokernels and $G$ preserves kernels.*

*Proof.* By (2.6.75) we see that $F$ and $G$ preserve zero morphisms. Therefore we are done by (2.6.142) since a cokernel of $f$ is just the coequaliser of $(f, 0)$. ∎

## 2.6.20 Normal, Regular and Extremal Morphisms

In a category which is not balanced, monic and epic are not always the "right" notions to characterise isomorphisms; the following refinements repair this.

**Definition 2.6.156 (Extremal Epimorphism / Monomorphism)**
*An epimorphism $e : X \to Y$ is **extremal** if whenever $e = m \circ f$ for some monomorphism $m$, then $m$ is an isomorphism.*

*Dually, a monomorphism $m : X \to Y$ is **extremal** if whenever $m = f \circ e$ for some epimorphism $e$, then $e$ is an isomorphism.*

**Proposition 2.6.157**
*Let $f : X \to Y$ be a morphism. Then the following are equivalent*

*a) $f$ is an isomorphism*
*b) $f$ is a monomorphism and an extremal epimorphism*
*c) $f$ is an extremal monomorphism and an epimorphism*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Being an **isomorphism** is a self-dual notion, and monic/epic swap under passing to $\mathcal{C}^{\mathrm{op}}$ (2.6.33), so condition c) is precisely condition b) applied in $\mathcal{C}^{\mathrm{op}}$. Hence it suffices to prove a) $\iff$ b); applying this to $\mathcal{C}^{\mathrm{op}}$ then gives a) $\iff$ c).

a) $\implies$ b) An isomorphism is trivially monic and epic; if $f$ is an isomorphism and $f = m \circ g$ then by (2.6.37), $m$ is an isomorphism, so $f$ is an extremal epimorphism.

b) $\implies$ a) Suppose $f$ is monic and an extremal epimorphism. The trivial factorisation $f = f \circ 1_X$ exhibits $f$ as a monomorphism through which $f$ factors, so by extremality $f$ is an isomorphism. ∎

© David Rufino CC BY-NC-ND 4.0

**Definition 2.6.158 (Regular Epimorphism / Monomorphism)**
*An epimorphism $e : X \to Y$ is **regular** if there exist morphisms $\alpha, \beta : W \rightrightarrows X$ such that $e$ is the coequaliser of $(\alpha, \beta)$.*

*Dually, a monomorphism $m : X \to Y$ is **regular** if there exist morphisms $\alpha, \beta : Y \rightrightarrows Z$ such that $m$ is the equaliser of $(\alpha, \beta)$.*

**Definition 2.6.159 (Normal Epimorphism / Monomorphism)**
*Let $\mathcal{C}$ be a pointed category (2.6.54). An epimorphism $e : X \to Y$ is **conormal** if $(e, Y)$ is the cokernel of some morphism $\alpha : W \to X$.*

*Dually, a monomorphism $m : X \to Y$ is **normal** if $(m, X)$ is the kernel of some morphism $\alpha : Y \to Z$.*

**Proposition 2.6.160 (Normal, Regular and Extremal Morphisms)**
*Let $\mathcal{C}$ be a pointed category. Then every normal monomorphism (resp. conormal epimorphism) is regular.*

*In a general category $\mathcal{C}$ every regular monomorphism (resp. epimorphism) is an extremal monomorphism (resp. epimorphism).*

*In summary*

$$\textbf{normal}\ \ (\textbf{conormal}) \implies \textbf{regular} \implies \textbf{extremal} \implies \textbf{monic}\ (\textbf{epic})$$

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

(Normal $\implies$ regular) If $e$ is a normal monomorphism, then by (2.6.150) $e$ is the equaliser of $(\alpha, 0)$ for some $\alpha$, so $e$ is regular.

(Regular $\implies$ extremal) Suppose $m : X \to Y$ is the equaliser of $\alpha, \beta : Y \rightrightarrows W$, and $m = h \circ e$ for some epimorphism $e : X \to Z$. Since $\alpha \circ m = \beta \circ m$ we get $(\alpha \circ h) \circ e = (\beta \circ h) \circ e$, and as $e$ is epic, $\alpha \circ h = \beta \circ h$. Hence $h$ is equalised by $(\alpha, \beta)$, so by the universal property of the equaliser there is a unique $k : Z \to X$ with $m \circ k = h$.




Then $m \circ (k \circ e) = h \circ e = m = m \circ 1_X$, and since equalisers are monic (2.6.135) we may cancel $m$ to get $k \circ e = 1_X$, so $e$ is split-monic. As $e$ is also epic, (2.6.36) gives that $e$ is an isomorphism, and $m$ is extremal.

Dually, every conormal epimorphism is regular and every regular epimorphism is extremal. ∎

For balanced categories we can reverse the implication

**Proposition 2.6.161 (Balanced Categories via Extremal Morphisms)**
*Let $\mathcal{C}$ be a category. Then the following are equivalent*

*a) $\mathcal{C}$ is balanced*
*b) every monomorphism in $\mathcal{C}$ is extremal*
*c) every epimorphism in $\mathcal{C}$ is extremal*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Being **balanced** is a self-dual notion: monic and epic swap under passing to $\mathcal{C}^{\mathrm{op}}$ (2.6.33) while isomorphisms remain isomorphisms, so $\mathcal{C}$ is balanced iff $\mathcal{C}^{\mathrm{op}}$ is. Likewise, by the "Dually" clause of (2.6.156), a morphism is an extremal epimorphism in $\mathcal{C}$ iff it is an extremal monomorphism in $\mathcal{C}^{\mathrm{op}}$. So it suffices to prove a) $\iff$ b); applying this to $\mathcal{C}^{\mathrm{op}}$ then gives a) $\iff$ c).

a) $\implies$ b) Suppose $\mathcal{C}$ is balanced and $m : X \to Y$ is monic. If $m = f \circ e$ for some epimorphism $e : X \to W$, then $e$ is also monic: if $eg_1 = eg_2$ for $g_1, g_2 : Z \to X$ then $f \circ e \circ g_1 = f \circ e \circ g_2$, that is $mg_1 = mg_2$, whence $g_1 = g_2$ as $m$ is monic. As $e$ is both monic and epic, balancedness gives that $e$ is an isomorphism, so $m$ is extremal.

b) $\implies$ a) Suppose every monomorphism is extremal and $f$ is both monic and epic. Then $f$ is an extremal monomorphism, so by (2.6.157) (with $f$ extremal monic and epic) $f$ is an isomorphism. So $\mathcal{C}$ is balanced. ∎

**Lemma 2.6.162**
*Let $\mathcal{C}$ be a pointed category and $f : X \to Y$ a morphism for which $\ker(\mathrm{coker}(f))$ exists. Then the following are equivalent*

© David Rufino CC BY-NC-ND 4.0

*a)* $f$ *is normal*
*b)* $f = \ker(\operatorname{coker}(f))$

*Dually, suppose* $\operatorname{coker}(\ker(f))$ *exists then the following are equivalent*

*a)* $f$ *is conormal*
*b)* $f = \operatorname{coker}(\ker(f))$

*Proof.* b) $\implies$ a) immediate.

a) $\implies$ b) By assumption $\pi : Y \to \operatorname{coker}(f)$ and $i : \ker(\pi) \to Y$ exists so we have a commutative diagram.




By definition of the cokernel $\pi \circ f = 0$ whence there is a unique $\phi : X \to \ker(\pi)$ such that $i \circ \phi = f$.

As $g \circ f = 0$ then by definition there is a unique $\theta : \operatorname{coker}(f) \to Z$ such that $\theta \circ \pi = g$.

Therefore $g \circ i = \theta \circ \pi \circ i = 0$. So by definition there is a unique $\psi : \ker(\pi) \to X$ such that $f \circ \psi = i$.

Then $i \circ \phi \circ \psi = f \circ \psi = i$ and since $i$ is a monomorphism (2.6.135) $\phi \circ \psi = 1_{\ker(\pi)}$. Similarly $f \circ \psi \circ \phi = i \circ \phi = f$ and since $f = \ker(g)$ is a monomorphism (2.6.135) $\psi \circ \phi = 1_X$. Therefore $X$ and $\ker(\pi)$ are isomorphic via $\phi$ and $\psi$, and since $i \circ \phi = f$, by (2.6.137) $X$ is also a kernel for $\pi$ as required.

Dually, $\mathcal{C}^{\mathrm{op}}$ is again pointed with the same zero object (2.6.54), monic/epic swap under passing to $\mathcal{C}^{\mathrm{op}}$ (2.6.33), and $\ker(f)$, $\operatorname{coker}(f)$ computed in $\mathcal{C}$ are precisely $\operatorname{coker}(f)$, $\ker(f)$ computed in $\mathcal{C}^{\mathrm{op}}$ (2.6.154). So applying the equivalence just proved to $f$ regarded as a morphism of $\mathcal{C}^{\mathrm{op}}$ shows $f$ is conormal iff $f = \operatorname{coker}(\ker(f))$. ∎

**Definition 2.6.163**
*We say a pointed category* $\mathcal{C}$ *is*

- ***normal*** *if every monomorphism is normal (i.e. a kernel of some morphism).*
- ***conormal*** *if every epimorphism is conormal (i.e. the cokernel of some morphism).*
- ***binormal*** *if it is both normal and conormal.*

**Corollary 2.6.164**
*A normal (resp. conormal, binormal) category is balanced.*

**Example 2.6.165**
*We will see that the categories* **AbGrp** *and* $A$**-Mod** *are binormal (and therefore every epi and mono is regular), whereas* **Grp** *is only conormal. All three are balanced.*

*We also see may verify directly that in* **Set** *every monomorphism and epimorphism is regular.*

**Proposition 2.6.166**
*Let* $\mathcal{C}$ *be a pointed category and* $f$ *a morphism. Then the following are equivalent*

*a)* $\ker(f) = 0$
*b)* $f \circ g = 0 \implies g = 0$

*Dually, the following are equivalent*

*a)* $\operatorname{coker}(f) = 0$
*b)* $g \circ f = 0 \implies g = 0$

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Write $f : X \to Y$ and $i : \ker(f) \to X$ for the kernel of $f$, so by (2.6.150), $f \circ i = 0$ and for every $i' : W \to X$ with $f \circ i' = 0$ there is a unique $e : W \to \ker(f)$ with $i \circ e = i'$.

a) $\implies$ b) Suppose $\ker(f) = 0$ and $f \circ g = 0$ for some $g : W \to X$. By the universal property there is a unique $e : W \to 0$ with $i \circ e = g$; as $i = 0_{0,X}$ (since $\ker(f) = 0$) and $e = 0_{W,0}$ (since 0 is terminal), $g = i \circ e = 0_{W,X} = 0$.

© David Rufino CC BY-NC-ND 4.0

b) $\implies$ a) Let $\iota : 0 \to X$ be the canonical morphism. Since 0 is initial, $f \circ \iota = 0$ automatically. Given any $i' : W \to X$ with $f \circ i' = 0$, the hypothesis gives $i' = 0$, that is $i' = \iota \circ 0_{W,0}$, and $0_{W,0} : W \to 0$ is the unique such morphism since 0 is terminal. So $(0, \iota)$ satisfies the universal property of $\ker(f)$ (2.6.150), whence $\ker(f) = 0$.

Dually, $\mathcal{C}^{\mathrm{op}}$ is again pointed with the same zero object (2.6.54), and $\ker(f)$, $\operatorname{coker}(f)$ computed in $\mathcal{C}$ are precisely $\operatorname{coker}(f)$, $\ker(f)$ computed in $\mathcal{C}^{\mathrm{op}}$ (2.6.154), so applying the above to $f$ regarded as a morphism of $\mathcal{C}^{\mathrm{op}}$ gives the dual equivalence. ■

## 2.6.21 Images and Coimages

In order to define exact sequence we need a categorical notion of image. We also define its dual.

**Definition 2.6.167 (Image)**
*Let $f : X \to Y$ be a morphism in $\mathcal{C}$. An **image** for $f$ is a triple $(\mathrm{Im}(f), e, m)$ such that $m$ is monic, we have a commutative triangle*




*and for any other factorisation $f = m' \circ e'$ with $m'$ monic there exists a unique morphism $g : \mathrm{Im}(f) \to T$ such that $m' \circ g = m$.*

*Note that any such $g$ satisfies $g \circ e = e'$, since $m' \circ (g \circ e) = m \circ e = f = m' \circ e'$ and $m'$ is monic.*

*The factorisations of $f$ through a monomorphism constitute a category $\mathrm{SubObjFactors}(f)$, with objects the triples $(T, e', m')$ as above and a morphism $(T, e', m') \to (T'', e'', m'')$ given by a morphism $g : T \to T''$ such that $m'' \circ g = m'$. So an image for $f$ is precisely an initial object (2.6.48) of $\mathrm{SubObjFactors}(f)$, and is therefore unique up to unique isomorphism.*

There is a dual notion

**Definition 2.6.168 (Coimage)**
*Let $f : X \to Y$ be a morphism in $\mathcal{C}$. A **coimage** for $f$ is a triple $(\mathrm{Coim}(f), e, m)$ such that $e$ is epic and we have a commutative triangle*




*and for any other factorisation $f = m' \circ e'$ with $e'$ epic there exists a unique morphism $g : T \to \mathrm{Coim}(f)$ such that $g \circ e' = e$.*

*Note that any such $g$ satisfies $m \circ g = m'$, since $(m \circ g) \circ e' = m \circ e = f = m' \circ e'$ and $e'$ is epic.*

*Dually, the factorisations of $f$ through an epimorphism constitute a category $\mathrm{QuotObjFactors}(f)$, with objects the triples $(T, e', m')$ as above and a morphism $(T, e', m') \to (T'', e'', m'')$ given by a morphism $g : T \to T''$ such that $g \circ e' = e''$. So a coimage for $f$ is precisely a terminal object (2.6.50) of $\mathrm{QuotObjFactors}(f)$, and is therefore unique up to unique isomorphism.*

© David Rufino CC BY-NC-ND 4.0

**Remark 2.6.169**
*We note that $(I, e, m)$ is an image in $\mathcal{C}$ if and only if $(I, m^{op}, e^{op})$ is a coimage in $\mathcal{C}^{op}$.*

**Lemma 2.6.170**
*Let $f : X \to Y$ be a morphism and $(I, e, m)$ an image. Suppose $\mathcal{C}$ has equalisers, then $e$ is epic.*

*Dually if $(I, e, m)$ is a coimage for $f : X \to Y$, and suppose $\mathcal{C}$ has coequalisers. Then $m$ is monic.*

*Proof.* Consider the diagram




and suppose $\alpha \circ e = \beta \circ e$. Let $K = \text{coeq}(\alpha, \beta)$, then by definition there is a morphism $g : X \to K$ such that $i \circ g = e$. Then $(m \circ i) \circ g = f$ is a factorisation with $m \circ i$ a monomorphism (2.6.135). Then by the universal property of images there is a morphism $j : \text{Im}(f) \to K$ such that $\left(m_f \circ i\right) \circ j = m$. As $m$ is monic $i \circ j = 1_{\text{Im}(f)}$ and so $i$ is split-epic and therefore an isomorphism (2.6.36). ■

**Lemma 2.6.171**
*Let $\mathcal{C}$ be a pointed category and $f : X \to Y$ a monomorphism. Then the triple $(X, 1_X, f)$ is an image for $f$.*

*Proof.* Denote the morphism by $f$ and suppose we have another factorisation $X \xrightarrow{\bar{f}} W \xrightarrow{i} Y$ with $i$ monic, then we have a commutative diagram




and we may show by the argument in (2.6.167) that the factorisation is unique. ■

**Lemma 2.6.172 (Images and Coimages are Preserved under Isomorphism)**
*Let $(T, e, m)$ be an image of $f : X \to Y$ and $\theta : T \xrightarrow{\sim} T'$ an isomorphism. Then $(T', \theta \circ e, m \circ \theta^{-1})$ is also an image of $f$.*

*Dually, if $(T, e, m)$ is a coimage of $f$ and $\theta : T \xrightarrow{\sim} T'$ an isomorphism, then $(T', \theta \circ e, m \circ \theta^{-1})$ is also a coimage of $f$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

For the image case $\theta$ determines a well-defined isomorphism in $\text{SubObjFactors}(f)$

$$(T, e, m) \xrightarrow{\sim} (T', \theta \circ e, m \circ \theta^{-1})$$

As isomorphisms preserve initial objects (2.6.49) we see that the right hand side is an image.

Dually, the coimage case follows by applying the image case to $f^{\text{op}}$ in $\mathcal{C}^{\text{op}}$ (2.6.169). ■

## 2.6.22 Epi-Mono Factorisations

We saw that under mild conditions image and coimage yield an epi-mono factorisation. Strengthening this factorisation can ensure functorial properties; concretely, in an algebraic category, upgrading an epimorphism to a regular epimorphism is exactly upgrading from the (possibly too weak) categorical notion of epimorphism to the familiar notion of a surjective homomorphism (2.6.195).

© David Rufino CC BY-NC-ND 4.0

**Definition 2.6.173**
*Let $\mathcal{C}$ be a category and $f : X \to Y$ a morphism. We say that $f = m \circ e$ is*

- *a **regular epi-mono factorisation** if $m$ is a monomorphism and $e$ is a regular epimorphism.*
- *an **epi-regular mono factorisation** if $m$ is a regular monomorphism and $e$ is an epimorphism*

*We note that $f = m \circ e$ is a regular epi-mono (resp. epi-regular mono) factorisation if and only if $f^{op} = e^{op} \circ m^{op}$ is an epi-regular mono (resp. regular epi-mono) factorisation.*

**Proposition 2.6.174 (Regular Epi-Mono Factorisation)**
*Let $f : X \to Y$ be a morphism with regular epi-mono factorisation $f = m \circ e$. Then $(I, e, m)$ is an image and every image is a regular epi-mono factorisation.*

*Dually, let $f : X \to Y$ be a morphism with epi-regular mono factorisation $f = m \circ e$. Then $(I, e, m)$ is a coimage and every coimage is an epi-regular mono factorisation.*

*Proof.* Consider the diagram




where $e$ is a coequaliser for $(\alpha, \beta)$, $m'$ is a monomorphism and $e'$ is a general morphism. Then we need to prove the existence of a unique morphism $(I, e, m) \to (I', e', m')$. Since $e$ coequalises $(\alpha, \beta)$, $e \circ \alpha = e \circ \beta$, so

$$f \circ \alpha = m \circ e \circ \alpha = m \circ e \circ \beta = f \circ \beta$$

and as $f = m' \circ e'$ with $m'$ monic, $m' \circ e' \circ \alpha = f \circ \alpha = f \circ \beta = m' \circ e' \circ \beta$ gives $e' \circ \alpha = e' \circ \beta$. Therefore by definition of the coequaliser there is a unique $\psi$ such that $\psi \circ e = e'$. Further

$$m' \circ \psi \circ e = m' \circ e' = f = m \circ e$$

and as $e$ is an epimorphism (2.6.135) we conclude that $m' \circ \psi = m$, as required for a morphism $(I, e, m) \to (I', e', m')$ in $\mathrm{SubObjFactors}(f)$ (2.6.167). If $\psi'$ also satisfies $m' \circ \psi' = m$, then $m' \circ \psi = m = m' \circ \psi'$, and since $m'$ is monic $\psi = \psi'$; so $\psi$ is unique.

Let $(I', e', m')$ be an image, so $m'$ is in particular monic and the above applies. As images are unique up to isomorphism (2.6.167) there is an isomorphism $\theta : I \xrightarrow{\sim} I'$ such that $m' \circ \theta = m$; then

$$m' \circ (\theta \circ e) = (m' \circ \theta) \circ e = m \circ e = f = m' \circ e'$$

and as $m'$ is monic, $\theta \circ e = e'$. Since $e$ is a coequaliser of $(\alpha, \beta)$ and $\theta$ is an isomorphism with $\theta \circ e = e'$, post-composing the coequalising cocone for $e$ with $\theta$ shows $e'$ is again a coequaliser of $(\alpha, \beta)$, as colimits are preserved under isomorphisms (2.6.90); hence $e'$ is regular (2.6.158), as required. ∎

**Proposition 2.6.175 (Regular Epi-Mono)**
**Set** *has regular epi-mono factorisations.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Let $f : X \to Y$ be a morphism in **Set**. Set $\mathrm{Im}(f) \coloneqq \{f(x) \mid x \in X\}$ and let $m : \mathrm{Im}(f) \hookrightarrow Y$ denote the inclusion, which is injective and therefore monic by (2.6.38). Let $e : X \to \mathrm{Im}(f)$ be the corestriction of $f$, so that $f = m \circ e$.

Define $R = \{(x, x') \in X \times X \mid f(x) = f(x')\}$ with projections $\alpha, \beta : R \rightrightarrows X$, $\alpha(x, x') = x$, $\beta(x, x') = x'$. We claim $e$ is the coequaliser of $(\alpha, \beta)$.

First $e \circ \alpha = e \circ \beta$: for $(x, x') \in R$ we have $f(x) = f(x')$ by definition of $R$, that is $e(x) = e(x')$.

Now let $\pi' : X \to Z$ satisfy $\pi' \circ \alpha = \pi' \circ \beta$, that is $f(x) = f(x') \implies \pi'(x) = \pi'(x')$ for all $x, x' \in X$. Since every $y \in \mathrm{Im}(f)$ is of the form $f(x)$ for some $x \in X$, we may define $\psi : \mathrm{Im}(f) \to Z$ by $\psi(y) = \pi'(x)$ for any such $x$; this is well-

© David Rufino CC BY-NC-ND 4.0

defined precisely because $f(x) = f(x') = y$ implies $\pi'(x) = \pi'(x')$. By construction $\psi \circ e = \pi'$, and since $e$ is surjective onto $\mathrm{Im}(f)$ by definition, any $\psi'$ with $\psi' \circ e = \pi'$ agrees with $\psi$ at every point of $\mathrm{Im}(f)$, so $\psi$ is the unique such factorisation.

Hence $e$ is a coequaliser of $(\alpha, \beta)$, so is a regular epimorphism (2.6.158). ∎

**Lemma 2.6.176 (Diagonal Fill-In Lemma)**
*Let $e : X \to I$ be a regular epimorphism and $m' : I' \to Y$ a monomorphism. Then for every commutative square*




*there is a unique $\theta$ such that $\theta \circ e = a$. Furthermore it satisfies $m' \circ \theta = m$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Since $e$ is a regular epimorphism, by (2.6.158) there are morphisms $\alpha, \beta : W \rightrightarrows X$ such that $e$ is the coequaliser of $(\alpha, \beta)$.

We have $e \circ \alpha = e \circ \beta$, so $m \circ e \circ \alpha = m \circ e \circ \beta$, whence by commutativity of the square $m' \circ a \circ \alpha = m' \circ a \circ \beta$. As $m'$ is monic we may cancel it to get $a \circ \alpha = a \circ \beta$.

By the universal property of the coequaliser $e$ there is therefore a unique morphism $\theta : I \to I'$ such that

$$\theta \circ e = a$$

We check that $\theta$ also satisfies the remaining identity: $m' \circ \theta \circ e = m' \circ a = m \circ e$, and since $e$ is an epimorphism (2.6.135) we may cancel it to get

$$m' \circ \theta = m$$

∎

**Proposition 2.6.177 (Regular Epi-Mono Factorisation is Functorial)**
*Suppose $\mathcal{C}$ has regular epi-mono factorisations $f = m \circ e : X \to Y$, $f' = m' \circ e' : X' \to Y'$ and morphisms $a : X \to X'$, $b : Y \to Y'$ such that $b \circ f = f' \circ a$. Then there is a unique morphism $\psi_{a,b} : I \to I'$ which makes the top square commute*




*Furthermore the whole diagram commutes and the construction is functorial: $\psi_{1_X, 1_Y} = 1_I$ (taking $f' = f$, $a = 1_X$, $b = 1_Y$), and given a further regular epi-mono factorisation $f'' = m'' \circ e'' : X'' \to Y''$ and morphisms $c : X' \to X''$, $d : Y' \to Y''$ such that $d \circ f' = f'' \circ c$ (with induced comparison morphism $\psi_{c,d} : I' \to I''$), we have $\psi_{c,d} \circ \psi_{a,b} = \psi_{c \circ a, d \circ b}$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Since $e$ is a regular epimorphism, by (2.6.158) there are morphisms $\alpha, \beta : W \rightrightarrows X$ such that $e$ is the coequaliser of $(\alpha, \beta)$.

We first show $e' \circ a$ coequalises $(\alpha, \beta)$. Since $e \circ \alpha = e \circ \beta$ we have $f \circ \alpha = m \circ e \circ \alpha = m \circ e \circ \beta = f \circ \beta$, and so by hypothesis $f' \circ a \circ \alpha = b \circ f \circ \alpha = b \circ f \circ \beta = f' \circ a \circ \beta$, that is $m' \circ e' \circ a \circ \alpha = m' \circ e' \circ a \circ \beta$. As $m'$ is monic we may cancel it to get $(e' \circ a) \circ \alpha = (e' \circ a) \circ \beta$.

© David Rufino CC BY-NC-ND 4.0

By the universal property of the coequaliser $e$ there is therefore a unique morphism $\psi_{a,b} : I \to I'$ such that

$$\psi_{a,b} \circ e = e' \circ a$$

We check that $\psi_{a,b}$ also satisfies the second required identity: $m' \circ \psi_{a,b} \circ e = m' \circ e' \circ a = f' \circ a = b \circ f = b \circ m \circ e$, and since $e$ is an epimorphism (2.6.135) we may cancel it to get

$$m' \circ \psi_{a,b} = b \circ m$$

so the diagram commutes.

Taking $f' = f$, $a = 1_X$, $b = 1_Y$ (a valid compatible pair since $1_Y \circ f = f = f \circ 1_X$), $1_I$ satisfies $1_I \circ e = e = e \circ 1_X$ and $m \circ 1_I = m = 1_Y \circ m$, so by the uniqueness established above $\psi_{1_X,1_Y} = 1_I$.

For the composition law, let $c : X' \to X''$, $d : Y' \to Y''$ satisfy $d \circ f' = f'' \circ c$, with induced comparison morphism $\psi_{c,d} : I' \to I''$. Then

$$\left(\psi_{c,d} \circ \psi_{a,b}\right) \circ e = \psi_{c,d} \circ (e' \circ a) = \left(\psi_{c,d} \circ e'\right) \circ a = e'' \circ c \circ a = e'' \circ (c \circ a)$$

Note $d \circ b \circ f = d \circ f' \circ a = f'' \circ c \circ a = f'' \circ (c \circ a)$, so $(c \circ a, d \circ b)$ is again a compatible pair, with induced comparison morphism $\psi_{c\circ a, d\circ b} : I \to I''$ satisfying $\psi_{c\circ a, d\circ b} \circ e = e'' \circ (c \circ a)$ by definition. Since both $\psi_{c,d} \circ \psi_{a,b}$ and $\psi_{c\circ a, d\circ b}$ satisfy this same equation, and $\psi_{c\circ a, d\circ b}$ is, by the uniqueness established above, the unique morphism $I \to I''$ doing so, we conclude

$$\psi_{c,d} \circ \psi_{a,b} = \psi_{c\circ a, d\circ b}$$

■

**Proposition 2.6.178 (Regular Epi-Mono Factorisation is Unique)**
*Let $f = m \circ e = m' \circ e'$ be two regular epi-mono factorisations, with $e : X \to I$, $e' : X \to I'$. Then there is a unique morphism $\theta : I \to I'$ such that $e' = \theta \circ e$. Furthermore it is an isomorphism and satisfies $m' \circ \theta = m$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Applying (2.6.177) to the factorisations $(I, e, m)$ and $(I', e', m')$ of $f$, with $f' = f$, $a = 1_X$, $b = 1_Y$, gives a unique morphism $\theta : I \to I'$ such that $\theta \circ e = e'$ and $m' \circ \theta = m$.

Symmetrically, applying (2.6.177) to the factorisations $(I', e', m')$ and $(I, e, m)$ of $f$ (i.e. with the roles of the two factorisations swapped) gives a unique morphism $\theta' : I' \to I$ such that $\theta' \circ e' = e$ and $m \circ \theta' = m'$.

By the composition law of (2.6.177), applied to the chain of factorisations $(I, e, m) \to (I', e', m') \to (I, e, m)$ (i.e. $f' = f$, $f'' = f$, with $c = 1_X$, $d = 1_Y$, and $\theta$, $\theta'$ as the two successive comparison morphisms), $\theta' \circ \theta$ is the comparison morphism from $(I, e, m)$ to itself, and so equals $1_I$ by the identity law of (2.6.177).

Symmetrically $\theta \circ \theta' = 1_{I'}$, so $\theta$ is an isomorphism with inverse $\theta'$. ■

## 2.6.23 Algebraic Categories

*Drafted by Claude Sonnet, reviewed by David Rufino.*

The categories of groups, rings and modules all share a common shape: their objects are sets equipped with some finitary operations, subject to some equational axioms. The following makes this precise, following the classical treatment of universal algebra [Bir35] [BS81] and its later categorical formulation [ARV10].

**Definition 2.6.179 (Signature)**
*A **signature** $\Omega$ is a set of **operation symbols**, each $\omega \in \Omega$ assigned a natural number $n_\omega$, its **arity**.*

**Definition 2.6.180 ($\Omega$-algebra)**
*Let $\Omega$ be a signature. An **$\Omega$-algebra** is a set $S$ together with, for each $\omega \in \Omega$, a function $\omega_S : S^{n_\omega} \to S$ (when $n_\omega = 0$ this is simply a distinguished element of $S$, since $S^0$ is a singleton).*

*A **homomorphism** of $\Omega$-algebras $f : S \to T$ is a function such that*

$$f\left(\omega_S\left(s_1, ..., s_{n_\omega}\right)\right) = \omega_T\left(f(s_1), ..., f\left(s_{n_\omega}\right)\right) \quad \forall \omega \in \Omega, s_1, ..., s_{n_\omega} \in S$$

*This determines a concrete category $\Omega$-**Alg** (2.6.9).*

**Example 2.6.181**
*Taking $\Omega = \{\cdot\,(2), {}^{-1}(1), e(0)\}$ (writing the arity of each symbol in brackets), $\Omega$-algebras are exactly sets equipped*

© David Rufino CC BY-NC-ND 4.0

*with a binary, a unary and a nullary operation; the subcategory of those satisfying the group axioms, (2.6.184) below, is the category of groups.*

**Definition 2.6.182 (Terms)**
*Let $\Omega$ be a signature and $X$ a set, whose elements we call **variables**. The set $T_\Omega(X)$ of **$\Omega$-terms** over $X$ is defined recursively by*

- *every $x \in X$ is a term;*
- *if $\omega \in \Omega$ and $t_1, ..., t_{n_\omega} \in T_\Omega(X)$ then $\omega\big(t_1, ..., t_{n_\omega}\big) \in T_\Omega(X)$.*

*Given an $\Omega$-algebra $S$ and a function $\theta : X \to S$, there is a unique **evaluation** $\overline{\theta} : T_\Omega(X) \to S$ such that $\overline{\theta}(x) = \theta(x)$ for $x \in X$ and*

$$\overline{\theta}\big(\omega\big(t_1, ..., t_{n_\omega}\big)\big) = \omega_S\big(\overline{\theta}(t_1), ..., \overline{\theta}\big(t_{n_\omega}\big)\big)$$

**Remark 2.6.183 (The term algebra)**
*$T_\Omega(X)$ is itself an $\Omega$-algebra, via $\omega_{T_\Omega(X)}\big(t_1, ..., t_{n_\omega}\big) := \omega\big(t_1, ..., t_{n_\omega}\big)$; with this structure, the defining recursion for $\overline{\theta}$ is exactly the condition that $\overline{\theta} : T_\Omega(X) \to S$ is a homomorphism, and it is the unique homomorphism $T_\Omega(X) \to S$ restricting to $\theta$ on $X$.*

*Consequently, for any homomorphism $h : S \to S'$ and $\theta : X \to S$, both $h \circ \overline{\theta}$ and $\overline{h \circ \theta}$ are homomorphisms $T_\Omega(X) \to S'$ restricting to $h \circ \theta$ on $X$, so by uniqueness*

$$h \circ \overline{\theta} = \overline{h \circ \theta}$$

**Definition 2.6.184 (Identity, Algebraic category)**
*Let $\Omega$ be a signature. An **$\Omega$-identity** is a pair $p, q \in T_\Omega(X)$ for some set of variables $X$, written $p = q$. An $\Omega$-algebra $S$ **satisfies** $p = q$ if $\overline{\theta}(p) = \overline{\theta}(q)$ for every $\theta : X \to S$ (2.6.182).*

*Given a set $E$ of $\Omega$-identities, the **algebraic category** determined by $(\Omega, E)$ – also called an **equational category** or **variety** in the literature – is the full subcategory of $\Omega$-**Alg** of those $\Omega$-algebras satisfying every identity in $E$, again concrete over* **Set**.

**Example 2.6.185**
**Grp**, **AbGrp**, **Rng** *are algebraic categories, taking $E$ to consist of the usual group, abelian group and ring axioms respectively, each expressed as an identity. By contrast, the category of fields is* not *an algebraic category: the axiom that every nonzero element has an inverse cannot be expressed as an identity holding at every element, since it fails at 0.*

**Example 2.6.186 ($A$-Mod as an algebraic category)**
*Fix a ring $A$. Take $\Omega = \{+, -, 0\} \cup \{\lambda_a\}_{a \in A}$, with $+$ binary, $-$ unary, $0$ nullary and each $\lambda_a$ unary; note $\Omega$ is infinite (one symbol per element of $A$) but every operation remains finitary. Take $E$ to consist of the abelian group axioms for $(+, -, 0)$ together with, for all $a, a' \in A$,*

$$\lambda_a(x + y) = \lambda_a(x) + \lambda_a(y), \quad \lambda_1(x) = x, \quad \lambda_{a+a'}(x) = \lambda_a(x) + \lambda_{a'}(x), \quad \lambda_{aa'}(x) = \lambda_a(\lambda_{a'}(x))$$

*Setting $\lambda_a(m) := a \cdot m$, these are exactly the $A$-module axioms, so $A$-**Mod** is the algebraic category determined by $(\Omega, E)$.*

**Example 2.6.187 ($\mathbf{Alg}_k$ as an algebraic category)**
*Fix a commutative ring $A$ (so $\mathbf{Alg}_k$ is the case $A = k$ a field). Take $\Omega = \{+, \times, -, 0, 1\} \cup \{\lambda_a\}_{a \in A}$, with $+, \times$ binary, $-$ unary, $0, 1$ nullary and each $\lambda_a$ unary, and take $E$ to consist of the ring axioms for $(+, \times, -, 0, 1)$ (*not *imposing commutativity of $\times$) together with the four identities above and, in addition,*

$$\lambda_a(x) = \lambda_a(1) \times x \quad \forall a \in A$$

*Given such an $\Omega$-algebra $B$, set $i_B(a) := \lambda_a(1)$. Additivity and unitality of $i_B$ are immediate from the identities above at $x = 1$, and multiplicativity follows from the last two identities,*

$$i_B(aa') = \lambda_{aa'}(1) = \lambda_a(\lambda_{a'}(1)) = \lambda_a(1) \times \lambda_{a'}(1) = i_B(a) i_B(a')$$

*so $i_B$ is a ring homomorphism, and the final identity reads $\lambda_a(b) = i_B(a) \cdot b$ for all $b \in B$: this is exactly an $A$-algebra structure on $B$. Conversely every $A$-algebra satisfies these identities, and a ring homomorphism commutes with every $\lambda_a$ if and only if it respects the structural morphisms (apply the final identity at $b = 1$), so $\mathbf{Alg}_A$, and in particular $\mathbf{Alg}_k$, is the algebraic category determined by $(\Omega, E)$.*

© David Rufino CC BY-NC-ND 4.0

**Remark 2.6.188**
*Birkhoff's theorem characterises the algebraic (equational) subcategories of* $\Omega$-**Alg** *as exactly those closed under homomorphic images, subalgebras and products [Bir35]; see [BS81] for a modern account, and [ARV10] for the categorical perspective via algebraic theories.*

**Proposition 2.6.189 (Properties of Algebraic Categories)**
*Let* $\mathcal{C}$ *be an algebraic category determined by* $(\Omega, E)$ *(2.6.184), with forgetful functor* $U : \mathcal{C} \to \mathbf{Set}$*. Then*

*a)* $U$ *reflects isomorphisms*
*b)* $U$ *creates limits and directed colimits*
*c)* $U$ *has a left adjoint (i.e. free algebras exist)*

*In particular* $U$ *also preserves limits and directed colimits by (2.6.87).*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Throughout, for $\omega \in \Omega$ we write $r = 1, ..., n_\omega$ for the arguments of $\omega$, reserving $i, j, k$ for diagram indices.

***U* creates limits.** Let $F : I \to \mathcal{C}$ be a diagram and suppose $L := \lim(U \circ F)$ exists in **Set**, with structure maps $\pi_i : L \to U(F(i))$; explicitly, by (2.6.85), $L = \left\{(x_i)_{i\in I} \in \prod_i U(F(i)) \;\middle|\; U(F(\alpha))(x_i) = x_j \quad \forall \alpha : i \to j\right\}$.

For $\omega \in \Omega$ and $x^{(1)}, ..., x^{(n_\omega)} \in L$, define $\omega_L\big(x^{(1)}, ..., x^{(n_\omega)}\big)_i := \omega_{F(i)}\Big(x_i^{(1)}, ..., x_i^{(n_\omega)}\Big)$ for each $i \in I$. This again lies in $L$: for $\alpha : i \to j$, since $F(\alpha)$ is a homomorphism, $F(\alpha)\Big(\omega_{F(i)}\Big(x_i^{(1)}, ...\Big)\Big) = \omega_{F(j)}\Big(F(\alpha)\Big(x_i^{(1)}\Big), ...\Big) = \omega_{F(j)}\Big(x_j^{(1)}, ...\Big)$, using that each $x^{(r)} \in L$. So $L$ is an $\Omega$-algebra and each $\pi_i$ is, by construction, a homomorphism; this is the unique such structure on $L$, since $\omega_L(...)$ is pinned down coordinatewise by the requirement that every $\pi_i$ be a homomorphism, and $L \subseteq \prod_i U(F(i))$.

Given $p = q \in E$ with variables in a set $X$, and $\theta : X \to L$: for each $i$, $\pi_i \circ \theta : X \to U(F(i))$, and by (2.6.183), $\pi_i\big(\overline{\theta}(t)\big) = \overline{\pi_i \circ \theta}(t)$ for every term $t$. As $F(i) \in \mathcal{C}$ satisfies $p = q$, $\overline{\pi_i\theta}(p) = \overline{\pi_i\theta}(q)$, so $\pi_i\big(\overline{\theta}(p)\big) = \pi_i\big(\overline{\theta}(q)\big)$ for every $i \in I$, hence $\overline{\theta}(p) = \overline{\theta}(q)$ as elements of $L \subseteq \prod_i U(F(i))$. So $L \in \mathcal{C}$.

Finally $(L, \{\pi_i\})$ is a limit for $F$ in $\mathcal{C}$: given a cone $(M, \{\mu_i\})$ in $\mathcal{C}$, the unique **Set**-map $\varphi : U(M) \to L$ with $\pi_i \circ \varphi = \mu_i$ (afforded by $L = \lim(U \circ F)$) is a homomorphism, since for $\omega \in \Omega$, $m^{(1)}, ..., m^{(n_\omega)} \in U(M)$ and every $i$,

$$\pi_i\big(\varphi\big(\omega_M\big(m^{(1)}, ...\big)\big)\big) = \mu_i\big(\omega_M\big(m^{(1)}, ...\big)\big) = \omega_{F(i)}\big(\mu_i\big(m^{(1)}\big), ...\big) = \omega_{F(i)}\big(\pi_i\varphi\big(m^{(1)}\big), ...\big) = \pi_i\big(\omega_L\big(\varphi\big(m^{(1)}\big), ...\big)\big)$$

using that $\mu_i$ is a homomorphism, so $\varphi\big(\omega_M\big(m^{(1)}, ...\big)\big) = \omega_L\big(\varphi\big(m^{(1)}\big), ...\big)$. As $\mathcal{C}$ is a full subcategory of $\Omega$-**Alg**, $\varphi$ is the unique morphism of $\mathcal{C}$ with $\pi_i \circ \varphi = \mu_i$, so $(L, \{\pi_i\})$ is a limit for $F$ in $\mathcal{C}$ with $U(L) = L$, as required.

***U* creates directed colimits.** Let $I$ be a directed preorder and $F : I \to \mathcal{C}$ a diagram, and suppose $L := \varinjlim(U \circ F)$ exists in **Set** (2.6.128), with structure maps $\mu_i : U(F(i)) \to L$.

For $\omega \in \Omega$ and $\ell^{(1)}, ..., \ell^{(n_\omega)} \in L$: as $I$ is directed and $n_\omega$ is finite, there is a common $i \in I$ and $y^{(1)}, ..., y^{(n_\omega)} \in U(F(i))$ with $\mu_i\big(y^{(r)}\big) = \ell^{(r)}$ for all $r$ (each $\ell^{(r)}$ is represented at some stage, and finitely many stages admit a common upper bound); define $\omega_L\big(\ell^{(1)}, ..., \ell^{(n_\omega)}\big) := \mu_i\big(\omega_{F(i)}\big(y^{(1)}, ..., y^{(n_\omega)}\big)\big)$. This is independent of the choice of $i$ and the $y^{(r)}$: given another such $i', \big(y'^{(r)}\big)$, for each $r$ there is, by (2.6.128), a stage $k_r \succeq i, i'$ with $F(i \to k_r)\big(y^{(r)}\big) = F(i' \to k_r)\big(y'^{(r)}\big)$; as $I$ is directed and there are finitely many $r$, there is a common $j \succeq k_1, ..., k_{n_\omega}$, and applying $F(k_r \to j)$ to each equality gives $F(i \to j)\big(y^{(r)}\big) = F(i' \to j)\big(y'^{(r)}\big)$ for every $r$ simultaneously. As $F(i \to j), F(i' \to j)$ are homomorphisms,

$$\mu_i\big(\omega_{F(i)}\big(y^{(1)}, ...\big)\big) = \mu_j\big(\omega_{F(j)}\big(F(i \to j)\big(y^{(1)}\big), ...\big)\big) = \mu_j\big(\omega_{F(j)}\big(F(i' \to j)\big(y'^{(1)}\big), ...\big)\big) = \mu_{i'}\big(\omega_{F(i')}\big(y'^{(1)}, ...\big)\big)$$

So $L$ is an $\Omega$-algebra and each $\mu_i$ is, by construction, a homomorphism; as the $\mu_i$ are jointly surjective (2.6.128), this is again the unique such structure.

Given $p = q \in E$ with (finitely many) variables $X$, and $\theta : X \to L$: as $X$ is finite and $I$ directed, there is $i \in I$ and $\theta_i : X \to U(F(i))$ with $\theta = \mu_i \circ \theta_i$ (as above); by (2.6.183), $\overline{\theta} = \mu_i \circ \overline{\theta_i}$, so $\overline{\theta}(p) = \mu_i\big(\overline{\theta_i}(p)\big) = \mu_i\big(\overline{\theta_i}(q)\big) = \overline{\theta}(q)$, using $F(i) \in \mathcal{C}$. So $L \in \mathcal{C}$.

Finally $(L, \{\mu_i\})$ is a colimit for $F$ in $\mathcal{C}$: given a cocone $(M, \{\nu_i\})$ in $\mathcal{C}$, the unique **Set**-map $\psi : L \to U(M)$ with $\psi \circ \mu_i = \nu_i$ is a homomorphism (for $\ell^{(r)} = \mu_i\big(y^{(r)}\big)$ at a common $i$, $\psi\big(\omega_L\big(\ell^{(1)}, ...\big)\big) = \psi\big(\mu_i\big(\omega_{F(i)}\big(y^{(1)}, ...\big)\big)\big) = \nu_i\big(\omega_{F(i)}\big(y^{(1)}, ...\big)\big) = \omega_M\big(\nu_i\big(y^{(1)}\big), ...\big) = \omega_M\big(\psi\big(\ell^{(1)}\big), ...\big)$, using that $\nu_i$ is a homomorphism), and is the unique morphism of $\mathcal{C}$ with this property as $\mathcal{C}$ is full in $\Omega$-**Alg**.

***U* reflects isomorphisms.** Let $f : X \to Y$ in $\mathcal{C}$ with $U(f)$ bijective. For $\omega \in \Omega$ and $y^{(1)}, ..., y^{(n_\omega)} \in U(Y)$, writing $x^{(r)} := U(f)^{-1}\big(y^{(r)}\big)$, since $f$ is a homomorphism $f\big(\omega_X\big(x^{(1)}, ...\big)\big) = \omega_Y\big(f\big(x^{(1)}\big), ...\big) = \omega_Y\big(y^{(1)}, ...\big)$, so applying $U(f)^{-1}$,

© David Rufino CC BY-NC-ND 4.0

$$U(f)^{-1}\big(\omega_Y(y^{(1)},\dots,y^{(n_\omega)})\big)=\omega_X\big(U(f)^{-1}(y^{(1)}),\dots,U(f)^{-1}(y^{(n_\omega)})\big)$$

so $U(f)^{-1}$ is a homomorphism $Y\to X$, hence a morphism of $\mathcal{C}$ (full subcategory), inverse to $f$ on underlying sets; as $U$ is faithful, it is a two-sided inverse to $f$ in $\mathcal{C}$, so $f$ is an isomorphism.

***U* has a left adjoint.** Fix a set $X$. Define a relation $\sim$ on $T_\Omega(X)$ by $t\sim t'$ if $\overline{\theta}(t)=\overline{\theta}(t')$ for every $A\in\mathcal{C}$ and $\theta: X\to U(A)$; this is manifestly an equivalence relation, and a congruence: if $t^{(r)}\sim t'^{(r)}$ for $r=1,\dots,n_\omega$, then for every $A,\theta$, $\overline{\theta}\big(\omega(t^{(1)},\dots)\big)=\omega_A\big(\overline{\theta}(t^{(1)}),\dots\big)=\omega_A\big(\overline{\theta}(t'^{(1)}),\dots\big)=\overline{\theta}\big(\omega(t'^{(1)},\dots)\big)$, so $\omega\big(t^{(1)},\dots,t^{(n_\omega)}\big)\sim\omega\big(t'^{(1)},\dots,t'^{(n_\omega)}\big)$.

Let $\mathrm{Free}(X):=T_\Omega(X)/\sim$, with $\omega_{\mathrm{Free}(X)}\big([t^{(1)}],\dots,[t^{(n_\omega)}]\big):=\big[\omega(t^{(1)},\dots,t^{(n_\omega)})\big]$ (well-defined as $\sim$ is a congruence), and let $\pi: T_\Omega(X)\to\mathrm{Free}(X)$, $t\mapsto[t]$, an (evidently surjective) homomorphism. Write $\iota: X\to U(\mathrm{Free}(X))$, $\iota(x):=[x]$.

**Free($X$) $\in\mathcal{C}$:** let $p=q\in E$ with variable set $Y$, and $\psi: Y\to U(\mathrm{Free}(X))$; choose $\tilde{\psi}: Y\to T_\Omega(X)$ with $\pi\circ\tilde{\psi}=\psi$ (possible as $\pi$ is surjective). Viewing $T_\Omega(X)$ as an $\Omega$-algebra (2.6.183), $\overline{\tilde{\psi}}: T_\Omega(Y)\to T_\Omega(X)$ is a homomorphism, and for every $A\in\mathcal{C}$, $\theta: X\to U(A)$, (2.6.183) gives $\overline{\theta}\circ\overline{\tilde{\psi}}=\overline{\overline{\theta}\circ\tilde{\psi}}$; as $\overline{\theta}\circ\tilde{\psi}: Y\to U(A)$ and $A\vDash p=q$, $\overline{\overline{\theta}\tilde{\psi}}(p)=\overline{\overline{\theta}\tilde{\psi}}(q)$, so $\overline{\theta}\Big(\overline{\tilde{\psi}}(p)\Big)=\overline{\theta}\Big(\overline{\tilde{\psi}}(q)\Big)$. As $A,\theta$ were arbitrary, $\overline{\tilde{\psi}}(p)\sim\overline{\tilde{\psi}}(q)$, that is $\pi\Big(\overline{\tilde{\psi}}(p)\Big)=\pi\Big(\overline{\tilde{\psi}}(q)\Big)$. Again by (2.6.183), $\pi\circ\overline{\tilde{\psi}}=\overline{\psi}$ (both extend $\pi\circ\tilde{\psi}=\psi$ on $Y$), so $\overline{\psi}(p)=\overline{\psi}(q)$. As $\psi$ was arbitrary, $\mathrm{Free}(X)\vDash p=q$; as $p=q\in E$ was arbitrary, $\mathrm{Free}(X)\in\mathcal{C}$.

**Universal property:** given $A\in\mathcal{C}$ and $\theta: X\to U(A)$, define $\hat{\theta}([t]):=\overline{\theta}(t)$; this is well-defined ($t\sim t'\implies\overline{\theta}(t)=\overline{\theta}(t')$, by definition of $\sim$) and a homomorphism, since $\hat{\theta}\big(\omega_{\mathrm{Free}(X)}\big([t^{(1)}],\dots\big)\big)=\hat{\theta}\big([\omega(t^{(1)},\dots)]\big)=\overline{\theta}\big(\omega(t^{(1)},\dots)\big)=\omega_A\big(\overline{\theta}(t^{(1)}),\dots\big)=\omega_A\big(\hat{\theta}([t^{(1)}]),\dots\big)$, and satisfies $U\big(\hat{\theta}\big)\circ\iota=\theta$. If $\varphi:\mathrm{Free}(X)\to A$ is any morphism of $\mathcal{C}$ with $U(\varphi)\circ\iota=\theta$, then $\varphi\circ\pi$ is a homomorphism $T_\Omega(X)\to A$ restricting to $\theta$ on $X$, so $\varphi\circ\pi=\overline{\theta}$ by (2.6.183); as $\pi$ is surjective, $\varphi=\hat{\theta}$. So $\mathrm{Free}(X)$ is free on $X$, and $X\mapsto\mathrm{Free}(X)$ is left adjoint to $U$ by (2.6.74). ∎

**Remark 2.6.190 (Limits are computed pointwise)**
*Unwinding the construction in the proof of (2.6.189): for a diagram $F: I\to\mathcal{C}$ admitting a limit, $U(\lim F)=\lim(U\circ F)$ (2.6.85), and each $\omega\in\Omega$ acts on $\lim F$ coordinatewise, $\omega_{\lim F}\big(x^{(1)},\dots\big)_i=\omega_{F(i)}\Big(x_i^{(1)},\dots\Big)$ for $i\in I$. In particular this applies to products.*

**Corollary 2.6.191 (Inverse Limits in Algebraic Categories)**
*Let $\mathcal{C}$ be an algebraic category determined by $(\Omega,E)$ (2.6.184), with forgetful functor $U:\mathcal{C}\to\mathbf{Set}$, $I$ a directed preorder, and $F: I^{op}\to\mathcal{C}$ an inverse system. Then $\varprojlim_i F(i)$ exists in $\mathcal{C}$, with*

$$U\Big(\varprojlim_i F(i)\Big)=\varprojlim_i U(F(i))=\left\{(x_i)_{i\in I}\in\prod_i U(F(i))\;\middle|\;U\big(\rho_{ij}\big)(x_j)=x_i\quad\forall i\prec j\right\}$$

*(2.6.130), and each $\omega\in\Omega$ acts coordinatewise: $\omega_{\varprojlim_i F(i)}\big(x^{(1)},\dots\big)_i=\omega_{F(i)}\Big(x_i^{(1)},\dots\Big)$ for every $i\in I$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

By (2.6.189), $U$ creates limits, and $\lim_{I^{op}}(U\circ F)=\varprojlim_i U(F(i))$ exists in **Set** for every diagram, being computed as in (2.6.130) (a special case of (2.6.85)); so $\varprojlim_i F(i):=\lim_{I^{op}}F$ exists in $\mathcal{C}$ too, with $U$ carrying it to this set-level description. The coordinatewise formula for $\omega\in\Omega$ is then the case $\lim F:=\varprojlim_i F(i)$ of (2.6.190). ∎

**Definition 2.6.192 (Subalgebra)**
*Let $A$ be an $\Omega$-algebra. A subset $S\subseteq U(A)$ is a **subalgebra** if $\omega_A\Big(s_1,\dots,s_{n_\omega}\Big)\in S$ for all $\omega\in\Omega$ and $s_1,\dots,s_{n_\omega}\in S$.*

**Proposition 2.6.193 (Subalgebra criterion)**
*Let $A$ be an $\Omega$-algebra and $S\subseteq U(A)$ a subalgebra (2.6.192). Then $S$ carries a unique $\Omega$-algebra structure making the inclusion $\iota: S\hookrightarrow A$ a homomorphism, namely $\omega_S:=\omega_A\,|_{S^{n_\omega}}$.*

*If moreover $A$ satisfies an identity $p=q$, then so does $S$; in particular, if $A$ lies in the algebraic category $\mathcal{C}$ determined by $(\Omega,E)$, then so does $S$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Existence and uniqueness of $\omega_S$ are immediate: $S$ being closed under $\omega_A$ means $\omega_A\,|_{S^{n_\omega}}$ takes values in $S$, and this is forced by the requirement that $\iota$ be a homomorphism.

Let $p=q$ be an identity with variables $Y$, satisfied by $A$, and $\theta: Y\to S$. Writing $\iota\circ\theta: Y\to U(A)$, (2.6.183) gives $\iota\Big(\overline{\theta}(p)\Big)=\overline{\iota\theta}(p)=\overline{\iota\theta}(q)=\iota\Big(\overline{\theta}(q)\Big)$, using that $A$ satisfies $p=q$; as $\iota$ is injective, $\overline{\theta}(p)=\overline{\theta}(q)$, so $S$ satisfies $p=q$. ∎

© David Rufino CC BY-NC-ND 4.0

**Proposition 2.6.194 (Injective $\iff$ Monic)**
*Let $\mathcal{C}$ be an algebraic category. Then a morphism $f : X \to Y$ is a monomorphism if and only if $U(f)$ is injective.*

*Proof.* If $U(f)$ is injective and $f \circ g = f \circ h$ then $U(f) \circ U(g) = U(f) \circ U(h)$ whence $U(g) = U(h)$ and by faithful-ness $g = h$ as required.

Conversely if $f : X \to Y$ is monic, consider the left adjoint $F$ to $U$ (2.6.189), then by definition we have a natural bijection

$$\Psi_X : \mathrm{Mor}(F(\{\star\}), X) \xrightarrow{\sim} U(X)$$

Suppose $U(f)(x) = U(f)(y)$ for $x, y \in U(X)$. Let $g, h$ be morphisms corresponding to $x$ and $y$ under this bijection, that is $x = \Psi_X(g)$ and $y = \Psi_X(h)$.

Then by naturality of $\Psi$

$$U(f)(x) = \Psi_Y(f \circ g)$$

and

$$U(f)(y) = \Psi_Y(f \circ h)$$

so by assumption $f \circ g = f \circ h$. As $f$ is monic we have $g = h$ whence $x = y$. Therefore $U(f)$ is injective. ∎

**Proposition 2.6.195 (Regular Epimorphism $\iff$ Surjective)**
*Let $\mathcal{C}$ be an algebraic category (2.6.184) and $e : X \to Y$ a morphism of $\mathcal{C}$. Then $e$ is a regular epimorphism if and only if $U(e)$ is surjective.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

($\Longrightarrow$) Since $e$ is regular, there are $\alpha, \beta : W \rightrightarrows X$ such that $e$ is the coequaliser of $(\alpha, \beta)$ (2.6.158).

**The image is a subalgebra.** Let $I \coloneqq U(e)(U(X)) \subseteq U(Y)$. For $\omega \in \Omega$ and $y_1, ..., y_{n_\omega} \in I$, writing $y_r = U(e)(x_r)$ for $x_r \in U(X)$, since $e$ is a homomorphism $\omega_Y(y_1, ...) = \omega_Y(U(e)(x_1), ...) = U(e)(\omega_X(x_1, ...)) \in I$; so $I$ is a subalgebra (2.6.192). By (2.6.193), $I$ carries an $\Omega$-algebra structure making the inclusion $\iota : I \hookrightarrow Y$ a homomorphism, and $I \in \mathcal{C}$ (as $Y \in \mathcal{C}$); as $\mathcal{C}$ is full in $\Omega$-**Alg**, $\iota$ is a morphism of $\mathcal{C}$.

Write $e' : X \to I$ for the corestriction of $e$, that is $U(e') \coloneqq U(e)$ regarded as a map into $I$; since $\omega_I$ is just $\omega_Y$ restricted to $I$, $e'$ is again a homomorphism, hence a morphism of $\mathcal{C}$, with $\iota \circ e' = e$ and $U(e')$ surjective onto $I$ by construction.

**$(I, e')$ is a coequaliser of $(\alpha, \beta)$.** Since $\iota \circ (e'\alpha) = e\alpha = e\beta = \iota \circ (e'\beta)$ and $\iota$ is monic (2.6.194), $e'\alpha = e'\beta$. Given $h : X \to Z$ with $h\alpha = h\beta$, the universal property of $e$ gives a unique $k : Y \to Z$ with $k \circ e = h$; then $k' \coloneqq k \circ \iota$ satisfies $k' \circ e' = k \circ \iota \circ e' = k \circ e = h$. If $k''$ also satisfies $k'' \circ e' = h$, then $U(k') \circ U(e') = U(h) = U(k'') \circ U(e')$; as $U(e')$ is surjective, $U(k') = U(k'')$, so $k' = k''$ ($U$ faithful). So $(I, e')$ is a coequaliser of $(\alpha, \beta)$ (2.6.140), in particular $e'$ is epic (2.6.135).

**$\iota$ is an isomorphism.** Applying the universal property of $e$ to $e'$ (which satisfies $e'\alpha = e'\beta$) gives a unique $\psi : Y \to I$ with $\psi \circ e = e'$. Then $(\iota \circ \psi) \circ e = \iota \circ e' = e = 1_Y \circ e$, and as $e$ is epic (2.6.135), $\iota \circ \psi = 1_Y$. Also $(\psi \circ \iota) \circ e' = \psi \circ e = e' = 1_I \circ e'$, and as $e'$ is epic, $\psi \circ \iota = 1_I$. So $\iota$ is an isomorphism, with inverse $\psi$.

So $U(\iota)$ is bijective; but $U(\iota)$ is, by construction, the inclusion $I \hookrightarrow U(Y)$, so $I = U(Y)$. As $I = U(e)(U(X))$ by definition, $U(e)$ is surjective.

($\Longleftarrow$) Suppose $U(e)$ is surjective.

**Construction of the quotient.** Write $\sim$ for the relation $a \sim b :\iff U(e)(a) = U(e)(b)$ on $U(X)$, an equivalence relation, and $Q \coloneqq U(X)/\sim$ with quotient map $\pi : U(X) \to Q$, $a \mapsto [a]$; $\pi$ is surjective by construction. For $\omega \in \Omega$ and $a_1, ..., a_{n_\omega}, b_1, ..., b_{n_\omega} \in U(X)$ with $a_r \sim b_r$ for every $r$: since $e$ is a homomorphism, $U(e)(\omega_X(a_1, ...)) = \omega_Y(U(e)(a_1), ...) = \omega_Y(U(e)(b_1), ...) = U(e)(\omega_X(b_1, ...))$, that is $\omega_X(a_1, ...) \sim \omega_X(b_1, ...)$; so $\sim$ is a congruence, and $\omega_Q\big([a_1], ..., [a_{n_\omega}]\big) \coloneqq \big[\omega_X\big(a_1, ..., a_{n_\omega}\big)\big]$ is well-defined, making $Q$ into an $\Omega$-algebra with $\pi$ a homomorphism.

**$Q \in \mathcal{C}$.** Let $p = q \in E$ with variables $W$, and $\theta : W \to Q$; as $\pi$ is surjective, choose $\tilde{\theta} : W \to U(X)$ with $\pi \circ \tilde{\theta} = \theta$. Since $X \in \mathcal{C}$, $X$ satisfies $p = q$, that is $\overline{\tilde{\theta}}(p) = \overline{\tilde{\theta}}(q)$ in $U(X)$; applying the homomorphism $\pi$ and (2.6.183) ($\pi \circ \overline{\tilde{\theta}} = \overline{\pi \circ \tilde{\theta}} = \overline{\theta}$),

$$\overline{\theta}(p) = \pi\Big(\overline{\tilde{\theta}}(p)\Big) = \pi\Big(\overline{\tilde{\theta}}(q)\Big) = \overline{\theta}(q)$$

© David Rufino CC BY-NC-ND 4.0

As $\theta$ was arbitrary, $Q$ satisfies $p = q$; as $p = q \in E$ was arbitrary, $Q \in \mathcal{C}$, and $\pi : X \to Q$ is a morphism of $\mathcal{C}$ (full subcategory of $\Omega$-**Alg**).

**The kernel relation is a subalgebra.** Let $R \coloneqq \{(a,b) \in U(X)^2 \mid a \sim b\}$. For $\omega \in \Omega$ and $(a_1,b_1), ..., \left(a_{n_\omega}, b_{n_\omega}\right) \in R$, since $e$ is a homomorphism, $U(e)(\omega_X(a_1,...)) = \omega_Y(U(e)(a_1),...) = \omega_Y(U(e)(b_1),...) = U(e)(\omega_X(b_1,...))$, so $(\omega_X(a_1,...), \omega_X(b_1,...)) \in R$; as $U(X \times X) = U(X)^2$ (2.6.190), $R$ is a subalgebra of $X \times X$ (2.6.192). By (2.6.193), $R \in \mathcal{C}$, and $\alpha \coloneqq \pi_1 \mid_R$, $\beta \coloneqq \pi_2 \mid_R$ (restrictions of the product projections) are morphisms of $\mathcal{C}$.

**$(Q, \pi)$ is a coequaliser of $(\alpha, \beta)$.** For $(a,b) \in R$, i.e. $a \sim b$, $\pi(\alpha(a,b)) = \pi(a) = [a] = [b] = \pi(\beta(a,b))$; as $U$ is faithful, $\pi \circ \alpha = \pi \circ \beta$. Suppose $h : X \to Z$ satisfies $h \circ \alpha = h \circ \beta$; then for every $(a,b) \in R$ (i.e. every $a \sim b$), $U(h)(a) = U(h)(b)$, so $\overline{h} : Q \to U(Z)$, $[a] \mapsto U(h)(a)$, is a well-defined map, and a homomorphism, since $\overline{h}(\omega_Q([a_1],...)) = \overline{h}([\omega_X(a_1,...)]) = U(h)(\omega_X(a_1,...)) = \omega_Z(U(h)(a_1),...) = \omega_Z\left(\overline{h}([a_1]),...\right)$, using that $h$ is a homomorphism. So $\overline{h}$ is a morphism of $\mathcal{C}$ with $\overline{h} \circ \pi = h$ ($U$ faithful). If $k : Q \to Z$ also satisfies $k \circ \pi = h$, then $U(k) \circ U(\pi) = U(h) = U\left(\overline{h}\right) \circ U(\pi)$; as $U(\pi)$ is surjective, $U(k) = U\left(\overline{h}\right)$, so $k = \overline{h}$ ($U$ faithful). So $(Q, \pi)$ is a coequaliser of $(\alpha, \beta)$ (2.6.140), in particular a regular epimorphism.

**Conclusion.** Define $\overline{e} : Q \to Y$, $[a] \mapsto U(e)(a)$; well-defined as $a \sim b \implies U(e)(a) = U(e)(b)$, and a homomorphism by the same computation as for $\pi$, hence a morphism of $\mathcal{C}$, with $\overline{e} \circ \pi = e$. As $\overline{e}([a]) = \overline{e}([b]) \implies U(e)(a) = U(e)(b) \implies a \sim b \implies [a] = [b]$, $U(\overline{e})$ is injective; as $e = \overline{e} \circ \pi$ and $U(e)$ is surjective (hypothesis), $U(\overline{e})$ is surjective. So $U(\overline{e})$ is bijective, hence $\overline{e}$ is an isomorphism, as $\mathcal{C}$ is algebraic and $U$ reflects isomorphisms (2.6.189).

Since $\pi$ is the coequaliser of $(\alpha, \beta)$ and $\overline{e}$ is an isomorphism, $e = \overline{e} \circ \pi$ is again a coequaliser of $(\alpha, \beta)$, as colimits are preserved under isomorphism (2.6.90); hence $e$ is a regular epimorphism (2.6.158). ■

**Proposition 2.6.196 (Algebraic Categories have Regular Epi-Mono Factorisations)**
*Let $\mathcal{C}$ be an algebraic category (2.6.184). Then $\mathcal{C}$ has regular epi-mono factorisations for every morphism (2.6.173).*

*More precisely, let $f : X \to Y$ be a morphism of $\mathcal{C}$ and $I \coloneqq U(f)(U(X)) \subseteq U(Y)$ the (underlying-set) image of $f$. Then $I$ is a subalgebra of $Y$, and $f = m \circ e$ is a regular epi-mono factorisation of $f$, where $e : X \to I$ is the corestriction of $f$ and $m : I \hookrightarrow Y$ is the inclusion.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

***I* is a subalgebra.** For $\omega \in \Omega$ and $y_1, ..., y_{n_\omega} \in I$, writing $y_r = U(f)(x_r)$ for $x_r \in U(X)$, since $f$ is a homomorphism $\omega_Y(y_1,...) = \omega_Y(U(f)(x_1),...) = U(f)(\omega_X(x_1,...)) \in I$; so $I$ is a subalgebra (2.6.192). By (2.6.193), $I$ carries a (unique) $\Omega$-algebra structure making the inclusion $m : I \hookrightarrow Y$ a homomorphism, with $I \in \mathcal{C}$ (as $Y \in \mathcal{C}$); as $\mathcal{C}$ is full in $\Omega$-**Alg**, $m$ is a morphism of $\mathcal{C}$, and monic by (2.6.194) (being injective).

Write $e : X \to I$ for the corestriction of $f$, that is $U(e) \coloneqq U(f)$ regarded as a map into $I$; since $\omega_I$ is just $\omega_Y$ restricted to $I$, $e$ is again a homomorphism, hence a morphism of $\mathcal{C}$, with $m \circ e = f$ and $U(e)$ surjective onto $I$ by construction. By (2.6.195), $e$ is therefore a regular epimorphism, so $f = m \circ e$ is a regular epi-mono factorisation of $f$ (2.6.173). ■

**Proposition 2.6.197 (Equalisers in Algebraic Categories)**
*Consider a commutative diagram in an algebraic category $\mathcal{C}$*

$$X \xrightarrow{\ e\ } Y \underset{d_1}{\overset{d_0}{\rightrightarrows}} Z$$

*Then $e$ is an equaliser for $(d_0, d_1)$ if and only if $U(e)$ is injective and $\mathrm{Im}(U(e)) = \{y \in U(Y) \mid U(d_0)(y) = U(d_1)(y)\}$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Write $F : \mathbf{2} \to \mathcal{C}$ for the diagram $d_0, d_1 : Y \rightrightarrows Z$, so that $e$ being an equaliser for $(d_0, d_1)$ means precisely that $(X, e)$ is a limit for $F$.

($\implies$) Suppose $(X, e)$ is a limit for $F$. Since $\mathcal{C}$ is algebraic, $U$ creates limits (2.6.189), so by (2.6.87), using that **Set** has all limits, $U$ preserves limits. Hence $(U(X), U(e))$ is a limit for $U \circ F$ in **Set**, that is, an equaliser for $(U(d_0), U(d_1))$. By (2.6.139), $U(e)$ is injective and $\mathrm{Im}(U(e)) = \{y \in U(Y) \mid U(d_0)(y) = U(d_1)(y)\}$, as required.

($\impliedby$) Suppose $U(e)$ is injective and $\mathrm{Im}(U(e)) = \{y \in U(Y) \mid U(d_0)(y) = U(d_1)(y)\}$.

By (2.6.139) the diagram $U(Y) \rightrightarrows U(Z)$ has an equaliser in **Set**, that is $\lim(U \circ F)$ exists. Since $U$ creates limits (2.6.189), $\lim F$ exists in $\mathcal{C}$; write $(X', e')$ for this limit, so that $e'$ is an equaliser for $(d_0, d_1)$. By the ($\implies$) direction applied to $e'$, $U(e')$ is injective and

$$\mathrm{Im}(U(e')) = \{y \in U(Y) \mid U(d_0)(y) = U(d_1)(y)\} = \mathrm{Im}(U(e))$$

© David Rufino CC BY-NC-ND 4.0

using the hypothesis on $e$ for the second equality.

As $d_0 \circ e = d_1 \circ e$, the universal property of the equaliser $(X', e')$ yields a unique morphism $\varphi : X \to X'$ with $e' \circ \varphi = e$.

We claim $U(\varphi)$ is a bijection. If $U(\varphi)(x) = U(\varphi)(y)$ then applying $U(e')$ gives $U(e)(x) = U(e')(U(\varphi)(x)) = U(e')(U(\varphi)(y)) = U(e)(y)$, so $x = y$ as $U(e)$ is injective; hence $U(\varphi)$ is injective. Given $x' \in U(X')$, we have $U(e')(x') \in \mathrm{Im}(U(e')) = \mathrm{Im}(U(e))$, so $U(e')(x') = U(e)(x)$ for some $x \in U(X)$, that is $U(e')(x') = U(e')(U(\varphi)(x))$, whence $x' = U(\varphi)(x)$ as $U(e')$ is injective; hence $U(\varphi)$ is surjective.

So $U(\varphi)$ is a bijection, hence an isomorphism in **Set**. Since $\mathcal{C}$ is algebraic, $U$ reflects isomorphisms (2.6.189), so $\varphi$ is an isomorphism in $\mathcal{C}$.

Since $(X', e')$ is an equaliser for $(d_0, d_1)$ and $\varphi : X \to X'$ is an isomorphism with $e' \circ \varphi = e$, by (2.6.137) $(X, e)$ is also an equaliser for $(d_0, d_1)$, as required. ∎

**Proposition 2.6.198 (Every Object has a Zero Element)**
*Let $\mathcal{C}$ be an algebraic pointed category. Then a zero object $\mathbf{0}$ consists of precisely one element, denoted by $0_{\mathbf{0}}$.*

*Every object $A$ has a unique element $0_A$ such that $U\big(0_{\mathbf{0},A}\big)(0) = 0_A$ for any (resp. all) zero objects $\mathbf{0}$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Since $\mathcal{C}$ is an algebraic category, $U$ creates limits (2.6.189), so by (2.6.87) $U$ preserves limits, and this holds unconditionally as the target category is **Set**. In particular $U$ preserves terminal objects, these being the limit over the empty diagram.

As $\mathbf{0}$ is a zero object it is in particular a terminal object of $\mathcal{C}$ (2.6.55), so $U(\mathbf{0})$ is a terminal object of **Set**. A terminal object of **Set** is a singleton, so $U(\mathbf{0})$ consists of precisely one element, which we denote $0_{\mathbf{0}}$.

Now fix an object $A$ and a zero object $\mathbf{0}$, and set

$$0_A \coloneqq U\big(0_{\mathbf{0},A}\big)(0_{\mathbf{0}}) \in U(A)$$

where $0_{\mathbf{0},A} : \mathbf{0} \to A$ is the unique morphism afforded by $\mathbf{0}$ being an initial object (2.6.48), and $0_{\mathbf{0}} \in U(\mathbf{0})$ is the unique element found above.

It remains to check that $0_A$ does not depend on the choice of zero object. Let $\mathbf{0}'$ be another zero object, with associated element $0'_A \coloneqq U\big(0_{\mathbf{0}',A}\big)(0_{\mathbf{0}'})$. Since $\mathbf{0}, \mathbf{0}'$ are both initial objects (2.6.55), by uniqueness of initial objects (2.6.48) there is a (unique) isomorphism $\varphi : \mathbf{0} \to \mathbf{0}'$. As $\mathrm{Mor}(\mathbf{0}, A)$ is a singleton, being initial, we have

$$0_{\mathbf{0}',A} \circ \varphi = 0_{\mathbf{0},A}$$

so applying $U$ and evaluating at $0_{\mathbf{0}}$

$$U\big(0_{\mathbf{0},A}\big)(0_{\mathbf{0}}) = U\big(0_{\mathbf{0}',A}\big)(U(\varphi)(0_{\mathbf{0}}))$$

Now $U(\varphi) : U(\mathbf{0}) \to U(\mathbf{0}')$ is a map between singletons, so it necessarily sends $0_{\mathbf{0}}$ to the unique element $0_{\mathbf{0}'}$ of $U(\mathbf{0}')$. Hence

$$0_A = U\big(0_{\mathbf{0},A}\big)(0_{\mathbf{0}}) = U\big(0_{\mathbf{0}',A}\big)(0_{\mathbf{0}'}) = 0'_A$$

so $0_A$ is independent of the choice of zero object, as required. ∎

**Proposition 2.6.199 (Kernels in Algebraic Categories)**
*Let $\mathcal{C}$ be an algebraic pointed category, and consider a diagram*

$$\ker(\alpha) \xrightarrow{i} Y \xrightarrow{\alpha} Z$$

*Then $(i, \ker(\alpha))$ is a kernel for $\alpha$ if and only if*

*a) $U(i)$ is injective*
*b) $\mathrm{Im}(U(i)) = \{y \in U(Y) \mid U(\alpha)(y) = 0_Z\}$*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Since $\mathcal{C}$ is pointed, by (2.6.150) $(i, \ker(\alpha))$ is a kernel for $\alpha$ if and only if $i$ is an equaliser for $\big(\alpha, 0_{Y,Z}\big)$, where $0_{Y,Z} : Y \to Z$ denotes the zero morphism (2.6.54).

As $\mathcal{C}$ is algebraic, (2.6.197) (applied with $d_0 = \alpha$, $d_1 = 0_{Y,Z}$) shows this holds if and only if $U(i)$ is injective and

© David Rufino CC BY-NC-ND 4.0

$$\mathrm{Im}(U(i)) = \{y \in U(Y) \mid U(\alpha)(y) = U(0_{Y,Z})(y)\}$$

It remains to identify $U(0_{Y,Z})(y)$ with $0_Z$ for every $y \in U(Y)$. Write $0_{Y,Z} = 0_{\mathbf{0},Z} \circ !_Y$, where $!_Y : Y \to \mathbf{0}$ and $0_{\mathbf{0},Z} : \mathbf{0} \to Z$ are the unique morphisms afforded by $\mathbf{0}$ being terminal, resp. initial (2.6.54). By (2.6.198), $U(\mathbf{0})$ is a singleton $\{0_{\mathbf{0}}\}$, so $U(!_Y)(y) = 0_{\mathbf{0}}$ for every $y \in U(Y)$, and therefore

$$U(0_{Y,Z})(y) = U(0_{\mathbf{0},Z})(U(!_Y)(y)) = U(0_{\mathbf{0},Z})(0_{\mathbf{0}}) = 0_Z$$

using the definition of $0_Z$ from (2.6.198). Hence

$$\{y \in U(Y) \mid U(\alpha)(y) = U(0_{Y,Z})(y)\} = \{y \in U(Y) \mid U(\alpha)(y) = 0_Z\}$$

so combining with the above, $(i, \ker(\alpha))$ is a kernel for $\alpha$ if and only if $U(i)$ is injective and $\mathrm{Im}(U(i)) = \{y \in U(Y) \mid U(\alpha)(y) = 0_Z\}$, as required. ∎

## 2.6.24 Exact Sequences

**Definition 2.6.200 (Exact Sequence)**
*Let $\mathcal{C}$ be a pointed category with an image of every morphism (2.6.167) – e.g. a pointed algebraic category (2.6.184), which has regular epi-mono factorisations (2.6.196), hence images (2.6.174).*

*Let $X \xrightarrow{f} Y \xrightarrow{g} Z$ be a pair of composable morphisms and $(\mathrm{Im}(f), e, m)$ an image of $f$. We say the pair is **exact at $Y$** if $\mathrm{Im}(f) = \ker(g)$, that is, $(\mathrm{Im}(f), m)$ is (literally) a kernel of $g$.*

*This does not depend on the choice of image of $f$: images are unique up to (unique) isomorphism (2.6.167), so given any other image $(\mathrm{Im}'(f), e', m')$ of $f$, there is an isomorphism $\theta : \mathrm{Im}(f) \to \mathrm{Im}'(f)$ with $m' \circ \theta = m$, and by (2.6.137) (applied to $\theta$ and to $\theta^{-1}$), $(\mathrm{Im}(f), m)$ is a kernel of $g$ if and only if $(\mathrm{Im}'(f), m')$ is.*

**Proposition 2.6.201 (Concrete Criterion, Algebraic Case)**
*Let $\mathcal{C}$ be a pointed algebraic category, and $\mathrm{Im}(f)$ the canonical (literal subalgebra) construction. Then the pair is exact at $Y$ (2.6.200) if and only if $f(X) = g^{-1}(\{0\})$, as subsets of $Y$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Write $I \coloneqq f(X) \subseteq Y$ for the canonical image of $f$ (2.6.196), a subalgebra of $Y$ with the literal inclusion $m : I \hookrightarrow Y$ as structure map, so $f = m \circ e$ (with $e$ the corestriction of $f$) is a regular epi-mono factorisation of $f$. By (2.6.200) (independence of representatives), we may use this particular representative: the pair is exact at $Y$ if and only if $(I, m)$ is a kernel of $g$, that is, by (2.6.150), $m$ is an equaliser of $(g, 0)$. By (2.6.197), since $m$ is (already) injective, this holds if and only if $\mathrm{Im}(m) = I$ equals $\{y \in Y \mid g(y) = 0\} = g^{-1}(\{0\})$. ∎

© David Rufino CC BY-NC-ND 4.0

## 2.7 Abelian Categories

Following [Fre64] we define an Abelian category without reference to additivity, however to make the presentation simpler we assume finite limits and colimits exist.

**Definition 2.7.1**
*We say a category $\mathcal{C}$ is Abelian if it*

*a) is pointed,*
*b) has finite limits and colimits, and*
*c) is binormal*

**Remark 2.7.2 (Abelian is Self-Dual)**
*If $\mathcal{C}$ is Abelian then so is $\mathcal{C}^{op}$. Indeed $\mathcal{C}^{op}$ is again pointed with the same zero object (2.6.54); finite limits in $\mathcal{C}$ become finite colimits in $\mathcal{C}^{op}$ and vice versa (2.6.92), so $\mathcal{C}^{op}$ again has finite limits and colimits; and $\mathcal{C}^{op}$ is binormal, since monic/epic swap under passing to $\mathcal{C}^{op}$ (2.6.33) while $\ker(f)/\mathrm{coker}(f)$ swap (2.6.154), so a monomorphism being a kernel in $\mathcal{C}$ corresponds to an epimorphism being a cokernel in $\mathcal{C}^{op}$ and vice versa.*

**Lemma 2.7.3**
*Let $\mathcal{C}$ be a normal pointed category and $m : I \hookrightarrow Y$ a monomorphism with cokernel $p : Y \to \mathrm{coker}(m)$. Then $f : X \to Y$ factors through $m$ if and only if $p \circ f = 0$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Suppose $f = m \circ g$ for some $g : X \to I$. Then $p \circ f = p \circ m \circ g = 0 \circ g = 0$, since $p \circ m = 0$ by definition of the cokernel (2.6.153).

Conversely, suppose $p \circ f = 0$. As $\mathcal{C}$ is normal and $m$ is monic, $m$ is a normal monomorphism (2.6.163), i.e. $(m, I)$ is the kernel of some morphism (2.6.159). By (2.6.162) it follows that $m = \ker(\mathrm{coker}(m)) = \ker(p)$. So by (2.6.150), $p \circ f = 0$ gives a (unique) morphism $g : X \to I$ such that $m \circ g = f$, that is, $f$ factors through $m$. ∎

**Lemma 2.7.4**
*Let $\mathcal{C}$ be an Abelian category and $f : X \to Y$ a morphism. Then the following are equivalent*

*a) $f$ is epic*
*b) $\mathrm{Im}(f) = Y$ (i.e. $(Y, f, 1_Y)$ is an image for $f$)*
*c) $\mathrm{coker}(f) = 0$*

*Dually, the following are equivalent*

*a) $f$ is monic*
*b) $\mathrm{Coim}(f) = X$ (i.e. $(X, 1_X, f)$ is a coimage for $f$)*
*c) $\ker(f) = 0$*

*Proof.* We only prove the first half, with the second following immediately by duality: since $\mathcal{C}$ is Abelian, so is $\mathcal{C}^{op}$ (2.7.2), and by (2.6.169) and (2.6.154) the conditions a)–c) are respectively dual to monic, $\mathrm{Coim}(f) = X$, and $\ker(f) = 0$.

a) $\implies$ c) This is (2.6.151).

c) $\implies$ b) Suppose $f = m \circ e$ where $m : I \hookrightarrow Y$ is a monomorphism. Consider the cokernel $p : Y \to \mathrm{coker}(m)$




Then $p \circ m = 0 \implies p \circ f = 0$. As $\mathrm{coker}(f) = 0$ there is a morphism $0 \to Z$ such that the bottom right triangle commutes, i.e. $p = 0$. Then $p \circ 1_Y = 0$ and so by (2.7.3) there exists $j : Y \to I$ such that $m \circ j = 1_Y$. As $m$ is monic this factorisation is unique and we conclude that $\mathrm{Im}(f) = Y$.

© David Rufino CC BY-NC-ND 4.0

b) $\implies$ a) Suppose that $\alpha \circ f = \beta \circ f$ and let $m : \mathrm{eq}(\alpha, \beta) \to Y$ be the equaliser, which is therefore monic (2.6.135).




By the universal property of the equaliser there is a (unique) factorisation $f = m \circ e$. Since $(Y, f, 1_Y)$ is the image of $f$ (hypothesis b)) and $(I, e, m)$ is another factorisation of $f$ through a monomorphism, by the universal property of the image (2.6.167) there is a unique morphism $j : Y \to I$ such that $m \circ j = 1_Y$. Therefore $m$ is split-epic so by (2.6.36) it is an isomorphism. Since $(I, m) = \mathrm{eq}(\alpha, \beta)$, by (2.6.136) we conclude $\alpha = \beta$. ∎

### 2.7.1 Images, Coimages and the First Isomorphism Theorem

In case of modules we have $\mathrm{coker}(f) = Y/\mathrm{Im}(f)$ and $\ker(\mathrm{coker}(f)) = \mathrm{Im}(f)$ so we might expect this relation to hold in the abstract setting.

**Proposition 2.7.5 ($\mathrm{Im}(f) = \ker(\mathrm{coker}(f))$ and $\mathrm{Coim}(f) = \mathrm{coker}(\ker(f))$)**
*Let $\mathcal{C}$ be a pointed category with kernels and cokernels and $f : X \to Y$ a morphism. Then there is a unique morphism $e : X \to \ker(\mathrm{coker}(f))$ making the following diagram commute*




*When $\mathcal{C}$ is Abelian $e$ is an epimorphism and $\mathrm{Im}(f) = (\ker(\mathrm{coker}(f)), e, m)$.*

*Dually, there is a unique morphism $m' : \mathrm{coker}(\ker(f)) \to Y$ such that the following diagram commutes*




*When $\mathcal{C}$ is Abelian $m'$ is a monomorphism and $\mathrm{Coim}(f) = (\mathrm{coker}(\ker(f)), e', m')$.*

*Proof.* Let $p : Y \to \mathrm{coker}(f)$ be the cokernel of $f$, so by (2.6.153) $p \circ f = 0$. Then by (2.6.150), applied to the kernel $m = \ker(p)$, $e$ exists and is unique.

Dually, let $i : \ker(f) \to X$ be the kernel of $f$, so by (2.6.150) $f \circ i = 0$. Then by (2.6.153), applied to the cokernel $e' = \mathrm{coker}(i)$, $m'$ exists and is unique.

Now suppose additionally that $\mathcal{C}$ is Abelian. We first show that $(\ker(\mathrm{coker}(f)), e, m) = \mathrm{Im}(f)$. By (2.6.135) $m$ is a monomorphism. Consider another factorisation $f = m'' \circ e''$ with $m''$ a monomorphism




Let $r := \mathrm{coker}(m'')$. Then $r \circ f = r \circ m'' \circ e'' = 0$. Therefore, by the universal property of $\mathrm{coker}(f)$ (2.6.153), there is a unique morphism $h : \mathrm{coker}(f) \to \mathrm{coker}(m'')$ such that $h \circ p = r$, and we deduce that $r \circ m = h \circ p \circ m = h \circ 0 = 0$. Therefore by (2.7.3) $m$ factors through $m''$ as required. Uniqueness follows because $m''$ is monic and therefore $(\ker(\mathrm{coker}(f)), e, m)$ is an image for $f$

Next we show $e$ is an epimorphism. By (2.7.4) applied to $e$ (with codomain $I := \ker(\mathrm{coker}(f))$), it suffices to show $\mathrm{Im}(e) = I$, i.e. that $(I, e, 1_I)$ is the image of $e$. Suppose $e = m_1 \circ e_1$ for some monomorphism $m_1 : I_1 \to I$. Then $f =$

© David Rufino CC BY-NC-ND 4.0

$m \circ e = (m \circ m_1) \circ e_1$ is a factorisation of $f$ through the monomorphism $m \circ m_1$, so by the universal property of the image $(I, e, m)$ of $f$ (2.6.167) (established above) there is a unique morphism $g : I \to I_1$ such that $(m \circ m_1) \circ g = m$; as $m$ is monic this gives $m_1 \circ g = 1_I$. This is precisely the factorisation required for $(I, e, 1_I)$ to be the image of $e$, so $\mathrm{Im}(e) = I$, and therefore $e$ is an epimorphism.

The remaining claims for the dual statement now follow by duality, applying the above to $f^{\mathrm{op}}$ in $\mathcal{C}^{\mathrm{op}}$, which is again Abelian (2.7.2): by (2.6.154), $\ker_{\mathcal{C}^{\mathrm{op}}}(\mathrm{coker}_{\mathcal{C}^{\mathrm{op}}}(f^{\mathrm{op}}))$ is identified with $\mathrm{coker}(\ker(f))$, matching (by uniqueness of the first paragraph, dualized) our $m'$ and $e'$, and by (2.6.33) and (2.6.169) the epic/image conclusions in $\mathcal{C}^{\mathrm{op}}$ translate to $m'$ monic and $(\mathrm{coker}(\ker(f)), e', m') = \mathrm{Coim}(f)$ in $\mathcal{C}$. ∎

In the category of modules we have $\mathrm{coker}(\ker(f)) = X/\ker(f)$ and a canonical isomorphism $X/\ker(f) \xrightarrow{\sim} \mathrm{Im}(f) = \ker(\mathrm{coker}(f))$, so we may expect something similar to hold true in the Abelian setting.

**Proposition 2.7.6 (First Isomorphism Theorem)**
*Let $\mathcal{C}$ be a pointed category with kernels and cokernels and $f : X \to Y$ a morphism. Then there is a unique morphism $\theta : \mathrm{coker}(\ker(f)) \to \ker(\mathrm{coker}(f))$ such that the following diagram commutes*

$$\begin{array}{ccc} X & \xrightarrow{\quad f \quad} & Y \\ \downarrow{\scriptstyle e'} & & \uparrow{\scriptstyle m} \\ \mathrm{coker}(\ker(f)) & \overset{\theta}{\dashrightarrow} & \ker(\mathrm{coker}(f)) \end{array}$$

*When $\mathcal{C}$ is an Abelian category then $\theta$ is an isomorphism. Furthermore both $(\mathrm{coker}(\ker(f)), e', m \circ \theta)$ and $(\ker(\mathrm{coker}(f)), \theta \circ e', m)$ are epi-mono factorisations of $f$ and both an image and coimage.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

By (2.7.5) there is a unique morphism $m' : \mathrm{coker}(\ker(f)) \to Y$ such that $f = m' \circ e'$.

Since $q \circ f = 0$, where $q : Y \to \mathrm{coker}(f)$ is the cokernel of $f$ (2.6.153), we have $q \circ m' \circ e' = q \circ f = 0$; as $e'$ is epic (2.6.135), $q \circ m' = 0$. So by the universal property of $m = \ker(q)$ (2.6.150), there is a unique morphism $\theta : \mathrm{coker}(\ker(f)) \to \ker(\mathrm{coker}(f))$ such that $m \circ \theta = m'$. Combining, $m \circ \theta \circ e' = m' \circ e' = f$, as required.

For uniqueness, suppose $\theta'$ also satisfies $m \circ \theta' \circ e' = f$. Then $m \circ (\theta \circ e') = f = m \circ (\theta' \circ e')$, and since $m$ is monic (2.6.135), $\theta \circ e' = \theta' \circ e'$; as $e'$ is epic (2.6.135), $\theta = \theta'$.

Suppose now $\mathcal{C}$ is Abelian, so that $f = m \circ e$ is the image factorisation and $f = m' \circ e'$ is the coimage factorisation (2.7.5). Since $m \circ \theta \circ e' = f = m \circ e$ and $m$ is monic (2.6.135), we may conclude that $\theta \circ e' = e$.

Since $m \circ \theta = m'$ is monic, $\theta$ is monic, and since $\theta \circ e' = e$ is epic, $\theta$ is epic. As $\mathcal{C}$ is Abelian it is binormal (2.7.1), hence balanced (2.6.164), so $\theta$, being both monic and epic, is an isomorphism.

Finally, applying (2.6.172) to the coimage $(\mathrm{coker}(\ker(f)), e', m')$, and dually to the image $(\ker(\mathrm{coker}(f)), e, m)$, via the isomorphism $\theta$ shows each factorisation is both an image and a coimage of $f$. ∎

**Proposition 2.7.7 (Epi-Mono Factorisation in an Abelian Category)**
*Let $\mathcal{C}$ be an Abelian category and $f : X \to Y$ a morphism. Then $f$ factors as $f = m \circ e$ for some epimorphism $e$ and monomorphism $m$*

$$\begin{array}{ccccc} X & & \xrightarrow{\quad f \quad} & & Y \\ & {\scriptstyle e}\searrow & & \nearrow{\scriptstyle m} & \\ & & I & & \end{array}$$

*and this factorisation is unique up to isomorphism: if $f = m_1 \circ e_1$ is another such factorisation, with $e_1 : X \to I_1$ epic and $m_1$ monic, then there is a unique morphism $\theta : I \to I_1$ such that the following diagram commutes, and $\theta$ is (moreover) an isomorphism*

© David Rufino CC BY-NC-ND 4.0




*Moreover, every epimorphism and every monomorphism of $\mathcal{C}$ is regular, so any such factorisation $f = m_1 \circ e_1$ is simultaneously a regular epi-mono and an epi-regular-mono factorisation of $f$, and hence $(I_1, e_1, m_1)$ is both an image and a coimage of $f$ (2.6.174).*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

**Regularity.** Since $\mathcal{C}$ is Abelian it is binormal (2.7.1), so every monomorphism is normal and every epimorphism is conormal (2.6.163), hence every monomorphism and every epimorphism of $\mathcal{C}$ is regular (2.6.160). In particular, for any epi-mono factorisation $f = m_1 \circ e_1$, $e_1$ is a regular epimorphism and $m_1$ is a regular monomorphism, so $f = m_1 \circ e_1$ is simultaneously a regular epi-mono and an epi-regular-mono factorisation of $f$, and by (2.6.174) (both halves), $(I_1, e_1, m_1)$ is both an image and a coimage of $f$.

**Existence.** By (2.7.5), taking $I \coloneqq \ker(\operatorname{coker}(f))$ with structure maps $e, m$, we obtain a factorisation $f = m \circ e$ with $m$ monic and, since $\mathcal{C}$ is Abelian, $e$ epic.

**Uniqueness up to isomorphism.** Let $f = m_1 \circ e_1$ be any other such factorisation, $e_1 : X \to I_1$. By the Regularity paragraph, $e$ and $e_1$ are regular epimorphisms, so $f = m \circ e$ and $f = m_1 \circ e_1$ are regular epi-mono factorisations of $f$ (2.6.173), and (2.6.178) gives a unique isomorphism $\theta : I \to I_1$ with $\theta \circ e = e_1$ and $m_1 \circ \theta = m$.

*Alternatively*, uniqueness may be shown directly from the universal property of the image (2.6.167), without invoking regular epi-mono factorisations at all.

**Uniqueness up to isomorphism.** Let $f = m_1 \circ e_1$ be any other epi-mono factorisation, $e_1 : X \to I_1$. Since $(I, e, m) = \operatorname{Im}(f)$ (2.7.5) and $m_1$ is monic, the universal property of the image (2.6.167) gives a unique morphism $\theta : I \to I_1$ such that $m_1 \circ \theta = m$, and (as noted there) $\theta \circ e = e_1$.

We show $\theta$ is an isomorphism. If $\theta k_1 = \theta k_2$ then $m_1 \theta k_1 = m_1 \theta k_2$, i.e. $m k_1 = m k_2$, and since $m$ is monic, $k_1 = k_2$; so $\theta$ is monic. Since $\theta \circ e = e_1$ is epic, $\theta$ is epic. As $\mathcal{C}$ is Abelian it is binormal (2.7.1), hence balanced (2.6.164), so $\theta$, being both monic and epic, is an isomorphism. ∎

## 2.7.2 Exact Sequences

**Lemma 2.7.8 (Cancellation for Kernels and Cokernels)**
*Let $\mathcal{C}$ be a pointed category with kernels and cokernels.*

*a) If $h = \psi \circ p$ with $\psi$ monic, then $(K, i)$ is a kernel of $p$ if and only if it is a kernel of $h$.*
*b) If $f = m \circ e$ with $e$ epic, then $(Q, q)$ is a cokernel of $m$ if and only if it is a cokernel of $f$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

a) For any $k$ into the common domain of $h, p$, $h \circ k = \psi \circ p \circ k$; since $\psi$ is monic, $\psi \circ p \circ k = 0 \iff p \circ k = 0$, so $h \circ k = 0 \iff p \circ k = 0$. In particular $p \circ i = 0 \iff h \circ i = 0$, and for any $i'$, $p \circ i' = 0 \iff h \circ i' = 0$; so the defining conditions of (2.6.150) hold for $(K, i)$ and $p$ if and only if they hold for $(K, i)$ and $h$, that is, $(K, i)$ is a kernel of $p$ if and only if it is a kernel of $h$.

b) Dually, for any $q'$ out of the common codomain of $f, m$, $q' \circ f = q' \circ m \circ e$; since $e$ is epic, $q' \circ m \circ e = 0 \iff q' \circ m = 0$, so $q' \circ f = 0 \iff q' \circ m = 0$; in particular $q \circ m = 0 \iff q \circ f = 0$, and for any $q'$, $q' \circ m = 0 \iff q' \circ f = 0$; so the defining conditions of (2.6.153) hold for $(Q, q)$ and $m$ if and only if they hold for $(Q, q)$ and $f$, that is, $(Q, q)$ is a cokernel of $m$ if and only if it is a cokernel of $f$. ∎

**Proposition 2.7.9 (Criteria for Exactness)**
*Let $\mathcal{C}$ be an Abelian category, $X \xrightarrow{f} Y \xrightarrow{g} Z$ a pair of composable morphisms, $f = m \circ e$ an epi-mono factorisation of $f$ and $g = m' \circ e'$ an epi-mono factorisation of $g$. Then the following are equivalent:*

*a) $m = \operatorname{Im}(f) = \ker(g)$ (i.e. exactness in the sense of (2.6.200));*

© David Rufino CC BY-NC-ND 4.0

*b)* $g \circ f = 0$*, and the canonical morphism* $\varphi : \mathrm{Im}(f) \to \ker(g)$ *is epic;*
*c)* $\mathrm{coker}(f) = \mathrm{Coim}(g) = e'$*.*
*d)* $g \circ f = 0$*, and the canonical morphism* $\varphi : \mathrm{coker}(f) \to \mathrm{Coim}(g)$ *is monic.*

*This does not depend on the choice of factorisation of either* $f$ *or* $g$*.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Write $k : \ker(g) \to Y$ for a kernel of $g$.

**Independence of factorisation of $f$.** By (2.6.200) (independence of representatives), applied with $\mathcal{C}$ Abelian – so every epi-mono factorisation of $f$ is (in particular) an image of $f$ (2.7.7) – exactness of the pair at $Y$, in particular the condition $m = \ker(g)$, does not depend on the choice of epi-mono factorisation of $f$: if $f = m_1 \circ e_1$ is another, then $m = \ker(g)$ if and only if $m_1 = \ker(g)$.

**Independence of factorisation of $g$.** If $g = m_1' \circ e_1'$ is another epi-mono factorisation of $g$, (2.7.7) gives an isomorphism $\phi : I' \to I_1'$ with $\phi \circ e' = e_1'$, so by (2.6.138) (applied to $\phi$ and to $\phi^{-1}$), $\mathrm{coker}(f) = e'$ if and only if $\mathrm{coker}(f) = e_1'$.

**Preliminary.** By (2.7.7), $(I', e', m')$ is both an image and a coimage of $g$; in particular $\mathrm{Coim}(g) = e'$.

a) $\Longrightarrow$ b) Taking $\varphi := 1_I$ gives $k \circ \varphi = k = m$; as $g \circ k = 0$ (2.6.150), $g \circ f = g \circ m \circ e = g \circ k \circ e = 0$. So (by uniqueness) $\varphi = 1_I$ is the morphism of b), and is epic, being an isomorphism.

b) $\Longrightarrow$ a) Since $m = k \circ \varphi$ and $m$ is monic, $\varphi$ is monic: if $\varphi \circ a = \varphi \circ b$ then $m \circ a = k \circ \varphi \circ a = k \circ \varphi \circ b = m \circ b$, so $a = b$. As $\varphi$ is also epic (b)), and $\mathcal{C}$ is Abelian hence balanced (2.6.164), $\varphi$ is an isomorphism; so $m = k \circ \varphi$ exhibits $m = \ker(g)$, that is, a).

a) $\Longrightarrow$ c) By (2.7.8) b) (as $e$ is epic), $\mathrm{coker}(f) = \mathrm{coker}(m)$; since $m = \ker(g)$ (a)), $\mathrm{coker}(m) = \mathrm{coker}(\ker(g)) = \mathrm{Coim}(g)$ (2.7.5), which by the Preliminary equals $e'$. So $\mathrm{coker}(f) = e'$, that is, c).

c) $\Longrightarrow$ a) By hypothesis $\mathrm{coker}(f) = e'$; since $g = m' \circ e'$, (2.7.8) a) (with $m'$ in the role of the monomorphism and $e' = \mathrm{coker}(f)$ in the role of $p$) gives $\ker(g) = \ker(\mathrm{coker}(f))$. By (2.7.5), $\mathrm{Im}(f) = \ker(\mathrm{coker}(f))$, i.e. (as $\mathrm{Im}(f) = (I, e, m)$) $m = \ker(\mathrm{coker}(f)) = \ker(g)$, that is, a).

c) $\Longleftrightarrow$ d) By duality: apply a) $\Longleftrightarrow$ b), established above for an arbitrary composable pair in an Abelian category, to $g^{\mathrm{op}} : Z \to Y$ and $f^{\mathrm{op}} : Y \to X$ in $\mathcal{C}^{\mathrm{op}}$ (Abelian, by (2.7.2)), using the epi-mono factorisation $g^{\mathrm{op}} = (e')^{\mathrm{op}} \circ (m')^{\mathrm{op}}$ of $g^{\mathrm{op}}$ (dual to $g = m' \circ e'$). This gives: $(e')^{\mathrm{op}} = \ker_{\mathcal{C}^{\mathrm{op}}}(f^{\mathrm{op}})$ if and only if $f^{\mathrm{op}} \circ g^{\mathrm{op}} = 0$ and the canonical morphism $\mathrm{Im}_{\mathcal{C}^{\mathrm{op}}}(g^{\mathrm{op}}) \to \ker_{\mathcal{C}^{\mathrm{op}}}(f^{\mathrm{op}})$ is epic in $\mathcal{C}^{\mathrm{op}}$.

Translating via (2.6.154) ($\ker_{\mathcal{C}^{\mathrm{op}}}(f^{\mathrm{op}})$ corresponds to $\mathrm{coker}(f)$), (2.6.169) ($\mathrm{Im}_{\mathcal{C}^{\mathrm{op}}}(g^{\mathrm{op}})$ corresponds to $\mathrm{Coim}(g)$), and (2.6.33) (epic in $\mathcal{C}^{\mathrm{op}}$ corresponds to monic in $\mathcal{C}$), while $f^{\mathrm{op}} \circ g^{\mathrm{op}} = (g \circ f)^{\mathrm{op}}$ vanishes if and only if $g \circ f = 0$: this reads exactly as $e' = \mathrm{coker}(f)$ if and only if $g \circ f = 0$ and the canonical morphism $\mathrm{coker}(f) \to \mathrm{Coim}(g)$ is monic, that is, c) $\Longleftrightarrow$ d).

**Independence of choice.** a) does not depend on the choice of epi-mono factorisation of $f$ (shown above), nor of $g$ (as a) only references the fixed kernel $k$, not the chosen factorisation of $g$); so, by the equivalences just established, neither do b), c), d), each being equivalent to a). ∎

**Remark 2.7.10**
*Conditions b)–d) give further equivalent characterisations of this same notion of exactness at* $Y$*, in particular the self-dual form* $\mathrm{coker}(f) = \mathrm{Coim}(g)$*.*

## 2.7.3 Biproducts

**Lemma 2.7.11**
*Let* $\mathcal{C}$ *be an Abelian category,* $X, Y \in \mathcal{C}$*, and* $\psi : X \coprod Y \to X \times Y$ *the canonical comparison morphism (2.6.104), where* $\pi_1, \pi_2$ *are the projections of* $X \times Y$ *and* $s_1, s_2$ *are the injections of* $X \coprod Y$*. Then* $\psi$ *is monic.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Write $X_1 := X$, $X_2 := Y$.

**Maps out of the coproduct.** By (2.6.93) let $q_1 := [1, 0] : X \coprod Y \to X$ and $q_2 := [0, 1] : X \coprod Y \to Y$, i.e. the morphisms induced by $q_k \circ s_i = \delta_{ki}$. Since $\pi_i \circ \psi \circ s_j = \delta_{ij} = q_i \circ s_j$ for all $j$, by the uniqueness clause of (2.6.93), $\pi_i \circ \psi = q_i$, that is, the following diagram commutes (which also displays the canonical injections $s_1, s_2$)

© David Rufino CC BY-NC-ND 4.0




**$\operatorname{coker}(s_1) = q_2$ and $\operatorname{coker}(s_2) = q_1$.** Indeed $q_1 s_2 = \delta_{12} = 0$ by construction. By (2.6.153), given $h : X \coprod Y \to Z$ with $h s_2 = 0$, we must produce a unique $g : X \to Z$ making the following diagram commute




By the uniqueness clause of (2.6.93), $h = [h s_1, h s_2]$ and, for any $g : X \to Z$, $g \circ [1, 0] = [g, 0]$ (both sides restrict to the same maps along $s_1, s_2$). So since $h s_2 = 0$, $h = [h s_1, 0] = (h s_1) \circ [1, 0] = (h s_1) \circ q_1$. Setting $g := h s_1$ then ensures the diagram commutes. Since $q_1 s_1 = \delta_{11} = 1_X$, $q_1$ is split-epic, hence epic (2.6.35), so $g$ is the unique such morphism. Therefore $\operatorname{coker}(s_2) = q_1$ as required. By the symmetric argument, $\operatorname{coker}(s_1) = q_2$.

**$s_1 = \ker(q_2)$ and $s_2 = \ker(q_1)$.** Since $q_1 s_1 = \delta_{11} = 1_X$, $s_1$ is split-monic, hence monic (2.6.35); similarly $s_2$ is monic (via $q_2 s_2 = \delta_{22} = 1_Y$). As $\mathcal{C}$ is Abelian it is binormal (2.7.1), so the monomorphism $s_2$ is normal (2.6.163), i.e. the kernel of some morphism (2.6.159), whence by (2.6.162), $s_2 = \ker(\operatorname{coker}(s_2)) = \ker(q_1)$. Similarly $s_1 = \ker(\operatorname{coker}(s_1)) = \ker(q_2)$.

**$\ker(\psi) = 0$.** Since $\pi_1, \pi_2$ jointly determine morphisms into $X \times Y$ (2.6.81), $\psi \circ k = 0$ iff $q_1 k = 0$ and $q_2 k = 0$, that is, $\ker(\psi)$ is the intersection, as subobjects of $X \coprod Y$, of $\ker(q_1) = s_2$ and $\ker(q_2) = s_1$. Informally, $\ker(q_1)$ is the "$Y$-summand" of $X \coprod Y$ (as $q_1$ is the identity on $X$ and kills $Y$) and $\ker(q_2)$ is the "$X$-summand", so this intersection should vanish just as the coordinate subspaces $X \times \{0\}$ and $\{0\} \times Y$ meet only at the origin inside $X \oplus Y$; we verify this directly.

Suppose $k : K \to X \coprod Y$ satisfies $\psi \circ k = 0$. Then $q_1 k = \pi_1 \psi k = 0$ and $q_2 k = \pi_2 \psi k = 0$. Since $s_2 = \ker(q_1)$ there is $\alpha : K \to Y$ with $s_2 \alpha = k$, and since $s_1 = \ker(q_2)$ there is $\beta : K \to X$ with $s_1 \beta = k$, so $s_2 \alpha = s_1 \beta$, that is, the following diagram commutes




Composing with $q_1$: $q_1 s_2 \alpha = q_1 s_1 \beta$ gives $0 = \beta$. Therefore $k = s_1 \beta = 0$.

By (2.6.166), $\ker(\psi) = 0$, so by (2.7.4), $\psi$ is monic. ∎

**Proposition 2.7.12 (Finite Biproducts Exist)**
*Let $\mathcal{C}$ be an Abelian category. Then $\mathcal{C}$ has all finite biproducts (2.6.103).*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

By (2.6.105) it suffices to show that binary biproducts exist. Fix $X, Y \in \mathcal{C}$; by (2.6.104) there is a unique comparison morphism $\psi : X \coprod Y \to X \times Y$, with $\pi_1, \pi_2$ the projections of $X \times Y$ and $s_1, s_2$ the injections of $X \coprod Y$, and it suffices to show $\psi$ is an isomorphism.

By (2.7.11), $\psi$ is monic. Since $\mathcal{C}^{\mathrm{op}}$ is again Abelian (2.7.2), in which $X \times Y$ (with $\pi_1, \pi_2$) and $X \coprod Y$ (with $s_1, s_2$) play the roles of the coproduct and product respectively (2.6.92), the canonical comparison morphism for $X, Y$ in $\mathcal{C}^{\mathrm{op}}$ is

© David Rufino CC BY-NC-ND 4.0

exactly $\psi^{\mathrm{op}}$, since both are characterised by the same defining equations $\pi_i \circ (-) \circ s_j = \delta_{ij}$. So applying (2.7.11) to $\mathcal{C}^{\mathrm{op}}$, $\psi^{\mathrm{op}}$ is monic in $\mathcal{C}^{\mathrm{op}}$, i.e. $\psi$ is epic in $\mathcal{C}$ (2.6.33).

As $\mathcal{C}$ is binormal it is balanced (2.6.164), so $\psi$, being both monic and epic, is an isomorphism. By (2.6.104), a biproduct $X \oplus Y$ therefore exists. ∎

**Proposition 2.7.13 ($\mathrm{Mor}(X,Y)$ is an Abelian Group)**
*Let $\mathcal{C}$ be an Abelian category and $X, Y \in \mathcal{C}$. Then every $f \in \mathrm{Mor}(X,Y)$ has an additive inverse; combined with (2.6.121) (applicable since $\mathcal{C}$ has all finite biproducts (2.7.12)), $(\mathrm{Mor}(X,Y), +)$ is an abelian group.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.* Following [Fre64] Theorem 2.39.

The underlying idea: in a category of modules, the inverse of the unitriangular matrix $\left(\begin{smallmatrix} 1 & 0 \\ f & 1 \end{smallmatrix}\right)$ is $\left(\begin{smallmatrix} 1 & 0 \\ -f & 1 \end{smallmatrix}\right)$, so the additive inverse of $f$ can be read off from the inverse of this matrix; we therefore try to show that the same matrix $M$ is an isomorphism in any Abelian category.

Fix $f : X \to Y$ and let $M := \left(\begin{smallmatrix} 1_X & 0 \\ f & 1_Y \end{smallmatrix}\right) : X \oplus Y \to X \oplus Y$ (2.6.108). By (2.6.122), matrices now multiply in the usual way, entry-by-entry, dot-product style.

*$M$ is monic.* For $g = \left(\begin{smallmatrix} g_1 \\ g_2 \end{smallmatrix}\right) : Z \to X \oplus Y$

$$M \circ g = \begin{pmatrix} 1_X & 0 \\ f & 1_Y \end{pmatrix} \circ \begin{pmatrix} g_1 \\ g_2 \end{pmatrix} = \begin{pmatrix} g_1 \\ f \circ g_1 + g_2 \end{pmatrix}$$

so $M \circ g = 0$ gives $g_1 = 0$ and $f \circ g_1 + g_2 = 0$; substituting, $f \circ 0 + g_2 = 0$, i.e. $g_2 = 0$, so $g = 0$. By (2.6.166), $\ker(M) = 0$, so by (2.7.4), $M$ is monic.

*$M$ is epic.* Dually, for $h = \left(\begin{smallmatrix} h_1 & h_2 \end{smallmatrix}\right) : X \oplus Y \to Z$

$$h \circ M = \begin{pmatrix} h_1 & h_2 \end{pmatrix} \circ \begin{pmatrix} 1_X & 0 \\ f & 1_Y \end{pmatrix} = \begin{pmatrix} h_1 + h_2 \circ f & h_2 \end{pmatrix}$$

so $h \circ M = 0$ gives $h_2 = 0$ and $h_1 + h_2 \circ f = 0$, i.e. $h_1 = 0$, so $h = 0$. Dually, $\mathrm{coker}(M) = 0$, so by (2.7.4), $M$ is epic.

*$M$ is invertible.* Since $\mathcal{C}$ is Abelian it is binormal (2.7.1), hence balanced (2.6.164), so $M$, being monic and epic, is an isomorphism.

*Extracting the inverse.* Write $M^{-1} = \left(\begin{smallmatrix} p & q \\ r & s \end{smallmatrix}\right)$. Then

$$\begin{pmatrix} 1_X & 0 \\ f & 1_Y \end{pmatrix} \circ \begin{pmatrix} p & q \\ r & s \end{pmatrix} = \begin{pmatrix} p & q \\ f \circ p + r & f \circ q + s \end{pmatrix} = \begin{pmatrix} 1_X & 0 \\ 0 & 1_Y \end{pmatrix}$$

gives $p = 1_X$, $q = 0$, $f \circ p + r = 0$ and $f \circ q + s = 1_Y$. Substituting $p = 1_X$ into the third equation gives $f + r = 0$: so $f$ has additive inverse $-f := r$. ∎

© David Rufino CC BY-NC-ND 4.0

## Chapter 3

# Algebra

Follows largely Lang with some Bourbaki.

© David Rufino CC BY-NC-ND 4.0

# 3.1 Magmas and Monoids

**Definition 3.1.1 (Magma)**
*Let $X$ be a set. A **law of composition** on $X \times X$ is a function*

$$\cdot : X \times X \to X$$

*and we typically write the composition of $x, y \in X$ as either*

$$x \cdot y$$

*or $xy$, or $x + y$ in the commutative case.*

*A pair $(X, \cdot)$ consisting of a set $X$ and law of composition on $X$ is called a **magma**.*

**Definition 3.1.2 (Magma/Monoid)**
*A **magma** $(X, \cdot)$ is said to be*

- ***associative** if $(x \cdot y) \cdot z = x \cdot (y \cdot z)$*
- ***commutative** if $x \cdot y = y \cdot x$ for all $x, y \in X$*
- ***unital** if there exists $e \in X$ such that $e \cdot x = x \cdot e = x$ for all $x \in X$. Such an $e$ is called an **identity**.*
- *a **monoid** if it is both **associative** and **unital***

**Proposition 3.1.3 (Identity is Unique)**
*A magma $(X, \cdot)$ has at most one element $e$ such that*

$$x \cdot e = e \cdot x = x$$

*for all $x \in X$.*

**Definition 3.1.4 (Invertible / Monoid)**
*Let $(X, \cdot)$ be a unital magma. An element $x \in X$ is **invertible** if there exists $y \in X$ such that*

$$x \cdot y = y \cdot x = e$$

**Proposition 3.1.5 (Inverses are unique)**
*Let $(X, \cdot)$ be a monoid. If $x \in X$ is invertible then its inverse is unique and denoted $x^{-1}$.*

*Proof.* Suppose that $xy = xy' = e = yx = y'x$. Then

$$xy = e \implies y'(xy) = y'e = y' \implies (y'x)y = y' \implies y = y'$$

■

**Definition 3.1.6 (Homomorphism)**
*Let $(X, \cdot)$, $(Y, \cdot)$ be magmas. Then a function $\varphi : X \to Y$ is said to be a **magma homomorphism** if it satisfies*

$$\varphi(x_1 \cdot x_2) = \varphi(x_1) \cdot \varphi(x_2) \quad \forall x_1, x_2 \in X$$

*If $(X, \cdot)$ and $(Y, \cdot)$ are unital then $\varphi$ is **unital** if*

$$\varphi(e_X) = e_Y$$

*If $(X, \cdot)$ and $(Y, \cdot)$ are monoids then $\varphi$ is a **monoid morphism** if it satisfies both these conditions.*

**Proposition 3.1.7 ($\mathbb{N}$ is an initial object)**
*Let $(X, \cdot)$ be a monoid and $x \in X$. Then there is a unique monoid morphism*

$$x^{(-)} : (\mathbb{N}, +) \to (X, \cdot)$$

*such that*

$$x^1 = x$$

*Furthermore if $\varphi : (X, \cdot) \to (Y, \cdot)$ is a monoid morphism then*

$$\varphi(x^n) = \varphi(x)^n$$

*for all $x \in X$ and $x \in \mathbb{N}$.*

© David Rufino CC BY-NC-ND 4.0

**Proposition 3.1.8 ($\mathbb{Z}$ is an initial object)**
*Let $(X, \cdot)$ be a monoid and $x \in X$ be an invertible element. Then there is a unique monoid morphism*

$$x^{(-)} : (\mathbb{Z}, +) \to (X, \cdot)$$

*such that*

$$x^1 = x$$

*and*

$$x^{-n} \text{ is the inverse of } x^n$$

*Furthermore if $\varphi : (X, \cdot) \to (Y, \cdot)$ is monoid morphism then*

$$\varphi(x^n) = \varphi(x)^n$$

*for all $x \in X$ invertible and $n \in \mathbb{Z}$.*

© David Rufino CC BY-NC-ND 4.0

## 3.2 Groups

**Definition 3.2.1 (Group)**
*A **group** is a **monoid** $(G, \cdot)$ in which every element is invertible.*

*A group $G$ is said to be **abelian** if the binary operation is **commutative**. In this case we typically write the group operation additively*

$$g + h$$

**Definition 3.2.2 (Subgroup, Normal Subgroup)**
*A **subgroup** $H \leq G$ is a subset with the following properties*

- $e_G \in H$
- $x, y \in H \implies xy \in H$
- $x \in H \implies x^{-1} \in H$

*A subgroup $H$ is said to be **normal** in $G$ if in addition it satisfies*

$$gHg^{-1} \coloneqq \{ghg^{-1} \mid g \in G\} = H$$

*for all $g \in G$. NB it is easily verified that in an abelian group every subgroup is normal.*

**Proposition 3.2.3 (Subgroup is a group)**
*Let $H$ be a subgroup of $(G, \cdot)$ then $(H, \cdot|_{H \times H})$ is a group.*

**Example 3.2.4**
*$\mathbb{Z}$ is an abelian group under addition. The subgroups are of the form $n\mathbb{Z}$.*

**Definition 3.2.5**
*Let $(G, \cdot)$ and $(H, \cdot)$ be groups. A function $\varphi : G \to H$ is a **group homomorphism** if*

- $\varphi(e_G) = e_H$
- $\varphi(x \cdot y) = \varphi(x) \cdot \varphi(y)$

*Define the **image** of $\varphi$ to be*

$$\mathrm{Im}(\varphi) = \{\varphi(g) \mid g \in G\}$$

*and the **kernel** to be*

$$\ker(\varphi) \coloneqq \{g \in G \mid \varphi(g) = e_H\}$$

*It may be verified that $\mathrm{Im}(\varphi)$ is a subgroup of $H$ and $\ker(\varphi)$ is a normal subgroup of $G$.*

**Proposition 3.2.6 (Raise to the $n$-th power)**
*Let $g \in G$ be a group element. Then there exist a unique group homomorphism*

$$g^{(-)} : (\mathbb{Z}, +) \to (G, \cdot)$$

*satisfying*

$$g^1 = g$$

*In other words such that*

$$g^0 = e_G$$

$$g^{n+m} = g^n \cdot g^m \quad \forall n, m \in \mathbb{Z}$$

**Proposition 3.2.7**
*Let $g \in G$ be a group element. Then*

$$(g^n)^m = g^{nm}$$

*for all integers $n, m \in \mathbb{Z}$.*

**Definition 3.2.8 (Order of an element)**
*For $g \in G$ define the **order** of $g$ to be $o(g) \coloneqq \inf\{n \geq 0 \mid g^n = e\}$ where $\inf \varnothing = \infty$.*

© David Rufino CC BY-NC-ND 4.0

*We say $g$ has **finite order** if $o(g) \neq \infty$.*

**Definition 3.2.9 (Subgroup generated by an element)**
*The **subgroup generated** by an element $g \in G$ is defined to be $\langle g \rangle = \{g^n \mid n \in \mathbb{Z}\}$*

*It may be shown that when $g$ has finite order $n$ we have*

$$\langle g \rangle = \{e, g, ..., g^{n-1}\}$$

*and in particular $\#\langle g \rangle = o(g)$.*

**Proposition 3.2.10 (Cosets)**
*Let $H$ be a subgroup of $G$. The following is an equivalence relation on $G$*

$$g_1 \underset{H}{\sim} g_2 \iff g_1 g_2^{-1} \in H$$

*and the equivalence classes are precisely the sets of the form*

$$gH = \{gh \mid h \in H\} = [g]_{\underset{H}{\sim}}$$

*for some $g \in G$. Such an equivalence class is called a **right coset** and we denote the set of right cosets by*

$$G/H$$

*Define the **index** of $H$ in $G$ by $[G : H] := \#G/H$. When $H$ is finite each equivalence class has order $\#H$.*

*We say $\{g_i \in G\}_{i \in I}$ is a set of **coset representatives** for $H$ if the corresponding equivalence classes $\{[g_i]\}_{i \in I}$ are pairwise disjoint and cover $G$.*

*Proof.* It's trivial to show that $\underset{H}{\sim}$ is an equivalence relation (precisely because $H$ is a subgroup). Therefore by (2.1.6) the equivalence classes form a partition which we denote $G/H$.

We claim that $[g_1] = g_1 H$. Then $g_2 \in [g_1] \iff g_1 \underset{H}{\sim} g_2 \iff g_2 \underset{H}{\sim} g_1 \iff g_2 g_1^{-1} \in H \iff g_2 \in g_1 H$, which shows that the sets are equal.

The translation map $\psi_g : G \to G$ given by $g' \to gg'$ is bijective (for it has a two-sided inverse equal to $\psi_{g^{-1}}$). So in particular restricts to a bijective map $H \to gH$. This shows that all the cosets have the same order. ∎

**Example 3.2.11**
*$d\mathbb{Z}$ is a subgroup of $\mathbb{Z}$ of index $d$. A set of coset representatives are $\{0, 1, ..., d-1\}$.*

**Corollary 3.2.12 (Lagrange's Theorem)**
*Let $H \leq G$ be a subgroup then*

$$\#G = [G : H] \times \#H$$

*More generally if $K \leq H$ then*

$$[G : K] = [G : H][H : K]$$

**Example 3.2.13**
*$d\mathbb{Z} \subseteq e\mathbb{Z} \iff d \mid e$ and $[e\mathbb{Z} : d\mathbb{Z}] = e/d$.*

**Proposition 3.2.14**
*Let $g \in G$ be an element of finite order. Then*

$$o(g) \mid \#G$$

*Furthermore*

$$g^n = e \iff o(g) \mid n$$

*Proof.* The first statement follows because the order $o(g)$ equals the order of the subgroup $\langle g \rangle$ generated by $g$.

Let $m = o(g)$ then by the division algorithm $n = qm + r$ for some $r < m$. Then $e = g^n = g^{qm} g^r = (g^m)^q g^r = e^q g^r = g^r$. By minimality we have $r = 0$ and $m \mid n$ as required. ∎

**Proposition 3.2.15 (Quotient Group)**
*Let $N$ be a normal subgroup $G$. Then the set of cosets*

© David Rufino CC BY-NC-ND 4.0

$$G/N$$

*forms a group under the binary operation*

$$g_1N \cdot g_2N \to (g_1g_2)N$$

*with identity $eN$.*

a) *There is a canonical surjective group homomorphism*

$$\pi : G \to G/N$$

$$g \to gN$$

*with kernel $N$.*

b) *Let $N \subseteq H$ be a subgroup then define the correseponding subgroup of $G/N$*

$$H/N \coloneqq \pi(H) = \{hN \mid h \in H\}.$$

c) *Let $\varphi : G \to G'$ be a homomorphism with $N \subseteq \ker(\varphi)$, then there exists a unique homomorphism $\tilde{\varphi}$ making the diagram commute*




*such that*

a) $\mathrm{Im}(\varphi) = \mathrm{Im}(\tilde{\varphi})$
b) $\ker(\tilde{\varphi}) = \ker(\varphi)/N$

**Corollary 3.2.16 (Isomorphism Theorem)**
*Let $\varphi : G \to H$ be a group homomorphism, then there is a canonical isomorphism*

$$G/\ker(\varphi) \xrightarrow{\sim} \mathrm{Im}(\varphi)$$

**Proposition 3.2.17 (Correspondence Theorem)**
*Let $\pi : G \to G'$ be a surjective homomorphism with $\ker(\varphi) = N$ then there is a bijective correspondence of subgroups*

$$\begin{aligned} \{H \leq G \mid N \subseteq H\} &\longleftrightarrow \{H' \leq G'\} \\ H &\to \pi(H) \\ \pi^{-1}(H') &\leftarrow H' \end{aligned}$$

*which preserves index, that is*

$$[G' : H'] = [G : H]$$

*Furthermore $\#H' = [H : N]$.*

**Proposition 3.2.18**
*Suppose $N \subset G$ and $K \subset N$ are normal subgroups. Then $K \subset G$ is normal and there is a canonical isomorphism*

$$(G/N)/(K/N) \xrightarrow{\sim} G/K$$

*given by*

$$\overline{g+N} \to g + K$$

### 3.2.1 Cyclic Groups

**Definition 3.2.19**
*A group $G$ is said to be **cyclic** if there is a surjective group homomorphism*

$$(\mathbb{Z}, +) \longrightarrow (G, \cdot)$$

*equivalently if there is $g \in G$ such that $\langle g \rangle = G$. Such a $g$ is called a **generator** for $G$.*

© David Rufino CC BY-NC-ND 4.0

**Proposition 3.2.20**
*Consider the additive group $(\mathbb{Z}, +)$. Then*

a) *Every subgroup is of the form $d\mathbb{Z}$ for $d \geq 0$ and is itself cyclic*

b) *When $d > 0$, then $Z/d\mathbb{Z}$ has a complete set of coset representatives*

$$S := \{0, 1, ..., d-1\}$$

c) *In particular $[Z : d\mathbb{Z}] = d$ when $d > 0$*

d) *$d\mathbb{Z} \subseteq e\mathbb{Z} \iff e \mid d$ and in this case $[e\mathbb{Z} : d\mathbb{Z}] = \frac{d}{e}$*

*Proof.* We prove each in turn

a) By (2.2.6) every subgroup is of the form $d\mathbb{Z}$. Multiplication map $[d] : \mathbb{Z} \to d\mathbb{Z}$ shows it is itself cyclic.
b) By the division algorithm (2.2.5) $S$ is a complete set. Given $i, j \in S$ we note that $|i - j| < d$. And $i \underset{d}{\sim} j \implies d \mid |i - j| \implies |i - j| = 0 \implies i = j$. Therefore the set $S$ consists of distinct coset representatives.
c) This is clear from the previous step
d) The first equivalence is clear. By (3.2.12)

$$[\mathbb{Z} : d\mathbb{Z}] = [\mathbb{Z} : e\mathbb{Z}][e\mathbb{Z} : d\mathbb{Z}]$$

and the result follows.

■

**Proposition 3.2.21**
*Let $G$ be a cyclic group. Then*

- *If $G$ is infinite it is isomorphic to $\mathbb{Z}$*
- *If $G$ is finite it is isomorphic to $\mathbb{Z}/n\mathbb{Z}$ for some $n > 0$*

*Proof.* By the previous Proposition the kernel of the homomorphism $\mathbb{Z} \to G$ is of the form $n\mathbb{Z}$ for $n \geq 0$. By (...) $G$ is isomorphic to $\mathbb{Z}/n\mathbb{Z}$. When $n = 0$ this is canonically isomorphic to $\mathbb{Z}$.

By the previous Proposition $\mathbb{Z}/n\mathbb{Z}$ is finite for $n > 0$ and therefore if $G$ is not finite we must have $n = 0$. Similarly If $G$ is finite then we must have $n > 0$. ■

We analyse the structure of finite cyclic groups in more detail. First recall the definition of **Euler's Totient Function**

**Definition 3.2.22 (Euler Totient Function)**
*Define the function*

$$\phi(n) = \#\{0 < d \leq n \mid (d, n) = 1\}$$

**Proposition 3.2.23 (Finite Cyclic Groups)**
*Consider a finite cyclic group $G$ of order $n$. Then*

a) *The order of $g^r$ is $\frac{n}{n,r}$ where $0 < r \leq n$.*
b) *There are $\phi(n)$ generators*
c) *For every $d \mid n$ there is a unique subgroup of order $\frac{n}{d}$ given by $\langle g^d \rangle$, which is cyclic.*
d) *For $d \mid n$ there are precisely $\phi(d)$ elements of order $d$*
e) *There are precisely $d$ elements of order dividing $d$*

*Proof.* We prove each in turn

a) $(g^r)^s = e_G \iff g^{rs} = e_G \overset{(3.2.14)}{\iff} n \mid rs \overset{(2.2.13)}{\iff} \frac{n}{n,r} \mid s$. Therefore $g^r$ has order $\frac{n}{n,r}$ as required.
b) Note $h$ is a generator iff $o(h) = n$. So $g^r$ is a generator iff $(n, r) = 1$ by the previous step. As $G = \{g, g^2, ..., g^n\}$ the result follows by definition of the totient function.
c) Recall there is a canonical surjective morphism $\pi : \mathbb{Z} \to G$ with kernel $n\mathbb{Z}$ and $\pi(1) = g$. By (3.2.17) the subgroups $H$ of $G$ correspond bijectively to subgroups $H'$ of $\mathbb{Z}$ containing $n\mathbb{Z}$, preserving the index. By (3.2.20) these are of the form $H' = d\mathbb{Z}$ for $d \mid n$, which correspond under $\pi$ to subgroups $H = \langle g^d \rangle$. Further $[G : \langle g^d \rangle] = [\mathbb{Z} : d\mathbb{Z}] = d$ whence $\#\langle g^d \rangle = \frac{n}{d}$. By definition $\langle g^d \rangle$ is cyclic.
d) Let $G[d]$ be the unique (cyclic) subgroup of order $d$. If $h$ has order $d$ then $\langle h \rangle$ has order $d$, and therefore by uniqueness is equal to $G[d]$. In particular $h \in G[d]$. Therefore by the previous part there are $\phi(d)$ elements of order $d$

© David Rufino CC BY-NC-ND 4.0

e) Suppose $h$ has order $e \mid d$. Both $G$ and $G[d]$ contain a unique subgroup of order $e$ and therefore by uniqueness this is simply $G[e] \subseteq G[d]$. Similarly by uniqueness $G[e] = \langle h \rangle$. Therefore $h \in G[d]$. Conversely suppose $h \in G[d]$ then $o(g) \mid d$ by (3.2.14). Therefore $G[d]$ consists of all the elements of order dividing $d$.

■

**Corollary 3.2.24**
*Let $n$ be a positive integer then*

$$n = \sum_{d|n} \phi(d)$$

*Proof.* Consider a cyclic group $G$ of order $n$. Every element has order dividing $n$ so the result follows from the previous Proposition by partitioning the group $G$ into subsets consisting of elements of equal order. ■

For an abelian group $G$ define the following subgroup

$$G[d] := \{g \in G \mid g^d = e\}.$$

We have shown for a cyclic group that $\#G[d] = d$ whenever $d \mid n$ and it is empty otherwise. We claim that this can be used to characterize cyclic groups. NB the following is adapted from this stackexchange answer

**Proposition 3.2.25 (Characterization of cyclic group)**
*Let $G$ be a finite abelian group such that $\#G[d] \leq d$ for all $d \mid n$. Then $G$ is cyclic.*

*Proof.* Let $n = \#G$ and $G_d$ be the subset of elements of order exactly $d$. Then we wish to show that $G_n$ is non-empty as any element of this set will be a generator. We actually show that $\#G_d = \phi(d) > 0$ whenever $d \mid n$.

Note that $G_d \subseteq G[d]$. If it's non-empty then for any $y \in G_d$, we have $\langle y \rangle$ is a subgroup of $G[d]$ of order $d$. As $\#G[d] \leq d$ we have $G[d] = \langle y \rangle$ is cyclic of order $d$. In other words $G_d$ is equal to the set of generators for $G[d]$. By the previous Proposition $G[d]$ has $\phi(d)$ generators. We conclude that for all $d \mid n$ we have $G_d$ is either empty or of order $\phi(d)$.

Therefore

$$n = \sum_{d|n} \#G_d \leq \sum_{d|n} \phi(d) = n$$

Therefore we must have equality everywhere and $\#G_d = \phi(d)$ as required. ■

**Example 3.2.26**
*Let $G = \mathbb{Z}/p\mathbb{Z} \times \mathbb{Z}/p\mathbb{Z}$ of order $p^2$. Then $G[p] = G$ so $\#G[p] = p^2 > p$.*

## 3.2.2 Group Actions

**Definition 3.2.27 (Group Action)**
*Let $G$ be a group and $S$ a set. A group action of $G$ on $S$ is a map*

$$G \times S \longrightarrow S$$

$$(g, s) \longrightarrow g \cdot s$$

*such that*

- $es = s$
- $g(hs) = (gh)s$

**Definition 3.2.28 (Faithful group action)**
*A group action $G$ on $S$ is faithful if*

$$gs = s \quad \forall s \in S \Longrightarrow g = e$$

**Definition 3.2.29 (Free group action)**
*A group action $G$ on $S$ is free if*

$$g \neq e \Longrightarrow gs \neq s \quad \forall s$$

**Definition 3.2.30 (Orbit/Stabilizer)**
*Let $G$ be a group with an action on $S$ and $s \in S$. Define the stabilizer subgroup*

© David Rufino CC BY-NC-ND 4.0

$$G_s := \{g \in G \mid gs = s\}$$

*and the orbit*

$$Gs := \{gs \mid g \in G\}$$

**Proposition 3.2.31 ($S$ is disjoint union of orbits)**
*Let $G$ be a group with an action on $S$. Then the following is an equivalence relation*

$$s \sim t \iff gs = t \text{ some } g \in G$$

*and the equivalence classes are precisely the orbits of elements of $S$ under $G$. Further $S$ is the disjoint union of orbits.*

**Remark 3.2.32**
*An action is free if and only if $G_s = \{e\}$ for all $s \in S$.*

**Proposition 3.2.33 (Orbit-Stabilizer Theorem)**
*Let $G$ be a group with an action on $S$. Given an element $s \in S$ there is a natural bijection*

$$G/G_s \longrightarrow Gs$$

*between the cosets of $G_s$ and the orbit $Gs$. In particular when $G$ is finite*

$$\#G = \#(Gs) \times \#G_s$$

*and when the action is free*

$$\#G = \#(Gs)$$

### 3.2.3 Symmetric Group

**Definition 3.2.34 (Symmetric Group)**
*Let $S_n$ denote the set of permutations (bijections) of $\{1, ..., n\}$.*

*Permutations $\sigma, \tau \in S_n$ are called **disjoint** if the supports are disjoint. Note disjoint permutations commute.*

**Definition 3.2.35 (Cycle)**
*Let $i_1, ..., i_r \in J_n$ be an ordered $r$-tuple, the permutation which maps*

$$i_k \to \begin{cases} i_{k+1} & k < r \\ i_1 & k = r \end{cases}$$

*is denoted by $(i_1\, i_2 \ldots i_r)$ and called a **cycle**.*

*A cycle with two elements $(i\, j)$ is called a **transposition**. Finally an **adjacent transposition** is one of the form $(i\, i+1)$.*

**Proposition 3.2.36**
*Let $\sigma \in S_n$. Then $\sigma$ may be represented by*

*a) a product of disjoint cycles, which is unique up to permutation of cycles.*
*b) a product of transpositions, the number of which is unique modulo 2*
*c) There is a well-defined group homomorphism*

$$\varepsilon : S_n \to \{-1, 1\}$$

*such that*

$$\varepsilon((i\, j)) = -1$$

*and $\varepsilon(1) = 1$.*
*d) A cycle $\sigma$ of length $r$ satisfies $\varepsilon(\sigma) = (-1)^{r+1}$*

**Proposition 3.2.37 (Adjacent Transpositions Generate the Symmetric Group)**
*Every $\sigma \in S_n$ is a product of adjacent transpositions $(i\, i+1)$, $1 \le i < n$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Every permutation is a product of transpositions, so it suffices to write each transposition $(i\, j)$ with $i < j$ as a product of adjacent ones, which we do by induction on $j - i$. If $j - i = 1$ there is nothing to show. If $j - i > 1$ then

© David Rufino CC BY-NC-ND 4.0

$$(i\,j) = (j-1\,j)(i\,j-1)(j-1\,j)$$

since conjugating $(i\,j-1)$ by $(j-1\,j)$ replaces $j-1$ with $j$, and $(i\,j-1)$ is a product of adjacent transpositions by induction. ∎

### 3.2.4 Shuffle Permutations

**Definition 3.2.38**
*Let $p+q=n$ be positive integers. We define the subgroup of* ***shuffle permutations***

$$\mathrm{Sh}(p,q) \coloneqq \{\sigma \in S_n \mid \sigma(1) < ... < \sigma(p) \text{ and } \sigma(p+1) < ... < \sigma(p+q)\}$$

*Similarly if $p+q+r=n$ are positive integers define the subgroup*

$$\mathrm{Sh}(p,q,r) \coloneqq \{\sigma \in S_n \mid \sigma(1) < ... < \sigma(p) \text{ and } \sigma(p+1) < ... < \sigma(p+q) \text{ and } \sigma(p+q+1) < ... < \sigma(p+q+r)\}$$

*Let $u : [1,p] \to [1,n]$ and $v : [1,q] \to [1,n]$ be injective maps such that $u([1,p]) \cap v([1,q]) = \varnothing$. Define $u \star v \in S_n$ by*

$$(u \star v)(i) \coloneqq \begin{cases} u(i) & 1 \le i \le p \\ v(i-p) & p < i \le p+q \end{cases}$$

*If $u, v$ are order preserving then $u \star v \in \mathrm{Sh}(p,q)$. Every shuffle permutation is of this form.*

**Lemma 3.2.39**
*Let $X, Y \subset \mathbb{N}$ be a finite sets of order $n$. There exists a unique order isomorphism*

$$u : X \to Y$$

**Proposition 3.2.40**
*Let $p+q=n$ be positive integers. There is a bijection*

$$\mathrm{Sh}(p,q) \to \{u : [1,p] \to [1,n] \text{ order preserinvg}\}$$

$$\sigma \to \sigma|_{[1,p]}$$

*Proof.* We provide an explicit inverse. For given $u$ the set $Y \coloneqq [1,n] \setminus u([1,p])$. As $u$ is order preserving it is injective, whence $\#Y = n - \#u([1,p]) = n-p = q$. Therefore there is an order isomorphism $v : [1,q] \to Y$ and we may define the shuffle permutation $\sigma \coloneqq u \star v$. ∎

**Proposition 3.2.41**
*Let $p+q=n$ be positive integers. There is an injective group homomorphism*

$$\begin{aligned} S_p \times S_q &\to S_n \\ (\sigma_1, \sigma_2) &\to \sigma_1 \star (\sigma_2 + p) \end{aligned}$$

*The image is the set $\sigma \in S_n$ for which $\sigma([1,p]) = [1,p]$ (equivalently $\sigma([p+1,q]) = \sigma([p+1,q])$.*

**Proposition 3.2.42**
*Let $p+q=n$ be positive integers. There is a bijection*

$$\mathrm{Sh}(p,q) \to S_n / (S_p \times S_q)$$

*Furthermore $\#\,\mathrm{Sh}(p,q) = \frac{(p+q)!}{p!q!}$. By a similar argument $\#\,\mathrm{Sh}(p,q,r) = \frac{(p+q+r)!}{p!q!r!}$.*

*Proof.* Let $\sigma \in S_n$. Then there exists an order isomorphism $u : [1,p] \to \sigma([1,p])$ and $v : [1,q] \to \sigma([p+1,p+q])$. Define $\sigma_1(i) \coloneqq u^{-1}(\sigma(i))$ and $\sigma_2(j) \coloneqq v^{-1}(\sigma(j+p))$. Then we may verify that $\sigma_1 \in S_p$, $\sigma_2 \in S_q$ and

$$(u \star v) \circ (\sigma_1 \star (\sigma_2 + p)) = \sigma$$

so the map is surjective.

Suppose $\sigma, \sigma' \in \mathrm{Sh}(p,q)$ are such that

$$\sigma = \sigma' \circ (\sigma_1 \star (\sigma_2 + p))$$

for $\sigma_1 \in S_p$ and $\sigma_2 \in S_q$. If $\sigma_1 \neq e$ then there exists $1 \le i < j \le p$ such that $\sigma_1(j) < \sigma_1(i)$. Then by monotonicity of $\sigma'$ we have $\sigma(j) = \sigma'(\sigma_1(j)) < \sigma'(\sigma_1(i)) = \sigma(i)$, which contradicts monotonicity of $\sigma$. So we conclude $\sigma_1 = e$ and similarly $\sigma_2 = e$. Therefore $\sigma = \sigma'$, and the map is injective. By the Orbit-Stabilizer theorem we have

© David Rufino CC BY-NC-ND 4.0

$$\#\operatorname{Sh}(p,q) = \#S_n / \#(S_p \times S_q)$$

from which the result follows. ∎

**Definition 3.2.43 (Circulant Permutation)**
*Let $n$ be an integer, and write $\rho_n \in S_n$ for the permutation*

$$\{1,...,n\} \to \{2,3,...,n,n+1\}$$

*Note that $\varepsilon(\rho_n) = (-1)^{n+1}$.*

**Proposition 3.2.44 (Symmetry of Shuffling)**
*Let $p+q=n$ be positive integers. Then there is a bijection*

$$\begin{aligned} \operatorname{Sh}(p,q) &\to \operatorname{Sh}(q,p) \\ \sigma &\to \sigma \circ \rho^p_{p+q} \end{aligned}$$

*Note that $\varepsilon\left(\rho^p_{p+q}\right) = (-1)^{pq}$ by (3.2.43).*

**Proposition 3.2.45 (Associativity of Shuffling)**
*The following map is a bijection*

$$\begin{aligned} \pi : \operatorname{Sh}(p+q,r) \times \operatorname{Sh}(p,q) &\to \operatorname{Sh}(p,q,r) \\ (\sigma_1,\sigma_2) &\to \sigma_1 \circ (\sigma_2 \star (1_r + p + q)) \end{aligned}$$

*Similarly so is*

$$\begin{aligned} \pi' : \operatorname{Sh}(p,q+r) \times \operatorname{Sh}(q,r) &\to \operatorname{Sh}(p,q,r) \\ (\sigma_1,\sigma_2) &\to \sigma_1 \circ \left(1_p \star (\sigma_2 + p)\right) \end{aligned}$$

*Proof.* It is straightforward to verify that $\pi(\sigma_1,\sigma_2)$ is injective and satisfies the ordering properties. Therefore $\pi$ is well-defined and we claim it is surjective.

Given $\sigma \in \operatorname{Sh}(p,q,r)$ let $u : [1,p+q] \to \sigma([1,p+q])$, $v : [1,r] \to \sigma([p+q+1,p+q+r])$ be the unique order isomorphisms. Define $\sigma_1 = u \star v$ and $\sigma_2 := u^{-1} \circ \sigma|_{[1,p+q]}$. We may verify that $\sigma_1 \in \operatorname{Sh}(p+q,r)$ and $\sigma_2 \in \operatorname{Sh}(p,q)$, and furthermore that $\pi(\sigma_1,\sigma_2) = \sigma$. By counting $\pi$ is injective.

For the second case let $u : [1,p] \to \sigma([1,p])$, $v : [1,q+r] \to \sigma([p+1,p+q+r])$ be the unique order isomorphisms. Define $\sigma_1 = u \star v$ and $\sigma_2(j) := v^{-1}(\sigma(j+p))$. The verification follows as before. ∎

## 3.2.5 Totally Ordered Abelian Group

**Definition 3.2.46 (Ordered Abelian Group)**
*An abelian group $(G,+)$ together with a total order $\leq$ is an **ordered abelian group** if it satisfies*

$$x \leq y \implies x + z \leq y + z \quad \forall x,y,z \in G$$

*Define $G^+ := \{g \in G \mid (0 \leq g) \wedge (g \neq 0)\}$ and $G^- := \{g \in G \mid (g \leq 0) \wedge (g \neq 0)\}$.*

## 3.2.6 Sums in an Abelian Group

**Proposition 3.2.47 (Finite Sums in an Abelian Group)**
*Let $G$ be an abelian group. For every integer $n \geq 1$ there is a unique map $\psi_n : G^n \to G$ such that $\psi_1 = 1_G$,*

$$\psi_n(x_1,...,x_n) = x_1 + \psi_{n-1}(x_2,...,x_n) \quad (n \geq 2)$$

*and $\psi_n$ is invariant under permutations, that is $\psi_n\left(x_{\sigma(1)},...,x_{\sigma(n)}\right) = \psi_n(x_1,...,x_n)$ for every $\sigma \in S_n$. Moreover:*

*a) $\psi_{m+r}(x_1,...,x_m,y_1,...,y_r) = \psi_m(x_1,...,x_m) + \psi_r(y_1,...,y_r)$ for $m,r \geq 1$*
*b) $\psi_n(0,...,0) = 0$ and $\psi_{n+1}(x_1,...,x_n,0) = \psi_n(x_1,...,x_n)$ for $n \geq 1$*
*c) $\psi_n(x_1+y_1,...,x_n+y_n) = \psi_n(x_1,...,x_n) + \psi_n(y_1,...,y_n)$ for $n \geq 1$*
*d) if $\varphi : G \to H$ is a group homomorphism to an abelian group $H$ then $\psi_n(\varphi(x_1),...,\varphi(x_n)) = \varphi(\psi_n(x_1,...,x_n))$*
*e) $\psi_n(-x_1,...,-x_n) = -\psi_n(x_1,...,x_n)$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

The conditions $\psi_1 = 1_G$ and $\psi_n(x_1,...,x_n) = x_1 + \psi_{n-1}(x_2,...,x_n)$ determine $\psi_n$ by induction on $n$, which gives uniqueness and shows that there is exactly one family of maps satisfying them. It remains to show that each $\psi_n$

© David Rufino CC BY-NC-ND 4.0

so defined is invariant under permutations; we do this by induction on $n$, the cases $n = 1$ and $n = 2$ being clear ($\psi_2(x_1, x_2) = x_1 + x_2 = x_2 + x_1$ as $G$ is abelian). Since $S_n$ is generated by the adjacent transpositions ((3.2.37)) it suffices to check invariance under $(i\, i+1)$ for $1 \leq i < n$, and let $n \geq 3$ with $\psi_{n-1}$ invariant.

If $i \geq 2$ the transposition only exchanges two entries of $(x_2, ..., x_n)$, so $\psi_n$ is unchanged since $\psi_{n-1}$ is invariant. If $i = 1$ then

$$\begin{aligned}\psi_n(x_2, x_1, x_3, ..., x_n) &= x_2 + (x_1 + \psi_{n-2}(x_3, ..., x_n)) && \text{by the recursion}\\ &= x_1 + (x_2 + \psi_{n-2}(x_3, ..., x_n)) && \text{by associativity and commutativity in } G\\ &= \psi_n(x_1, x_2, x_3, ..., x_n) && \text{by the recursion}\end{aligned}$$

For a) we use induction on $m$. For $m = 1$ this is the recursion defining $\psi_{1+r}$, since $\psi_1 = 1_G$. For $m > 1$, by the recursion, the induction hypothesis and associativity,

$$\psi_{m+r}(x_1, ..., x_m, y_1, ..., y_r) = x_1 + (\psi_{m-1}(x_2, ..., x_m) + \psi_r(y_1, ..., y_r)) = \psi_m(x_1, ..., x_m) + \psi_r(y_1, ..., y_r)$$

Next, $\psi_n(0, ..., 0) = 0 + \psi_{n-1}(0, ..., 0) = 0$ by induction on $n$, and by invariance and the recursion $\psi_{n+1}(x_1, ..., x_n, 0) = \psi_{n+1}(0, x_1, ..., x_n) = 0 + \psi_n(x_1, ..., x_n)$, which proves b). The remaining properties are proved by induction on $n$, the case $n = 1$ being clear. For c), writing $x' = (x_2, ..., x_n)$ and $y' = (y_2, ..., y_n)$,

$$\psi_n(x + y) = (x_1 + y_1) + \psi_{n-1}(x' + y') = (x_1 + \psi_{n-1}(x')) + (y_1 + \psi_{n-1}(y')) = \psi_n(x) + \psi_n(y)$$

using the induction hypothesis and commutativity. For d), $\psi_n(\varphi(x)) = \varphi(x_1) + \psi_{n-1}(\varphi(x')) = \varphi(x_1) + \varphi(\psi_{n-1}(x')) = \varphi(\psi_n(x))$ since $\varphi$ is additive. Finally e) is d) for the homomorphism $x \mapsto -x$ of $G$. ∎

**Definition 3.2.48 (Sum of a Finite Family)**
*Let $G$ be an abelian group, written additively. For a family $(g_i)_{i \in I}$ in $G$ indexed by a finite set $I$ with $n \geq 1$ elements, define*

$$\sum_{i \in I} g_i := \psi_n\left(g_{e(1)}, ..., g_{e(n)}\right)$$

*for any enumeration of $I$, that is a bijection $e : \{1, ..., n\} \to I$, where $\psi_n$ is as in (3.2.47), and define the sum over the empty set to be 0. If $g_i = 0$ for all $i \in I$ then $\sum_{i \in I} g_i = 0$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

If $e, e' : \{1, ..., n\} \to I$ are two enumerations of $I$ then $\sigma := e^{-1} \circ e' \in S_n$ and $g_{e'(k)} = g_{e(\sigma(k))}$, so the sum is independent of the enumeration since $\psi_n$ is invariant under permutations by (3.2.47). If $g_i = 0$ for all $i \in I$ then every entry of $\psi_n\left(g_{e(1)}, ..., g_{e(n)}\right)$ is 0, so the sum is 0 since $\psi_n(0, ..., 0) = 0$ by the same proposition. ∎

**Proposition 3.2.49 (Sum over a Disjoint Union)**
*Let $G$ be an abelian group and $(g_i)_{i \in I \cup J}$ a family in $G$ indexed by the union of two disjoint finite sets $I$ and $J$. Then*

$$\sum_{i \in I \cup J} g_i = \sum_{i \in I} g_i + \sum_{i \in J} g_i$$

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

If $I$ or $J$ is empty this is clear, so let $|I| = m \geq 1$ and $|J| = r \geq 1$. Now let $e_I : \{1, ..., m\} \to I$ and $e_J : \{1, ..., r\} \to J$ be enumerations. Since $I$ and $J$ are disjoint, $e : \{1, ..., m + r\} \to I \cup J$ with $e(k) = e_I(k)$ for $k \leq m$ and $e(k) = e_J(k - m)$ for $k > m$ is an enumeration of $I \cup J$, and by (3.2.47) (the identity for $\psi_{m+r}$) and independence of the enumeration ((3.2.48)) $\sum_{i \in I \cup J} g_i = \psi_{m+r}\left(g_{e(1)}, ..., g_{e(m+r)}\right) = \psi_m\left(g_{e_I(1)}, ..., g_{e_I(m)}\right) + \psi_r\left(g_{e_J(1)}, ..., g_{e_J(r)}\right) = \sum_{i \in I} g_i + \sum_{i \in J} g_i$. ∎

**Definition 3.2.50 (Sum of an Almost Zero Family)**
*Let $G$ be an abelian group, written additively. Let $(g_i)_{i \in I}$ be a family in $G$ indexed by an arbitrary set $I$ for which all but finitely many $g_i$ are zero. Define*

$$\sum_{i \in I} g_i := \sum_{i \in J} g_i$$

*where $J \subseteq I$ is any finite subset containing every $i$ with $g_i \neq 0$, and the sum over the finite set $J$ is as in (3.2.48).*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Such a $J$ exists since all but finitely many $g_i$ vanish. If $J, J'$ are two finite subsets of $I$ each containing every $i$ with $g_i \neq 0$, then $g_i = 0$ for all $i \in (J \cup J') \setminus J$ and for all $i \in (J \cup J') \setminus J'$, so by (3.2.49) and (3.2.48) (a sum of zero

© David Rufino CC BY-NC-ND 4.0

terms is 0) $\sum_{i\in J\cup J'} g_i = \sum_{i\in J} g_i + \sum_{i\in J'\setminus J} g_i = \sum_{i\in J} g_i$, and likewise $\sum_{i\in J\cup J'} g_i = \sum_{i\in J'} g_i$, and the definition does not depend on the choice of $J$. ∎

**Proposition 3.2.51 (Sums of Almost Zero Families)**
*Let $G$ be an abelian group and $(g_i)_{i\in I}$, $(h_i)_{i\in I}$ families in $G$ indexed by the same set $I$, with all but finitely many terms zero. Then $(g_i + h_i)_{i\in I}$ and $(-g_i)_{i\in I}$ are again families with all but finitely many terms zero, and*

*a) $\sum_{i\in I}(g_i + h_i) = \sum_{i\in I} g_i + \sum_{i\in I} h_i$*
*b) $\sum_{i\in I}(-g_i) = -\sum_{i\in I} g_i$, and consequently $\sum_{i\in I}(g_i - h_i) = \sum_{i\in I} g_i - \sum_{i\in I} h_i$.*

*Moreover, if $\varphi : G \to H$ is a group homomorphism to an abelian group $H$ (for instance multiplication by an element $r$ of a ring $R$ on the additive group $G = R$), then $(\varphi(g_i))_{i\in I}$ again has all but finitely many terms zero, and*

*a) $\sum_{i\in I} \varphi(g_i) = \varphi\left(\sum_{i\in I} g_i\right)$.*

*Finally, if $I = \bigsqcup_{\lambda\in\Lambda} I_\lambda$ is a partition of $I$ into disjoint subsets then each $(g_i)_{i\in I_\lambda}$ and the family $\left(\sum_{i\in I_\lambda} g_i\right)_{\lambda\in\Lambda}$ again have all but finitely many terms zero, and*

$$\sum_{i\in I} g_i = \sum_{\lambda\in\Lambda}\sum_{i\in I_\lambda} g_i$$

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Let $J \subseteq I$ be a finite subset containing every $i$ with $g_i \neq 0$ or $h_i \neq 0$. For $i \notin J$ we have $g_i + h_i = 0$ and $-g_i = 0$, so both families have all but finitely many terms zero, and $J$ is a valid choice of finite subset in the definition of all of the sums involved. Therefore

$$\sum_{i\in I}(g_i + h_i) = \sum_{i\in J}(g_i + h_i) = \sum_{i\in J} g_i + \sum_{i\in J} h_i = \sum_{i\in I} g_i + \sum_{i\in I} h_i$$

$$\sum_{i\in I}(-g_i) = \sum_{i\in J}(-g_i) = -\sum_{i\in J} g_i = -\sum_{i\in I} g_i$$

using additivity and compatibility with negation of finite sums ((3.2.47)). Next, $g_i - h_i = g_i + (-h_i)$, so the last claim of b) follows from the two parts already shown. Finally for $i \notin J$ we have $\varphi(g_i) = \varphi(0) = 0$, so $(\varphi(g_i))_{i\in I}$ has all but finitely many terms zero and $J$ is again a valid choice of finite subset. Then

$$\sum_{i\in I} \varphi(g_i) = \sum_{i\in J} \varphi(g_i) = \varphi\left(\sum_{i\in J} g_i\right) = \varphi\left(\sum_{i\in I} g_i\right)$$

since finite sums commute with group homomorphisms ((3.2.47)). For the last claim, let $J \subseteq I$ be finite containing every $i$ with $g_i \neq 0$, and let $\Lambda_J := \{\lambda \in \Lambda \mid I_\lambda \cap J \neq \varnothing\}$, which is finite. For $\lambda \in \Lambda_J$ the set $J \cap I_\lambda$ is a valid choice for the subfamily, so $\sum_{i\in I_\lambda} g_i = \sum_{i\in J\cap I_\lambda} g_i$, while for $\lambda \notin \Lambda_J$ all terms of the subfamily are zero and $\sum_{i\in I_\lambda} g_i = 0$. So $\left(\sum_{i\in I_\lambda} g_i\right)_{\lambda\in\Lambda}$ is almost zero with $\Lambda_J$ a valid choice, and since $J$ is the disjoint union of the finitely many $J \cap I_\lambda$, $\lambda \in \Lambda_J$, induction on $|\Lambda_J|$ using (3.2.49) gives

$$\sum_{i\in I} g_i = \sum_{i\in J} g_i = \sum_{\lambda\in\Lambda_J}\sum_{i\in J\cap I_\lambda} g_i = \sum_{\lambda\in\Lambda}\sum_{i\in I_\lambda} g_i$$

∎

**Lemma 3.2.52 (Reindexing and Iterated Sums)**
*Let $G$ be an abelian group.*

*a) If $(g_i)_{i\in I}$ has all but finitely many terms zero and $\sigma : I' \to I$ is a bijection, then $\left(g_{\sigma(i')}\right)_{i'\in I'}$ has all but finitely many terms zero and $\sum_{i'\in I'} g_{\sigma(i')} = \sum_{i\in I} g_i$.*
*b) If $\left(g_{(i,k)}\right)_{(i,k)\in I\times K}$ has all but finitely many terms zero then so do each row $\left(g_{(i,k)}\right)_{i\in I}$ and $\left(\sum_{i\in I} g_{(i,k)}\right)_{k\in K}$, and*

$$\sum_{(i,k)\in I\times K} g_{(i,k)} = \sum_{k\in K}\sum_{i\in I} g_{(i,k)}$$

*and likewise with the roles of $i$ and $k$ exchanged.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

© David Rufino CC BY-NC-ND 4.0

a) Let $J \subseteq I$ be finite containing every $i$ with $g_i \neq 0$. Then $\sigma^{-1}(J)$ is finite and contains every $i'$ with $g_{\sigma(i')} \neq 0$, so the reindexed family has all but finitely many terms zero, and

$$\sum_{i' \in I'} g_{\sigma(i')} = \sum_{i' \in \sigma^{-1}(J)} g_{\sigma(i')} = \sum_{i \in J} g_i = \sum_{i \in I} g_i$$

since $\sigma$ restricts to a bijection $\sigma^{-1}(J) \to J$ and finite sums do not depend on the ordering of the terms.

b) Let $S \subseteq I \times K$ be finite containing every $(i,k)$ with $g_{(i,k)} \neq 0$, and let $I_0 \subseteq I$, $K_0 \subseteq K$ be its (finite) images under the two projections. Each row $\left(g_{(i,k)}\right)_{i \in I}$ has all its nonzero terms in $I_0$, and for $k \notin K_0$ it is identically zero. Hence $\sum_{i \in I} g_{(i,k)} = \sum_{i \in I_0} g_{(i,k)}$, which is 0 for $k \notin K_0$, so $\left(\sum_{i \in I} g_{(i,k)}\right)_{k \in K}$ has all but finitely many terms zero and $K_0$ is a valid choice of finite subset for its sum. Therefore

$$\sum_{k \in K} \sum_{i \in I} g_{(i,k)} = \sum_{k \in K_0} \sum_{i \in I_0} g_{(i,k)} = \sum_{(i,k) \in I_0 \times K_0} g_{(i,k)} = \sum_{(i,k) \in I \times K} g_{(i,k)}$$

since $I_0 \times K_0$ is a finite set containing every $(i,k)$ with $g_{(i,k)} \neq 0$. The other order is identical. ■

© David Rufino CC BY-NC-ND 4.0

# 3.3 Rings and Modules

## 3.3.1 Commutative Rings

**Definition 3.3.1 (Ring)**
*A ring consists of a triple $(A, +, \cdot)$ where $A$ is a set and $+$ and $\cdot$ are laws of composition ("additive" and "multiplicative" respectively) such that the following holds*

- $(A, +)$ *is an* ***abelian group****, whose identity element we refer to as* $0_A$*.*
- $(A, \cdot)$ *is a* ***monoid****, whose identity element we refer to as* $1_A$
- $+$ *and* $\cdot$ *satisfy the* ***distributive property****, that is for all* $x, y, z \in A$

$$x \cdot (y + z) = x \cdot y + x \cdot z$$

$$(x + y) \cdot z = x \cdot z + y \cdot z$$

*For $x \in A$ we write the additive inverse as $-x$, and abbreviate multiplication $x \cdot y =: xy$.*

*We say that $A$ is a* ***zero-ring*** *(or trivial) if $0_A = 1_A \iff A = \{0\}$.*

*$A$ is* ***commutative*** *if in addition $xy = yx$ i.e. $(A, \cdot)$ is abelian.*

**Example 3.3.2**
*The set of integers $\mathbb{Z}$ (see the Integers subsection) is the canonical example of a ring with operations of addition and multiplication.*

**Definition 3.3.3 (Subring)**
*A* ***subring*** *of a ring $A$ is a subset $B$ such that*

- $0_A, 1_A \in B$
- $x \in B \implies -x \in B$
- $x, y \in B \implies x + y \in B$
- $x, y \in B \implies x \cdot y \in B$

*Then $(B, +|_{B\times B}, \cdot\,|_{B\times B})$ is a ring.*

**Definition 3.3.4 (Multiplicative set)**
*A subset $S \subset A$ is said to be* ***multiplicative*** *if*

- $1 \in S$
- $x, y \in S \implies xy \in S$

*Further it is said to be* ***saturated*** *if in addition*

$$x, y \in S \iff xy \in S$$

**Definition 3.3.5 (Integral Domain)**
*A commutative ring $A$ is said to be an* ***integral domain*** *if it is not a zero-ring and it is* cancellative*, that is*

$$ab = ac, a \neq 0 \implies b = c.$$

**Definition 3.3.6 (Reduced)**
*A commutative ring $A$ is said to be* ***reduced*** *if for all $a \in A$*

$$a^n = 0 \implies a = 0$$

**Definition 3.3.7 (Unit / Group of Units)**
*An element $0 \neq a$ of a ring $A$ is called a* ***unit*** *if it has a two-sided multiplicative inverse.*

*For $A$ not a zero-ring, the set of units $A^\star$ forms a group under multiplication, called the* ***group of units****.*

**Definition 3.3.8 (Field)**
*A* ***field*** *$K$ is a commutative non-zero ring such that every non-zero element is a unit, so that $K^\star$ is a group under multiplication and $K^\star = K \setminus \{0\}$.*

**Proposition 3.3.9**
*Every subring of a field $K$ is an integral domain.*

© David Rufino CC BY-NC-ND 4.0

*Proof.* Suppose $A \subset K$ is a subring. Suppose that $a, b \in A$ such that $ab = 0$. Suppose $a \neq 0$ then $a^{-1}ab = 0 \implies b = 0$. ∎

Note we have the implications

**Corollary 3.3.10**
*Let $A$ be a ring then we have the following implications*

$$\textbf{field} \implies \textbf{integral domain} \implies \textbf{reduced}$$

**Definition 3.3.11 (Ring homomorphism)**
*A **ring homomorphism** $\varphi : A \to B$ is a mapping which is both a multiplicative (monoid) and additive (group) homomorphism*

- $\varphi(0_A) = 0_B$
- $\varphi(1_B) = 1_B$
- $\varphi(x + y) = \varphi(x) + \varphi(y)$
- $\varphi(xy) = \varphi(x)\varphi(y)$

*The **kernel** of $\varphi$ is defined to be*

$$\ker(\varphi) = \{a \mid \varphi(a) = 0_B\}$$

**Definition 3.3.12 (Ideal)**
*A (two-sided) **ideal** $\mathfrak{a}$ of a ring $A$ is a subset of $A$ which is an additive subgroup and closed under multiplication by $A$ :*

- $0_A \in \mathfrak{a}$
- $x, y \in \mathfrak{a} \implies x + y \in \mathfrak{a}$
- $x \in \mathfrak{a} \implies -x \in \mathfrak{a}$
- $x \in \mathfrak{a}, a \in A \implies ax, xa \in \mathfrak{a}$

*$\mathfrak{a}$ is said to be **proper** if $\mathfrak{a} \neq A$.*

**Lemma 3.3.13 (Proper ideal)**
*An ideal $\mathfrak{a}$ is proper if and only if $1 \notin \mathfrak{a}$ if and only if $\mathfrak{a} \cap A^\star = \varnothing$.*

*Alternatively $\mathfrak{a} = A$ if and only if $1 \in \mathfrak{a}$ if and only if $\mathfrak{a} \cap A^\star \neq \varnothing$.*

**Proposition 3.3.14**
*Let $\varphi : A \to B$ be a ring homomorphism, then*

*a) The kernel $\ker(\varphi)$ is a two-sided ideal of $A$*
*b) The image $\varphi(A)$ is a subring of $B$*
*c) $\varphi$ is injective if and only if $\ker(\varphi) = \{0\}$*

**Proposition 3.3.15 (Krull's Theorem)**
*Let $A$ be a ring and $\mathfrak{a}$ a proper ideal. Then it is contained in a proper maximal ideal $\mathfrak{m}$.*

*In particular any non-unit $a \notin A^\star$ is contained in a maximal ideal.*

**Proposition 3.3.16 (Criteria to be a Field)**
*Let $A$ be a ring. Then the following are equivalent*

*a) $A$ is a field*
*b) $A$ is not the zero-ring and the only proper ideal is $\{0\}$.*

*Proof.* a) $\implies$ b). By definition $A$ is not the zero-ring. Let $\mathfrak{a}$ be a proper ideal. By (3.3.13) $\mathfrak{a} \cap A^\star = \varnothing \implies \mathfrak{a} = \{0\}$ as required.

b) $\implies$ a). Suppose $0 \neq a \in A$ then the ideal $Aa = (a)$ is either $\{0\}$ or $A$. However $a = 1 \cdot a \in (a)$ which implies $(a) \neq \{0\}$ and therefore $(a) = A$. In particular there exists $a^{-1} \in A$ such that $a^{-1}a = 1_A$ and $a$ is invertible. ∎

© David Rufino CC BY-NC-ND 4.0

### 3.3.2 Sums in a Ring

**Proposition 3.3.17 (Distributivity for Sums in a Ring)**
*Let $A$ be a ring, $r \in A$ and $(a_i)_{i\in I}$ a family in $A$ with all but finitely many terms zero. Then $(ra_i)_{i\in I}$ and $(a_i r)_{i\in I}$ again have all but finitely many terms zero, and*

$$r\sum_{i\in I} a_i = \sum_{i\in I} ra_i \qquad \left(\sum_{i\in I} a_i\right) r = \sum_{i\in I} a_i r$$

*If $(b_j)_{j\in J}$ is another such family then $(a_i b_j)_{(i,j)\in I\times J}$ has all but finitely many terms zero, and*

$$\left(\sum_{i\in I} a_i\right)\left(\sum_{j\in J} b_j\right) = \sum_{(i,j)\in I\times J} a_i b_j = \sum_{i\in I}\sum_{j\in J} a_i b_j$$

*and likewise with the order of summation exchanged.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

By the distributive property $x \mapsto rx$ and $x \mapsto xr$ are homomorphisms of the additive group of $A$, so the first claims are (3.2.51) for these homomorphisms. If $a_i b_j \neq 0$ then $a_i \neq 0$ and $b_j \neq 0$, so $a_i b_j$ vanishes outside a product of two finite sets. Write $b := \sum_{j\in J} b_j$. Then by the first part applied to $b$ and then to each $a_i$

$$\left(\sum_{i\in I} a_i\right) b = \sum_{i\in I} a_i b = \sum_{i\in I} a_i \sum_{j\in J} b_j = \sum_{i\in I}\sum_{j\in J} a_i b_j = \sum_{(i,j)\in I\times J} a_i b_j$$

where the last equality is (3.2.52). The other order of summation is identical. ∎

### 3.3.3 Modules I

**Definition 3.3.18 (Module)**
*Let $A$ be a ring. A **left $A$-module** $(M, +, \cdot)$ is an abelian group $(M, +)$ together with a "multiplication" operation*

$$\cdot : A \times M \to M$$

*which satisfies the following properties*

- $(a \times_A a') \cdot x = a \cdot (a' \cdot x)$
- $(a +_A a') \cdot x = a \cdot x + a' \cdot x$
- $a \cdot (x + y) = a \cdot x + a \cdot y$

*Similarly a **right $A$-module** $(M, +, \cdot)$ is an abelian group $(M, +)$ together with a multiplication operation*

$$\cdot : M \times A \to M$$

- $(a \times_A a') \cdot x = (x \cdot a') \times_A a$
- $(a +_A a') \cdot x = a \cdot x + a' \cdot x$
- $a \cdot (x + y) = a \cdot x + a \cdot y$

*Considering the first property $M$ is a left $A$-module iff it is a right $A^{op}$ module in the obvious way. Therefore in the usual case that $A$ is commutative the concepts coincide and we may speak simply of an **$A$-module**, though we almost always write the action on the left.*

*Similarly almost all results for left $A$-modules carry over unchanged for right $A$-modules. In this case we may simply by refer to $A$-modules rather than state the result for both cases separately.*

**Definition 3.3.19 (Submodule)**
*Let $(M, +, \cdot)$ be a left $A$-module. Then a subset $N \subset M$ is called an $A$-**submodule** if*

- *$N$ is a subgroup of $(M, +)$*
- *$m \in N, a \in A \implies am \in N$*

*Then $(N, +|_{N\times N}, \cdot|_{A\times N})$ is a left $A$-module. Similar definition applies for a right $A$-module.*

**Definition 3.3.20 (Module homomorphism)**
*Let $(M, +, \cdot), (N, +, \cdot)$ be left $A$-modules. A function $f : M \to N$ is an $A$-module homomorphism if*

© David Rufino CC BY-NC-ND 4.0

- *It is an (additive) group homomorphism* $(M,+) \to (N,+)$.
- *It is A-linear;* $\forall a \in A, m \in M \quad f(a \cdot m) = a \cdot f(m)$

*It may be verified that* $f$ *is bijective if and only if it's an isomorphism. In this way we have the following categories*

- $A$-**Mod** *the category of modules over a* commutative *ring* $A$
- ${}_A$**Mod** *the category of left* $A$*-modules*
- **Mod**$_A$ *the category if right* $A$*-modules*

**Definition 3.3.21 (Kernel and Image)**
*The* ***kernel*** *of a module homomorphism* $f$ *is given by*

$$\ker(f) := \{m \in M \mid f(m) = 0\}$$

*and the* ***image*** *is given by*

$$\mathrm{Im}(f) = f(M)$$

**Example 3.3.22 (Trivial Examples)**
*A ring* $A$ *is a left* $A$*-module over itself, denoted* $A_s$.

**Definition 3.3.23 (Restriction of Scalars)**
*Let* $\varphi : A \to B$ *a ring homomorphism and* $M$ *a* $B$*-module. Then we may consider* $M$ *as an* $A$*-module in the obvious way. Denote this by* $[M]_\varphi$.

**Proposition 3.3.24 (Submodules constitute a lattice)**
*Let* $M$ *be an* $A$*-module then the collection* $\mathrm{SubMod}(M)$ *of* $A$*-submodules form a complete sub-lattice of* $\mathcal{P}(M)$ *with meet and join given by*

$$\bigwedge_{i\in I} N_i = \bigcap_{i\in I} N_i$$

*and (the* ***internal sum****)*

$$\bigvee_{i\in I} N_i = \bigcap_{N_i \subseteq N \leq M} N =: \sum_{i\in I} N_i = \left\{ \sum_{j\in J} n_j \;\middle|\; n_j \in N_j \quad \#J < \infty \right\}$$

*Moreover it is the image of the closure operator* $\langle - \rangle : \mathcal{P}(M) \to \mathcal{P}(M)$ *given by*

$$\langle X \rangle = \bigcap_{X \subseteq N} N = \left\{ \sum_j a_j x_j \;\middle|\; x_j \in X \right\}$$

*Proof.* The $A$-submodules of $M$ naturally form a Moore family of subsets of $M$. By (2.1.41) they form a complete sub-lattice with the given form of meet and join. Furthermore it is the image of the given closure operator. The only non-trivial statement is the explicit form of $\sum_{i\in I} N_i$ TODO. ∎

**Lemma 3.3.25**
*Let* $M$ *be a module. Then*

*a)* $\langle \bigcup_{i\in I} X_i \rangle = \sum_{i\in I} \langle X_i \rangle$
*b)* $\langle \bigcup_{i\in I} N_i \rangle = \sum_{i\in I} N_i$
*c)* $N_1 \subseteq N_2 \implies N_1 + N_2 = N_2$

*Proof.*
a) This follows from (2.1.44) applied to the closure operator $\langle - \rangle$
b) This follows from a) because $N_i = \langle N_i \rangle$
c) This follows from b) because $N_1 \cup N_2 = N_2$

∎

**Definition 3.3.26 (Hom Sets)**
*A* ***module homomorphism*** $\varphi : M \to N$ *is an additive group homomorphism which commutes with the* $A$ *action*

$$\varphi(am) = a\varphi(m) \quad \forall a \in A \, m \in M$$

*Denote the abelian group of* $A$*-module homomorphisms*

© David Rufino CC BY-NC-ND 4.0

$$\mathrm{Hom}_A(M,N)$$

*and the endomorphism ring*

$$\mathrm{End}_A(M) \coloneqq \mathrm{Hom}_A(M,M)$$

*When $A$ is commutative then these have natural $A$-module and $A$-algebra structures respectively.*

### 3.3.4 Operations on Ideals

For this section we assume $A$ is a commutative ring.

**Definition 3.3.27 (Product of ideal and module)**
*Let $M$ be an $A$-module and $\mathfrak{a} \triangleleft A$ an ideal. Define*

$$\mathfrak{a}M = \langle \mathfrak{a} \cdot M \rangle = \left\{ \sum_{i=1}^{n} a_i m_i \;\middle|\; a_i \in \mathfrak{a} \quad m_i \in M \right\}$$

**Proposition 3.3.28 (Lattice of Ideals)**
*Let $A$ be a ring and $\mathcal{I}(A)$ the set of ideals. Then $\mathcal{I}(A)$ forms a complete lattice ordered by inclusion with join and meets given by*

$$\bigwedge_{i\in I} \mathfrak{a}_i = \bigcap_{i\in I} \mathfrak{a}_i$$

*and*

$$\bigvee_{i\in I} \mathfrak{a}_i = \bigcap_{\mathfrak{a}_i \subseteq \mathfrak{a}} \mathfrak{a} =: \sum_i \mathfrak{a}_i \coloneqq \left\{ \sum_i a_i \;\middle|\; a_i \in \mathfrak{a}_i \right\}$$

*This induces a corresponding closure operator*

$$\langle - \rangle : \mathcal{P}(A) \to \mathcal{I}(A)$$

*given by*

$$\langle X \rangle \coloneqq \bigcap_{X \subseteq \mathfrak{a}} \mathfrak{a} = \left\{ \sum_j a_j x_j \;\middle|\; a_j \in A \quad x_j \in X \right\}$$

**Proposition 3.3.29**
*Let $A$ be a ring and $\mathfrak{a}_i$ a family of ideals. Then*

$$\langle \bigcup_{i\in I} \mathfrak{a}_i \rangle = \sum_{i\in I} \mathfrak{a}_i$$

**Definition 3.3.30 (Product of ideals)**
*The product of two ideals $\mathfrak{ab}$ is*

$$\mathfrak{ab} = \left\{ \sum_{i=1}^{n} a_i b_i \;\middle|\; a_i \in \mathfrak{a} \quad b_i \in \mathfrak{b} \right\}$$

*and is itself an ideal.*

**Definition 3.3.31 (Coprime)**
*We say two elements $x, y$ of a commutative ring $A$ are **co-prime** if $(x,y) = (1) \iff ax + by = 1$ for some $a, b \in A$*

*We say a family of ideals $\{\mathfrak{a}_i\}_{i\in I}$ are co-prime if $\sum_{i\in I} \mathfrak{a}_i = A$.*

**Definition 3.3.32 (Principal Ideal)**
*A **principal ideal** is an ideal generated by a single element*

$$(a) \coloneqq \langle \{a\} \rangle = Aa$$

**Lemma 3.3.33**
*A principal ideal $(a)$ is proper if and only if $a \notin A^\star$*

© David Rufino CC BY-NC-ND 4.0

**Definition 3.3.34 (Maximal Ideal)**
*An ideal $\mathfrak{m} \triangleleft A$ is **maximal** if it is both proper and not contained in another proper ideal.*

**Definition 3.3.35 (Prime Ideal)**
*An ideal $\mathfrak{p} \triangleleft A$ is **prime** if it is both proper and satisfies the following property*

$$xy \in \mathfrak{p} \implies x \in \mathfrak{p} \vee y \in \mathfrak{p}$$

**Definition 3.3.36 (Radical Ideal)**
*An ideal $\mathfrak{a} \triangleleft A$ is **radical** if it satisfies the following property*

$$x^n \in \mathfrak{a} \implies x \in \mathfrak{a}$$

**Proposition 3.3.37 (Maximal ideals exist)**
*Let $A$ be a ring and $\mathfrak{a} \triangleleft A$ a proper ideal. Then it is contained in some maximal ideal $\mathfrak{m}$.*

*In particular there always exists a maximal ideal by considering $\mathfrak{a} = (0)$.*

*Proof.* Simple application of Zorn's Lemma. ∎

**Proposition 3.3.38 (Properties of prime ideals)**
*Let $\mathfrak{p}$ be a prime ideal and $\mathfrak{a}$, $\mathfrak{b}$ be ideals then the following are equivalent*

*a)* $\mathfrak{a} \subseteq \mathfrak{p}$ or $\mathfrak{b} \subseteq \mathfrak{p}$
*b)* $\mathfrak{a} \cap \mathfrak{b} \subseteq \mathfrak{p}$
*c)* $\mathfrak{ab} \subseteq \mathfrak{p}$

*Proof.* a) $\implies$ b) Follows because $\mathfrak{a} \cap \mathfrak{b} \subseteq \mathfrak{a}$

b) $\implies$ c) Follows because $\mathfrak{ab} \subseteq \mathfrak{a} \cap \mathfrak{b}$

c) $\implies$ a) If $\mathfrak{a} \nsubseteq \mathfrak{p}$, then choose $a \in \mathfrak{a} \setminus \mathfrak{p}$. By hypothesis $a\mathfrak{b} \subseteq \mathfrak{p}$ and since $\mathfrak{p}$ is prime $\mathfrak{b} \subseteq \mathfrak{p}$. ∎

**Corollary 3.3.39 (Ideal version of primality)**
*Let $\mathfrak{p}$ be a proper ideal. Then $\mathfrak{p}$ is prime if and only if the following condition holds for all ideals $\mathfrak{a}$, $\mathfrak{b}$*

$$\mathfrak{ab} \subseteq \mathfrak{p} \implies \mathfrak{a} \subseteq \mathfrak{p} \text{ or } \mathfrak{b} \subseteq \mathfrak{p}$$

*In particular for all $k > 0$ we have*

$$\mathfrak{a} \subseteq \mathfrak{p} \iff \mathfrak{a}^k \subseteq \mathfrak{p}$$

*Proof.* One direction has been shown in (3.3.38). Conversely suppose $fg \in \mathfrak{p}$ then apply the condition to the ideals $(f)$ and $(g)$ we find $f \in \mathfrak{p}$ or $g \in \mathfrak{p}$. ∎

**Lemma 3.3.40 (Prime ideals are meet-prime)**
*Let $\mathfrak{p}$ be a prime ideal. Then*

$$\bigcap_{i=1}^{n} \mathfrak{a}_i \subseteq \mathfrak{p} \implies \mathfrak{a}_i \subseteq \mathfrak{p} \text{ some } i = 1...n$$

*in other words $\mathfrak{p}$ is meet-prime in the lattice of ideals.*

*Proof.* Suppose $\mathfrak{a}_i \nsubseteq \mathfrak{p}$ for all $i$ then there exists $x_i \in \mathfrak{a}_i \setminus \mathfrak{p}$. Then $x_1...x_n \in \bigcap_{i=1}^{n} \mathfrak{a}_i \subseteq \mathfrak{p}$ by hypothesis, so by primality $x_i \in \mathfrak{p}$ for some $i$, a contradiction. ∎

**Lemma 3.3.41 (Generate prime ideals)**
*Let $A$ be a ring, $S$ a multiplicative set and $\mathfrak{b} \triangleleft A$ such that $\mathfrak{b} \cap S = \varnothing$ then*

$$\mathcal{I} = \{\mathfrak{a} \mid \mathfrak{b} \subseteq \mathfrak{a} \quad \mathfrak{a} \cap S = \varnothing\}$$

*has a maximal element, which is prime.*

*Proof.* Since $\mathfrak{b} \in \mathcal{I}$ it is non-empty. By Zorn's Lemma it has a maximal element, $\mathfrak{p}$. We claim it is prime, for suppose $xy \in \mathfrak{p}$ and $x, y \notin \mathfrak{p}$. Then by maximality $\mathfrak{p} + (x)$ and $\mathfrak{p} + (y)$ intersect $S$. Therefore $S$ intersects $(\mathfrak{p} + (x))(\mathfrak{p} + (y)) \subseteq \mathfrak{p}$, a contradiction. ∎

**Definition 3.3.42 (Minimal prime)**
*Let $A$ be a ring and $\mathfrak{a} \triangleleft A$ a proper ideal. A prime ideal $\mathfrak{p}$ is a **minimal prime over** $\mathfrak{a}$ if it contains $\mathfrak{a}$, and every other such prime ideal contains $\mathfrak{p}$.*

© David Rufino CC BY-NC-ND 4.0

*We say it is simply a* ***minimal prime*** *if it is minimal over* $(0)$.

**Proposition 3.3.43 (Prime ideals are chain complete)**
*Let* $\{\mathfrak{p}_i\}_{i\in I}$ *be a* ***chain*** *of prime ideals, then* $\bigcap_i \mathfrak{p}_i$ *and* $\bigcup_i \mathfrak{p}_i$ *are prime ideals.*

*Proof.* By (3.3.37) $\bigcup_i \mathfrak{p}_i$ is an ideal, and it's easily verified to be prime. Clearly $\bigcap_i \mathfrak{p}_i$ is an ideal. Suppose $a, b \notin \bigcap_i \mathfrak{p}_i$ then $a \notin \mathfrak{p}_j$ and $b \notin \mathfrak{p}_k$ with $j \leq k$. Then $b \notin \mathfrak{p}_j$, and $ab \notin \mathfrak{p}_j$ by primality, whence $ab \notin \bigcap_i \mathfrak{p}_i$. ∎

**Corollary 3.3.44 (Minimal primes exist)**
*Let* $A$ *be a ring and* $\mathfrak{a} \lhd A$ *be a proper ideal contained in a prime ideal* $\mathfrak{p}$. *Then there exists a minimal prime over* $\mathfrak{a}$ *contained in* $\mathfrak{p}$.

*In particular there always exists a minimal prime over* $\mathfrak{a}$ *and every prime ideal contains a minimal prime ideal.*

*Proof.* We may use (3.3.43) together with Zorn's Lemma. ∎

**Proposition 3.3.45**
*Let* $A$ *be a ring. Then the set* $\operatorname{Rad}(A)$ *of radical ideals forms a complete sub-lattice of the lattice of ideals* $\mathcal{I}(A)$. *This induces a closure operator*

$$\sqrt{-} : \mathcal{I}(A) \to \operatorname{Rad}(A)$$

*given by*

$$\sqrt{\mathfrak{a}} \coloneqq \bigcap_{\mathfrak{a}\subseteq\mathfrak{r}} \mathfrak{r} = \{x \mid x^n \in \mathfrak{a} \quad n > 0\}$$

*The "join" is given by*

$$\bigvee_{i\in I} \mathfrak{a}_i = \sqrt{\sum_i \mathfrak{a}_i}$$

*In particular*

*a)* $\mathfrak{a} \subseteq \sqrt{\mathfrak{a}}$
*b)* $\mathfrak{a} \subseteq \mathfrak{b} \implies \sqrt{\mathfrak{a}} \subseteq \sqrt{\mathfrak{b}}$
*c)* $\sqrt{\sqrt{\mathfrak{a}}} = \sqrt{\mathfrak{a}}$

*Proof.* The set of radical ideals is closed under arbitrary intersections (which are meets in the lattice $\mathcal{I}(A)$). Therefore by (2.1.41) it forms a complete sub-lattice with meet given by intersection of ideals.

It also shows that $\sqrt{-}$ as defined is a closure operator with image $\operatorname{Rad}(A)$, which demonstrates the required properties.

Finally we just need to show that $I' \coloneqq \{x \mid x^n \in \mathfrak{a} \quad n > 0\}$ is equal to $\sqrt{\mathfrak{a}}$. Firstly it's an ideal for if $x, y \in I'$ then $x^n \in \mathfrak{a}$ and $y^m \in \mathfrak{a}$, so we may show that $(x+y)^{n+m} \in \mathfrak{a}$ whence $x + y \in I'$. Similarly $a \in A$ and $x \in I'$ implies $(ax)^n = a^n x^n \in I'$. It's radical for suppose $x^m \in I'$ then $x^{mn} = (x^m)^n \in \mathfrak{a}$ by definition whence $x \in I'$. As it contains $\mathfrak{a}$ we find that $\sqrt{\mathfrak{a}} \subseteq I'$. Let $\mathfrak{r}$ be another radical ideal containing $\mathfrak{a}$ then $x \in I' \implies x^n \in \mathfrak{a} \implies x^n \in \mathfrak{r} \implies x \in \mathfrak{r}$. Therefore the reverse inclusion follows. ∎

**Proposition 3.3.46 (Prime Nullstellensatz)**
*The radical of an ideal satisfies*

$$\sqrt{\mathfrak{a}} = \bigcap_{\mathfrak{a}\subseteq\mathfrak{p}:\mathfrak{p}\text{ prime}} \mathfrak{p}$$

*Further the intersection may be taken over all* minimal *primes over* $\mathfrak{a}$.

*Proof.* Suppose $x \in \sqrt{\mathfrak{a}}$ and $\mathfrak{p} \supseteq \mathfrak{a}$. Then $x^n \in \mathfrak{p} \implies x \in \mathfrak{p}$. Therefore $\sqrt{\mathfrak{a}} \subseteq \bigcap_{\mathfrak{a}\subseteq\mathfrak{p}} \mathfrak{p}$. Conversely suppose $x \notin \sqrt{\mathfrak{a}}$ then $S \coloneqq \{1, x, x^2, \ldots\}$ is a proper multiplicative set such that $S \cap \mathfrak{a} = \varnothing$. By (3.3.41) there is a prime ideal $\mathfrak{p}$ containing $\mathfrak{a}$ which does not intersect $S$. Therefore $x \notin$ RHS as required. ∎

**Proposition 3.3.47 (Properties of Radical Ideals)**
*Let* $\mathfrak{a}$, $\mathfrak{b}$ *be ideals then*

*a)* $\sqrt{\mathfrak{a}^k} = \sqrt{\mathfrak{a}}$ *for* $k > 0$
*b)* $\sqrt{\sum_i \mathfrak{a}_i} = \sqrt{\sum_i \sqrt{\mathfrak{a}_i}}$
*c)* $\sqrt{\mathfrak{a}} = A \iff \mathfrak{a} = A$
*d)* $\sum_i \mathfrak{a}_i = A \iff \sum_i \sqrt{\mathfrak{a}_i} = A$
*e)* $\sum_{i=1}^n \mathfrak{a}_i^{k_i} = A \iff \sum_{i=1}^n \mathfrak{a}_i = A \quad k_i > 0.$

© David Rufino CC BY-NC-ND 4.0

*Proof.*

a) This may be shown by direct calculation or combining (3.3.46) and (3.3.38).
b) This follows by applying (2.1.44) to the closure operator $\sqrt{-}$.
c) $\sqrt{\mathfrak{a}} = A \iff 1 \in \sqrt{\mathfrak{a}} \iff 1 \in \mathfrak{a} \iff \mathfrak{a} = A$
d) This follows from combining c) and b)
e) This follows from d), b) and a)

■

**Lemma 3.3.48 (Ideal Finitely Generated by Nilpotents is Nilpotent)**
*Let $A$ be a ring with ideals $\mathfrak{a}, \mathfrak{b}$ such that $\sqrt{\mathfrak{a}} \subseteq \mathfrak{b}$ and $\mathfrak{b}$ finitely generated. Then there exists an integer $n > 0$ such that $\mathfrak{b}^n \subseteq \mathfrak{a}$.*

**Definition 3.3.49 (Extended and contracted ideals)**
*Let $\varphi : A \to B$ be a homomorphism and $\mathfrak{a}$ (resp. $\mathfrak{b}$) be an ideal of $A$ (resp. $B$). Define the **contraction** (resp. **extension**) ideals as follows*

$$\mathfrak{b}^c := \varphi^{-1}(\mathfrak{b})$$
$$\mathfrak{a}^e := \varphi(\mathfrak{a})B := \langle \varphi(\mathfrak{a}) \rangle = \left\{ \sum_i b_i \varphi(a_i) \;\middle|\; a_i \in \mathfrak{a} \right\}$$

*An ideal is said to be **contracted** (resp. **extended**) if it is of the form $\mathfrak{b}^c$ (resp. $\mathfrak{a}^e$)*

**Proposition 3.3.50 (Operations on ideals)**
*Let $\varphi : A \to B$ a ring homomorphism and $\mathfrak{a} \lhd A$, $\mathfrak{b} \lhd B$ ideals then*

*a) $\mathfrak{b}^c \lhd A$ and $\mathfrak{a}^e \lhd B$*
*b) $\mathfrak{b}^c$ proper if and only if $\mathfrak{b}$ is proper*
*c) $\mathfrak{b}^{ce} \subseteq \mathfrak{b}$ and $\mathfrak{a} \subseteq \mathfrak{a}^{ec}$*
*d) $\mathfrak{a}^{ece} = \mathfrak{a}^e$ and $\mathfrak{b}^{cec} = \mathfrak{b}^c$*
*e) $\mathfrak{b}^{ce} = \mathfrak{b} \iff \mathfrak{b}$ is an extended ideal $\iff \mathfrak{b} \subseteq \mathfrak{b}^{ce}$*
*f) $\mathfrak{a}^{ec} = \mathfrak{a} \iff \mathfrak{a}$ is a contracted ideal $\iff \mathfrak{a}^{ec} \subseteq \mathfrak{a}$*
*g) $\sqrt{\mathfrak{b}^c} = \left(\sqrt{\mathfrak{b}}\right)^c$*
*h) $\left(\sqrt{\mathfrak{b}^c}\right)^e \subseteq \sqrt{\mathfrak{b}}$ with equality when $\varphi$ is surjective*

*When $\varphi$ is surjective every ideal $\mathfrak{b} \lhd B$ is extended, and the contracted ideals are precisely the ideals containing $\ker(\varphi)$.*

*Proof.* We prove each in turn

a)-c) Straightforward

d) By the previous step $\mathfrak{b}^{ce} \subseteq \mathfrak{b} \implies (\mathfrak{b}^{ce})^c \subseteq \mathfrak{b}^c$, similarly $\mathfrak{b}^c \subseteq (\mathfrak{b}^c)^{ec}$. The other relation is similar.

e)-f) These follow from c) and d)

g) $x \in \left(\sqrt{\mathfrak{b}}\right)^c \iff \varphi(x) \in \sqrt{\mathfrak{b}} \iff \varphi(x)^n \in \mathfrak{b} \iff \varphi(x^n) \in \mathfrak{b} \iff x^n \in \mathfrak{b}^c \iff x \in \sqrt{\mathfrak{b}^c}$

h) By c) and g) we find $\left(\sqrt{\mathfrak{b}^c}\right)^e = \left(\sqrt{\mathfrak{b}}\right)^{ce} \subseteq \sqrt{\mathfrak{b}}$. We will show that when $\varphi$ is surjective every ideal is extended, in which case the equality follows from e).

Suppose that $\varphi$ is surjective. Then by e) we only need to show that $\mathfrak{b} \subseteq \mathfrak{b}^{ce}$ for every ideal $\mathfrak{b}$. Let $y \in \mathfrak{b}$ then $y = \varphi(x)$, whence $x \in \mathfrak{b}^c$ and $y \in \mathfrak{b}^{ce}$. ■

**Proposition 3.3.51**
*Let $\varphi : A \to B$ be a ring homomorphism and $\mathfrak{a}_1, ..., \mathfrak{a}_n$ ideals of $A$. Then*

$$\left(\sum_{i=1}^n \mathfrak{a}_i\right)^e = \sum_{i=1}^n \mathfrak{a}_i^e$$

*Similarly*

$$\left(\prod_{i=1}^n \mathfrak{a}_i\right)^e = \prod_{i=1}^n \mathfrak{a}_i^e$$

*Proof.* Evidently $\varphi(\mathfrak{a}_i) \subseteq \varphi\left(\sum_{i=1}^n \mathfrak{a}_i\right) \subseteq \left(\sum_{i=1}^n \mathfrak{a}_i\right)^e$ and therefore $\mathfrak{a}_i^e \subseteq \left(\sum_{i=1}^n \mathfrak{a}_i\right)^e \implies \sum_{i=1}^n \mathfrak{a}_i^e \subseteq \left(\sum_{i=1}^n \mathfrak{a}_i\right)^e$.

© David Rufino CC BY-NC-ND 4.0

Similarly $\varphi\left(\sum_{i=1}^n \mathfrak{a}_i\right) \subseteq \sum_{i=1}^n \varphi(\mathfrak{a}_i) \subseteq \sum_{i=1}^n \mathfrak{a}_i^e$ whence the reverse inclusion follows.

The second statement follows similarly. ■

**Corollary 3.3.52**
*Let $\varphi : A \to B$ be a ring homomorphism then extension and contraction constitute a monotone Galois connection*

$$\{\mathfrak{a} \lhd A\} \longleftrightarrow \{\mathfrak{b} \lhd B\}$$

*and therefore is order-preserving and satisfies the adjoint property*

$$\mathfrak{a} \subseteq \mathfrak{b}^c \iff \mathfrak{a}^e \subseteq \mathfrak{b}$$

*is satisfied.*

*Proof.* Extension and contraction satisfy conditions of (2.1.50) by (3.3.50) c) and d) ■

**Corollary 3.3.53**
*Let $\varphi : A \to B$ be a ring homomorphism then there is a order-preserving bijection between "contracted" and "extended ideals"*

$$\{\mathfrak{a} \lhd A \mid \mathfrak{a} \text{ contracted}\} \longleftrightarrow \{\mathfrak{b} \lhd B \mid \mathfrak{b} \text{ extended}\}$$

*which restricts to proper ideals.*

*Proof.* We've shown that $\mathfrak{a}$ (resp. $\mathfrak{b}$) is contracted (resp. extended) if and only if the given maps are mutually inverse. Note that $\mathfrak{b}$ is proper implies $\mathfrak{b}^c$ is proper. Furthermore $\mathfrak{b}^c$ proper implies $\mathfrak{b}^{ce} \subseteq \mathfrak{b}$ is proper. Therefore it restricts to proper ideals. ■

**Proposition 3.3.54 (Inverse image of maximal / prime ideals)**
*Let $\varphi : A \to B$ be a morphism then*

- *$\mathfrak{q} \lhd B$ prime $\implies \varphi^{-1}(\mathfrak{q})$ prime*
- *$\mathfrak{n} \lhd B$ maximal and $\varphi$ surjective $\implies \varphi^{-1}(\mathfrak{n})$ is maximal*

**Proposition 3.3.55**
*Consider maps $\varphi : A \to B$ and $\psi : B \to C$ and an ideal $\mathfrak{a} \lhd A$. Then extension of ideals is transitive, that is*

$$\psi(\varphi(\mathfrak{a})B)C = (\psi \circ \varphi)(\mathfrak{a})C$$

## 3.3.5 Quotient Rings

**Proposition 3.3.56 (Quotient Ring)**
*Let $(A, +, \cdot)$ be a ring and $\mathfrak{a}$ an ideal. As $\mathfrak{a}$ is an additive subgroup we may consider the quotient group $(A/\mathfrak{a}, +)$. For an element $a \in A$ write $a + \mathfrak{a}$ for the coset $[a]_\mathfrak{a} \in A/\mathfrak{a}$. There is a well-defined multiplicative law of composition*

$$\cdot : A/\mathfrak{a} \times A/\mathfrak{a} \to A/\mathfrak{a}$$

$$(a + \mathfrak{a}) \cdot (b + \mathfrak{a}) \to (a \cdot b + \mathfrak{a})$$

*which makes $(A/\mathfrak{a}, +, \cdot)$ into a ring. Further there is a canonical surjective ring homomorphism*

$$\pi : A \to A/\mathfrak{a}$$

*with the following properties*

- $\ker(\pi) = \mathfrak{a}$
- *Every morphism $\varphi : A \to B$ such that $\mathfrak{a} \subseteq \ker(\varphi)$, factors uniquely through $\pi$.*




- $\ker(\tilde{\varphi}) = \ker(\varphi)/\mathfrak{a}$
- *$\tilde{\varphi}$ is injective if and only if* $\ker(\varphi) = \mathfrak{a}$
- *$\tilde{\varphi}$ is surjective if and only if $\varphi$ is surjective*

© David Rufino CC BY-NC-ND 4.0

*For an ideal $\mathfrak{b} \supseteq \mathfrak{a}$ define the corresponding quotient ideal*

$$\mathfrak{b}/\mathfrak{a} := \{b + \mathfrak{a} \mid b \in \mathfrak{b}\} = \pi(\mathfrak{b})$$

*This induces a bijective, order-preserving correspondence of ideals*

$$\{\mathfrak{b}' \lhd A/\mathfrak{a}\} \underset{\pi(-)}{\overset{\pi^{-1}(-)}{\rightleftarrows}} \{\mathfrak{b} \lhd A \mid \mathfrak{a} \subseteq \mathfrak{b}\}$$

*under which maximal (resp. prime, radical) ideals of $A$ containing $\mathfrak{a}$ correspond to maximal (resp. prime, radical) ideals of $A/\mathfrak{a}$.*

**Corollary 3.3.57 (Isomorphism Theorem)**
*Let $\varphi : A \to B$ be a ring homomorphism. Then this induces a canonical isomorphism*

$$A/\ker(\varphi) \cong \varphi(A) \subset B$$

**Corollary 3.3.58 (Second Isomorphism Theorem)**
*Let $\mathfrak{b}$,$\mathfrak{a}$ be ideals then there is a unique morphism making the diagram commute*

$$\begin{array}{ccc} A & \xrightarrow{\pi} & A/\mathfrak{a} \\ {\scriptstyle\pi}\downarrow & \swarrow & \downarrow{\scriptstyle\pi} \\ A/(\mathfrak{a}+\mathfrak{b}) & \xrightarrow{\sim} & (A/\mathfrak{a})/((\mathfrak{a}+\mathfrak{b})/\mathfrak{a}) \end{array}$$

*which is in fact an isomorphism. If $\mathfrak{a} + \mathfrak{b} \subseteq \mathfrak{c}$ this restricts to an isomorphism of $A/\mathfrak{b}$-modules*

$$\begin{array}{ccc} \mathfrak{c} & \xrightarrow{\pi} & \mathfrak{c}/\mathfrak{a} \\ {\scriptstyle\pi}\downarrow & \swarrow & \downarrow{\scriptstyle\pi} \\ \mathfrak{c}/(\mathfrak{a}+\mathfrak{b}) & \xrightarrow{\sim} & (\mathfrak{c}/\mathfrak{a})/((\mathfrak{a}+\mathfrak{b})/\mathfrak{a}) \end{array}$$

**Proposition 3.3.59 (Criteria for Maximal, Prime and Reduced)**
*Let $\mathfrak{a} \lhd A$ then $\mathfrak{a}$ is*

- ***maximal** if and only if $A/\mathfrak{a}$ is a field*
- ***prime** if and only if $A/\mathfrak{a}$ is an integral domain*
- ***radical** if and only if $A/\mathfrak{a}$ is reduced*

*Proof.* Suppose $\mathfrak{a}$ is maximal then it is by definition proper so $A/\mathfrak{a}$ is not the zero-ring. By (3.3.56) then $A/\mathfrak{a}$ has no proper non-zero ideals and so by (3.3.16) is a field.

Conversely if $A/\mathfrak{a}$ is a field it is by definition not the zero ring, and so by (3.3.56) $\mathfrak{a}$ is proper. Furthermore by the same result $\mathfrak{a}$ is maximal.

Suppose $\mathfrak{a}$ is prime and $\overline{x} \cdot \overline{y} = 0$. Then by definition $\overline{x \cdot y} = 0 \implies x \cdot y \in \mathfrak{a} \implies x \in \mathfrak{a}$ or $y \in \mathfrak{a} \implies \overline{x} = 0$ or $\overline{y} = 0$. This shows that $A/\mathfrak{a}$ is an integral domain. The converse is similar. ∎

**Corollary 3.3.60**
*Let $\mathfrak{a} \lhd A$ be a proper ideal, then the following implications hold*

$$\textbf{maximal} \implies \textbf{prime} \implies \textbf{radical}$$

*Proof.* This follows by combining (3.3.10) and (3.3.59). ∎

**Corollary 3.3.61 (Field Morphisms are injective)**
*Let $\varphi : k \to B$ be a homomorphism from a field to a non-zero ring. Then $\varphi$ is injective.*

*Proof.* $\ker(\varphi)$ is an ideal. As $\varphi(1_k) = 1_B$ and $0_B \neq 1_B$ then $\ker(\varphi) \neq k$. Since the only ideals are $(0)$ and $k$ we see $\ker(\varphi) = \{0\}$ and $\varphi$ is injective. ∎

© David Rufino CC BY-NC-ND 4.0

### 3.3.6 Chinese Remainder Theorem

**Proposition 3.3.62 (Chinese Remainder Theorem)**
*Let $A$ be a commutative ring with ideals $\mathfrak{a}_1, ..., \mathfrak{a}_n$ which are pairwise coprime (i.e. $\mathfrak{a}_i + \mathfrak{a}_j = A$ for all $1 \leq i < j \leq n$). Then there is a canonical isomorphism*

$$A/\bigcap_{i=1}^{n} \mathfrak{a}_i \longrightarrow A/\mathfrak{a}_1 \times ... \times A/\mathfrak{a}_n$$

*Further we have the following identity*

$$\bigcap_{i=1}^{n} \mathfrak{a}_i = \prod_{i=1}^{n} \mathfrak{a}_i$$

*Proof.* Consider the canonical ring homomorphism

$$\varphi : A \longrightarrow A/\mathfrak{a}_1 \times ... \times A/\mathfrak{a}_n.$$

Trivially it has kernel $\bigcap_{i=1}^{n} \mathfrak{a}_i$. Therefore by (3.3.56) it is sufficient to show it is surjective. First we show that $e_1 := (1, 0, ..., 0)$ is in the image of $\varphi$. For we have $x_i + y_i = 1$ for $x_i \in \mathfrak{a}_1$, $y_i \in \mathfrak{a}_i$ and $1 < i \leq n$. Take $f_1 = y_2 ... y_n \in \bigcap_{i=2}^{n} \mathfrak{a}_i$; since $y_i = 1 - x_i$, $f_1 = \prod_{i=2}^{n}(1 - x_i) \in 1 + \mathfrak{a}_1$ (as each $x_i \in \mathfrak{a}_1$), so combined with $f_1 \in \bigcap_{i=2}^{n} \mathfrak{a}_i$, $\varphi(f_1) = e_1$ as required.

Similarly there exists $f_i$ such that $\varphi(f_i) = (0, ..., 1, ..., 0) =: e_i$. Finally for $\overline{x_1} \in A/\mathfrak{a}_1, ..., \overline{x_n} \in A/\mathfrak{a}_n$ then

$$\varphi(f_1 x_1 + ... + f_n x_n) = (\overline{x_1}, ..., \overline{x_n})$$

For the final statement we may assume by induction that

$$\bigcap_{i=1}^{n-1} \mathfrak{a}_i = \prod_{i=1}^{n-1} \mathfrak{a}_i$$

We have equations

$$x_i + y_i = 1 \quad x_i \in \mathfrak{a}_i, y_i \in \mathfrak{a}_n$$

Then

$$x_1 ... x_{n-1} = \prod_{i=1}^{n-1}(1 - y_i) \in 1 + \mathfrak{a}_n$$

so we may conclude

$$\bigcap_{i=1}^{n-1} \mathfrak{a}_i + \mathfrak{a}_n = A.$$

This then reduces to the case $n = 2$. Suppose $x \in \mathfrak{a} \cap \mathfrak{b}$, by assumption $1 = y + z$ for $y \in \mathfrak{a}$ and $z \in \mathfrak{b}$. Then $x = x(y + z) \in \mathfrak{ab}$, so $\mathfrak{a} \cap \mathfrak{b} \subset \mathfrak{ab}$. The reverse inclusion is obvious. ∎

**Proposition 3.3.63 (Chinese Remainder Theorem II)**
*Let $A$ be a commutative ring with $\mathfrak{a}_1, ..., \mathfrak{a}_n$ co-prime ideals. For any positive integers $k_1, ..., k_n$ there is a canonical isomorphism*

$$A/\bigcap_{i=1}^{n} \mathfrak{a}_i^{k_i} \longrightarrow A/\mathfrak{a}_1^{k_1} \times ... \times A/\mathfrak{a}_n^{k_n}$$

*and*

$$\bigcap_{i=1}^{n} \mathfrak{a}_i^{k_i} = \prod_{i=1}^{n} \mathfrak{a}_i^{k_i}$$

*Proof.* By (3.3.47) the ideals $\mathfrak{a}_1^{k_1}, ..., \mathfrak{a}_n^{k_n}$ are pairwise co-prime. Therefore we may apply (3.3.62). ∎

### 3.3.7 Irreducible and Reduced rings

We say an element $x$ is nilpotent if $x^n = 0$. By (3.3.45) these form an ideal.

**Definition 3.3.64 (Nilradical)**
*Define the **nilradical** to be the set (ideal) of nilpotents*

© David Rufino CC BY-NC-ND 4.0

$$N(A) := \sqrt{(0)} \overset{(3.3.46)}{=} \bigcap_{\mathfrak{p}} \mathfrak{p}$$

*Clearly $A$ is reduced if and only if $N(A) = \{0\}$.*

We also make the following definition

**Definition 3.3.65 (Irreducible)**
*Let $A$ be a ring. We say $A$ is **irreducible** if $N(A)$ is prime.*

The notion of irreducible ring is related to the notion of minimal primes

**Proposition 3.3.66**
*A ring $A$ is irreducible if and only if it has a unique minimal prime ideal. In this case it is equal to $N(A)$.*

*Proof.* First we note that every prime ideal contains $N(A)$. If $A$ is irreducible then by definition $N(A)$ is prime and it is therefore the unique minimal prime ideal.

Conversely if $\mathfrak{p}_0$ is the unique minimal prime ideal then by (3.3.44) it is contained in every prime ideal. Therefore (3.3.64) $N(A) = \bigcap_{\mathfrak{p}} \mathfrak{p} = \mathfrak{p}_0$ is prime and $A$ is irreducible. ∎

**Proposition 3.3.67 (Integral Domain $\iff$ Reduced and Irreducible)**
*Let $A$ be a ring. Then the following are equivalent*

*a) $A$ is an integral domain*
*b) $A$ is reduced and has a unique minimal prime*
*c) $(0)$ is prime*
*d) $A$ is reduced and irreducible.*

The following may be useful

**Proposition 3.3.68**
*Let $A$ be a ring then the sum of an invertible and nilpotent element is again invertible*

$$A^\star + N(A) \subseteq A^\star$$

*Proof.* For if

$$a = u + n = u(1 - (u^{-1})(-n))$$

with $n \in N(A)$ and $u \in A^\star$. Then way may reduce to the case $1 - n$ and observe that

$$(1-n)^{-1} = \sum_{i=0}^{\infty} n^i$$

which by assumption is a finite sum. ∎

## 3.3.8 Algebra over a Commutative Ring

For what follows let $A$ be a commutative ring.

**Definition 3.3.69 (Algebra (over a commutative ring))**
*An algebra over $A$ (or an $A$-algebra) is a pair $(i_B, B)$ where $B$ is a (not necessarily commutative) ring and $i_B : A \to B$ is a ring homomorphism.*

*We call $i_B$ the structural morphism and write $a \cdot b := i_B(a)b$*

*Morphisms of $A$-algebras are the ring homomorphisms $\varphi : B \to C$ such that $\varphi \circ i_B = i_C$. This then constitutes a category $\mathbf{Alg}_A$.*

*If $k$ is a field an algebra over $k$ is referred to as a $k$-algebra.*

**Definition 3.3.70 (Sub-algebra)**
*Let $(i_B, B)$ be an $A$-algebra. A sub-algebra $C$ is a subring $C$ of $B$ for which*

$$a \in A \quad c \in C \implies i_B(a)c \in C$$

**Example 3.3.71 (Algebra over commutatve sub-ring)**
*If $A \subset B$ is a commutative sub-ring, then $B$ is naturally a $A$-algebra.*

*The polynomial ring $A[X]$ is naturally an $A$-algebra*

© David Rufino CC BY-NC-ND 4.0

**Definition 3.3.72 (Algebra generated by a set)**
*Let $B$ be an $A$-algebra. The collection of $A$-subalgebras forms a Moore family. Therefore by (2.1.41) there is a canonical closure operator*

$$A[-] : \mathcal{P}(B) \to \mathrm{SubAlg}_A(B)$$

*which we denote by $A[S]$ for $S \subset B$. A more explicit characterization when $S$ is finite is given in the Polynomial Rings in Many Variables subsection. More generally we have*

$$A[S] = \bigcup_{S' \subset S | S' \text{ finite}} A[S']$$

**Proposition 3.3.73**
*Let $B = A[S]$ be an $A$-algebra and $\mathfrak{a}$ a sub-$A$-module of $B$. Then $\mathfrak{a}$ is an ideal if and only if*

$$s \in S \implies s\mathfrak{a} \subseteq \mathfrak{a}$$

*Proof.* One direction is obvious. Suppose the condition given holds, and define

$$B' := \{b \in B \mid b\mathfrak{a} \subseteq \mathfrak{a}\}$$

Then clearly $S \subseteq B'$. It's easy to show that $B'$ is a sub-$A$-algebra of $B$, so $B' = B$ and $\mathfrak{a}$ is an ideal. ∎

## 3.3.9 Finite-type Algebras

**Definition 3.3.74 (Finite algebra)**
*An $A$-algebra $B$ is* finite *if it is finite as an $A$-module.*

**Definition 3.3.75 (Finitely generated algebra)**
*An $A$-algebra $B$ is finitely generated (or of finite type) if there exists an integer $n \in \mathbb{N}$ and a surjection of $A$-algebras*

$$A[X_1, ..., X_n] \to B$$

*the images of $X_i$ are the generators.*

**Proposition 3.3.76**
*Let $C$ be a finitely-generated $B$ algebra and $B$ a finitely-generated $A$-algebra. Then $C$ is finitely-generated as an $A$-algebra.*

*Proof.* Suppose $C = B[c_1, ..., c_n]$. Then by definition $c_i = F_i(b_1, ..., b_m)$ for some $F_i \in A[X_1, ..., X_m]$. Then we may take as a generating set

$$b_1^{\lambda_1} ... b_m^{\lambda_m}$$

where $\lambda \in \mathbb{N}^m$ runs through all the exponents of non-zero coefficients of the $F_i$. ∎

## 3.3.10 Bimodules

For applications it is often useful to have multiple rings acting on a single module ("bimodule"). For example an $A$-algebra $B$ naturally has an action from both $A$ and $B$. It may also allow us to generalise results which would otherwise only hold in the commutative case.

**Definition 3.3.77 (Bimodule)**
*Let $A, B$ be rings. We say an abelian group $M$ is a $(A, B)$-**bimodule** if it is both a left $A$-module and a right $B$-module for which the two actions commute*

$$a \cdot (m \cdot b) = (a \cdot m) \cdot b \quad \forall a \in A, b \in B, m \in M$$

*We may denote this by ${}_AM_B$ in order to emphasise the actions. If homomorphisms are defined in the obvious way then these constitute a category ${}_A\mathbf{Mod}_B$. Recall that every abelian group is automatically both a left and right $\mathbb{Z}$-module with the following action for $N \in \mathbb{Z}$*

$$N \cdot m = m \cdot N := \begin{cases} \underbrace{m + ... + m}_{N \text{ times}} & N \geq 0 \\ -(m \cdot (-N)) & N < 0 \end{cases}$$

*Therefore we have the following generalisations*

- *A right $B$-module is precisely a $(\mathbb{Z}, B)$-bimodule*

© David Rufino CC BY-NC-ND 4.0

- *A left $A$-module is precisely a $(A,\mathbb{Z})$-bimodule*
- *When $A$ is commutative an $A$-module has a well-defined $(A,A)$-bimodule structure by defining*

$$a\cdot m\cdot a' := a'a\cdot m = aa'\cdot m$$

  *in other words $A$-**Mod** is (equivalent to) a full subcategory of ${}_A\mathbf{Mod}_A$.*
- *A ring $A$ has an obvious $(A,A)$-bimodule structure by associativity of the multiplication operation*

*We will understand by the notation $N_B$ that $N$ is a $(\mathbb{Z},B)$-bimodule and by ${}\{\}_A M$ that $M$ is a $(A,\mathbb{Z})$-bimodule.*

We generalize slightly the consideration of the third bullet point

**Proposition 3.3.78 ($(A,A)$-Bimodule $\equiv$ $A$-module)**
*Let $\varphi : A\to B$ be a homomorphism of commutative rings and $Z$ a $B$-module. Then $Z$ is naturally a $(B,A)$-module with the following action*

$$b\cdot m\cdot a := \varphi(a)b\cdot m \qquad (\star)\quad \forall a\in A, b\in B, m\in Z$$

*Suppose $M$ is another $(B,A)$-bimodule satisfying $(\star)$ then we may identify the $(B,A)$-bimodule homomorphisms and the left $B$-module homomorphisms.*

$$\mathrm{Hom}(M,\,{}_BZ_A)\xrightarrow{\sim}\mathrm{Hom}({}_BM,\,{}_BZ)$$

*In particular when $\varphi = 1_A$ and $B=A$ we may identify $(A,A)$-bimodules and $A$-modules, as well as the corresponding homomorphisms. Explicitly for $M,Z$ $A$-modules these have well-defined $(A,A)$-bimodule structures and there is a bijection*

$$\mathrm{Hom}({}_AM_A,\,{}_AZ_A)\xrightarrow{\sim}\mathrm{Hom}(M,Z)$$

*Proof.* We may show that the right $A$-module structure on $Z$ is well-defined because $B$ is commutative and using the associativity of the $B$-module action. Similarly the associativity of the $B$-module action ensures that the actions commute and so form a $(B,A)$-bimodule structure.

Suppose $\theta : M\to Z$ is a (left) $B$-module homomorphism. Then

$$\theta(m\cdot a)=\theta(\varphi(a)\cdot m)=\varphi(a)\theta(m)=\theta(m)\cdot a$$

so it is $(B,A)$-bilinear as required. The converse is clear, so we see that the sets are equal as subsets of the set of abelian group homomorphisms.

The case $\varphi = 1_A$ is immediate. ∎

We generalize the notion of "Hom-Set"

**Definition 3.3.79 (Hom Functors)**
*Let $A,B,C$ be arbitrary rings. Then we have an enriched hom functor for ${}_A\mathbf{Mod}_B$*

$$\mathrm{Hom} : {}_A\mathbf{Mod}_B^{op}\times{}_A\mathbf{Mod}_B\to\mathbf{AbGrp}$$

*In order to generalise the Tensor-Hom adjunction we consider the following functors ("right-linear" and "left-linear" respectively)*

$$\mathrm{RHom} : {}_B\mathbf{Mod}_A^{op}\times{}_C\mathbf{Mod}_A\to{}_C\mathbf{Mod}_B$$
$$\mathrm{LHom} : {}_A\mathbf{Mod}_B^{op}\times{}_A\mathbf{Mod}_C\to{}_B\mathbf{Mod}_C$$

*For example if $\psi\in\mathrm{RHom}({}_BM_C,\,{}_AN_C)$ then the $(A,B)$-bimodule action is defined to be*

$$(a\psi b)(m) := a\psi(bm)$$

*and similarly if $\varphi\in\mathrm{LHom}({}_AM_B,\,{}_AN_C)$ then*

$$(b\varphi c)(m) := \varphi(mb)c$$

*which we may verify satisfies the axioms of a bimodule. In order to standardize notation we may write $\mathrm{Hom}_A$ in place of* RHom *and* LHom.

*If $B=C=\mathbb{Z}$ then we recover the simple case (3.3.26).*

© David Rufino CC BY-NC-ND 4.0

*If $A$ is a commutative ring then we've observed that $A$-**Mod** is a full subcategory of ${}_A\mathbf{Mod}_A$ and the functors* RHom, LHom *become equal to the simple case (3.3.26) when restricted to this subcategory.*

### 3.3.11 Module Direct Product and Sum

**Definition 3.3.80 (External Direct Product / Sum)**
*Let $A$ be a ring and $\{M_i\}_{i\in I}$ a family of $A$-modules. Define the **external direct product** as the set of ordered tuples indexed over $I$*

$$\prod_{i\in I} M_i \coloneqq \left\{ (m_i)_{i\in I} \;\middle|\; m_i \in M_i \right\}$$

*with the obvious module operations. The **external direct sum** is the subset of tuples for which all but finitely many elements are zero. We denote this as follows*

$$\bigoplus_{i\in I} M_i$$

*Clearly when $I$ is finite then these are equal.*

**Proposition 3.3.81 (Categorical Product)**
*Let $\{M_i\}_{i\in I}$ be a family of left $A$-modules and consider the family of projections*

$$\pi_i : \prod_{i\in I} M_i \to M_i$$

*For $Z$ an $(A,C)$-bimodule there is a natural isomorphism of left $C$-modules*

$$\begin{aligned} \mathrm{Hom}_A\left(Z, \prod_{i\in I} M_i\right) &\xrightarrow{\sim} \prod_{i\in I} \mathrm{Hom}_A(Z, M_i) \\ \theta &\to (\pi_i \circ \theta)_{i\in I} \end{aligned}$$

*This in particular includes the case $A = C$ is commutative.*

**Proposition 3.3.82 (Categorical Coproduct)**
*Let $\{M_i\}_{i\in I}$ be a family of left $A$-modules and consider the family of inclusions*

$$u_i : M_i \to \bigoplus_{i\in I} M_i$$

*For $Z$ a $(A,C)$-bimodule there is a natural isomorphism of right $C$-modules*

$$\begin{aligned} \mathrm{Hom}_A\left(\bigoplus_{i\in I} M_i, Z\right) &\xrightarrow{\sim} \prod_{i\in I} \mathrm{Hom}_A(M_i, Z) \\ \theta &\to (\theta \circ u_i)_{i\in I} \end{aligned}$$

*This in particular includes the case $A = C$ is commutative.*

**Corollary 3.3.83**
*Let $(M_i)_{i\in I}$ and $(N_j)_{j\in J}$ be families of left $A$-modules. Then there is an isomorphism of abelian groups*

$$\begin{aligned} \mathrm{Hom}_A\left(\bigoplus_{i\in I} M_i, \prod_{j\in J} N_j\right) &\cong \prod_{(i,j)\in I\times J} \mathrm{Hom}_A(M_i, N_j) \\ \psi &\to (\pi_j \circ \psi \circ u_i)_{(i,j)} \end{aligned}$$

*where $\pi_j : \prod_{j\in J} N_j \to N_j$ and $u_i : M_i \to \bigoplus_{i\in I} M_i$ are the canonical projections and injections respectively.*

*These maps are $A$-linear if $A$ is commutative.*

**Definition 3.3.84 (Free Module)**
*An $A$-module $M$ is*

- ***free** if it is isomorphic to*

© David Rufino CC BY-NC-ND 4.0

$$\bigoplus_{i\in A} A =: A^{(I)}$$

*for some indexing set $I$*

- ***finite free*** *if it is free with respect to a* finite *indexing set $I$*

*Under the isomorphism $M \to \bigoplus_{i\in I} A$ the set of elements $\{m_i\}_{i\in I}$ corresponding to the standard basis vectors $e_i$ is called a* ***basis*** *for $M$.*

*We say that $M$ has* ***rank*** *$n$ if there exists a basis of order $n$. NB rank is not necessarily unique.*

**Proposition 3.3.85**
*Let $M$ be an $(A, C)$-bimodule then there is a canonical isomorphism of right $C$-modules*

$$\begin{aligned} \mathrm{Hom}_A(A, M) &\cong M \\ \theta &\to \theta(1_A) \end{aligned}$$

*This in particular includes the case $A = C$ is commutative.*

**Proposition 3.3.86**
*Let $M$ be a free left $A$-module with basis $(m_i)_{i\in I}$ and $N$ an $(A, C)$-bimodule then there is an isomorphism of right $C$-modules*

$$\begin{aligned} \mathrm{Hom}_A(M, N) &\cong \prod_{i\in I} N \\ \theta &\to (\theta(m_i))_{i\in I} \end{aligned}$$

*This in particular includes the case $A = C$ is commutative.*

## 3.3.12 Free Modules

**Definition 3.3.87 (Faithful Module)**
*We say an $A$-module $M$ is* ***faithful*** *if*

$$am = 0 \quad \forall m \in M \implies a = 0$$

**Definition 3.3.88 (Linearly Independent, Spanning and Basis)**
*Let $M$ be an $A$-module and $S \subset M$ a set. We say $S$ is*

- ***spanning*** *if $\langle S \rangle = M$*
- ***linearly independent*** *if for every finite subset $\{s_1, ..., s_n\} \subseteq S$ with $s_i$ distinct we have*

$$\sum_{i=1}^{n} a_i s_i = 0 \implies a_i = 0 \quad 1 \leq i \leq n$$

- *a* ***basis*** *if it is both spanning and linearly independent*

**Definition 3.3.89 (Finite Module)**
*An $A$-module $M$ is* ***finite*** *if there exists a finite spanning set.*

**Definition 3.3.90 (Minimal spanning set)**
*Let $M$ be an $A$-module. Then $S \subset M$ is a* ***minimal spanning set*** *if it generates $M$ and no proper subset does so.*

**Definition 3.3.91 (Free Module)**
*Let $M$ be an $A$-module. We say that $M$ is a* ***free module*** *over $A$ if it has a basis.*

**Proposition 3.3.92 (Free $A$-module is an external sum of $A$)**
*An $A$-module $M$ is free if and only if it is isomorphic to $\bigoplus_{i\in I} A$ for some $I$. The isomorphism is given by*

$$\sum_{i\in I} a_i m_i \to (a_i)_{i\in I}$$

**Definition 3.3.93 (Internal Direct Sum Module)**
*An $A$-module $M$ is an* ***internal direct sum*** *of submodules $\{M_i\}_{i\in I}$ the canonical mapping of the external direct sum*

$$\bigoplus_{i\in I} M_i \to M$$

© David Rufino CC BY-NC-ND 4.0

*is an isomorphism.*

**Proposition 3.3.94**
*Let $M$ be an $A$-module and $\{M_i\}_{i\in I}$ a family of submodules. Then the following are equivalent*

*a) $\sum_{i\in I} M_i$ is the internal direct sum of the family $\{M_i\}_{i\in I}$*
*b) The relation $\sum_{i\in I} m_i = 0$ implies $m_i = 0$ for all $i \in I$*
*c) For any $i \in I$ we have $M_i \cap \left(\sum_{k\neq i} M_k\right) = \{0\}$*

**Proposition 3.3.95**
*We say that $M_1, M_2$ are **supplementary** submodules of $M$ if $M$ is the internal direct sum of $M_1$ and $M_2$.*

*We say that $M_1$ is a **direct factor** of $M$ if it is supplementary to another submodule.*

## 3.3.13 Exact Sequences

**Definition 3.3.96 (Vector space)**
*If $k$ is a field and $V$ a $k$-module, then we say $V$ is a **vector space** over $k$.*

**Remark 3.3.97**
*We will see that every vector space is free and every $k$-submodule is a direct factor.*

**Proposition 3.3.98 (Kernel)**
*Let $\varphi : M \to N$ be an $A$-module homomorphism, then the **kernel** of $\varphi$*

$$\ker(\varphi) \coloneqq \{m \in M \mid \varphi(m) = 0\}$$

*is an $A$-submodule of $M$. Observe $\varphi$ is injective iff $\ker(\varphi) = 0$.*

**Proposition 3.3.99 (Image)**
*Let $\varphi : M \to N$ be an $A$-module homomorphism then the image*

$$\mathrm{Im}(\varphi) = \{\varphi(m) \mid m \in M\}$$

*is an $A$-submodule of $N$.*

**Definition 3.3.100 (Quotient Module)**
*Let $N \subseteq M$ be an $A$-submodule then define the **quotient module** $M/N$ to be the quotient group with an action of $A$ given by*

$$a(m + N) = (am + N)$$

*When $N \subseteq P \subseteq M$ is a sequence of submodules then define the $A$-submodule $P/N$ of $M/N$ by*

$$P/N \coloneqq \{p + N \mid p \in P\}$$

**Proposition 3.3.101 (Quotient Module Properties)**
*Let $N \subseteq M$ be an $A$-submodule then there is a canonical surjective morphism*

$$\pi : M \to M/N$$

*with the following properties*

*a) $\pi(m) = m + N$*
*b) $\ker(\pi) = N$*
*c) Every homomorphism $\psi : M \to P$ such that $N \subseteq \ker(\psi)$, factors uniquely through $\pi$*




*Furthermore there is a bijection of $A$-submodules*

$$\{P' \subseteq M/N\} \leftrightarrow \{P \mid N \subseteq P \subseteq M\}$$

*given by $P' = P/N$. In the situation above $\ker(\tilde{\psi}) = \ker(\psi)/N$. In particular if $\ker(\psi) = N$ then $\tilde{\psi}$ is injective.*

© David Rufino CC BY-NC-ND 4.0

**Corollary 3.3.102**
*Let $\psi : M \to N$ be an $A$-module homomorphism, then this induces an isomorphism*

$$M/\ker(\psi) \cong \mathrm{Im}(\psi)$$

**Definition 3.3.103 (Exact Sequence)**
*Let $N \xrightarrow{\varphi} M \xrightarrow{\psi} P$ be an sequence of $A$-module homomorphisms. We say it is **exact** if*

$$\mathrm{Im}(\varphi) = \ker(\psi)$$

*It is equivalent to the following two conditions*

*a)* $\psi \circ \varphi = 0$
*b)* $\psi(m) = 0 \implies m = \varphi(n)$ *for some* $n \in N$.

*An exact sequence of the form*

$$0 \to N \to M \to P \to 0$$

*is said to be **short-exact**.*

**Remark 3.3.104**
*There are a few trivial observations*

- $0 \to M \to N$ *is exact if and only if the map* $M \to N$ *is injective*
- $M \to N \to 0$ *is exact if and only if the map* $M \to N$ *is surjective.*

**Proposition 3.3.105 (Isomorphism induced by short-exact sequence)**
*Let $N \subseteq M$ be a $A$-submodule then there is a canonical short-exact sequence*

$$0 \to N \to M \to M/N \to 0$$

*Conversely suppose we have a short exact sequence*

$$0 \to N \xrightarrow{i} M \xrightarrow{\pi} P \to 0$$

*then this induces an isomorphism*

$$M/i(N) \cong P$$

*If $N$ is a submodule of $M$ then we would simply write $M/N \cong P$.*

**Proposition 3.3.106 (Third Isomorphism Theorem)**
*Let $N \subseteq N' \subseteq M$ be a chain of modules then there is a short-exact sequence*

$$0 \to N'/N \to M/N \to M/N' \to 0$$

*which then induces an isomorphism*

$$(M/N)/(N'/N) \cong M/N'$$

**Proposition 3.3.107 (Product of ideal and quotient module)**
*Let $N$ be a submodule of $M$ and $\mathfrak{a} \lhd A$ an ideal. Then*

$$\mathfrak{a}(M/N) = (N + \mathfrak{a}M)/N$$

**Proposition 3.3.108 (Induced module)**
*Let $M$ be an $A$-module and $\mathfrak{a}$ an ideal such that $\mathfrak{a}M = 0$, then $M$ is naturally an $A/\mathfrak{a}$-module with action given by*

$$\overline{a} \cdot m := a \cdot m$$

### 3.3.14 Dual Module

**Proposition 3.3.109 (Dual Module)**
*Let $M$ be a left (resp. right) $A$-module. Then the set*

$$M^\vee := \mathrm{Hom}_A(M, A)$$

*is canonically a* right *(resp.* left*) $A$-module.*

© David Rufino CC BY-NC-ND 4.0

*If $M$ is a finite free left (resp. right) $A$-module with basis $\{v_1, ..., v_n\}$ then $M^\vee$ is a finite free* right *(resp.* left*) $A$-module with basis $\{v_1^\vee, ..., v_n^\vee\}$ where these are the unique homomorphisms satisfying*

$$v_i^\vee(v_j) = \delta_{ij}$$

*Moreover every basis of $M^\vee$ is of this form.*

**Proposition 3.3.110 (Hom-Set is free)**
*Let $M$ be a finite free left $A$-module with basis $\{v_1, ..., v_n\}$ and $N$ a $(A, B)$-bimodule. Then there is an isomorphism of right $B$-modules*

$$\begin{aligned} \mathrm{Hom}_A(M, N) \cong & \bigoplus_{i=1}^{n} N \\ \theta \to & (\theta(v_i))_{i \in I} \end{aligned}$$

*If $A$ is commutative and $N$ is a finite-free $A$-module with basis $\{w_1, ..., w_m\}$ then there is further an isomorphism of $A$-modules*

$$\begin{aligned} \mathrm{Hom}_A(M, N) \cong & \bigoplus_{i,j=1}^{n,m} A \\ \theta \to & \left(w_j^\vee(\theta(v_i))\right)_{i,j} \end{aligned}$$

*where $w_1^\vee, ..., w_m^\vee$ is the dual basis for $N^\vee$. In particular $\mathrm{Hom}_A(M, N)$ is a finite-free $A$-module with basis*

$$\left\{w_j v_i^\vee\right\}_{i,j}$$

**Definition 3.3.111 (Dual Functor)**
*Let $A$ be a commutative ring and $\varphi : M \to N$ an $A$-module homomorphism. Define the dual homomorphism $\varphi^\vee : N^\vee \to M^\vee$ by*

$$\varphi^\vee(\psi) := \psi \circ \varphi$$

*This determines a contravariant functor*

$$(-)^\vee : A\textbf{-Mod} \to A\textbf{-Mod}$$

**Corollary 3.3.112 (Double Dual Natural Isomorphism)**
*Let $A$ be a commutative ring and $M$ a finite free $A$-module then the canonical $A$-module homomorphism*

$$\begin{aligned} \eta : M \longrightarrow & \quad M^{\vee\vee} \\ x \to & \quad (\varphi \to \varphi(x)) \end{aligned}$$

*is an isomorphism, which is natural in $M$.*

**Corollary 3.3.113**
*The contravariant functor $(-)^\vee : \mathbf{FiniteFreeMod}_A \to \mathbf{FiniteFreeMod}_A$ is an equivalence of categories and therefore full and faithful.*

*Proof.* Use the dual isomorphism $\eta$ together with (2.6.29) and (2.6.28). ∎

## 3.3.15 Matrices

For this section we assume that $A$ is a commutative ring. In this context $A^n := A \times ... \times A$ is a finite free module with basis $e_1, ..., e_n$. Matrices are concrete realisations of linear maps of finite free modules.

**Proposition 3.3.114**
*Let $M, N$ be finite free $A$-modules with ordered bases $\mathcal{B} := \{v_1, ..., v_n\}$, $\mathcal{B}' := \{w_1, ..., w_m\}$ respectively. Let $\varphi : M \to N$ be an $A$-linear map and define $[\varphi]$ to be the unique $m \times n$ matrix satisfying the following condition*

$$\varphi(v_i) = \sum_{j=1}^{m} [\varphi]_{ji} w_j \quad \forall i = 1...n$$

*(**notice sum over the first index**). Then it is also the unique matrix ensuring that the following diagram commutes*

© David Rufino CC BY-NC-ND 4.0

$$\begin{array}{ccc} M & \xrightarrow{\varphi} & N \\ \uparrow \sim & & \uparrow \sim \\ A^n & \xrightarrow{x \to [\varphi]x} & A^m \end{array}$$

*where the bottom arrow is given by usual multiplication of a matrix by a column vector*

**Proposition 3.3.115 (Matrices as linear maps)**
*Let $M$, $N$ be free $A$-modules with ordered bases $\mathcal{B} := \{v_1, ..., v_n\}$, $\mathcal{B}' := \{w_1, ..., w_m\}$ respectively. Then there are mutually inverse isomorphisms of $A$-modules*

$$\begin{aligned} \mathrm{Mat}_{m\times n}(A) &\longleftrightarrow \mathrm{Hom}_A(M, N) \\ E &\to \hat{E} \\ [\varphi]^{\mathcal{B}}_{\mathcal{B}'} &\leftarrow \varphi \end{aligned}$$

*where*

$$\begin{aligned} \hat{E}\left(\sum_{i=1}^{n} \lambda_i v_i\right) &:= \sum_{j=1}^{m}\left(\sum_{i=1}^{n} E_{ji}\lambda_i\right) w_j \\ \varphi(v_i) &= \sum_{j=1}^{m} [\varphi]_{ji} w_j \end{aligned}$$

*If we further consider a free $A$-module $P$ with ordered bases $\mathcal{B}'' = \{u_1, ..., u_p\}$ then*

$$\begin{aligned} \hat{E} \circ \hat{F} &= \widehat{EF} \\ [\psi \circ \varphi]^{\mathcal{B}}_{\mathcal{B}''} &= [\psi]^{\mathcal{B}'}_{\mathcal{B}''}[\varphi]^{\mathcal{B}}_{\mathcal{B}'} \end{aligned}$$

*Observe that*

$$[1_M]^{\mathcal{B}}_{\mathcal{B}} = I_n$$

*Furthermore there is an isomorphism of $A$-algebras*

$$\begin{aligned} \mathrm{End}_A(M) &\longleftrightarrow \mathrm{Mat}_{n,n}(A) \\ \varphi &\to \left(v_i^\vee(\varphi(v_j))\right)_{ij} \\ \sum_{ij} E_{ij} v_i v_j^\vee &\leftarrow E \end{aligned}$$

**Corollary 3.3.116**
*Matrix multiplication is associative. In particular*

$$(EF)v = E(Fv)$$

*Proof.* We may consider the free $A$-modules $A^n$, $A^m$ and $A^p$ with canonical bases. The result follows because function composition is associative and $\hat{\cdot}$ is injective. ∎

**Corollary 3.3.117**
*There is an isomorphism of $A$-modules*

$$\begin{aligned} \mathrm{Mat}_{m\times n}(A) &\longleftrightarrow \mathrm{Hom}_A(A^n, A^m) \\ E &\to (v \to Ev) \end{aligned}$$

*and further an isomorphism of $A$-algebras*

$$\mathrm{Mat}_{n,n}(A) \longleftrightarrow \mathrm{End}_A(A^n)$$

**Corollary 3.3.118**
*Let $M$ be a finite free $A$-module with bases $\mathcal{B}, \mathcal{B}'$ and $\varphi \in \mathrm{End}_A(M)$. Then $\varphi$ is an isomorphism if and only if $[\varphi]^{\mathcal{B}}_{\mathcal{B}'} \in \mathrm{Mat}_{n\times n}(A)$ is invertible.*

© David Rufino CC BY-NC-ND 4.0

**Corollary 3.3.119 (Change of basis)**
*Let $M$ be a finite free $A$-module and $\mathcal{B}, \mathcal{B}'$ bases then*

$$[1_M]^{\mathcal{B}}_{\mathcal{B}'} = \left([1_M]^{\mathcal{B}'}_{\mathcal{B}}\right)^{-1}$$

*and*

$$[\varphi]^{\mathcal{B}'}_{\mathcal{B}'} = P[\varphi]^{\mathcal{B}}_{\mathcal{B}}P^{-1}$$

*where*

$$P := [1_M]^{\mathcal{B}}_{\mathcal{B}'}$$

*is invertible.*

**Definition 3.3.120 (Transpose)**
*Let $E$ be an $m \times n$ matrix in $A$, then define the **transpose** of $E$ to be the $n \times m$ matrix $E^t$ where*

$$\left(E^t\right)_{ij} := E_{ji}$$

**Proposition 3.3.121**
*Let $M, N$ be finite-free $A$-modules with bases $\mathcal{B} = \{v_1, ..., v_n\}$ and $\mathcal{B}' = \{w_1, ..., w_m\}$. Let $\varphi : M \to N$ be an $A$-module homomorphism and $\varphi^\vee : N^\vee \to M^\vee$ the dual homomorphism then*

$$[\varphi^\vee]^{\mathcal{B}'^\vee}_{\mathcal{B}^\vee} = \left([\varphi]^{\mathcal{B}}_{\mathcal{B}'}\right)^t$$

*Similarly if $E$ is an $m \times n$ matrix over $A$ then*

$$\hat{E}^\vee = \widehat{E^t}$$

*where the right hand side is understood to be with respect to the dual bases.*

**Corollary 3.3.122**
*Let $E, F$ be matrices then*

$$(EF)^t = (FE)^t$$

### 3.3.16 Vector Spaces

**Definition 3.3.123 (Vector Space)**
*A **vector space** $V$ over $k$ is simply a $k$-module.*

*A $k$-submodule is referred to as a **subspace***

*A $k$-module homomorphism is referred to as a **linear map***

*A vector space is finite-dimensional if it is finite as a $k$-module.*

The main result on vector spaces is that bases exist and all have the same cardinality. Recall that $\langle\cdot\rangle$ is a closure operator. We show that $(V, \langle\cdot\rangle)$ determines a matroid so that we may appeal to results in the Matroids section. First we need to show that the notions of independence coincide

**Proposition 3.3.124 (Equivalent definitions of linear independence)**
*Let $V$ be a vector space and $S \subset V$. Then the following are equivalent*

*a) $S$ is linearly independent*
*b) No proper subset $S' \subset S$ satisfies $\langle S' \rangle = \langle S \rangle$*
*c) **Matroid Independence** $x \in S \implies x \notin \langle S \setminus \{x\} \rangle$*

*Further $S$ is independent if and only if every finite subset of $S$ is.*

*Proof.* b) $\iff$ c) This is b).

a) $\implies$ c). If $x \in \langle S \setminus \{x\} \rangle$ then it's clear that $S$ is not linearly independent.

c) $\implies$ a). Suppose we have a linear relationship

© David Rufino CC BY-NC-ND 4.0

$$0 = \sum_{i=1}^{n} \lambda_i v_i \quad v_i \in S$$

By renumbering assume that $\lambda_1 \neq 0$, then rearrange to show $v_1 \in \langle S \setminus \{v_1\} \rangle$, contradicting the hypothesis. ∎

We show that $(V, \langle \cdot \rangle)$ satisfies the Exchange Property and therefore constitutes a matroid.

**Proposition 3.3.125 (Exchange Property)**
*Let $V$ be a vector space and $S \subset V$. Then*

$$y \in \langle S \cup \{x\} \rangle \setminus \langle S \rangle \implies x \in \langle S \cup \{y\} \rangle$$

*Proof.* Suppose $y$ is as given, then

$$y = \lambda x + \sum_i \lambda_i s_i \quad s_i \in S$$

The case $\lambda = 0$ is obvious. Otherwise we may rearrange to find

$$x = \lambda^{-1} y - \sum_i \lambda^{-1} \lambda_i s_i$$

whence $x \in \langle S \cup \{y\} \rangle$. ∎

Therefore we have the following

**Proposition 3.3.126 (Vector Spaces are Free)**
*Every vector space has a basis, and in the finite-dimensional case every basis is finite of the same size. We denote this by $\dim_k V$.*

*More generally every linearly independent set is contained in a basis (so has order at most $\dim_k V$) and every spanning set contains a basis (so has order at least $\dim_k V$)*

*Proof.* Follows from (2.3.7) and (2.3.12). The final statement follows from (2.3.13). ∎

**Proposition 3.3.127 (Basis Criteria)**
*Let $V$ be a vector space with $n = \dim_k V$ and $\mathcal{B} \subseteq V$. Then TFAE*

*a) $\mathcal{B}$ is a basis*
*b) $\mathcal{B}$ is linearly independent and $\#\mathcal{B} \geq \dim_k V$*
*c) $\mathcal{B}$ is spanning and $\#\mathcal{B} \leq \dim_k V$*

*Proof.* The equivalence of a), b) and c) follows from (2.3.14). ∎

**Proposition 3.3.128**
*A vector space $V = \{0\}$ if and only if $\dim_k V = 0$*

**Proposition 3.3.129 (Image of a basis)**
*Let $\varphi : V \to W$ be a linear map*

*a) If $S$ is linearly-independent and $\varphi$ is injective, then $\varphi(S)$ is linearly-independent*
*b) If $\Gamma$ is spanning then ($\varphi$ is surjective $\iff$ $\varphi(\Gamma)$ is spanning)*
*c) If $\mathcal{B}$ is a basis then ($\varphi$ is an isomorphism $\iff$ $\varphi(\mathcal{B})$ is a basis and $\varphi$ injective on $\mathcal{B}$)*

*Proof.*

a) Suppose $\sum_i \lambda_i \varphi(s_i) = 0 \implies \varphi\left(\sum_i \lambda_i s_i\right) = 0$. As $\varphi$ is injective this implies $\sum_i \lambda_i s_i = 0 \implies \lambda_i = 0$.
b) If $\varphi$ is surjective then for $w \in W$ we have $\varphi(v) = w$ for some $v \in V$. By hypothesis $v = \sum_i \lambda_i \gamma_i$ and $w = \sum_i \varphi(\lambda_i)$. Conversely given $w \in W$ by hypothesis $w = \sum_i \lambda_i \varphi(\gamma_i) = \varphi\left(\sum_i \lambda_i \gamma_i\right)$ and $\varphi$ is surjective as required.
c) Suppose $\varphi$ is isomorphism, then it's surely injective on $\mathcal{B}$ and by a),b) $\varphi(\mathcal{B})$ is a basis. Conversely if $\varphi(\mathcal{B}) =: \mathcal{B}'$ is a basis then by b) $\varphi$ is surjective. Suppose $\varphi(v) = 0$. Then by hypothesis $v = \sum_i \lambda_i v_i$ for $v_i \in \mathcal{B}$ and $0 = \varphi(v) = \sum_i \lambda_i \varphi(v_i)$. By hypothesis $\varphi(v_i)$ are distinct elements of the basis $\mathcal{B}'$ and therefore $\lambda_i = 0$ and $v = 0$. Therefore $\varphi$ is injective and hence bijective.

∎

**Corollary 3.3.130 (Dimension is an invariant)**
*Dimension is preserved under isomorphism. More generally for $\varphi : V \to W$ we have*

© David Rufino CC BY-NC-ND 4.0

$$\varphi \text{ injective} \implies \dim_k V \leq \dim_k W$$

$$\varphi \text{ surjective} \implies \dim_k V \geq \dim_k W$$

**Proposition 3.3.131**
*Let $W \subseteq V$ be finite-dimensional vector spaces then the dimension of the quotient module satisfies*

$$\dim_k V/W = \dim_k V - \dim_k W$$

*Proof.* Let $\{v_1, ..., v_m\}$ be a basis of $W$, then there exists a basis $\{v_1, ..., v_m, v_{m+1}, ..., v_n\}$ of $V$ containing the first by (3.3.126). We claim that

$$\{[v_{m+1}], ..., [v_n]\}$$

is a basis for $V/W$, and the result follows. For given $[v] \in V/W$ then

$$v = \sum_{i=1}^{n} \lambda_i v_i$$

since the basis is spanning. We have

$$v - \sum_{i=m+1}^{n} \lambda_i v_i \in W$$

therefore

$$[v] = \left[\sum_{i=m+1}^{n} \lambda_i v_i\right] = \sum_{i=m+1}^{n} \lambda_i [v_i]$$

and the given set is spanning. Similarly suppose

$$\sum_{i=m+1}^{n} \lambda_i [v_i] = 0$$

then by definition $\sum_{i=m+1}^{n} \lambda_i v_i \in W$. Therefore

$$\sum_{i=m+1}^{n} \lambda_i v_i = \sum_{i=1}^{m} \lambda_i v_i$$

and since $v_i$ are linearly independent we must have $\lambda_i = 0$. ∎

**Proposition 3.3.132 (Injective Criteria)**
*Let $\varphi : V \to W$ be a linear map then*

$$\varphi \text{ injective} \iff \ker(\varphi) = \{0\} \iff \dim_k \ker(\varphi) = 0$$

*Proof.* Note for any linear map $\varphi$ we have $\varphi(0) = 0$. Therefore $\varphi$ injective clearly shows $\ker(\varphi) = \{0\}$. Conversely suppose $\ker(\varphi) = 0$ and $\varphi(v) = \varphi(w)$. Then $\varphi(v - w) = 0 \implies v - w = 0 \implies v = w$ as required. ∎

**Definition 3.3.133 (Rank)**
*Let $\varphi : V \to W$ be a linear map then define*

$$\operatorname{rank}_k(\varphi) := \dim_k(\operatorname{Im}(\varphi))$$

**Proposition 3.3.134 (Surjective Criteria)**
*Let $\varphi : V \to W$ be a linear map with $W$ finite-dimensional then*

$$\varphi \text{ surjective} \iff \operatorname{rank}_k(\varphi) = \dim_k W$$

*Proof.* This follows directly from (2.3.15). ∎

**Proposition 3.3.135 (Isomorphism Theorem / Rank-Nullity)**
*Let $\varphi : V \to W$ be a linear map then this induces an isomorphism*

$$V/\ker(\varphi) \xrightarrow{\sim} \operatorname{Im}(\varphi)$$

*in particular when $V$ is finite-dimensional*

© David Rufino CC BY-NC-ND 4.0

$$\dim_k V = \dim_k \ker(\varphi) + \operatorname{rank}_k(\varphi)$$

**Corollary 3.3.136 (Isomorphism Criteria)**
*Let $V, W$ vector spaces with $W$ finite-dimensional. A linear map $\varphi : V \to W$ is an isomorphism if and only if any* two *of the following are satisfied*

*a)* $\dim_k \ker(\varphi) = 0 \iff \varphi$ *injective*
*b)* $\dim_k V = \dim_k W$
*c)* $\operatorname{rank}_k(\varphi) = \dim_k W \iff \varphi$ *surjective*

*Proof.* The rank-nullity equation ensures that if any two hold the third is automatically satisfied. In this case $\varphi$ is isomorphism as required. ∎

**Corollary 3.3.137 (Endomorphism Isomorphism Criteria)**
*Let $V$ be a finite-dimensional vector space and $\varphi : V \to V$ then TFAE*

*a)* $\varphi$ *is injective*
*b)* $\varphi$ *is surjective*
*c)* $\varphi$ *is an isomorphism*

*Proof.* For the equivalence of a), b) and c) we may use the previous result with $W = V$ and note $\dim_k W = \dim_k V$ is automatically satisfied. ∎

**Proposition 3.3.138 (Internal Direct Sum)**
*Let $U_1, U_2$ be two subspaces of $V$ then TFAE*

*a)* $U_1 \cap U_2 = \{0\}$ *and* $V = U_1 + U_2$
*b) Every* $v \in V$ *may be written uniquely as* $u_1 + u_2$ *for* $u_i \in U_i$.

*and we say $V = U_1 \oplus U_2$ is an* ***internal direct sum*** *and $U_2$ is a* ***supplementary subspace*** *for $U_1$.*

**Proposition 3.3.139 (Subspaces are direct factors)**
*Every subspace $U$ has a supplementary subspace $U'$ such that*

$$V = U \oplus U'$$

*Proof.* Let $\mathcal{B}_1$ be a basis for $U$ and extend to a basis $\mathcal{B}$ and define $\mathcal{B}_2 := \mathcal{B} \setminus \mathcal{B}_1$. Then it's easy to show that $U' = \langle \mathcal{B}_2 \rangle$ is a supplementary subspace. ∎

**Proposition 3.3.140 (Dimension formula for direct sums)**
*Suppose $V = U_1 \oplus U_2$, $\mathcal{B}_1$ is a basis for $U_1$ and $\mathcal{B}_2$ is a basis for $U_2$. Then $\mathcal{B}_1 \cap \mathcal{B}_2 = \varnothing$ and $\mathcal{B}_1 \cup \mathcal{B}_2$ is a basis for $V$. In particular*

$$\dim_k V = \dim_k U_1 + \dim_k U_2$$

#### 3.3.16.1 Dual Space

**Definition 3.3.141 (Dual Space)**
*Let $V$ be a $k$-vector space and define the* ***dual space*** *to be*

$$V^\vee := \operatorname{Hom}_k(V, k)$$

*This is an abelian group and even a $k$-vector space under the obvious operations. The construction $V \to V^\vee$ determines a contravariant functor*

$$(-)^\vee : \mathbf{Vect}_k \to \mathbf{Vect}_k$$

**Definition 3.3.142 (Annihilator)**
*Let $V$ be a vector space and $U \subseteq V$ a subspace. Define the* ***annihilator*** *of $U$ by*

$$U^\circ = \{\theta \in V^\vee \mid \theta(u) = 0 \quad \forall u \in U\}$$

*This is a linear subspace of $V^\vee$.*

**Proposition 3.3.143 (Dimension formula for annihilators)**
*There is a canonical isomorphism by restriction*

$$V^\vee / U^\circ \to U^\vee$$

© David Rufino CC BY-NC-ND 4.0

*In particular when $V$ is a finite-dimensional vector space then*

$$\dim_k V = \dim_k U + \dim_k U^\circ$$

*Proof.* Let $W$ be a supplementary subspace and consider the morphism $V = U \oplus W \xrightarrow{\pi_U} U$. Then $(\theta \circ \pi_U)|_U = \theta$ so the restriction map is surjective. Clearly the kernel is $U^\circ$. The dimension formula follows from (3.3.135) and (3.3.109). ■

**Corollary 3.3.144 (Dual rank = rank)**
*Let $\varphi : V \to W$ be a linear map and $\varphi^\vee : W^\vee \to V^\vee$ then*

$$\begin{aligned} \ker(\varphi^\vee) &= \operatorname{Im}(\varphi)^\circ \\ \operatorname{Im}(\varphi^\vee) &\subseteq \ker(\varphi)^\circ \end{aligned}$$

*In the finite-dimensional case $\operatorname{Im}(\varphi^\vee) = \ker(\varphi)^\circ$ and*

$$\begin{aligned} \dim_k \ker(\varphi^\vee) &= \dim_k W - \operatorname{rank}_k(\varphi) \\ \operatorname{rank}_k(\varphi^\vee) &= \operatorname{rank}_k(\varphi) \end{aligned}$$

*Proof.* Note $\ker(\varphi^\vee) = \operatorname{Im}(\varphi)^\circ$ and $\operatorname{Im}(\varphi^\vee) \subseteq \ker(\varphi)^\circ$ by the definitions.

Consider the finite-dimensional case. By (3.3.143)

$$\dim_k \ker(\varphi^\vee) = \dim_k \operatorname{Im}(\varphi)^\circ = \dim_k W - \operatorname{rank}_k(\varphi)$$

By rank-nullity applied to $\varphi^\vee$ and $\dim_k W = \dim_k W^\vee$ we deduce

$$\operatorname{rank}_k(\varphi^\vee) = \operatorname{rank}_k(\varphi).$$

By (3.3.143) and rank-nullity applied to $\varphi$

$$\dim_k \ker(\varphi)^\circ = \dim_k V - \dim_k \ker(\varphi) = \operatorname{rank}_k(\varphi).$$

Finally by (2.3.15) $\operatorname{Im}(\varphi^\vee) = \ker(\varphi)^\circ$. ■

From this it follows that taking duals reflects and preserve isomorphisms

**Corollary 3.3.145 ($(-)^\vee$ reflects isomorphisms)**
*Let $\varphi : V \to W$ be a linear map of finite-dimensional spaces then*

*a) $\varphi$ is injective if and only if $\varphi^\vee$ is surjective*
*b) $\varphi$ is surjective if and only if $\varphi^\vee$ is injective*
*c) $\varphi$ is iso if and only if $\varphi^\vee$ is iso*

*Proof.* Note by (3.3.144) we have $\operatorname{rank}_k(\varphi) = \operatorname{rank}_k(\varphi^\vee)$ and by (3.3.109) $\dim_k V^\vee = \dim_k V$.

$\varphi$ is surjective $\iff \operatorname{rank}_k(\varphi) = \dim_k W = \operatorname{rank}_k(\varphi^\vee) \overset{(3.3.135)}{\iff} \dim_k \ker(\varphi^\vee) = 0 \iff \ker(\varphi^\vee) = \{0\}$

$\varphi$ is injective $\iff \dim_k \ker(\varphi) = 0 \overset{(3.3.135)}{\iff} \operatorname{rank}_k(\varphi) = \dim_k V \iff \dim_k V^\vee = \operatorname{rank}_k(\varphi^\vee) \iff \varphi^\vee$ is surjective.

The last point may be deduced from the first two, or the fact that $(-)^\vee$ is full and faithful (3.3.113) and category-theoretic result (2.6.42). ■

### 3.3.16.2 Bilinear Pairings

**Definition 3.3.146 (Bilinear maps)**
*Let $V, W$ be vector spaces a bilinear map $\psi$ is a map*

$$\psi : V \times W \to k$$

*which is $k$-linear in each variable. We denote the set of bilinear maps as*

$$\operatorname{Bilin}_k(V, W)$$

**Proposition 3.3.147 (Matrix Representation)**
*Let $V, W$ be finite-dimensional vector spaces with bases $\{v_1, ..., v_n\}$ and $\{w_1, ..., w_m\}$ then there is a canonical isomorphism*

$$\begin{aligned} \operatorname{Bilin}_k(V, W) &\xrightarrow{\sim} \operatorname{Mat}_{n,m}(k) \\ \psi &\to \left(\psi(v_i, w_j)\right)_{ij} \end{aligned}$$

© David Rufino CC BY-NC-ND 4.0

*In particular a bilinear map $\psi$ is determined uniquely by the values $\psi(v_i, w_j)$.*

**Proposition 3.3.148 (Dual maps)**
*Let $V$ and $W$ be vector spaces, then there is a natural bijection*

$$\begin{array}{ccccc} \mathrm{Mor}_k(V, W^\vee) & \leftrightarrow & \mathrm{Bilin}_k(V, W) & \leftrightarrow & \mathrm{Mor}_k(W, V^\vee) \\ \psi_L & \leftarrow & \psi & & \\ & & \psi & & \rightarrow \psi_R \end{array}$$

*where*

$$\psi_L(v)(w) = \psi(v, w) = \psi_R(w)(v)$$

*When $V, W$ are finite-dimensional then $\psi_L$ is an isomorphism if and only if $\psi_R$ is an isomorphism. In this case we say $\psi$ is a perfect pairing. More generally*

$$\mathrm{rank}_k(\psi_L) = \mathrm{rank}_k(\psi_R)$$

*Proof.* The bijections stated are obvious. One may show that $\psi_L = \psi_R^\vee \circ \eta_V$ where $\eta_V : V \to V^{\vee\vee}$ is the dual isomorphism. Therefore $\psi_L$ is an isomorphism if and only if $\psi_R^\vee$ is an isomorphism, and by (3.3.145) if and only if $\psi_R$ is an isomorphism. Since $\eta_V$ is surjective we have $\mathrm{rank}_k(\psi_L) = \mathrm{rank}_k(\psi_R^\vee) = \mathrm{rank}_k(\psi_R)$, by (3.3.144). ∎

**Proposition 3.3.149**
*Let $\psi : V \times V \to k$ be a bilinear map. Then the following are equivalent*

*a) $\psi$ is a perfect pairing*
*b) $\psi_L$ is an isomorphism*
*c) $\psi_R$ is an isomorphism*
*d) $\psi_L$ is injective*
*e) $\psi_R$ is injective*
*f) $\psi_L$ is surjective*
*g) $\psi_R$ is surjective.*

*Proof.* We may use the fact $\dim_k V = \dim_k V^\vee$ and (3.3.136) to show b) ⟺ d) ⟺ f) and c) ⟺ e) ⟺ g). We've already shown that a) ⟺ b) ⟺ c). ∎

**Definition 3.3.150 (Orthogonal Complement)**
*Let $\psi : V \times W \to k$ be a perfect pairing of finite-dimensional vector spaces. Suppose $U \subset V$ is a subspace then define the **orthogonal complement***

$$U^\perp := \{w \in W \mid \psi(v, w) = 0 \quad \forall v \in U\}$$

**Proposition 3.3.151**
*Let $\psi : V \times W \to k$ be a perfect pairing of finite-dimensional vector spaces and $U \subset V$ a subspace. Then*

$$\dim_k U + \dim_k U^\perp = \dim_k V$$

*Indeed $\psi_R$ induces an isomorphism $U^\perp \to U^\circ$.*

*Proof.* We claim that $\psi_R(U^\perp) = U^\circ$. For if $w \in U^\perp$ then $\psi_R(w)(v) = \psi(w, v) = 0$ for all $v \in U$, and so $\psi_R(w) \in U^\circ$. Conversely given $\theta \in U^\circ$, as $\psi_R$ is surjective, there is $w \in W$ such that $\psi_R(w) = \theta$. By definition $w \in U^\perp$ as required.

As $\psi_R$ is injective then $\dim_k U^\perp \overset{(3.3.130)}{=} \dim_k U^\circ \overset{(3.3.143)}{=} \dim_k V/U = \dim_k V - \dim_k U$. ∎

**Remark 3.3.152**
*In the case $V = W$, then it's not necessarily true that $U \cap U^\perp = \{0\}$, and so $U^\perp$ is not necessarily a complementary subspace.*

*The classic example is the perfect pairing on $\mathbb{R}^n$ induced by $vDv^T$ for a real diagonal matrix $D$. Then it's true in general if and only if $D$ is positive-definite.*

**Proposition 3.3.153 (Quotients are dual to subspaces)**
*Let $\psi : V \times W \to k$ be a perfect pairing of finite-dimensional vector spaces. Suppose $U \subset V$ is a subspace, then there is a canonical perfect pairing*

$$\psi' : V/U \times U^\perp \to k$$

© David Rufino CC BY-NC-ND 4.0

*given by*

$$\psi'(v+U,w)=\psi(v,w)$$

*Proof.* The given map is well defined, for suppose $v_1+U=v_2+U$ then $v_1-v_2\in U\implies\psi(v_1-v_2,w)=0\quad\forall w\in U^\perp\implies\psi(v_1,w)=\psi(v_2,w)$ as required. It's clearly $k$-bilinear.

It's clear that $\psi'_R$ is injective, because $\psi'_R(w)=0_{V/U}\implies\psi_R(w)=0_V\implies w=0$.

By the previous Proposition $\dim_k U^\perp=\dim_k V/U$. Therefore by (3.3.136) $\psi'_R$ is an isomorphism and $\psi'$ is perfect.∎

## 3.3.17 Inverse Limits

We record the basic construction of an inverse limit of a tower of rings, modules, or algebras, as a concrete model for the categorical inverse limit (2.6.129), together with its functoriality and its compatibility with products; these are used repeatedly to build completions later on.

**Definition 3.3.154 (Inverse Limit of a Tower)**
*Let $\mathcal{C}$ be one of* **Ring**, *$A$-***Mod** *(for a fixed ring $A$), or $\mathrm{Alg}_A$ (for a fixed ring $A$, so in particular* $\mathbf{Alg}_k$*) – each an algebraic category (2.6.184) (via the usual axioms; (2.6.185), (2.6.186), (2.6.187) respectively). Let $\cdots\twoheadrightarrow M_2\twoheadrightarrow M_1\twoheadrightarrow M_0$ be a sequence of morphisms $\rho_n:M_n\to M_{n-1}$ of $\mathcal{C}$ $(n\geq 1)$. Define*

$$\varprojlim_n M_n:=\left\{(m_n)_{n\geq0}\in\prod_{n\geq0}M_n\;\middle|\;\rho_n(m_n)=m_{n-1}\text{ for every }n\geq1\right\},$$

*a sub-object of $\prod_n M_n$ in $\mathcal{C}$ (a subring, submodule, or subalgebra respectively, the ambient operations acting componentwise), with the evident projections $\lambda_n:\varprojlim_m M_m\to M_n$. Writing $\rho_{n,n}:=1_{M_n}$ and $\rho_{n,m}:=\rho_{m+1}\circ\cdots\circ\rho_n:M_n\to M_m$ for $n>m$ (composites of consecutive $\rho_k$'s), every $(m_n)_{n\geq0}\in\varprojlim_n M_n$ satisfies the* ***repeated compatibility***

$$\rho_{n,m}(m_n)=m_m\quad\text{for all }n\geq m,$$

*not just for consecutive $n,m$: by induction on $n-m$, composing the defining relations $\rho_k(m_k)=m_{k-1}$ for $m<k\leq n$.*

**Lemma 3.3.155 (This Realises the Categorical Inverse Limit)**
*In the setting of (3.3.154), regard $I:=\mathbb{N}$ (usual order) as a directed preorder and the tower as an inverse system $F:I^{op}\to\mathcal{C}$, $F(n):=M_n$, with restriction maps $\rho_{n,m}$ as in (3.3.154). Then $\left(\varprojlim_n M_n,\{\lambda_n\}\right)$ is an inverse limit for $F$ in the sense of (2.6.129).*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

$\mathcal{C}$ is an algebraic category (2.6.184), in each of the three cases (as recalled in (3.3.154)). By (2.6.191), $\varprojlim_n F(n)$ exists in $\mathcal{C}$, with underlying set $\left\{(m_n)_n\in\prod_n M_n:\rho_{n,m}(m_n)=m_m\text{ for all }n\geq m\right\}$ and the ambient operations acting coordinatewise; by the repeated compatibility (3.3.154), this defining condition holds for all $n\geq m$ as soon as it holds for consecutive $n=m+1$, so this is exactly the sub-object $\varprojlim_n M_n$ of (3.3.154), with the same (componentwise) structure and projections. So $\left(\varprojlim_n M_n,\{\lambda_n\}\right)$ is this categorical inverse limit. ∎

**Corollary 3.3.156 (Functoriality of the Inverse Limit)**
*Let $\mathcal{C}$ be as in (3.3.154), $(A_n,\rho_n)_{n\geq0}$ and $(B_n,\tau_n)_{n\geq0}$ towers in $\mathcal{C}$, and for each $n\geq0$ let $\sigma_n:A_n\to B_n$ be a morphism of $\mathcal{C}$ such that, for every $n\geq1$, the square*

$$\begin{array}{ccc} A_n & \xrightarrow{\sigma_n} & B_n \\ {\scriptstyle\rho_n}\downarrow & & \downarrow{\scriptstyle\tau_n} \\ A_{n-1} & \xrightarrow[\sigma_{n-1}]{} & B_{n-1} \end{array}$$

*commutes, i.e. $\tau_n\circ\sigma_n=\sigma_{n-1}\circ\rho_n$. Then*

$$\varprojlim\sigma:\varprojlim_n A_n\longrightarrow\varprojlim_n B_n,\quad(a_n)_n\mapsto(\sigma_n(a_n))_n$$

*is a well-defined morphism of $\mathcal{C}$; and if every $\sigma_n$ is an isomorphism, so is $\varprojlim\sigma$.*

© David Rufino CC BY-NC-ND 4.0

*Proof.* By (3.3.155), $\lim_{\leftarrow n} A_n$ and $\lim_{\leftarrow n} B_n$ (with their projections) are inverse limits, in the sense of (2.6.129), of the inverse systems $F, G : \mathbb{N}^{\mathrm{op}} \to \mathcal{C}$ built from the two towers; the commuting squares above are exactly the naturality hypothesis of (2.6.132) for the family $\sigma = (\sigma_n)_n$ (composing squares for consecutive $n$ gives it for the composite restriction maps $\rho_{n,m}$, $\tau_{n,m}$, $n > m$, as needed there). That corollary gives the induced morphism $\lim_{\leftarrow n} \sigma_n$, agreeing on each coordinate with $\sigma_n$ by construction of the induced map on limits-as-sets (2.6.130) – that is, the map $(a_n)_n \to (\sigma_n(a_n))_n$ above – and an isomorphism whenever every $\sigma_n$ is. ∎

**Proposition 3.3.157 (Inverse Limits Commute with Products)**
*Let $\mathcal{C}$ be as in (3.3.154), $K$ a set, and for each $k \in K$ a tower $(M_{k,n}, \rho_{k,n})_{n\geq 0}$ in $\mathcal{C}$. Then $\left(\prod_{k\in K} M_{k,n}\right)_{n\geq 0}$, with restriction maps $\prod_{k\in K} \rho_{k,n} : \prod_k M_{k,n} \to \prod_k M_{k,n-1}$, is again a tower in $\mathcal{C}$, and*

$$\Phi : \prod_{k\in K} \lim_{\leftarrow n} M_{k,n} \longrightarrow \lim_{\leftarrow n} \left(\prod_{k\in K} M_{k,n}\right), \quad \Phi\big((x_k)_{k\in K}\big) := \Big(\big(x_{k,n}\big)_{k\in K}\Big)_{n\geq 0} \quad \big(x_k = (x_{k,n})_{n\geq 0}\big)$$

*is a well-defined isomorphism of $\mathcal{C}$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Write $F_k : \mathbb{N}^{\mathrm{op}} \to \mathcal{C}$ for the inverse system of the tower $(M_{k,n})_n$, so $\left(\lim_{\leftarrow n} M_{k,n}, \{\lambda_{k,n}\}_n\right)$ realises $\lim_{\leftarrow} F_k$ (3.3.155), and likewise $\left(\lim_{\leftarrow n}\left(\prod_k M_{k,n}\right), \{\Lambda_n\}_n\right)$ realises $\lim_{\leftarrow}\left(\prod_k F_k\right)$, where $\Lambda_n$ is projection onto the $n$-th coordinate. Since $\prod_{k\in K} \lim_{\leftarrow n} M_{k,n}$ (with the evident coordinate projections $p_k$) is the categorical product of the $\lim_{\leftarrow n} M_{k,n}$ in $\mathcal{C}$, (2.6.133) gives that $\lim_{\leftarrow n}\left(\prod_k M_{k,n}\right)$ exists (as it already does, here) and that there is a *unique morphism* $\Phi'$ (of $\mathcal{C}$, not assumed a priori to be an isomorphism) such that

$$\pi_{k,n} \circ \Lambda_n \circ \Phi' = \lambda_{k,n} \circ p_k \quad \text{for all } k \in K, n \geq 0,$$

where $\pi_{k,n} : \prod_{k'} M_{k',n} \to M_{k,n}$ is the product projection; and that this $\Phi'$ is an isomorphism.

The explicit $\Phi$ above satisfies exactly this defining property: for $(x_k)_k \in \prod_k \lim_{\leftarrow n} M_{k,n}$ (so $x_k = (x_{k,n})_n$), both $\pi_{k,n}\big(\Lambda_n\big(\Phi\big((x_k)_k\big)\big)\big)$ and $\lambda_{k,n}\big(p_k\big((x_k)_k\big)\big)$ equal $x_{k,n}$. By uniqueness of the morphism with this property, $\Phi = \Phi'$; as $\Phi'$ is an isomorphism, so is $\Phi$. ∎

**Corollary 3.3.158 (Cofinal Reindexing of a Tower)**
*Let $\mathcal{C}$ be as in (3.3.154), $(M_n, \rho_n)_{n\geq 0}$ a tower in $\mathcal{C}$, and $n_0 < n_1 < n_2 < \cdots$ a strictly increasing sequence in $\mathbb{N}$. Then, writing $\tilde{\rho}_N := \rho_{n_{N+1},n_N} : M_{n_{N+1}} \to M_{n_N}$ for the composite restriction map of (3.3.155), $\left(M_{n_N}, \tilde{\rho}_N\right)_{N\geq 0}$ is again a tower in $\mathcal{C}$, and*

$$(m_n)_{n\geq 0} \mapsto \left(m_{n_N}\right)_{N\geq 0}$$

*defines an isomorphism $\lim_{\leftarrow n} M_n \xrightarrow{\sim} \lim_{\leftarrow N} M_{n_N}$ of $\mathcal{C}$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Regard $I := \mathbb{N}$ as a directed preorder and, per (3.3.155), the tower $(M_n)_n$ as an inverse system $F : I^{\mathrm{op}} \to \mathcal{C}$ realising $\lim_{\leftarrow n} M_n$. Let $I' := \{n_N : N \geq 0\} \subset I$, a full subcategory: it is directed (a subset of the totally ordered set $I$), and cofinal, as $n_N \to \infty$ so every $n \in I$ satisfies $n \leq n_N$ for some $N$ (2.6.126). The order-preserving bijection $\sigma : \mathbb{N} \to I'$, $\sigma(N) := n_N$, is an isomorphism of the directed preorders $\mathbb{N}$ and $I'$, and $F\sigma : \mathbb{N}^{\mathrm{op}} \to \mathcal{C}$ is exactly the tower $\left(M_{n_N}, \tilde{\rho}_N\right)_N$ (its restriction map for $N < N+1$ is $F(\sigma(N) \to \sigma(N+1)) = \rho_{n_{N+1},n_N} = \tilde{\rho}_N$, and composites thereof match for $N < N'$); so, by (2.6.98) (applied to $\sigma$) together with (3.3.155) applied to the tower $\left(M_{n_N}, \tilde{\rho}_N\right)_N$, $\left(\lim_{\leftarrow n} M_n, \{\lambda_{n_N}\}_{N\geq 0}\right)$ realises $\lim_{\leftarrow i\in I'} F(i)$, and $\left(\lim_{\leftarrow N} M_{n_N}, \{\lambda'_N\}_{N\geq 0}\right)$ (the latter's own projections) also realises it; identifying these two cones over $F|_{I'^{\mathrm{op}}}$ via the (unique, by terminality) comparison isomorphism between them, which sends $(m_n)_{n\geq 0} \in \lim_{\leftarrow n} M_n$ (viewed as a cone-point of the first) to the compatible sequence $\left(m_{n_N}\right)_{N\geq 0} \in \lim_{\leftarrow N} M_{n_N}$ agreeing with it on projections $\lambda_{n_N} = \lambda'_N \circ$ (this isomorphism).

By (2.6.131) (as $I'$ is cofinal in $I$), the canonical morphism of cones $\lim_{\leftarrow i\in I} F(i) \to \lim_{\leftarrow i\in I'} F(i)$ is an isomorphism, and by construction (restriction of the limit cone's own projections $\lambda_n$ to $i = n_N \in I'$) it sends $(m_n)_{n\geq 0} \to \left(m_{n_N}\right)_{N\geq 0}$ under the identification of the previous paragraph. Composing the two isomorphisms of this and the previous paragraph gives the stated isomorphism $\lim_{\leftarrow n} M_n \xrightarrow{\sim} \lim_{\leftarrow N} M_{n_N}$, $(m_n)_n \to \left(m_{n_N}\right)_N$. ∎

© David Rufino CC BY-NC-ND 4.0

# 3.4 Tensor Products

## 3.4.1 Commutative Tensor Product

We consider the "Tensor Product" of modules over a single commutative ring as an important special case, and as a guide to the general bimodule case.

In order to streamline the (typically tedious) verification of standard properties we make heavy use of the notion of **representable bifunctor** (2.6.65) however all the results may be proved in an essentially identical "low-tech" way.

First we define the so-called "Internal Hom Functor"

**Definition 3.4.1 (Internal Hom)**
*Let $A$ be a commutative ring and $M, N$ be $A$-modules. The set of $A$-linear homomorphisms $M \to N$ is itself an $A$-module. Therefore we have an enriched hom functor*

$$\mathrm{Hom} : A\text{-}\mathbf{Mod}^{op} \times A\text{-}\mathbf{Mod} \to A\text{-}\mathbf{Mod}$$

*where*

$$(a \cdot \psi)(m) \coloneqq a \cdot \psi(m) = \psi(a \cdot m)$$

*and in particular we have a bijection*

$$\mathrm{Mor}(M, N) \xrightarrow{\sim} \mathrm{Nat}(\mathrm{Hom}(N, -), \mathrm{Hom}(M, -))$$

**Definition 3.4.2 (Bilinear Map)**
*Let $M, N, Z$ be $A$-modules. Then a map*

$$\psi : M \times N \to Z$$

*is **bilinear** if it satisfies*

- ***additive*** $\psi(m + m', n) = \psi(m, n) + \psi(m', n)$ *and* $\psi(m, n + n') = \psi(m, n) + \psi(m, n')$
- ***$A$-linear*** $\psi(am, n) = \psi(m, an) = a\psi(m, n)$ *for all* $a \in A$

*This yields a bifunctor which we denote by*

$$\mathrm{Bilin}(-, -; -) : (A\text{-}\mathbf{Mod} \times A\text{-}\mathbf{Mod})^{op} \times A\text{-}\mathbf{Mod} \to A\text{-}\mathbf{Mod}$$

**Remark 3.4.3 ($A$-linearity)**
*The $A$-module structure on the image of* Hom *and* Bilin *is only well-defined because $A$ is commutative. For given $\psi : M \to N$ an $A$-linear map then for $(a \cdot \psi)$ to be $A$-linear we may verify this requires*

$$a'a\psi(m) = (a\psi)(a'm) = aa'\psi(m)$$

*which in general does not hold unless $A$ is commutative.*

**Proposition 3.4.4 (Existence of Tensor Product)**
*The bifunctor* $\mathrm{Bilin}(-, -; -)$ *is representable by a functor*

$$\otimes_A : A\text{-}\mathbf{Mod} \times A\text{-}\mathbf{Mod} \to A\text{-}\mathbf{Mod}$$

*for which there exists an isomorphism natural in $M, N, Z$*

$$\Phi : \mathrm{Hom}(M \otimes N, Z) \cong \mathrm{Bilin}(M, N; Z)$$

*For every pair $M, N$ there is a distinguished bilinear map*

$$i : M \times N \to M \otimes N$$

*through which every bilinear map $M \times N \to Z$ factors uniquely. We denote the **elementary tensor** by*

$$m \otimes n \coloneqq i(m, n)$$

*Further for morphisms $f : M \to M'$ and $g : N \to N'$ there is a unique morphism*

$$(f \otimes g) : M \otimes N \to M' \otimes N'$$

*such that*

© David Rufino CC BY-NC-ND 4.0

$$(f \otimes g)(m \otimes n) = f(m) \otimes g(n)$$

*Proof.* In light of (2.6.66) it is sufficient to show that $\text{Bilin}(M, N; -)$ is representable for fixed $M, N$ and the rest of the properties follow, observing that $i = \Phi(1_{M \otimes N})$.

For let $F$ be the free abelian group on $M \times N$ and let $S \subset F$ be the subgroup generated by elements of the form

- $(m + m', n) - (m, n) - (m', n)$
- $(m, n' + n) - (m, n') - (m, n)$
- $(am, n) - a(m, n)$
- $(m, n) - (m, an)$

Observe there is a canonical map $M \times N \to F$. Suppose that $Z$ is an abelian group and $\psi : M \times N \to Z$ is a bilinear map. Then we may extend by linearity to a group homomorphism $\psi : F \to Z$. By construction $S \subseteq \ker(\psi)$ whence there is a unique group homomorphism $\hat{\psi} : F/S \to Z$ such that $\hat{\psi} \circ i = \psi$ where $i : M \times N \to F/S$ is the canonical inclusion. This shows that $(F/S, i)$ is a universal element for $\text{Bilin}(-, -; -)$ as we required and we denote this by $M \otimes N$. The representation exists as a functor of sets by (2.6.62), and $\Phi$ is clearly an isomorphism of $A$-modules because

$$\Phi(f + g) = (f + g) \circ i = f \circ i + g \circ i = \Phi(f) + \Phi(g)$$

and

$$\Phi(\lambda f) = (\lambda f) \circ i = \lambda(f \circ i) = \lambda \Phi(f)$$

■

**Lemma 3.4.5 (Currying Lemma)**
*Let $M, N, Z$ be $A$-modules then there are natural isomorphisms*

$$\begin{array}{ccccccc} \text{Bilin}(M, N, Z) & \cong & \text{Hom}(M, \text{Hom}(N, Z)) & \cong & \text{Hom}(N, \text{Hom}(M, Z)) & \cong & \text{Bilin}(N, M, Z) \\ \psi & & m \to (n \to \psi(m, n)) & & n \to (m \to \psi(m, n)) & & (n, m) \to \psi(m, n) \end{array}$$

This gives the following important result (which may be seen as an adjoint relationship)

**Proposition 3.4.6 (Tensor-Hom Adjunction)**
*Let $M, N, Z$ be $A$-modules then we have the following natural isomorphism of functors*

$$\begin{array}{ccc} \text{Hom}(M \otimes N, Z) & \cong & \text{Hom}(M, \text{Hom}(N, Z)) \\ & & \theta \to (m \to (n \to \theta(m \otimes n))) \end{array}$$

We may also use the Currying Lemma (3.4.5) to demonstrate symmetry of the tensor product

**Proposition 3.4.7 (Symmetry of Tensor Product)**
*There is a natural isomorphism*

$$\begin{array}{ccc} M \otimes N & \cong & N \otimes M \\ m \otimes n & \to & n \otimes m \end{array}$$

*Proof.* We observe that there is a natural isomorphism of functors which the tensor products represent

$$\text{Bilin}(M, N; Z) \cong \text{Bilin}(N, M; Z)$$

so the result follows from (2.6.68). ■

**Proposition 3.4.8 (Associativity of Tensor Product)**
*There is a natural isomorphism of $A$-modules*

$$\begin{array}{ccc} (M \otimes N) \otimes P & \cong & M \otimes (N \otimes P) \\ (m \otimes n) \otimes p & \to & m \otimes (n \otimes p) \end{array}$$

*Proof.* We may make repeated use of the Tensor-Hom adjunction to exhibit natural isomorphisms

© David Rufino CC BY-NC-ND 4.0

$$\begin{aligned}\mathrm{Hom}((M\otimes N)\otimes P, Z) &\cong \mathrm{Hom}(M\otimes N, \mathrm{Hom}(P,Z))\\ &\cong \mathrm{Hom}(M, \mathrm{Hom}(N, \mathrm{Hom}(P,Z)))\\ &\cong \mathrm{Hom}(M, \mathrm{Hom}(N\otimes P, Z))\\ &\cong \mathrm{Hom}(M\otimes(N\otimes P), Z)\end{aligned}$$

The required isomorphism is then a consequence of (2.6.68). For explicit form set $Z = M\otimes(N\otimes P)$ and consider the identity map $1_Z$. We see tracing back under these natural isomorphisms this corresponds to the given map. ■

### 3.4.2 Bimodule Tensor Product

To develop the tensor product in general it is convenient to work with bimodules. We make heavy use of the notion if bimodule hom-sets (3.3.79).

**Definition 3.4.9 (Bilinear maps)**
*Consider the bimodules ${}_AM_B$, ${}_BN_C$. We say that a map*

$$\psi : {}_AM_B \times {}_BN_C \to {}_AZ_C$$

*is $(A,C)$-**bilinear** if it satisfies all of the following conditions*

- ***additive*** $\psi(m+m', n+n') = \psi(m,n)+\psi(m',n)+\psi(m,n')+\psi(n,n')$
- ***balanced*** $\psi(mb,n) = \psi(m,bn)$
- ***A-linear*** $\psi(am,n) = a\psi(m,n)$
- ***C-linear*** $\psi(m,nc) = \psi(m,n)c$

*Extending the notation from earlier we denote the set of additive, balanced and $(A,C)$-bilinear maps by* $\mathrm{Bilin}(M,N;Z)$. *This determines a functor*

$$\mathrm{Bilin} : ({}_A\mathbf{Mod}_B \times {}_B\mathbf{Mod}_C)^{op} \times {}_A\mathbf{Mod}_C \to \mathbf{AbGrp}$$

*where addition is defined pointwise. We may also consider the functor of only additive and balanced maps*

$$\mathrm{Balan} : (\mathbf{Mod}_B \times {}_B\mathbf{Mod})^{op} \times \mathbf{AbGrp} \to \mathbf{AbGrp}$$

*and the functor*

$$\mathrm{RBilin} : ({}_B\mathbf{Mod}_C \times {}_C\mathbf{Mod}_D)^{op} \times {}_A\mathbf{Mod}_D \to {}_A\mathbf{Mod}_B$$

*where we drop the $B$-linear requirement but also introduce an extra $(A,B)$-module structure on the maps, as follows*

$$(a\psi b)(m,n) \coloneqq a\psi(bm,n)$$

**Proposition 3.4.10 (Existence of Bimodule Tensor Product)**
*Let $A, B, C$ be arbitrary rings*

- *The bifunctor* $\mathrm{Balan}(M,N;-)$ *is represented by the **balanced tensor product***

  $$\otimes_B : \mathbf{Mod}_B \times {}_B\mathbf{Mod} \to \mathbf{AbGrp}$$

  *with natural isomorphism for $Z$ an Abelian group*

  $$\begin{aligned}\Phi : \mathrm{Hom}(M_B \otimes_B {}_BN, Z) &\cong \mathrm{Balan}(M_B, {}_BN; Z)\\ \theta &\to \theta\circ i\end{aligned}$$

  *where $i$ is a universal balanced map*

  $$i : M\times N \to M_B\otimes_B {}_BN$$

  *The tensor product is generated by **elementary tensors** of the form $m\otimes n \coloneqq i(m,n)$.*

- *The bifunctor* $\mathrm{Bilin}(M,N;-)$ *is represented by the balanced tensor product $M\otimes_B N$ with an additional $(A,C)$-bimodule structure given by*

  $$a\cdot(m\otimes n)\cdot c \coloneqq (a\cdot m)\otimes(n\cdot c)$$

  *This determines a bifunctor*

  $$\otimes_B : {}_A\mathbf{Mod}_B \times {}_B\mathbf{Mod}_C \to {}_A\mathbf{Mod}_C$$

© David Rufino CC BY-NC-ND 4.0

*and a natural isomorphism of Abelian groups*

$$\begin{aligned}\mathrm{Hom}({}_AM_B \otimes_B {}_BN_C, Z) &\cong \mathrm{Bilin}({}_AM_B, {}_BN_C; {}_AZ_C)\\ \theta &\to \theta \circ i\end{aligned}$$

*In either case, if $f : M \to M'$ and $g : N \to N'$ are morphisms then there is a unique morphism*

$$(f \otimes g) : (M \otimes N) \to (M' \otimes N')$$

*such that*

$$(f \otimes g)(m \otimes n) = f(m) \otimes g(n)$$

*Furthermore for maps $f' : M' \to M''$ and $g' : N' \to N''$ the following property is satisfised*

$$(f' \otimes g') \circ (f \otimes g) = (f' \circ f) \otimes (g' \circ g)$$

*Proof.* In light of (2.6.65) we only need to show that the functors are representable for fixed $M, N$, as the functoriality will follow immediately by abstract nonsense.

We first show that $\mathrm{Balan}(-,-;-)$ is representable. For let $F$ be the free abelian group on $M \times N$ and let $S \subset F$ be the subgroup generated by elements of the form

- $(m + m', n) - (m, n) - (m', n)$
- $(m, n' + n) - (m, n') - (m, n)$
- $(mb, n) - (m, bn)$

Observe there is a canonical map $M \times N \to F$. Suppose that $Z$ is an abelian group and $\psi : M \times N \to Z$ is a balanced additive map. Then we may extend by linearity to a group homomorphism $\psi : F \to Z$. By construction $S \subseteq \ker(\psi)$ whence there is a unique group homomorphism $\hat{\psi} : F/S \to Z$ such that $\hat{\psi} \circ i = \psi$ where $i : M \times N \to F/S$ is the canonical inclusion. This shows that $(F/S, i)$ is a universal element for $\mathrm{Balan}(-,-;-)$ and we denote this by $M_B \otimes_B {}_BN$. Then (2.6.62) exhibits $\Phi$ as a natural isomorphism of sets, which may readily observed to be additive.

In order to define the $(A, C)$-bimodule structure in the second case let $\psi_a : M \to M$ denote multiplication by $a$ and $\psi_c : N \to N$ multiplication by $c$. We note these are both $B$-module homomorphisms by definition so we may define the action by functoriality of the balanced tensor product case

$$\begin{aligned}a \cdot v &:= (\psi_a \otimes 1_N)(v) \quad \forall v \in M \otimes N\\ v \cdot c &:= (1_M \otimes \psi_c)(v) \quad \forall v \in M \otimes N\end{aligned}$$

These are $B$-linear by construction and we may demonstrate $(A, C)$-bimodule actions are associative and commute by using the functoriality of the tensor product

$$\begin{aligned}(\psi_{a'} \otimes 1_N) \circ (\psi_a \otimes 1_N) = (\psi_{a'} \circ \psi_a) \otimes 1_N &= (\psi_{a'a}) \otimes 1_N\\ (1_M \otimes \psi_c) \circ (1_M \otimes \psi_{c'}) = 1_M \otimes (\psi_c \circ \psi_{c'}) &= 1_M \otimes \psi_{cc'}\\ (\psi_a \otimes 1_N) \circ (1_M \otimes \psi_c) = \psi_a \otimes \psi_c &= (1_M \otimes \psi_c) \otimes (\psi_a \otimes 1_N)\end{aligned}$$

Further by construction the canonical map $i$ is $(A, C)$-bilinear. Consider any $(A, C)$-bilinear map $\psi : M \times N \to Z$ then by the previous part there is a unique group homomorphism $\hat{\psi} : M \otimes_B N \to Z$ such that

$$\hat{\psi}(m \otimes n) := \psi(m, n)$$

which we see by construction (and linearity) is $(A, C)$-bilinear. It is clearly the unique such map so that we may conclude $\mathrm{Bilin}(-,-;-)$ has universal element $({}_AM_B \otimes_B {}_BN_C, i)$. The (2.6.62) exhibits $\Phi$ as a natural isomorphism of sets which may be readily exhibited to be additive. ∎

**Lemma 3.4.11 (Currying Lemma)**
*Let $M, N, P, Y, Z$ be bimodules. Then there is an isomorphism of abelian groups and $(A, B)$-bimodules respectively*

$$\begin{aligned}\mathrm{Bilin}({}_AM_B, {}_BN_C; {}_AY_C) &\cong \mathrm{Hom}({}_AM_B, \mathrm{RHom}({}_BN_C, {}_AY_C))\\ &\cong \mathrm{Hom}({}_BN_C, \mathrm{LHom}({}_AM_B, {}_AY_C))\\ \mathrm{RBilin}({}_BN_C, {}_CP_D; {}_AZ_D) &\cong \mathrm{RHom}({}_BN_C, \mathrm{RHom}({}_CP_D, {}_AZ_D))\end{aligned}$$

*contravariant in $M, N, P$ and covariant in $Y$ and $Z$.*

© David Rufino CC BY-NC-ND 4.0

*Proof.* The maps on abelian groups are clear, namely given a bi-additive map $\psi$ we have a well-defined homomorphism of abelian groups

$$m \to (n \to \psi(m,n))$$

The verification that the maps are well-defined and bijective is tedious but mechanical. ∎

**Proposition 3.4.12 (Tensor-Hom Adjunction)**
*Let $M, N, P, Y, Z$ be bimodules then*

$$\begin{aligned} \mathrm{Hom}({}_AM_B \otimes {}_BN_C, {}_AY_C) &\cong \mathrm{Hom}({}_AM_B, \mathrm{RHom}({}_BN_C, {}_AY_C)) \\ &\cong \mathrm{Hom}({}_BN_C, \mathrm{LHom}({}_AM_B, {}_AY_C)) \\ \mathrm{RHom}({}_BN_C \otimes {}_CP_D, {}_AZ_D) &\cong \mathrm{RHom}({}_BN_C, \mathrm{RHom}({}_CP_D, {}_AZ_D)) \end{aligned}$$

*where the first is an isomorphism of Abelian Groups and the second of $(A,B)$-bimodules. Further the isomorphisms are contravariant in $M, N, P$ and covariant in $Y$ and $Z$.*

*Proof.* This first two isomorphism follow directly from the Currying Lemma and universal property for bimodule tensor product.

The final isomorphism follows from the Currying Lemma if we show that there is an isomorphism

$$\begin{array}{ccc} \mathrm{RHom}({}_BN_C \otimes {}_CP_D, {}_AZ_D) & \xrightarrow{\sim} & \mathrm{RBilin}({}_BN_C, {}_CP_D; {}_AZ_D) \\ \downarrow & & \downarrow \\ \mathrm{Hom}(N_C \otimes {}_CP, Z) & \xrightarrow{\sim} & \mathrm{Balan}(N_C, {}_CP; Z) \end{array}$$

where the bottom is simply the universal property of the balanced tensor product and the top we require to also be $(A,B)$-bilinear. Recall the underlying sets are the same and the horizontal maps are given by $\theta \to \theta \circ i$ in both cases. It's clear that the map is well-defined and injective. Further it's surjective because if $\theta$ is $(B,D)$-bilinear on elementary tensors it is $(B,D)$-bilinear everywhere. Finally we need to show that the isomorphism is $(A,B)$-bilinear as follows

$$(a(\theta \circ i)b)(n,p) = a(\theta \circ i)(bn,p) = a\theta((bn) \otimes p) = a\theta(b(n \otimes p)) = (a\theta b)(n \otimes p)$$

∎

**Proposition 3.4.13 (Associativity of Tensor Product)**
*Let ${}_AM_B$, ${}_BN_C$, ${}_CP_D$ be bimodules then there is a natural isomorphism of $(A,D)$-bimodules*

$$\begin{aligned} (M \otimes_B N) \otimes_C P &\cong M \otimes_B (N \otimes_C P) \\ (m \otimes n) \otimes p &\to m \otimes (n \otimes p) \end{aligned}$$

*Proof.* This follows much as in the commutative case. Let $Z$ be an $(A,D)$-bimodule. Then by the Tensor-Hom adjunction there are natural isomorphisms of Abelian groups

$$\begin{aligned} \mathrm{Hom}(({}_AM_B \otimes {}_BN_C) \otimes {}_CP_D, {}_AZ_D) &\cong \mathrm{Hom}({}_AM_B \otimes {}_BN_C, \mathrm{RHom}({}_CP_D, {}_AZ_D)) \\ &\cong \mathrm{Hom}({}_AM_B, \mathrm{RHom}({}_BN_C, \mathrm{RHom}({}_CP_D, {}_AZ_D))) \\ &\cong \mathrm{Hom}({}_AM_B, \mathrm{RHom}({}_BN_C \otimes_C {}_CP_D, {}_AZ_D)) \\ &\cong \mathrm{Hom}({}_AM_B \otimes_B ({}_BN_C \otimes_C {}_CP_D), {}_AZ_D) \end{aligned}$$

The required isomorphism is then a consequence of (2.6.68). For explicit form set $Z = M \otimes (N \otimes P)$ and consider the identity map $1_Z$. We see tracing back under these natural isomorphisms this corresponds to the given map. ∎

### 3.4.3 Extensions of Scalars

We observe that for a ring homomorphism $\varphi : A \to B$ we have on $B$ a natural $(A,B)$-bimodule and $(B,A)$-bimodule structure. Using the tensor product then we may use this to extend the coefficients from $A$ to $B$. We first prove some elementary lemmas

**Lemma 3.4.14 (Unit Hom)**
*Let $\varphi : B \to C$ be a ring homomorphism and $Z$ a $(A,C)$-bimodule. Then there is a natural isomorphism of $(A,B)$-bimodules*

© David Rufino CC BY-NC-ND 4.0

$$\begin{aligned}\mathrm{RHom}({}_BC_C, {}_AZ_C) &\cong {}_AZ_B\\ \theta &\to \theta(1)\\ z(-) &\leftarrow z\end{aligned}$$

*and similarly let $\varphi : A \to B$ be a ring homomorphism and $Y$ a $(B,C)$-bimodule then there is a natural isomorphism of $(A,C)$-bimodules*

$$\begin{aligned}\mathrm{LHom}({}_BB_A, {}_BY_C) &\cong {}_AY_C\\ \theta &\to \theta(1)\end{aligned}$$

*Proof.* Observe

$$(a\theta b)(1) = a\theta(\varphi(b)) = a\theta(1)\varphi(b)$$

so the left to right map is an $(A,B)$-bimodule homomorphism. Further $\theta(c) = \theta(1)c$ so the maps are mutual inverses.■

**Lemma 3.4.15 (Unit Hom (Commutative Ring Case))**
*Let $\varphi : A \to B$ be a homomorphism of commutative rings and $Z$ a $B$-module. Then there is a natural isomorphism of $B$-modules*

$$\begin{aligned}\mathrm{Hom}(B, Z) &\cong Z\\ \theta &\to \theta(1)\end{aligned}$$

**Proposition 3.4.16 (Extension of Scalars)**
*Let $M$ be an $(A,C)$-bimodule and $\varphi : A \to B$ a ring homomorphism. We may define a $(B,C)$-module*

$$M_{(B)} \coloneqq {}_BB_A \otimes_A M$$

*For $Z$ a $(B,C)$-bimodule there is a natural isomorphism of abelian groups*

$$\begin{aligned}\mathrm{Hom}\Big(M_{(B)}, Z\Big) &\xrightarrow{\sim} \mathrm{Hom}(M, {}_AZ_C)\\ \psi &\to m \to \psi(1 \otimes m)\\ b \otimes m \to b \cdot \phi(m) &\leftarrow \phi\end{aligned}$$

*In otherwords we have an adjunction (2.6.69)*

$${}_A\mathbf{Mod}_C \longleftrightarrow {}_B\mathbf{Mod}_C$$

*Setting $C = \mathbb{Z}$ yields an adjunction for left modules*

$${}_A\mathbf{Mod} \longleftrightarrow {}_B\mathbf{Mod}$$

*In particular for $M$ a left $A$-module we have an isomorphism of abelian groups*

$$\mathrm{Hom}\Big(M_{(B)}, B\Big) \xrightarrow{\sim} \mathrm{Hom}_A(M, B)$$

*which is $B$-linear when $B$ is commutative.*

*Proof.* By the tensor-hom adjunction (3.4.12) and (3.4.14) there is a natural isomorphism

$$\mathrm{Hom}(B \otimes_A M, Z) \cong \mathrm{Hom}(M, \mathrm{LHom}({}_BB_A, {}_BZ_C)) \cong \mathrm{Hom}(M, {}_AZ_C)$$

■

**Proposition 3.4.17**
*Let $\varphi : A \to B$ be a homomormorphism of commutative rings and $\varphi : M \to N$ a homomorphism of $A$-modules. Then there is a unique $B$-module homomorphism $\varphi_{(B)} : M_{(B)} \to N_{(B)}$ such that*

$$\varphi_{(B)}(1 \otimes m) = 1 \otimes \varphi(m)$$

*Proof.* The homomorphism $\varphi_{(B)} = 1_B \otimes \psi$ exists by functoriality of the tensor product (3.4.4). It clearly satisfies the given property, and as the tensor product is generated by elementary tensors it is evidentally unique. ■

**Proposition 3.4.18 (Tensor Unit)**
*Let $M$ be an $(A,C)$-bimodule then there is a natural isomorphism of $(A,C)$-bimodules*

© David Rufino CC BY-NC-ND 4.0

$$\begin{aligned} A \otimes_A M &\cong M \\ a \otimes m &\to a \cdot m \\ 1 \otimes m &\leftarrow m \end{aligned}$$

*Similarly there is a natural isomorphism of $(A, C)$-bimodules*

$$M \otimes_C C \cong M$$

*Proof.* By (3.4.16) with $\phi = 1_A$ there is a natural isomorphism

$$\mathrm{Hom}(A \otimes_A M, -) \cong \mathrm{Hom}(M, -)$$

and so natural isomorphism follows from (2.6.68).

We may deduce that there is a natural isomorphism of $(C^{\mathsf{op}}, A^{\mathsf{op}})$-bimodules

$$C^{\mathsf{op}} \otimes_{C^{\mathsf{op}}} M^{\mathsf{op}} \cong M^{\mathsf{op}}$$

which amounts to a natural isomorphism of $(A, C)$-bimodules

$$M \otimes_C C \cong M$$

■

**Proposition 3.4.19 (Transitivity of Extension of Scalars)**
*Let $M$ be a $(A, D)$-bimodule and $\varphi : A \to B$, $\psi : B \to C$ homomorphisms of commutative rings. Then there is an isomorphism of $(C, D)$-bimodules*

$$\begin{aligned} C \otimes_B (B \otimes_A M) &\xrightarrow{\sim} {}_C C_A \otimes_A M \\ c \otimes (b \otimes m) &\to (c\psi(b)) \otimes m \end{aligned}$$

*Proof.* By associativity (3.4.13) and (3.4.18) there is a natural isomorphism

$$C \otimes_B (B \otimes_A M) \cong (C \otimes_B B) \otimes_A M \cong C \otimes_A M$$

■

**Proposition 3.4.20 (Transitivity of Extension of Scalars (Commutative Case))**
*Let $M$ be an $A$-module and $\phi : A \to B$ and $\psi : B \to C$ homomorphisms of commutative rings. Then there is an isomorphism of $C$-modules*

$$C \otimes_B (B \otimes_A M) \cong {}_C C_A \otimes_A M$$

*Proof.* Regarding $M$ as an $(A, A)$-bimodule we have an isomorphism of $(C, A)$-bimodules by (3.4.19) which is a-fortiori a $C$-module isomorphism. ■

## 3.4.4 Tensor Product Commutes with Direct Sum

**Proposition 3.4.21 (Tensor Product Commutes with Sum)**
*Let $(M_i)_{i \in I}$ be a family of $(A, B)$-bimodules and $(N_j)_{j \in J}$ a family of $(B, C)$-bimodules. Then there is an isomorphism of $(A, C)$-bimodules*

$$\left( \bigoplus_{i \in I} M_i \right) \otimes_B \left( \bigoplus_{j \in J} N_j \right) \cong \bigoplus_{(i,j) \in I \times J} (M_i \otimes_B N_j)$$

**Corollary 3.4.22**
*Suppose $M$ is an $(A, B)$-bimodule and $N$ is a $(B, C)$-bimodule and $M' \subseteq M$ and $N' \subseteq N$ are direct factors. Then the canonical map is injective*

$$M' \otimes_B N' \hookrightarrow M \otimes_B N$$

**Corollary 3.4.23**
*Let $N$ be **free** left $A$-module with basis $(n_i)_{i \in I}$ and $M$ a $(B, A)$-bimodule. Then there is an isomorphism of left $B$-modules*

© David Rufino CC BY-NC-ND 4.0

$$M \otimes_A N \cong \bigoplus_{i \in I} M$$
$$m \otimes \sum_i a_i n_i \to (m \cdot a_i)_{i \in I}$$
$$\sum_i (m_i \otimes n_i) \leftarrow (m_i)_{i \in I}$$

*When $A$ is commutative then this is a $(B, A)$-bimodule isomorphism. Further when $M$ is an $A$-module then this is an isomorphism of $A$-modules.*

*Proof.* By (3.4.21) and (3.4.18)

$$M \otimes_A N \cong M \otimes_A \left( \bigoplus_{i \in I} A \right) \cong \bigoplus_{i \in I} (M \otimes_A A) \cong \bigoplus_{i \in I} M$$
$$m \otimes \sum_i a_i n_i \to \quad m \otimes (a_i)_{i \in I} \to \quad (m \otimes a_i)_{i \in I} \to (m \cdot a_i)_{i \in I}$$

■

**Corollary 3.4.24**
*If $M, N$ are free $A$-modules with bases $\{m_i\}_{i \in I}$ and $\{n_j\}_{j \in J}$ then $M \otimes_A N$ is a free $A$-module with basis $\{m_i \otimes n_j\}_{(i,j) \in I \times J}$.*

**Corollary 3.4.25 (Free modules are flat)**
*Let $N$ be a free left $A$-module and $i : M' \to M$ an injective map of $(B, A)$-bimodules. Then the corresponding map*

$$i \otimes 1_N : M' \otimes_A N \to M \otimes_A N$$

*is injective.*

**Corollary 3.4.26 (Extension of Scalars (Free Module))**
*Let $M$ be **free** left $A$-module with basis $\{m_i\}_{i \in I}$ and $\varphi : A \to B$ a ring homomorphism. Then there is a canonical isomorphism of left $B$-modules*

$$M_{(B)} \cong \bigoplus_{i \in I} B$$
$$b \otimes \sum_i a_i m_i \to (b\varphi(a_i))_{i \in I}$$

*In particular $\{1 \otimes m_i\}_{i \in I}$ is a basis for $M_{(B)}$. Further when $Z$ is a left $B$-module then there is an isomorpihsm of abelian groups*

$$\begin{array}{ccc} \mathrm{Hom}_A(M, Z) & \cong & \mathrm{Hom}_B\left(M_{(B)}, Z\right) \cong \prod_{i \in I} Z \\ \theta & & \to \quad (\theta(m_i))_{i \in I} \end{array}$$

*which is $B$-linear when $B$ is commutative.*

*When $M$ is a finite free $A$-module of rank $n$ (with basis $\{v_i\}$) then $M_{(B)}$ is a finite free $B$-module of rank $n$ (with basis $\{1 \otimes v_i\}$). When $B$ is commutative, $M_{(B)}^\vee$ is also a finite free $B$-module of rank $n$.*

*Proof.* The first isomorphism follows directly from (3.4.23). Then by (3.3.83) there is an isomorphism

$$\mathrm{Hom}\left(M_{(B)}, Z\right) \cong \mathrm{Hom}\left( \bigoplus_{i \in I} B, Z \right) \cong \prod_{i \in I} \mathrm{Hom}(B, Z) \cong \prod_{i \in I} Z$$

■

We may generalize (3.3.110)

**Proposition 3.4.27 (Extension of Scalars (Hom-Set))**
*Let $\varphi : A \to B$ be a homomorphism of commutative rings, $M$ a finite-free $A$-module and $N$ a finite-free $B$-module. Then*

$$\mathrm{Hom}_A(M, N) \cong \mathrm{Hom}_B\left(M_{(B)}, N\right)$$

© David Rufino CC BY-NC-ND 4.0

*is a finite-free $B$-module. More precisely suppose $M$ has basis $\{v_1, ..., v_n\}$ and $N$ has basis $\{w_1, ..., w_m\}$ then* $\mathrm{Hom}_A(M,N)$ *has basis*

$$\left\{w_j v_i^\vee\right\}_{i,j}$$

*Proof.* By (3.4.26) there are isomorphisms of $B$-modules

$$\begin{aligned}\mathrm{Hom}_A(M,N) \cong \mathrm{Hom}_B\left(M_{(B)}, N\right) \cong \bigoplus_{i=1}^{n} N \cong \bigoplus_{i=1}^{n}\bigoplus_{j=1}^{m} B\\ \theta \to \qquad 1\otimes\theta \to (\theta(v_i))_i \to \left(w_j^\vee(\theta(v_i))\right)_{i,j}\end{aligned}$$

Under this isomorphism the standard basis of the right hand side corresponds to the set $\left\{w_j v_i^\vee\right\}_{i,j}$ whence these constitute a basis. ∎

## 3.4.5 Tensor Product Exact Sequences

**Proposition 3.4.28 (Tensor Product is Right Exact)**
*Consider an exact sequence of $A$-modules (i.e. such that $\beta$ is surjective and* $\ker(\beta) = \mathrm{Im}(\alpha)$*)*

$$M \xrightarrow{\alpha} N \xrightarrow{\beta} P \to 0$$

*Then the corresponding sequence of $A$-modules*

$$M \otimes_A Z \xrightarrow{\alpha\otimes 1_Z} N \otimes_A Z \xrightarrow{\beta\otimes 1_Z} P \otimes_A Z \to 0$$

*is exact. Similarly the sequence*

$$Z \otimes_A M \xrightarrow{1_Z\otimes\alpha} Z \otimes_A N \xrightarrow{1_Z\otimes\beta} Z \otimes_A P \to 0$$

*is exact.*

*Proof.* Trivially the submodule $\mathrm{Im}(\beta\otimes 1_Z)$ contains elementary tensors and therefore equals $P\otimes_A Z$. In otherwords $\beta\otimes 1_Z$ is surjective. Furthermore by functoriality $(\beta\otimes 1_Z)\circ(\alpha\otimes 1_Z) = (\beta\circ\alpha)\otimes 1_Z$ which is zero on elementary tensors and therefore zero everywhere. In particular $\mathrm{Im}(\alpha\otimes 1_Z) \subseteq \ker(\beta\otimes 1_Z)$. Consider a commutative diagram

$$\begin{array}{ccc} P\times Z & \xrightarrow{\psi} & (N\otimes_A Z)/\mathrm{Im}(\alpha\otimes 1_Z) \\ & \searrow & \downarrow{\scriptstyle \overline{\beta\otimes 1_Z}} \\ & & P\otimes_A Z \end{array}$$

where $\psi$ is yet to be defined. For a given $p\in P$ and $z\in Z$ consider the subset

$$\mathcal{I}_{p,z} \coloneqq \{n\otimes z \mid n\in N, \beta(n) = p\} \subset N\otimes_A Z$$

Observe that any $\psi$ making the diagram commute satisfies $\psi(p,z)\in\overline{\mathcal{I}_{p,z}}$, and we aim to show this set is a singleton.

As $\beta$ is surjective $\mathcal{I}_{p,z}$ is non-empty. First we show that $x,y\in\mathcal{I}_{p,z} \implies x-y\in\mathrm{Im}(\alpha\otimes 1_Z)$. For if $\beta(n)=\beta(n')$ then $\beta(n-n')=0$ by definition $n-n'=\alpha(m)$ for some $m\in M$. It follows that $n\otimes z - n'\otimes z = \alpha(m)\otimes z = (\alpha\otimes 1_Z)(m\otimes z)$. This shows that $\overline{\mathcal{I}_{p,z}}$ is a singleton, whose element we define to be $\psi(p,z)$. Then $\psi$ is the unique map such that

$$\psi(\beta(n), z) = \overline{n\otimes z}$$

for all $n\in N$ and $z\in Z$. Clearly $\psi$ is $A$-bilinear and therefore there is a map $\sigma : P\otimes_A Z \to (N\otimes_A Z)/\mathrm{Im}(\alpha\otimes 1_Z)$ such that $\sigma(\beta(n)\otimes z) = \overline{n\otimes z}$. By linearity $\sigma\circ\overline{\beta\otimes 1_Z}$ is the identity which shows $\overline{\beta\otimes 1_Z}$ is injective. Therefore $\ker(\beta\otimes 1_Z) = \mathrm{Im}(\alpha\otimes 1_Z)$ as required (for $x\in\ker(\beta\otimes 1_Z) \implies (\beta\otimes 1_Z)(x) = 0 \implies \left(\overline{\beta\otimes 1_Z}\right)(\overline{x}) = 0 \implies \overline{x} = 0 \implies x\in\mathrm{Im}(\alpha\otimes 1_Z)$). ∎

**Proposition 3.4.29 (Quotient of a Tensor Product)**
*Consider exact sequences*

$$0 \to M' \xrightarrow{i} M \xrightarrow{\pi_1} M/M' \to 0$$

© David Rufino CC BY-NC-ND 4.0

$$0 \to N' \xrightarrow{j} N \xrightarrow{\pi_2} N/N' \to 0$$

*Then there is an exact sequence*

$$\begin{array}{rccc} (M' \otimes N) \oplus (M \otimes N') & \xrightarrow{\alpha} & M \otimes N & \xrightarrow{\pi_1 \otimes \pi_2} M/M' \otimes N/N' \to 0 \\ (m' \otimes n, m \otimes n') & \to & i(m') \otimes n + m \otimes j(n') & \to \quad \overline{m} \otimes \overline{n} \to 0 \end{array}$$

*In particular there is a canonical isomorphism*

$$(M \otimes N)/(M' \otimes N + M \otimes N') \cong M/M' \otimes N/N'$$

*Proof.* Observe $\pi_1 \otimes \pi_2 = \left(\pi_1 \otimes 1_{N/N'}\right) \circ (1_M \otimes \pi_2)$ by comparison on elementary tensors. By (3.4.28) each of these maps is surjective, and so the composite is surjective. Evidently $(\pi_1 \otimes \pi_2) \circ (i \otimes 1_N) = 0$ and $(\pi_1 \otimes \pi_2) \circ (1_M \otimes j) = 0$ by checking elementary tensors. This in particular means that $(\pi_1 \otimes \pi_2) \circ \alpha = 0$ where $\alpha$ is the first map. Therefore $\mathrm{Im}(\alpha) \subseteq \ker(\pi_1 \otimes \pi_2)$ and it suffices to demonstrate the reverse inclusion. Suppose $z \in \ker(\pi_1 \otimes \pi_2)$. This means precisely that $(1_M \otimes \pi_2)(z) \in \ker\left(\pi_1 \otimes 1_{N/N'}\right) = \mathrm{Im}\left(i \otimes 1_{N/N'}\right)$ by (3.4.28). Therefore $(1_M \otimes \pi_2)(z) = \left(i \otimes 1_{N/N'}\right)(y)$ for $y \in M' \otimes N/N'$. By (3.4.28) again $y = (1_{M'} \otimes \pi_2)(x)$ for $x \in M' \otimes N$ and $(1_M \otimes \pi_2)(z) = (i \otimes \pi_2)(x)$. Therefore $(1_M \otimes \pi_2)(z - (i \otimes 1_N)(x)) = 0$ and $z - (i \otimes 1_N)(x) = (1_M \otimes j)(w)$ for some $w \in M \otimes N'$. Therefore we conclude $z \in \mathrm{Im}(\alpha)$ as required. ∎

## 3.4.6 Vector Space Tensor Product

Recall that vector spaces are free, every linearly independent set may be extended to a basis and every subspace is a direct factor. This simplifies the structure of tensor product.

**Proposition 3.4.30**
*Let $V, W$ be $k$-vector spaces with subspaces $V'$ and $W'$. Then the canonical $k$-module homomorphism*

$$V' \otimes_k W' \to V \otimes_k W$$

*is injective.*

*Proof.* The homomorphism exists by (3.4.4). By (3.3.139) both $V'$ and $W'$ are direct factors. Therefore the map is injective by (3.4.22). ∎

**Proposition 3.4.31**
*Let $V, W$ be $k$-modules. Suppose $\{v_i\}_{i \in I}$ and $\{w_j\}_{j \in J}$ are linearly independent (resp. bases) then $\{v_i \otimes w_j\}_{(i,j) \in I \times J}$ is a $k$-linearly independent subset (resp. a $k$-basis) of $V \otimes_k W$.*

*Proof.* The case that they are bases is simply (3.4.24). For the general case we may use (3.3.126) to extend to bases, then a-fortiori the given set, being a subset of a basis, is linearly independent. ∎

## 3.4.7 Algebra Tensor Product

Let $A$ be a commutative ring and revert to the case of commutative tensor product described in (3.4.4). We show that given two $A$-algebras $B$, $C$ the tensor product naturally forms an algebra. We first prove some preliminary results.

**Lemma 3.4.32**
*Let $B$ be a commutative $A$-algebra. Then there exists a unique homomorphism of $A$-modules*

$$B \otimes_A B \to B$$

*such that*

$$b \otimes b' \to bb'$$

*Proof.* Multiplication is bilinear so the map exists by universal property. ∎

**Lemma 3.4.33**
*Let $B, C$ be commutative $A$-algebras and define the $A$-module $X \coloneqq B \otimes_A C$. Then there is a unique homomorphism of $A$-modules*

$$m : X \otimes_A X \to X$$

*such that*

$$(b \otimes c) \otimes (b' \otimes c') \to (bb') \otimes (cc')$$

© David Rufino CC BY-NC-ND 4.0

*Define* $1_X := (1 \otimes 1)$ *then it satisfies*

*a)* $m(1_X \otimes x) = m(x \otimes 1_X) = x$
*b)* $m(x \otimes y) = m(y \otimes x)$
*c)* $m(x \otimes m(y \otimes z)) = m(m(x \otimes y) \otimes z)$
*d)* $m((x + y) \otimes z) = m(x \otimes z) + m(y \otimes z)$
*e)* $m(x \otimes (y + z)) = m(x \otimes y) + m(x \otimes z)$

*for* $x, y, z \in X$.

*Proof.* By associativity and commutativity of the tensor product there is an $A$-module isomorphism

$$X \otimes_A X \cong (B \otimes_A B) \otimes_A (C \otimes_A C)$$
$$(b \otimes c) \otimes (b' \otimes c') \to (b \otimes b') \otimes (c \otimes c')$$

composing with $m_B \otimes m_C$ where $m_B : (B \otimes_A B) \to B$ and $m_C : (C \otimes_A C) \to C$ are the maps given in (3.4.32), yields the required map $m$.

The properties may be verified on tensors of the form $x_i = (b_i \otimes c_i)$ and $y_j = \left(b'_j \otimes c'_j\right)$. The results follow from linearity, since $\left(\sum_i x_i\right) \otimes \left(\sum_j y_j\right) = \sum_{i,j}(x_i \otimes y_j)$. ∎

**Proposition 3.4.34 (Algebra Tensor Product)**
*Let* $B, C$ *be commutative* $A$ *algebras. Then the* $A$*-module* $B \otimes_A C$ *has a unique commutative ring structure with unit* $1 \otimes 1$ *and multiplication which satisfies*

$$\cdot : (B \otimes_A C) \times (B \otimes_A C) \to B \otimes_A C$$
$$(b \otimes c) \cdot (b' \otimes c') := (bb') \otimes (cc')$$

*and may be extended by linearity. Further there is an ring homomorphism*

$$i_A : A \to B \otimes_A C$$
$$a \to (a \otimes 1) = (1 \otimes a)$$

*making* $B \otimes_A C$ *into an* $A$*-algebra.*

*Proof.* Let $X := B \otimes_A C$ and consider the map $m : X \otimes_A X \to X$ defined in (3.4.33). Define $x \cdot y := m(x \otimes y)$ then the properties of $m$ ensure that this satisfies the properties of a ring. ∎

**Proposition 3.4.35 (Criteria to be an algebra homomorphism)**
*Let* $B, C, Z$ *be commutative* $A$*-algebras and let* $\phi : B \otimes_A C \to Z$ *be an* $A$*-module homomorphism. Then the following are equivalent*

*a)* $\phi$ *is an* $A$*-algebra homomorphism*
*b)* $\phi((b \otimes c) \cdot (b' \otimes c')) = \phi(b \otimes c) \cdot \phi(b' \otimes c')$

*Similarly suppose* $\psi : Z \to B \otimes_A C$ *is an* $A$*-module homomorphism and* $Z$ *is generated as an* $A$*-module by* $S \subset Z$*. Then the following equivalent*

*a)* $\psi$ *is an* $A$*-algebra homomorphism*
*b)* $\psi(ss') = \psi(s)\psi(s')$

*Proof.* In each case $a) \implies b)$ is obvious. For the converse we simply need to show that $\phi$ and $\psi$ are in general multiplicative. This follows by linearity and because every element of the tensor product is a linear combination of elementary tensors. ∎

**Proposition 3.4.36 (Functoriality)**
*Let* $\phi : B \to B'$ *and* $\psi : C \to C'$ *be* $A$*-algebra homomorphisms. The* $A$*-module homomorphism*

$$\phi \otimes \psi : B \otimes_A C \to B' \otimes_A C'$$
$$b \otimes c \to \phi(b) \otimes \phi(c)$$

*is an* $A$*-algebra homomorphism.*

*Proof.* The $A$-module homomorphism exists by (3.4.4). It is in $A$-algebra homomorphism by (3.4.35). ∎

**Proposition 3.4.37 (Tensor Product is Coproduct)**
*Let* $B, C$ *be commutative* $A$*-algebras then there are* $A$*-algebra homomorphisms*

© David Rufino CC BY-NC-ND 4.0

$$\begin{aligned} i_B : B &\to B \otimes_A C \\ b &\to b \otimes 1 \\ i_C : C &\to B \otimes_A C \\ c &\to 1 \otimes c \end{aligned}$$

*which are natural in $B$ and $C$ respectively. In particular $B \otimes_A C$ is both a $C$-algebra and a $B$-algebra.*

*Furthermore for $Z$ an $A$-algebra there is a bijection*

$$\begin{aligned} \mathrm{AlgHom}_A(B \otimes_A C, Z) &\cong \mathrm{AlgHom}_A(B, Z) \times \mathrm{AlgHom}_A(C, Z) \\ \psi &\to (\psi \circ i_B, \psi \circ i_C) \end{aligned}$$

*which is natural in $Z$.*

*Proof.* The existence of $i_B$ and $i_C$ is easily demonstrated using universal property of tensor product. Observe that in general

$$(\psi \circ i_B)(b) = \psi(b \otimes 1_C)$$

$$(\psi \circ i_C)(c) = \psi(1_B \otimes c)$$

Furthermore the map is clearly well-defined. It's injective because if $\psi$ and $\psi'$ have the same image then $\psi(b \otimes 1) = \psi'(b \otimes 1)$ and $\psi(1 \otimes c) = \psi'(1 \otimes c)$ whence $\psi(b \otimes c) = \psi'(b \otimes c)$. By linearity they are everywhere equal.

To show that the map is a bijection we construct a two-sided inverse. For given $A$-algebra homomorphisms $f : B \to Z$ and $g : C \to Z$ then $(b, c) \to (f(b)g(c))$ is $A$-bilinear and so corresponds to an $A$-module homomorphism

$$\begin{aligned} \psi_{f,g} : B \otimes_A C &\to Z \\ b \otimes c &\to f(b)g(c) \end{aligned}$$

and by (3.4.35) this is an algebra homomorphism. Clearly $\psi \circ i_B = f$ and $\psi \circ i_C = g$. Conversely let $\psi : B \otimes_A C \to Z$ be an algebra homomorphism and we define $f(b) := \psi(b \otimes 1_C)$ and $g(c) := \psi(1_C \otimes c)$. Then $f(b)g(c) = \psi(b \otimes c)$ which shows that $\psi_{f,g}$ agrees with $\psi$ on elementary tensors, and therefore is identically equal. This completes the demonstration that $\psi_{f,g}$ is a two-sided inverse to the given map.

Naturality in $Z$ is straightforward. ∎

**Proposition 3.4.38 (Extension of Scalars)**
*Let $B, C$ be commutative $A$-algebras. Then define the $C$-algebra*

$$B_{(C)} := C \otimes_A B$$

*There is a natural isomorphism for any $C$-algebra $Z$*

$$\begin{aligned} \mathrm{AlgHom}_C\left(B_{(C)}, Z\right) &\cong \mathrm{AlgHom}_A(B, Z) \\ \psi &\to \psi \circ i_B \end{aligned}$$

*Proof.* We consider the commutative diagram obtained from (3.4.37)

$$\begin{array}{ccc} \mathrm{AlgHom}_A\left(B_{(C)}, Z\right) & \xrightarrow{\sim} & \mathrm{AlgHom}_A(B, Z) \times \mathrm{AlgHom}_A(C, Z) \\ \cup & & \cup \\ \mathrm{AlgHom}_C\left(B_{(C)}, Z\right) & \xrightarrow{\sim} & \mathrm{AlgHom}_A(B, Z) \times \{i_{CZ}\} \end{array}$$

An $A$-algebra homomorphism $\phi : B_{(C)} \to Z$ is a $C$-algebra homomorphism precisely when $\phi \circ i_C = i_{CZ}$ so that the bottom arrow is well-defined and bijective as required. ∎

**Proposition 3.4.39 (Transitivity of Extension of Scalars)**
*Let $B, C$ be commutative $A$-algebras and $D$ a commutative $C$-algebra. Then there is an isomorphism of $D$-algebras*

$$\begin{aligned} D \otimes_C (C \otimes_A B) &\cong D \otimes_A B \\ d \otimes (c \otimes b) &\to (dc) \otimes b \end{aligned}$$

*Proof.* Let $Z$ be a $D$-algebra then there is a natural isomorphism of functors

© David Rufino CC BY-NC-ND 4.0

$$\begin{aligned}\mathrm{AlgHom}_D(D \otimes_C (C \otimes_A B), Z) &\cong \mathrm{AlgHom}_C(C \otimes_A B, Z)\\ &\cong \mathrm{AlgHom}_A(B, Z)\\ &\cong \mathrm{AlgHom}_D(D \otimes_A B, Z)\end{aligned}$$

Consider the case $Z = D \otimes_A B$ and the identity map yields the required isomorphism (by the Yoneda Lemma). ∎

**Proposition 3.4.40**
*Let $B$ be a commutative $A$-algebra then there is a natural isomorphism of $A$-algebras*

$$\begin{aligned}u_B : B &\xrightarrow{\sim} B_{(A)}\\ b &\to 1 \otimes b\\ ab &\leftarrow a \otimes b\end{aligned}$$

*Furthermore for $C$ a commutative $A$-algebra there is a commutative diagram*

$$\begin{array}{ccc} B & \xrightarrow{u_B} & A \otimes_A B \\ & \searrow^{u_B} & \downarrow{\scriptstyle i_A \otimes 1_B} \\ & & C \otimes_A B \end{array}$$

*Proof.* By (3.4.38) there is an bijection natural in $Z$

$$\begin{aligned}\mathrm{AlgHom}_A\left(B_{(A)}, Z\right) &\xrightarrow{\sim} \mathrm{AlgHom}_A(B, Z)\\ \psi &\to \psi(1 \otimes -)\end{aligned}$$

which by the Yoneda Lemma (2.6.58) yields the required isomorphism (given by the image of $1_{B_{(A)}}$). ∎

**Proposition 3.4.41 (Structural Morphisms are Injective)**
*Let $B, C$ be commutative $A$-algebras such that the structural morphisms are injective and $A$ is a direct factor of $B$ and $C$ (e.g. if $A = k$ is a field). Then the canonical maps (3.4.37)*

$$B \to B \otimes_A C$$

$$C \to B \otimes_A C$$

*are injective.*

*Proof.* By (3.4.22) the canonical map

$$B \otimes_A A \to B \otimes_A C$$

is injective. By (3.4.40) we have a canonical isomorphism $B \cong B \otimes_A A$ and the composite is simply the canonical map $B \to B \otimes_A C$. ∎

**Proposition 3.4.42 (Extension of Ideals under Tensor Product)**
*Let $B, C$ be commutative $A$-algebras and $\mathfrak{b} \lhd B$ and $\mathfrak{c} \lhd C$ ideals. Then $\mathfrak{b}^e$ (resp. $\mathfrak{c}^e$) is the image of the $A$-module $\mathfrak{b} \otimes_A C$ (resp. $B \otimes_A \mathfrak{c}$) in $B \otimes_A C$. Furthermore there is a canonical $A$-algebra isomorphism*

$$(B \otimes_A C)/(\mathfrak{b}^e + \mathfrak{c}^e) \cong B/\mathfrak{b} \otimes_A C/\mathfrak{c}$$

*Proof.* Let $\phi : B \to B \otimes_A C$ be the structural morphism then for $b \in \mathfrak{b}$ we have $b \otimes c = (c \otimes 1)(b \otimes 1) = (1 \otimes c)\phi(b)$ whence $\phi(\mathfrak{b}) \subseteq \mathfrak{b} \otimes_A C \subseteq \mathfrak{b}^e$. Evidently $\mathfrak{b} \otimes_A C$ is an ideal and therefore we see $\mathfrak{b} \otimes_A C = \mathfrak{b}^e$ as required. The isomorphism follows from (3.4.29). ∎

© David Rufino CC BY-NC-ND 4.0

# 3.5 Multilinear Algebra

## 3.5.1 Multilinear Maps and Determinants

**Remark 3.5.1**
*The purpose of this section is to define the determinant, rather than develop any further theory. The approach is a little non-standard as we take an abstract module-theoretic approach as in [Bou98a], but avoid use of the tensor algebra. The connection is there are natural isomorphisms (i.e. the tensor algebra represents the functor $L^n_a(M;-)$)*

$$L^n_a(M;N) \xrightarrow{\sim} \mathrm{Hom}(\Lambda^n(M), N)$$

*and*

$$\Lambda^n(M)^\vee \xrightarrow{\sim} \Lambda^n(M^\vee)$$

*when $M$ is finite free. Therefore the exterior product we define really corresponds to the exterior product on (the tensor algebra of) $M^\vee$.*

**Definition 3.5.2 (Multilinear Map)**
*Let $M_1, ..., M_n, N$ be $A$-modules then a map*

$$\psi : M_1 \times ... \times M_n \to N$$

*is $A$-**multilinear** if it is $A$-linear in each variable, whilst fixing the other variables at any value.*

**Definition 3.5.3 (Bilinear form)**
*Let $M, N$ be $A$-modules then $\psi : M \times N \to A$ is a **bilinear form** if it is $A$-multilinear.*

*Denote the set of such bilinear pairings by $\mathrm{Bilin}_A(M,N)$. It is naturally an $A$-module.*

**Proposition 3.5.4**
*Let $M, N$ be $A$-modules then there is a natural bijection*

$$\begin{aligned} \mathrm{Hom}_A(M, \mathrm{Hom}_A(N,A)) &\leftrightarrow \mathrm{Bilin}_A(M,N) \leftrightarrow \mathrm{Hom}_A(N, \mathrm{Hom}_A(M,A)) \\ \psi_L &\leftarrow \psi \\ \psi &\qquad\qquad \to \psi_R \end{aligned}$$

*where*

$$\psi_L(m)(n) = \psi(m,n) = \psi_R(n)(m)$$

**Definition 3.5.5 (Alternating map)**
*An $A$-multilinear map $f : M^n \to N$ is **alternating** if*

$$f(x_1, ..., x_n) = 0$$

*whenever $x_i = x_{i+1}$ for some $i = 1..., n-1$.*

*Denote by $L^n_a(M,N)$ the set of such alternating maps, and $L^n_a(M) := L^n_a(M,A)$ the set of **alternating forms**. These are clearly $A$-modules.*

**Proposition 3.5.6 (Functorial Properties)**
*Let $M$ be an $A$-module and $L^k_a(M)$ the set of $k$-alternating forms. Then it is a contravariant functor in $M$, that is if $g : M \to N$ then there is a well-defined map*

$$\begin{aligned} L^k_a(g) : L^k_a(N) &\to L^k_a(M) \\ \psi &\to \psi \circ g^{(k)} \end{aligned}$$

*such that $L^k_a(g \circ h) = L^k_a(h) \circ L^k_a(g)$*

**Lemma 3.5.7**
*Let $M$ be an $A$-module and $\psi \in L^{p-1}_a(M;A)$. Then*

$$\psi'(x_1, ..., x_p) := \sum_{j=1}^{p} (-1)^{j+1} \psi(x_1, ..., \hat{x_j}, ..., x_p) x_j$$

*is an element of $L^p(M;M)$.*

© David Rufino CC BY-NC-ND 4.0

*Proof.* It is clear that $\psi'$ is multilinear, so we only need to show it is alternating. Suppose that $x_i = x_{i+1}$. then the only non-zero terms are where one of $x_i$ and $x_{i+1}$ are omitted, hence

$$\psi'(x_1, ..., x_p) = (-1)^i \psi(x_1, ..., \hat{x_i}, ..., x_p) x_i + (-1)^{i+1} \psi(x_1, ..., \widehat{x_{i+1}}, ..., x_p) x_{i+1} = 0$$

as required. ∎

**Corollary 3.5.8 (Exterior Product of Alternating Forms)**
*Let $M$ be an $A$-module, $\theta \in L_a^1(M)$, $\psi \in L_a^{n-1}(M)$ then the function*

$$\theta \wedge \psi : M^p \to A$$
$$(x_1, ..., x_n) \to \sum_{j=1}^{n} (-1)^{j+1} \psi(x_1, ..., \hat{x_j}, ..., x_p) \theta(x_j)$$

*is an alternating form. We denote this by $\theta \wedge \psi$ and refer to it as either the **wedge product** or **exterior product**, and determines a bilinear map*

$$\wedge : L_a^1(M) \times L_a^{n-1}(M) \to L_a^n(M)$$

**Lemma 3.5.9**
*Let $f : M^n \to N$ be an alternating map then*

$$f\left(x_{\sigma(1)}, ..., x_{\sigma(n)}\right) = \varepsilon(\sigma) f(x_1, ..., x_n)$$

*for any permutation $\sigma \in S_n$.*

*Furthermore if any of the $x_i$ are equal then $f(x_1, ..., x_n) = 0$*

*Proof.* A permutation $\sigma$ may be represented as a product of adjacent transpositions (...) therefore it's enough to demonstrate the case $\sigma = (i \quad i+1)$. This follows directly from the definition because

$$0 = f(x+y, x+y) = f(x,x) + f(y,x) + f(x,y) + f(y,y) = f(x,y) + f(y,x)$$

Suppose $x_i = x_j$, then we may apply the first result to the transposition $\sigma = (i\ j)$ to see that $f(x_1, ..., x_n) = 0$. ∎

**Lemma 3.5.10**
*Let $\psi : M^p \to N$ be a multilinear map. Suppose $v_1, ..., v_n \in M$ and $w_1, ..., w_p \in M$ such that*

$$w_i = \sum_{j=1}^{n} a_{ij} v_j \quad \forall i = 1...p \text{ and some } a_{ij} \in A$$

*Then*

$$\psi(w_1, ..., w_p) = \sum_{\sigma} a_{1\sigma(1)} ... a_{p\sigma(p)} \psi\left(v_{\sigma(1)}, ..., v_{\sigma(p)}\right)$$

*where $\sigma$ runs over all functions $[1, p] \to [1, n]$.*

**Lemma 3.5.11**
*Let $\psi : M^n \to N$ be an alternating map. Suppose $v_1, ..., v_n \in M$ and $w_1, ..., w_n \in M$ such that*

$$w_i = \sum_{j=1}^{n} a_{ij} v_j \quad \forall i = 1...n \text{ and some } a_{ij} \in A$$

*then*

$$\psi(w_1, ..., w_n) = \sum_{\sigma \in S_n} \varepsilon(\sigma) a_{1\sigma(1)} ... a_{n\sigma(n)} \psi(v_1, ..., v_n)$$

*Proof.* By (3.5.10)

$$\psi(w_1, ..., w_n) = \sum_{\sigma} a_{1\sigma(1)} ... a_{n\sigma(n)} \psi\left(v_{\sigma(1)}, ..., v_{\sigma(n)}\right)$$

where $\sigma$ ranges over all maps from $\{1, ..., n\}$ to itself. If $\sigma$ is not a permutation, then it must not be injective and by (3.5.9) the corresponding term is zero. Therefore we may restrict to the case $\sigma \in S_n$. By the first part of (3.5.9) the result follows. ∎

© David Rufino CC BY-NC-ND 4.0

**Proposition 3.5.12**
*Let $M$ be a finite free $A$-module of rank $n$. Then the space of alternating forms $L_a^p(M;N) = 0$ for any integer $p > n$.*

*Proof.* We may use (3.5.10) and observe that at least one basis element must occur at least twice in the expansion, and so is zero by (3.5.9). ∎

**Proposition 3.5.13 (Existence and Uniqueness of Determinant Corresponding to a Basis)**
*Let $M$ be a finite free $A$-module with basis $\{v_1, ..., v_n\}$. Then there is a unique alternating form $\Delta_v \in L_a^n(M)$ such that $\Delta_v(v_1, ..., v_n) = 1$.*

*Furthermore $L_a^n(M)$ is a free module of rank 1 with basis $\Delta_v$.*

*More precisely every $\Delta \in L_a^n(M)$ satisfies $\Delta = \Delta(v_1, ..., v_n)\Delta_v$. Further every $\Delta$ also satisifes **Leibniz' formula***

$$\Delta(x_1, ..., x_n) = \sum_{\sigma \in S_n} \varepsilon(\sigma) v_{\sigma(1)}^\vee(x_1)...v_{\sigma(n)}^\vee(x_n)\Delta(v_1, ..., v_n)$$

*Proof.* Let $\{v_1^\vee, ..., v_n^\vee\}$ be the canonical dual basis of $M^\vee$. Then we may use the wedge product (3.5.8) to construct $\Delta_v$ inductively

$$\Delta_v := v_1^\vee \wedge ... \wedge v_n^\vee$$

(using "right associativity" to ensure this expression is meaningful as the left operand must always be a linear functional). It is then clear that $\Delta_v(v_1, ..., v_n) = 1$. We may apply (3.5.11) to obtain Leibniz' formula for both $\Delta_v$ and $\Delta$, which shows that $\Delta = \Delta(v_1, ..., v_n)\Delta_v$. ∎

**Corollary 3.5.14 (Existence and Uniqueness of Determinant)**
*For every $n \geq 1$ there is a unique $D_n \in L_a^n(A^n)$ such that $D_n(e_1, ..., e_n) = 1$, given by the **Leibniz Formula***

$$D_n(v_1, ..., v_n) = \sum_{\sigma \in S_n} \varepsilon(\sigma) v_{1\sigma(1)}...v_{n\sigma(n)}.$$

*Further $L_a^n(A^n)$ is a free $A$-module of rank 1 generated by $D_n$. Explicitly every $\Delta \in L_a^n(A^n)$ satisfies*

$$\Delta = \Delta(e_1, ..., e_n)D_n$$

*Further $D_n$ satisfies the **Laplace Expansion Formula** that is for all $i, j = 1...n$ we have the relationship*

$$D_n(v_1, ..., v_n) = \sum_{j=1}^{n} (-1)^{i+j} v_{ji} D_{n-1}\Big(\pi_i(v_1), ..., \widehat{\pi_i(v_j)}, ..., \pi_i(v_n)\Big)$$

*where $\pi_i : A^n \to A^{n-1}$ drops the $i$-th element.*

*Proof.* The first two statements follow from (3.5.13) in the case $M = A^n$.

The function $(-1)^{i+1} D_{n-1}(\pi_i(v_1), ..., \pi_i(v_{n-1}))$ is evidentally multilinear and alternating, that is an element of $L^{n-1}(A^n)$. Forming the wedge product with $e_i^\vee$ we obtain an element $\Delta$ of $L^n(A^n)$, which has the form given by the Laplace Expansion. Furthermore

$$\Delta(e_1, ..., e_n) = (-1)^{i+1} e_i^\vee(e_i)(-1)^{i+1} D_{n-1}\Big(\pi_i(e_n), ..., \widehat{\pi_i(e_i)}, ..., \pi_i(e_n)\Big) = 1.$$

whence by (3.5.13) $\Delta = D_n$ as required. ∎

**Proposition 3.5.15 (Rank is Unique)**
*Let $M$ be a finite free $A$-module. Then the rank is unique.*

*Proof.* This follows from (3.5.12) and (3.5.13). Explicitly if $M$ has bases of order $p$ and $q$ with $p < q$, then we deduce $L_a^q(M)$ is both zero and non-zero a contradiction. ∎

**Definition 3.5.16 (Determinant of a Module)**
*Let $M$ be a finite free $A$-module of rank $n$, then we say a generator for $L_a^n(M)$ is a **determinant** and $\Delta_v$ is the **determinant** corresponding to the basis $v_1, ..., v_n$.*

*The determinant for $A^n$ corresponding to the standard basis $e_1, ..., e_n$ is called the **standard determinant** for $A^n$, and denoted by $D_n$.*

**Corollary 3.5.17 (Determinant of an endomorphism)**
*Let $M$ be a finite free $A$-module of rank $n$ and $f \in \mathrm{End}_A(M)$ an endomorphism. Then the corresponding linear map*

© David Rufino CC BY-NC-ND 4.0

$$L_a^n(f) : L_a^n(M) \to L_a^n(M)$$

*satisfies*

$$L_a^n(f)(\psi) = \det(f)\psi$$

*for a unique* $\det(f) \in A$*, which we call the* ***determinant*** *of* $f$*. We have the following properties*

$$\begin{aligned} \det(f \circ g) &= \det(f)\det(g) \\ \det(1_M) &= 1_A \\ \det(f) &= \Delta_v(f(v_1), ..., f(v_n)) \end{aligned}$$

*for* $\Delta_v$ *any generator for* $L_a^n(M)$ *corresponding to a basis* $v_1, ..., v_n$*.*

*Proof.* Let $\Delta$ be a generator then $L_a^n(f)(\Delta) = \Delta_v(f(v_1), ..., f(v_n))\Delta$ by (3.5.13). Clearly $\det(f) \coloneqq \Delta_v(f(v_1), ..., f(v_n))$ then satisfies the equation for all such $\psi = a\Delta$. It's unique because $L_a^n(M)$ is a free module, and the properties follow from uniqueness. ∎

We may use this to define the determinant of a matrix

**Definition 3.5.18 (Determinant of a Matrix)**
*Let* $E \in \mathrm{Mat}_{n\times n}(A)$ *then we define the* ***determinant*** *of* $E$ *to be simply* $\det\left(\hat{E}\right)$*.*

**Proposition 3.5.19 (Expressions for Matrix Determinant)**
*Let* $E \in \mathrm{Mat}_{n\times n}(A)$ *and denote by* $v_1, ..., v_n$ *the column vectors of* $E$*. Then*

$$\det(E) = D_n(v_1, ..., v_n)$$

*where* $D_n$ *is the standard determinant. Further there is the Leibniz Formula*

$$\det(E) = \sum_{\sigma \in S_n} \varepsilon(\sigma) \prod_{i=1}^{n} E_{i\sigma(i)}$$

*and Laplace Expansion Formula*

$$\det(E) = \sum_{j=1}^{n} (-1)^{i+j} E_{ij} \det\left(E_{(ij)}\right)$$

*where* $E_{(ij)}$ *is the* $(n-1) \times (n-1)$ *matrix with the* $i$*-th row and* $j$*-th column dropped.*

*Proof.* Let $e_1, ..., e_n$ be the standard basis of $A^n$ then by definition we have $\hat{E}(e_i) = v_i$. Recall by (3.5.14) that $D_n$ is a generator of $L_a^n(M)$ and so by (3.5.17) we have $\det\left(\hat{E}\right) = D_n(v_1, ..., v_n)$.

The Leibniz Formula and Laplace Expansion Formula also follows directly from (3.5.14). ∎

**Corollary 3.5.20 (Properties of Matrix Determinant)**
*The determinant satisfies a number of properties*

*a)* $\det(EF) = \det(E)\det(F) = \det(F)\det(E)$
*b)* $\det(I_n) = 1$
*c)* $\det(PEP^{-1}) = \det(E)$
*d)* $\det(E^t) = \det(E)$

*Proof.* a)-c) follow from (3.3.115) and (3.5.17). Explicitly

$$\det(EF) = D\left(\widehat{EF}\right) = D\left(\hat{E} \circ \hat{F}\right) = D\left(\hat{E}\right)D\left(\hat{F}\right) = \det(E)\det(F)$$

and

$$\det(I_n) = D\left(\hat{I}_n\right) = D(\mathbf{1}_{A^n}) = 1_A$$

d) can be shown by using Leibniz formula and observing that $\sigma \to \sigma^{-1}$ simply permutes the elements of $S_n$. ∎

**Proposition 3.5.21**
*Let* $M$ *be a finite free* $A$*-module and* $f \in \mathrm{End}_A(M)$*. Let* $(e_1, ..., e_n)$ *be a basis and* $\Delta_e$ *the corresponding determinant. Define*

© David Rufino CC BY-NC-ND 4.0

$$f^{\mathrm{ad}}(x) \coloneqq \sum_{j=1}^{n} \Delta_e\big(f(e_1), ..., f\big(e_{j-1}\big), x, f\big(e_{j+1}\big), ..., f(e_n)\big) e_j.$$

*Then*

$$f \circ f^{\mathrm{ad}} = f^{\mathrm{ad}} \circ f = \det(f) \mathbf{1}_M$$

*Furthermore the matrix of $f^{\mathrm{ad}}$ with respect to the given basis satisfies*

$$\big[f^{\mathrm{ad}}\big]_{ij} = (-1)^{i+j} \det\Big(\big[f\big]_{(ji)}\Big)$$

*where $M_{(ji)}$ denotes the matrix $M$ with the $j$-th row and $i$-th column removed.*

*Proof.* We may consider by (3.5.7) the alternating map $\psi \in L_a^{n+1}(M; M)$ given by

$$\psi(x_1, ..., x_{n+1}) \coloneqq \sum_{j=1}^{n+1} (-1)^{j+1} \Delta_e\big(x_1, ..., \hat{x}_j, ..., x_{n+1}\big) x_j$$

By (3.5.12) necessarily $\psi$ is identically zero. Evaluating at $(x, f(e_1), ..., f(e_n))$ we find that

$$0 = \det(f) x + \sum_{j=1}^{n} (-1)^j \Delta_e\Big(x, f(e_1), ..., \widehat{f\big(e_j\big)}, ..., f(e_n)\Big) f\big(e_j\big)$$

whence

$$\begin{aligned} \det(f) x &= \sum_{j=1}^{n} (-1)^{j+1} \Delta_e\Big(x, f(e_1), ..., \widehat{f\big(e_j\big)}, ..., f(e_n) f\big(e_j\big)\Big) \\ &= \sum_{j=1}^{n} \Delta_e(f(e_1), ..., x, ..., f(e_n)) f\big(e_j\big) \end{aligned}$$

where we have used (3.5.9).

Therefore using the definition of $f^{\mathrm{ad}}(x)$ we find that $f\big(f^{\mathrm{ad}}(x)\big) = \det(f) x$ as required. For the second relation, first observe that (3.3.109)

$$x = \sum_{k=1}^{n} e_k^\vee(x) e_k$$

and so applying $f(-)$

$$f(x) = \sum_{k=1}^{n} e_k^\vee(x) f(e_k)$$

Therefore we may calculate directly

$$\begin{aligned} f^{\mathrm{ad}}(f(x)) &= \sum_{j=1}^{n} \Delta_e\big(f(e_1), ..., f\big(e_{j-1}\big), f(x), f\big(e_{j+1}\big), ..., f(e_n)\big) e_j \\ &= \sum_{j=1}^{n} \Delta_e(f(e_1), ..., f(e_n)) e_j^\vee(x) e_j \\ &= \det(f) \sum_{j=1}^{n} e_j^\vee(x) e_j \\ &= \det(f) x \end{aligned}$$

as required.

For the final statement we may assume that $M = A^n$ with $(e_1, ..., e_n)$ the standard basis and $D_n = \Delta_e$. Then by (3.5.14), (3.5.18) we have

$$\begin{aligned} \big[f^{\mathrm{ad}}\big]_{ij} &= D_n\big(f(e_1), ..., f(e_{i-1}), e_j, f(e_{i+1}), ..., f(e_n)\big) \\ &= (-1)^{i+j} D_{n-1}\Big(\pi_j(f(e_1)), ..., \pi_j\widehat{(f(e_i))}, ..., \pi_j(f(e_n))\Big) \\ &= (-1)^{i+j} \det\Big(\big[f\big]_{(ji)}\Big) \end{aligned}$$

© David Rufino CC BY-NC-ND 4.0

■

**Corollary 3.5.22**
*Let $M$ be a finite free $A$-module. Then $f \in \mathrm{End}_A(M)$ is an isomorphism if and only if $D(f) \in A^\star$.*

**Corollary 3.5.23**
*Let $E \in \mathrm{Mat}_{n\times n}(A)$ and define the adjugate matrix*

$$E^{\mathrm{ad}}_{ij} \coloneqq (-1)^{i+j} \det\left(E_{(ji)}\right)$$

*Then*

$$E^{\mathrm{ad}}E = EE^{\mathrm{ad}} = \det(E)I_n$$

*Proof.* By definition and (3.5.21) we have that $\widehat{E^{\mathrm{ad}}} = \hat{E}^{\mathrm{ad}}$. The identities of endomorphisms are proven in (3.5.21), and we deduce the corresponding identities of matrices by (3.3.117). ■

## 3.5.2 Trace Map

**Proposition 3.5.24 (Existence and Uniqueness of the Trace Map)**
*Let $M$ be an $A$-module, then there exists a well-defined $A$-bilinear map*

$$\begin{aligned} M \otimes_A M^\vee &\to \mathrm{End}_A(M) \\ m \otimes \theta &\to \theta(-)\cdot m \cdot \end{aligned}$$

*When $M$ is a finite free $A$-module this is bijective, and there exists a unique $A$-linear map $\mathrm{Tr} : \mathrm{End}_A(M) \to A$ making the following diagram commute*

$$\begin{array}{ccc} M \otimes_A M^\vee & \xrightarrow{\sim} & \mathrm{End}_A(M) \\ & \searrow & \downarrow \mathrm{Tr} \\ & & A \end{array}$$

*where the diagonal map is simply evaluation*

$$m \otimes \theta \to \theta(m)$$

*Equivalently it is the unique $A$-linear map satisfying*

$$\mathrm{Tr}(\theta(-)\cdot m) = \theta(m)$$

*for all $m \in M$ and $\theta \in M^\vee$.*

*Finally we have the identity*

$$\mathrm{Tr}(\phi) = \mathrm{Tr}([\phi]) = \sum_{i=1}^{n} [\phi]_{ii}$$

*for all bases $\mathcal{B}$ of $M$.*

*Proof.* There exists an obvious $A$-bilinear map $M \times M^\vee \to \mathrm{End}_A(M)$ so the map exists by the universal property. If $M$ is finite free then it has a basis $\{v_1, ..., v_n\}$. By (3.3.109) $M^\vee$ has basis $\{v_1^\vee, ..., v_n^\vee\}$ and by (...) $M \otimes_A M^\vee$ has basis $\left\{v_i \otimes v_j^\vee\right\}$. By (3.3.110) this maps to a basis of $\mathrm{End}_A(M)$ and so it is an isomorphism.

Similarly the map $M \times M^\vee \to A$ is $A$-bilinear, and so the evaluation map $M \otimes_A M^\vee \to A$ exists. In light of the isomorphism then Tr exists and satisfies the required uniqueness property. ■

**Proposition 3.5.25 (Trace is Symmetric)**
*Let $\phi, \psi : M \to M$ be endormophisms of a finite free $A$-module. Then*

$$\mathrm{Tr}(\phi \circ \psi) = \mathrm{Tr}(\psi \circ \phi)$$

*Proof.* We may verify that the mapping

$$\begin{aligned} \mathrm{End}_A(M) \times \mathrm{End}_A(M) &\to A \\ (\phi, \psi) &\to \mathrm{Tr}(\phi \circ \psi) \end{aligned}$$

© David Rufino CC BY-NC-ND 4.0

is $A$-bilinear. Therefore it is sufficient to show the relationship holds for rank-one endomorphisms $\phi = \theta(-) \cdot m$ and $\psi = \iota(-) \cdot n$, as these generate $\mathrm{End}_A(M)$. Then we have

$$\mathrm{Tr}(\phi \circ \psi) = \mathrm{Tr}(\theta(\iota(-) \cdot n) \cdot m) = \mathrm{Tr}((\iota(-)\theta(n)) \cdot m) = \iota(m)\theta(n)$$

and by symmetry the result follows. ∎

Recall from (...) that every endomorphism $\phi : M \to M$ determines a family of endomorphisms $L^r_a(\phi) : L^r_a(M) \to L^r_a(M)$ for all integers $1 \leq r \leq n$, where $n$ is the rank of $M$.

**Proposition 3.5.26 (Determinant as a Trace)**
*Let $M$ be a finite-free $A$ module of rank $n$ and $\phi : M \to M$ an endomorphism. Then*

$$\det(\phi) = \mathrm{Tr}(L^n_a(\phi))$$

*and*

$$\mathrm{Tr}(\phi) = \mathrm{Tr}(L^1_a(\phi))$$

**Definition 3.5.27**
*Let $K$ be a subset of $[1,n]$ of order $k$. Then we denote by $\sigma_K : [1,k] \to K$ the unique order preserving bijection.*

*Denote by $\mathfrak{F}_k(n)$ the family of subsets of $[1,n]$ of order $k$.*

*If $X$ is an $m \times n$ matrix and $K \subset [1,m]$, $H \subset [1,n]$, then we denote by $X_{H,K}$ the $\#K \times \#H$ matrix given by*

$$\left(X_{H,K}\right)_{ij} = X_{\sigma_K(i),\sigma_H(j)}$$

**Lemma 3.5.28**
*Let $M$ be an $A$-module with subset $\{v_1, ..., v_n\}$ and $\psi : M^k \to N$ be an alternating map. Suppose that $x_1, ..., x_k$ satisfy*

$$x_i = \sum_{j=1}^{n} A_{ji} v_j.$$

*for some $n \times k$ matrix $A$. Then*

$$\psi(x_1, ..., x_k) = \sum_{K \subset \mathfrak{F}_k(n)} \det\left(A_{K,[1,k]}\right)\psi(v_K)$$

*where $v_K := \left(v_{\sigma_K(1)}, ..., v_{\sigma_K(k)}\right)$.*

*Proof.*

$$\begin{aligned}
\psi(x_1, ..., x_k) &= \sum_{i_1,...,i_k} a_{i_1 1} ... a_{i_k k} \psi\left(v_{i_1}, ..., v_{i_k}\right) \\
&= \sum_{K \subset \mathfrak{F}_k(n)} \sum_{\sigma \in S_k} a_{\sigma_K(\sigma(1))1} ... a_{\sigma_K(\sigma(k))k} \psi\left(v_{\sigma_K(\sigma(1))}, ..., v_{\sigma_K(\sigma(k))}\right) \\
&= \sum_{K \subset \mathfrak{F}_k(n)} \sum_{\sigma \in S_k} \varepsilon(\sigma) a_{\sigma_K(\sigma(1))1} ... a_{\sigma_K(\sigma(k))k} \psi\left(v_{\sigma_K(1)}, ..., v_{\sigma_K(k)}\right) \\
&= \sum_{K \subset \mathfrak{F}_k(n)} \det\left(A_{K,[1,k]}\right)\psi(v_K)
\end{aligned}$$

∎

**Proposition 3.5.29**
*Let $M$ be a finite free $A$-module with $v_1, ..., v_n$ a basis. For every subset $K \subset [1,n]$ of order $k$ denote*

$$v^\vee_K := v_{\sigma_K(1)} \wedge ... \wedge v_{\sigma_K(k)}.$$

*Then $L^k_a(M)$ has basis $\left\{v^\vee_K\right\}_{K \in \mathfrak{F}_k(n)}$. Furthermore we have the identity*

$$v^\vee_K(v_{K'}) = \delta_{KK'}$$

*and each basis element is uniquely determined by this relationship.*

© David Rufino CC BY-NC-ND 4.0

**Proposition 3.5.30**
*Let $M$ be a finite free $A$-module with basis $\mathcal{B} = \{v_1, ..., v_n\}$. Let $\phi : M \to M$ be an $A$-module endomorphism. Then for all $K \in \mathfrak{F}_k(n)$*

$$L_a^k(\phi)(v_K^\vee) = \sum_{H \in \mathfrak{F}_k(n)} \det\big(X_{K,H}\big) v_H^\vee$$

*where $X = [\phi]_{\mathcal{B}}$.*

*In particular we see that*

$$\mathrm{Tr}\big(L_a^k(\phi)\big) = \sum_{K \in \mathfrak{F}_k(n)} \det\big(X_{K,K}\big)$$

*Proof.* We have for $H \in \mathfrak{F}_k(n)$

$$\begin{aligned} L_a^k(\phi)(v_K^\vee)(v_H) &= v_K^\vee\Big(\phi\big(v_{\sigma_H(1)}\big), ..., \phi\big(v_{\sigma_H(k)}\big)\Big) \\ &= \sum_{L \subset \mathfrak{F}_k(n)} \det\big(X_{L,H}\big) v_K^\vee(v_L) \\ &= \det\big(X_{K,H}\big) \end{aligned}$$

By (3.5.29) then both sides agree on $v_H$ for all $H \in \mathfrak{F}_k(n)$, which shows they are equal by (3.5.28). ∎

**Proposition 3.5.31**
*Let $M$ be a finite-free $A$ module of rank $n$ and $\phi : M \to M$ an endomorphism. For $a, b \in A$ we have*

$$\det(a1_M + b\phi) = \sum_{k=0}^{n} \mathrm{Tr}\big(L_a^k(\phi)\big) a^{n-k} b^k$$

*Proof.* To calculate the left hand side consider a basis $\{v_1, ...v_n\}$ for $M$ and the corresponding determinant $\Delta_v$. Then

$$\det(a1_M + b\phi) = \Delta_v(av_1 + b\phi(v_1), ..., av_n + b\phi(v_n))$$

Expand to find that this equals

$$\sum_{k=0}^{n} a^{n-k} b^k \sum_{K \in \mathfrak{F}_k(n)} \Delta_v(x_{1K}, ..., x_{nK})$$

where $x_{iK} = \phi(v_i)$ if $i \in K$ and $v_i$ otherwise.

Define $\tau_K$ to be the unique shuffle permutation of type $(k, n-k)$ (3.2.38) such that $\tau_K\,|_{[1,k]} = \sigma_K$. Denote the inner term of the sum by $z_K$. Then we have by (3.5.9)

$$z_K = \varepsilon(\tau_K)\Delta_v\Big(\phi\big(v_{\sigma_K(1)}\big), ..., \phi\big(v_{\sigma_K(k)}\big), v_{\sigma_H(1)}, ..., v_{\sigma_H(n-k)}\Big)$$

where $H$ is the complement of $K$ in $[1, n]$. Then by considering $\Delta_v$ to be an alternating form in the first $k$ arguments, we may apply (3.5.28) to find

$$z_K = \varepsilon(\tau_K) \sum_{L \subset \mathfrak{F}_k(n)} \det\big(X_{L,K}\big)\Delta_v\Big(v_{\sigma_L(1)}, ..., v_{\sigma_L(k)}, v_{\sigma_H(1)}, ..., v_{\sigma_H(n-k)}\Big)$$

The summand is non-zero if and only if $L \cap H = \varnothing$, which occurs if and only if $L = K$. Therefore we conclude that

$$\begin{aligned} z_K &= \varepsilon(\tau_K)\det\big(X_{K,K}\big)\Delta_v\Big(v_{\tau_K(1)}, ..., v_{\tau_K(n)}\Big) \\ &= \det\big(X_{K,K}\big)\Delta_v(v_1, ..., v_n) \\ &= \det\big(X_{K,K}\big) \end{aligned}$$

The result then follows easily from (3.5.30). ∎

### 3.5.3 Block Matrices

**Proposition 3.5.32**
*Let $A = \mathbb{Z}[X_{11}, ..., X_{nn}]$ be a polynomial ring in $n^2$ indeterminates. Consider the matrix $\big(X_{ij}\big) \in \mathrm{Mat}_{n\times n}(A)$ and let $D_n$ denote the determinant of this matrix.*

*Then for any commutative ring $B$ and matrix $M \in \mathrm{Mat}_{n\times n}(B)$ we have*

© David Rufino CC BY-NC-ND 4.0

$$\det(B) = D_n(B_{11}, ..., B_{1n}, B_{21}, ..., B_{nn})$$

*where the right hand side corresponds the obvious evaluation homomorphism $A \to B$.*

**Proposition 3.5.33 (Block Matrix Determinant)**
*Suppose $M_{ij}$ $(1 \leq i \leq n, 1 \leq j \leq n)$ are $m \times m$ matrices with coefficients in a ring $A$ and which commute pairwise. Form the matrix $M$ with coefficients in $A$ as follows*

$$M = \begin{pmatrix} M_{11} & ... & M_{1n} \\ \vdots & & \vdots \\ M_{n1} & ... & M_{nn} \end{pmatrix}$$

*Then*

$$\det(M) = \det(D_n(M_{11}, ..., M_{1n}, M_{21}, ..., M_{nn}))$$

*Proof.* We observe without proof that there is a "block" correspondence

$$\mathrm{Mat}_{n\times n}(\mathrm{Mat}_{m\times m}(A)) \leftrightarrow \mathrm{Mat}_{(nm)\times(nm)}(A)$$

which commutes with matrix multiplication.

Let $B$ denote the commutative subring of $\mathrm{Mat}_{m\times m}(A)$ generated by the matrices $M_{ij}$. Then we may regard $M$ as an element of $\mathrm{Mat}_{n\times n}(B)$, and denote this by $\widetilde{M}$ to disambiguate. As $B$ is commutative we my consider the adjugate matrix $\widetilde{M}^{\mathrm{ad}}$. Further consider the matrix $M'$ given in block form by

$$M' = \begin{pmatrix} \left(\widetilde{M}^{\mathrm{ad}}\right)_{11} & 0 & ... & 0 \\ \left(\widetilde{M}^{\mathrm{ad}}\right)_{21} & I_m & ... & 0 \\ \vdots & \vdots & \ddots & \vdots \\ \left(\widetilde{M}^{\mathrm{ad}}\right)_{n1} & 0 & ... & I_m \end{pmatrix}$$

Recall by (3.5.23) and (3.5.32) that

$$\widetilde{M}^{\mathrm{ad}}_{11} = D_{n-1}(M_{22}, ..., M_{2n}, M_{31}, ..., M_{nn})$$

and

$$\widetilde{M}\widetilde{M}^{\mathrm{ad}} = \begin{pmatrix} \det\left(\widetilde{M}\right) & ... & 0 \\ \vdots & \ddots & \vdots \\ 0 & ... & \det\left(\widetilde{M}\right) \end{pmatrix}$$

as elements of $\mathrm{Mat}_{n\times n}(B)$. Therefore computing the matrix product block-wise we find

$$MM' = \begin{pmatrix} \det\left(\widetilde{M}\right) & M_{12} & ... & M_{1n} \\ 0 & M_{22} & ... & M_{2n} \\ \vdots & \vdots & & \vdots \\ 0 & M_{n2} & ... & M_{nn} \end{pmatrix}$$

Then we may show using the Leibniz Expansion that

$$\det(M) \cdot \det(M') = \det(MM') = \det\left(\det\left(\widetilde{M}\right)\right) \cdot \det\left(M_{[n+1,mn,n+1,mn]}\right)$$

By induction on $n$ we may then assume $\det\left(M_{[n+1,mn,n+1,mn]}\right) = \det(D_{n-1}(M_{22}, ..., M_{2n}, ..., M_{nn})) = \det\left(\left(\widetilde{M}^{\mathrm{ad}}\right)_{11}\right)$. Similarly we may argue by the Leibniz expansion that $\det(M') = \det\left(\left(\widetilde{M}^{\mathrm{ad}}\right)_{11}\right)$. When this is not a zero-divisor in $A$ then we conclude that

$$\det(M) = \det\left(\det\left(\widetilde{M}\right)\right) = \det(D_n(M_{11}, ..., M_{1n}, ..., M_{nn}))$$

as required.

In order to handle the case where $\det\left(\left(\widetilde{M}^{\mathrm{ad}}\right)_{11}\right)$ is a zero divisor we may replace the ring $A$ with the polynomial ring $A[Z]$ and consider the matrix $N$ for which

© David Rufino CC BY-NC-ND 4.0

$$\tilde{N} = \widetilde{M} + (Z \cdot I_m) \cdot I_n.$$

Then the submatrices of $N$ still commute pairwise. Further $\det\Big(\big(\tilde{N}^{\mathrm{ad}}\big)_{11}\Big)$ is a polynomial of non-zero degree in $Z$ whose leading coefficient is 1, and therefore is not a zero-divisor. We conclude from the previous case that

$$\det(N) = \det(D_n(N_{11}, ..., N_{1n}, ..., N_{nn}))$$

Finally we may apply the ring homomorphism $A[Z] \to A$ (evaluation at 0) to deduce the general case. ∎

### 3.5.4 Exterior Product

**Proposition 3.5.34**
*Let $M$ be an $A$-module. There is a bilinear map*

$$\wedge : L_a^p(M) \times L_a^q(M) \to L_a^{p+q}(M)$$

*given by*

$$(f \wedge g)(x_1, ..., x_{p+q}) := \sum_{\sigma \in \mathrm{Sh}(p,q)} \varepsilon(\sigma) f\big(x_{\sigma(1)}, ..., x_{\sigma(p)}\big) g\big(x_{\sigma(p+1)}, ..., x_{\sigma(p+q)}\big)$$

*where the sum is taken over shuffle permutations (3.2.38).*

*Proof.* Suppose that $x_i = x_j$ then we require to prove that $(f \wedge g)(x_1, ..., x_{p+q}) = 0$ in order to demonstrate that it is alternating. If $1 \leq i, j \leq p$ then this follows from (3.5.9). Similarly if $p < i, j \leq p + q$. So we may consider the case $i \leq p < j$. Consider the family of right cosets $\mathrm{Sh}(p,q)/\{e, (i,j)\}$, and let $\Sigma$ be some coset representatives. Then by (3.2.10)

$$\mathrm{Sh}(p,q) = \bigsqcup_{\sigma \in \Sigma} \{\sigma, (i,j)\sigma\}$$

Therefore the sum is equal to

$$\sum_{\sigma \in \Sigma} \varepsilon(\sigma) \big[ f(x_\sigma) g(x_\sigma) - f\big(x_{(i\,j)\sigma}\big) g\big(x_{((i\,j)\sigma)}\big) \big]$$

so when $x_i = x_j$ this sum is zero. ∎

**Proposition 3.5.35 (Basic Properties of Exterior Product)**
*Let $M$ be an $A$-module. Suppose $f \in L_a^p(M)$, $g \in L_a^q(M)$ and $h \in L_a^r(M)$. Then the following properties hold*

*a)* $f \wedge g = (-1)^{pq} g \wedge f$
*b)* $(f \wedge g) \wedge h = f \wedge (g \wedge h)$

*Proof.*

a) By definition and (3.2.44)

$$\begin{aligned}(f \wedge g)(x) &= \sum_{\sigma \in \mathrm{Sh}(p,q)} \varepsilon(\sigma) f\big(x_{\sigma(1)}, ..., x_{\sigma(p)}\big) g\big(x_{\sigma(p+1)}, ..., x_{\sigma(p+q)}\big) \\ &= \sum_{\sigma \in \mathrm{Sh}(q,p)} (-1)^{pq} \varepsilon(\sigma) f\big(x_{\sigma(p+1)}, ..., x_{\sigma(p+q)}\big) g\big(x_{\sigma(1)}, ..., x_{\sigma(p)}\big) \\ &= (-1)^{pq}(g \wedge f)\end{aligned}$$

b) Similarly by definition

$$\begin{aligned}((f \wedge g) \wedge h)(x_1, ..., x_{p+q+r}) &= \sum_{\sigma \in \mathrm{Sh}(p+q,r)} \varepsilon(\sigma)(f \wedge g)\big(x_{\sigma(1)}, ..., x_{\sigma(p+q)}\big) h\big(x_{\sigma(p+q+1)}, ..., x_{\sigma(p+q+r)}\big) \\ &= \sum_{\sigma \in \mathrm{Sh}(p+q,r)} \sum_{\tau \in \mathrm{Sh}(p,q)} \varepsilon(\sigma)\varepsilon(\tau) f\big(x_{\sigma(\tau(1))}, ..., x_{\sigma(\tau(p))}\big) g\big(x_{\sigma(\tau(p+1))}, ..., x_{\sigma(\tau(p+q))}\big) \\ &\qquad h\big(x_{\sigma(p+q+1)}, ..., x_{\sigma(p+q+r)}\big) \\ &\overset{(3.2.45)}{=} \sum_{\sigma \in \mathrm{Sh}(p,q,r)} \varepsilon(\sigma) f\big(x_{\sigma(1)}, ..., x_{\sigma(p)}\big) g\big(x_{\sigma(p+1)}, ..., x_{\sigma(p+q)}\big) h\big(x_{\sigma(p+q+1)}, ..., x_{\sigma(p+q+r)}\big)\end{aligned}$$

and symmetrically

© David Rufino CC BY-NC-ND 4.0

$$\begin{aligned}(f\wedge(g\wedge h))(x_1,\dots,x_{p+q+r}) &= \sum_{\sigma\in \mathrm{Sh}(p,q+r)} \varepsilon(\sigma) f\left(x_{\sigma(1)},\dots,x_{\sigma(p)}\right)(g\wedge h)\left(x_{\sigma(p+1)},\dots,x_{\sigma(p+q+r)}\right)\\ &= \sum_{\sigma\in \mathrm{Sh}(p,q+r)}\sum_{\tau\in \mathrm{Sh}(q,r)} \varepsilon(\sigma)\varepsilon(\tau) f\left(x_{\sigma(1)},\dots,x_{\sigma(p)}\right) g\left(x_{\sigma(\tau(p+1))},\dots,x_{\sigma(\tau(p+q))}\right)\\ &\qquad h\left(x_{\sigma(\tau(p+q+1))},\dots,x_{\sigma(\tau(p+q+r))}\right)\\ &\overset{(3.2.45)}{=} \sum_{\sigma\in \mathrm{Sh}(p,q,r)} \varepsilon(\sigma) f\left(x_{\sigma(1)},\dots,x_{\sigma(p)}\right) g\left(x_{\sigma(p+1)},\dots,x_{\sigma(p+q)}\right) h\left(x_{\sigma(p+q+1)},\dots,x_{\sigma(p+q+r)}\right)\end{aligned}$$

therefore we conclude that the expressions are equal.

■

## 3.5.5 Matrix Rank

**Definition 3.5.36 (Column and Row Rank)**
*Let $k$ be a field and $E$ an $m\times n$ a matrix over $k$. Consider the canonical vector spaces $k^n$ and $k^m$. Then define the **column rank** of $E$ to be*

$$\mathrm{rank}_k\left(\hat{E}\right)$$

*and the **row rank** of $E$ to be*

$$\mathrm{rank}_k\left(\widehat{E^t}\right)$$

**Proposition 3.5.37 (Row Rank = Column Rank)**
*Let $E$ be a matrix over $k$, then row rank and column rank are equal, and denote this by $\mathrm{rk}(E)$.*

*It is also the maximal number of linearly independent rows, or columns, and furthermore $\mathrm{rk}(E)\leq \min(m,n)$.*

*We say $E$ is **full rank** if $\mathrm{rk}(E)=\min(m,n)$.*

*Proof.* By (3.3.144) $\mathrm{rank}_k\left(\hat{E}\right)=\mathrm{rank}_k\left(\hat{E}^\vee\right)$ and by (3.3.121) this equals $\mathrm{rank}_k\left(\widehat{E^t}\right)$ as required.

The columns (resp. rows) clearly span $\mathrm{Im}\left(\hat{E}\right)$ (resp. $\mathrm{Im}\left(\widehat{E^t}\right)$). By (3.3.126) there are $r:=\mathrm{rk}(E)$ columns (resp. rows) constituting a basis, and therefore linearly independent. For any other subset of linearly independent columns (resp. rows) we must have the order is less than $r$ by (3.3.126). Therefore $\mathrm{rk}(E)$ is the maximal number of linearly independent rows or columns. ■

**Proposition 3.5.38 (Criteria for Full Rank Square Matrix)**
*Let $E$ be an $n\times n$ matrix over a field $k$. Then the following are equivalent*

*a) $E$ is invertible*
*b) $\mathrm{rk}(E)=n$ (i.e. $\hat{E}$ is surjective or $E$ is full-rank)*
*c) $Ev=0\implies v=0$ for all column vectors $v$ (i.e. $\hat{E}$ is injective).*
*d) The columns of $E$ are linearly independent*
*e) $\det(E)\neq 0$*

*Finally $E$ is full rank if and only if $E^t$ is full rank.*

*Proof.* Consider $k^n$ with canonical basis, then by (3.3.115) $E$ is invertible if and only if $\hat{E}$ is an isomorphism. By definition $\mathrm{rk}(E)=\mathrm{rank}_k\left(\hat{E}\right)$. Furthermore c) is equivalent to $\hat{E}$ being injective, and is also equivalent to d). Therefore the equivalence follows from (3.3.137).

Finally it's clear from either a), b) or e) that this property is self-dual. ■

**Definition 3.5.39 (Minor of a matrix)**
*Let $E$ be an $m\times n$ matrix, we say a **$k$-minor** (for $k\leq\min(m,n)$) is the determinant of a $k\times k$ submatrix obtained by deleting $m-k$ rows and $n-k$ columns.*

**Proposition 3.5.40 (Criteria for rank)**
*Let $E$ be an $m\times n$ matrix over $k$. Then the following are equivalent*

*a) $\mathrm{rk}(E)\geq r$*
*b) There exists an $r\times r$ sub-matrix with full rank*
*c) There exists a non-zero $r$-minor*

© David Rufino CC BY-NC-ND 4.0

*Proof.* We see b) $\iff$ c) by (3.5.38) e).

Suppose b) holds, then a-fortiori $E$ has $r$ linearly independent columns. Therefore by (3.5.37) $\mathrm{rk}(E) \geq r$ and a) holds.

Conversely suppose $\mathrm{rk}(E) \geq r$ then by (3.5.37) there are certainly $r$ linearly independent columns. We consider the $m \times r$ sub-matrix $E'$ consisting of these columns. By (3.5.37) $\mathrm{rk}(E') = r$ and there are $r$ linearly independent rows. Choosing these rows yields an $r \times r$ submatrix $E''$ which has $r$ linearly independent rows, and so by (3.5.38) is full rank as required. ■

**Corollary 3.5.41**
*Let $E$ be an $m \times n$ matrix over $k$ and $r$ an integer. Then the following are equivalent*

*a)* $\mathrm{rk}(E) = r$
*b) $r$ is the maximal dimension of a full-rank square sub-matrix*
*c) $r$ is the maximal dimension of a non-zero minor*

© David Rufino CC BY-NC-ND 4.0

# 3.6 Localisation

Algebraically, localisation can be seen as enlargening a ring to include inverses. In terms of the ideal structure this means removing (proper) ideals which contain the newly inverted elements. Geometrically ideals correspond to points/subsets, so localisation may be viewed as reducing the set of interest.

Recall the definition of multiplicative set. Some rather canonical examples are as follows

**Example 3.6.1**
*The set $S_f = \{1, f, f^2, ...\}$ is m.c. but not necessarily saturated. As an example consider $A = \mathbb{Z}$ and $S_n = \{1, n, n^2, ...\}$ for $n$ composite. Then $pq \in S_n$ but $p \notin S_n$.*

*We denote the localisation of $A$ at $S_f$ by $A_f$ (or alternatively $A[1/f]$).*

**Example 3.6.2**
*If $\mathfrak{p} \lhd A$ is a prime ideal, then $A \setminus \mathfrak{p}$ is a saturated multiplicative set. More generally, we show later that $S$ is a saturated multiplicative set if and only if it's of the form*

$$A \setminus \bigcup_i \mathfrak{p}_i$$

*for some family of prime ideals. We denote the localisation of $A$ at $\mathfrak{p}$ by $A_{\mathfrak{p}}$.*

## 3.6.1 Rings

**Definition 3.6.3 (Localisation of a ring)**
*Let $A$ be a ring and $S$ a multiplicative set. Define the set*

$$S^{-1}A = \left\{ \frac{a}{s} \;\middle|\; a \in A \, s \in S \right\}$$

*under the equivalence relation*

$$\frac{a}{s} = \frac{b}{t} \iff u(at - bs) = 0 \quad \text{some } u \in S.$$

*then this is a ring in the obvious way*

**Proposition 3.6.4**
*The set $S^{-1}A$ is a ring under the obvious ring operations. It is non-zero precisely when $S$ is proper. There is a canonical homomorphism*

$$i_S : A \to S^{-1}A$$
$$a \to \left[\frac{a}{1}\right]$$

*a)* $\ker(i_S) = \mathrm{Ann}(S)$
*b)* $S^{-1}A$ *is the zero-ring if and only if* $0 \in S$ *if and only if there exists* $s, t \in S$ *such that* $st = 0$.
*c)* $i_S(s)$ *is invertible for all* $s \in s$
*d)* $i_S$ *is injective if and only if* $S$ *has no zero-divisors*
*e) This is an isomorphism if and only if $S \subseteq A^\star$ already consists only of invertible elements (e.g. $S = \{1\}$).*

*Proof.*
a) This follows by the definitions
b) $\frac{1}{1} = \frac{0}{0} \iff s = 0$ for some $s \in s$ by the definitions
c) $\frac{s}{1} \cdot \frac{1}{s} = \frac{s}{s} = \frac{1}{1}$
d) This follows from the first part.
e) If $S \subseteq A^\star$ then it contains no zero-divisors and $i_S$ is injective. Further it's clear that $\frac{a}{s} = \frac{as^{-1}}{1}$ so that the map is surjective. Similarly if the map is bijective $S$ does not contain zero-divisors and $\frac{1}{s}$ is in the image. Therefore there is $a$ such that $tas = 1$ for some $t$, which implies $s$ is invertible.

■

Note when $A$ is an integral domain and $S$ is proper then the equivalence relation may be weakened to $at - bs = 0$.

**Proposition 3.6.5 (Universal Property)**
*Let $\phi : A \to B$ be a ring homomorphism and $S$ a multiplicative set. Then*

© David Rufino CC BY-NC-ND 4.0

*a) There is a unique morphism $\tilde{\phi}$ making the diagram commute*




*if and only if $\phi(S) \subseteq B^\star$. In this case it's given by*

$$\tilde{\phi}\left(\frac{a}{s}\right) = \phi(a)\phi(s)^{-1}$$

*b)* $\ker(\tilde{\phi}) = S^{-1}\ker(\phi)$

*Proof.*

a) If $\tilde{\phi}$ exists then $1 = \tilde{\phi}(1) = \tilde{\phi}(\frac{s}{1}\cdot\frac{1}{s}) = \tilde{\phi}(\frac{s}{1})\tilde{\phi}(\frac{1}{s}) = \phi(s)\tilde{\phi}(\frac{1}{s})$. Which shows that $\phi(S) \subseteq B^\star$ and $\phi(s)^{-1} = \tilde{\phi}(\frac{1}{s})$.
Conversely suppose $\phi(S) \subseteq B^\star$ then we claim that the given mapping is well-defined. For

$$\frac{a}{s} = \frac{a'}{s'} \implies s''(s'a - sa') = 0 \implies \phi(s'')\phi(s')\phi(a) = \phi(s'')\phi(s)\phi(a')$$

Multiply by the appropriate inverses to find

$$\phi(a)\phi(s)^{-1} = \phi(a')\phi(s')^{-1}$$

It's clearly a multiplicative homomorphism. Further it's additive because

$$\begin{aligned}\tilde{\phi}\left(\frac{a}{s} + \frac{b}{t}\right) &= \tilde{\phi}\left(\frac{at+bs}{st}\right)\\ &= \phi(at+bs)\phi(st)^{-1}\\ &= \phi(a)\phi(t)\phi(s)^{-1}\phi(t)^{-1} + \phi(b)\phi(s)\phi(s)^{-1}\phi(t)^{-1}\\ &= \phi(a)\phi(s)^{-1} + \phi(b)\phi(t)^{-1}\\ &= \tilde{\phi}\left(\frac{a}{s}\right) + \tilde{\phi}\left(\frac{b}{t}\right)\end{aligned}$$

b) Suppose $\tilde{\phi}(\frac{a}{s}) = 0$ then clearly $a \in \ker(\phi) \implies \frac{a}{s} \in S^{-1}\ker(\phi)$. The converse is clear.

■

In the case that $A$ is an integral domain then generally everything becomes a lot simpler.

**Example 3.6.6 (Quotient Field)**
*Let $A$ be an integral domain then $A \setminus 0 = A^\star$ and we define the* ***quotient field***

$$\mathrm{Frac}(A) \coloneqq (A \setminus 0)^{-1}A$$

**Proposition 3.6.7 (Quotient Field contains all localisation)**
*Let $A$ be an integral domain, and* $\mathrm{Frac}(A)$ *the quotient field. Define another model for $S^{-1}A$ as follows*

$$S^{-1}A \coloneqq \left\{\frac{a}{s} \in \mathrm{Frac}(A) \;\middle|\; a \in A\, s \in S\right\}$$

*The canonical map $A \to S^{-1}A \subset \mathrm{Frac}(A)$ is injective, and satisfies the universal property for localisation.*

*Proof.* It's injective because $A$ has no zero-divisors. That it satisfies the universal property is very similar as before.■

**Proposition 3.6.8 (Directed Limit)**
*Let $S_i$ be a family of multiplicatively closed sets directed by inclusion, such that $S = \bigcup_i S_i$ is multiplicatively closed. Then there is a canonical isomorphism*

$$\varinjlim_i S_i^{-1}A \to S^{-1}A$$

*induced by the canonical maps*

© David Rufino CC BY-NC-ND 4.0

$$S_i^{-1}A \to S^{-1}A$$

*Proof.* The canonical maps $i_{S_iS}$ induce a unique morphism

$$\varinjlim_i S_i^{-1}A \to S^{-1}A$$
$$[a_i/s_i] \to a_i/s_i$$

by the universal property. An element on the right hand side is written $a/s$ for some $s \in S$. By hypothesis $s \in S_i$ for some $i$, therefore it is surjective. Suppose we have two elements $[a_i/s_i]$ and $[a_j/s_j]$ on the left hand side which become equal in $S^{-1}A$. Then by definition $s_k(s_ja_i - a_js_i) = 0$ for some $s_k \in S_k$. Since it's a directed system we can find $S_l$ containing $S_i, S_j, S_k$. Then by definition $a_i/s_i = a_j/s_j$ in $S_l^{-1}A$ and we see that $[a_i/s_i] = [a_j/s_j]$. Therefore the given morphism is also injective as required. ∎

**Lemma 3.6.9 (Localisation Respects Contraction of Multiplicative Sets)**
*Let $A \subseteq B$ be subrings of a common ring $C$, and $T \subset B$ a multiplicative set. Writing $S := T \cap A$ (a multiplicative set of $A$), and realising both localisations as subrings of $C$ (or of a common ring of fractions), $S^{-1}A \subseteq T^{-1}B$.*

*In particular, for $\mathfrak{q}$ a prime ideal of $B$ and $\mathfrak{p} := \mathfrak{q} \cap A$, taking $T := B \setminus \mathfrak{q}$ (so $S = A \setminus \mathfrak{p}$) gives $A_\mathfrak{p} \subseteq B_\mathfrak{q}$.*

*Proof.* For $a/s \in S^{-1}A$ ($a \in A$, $s \in S \subseteq T$), viewing $a, s \in B$ (as $A \subseteq B$), $s \in T$ directly gives $a/s \in T^{-1}B$. ∎

Combining the previous two results, for a directed union of subrings rather than a single fixed ring:

**Lemma 3.6.10 (Localisation Commutes with Directed Unions of Subrings)**
*Let $(A_i)_{i\in I}$ be a directed system of subrings of a common ring $C$, with union $A := \bigcup_i A_i$, and $T \subset A$ a multiplicative set. Writing $T_i := T \cap A_i$, and realising all localisations as subrings of $C$ (or of a common ring of fractions),*

$$T^{-1}A = \bigcup_i T_i^{-1}A_i$$

*In particular, for $\mathfrak{p} \lhd A$ a prime ideal and $\mathfrak{p}_i := \mathfrak{p} \cap A_i$, taking $T := A \setminus \mathfrak{p}$ (so $T_i = A_i \setminus \mathfrak{p}_i$) gives $A_\mathfrak{p} = \bigcup_i (A_i)_{\mathfrak{p}_i}$.*

*Proof.* If $a/s \in T^{-1}A$ ($a \in A$, $s \in T$), then $a, s \in A_i$ for some $i$ (directed union), with $s \in T \cap A_i = T_i$, so $a/s \in T_i^{-1}A_i$. Conversely if $a/s \in T_i^{-1}A_i$ ($a, s \in A_i$, $s \in T_i \subseteq T$), then viewing $a, s \in A_i \subset A$, $s \in T$, so $a/s \in T^{-1}A$. ∎

## 3.6.2 Modules

**Definition 3.6.11 (Localisation of a module)**
*Let $A$ be a ring with $S$ multiplicative set and $M$ an $A$-module. Then we define*

$$S^{-1}M = \left\{ \frac{m}{s} \;\middle|\; m \in M \right\}$$

*under the obvious equivalence relation. This is then an $S^{-1}A$-module in the obvious way.*

**Definition 3.6.12 (Localisation of a sub-module)**
*Let $M$ be an $A$-module and $N \subseteq M$ a sub-$A$-module then define*

$$S^{-1}N = \left\{ \frac{n}{s} \;\middle|\; n \in M\ s \in S \right\} \subseteq S^{-1}M$$

**Proposition 3.6.13**
*$S^{-1}(-)$ constitutes a functor $A$-**Mod** $\to S^{-1}A$-**Mod**. More precisely there is a unique morphism $\psi$ making the following diagram commute as $A$-module morphisms*

$$\begin{array}{ccc} N & \xrightarrow{\psi} & M \\ \downarrow{\scriptstyle i_S} & & \downarrow{\scriptstyle i_S} \\ S^{-1}N & \xrightarrow{S^{-1}(\psi)} & S^{-1}M \end{array}$$

*where $S^{-1}(\psi)$ is in fact an $S^{-1}A$-module morphism.*

*It is an exact functor; for an exact sequence*

$$N \to M \to P$$

*the corresponding sequence of $S^{-1}A$-module morphisms*

© David Rufino CC BY-NC-ND 4.0

$$S^{-1}N \to S^{-1}M \to S^{-1}P$$

*is exact. If $N$ is a submodule of $M$ then we may regard $S^{-1}N$ as a submodule of $S^{-1}M$.*

**Proposition 3.6.14 (Localisation commutes with quotients)**
*There is a commutative diagram of $A$-module morphisms*

$$\begin{array}{ccccccccc} 0 & \longrightarrow & N & \xrightarrow{i} & M & \xrightarrow{\pi} & M/N & \longrightarrow & 0 \\ & & \downarrow{\scriptstyle i_S} & & \downarrow{\scriptstyle i_S} & & \downarrow & & \\ 0 & \longrightarrow & S^{-1}N & \longrightarrow & S^{-1}M & \xrightarrow{S^{-1}(\pi)} & S^{-1}(M/N) & \longrightarrow & 0 \end{array}$$

*with exact rows and the bottom row consists of $S^{-1}A$-module morphisms. This induces an isomorphism of $S^{-1}A$-modules.*

$$S^{-1}M/S^{-1}N \cong S^{-1}(M/N)$$

**Proposition 3.6.15**
*Suppose $N \subseteq N' \subseteq M$ then there is a canonical short-exact sequence of $S^{-1}A$-modules*

$$0 \to S^{-1}(N'/N) \to S^{-1}(M/N) \to S^{-1}(M/N') \to 0$$

*which induces an isomorphism*

$$S^{-1}(M/N)/S^{-1}(N'/N) \cong S^{-1}(M/N')$$

**Proposition 3.6.16**
*Let $M$ be a finitely-generated $A$-module. Then*

$$S^{-1}M = 0 \iff sM = 0 \text{ some } s \in S$$

### 3.6.3 Ideals

Recall the notion of extended and contracted ideals in Definition (3.3.49).

**Definition 3.6.17 (Localisation of an ideal)**
*Let $A$ be a ring and $S$ a multiplicative set and $\mathfrak{a} \lhd A$ define*

$$S^{-1}\mathfrak{a} := \left\{ \frac{a}{s} \;\middle|\; a \in \mathfrak{a} \right\}$$

*then this is an ideal of $S^{-1}A$.*

**Proposition 3.6.18 (Extension and Contraction)**
*Let $A$ be a ring with multiplicative set $S$ and canonical morphism $i_S : A \to S^{-1}A$.*

*a)* $\mathfrak{a}^e = i_S(\mathfrak{a})S^{-1}A = \left\{ \frac{a}{s} \mid a \in \mathfrak{a}, s \in S \right\} = S^{-1}\mathfrak{a}$

*b)* $\mathfrak{b}^c = \left\{ a \mid \frac{a}{1} \in \mathfrak{b} \right\}$

*c) An ideal $\mathfrak{a}$ in $A$ satisfies*

$$\mathfrak{a}^{ec} = \bigcup_{s \in S} (\mathfrak{a} : s) = \{ a \in A \mid as \in \mathfrak{a} \text{ some } s \in S \}$$

*In particular $\mathfrak{a}$ is contracted if and only if $as \in \mathfrak{a} \wedge s \in S \implies a \in \mathfrak{a}$*

*d)* $\mathfrak{b}$ proper $\iff \mathfrak{b}^c$ proper $\iff \mathfrak{b}^c \cap S = \varnothing$

*e)* $\mathfrak{a}^e$ proper $\iff \mathfrak{a} \cap S = \varnothing$

*f) Every ideal $\mathfrak{b} \lhd S^{-1}A$ is extended (equiv. $\mathfrak{b} = \mathfrak{b}^{ce} = S^{-1}\mathfrak{b}^c$).*

*g) A prime ideal $\mathfrak{p}$ is contracted if and only if $\mathfrak{p} \cap S = \varnothing$. In this case $\mathfrak{p}^e$ is prime.*

*h)* $\mathfrak{q}$ prime $\implies \mathfrak{q}^c$ *is prime and satisfies* $\mathfrak{q}^c \cap S = \varnothing$.

*Proof.*

© David Rufino CC BY-NC-ND 4.0

a) $S^{-1}\mathfrak{a}$ is an additive subgroup because $\frac{a_1}{s_1}+\frac{a_2}{s_2}=\frac{a_1s_2+a_2s_1}{s_1s_2}$. It contains $i_S(\mathfrak{a})$ and is closed under multiplication by $A$, therefore $\mathfrak{a}^e \subseteq S^{-1}\mathfrak{a}$. Similarly as $\mathfrak{a}^e$ is an ideal containing $i_S(\mathfrak{a})$, we have $\frac{a}{s}=\frac{1}{s}\cdot\frac{a}{1}\in\mathfrak{a}^e$, i.e. $S^{-1}\mathfrak{a}\subseteq\mathfrak{a}^e$ as required.

b) This is clear

c) Observe that

$$\begin{aligned}\mathfrak{a}^{\mathrm{ec}} &= \left\{a\in A \;\middle|\; \frac{a}{1}\in\mathfrak{a}^e\right\}\\ &= \left\{a\in A \;\middle|\; \frac{a}{1}=\frac{a'}{s} \quad a'\in\mathfrak{a}\, s\in S\right\}\\ &= \{a\in A \mid sa\in\mathfrak{a} \text{ some } s\in S\}\end{aligned}$$

By (3.3.50) an ideal $\mathfrak{a}$ is contracted if and only if $\mathfrak{a}=\mathfrak{a}^{\mathrm{ec}}$. Furthermore it always satisfies $\mathfrak{a}^{\mathrm{ec}}\subseteq\mathfrak{a}$. The reverse inclusion is precisely the condition given.

d) This first equivalence is true in general, see (3.3.50). Clearly $\mathfrak{b}^c=A\implies\mathfrak{b}^c\cap S\neq\varnothing$. Similarly if $S\cap\mathfrak{b}^c\neq\varnothing$ then $s\in\mathfrak{b}^c\implies\frac{s}{1}\in\mathfrak{b}\implies 1\in\mathfrak{b}\implies 1\in\mathfrak{b}^c$.

e) By d) $\mathfrak{a}^e$ is proper if and only if $\mathfrak{a}^{\mathrm{ec}}$ is proper. By c) we see $1\in\mathfrak{a}^{\mathrm{ec}}$ if and only if $S\cap\mathfrak{a}\neq\varnothing$ and the result follows.

f) By (3.3.50) we need only show $\mathfrak{b}^{\mathrm{ce}}\subseteq\mathfrak{b}$. Note $\frac{a}{s}\in\mathfrak{b}^{\mathrm{ce}}\implies\frac{a}{s}=\frac{a'}{s'}$ with $a'\in\mathfrak{b}^c$. By 2. $\frac{a'}{1}\in\mathfrak{b}$ and therefore so is $\frac{a}{s}=\frac{a'}{s'}=\frac{a'}{1}\cdot\frac{1}{s'}\in\mathfrak{b}$ as required.

g) If $\mathfrak{p}\cap S=\varnothing$ then by primality it automatically satisfies the conditions in c) and is therefore contracted. Conversely if a prime ideal $\mathfrak{p}$ is contracted then $\mathfrak{p}=\mathfrak{q}^c$. It is by definition proper so by d) it satisfies $\mathfrak{p}\cap S=\varnothing$ as required.

Suppose $\mathfrak{p}$ is prime and $\frac{a}{s}\cdot\frac{b}{t}\in\mathfrak{p}^e$ then $\frac{ab}{st}=\frac{x}{u}$ for $x\in\mathfrak{p}\implies v(abu-xst)=0\implies uvab\in\mathfrak{p}\implies a\in\mathfrak{p}$ or $b\in\mathfrak{p}$. Therefore $\frac{a}{s}\in\mathfrak{p}^e$ or $\frac{b}{t}\in\mathfrak{p}^e$ as required.

h) Generically $\mathfrak{q}^c$ is a contracted prime ideal and we've already shown in d) that $\mathfrak{q}^c\cap S=\varnothing$.

■

**Corollary 3.6.19 (Ideal Structure Localisation)**
*Let $A$ be a ring and $S$ a multiplicative set then there is an order-preserving bijection of proper ideals*

$$\{\mathfrak{a}\lhd A \mid \mathfrak{a} \text{ contracted}\}\leftrightarrow\{\mathfrak{b}\lhd S^{-1}A\}$$

*which restricts to a bijection of prime ideals*

$$\{\mathfrak{p}\lhd A \mid \mathfrak{p}\cap S=\varnothing\}\leftrightarrow\{\mathfrak{q}\lhd S^{-1}A\}$$

*Proof.* From (3.6.18) every ideal of $S^{-1}A$ is extended. Therefore the bijection of proper ideals follows from (3.3.53). For prime ideals each direction is well-defined by (3.6.18) g). ■

## 3.6.4 Change of Rings

For what follows it is useful to have the concept of saturation of a multiplicatively closed set. Essentially taking the saturation $\overline{S}$ of $S$ doesn't change the ring $S^{-1}A$.

**Proposition 3.6.20 (Saturation)**
*Let $A$ be a ring and $S$ a multiplicatively closed set. Then the following sets are equal*

*a)* $(i_S)^{-1}\left((S^{-1}A)^\star\right)$
*b)* $\{x\in A \mid ax\in S \text{ for some } a\in A\}$
*c)* $\bigcap_{T\supseteq S:T \text{ saturated}} T$

*which we denote by $\overline{S}$. We have the following properties*

- *$\overline{S}$ is saturated.*
- *$S$ is saturated if and only if $S=\overline{S}$*
- *$\overline{\overline{S}}=\overline{S}$.*

*Proof.* Note $x\in(i_S)^{-1}\left((S^{-1}A)^\star\right)\implies\frac{x}{1}\cdot\frac{b}{t}=1\implies s(xb-t)=0\implies(sb)x\in S$. Similarly if $ax\in S$ then $\frac{x}{1}\cdot\frac{a}{ax}=1$.

© David Rufino CC BY-NC-ND 4.0

It's clear from b) that the set thus defined is saturated and multiplicatively closed. Let $T$ be another saturated multiplicatively closed set containing $S$ and suppose $ax \in S \implies ax \in T \implies x \in T$, so we find that the sets are equal.

We've proved that $\overline{S}$ is saturated. Clearly $S = \overline{S}$ implies $S$ is saturated. Conversely if $S$ is saturated then by c) we have $\overline{S} \subseteq S$, and clearly $S \subseteq \overline{S}$. The final part follows easily. ∎

We also give another characterization of saturated multiplicatively closed subsets

**Proposition 3.6.21**
*Let $A$ be a ring and $S$ a multiplicatively closed subset. Then*

$$\overline{S} = A \setminus \bigcup_{\mathfrak{p} \cap S = \varnothing} \mathfrak{p}$$

*Proof.* Denote the right hand side by $T$. Then clearly $S \subseteq T$ and as noted before in (3.6.2) $T$ is saturated. Therefore $\overline{S} \subseteq T$ by (3.6.20) c).

Conversely suppose $a \notin \overline{S}$. Consider the principal ideal $(a)$ then $(a) \cap S = \varnothing$ (because $ab \in S \implies a \in \overline{S}$ by (3.6.20) b)). Therefore by (3.3.41) there is a prime ideal $\mathfrak{p}$ containing $a$ which does not intersect $S$. Therefore $a \notin T$. Contrapositively $T \subseteq \overline{S}$ as required. ∎

**Proposition 3.6.22 (Change of Rings)**
*Let $\phi : A \to B$ be a ring homomorphism, $S, T$ corresponding multiplicative subsets. Then*

- *There exists a morphism $\tilde{\phi}$ making the diagram commute*

$$\begin{array}{ccc} A & \xrightarrow{\phi} & B \\ \downarrow{\scriptstyle i_S} & & \downarrow{\scriptstyle i_T} \\ S^{-1}A & \overset{\tilde{\phi}}{\dashrightarrow} & T^{-1}B \end{array}$$

  *if and only if $\phi(S) \subseteq \overline{T}$. In this case it is unique and given by*

$$\tilde{\phi}\left(\frac{a}{s}\right) = \frac{\phi(a)b'}{\phi(s)b'}$$

  *where $b' \in B$ is any $b'$ such that $\phi(s)b' \in T$.*

- *If in addition $T \subseteq \phi\left(\overline{S}\right)$ then $\phi$ injective (resp. surjective, bijective) implies $\tilde{\phi}$ is injective (resp. surjective, bijective)*

- *Further $\phi$ surjective $\implies \ker\left(\tilde{\phi}\right) = S^{-1}\ker(\phi)$.*

*Proof.*

- If $\tilde{\phi}$ is well-defined, then $i_T(\phi(S)) = \tilde{\phi}(i_S(S)) \subseteq \tilde{\phi}\left(\left(S^{-1}A\right)^\star\right) \subseteq \left(T^{-1}B\right)^\star$, which implies $\phi(S) \subseteq i_T^{-1}\left(\left(T^{-1}B\right)^\star\right) = \overline{T}$.

  Conversely if $\phi(S) \subseteq \overline{T}$ then $(i_T \circ \phi)(S) \subseteq \left(T^{-1}B\right)^\star$ therefore by (3.6.5) the morphism exists making the diagram commute.

  Note that

$$\tilde{\phi}\left(\frac{a}{s}\right) = \tilde{\phi}(i_S(a)i_S(s)^{-1}) = \tilde{\phi}(i_S(a))\tilde{\phi}(i_S(s))^{-1}$$

  so it is uniquely defined by the commutativity condition. Note that given $s \in S$, by (3.6.20) there exists $b' \in B$ such that $\phi(s)b' \in T$. In this case it's clear that $i_T(\phi(s))^{-1} = \frac{b'}{\phi(s)b'}$ from which the explicit form results.

- Suppose $T \subseteq \phi\left(\overline{S}\right)$ and $\phi$ is injective. Then $\tilde{\phi}\left(\frac{a}{s}\right) = 0 \implies t\phi(a) = 0$ for $t \in T$. Then there exists $s' \in \overline{S}$ and $x \in A$ such that $xs' \in S$ and $\phi(s') = t$. Therefore $\phi(as') = 0 \implies as' = 0 \implies a(xs') = 0 \implies \frac{a}{s} = 0$ as required.

  Similarly if $\phi$ is surjective and given $\frac{b}{t} \in T^{-1}B$ there exists $a \in A$ such that $\phi(a) = b$ and $s \in \overline{S}$ such that $\phi(s) = t$. Then $xs \in S$, $\phi(xs) \in \overline{T}$ and $\phi(yxs) \in T$ for some $x, y \in A$. Finally

$$\tilde{\phi}\left(\frac{axy}{sxy}\right) = \frac{\phi(axy)}{\phi(sxy)} = \frac{b}{t}$$

© David Rufino CC BY-NC-ND 4.0

as required.

- TODO

■

**Corollary 3.6.23**
*Let $A \xrightarrow{\phi} B \xrightarrow{\psi} C$ be a sequence of homomorphisms and $S, T, U$ be multiplicative sets such that $\phi(S) \subseteq \overline{T}$ and $\psi(T) \subseteq \overline{U}$, then in the notation of the previous Proposition*

$$\tilde{\psi} \circ \tilde{\phi} = \widetilde{\psi \circ \phi}$$

*Proof.* This follows from the uniqueness condition in (3.6.22). ■

**Corollary 3.6.24 (Localisation Maps)**
*Let $A$ be a ring and $S, T$ two multiplicative sets. Then TFAE*

- *There exists $i_{ST} : S^{-1}A \to T^{-1}A$ such that $i_{ST} \circ i_S = i_T$*
- *$S \subseteq \overline{T}$*

*In this case $i_{ST}$ is the unique such map. We have the transitivity relationships*

$$i_{TU} \circ i_{ST} = i_{SU}$$

$$i_{SS} = \mathbf{1}_{S^{-1}A}$$

*and furthermore $i_{ST}$ is an isomorphism if and only if $\overline{S} = \overline{T}$. In particular $i_{S\overline{S}}$ is an isomorphism.*

*Finally* $\ker(i_{ST}) = S^{-1}\ker(i_T)$.

*Proof.* This existence of $i_{ST}$ follows from (3.6.22) when considering the map $\phi = 1_A$. The transitivity and reflexive relationships follow from (3.6.23). ■

**Corollary 3.6.25 (Localisation commutes with quotient)**
*Let $A$ be a ring, $\mathfrak{a}$ an ideal and $S$ a multiplicative set. Then there exists a unique morphism making the diagram commute*

$$\begin{array}{ccc} A & \xrightarrow{\pi} & A/\mathfrak{a} \\ \downarrow{\scriptstyle i_S} & & \\ S^{-1}A & & \downarrow{\scriptstyle i_{\pi(S)}} \\ \downarrow{\scriptstyle \pi} & & \\ S^{-1}A/S^{-1}\mathfrak{a} & \xrightarrow{\sim} & \pi(S)^{-1}(A/\mathfrak{a}) \end{array}$$

*which is an isomorphism, and determined by*

$$\frac{a}{s} + S^{-1}\mathfrak{a} \to \frac{a+\mathfrak{a}}{s+\mathfrak{a}}$$

*Note that $S \cap \mathfrak{a} \neq \varnothing \iff S^{-1}A/S^{-1}\mathfrak{a} = 0 \iff \pi(S)^{-1}(A/\mathfrak{a}) = 0$.*

*When $\mathfrak{b} \supseteq \mathfrak{a}$ this restricts to a commutative diagram of $A$-modules*

$$\begin{array}{ccc} \mathfrak{b} & \xrightarrow{\pi} & \mathfrak{b}/\mathfrak{a} \\ \downarrow{\scriptstyle i_S} & & \\ S^{-1}\mathfrak{b} & & \downarrow{\scriptstyle i_{\pi(S)}} \\ \downarrow{\scriptstyle \pi} & & \\ S^{-1}\mathfrak{b}/S^{-1}\mathfrak{a} & \xrightarrow{\sim} & \pi(S)^{-1}(\mathfrak{b}/\mathfrak{a}) \end{array}$$

*and the bottom arrow is still an isomorphism of $S^{-1}A/S^{-1}\mathfrak{a}$-modules.*

© David Rufino CC BY-NC-ND 4.0

**Corollary 3.6.26 (Localisation commutes with quotient II)**
*Let $A$ be a ring, $\mathfrak{a}$ an ideal and $S$ a multiplicative set. Then there exists a unique morphism making the diagram commute*

$$\begin{array}{ccc} A & \xrightarrow{\pi} & A/\mathfrak{a} \\ \downarrow{\scriptstyle i_S} & & \vdots \\ S^{-1}A & \xrightarrow{\pi} & S^{-1}A/S^{-1}\mathfrak{a} \end{array}$$

*given by*

$$a + \mathfrak{a} \to \frac{a}{1} + S^{-1}\mathfrak{a}$$

*and it is an isomorphism precisely when every $s \in S$ is co-prime to $\mathfrak{a}$, i.e.*

$$(s) + \mathfrak{a} = A \quad \forall s \in S.$$

*When $\mathfrak{b} \supseteq \mathfrak{a}$ this restricts to a commutative diagram*

$$\begin{array}{ccc} \mathfrak{b} & \xrightarrow{\pi} & \mathfrak{b}/\mathfrak{a} \\ \downarrow{\scriptstyle i_S} & & \vdots \\ S^{-1}\mathfrak{b} & \xrightarrow{\pi} & S^{-1}\mathfrak{b}/S^{-1}\mathfrak{a} \end{array}$$

*which is an $A/\mathfrak{a}$-module morphism, and is an isomorphism when the condition (...) holds.*

*Proof.* The condition given for being an isomorphism is equivalent to $\pi(S) \subset (A/\mathfrak{a})^\star$, which is equivalent to $i_{\pi(S)}$ being an isomorphism. Therefore we may apply (3.6.25). ∎

**Corollary 3.6.27**
*Let $A$ be a ring, $\mathfrak{m}$ a maximal ideal and $S$ a multiplicative set such that $S \cap \mathfrak{m} = \varnothing$. Then there is an isomorphism*

$$A/\mathfrak{m} \to S^{-1}A/S^{-1}\mathfrak{m}$$

*In particular $S^{-1}\mathfrak{m}$ is a maximal ideal of $S^{-1}A$.*

**Proposition 3.6.28 (Transitivity)**
*Let $S \subset T$ be multiplicative subsets of $A$ and let*

$$i_S : A \to S^{-1}A$$

*be the localisation at $S$. Define $T_S := i_S(T)$. Then $T_S$ is multiplicative and there is a canonical isomorphism*

$$T^{-1}A \to (T_S)^{-1}(S^{-1}A) \to \left(\overline{T_S}\right)^{-1}(S^{-1}A)$$

*Furthermore if $T \subseteq U$ then $T_S \subseteq U_S$ there is a commutative diagram*

$$\begin{array}{ccccc} & & S^{-1}A & & \\ & \swarrow^{i_{ST}} & \downarrow{\scriptstyle i_{T_S}} & \searrow^{i_{\overline{T_S}}} & \\ T^{-1}A & \xrightarrow{\sim} & (T_S)^{-1}(S^{-1}A) & \xrightarrow{\sim} & \left(\overline{T_S}\right)^{-1}(S^{-1}A) \\ \downarrow{\scriptstyle i_{TU}} & & \downarrow & & \downarrow \\ U^{-1}A & \xrightarrow{\sim} & (U_S)^{-1}(S^{-1}A) & \xrightarrow{\sim} & \left(\overline{U_S}\right)^{-1}(S^{-1}A) \end{array}$$

## 3.6.5 Algebras

**Lemma 3.6.29 (Localisation is a Base Change)**
*Let $A$ be a ring, $S$ a multiplicative subset, and $M$ an $A$-module (3.6.11). Then there is a natural isomorphism of $S^{-1}A$-modules*

© David Rufino CC BY-NC-ND 4.0

$$S^{-1}A \otimes_A M \xrightarrow{\sim} S^{-1}M, \quad \frac{a}{s} \otimes m \mapsto \frac{am}{s}.$$

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

The map $\Phi : S^{-1}A \times M \to S^{-1}M$, $\Phi\left(\frac{a}{s}, m\right) := \frac{am}{s}$, is well-defined: if $\frac{a}{s} = \frac{a'}{s'}$ in $S^{-1}A$ then $u(as' - a's) = 0$ in $A$ for some $u \in S$, hence $u(as' - a's)m = 0$ in $M$, i.e. $u(s'(am) - s(a'm)) = 0$, i.e. $\frac{am}{s} = \frac{a'm}{s'}$ in $S^{-1}M$. $\Phi$ is clearly bi-additive, and $A$-balanced since $\Phi\left(\frac{a}{s} \cdot b, m\right) = \frac{abm}{s} = \Phi\left(\frac{a}{s}, bm\right)$ for $b \in A$; so by the universal property of the tensor product (3.4.4) it induces a group homomorphism

$$\phi : S^{-1}A \otimes_A M \to S^{-1}M, \quad \frac{a}{s} \otimes m \mapsto \frac{am}{s},$$

which is visibly $S^{-1}A$-linear.

Conversely define $\psi : S^{-1}M \to S^{-1}A \otimes_A M$ by $\psi\left(\frac{m}{s}\right) := \frac{1}{s} \otimes m$. This is well-defined: if $\frac{m}{s} = \frac{m'}{s'}$ in $S^{-1}M$ then $u(s'm - sm') = 0$ for some $u \in S$, and then, using $\frac{1}{s} \otimes m = \frac{us'}{uss'} \otimes m = \frac{1}{uss'} \otimes us'm$ (moving $us' \in A$ across the tensor),

$$\frac{1}{s} \otimes m - \frac{1}{s'} \otimes m' = \frac{1}{uss'} \otimes us'm - \frac{1}{uss'} \otimes usm' = \frac{1}{uss'} \otimes u(s'm - sm') = 0.$$

Then $\phi\left(\psi\left(\frac{m}{s}\right)\right) = \phi\left(\frac{1}{s} \otimes m\right) = \frac{m}{s}$, so $\phi \circ \psi = \mathbf{1}$. Also $\psi\left(\phi\left(\frac{a}{s} \otimes m\right)\right) = \psi\left(\frac{am}{s}\right) = \frac{1}{s} \otimes am = \frac{a}{s} \otimes m$ (again moving $a$ across the tensor), and since elementary tensors generate $S^{-1}A \otimes_A M$, $\psi \circ \phi = \mathbf{1}$. Hence $\phi$ is an isomorphism, with inverse $\psi$. ∎

**Corollary 3.6.30 (Localisation of an Algebra is a Base Change)**
*Let $A$ be a ring, $S$ a multiplicative subset, and $B$ an $A$-algebra. Then $S^{-1}B$ is naturally an $S^{-1}A$-algebra (via $S^{-1}A \to S^{-1}B$, $\frac{a}{s} \mapsto \frac{ab_0}{s}$ where $b_0 \in B$ is the image of $1_A$, the unique extension of $A \to B \to S^{-1}B$ along $i_S$ (3.6.5), as $i_S(S) \subset (S^{-1}B)^\star$), and the isomorphism of $S^{-1}A$-modules furnished by (3.6.29) (with $M := B$),*

$$S^{-1}A \otimes_A B \xrightarrow{\sim} S^{-1}B, \quad \frac{a}{s} \otimes b \mapsto \frac{ab}{s},$$

*is moreover an isomorphism of rings, where $S^{-1}A \otimes_A B$ carries the ring structure of (3.4.34).*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Write $\phi$ for the $S^{-1}A$-module isomorphism of (3.6.29). It remains only to check $\phi$ is multiplicative on elementary tensors, since it is already additive:

$$\phi\left(\left(\frac{a}{s} \otimes b\right) \cdot \left(\frac{a'}{s'} \otimes b'\right)\right) = \phi\left(\frac{aa'}{ss'} \otimes bb'\right) = \frac{aa'bb'}{ss'} = \frac{a}{s} \cdot b \cdot \frac{a'}{s'} \cdot b' = \phi\left(\frac{a}{s} \otimes b\right) \cdot \phi\left(\frac{a'}{s'} \otimes b'\right).$$

As every element of $S^{-1}A \otimes_A B$ is a sum of elementary tensors, and $\phi$ is additive, $\phi$ is a ring homomorphism; being also a bijection, it is a ring isomorphism. ∎

**Corollary 3.6.31 (Localisation of a Domain is a Base Change to the Fraction Field)**
*Let $A \subseteq B$ be an inclusion of integral domains, $K := \mathrm{Frac}(A)$, and $S := A \setminus \{0\}$, so that $S^{-1}B$ is naturally a subring of $\mathrm{Frac}(B)$ (as $S \subset B \setminus \{0\}$ too). Then there is a natural isomorphism of rings*

$$B \otimes_A K \xrightarrow{\sim} S^{-1}B, \quad b \otimes x \mapsto bx,$$

*where the right-hand side is computed in $\mathrm{Frac}(B)$, regarding $B, K \subseteq \mathrm{Frac}(B)$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

By (3.6.30) (applied to the $A$-algebra $B$, noting $S^{-1}A = K$) there is a ring isomorphism $K \otimes_A B \xrightarrow{\sim} S^{-1}B$, $\frac{a}{s} \otimes b \mapsto \frac{ab}{s}$. Composing with the symmetry isomorphism $B \otimes_A K \cong K \otimes_A B$, $b \otimes x \mapsto x \otimes b$ (3.4.7) (which is also a ring isomorphism, since swapping factors clearly preserves the componentwise multiplication of (3.4.34)), yields a ring isomorphism $B \otimes_A K \xrightarrow{\sim} S^{-1}B$ sending $b \otimes \frac{a}{s} \mapsto \frac{ab}{s}$; taking $a = 1$ this is $b \otimes x \mapsto x \cdot b = bx$ for $x = \frac{1}{s}$, and since $x = \frac{a}{s} = a \cdot \frac{1}{s}$ for general $x \in K$, bilinearity gives $b \otimes x = ab \otimes \frac{1}{s} \mapsto \frac{ab}{s} = (ab) \cdot \frac{1}{s} = b \cdot \left(a \cdot \frac{1}{s}\right) = bx$ on every elementary tensor.

Finally $\frac{ab}{s} \in S^{-1}B \subseteq \mathrm{Frac}(B)$, computed there, agrees with the product $bx$ computed in $\mathrm{Frac}(B)$ (as $B, K \subseteq \mathrm{Frac}(B)$). ∎

**Remark 3.6.32**
*(3.6.30) recovers (3.6.41) (relative localisation at a prime) as the special case $S := A \setminus \mathfrak{p}$ of a general ring homomor-*

© David Rufino CC BY-NC-ND 4.0

*phism $\phi : A \to B$: the module localisation $S^{-1}B$ (with $B$ regarded as an $A$-module via $\phi$) is exactly $\phi(S)^{-1}B = B_{\mathfrak{p}}$, since the two defining equivalence relations on fractions agree termwise.*

### 3.6.6 Localisation at an element

**Definition 3.6.33 (Localisation at an element)**
*Let $A$ be a ring and $f \in A$. Then define*

$$S_f = \{1, f, ..., f^n, ...\}$$

*and*

$$A_f \coloneqq \left(S_f\right)^{-1} A$$

*We have canonical maps*

$$i_f : A \to A_f$$

*given by $i_f \coloneqq i_{S_f}$.*

*Note that $A_f$ is a zero ring iff $f$ is nilpotent.*

**Proposition 3.6.34 (Transition maps for localisation at an element)**
*Let $A$ be a ring and $f, g \in A$ then*

$$S_f \subseteq \overline{S_g} \iff f \mid g^N \text{ some N} > 0$$

*in this case define $i_{fg} = i_{S_f S_g}$ to be the unique morphism such that $i_{fg} \circ i_f = i_g$ (3.6.24). In addition if $h \in A$ and $S_g \subseteq \overline{S_h}$ we have a commutative diagram*




*Furthermore $\overline{S_1} = A^\star$ and $i_1$ is an isomorphism.*

**Proposition 3.6.35 (Transitivity of localizing at elements)**
*Let $A$ be a ring and $f, g \in A$ such that $S_f \subseteq \overline{S_g}$. There is a canonical isomorphism*

$$A_g \xrightarrow{\sim} \left(A_f\right)_{g/1}$$

*Furthermore $S_g \subseteq \overline{S_h} \implies S_{g/1} \subseteq \overline{S_{h/1}}$ and there is a commutative diagram*

$$\begin{array}{ccc}
\left(A_f\right)_1 & \xrightarrow{\sim} & A_f \\
\downarrow{\scriptstyle i_{1(g/1)}} & & \downarrow{\scriptstyle i_{g/1}} \\
\left(A_f\right)_{g/1} & \xrightarrow{\sim} & A_g \\
\downarrow & & \downarrow{\scriptstyle i_{gh}} \\
\left(A_f\right)_{h/1} & \xrightarrow{\sim} & A_h
\end{array}$$

*with the horizontal arrows isomorphisms and the vertical arrows are well-defined.*

*Proof.* The existence of the isomorphism is from (3.6.28) as $i_f\left(S_g\right) = S_{g/1}$. The second statement follows because $g \mid h^N \implies g/1 \mid h^N/1$ trivially. ∎

**Proposition 3.6.36**
*Let $A$ be a ring and $f \in A$. Then localisation map $A \to A_f$ induces an order-preserving correspondence of prime ideals*

$$\left\{\mathfrak{p} \lhd A \mid f \notin \mathfrak{p}\right\} \leftrightarrow \left\{\mathfrak{p} \lhd A_f\right\}$$

© David Rufino CC BY-NC-ND 4.0

*Proof.* The correspondence of prime ideals follows from (3.6.19) once we observe that $S_f \cap \mathfrak{p} = \varnothing \iff f \notin \mathfrak{p}$. ∎

**Proposition 3.6.37**
*Let $A$ be a ring, $S$ a multiplicatively closed subset and $g \in S$. Then the canonical map $A_g \to S^{-1}A$ has kernel $\ker(i_S)_g$. In particular when $\ker(i_S)$ is finitely generated it is injective iff $g^r \ker(i_S) = 0$ for some $r > 0$.*

*When $A$ is Noetherian then there always exists $g$ such $A_g \to S^{-1}A$ is injective, and any multiple of $g$ also satisfies this property.*

*Proof.* The first statement follows from (3.6.4).

When $A$ is Noetherian then $\ker(i_S) = (a_1, ..., a_n)$ for some $a_i \in A$. By definition there exists some $s_i$ such that $s_i a_i = 0$, whence we may take $g = s_1 \cdots s_n$. ∎

**Proposition 3.6.38**
*Let $A$ be a ring, $S$ a multiplicatively closed subset and $M_1, ..., M_n$ a family of $A$-modules. Then there is an isomorphism*

$$S^{-1}\left(\bigoplus_{i=1}^{n} M_i\right) \xrightarrow{\sim} \bigoplus_{i=1}^{n} S^{-1}M_i$$

## 3.6.7 Localisation at a prime ideal

**Definition 3.6.39 (Localisation at a prime ideal)**
*Let $A$ be a ring and $\mathfrak{p} \lhd A$ a prime ideal. Then $S := A \setminus \mathfrak{p}$ is a saturated multiplicatively closed subset, and we define*

$$A_{\mathfrak{p}} := (A \setminus \mathfrak{p})^{-1}A$$

*For an ideal $\mathfrak{a} \lhd A$ write the extended ideal*

$$\mathfrak{a}A_{\mathfrak{p}} := S^{-1}\mathfrak{a} = \mathfrak{a}^e.$$

**Definition 3.6.40 (Relative localisation at a prime ideal)**
*Let $\phi : A \to B$ be a ring homomorphism and $\mathfrak{p} \lhd A$ a prime ideal. Define*

$$B_{\mathfrak{p}} := \phi(A \setminus \mathfrak{p})^{-1}B$$

*For an ideal $\mathfrak{a} \lhd A$ write*

$$\mathfrak{a}B_{\mathfrak{p}} := (\phi(\mathfrak{a})B)B_{\mathfrak{p}}$$

*Observe $B_{\mathfrak{p}} = 0 \iff \mathfrak{p} \subsetneq \ker(\phi)$, so we would typically assume $\ker(\phi) \subseteq \mathfrak{p}$.*

**Proposition 3.6.41 (Relative Localisation as a Tensor Product)**
*Let $\phi : A \to B$ be a ring homomorphism (so $B$ is an $A$-algebra) and $\mathfrak{p} \lhd A$ a prime ideal, $S := A \setminus \mathfrak{p}$. Then $B_{\mathfrak{p}}$ is an $A_{\mathfrak{p}}$-algebra via $\frac{a}{s} \mapsto \phi(a)/\phi(s)$ (well-defined as $\phi(S) \subseteq B_{\mathfrak{p}}^{\star}$ already), and there is an isomorphism of $A_{\mathfrak{p}}$-algebras*

$$\Psi : B \otimes_A A_{\mathfrak{p}} \xrightarrow{\sim} B_{\mathfrak{p}}, \quad b \otimes \frac{a}{s} \mapsto \frac{\phi(a)b}{\phi(s)},$$

*with inverse $\Theta(b/\phi(s)) := b \otimes \left(\frac{1}{s}\right)$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

**$\Psi$ is well-defined.** For fixed $\frac{a}{s} \in A_{\mathfrak{p}}$, $b \mapsto \phi(a)b/\phi(s)$ is $A$-linear in $b$ ($B_{\mathfrak{p}}$ being an $A$-module via $\phi$); for fixed $b \in B$, if $\frac{a}{s} = \frac{a'}{s'}$ in $A_{\mathfrak{p}}$ then $s''(as' - a's) = 0$ for some $s'' \in S$, so applying $\phi$ and multiplying by $b$, $\phi(s'')(\phi(a)\phi(s') - \phi(a')\phi(s))b = 0$ in $B$, giving $\phi(a)b/\phi(s) = \phi(a')b/\phi(s')$ in $B_{\mathfrak{p}}$; and $\frac{a}{s} \mapsto \phi(a)b/\phi(s)$ is $A$-linear in $\frac{a}{s}$ (direct check, as in (3.6.5)). So the map $B \times A_{\mathfrak{p}} \to B_{\mathfrak{p}}$, $\left(b, \frac{a}{s}\right) \mapsto \phi(a)b/\phi(s)$, is $A$-bilinear, and induces $\Psi$ by the universal property of the tensor product (3.4.4).

**$\Theta$ is well-defined.** Write $a \cdot b := \phi(a)b$ for the $A$-module structure on $B$. If $b/\phi(s) = b'/\phi(s')$ in $B_{\mathfrak{p}}$ then $s'' \cdot (s' \cdot b) = s'' \cdot (s \cdot b')$ in $B$ for some $s'' \in S$; using $(a \cdot b) \otimes x = b \otimes (ax)$ (the defining relation of $\otimes_A$) twice,

$$b \otimes \frac{1}{s} = (s''s') \cdot b \otimes \frac{1}{s''s's} = (s'' \cdot (s' \cdot b)) \otimes \frac{1}{s''s's} = (s'' \cdot (s \cdot b')) \otimes \frac{1}{s''ss'} = b' \otimes \frac{1}{s'},$$

so $\Theta$ is well-defined on $B_{\mathfrak{p}}$.

© David Rufino CC BY-NC-ND 4.0

**$\Theta$ and $\Psi$ are mutually inverse.** $\Psi\Theta(b/\phi(s)) = \Psi\left(b \otimes \frac{1}{s}\right) = \phi(1)b/\phi(s) = b/\phi(s)$. And $\Theta\Psi\left(b \otimes \frac{a}{s}\right) = \Theta(\phi(a)b/\phi(s)) = (\phi(a)b) \otimes \left(\frac{1}{s}\right) = b \otimes \left(\frac{a}{s}\right)$ (the same tensor relation as above), so $\Theta\Psi$ is the identity on simple tensors, hence (by linearity) on all of $B \otimes_A A_\mathfrak{p}$.

**$\Psi$ is $A_\mathfrak{p}$-linear.** For $\frac{a'}{s'} \in A_\mathfrak{p}$,

$$\Psi\left(\frac{a'}{s'} \cdot \left(b \otimes \frac{a}{s}\right)\right) = \Psi\left(b \otimes \frac{aa'}{ss'}\right) = \frac{\phi(aa')b}{\phi(ss')} = \frac{\phi(a')}{\phi(s')} \cdot \frac{\phi(a)b}{\phi(s)} = \frac{a'}{s'} \cdot \Psi\left(b \otimes \frac{a}{s}\right),$$

using the $A_\mathfrak{p}$-algebra structure on $B_\mathfrak{p}$ displayed above; so $\Psi$ is $A_\mathfrak{p}$-linear on simple tensors, hence (by linearity) on all of $B \otimes_A A_\mathfrak{p}$.

**Multiplicativity.** By the algebra tensor product structure on $B \otimes_A A_\mathfrak{p}$, $\left(b \otimes \frac{a}{s}\right)\left(b' \otimes \frac{a'}{s'}\right) = bb' \otimes \frac{aa'}{ss'}$, and

$$\Psi\left(\left(b \otimes \frac{a}{s}\right)\left(b' \otimes \frac{a'}{s'}\right)\right) = \frac{\phi(aa')bb'}{\phi(ss')} = \frac{\phi(a)b}{\phi(s)} \cdot \frac{\phi(a')b'}{\phi(s')} = \Psi\left(b \otimes \frac{a}{s}\right)\Psi\left(b' \otimes \frac{a'}{s'}\right),$$

so $\Psi$ is multiplicative on simple tensors, hence (by linearity) a ring homomorphism – and so, with $A_\mathfrak{p}$-linearity above, an isomorphism of $A_\mathfrak{p}$-algebras. ∎

**Proposition 3.6.42**
*Let $A$ be a ring and $\mathfrak{p}$ a prime ideal. Consider the localisation $A \to A_\mathfrak{p}$. Then there is an order-preserving bijection between (prime) ideals contained in $\mathfrak{p}$ and (prime) ideals of $A_\mathfrak{p}$*

$$\{\mathfrak{q} \lhd A \mid \mathfrak{q} \subseteq \mathfrak{p}\} \leftrightarrow \{\mathfrak{q} \lhd A_\mathfrak{p}\}$$
$$\mathfrak{q} \to \mathfrak{q}A_\mathfrak{p}$$

*In particular $A_\mathfrak{p}$ is a local ring with unique maximal ideal $\mathfrak{p}A_\mathfrak{p}$.*

*Proof.* Clearly $\mathfrak{q} \cap (A \setminus \mathfrak{p}) = \varnothing \iff \mathfrak{q} \subseteq \mathfrak{p}$, so the result follows from (3.6.19). ∎

**Proposition 3.6.43**
*Let $A$ be a ring and $\mathfrak{p}$ a prime ideal. Then the map*

$$\mathrm{Frac}(A/\mathfrak{p}) \xrightarrow{\sim} A_\mathfrak{p}/\mathfrak{p}A_\mathfrak{p}$$
$$\frac{\overline{a}}{\overline{b}} \to \frac{\overline{a/1}}{\overline{b/1}} = \overline{\left(\frac{a}{b}\right)}$$

*is a ring isomorphism*

*Proof.* Consider the ring map $A \to A_\mathfrak{p} \to A_\mathfrak{p}/\mathfrak{p}A_\mathfrak{p}$. Then this has kernel $\mathfrak{p}^{\mathrm{ec}} = \mathfrak{p}$ (3.6.18), so may be factored as an injective map $A/\mathfrak{p} \hookrightarrow A_\mathfrak{p}/\mathfrak{p}A_\mathfrak{p}$. By (3.6.42) $A_\mathfrak{p}/\mathfrak{p}A_\mathfrak{p}$ is a field so by universal property of localisation (3.6.5) this extends to an injective map $\mathrm{Frac}(A/\mathfrak{p}) \hookrightarrow A_\mathfrak{p}/\mathfrak{p}A_\mathfrak{p}$ of the given form. We may verify directly that it is surjective. ∎

**Proposition 3.6.44 (Localisation at prime is direct limit of localisation at an element)**
*Let $A$ be a ring and $\mathfrak{p}$ a prime ideal then*

$$S_\mathfrak{p} := A \setminus \mathfrak{p} = \bigcup_{f \in A \setminus \mathfrak{p}} \overline{S_f}$$

*Therefore there are canonical morphisms (for $f \notin \mathfrak{p}$)*

$$i_{S_f S_\mathfrak{p}} : A_f \to A_\mathfrak{p}$$

*Furthermore the family of multiplicatively closed sets $\{S_f\}_{f \notin \mathfrak{p}}$ (resp. rings $\{A_f\}_{f \notin \mathfrak{p}}$) form a directed system under the relation $S \prec T \iff S \subseteq \overline{T}$. Therefore we have a canonical ring homomorphism*

$$\varinjlim_{f \notin \mathfrak{p}} A_f \to A_\mathfrak{p}$$

*which is an isomorphism.*

*Proof.* As $A \setminus \mathfrak{p}$ is a saturated multiplicatively closed set we have $f \in A \setminus \mathfrak{p} \iff S_f \subseteq A \setminus \mathfrak{p} \iff \overline{S_f} \subseteq A \setminus \mathfrak{p}$. Therefore the expression for $S_\mathfrak{p}$ follows.

The family of multiplicatively closed subsets is a directed system because $S_f \subseteq \overline{S_{fg}}$. To see this note $fg \in \overline{S_{fg}} \implies f \in \overline{S_{fg}} \implies S_f \subseteq \overline{S_{fg}}$.

© David Rufino CC BY-NC-ND 4.0

The final isomorphism follows because we can decompose it into two maps

$$\varinjlim_{f\notin\mathfrak{p}} A_f \xrightarrow{\sim} \varinjlim_{f\notin\mathfrak{p}} \overline{S_f}^{-1}A \xrightarrow{\sim} A_{\mathfrak{p}}$$

The first is an isomorphism by (3.6.24) and the second by (3.6.8), in light of the first statement. ∎

**Proposition 3.6.45 (Quotient commutes with Localisation at Prime ideal)**
*Let $A$ be a ring, $\mathfrak{a}$ an ideal and $\mathfrak{p}\supset\mathfrak{a}$ a prime ideal. Then there is an isomorphism of non-zero rings*

$$A_{\mathfrak{p}}/\mathfrak{a}_{\mathfrak{p}} \cong (A/\mathfrak{a})_{\mathfrak{p}/\mathfrak{a}}$$
$$\frac{a}{s}+\mathfrak{a}_{\mathfrak{p}} \to \frac{a+\mathfrak{a}}{s+\mathfrak{a}}$$

*Proof.* Let $\pi: A\to A/\mathfrak{a}$ be the quotient map then we claim $\pi(A\setminus\mathfrak{p})=\pi(A)\setminus\pi(\mathfrak{p})$. As $\pi$ is surjective we only need to show that $\pi^{-1}(\pi(\mathfrak{p}))=\mathfrak{p}$. That is if $\pi(a)=\pi(b)$ and $b\in\mathfrak{p}$ then $a\in\mathfrak{p}$. However this follows by assumption because $\mathfrak{a}\subset\mathfrak{p}$. Therefore we may apply (3.6.25). ∎

**Proposition 3.6.46 (Minimal Primes consist of Zero Divisors)**
*Let $\mathfrak{p}$ be a proper prime ideal minimal over $\mathfrak{a}$. Then for every $x\in\mathfrak{p}$ there exists $a\in A\setminus\mathfrak{a}$ such that $xa\in\mathfrak{a}$.*

*In particular a minimal prime ideal consists of zero-divisors.*

*Proof.* Observe that $\mathfrak{p}/\mathfrak{a}$ is a proper minimal prime ideal of $A/\mathfrak{a}$ by (3.3.56). Then the first statement is equivalent to showing every $x\in\mathfrak{p}/\mathfrak{a}$ is a zero divisor and we may reduce to the case $\mathfrak{a}=(0)$.

Consider the ring $A_{\mathfrak{p}}$, it has a unique maximal ideal $\mathfrak{p}A_{\mathfrak{p}}$ which is also minimal by (3.6.42). Every other prime ideal is contained in $\mathfrak{p}A_{\mathfrak{p}}$ by (3.3.37), and therefore equal by minimality. Therefore $\sqrt{(0)A_{\mathfrak{p}}}=\mathfrak{p}A_{\mathfrak{p}}$ by (3.3.46). Suppose $x\in\mathfrak{p}$ then $\frac{x^n}{1}=0$ and $tx^n\in\mathfrak{a}$ for $t\notin\mathfrak{p}$. Choose $n$ minimal subject to this condition. Then $n\geq 1$ because $t\notin\mathfrak{a}$, and so $a=tx^{n-1}$ has the required properties. ∎

## 3.6.8 Finiteness Results

**Lemma 3.6.47**
*Let $M$ be an $A$-module, $N$ a submodule and $f_1,...,f_r\in A$ such that*

*a) $N_{f_i}\to M_{f_i}$ is surjective for all $i=1...r$*
*b) $A=(f_1,...,f_r)$ as an ideal*

*then $M=N$.*

*Proof.* Consider the $A$-module $P:=M/N$. Suppose $P\neq 0$, then $0\neq p\in P$ and $\mathrm{Ann}(p)$ is a proper ideal of $A$. In particular $f_i\notin\mathrm{Ann}(p)$ for some $i$. This shows $P_{f_i}\neq 0$. From (3.6.14) we have $P_{f_i}\xrightarrow{\sim} M_{f_i}/N_{f_i}$ and so $N_{f_i}\neq M_{f_i}$ a contradiction. ∎

**Proposition 3.6.48**
*Let $M$ be an $A$-module. Then*

*a) $M$ finitely-generated $\implies M_f$ is a finitely generated $A_f$-module for all $f\in A$*
*b) $(f_1,...,f_r)=A$ and $M_{f_i}$ a finitely generated $A_{f_i}$-module for all $i=1...r\implies M$ is a finitely-generated $A$-module*

*Proof.*
a) Let $m_1,...m_n\in M$ be a generating set. Then it is easy to verify that $m_1/1,...,m_n/1$ is a generating set for $M_f$.
b) For each $i=1...r$, let $m_{ij}$ be such that $m_{ij}/1$ generates $M_{f_i}$ as an $A_{f_i}$-module. Let $N$ be the $A$-submodule of $M$ generated by the $m_{ij}$. Then $N_{f_i}=M_{f_i}$. We conclude by (3.6.47) that $N=M$ is finitely generated as an $A$-module.

∎

© David Rufino CC BY-NC-ND 4.0

## 3.7 Monoid Ring

**Definition 3.7.1 (Monoid Ring)**
*Let $A$ be a commutative ring and $G$ be a commutative monoid. We define the **monoid ring** to be the free abelian group*

$$A[G] := \bigoplus_{g \in G} A$$

*We may write a general element as a formal sum*

$$\sum_{g \in G} a_g g$$

*where all but finitely many $a_g$ are zero. Multiplication is defined extending the group operation. More precisely*

$$\left(\sum_{g \in G} a_g g\right) \cdot \left(\sum_{g \in G} b_g g\right) = \sum_{g \in G} \left( \sum_{(h,k) | h+k=g} a_h b_k g \right)$$

*We may verify that the identity is simply $e$ and the multiplication is both associative and distributive.*

*When $G$ is an abelian group we say that $A[G]$ is the **group ring**. There is a canonical map $A \to A[G]$ making $A[G]$ into an $A$-algebra.*

**Proposition 3.7.2**
*Let $A, B$ be commutative rings and $G$ a commutative monoid. Let $\phi : A \to B$ be a ring homomorphism and $\alpha : G \to (B, \times)$ a monoid homomorphism. Then there is a unique homomorphism*

$$\phi \ltimes \alpha : A[G] \to B$$

*such that $(\phi \ltimes \alpha)(ae) = \phi(a)$ and $(\phi \ltimes \alpha)(1_A g) = \alpha(g)$.*

*Proof.* Define

$$(\phi \ltimes \alpha)\left(\sum_{g \in G} a_g g\right) = \sum_{\substack{g \in G \\ a_g \neq 0}} \phi(a_g)\alpha(g)$$

■

© David Rufino CC BY-NC-ND 4.0

# 3.8 Polynomial Rings in One Variable

**Definition 3.8.1 (Polynomial Ring)**
*Let $A$ be a commutative ring. Define the polynomial ring $A[X]$ to be the monoid ring $A[\mathbb{N}]$ consisting of formal*

$$f(X) = \sum_{i=0}^{\infty} a_i X^i$$

*such that only finitely many $a_i$ are non-zero. Define degree in the obvious way*

$$\deg(f) = \inf\{n \mid m > n \implies a_m = 0\} < \infty$$

*and the leading coefficient to be $\ell(f) := a_{\deg(f)}$. By convention $\deg(0) = -\infty$.*

**Definition 3.8.2 (Monic polynomial)**
*Let $f \in A[X]$. We say $f$ is **monic** if the leading coefficient, $\ell(f)$, is $1$.*

**Lemma 3.8.3**
*If $A$ is an integral domain then for elements $f, g \in A[X]$*

$$\deg(fg) = \deg(f) + \deg(g)$$

$$\ell(fg) = \ell(f)\ell(g)$$

*Further $A[X]$ is an integral domain.*

**Proposition 3.8.4 (Nilpotent and Invertible Polynomials)**
*Let $A$ be a ring then*

*a) $N(A[X]) = N(A)[X] \subset A[X]$*
*b) $A[X]^\star = A^\star + XN(A)[X]$*

*Proof.* Suppose $a \in N(A)$ then clearly $aX^i$ is nilpotent. Therefore $N(A)[X] \subseteq N(A[X])$ since the nilradical is an ideal. Conversely suppose $f \in A[X]$ is nilpotent, i.e. $f^n = 0$. For any prime ideal $\mathfrak{p} \lhd A$ we find that $\overline{f}^n = 0$ as an element of $(A/\mathfrak{p})[X]$. As $A/\mathfrak{p}$ is an integral domain we have by the previous Lemma $\overline{f} = 0$. As $\mathfrak{p}$ was arbitrary and $N(A) = \bigcap \mathfrak{p}$ we see that $f \in N(A)[X]$ as required.

Suppose $f \in A[X]^\star$ and $fg = 1$, then clearly the constant term of $f$ must be invertible. Reduce modulo $\mathfrak{p}$ to find $\deg\left(\overline{f}\right) + \deg(\overline{g}) = 0 \implies \deg\left(\overline{f}\right) = \deg(\overline{g}) = 0$, which means $\overline{f}$ is a constant polynomial. As $\mathfrak{p}$ was arbitrary we see again that the other coefficients of $f$ must be nilpotent as required. Therefore $A[X]^\star \subseteq A^\star + XN(A)[X]$. Conversely by (3.3.68) and a) we have $A^\star + XN(A)[X] \subseteq A[X]^\star + N(A[X]) \subseteq A[X]^\star$ so the result follows. ∎

It satisfies the following universal property

**Proposition 3.8.5 (Evaluation at a point)**
*Consider an $A$-algebra $B$ and $b \in B$. Then there exists a unique $A$-algebra homomorphism*

$$\mathrm{ev}_b : A[X] \to B$$

*such that $\mathrm{ev}_b(X) = b$. We write $p(b) = \mathrm{ev}_b(p)$. It is given by*

$$p(b) = \sum_{k=0}^{\deg(p)} i_B(a_k) b^k$$

*The image of $\mathrm{ev}_p$ is equal to $A[b]$ the smallest sub-$A$-algebra generated by $b$. For any morphism $\phi : B \to C$ such that $\phi(b) = c$ we have*

$$\phi \circ \mathrm{ev}_b = \mathrm{ev}_c$$

**Remark 3.8.6**
*In categorical jargon $A[X]$ is an initial object in the category of pointed $A$-algebras.*

**Proposition 3.8.7 (Base Change of the Polynomial Ring)**
*Let $\phi : A \to B$ be a homomorphism of commutative rings. Then there is a natural isomorphism of $B$-algebras*

© David Rufino CC BY-NC-ND 4.0

$$A[X]_{(B)} \xrightarrow{\sim} B[X]$$
$$b \otimes \sum_i a_i X^i \to \sum_i b\phi(a_i)X^i$$

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

For any $B$-algebra $Z$ (in particular an $A$-algebra via $\phi$), combining (3.4.38) with the universal property of $A[X]$ (3.8.5) (applied to the underlying $A$-algebra of $Z$) gives a bijection natural in $Z$

$$\mathrm{AlgHom}_B\big(A[X]_{(B)}, Z\big) \cong \mathrm{AlgHom}_A(A[X], Z) \cong Z, \quad \psi \to \psi(1 \otimes X)$$

which is exactly the universal property of $B[X]$ (3.8.5) (applied with base ring $B$); by the Yoneda Lemma (2.6.58), this yields the required isomorphism (the image of $1_{A[X]_{(B)}}$ under it, sending $X \to 1 \otimes X$). ∎

**Proposition 3.8.8 (Evaluation commutes with algebra homomorphism)**
*Let $\phi : B \to C$ be a homomorphism of $A$-algebras and $p \in A[X]$ then*

$$\phi(p(b)) = p(\phi(b))$$

**Proposition 3.8.9 (Formal Derivative)**
*Let $A$ be a ring. Then there exists a unique $A$-module homomorphism*

$$(-)' : A[X] \to A[X]$$

*such that*

$$(X^r)' = \begin{cases} 0 & r = 0 \\ rX^{r-1} & r > 0 \end{cases}$$

*It satisfies the product rule*

$$(fg)' = f'g + fg'$$

*for all $f, g \in A[X]$. Define recursively $f^{(0)} = f$ and $f^{(n)} := f^{(n-1)'}$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

By (3.8.1), $A[X]$ is the free $A$-module on $\{1, X, X^2, ...\}$, so $(-)'$ exists and is unique.

Suppose $f(X) = \sum_{i=0}^n a_i X^i$ and $g(X) = \sum_{i=0}^m b_i X^i$. Then

$$(fg)(X) = \sum_{i=0}^{n+m} \left( \sum_{k+l=i} a_k b_l \right) X^i$$

and

$$\begin{aligned}
f'(X) &= \sum_{i=0}^{n-1} (i+1) a_{i+1} X^i \\
g'(X) &= \sum_{i=0}^{m-1} (i+1) b_{i+1} X^i \\
(fg)'(X) &= \sum_{i=0}^{n+m-1} (i+1) \left( \sum_{k+l=i+1} a_k b_l \right) X^i \\
&= \sum_{i=0}^{n+m-1} \left( \sum_{k+l=i+1} (k+l) a_k b_l \right) X^i \\
&+ \sum_{i=0}^{n+m-1} \left( \sum_{k+l=i} (k+1) a_{k+1} b_l \right) X^i + \sum_{i=0}^{n+m-1} \left( \sum_{k+l=i} (l+1) a_k b_{l+1} \right) X^i \\
&= f'(X)g(X) + f(X)g'(X)
\end{aligned}$$

∎

**Corollary 3.8.10 (Taylor Expansion mod $h^2$)**
*Let $A$ be a ring, $f \in A[X]$, and $a, h \in A$. Then*

© David Rufino CC BY-NC-ND 4.0

$$f(a+h) - f(a) - f'(a)h \in (h^2) \lhd A.$$

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Write $f(X) = \sum_{i=0}^{d} c_i X^i$. We first show, by induction on $i \geq 0$, that $(a+h)^i - a^i - ia^{i-1}h \in (h^2)$ (reading $a^{-1}h := 0$ for $i = 0$). The cases $i = 0, 1$ are immediate equalities ($1 - 1 - 0 = 0$, $(a+h) - a - h = 0$). Given $(a+h)^i = a^i + ia^{i-1}h + h^2 z$ for some $z \in A$:

$$(a+h)^{i+1} = (a+h)(a^i + ia^{i-1}h + h^2 z) = a^{i+1} + (i+1)a^i h + h^2(az + ia^{i-1} + hz),$$

so $(a+h)^{i+1} - a^{i+1} - (i+1)a^i h \in (h^2)$, completing the induction. Multiplying the case-$i$ congruence by $c_i$ and summing over $i = 0, ..., d$ (as $(h^2)$ is an ideal) gives $f(a+h) - f(a) - f'(a)h \in (h^2)$, noting $f'(a) = \sum_i c_i \cdot ia^{i-1}$ by (3.8.9) evaluated at $a$ (3.8.5). ∎

**Lemma 3.8.11 (Shifted Powers Form a Basis)**
*Let $k$ be a field, $\theta \in k$, and $n \geq 1$ an integer. Then $1, (X-\theta), ..., (X-\theta)^{n-1}$ is a $k$-basis for the space of polynomials in $k[X]$ of degree $< n$.*

*Proof.* Using (3.8.5) we may apply an isomorphism $X \to X - \theta$, from which the result follows immediately. ∎

**Definition 3.8.12 (Conjugate polynomial)**
*Let $\phi : A \to B$ be a homomorphism and $f \in A[X]$, then define*

$$f^\phi(X) := \sum_{i=0}^{n} \phi(a_i) X^i$$

*It induces a ring homomorphism*

$$A[X] \to B[X]$$

*and has the property that*

$$f^\phi(\phi(a)) = \phi(f(a))$$

**Proposition 3.8.13 (Division Algorithm I)**
*Let $A$ be an integral domain and $f(X) \in A[X]$ a polynomial and $g(X) \in A[X]$ a non-zero monic polynomial. Then there exists unique polynomials $q(X)$ and $r(X)$ such that*

- $f(X) = q(X)g(X) + r(X)$
- $\deg(r) < \deg(g)$

*In particular when $\deg(g) = 1$ then $r \in A$.*

*Proof.* If $\deg(f) < \deg(g)$ then $q = 0$ and $r = f$. Otherwise assume $n = \deg(f) \geq \deg(g) = m$ and proceed by induction on $n$. Note that since $g$ is monic then we have $f - \ell(f)gX^{n-m}$ has degree $n-1$, so by induction

$$f - \ell(f)gX^{n-m} = q'g + r$$

with $\deg(r) < \deg(g)$. Therefore

$$f = (q' + \ell(f)X^{n-m})g + r$$

as required.

Suppose $qg + r = q' + r'$ then $(q - q')g = (r' - r)$. This implies that $\deg((q-q')g) < \deg(g)$ which implies $(q-q')g = 0 \implies q = q'$ and $r = r'$. This demonstrates uniqueness. ∎

**Proposition 3.8.14**
*Let $B$ be an $A$-algebra and $f \in B$ not nilpotent. Then there is a canonical isomorphism*

$$B[X]/(1 - Xf) \to B_f$$
$$\overline{P(X)} \to P(1/f)$$

*In particular if $B$ is a finitely-generated $A$-algebra, then so is $B_f$.*

*Proof.* By (3.8.5) there is a $A$-algebra homomorphism $\phi : B[X] \to B_f$ such that $\phi(X) = \frac{1}{f}$. Evidently $(1 - Xf) \subset \ker(\phi)$ so by (...) this induces an $A$-algebra homomorphism $B'_f := B[X]/(1-Xf) \to B_f$. Evidently $\overline{X} = \overline{f}^{-1}$ in $B'_f$, so

© David Rufino CC BY-NC-ND 4.0

by (3.6.5) there is a map $\psi : B_f \to B[X]/(1 - Xf)$ given by $\frac{b}{f^r} \to \overline{bX^r}$. One may verify that $\psi$ is a two-sided inverse for $\phi$, and so is an isormophism.

The final statement follows from (3.3.76). ■

**Corollary 3.8.15**
*For $k$ a field, the units of $k[X]$ are precisely the non-zero polynomials of degree 0.*

*Proof.* As $k$ is a field, $N(k) = 0$, so by (3.8.4) $k[X]^\star = k^\star + X \cdot 0 = k^\star$, that is to say the non-zero constants. ■

**Proposition 3.8.16 (Division Algorithm over a field)**
*For $k$ a field consider the polynomial ring $k[X]$. For every pair of polynomials $f(X)$, $g(X)$ there exists unique polynomials $q(X)$ and $r(X)$ such that*

$$f(X) = q(X)g(X) + r(X)$$

*and* $\deg(r) < \deg(g)$.

*Proof.* Apply (3.8.13) to $g/\ell(g)$, and multiply by $\ell(g)$ again. ■

© David Rufino CC BY-NC-ND 4.0

## 3.9 Laurent Polynomials

**Definition 3.9.1 (Laurent polynomial ring)**
*Let $A$ be a commutative ring. Define the* ***laurent polynomial ring*** $A[X, X^{-1}]$ *to be the group ring $A[\mathbb{Z}]$. Denote an element as a formal sum*

$$f(X) \coloneqq \sum_{i=-\infty}^{\infty} a_i X^i$$

**Proposition 3.9.2**
*Let $B$ be an $A$-algebra and $b \in B^\star$ a unit. Then there is a unique $A$-algebra homomorphism $\mathrm{ev}_b : A[X, X^{-1}] \to B$ such that*

$$\mathrm{ev}_b(X) = b$$

*If the structural morphism $A \to B$ is injective, so is $\mathrm{ev}_b$.*

**Proposition 3.9.3**
*Let $A$ be a commutative ring. Then there is an isomorphism of rings*

$$\begin{aligned} A[X, X^{-1}] &\xrightarrow{\sim} A[X]_X \\ \sum_{i=-r}^{\infty} a_i X^i &\to \frac{\sum_{i=0}^{\infty} a_{i-r} X^i}{X^r} \quad r = \min\{k \mid a_k \neq 0\} \vee 0 \\ \sum_{i=-r}^{\infty} a_{i+r} X^i &\leftarrow \frac{\sum_{i=0}^{\infty} a_i X^i}{X^r} \end{aligned}$$

*Proof.* Denote the maps by $\psi$ and $\phi$. Then $\psi$ exists by (3.9.2) and $\phi$ exists by the universal property of localisation.■

© David Rufino CC BY-NC-ND 4.0

## 3.10 Formal Power Series and Laurent Series

**Definition 3.10.1 (Formal Laurent Series)**
*Let $A$ be a commutative ring. Define the **formal Laurent series ring** $A((X))$ to be the set of sequences $(c_n)_{n\in\mathbb{Z}}$ in $A$ such that there exists $N$ for which $n \leq N \implies c_n = 0$. Addition is defined pointwise with product defined as follows*

$$\left(\sum_{n\in\mathbb{Z}} c_n X^n\right)\left(\sum_{n\in\mathbb{Z}} d_n X^n\right) = \sum_{n\in\mathbb{Z}}\left(\sum_{k\in\mathbb{Z}} c_{n-k}d_k\right)X^n$$

*where the inner sum is a sum of a family with all but finitely many terms zero. That this is well-defined and makes $A((X))$ into a commutative ring is shown in (3.10.3).*

**Definition 3.10.2 (Order of a Formal Laurent Series)**
*Let $A$ be a commutative ring. For $0 \neq h = \sum_n c_n X^n \in A((X))$ define the **order** $v(h) := \min\{n \mid c_n \neq 0\}$, and $v(0) := \infty$. The minimum exists since the $c_n$ vanish for $n$ sufficiently negative and $h \neq 0$. We call $c_{v(h)}$ the **leading coefficient** of $h$.*

**Proposition 3.10.3 ($A((X))$ is a Commutative Ring)**
*Let $A$ be a commutative ring and $c, d \in A((X))$. For every $n \in \mathbb{Z}$ the family $(c_{n-k}d_k)_{k\in\mathbb{Z}}$ has all but finitely many terms zero, so $(cd)_n := \sum_{k\in\mathbb{Z}} c_{n-k}d_k$ is defined, and $(cd)_n = 0$ for $n < v(c) + v(d)$. In particular $cd \in A((X))$. Moreover $A((X))$ is a commutative ring under pointwise addition and this product.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

**The product is well-defined.** Let $c = (c_n)_{n\in\mathbb{Z}}$ and $d = (d_n)_{n\in\mathbb{Z}}$ be elements of $A((X))$. If $c = 0$ or $d = 0$ every term below is zero and the product is 0, so assume $v(c), v(d)$ are finite. Fix $n \in \mathbb{Z}$. If the term $c_{n-k}d_k$ is nonzero then $n - k \geq v(c)$ and $k \geq v(d)$, that is

$$v(d) \leq k \leq n - v(c)$$

so only finitely many $k \in \mathbb{Z}$ give a nonzero term. Hence $(c_{n-k}d_k)_{k\in\mathbb{Z}}$ has all but finitely many terms zero, and the inner sum $e_n := \sum_{k\in\mathbb{Z}} c_{n-k}d_k$ is defined by (3.2.50). Moreover if $n < v(c) + v(d)$ then there is no integer $k$ with $v(d) \leq k \leq n - v(c)$, so every term vanishes and $e_n = 0$. Therefore $(e_n)_{n\in\mathbb{Z}}$ is a sequence in $A$ vanishing for $n < v(c) + v(d)$, that is an element of $A((X))$.

**Ring axioms.** Write $(cd)_n := \sum_{k\in\mathbb{Z}} c_{n-k}d_k$ for the coefficients of the product, an element of $A((X))$ by the first part.

**Addition.** For $c, d \in A((X))$ we have $c_n + d_n = 0$ and $-c_n = 0$ for $n < \min(v(c), v(d))$, so $c + d$ and $-c$ lie in $A((X))$, as does the zero sequence. So $A((X))$ is a subgroup of the abelian group $A^{\mathbb{Z}}$ under pointwise addition.

**Identity.** Let $1 := (\delta_{n,0})_{n\in\mathbb{Z}}$, which vanishes for $n \leq -1$ and so lies in $A((X))$. For $c \in A((X))$ the family $(\delta_{n-k,0}c_k)_{k\in\mathbb{Z}}$ has $k = n$ as its only possibly nonzero term, so $(1c)_n = c_n$.

**Commutativity.** The map $k \mapsto n - k$ is a bijection of $\mathbb{Z}$, so by (3.2.52) and commutativity of $A$,

$$(cd)_n = \sum_{k\in\mathbb{Z}} c_{n-k}d_k = \sum_{k\in\mathbb{Z}} c_k d_{n-k} = \sum_{k\in\mathbb{Z}} d_{n-k}c_k = (dc)_n$$

**Distributivity.** For $c, d, d' \in A((X))$, the families $(c_{n-k}d_k)_{k\in\mathbb{Z}}$ and $(c_{n-k}d'_k)_{k\in\mathbb{Z}}$ have all but finitely many terms zero, so by (3.2.51)

$$(c(d + d'))_n = \sum_{k\in\mathbb{Z}} c_{n-k}(d_k + d'_k) = \sum_{k\in\mathbb{Z}} c_{n-k}d_k + \sum_{k\in\mathbb{Z}} c_{n-k}d'_k = (cd)_n + (cd')_n$$

and the other distributive law follows by commutativity.

**Associativity.** Let $c, d, e \in A((X))$ and fix $n \in \mathbb{Z}$; if any of them is 0 both sides are 0, so assume $v(c), v(d), v(e)$ are finite. Set $g(j, k) := c_{n-j-k}d_j e_k$ for $(j, k) \in \mathbb{Z} \times \mathbb{Z}$. If $g(j, k) \neq 0$ then $j \geq v(d)$, $k \geq v(e)$ and $n - j - k \geq v(c)$, so $v(d) \leq j \leq n - v(c) - v(e)$ and $v(e) \leq k \leq n - v(c) - v(d)$; hence $g$ has all but finitely many terms zero. Recall that $(cd)_m = \sum_{j\in\mathbb{Z}} c_{m-j}d_j$ and $(de)_m = \sum_{k\in\mathbb{Z}} d_{m-k}e_k$. Then

© David Rufino CC BY-NC-ND 4.0

$$\begin{aligned}((cd)e)_n &= \sum_{k\in\mathbb{Z}} (cd)_{n-k}e_k && \text{by definition of the product}\\ &= \sum_{k\in\mathbb{Z}}\left(\sum_{j\in\mathbb{Z}} c_{n-k-j}d_j\right)e_k && \text{by definition of } (cd)_{n-k}\\ &= \sum_{k\in\mathbb{Z}}\sum_{j\in\mathbb{Z}} g(j,k) && \text{by (3.3.17)}\\ &= \sum_{(j,k)\in\mathbb{Z}\times\mathbb{Z}} g(j,k) && \text{by (3.2.52)}\end{aligned}$$

and on the other side

$$\begin{aligned}(c(de))_n &= \sum_{l\in\mathbb{Z}} c_{n-l}(de)_l && \text{by definition of the product}\\ &= \sum_{l\in\mathbb{Z}} c_{n-l}\sum_{k\in\mathbb{Z}} d_{l-k}e_k && \text{by definition of } (de)_l\\ &= \sum_{l\in\mathbb{Z}}\sum_{k\in\mathbb{Z}} g(l-k,k) && \text{by (3.3.17)}\\ &= \sum_{(l,k)\in\mathbb{Z}\times\mathbb{Z}} g(l-k,k) && \text{by (3.2.52)}\\ &= \sum_{(j,k)\in\mathbb{Z}\times\mathbb{Z}} g(j,k) && \text{by (3.2.52) for the bijection } (l,k)\mapsto(l-k,k)\end{aligned}$$

Hence $((cd)e)_n = (c(de))_n$ for all $n$. ∎

**Lemma 3.10.4 (Valuation of a Sum and a Product)**
*Let $A$ be a commutative ring and $c, d \in A((X))$. Then $v(1) = 0$, $v(-c) = v(c)$, and*

*a) $v(c+d) \geq \min(v(c), v(d))$*
*b) $v(cd) \geq v(c) + v(d)$*
*c) if $A$ is an integral domain then so is $A((X))$, and $v(cd) = v(c) + v(d)$*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Clearly $v(1) = 0$ and $v(-c) = v(c)$. If $n < \min(v(c), v(d))$ then $c_n = d_n = 0$, so $(c+d)_n = 0$, which gives a). For b), if $c = 0$ or $d = 0$ then $cd = 0$ and both sides are $\infty$. Otherwise $(cd)_n = 0$ for $n < v(c) + v(d)$ by (3.10.3), which means $v(cd) \geq v(c) + v(d)$. For c), let $c, d \neq 0$ with $m := v(c)$ and $r := v(d)$. A term $c_{m+r-k}d_k$ of $(cd)_{m+r}$ is nonzero only if $k \geq r$ and $m + r - k \geq m$, that is $k = r$, so $(cd)_{m+r} = c_m d_r \neq 0$ as $A$ is a domain, while $(cd)_n = 0$ for $n < m + r$. Hence $cd \neq 0$ and $v(cd) = m + r$. ∎

**Definition 3.10.5 (Formal Power Series)**
*Let $A$ be a commutative ring. Define the **formal power series ring** $A[[X]]$ to be the subring of $A((X))$ consisting of those sequences $(c_n)_{n\in\mathbb{Z}}$ for which $n < 0 \implies c_n = 0$, or equivalently those $h \in A((X))$ for which $v(h) \geq 0$.*

*Proof that $A[[X]]$ is a subring of $A((X))$. Drafted by Claude Sonnet, reviewed by David Rufino.*

$A[[X]] = \{h \in A((X)) \mid v(h) \geq 0\}$, so by (3.10.4) it contains 1 and is closed under negation, addition and multiplication. ∎

**Definition 3.10.6 (Formal Derivative)**
*Let $A$ be a commutative ring. For $f = \sum_n c_n X^n \in A((X))$, define $f' := \sum_n n c_n X^{n-1} \in A((X))$, that is $(f')_n = (n+1)c_{n+1}$.*

**Lemma 3.10.7 (Valuation of a Derivative)**
*Let $A$ be a commutative ring and $f \in A((X))$. Then $v(f') \geq v(f) - 1$. In particular if $f \in A[[X]]$ then $f' \in A[[X]]$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

If $n < v(f) - 1$ then $n + 1 < v(f)$, so $c_{n+1} = 0$ and $(f')_n = (n+1)c_{n+1} = 0$. This gives $v(f') \geq v(f) - 1$. If $f \in A[[X]]$ then $v(f) \geq 0$ and $(f')_n = 0$ for $n < -1$; also $(f')_{-1} = 0 \cdot c_0 = 0$, so $f' \in A[[X]]$. ∎

**Definition 3.10.8 (Summable Family in $A((X))$)**
*Let $A$ be a commutative ring. A family $(h_i)_{i\in I}$ in $A((X))$ is **summable** if for every $N \in \mathbb{Z}$ all but finitely many $i \in I$ satisfy $v(h_i) \geq N$. Its **sum** is the formal Laurent series*

© David Rufino CC BY-NC-ND 4.0

$$\sum_{i\in I} h_i := \sum_{n\in\mathbb{Z}} \left( \sum_{i\in I} (h_i)_n \right) X^n$$

*where $(h_i)_n$ denotes the coefficient of $X^n$ in $h_i$.*

*Proof that the sum is well-defined. Drafted by Claude Sonnet, reviewed by David Rufino.*

Write $c_{i,n} := (h_i)_n$ and fix $n \in \mathbb{Z}$. By hypothesis (with $N = n+1$) the set $\{i \in I \mid v(h_i) \leq n\}$ is finite, and it contains every $i$ with $c_{i,n} \neq 0$. So $(c_{i,n})_{i\in I}$ has all but finitely many terms zero and $s_n := \sum_{i\in I} c_{i,n}$ is defined by (3.2.50). Let $J_0 := \{i \in I \mid v(h_i) \leq 0\}$, which is finite, and $N_0 := \min\big(0, \min_{i\in J_0} v(h_i)\big)$ (with $N_0 := 0$ if $J_0$ is empty). If $n < N_0$ then $v(h_i) > n$ for every $i$, so every $c_{i,n} = 0$ and $s_n = 0$. Hence $(s_n)_{n\in\mathbb{Z}}$ lies in $A((X))$. ∎

**Proposition 3.10.9 (Sums of Summable Families)**
*Let $A$ be a commutative ring and $(h_i)_{i\in I}$ a summable family in $A((X))$. Then*

*a) If $v(h_i) \geq N$ for every $i \in I$ then $v\big(\sum_{i\in I} h_i\big) \geq N$. In particular if every $h_i \in A[[X]]$ then $\sum_{i\in I} h_i \in A[[X]]$.*
*b) For a bijection $\sigma : I' \to I$ the family $\big(h_{\sigma(i')}\big)_{i'\in I'}$ is summable and $\sum_{i'\in I'} h_{\sigma(i')} = \sum_{i\in I} h_i$.*
*c) The family $(h_i')_{i\in I}$ is summable and $\big(\sum_{i\in I} h_i\big)' = \sum_{i\in I} h_i'$.*
*d) For any $g \in A((X))$ the family $(gh_i)_{i\in I}$ is summable and $g \cdot \sum_{i\in I} h_i = \sum_{i\in I} gh_i$.*
*e) If $(k_i)_{i\in I}$ is another summable family then $(h_i + k_i)_{i\in I}$ is summable and $\sum_{i\in I}(h_i + k_i) = \sum_{i\in I} h_i + \sum_{i\in I} k_i$.*
*f) If $I = \bigsqcup_{\lambda\in\Lambda} I_\lambda$ is a partition of $I$ into disjoint subsets then each $(h_i)_{i\in I_\lambda}$ and $\big(\sum_{i\in I_\lambda} h_i\big)_{\lambda\in\Lambda}$ is summable, and $\sum_{i\in I} h_i = \sum_{\lambda\in\Lambda} \sum_{i\in I_\lambda} h_i$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Write $c_{i,n} := (h_i)_n$ and $s_n := \sum_{i\in I} c_{i,n} = \big(\sum_{i\in I} h_i\big)_n$.

**a)** If $n < N$ then $c_{i,n} = 0$ for every $i$, so $s_n = 0$, as $J = \varnothing$ is a valid choice in (3.2.50). Hence $v\big(\sum_{i\in I} h_i\big) \geq N$, and taking $N = 0$ gives the statement for $A[[X]]$.

**b)** The reindexed family clearly meets the summability condition. The family $(h_i)_{i\in I}$ itself is not almost zero in $A((X))$, but for each $n$ the coefficient family $(c_{i,n})_{i\in I}$ in $A$ is, so the statement holds coefficientwise by (3.2.52).

**c)** Since $v(h_i') \geq v(h_i) - 1$ by (3.10.7) the family $(h_i')_{i\in I}$ is also summable, so $\sum_{i\in I} h_i'$ is defined. Using $(h')_n = (n+1)h_{n+1}$, for every $n \in \mathbb{Z}$

$$\begin{aligned}
\left( \left( \sum_{i\in I} h_i \right)' \right)_n &= (n+1)s_{n+1} = (n+1) \sum_{i\in I} c_{i,n+1} && \text{by definition of the sum} \\
&= \sum_{i\in I} (n+1) c_{i,n+1} && \text{by (3.3.17)} \\
&= \sum_{i\in I} (h_i')_n = \left( \sum_{i\in I} h_i' \right)_n && \text{by definition of the sum}
\end{aligned}$$

**d)** Let $g \in A((X))$. If $g = 0$ both sides are 0, so assume $v(g)$ is finite. By (3.10.4) $v(gh_i) \geq v(g) + v(h_i)$, so $(gh_i)_{i\in I}$ is summable and $\sum_{i\in I} gh_i$ is defined. Fix $n \in \mathbb{Z}$ and set $a(k,i) := g_{n-k} c_{i,k}$ for $(k,i) \in \mathbb{Z} \times I$. If $a(k,i) \neq 0$ then $k \leq n - v(g)$ and $v(h_i) \leq k$, so $i$ lies in the finite set $\{i \in I \mid v(h_i) \leq n - v(g)\}$ and $v(h_i) \leq k \leq n - v(g)$; hence $a$ has all but finitely many terms zero. Then

$$\begin{aligned}
\left( g \sum_{i\in I} h_i \right)_n &= \sum_{k\in\mathbb{Z}} g_{n-k} s_k = \sum_{k\in\mathbb{Z}} g_{n-k} \sum_{i\in I} c_{i,k} && \text{by definition of the product and the sum} \\
&= \sum_{k\in\mathbb{Z}} \sum_{i\in I} a(k,i) && \text{by (3.3.17)} \\
&= \sum_{i\in I} \sum_{k\in\mathbb{Z}} a(k,i) && \text{by (3.2.52)} \\
&= \sum_{i\in I} (gh_i)_n = \left( \sum_{i\in I} gh_i \right)_n && \text{by definition of the product and the sum}
\end{aligned}$$

**e)** By (3.10.4) a), $v(h_i + k_i) \geq \min(v(h_i), v(k_i))$, so for every $N$ all but finitely many $i$ satisfy $v(h_i + k_i) \geq N$ and $(h_i + k_i)_{i\in I}$ is summable. The identity holds in each coefficient by (3.2.51).

© David Rufino CC BY-NC-ND 4.0

**f)** The subfamilies are summable. For $N \in \mathbb{Z}$ the set $F_N := \{i \in I \mid v(h_i) < N\}$ is finite, so only finitely many $\lambda$ have $I_\lambda \cap F_N \neq \varnothing$, and for every other $\lambda$ we have $v\left(\sum_{i\in I_\lambda} h_i\right) \geq N$ by a). Hence $\left(\sum_{i\in I_\lambda} h_i\right)_{\lambda\in\Lambda}$ is summable. The identity holds in each coefficient by (3.2.51) applied to $\left(c_{i,n}\right)_{i\in I}$. ∎

**Proposition 3.10.10 (Units of a Formal Power Series Ring)**
*Let $A$ be a commutative ring and $u \in A[[X]]$, with constant coefficient $u_0$ (the coefficient of $X^0$). Then $u$ is a unit of $A[[X]]$ if and only if $u_0 \in A^\star$. In particular, when $A = k$ is a field, $u \in k[[X]]^\star \iff u_0 \neq 0$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Suppose $u_0 \in A^\star$ and set $w := 1 - u_0^{-1}u \in A[[X]]$. Its constant coefficient is 0, that is $v(w) \geq 1$, so $v(w^i) \geq i$ for all $i \geq 0$ by (3.10.4). Hence $\left(w^i\right)_{i\geq 0}$ is summable (3.10.8) and $s := \sum_{i\geq 0} w^i \in A[[X]]$ by (3.10.9). Then

$$\begin{aligned} s &= 1 + \sum_{i\geq 1} w^i && \text{splitting off } i = 0 \text{ by (3.10.9) } f) \\ &= 1 + \sum_{i\geq 0} w^{i+1} && \text{reindexing } i \mapsto i+1 \text{ by (3.10.9) } b) \\ &= 1 + ws && \text{distributivity, by (3.10.9) } d) \end{aligned}$$

so $(1-w)s = 1$. Since $u_0^{-1}u = 1 - w$, this says $u \cdot (u_0^{-1}s) = 1$, so $u_0^{-1}s$ is an inverse of $u$.

Conversely, if $u$ is a unit with inverse $t \in A[[X]]$ then $(ut)_0 = \sum_{k\in\mathbb{Z}} u_{-k}t_k = u_0t_0$, as $u_{-k} = 0$ for $k > 0$ and $t_k = 0$ for $k < 0$. So $u_0t_0 = 1$ and $u_0 \in A^\star$. When $A = k$ is a field, $u_0 \in k^\star$ is equivalent to $u_0 \neq 0$. ∎

**Proposition 3.10.11 (Units of a Formal Laurent Series Ring)**
*Let $A$ be an integral domain and $0 \neq h \in A((X))$ with $m := v(h)$. The following are equivalent*

*a) $h$ is a unit of $A((X))$*
*b) $h = X^m u$ for a unit $u \in A[[X]]^\star$*
*c) the leading coefficient $h_m$ of $h$ is a unit of $A$*

*In particular when $A = k$ is a field, $k((X))$ is a field.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

c) $\implies$ b): $u := X^{-m}h \in A[[X]]$ has constant coefficient $u_0 = h_m \in A^\star$, so $u$ is a unit of $A[[X]]$ by (3.10.10).

b) $\implies$ a): $X^{-m}u^{-1}$ is an inverse of $h = X^m u$.

a) $\implies$ c): let $hh' = 1$ in $A((X))$. Then $v(h) + v(h') = v(1) = 0$ by (3.10.4) c), so $v(h') = -m$, and the coefficient of $X^0$ in $hh'$ is $h_m h'_{-m} = 1$, so $h_m \in A^\star$. ∎

**Lemma 3.10.12 ($e$-th Roots of Units in a Formal Power Series Ring)**
*Let $k$ be a field, $e \geq 1$ an integer invertible in $k$ (i.e. $\mathrm{char}(k) \nmid e$), and $u \in k[[X]]^\star$. If $u(0)$ has an $e$-th root in $k$ (automatic if $k$ is algebraically closed), then $u$ has an $e$-th root in $k[[X]]^\star$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Let $\lambda \in k$ with $\lambda^e = u(0)$; replacing $u$ by $\lambda^{-e}u$ (whose $e$-th root, if found, gives one for $u$ after multiplying by $\lambda$), we may assume $u(0) = 1$. Construct $y = 1 + \sum_{n\geq 1} y_n X^n$ recursively: set $y^{(0)} := 1$, and given $y^{(n-1)} := 1 + \sum_{j=1}^{n-1} y_j X^j$ with $\left(y^{(n-1)}\right)^e \equiv u \pmod{X^n}$ (trivial for $n = 1$, as $u(0) = 1$), write $\left(y^{(n-1)}\right)^e = u + c_n X^n + O(X^{n+1})$ for some $c_n \in k$; by the binomial theorem,

$$\left(y^{(n-1)} + y_n X^n\right)^e \equiv \left(y^{(n-1)}\right)^e + e\left(y^{(n-1)}\right)^{e-1} y_n X^n \pmod{X^{n+1}}$$

(every other term has order $\geq 2n \geq n+1$), and $\left(y^{(n-1)}\right)^{e-1} \equiv 1 \pmod{X}$ (as $y^{(n-1)}(0) = 1$), so the coefficient of $X^n$ on the right is $c_n + ey_n$. As $e$ is invertible in $k$, setting $y_n := -c_n/e$ gives $y^{(n)} := y^{(n-1)} + y_n X^n$ with $\left(y^{(n)}\right)^e \equiv u \pmod{X^{n+1}}$, continuing the induction. The resulting $y := 1 + \sum_{n\geq 1} y_n X^n \in k[[X]]$ satisfies $y^e = u$ (agreeing with $u$ to every order), and is a unit as $y(0) = 1 \neq 0$ (3.10.10). ∎

**Proposition 3.10.13 ($k((X))$ is a Field of Fractions for $k[[X]]$)**
*Let $k$ be a field. Then every element of the field $k((X))$ is $a/b$ for $a, b \in k[[X]]$, $b \neq 0$ – that is, $k((X))$ is a field of fractions for the subring $k[[X]] \subset k((X))$. Moreover, the order function $v : k((X))^\star \to \mathbb{Z}$ is a group homomorphism under multiplication: $v(fg) = v(f) + v(g)$, and $v(f^{-1}) = -v(f)$.*

*Proof.* By (3.10.11), as $k$ is a field every $c \neq 0$ in $k((X))$ is $c = X^m u$ with $m = v(c)$ and $u \in k[[X]]^\star$, which is $uX^m \in k[[X]]$ if $m \geq 0$ and $u/X^{-m}$ otherwise, in either case a ratio of two elements of $k[[X]]$.

© David Rufino CC BY-NC-ND 4.0

By (3.10.4) c), as $k$ is a domain, $v(fg) = v(f) + v(g)$ for $f, g \neq 0$. Applying this to $f, f^{-1}$: $0 = v(1) = v(ff^{-1}) = v(f) + v(f^{-1})$, giving $v(f^{-1}) = -v(f)$. ∎

**Proposition 3.10.14 (Formal Power Series Ring is a DVR)**
*Let $k$ be a field. Then $k[[X]]$ is an integral domain, not a field, in which every nonzero element $a$ can be written uniquely as $a = X^{v(a)}u$ with $u \in k[[X]]^\star$ (equivalently, $u(0) \neq 0$ (3.10.10)).*

*Using the criteria (3.31.2)e) $k[[X]]$ is a discrete valuation ring with valuation precisely the order function $v$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

$k[[X]]$ is a domain, as a subring of the domain $k((X))$ ((3.10.4) c)), and not a field, as $X \neq 0$ has no inverse ($v(Xg) = 1 + v(g) \neq 0$ for every $g \in k[[X]]$ by the same result). Given $0 \neq a \in k[[X]]$, by (3.10.11) $a = X^{v(a)}u$ with $u \in k[[X]]^\star$, where $v(a) \geq 0$: this is exactly the uniformised-element criterion (3.31.2)e), with $\pi := X$. If also $a = X^n u'$ with $u' \in k[[X]]^\star$ then $v(u') = 0$ and $v(a) = n + v(u') = n$ by (3.10.4) c), so the exponent is $v(a)$ and then $u' = u$ as $k((X))$ is a domain. ∎

**Proposition 3.10.15 (Product Rule)**
*Let $A$ be a commutative ring and $f, g \in A((X))$. Then $(fg)' = f'g + fg'$.*

*Proof.* Write $f = \sum_i a_i X^i$, $g = \sum_j b_j X^j$. The coefficient of $X^{n-1}$ in $(fg)'$ is $n$ times the coefficient of $X^n$ in $fg$, that is $n\sum_{i+j=n} a_i b_j$. The coefficient of $X^{n-1}$ in $f'g$ is $\sum_{i+j=n}(ia_i)b_j$ (as $f'$ has coefficient $ia_i$ at degree $i-1$), and in $fg'$ is $\sum_{i+j=n} a_i(jb_j)$. Summing,

$$\sum_{i+j=n} ia_i b_j + \sum_{i+j=n} a_i j b_j = \sum_{i+j=n} (i+j)a_i b_j = n \sum_{i+j=n} a_i b_j$$

matching the coefficient of $X^{n-1}$ in $(fg)'$. As $n$ was arbitrary, $(fg)' = f'g + fg'$. ∎

**Lemma 3.10.16 (Power Rule)**
*Let $A$ be a commutative ring and $\psi \in A((X))$. Then $(\psi^n)' = n\psi^{n-1}\psi'$ for every $n \geq 0$. If moreover $\psi$ is a unit of $A((X))$, this also holds for every $n < 0$.*

*Proof.* **$n \geq 0$.** Induction on $n$: the case $n = 0$ is $0 = 0$ ($\psi^0 = 1$ is constant). Given $(\psi^n)' = n\psi^{n-1}\psi'$, the product rule (3.10.15) gives

$$(\psi^{n+1})' = (\psi \cdot \psi^n)' = \psi'\psi^n + \psi(\psi^n)' = \psi'\psi^n + n\psi^n\psi' = (n+1)\psi^n\psi'$$

**$n < 0$, $\psi$ a unit.** Differentiating $\psi\psi^{-1} = 1$ with the product rule, $0 = \psi'\psi^{-1} + \psi(\psi^{-1})'$, so $(\psi^{-1})' = -\psi^{-2}\psi'$ – the case $n = -1$. For $n \leq -2$, write $n = -r$, $r \geq 2$: applying the $n \geq 0$ case just shown to the element $\psi^{-1} \in A((X))$ (no hypothesis on its order was used there),

$$(\psi^{-r})' = \left((\psi^{-1})^r\right)' = r(\psi^{-1})^{r-1}(\psi^{-1})' = r(\psi^{-1})^{r-1}(-\psi^{-2}\psi') = -r\psi^{-r-1}\psi' = n\psi^{n-1}\psi'$$

∎

**Proposition 3.10.17 (Substitution into a Formal Laurent Series)**
*Let $A$ be an integral domain, $f = \sum_{n \geq n_0} c_n X^n \in A((X))$ and $\psi \in XA[[X]]$ (that is, $\psi(0) = 0$). Suppose that either*

- *$f \in A[[X]]$, that is $n_0 \geq 0$, or*
- *$\psi \in A((X))^\star$, that is $\psi \in X^m A[[X]]^\star$ for some $m > 0$ (3.10.11)*

*Then*

$$(f \circ \psi)(X) := \sum_{n \geq n_0} c_n \psi(X)^n$$

*is a well-defined element of $A((X))$ (with $\psi^n$, $n < 0$, the power in the ring $A((X))$), and $f \circ \psi \in A[[X]]$ if $f \in A[[X]]$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

As $\psi \in XA[[X]]$ has no terms of degree $\leq 0$ we have $v(\psi) \geq 1$, so $v(\psi^n) \geq n$ for every $n \geq 0$ by (3.10.4) b). Negative powers of $\psi$ only occur for the finitely many $n$ with $n_0 \leq n < 0$. Hence for every $N$ all $n \geq \max(N, 0)$ satisfy $v(c_n\psi^n) \geq n \geq N$, and only the finitely many $n$ with $n_0 \leq n < \max(N, 0)$ can fail this, so $(c_n\psi^n)_{n \geq n_0}$ is summable (3.10.8) and its sum lies in $A((X))$. If $f \in A[[X]]$ then every $c_n\psi^n \in A[[X]]$ and the sum lies in $A[[X]]$ by (3.10.9) a). ∎

© David Rufino CC BY-NC-ND 4.0

**Proposition 3.10.18 (Chain Rule)**
*In the setting of (3.10.17), $(f \circ \psi)' = (f' \circ \psi) \cdot \psi'$.*

*Proof.* By the power rule ((3.10.16), which for $n < 0$ requires that $\psi$ is a unit of $A((X))$, as it is then), $(c_n\psi^n)' = c_n(\psi^n)' = nc_n\psi^{n-1}\psi'$ for every $n \geq n_0$, using linearity of $(-)'$ (immediate from (3.10.6), as $c_n \in A$ is a constant). The term $n = 0$ is 0, and is simply omitted if $\psi$ is not a unit. Then, by termwise differentiation (3.10.9) c) of the summable family $(c_n\psi^n)_{n\geq n_0}$ ((3.10.17)'s proof),

$$(f \circ \psi)' = \sum_{n\geq n_0} (c_n\psi^n)' = \sum_{n\geq n_0} nc_n\psi^{n-1}\psi'$$

The family $(nc_n\psi^{n-1})_{n\geq n_0}$ is summable by the same order bound as $(c_n\psi^n)_{n\geq n_0}$ ((3.10.17)'s proof, shifted down one degree), so distributivity ((3.10.9) d)) gives

$$\sum_{n\geq n_0} nc_n\psi^{n-1}\psi' = \left(\sum_{n\geq n_0} nc_n\psi^{n-1}\right)\psi'$$

Reindexing $m = n - 1$, $\sum_{n\geq n_0} nc_n\psi^{n-1} = \sum_{m\geq n_0-1}(m+1)c_{m+1}\psi^m = f' \circ \psi$, giving $(f \circ \psi)' = (f' \circ \psi) \cdot \psi'$. ∎

**Proposition 3.10.19 (Substitution is a Ring Homomorphism)**
*Let $A$ be an integral domain and $\psi \in XA[[X]]$. Then $f \mapsto f \circ \psi$ is a ring homomorphism $A[[X]] \to A[[X]]$. If moreover the leading coefficient of $\psi$ is a unit (so $\psi \in A((X))^\star$ by (3.10.11)) then $f \mapsto f \circ \psi$ is a ring homomorphism $A((X)) \to A((X))$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

The arguments below apply to $f, g, h$ in $A[[X]]$ in the first case and in $A((X))$ in the second.

**Identity.** $1 \circ \psi = \psi^0 = 1$.

**Additivity.** Write $f = \sum_{n\geq n_0} c_nX^n$, $h = \sum_{n\geq m_0} d_nX^n$, and set $p_0 := \min(n_0, m_0)$, extending both coefficient sequences by $c_n := 0$ (resp. $d_n := 0$) for $p_0 \leq n < n_0$ (resp. $< m_0$); this changes neither series, nor the sums below, as it only adds zero terms. Then $f + h = \sum_{n\geq p_0}(c_n + d_n)X^n$, and the families $(c_n\psi^n)_{n\geq p_0}$ and $(d_n\psi^n)_{n\geq p_0}$ are summable by (3.10.17). Hence

$$\begin{aligned}(f+h)\circ\psi &= \sum_{n\geq p_0}(c_n+d_n)\psi^n = \sum_{n\geq p_0}(c_n\psi^n + d_n\psi^n) && \text{by distributivity in } A((X))\\ &= \sum_{n\geq p_0} c_n\psi^n + \sum_{n\geq p_0} d_n\psi^n = f\circ\psi + h\circ\psi && \text{by (3.10.9) } e)\end{aligned}$$

**Multiplicativity.** Write $f = \sum_{i\geq i_0} c_iX^i$, $g = \sum_{j\geq j_0} d_jX^j$, so $f \circ \psi = \sum_i c_i\psi^i$ and $g \circ \psi = \sum_j d_j\psi^j$, and let $P := \mathbb{Z}_{\geq i_0} \times \mathbb{Z}_{\geq j_0}$ and $a(i,j) := c_id_j\psi^{i+j}$ for $(i,j) \in P$. We have $v(a(i,j)) \geq i + j$ whenever $i + j \geq 0$ as $v(\psi^n) \geq n$ for $n \geq 0$ ((3.10.17)'s proof), and for every $N$ only the finitely many pairs with $i + j < \max(N, 0)$ can fail $v(a(i,j)) \geq N$, so $(a(i,j))_{(i,j)\in P}$ is summable. Then

$$\begin{aligned}(f\circ\psi)(g\circ\psi) &= \sum_{j\geq j_0}(f\circ\psi)d_j\psi^j && \text{by (3.10.9) } d)\\ &= \sum_{j\geq j_0}\sum_{i\geq i_0} c_id_j\psi^{i+j} && \text{by (3.10.9) } d)\\ &= \sum_{(i,j)\in P} a(i,j) && \text{by (3.10.9) } f) \text{ for the partition of } P \text{ by } j\end{aligned}$$

and, grouping instead by $N := i + j$, whose fibres are finite,

$$\begin{aligned}(fg)\circ\psi &= \sum_N\left(\sum_{i+j=N} c_id_j\right)\psi^N = \sum_N\sum_{i+j=N} c_id_j\psi^N && \text{by (3.3.17)}\\ &= \sum_{(i,j)\in P} a(i,j) && \text{by (3.10.9) } f) \text{ for the partition of } P \text{ by } i+j\end{aligned}$$

so $(f \circ \psi)(g \circ \psi) = (fg) \circ \psi$. ∎

© David Rufino CC BY-NC-ND 4.0

## 3.11 Polynomial Rings in Many Variables

**Definition 3.11.1 (Polynomial ring in many variables)**
*Let $A$ be a ring then the polynomial ring $A[X_1, ..., X_n]$ is the monoid ring $A[\mathbb{N}^n]$ (3.7.1) consisting of formal sums of monomials*

$$f(X_1, ..., X_n) = \sum_{v \in \mathbb{N}^n} f_v X_1^{v_1} \cdots X_n^{v_n}$$

*where $f_v \in A$ and only finitely many coefficients are non-zero. Addition is defined in the obvious way.*

*We may canonically regard $A$, $A[X_i]$ and $A[X_1, ..., X_i]$ as subrings in the obvious way.*

*Define $\deg(f, i)$ to be the maximal power of $X_i$ with a non-zero coefficient.*

**Remark 3.11.2**
*It may be useful for certain induction arguments to write*

$$A[X_1, ..., X_n] = A$$

*when $n = 0$.*

**Proposition 3.11.3 (Evaluation Homomorphism)**
*$A[X_1, ..., X_n]$ satisfies the following universal property. Given any $A$-algebra $B$ and points $(b_1, ..., b_n)$ there exists a ring homomorphism*

$$\phi_b : A[X_1, ..., X_n] \to B$$

*such that*

$$\phi_b(X_i) = \phi(b_i)$$

*given by*

$$\phi_b\left(\sum_v a_v X_1^{v_1} \cdots X_n^{v_n}\right) = \sum_v i_B(a_v)\phi(b_1)^{v_1} \cdots \phi(b_n)^{v_n}$$

*In other words it is an initial object in the category of $n$-pointed $A$-algebras. Furthermore*

$$\mathrm{Im}(\phi_b) = A[b_1, ..., b_n]$$

**Lemma 3.11.4 (Iterated polynomial ring)**
*Given $f \in A[X_1, ..., X_n]$ and let $N = \deg(f, n)$ then there exist unique polynomials $g_i \in A[X_1, ..., X_{n-1}]$ such that*

$$f = \sum_{i=0}^{N} g_i X_n^i$$

*in other words there is a canonical isomorphism*

$$\psi : A[X_1, ..., X_{n-1}][X_n] \to A[X_1, ..., X_n]$$

*under which $\deg(f) = \deg(\psi(f); n)$.*

**Proposition 3.11.5 (Homogenous grading)**
*Consider $R = k[X_1, ..., X_n]$ and $x \in k^n$. Then there is a direct sum of $k$-submodules*

$$R = \bigoplus_{n \geq 0} R^{n,x}$$

*where*

$$R^{n,x} = \left\{ \sum_{|\alpha|_1 = n} \lambda_\alpha (X_1 - x_1)^{\alpha_1} \cdots (X_n - x_n)^{\alpha_n} \;\middle|\; \lambda_\alpha \in k \quad \alpha \in \mathbb{N}^n \right\}$$

*and every $F \in R$ may be written uniquely as*

$$F(X) = F(x) + F^{(1,x)}(X) + ... + F^{(n,x)}(X) + ...$$

© David Rufino CC BY-NC-ND 4.0

*with* $F^{(n,x)} \in R^{n,x}$. *Note that*

$$\ker(\mathrm{ev}_x) =: M_x = \bigoplus_{n\geq 1} R^{n,x} = (X_1 - x_1, ..., X_n - x_n)$$

*and*

$$M_x^k = \bigoplus_{n\geq k} R^{n,x}$$

*Finally there is a canonical isomorphism*

$$k[X_1, ..., X_n]^{1,x} \cong M_x/M_x^2$$

*Proof.* By Proposition (...) there is a $k$-algebra homomorphism $\rho_x : R \to R$ given by $X_i \to X_i + x_i$. It is an isomorphism with two-sided inverse $\rho_{-x}$. Let $F \in R$ then

$$\rho_x(F) = \sum_{n=0}^{\infty}\left(\sum_{|\alpha|_1=n} \lambda_\alpha X_1^{\alpha_1}\cdots X_n^{\alpha_n}\right)$$

whence applying $\rho_{-x}$

$$\begin{aligned} F(X) &= \sum_{n=0}^{\infty} F^{(n)}(X) \\ F^{(n)}(X) &= \sum_{|\alpha|_1=n} \lambda_\alpha (X_1 - x_1)^{\alpha_1}\cdots(X_n - x_n)^{\alpha_n} \end{aligned}$$

as required. The coefficients $\lambda_\alpha$ are seen to be uniquely determined by applying $\rho_x$. Therefore the internal sum is direct. Finally evaluate at $x$ to find $F^{(0)} = F(x)$. The statement regarding $M_x$ is straightforward. And because $R^{n,x} \cdot R^{m,x} \subseteq R^{n+m,x}$ the statement regarding $M_x^k$ follows by induction. ∎

**Definition 3.11.6 (Projection to linear terms)**
*Given* $\mathfrak{a} \lhd k[X_1, ..., X_n]$ *and* $x \in k^n$ *define*

$$\mathfrak{a}^{(i,x)} = \left\{F^{(i,x)} \;\middle|\; F \in \mathfrak{a}\right\}$$

The following is useful

**Lemma 3.11.7**
*Let* $A$ *be a* $k$*-algebra and* $F \in k[X_1, ..., X_n]$ *and* $G_1, ..., G_n \in k[Y_1, ..., Y_m]$ *polynomials. For* $\lambda_1, ..., \lambda_m \in A$ *we have*

$$F(G_1, ..., G_n)(\lambda_1, ..., \lambda_m) = F\Big(G_1(\lambda_1, ..., \lambda_m), ..., G_{n(\lambda_1,...,\lambda_m)}\Big)$$

© David Rufino CC BY-NC-ND 4.0

## 3.12 Δ-Graded Rings

References:

- [Bou98a] (Chap. II §11)

Let $\Delta$ be a commutative monoid. By convention we write the monoid operation additively and 0 for the identity. Further we assume it is **cancellable**: $\lambda + \mu = \lambda + \mu' \implies \mu = \mu'$. Typically $\Delta = \mathbb{Z}$, but for example we may also consider $\mathbb{N}^k$ for some positive integer $k$.

**Definition 3.12.1 (Graded Abelian Groups)**
*Let $\Delta$ be a cancellable commutative monoid. An abelian group $G$ is $\Delta$-**graded** if it may be written as internal direct sum (3.3.93) of $\mathbb{Z}$-modules*

$$G = \bigoplus_{\lambda \in \Delta} G_\lambda$$

*for subgroups $G_\lambda \leq G$. In particular the subgroups $G_\lambda$ are disjoint. This means for every $g \in G$ there is a unique decomposition into a finite sum*

$$g = \sum_{\lambda \in \Delta} g_\lambda$$

*If $g \in G_\lambda$ then we say $g$ is **homogenous**. In this case we write $\delta(g) \coloneqq \lambda$ for the **degree** of $g$.*

*We may write $\pi_\lambda : G \to G_\lambda$ and $i_\lambda : G_\lambda \to G$ for projection and inclusion respectively. For $g \in G$ we may abbreviate $\pi_\lambda(g)$ by $g_\lambda$.*

**Definition 3.12.2 (Graded Ring)**
*Let $\Delta$ be a cancellable commutative monoid. A commutative ring $A$ is **graded** of type $\Delta$ if the additive group $(A, +)$ is $\Delta$-graded, and the multiplication is compatible in the sense that*

$$a \in A_\lambda, b \in A_\mu \implies a \cdot b \in A_{\lambda+\mu}$$

**Lemma 3.12.3**
*Let $A$ be a $\Delta$-ring. Then $A_0$ is a subring.*

*In the case $\Delta = \mathbb{N}$ the additive subgroups*

$$A_{\geq k} \coloneqq \sum_{l \geq k} A_l$$

*are homogenous ideals. We write $A_+ \coloneqq A_{\geq 1}$ which we call the **irrelevant ideal**.*

*Proof.* We need only show $1 \in A_0$. By hypothesis there is a unique decomposition

$$1 = \sum_{\lambda \in \Delta} e_\lambda \quad e_\lambda \in A_\lambda$$

For every $x \in A_\mu$ we have

$$x = x \cdot 1 = \sum_{\lambda \in \Delta} x \cdot e_\lambda$$

By the cancellable assumption we have $\lambda \neq 0 \implies \lambda + \mu \neq \mu$. Therefore $x \cdot e_0 = x$. By distributivity this then holds for all $x \in A$ and in particular $1 = 1 \cdot e_0 = e_0$. ∎

**Definition 3.12.4 (Graded Module)**
*Let $A$ be a $\Delta$-graded ring and $M$ a left (resp. right) $A$-module which is a $\Delta$-graded as an abelian group. Then $M$ is a **graded module** if the $A$-module structure is compatible with the grading, namely*

$$a \in A_\lambda, m \in M_\mu \implies a \cdot m \in M_{\lambda+\mu}$$

*Elements of $M_\lambda$ are known as homogenous.*

**Definition 3.12.5 (Graded Homomorphisms)**
*Let $A, B$ be $\Delta$-graded rings and $\phi : A \to B$ a ring homomorphism. It is a **graded ring homomorphism** if*

$$\phi(A_\lambda) \subseteq B_\lambda \quad \forall \lambda \in \Delta$$

© David Rufino CC BY-NC-ND 4.0

*Let $M, M'$ be graded $A$-modules and $\phi : M \to M'$ an $A$-module homomorphism. Then $\phi$ is a* ***graded module homomorphism*** *if*

$$\phi(M_\lambda) \subseteq M'_\lambda \quad \forall \lambda \in \Delta$$

**Definition 3.12.6 (Graded Submodules)**
*Let $M$ be a graded $A$-module, and $N \leq M$ an $A$-submodule. Then the following are equivalent*

*a)* $n \in N, \lambda \in \Delta \implies \pi_\lambda(n) \in N$
*b)* $N$ *is generated by homogenous elements*
*c)* $N$ *is equal to the internal direct sum*

$$\bigoplus_{\lambda \in \Delta} N \cap M_\lambda$$

*We say such a submodule is* ***graded*** *or* ***homogenous*** *and we write $N_\lambda \coloneqq N \cap M_\lambda$.*

**Proposition 3.12.7**
*Let $M$ be a graded $A$-module and $\{N_\alpha\}_{\alpha \in I}$ be a family of submodules. Then both*

$$\sum_{\alpha \in I} N_\alpha$$

*and*

$$\bigcap_{\alpha \in I} N_\alpha$$

*are graded submodules.*

**Proposition 3.12.8**
*Let $M$ be a finitely-generated graded $A$-module. Then $M$ has a finite generating set of homogenous elements.*

**Definition 3.12.9 (Homogenous Ideal)**
*Let $A$ be a commutative graded ring. An ideal $\mathfrak{a} \lhd A$ which is homogenous as an $A$-submodule of $A$ is a* ***homogenous ideal****. We write $\mathfrak{a}_\lambda \coloneqq \mathfrak{a} \cap A$, then by definition*

$$\mathfrak{a} = \bigoplus_{\lambda \in \Delta} \mathfrak{a}_\lambda$$

*Note if $A$ is Noetherian then every homogenous ideal has a finite generating set of homogenous elements.*

**Lemma 3.12.10**
*Let $\phi : A \to B$ be a graded ring and $\mathfrak{b} \lhd B$ a homogenous ideal. Then $\phi^{-1}(\mathfrak{b})$ is homogenous.*

**Example 3.12.11 (Polynomial Ring)**
*The polynomial ring $k[X_0, ..., X_n]$ is a graded ring over $\mathbb{N}$ with*

$$k[X_0, ..., X_n]_d = k\langle \{ X_1^{\alpha_1} \cdot ... \cdot X_n^{\alpha_n} \mid \alpha_1 + ... + \alpha_n = d \} \rangle$$

*Furthermore*

$$\bigoplus_{i \geq d} k[X_0, ..., X_n]_i = (X_0, ..., X_n)^d$$

**Proposition 3.12.12 (Quotient by a Homogenous Ideal)**
*Suppose $\Delta$ is a commutative cancellable monoid. Let $A$ be a $\Delta$-graded ring and $\mathfrak{a} \lhd A$ a homogenous ideal. Then $A/\mathfrak{a}$ is a $\Delta$-graded ring by considering the internal direct sum of abelian groups*

$$A/\mathfrak{a} = \bigoplus_{\lambda \in \Delta} (A/\mathfrak{a})_\lambda$$
$$(A/\mathfrak{a})_\lambda \coloneqq \{ \overline{a} \mid a \in A_\lambda \}$$

*Furthermore there is an isomorphism of $A$-modules*

$$(A/\mathfrak{a})_\lambda \xrightarrow{\sim} A_\lambda/(\mathfrak{a} \cap A_\lambda)$$

*For every homogenous ideal $\mathfrak{b} \supseteq \mathfrak{a}$ the quotient ideal $\mathfrak{b}/\mathfrak{a}$ is homogenous with*

© David Rufino CC BY-NC-ND 4.0

$$\mathfrak{b}/\mathfrak{a} = \bigoplus_{\lambda\in\Delta} (\mathfrak{b}/\mathfrak{a})_\lambda$$

*Proof.* Evidently the $(A/\mathfrak{a})_\lambda$ are abelian subgroups and span $A/\mathfrak{a}$. Observe that

$$0 = \sum_{\lambda\in\Delta} \overline{a_\lambda} \Longrightarrow \sum_{\lambda\in\Delta} a_\lambda \in \mathfrak{a}$$

Then as $\mathfrak{a}$ is homogenous we conclude $a_\lambda \in \mathfrak{a}$ and therefore $\overline{a_\lambda} = 0$. Therefore the family $(A/\mathfrak{a})_\lambda$ satisfy the criteria of (3.3.94) and the direct sum is well-defined. Evidently the ring multiplication on $A/\mathfrak{a}$ is compatible with the given $\Delta$-grading.

The isomorphism is induced by $\pi_\lambda$.

Suppose that $\overline{b} \in \mathfrak{b}/\mathfrak{a}$ then we require to prove that $\left(\overline{b}\right)_\lambda \in (\mathfrak{b}/\mathfrak{a})$. By assumption $b_\lambda \in \mathfrak{b}$ so $\overline{b_\lambda} \in \mathfrak{b}/\mathfrak{a}$. However by the first part $\overline{b_\lambda} = \left(\overline{b}\right)_\lambda$. ∎

**Proposition 3.12.13 (Localization at Homogenous Elements)**
*Let $\Delta$ be an abelian group, $A$ a $\Delta$-graded ring and $S \subset A$ a multiplicative set of* ***homogenous*** *elements. Then we may define a $\Delta$-gradation on $S^{-1}A$ by*

$$\left(S^{-1}A\right)_\lambda := \left\{ \frac{a}{s} \;\middle|\; \delta(a) - \delta(s) = \lambda \right\}$$

*For a homogenous ideal $\mathfrak{a} \lhd A$ then the ideal $S^{-1}\mathfrak{a}$ is homogenous.*

*Proof.* We first show that $\left(S^{-1}A\right)_\lambda$ is an additive subgroup, for given homogenous $a, a' \in A$ and $s, s' \in s$ by definition

$$\frac{a}{s} + \frac{a'}{s'} = \frac{as' + a's}{ss'}$$

Suppose $\delta(a) - \delta(s) = \lambda = \delta(a') - \delta(s')$ then $\delta(as') = \delta(a's)$. Therefore $as' + a's$ is homogenous and $\delta(as' + a's) = \delta(as') = \lambda + \delta(s) + \delta(s') = \lambda + \delta(ss')$ whence $\left(S^{-1}A\right)_\lambda$ is an additive subgroup. Similarly for the multiplicative structure suppose $\delta(a) = \delta(s) + \lambda$ and $\delta(a') = \delta(s') + \lambda'$ then

$$\delta(aa') = \delta(a) + \delta(a') = \delta(\lambda) + \delta(\lambda') + (\lambda + \lambda') = \delta(\lambda\lambda') + (\lambda + \lambda')$$

so that $\frac{a}{s} \cdot \frac{a'}{s'} \in \left(S^{-1}A\right)_{\lambda+\lambda'}$ as required. Evidently $S^{-1}A$ is the internal sum of $\left(S^{-1}A\right)_\lambda$. To show it is direct suppose that

$$0 = \sum_{i=1}^n \frac{a_i}{s_i} \quad \delta(a_i) = \delta(s_i) + \lambda_i$$

for distinct $\lambda_1, ..., \lambda_n$. Then

$$0 = \sum_{i=1}^n a_i t \prod_{j\neq i} s_j$$

for some $t \in S$. Furthermore

$$\delta\left(a_i t \prod_{j\neq i} s_j\right) = \delta(a_i) + \sum_{j\neq i} \delta\left(s_j\right) + \delta(t) = \lambda_i + \sum_{j=1}^n \delta\left(s_j\right) + \delta(t)$$

As $\Delta$ is cancellable the elements $\hat{\lambda}_i := \lambda_i + \sum_{j=1}^n \delta\left(s_j\right) + \delta(t)$ are distinct. By assumption we conclude that $a_i t \prod_{j\neq i} s_j = 0$ and so $\frac{a_i}{s_i} = 0$. Therefore $S^{-1}A$ is the direct sum of $\left(S^{-1}A\right)_\lambda$ by (3.3.94).

For $\frac{a}{s} \in S^{-1}\mathfrak{a}$ we wish to show that $\left(\frac{a}{s}\right)_\lambda \in S^{-1}\mathfrak{a}$ for all $\lambda \in \Delta$. Here $a \in \mathfrak{a}$ so by assumption $a_\lambda \in \mathfrak{a}$ for all $\lambda \in \Delta$ and $a = \sum_{\lambda\in\Delta} a_\lambda$ so

$$\frac{a}{s} = \sum_{\lambda\in\Delta} \frac{a_\lambda}{s} = \sum_{\lambda\in\Delta} \frac{a_{\lambda+\delta(s)}}{s}$$

and so $\left(\frac{a}{s}\right)_\lambda = \frac{a_{\lambda+\delta(s)}}{s} \in S^{-1}\mathfrak{a}$ as required. ∎

Recall for any $\Delta$-graded ring $A$ the additive subgroup $A_0$ is also a subring.

© David Rufino CC BY-NC-ND 4.0

**Definition 3.12.14**
*Let $\Delta$ be a commutative abelian group, $A$ a $\Delta$-graded ring and $S$ be a multiplicative subset of homogenous elements. Define the subring of $S^{-1}A$*

$$A_{(S)} \coloneqq \left(S^{-1}A\right)_0 = \left\{\frac{a}{s} \;\middle|\; \delta(s) = \delta(a)\right\}$$

*Similarly for $\mathfrak{a}$ a homogenous ideal define the homogenous ideal of $S^{-1}A$*

$$\mathfrak{a}_{(S)} \coloneqq \left\{\frac{a}{s} \;\middle|\; a \in \mathfrak{a}, s \in S, \delta(s) = \delta(a)\right\}$$

*In the case $\mathfrak{p}$ is a homogenous prime ideal define the local ring*

$$A_{(\mathfrak{p})} \coloneqq A_{(\{x \notin \mathfrak{p}\})}$$

*and for $f \in A$ homogenous define*

$$A_{(f)} \coloneqq A_{(\{1, f, f^2, \ldots\})}$$

**Lemma 3.12.15**
*Let $\Delta$ be an abelian group, $A$ a $\Delta$-graded ring and $\mathfrak{a} \lhd A$ a homogenous ideal. Then*

$$\mathfrak{a}_{(S)} = A_{(S)} \cap S^{-1}\mathfrak{a}$$

*as subsets of $S^{-1}A$.*

**Lemma 3.12.16**
*Let $\Delta$ be an abelian group, $A$ a $\Delta$-graded ring, $a \in A_\mu$ homogenous and $x \in A$. Then*

$$(ax)_\lambda = a \cdot x_{\lambda - \mu}$$

*In particular $ax$ is homogenous iff $x$ is homogenous.*

**Proposition 3.12.17 (Homogenous localisation comutes with quotients)**
*Let $\Delta$ be an abelian group, $A$ a $\Delta$-graded ring, $\mathfrak{a}$ a homogenous ideal and $S \subset A$ a multiplicatively closed set of homogenous elements such that $S \cap \mathfrak{a} = \varnothing$. Then there is a canonical isomorphism of $\Delta$-graded rings*

$$\begin{aligned}(S^{-1}A)/(S^{-1}\mathfrak{a}) &\xrightarrow{\sim} \pi(S)^{-1}(A/\mathfrak{a}) \\ \frac{a}{s} + S^{-1}A &\to \frac{a + \mathfrak{a}}{s + \mathfrak{a}}\end{aligned}$$

*where $\pi : A \to A/\mathfrak{a}$ is the canonical surjection. In particular there is an isomorphism of rings*

$$A_{(S)}/\mathfrak{a}_{(S)} \xrightarrow{\sim} \left(S^{-1}A/S^{-1}\mathfrak{a}\right)_0 \xrightarrow{\sim} (A/\mathfrak{a})_{(\pi(S))}$$

*Proof.* The first isomorphism of rings is from (3.6.25), and it is obviously compatible with the $\Delta$-grading. For suppose $x \in (S^{-1}A)/(S^{-1}\mathfrak{a})$ with $\delta(x) = \lambda$. Then $x = \frac{a}{s}$ with $\delta(a) - \delta(s) = \lambda$. By definition $\delta(a + \mathfrak{a}) - \delta(s + \mathfrak{a}) = \lambda$ as required.

The third isomorphism follows immediately. For the second consider the ring homomorphism

$$\phi : A_{(S)} \hookrightarrow S^{-1}A \to S^{-1}A/S^{-1}\mathfrak{a}$$

Then evidently the image is $\left(S^{-1}A/S^{-1}\mathfrak{a}\right)_0$ as the quotient map is graded. The kernel is $A_{(S)} \cap S^{-1}\mathfrak{a}$ which equals $\mathfrak{a}_{(S)}$ by (3.12.15). ∎

**Proposition 3.12.18**
*Let $\Delta$ be an abelian group, $A$ a $\Delta$-graded ring, $\mathfrak{a}$ a homogenous ideal and $\mathfrak{p} \supseteq \mathfrak{a}$ a homogenous prime ideal. Then there is a canonical isomorphism of $\Delta$-graded rings*

$$\begin{aligned}A_{\mathfrak{p}}/\mathfrak{a}_{\mathfrak{p}} &\xrightarrow{\sim} (A/\mathfrak{a})_{\mathfrak{p}/\mathfrak{a}} \\ \frac{a}{s} + \mathfrak{a}_{\mathfrak{p}} &\to \frac{a + \mathfrak{a}}{s + \mathfrak{a}}\end{aligned}$$

*In particular there is an isomorphism of rings*

© David Rufino CC BY-NC-ND 4.0

$$A_{(\mathfrak{p})}/\mathfrak{a}_{(\mathfrak{p})} \xrightarrow{\sim} \left(A_{\mathfrak{p}}/\mathfrak{a}_{\mathfrak{p}}\right)_0 \xrightarrow{\sim} (A/\mathfrak{a})_{(\mathfrak{p}/\mathfrak{a})}$$

*given by*

$$\frac{a}{s} + \mathfrak{a}_{(\mathfrak{p})} \to \frac{a+\mathfrak{a}}{s+\mathfrak{a}}$$

*Proof.* Follows from (3.12.17) and the observation that $A/\mathfrak{a} \setminus \mathfrak{p}/\mathfrak{a} = \pi(A \setminus \mathfrak{p})$. ■

© David Rufino CC BY-NC-ND 4.0

# 3.13 Graded Rings

**Definition 3.13.1 (Graded Ring over $\mathbb{Z}$)**
*A ring $A$ (resp. $A$-module $M$) is said to be $\mathbb{Z}$-**graded** (or simply **graded**) if it is graded in the sense of (3.12.2) with gradation group $\Delta = \mathbb{Z}$.*

*If $i < 0 \implies A_i = 0$ (resp. $M_i = 0$) then we say $A$ (resp. $M$) is **positively graded**.*

*Note in this case that we have the multiplication formula*

$$(a \cdot b)_i = \sum_{j+k=i} a_j \cdot b_k$$

**Definition 3.13.2**
*For an element $a \in A$ we define the **degree** to be*

$$\deg(a) := \max\{i \mid a_i \neq 0\}$$

**Definition 3.13.3 (Essential and Irrelevant Ideals)**
*Let $A$ be a positively graded ring. Then we say a homogenous ideal $\mathfrak{a}$ is **irrelevant** if the following condition holds*

$$\exists n_0 \text{ s.t. } n \geq n_0 \implies \mathfrak{a}_n = A_n$$

*In particular $A_+$ is irrelevant. Otherwise we say that it is **essential**.*

**Proposition 3.13.4 (Radical of Homogenous Ideal is Homogenous)**
*Let $A$ be a graded ring and $\mathfrak{a}$ a homogenous ideal. Then $\sqrt{\mathfrak{a}}$ is also a homogenous ideal.*

*Proof.* Define $\mathfrak{r} := \sqrt{\mathfrak{a}}$ and fix $x \in \mathfrak{r}$. Suppose

$$x = \sum_{i=-\infty}^{\infty} x_i$$

where $\delta(x_i) = i$ and $x^k \in \mathfrak{r}$. Then

$$x^k = \sum_{i=-\infty}^{\infty} x_i^k$$

where $\delta(x_i^k) = ki$. Without loss of generality we may assume $k > 0$ so by assumption $x_i^k \in \mathfrak{r}$ for all $i$ and therefore $x_i \in \sqrt{\mathfrak{r}}$. ∎

**Proposition 3.13.5**
*Let $A$ be a graded ring and $\mathfrak{p}$ a homogenous ideal satisfying the following property*

$$a \in A_n, b \in A_m, ab \in \mathfrak{p} \implies a \in \mathfrak{p} \text{ or } b \in \mathfrak{p}$$

*Then $\mathfrak{p}$ is a prime ideal.*

*Proof.* Suppose $\mathfrak{p}$ is not prime. Then there exists $a, b \in A$ such that $ab \in \mathfrak{p}$ and $a, b \notin \mathfrak{p}$. Write $a = \sum_j a_j$ and $b = \sum_k b_k$. Then

$$ab = \sum_{i=-\infty}^{\infty} \left( \sum_{(j,k)|j+k=i} a_j b_k \right)$$

As $\mathfrak{p}$ is assumed homogenous then for each $i$ we have

$$(ab)_i := \sum_{(j,k)|j+k=i} a_j b_k \in \mathfrak{p}_i$$

Let $j'$ (resp. $k'$) be the largest integer such that $a_{j'} \notin \mathfrak{p}$ (resp. $b_{k'} \notin \mathfrak{p}$). For all other pairs $j''$, $k''$ such that $j' + k' = j'' + k''$ we have by maximality $a_{j''} b_{k''} \in \mathfrak{p}_i$. Therefore we conclude $a_{j'} b_{k'} \in \mathfrak{p}$, which contradicts the original assumption. ∎

**Proposition 3.13.6**
*Let $A$ be a positively graded ring. Then the following are equivalent*

*a) $A_+$ is a finitely-generated $A$-module*
*b) $A$ is a finitely-generated $A_0$-algebra*

© David Rufino CC BY-NC-ND 4.0

*More precisely for homogenous elements $x_0, ..., x_n \in A_+$ we have*

$$A_+ = \sum_{i=0}^{n} Ax_i \iff A = A_0[x_0, ..., x_n]$$

*In this case we say that $A$ is **finitely generated**. Further*

$$A_+^m = (x_0, ..., x_n)^m$$

*for all integers $m > 0$.*

**Proposition 3.13.7 (Criteria for Irrelevant ideals)**
*Let $A$ be a positively graded ring which is finitely generated and $\mathfrak{a}$ a homogenous ideal. Then the following are equivalent*

*a) $\mathfrak{a}$ is irrelevant (i.e. $n \geq n_0 \implies \mathfrak{a}_n = A_n$)*
*b) $A_+ \subset \sqrt{\mathfrak{a}}$*

*Proof.* a) $\implies$ b) It is sufficient to show that $i > 0 \implies A_i \subset \sqrt{\mathfrak{a}}$. Given $x \in A_i$ then by the Archimedean property there exists $k > 0$ such that $ik > n_0$. Therefore by assumption $x^k \in \mathfrak{a}$ and $x \in \sqrt{\mathfrak{a}}$.

b) $\implies$ a) TODO. ∎

**Proposition 3.13.8**
*Let $A$ be a graded ring and $f \in A_1$. Then there is an isomorphism*

$$\begin{aligned} A_{(f)}[X, X^{-1}] &\xrightarrow{\sim} A_f \\ \sum_{i\in\mathbb{Z}} \frac{a_{n_i}}{f^{n_i}} X^i &\longrightarrow \sum_{i\in\mathbb{Z}} \frac{a_{n_i}}{f^{n_i - i}} \quad a_{n_i} \in A \text{ homogenous of degree } n_i \\ \sum_{i\in\mathbb{Z}} \frac{a_i}{f^i} X^{i-r} &\longleftarrow \frac{\sum_{i\in\mathbb{Z}} a_i}{f^r} \end{aligned}$$

*Proof.* Denote the maps by $\psi, \phi$, then $\psi$ exists and is a ring homomorphism by (3.9.2) satisfying $\psi(X) = f$ and whence $\psi(X^i) = f^i$. To show $\psi$ is well-defined first consider the map

$$\begin{aligned} \psi' : A &\to A_{(f)}[X, X^{-1}] \\ \sum_{i\in\mathbb{Z}} a_i &\to \sum_{i\in\mathbb{Z}} \frac{a_i}{f^i} X^i \end{aligned}$$

which is evidently a well-defined homomorphism of abelian groups. Furthermore

$$\left(\sum_{i\in\mathbb{Z}} \frac{a_i}{f^i} X^i\right)\left(\sum_{i\in\mathbb{Z}} \frac{b_i}{f^i} X^i\right) = \sum_{i\in\mathbb{Z}} \left(\sum_{j+k=i} \frac{a_j b_k}{f^{j+k}}\right) X^i = \sum_{i\in\mathbb{Z}} \left(\frac{1}{f^i} \sum_{j+k=i} a_j b_k\right) X^i$$

and so the map is a ring-homomorphism by the multiplication formula in $A$. By the universal property of localisation the ring homomorphism $\phi$ exists with $\phi(f) = X$ and hence $\phi(f^i) = X^i$. We have

$$\phi(\psi(X)) = \phi(f) = X$$

$$\phi\left(\psi\left(\frac{a_j}{f^j}\right)\right) = \phi\left(\frac{a_j}{f^j}\right) = \frac{a_j}{f^j}$$

so by linearity $\phi \circ \psi = \mathbf{1}$. Similarly

$$\psi\left(\phi\left(\frac{a_i}{f^r}\right)\right) = \psi\left(\frac{a_i}{f^i} X^{i-r}\right) = \frac{a_i}{f^i} f^{i-r} = \frac{a_i}{f^r}$$

so by linearity $\psi \circ \phi = \mathbf{1}$. ∎

© David Rufino CC BY-NC-ND 4.0

## 3.14 Filtered Rings and Completions

Both DVRs and complete DVRs sit naturally in the category of filtered rings developed in this subsection: a DVR such as $A_{(p)}$ is filtered by the powers of its maximal ideal, and a complete DVR such as $\mathbb{Z}_p$ or $k[[X]]$ is exactly such a filtered ring whose canonical map to its own completion happens to be an isomorphism (3.14.20). Working at this level of generality – filtered rings and continuous homomorphisms between them, rather than DVRs and local homomorphisms specifically – gives a category closed under more operations than DVRs themselves support: for instance, an arbitrary product of filtered rings is again filtered, with completion computed factorwise (3.14.22), even though a product of DVRs is essentially never itself a DVR. This is the kind of fact the filtered-ring framework makes easy to state: the completion of a semi-local Dedekind domain (such as the integral closure $\mathcal{O}_L$ of a DVR in a finite extension) agrees with the product of the completions at its finitely many primes, by the Chinese Remainder Theorem factorwise – though the trace/residue theory developed later avoids needing this, working directly with Laurent series instead (see the Extensions of Power Series Rings section).

Filtered rings, at this level of generality, are the algebraic core of the **admissible rings** of [Gro60] ($0_{\mathrm{I}}$, §7) (linearly topologized rings with a fundamental system of open ideals, not necessarily powers of a single ideal).

**Definition 3.14.1 (Filtered Ring)**
*A **filtered ring** $(A, \mathfrak{a}_\bullet)$ is a ring $A$ together with a decreasing chain of ideals $\mathfrak{a}_\bullet = (\mathfrak{a}_n)_{n\in\mathbb{Z}}$ of $A$, that is $\mathfrak{a}_{n+1} \subseteq \mathfrak{a}_n$ for every $n \in \mathbb{Z}$, with $\mathfrak{a}_n = A$ for every $n < 0$. We say $(A, \mathfrak{a}_\bullet)$ is **separated** if $\bigcap_n \mathfrak{a}_n = 0$.*

**Definition 3.14.2 (Filtered Module)**
*Let $A$ be a ring and $M$ an $A$-module. A **filtration** of $M$ (as an $A$-module) is a decreasing chain $\mathcal{M}_\bullet = (\mathcal{M}_n)_{n\in\mathbb{Z}}$ of $A$-submodules of $M$, that is $\mathcal{M}_{n+1} \subseteq \mathcal{M}_n$ for every $n \in \mathbb{Z}$. We call $(M, \mathcal{M}_\bullet)$ a **filtered $A$-module**, and say it is **separated** if $\bigcap_n \mathcal{M}_n = 0$, and **exhaustive** if $\bigcup_n \mathcal{M}_n = M$.*

*Every positively filtered $A$-module (3.14.3) is exhaustive, trivially ($\mathcal{M}_n = M$ already for $n < 0$); the fractional-ideal filtration of $K = \mathrm{Frac}(A)$, $A$ a DVR, is a genuinely exhaustive example that is not positively filtered.*

*When $(A, \mathfrak{a}_\bullet)$ is itself a filtered ring, every $A$-module $M$ carries a canonical filtration $\mathfrak{a}_\bullet M := (\mathfrak{a}_n M)_{n\in\mathbb{Z}}$, the **induced filtration** (with $\mathfrak{a}_n M = M$ for $n < 0$, as $\mathfrak{a}_n = A$ there; so $(M, \mathfrak{a}_\bullet M)$ is positively filtered, hence exhaustive).*

**Definition 3.14.3 (Positively Filtered)**
*A filtered ring (3.14.1), or more generally a filtered $A$-module $(M, \mathcal{M}_\bullet)$ (3.14.2), is **positively filtered** if $\mathcal{M}_n$ equals the whole module for every $n < 0$. Taking $M := A$, $\mathcal{M}_n := \mathfrak{a}_n$ exhibits every filtered ring as a positively filtered $A$-module over itself.*

**Definition 3.14.4 ($\mathfrak{a}$-adic Filtration)**
*Let $A$ be a ring and $\mathfrak{a} \triangleleft A$ an ideal. The **$\mathfrak{a}$-adic filtration** is $\mathfrak{a}_n := \mathfrak{a}^{n+1}$ for $n \geq 0$ and $\mathfrak{a}_n := A$ for $n < 0$, a filtration of $A$ (3.14.1).*

**Definition 3.14.5 (Continuous Homomorphism)**
*Let $(A, \mathfrak{a}_\bullet)$, $(B, \mathfrak{b}_\bullet)$ be filtered rings. A ring homomorphism $\phi : A \to B$ is **continuous** if, for every $m \geq 0$, there is $n \geq 0$ with $\phi(\mathfrak{a}_n) \subseteq \mathfrak{b}_m$.*

**Remark 3.14.6**
*Continuity is exactly continuity of $\phi$ at $0$ in the usual metric-space sense, for the metrics $d_A(x,y) := 2^{-\sup\{n : x-y\in\mathfrak{a}_n\}}$ on $A$ and $d_B(x,y) := 2^{-\sup\{n : x-y\in\mathfrak{b}_n\}}$ on $B$ (with $\sup\varnothing := -1$, so $d_A, d_B$ vanish exactly on equal pairs): a ball of radius $2^{-m}$ about $0 \in B$ is exactly $\mathfrak{b}_{m+1}$, so "$\phi(\mathfrak{a}_{n+1}) \subseteq \mathfrak{b}_{m+1}$ for some $n$, for every $m$" is exactly continuity of $\phi$ at $0$. As $d_A, d_B$ are translation-invariant ($d(x,y) = d(x-y, 0)$, immediate as $\mathfrak{a}_n, \mathfrak{b}_n$ are subgroups), this already gives continuity of $\phi$ at every point of $A$: given $x_0 \in A$ and $\varepsilon > 0$, pick $m$ with $2^{-m} \leq \varepsilon$ and $n$ with $\phi(\mathfrak{a}_{n+1}) \subseteq \mathfrak{b}_{m+1}$; then $d_A(x, x_0) < 2^{-n}$ gives $x - x_0 \in \mathfrak{a}_{n+1}$, so $\phi(x) - \phi(x_0) = \phi(x - x_0) \in \mathfrak{b}_{m+1}$, that is $d_B(\phi(x), \phi(x_0)) \leq 2^{-m} \leq \varepsilon$. (This is the same translation-invariance trick used for the identity map in (3.14.13), which compares two filtrations of a single ring this way.)*

For the remainder of this subsection, fix filtered rings $(A, \mathfrak{a}_\bullet)$, $(A', \mathfrak{a}'_\bullet)$, a ring homomorphism $\rho : A \to A'$, and filtered $A$-, $A'$-modules $(M, \mathcal{M}_\bullet)$, $(M', \mathcal{M}'_\bullet)$ (3.14.2): this notation is reused without restatement in every definition and proposition below phrased in terms of a map $\psi : M \to M'$.

**Definition 3.14.7 (Continuous Homomorphism, Module Case)**
*An additive map $\psi : M \to M'$ is a **$\rho$-linear map** (or **$\rho$-homomorphism**) if $\psi(a \cdot x) = \rho(a) \cdot \psi(x)$ for every $a \in A$, $x \in M$, and is **continuous** if, for every $m \in \mathbb{Z}$, there is $n \in \mathbb{Z}$ with $\psi(\mathcal{M}_n) \subseteq \mathcal{M}'_m$.*

© David Rufino CC BY-NC-ND 4.0

*When $A = A'$ and $\rho = \mathrm{id}_A$, a continuous $\rho$-homomorphism is exactly an ordinary continuous $A$-module homomorphism.*

**Definition 3.14.8 (Dense Homomorphism)**
*Let $(A, \mathfrak{a}_\bullet)$, $(B, \mathfrak{b}_\bullet)$ be filtered rings. A ring homomorphism $\phi : A \to B$ is **dense** if $B = \phi(A) + \mathfrak{b}_n$ for every $n \geq 0$.*

**Definition 3.14.9 (Dense Homomorphism, Module Case)**
*A $\rho$-homomorphism $\psi : M \to M'$ (3.14.7) is **dense** if $M' = \psi(M) + \mathcal{M}'_n$ for every $n \geq 0$. (Taking $M := A$, $M' := A'$ as filtered $A$-, $A'$-modules over themselves and $\psi := \rho$ recovers (3.14.8).)*

**Definition 3.14.10 (Taut Homomorphism)**
*Let $(A, \mathfrak{a}_\bullet)$, $(B, \mathfrak{b}_\bullet)$ be filtered rings and $\phi : A \to B$ a ring homomorphism with $\phi(\mathfrak{a}_n) \subseteq \mathfrak{b}_n$ for every $n \geq 0$. Then $\phi$ is **taut** if there is a strictly increasing sequence $n_0 < n_1 < n_2 < \cdots$ in $\mathbb{N}$ such that*

$$B = \phi(A) + \mathfrak{b}_{n_0} \qquad \text{and} \qquad \mathfrak{b}_{n_N} = \phi\big(\mathfrak{a}_{n_N}\big) + \mathfrak{b}_{n_{N+1}} \text{ for every } N \geq 0.$$

**Definition 3.14.11 (Taut Homomorphism, Module Case)**
*A $\rho$-homomorphism $\psi : M \to M'$ with $\psi(\mathcal{M}_n) \subseteq \mathcal{M}'_n$ for every $n \geq 0$ is **taut** if there is a strictly increasing sequence $n_0 < n_1 < n_2 < \cdots$ in $\mathbb{N}$ such that*

$$M' = \psi(M) + \mathcal{M}'_{n_0} \qquad \text{and} \qquad \mathcal{M}'_{n_N} = \psi\big(\mathcal{M}_{n_N}\big) + \mathcal{M}'_{n_{N+1}} \text{ for every } N \geq 0.$$

*(Taking $M := A$, $M' := A'$ as filtered $A$-, $A'$-modules over themselves and $\psi := \rho$ recovers (3.14.10).)*

**Remark 3.14.12**
*The name is borrowed, by analogy, from the **taut** ring maps of linearly topologized rings in [Stacks] (Tag 0AMX) – itself flagged there as nonstandard terminology – where an analogous (though not identical) condition, combined with density and completeness, is used to conclude surjectivity via an open mapping theorem ([Stacks] Tag 0APT) rather than the direct successive-approximation argument used here.*

**Proposition 3.14.13 (Refinement Criterion)**
*Let $\mathfrak{a}_\bullet$, $\mathfrak{a}'_\bullet$ be filtrations of a ring $A$ (3.14.1), with metrics $d_1(x,y) := 2^{-\sup\{n : x-y \in \mathfrak{a}_n\}}$ and $d_2(x,y) := 2^{-\sup\{n : x-y \in \mathfrak{a}'_n\}}$ on $A$ ($\sup \varnothing := -1$). The following are equivalent:*

*a) $1_A : (A, \mathfrak{a}_\bullet) \to (A, \mathfrak{a}'_\bullet)$ is continuous (3.14.5)*
*b) $\mathfrak{a}_\bullet$ is **cofinal** in $\mathfrak{a}'_\bullet$: for every $m \geq 0$ there is $n \geq 0$ with $\mathfrak{a}_n \subseteq \mathfrak{a}'_m$;*
*c) $1_A : (A, d_1) \to (A, d_2)$ is continuous at $0$ in the metric-space sense;*
*d) the topology on $A$ induced by $d_2$ is weaker than the one induced by $d_1$: every $d_2$-open subset of $A$ is $d_1$-open.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

a) $\iff$ b) is (3.14.5) applied to $\phi := 1_A$.

b) $\implies$ c): given $\varepsilon > 0$, pick $m \geq 0$ with $2^{-m} \leq \varepsilon$; by b) at $m+1$, pick $n \geq 0$ with $\mathfrak{a}_n \subseteq \mathfrak{a}'_{m+1}$, and set $\delta := 2^{-(n-1)}$ (any $\delta > 0$ if $n = 0$). If $d_1(x,0) < \delta$ then $x \in \mathfrak{a}_n \subseteq \mathfrak{a}'_{m+1}$, so $d_2(x,0) \leq 2^{-m} \leq \varepsilon$.

c) $\implies$ b): given $m \geq 0$, apply c) with $\varepsilon := 2^{-m}$ to get $\delta > 0$ with $d_1(x,0) < \delta \implies d_2(x,0) \leq 2^{-m}$; picking $n \geq 0$ with $2^{-(n-1)} < \delta$ (any $n \geq 1$ if $\delta$ exceeds every such bound) gives $\mathfrak{a}_n \subseteq \{x : d_1(x,0) < \delta\} \subseteq \{x : d_2(x,0) \leq 2^{-m}\} = \mathfrak{a}'_m$.

c) $\implies$ d): both $d_1, d_2$ are translation-invariant ($d_i(x+z, y+z) = d_i(x,y)$, immediate as $(x+z)-(y+z) = x-y$), so continuity of $1_A : (A, d_1) \to (A, d_2)$ at $0$ gives continuity at every point $x_0 \in A$: given $\varepsilon > 0$, take $\delta > 0$ as in c); if $d_1(x, x_0) < \delta$ then $d_1(x - x_0, 0) < \delta$, so $d_2(x - x_0, 0) < \varepsilon$, that is $d_2(x, x_0) < \varepsilon$. Continuity of $1_A$ at every point is exactly d) (the preimage, under the identity map, of a $d_2$-open set is itself, and it is $d_1$-open).

d) $\implies$ c): a $d_2$-ball around $0$ is $d_2$-open, hence $d_1$-open by d), hence contains a $d_1$-ball around $0$ (as $0$ lies in it); this is exactly continuity of $1_A$ at $0$. ∎

**Proposition 3.14.14 (Equivalence Criterion)**
*Let $\mathfrak{a}_\bullet$, $\mathfrak{a}'_\bullet$ be filtrations of a ring $A$ (3.14.1), with metrics $d_1$, $d_2$ as in (3.14.13). The following are equivalent:*

*a) $1_A : (A, \mathfrak{a}_\bullet) \to (A, \mathfrak{a}'_\bullet)$ is **bicontinuous**, that is $1_A$ is continuous both as a map $(A, \mathfrak{a}_\bullet) \to (A, \mathfrak{a}'_\bullet)$ and as a map $(A, \mathfrak{a}'_\bullet) \to (A, \mathfrak{a}_\bullet)$ (3.14.5)*
*b) $\mathfrak{a}_\bullet$ and $\mathfrak{a}'_\bullet$ are cofinal in each other (3.14.13)*
*c) $d_1$ and $d_2$ are **equivalent** metrics on $A$: $1_A$ is bicontinuous at $0$ as a map $(A, d_1) \to (A, d_2)$;*
*d) $d_1$ and $d_2$ induce the* same *topology on $A$.*

© David Rufino CC BY-NC-ND 4.0

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Applying (3.14.13) to the pair $(\mathfrak{a}_\bullet, \mathfrak{a}'_\bullet)$ gives the equivalence of "$\mathfrak{a}_\bullet$ cofinal in $\mathfrak{a}'_\bullet$", "$1_A : (A, \mathfrak{a}_\bullet) \to (A, \mathfrak{a}'_\bullet)$ continuous", "$1_A : (A, d_1) \to (A, d_2)$ continuous at 0", and "every $d_2$-open set is $d_1$-open"; applying it to the reversed pair $(\mathfrak{a}'_\bullet, \mathfrak{a}_\bullet)$ gives the equivalence of the four corresponding statements with the roles of $1, 2$ exchanged. Conjoining each pair of equivalences (an "if and only if" together with its reverse) gives exactly a) $\iff$ b) $\iff$ c) $\iff$ d), the last as "every $d_2$-open set is $d_1$-open, and every $d_1$-open set is $d_2$-open" is exactly the two topologies having the same open sets. ∎

**Lemma 3.14.15 (Continuity Criterion for Ideal-Power Filtrations)**
*Let $A$, $B$ be rings with the $\mathfrak{a}$-adic and $\mathfrak{b}$-adic filtrations (3.14.4), for an ideal $\mathfrak{a} \lhd A$ and a* radical *ideal $\mathfrak{b} \lhd B$ (3.3.36). Then a ring homomorphism $\phi : A \to B$ is continuous (3.14.5) if and only if $\phi(\mathfrak{a}) \subseteq \mathfrak{b}$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

If $\phi(\mathfrak{a}) \subseteq \mathfrak{b}$: for every $k \geq 1$, $\phi\big(\mathfrak{a}^k\big)$ is generated (as an ideal of $B$) by products of $k$ elements of $\phi(\mathfrak{a}) \subseteq \mathfrak{b}$, hence $\phi\big(\mathfrak{a}^k\big) \subseteq \mathfrak{b}^k$; so, given $m$, taking $n := m$ gives $\phi(\mathfrak{a}_n) = \phi\big(\mathfrak{a}^{n+1}\big) \subseteq \mathfrak{b}^{m+1} = \mathfrak{b}_m$: $\phi$ is continuous.

Conversely, suppose $\phi$ is continuous; taking $m := 0$ gives some $n$ with $\phi\big(\mathfrak{a}^{n+1}\big) \subseteq \mathfrak{b}$. If $\phi(\mathfrak{a})\neg \subseteq \mathfrak{b}$, pick $a \in \mathfrak{a}$ with $\phi(a) \notin \mathfrak{b}$; as $\mathfrak{b}$ is radical, its defining property $x^{n+1} \in \mathfrak{b} \implies x \in \mathfrak{b}$ contraposes to $\phi(a) \notin \mathfrak{b} \implies \phi(a)^{n+1} \notin \mathfrak{b}$. But $\phi(a)^{n+1} = \phi(a^{n+1}) \in \phi\big(\mathfrak{a}^{n+1}\big) \subseteq \mathfrak{b}$, as $a^{n+1} \in \mathfrak{a}^{n+1}$: a contradiction. So $\phi(\mathfrak{a}) \subseteq \mathfrak{b}$. ∎

**Definition 3.14.16 (Completion of a Filtered Ring)**
*Let $(A, \mathfrak{a}_\bullet)$ be a filtered ring. For $n \geq m \geq 0$, as $\mathfrak{a}_n \subseteq \mathfrak{a}_m$, (3.3.56) gives a unique ring homomorphism $\rho_{n,m} : A/\mathfrak{a}_n \to A/\mathfrak{a}_m$ making*

$$\begin{array}{ccc} A & \twoheadrightarrow & A/\mathfrak{a}_n \\ & \searrow & \downarrow{\scriptstyle \rho_{n,m}} \\ & & A/\mathfrak{a}_m \end{array}$$

*commute; in particular it is surjective. These satisfy $\rho_{n,n} = 1_{A/\mathfrak{a}_n}$ and, for $n \geq k \geq m \geq 0$,*

$$\rho_{k,m} \circ \rho_{n,k} = \rho_{n,m}$$

*(both sides make the outer triangle $A \twoheadrightarrow A/\mathfrak{a}_n \to A/\mathfrak{a}_m$ commute, so agree by the uniqueness clause of (3.3.56)). Specialising to $(n, m) := (n+1, n)$ gives reduction maps $\rho_{n+1,n} : A/\mathfrak{a}_{n+1} \twoheadrightarrow A/\mathfrak{a}_n$, which form a tower (3.3.154) the* ***completion*** *of $(A, \mathfrak{a}_\bullet)$ is*

$$\hat{A} := \varprojlim_n A/\mathfrak{a}_n,$$

*with projections $\lambda_n : \hat{A} \to A/\mathfrak{a}_n$ (3.3.154) and the canonical ring homomorphism*

$$\iota : A \to \hat{A}, \quad a \mapsto (a \bmod \mathfrak{a}_n)_{n \geq 0}.$$

*Consequently, every $x = (a_n)_{n \geq 0} \in \hat{A}$ satisfies the* ***repeated compatibility*** *$\rho_{n,m}(a_n) = a_m$ for all $n \geq m \geq 0$, not just consecutive $n = m + 1$: by induction on $n - m$, composing the defining consecutive relations $\rho_{k,k-1}(a_k) = a_{k-1}$ for $m < k \leq n$.*

*Moreover $\ker \lambda_\bullet := (\ker \lambda_n)_{n \geq 0}$ is itself a decreasing chain of ideals of $\hat{A}$ – for $x = (\overline{a}_n)_n$, $\lambda_n(x) = 0$ forces $\lambda_{n-1}(x) = 0$ too (the further reduction of $\overline{a}_n = 0$ is 0) – so $\big(\hat{A}, \ker \lambda_\bullet\big)$ is again a filtered ring (3.14.1).*

**Definition 3.14.17 (Completion of a Filtered Module)**
*Let $A$ be a ring and $(M, \mathcal{M}_\bullet)$ a filtered $A$-module (3.14.2). For $n \geq m \geq 0$, as $\mathcal{M}_n \subseteq \mathcal{M}_m$, (3.3.101) gives a unique $A$-module homomorphism making*

$$\begin{array}{ccc} M & \twoheadrightarrow & M/\mathcal{M}_n \\ & \searrow & \downarrow \\ & & M/\mathcal{M}_m \end{array}$$

*commute; in particular it is surjective. Specialising to $(n, m) := (n+1, n)$ gives reduction maps $M/\mathcal{M}_{n+1} \twoheadrightarrow M/\mathcal{M}_n$, which form a tower of $A$-modules (3.3.154) the* ***completion*** *of $(M, \mathcal{M}_\bullet)$ is the $A$-module*

$$\widehat{M} := \varprojlim_n M/\mathcal{M}_n,$$

© David Rufino CC BY-NC-ND 4.0

*with projections* $\lambda_n : \widehat{M} \to M/\mathcal{M}_n$ *(3.3.154) and the canonical* $A$*-module homomorphism*

$$\iota : M \to \widehat{M}, \quad v \mapsto (v \bmod \mathcal{M}_n)_{n\geq 0}.$$

*Moreover* $\ker \lambda_\bullet := (\ker \lambda_n)_{n\geq 0}$ *is itself a decreasing chain of* $A$*-submodules of* $\widehat{M}$ *– for* $x = (\overline{v}_n)_n$*,* $\lambda_n(x) = 0$ *forces* $\lambda_{n-1}(x) = 0$ *too (the further reduction of* $\overline{v}_n = 0$ *is* $0$*).*

*Taking* $M := A$ *as a filtered* $A$*-algebra over itself (3.14.3),* $\mathcal{M}_n = \mathfrak{a}_n$ *is an ideal of* $M = A$*, so each* $M/\mathcal{M}_n$ *is also a ring and* $\iota$ *a ring homomorphism ((3.3.56) giving the same reduction maps as ring homomorphisms): this recovers exactly the completion of the filtered ring* $(A, \mathfrak{a}_\bullet)$ *(3.14.16).*

**Corollary 3.14.18 (Completion via a Cofinal Sequence)**
*Let* $(M, \mathcal{M}_\bullet)$ *be a filtered* $A$*-module, with completion* $\widehat{M}$*,* $\lambda_n$ *(3.14.17), and* $n_0 < n_1 < n_2 < \cdots$ *a strictly increasing sequence in* $\mathbb{N}$*, with projections* $\lambda'_N$ *of* $\varprojlim_N M/\mathcal{M}_{n_N}$*. Then*

$$\theta : \widehat{M} \xrightarrow{\sim} \varprojlim_N M/\mathcal{M}_{n_N}, \quad \theta\Big((\overline{v}_n)_{n\geq 0}\Big) = \big(\overline{v}_{n_N}\big)_{N\geq 0},$$

*is a bicontinuous* $A$*-module isomorphism, with* $\lambda'_N \circ \theta = \lambda_{n_N}$ *for every* $N \geq 0$*. (When* $M := A$*, this is a bicontinuous ring isomorphism.)*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

That $\theta$ is an $A$-module isomorphism with $\lambda'_N \circ \theta = \lambda_{n_N}$ for every $N \geq 0$ is (3.3.158), applied to the tower $(M/\mathcal{M}_n)_{n\geq 0}$ of $A$-modules (3.14.17) and this sequence. As $\theta$ is bijective, this gives $\theta\big(\ker \lambda_{n_N}\big) = \ker \lambda'_N$ for every $N \geq 0$.

**Continuity of $\theta$.** Given $N \geq 0$, taking $n := n_N$: $\theta(\ker \lambda_n) = \ker \lambda'_N$.

**Continuity of $\theta^{-1}$.** Given $n \geq 0$: as $(n_N)_N$ is unbounded, choose $N$ with $n_N \geq n$, so $\ker \lambda_{n_N} \subseteq \ker \lambda_n$ ((3.14.17), the filtration $\ker \lambda_\bullet$ decreasing); then $\theta^{-1}(\ker \lambda'_N) = \ker \lambda_{n_N} \subseteq \ker \lambda_n$. ∎

**Definition 3.14.19 ($\mathfrak{a}$-adic Completion)**
*Let* $A$ *be a ring and* $\mathfrak{a} \lhd A$ *an ideal. The* ***$\mathfrak{a}$-adic completion*** *of* $A$ *is the completion (3.14.16) of* $A$ *for the* $\mathfrak{a}$*-adic filtration (3.14.4):*

$$\hat{A}_\mathfrak{a} := \hat{A} = \varprojlim_n A/\mathfrak{a}^{n+1},$$

*with projections* $\lambda_n : \hat{A}_\mathfrak{a} \to A/\mathfrak{a}^{n+1}$ *and canonical ring homomorphism*

$$\iota_\mathfrak{a} : A \to \hat{A}_\mathfrak{a}, \quad a \mapsto (a \bmod \mathfrak{a}^{n+1})_{n\geq 0},$$

*as in (3.14.16).*

**Proposition 3.14.20 (Completeness Criterion)**
*Let* $(M, \mathcal{M}_\bullet)$ *be a* separated *filtered* $A$*-module (3.14.2), and* $\widehat{M}$*,* $\iota$ *as in (3.14.17). Then the following are equivalent:*

*a)* $\iota : M \to \widehat{M}$ *is surjective;*
*b) every sequence* $(\overline{v}_n)_{n\geq 0}$*,* $\overline{v}_n \in M/\mathcal{M}_n$*, compatible under reduction (*$\overline{v}_n \equiv \overline{v}_{n-1} \bmod \mathcal{M}_{n-1}$ *for* $n \geq 1$*), equals* $(v \bmod \mathcal{M}_n)_n$ *for some* $v \in M$*;*
*c) for some strictly increasing sequence* $n_0 < n_1 < n_2 < \cdots$ *in* $\mathbb{N}$ *(equivalently, for every such sequence), every sequence* $(\overline{v}_N)_{N\geq 0}$*,* $\overline{v}_N \in M/\mathcal{M}_{n_N}$*, compatible under reduction, equals* $\big(v \bmod \mathcal{M}_{n_N}\big)_N$ *for some* $v \in M$*.*

*When these hold,* $\iota$ *is an* $A$*-module isomorphism, and we say* $(M, \mathcal{M}_\bullet)$ *is* ***complete****. (When* $M := A$*,* $\iota$ *is a ring isomorphism, and we say the filtered ring* $(A, \mathfrak{a}_\bullet)$ *is* ***complete****.)*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

By (3.3.154), $\widehat{M} = \varprojlim_n M/\mathcal{M}_n$ consists precisely of the compatible sequences $(\overline{v}_n)_{n\geq 0}$ of b); and $\iota(v) = (v \bmod \mathcal{M}_n)_n$ by definition. So b) says exactly that every element of $\widehat{M}$ lies in the image of $\iota$, that is, a).

a) $\iff$ c): fix a strictly increasing sequence $n_0 < n_1 < n_2 < \cdots$ in $\mathbb{N}$. By (3.14.18), $(\overline{v}_n)_{n\geq 0} \mapsto \big(\overline{v}_{n_N}\big)_{N\geq 0}$ is an isomorphism $\widehat{M} \xrightarrow{\sim} \varprojlim_N M/\mathcal{M}_{n_N}$; composing with $\iota$, this sends $v \in M$ to $\big(v \bmod \mathcal{M}_{n_N}\big)_N$, so identifies the image of $\iota$ with the image of $v \mapsto \big(v \bmod \mathcal{M}_{n_N}\big)_N$. Hence $\iota$ is surjective iff this latter map is, that is, iff c) holds for this particular sequence $(n_N)_N$; as $(n_N)_N$ was an arbitrary strictly increasing sequence, a) is equivalent to c) holding for some such sequence, and also to it holding for every such sequence.

© David Rufino CC BY-NC-ND 4.0

Finally, $\ker\iota=\bigcap_n\mathcal{M}_n=0$ ($(M,\mathcal{M}_\bullet)$ being separated): $\iota(v)=0$ iff $v \bmod \mathcal{M}_n=0$ for every $n$, that is $v\in\mathcal{M}_n$ for every $n$. So $\iota$ is injective, hence, combined with a), an $A$-module isomorphism (a ring isomorphism when $M:=A$, as $\iota$ is then also a ring homomorphism (3.14.16)). ∎

**Proposition 3.14.21 (Completeness Criterion, $\mathfrak{a}$-adic Case)**
*Let $A$ be a ring, $\mathfrak{a}\triangleleft A$ an ideal with $\bigcap_n\mathfrak{a}^n=0$ (that is, $A$ is separated for the $\mathfrak{a}$-adic filtration), and $\hat{A}_\mathfrak{a}$, $\iota_\mathfrak{a}$ as in (3.14.19). This is the case $B:=A$, $\mathcal{M}_n:=\mathfrak{a}_n:=\mathfrak{a}^{n+1}$ of (3.14.20), giving: the following are equivalent:*

- *a) $\iota_\mathfrak{a}:A\to\hat{A}_\mathfrak{a}$ is surjective;*
- *b) every sequence $(\overline{a}_n)_{n\geq0}$, $\overline{a}_n\in A/\mathfrak{a}^{n+1}$, compatible under reduction ($\overline{a}_n\equiv\overline{a}_{n-1} \bmod \mathfrak{a}^n$ for $n\geq1$), equals $(a \bmod \mathfrak{a}^{n+1})_n$ for some $a\in A$;*
- *c) for some strictly increasing sequence $n_0<n_1<n_2<\cdots$ in $\mathbb{N}$ (equivalently, for every such sequence), every sequence $(\overline{a}_N)_{N\geq0}$, $\overline{a}_N\in A/\mathfrak{a}^{n_N+1}$, compatible under reduction, equals $(a \bmod \mathfrak{a}^{n_N+1})_N$ for some $a\in A$.*

*When these hold, $\iota_\mathfrak{a}$ is an isomorphism, and we say $A$ is **$\mathfrak{a}$-adically complete**.*

**Proposition 3.14.22 (Products of Filtered Modules)**
*Let $A$ be a ring, $K$ a set, and $(M_k,\mathcal{M}_{k,\bullet})_{k\in K}$ a family of filtered $A$-modules (3.14.2), with completions $\widehat{M}_k$, $\lambda_{k,n}$ (3.14.17). Then $\mathcal{M}_\bullet:=\left(\prod_{k\in K}\mathcal{M}_{k,n}\right)_{n\in\mathbb{Z}}$ is a filtration of the $A$-module $M:=\prod_{k\in K}M_k$, with completion $\widehat{M}$, $\lambda_n$. There is a unique $A$-module homomorphism $\Psi$ making*

$$\begin{array}{ccc}\prod_{k\in K}\widehat{M}_k & \xrightarrow{\ \Psi\ } & \widehat{\left(\prod_{k\in K}M_k\right)}\\ \Big\downarrow{\scriptstyle \prod_k\lambda_{k,n}} & & \Big\downarrow{\scriptstyle \lambda_n}\\ \prod_{k\in K}(M_k/\mathcal{M}_{k,n}) & \xleftarrow[\sim]{\ \overline{q}_n\ } & \left(\prod_{k\in K}M_k\right)/\mathcal{M}_n\end{array}$$

*commute for every $n\geq0$, where $\overline{q}_n\left((m_k)_k \bmod \mathcal{M}_n\right):=\left(m_k \bmod \mathcal{M}_{k,n}\right)_k$; explicitly, writing $\hat{x}_k=(\overline{m}_{k,n})_{n\geq0}$ for $\hat{x}_k\in\widehat{M}_k$,*

$$\Psi\left((\hat{x}_k)_k\right)=\left((m_{k,n})_k \bmod \mathcal{M}_n\right)_{n\geq0}$$

*for any representatives $m_{k,n}\in M_k$ of $\overline{m}_{k,n}$; and $\Psi$ is a bicontinuous $A$-module isomorphism, for the product filtration $\left(\prod_k\ker\lambda_{k,n}\right)_n$ of $\prod_k\widehat{M}_k$ and $\ker\lambda_\bullet$ of $\widehat{M}$. (When $K$ is* finite *and each $M_k:=A_k$ is taken to be a filtered ring over itself, $\Psi$ is a bicontinuous ring isomorphism, recovering the completion of the product of filtered rings $(A_k,\mathfrak{a}_{k,\bullet})_{k\in K}$ – exhaustiveness of the product filtration, $\bigcup_n\prod_k\mathfrak{a}_{k,n}=\prod_kA_k$, using finiteness of $K$ (taking, for each $(a_k)_k$, $n:=\min_k n_k$ over the finitely many $n_k$ with $a_k\in\mathfrak{a}_{k,n_k}$); for infinite $K$, exhaustiveness can fail even if every factor is exhaustive, as no single $n$ need work for infinitely many coordinates at once.)*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

**$\mathcal{M}_\bullet$ is a filtration of $\prod_k M_k$.** Each $\mathcal{M}_n:=\prod_k\mathcal{M}_{k,n}$ is an $A$-submodule of $\prod_kM_k$ (a product of submodules, coordinatewise), and $\mathcal{M}_{n+1}\subseteq\mathcal{M}_n$ as $\mathcal{M}_{k,n+1}\subseteq\mathcal{M}_{k,n}$ for every $k$.

**A levelwise isomorphism.** For each $n$, the coordinatewise reduction $q_n:\prod_kM_k\to\prod_k(M_k/\mathcal{M}_{k,n})$, $(m_k)_k\mapsto(m_k \bmod \mathcal{M}_{k,n})_k$, is a surjective $A$-module homomorphism with kernel $\mathcal{M}_n$, so induces an isomorphism $\overline{q}_n:\left(\prod_kM_k\right)/\mathcal{M}_n\xrightarrow{\sim}\prod_k(M_k/\mathcal{M}_{k,n})$ (3.3.102), compatible with reduction as $n$ varies (both sides' reduction maps being coordinatewise reduction). So, by (3.3.156), $\varprojlim\overline{q}:\widehat{\left(\prod_kM_k\right)}\xrightarrow{\sim}\varprojlim_n\left(\prod_k(M_k/\mathcal{M}_{k,n})\right)$ is an isomorphism.

**Composing with (3.3.157).** That proposition (applied to the towers $(M_k/\mathcal{M}_{k,n})_n$ of $A$-modules) gives an isomorphism $\Phi:\prod_k\widehat{M}_k\xrightarrow{\sim}\varprojlim_n\left(\prod_k(M_k/\mathcal{M}_{k,n})\right)$. Set $\Psi:=\left(\varprojlim\overline{q}\right)^{-1}\circ\Phi$, an isomorphism.

**$\Psi$ has the stated property.** By the explicit formulas of (3.3.156) and (3.3.157) for the projections of $\varprojlim\overline{q}$ and $\Phi$ respectively: for $(\hat{x}_k)_k\in\prod_k\widehat{M}_k$, the $n$-th coordinate of $\Phi\left((\hat{x}_k)_k\right)$ is $(\lambda_{k,n}(\hat{x}_k))_k\in\prod_k(M_k/\mathcal{M}_{k,n})$, and this equals $\overline{q}_n\left(\lambda_n\left(\Psi\left((\hat{x}_k)_k\right)\right)\right)$ ($\varprojlim\overline{q}$'s $n$-th coordinate being $\overline{q}_n\circ\lambda_n$); so $\overline{q}_n\left(\lambda_n\left(\Psi\left((\hat{x}_k)_k\right)\right)\right)=(\lambda_{k,n}(\hat{x}_k))_k=(\overline{m}_{k,n})_k$: the square commutes, for every $n$. Unwinding $\overline{q}_n^{-1}$ (3.3.102), $\lambda_n\left(\Psi\left((\hat{x}_k)_k\right)\right)=(m_{k,n})_k \bmod \mathcal{M}_n$ for any representatives $m_{k,n}$ of $\overline{m}_{k,n}$, giving the stated explicit formula for $\Psi$.

© David Rufino CC BY-NC-ND 4.0

**Uniqueness.** $\left(\widehat{\prod_k M_k}\right) = \varprojlim_n \left(\prod_k M_k\right)/\mathcal{M}_n$ is universal for compatible families of morphisms into the tower $\left(\left(\prod_k M_k\right)/\mathcal{M}_n\right)_n$ (3.3.154): an $A$-module homomorphism $\Psi'$ making the square commute for every $n$ is exactly a compatible family of morphisms $\prod_k \widehat{M}_k \to \left(\prod_k M_k\right)/\mathcal{M}_n$ (namely $\overline{q}_n^{-1} \circ \prod_k \lambda_{k,n}$, compatible as $n$ varies since the $\lambda_{k,n}$ and $\overline{q}_n$ are), and that universal property gives a unique such morphism: so $\Psi' = \Psi$.

**Bicontinuity.** Rearranging the commuting square, $\lambda_n \circ \Psi = \overline{q}_n^{-1} \circ \prod_k \lambda_{k,n}$ for every $n \geq 0$. For continuity of $\Psi$: given $n$, if $(\hat{x}_k)_k \in \prod_k \ker \lambda_{k,n}$ then $\lambda_n\left(\Psi\left((\hat{x}_k)_k\right)\right) = \overline{q}_n^{-1}\left((\lambda_{k,n}(\hat{x}_k))_k\right) = \overline{q}_n^{-1}(0) = 0$, so $\Psi\left(\prod_k \ker \lambda_{k,n}\right) \subseteq \ker \lambda_n$ (same index $n$). For continuity of $\Psi^{-1}$: given $n$, if $x \in \ker \lambda_n$, applying $\overline{q}_n$ to $\lambda_n(x) = 0$ gives $0 = \overline{q}_n(\lambda_n(x)) = \left(\lambda_{k,n}(\Psi^{-1}(x)_k)\right)_k$ (the displayed equation, applied to $\Psi^{-1}(x)$ and rearranged), so $\lambda_{k,n}(\Psi^{-1}(x)_k) = 0$ for every $k$, that is $\Psi^{-1}(x) \in \prod_k \ker \lambda_{k,n}$: $\Psi^{-1}(\ker \lambda_n) \subseteq \prod_k \ker \lambda_{k,n}$ (same index $n$). ∎

**Lemma 3.14.23 (Continuity into a Finite Product)**
*Let $(A, \mathfrak{a}_\bullet)$ be a filtered ring, $(N, \mathcal{N}_\bullet)$ and $(M_i, \mathcal{M}_{i,\bullet})_{i=1}^n$ filtered $A$-modules (3.14.2), and equip $\prod_{i=1}^n M_i$ with the product filtration (3.14.22). Then an $A$-module homomorphism $\phi : N \to \prod_{i=1}^n M_i$ is continuous (3.14.7) (with $\rho := \mathrm{id}_A$) if and only if each $\phi_i := \pi_i \circ \phi : N \to M_i$ is continuous, where $\pi_i$ is the $i$-th projection. (Taking $N := A$ and each $M_i$ as filtered rings over themselves recovers the corresponding test for (3.14.5).)*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

If $\phi$ is continuous, then for each $J$ there is $L$ with $\phi(\mathcal{N}_L) \subseteq \prod_i \mathcal{M}_{i,J}$; applying $\pi_i$, $\phi_i(\mathcal{N}_L) \subseteq \mathcal{M}_{i,J}$, so each $\phi_i$ is continuous.

Conversely, suppose each $\phi_i$ is continuous. Fix $J$: for each $i = 1, \ldots, n$ there is $m_i$ with $\phi_i\left(\mathcal{N}_{m_i}\right) \subseteq \mathcal{M}_{i,J}$; as there are finitely many $i$, $L := \max_i m_i$ satisfies $\mathcal{N}_L \subseteq \mathcal{N}_{m_i}$ for every $i$ (the filtration being decreasing), so $\phi_i(\mathcal{N}_L) \subseteq \mathcal{M}_{i,J}$ for every $i$, that is $\phi(\mathcal{N}_L) \subseteq \prod_i \mathcal{M}_{i,J}$: $\phi$ is continuous. ∎

**Lemma 3.14.24 (Projections from the Completion are Surjective)**
*Let $(M, \mathcal{M}_\bullet)$ be a filtered $A$-module, and $\widehat{M}$, $\lambda_n$ as in (3.14.17). Then each $\lambda_n$ is surjective. (In particular, this applies to $\hat{A}_\mathfrak{a}$, $\lambda_n$ as in (3.14.19), the case $M := A$, $\mathfrak{a}_n := \mathfrak{a}^{n+1}$.)*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Given $\overline{v}_n \in M/\mathcal{M}_n$, choose any representative $v_n \in M$. As each reduction $M/\mathcal{M}_{m+1} \twoheadrightarrow M/\mathcal{M}_m$ is surjective (3.14.17), recursively choose $v_{m+1} \in M$ reducing to (the class of) $v_m$ for every $m \geq n$; for $m < n$, let $\overline{v}_m$ be the reduction of $v_n$. This gives a compatible sequence $(\overline{v}_m)_{m \geq 0} \in \widehat{M}$ with $\lambda_n\left((\overline{v}_m)_m\right) = \overline{v}_n$. ∎

**Proposition 3.14.25 (Completion Matches the Filtration)**
*Let $(M, \mathcal{M}_\bullet)$ be a filtered $A$-module, with completion $\widehat{M}$, $\iota$, $\lambda_n$ (3.14.17). For every $n \geq 0$, $\iota(\mathcal{M}_n) \subseteq \ker \lambda_n \subseteq \ker \lambda_{n-1}$ (the latter for $n \geq 1$), and, writing $\overline{\pi}_n : M \to M/\mathcal{M}_n$ and $\pi_n : \widehat{M} \to \widehat{M}/\ker \lambda_n$ for the quotient maps, there is a unique $A$-module homomorphism $\nu_n : M/\mathcal{M}_n \to \widehat{M}/\ker \lambda_n$ (3.3.101) such that the square*

$$\begin{array}{ccc} M & \xrightarrow{\ \iota\ } & \widehat{M} \\ {\scriptstyle \overline{\pi}_n}\downarrow & & \downarrow{\scriptstyle \pi_n} \\ M/\mathcal{M}_n & \xrightarrow[\nu_n]{} & \widehat{M}/\ker \lambda_n \end{array}$$

*commutes, that is $\nu_n \circ \overline{\pi}_n = \pi_n \circ \iota$; and $\nu_n$ is an isomorphism, with inverse $\mu_n := \nu_n^{-1}$ satisfying $\mu_n \circ \pi_n = \lambda_n$. Moreover, writing $\rho_n : M/\mathcal{M}_n \to M/\mathcal{M}_{n-1}$ for the natural reduction (well-defined by (3.3.101), as $\mathcal{M}_n \subseteq \mathcal{M}_{n-1}$) and $\hat{\rho}_n : \widehat{M}/\ker \lambda_n \to \widehat{M}/\ker \lambda_{n-1}$ for the one induced by $\ker \lambda_n \subseteq \ker \lambda_{n-1}$, $(\nu_n)_{n \geq 0}$ is natural in $n$: for every $n \geq 1$, the square*

$$\begin{array}{ccc} M/\mathcal{M}_n & \xrightarrow[\sim]{\nu_n} & \widehat{M}/\ker \lambda_n \\ {\scriptstyle \rho_n}\downarrow & & \downarrow{\scriptstyle \hat{\rho}_n} \\ M/\mathcal{M}_{n-1} & \xrightarrow[\sim]{\nu_{n-1}} & \widehat{M}/\ker \lambda_{n-1} \end{array}$$

*commutes, that is $\hat{\rho}_n \circ \nu_n = \nu_{n-1} \circ \rho_n$ (equivalently, $\rho_n \circ \mu_n = \mu_{n-1} \circ \hat{\rho}_n$). (When $M := A$, $\nu_n$, $\mu_n$ are ring isomorphisms, and moreover the stronger inclusion $\mathfrak{a}_n \hat{A} \subseteq \ker \lambda_n$ holds: for $c \in \mathfrak{a}_n$, $y \in \hat{A}$, $\lambda_n(\iota(c)y) = \lambda_n(\iota(c))\lambda_n(y) = 0$, using that $\hat{A}$ is then a ring and $\lambda_n$ a ring homomorphism.)*

© David Rufino CC BY-NC-ND 4.0

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

**$\iota(\mathcal{M}_n) \subseteq \ker\lambda_n \subseteq \ker\lambda_{n-1}$.** For $c \in \mathcal{M}_n$, $\lambda_n(\iota(c)) = c \bmod \mathcal{M}_n = 0$ by definition of $\iota$, giving $\iota(\mathcal{M}_n) \subseteq \ker\lambda_n$. For $n \geq 1$: if $x = (\overline{v}_m)_m$ has $\lambda_n(x) = \overline{v}_n = 0$, then $\overline{v}_{n-1}$, being the reduction of $\overline{v}_n$ modulo $\mathcal{M}_{n-1}$ (as $(\overline{v}_m)_m$ is compatible under reduction), is also 0; so $\lambda_{n-1}(x) = 0$, giving $\ker\lambda_n \subseteq \ker\lambda_{n-1}$.

**$\nu_n$.** As $\iota(\mathcal{M}_n) \subseteq \ker\lambda_n$ (above), the composite $\pi_n \circ \iota : M \to \widehat{M}/\ker\lambda_n$ kills $\mathcal{M}_n = \ker\overline{\pi}_n$, so (3.3.101) gives the unique $A$-module homomorphism $\nu_n : M/\mathcal{M}_n \to \widehat{M}/\ker\lambda_n$ with $\nu_n \circ \overline{\pi}_n = \pi_n \circ \iota$, that is $\nu_n(v \bmod \mathcal{M}_n) = \pi_n(\iota(v))$ for $v \in M$.

$\nu_n$ is injective: if $\nu_n(v \bmod \mathcal{M}_n) = 0$, that is $\iota(v) \in \ker\lambda_n$, then $0 = \lambda_n(\iota(v)) = v \bmod \mathcal{M}_n$ (by definition of $\iota$).

$\nu_n$ is surjective, and $\pi_n = \nu_n \circ \lambda_n$: given $x \in \widehat{M}$, let $v \in M$ be any representative of $\lambda_n(x) \in M/\mathcal{M}_n$ ($\overline{\pi}_n$ being surjective); then $\lambda_n(\iota(v)) = v \bmod \mathcal{M}_n = \lambda_n(x)$, so $x - \iota(v) \in \ker\lambda_n$, giving $\pi_n(x) = \pi_n(\iota(v)) = \nu_n(v \bmod \mathcal{M}_n) = \nu_n(\lambda_n(x))$; in particular every $\pi_n(x)$ is of the form $\nu_n(-)$.

So $\nu_n$ is an isomorphism. **$\mu_n$.** Set $\mu_n := \nu_n^{-1}$; then $\mu_n \circ \pi_n = \mu_n \circ \nu_n \circ \lambda_n = \lambda_n$ (using $\pi_n = \nu_n \circ \lambda_n$, just shown), as stated.

**Naturality.** $\hat{\rho}_n$ is well-defined since $\ker\lambda_n \subseteq \ker\lambda_{n-1}$, and $\hat{\rho}_n \circ \pi_n = \pi_{n-1}$, $\rho_n \circ \overline{\pi}_n = \overline{\pi}_{n-1}$ (the standard compatibility of quotient maps). For $v \in M$: $\hat{\rho}_n(\nu_n(\overline{\pi}_n(v))) = \hat{\rho}_n(\pi_n(\iota(v))) = \pi_{n-1}(\iota(v)) = \nu_{n-1}(\overline{\pi}_{n-1}(v)) = \nu_{n-1}(\rho_n(\overline{\pi}_n(v)))$; as $\overline{\pi}_n$ is surjective, this gives $\hat{\rho}_n \circ \nu_n = \nu_{n-1} \circ \rho_n$. Inverting (composing with $\mu_n = \nu_n^{-1}$ on the right and $\mu_{n-1} = \nu_{n-1}^{-1}$ on the left) gives the equivalent $\rho_n \circ \mu_n = \mu_{n-1} \circ \hat{\rho}_n$. ∎

**Proposition 3.14.26 (The Completion is Complete)**
*Let $(M, \mathcal{M}_\bullet)$ be a filtered $A$-module, with completion $\widehat{M}$, $\iota$, $\lambda_n$ (3.14.17), filtered by $\ker\lambda_\bullet$. Then $\left(\widehat{M}, \ker\lambda_\bullet\right)$ is separated (3.14.2) and complete (3.14.20). (When $M := A$ is a filtered ring, $\left(\hat{A}, \ker\lambda_\bullet\right)$ is moreover again a filtered ring (3.14.16).)*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Write $\pi_n : \widehat{M} \to \widehat{M}/\ker\lambda_n$ for the quotient maps and $\rho_n : M/\mathcal{M}_n \to M/\mathcal{M}_{n-1}$, $\hat{\rho}_n : \widehat{M}/\ker\lambda_n \to \widehat{M}/\ker\lambda_{n-1}$ for the reduction maps. By (3.14.25), there is an isomorphism $\mu_n : \widehat{M}/\ker\lambda_n \xrightarrow{\sim} M/\mathcal{M}_n$ with $\mu_n \circ \pi_n = \lambda_n$, natural in $n$: $\rho_n \circ \mu_n = \mu_{n-1} \circ \hat{\rho}_n$.

**Separated.** If $x = (\overline{v}_n)_{n\geq 0} \in \widehat{M}$ (a compatible sequence, $\overline{v}_n \in M/\mathcal{M}_n$) has $\lambda_n(x) = 0$ for every $n$, that is $\overline{v}_n = 0$ for every $n$, then $x = 0$; so $\bigcap_n \ker\lambda_n = 0$, that is $\left(\widehat{M}, \ker\lambda_\bullet\right)$ is separated (3.14.2).

**Complete** (item b)). By the above, (3.14.20) applies to $\left(\widehat{M}, \ker\lambda_\bullet\right)$; we verify its item b). Let $(\overline{x}_n)_{n\geq 0}$, $\overline{x}_n \in \widehat{M}/\ker\lambda_n$, be compatible under reduction (that is $\hat{\rho}_n(\overline{x}_n) = \overline{x}_{n-1}$ for $n \geq 1$). Writing $\overline{v}_n := \mu_n(\overline{x}_n) \in M/\mathcal{M}_n$: for $n \geq 1$, naturality gives $\rho_n(\overline{v}_n) = \rho_n(\mu_n(\overline{x}_n)) = \mu_{n-1}(\hat{\rho}_n(\overline{x}_n)) = \mu_{n-1}(\overline{x}_{n-1}) = \overline{v}_{n-1}$: the sequence $(\overline{v}_n)_n$ is compatible. So $(\overline{v}_n)_n \in \widehat{M}$ itself, with $\lambda_n(\overline{v}_\bullet) = \overline{v}_n$ for every $n$ (its own $n$-th coordinate); then $\mu_n(\pi_n(\overline{v}_\bullet)) = \lambda_n(\overline{v}_\bullet) = \overline{v}_n = \mu_n(\overline{x}_n)$, and $\mu_n$ injective gives $\pi_n(\overline{v}_\bullet) = \overline{x}_n$: the compatible sequence $(\overline{x}_n)_n$ arises from $\overline{v}_\bullet \in \widehat{M}$. So $\left(\widehat{M}, \ker\lambda_\bullet\right)$ is complete.∎

**Proposition 3.14.27 (Functoriality of Completion)**
*With completions $\widehat{M}$, $\iota_M$, $\lambda_\bullet^M$ and $\widehat{M'}$, $\iota_{M'}$, $\lambda_\bullet^{M'}$ (3.14.17), let $\psi : M \to M'$ be a continuous $\rho$-homomorphism (3.14.7). Then there is a unique continuous $\rho$-homomorphism $\hat{\psi} : \widehat{M} \to \widehat{M'}$ (for the filtrations $\ker\lambda_\bullet^M$, $\ker\lambda_\bullet^{M'}$) with $\hat{\psi} \circ \iota_M = \iota_{M'} \circ \psi$.*

*Moreover this is functorial: $\widehat{\mathrm{id}_M} = \mathrm{id}_{\widehat{M}}$ (taking $\rho := \mathrm{id}_A$); and, for a further filtered $A''$-module $(N, \mathcal{N}_\bullet)$, ring homomorphism $\rho' : A' \to A''$, and continuous $\rho'$-homomorphism $\chi : M' \to N$ (so $\chi \circ \psi$ is a continuous $(\rho' \circ \rho)$-homomorphism), $\widehat{\chi \circ \psi} = \hat{\chi} \circ \hat{\psi}$.*

*Moreover (**surjectivity criterion**), if $\psi$ is taut (3.14.11) then $\hat{\psi}$ is surjective.*

*Moreover (**injectivity criterion**), if $\psi(\mathcal{M}_n) \subseteq \mathcal{M}'_n$ for every $n \geq 0$ and the induced map $\overline{\psi}_{n_J} : M/\mathcal{M}_{n_J} \to M'/\mathcal{M}'_{n_J}$ (3.3.101) is injective for every $J \geq 0$, for some strictly increasing sequence $n_0 < n_1 < n_2 < \cdots$, then $\hat{\psi}$ is injective.*

*(Taking $M := A$, $M' := A'$ as filtered $A$-, $A'$-modules over themselves (3.14.3) and $\psi := \rho$ recovers (3.14.5) and the ring-level statement of this proposition exactly.)*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Write $\overline{\pi}_n^M : M \to M/\mathcal{M}_n$, $\overline{\pi}_m^{M'} : M' \to M'/\mathcal{M}'_m$ for the canonical surjections, and $\rho^M_{n,n'} : M/\mathcal{M}_n \to M/\mathcal{M}_{n'}$ ($n \geq n'$) for $M$'s reduction maps. Throughout, regard $M'/\mathcal{M}'_m$ as an $A$-module via restriction of scalars along $\rho$ ($a \cdot x := \rho(a) \cdot x$); a $\rho$-linear map out of $M$ (or a quotient of $M$) is then exactly an $A$-linear map into $M'/\mathcal{M}'_m$ for this restricted structure, so (3.3.101) applies to it unchanged.

© David Rufino CC BY-NC-ND 4.0

**A compatible family of maps into the levels of $\widehat{M'}$.** For each $m \geq 0$, by continuity of $\psi$, choose $n(m) \geq 0$ with $\psi\big(\mathcal{M}_{n(m)}\big) \subseteq \mathcal{M}'_m$; (3.3.101) then gives a unique $A$-linear (that is, $\rho$-linear) map $\overline{\psi}_{n(m),m} : M/\mathcal{M}_{n(m)} \to M'/\mathcal{M}'_m$ with $\overline{\psi}_{n(m),m} \circ \overline{\pi}^M_{n(m)} = \overline{\pi}^{M'}_m \circ \psi$. Set $\psi_m := \overline{\psi}_{n(m),m} \circ \lambda^M_{n(m)} : \widehat{M} \to M'/\mathcal{M}'_m$.

These are compatible as $m$ varies: fix $m$, write $\tau_m : M'/\mathcal{M}'_{m+1} \to M'/\mathcal{M}'_m$ for the reduction, and set $n := \max(n(m), n(m+1))$. As $\mathcal{M}_n \subseteq \mathcal{M}_{n(m)}, \mathcal{M}_{n(m+1)}$, (3.3.101) gives $\overline{\psi}_{n,m} : M/\mathcal{M}_n \to M'/\mathcal{M}'_m$, $\overline{\psi}_{n,m+1} : M/\mathcal{M}_n \to M'/\mathcal{M}'_{m+1}$ (the same defining property, through the finer level $n$); by uniqueness there, $\overline{\psi}_{n(m),m} \circ \rho^M_{n,n(m)} = \overline{\psi}_{n,m}$, $\overline{\psi}_{n(m+1),m+1} \circ \rho^M_{n,n(m+1)} = \overline{\psi}_{n,m+1}$, and $\tau_m \circ \overline{\psi}_{n,m+1} = \overline{\psi}_{n,m}$ (all being the map $M/\mathcal{M}_n \to M'/\mathcal{M}'_m$ induced by $\psi$). As $\lambda^M_{n(m)} = \rho^M_{n,n(m)} \circ \lambda^M_n$ and $\lambda^M_{n(m+1)} = \rho^M_{n,n(m+1)} \circ \lambda^M_n$ (compatibility of $\widehat{M}$'s own projections),

$$\tau_m \circ \psi_{m+1} = \tau_m \circ \overline{\psi}_{n(m+1),m+1} \circ \rho^M_{n,n(m+1)} \circ \lambda^M_n = \tau_m \circ \overline{\psi}_{n,m+1} \circ \lambda^M_n = \overline{\psi}_{n,m} \circ \lambda^M_n = \overline{\psi}_{n(m),m} \circ \rho^M_{n,n(m)} \circ \lambda^M_n = \psi_m.$$

**Existence.** By the universal property of $\widehat{M'} = \varprojlim_m M'/\mathcal{M}'_m$ as an inverse limit (2.6.129), the compatible family $(\psi_m)_m$ gives a unique $A$-linear (that is, $\rho$-linear) map $\hat{\psi} : \widehat{M} \to \widehat{M'}$ with $\lambda^{M'}_m \circ \hat{\psi} = \psi_m$ for every $m$. (If $M := A$, $M' := A'$ and $\psi$ is moreover a ring homomorphism with $\psi(\mathfrak{a}_n) \subseteq \mathfrak{a}'_{n(m)}$ at the same witnessing indices, $\hat{\psi}$ is a ring homomorphism too, by the same construction run with $\overline{\psi}_{n(m),m}$ the induced ring homomorphisms.)

For $v \in M$: $\lambda^{M'}_m\big(\hat{\psi}(\iota_M(v))\big) = \psi_m(\iota_M(v)) = \overline{\psi}_{n(m),m}\big(v \bmod \mathcal{M}_{n(m)}\big) = \psi(v) \bmod \mathcal{M}'_m = \lambda^{M'}_m(\iota_{M'}(\psi(v)))$; as $m$ was arbitrary, $\hat{\psi}(\iota_M(v)) = \iota_{M'}(\psi(v))$: the square commutes.

For continuity of $\hat{\psi}$: fix $m$; for $x \in \ker \lambda^M_{n(m)}$, $\lambda^{M'}_m\big(\hat{\psi}(x)\big) = \psi_m(x) = \overline{\psi}_{n(m),m}(0) = 0$, so $\hat{\psi}(x) \in \ker \lambda^{M'}_m$: taking $n := n(m)$ verifies $\hat{\psi}(\ker \lambda^M_n) \subseteq \ker \lambda^{M'}_m$.

**Uniqueness.** Suppose $\hat{\psi}_1, \hat{\psi}_2 : \widehat{M} \to \widehat{M'}$ are both continuous $\rho$-homomorphisms with $\hat{\psi}_i \circ \iota_M = \iota_{M'} \circ \psi$. Fix $x \in \widehat{M}$ and $m \geq 0$; choose $n_i$ with $\hat{\psi}_i\big(\ker \lambda^M_{n_i}\big) \subseteq \ker \lambda^{M'}_m$ $(i = 1, 2)$, and set $n := \max(n_1, n_2)$, so $\ker \lambda^M_n \subseteq \ker \lambda^M_{n_i}$ (3.14.17) and $\hat{\psi}_i(\ker \lambda^M_n) \subseteq \ker \lambda^{M'}_m$ for both $i$. By surjectivity of $\lambda^M_n$ (3.14.24), choose $v \in M$ with $\lambda^M_n(\iota_M(v)) = \lambda^M_n(x)$, i.e. $\iota_M(v) - x \in \ker \lambda^M_n$. Then $\hat{\psi}_i(\iota_M(v)) - \hat{\psi}_i(x) = \hat{\psi}_i(\iota_M(v) - x) \in \ker \lambda^{M'}_m$, and $\hat{\psi}_i(\iota_M(v)) = \iota_{M'}(\psi(v))$; so $\hat{\psi}_i(x) \equiv \iota_{M'}(\psi(v)) \bmod \ker \lambda^{M'}_m$ for $i = 1, 2$, giving $\hat{\psi}_1(x) - \hat{\psi}_2(x) \in \ker \lambda^{M'}_m$. As $m \geq 0$ was arbitrary, $\hat{\psi}_1(x) - \hat{\psi}_2(x) \in \bigcap_m \ker \lambda^{M'}_m = 0$ (3.14.26) ($\widehat{M'}$ always separated); so $\hat{\psi}_1(x) = \hat{\psi}_2(x)$.

**Identities.** $\mathrm{id}_{\widehat{M}}$ is continuous and satisfies $\mathrm{id}_{\widehat{M}} \circ \iota_M = \iota_M \circ \mathrm{id}_M$; by the uniqueness just shown (applied to $\psi := \mathrm{id}_M$, $\rho := \mathrm{id}_A$), $\widehat{\mathrm{id}_M} = \mathrm{id}_{\widehat{M}}$.

**Composition.** $\hat{\chi} \circ \hat{\psi} : \widehat{M} \to \widehat{N}$ is continuous (a composite of continuous maps) and is a $(\rho' \circ \rho)$-homomorphism (a composite of $\rho'$- and $\rho$-linear maps), satisfying $\big(\hat{\chi} \circ \hat{\psi}\big) \circ \iota_M = \hat{\chi} \circ \iota_{M'} \circ \psi = \iota_N \circ \chi \circ \psi$; by uniqueness (applied to $\chi \circ \psi$), $\widehat{\chi \circ \psi} = \hat{\chi} \circ \hat{\psi}$.

**Surjectivity criterion.** Suppose $\psi$ is taut (3.14.11), witnessed by a strictly increasing sequence $n_0 < n_1 < n_2 < \cdots$ with $M' = \psi(M) + \mathcal{M}'_{n_0}$ and $\mathcal{M}'_{n_J} = \psi\big(\mathcal{M}_{n_J}\big) + \mathcal{M}'_{n_{J+1}}$ for every $J \geq 0$; in particular $\psi(\mathcal{M}_n) \subseteq \mathcal{M}'_n$ for every $n \geq 0$, so, as above, $\overline{\psi}_n := \overline{\psi}_{n,n}$ is defined for every $n$, and $\hat{\psi} = \varprojlim \overline{\psi}$ by (3.3.156), taking $n(m) := m$ throughout. By (3.14.18) (applied to $M$ and to $M'$, with this sequence), $\widehat{M} \xrightarrow{\sim} \varprojlim_J M/\mathcal{M}_{n_J}$ and $\widehat{M'} \xrightarrow{\sim} \varprojlim_J M'/\mathcal{M}'_{n_J}$ (this identification of $\widehat{M'}$ with the compatible sequences of the subsequence needing no completeness of $M'$, only its definition as an inverse limit), identifying $\hat{\psi}$ with $\varprojlim_J \overline{\psi}_{n_J}$; so it suffices to show this restricted map hits every compatible sequence $\overline{v}' = \big(\overline{v}'_{n_J}\big)_{J \geq 0}$, $\overline{v}'_{n_J} \in M'/\mathcal{M}'_{n_J}$. We build, together with representatives $v \in M$, a compatible sequence $\big(\overline{v}_{n_J}\big)_{J \geq 0}$, $\overline{v}_{n_J} \in M/\mathcal{M}_{n_J}$, with $\overline{\psi}_{n_J}\big(\overline{v}_{n_J}\big) = \overline{v}'_{n_J}$ for every $J$.

Choose any representative $v' \in M'$ of $\overline{v}'_{n_0}$. By $M' = \psi(M) + \mathcal{M}'_{n_0}$, write $v' = \psi(v) + d$ for some $v \in M$, $d \in \mathcal{M}'_{n_0}$; then $\psi(v) \equiv v' \bmod \mathcal{M}'_{n_0}$, that is $\overline{\psi}_{n_0}\big(\overline{v}_{n_0}\big) = \overline{v}'_{n_0}$ for $\overline{v}_{n_0} := \overline{\pi}^M_{n_0}(v)$. Inductively, given $v \in M$ with $\overline{\psi}_{n_J}\big(\overline{\pi}^M_{n_J}(v)\big) = \overline{v}'_{n_J}$ $(J \geq 0)$: choose a representative $c' \in M'$ of $\overline{v}'_{n_{J+1}}$; as $\psi(v)$ and $c'$ have the same reduction mod $\mathcal{M}'_{n_J}$ (namely $\overline{\psi}_{n_J}\big(\overline{\pi}^M_{n_J}(v)\big) = \overline{v}'_{n_J}$ = the reduction of $\overline{v}'_{n_{J+1}}$, as $\overline{v}'$ is compatible), $\psi(v) - c' \in \mathcal{M}'_{n_J}$. By $\mathcal{M}'_{n_J} = \psi\big(\mathcal{M}_{n_J}\big) + \mathcal{M}'_{n_{J+1}}$, write $\psi(v) - c' = \psi(c) + e$ for some $c \in \mathcal{M}_{n_J}$, $e \in \mathcal{M}'_{n_{J+1}}$; then $\psi(v - c) - c' = -e \in \mathcal{M}'_{n_{J+1}}$, that is $\overline{\psi}_{n_{J+1}}\big(\overline{\pi}^M_{n_{J+1}}(v - c)\big) = \overline{v}'_{n_{J+1}}$. As $c \in \mathcal{M}_{n_J}$, $\overline{\pi}^M_{n_J}(v - c) = \overline{\pi}^M_{n_J}(v)$, so replacing $v$ by $v - c$ preserves the sequence built so far (compatible at every earlier level) while extending it to level $n_{J+1}$; continuing the induction, and setting $\overline{v}_{n_J} := \overline{\pi}^M_{n_J}(v)$ at each stage for the (eventually fixed) representative $v$ at that stage, gives the required compatible sequence.

**Injectivity criterion.** Suppose $\psi(\mathcal{M}_n) \subseteq \mathcal{M}'_n$ for every $n \geq 0$ and $\overline{\psi}_{n_J}$ is injective for every $J \geq 0$, for some strictly increasing sequence $n_0 < n_1 < n_2 < \cdots$. As above (taking $n(m) := m$ throughout), $\hat{\psi} = \varprojlim \overline{\psi}$ (3.3.156), and, by (3.14.18) (applied to $M$ and to $M'$, with this sequence), $\widehat{M} \xrightarrow{\sim} \varprojlim_J M/\mathcal{M}_{n_J}$ and $\widehat{M'} \xrightarrow{\sim} \varprojlim_J M'/\mathcal{M}'_{n_J}$, identifying $\hat{\psi}$ with $\varprojlim_J \overline{\psi}_{n_J}$. If $x = (\overline{v}_n)_{n \geq 0} \in \widehat{M}$ has $\hat{\psi}(x) = 0$, then (under this identification) $\overline{\psi}_{n_J}\big(\overline{v}_{n_J}\big) = 0$ for every $J$, so

© David Rufino CC BY-NC-ND 4.0

$\overline{v}_{n_J} = 0$ for every $J$ ($\overline{\psi}_{n_J}$ injective); as the identification $\widehat{M} \xrightarrow{\sim} \varprojlim_J M/\mathcal{M}_{n_J}$, $x \mapsto \left(\overline{v}_{n_J}\right)_J$, is an isomorphism, $x = 0$. So $\hat{\psi}$ is injective. ∎

**Proposition 3.14.28 (Universal Property of Completion)**
*With completion $\widehat{M}$, $\iota$, $\lambda_n$ (3.14.17), suppose moreover $(M', \mathcal{M}'_\bullet)$ is* complete *(3.14.20), with $\iota_{M'} : M' \xrightarrow{\sim} \widehat{M}'$ its own completion, and let $\psi : M \to M'$ be a continuous $\rho$-homomorphism (3.14.7). This is (3.14.27), composed with $\iota_{M'}^{-1}$: there is a unique continuous $\rho$-homomorphism $\hat{\psi}$ (for the filtration $\ker \lambda_\bullet$ of $\widehat{M}$) making*

$$\begin{array}{ccc} M & \xrightarrow{\psi} & M' \\ {\scriptstyle\iota}\downarrow & \nearrow_{\hat{\psi}} & \\ \widehat{M} & & \end{array}$$

*commute.*

*Moreover (**surjectivity criterion**), if $\psi$ is taut (3.14.11) then $\hat{\psi}$ is surjective.*

*Moreover (**injectivity criterion**), if $\psi(\mathcal{M}_n) \subseteq \mathcal{M}'_n$ for every $n \geq 0$ with $\overline{\psi}_{n_J} : M/\mathcal{M}_{n_J} \to M'/\mathcal{M}'_{n_J}$ injective for every $J \geq 0$, for some strictly increasing sequence $n_0 < n_1 < n_2 < \cdots$, then $\hat{\psi}$ is injective.*

*(Taking $M := A$, $M' := A'$ as filtered $A$-, $A'$-modules over themselves and $\psi := \rho$ recovers the ring-level statement exactly.)*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

By (3.14.27), there is a unique continuous $\rho$-homomorphism $\Psi : \widehat{M} \to \widehat{M}'$ with $\Psi \circ \iota = \iota_{M'} \circ \psi$; set $\hat{\psi} := \iota_{M'}^{-1} \circ \Psi$, continuous as a composite, with $\hat{\psi} \circ \iota = \iota_{M'}^{-1} \circ \Psi \circ \iota = \iota_{M'}^{-1} \circ \iota_{M'} \circ \psi = \psi$: the triangle commutes.

**Uniqueness.** If $\hat{\psi}_1, \hat{\psi}_2 : \widehat{M} \to M'$ are both continuous $\rho$-homomorphisms with $\hat{\psi}_i \circ \iota = \psi$, then $\iota_{M'} \circ \hat{\psi}_i : \widehat{M} \to \widehat{M}'$ are both continuous with $\left(\iota_{M'} \circ \hat{\psi}_i\right) \circ \iota = \iota_{M'} \circ \psi$; by uniqueness in (3.14.27), $\iota_{M'} \circ \hat{\psi}_1 = \iota_{M'} \circ \hat{\psi}_2$, and as $\iota_{M'}$ is injective, $\hat{\psi}_1 = \hat{\psi}_2$.

**Surjectivity criterion.** If $\psi$ is taut, (3.14.27)'s own surjectivity criterion gives $\Psi$ surjective, hence $\hat{\psi} = \iota_{M'}^{-1} \circ \Psi$ surjective ($\iota_{M'}^{-1}$ an isomorphism).

**Injectivity criterion.** Under the stated hypotheses, (3.14.27)'s own injectivity criterion gives $\Psi$ injective, hence $\hat{\psi} = \iota_{M'}^{-1} \circ \Psi$ injective. ∎

**Remark 3.14.29**
*Density (3.14.8) is exactly density in the usual topological sense for the metric $d(x, y) := 2^{-\sup\{n : x - y \in \mathfrak{b}_n\}}$ on $B$ (with $\sup \varnothing := -1$, so $d(x, y) = 0$ iff $x = y$): a ball of radius $2^{-n}$ about $b \in B$ is exactly $b + \mathfrak{b}_{n+1}$, so $\phi(A)$ meets every such ball iff $B = \phi(A) + \mathfrak{b}_{n+1}$ for every $n$, that is $\phi$ is dense.*

**Proposition 3.14.30 (Completion Matches the Filtration, $\mathfrak{a}$-adic Case)**
*Let $A$ be a ring, $\mathfrak{a} \triangleleft A$ an ideal, and $\hat{A}_\mathfrak{a}$, $\iota_\mathfrak{a}$, with projections $\lambda_n : \hat{A}_\mathfrak{a} \to A/\mathfrak{a}^{n+1}$, as in (3.14.19). This is the case $\mathfrak{a}_n := \mathfrak{a}^{n+1}$ of (3.14.25): for every $n \geq 0$, $\mathfrak{a}^{n+1}\hat{A}_\mathfrak{a} \subseteq \ker \lambda_n \subseteq \ker \lambda_{n-1}$ (the latter for $n \geq 1$), there is a unique isomorphism $\nu_n : A/\mathfrak{a}^{n+1} \xrightarrow{\sim} \hat{A}_\mathfrak{a}/\ker \lambda_n$ making the analogous square commute, with inverse $\mu_n$ satisfying $\mu_n \circ \pi_n = \lambda_n$, and $(\nu_n)_{n \geq 0}$ is natural in $n$ with respect to the reductions $A/\mathfrak{a}^{n+1} \to A/\mathfrak{a}^n$ and $\hat{A}_\mathfrak{a}/\ker \lambda_n \to \hat{A}_\mathfrak{a}/\ker \lambda_{n-1}$.*

**Corollary 3.14.31 (Adic Completion via any Exponent)**
*In the setting of (3.14.19), for every integer $e \geq 1$, there is a unique continuous ring homomorphism*

$$\theta_{\mathfrak{a},e} : \hat{A} \xrightarrow{\sim} \varprojlim_N A/\mathfrak{a}^{Ne}$$

*compatible with projections, given explicitly by*

$$\theta_{\mathfrak{a},e}\left((\overline{a}_n)_{n \geq 0}\right) = (\overline{a}_{Ne-1})_{N \geq 1}.$$

*which is furthermore a bicontinuous isomorphism.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

This is (3.14.18) (for the $\mathfrak{a}$-adic filtration $\mathfrak{a}_n := \mathfrak{a}^{n+1}$ (3.14.4)), applied to the sequence $n_N := Ne - 1$ ($N \geq 1$, so $\mathfrak{a}_{n_N} = \mathfrak{a}^{Ne}$ exactly). ∎

© David Rufino CC BY-NC-ND 4.0

**Remark 3.14.32**
*If $A$ is a $k$-algebra, for some ring $k$, then each $A/\mathfrak{a}^{n+1}$ is again a $k$-algebra, and the tower of (3.14.19) is a tower in* $\mathbf{Alg}_k$ *(3.3.154) so $\hat{A}_{\mathfrak{a}}$ is itself a $k$-algebra, and $\iota_{\mathfrak{a}}$ (running the same construction in* $\mathbf{Alg}_k$ *rather than* **Ring***) is a $k$-algebra homomorphism. Likewise, in (3.14.31), each $A/\mathfrak{a}^{Ne}$ is a $k$-algebra and $\theta_{\mathfrak{a},e}$ is a $k$-algebra isomorphism.*

## 3.14.1 Hensel's Lemma

**Lemma 3.14.33 (Convergent Sums in a Complete Ring)**
*Let $A$ be a ring and $\mathfrak{a} \triangleleft A$ an ideal such that $A$ is $\mathfrak{a}$-adically complete (3.14.21). Let $(y_i)_{i\geq 0}$ be a sequence in $A$ with $y_i \in \mathfrak{a}^i$ for every $i \geq 0$. Then there is a unique $s \in A$ – written $s =: \sum_{i\geq 0} y_i$ – with*

$$s \equiv \sum_{i=0}^{N} y_i \mod \mathfrak{a}^{N+1} \qquad \text{for every } N \geq 0.$$

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Set $s_N := \sum_{i=0}^{N} y_i \in A$ for $N \geq 0$. As $s_N - s_{N-1} = y_N \in \mathfrak{a}^N$ for $N \geq 1$, $(\overline{s}_N)_{N\geq 0}$, $\overline{s}_N := s_N \bmod \mathfrak{a}^{N+1}$, is compatible under reduction; by (3.14.21) item b) ($A$ being $\mathfrak{a}$-adically complete), there is $s \in A$ with $s \equiv s_N \bmod \mathfrak{a}^{N+1}$ for every $N \geq 0$: existence. If $s, s'$ both satisfy this, $s - s' \in \bigcap_N \mathfrak{a}^{N+1} = 0$ (separated): uniqueness. ∎

**Lemma 3.14.34 (Units of a Complete Ring)**
*Let $A$ be a ring and $\mathfrak{a} \triangleleft A$ an ideal such that $A$ is $\mathfrak{a}$-adically complete (3.14.21). If $u \in A^\star$ and $x \in \mathfrak{a}$, then $u + x \in A^\star$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

As $u + x = u(1 + u^{-1}x)$ and $u^{-1}x \in \mathfrak{a}$ ($\mathfrak{a}$ an ideal), it suffices to treat $u = 1$: show $1 + y \in A^\star$ for $y \in \mathfrak{a}$.

As $(-y)^i \in \mathfrak{a}^i$ for every $i \geq 0$, (3.14.33) gives $s := \sum_{i\geq 0} (-y)^i \in A$ with $s \equiv s_N := \sum_{i=0}^{N} (-y)^i \bmod \mathfrak{a}^{N+1}$ for every $N \geq 0$.

For every $N \geq 0$: $(1 + y)s_N = \sum_{i=0}^{N} (-y)^i + \sum_{i=0}^{N} y(-y)^i = \sum_{i=0}^{N} (-y)^i - \sum_{i=1}^{N+1} (-y)^i = 1 - (-y)^{N+1}$ (telescoping), and likewise $s_N(1 + y) = 1 - (-y)^{N+1}$ ($y$ commuting with its own powers); so

$$(1 + y)s - 1 = (1 + y)(s - s_N) + ((1 + y)s_N - 1) = (1 + y)(s - s_N) - (-y)^{N+1} \in \mathfrak{a}^{N+1}$$

($\mathfrak{a}^{N+1}$ an ideal, $s - s_N \in \mathfrak{a}^{N+1}$, $(-y)^{N+1} \in \mathfrak{a}^{N+1}$), and likewise $s(1 + y) - 1 \in \mathfrak{a}^{N+1}$. As $N \geq 0$ was arbitrary, $(1 + y)s - 1, s(1 + y) - 1 \in \bigcap_N \mathfrak{a}^{N+1} = 0$ (separated): $1 + y \in A^\star$, with inverse $s$. ∎

**Lemma 3.14.35 (Hensel's Lemma)**
*Let $A$ be a ring and $\mathfrak{a} \triangleleft A$ an ideal such that $A$ is $\mathfrak{a}$-adically complete (3.14.21). Let $f \in A[X]$, with formal derivative $f' \in A[X]$ (3.8.9) (writing $f(x)$, $f'(x)$ for their evaluations at $x \in A$ (3.8.5)). If $a_0 \in A$ has $f(a_0) \in \mathfrak{a}$ and $f'(a_0) \in A^\star$, then there is $a \in A$ with $f(a) = 0$ and $a \equiv a_0 \bmod \mathfrak{a}$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

**Taylor expansion.** By (3.8.10), $f(a + h) - f(a) - f'(a)h \in (h^2)$ for every $a, h \in A$. In particular (taking $a := y$, $h := x - y$ for $x, y \in A$), $f(x) - f(y) \in (x - y)$.

**Construction.** Define inductively $a_{n+1} := a_n - f'(a_n)^{-1} f(a_n)$ for $n \geq 0$ (well-defined once $f'(a_n) \in A^\star$, shown below).

**Claim: $f(a_n) \in \mathfrak{a}^{2^n}$, $f'(a_n) \in A^\star$, and $a_{n+1} \equiv a_n \bmod \mathfrak{a}^{2^n}$, for every $n \geq 0$.** Base case $n = 0$: given. Inductively, suppose $f(a_n) \in \mathfrak{a}^{2^n} =: \mathfrak{a}^m$ and $f'(a_n) \in A^\star$; write $u_n := f'(a_n)^{-1}$, $h_n := -u_n f(a_n) \in \mathfrak{a}^m$ ($\mathfrak{a}^m$ an ideal), so $a_{n+1} = a_n + h_n \equiv a_n \bmod \mathfrak{a}^m$. By the Taylor expansion,

$$f(a_{n+1}) \equiv f(a_n) + f'(a_n)h_n = f(a_n) - f'(a_n)u_n f(a_n) = f(a_n)(1 - f'(a_n)u_n) = 0 \mod (h_n^2)$$

(as $f'(a_n)u_n = 1$ exactly), so $f(a_{n+1}) \in (h_n^2) \subseteq \mathfrak{a}^{2m} = \mathfrak{a}^{2^{n+1}}$ ($h_n \in \mathfrak{a}^m$, $\mathfrak{a}^m$ an ideal). As $a_{n+1} \equiv a_n \bmod \mathfrak{a}^m \subseteq \mathfrak{a}$ (as $m = 2^n \geq 1$), and (telescoping) $a_n \equiv a_0 \bmod \mathfrak{a}$ for every $n$, the Taylor expansion's consequence $f'(a_{n+1}) - f'(a_0) \in (a_{n+1} - a_0) \subseteq \mathfrak{a}$ gives, by (3.14.34) ($u := f'(a_0) \in A^\star$, $x := f'(a_{n+1}) - f'(a_0) \in \mathfrak{a}$), $f'(a_{n+1}) = u + x \in A^\star$. This completes the induction.

**Existence of $a$.** The sequence $n_N := 2^N - 1$ is strictly increasing, and the Claim gives $a_N \equiv a_{N-1} \bmod \mathfrak{a}^{2^{N-1}} = \mathfrak{a}^{n_{N-1}+1}$ for every $N \geq 1$, so $(\overline{a}_N)_{N\geq 0}$, $\overline{a}_N := a_N \bmod \mathfrak{a}^{n_N+1}$, is compatible under reduction; by (3.14.21) item c), there is $a \in A$ with $a \equiv a_N \bmod \mathfrak{a}^{2^N}$ for every $N \geq 0$.

© David Rufino CC BY-NC-ND 4.0

**$a$ is the desired root.** For every $N \geq 0$: $a - a_N \in \mathfrak{a}^{2^N}$, so, by the Taylor expansion's consequence, $f(a) - f(a_N) \in (a - a_N) \subseteq \mathfrak{a}^{2^N}$; as also $f(a_N) \in \mathfrak{a}^{2^N}$ (the Claim), $f(a) \in \mathfrak{a}^{2^N}$, for every $N \geq 0$. So $f(a) \in \bigcap_N \mathfrak{a}^{2^N} = \bigcap_n \mathfrak{a}^n = 0$ (separated), giving $f(a) = 0$; and, at $N = 0$, $a \equiv a_0 \bmod \mathfrak{a}^{2^0} = \mathfrak{a}$. ■

**Corollary 3.14.36 (Idempotent Lifting)**
*Let $A$ be a ring and $\mathfrak{a} \triangleleft A$ an ideal such that $A$ is $\mathfrak{a}$-adically complete (3.14.21). Then every idempotent $\overline{e} \in A/\mathfrak{a}$ lifts to an idempotent $e \in A$, that is $e^2 = e$ and $e \equiv \overline{e} \bmod \mathfrak{a}$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Pick $a_0 \in A$ representing $\overline{e}$, so $f(a_0) := a_0^2 - a_0 \in \mathfrak{a}$ for $f(X) := X^2 - X \in A[X]$, with $f'(X) = 2X - 1$ (3.8.9). As $\overline{e}^2 = \overline{e}$, $(2a_0 - 1)^2 = 4a_0^2 - 4a_0 + 1 = 4(a_0^2 - a_0) + 1 \equiv 1 \bmod \mathfrak{a}$ (writing $4(a_0^2 - a_0) = 4f(a_0) \in \mathfrak{a}$), so $2a_0 - 1 \in A^\star$ by (3.14.34) (applied with $u := 1$). So (3.14.35) gives $e \in A$ with $e^2 - e = 0$ and $e \equiv a_0 \equiv \overline{e} \bmod \mathfrak{a}$: the desired lift. ■

© David Rufino CC BY-NC-ND 4.0

## 3.15 Chain Conditions

**Definition 3.15.1 (Noetherian / Artinian / Finite Modules)**
*We say an $A$-module $M$ is **Noetherian** if it satisfies the **ascending chain condition**, namely any ascending chain of submodules*

$$M_0 \subseteq M_1 \subseteq \cdots \subseteq M$$

*eventually stabilizes, i.e $M_n = M_{n+1}$ for all $n \geq N$.*

*Similarly we say an $A$-module $M$ is **Artinian** if it satisfies the **descending chain condition**, namely any descending chain of submodules*

$$M \supseteq M_0 \supseteq M_1 \supseteq \cdots$$

*eventually stabilizes.*

**Definition 3.15.2 (Noetherian / Artinian Ring)**
*We say a ring $A$ is **Noetherian** (resp. Artinian) if it is Noetherian (resp. Artinian) as an $A$-module.*

The following is useful

**Proposition 3.15.3 (Noetherian criterion)**
*Let $M$ be an $A$-module. The following are equivalent*

*a) $M$ is Noetherian*
*b) Every submodule $N \subseteq M$ is finitely-generated*
*c) Every set of submodules has a maximal element*

**Proposition 3.15.4**
*Let $M$ be an $A$-module and $A$ a Noetherian ring. Then $M$ is Noetherian if and only if it is finitely generated as an $A$-module.*

**Proposition 3.15.5 (Restriction of Scalars preserves finiteness)**
*Let $\phi : A \to B$ be a finite $A$-algebra and $M$ a (3.3.89) $B$-module. Then $[M]_\phi$ is a (3.3.89) $A$-module.*

*Proof.* We suppose that $M$ is generated by $m_1, ..., m_n$, and $B$ is generated by $b_1, ..., b_m$. Then we claim that the elements $b_i m_j$ generate $[M]_\phi$. ∎

**Proposition 3.15.6**
*Let $A$ be a Noetherian ring and $\mathfrak{a} \triangleleft A$ an ideal. Then $A/\mathfrak{a}$ is Noetherian.*

*Proof.* Consider an increasing sequence of ideals $\mathfrak{a}_i \triangleleft A/\mathfrak{a}$. Then by (3.3.56) this corresponds to an increasing sequence of ideals $\mathfrak{a}_i' \triangleleft A$ containing $\mathfrak{a}$. As $A$ is Noetherian, this sequence eventually stabilizes. Again by (3.3.56) the original sequence stabilizes. ∎

**Proposition 3.15.7 (Localisation preserves Noetherian)**
*Let $A$ be a Noetherian ring and $S \subset A$ a multiplicative set. Then $S^{-1}A$ is Noetherian.*

*Proof.* Let $\mathfrak{b}_0 \subseteq \mathfrak{b}_1 \subseteq \cdots$ be an ascending chain of ideals of $S^{-1}A$. Contracting gives an ascending chain of ideals $\mathfrak{b}_0^c \subseteq \mathfrak{b}_1^c \subseteq \cdots$ of $A$, which stabilizes as $A$ is Noetherian. As every ideal of $S^{-1}A$ is extended (3.6.18), $\mathfrak{b}_i = \mathfrak{b}_i^{ce}$ for each $i$, so the original chain stabilizes too. ∎

**Proposition 3.15.8**
*Let $A$ be a Noetherian ring then every finitely-generated $A$-algebra is Noetherian.*

*In particular $A[X_1, ..., X_n]$ is Noetherian.*

*Proof.* By (3.15.6) it's enough to show that $A[X_1, ..., X_n]$ is Noetherian. By induction and (3.11.4) it's enough to consider the case $n = 1$. Let $\mathfrak{a} \triangleleft A[X]$ then by (3.15.3) it's enough to show that $\mathfrak{a}$ is finitely-generated.

Define

$$\tilde{\mathfrak{a}}_i := \{\ell(f) \mid f \in \mathfrak{a} \text{ s.t. } \deg(f) = i\}$$

Then clearly $\tilde{\mathfrak{a}}_i \triangleleft A$ is an ideal. This is an increasing sequence of ideals, and so it stabilizes for $i \geq d$ for some $d > 0$. Furthermore each ideal is finitely generated

$$\tilde{\mathfrak{a}}_i := \left(c_{i1}, ..., c_{in(i)}\right)$$

© David Rufino CC BY-NC-ND 4.0

where $c_{ij} = \ell(f_{ij})$ for polynomials $f_{ij} \in \mathfrak{a}$ of degree $i$. We claim that

$$\mathfrak{a} = (f_{ij})_{i \leq d; j \leq n(i)}$$

Denote the right hand side by $\mathfrak{b}$, then clearly $\mathfrak{b} \subseteq \mathfrak{a}$. We show by induction on $m = \deg(f)$ that $f \in \mathfrak{a} \implies f \in \mathfrak{b}$. Let $m' \coloneqq \min(m, d)$. Then by assumption $\ell(f) \in \tilde{\mathfrak{a}}_m = \tilde{\mathfrak{a}}_{m'}$ so there exists $\lambda_j \in A$ such that

$$\ell(f) = \sum_{j=1}^{n(m')} c_{m'j} \lambda_j$$

Consider the decomposition

$$f = \left( f - \sum_{j=1}^{n(m')} \lambda_j f_{m'j} X^{m-m'} \right) + \sum_{j=1}^{n(m')} \lambda_j f_{m'j} X^{m-m'}$$

The first term has strictly smaller degree than $f$, so by the inductive hypothesis lies in $\mathfrak{b}$. Therefore $f \in \mathfrak{b}$ as required.■

© David Rufino CC BY-NC-ND 4.0

## 3.16 Principal Ideal Domains

**Definition 3.16.1 (Principal Ideal Domain)**
*An integral domain $A$ is a **principal ideal domain** (or **PID**) if every ideal $\mathfrak{a}$ is principal.*

**Proposition 3.16.2**
*A PID is Noetherian.*

*Proof.* Suppose we have an ascending chain of ideals

$$\mathfrak{a}_1 \subseteq \mathfrak{a}_2 \subseteq \cdots \subseteq \mathfrak{a}_n \subseteq \cdots$$

Clearly the union is again an ideal, which is also principal of the form $(a)$. We must have $a \in \mathfrak{a}_n$ for some $n$, whence it terminates after $n$. ∎

**Proposition 3.16.3 (Integers form a PID)**
*$\mathbb{Z}$ is a PID.*

*Proof.* This follows from the well-ordering principle. Let $\mathfrak{a}$ be an ideal with minimal positive element $d$. We claim $\mathfrak{a} = (d)$. By the division algorithm (or apply well-ordering principle to the coset $x + (d)$), for every $x \in \mathfrak{a}$ there is $0 \leq r < d$ and $q \in \mathbb{Z}$ such that

$$x = qd + r.$$

Clearly $r \in \mathfrak{a}$, whence by minimality $r = 0$ as required. ∎

**Proposition 3.16.4**
*Let $k$ be a field then the polynomial ring $k[X]$ is a PID*

*Proof.* Let $\mathfrak{a} \triangleleft k[X]$ be an ideal. Choose $f \in \mathfrak{a}$ to have minimal degree, then we claim $\mathfrak{a} = (f)$. For $g \in \mathfrak{a}$ we have by (3.8.13) $g = qf + r$ for $\deg(r) < \deg(f)$. Clearly $r \in \mathfrak{a}$, so by minimality $r = 0$ and the result follows. ∎

**Lemma 3.16.5 (Co-prime elements in a PID)**
*Let $A$ be a PID, then $x, y$ are coprime if and only if they have no non-invertible common divisors.*

*Proof.* First suppose $(x, y) = A$, then $ax + by = 1$ and any common divisor $d$ must divide 1 and therefore be invertible.

Conversely suppose $(x, y) \neq (1)$, since $A$ is a PID it must equal $(d)$ for some non-invertible $d$ which is then a common divisor. ∎

**Proposition 3.16.6**
*Let $A$ be a principal ideal domain, $M$ a free $A$-module with rank $m$ and $N \subset M$ an $A$-submodule. Then $N$ is free with rank $n \leq m$.*

*Proof.* Recall (...) that the rank of a finite free $A$-module is well-defined.

We proceed by induction on $m$. Suppose that $m = 1$ then there is an isomorphism of $A$-modules $A \xrightarrow{\sim} M$. Therefore under this isomorphism $N$ corresponds to an ideal, say $(a)$. If $a = 0$ then $N = 0$ is by definition free of rank 0. If $a \neq 0$ then we may consider the map of $A$-modules

$$\begin{aligned} A &\to (a) \\ x &\to ax \end{aligned}$$

it is by definition surjective, and injective by cancellation and so an isomorphism. This shows that $N$ is free of rank 1.

Suppose then $x_1, ..., x_m$ is a basis for $M$, and let $M'$ be the free submodule generated by $x_1, ..., x_{m-1}$, and $M''$ the rank-1 free module generated by $x_m$. Then

$$M \coloneqq M' \oplus M''$$

and we may show that $N$ is the internal direct sum

$$N = (N \cap M') \oplus (N \cap M'').$$

Then by the induction hypothesis we find $N$ is free of rank at most $m - 1 + 1 = m$. ∎

**Proposition 3.16.7**
*Let $A$ be a principal ideal domain and $M$ a finitely generated, torsion-free $A$-module. Then $M$ is a finite free $A$-module.*

© David Rufino CC BY-NC-ND 4.0

*Proof.* Let $\{m_1, ..., m_n\}$ be a generating set for $M$, and $\{v_1, ..., v_r\}$ an arbitrary linearly independent subset of $M$. Then by definition we have

$$v_i = \sum_{j=1}^{n} a_{ij} m_j \quad a_{ij} \in A$$

Consider the elements of $A^n$ given by

$$w_i := (a_{i1}, ..., a_{in})$$

Then we can see that $\{w_1, ..., w_r\}$ is an $A$-linearly independent subset of $A^n$, and therefore a $K$-linearly independent subset of $K^n$ for $K = \mathrm{Frac}(A)$. By (...) we see that $r \leq n$.

Therefore we may choose $\{v_1, ..., v_r\}$ to be a maximal linearly independent subset and define $M' := \langle v_1, ..., v_r \rangle$. For each generator $m_i$, by maximality of the subset there must exist an $a_i \neq 0$ such that

$$a_i m_i \in M'$$

Define $a := a_1 ... a_n$ then this satisfies $aM \subseteq M'$, since the $m_i$ generate $M$. As $A$ is an integral domain we have a series of injections

$$M \hookrightarrow aM \subseteq M' \xrightarrow{\sim} A^r$$

Therefore we are done by (3.16.6). ∎

© David Rufino CC BY-NC-ND 4.0

## 3.17 Factorisation

For this section we assume $A$ is a commutative integral domain.

**Definition 3.17.1 (Associates)**
*We say two non-zero elements $x$ and $y$ are **associates** if $x = uy$ for some $u \in A^\star$. We write $x \sim y$.*

*Note this is an equivalence relation on $A \setminus \{0\}$.*

**Lemma 3.17.2**
*Let $A$ be a ring and $x, y \in A$ non-zero elements then the following are equivalent*

- $x \mid y$
- $(y) \subseteq (x)$
- $y \in (x)$

**Lemma 3.17.3**
*Let $A$ be a ring and $x, y \in A$ non-zero elements then the following are equivalent*

- $x \mid y$ *and* $y \mid x$
- $(x) = (y)$

*If $A$ is an integral domain this is equivalent to $x \sim y$.*

**Definition 3.17.4 (Irreducible element)**
*We say $0 \neq x$ is **irreducible** if it is not invertible and $x = ab \implies a$ a unit or $b$ a unit.*

*Equivalently if $y \mid x$ implies either $y$ is a unit or $y \sim x$.*

**Definition 3.17.5 (Prime element)**
*We say $0 \neq p$ is prime if $p \mid ab \implies p \mid a$ or $p \mid b$.*

**Example 3.17.6**
*The units of $\mathbb{Z}$ are $\{-1, 1\}$ so each equivalence class is of the form $\{n, -n\}$.*

**Example 3.17.7**
*A number $p \in \mathbb{Z}$ is prime in the traditional sense exactly when it is irreducible. It is of course also prime in the ring-theoretic sense but this requires proof (see (2.2.13)).*

The concept of associates is important to unique factorization, because we may only hope to have unique factorization upto multiplication by a unit.

**Lemma 3.17.8**
*If $x \sim y$ are associates then $x$ is irreducible iff $y$ is*

*Proof.* Suppose $x \sim y$ and $x$ irreducible. If $y = ab$ then $x = abu \implies a$ a unit or $bu$ a unit $\implies b$ a unit. Therefore $y$ is irreducible as required. ∎

**Lemma 3.17.9 (Criterion for primality)**
*Suppose $0 \neq p \in \mathbb{Z}$. Then $p$ is prime if and only if $(p)$ is a prime ideal*

*Proof.* Note $x \mid y \iff y \in (x)$. So in particular if $p$ is prime then $xy \in (p) \implies p \mid xy \implies p \mid x$ or $p \mid y \implies x \in (p)$ or $y \in (p)$, whence $(p)$ is prime.

Conversely if $(p)$ is prime, then $p \mid xy \implies xy \in (p) \implies x \in (p)$ or $y \in p \implies p \mid x$ or $p \mid y$, so that $p$ is prime. ∎

**Lemma 3.17.10 (Criterion for irreducibility)**
*Let $A$ be an integral domain. Then $f$ is irreducible if and only if $(f)$ is maximal amongst proper principal ideals.*

*Proof.* Suppose $f$ is irreducible and $(f) \subseteq (g)$. Then $f = ag$ with either $a$ a unit or $g$ a unit. If $a$ is a unit then $(f) = (g)$, and if $g$ is a unit $(g) = A$. So the result follows.

Conversely suppose $f = ab$, then $f \in (a) \implies (f) \subseteq (a)$. Then by hypothesis either $(a) = (f)$ or $(a) = A$. In the second case $a$ is a unit. In the first case then $f \mid a \implies bf \mid f \implies b \mid 1$ whence $b$ is a unit. ∎

**Proposition 3.17.11 (Primes are Irreducible)**
*Let $A$ be an integral domain then $p$ prime $\implies p$ irreducible*

© David Rufino CC BY-NC-ND 4.0

*Proof.* Suppose $b \mid p$ then $p = ab$ and $a \mid p$. By hypothesis $p \mid a$ or $p \mid b$. If $p \mid a$ (resp. $b$) then by (3.17.3) $p \sim a$ (resp. $b$) as required. ∎

**Definition 3.17.12 (Unique Factorisation Domain (UFD) or Factorial Ring)**
*We say an integral domain $A$ is factorial (or a UFD) if every element $0 \neq a$ may be represented as*

$$a = u \prod_{i=1}^{n} p_i$$

*for $u$ a unit and $p_i$ irreducible, and moreover this is unique in the sense that given another factorization*

$$a = u' \prod_{i=1}^{m} p_i'$$

*we have $n = m$ and $p_i \sim p'_{\psi(i)}$, for $\psi$ a permutation on $\{1, ..., n\}$.*

Furthermore it may be convenient in applications to count the multiplicities

**Proposition 3.17.13 (Factorisation with multiplicities)**
*Let $A$ be a UFD, then for every element $0 \neq a \in A$ there is a factorization of the form*

$$a = u \prod_{i=1}^{n} p_i^{r_i}$$

*where $r_i > 0$ and none of the $p_i$ are associate to each other. Furthermore this is essentially unique in the sense that given another such factorization we have $n = n'$, $r_i = r'_{\sigma(i)}$ and $p_i \sim p'_{\sigma(i)}$ for some permutation $\sigma \in S_n$.*

*Proof.* Given a factorization into irreducible elements

$$a = u \prod_{i=1}^{n} p_i$$

Consider a representative set of irreducibles $q_1, ..., q_m$ (under the equivalence relation $x \sim y$). Then we have $p_i = q_{\pi(i)} u_i$ for some units $u_i$ and mapping $\pi : \{1, ..., n\} \to \{1, ..., m\}$. Let $r_j = \#\pi^{-1}(j)$. Then we have that the set of irreducibles $\{p_1, ..., p_n\}$ is the *disjoint* union of the set of equivalence classes with representatives $q_j$. Therefore

$$a = u \prod_{j=1}^{m} \prod_{p \sim q_j} p = u \prod_{j=1}^{m} \prod_{i:\pi(i)=j} u_i q_j = \left( u \prod_{j=1}^{m} \prod_{i:\pi(i)=j} u_i \right) \prod_{j=1}^{m} q_j^{r_j}$$

as required. Suppose we have two factorizations

$$u \prod_{i=1}^{n} p_i^{r_i} = u' \prod_{i=1}^{m} (p_i')^{r_i'}$$

Let $I$ be the indexing set of $p_i$ and $J$ the set of $p_j'$. By unique factorization there must be mappings $\sigma : I \to J$ such that $p_i \sim p'_{\sigma(i)}$, and $\tau : J \to I$ such that $p_j' \sim p_{\tau(j)}$. Which means that $p_i \sim p_{\tau(\sigma(i))}$ and $p_j' \sim p'_{\sigma(\tau(j))}$. Since none are associate to each other we see that $\tau$ and $\sigma$ are mutual inverses, whence $m = n$ and we may regard $\sigma \in S_n$. In the unique factorization $p_i$ appears $r_i$ times and $p'_{\sigma(i)}$ appears $r'_{\sigma(i)}$ times. Since $p_i$ is associate to $p'_{\sigma(i)}$ it is not associate to any $p_j'$ for $j \neq \sigma(i)$. Unique factorization shows that $r_i = r'_{\sigma(i)}$. ∎

**Definition 3.17.14**
*Let $A$ be a UFD and $x \in A$ a non-zero, non-unit such that*

$$x \sim \prod_{i=1}^{n} p_i^{r_i}$$

*is an (almost) unique factorization into irreducibles. Then for $p \in A$ an irreducible define*

$$v_p(x) := \begin{cases} r_i & p \sim p_i \\ 0 & \text{otherwise} \end{cases}$$

*If $x \in A$ is a unit then simply define $v_p(x) = 0$ for all $p$.*

**Lemma 3.17.15**
*Let $A$ be a UFD then the following are equivalent*

© David Rufino CC BY-NC-ND 4.0

*a)* $x \mid y$
*b)* $(y) \subseteq (x)$
*c)* $v_p(x) \leq v_p(y)$ *for all* $p$ *irreducible*

*In particular* $x \sim y \iff (x) = (y) \iff v_p(x) = v_p(y)$ *for all* $p$ *irreducible.*

**Definition 3.17.16 (Atomic Ring)**
*We say that* $A$ *is* ***atomic*** *if every element has a (not necessarily unique) decomposition into irreducible elements.*

**Definition 3.17.17 (Ascending Chain Condition for Principal Ideals (ACCP))**
*We say a ring* $A$ *satisfies* ***ACCP*** *if every ascending chain of principal ideals eventually stabilizes.*

*Note every Noetherian ring satisfies this condition.*

**Proposition 3.17.18**
*An integral domain* $A$ *satisfying ACCP is atomic.*

*In particular a Noetherian ring is atomic.*

*Proof.* Suppose a ring $A$ is not atomic, then choose any non-unit $x_1 \in A$. By repeated application (3.17.10) it's possible to construct a strictly ascending sequence of proper principal ideals

$$(x_1) \subsetneq (x_2) \subsetneq ... \subsetneq (x_n) \subsetneq ...$$

therefore $A$ does not satisfy ACCP. ■

**Remark 3.17.19**
*The converse is not in general true (...) but see (3.17.21) for a partial converse.*

We show a simple criterion for a ring to be a UFD.

**Definition 3.17.20**
*We say an integral domain* $A$ *is* ***AP*** *if* $p$ *irreducible* $\implies$ $p$ *prime.*

Roughly speaking, "atomic" ensures the existence of factorization and "AP" ensures the uniqueness.

**Proposition 3.17.21 (Atomic + AP $\iff$ UFD)**
*Let* $A$ *be an integral domain. The following are equivalent*
*a)* $A$ *is a UFD*
*b)* $A$ *is atomic and AP*
*c)* $A$ *satisfies ACCP and is AP*

*Proof.* *a)* $\implies$ *c)*. Suppose $A$ is a UFD. If $p$ is an irreducible element and $p \mid ab$, then by uniqueness it must appear in the irreducible factorization of either $a$ or $b$. Therefore $p$ is prime. If we have an ascending chain of principal ideals $(x_1) \subseteq (x_2)$... then by (3.17.15) we have $v_p(x_i)$ is a decreasing sequence for all irreducible $p$ occuring in the factorization of $x_1$. Furthermore $\max_p v_p(x_i)$ is a finite decreasing sequence. Choose $i = N$ such that $\max_p v(x_i)$ is minimal, then all these sequences must stabilise for $i \geq N$ and therefore by (3.17.15) the chain of principal ideals also stabilises.

*c)* $\implies$ *b)*. This is (3.17.18).

*b)* $\implies$ *a)*. We require to show that factorization into irreducibles is unique up to associates. Suppose

$$\prod_{i=1}^{n} p_i \sim \prod_{j=1}^{m} p'_j$$

By convention an empty product is 1 and by hypothesis all the elements are in fact prime. If $n = 0$, then since $p'_j$ is irreducible, it is not a unit and hence $m = 0$. Otherwise consider $p_1$, then $p_1 \mid$ RHS, so by definition of prime we must have $p_1 \mid p'_j$ for some $j$. Since $p'_j$ is irreducible and $p_1$ is not a unit, we have $p_1 \sim p'_j$. Since $A$ is integral we may cancel these two to obtain an equivalence of smaller degree and we may proceed by induction. ■

**Lemma 3.17.22 (PID is an AP-domain)**
*Let* $A$ *be a PID and* $a \in A$. *Then the following are equivalent*

*a)* $a$ *is prime*
*b)* $a$ *is irreducible*
*c)* $(a)$ *is maximal*

*Proof.* *a)* $\implies$ *b)* This holds for an arbitrary integral domain (3.17.11).

© David Rufino CC BY-NC-ND 4.0

$b) \iff c)$ as all ideals are principal this follows from (3.17.10)

$c) \implies a)$ By (3.3.60) $(a)$ is prime, whence $a$ is prime by (3.17.9) ∎

**Proposition 3.17.23**
*A PID is Noetherian UFD.*

*Furthermore ($f$ is irreducible $\iff$ $f$ is prime), and every prime ideal is maximal.*

*Proof.* $A$ is Noetherian by (3.16.2) and atomic by (3.17.18). And by (3.17.22) an irreducible element is prime. Therefore we are done by (3.17.21). ∎

If we take a suitable fixed set of irreducible elements we can obtain completely unique factorization

**Definition 3.17.24**
*Let $A$ be a ring we say $\mathcal{P}$ is a **representative set of irreducible elements** if*

- *No two elements $p, q \in \mathcal{P}$ are associate*
- *Every irreducible element $p \in A$ is associate to (precisely) one in $\mathcal{P}$*

**Example 3.17.25**
*For $\mathbb{Z}$ the* positive *primes are a canonical set of irreducible elements.*

**Proposition 3.17.26**
*Let $A$ be a UFD and $K = \operatorname{Frac}(A)$ then for every irreducible $p \in A$ there is a unique map*

$$v_p : K^\star \to \mathbb{Z}$$

*such that*

- $u \in A^\star \implies v_p(u) = 0$
- $v_p(xy) = v_p(x) + v_p(y)$
- $v_p(p) = 1$

*Furthermore let $\mathcal{P}$ be a set of irreducible representatives then we have a group isomorphism*

$$\begin{aligned} K^\star/A^\star &\xrightarrow{\sim} \bigoplus_{p\in\mathcal{P}} \mathbb{Z} \\ x &\longrightarrow \left(v_p(x)\right)_{p\in\mathcal{P}} \\ \prod_{p\in\mathcal{P}} p^{n_p} &\longleftarrow \left(n_p\right)_{p\in\mathcal{P}} \end{aligned}$$

*Finally $x \in A \iff v_p(x) \geq 0$ for all $p \in \mathcal{P}$.*

*Proof.* By (3.17.13) there is a well-defined map $v_p : A \setminus \{0\} \to \mathbb{Z}$ satisfying the given properties. We claim that

$$\begin{aligned} v_p : K^\star &\to \mathbb{Z} \\ xy^{-1} &\to v_p(x) - v_p(y) \end{aligned}$$

is well-defined. For suppose $xy^{-1} = wz^{-1}$ then $zx = wy$ whence $v_p(z) + v_p(x) = v(w) + v(y)$ and therefore $v_p(xy^{-1}) = v_p(wz^{-1})$.

It's clear the multiplicative property also holds and so $\phi$ is well-defined (since by definition $A^\star$ is in the kernel of $v_p$). Denote by $\phi, \psi$ the proposed maps. By definition of unique factorization $\phi$ and $\psi$ are mutual inverses when restricted as follows

$$(A \setminus \{0\})/A^\star \longleftrightarrow \bigoplus_{p\in\mathcal{P}} \mathbb{Z}_{\geq 0}$$

We also observe that $\phi(x^{-1}) = -\phi(x)$ and $\psi(-n) = \psi(n)^{-1}$, so then it's easy to demonstrate they are mutually inverse over the whole domain.

Suppose $v_p(xy^{-1}) \geq 0$ then $v_p(x) \geq v_p(y)$. If this holds for all $p \in \mathcal{P}$, then by (3.17.15) $y \mid x$ as elements of $A$ whence by definition $xy^{-1} \in A$ as required. ∎

**Proposition 3.17.27**
*Suppose $A$ is an integral domain satisfying ACCP then so is $A[X]$.*

© David Rufino CC BY-NC-ND 4.0

*Proof.* Suppose we have an ascending chain of principal ideals

$$(f_1) \subseteq (f_2) \subseteq ...(f_n)...$$

Without loss of generality the $f_i$ are non-zero. Then $f_{i+1} \mid f_i$ and by (...) $\deg(f_{i+1}) \leq \deg(f_i)$. Choose $N$ such that $\deg(f_i)$ is minimal, and define $a_i := \ell(f_i) \in A$. Then by (3.8.3) $a_{i+1} \mid a_i$ for $i \geq N$ as elements of $A$. Therefore we have an increasing sequence of principal ideals

$$(a_N) \subseteq (a_{N+1}) \subseteq ...$$

which by hypothesis stabilizes, that is $a_i \sim a_j$ for all $i, j \geq M$ for some $M \geq N$. For $i \geq M$ we have $ua_i = a_{i+1}$, consider $uf_i - f_{i+1} \in (f_i)$. This has degree strictly smaller than $N$, and therefore by minimality must be 0. In particular $f_i \sim f_{i+1}$ and $(f_i) = (f_{i+1})$. ∎

**Lemma 3.17.28**
*Suppose $p \in A$ is prime, then it is prime as an element of $A[X]$.*

**Lemma 3.17.29 (Nagata's Criterion)**
*Let $A$ be a ring and $S$ a multiplicative subset generated by prime elements and units. Let $f \in A$ be irreducible or a unit, then*

*a) $\frac{f}{1}$ is irreducible or a unit in $S^{-1}A$*
*b) $\frac{f}{1}$ prime or a unit in $S^{-1}A \implies f$ is a prime or a unit in $A$.*

*Furthermore if $S^{-1}A$ is AP then so is $A$.*

*Proof.* Note the condition on $S$ means every $a \in S$ satisfies $a \sim p_1...p_r$ for primes $p_i \in A$.

a) Suppose $\frac{f}{1} = \frac{g}{a}\frac{h}{b}$ for $f, g, h \in A$ and $a, b \in S$, then $abf = gh$. Further $ab \sim p_1...p_r$. Then $p_i \mid a$ or $p_i \mid b$, whence we can find $f \sim g'h'$ where $g = cg'$, $h = dh'$ and $c, d \in S$. As $f$ is irreducible (or a unit), then for example $g'$ is invertible, in which case $\frac{g}{a} = \frac{cg'}{a}$ is invertible. Therefore $\frac{f}{1}$ is either irreducible or a unit.
b) The case $f$ a unit is clear, so assume that $f$ is irreducible. Suppose $\frac{f}{1}$ is prime or a unit and $f \mid gh$. Then $\frac{f}{1} \mid \frac{g}{1}\frac{h}{1}$ and for example $\frac{f}{1} \mid \frac{g}{1}$. Therefore $ff' = gp_1...p_r$ for some $f' \in A$ and $p_1, ..., p_r \in A$ prime. If $p_i \mid f$ for some $i$ then by irreducibility we have $p_i \sim f$, and we see that $f$ is prime. Otherwise $p_i \mid f'$ for all $i$ and we find $f \mid g$. Therefore $f$ is prime as required.

The last statement follows immediately from the previous two results. ∎

## 3.17.1 Polynomials over a Field

In this section we consider the polynomial ring with coefficients in a field, $k[X]$.

**Proposition 3.17.30 (Associate polynomials)**
*The following are equivalent for $0 \neq f, g$*

- $f \sim g$
- $f = \lambda g$ *for* $\lambda \neq 0$
- $f \mid g$ *and* $g \mid f$

**Proposition 3.17.31**
*A polynomial $f \in k[X]$ is associate to precisely one monic polynomial $g$. If $f$ is irreducible so is $g$.*

*Proof.* **TODO** ∎

**Proposition 3.17.32 (Polynomial ring is a PID)**
*Let $k$ be a field, then $k[X]$ is a PID, and therefore a Noetherian UFD.*

*Proof.* Let $(0) \neq \mathfrak{a}$ be an ideal and let $f \in \mathfrak{a}$ be a polynomial of minimal degree. We may assume it is monic. Any $g \in \mathfrak{a}$ may be represented as $f = qg + r$ by the division algorithm. Clearly $r \in \mathfrak{a}$, therefore by minimality $r = 0$, whence $g \in (f)$. ∎

**Proposition 3.17.33 (Unique Factorisation of Polynomials)**
*For the ring $k[X]$ the set of irreducible monic polynomials constitutes a representative set (3.17.24). Therefore we have a unique factorization*

$$f = \ell(f) \prod_{p \text{ irreducible monic}} p^{v_p(f)}$$

© David Rufino CC BY-NC-ND 4.0

*such that*

$$v_p(fg) = v_p(f) + v_p(g)$$

*Proof.* (3.17.31) shows that the irreducible monic polynomials constitute a representative set. Therefore the result follows from (3.17.26). Let $u$ be the unit appearing in the factorization, it must be an element of $k$. Compare leading coefficients to see that $u = \ell(f)$. ∎

**Lemma 3.17.34 (Roots and Multiplicity)**
*For $f \in k[X]$ a non-constant polynomial and $\alpha \in k$ we have*

$$f(\alpha) = 0 \iff (X - \alpha) \mid f \iff v_{X-\alpha}(f) > 0$$

*In this case $r := v_{X-\alpha}(f)$ is the multiplicity of the root $\alpha$, and observe*

$$f(X) = \ell(f)(X - \alpha)^r g(X)$$

*with $g(\alpha) \neq 0$ (equivalently $v_{X-\alpha}(g) = 0$).*

*Proof.* The right to left implication is obvious. Conversely by the division algorithm we may write

$$f(X) = f(\alpha) + (X - \alpha)Q(X)$$

Then if $f(\alpha) = 0$ we clearly have $v_{X-\alpha}(f) > 0$. Finally we may construct

$$g(X) = \prod_{p \neq (X-\alpha)} p^{v_p(f)}$$

It's clear that for every $p$ appearing in the product $p(\alpha) \neq 0$ because otherwise we would have $(X - \alpha) \mid p$ and by irreducibility $(X - \alpha) = p$. Therefore $g(\alpha) \neq 0$ as required. ∎

**Corollary 3.17.35**
*Let $K/k$ be a field extension and $0 \neq f \in k[X]$. Then $f$ has at most $\deg(f)$ roots in $K$. In particular a polynomial of degree $< n$ with $n$ distinct roots in $K$ is zero.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Identify $f$ with its image in $K[X]$, which has the same degree. Let $\alpha_1, ..., \alpha_m$ be the distinct roots of $f$ in $K$ and $r_i := v_{X-\alpha_i}(f) \geq 1$ their multiplicities in $K[X]$ (3.17.34). Then $f = \ell(f) \prod_{i=1}^{m} (X - \alpha_i)^{r_i} g$ where $g$ is the product of the remaining irreducible factors of $f$ in $K[X]$, which is a constant when $f$ splits completely in $K$ (3.19.8). Therefore $\deg(f) = \sum_i r_i + \deg(g) \geq m$. ∎

**Lemma 3.17.36 (Lagrange Interpolation)**
*Let $k$ be a field, $y_1, ..., y_n \in k$ distinct and $h \in k[X]$ with $\deg(h) < n$. Then*

$$h(X) = \sum_{l=1}^{n} h(y_l) \prod_{k \neq l} \frac{X - y_k}{y_l - y_k}.$$

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

The difference of the two sides has degree $< n$ and vanishes at each $y_j$, as the $l$-th term of the sum vanishes at $y_j$ for $l \neq j$ and equals $h(y_j)$ at $y_j$ for $l = j$. It has $n$ roots so is zero by (3.17.35). ∎

**Proposition 3.17.37 (Criteria for Multiple Roots)**
*Let $f(X) \in k[X]$ be a polynomial, $r$ a positive integer and suppose that either $\mathrm{char}(k) = 0$ or $r < \mathrm{char}(k)$. Then $\alpha \in k$ is a root of $f$ with multiplicity $r$ precisely when*

$$f(\alpha) = f^{(1)}(\alpha) = ... = f^{(r-1)}(\alpha) = 0$$

*and $f^{(r)}(\alpha) \neq 0$.*

*Therefore the multiple roots are precisely the common roots of $f(X)$ and $f'(X)$ (irrespective of the characteristic).*

*Proof.* Note that by (3.17.34) and (3.8.9)

$$f^{(1)}(X) = (X - \alpha)^{r-1}[rg(X) + (X - \alpha)g'(X)]$$

with $g(\alpha) \neq 0$ and $r$ the multiplicity of the root. If $r = 1$, then $f^{(1)}(\alpha) = g(\alpha) \neq 0$ as required. If $r > 1$, then $f^{(1)}(X)$ has $\alpha$ as a root of multiplicity $r - 1$, so it follows by induction.

© David Rufino CC BY-NC-ND 4.0

The second statement is simply the case $r = 1$. ∎

**Definition 3.17.38 (Separable Polynomial)**
*A polynomial $f \in k[X]$ is **separable** if $f$ and $f'$ are co-maximal, that is $(f, f') = (1)$. Otherwise it is **inseparable**.*

**Proposition 3.17.39**
*A separable polynomial $f \in k[X]$ has no multiple roots (and $f$ and $f'$ have no common roots) in any extension field $K/k$.*

*Proof.* Since $(f, f') = (1)$ we have $af + bf' = 1$ for some $a, b \in k[X]$. Clearly $f$ and $f'$ can have no common roots, and therefore $f$ has no multiple roots by (3.17.37). ∎

**Proposition 3.17.40**
*Suppose $f, g \in k[X]$, $g$ is separable and $f \mid g$, then $f$ is separable.*

*Proof.* Suppose $f$ is not separable then by (3.16.5) $f$ and $f'$ have a common divisor $d$ such that $\deg(d) > 0$. Since $g = fh$, so $g' = f'h + fh'$. Therefore $d$ is also a non-trivial common divisor of $g$ and $g'$ contradicting (3.16.5). ∎

## 3.17.2 Polynomial Ring is a UFD

We prove the following result, first by Nagata's Criterion and again by Gauss' Lemma.

**Proposition 3.17.41**
*Suppose $A$ is a UFD, then so is $A[X]$.*

*Proof.* By (3.17.21) we need to show that $A[X]$ satisfies ACCP and is AP. The first follows from (3.17.27).

Let $S = A \setminus \{0\}$ the set of non-zero elements. Let $K = \mathrm{Frac}(A)$. If we regard $A[X]$ as a subring of $K[X]$ then we claim $S^{-1}(A[X]) = K[X]$; this follows by multiplying an element of $K[X]$ by the product of denominators of all the coefficients. Furthermore as $A$ is a UFD (and by (3.17.21)) $S$ is generated by prime elements and units. As $K$ is a field, $K[X]$ is a PID (3.17.32), so by (3.17.23) it is a UFD, and so in particular is AP. Therefore by (3.17.29) $A[X]$ is AP as required. ∎

For Gauss' Lemma we require to introduce some notation first.

**Definition 3.17.42 (Primitive polynomials)**
*Let $A$ be a UFD, $K := \mathrm{Frac}(A)$ and $f \in K[X]$ a polynomial given by*

$$f(X) = \sum_{i=0}^{n} a_i X^i$$

*Define the **content** of $f$ by*

$$c(f) = \prod_{p \in \mathcal{P}} p^{\min_i v_p(a_i)} \in K^\star$$

*where the product is taken over a representative set of primes for $A$. Changing the set of representatives only changes the value of $c(f)$ by multiplication of a unit.*

*We say that $f$ is **primitive** precisely when $c(f) = 1$*

**Lemma 3.17.43 (Gauss' Lemma I)**
*Let $A$ be a UFD and $f \in K[X]$ where $K = \mathrm{Frac}(A)$. Then the content $c(f)$ satisfies the following properties*

*a) $f \in A[X] \iff c(f) \in A$*
*b) $f$ is primitive $\iff \min_i v_p(a_i) = 0 \quad \forall p$ prime and in this case $f \in A[X]$*
*c) $c(\lambda f) = \lambda c(f)$ for $\lambda \in K^\star$ up to multiplication by a unit in $A$*
*d) $f/c(f) \in A[X]$ is primitive*
*e) $f, g$ primitive $\implies fg$ primitive*
*f) $c(fg) = c(f)c(g)$ for all $f, g \in K[X]$*

*Proof.* We prove each in turn

a) $f \in A[X] \iff a_i \in A \quad \forall i \overset{(3.17.26)}{\iff} v_p(a_i) \geq 0 \, \forall i \, \forall p \in \mathcal{P} \iff \min_i v_p(a_i) \geq 0 \, \forall p \in \mathcal{P} \overset{(3.17.26)}{\iff} c(f) \in A$
b) $c(f) = 1 \iff v_p(c(f)) = 0 \quad \forall p \iff \min_i v_p(a_i) = 0 \quad \forall p$. Further $v_p(a_i) \geq 0$ whence $f \in A[X]$ by (3.17.26).
c) Note $v_p(\lambda a_i) = v_p(\lambda) + v_p(a_i) \implies v_p(c(\lambda f)) = \min_i v_p(\lambda a_i) = v_p(\lambda) + \min_i v_p(a_i) = v_p(\lambda) + v_p(c(f))$. Whence by (3.17.26) we see $c(\lambda f)$ and $\lambda c(f)$ are equal up to multiplication by a unit in $A$.
d) Clear by $b$).

© David Rufino CC BY-NC-ND 4.0

e) Consider $p$ prime and the reduction $\bar{\cdot}: A[X] \to (A/(p))[X]$. Then by assumption $\overline{f}$ and $\overline{g}$ are non-zero. Furthermore $(A/(p))[X]$ is an integral domain so that $\overline{f \cdot g} = \overline{f} \cdot \overline{g} \neq 0$. Therefore $p$ does not divide all the coefficients of $f \cdot g$. As $p$ was arbitrary this shows that $f \cdot g$ is primitive.

f) We may reduce to $e$) by dividing by the content.

■

**Lemma 3.17.44 (Gauss' Lemma II)**
*Let $A$ be a UFD and $f \in A[X]$. Then the following are equivalent*

*a) $f$ is irreducible in $A[X]$*
*b) $f$ is an irreducible element of $A$ or ($f$ is primitive and irreducible in $K[X]$)*

*where $K = \mathrm{Frac}(A)$.*

*In particular if $f \in K[X]$ is irreducible then $f/c(f)$ is irreducible in $A[X]$.*

*Proof.* $b) \implies a)$. It's clear that an irreducible element of $A$ remains irreducible in $A[X]$. Suppose $0 \neq f \in A[X]$ is primitive and irreducible in $K[X]$. Then as $f \notin K[X]^\star$ we have $\deg(f) > 0$. Suppose $f = gh$ in $A[X]$ then by irreducibility $\deg(g) = 0$ or $\deg(h) = 0$. Further $1 = c(f) = c(g)c(h)$ by Gauss' Lemma, so one of $h$ and $g$ must lie in $A^\star = A[X]^\star$. Therefore $f$ is irreducible in $A[X]$ as required.

$a) \implies b)$. If $\deg(f) = 0$ then clearly $f$ is an irreducible element of $A$. So assume $\deg(f) > 0$. Observe $f = c(f) \cdot (f/c(f))$ so by irreducibility we require $c(f) = 1$ and therefore $f$ is primitive. Suppose $f = gh$ in $K[X]$ then $1 = c(g)c(h)$ and therefore

$$f = \frac{g}{c(g)} \frac{h}{c(h)}$$

is a decomposition in $A[X]$ which shows one of $g, h$ has degree 0. Therefore $f$ is irreducible in $K[X]$ as required. ■

**Proposition 3.17.45**
*Let $A$ be a UFD. Then so is $A[X]$.*

*Proof.* For $0 \neq f \in A[X]$ we may use irreducible factorisation in $K[X]$ to find

$$f = \lambda \pi_1 ... \pi_n$$

where $\lambda \in K$ and $\pi_i \in K[X]$ are irreducible. Replace $\pi_i \to \pi_i / c(\pi_i)$ then may assume that $\pi_i$ are irreducible in $A[X]$ by (3.17.44). Furthermore

$$c(f) = \lambda c(\pi_i) ... c(\pi_n) = \lambda$$

which shows $\lambda \in A$. As $A$ is a UFD then $\lambda$ may be decomposed into irreducibles of $A$, which by (3.17.44) are irreducible in $A[X]$. Therefore $A[X]$ is atomic.

Suppose that $f \in A[X]$ is irreducible we require to show it is prime. The case $\deg(f) = 0$ follows from the AP property of $A$, so assume $\deg(f) > 0$. Suppose $f \mid gh$ then as $K[X]$ is a UFD we see that, without loss of generality, $qf = g$ for $q \in K[X]$. Then $c(qf) = c(q)c(f) = c(q) = c(g) \in A$, so $q \in A[X]$ by (3.17.43) and we see $f \mid g$ in $A[X]$. This shows that $A[X]$ is AP and therefore a UFD by (3.17.21). ■

**Corollary 3.17.46**
*Suppose $A$ is a UFD. Then $A[X_1, ..., X_n]$ is a UFD.*

© David Rufino CC BY-NC-ND 4.0

## 3.18 Cayley-Hamilton Theorem

**Definition 3.18.1 (Characteristic Polynomial of a Matrix)**
*For a matrix $E \in \mathrm{Mat}_n(A)$ define the characteristic polynomial by*

$$P_E(X) := \det(X \cdot I_n - E)$$

*working in $\mathrm{Mat}_n(A[X])$. This is a monic polynomial in $A[X]$ and satisfies $P_E(X) = P_{E^t}(X)$ (3.5.20).*

**Definition 3.18.2 (Characteristic Polynomial of an endomorphism of a free module)**
*Let $M$ be a finite free $A$-module. Let $\mathcal{B}$ be a basis for $M$, $\phi : M \to M$ an $A$-module endomorphism and $[\phi]$ the matrix representation of $\phi$ with respect to $\mathcal{B}$. Define the characteristic polynomial of $\phi$ by*

$$P_\phi(X) := P_{[\phi]}(X)$$

*This is independent of the basis $\mathcal{B}$ by (3.5.20).*

**Lemma 3.18.3**
*Suppose $M = \langle m_1, ..., m_n \rangle$ is a finitely generated $A$-module and $\mathfrak{a} \triangleleft A$ an ideal, then*

$$\mathfrak{a}M = \mathfrak{a}m_1 + ... + \mathfrak{a}m_n$$

*That is every $m \in \mathfrak{a}M$ may be written as*

$$m = \sum_{i=1}^{n} a_i m_i \quad \text{for some } a_i \in \mathfrak{a}$$

*Proof.* By hypothesis

$$m = \sum_{j=1}^{n} b_j m'_j \quad \text{for some } m'_j \in M, b_j \in \mathfrak{a}$$

Furthermore by the finite-generation hypothesis for $j = 1...n$

$$m'_j = \sum_{i=1}^{n} c_{ji} m_i \quad \text{for some } c_{ji} \in A.$$

Therefore

$$m = \sum_{i=1}^{n} \left( \sum_{j=1}^{n} b_j c_{ji} \right) m_i$$

whence $a_i := \sum_{j=1}^{n} b_j c_{ji}$ lie in $\mathfrak{a}$ and give the required coefficients. ∎

**Theorem 3.18.4 (Cayley-Hamilton)**
*Let $M$ be a finitely generated $A$-module and $\phi \in \mathrm{End}_A(M)$. Then there exists a monic polynomial $P(X) \in A[X]$ such that $P(\phi) = 0$.*

*This may be strengthened in the following ways*

*a) If $M$ is a finite free $A$-module then $P$ may be taken to be the characteristic polynomial $P_\phi(X)$.*
*b) If $\phi(M) \subseteq \mathfrak{a}M$ for some ideal $\mathfrak{a} \triangleleft A$, then the non-leading coefficients of $P(X)$ may be chosen to be in $\mathfrak{a}$.*

*Proof.* First since $\mathrm{End}_A(M)$ is an $A$-algebra there is a canonical evaluation morphism

$$\mathrm{ev}_\phi : A[X] \to \mathrm{End}_A(M)$$

and the meaning of $P(\phi)$ is simply $\mathrm{ev}_\phi(P)$.

Let $\{m_1, ..., m_n\}$ be a generating set, then by definition

$$\phi(m_i) = \sum_{j=1}^{n} E_{ij} m_j \quad i = 1...n$$

for some $E \in \mathrm{Mat}_n(A)$. Consider the matrix

$$B(X) = XI_n - E \in \mathrm{Mat}_n(A[X])$$

Then we may define $B(\phi) := B(X)^{\mathrm{ev}_\phi} \in \mathrm{Mat}_n(\mathrm{End}_A(M))$ pointwise, so given by

© David Rufino CC BY-NC-ND 4.0

$$B(\phi)_{ij} = \delta_{ij}\phi - E_{ij}1_M \quad i,j = 1...n$$

By definition

$$\sum_j B(\phi)_{ij}m_j = \phi(m_i) - \sum_j E_{ij}m_j = 0$$

Formally we have a group action

$$\begin{aligned}\mathrm{Mat}_n(\mathrm{End}_A(M)) \times M^n &\to M^n \\ F \cdot \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix} &\to \begin{pmatrix} \sum_{j=1}^n F_{1j}(x_j) \\ \vdots \\ \sum_{j=1}^n F_{nj}(x_j) \end{pmatrix}\end{aligned}$$

such that $(EF)v = E(Fv)$ (check). And we have shown that

$$B(\phi)\begin{pmatrix} m_1 \\ \vdots \\ m_n \end{pmatrix} = \begin{pmatrix} 0 \\ \vdots \\ 0 \end{pmatrix}$$

Using (3.5.21), premultiply by the adjugate matrix to show that

$$\det(B(\phi))I_n\begin{pmatrix} m_1 \\ \vdots \\ m_n \end{pmatrix} = \begin{pmatrix} 0 \\ \vdots \\ 0 \end{pmatrix}$$

and $\det(B(\phi)) \in \mathrm{End}_A(M)$ annihilates $m_1, ..., m_n$ and therefore $M$.

Finally we claim that $P(X) := \det(B(X)) \in A[X]$ is a suitable monic polynomial. We see that

$$P(\phi) := \mathrm{ev}_\phi(\det(B(X))) = \det(B(X)^{\mathrm{ev}_\phi}) = \det(B(\phi)) = 0$$

When $M$ is a finite free $A$-module then we may choose $\{m_1, ..., m_n\}$ to be a basis, and then the matrix $E$ equals $[\phi]^T$ as required.

Finally when $\phi(M) \subseteq \mathfrak{a}M$ then (3.18.3) shows we may choose the coefficients $E_{ij}$ to be in $\mathfrak{a}$. It's clear that $P(X)$ then has non-leading coefficients in $\mathfrak{a}$. ∎

## 3.18.1 The $A[X]$-module associated to an endomorphism

**Proposition 3.18.5**
*Let $\phi : M \to M$ be an $A$-module endomorphism. Then there is a well-defined $A[X]$-module structure defined by*

$$\begin{aligned}A[X] \times M &\to M \\ (P, m) &\to P(\phi)(m)\end{aligned}$$

*where $P = \sum_{i=0}^n a_i X^i$. We denote this by $M_\phi$.*

*With this notation then $\phi$ is itself an $A[X]$-module endomorphism of $M_\phi$.*

*Proof.* Let $P, Q \in A[X]$ be two polynomials. Then we need to show that

$$(PQ)(\phi) = P(\phi) \circ Q(\phi)$$

which amounts to saying that $A[X] \to \mathrm{End}_A(M)$ is an $A$-algebra homomorphism.

The final statement follows because $X$ commutes with any polynomial in $A[X]$. More explicitly

$$\phi(P \cdot m) = \phi(P(\phi)(m)) = (XP)(\phi)(m) = (PX)(\phi)(m) = P(\phi)(\phi(m)) = P \cdot (\phi(m))$$

∎

**Proposition 3.18.6**
*Let $M$ be an $A$-module and $\phi \in \mathrm{End}_A(M)$. Then is an $A[X]$-linear map*

$$\begin{aligned}\tilde{\phi} : A[X] \otimes_A M &\to M_\phi \\ P \otimes m &\to P(\phi)(m)\end{aligned}$$

© David Rufino CC BY-NC-ND 4.0

*and similarly*

$$1 \otimes \phi : A[X] \otimes_A M \to A[X] \otimes_A M$$
$$P \otimes m \to P \otimes (\phi(m))$$

*Proof.* By the universal property $\tilde{\phi}$ exists and is $A$-linear. To show it is $A[X]$-linear observe that

$$\tilde{\phi}(Q(P \otimes m)) = \tilde{\phi}((QP) \otimes m) = (QP)(\phi)(m) = Q(\phi)(P(\phi)(m)) = Q \cdot P(\phi)(m) = Q \cdot \tilde{\phi}(P \otimes m)$$

and so the result follows by linearity.

The second exists by (3.4.17). ∎

**Proposition 3.18.7**
*Let $M$ be a finite free $A$-module and $\phi \in \mathrm{End}_A(M)$. Then the characteristic polynomial $P_\phi(X)$ equals the determinant of the $A[X]$-linear map*

$$(X - 1 \otimes \phi) : A[X] \otimes_A M \to A[X] \otimes_A M$$

*Furthermore*

$$P_\phi(X) = X^n + \sum_{j=1}^{n} (-1)^j \operatorname{Tr}(L_a^j(\phi)) X^{n-j}$$

*and in particular*

$$P_\phi(X) = X^n - \operatorname{Tr}(\phi) X^{n-1} + ... + (-1)^n \det(\phi)$$

*Proof.* Let $\mathcal{B} := \{v_1, ..., v_n\}$ be a basis for $M$, then by (3.4.24) $\{1 \otimes v_1, ..., 1 \otimes v_n\}$ is an $A[X]$-basis for $A[X] \otimes_A M$. Let $E = [\phi]$ be the matrix associated to the basis $\mathcal{B}$ then by definition

$$\phi(v_i) = \sum_{j=1}^{n} E_{ji} v_j$$

and

$$(X - 1 \otimes \phi)(1 \otimes v_i) = X \otimes v_i - 1 \otimes \phi(v_i) = X(1 \otimes v_i) - \sum_{j=1}^{n} E_{ji}(1 \otimes v_j)$$

This shows that

$$[X - 1 \otimes \phi]_{ij} = X\delta_{ij} - E_{ij}$$

and so $P_\phi(X) = \det(XI_n - E) = \det(X - 1 \otimes \phi)$ as required.

Then by (3.5.31) we have

$$P_\phi(X) = X^n + \sum_{j=1}^{n} (-1)^j \operatorname{Tr}(L_a^j(1 \otimes \phi)) X^{n-j}$$

The matrix of $1 \otimes \phi$ coincides with that of $\phi$ and so we deduce the second relationship. The final statement follows from (...). ∎

© David Rufino CC BY-NC-ND 4.0

# 3.19 Fields and Galois Theory

## 3.19.1 Prime Fields

Recall that for $p$ prime the quotient ring $\mathbb{Z}/p\mathbb{Z}$ is a field (3.16.3), (3.17.22), (3.3.59).

**Definition 3.19.1 (Finite field of order $p$)**
*Denote by $\mathbb{F}_p$ the field $\mathbb{Z}/p\mathbb{Z}$ of order $p$.*

**Definition 3.19.2 (Rational Integers)**
*We denote by $\mathbb{Q}$ the field of **rational numbers** defined to be the quotient field of the integers,* $\mathrm{Frac}(\mathbb{Z})$ *(see (3.6.6)).*

**Definition 3.19.3 (Prime Field)**
*We say that a field $k$ is a **prime field** if it is isomorphic to one of $\mathbb{F}_p$ or $\mathbb{Q}$.*

*Note none of these fields are mutually isomorphic by considering cardinalities.*

**Proposition 3.19.4 (Prime Subfield Exists)**
*Let $k$ be a field. Then $k$ contains a **prime subfield**. It is the smallest subfield contained in $k$.*

*Proof.* By (...) there is a unique ring homomorphism

$$\phi : \mathbb{Z} \to k$$

As $k$ is not the zero-ring then $\phi(1) = 1 \neq 0$ and $\ker(\phi)$ is a proper ideal. Then $\mathrm{Im}(\phi)$ is an integral domain by (3.3.9). By (3.3.57) $\mathbb{Z}/\ker(\phi) \cong \mathrm{Im}(\phi)$ and therefore by (3.3.59) $\ker(\phi) =: \mathfrak{p}$ is a prime ideal.

By (3.16.3) the ideal $\mathfrak{p}$ is principal. Suppose $\mathfrak{p} = (0)$. By (3.6.5) $\phi$ extends to a homomorphism $\phi' : \mathbb{Q} \to k$ whose kernel is zero and therefore injective (...). In particular $\mathrm{Im}(\phi')$ is a prime subfield being isomorphic to $\mathbb{Q}$.

Otherwise by (3.17.9) $\mathfrak{p} = (p)$ for $p$ a prime number and we have already observed that $\mathbb{F}_p = \mathbb{Z}/p\mathbb{Z} \cong \mathrm{Im}(\phi)$ is a prime subfield.

Let $k'$ be any subfield of $k$, then by induction we may show that $\mathrm{Im}(\phi) \subset k'$. In the case $\ker(\phi) = \{0\}$ then we may also show $\mathrm{Im}(\phi') \subset k'$. Therefore $k'$ contains the prime subfield, whence it is the smallest subfield. ∎

**Definition 3.19.5 (Characteristic)**
*Let $k$ be a field. Then we define the **characteristic** of $k$ to be $p$ if the prime subfield is isomorphic to $\mathbb{F}_p$ or $0$ if the prime subfield is isomorphic to $\mathbb{Q}$. This is denoted* $\mathrm{char}(k)$.

*If $A$ is a $k$-algebra then we define the characteristic of $A$ to be that of $k$. We also define the **characteristic exponent** of $A$ to be*

$$\begin{cases} 1 & \text{if } \mathrm{char}(A) = 0 \\ p & \text{if } \mathrm{char}(A) = p \end{cases}$$

**Proposition 3.19.6 (Frobenius Homomorphism)**
*Let $A$ be a $k$-algebra and suppose* $\mathrm{char}(A) = p$. *Then the mapping*

$$a \to a^p$$

*is a ring homomorphism which we call the **Frobenius map**. In particular*

$$(a+b)^p = a^p + b^p \quad \forall a, b \in A$$

**Definition 3.19.7**
*Let $A$ be a $k$-algebra with characteristic exponent $p$. We say that $A$ is **perfect** if $A$ is reduced and $A^p = A$. When $p > 1$ this is equivalent to the Frobenius homomorphism being an isomorphism.*

## 3.19.2 Separability

We consider separability of a polynomial, and how it behaves under field extension.

**Definition 3.19.8 (Splitting Polynomial)**
*Let $K/k$ be a field extension and $f \in k[X]$. By abuse of notation we may also identify $f$ with its image in $K[X]$. We say a polynomial $f$ **splits completely** in $K$ if the irreducible factorization of $f$ in $K[X]$ is*

$$f = \ell(f) \prod_{i=1}^{n} (X - \alpha_i)^{r_i}$$

© David Rufino CC BY-NC-ND 4.0

*where $\alpha_i$ are the distinct roots of $f$ in $K$ and $r_i := v_{X-\alpha_i}(f)$ are the multiplicities. Equivalently $f$ splits in $K$ if*

$$p \in K[X] \text{ irreducible} \wedge \deg(p) > 1 \implies v_p(f) = 0$$

*Observe that the number of roots counting multiplicities is* $\deg(f)$

$$\deg(f) = \sum_{i=1}^{n} v_{X-\alpha_i}(f)$$

**Corollary 3.19.9**
*Let $K/k$ be a field extension and $f, g \in k[X]$. Suppose $g \mid f$ (in $k[X]$) and $f$ splits completely in $K$. Then so does $g$.*

*Proof.* As $k[X] \subset K[X]$, $g \mid f$ in $K[X]$ too. By assumption the irreducible factorization of $f$ in $K[X]$ consists of polynomials of degree 1. Consider the irreducible factorization of $g = \prod_{i=1}^{n} g_i$ in $K[X]$, then by unique factorization (3.17.33) (applied to $K[X]$) each $g_i$ must appear in the factorization of $f$, that is to say $g$ splits completely in $K$. ■

**Lemma 3.19.10 (Coprime Polynomials have no Common Roots)**
*Let $K/k$ be a field extension and $f, g \in k[X]$ with $f \neq 0$ splitting completely in $K$. Then $f$ and $g$ are coprime if and only if they have no common root in $K$. In particular this holds whenever $f$ and $g$ both split completely in $K$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Suppose $f, g$ are coprime, so $af + bg = 1$ for some $a, b \in k[X]$. If $\alpha \in K$ were a common root then evaluating at $\alpha$ gives $0 = 1$, a contradiction.

Conversely suppose $f, g$ are not coprime. As $k[X]$ is a PID (3.17.32), by (3.16.5) they have a common non-invertible divisor $d$, and $\deg(d) > 0$ as the invertible elements of $k[X]$ are the nonzero constants (3.8.15). Since $d \mid f$ and $f$ splits completely in $K$, so does $d$ (3.19.9); as $\deg(d) > 0$ it has a root $\alpha \in K$, which is a common root of $f$ and $g$ as $d$ divides both. ■

We can provide a partial converse to (3.17.39) by working in a large enough extension field

**Proposition 3.19.11 (Separability)**
*Let $K/k$ be a field extension and $f \in k[X]$ a polynomial which splits completely in $K$. Then TFAE*

*a) $f$ is separable*
*b) $f$ has no multiple roots in $K$*
*c) $f$ and $f'$ have no common roots in $K$*
*d) $f$ has $\deg(f)$ distinct roots in $K$*

*Proof.* Using the formula

$$\deg(f) = \sum_{i=1}^{n} v_{X-\alpha_i}(f)$$

we see easily that $d) \iff b)$. By (3.17.37) $b) \iff c)$. As $f$ splits completely in $K$, (3.19.10) shows that $f$ and $f'$ are coprime, that is $f$ is separable (3.17.38), if and only if they have no common root in $K$: $a) \iff c)$. ■

## 3.19.3 Field Extensions

**Definition 3.19.12 (Field Extension)**
*Let $k$ be a field. A field extension $K/k$ is a $k$-algebra $K$ which is also a field. Every field $K$ is an extension over its prime subfield (or, over $\mathbb{Q}$ or $\mathbb{F}_p$).*

*We typically denote the structural morphism by $i_{kK} : k \to K$, and it is automatically injective (3.3.61). We may write $(K/k, i_{kK})$ if we need to stress the relevance of the structural morphism to the argument.*

*These objects form a category* $\mathbf{Field}_k$ *in the obvious way. The morphisms may be called $k$-embeddings and we denote them by*

$$\mathrm{Mor}_k(K, L) := \{\psi : K \to L \mid \psi \circ i_{kK} = i_{kL}\}$$

*and the set of automorphisms by*

$$\mathrm{Aut}(K/k).$$

*Observe every extension $K/k$ may be viewed as a $k$-vector space so we define the degree of an extension field to be the vector space dimension*

© David Rufino CC BY-NC-ND 4.0

$$[K:k] := \dim_k K$$

**Definition 3.19.13 (Finite field extension)**
*A field extension $K/k$ is finite if $[K:k] < \infty$*

**Definition 3.19.14 (Tower of Field Extensions)**
*We may also consider a "tower" of extensions*

$$K_n/\cdots/K_0 = k$$

*with embeddings $i_{K_iK_{i+1}} : K_i \to K_{i+1}$, with the picture that these usually correspond to inclusions. We may consider an extension $K_i/K_j$ for $j < i$. Typically if we have a family of morphisms*

$$\sigma_i : K_i \to M$$

*they would commute with these embeddings. In particular we may abuse notation by defining $\sigma_i|_{K_j} = \sigma_i \circ i_{K_{i-1}K_i} \circ \cdots \circ i_{K_jK_{j+1}}$.*

**Proposition 3.19.15**
*Let $K/k$ be a finite field extension and $V$ a finite-dimensional vector space over $K$. Then $V$ is naturally a finite-dimensional $k$-vector space with*

$$\dim_k V = [K:k] \dim_K V$$

*More precisely if $K/k$ has basis $\{\lambda_1, ..., \lambda_r\}$ and $V$ has $K$-basis $\{v_1, ..., v_n\}$, then the following is a $k$-basis for $V$*

$$\left\{\lambda_i v_j\right\}_{i,j}.$$

**Proposition 3.19.16**
*Let $L/K$ and $K/k$ be two finite extensions with basis $\{l_1, ..., l_n\}$ and $\{k_1, ..., k_m\}$. Then $L/k$ has basis $\left\{l_i k_j\right\}_{i,j}$. In particular*

$$[L:k] = [L:K][K:k]$$

**Corollary 3.19.17**
*Let $K = K_n/\cdots/K_0 = k$ be a tower of finite extensions then*

$$[K:k] = \prod_{i=1}^{n} [K_i : K_{i-1}]$$

**Lemma 3.19.18**
*Let $K/k$ be a field extension and $f, g \in k[X]$. Then $g \mid f$ in $k[X]$ if and only if $i_{kK}(g) \mid i_{kK}(f)$.*

*Proof.* One implication is obvious. For the converse we assume wlog that $k \subset K$ and suppose $f = gh$ for $h \in K[X]$. We may apply the division algorithm (3.8.16) in $k[X]$ to find $f = gq + r$ for $q, r \in k[X]$ and $\deg(r) < \deg(g)$. Then $r = g(h - q)$ and comparing degrees we conclude that $h = q$ and $r = 0$. This shows $f = gq$ and $g \mid f$ as elements of $k[X]$. ∎

**Definition 3.19.19 (Evaluation homomorphism)**
*Let $K/k$ be a field extension and $\alpha \in K$. There is a canonical homomorphism*

$$\mathrm{ev}_\alpha : k[X] \to K$$

$$\sum_{i=0}^{n} a_i X^i \to \sum_{i=0}^{n} i_{kK}(a_i)\alpha^i$$

*which we write as $f(\alpha)$. We say $\alpha \in K$ is a root of $f(X)$ if $f(\alpha) = 0$.*

**Proposition 3.19.20 (Morphisms commute with evaluation)**
*Let $\sigma : K/k \to L/k$ be a morphism of field extensions then*

$$\sigma(p(\alpha)) = p(\sigma(\alpha))$$

*for all $p \in k[X]$. In particular $\alpha$ is a root of $p \iff \sigma(\alpha)$ is a root of $p$.*

*Proof.* This is just a specific case of (3.8.8). The last statement is obvious, because $\sigma$ is injective (3.3.61). ∎

© David Rufino CC BY-NC-ND 4.0

**Definition 3.19.21 (Subalgebra generated by a set)**
*Let $K/k$ be a field extension and $S \subset K$. Define*

$$k[S] := \bigcap_{\substack{A \subset K/k \\ S \subset A}} A$$

*where the intersection is taken over all $k$-subalgebras. It is the smallest $k$-subalgebra of $K/k$.*

**Proposition 3.19.22 (Subalgebra generated by directed family)**
*Let $K/k$ be a field extension and $\{S_i\}_{i \in I}$ a family of directed subsets of $K$. Then*

$$k\left[\bigcup_{i \in I} S_i\right] = \bigcup_{i \in I} k[S_i]$$

*In particular for any set $S$ we have*

$$k[S] = \bigcup_{\substack{S' \subset S \\ S' \text{ finite}}} k[S']$$

**Proposition 3.19.23 (Subalgebra generated by a set)**
*Let $K/k$ be a field extension and $S \subset K$ a finite subset. Write $S = \{\alpha_1, ..., \alpha_n\}$ then*

$$k[S] = \{p(\alpha_1, ..., \alpha_n) \mid p \in k[X_1, ..., X_n]\} = \mathrm{Im}(\mathrm{ev}_\alpha)$$

**Lemma 3.19.24 (Trivial result)**
*For $S, T \subset K/k$ finite*

- $S \subset T \implies k[S] \subseteq k[T]$
- $k[S][T] = k[S \cup T]$

**Definition 3.19.25**
*Let $K/k$ be a field extension and $S \subset K$. Define*

$$k(S) := \bigcap_{\substack{K' \subset K/k \\ S \subset K'}} K'$$

*It is the smallest subfield of $K/k$ containing $S$.*

**Proposition 3.19.26**
*Let $K/k$ be a field extension and $\{S_i\}_{i \in I}$ be a family of directed subsets of $K$. Then*

$$k\left(\bigcup_{i \in I} S_i\right) = \bigcup_{i \in I} k(S_i)$$

*In particular for any set $S$ we have*

$$k(S) = \bigcup_{\substack{S' \subset S \\ \text{finite}}} k(S')$$

**Proposition 3.19.27**
*Let $K/k$ be a field extension and $S \subset K$ be a finite subset. Write $S = \{\alpha_1, ..., \alpha_n\}$ then*

$$k(S) := \left\{ \frac{p(\alpha_1, ..., \alpha_n)}{q(\alpha_1, ..., \alpha_n)} \;\middle|\; p, q \in k[X_1, ..., X_n], \quad q(\alpha_1, ..., \alpha_n) \neq 0 \right\}$$

**Lemma 3.19.28 (Trivial result)**
*For $S, T \subset K/k$*

- $S \subset T \implies k(S) \subseteq k(T)$
- $k(S)(T) = k(S \cup T)$

**Lemma 3.19.29**
*If $S \subset K$ and $k[S]$ is a field then $k[S] = k(S)$*

*Proof.* Generically $k[S] \subset k(S)$ from the definition. As $k[S]$ is a field then $k(S) \subset k[S]$. ∎

© David Rufino CC BY-NC-ND 4.0

**Lemma 3.19.30 (Image of f.g. field extension)**
*Let $K/k$ be a field extension and $S \subset K$ a subset. If $\sigma : K/k \to L/k$ is a morphism them*

$$\sigma(k(S)) = k(\sigma(S))$$

**Proposition 3.19.31 (Uniqueness of morphisms on a generating set)**
*Let $K/k$ be a field extension and $S \subset K$. If $\sigma, \sigma' : k(S)/k \to L/k$ are morphisms of field extensions such that $\sigma|_S = \sigma'|_S$. Then $\sigma = \sigma'$.*

**Definition 3.19.32 (Simple (Algebraic) Extension)**
*A field extension $K/k$ is **simple** if $K = k(\{\alpha\}) =: k(\alpha)$ for some $\alpha \in K$. It is a **simple algebraic** extension if $\alpha$ is also algebraic over $k$.*

**Definition 3.19.33 (Algebraic Element)**
*We say an element $\alpha \in K/k$ is **algebraic** if it is a root of a polynomial $f \in k[X]$ (i.e. $\alpha$ is integral, since we can always ensure $f$ is monic). Otherwise we say that $x \in K$ is **transcendental**.*

*We say $K/k$ is an **algebraic extension** if every element $\alpha \in K$ is algebraic over $k$.*

**Proposition 3.19.34 (Finite ⇒ algebraic)**
*A finite extension $K/k$ is algebraic.*

*Proof.* Suppose $n = \dim_k K$. The set $\{1, \alpha, \alpha^2, ..., \alpha^n\}$ is linearly dependent by (2.3.13). Therefore there is a non-zero polynomial with $\alpha$ as a root. ∎

**Proposition 3.19.35 (Endomorphisms are automorphisms)**
*Let $\sigma \in \mathrm{Mor}_k(K, K)$ be an endomorphism of an algebraic extension. Then it is an isomorphism. In other words*

$$\mathrm{Mor}_k(K, K) = \mathrm{Aut}(K/k)$$

*Proof.* As field morphisms are injective (3.3.61) we only need to show that $\sigma$ is surjective. Given $\alpha \in K$ let $T$ denote the set of roots of $m_\alpha \in k[X]$ in $K$. Note by (3.17.35) $T$ is finite. Further by (3.19.20) $\sigma$ maps $T$ to itself. Since $\sigma$ is injective it is also surjective on $T$. In particular $\alpha$ is in the image of $\sigma$ as required. ∎

## 3.19.4 Algebraic Extensions

**Proposition 3.19.36 (Minimal Polynomial)**
*If $\alpha \in K/k$ is algebraic then there is a unique monic, irreducible polynomial $m_{\alpha,k}(X) \in k[X]$ such that $m_{\alpha,k}(\alpha) = 0$. This is called the minimal polynomial of $\alpha$ over $k$ and $\big(m_{\alpha,k}\big) = \ker(\mathrm{ev}_\alpha)$.*

*In particular any polynomial $f(X) \in k[X]$ which has $\alpha$ as a root, satisfies $m_{\alpha,k}(X) \mid f(X)$.*

*Proof.* Let $\mathfrak{a} = \ker(\mathrm{ev}_\alpha)$. Since $k[X]$ is a PID it is of the form $\big(m_{\alpha,k}\big)$. As $\alpha$ is algebraic it is non-zero. $m_{\alpha,k}(X)$ cannot be a constant, and therefore is not a unit.

We claim $m_{\alpha,k}$ is irreducible. If $m_{\alpha,k}(X) = p(X)q(X)$ then $p, q$ are non-zero and either $p(\alpha) = 0$ or $q(\alpha) = 0$. If $p(\alpha) = 0$ then $m_{\alpha,k} \mid p$. As $p \mid m_{\alpha,k}$ by (3.17.30) $m_{\alpha,k} = \lambda p$. In particular $\deg\big(m_{\alpha,k}(X)\big) = \deg(p(X))$ so $\deg(q(X)) = 0$ and $q(X)$ is a unit (3.8.15). Therefore by definition $m_{\alpha,k}(X)$ is irreducible.

Dividing by the leading coefficient we may assume that this polynomial is monic. Suppose $m'(X)$ is another such irreducible monic polynomial. Then $m_\alpha \mid m'$. Since $m_\alpha$ is not a unit, by definition of irreducible $m' \sim m_\alpha$ whence $m' = \lambda m_\alpha$. Compare leading coefficients to find $\lambda = 1$ and $m' = m_\alpha$. ∎

**Lemma 3.19.37**
*Let $K/E/k$ be extensions and $\alpha \in K$ algebraic over $k$. Then $\alpha$ is algebraic over $E$.*

**Definition 3.19.38 (Conjugate elements)**
*Two elements $\alpha, \beta \in K$ are said to be **conjugate elements** if they have the same minimal polynomial.*

*NB it's necessary and sufficient that $m_{\alpha,k}(\beta) = 0$.*

**Proposition 3.19.39**
*Let $\sigma : K/k \to L/k$ be a field morphism and $\alpha \in K$. Then $m_{\alpha,k}(X) = m_{\sigma(\alpha),k}(X)$.*

*Proof.* This follows from (3.19.20). ∎

Given an irreducible polynomial $f \in k[X]$ it's possible to construct an extension field $K/k$ which has at least one root, as follows.

© David Rufino CC BY-NC-ND 4.0

**Proposition 3.19.40 (Construct simple extension)**
*Let $f \in k[X]$ be an irreducible polynomial. Then $(f)$ is maximal and $K := k[X]/(f)$ is a field extension with canonical structural morphism. Define $\alpha := X + (f)$*

- $f(\alpha) = 0$
- $K = k(\alpha)$ *is a simple field extension and* $k(\alpha) = k[\alpha]$
- $m_\alpha = f/\ell(f)$ *and* $\deg(m_\alpha) = \deg(f) =: n$
- $K$ *is a finite-dimensional $k$-vector space with basis*

$$\{1, \alpha, ..., \alpha^{n-1}\}$$

.

**Example 3.19.41**
*Take $k = \mathbb{R}$, $f(X) = X^2 + 1$, then $\mathbb{C}/\mathbb{R} = \mathbb{R}[i] = \mathbb{R}[X]/(X^2+1)$.*

*Proof.* Consider the structural morphism $i : k \to k[X]$ and canonical surjective homomorphism

$$\pi : k[X] \to k[X]/(f)$$

and $\alpha = X + (f) = \pi(X)$. As $k[X]$ is a PID, $f$ irreducible implies $(f)$ maximal by (3.17.22) so $K$ is a field by (3.3.59). The composition $\pi \circ i$ makes $K$ into a $k$-algebra and hence a field extension. Furthermore $\pi$ is then by definition a $k$ -algebra homomorphism.

Since $\pi$ is surjective every $\beta \in K$ is represented as $\pi(p(X)) \overset{(3.19.20)}{=} p(\pi(X)) = p(\alpha)$. By (3.19.23) we see $K = k[\alpha]$. Since $K$ is a field then $K = k[\alpha] = k(\alpha)$ is simple by (3.19.29).

Similarly $f(\alpha) = f(\pi(X)) \overset{(3.19.20)}{=} \pi(f(X)) = 0$, so $\alpha$ is a root of $f$. By (3.17.30) $f/\ell(f)$ is irreducible and by uniqueness in (3.19.36) we have $m_\alpha = f/\ell(f)$.

Given $\beta = p(\alpha)$, the division algorithm (...) yields

$$p(X) = q(X)f(X) + r(X)$$

with $\deg(r) < \deg(f) = n$. Therefore $\beta = r(\alpha)$ and the given set is spanning. A non-trivial linear dependence yields a non-zero polynomial $g(X)$ such that $g(\alpha) = 0$ and $\deg(g) < \deg(f)$. But by definition of the minimal polynomial $m_\alpha \mid g$, which is a contradiction by comparing degrees. Therefore the given set is linearly independent and hence a basis. ∎

Conversely any simple algebraic extension is obtained in this way, as follows

**Proposition 3.19.42 (Simple extension)**
*Let $k(\alpha)/k$ be a simple extension. Then there is a canonical isomorphism of $k$-algebras*

$$k[X]/(m_\alpha) \longrightarrow k(\alpha)$$

*under which $X + (m_\alpha) \to \alpha$. Further $k(\alpha)$ is a finite-dimensional vector space with basis*

$$\{1, \alpha, ..., \alpha^{n-1}\}$$

*where $n = \deg(m_\alpha) = [k(\alpha) : k]$ and $k(\alpha) = k[\alpha]$.*

*Proof.* By (3.19.36), Definition (3.19.23) and (3.3.57) there is a canonical isomorphism $k[X]/(m_\alpha) \to k[\alpha]$ of $k$-algebras induced by the evaluation homomorphism $\mathrm{ev}_\alpha : k[X] \to K$. (3.19.40) shows that the image of this isomorphism, $k[\alpha]$, is a field, whence $k[\alpha] = k(\alpha)$ by (3.19.29). Since a $k$-algebra isomorphism is a-fortiori a $k$-vector space isomorphism it maps a basis to a basis. The result follows from (3.19.40) as the basis thus defined is the image of the basis in the proposition under the specified isomorphism. ∎

**Definition 3.19.43 (Degree of an algebraic element)**
*Let $K/k$ be an algebraic extension and $\alpha \in K$. Then define*

$$\deg_k(\alpha) := \deg m_{\alpha,k} = [k(\alpha) : k]$$

We may show the following

© David Rufino CC BY-NC-ND 4.0

**Proposition 3.19.44 (Finitely generated by algebraic $\implies$ finite and algebraic)**
*Let $K = k(\alpha_1, ..., \alpha_n)/k$ be a field extension such that $\alpha_i$ is algebraic. Then $K/k$ is a finite algebraic extension. Furthermore*

$$k[\alpha_1, ..., \alpha_n] = k(\alpha_1, ..., \alpha_n)$$

*In particular a finitely-generated algebraic extension is finite.*

*Proof.* We write $K_i = k(\alpha_1, ..., \alpha_i)$. Then we have a tower

$$K = K_n/.../K_0 = k$$

such that $K_i = K_{i-1}(\alpha_i)$ is a simple algebraic extension. By (3.19.42) $K_i/K_{i-1}$ is finite. Therefore by (3.19.17) $K/k$ is finite. By (3.19.34) it's also algebraic. For the second statement we may proceed inductively. Note we have

$$k[\alpha_1, ..., \alpha_{i+1}] \overset{(3.19.24)}{=} k[\alpha_1, ..., \alpha_i][\alpha_{i+1}] = k(\alpha_1, ..., \alpha_i)[\alpha_{i+1}] \overset{(3.19.42)}{=} k(\alpha_1, ..., \alpha_i)(\alpha_{i+1}) \overset{(3.19.28)}{=} k(\alpha_1, ..., \alpha_{i+1})$$

The second equality is simply the inductive hypothesis. ∎

**Corollary 3.19.45**
*Let $K/k$ be a field extension then the algebraic elements form a subfield.*

*Proof.* For any two algebraic elements $\alpha, \beta \in K$ we have $k(\alpha, \beta)$ is an algebraic extension. ∎

The following is useful for reducing to cases of finite extensions where counting arguments work.

**Lemma 3.19.46 (Reduce to finite extensions)**
*Let $K/E/k$ be a tower with $E$ algebraic over $k$. For every $\alpha \in K$ algebraic over $E$, there is some subfield $E_0 \subset E$ such that*

- *$E_0/k$ is finite*
- *$\alpha$ is algebraic over $E_0$*
- *$m_{\alpha,E} = m_{\alpha,E_0}$*

*Therefore $\alpha$ is algebraic over $k$ iff it is algebraic over $E$ and $m_{\alpha,E_0} \mid m_{\alpha,k}^{i_{kE}}$.*

*Proof.* Suppose

$$m_{\alpha,E}(X) = a_0 + a_1 X + ...a_n X^n$$

Then define $E_0 = i_{kE}(k)(a_0, ..., a_n)$. By (3.19.44) $E_0/k$ is finite. Clearly $\alpha$ is algebraic over $E_0$ as it is a root of $m_{\alpha,E}$. By (3.19.36) $m_{\alpha,E_0} \mid m_{\alpha,E}$ as elements of $E_0[X]$ and $m_{\alpha,E} \mid m_{\alpha,E_0}$. Therefore $m_{\alpha,E_0} = m_{\alpha,E}$.

By (3.19.42) $E_0(\alpha)/E$ is finite, therefore $E_0(\alpha)/k$ is finite. By (3.19.34) $E_0(\alpha)/k$ is algebraic, whence $\alpha$ is algebraic over $k$. The last statement follows from (3.19.36) again. ∎

**Corollary 3.19.47 (Algebraic Extensions are Directed Unions of Finite Subextensions)**
*Let $E/k$ be algebraic. Then the finite subextensions of $E/k$ are directed under inclusion, and $E$ is their union.*

*Proof.* Given $\alpha \in E$, taking $K = E$ in (3.19.46) (with $\alpha$ trivially algebraic over $E$) gives a finite subextension $E_0 \subset E$ over which $\alpha$ is algebraic; then $E_0(\alpha)/E_0$ is finite (3.19.42), hence $E_0(\alpha)/k$ is finite too, a finite subextension of $E$ containing $\alpha$. Hence $E$ is the union of its finite subextensions. Given two finite subextensions $E_1, E_2 \subset E$, their compositum $E_1E_2 \subset E$ is finitely-generated and algebraic over $k$, hence finite (3.19.44), and contains both, so the finite subextensions are directed under inclusion. ∎

**Corollary 3.19.48**
*$K/E$ and $E/k$ are both algebraic if and only if $K/k$ is.*

*Proof.* One direction is (3.19.37). The converse follows from the previous result. ∎

We may prove the first lifting theorem

**Proposition 3.19.49 (Lifting to simple extensions)**
*Let $k(\alpha)/k$ be a simple algebraic extension and $L/k$ a field extension such that $m_{\alpha,k}$ has a root in $L$. Then there exists a morphism $\sigma : k(\alpha)/k \to L/k$.*

*More precisely there is a bijective mapping*

© David Rufino CC BY-NC-ND 4.0

$$\begin{aligned}\mathrm{Mor}_k(k(\alpha), L) &\longrightarrow \{\beta \in L \mid m_{\alpha,k}(\beta) = 0\} \\ \sigma &\to \ \sigma(\alpha)\end{aligned}$$

*and* $\sigma(k(\alpha)) = k(\sigma(\alpha))$*. In particular if* $m_{\alpha,k}$ *is separable and splits completely in* $L$ *then there are precisely* $\deg(m_\alpha) \overset{(3.19.42)}{=} [k(\alpha) : k]$ *such extensions.*

*Proof.* Observe $m_{\alpha,k}(\sigma(\alpha)) \overset{(3.19.20)}{=} \sigma\big(m_{\alpha,k}(\alpha)\big) = 0$. Therefore the mapping is well-defined. By (3.19.31) it is injective. We claim it is also surjective. By (3.19.42) there is a $k$-algebra isomorphism

$$k[X]/\big(m_{\alpha,k}\big) \longrightarrow k(\alpha)$$

Similarly for $\beta \in T$ there is a $k$-algebra isomorphism

$$k[X]/\big(m_{\beta,k}\big) \longrightarrow k(\beta)$$

We are done if $m_{\alpha,k} = m_{\beta,k}$. But $m_{\alpha,k}$ is monic, irreducible and has $\beta$ as a root. So this follows from uniqueness of the minimal polynomial in (3.19.36). The final statement follows from (3.19.11) ∎

We may use this to generalize to arbitrary extensions, but we require that the minimal polynomials split completely in order for the inductive step to work.

**Proposition 3.19.50 (Generic Lifting Theorem)**
*Let* $K/k$ *be an algebraic field extension such that* $K = k\big(\{\alpha_i\}_{i\in I}\big)$ *and* $L/k$ *a field extension such that* $m_{\alpha_i,k}(X)$ *splits completely in* $L$ *for all* $i \in I$*.*

*Then there exists a morphism* $\sigma : K/k \to L/k$*.*

*Furthermore given* $\alpha \in K$ *and* $\beta \in L$ *any root of* $m_{\alpha,k}(X)$ *we may choose* $\sigma$ *such that* $\sigma(\alpha) = \beta$*.*

*Proof.* If $K/k$ is finite then we may proceed by induction on $[K : k]$, using (3.19.49) and applying a similar argument to below.

For the general case we may consider the poset of morphisms $\sigma : K'/k \to L/k$ for subfields $K'/k \subset K/k$ ordered by consistency. It is non-empty by considering $K' = i_{kK}(k)$. By Zorn's Lemma it has a maximal element, $(K', \sigma')$. It's enough to show that $K' = K$.

If $\alpha_i \in K'$ for all $i \in I$ then $K' = K$ and we are done. Otherwise choose $\alpha = \alpha_i \notin K'$. By (...) $m_{\alpha,K'}(X) \mid m_{\alpha,k}(X)$. By (3.19.9) $m_{\alpha,K'}(X)$ splits in $L$ (because $m_{\alpha,k}(X)$ does). Therefore by (3.19.49) there is a morphism $\sigma : K'(\alpha)/K' \to (L/K', \sigma')$. Note that by definition $\sigma|_{K'} = \sigma'$ and $K' \subsetneq K'(\alpha)$, contradicting maximality.

For the final part we may consider the poset consisting only of morphisms such that $\sigma(\alpha) = \beta$. By (3.19.49) the poset is non-empty, and the same argument works. ∎

## 3.19.5 Galois Theory Summary

**Definition 3.19.51 (Separable, Normal and Galois)**
*Let* $K/k$ *be an algebraic extension. We say that* $K/k$ *is*

- ***Normal*** *if every minimal polynomial* $m_{\alpha,k} \in k[X]$ *splits completely in* $K$ *(iff every irreducible polynomial* $f \in k[X]$ *with at least one root in* $K$ *splits completely in* $K$*)*
- ***Separable*** *if every minimal polynomial* $m_{\alpha,k} \in k[X]$ *is separable.*
- ***Galois*** *if it is both normal and separable (iff* $m_{\alpha,k}$ *has* $\deg\big(m_{\alpha,k}\big)$ *distinct roots in* $K$*, see (3.19.11)).*

*In the case of a Galois extension we denote the group of automorphisms by* $\mathrm{Gal}(K/k)$*.*

To summarize the main results

a) The group of automorphism of a normal extension $K/k$ acts transitively on the roots of a given irreducible polynomial.
b) For $K/k$ finite we have $\#\,\mathrm{Aut}(K/k) \leq [K : k]$ with equality if and only if $K/k$ is Galois.
c) An algebraic extension $K/k$ is automatically separable whenever either $\mathrm{char}(k) = 0$ or $k$ is finite.
d) When $K/k$ is finite and Galois then we have an order-reversing bijection between subfields and subgroups

$$\begin{aligned}\{H \leq \mathrm{Gal}(K/k)\} &\longleftrightarrow \{F \subseteq K\} \\ H &\longrightarrow K^H := \{x \in K \mid h(x) = x \quad \forall h \in H\} \\ \mathrm{Gal}(K/F) &\longleftarrow F\end{aligned}$$

© David Rufino CC BY-NC-ND 4.0

### 3.19.6 Splitting Fields and Algebraic Closure

In this section we discuss splitting fields, which are the "smallest" extensions in which a given set of polynomials split completely. The fundamental result is that splitting fields are precisely the Normal extensions. Further we discuss the algebraic closure, in which every polynomial splits and in which every algebraic extension (normal or otherwise) may be embedded.

**Definition 3.19.52 (Splitting field)**
*Let $S \subset k[X]$ a family of polynomials. We say that $K/k$ is a **splitting field** for $S$ if*

- *Every polynomial $f \in S$ splits completely in $K$*
- *$K$ is generated by the roots of all the polynomials in $S$*

*Note that by (3.19.44) $K/k$ is necessarily algebraic, and if $S$ is finite then so is $K/k$.*

**Definition 3.19.53 (Set of roots)**
*Let $K/k$ be a field extension and $f \in k[X]$. Then define*

$$T_{f,K} := \{\beta \in K \mid f(\beta) = 0\}$$

**Proposition 3.19.54 (Splitting field is minimal)**
*Let $S \subset k[X]$ be a family of polynomials which split completely in $K/k$. Then the following are equivalent*

- *$K$ is generated by the roots of every polynomial $f \in S$, that is*

$$K = k\left(\bigcup_{f\in S} T_{f,K}\right)$$

- *Any subfield $K' \subset K$ in which $S$ splits completely is equal to $K$*

*Proof.* For all $f \in S$ there is a factorisation

$$f = \prod_{\alpha \in T_{f_i,K}} (X - \alpha)$$

Suppose $K = k\left(\bigcup_{f\in S} T_{f,K}\right)$ and let $K'$ be a subfield in which all $f \in S$ split completely. Then by unique factorization in $K[X]$ we have $T_{f,K} \subset K'$ for all $f \in S$ and therefore $K' = K$.

Conversely it's clear that $S$ splits completely in $k\left(\bigcup_{f\in S} T_{f,K}\right)$, therefore by hypothesis this equals $K$. ∎

**Lemma 3.19.55**
*Let $\sigma : K/k \to L/k$ be a morphism and $f(X) \in k[X]$ a polynomial. Then*

- *$\sigma$ induces an injective map on the roots $T_{f,K} \to T_{f,L}$*
- *$f$ splits completely in $K$ iff $f$ splits completely in $\sigma(K)$. In this case the above map is a bijection*

**Proposition 3.19.56 (Image of a splitting field is fixed)**
*Let $K/k$ be a splitting field for $S$ and $\sigma : K/k \to L/k$ a morphism. Then $S$ splits completely in $L$. Any such $\sigma$ satisfies*

$$\sigma(K) = k\left(\bigcup_{f\in S} T_{f,L}\right)$$

*Proof.* Clearly by (3.19.55) $S$ splits completely in $L$.

By the same result $\sigma$ induces a bijection $T_{f,K} \longleftrightarrow T_{f,L}$. Therefore $\sigma(K) = \sigma\left(k\left(\bigcup_{f\in S} T_{f,K}\right)\right) = k\left(\sigma\left(\bigcup_{f\in S} T_{f,K}\right)\right) = k\left(\bigcup_{f\in S} T_{f,L}\right)$ by (3.19.30). ∎

**Proposition 3.19.57 (Uniqueness of Splitting Fields)**
*Let $S \subset k[X]$ be a family of polynomials. Let $K/k$ be a splitting field for $S$ and $L/k$ an extension in which $S$ splits completely.*

*Then there exists a morphism $\sigma : K/k \to L/k$. Let $\alpha \in K$ and $\beta \in L$ be conjugate elements, then we may choose $\sigma$ such that $\sigma(\alpha) = \beta$.*

*Furthermore any two splitting fields are isomorphic.*

© David Rufino CC BY-NC-ND 4.0

*Proof.* By assumption $K$ is generated by the roots $\alpha_{ij}$ of $f_i \in S$. For each $\alpha_{ij}$ we therefore have $m_{\alpha_{ij},k}(X) \mid f_i(X)$ and $m_{\alpha_{ij},k}(X)$ splits completely in $L$ by (3.19.9). Therefore the morphism $\sigma : K/k \to L/k$ exists by (3.19.50).

Note by (3.19.56) $S$ splits in $\sigma(K) = k\left(\bigcup_{f\in S} T_{f,L}\right)$. If $L$ is also a splitting field for $S$ then $L = \sigma(K)$ by (3.19.54) and therefore $\sigma$ is an isomorphism as required. ∎

**Proposition 3.19.58 (Algebraically Closed)**
*A field $M$ is algebraically closed if one of the following equivalent conditions holds*

- *Every algebraic extension $M'/M$ is trivial*
- *Every non-constant polynomial in $M[X]$ has a root in $M$*
- *Every non-constant polynomial in $M[X]$ splits in $M$*

*NB in this case $M$ is also normal.*

**Definition 3.19.59 (Algebraic Closure)**
*An **algebraic closure** $\overline{k}$ of $k$ is a field extension $\overline{k}/k$ which is algebraic and for which $\overline{k}$ is algebraically closed.*

**Proposition 3.19.60 (Existence of Algebraic Closure)**
*Given a field $k$ there exists an algebraic closure $\overline{k}/k$*

**Proposition 3.19.61 (Algebraic extensions embed into Algebraic Closure)**
*Let $K/k$ be an algebraic extension and $M/k$ be field such that every polynomial $f \in k[X]$ splits completely then there exists a morphism $\sigma : K/k \to M/k$.*

*In particular this holds when $M$ is algebraically closed.*

*Proof.* A straightforward application of (3.19.50) since every $m_{\alpha,k}(X)$ splits in $M$. ∎

**Corollary 3.19.62 (Uniqueness of algebraic closure)**
*An algebraic closure $\overline{k}$ of $k$ is unique up to (non-unique) isomorphism.*

More generally we may show the existence of smaller splitting fields

**Proposition 3.19.63 (Existence of Splitting Field)**
*Given a field $k$ and family of polynomials $S \subset k[X]$ then there exists a splitting field $K$.*

*When $S = \{f\}$ then this can be chosen such that $[K : k] \leq n!$ where $n = \deg(f)$.*

*Proof.* We may take the subfield of $\overline{k}$ generated by the roots of polynomials in $S$.

In the case $S$ is finite it is possible to avoid the use of $\overline{k}$. First reduce to the case of a single polynomial $S = \{f\}$ and proceed by induction on $\deg(f)$. The inductive step may be demonstrated using (3.19.40). ∎

**Remark 3.19.64**
*Note if $K/k$ is an algebraic extension then (3.19.61) shows that we may construct an embedding $K \to \overline{k}$ commuting with $k \to \overline{k}$.*

*In general given a tower of algebraic extensions*

$$K = k_n/.../k_0 = k$$

*we will assume the existence of compatible embeddings $i_{k_i} : k_i \to \overline{k}$ such that $i_{k_{i+1}} \circ i_{k_i,k_{i+1}} = i_{k_i}$.*

## 3.19.7 Normal Extensions

Recall that an algebraic extension $K/k$ is normal if all minimal polynomials split completely. They are in some sense "closed". Furthermore $\overline{k}/k$ is clearly normal and results about $\overline{k}$ can often be generalized to normal fields $L/k$. We also show that an extension is normal iff it is a splitting field.

**Lemma 3.19.65**
*Let $L/K/k$ be a tower of algebraic extensions and $\alpha \in L$. If $m_{\alpha,k}(X)$ splits completely in $L$ so does $m_{\alpha,K}(X)$. In particular*

$$L/k \text{ normal} \implies L/K \text{ normal}$$

*Proof.* Note $m_{\alpha,K}(X) \mid m_{\alpha,k}^{i_{kK}}(X)$ as elements of $K[X]$ by (3.19.36). Apply $i_{KL}$ and then we may use (3.19.9). ∎

© David Rufino CC BY-NC-ND 4.0

**Proposition 3.19.66 (Conjugate elements in Normal Extensions)**
*Let $L/k$ be a* normal *extension (e.g. $L = \overline{k}$) and $\alpha, \beta \in L$ elements with the same minimal polynomial $m_\alpha(X) = m_\beta(X)$. Then there exists $\sigma \in \mathrm{Aut}(L/k)$ such that*

$$\sigma(\alpha) = \beta$$

*Proof.* Apply (3.19.50) with $K = L$. ∎

**Proposition 3.19.67 (Normal Criteria)**
*Let $L/K/k$ be a tower of extensions such that $L/k$ is normal (e.g. $L = \overline{k}$). Then the following are equivalent*

- ***NOR1**: For any $\sigma \in \mathrm{Mor}_k(K, L)$ we have $\sigma(K) = i_{KL}(K)$.*
- ***NOR2**: $K/k$ is the splitting field of some family of polynomials $f_i \in k[X]$.*
- ***NOR3**: $K/k$ is normal*

*Proof.* Clearly $3 \implies 2$, for $K$ is the splitting field of all the minimal polynomials of elements in $K$.

$2 \implies 1$. This is (3.19.56).

$1 \implies 3$. Consider any $\alpha \in K$ with minimal polynomial $m_{\alpha,k}(X)$. By definition $m_{\alpha,k}(X)$ splits completely in $L$ because it has a root $\alpha_1 = i_{KL}(\alpha)$. Denote the roots by $\alpha_1, ..., \alpha_r$. By (3.19.66) there is $\sigma_j \in \mathrm{Aut}(L/k)$ such that $\sigma_j(\alpha_1) = \alpha_j$. By hypothesis we have $\alpha_j \in (\sigma_j \circ i_{KL})(K) = i_{KL}(K)$ whence there exists $\alpha_j' \in K$ such that $i_{KL}(\alpha_j') = \alpha_j$. By (3.19.55) $m_{\alpha,k}(X)$ splits completely in $K$. Therefore $K/k$ is normal as required. ∎

**Corollary 3.19.68 (Splitting fields are normal)**
*An algebraic extension $K/k$ is normal if and only if it is a splitting field.*

*Proof.* We may apply the previous Proposition with $L = \overline{k}$.

We may prove a splitting field is normal more directly (without recourse to $\overline{k}$ or Zorn's Lemma in the finite case). Suppose $K/k$ is a splitting field for $S \subset k[X]$. Consider $\alpha \in K$ with minimal polynomial $m_{\alpha,k}(X)$. Let $(L/K, i_{KL})$ be a splitting field for $m_{\alpha,k}(X)$ (as a polynomial in $K[X]$, NB may not be irreducible).

Let $\beta \in L$ be another root of $m_{\alpha,k}(X)$. Observe that $S$ splits in $L$, so by (3.19.57) there is a morphism $\sigma : K/k \to L/k$ with $\sigma(\alpha) = \beta$. By (3.19.56) we have $i_{KL}(K) = \sigma(K)$ whence $\beta \in i_{KL}(K)$. As $\beta$ was an arbitrary root of $m_{\alpha,k}(X)$ we see it splits completely in $i_{KL}(K)$. Finally by (3.19.55) $m_{\alpha,k}(X)$ splits completely in $K$. As $\alpha$ was arbitrary then $K/k$ is normal. ∎

**Proposition 3.19.69 (Extension to normal overfield)**
*Let $L/K/k$ be a tower of algebraic field extensions with $L/k$ normal (e.g. $L = \overline{k}$) then there is a canonical surjection*

$$\begin{aligned} \mathrm{Mor}_k(i_{KL}, L) : \mathrm{Aut}(L/k) &\to \mathrm{Mor}_k(K, L) \\ \sigma &\to \sigma \circ i_{KL} \end{aligned}$$

*When $i_{KL}$ is inclusion then this is simply the restriction to $K$. The kernel is precisely $\mathrm{Aut}(L/K)$.*

*Proof.* Given $\tilde{\sigma} \in \mathrm{Mor}_k(K, L)$, apply (3.19.50) to construct a morphism $\sigma : (L/K, i_{KL}) \to (L/K, \tilde{\sigma})$. The hypotheses apply because the minimal polynomial $m_{\alpha,K}(X)$ with respect to either extension divides the minimal polynomial $m_{\alpha,k}^{i_{kK}}(X)$ which by assumption splits completely in $L$. By (3.19.35) it is an automorphism. Furthermore

$$\sigma \circ i_{kL} = \sigma \circ i_{KL} \circ i_{kK} = \tilde{\sigma} \circ i_{kK} = i_{kL}$$

whence $\sigma \in \mathrm{Aut}(L/k)$ as required. ∎

**Corollary 3.19.70 (Lifting inside normal overfield)**
*Let $L/K/F/k$ be a tower of field extensions with $L/k$ normal, then there is a surjection*

$$\mathrm{Mor}_k(i_{FK}, L) : \mathrm{Mor}_k(K, L) \to \mathrm{Mor}_k(F, L)$$

*Proof.* Note that $\mathrm{Mor}_k(i_{FK}, L) \circ \mathrm{Mor}_k(i_{KL}, L) = \mathrm{Mor}_k(i_{FL}, L)$. By (3.19.69) this composition is surjective, whence the result follows. ∎

**Corollary 3.19.71 (Quotient of automorphism group)**
*Let $L/K/k$ be a tower of extensions such that $L/k$ and $K/k$ are normal. Then $L/K$ is normal and there is an isomorphism of groups.*

© David Rufino CC BY-NC-ND 4.0

$$\begin{aligned} \operatorname{Aut}(L/k)/\operatorname{Aut}(L/K) &\longrightarrow \operatorname{Aut}(K/k) \\ \sigma \to\ & i_{KL}^{-1} \circ \sigma \circ i_{KL} \end{aligned}$$

*Proof.* The given map is well-defined by (3.19.67) since $(\sigma \circ i_{KL})(K) = i_{KL}(K)$, and $\sigma$ fixes $K$ precisely when $\sigma \circ i_{KL} = i_{KL}$, that is $\sigma \in \operatorname{Aut}(L/K)$. Given $\tau \in \operatorname{Aut}(K/k)$, by (3.19.69) there exists $\sigma \in \operatorname{Aut}(L/k)$ such that $\sigma \circ i_{KL} = i_{KL} \circ \tau$. This shows the given map is surjective. The result follows from the group isomorphism theorem. ∎

**Definition 3.19.72 (Normal Closure)**
*Let $K/k$ be an algebraic extension. Then an algebraic extension $L/K$ is a **normal closure** for $K/k$ if*

- *$L/k$ is normal*
- *No proper subfield $i_{KL}(K) \subseteq L' \subsetneq L$ is normal over $k$*

**Proposition 3.19.73 (Existence and Uniqueness of Normal Closure)**
*Let $K/k$ be an algebraic extension. Then a normal closure $L/K$ exists and is unique up to isomorphism. Indeed it is the splitting field for all the minimal polynomials $\{m_{\alpha,k}(X) \mid \alpha \in K\}$ over $k$.*

*Furthermore if $K/k$ is finite then so is $L/k$*

*Proof.* Suppose $K = k(\{\alpha_i\}_{i\in I})$. Let $L/k$ be the splitting field for $S = \{m_{\alpha_i,k}(X)\}_{i\in I}$. By (3.19.68) $L/k$ is normal and by (3.19.50) there is a morphism $\sigma : K/k \to L/k$, so we may consider it as an extension $(L/K, \sigma)$. Suppose $i_{KL}(K) \subset L' \subset L$ is normal. As $i_{KL}(\alpha_i) \in L'$, by definition $S$ splits in $L'$ and therefore $L' = L$ by (3.19.54). Therefore $L/K$ is a normal closure as required.

If $K/k$ is finite then we may choose $I$ to be finite, and therefore $L/k$ is finite.

Let $L/K$ be an arbitrary normal closure, then we claim $L/k$ is a splitting field for $S' := \{m_{\alpha,k}(X) \mid \alpha \in K\}$. Clearly $S'$ splits in $L/k$, because each has a root in $L$. Let $L'/k$ be the subfield generated by roots of $S'$. Then it is a splitting field and therefore normal by (3.19.68). By assumption $L' = L$ and therefore $L/k$ is the splitting field for $S'$. Uniqueness follows from the uniqueness of splitting fields (3.19.57). ∎

## 3.19.8 Separability (Algebraic Case)

We follow Lang and not only characterize separability but define a "separability degree" which equals the extension degree if and only if it's separable. The proofs are somewhat technical, especially in light of the fact most base fields will be perfect.

**Definition 3.19.74 (Separable Element)**
*We say $\alpha \in K/k$ is **separable** over $k$ if it is algebraic and $m_{\alpha,k}(X)$ is a separable polynomial.*

*We say $K/k$ is **separable algebraic** (or just separable if $K/k$ is assumed to be algebraic) if every $\alpha \in K$ is **separable**.*

**Lemma 3.19.75**
*$\alpha \in K/k$ is separable if and only if it is a root of a separable polynomial in $k[X]$.*

*In particular $\alpha$ separable over $k$ implies it is separable over any subfield $i_{kK}(k) \subset E \subset K$.*

*Proof.* One direction is obvious. Conversely suppose $f(\alpha) = 0$ with $f$ separable. Then $m_{\alpha,k} \mid f$ so the result follows from (3.17.40). ∎

**Proposition 3.19.76 (Separability Degree)**
*Let $K/k$ be an algebraic extension and $L/K$ an extension such that $L/k$ is normal (e.g. $L = \overline{k}$ or $L$ is a normal closure). Then define the **separability degree***

$$[K : k]_s := \#\operatorname{Mor}_k(K, L)$$

*This is independent of the choice of $L/K$.*

*Proof.* Given such an $L$, let $L'/K$ be the intersection of all subfields of $L/K$ normal over $k$. This is a normal closure of $K/k$. Let $\sigma \in \operatorname{Mor}_k(K, L)$ and $\alpha \in K$. Then $\sigma(\alpha)$ is a root of $m_{\alpha,k}(X)$ along with $i_{KL}(\alpha) \in L'$. As $L'$ is normal we have $\sigma(\alpha) \in L'$. Therefore without loss of generality we may replace $L$ with $L'$. As the normal closure is unique up to isomorphism the degree is well-defined. ∎

First we prove a key lemma regarding simple extension

**Lemma 3.19.77 (Separability Degree of Simple Extension)**
*If $k(\alpha)/k$ is a simple extension and $L/k$ normal overfield then*

© David Rufino CC BY-NC-ND 4.0

$$[k(\alpha):k]_s = \#\{\text{roots of } \ m_\alpha \ \text{ in } \ L\} \leq \deg(m_\alpha) = [k(\alpha):k]$$

*Furthermore equality holds iff $\alpha$ is separable over $k$.*

*Proof.* The first equality follows from (3.19.49), the final equality from (3.19.42). The inequality follows from (3.17.35). The final statement follows from (3.19.11). ∎

The main results of this section are the following

**Proposition 3.19.78 (Separability Degree)**
*Let $L/K/F/k$ be a tower of finite extensions with $L/k$ normal. Then the restriction map*

$$\mathrm{Mor}_k(i_{FK}, L) : \mathrm{Mor}_k(K, L) \to \mathrm{Mor}_k(F, L)$$

*is surjective with finite fibers of order $[K:F]_s$. In particular we have*

$$[K:k]_s = [K:F]_s[F:k]_s$$

*Further $[K:k]_s \leq [K:k]$ with equality if and only if $K/k$ is separable.*

*Proof.* For a tower $L/K/F/k$ with $L/k$ normal, consider the restriction map

$$\psi := \mathrm{Mor}_k(i_{FK}, L) : \mathrm{Mor}_k(K, L) \to \mathrm{Mor}_k(F, L)$$

It is surjective by (3.19.70). Consider $\sigma \in \mathrm{Mor}_k(F, L)$ and the fibre $\psi^{-1}(\sigma) = \mathrm{Mor}_F(K, (L/F, \sigma))$. This has order equal to $\#\psi^{-1}(\sigma) = [K:F]_s$ for all $\sigma$, because by (3.19.76) this does not depend on the embedding $i_{FL}$. As $\mathrm{Mor}_k(K, L)$ is equal to the disjoint union of all the fibres, then the multiplicativity result follows.

It is possible to decompose $K/k$ as a tower of simple extensions

$$K = K_n/.../K_0 = k$$

with $K_i = K_{i-1}(\alpha_i)$. By (3.19.77) we have

$$[K_i : K_{i-1}]_s \leq [K_i : K_{i-1}]$$

with equality iff $\alpha_i$ separable over $K_{i-1}$. By multiplicativity the inequality follows.

If $K/k$ is separable then by (3.19.75) $\alpha_i$ is separable over $K_{i-1}$ and we have $[K_i : K_{i-1}]_s = [K_i : K_{i-1}]$ and $[K:k]_s = [K:k]$ by multiplicativity. Conversely if $[K:k]_s = [K:k]$ then $[K_i : K_{i-1}]_s = [K_i : K_{i-1}]$ and $\alpha_i$ is separable over $K_{i-1}$. Since the choice of $\alpha_1$ was arbitrary we see that $K/k$ is separable. ∎

**Proposition 3.19.79 (Towers of Separable Extensions)**
*Consider a tower of algebraic extensions $K/E/k$. Then $K/E$ and $E/k$ is separable iff $K/k$ is.*

*Proof.* $K/k$ separable $\implies$ $K/E$ and $E/k$ separable follows from (3.19.75).

Conversely the finite case follows from (3.19.78) by multiplicativity. For the general case, consider $\alpha \in K$. Then (3.19.46) shows the existence of a finite subextension $E_0/k$ of $E$ such that $m_{\alpha,E} = m_{\alpha,E_0}$. Therefore $\alpha$ is separable over $E_0$. By (3.19.77) we see that $[E_0(\alpha) : E_0]_s = [E_0(\alpha) : E_0]$. As $E/k$ is separable a-fortiori $E_0/k$ is separable so by (3.19.78) $[E_0 : k]_s = [E_0 : k]$. By multiplicativity $[E_0(\alpha) : k]_s = [E_0(\alpha) : k]$ and the same result again shows that $E_0(\alpha)/k$ is separable. In particular $\alpha$ is separable over $k$ as required. ∎

**Proposition 3.19.80**
*An algebraic extension $K = k(\alpha_1, ..., \alpha_n)/k$ is separable iff $\alpha_i$ are.*

*Proof.* Let $K = k(\alpha_1, ..., \alpha_n)$. Then we may construct a tower of finite (simple) extensions

$$K = K_n/.../K_0 = k$$

with $K_i = k(\alpha_1, ..., \alpha_i)$ and $K_i = K_{i-1}(\alpha_i)$. By (3.19.75) $\alpha_i$ is separable over $K_{i-1}$. Therefore $[K_i : K_{i-1}]_s = [K_i : K_{i-1}]$ by (3.19.77) and $[K:k]_s = [K:k]$ by multiplicativity. (3.19.78) shows that $K/k$ is separable. ∎

**Proposition 3.19.81**
*Let $K/k$ be a splitting field for finitely many separable polynomials $f_i$. Then $K/k$ is finite and Galois.*

*Proof.* By definition $K$ is generated by the (finitely) many roots $\alpha_{ij}$ and so is finite by (3.19.44) and normal by (3.19.68). By (3.19.75) each $\alpha_{ij}$ is separable and so by (3.19.80) $K/k$ is separable. ∎

**Proposition 3.19.82**
*Let $K/k$ be a finite separable extension and $L/k$ a normal closure. Then $L/k$ is separable.*

© David Rufino CC BY-NC-ND 4.0

*Proof.* By (3.19.73) $L/k$ is the splitting field of finitely many minimal polynomials $m_{\alpha_i,k}(X)$ for $\alpha_i \in K$. By (3.19.80) the $\alpha_i$ are separable and therefore so are $m_{\alpha_i,k}(X)$. By (3.19.81) we see that $L/k$ is separable. ∎

**Proposition 3.19.83 (Equivalent Definition of Separability)**
*Let $K/k$ be an algebraic extension. TFAE*

*a) $K/k$ is separable*
*b) $E/k$ is separable for every finite subextension*
*c) $[E:k]_s = [E:k]$ for every finite subextension*

*Proof.* $a) \implies b)$ is trivial and $b) \iff c)$ follows from (3.19.78). We need only show $b) \implies a)$.

Consider $\alpha \in K$. Then by (3.19.46) there exists a finite subextension $E/k$ such that $\alpha$ is algebraic over $E$. Therefore $E(\alpha)/k$ is finite, and by assumption $E(\alpha)/k$ separable as required. ∎

## 3.19.9 Purely Inseparable Extensions and Separable Closure

In what follows we let $p$ be the characteristic exponent of $k$. In other words when $\operatorname{char}(k) = 0$ then $p = 1$ and the statements are trivial.

**Definition 3.19.84 (Purely Inseparable Element)**
*An element $x \in K/k$ is **purely inseparable** (or p-radical) if there exists an integer $n \geq 0$ such that $x^{p^n} \in k$. The **height** of $x$ is the least such integer.*

*We say an extension $K/k$ is **purely inseparable** if every $x \in K/k$ is purely inseparable.*

*Further $K/k$ is **purely inseparable of height $\leq n$** if additionally every $x \in K$ has height at most $n$.*

**Lemma 3.19.85**
*Let $a \in k$. For every integer $e \geq 0$ the polynomial $f(X) := X^{p^e} - a$ has at most one root in any extension field $K/k$.*

*Proof.* Suppose that $b$ is a root, then by (3.19.6) we have $f(X) = (X-b)^{p^e}$. Then evidently $b$ is the unique root. ∎

**Lemma 3.19.86**
*Let $a \in k \setminus k^p$. Then for every integer $e \geq 0$ the polynomial $X^{p^e} - a$ is irreducible in $k[X]$.*

*Proof.* Let $K/k$ be a splitting field for $f(X) = X^{p^e} - a$, $b \in K$ a root and $g(X)$ be the minimal polynomial of $b$. By (3.19.6) $f(X) = (X-b)^{p^e}$ in $K[X]$. Suppose $\pi$ is a monic irreducible factor of $f$ in $k[X]$ then by unique factorisation in $K[X]$ we have $\pi(X) = (X-b)^r$ for some $r$. In particular $\pi$ has $b$ as a root and therefore is divisible by $g$. By irreducibility we conclude every irreducible factor is equal to $g$ and the irreducible factorization of $f$ is $g^s$. Furthermore $p^e = rs$. Consequently both $r$ and $s$ are $p$th powers, and we have

$$\pi = (X-b)^{p^d}$$
$$f = \pi^{p^{e-d}}$$

for some $0 \leq d \leq e$. In particular $b^{p^d} \in k$ and $a = b^{p^{e-d}}$. By assumption $a$ is not a $p$-th power and so we must have $d = e$, and $f = \pi$ is irreducible. ∎

**Proposition 3.19.87**
*Let $x \in K/k$ be purely inseparable of height $e$. Then the minimal polynomial of $x$ is $X^{p^e} - x^{p^e}$. Furthermore*

$$[k(x):k] = p^e$$
$$[k(x):k]_s = 1$$

*More precisely for every field $L/k$ there is at most one morphism $k(x)/k \to L/k$.*

*Proof.* By definition $x^{p^e} \in k$ is not a $p$-th power. Therefore the given polynomial is irreducible by (3.19.86) and therefore is the minimal polynomial by uniqueness. The first relation follows from (3.19.42) and the second from (3.19.77). ∎

**Corollary 3.19.88**
*Let $x \in K/k$. Then the following are equivalent*

*a) $x$ is purely inseparable*
*b) $m_x(X) = (X-x)^r$ for some $r$*

*In this case we must have $r = p^n$ where $n$ is the height of $x$.*

© David Rufino CC BY-NC-ND 4.0

**Corollary 3.19.89**
*Suppose $x \in K/k$ is both separable and purely inseparable. Then $x \in k$.*

*Proof.* The minimal polynomial is $X^{p^e} - x^{p^e} = (X-x)^{p^e}$. Therefore by (3.19.11) $e = 0$ which means precisely $x \in k$. ∎

We first establish the following two results, which will be used below.

**Lemma 3.19.90 (Inseparable Polynomials)**
*Let $f \in k[X]$ be a non-constant polynomial and $p$ the characteristic exponent of $k$. Then $f' = 0 \iff p > 1$ and $f \in k[X^p]$.*

**Proposition 3.19.91 (Separable Irreducible Polynomials)**
*Let $f \in k[X]$ be an irreducible polynomial and $p$ the characteristic exponent of $k$. Then the following are equivalent*

*a) $f$ is separable (i.e. $f$ and $f'$ are co-maximal)*
*b) $f' \neq 0$*
*c) $p = 1$ or $f \notin k[X^p]$*

*Proof.* Note by assumption $f$ is not a unit and therefore not constant.

$a) \implies b)$ Suppose $f' = 0$ then $(f, f') = (f) \neq k[X]$ and therefore $f$ is not separable.

$b) \implies a)$ Suppose $f$ is not separable. By (...) $(f)$ is maximal and so we must have $f' \in (f)$. As $\deg(f') < \deg(f)$ this implies $f' = 0$.

$b) \iff c)$ This is just the contrapositive of (3.19.90). ∎

**Proposition 3.19.92 (Relative Separable Closure)**
*Let $K/k$ be a field extension. The subextension $K_s/k \subset K/k$ of separable elements form a separable algebraic subfield. Furthermore if $K/k$ is algebraic then $K/K_s$ is purely inseparable and the restriction map*

$$\mathrm{Mor}_k(K, L) \to \mathrm{Mor}_k(K_s, L)$$

*is bijective for every normal extension $L/K$. In particular when $K_s/k$ is finite then we have the relation*

$$[K_s : k] = [K_s : k]_s = [K : k]_s$$

*Proof.* Suppose $\alpha, \beta \in K$ are separable algebraic then the field $k(\alpha, \beta)$ is separable algebraic by (3.19.80). As this contains $\alpha \pm \beta$ and $\alpha\beta$ then $K_s$ is a subfield which is by definition separable. Then by (3.19.78) we have $[K_s : k] = [K_s : k]_s$ when this is finite.

For $x \in K$ let $f(X)$ be the minimal polynomial over $k$. There exists some integer $m \geq 0$ such that $f(X) \in k[X^{p^m}]$ but not in $k\left[X^{p^{m+1}}\right]$. In other words $f(X) = g(X^{p^m})$ and $g(X) \notin k[X^p]$. As $f$ is irreducible so is $g$, and is therefore the minimal polynomial of $x^{p^m}$. By (3.19.91) $g$ is separable and therefore $x^{p^m} \in K_s$. This shows that $K/K_s$ is purely inseparable.

Let $L/K/k$ be tower such that $L/k$ is normal. Consider the mapping

$$\mathrm{Mor}_k(K, L) \to \mathrm{Mor}_k(K_s, L)$$

obtained by restriction. By (3.19.70) it is surjective. Consider $\psi : K_s \to L$ and $\hat{\psi} : K \to L$ an extension. For every $x \in K$ we have $\hat{\psi}|_{K_s(x)} = i_{K_s(x)L}$ by (3.19.87). This shows that $\hat{\psi}$ is unique and the mapping is bijective. ∎

**Definition 3.19.93 (Degree of Inseparability)**
*Let $K/k$ be a finite extension. Then in light of (3.19.92) we may define the **degree of inseparability** to be the integer*

$$[K : k]_i := \frac{[K : k]}{[K : k]_s}$$

*By (3.19.16) and (3.19.78) it is multiplicative in the sense that for a tower of finite extensions*

$$[L : k]_i = [L : K]_i [K : k]_i$$

*Further by (3.19.78) $K/k$ is separable if and only if $[K : k]_i = 1$.*

We may refine this further by showing that $[K : k]_i$ is always a power of the characteristic exponent.

© David Rufino CC BY-NC-ND 4.0

**Proposition 3.19.94**
*Let $k(\alpha)/k$ be a simple algebraic extension with characteristic exponent $p$. There exists a unique integer $r \geq 0$ such that every root of $m_{\alpha,k}(X)$ has multiplicity $p^r$ (in any splitting field).*

*Further*

$$[k(\alpha) : k]_i = p^r$$

*and $m_{\alpha,k}(X)$ has $[k(\alpha) : k]_s$ distinct roots.*

*Proof.* Let $L/k(\alpha)/k$ be a normal overfield, and let $\alpha_1, ..., \alpha_s$ be the distinct roots of $m_{\alpha,k}(X)$ with $\alpha_1 = \alpha$. Then we have a factorisation in $L[X]$ given by

$$m_{\alpha,k}(X) = \prod_{i=1}^{s} (X - \alpha_i)^{m_i}$$

for some integers $m_i > 0$.

For every $1 \leq i \leq n$ there exists an automorphism $\sigma_i \in \mathrm{Aut}(L/k)$ such that $\sigma_i(\alpha) = \alpha_i$ (3.19.50). By definition $\sigma_i$ fixes $m_{\alpha,k}(X)$ so by unique factorisation we deduce that $m_i = m_1$, so all the roots have the same multiplicity $m$.

If $m = 1$ then by (3.17.39) $m_{\alpha,k}(X)$ is separable. Otherwise by (3.19.91) there is a polynomial $g(X)$ such that $f(X) = g(X^p)$ which has $\alpha^p$ as a root. We may continue inductively to find an integer $r \geq 0$ such that $f(X) = g(X^{p^r})$ and $g(X)$ is separable. Then $\alpha^{p^r}$ is separable. By definition $\alpha$ is purely inseparable over $k(\alpha^{p^r})$ so by (3.19.87)

$$\left[k(\alpha) : k(\alpha^{p^r})\right]_i = p^r$$

By (3.19.78) we have

$$\left[k(\alpha^{p^r}) : k\right]_i = 1$$

and by multiplicativity $[k(\alpha) : k]_i = p^r$. By (3.19.77) the number of distinct roots of $m_{\alpha,k}(X)$ is $[k(\alpha) : k]_s$. Comparing degrees we have $\deg\big(m_{\alpha,k}(X)\big) = [k(\alpha) : k] = [k(\alpha) : k]_s \cdot m$ whence $m = [k(\alpha) : k]_i = p^r$ as required. ∎

**Proposition 3.19.95**
*Let $K/k$ be a finite field extension with characteristic exponent $p$. Then $[K : k]_i = p^r$ for some integer $r \geq 0$.*

*$K/k$ is separable if and only if $p = 1$ or $r = 0$.*

*Proof.* We may decompose $K/k$ into a tower of simple extensions, and the result follows from multiplicativity and (3.19.94). ∎

## 3.19.10 Separable closure

**Proposition 3.19.96 (Relatively Separably Closed)**
*Let $K/k$ be an extension. The following are equivalent*

a) *Every separable $x \in K/k$ lies in $k$*
b) *Every separable algebraic subextension is trivial*
c) *$K_s/k$ is trivial*

*In this case we say $K/k$ is **relatively separably closed**.*

*If $K/k$ is algebraic then this is equivalent to being purely inseparable.*

*Proof.* The final statement follows from (3.19.92). ∎

**Definition 3.19.97**
*We say a field $k$ is **separably closed** if every separable algebraic extension is trivial.*

*We say an extension $k^{\mathrm{sep}}/k$ is a **separable closure** if it is separable algebraic and separably closed.*

**Proposition 3.19.98**
*Let $\overline{k}/k$ be an algebraic closure, then the subextension $\overline{k}_s/k$ is a separable closure for $k$.*

*Furthermore every separable closure is isomorphic to $\overline{k}_s/k$ and $\overline{k}/\overline{k}_s$ is purely inseparable.*

*Proof.* By (3.19.92) $\overline{k}_s/k$ is separable and algebraic. Suppose $K/\overline{k}_s$ is separable and algebraic then by (...) there exists an embedding $i : K/\overline{k}_s \to \overline{k}/\overline{k}_s$. As $\overline{k}_s$ is relatively separably closed in $\overline{k}$ we see that $i(K) = \overline{k}_s$ and therefore $K = \overline{k}_s$.

© David Rufino CC BY-NC-ND 4.0

Let $k^{\text{sep}}/k$ be a separable closure, then by (3.19.61) there is an embedding $\varphi : k^{\text{sep}}/k \to \overline{k}/k$. Then by definition $\varphi(k^{\text{sep}}) \subset \overline{k}_s$, so this defines an extension $\overline{k}_s/k^{\text{sep}}$ which is by definition separable. By assumption it is trivial which means precisely $\varphi(k^{\text{sep}}) = \overline{k}_s$.

It follows from (3.19.92) that $\overline{k}/\overline{k}_s$ is purely inseparable. ∎

**Proposition 3.19.99**
*Let $k^{\text{sep}}/k$ be a separable closure and $k'/k$ a separable algebraic extension. Then there exists a morphism*

$$k'/k \to k^{\text{sep}}/k$$

*Proof.* By (3.19.98) we may identify $k^{\text{sep}}/k$ with $\overline{k}_s/k$. By (3.19.61) there exists a morphism $k'/k \to \overline{k}/k$ which by definition has image in $\overline{k}_s/k$. ∎

## 3.19.11 Perfect Fields

**Proposition 3.19.100 (Perfect Closure)**
*Let $k$ be a field with characteristic exponent $p$ contained in an algebraically closed field $\Omega$ (e.g. $\overline{k}$). Define*

$$k^{p^{-n}} := \{x \in \Omega \mid x^{p^n} \in k\}$$

*Then $k^{p^{-n}}$ is a subfield of $\Omega/k$, which is purely inseparable of height $\leq n$. Furthermore it is a splitting field for the family of polynomials $\{X^{p^r} - a \mid a \in k, r \leq n\}$ which characterizes it up to isomorphism. Furthermore*

$$k^{p^{-\infty}} := \{x \in \Omega \mid x^{p^n} \in k \text{ some } n \geq 1\} = \bigcup_{n \geq 1} k^{p^{-n}}$$

*is the smallest perfect subfield of $\Omega$ containing $k$ and is also a splitting field for the family of polynomials $\{X^{p^n} - a \mid a \in k, n \in \mathbb{N}\}$.*

*Proof.* Suppose $\alpha, \beta \in k^{p^{-n}}$ then applying the Frobenius homomorphism (...) we see $(\alpha + \beta)^{p^n} = \alpha^{p^n} + \beta^{p^n}$ and so $k^{p^{-n}}$ is an additive subgroup. Furthermore $(\alpha\beta)^{p^n} = \alpha^{p^n}\beta^{p^n}$ whence it is a multiplicative subgroup as this demonstrates $(\alpha^{-1})^{p^n} = (\alpha^{p^n})^{-1} \in k$. Therefore it is a subfield which is by definition purely inseparable of height $\leq n$.

By definition $k^{p^{-n}}$ is precisely the set of roots of the given polynomials and it is therefore the corresponding splitting field by (...). This also shows that $n \leq m \implies k^{p^{-n}} \subseteq k^{p^{-m}}$ which shows that the union $k^{p^{-\infty}}$ is a subfield of $\Omega$.

To demonstrate it's perfect we need to show every element $x \in k^{p^{-n}}$ has a $p$-th root. By definition $x^{p^n} \in k$ and there exists a root $\alpha$ to the polynomial $X^{p^{n+1}} - x^{p^n}$ in $k^{p^{-(n+1)}}$. Therefore $(\alpha^p - x)^{p^n} = 0$ whence $\alpha^p = x$ as required.

Let $k'$ be another perfect subfield containing $k$. We claim by induction that $k^{p^{-n}} \subseteq k'$. For given $\alpha \in k^{p^{-(n+1)}}$ by definition $\alpha^p \in k^{p^{-n}}$ whence $\alpha^p \in k'$ by the inductive hypothesis, and by assumption $\alpha \in k'$. Therefore $k^{p^{-\infty}}$ is the smallest perfect subfield. Furthermore by definition it is the set of roots of the given family of polynomials and therefore the corresponding splitting field. ∎

**Proposition 3.19.101**
*Let $k'/k$ be a purely inseparable extension of height $\leq n$, then there exists an embedding*

$$k'/k \to k^{p^{-n}}/k$$

*If $k'/k$ is purely inseparable and $K/k$ is perfect then there also exists an embedding*

$$k'/k \to K/k$$

*Proof.* For the first part let $\alpha \in k'$ then by (3.19.87) the minimal polynomial of $\alpha$ is $g(X) = X^{p^e} - \alpha^{p^e}$ for $e \leq n$ which splits completely in $k^{p^{-n}}$ by (3.19.100). Then the embedding exists by (3.19.50).

Suppose $K/k$ is perfect then we claim that $X^{p^n} - a$ splits completely for every $a \in K$. By induction there is $b \in K$ such that $b^{p^n} = a$. Then $X^{p^n} - a = (X - b)^{p^n}$ splits completely. In particular for every $\alpha \in k'$ the minimal polynomial splits completely in $K/k$ and the embedding follows from (3.19.50). ∎

Recall a perfect field $k$ satisfies $k^p = k$ where $p$ is the characteristic exponent. We show that in this case there are no inseparable algebraic extensions. First we show that all finite fields are perfect.

**Proposition 3.19.102 (Finite fields are perfect)**
*Any finite field is perfect.*

*Proof.* The Frobenius homomorphism is injective and therefore surjective by counting. ∎

© David Rufino CC BY-NC-ND 4.0

**Proposition 3.19.103 (Perfect field criteria)**
*Let $k$ be a field with characteristic exponent $p$. Then the following are equivalent*

*a) Every irreducible polynomial in $k[X]$ is separable*
*b) Every algebraic extension $K/k$ is separable*
*c) $\overline{k}/k$ is separable*
*d) $k^p = k$*

*In particular every field of characteristic zero is perfect, and an algebraic extension of a perfect field is separable.*

*Proof.* We prove each in turn

$a) \implies b)$ Minimal polynomials are irreducible by (3.19.36) and therefore are separable by hypothesis.

$b) \iff c)$ One direction is automatic as $\overline{k}$ is algebraic. On the other hand every algebraic extension is isomorphic to a subfield of $\overline{k}$, so the implication follows from (3.19.79).

$b) \implies d)$ We need only show $k^p = k$ in the case $p > 1$. If there exists $a \in k \setminus k^p$ then by (3.19.86) the polynomial $f(X) := X^p - a$ is irreducible. Then it's clear that the field extension $K := k[X]/(X^p - a)$ is not separable. For the minimal polynomial of $\overline{X}$ is $f$ which by (3.19.91) is not separable. This contradicts the assumption that $K/k$ is separable.

$d) \implies a)$ Suppose that $f$ is irreducible but not separable. Then by (3.19.91) $\operatorname{char}(k) = p > 1$ and $f \in k[X^p]$. By assumption all the coefficients are $p$-th powers and therefore $f = h^p$ by (3.19.6) for some $h \in k[X]$. This contradicts irreducibility of $f$, and so $f$ must be separable. ∎

## 3.19.12 Applications of Separability

**Definition 3.19.104 (Bounds on $\operatorname{Aut}(K/k)$)**
*Let $K/k$ be an algebraic extension and $L/K$ an extension such that $L/k$ is normal. Then there is a natural injection*

$$\begin{aligned} \operatorname{Mor}_k(K, i_{KL}) : \operatorname{Aut}(K/k) &\to \operatorname{Mor}_k(K, L) \\ \sigma &\to i_{KL} \circ \sigma \end{aligned}$$

*In particular in the case $[K : k] < \infty$*

$$\#\operatorname{Aut}(K/k) \leq [K : k]_s \leq [K : k] < \infty$$

*If $i_{KL}$ is inclusion, then we may regard $\operatorname{Aut}(K/k)$ as a subset of $\operatorname{Mor}_k(K, L)$*

As an application of the concept of separability degree we prove

**Proposition 3.19.105 (Primitive Element Theorem)**
*Let $K/k$ be a finite separable extension of $k$ then $K = k(\alpha)$ is simple.*

*Proof.* We only prove the case $k$ is infinite. The finite case can be proven separately by showing that the $K^\star$ is cyclic.

Consider the set $\operatorname{Mor}_k\left(K, \overline{k}\right) = \{\sigma_1, ..., \sigma_n\}$ which by (3.19.83) has order $n = [K : k]$. By induction we can assume that $K = k(\alpha, \beta)$. We claim that there exists $0 \neq c \in k$ such that $\sigma_i(\alpha + c\beta)$ are all distinct. In this case we clearly have $\#\operatorname{Mor}_k\left(k(\alpha + c\beta), \overline{k}\right) \geq n$ so by the same result $[k(\alpha + c\beta) : k] \geq [k(\alpha + c\beta) : k]_s \geq n$ whence $k(\alpha + c\beta) = K$ (by (2.3.15)).

We have $\sigma_i(\alpha + c\beta) = \sigma_j(\alpha + c\beta) \iff c\big(\sigma_i(\beta) - \sigma_j(\beta)\big) = \big(\sigma_i(\alpha) - \sigma_j(\alpha)\big)$. Therefore consider the polynomial

$$f(X) = \prod_{i \neq j} \Big(X\big(\sigma_i(\beta) - \sigma_j(\beta)\big) - \big(\sigma_i(\alpha) - \sigma_j(\alpha)\big)\Big)$$

Then the embeddings are distinct precisely when $f(c) \neq 0$. Since $f(X)$ has at most finitely many roots and $k$ is infinite, there must exist such a $c$. ∎

## 3.19.13 Normal Extensions II

We provide some more straightforward criteria based on $[K : k]_s$

**Proposition 3.19.106 (Normal Criteria II)**
*Let $L/K/k$ be a tower of algebraic extensions such that $L/k$ is normal (e.g. $L = \overline{k}$). Then $K/k$ is normal if and only if the embedding*

$$\operatorname{Mor}_k(K, i_{KL}) : \operatorname{Aut}(K/k) \to \operatorname{Mor}_k(K, L)$$

© David Rufino CC BY-NC-ND 4.0

*is a bijection. In particular if $K/k$ is finite, then it is normal if and only if*

$$\#\operatorname{Aut}(K/k) = [K:k]_s$$

*Proof.* Suppose $K/k$ is normal and consider $\sigma \in \operatorname{Mor}_k(K,L)$. Then by **NOR1** $\sigma(K) = i_{KL}(K)$ and we may define $\tau := i_{KL}^{-1} \circ \sigma$ with $\tau \in \operatorname{Aut}(K/k)$. The converse is similar.

For the final part we've already observed (3.19.104) that in the finite case $\#\operatorname{Aut}(K/k) \leq [K:k]_s = \#\operatorname{Mor}_k(K,L) \leq [K:k] < \infty$. Therefore the embedding $\operatorname{Mor}_k(K, i_{KL})$ is a bijection precisely when the orders are the same. ∎

**Corollary 3.19.107 (Galois Criteria)**
*Let $K/k$ be a finite extension. Then*

$$\#\operatorname{Aut}(K/k) \leq [K:k]_s \leq [K:k]$$

*with equalities if and only if $K/k$ is Galois.*

*Proof.* We've seen the inequalities (3.19.104)

$$\#\operatorname{Aut}(K/k) \leq [K:k]_s \leq [K:k] < \infty$$

with equality if and only if $K/k$ is both normal (3.19.106) and separable (3.19.78) ∎

## 3.19.14 Finite Fields

A finite field $K$ necessarily has positive characteristic $p$, and therefore the prime subfield is isomorphic to the field $\mathbb{F}_p := \mathbb{Z}/p\mathbb{Z}$. We list some necessary properties of a finite field

**Proposition 3.19.108 (Properties of finite fields)**
*Every finite field $K$ is a finite-dimensional vector space over its prime subfield $\mathbb{F}_p$. Define $n = \left[K:\mathbb{F}_p\right]$.*

- $\#K = p^n$
- *$K$ is a splitting field for $X^{p^n} - X \in \mathbb{F}_p[X]$, and indeed is equal to the set of roots*
- *The multiplicative group of units $K^\star$ is cyclic.*
- *$K/\mathbb{F}_p$ is simple*

*Proof.* Since $K/\mathbb{F}_p$ is a finite-dimensional vector space it must have order $p^n$.

The group of units has order $p^n - 1$, so by Lagrange's theorem every non-zero element satisfies $X^{p^n-1} - 1 = 0$, so therefore every element satisfies $X^{p^n} - X = 0$. Since this polynomial can have at most $p^n$ roots (3.17.35) the roots are exactly all the elements of $K$.

We note again that $X^d - 1$ has at most $d$ roots by (3.17.35). Therefore the fact $K^\star$ is cyclic follows from (3.2.25). ∎

**Proposition 3.19.109 (Frobenius morphism)**
*Given any field $K/\mathbb{F}_p$ the Frobenius map*

$$\phi : x \to x^p$$

*is an injective field homomorphism. In particular when $K$ is finite (or even algebraic) it is an automorphism over $\mathbb{F}_p$.*

*Proof.* The only non-trivial step is showing

$$(x+y)^p = x^p + y^p$$

which follows from elementary calculations on binomial coefficients.

For the final statement use (3.19.35). ∎

Further we can show existence and uniqueness of finite fields.

**Proposition 3.19.110 (Existence and uniqueness of finite fields)**
*Consider the algebraic closure $\overline{\mathbb{F}_p}$ and let $\mathbb{F}_{p^n}$ denote the splitting field of $f(X) = X^{p^n} - X$ in $\overline{\mathbb{F}_p}$. Then*

- *$\mathbb{F}_{p^n}$ is equal to the set of roots of $X^{p^n} - X$*
- *It is the unique subfield of order $p^n$ and every finite field of order $p^n$ is isomorphic to this.*
- *$\mathbb{F}_{p^m} \subset \mathbb{F}_{p^n} \iff m \mid n$*

*Proof.* By (3.19.109) the set of roots of $f(X)$ forms a subfield of $\overline{\mathbb{F}_p}$.

© David Rufino CC BY-NC-ND 4.0

Furthermore $f'(X) = -1$ so $f(X)$ is separable because clearly $(f, f') = 1$. Therefore by (3.19.11) $f(X)$ has $p^n$ distinct roots and the splitting field of $f(X)$ is exactly the set of roots.

Furthermore every subfield of order $p^n$ must satisfy this polynomial by Lagrange's Theorem (3.2.12), so it is the unique such subfield.

Since every algebraic extension of $\mathbb{F}_p$ is isomorphic to a subfield of $\overline{\mathbb{F}_p}$ it's also the unique algebraic extension of order $p^n$ up to isomorphism.

Clearly if $\mathbb{F}_{p^m} \subset \mathbb{F}_{p^n}$ then $\left[\mathbb{F}_{p^n} : \mathbb{F}_p\right] = \left[\mathbb{F}_{p^n} : \mathbb{F}_{p^m}\right]\left[\mathbb{F}_{p^m} : \mathbb{F}_p\right]$, so we must have $m \mid n$. Conversely if $\alpha \in \mathbb{F}_{p^m}$ then $\alpha^{p^m} = \alpha \implies \alpha^{p^{rm}} = \alpha$ for all $r > 0$, so $\alpha \in \mathbb{F}_{p^n}$. ∎

It is usually most convenient to work in $\overline{\mathbb{F}_p}$ and consider the subfields of order $p^n$. We've seen in (3.19.102) that every finite field $\mathbb{F}_q \coloneqq \mathbb{F}_{p^n}$ is perfect and therefore every algebraic extension is separable (3.19.103). In fact we may show that every finite extension is Galois.

**Proposition 3.19.111**
*The field extension $\mathbb{F}_{p^n}/\mathbb{F}_p$ is Galois with*

$$\mathrm{Gal}\left(\mathbb{F}_{p^n}/\mathbb{F}_p\right) = \langle \phi \rangle$$

*cyclic of order $n$ generated by the Frobenius automorphism.*

*Proof.* Let $G = \mathrm{Aut}\left(\mathbb{F}_{p^n}/\mathbb{F}_p\right)$. We've observed that $\phi \in G$. Let $d = o(\phi)$, then we wish to prove that $n = d$. Certainly Lagrange's Theorem applied to the multiplicative group $\mathbb{F}_{p^n}^\star$ implies $\phi^n = 1$. Therefore $d \mid n$ by (3.2.12) applied to $G$. By definition $\phi^d = e$, so every $\alpha \in \mathbb{F}_{p^n}$ satisfies the polynomial $X^{p^d} - X = 0$. This polynomial has at most $p^d$ roots (3.17.35) so we must have $d \geq n$, and therefore $d = n$. Clearly $\phi$ generates a cyclic subgroup of order $n$. However by (3.19.107) $G$ has at most order $n$, whence $G = \langle \phi \rangle$ as required. Furthermore by the same result $\mathbb{F}_{p^n}/\mathbb{F}_p$ is Galois. ∎

**Proposition 3.19.112 (Subfields of $\mathbb{F}_{p^n}$)**
*Consider the field extension $\mathbb{F}_{p^n}/\mathbb{F}_p$. Then it has a unique subfield of order $p^m$ if $m \mid n$ (and no such subfield otherwise). In this case $\mathbb{F}_{p^n}/\mathbb{F}_{p^m}$ is Galois and*

$$\mathrm{Gal}\left(\mathbb{F}_{p^n}/\mathbb{F}_{p^m}\right) = \langle \phi^m \rangle$$

*and in particular has order $n/m$.*

*Proof.* We've already shown that $\mathbb{F}_{p^n}$ has a unique subfield of order $p^m$, by assuming an embedding in $\overline{\mathbb{F}_p}$. Let $H = \mathrm{Aut}\left(\mathbb{F}_{p^n}/\mathbb{F}_{p^m}\right)$. Note $\phi^m$ has order $n/m$. Furthermore from (3.19.110) every element of $\mathbb{F}_{p^m}$ satisfies $X^{p^m} - X$. In other words $\phi^m$ fixes $\mathbb{F}_{p^m}$ and $\phi^m \in H$. Therefore $\langle \phi^m \rangle \leq H$ and $\#H \geq n/m$. By (3.19.107) $\#H \leq \left[\mathbb{F}_{p^n} : \mathbb{F}_{p^m}\right] = n/m$, whence we have equality and so the extension is Galois by (3.19.107) and $H = \langle \phi^m \rangle$. ∎

**Lemma 3.19.113**
*Let $x \in \mathbb{F}_{p^n}$. Then* $\deg(x) = n \iff \mathrm{Fix}(x) = \{1\} \subset \mathrm{Gal}\left(\mathbb{F}_{p^n}/\mathbb{F}_p\right)$.

*In other words* $\mathrm{Gal}\left(\mathbb{F}_{p^n}/\mathbb{F}_p\right)$ *acts freely on the elements of degree $n$.*

*Proof.* Observe that $\deg(x) = n \iff \left[\mathbb{F}_p(x) : \mathbb{F}_p\right] = n \iff \mathbb{F}_p(x) = \mathbb{F}_{p^n} \overset{(3.19.115)}{\iff} \mathrm{Gal}\left(\mathbb{F}_{p^n}/\mathbb{F}_p(x)\right) = \{1\}$ which is equivalent to the statement given. ∎

## 3.19.15 Galois Theory

We've seen that for $K/k$ a finite extension

$$\#\,\mathrm{Aut}(K/k) \leq [K : k]_s \leq [K : k]$$

with equality if and only if $K/k$ is Galois, by (3.19.107).

**Remark 3.19.114**
*If $k$ is perfect then $\overline{k}/k$ is Galois.*

The main result of Galois Theory is

**Proposition 3.19.115**
*Let $K/k$ be a finite Galois extension then there is an order-reversing bijection between subgroups and subfields*

© David Rufino CC BY-NC-ND 4.0

$$\begin{aligned}\{H \leq \mathrm{Gal}(K/k)\} &\longleftrightarrow \{F/k \subseteq K/k\} \\ H &\longrightarrow K^H := \{x \in K \mid h(x) = x \quad \forall h \in H\} \\ \mathrm{Gal}(K/F) &\longleftarrow F\end{aligned}$$

*Proof.* This is proved in a series of Propositions in the rest of this section. Firstly we show it is well-defined in (3.19.116). The maps are mutual inverses by (3.19.117) and (3.19.118). ∎

Such an order reversing map is usually called an (antitone) Galois connection, as the first such type arose from Galois Theory. Note it is well-defined because of the following proposition.

**Proposition 3.19.116**
*If $K/k$ is Galois and $F \subset K/k$ then $K/F$ is Galois.*

*Proof.* This follows from (3.19.65) and (3.19.79). ∎

**Proposition 3.19.117 (Fixed field of Galois group)**
*If $K/k$ is Galois and $F \subset K/k$ a subfield then*

$$K^{\mathrm{Gal}(K/F)} = F$$

*Proof.* Clearly $F \subseteq K^{\mathrm{Gal}(K/F)}$. Conversely given $\alpha \in K \setminus F$, then $\deg(m_{\alpha,F}) > 1$. Since $\alpha$ is separable it must have another root $\beta \in K$. By (3.19.66) there is an element $\sigma \in \mathrm{Gal}(K/F)$ such that $\sigma(\alpha) = \beta$. In other words $\alpha \notin K^{\mathrm{Gal}(K/F)}$, which shows the reverse inclusion. ∎

**Proposition 3.19.118**
*Let $K/k$ be a field extension and $H \subseteq \mathrm{Aut}(K/k)$ a finite subgroup then $K/K^H$ is a finite Galois extension with*

$$\mathrm{Gal}(K/K^H) = H$$

*and order equal to $[K : K^H]$. When $K/k$ is finite then $H$ is automatically finite.*

*Proof.* Firstly observe that trivially $H \subseteq \mathrm{Aut}(K/K^H)$. If we know that $[K : K^H] < \infty$, then by (3.19.107) we have

$$\#H \leq \#\,\mathrm{Aut}(K/K^H) \leq [K : K^H]_s \leq [K : K^H]$$

We can prove equality everywhere if we show that $[K : K^H] \leq \#H$, which is shown either by (3.19.119) or (3.19.120). Note equality also shows that $K/K^H$ is finite Galois by the same result.

Finally when $K/k$ is finite, then $\#\,\mathrm{Aut}(K/k) < \infty$. So in this case $H$ is always finite. ∎

We present two approaches to showing the inequality $[K : K^H] \leq \#H$. The first uses independence of characters style argument, and the second which is more straightforward uses the action of $H$ to show that every element has degree at most $\#H$ (Artin).

**Lemma 3.19.119 (Bound degree of fixed field I)**
*Let $K/k$ be a field extension and $H \subset \mathrm{Aut}(K/k)$ a finite subgroup. Then*

$$[K : K^H] \leq \#H$$

*Proof.* Let $H = \{\sigma_1, ..., \sigma_n\}$ with $\sigma_1 = \mathrm{id}$ and $\{\alpha_1, ..., \alpha_m\}$ a $K^H$-linearly independent subset of $K$.

Consider the vector space $K^n$ and the elements $\hat{\alpha}_j = \left(\sigma_1(\alpha_j), ..., \sigma_{n(\alpha_j)}\right)$ for $j = 1...m$. It's enough to show that these are linearly independent over $K$, as that implies $m \leq n$ by (3.3.126). Therefore $K/K^H$ is finite dimensional with dimension at most $n = \#H$.

Let $S(K) := \left\{v \in K^m \;\middle|\; \sum_{j=1}^m v_j\hat{\alpha}_j = 0\right\}$, we aim to show that $S(K) = \{0\}$. If we also consider $S(K^H)$, any non-zero elements will be a $K^H$ linear-dependence for $\alpha_1, ..., \alpha_m$ by considering the first component ($\sigma_1 = \mathrm{id}$). Therefore by linear independence of $\alpha_i$ we see $S(K^H) = \{0\}$. So it's enough to show that $S(K) \neq \{0\} \implies S(K^H) \neq \{0\}$, to prove $S(K) = \{0\}$ by contradiction.

First observe that $K^\star$ and $H$ both act on $S(K)$ component-wise. The first by multiplication and the second by application. This is well-defined because $v \in S(K)$ if and only if

$$\sum_j v_j\sigma(\alpha_j) = 0 \quad \forall \sigma \in H.$$

Apply $\tau$ to obtain

© David Rufino CC BY-NC-ND 4.0

$$\sum_j \tau(v_j)(\tau\circ\sigma)(\alpha_j) = 0 \quad \forall \sigma \in H$$

and since multiplication by $\tau$ permutes $H$ we see $\tau(v) \in S(K)$ as required.

If there exists $0 \neq v \in S(K)$, consider $v$ with a minimal number of non-zero components. By scaling we can assume $\lambda v$ has at least one component in $K^H$. The vector $\tau(\lambda v) - \lambda v$ then has at least one fewer non-zero components, so by minimality must be zero. Since $\tau$ was arbitrary we see $0 \neq \lambda v \in S(K^H)$ as required. ∎

**Lemma 3.19.120 (Bound degree of fixed field II)**
*Let $K/k$ be a field extension and $H$ a finite subgroup of $\mathrm{Aut}(K/k)$. Then $K/K^H$ is finite separable, and simple, with $[K : K^H] \leq \#H$*

*Proof.* We show that $K/K^H$ is separable and every element has degree at most $\#H$. For any $\alpha \in K$, consider the orbit $\alpha^H = \{\sigma(\alpha) \mid \sigma \in H\}$, which is of order at most $\#H$. Then the polynomial

$$f(X) = \prod_{\beta \in \alpha^H} (X - \beta)$$

has $\alpha$ as a root and is separable by (3.19.11). Furthermore $f^\tau = f$ because $\tau$ permutes $\alpha^H$ (it's injective and hence bijective). Therefore $f \in K^H[X]$ and $m_{\alpha,K^H} \mid f$. We see that $\alpha$ has degree at most $\#H$ and is separable by (3.17.40).

If $K/k$ is finite, then a-fortiori $K/K^H$ is finite, so we may apply the Primitive Element Theorem (3.19.105) directly to show the result.

More generally let $K^H(\alpha)$ be a simple subfield of $K$ of maximal degree. This exists because the degree of $\alpha$ is bounded above by $\#H$. We claim $K^H(\alpha) = K$, for if not then $K^H \subseteq K^H(\alpha) \subsetneq K^H(\alpha, \beta)$ is a finite separable extension of $K^H$, whence it must be simple by the Primitive Element Theorem (3.19.105), contradicting maximality. Finally the degree of $[K : K^H]$ is the degree of $\alpha$, which we've seen is bounded above by $\#H$. ∎

Now we may demonstrate straightforward criteria for subfield to be normal

**Proposition 3.19.121**
*Let $K/k$ be a finite Galois extension and $k \subset F \subset K$ a subfield.*

*Then $F/k$ is Galois if and only if $\mathrm{Gal}(K/F) \lhd \mathrm{Gal}(K/k)$ is normal. In this case we have a canonical isomorphism*

$$\mathrm{Gal}(K/k)/\mathrm{Gal}(K/F) \to \mathrm{Gal}(F/k)$$

*Proof.* Recall from (3.19.67) we have $F/k$ is normal iff $\sigma(F) = F$ for all $\sigma \in \mathrm{Gal}(K/k)$. Recall $K/F$ is normal for all subfields $F$. Furthermore, we observe that

$$\mathrm{Gal}(K/\sigma(F)) = \sigma\,\mathrm{Gal}(K/F)\sigma^{-1}$$

By the correspondence (3.19.115) $\mathrm{Gal}(K/F) = \mathrm{Gal}(K/F') \iff F = F'$. Therefore

$$\begin{aligned} F/k \text{ normal} &\iff \sigma(F) = F \quad \forall \sigma \in \mathrm{Gal}(K/k) \\ &\iff \mathrm{Gal}(K/\sigma(F)) = \mathrm{Gal}(K/F) \quad \forall \sigma \in \mathrm{Gal}(K/k) \\ &\iff \sigma\,\mathrm{Gal}(K/F)\sigma^{-1} = \mathrm{Gal}(K/F) \quad \forall \sigma \in \mathrm{Gal}(K/k) \\ &\iff \mathrm{Gal}(K/F) \lhd \mathrm{Gal}(K/k) \end{aligned}$$

The result then follows from (3.19.71). ∎

## 3.19.16 Norm and Trace

In what follows we assume that $K$ is a commutative ring and $A$ is a $K$-algebra which is finite free as a $K$-module (free is automatic when $K$ is a field).

**Definition 3.19.122 (Norm, Trace and Characteristic Polynomial)**
*For $\alpha \in A$ consider the $K$-linear multiplication map*

$$\mathrm{mult}_{\alpha,K} : A \to A$$

*and define*

*a)* $\mathrm{Tr}_{A/K}(\alpha) := \mathrm{Tr}(\mathrm{mult}_\alpha) \in K$
*b)* $\mathrm{Norm}_{A/K}(\alpha) := \det(\mathrm{mult}_\alpha) \in K$
*c)* $\mathrm{Pc}_{A/K}(\alpha; X) := \chi_{\mathrm{mult}_\alpha} = \det\left(\mathrm{mult}_{X-\alpha,K[X]}\right) = \det(X \cdot I_n - [\mathrm{mult}_\alpha]) \in K[X]$ *the characteristic polynomial.*

© David Rufino CC BY-NC-ND 4.0

**Proposition 3.19.123**
*With the notation as in (3.19.122) we have the following properties*

*a)* $\mathrm{Tr}(\alpha+\beta) = \mathrm{Tr}(\alpha) + \mathrm{Tr}(\beta)$
*b)* $\mathrm{Tr}(\alpha\beta) = \mathrm{Tr}(\beta\alpha)$
*c)* $\mathrm{Norm}(\alpha\beta) = \mathrm{Norm}(\alpha)\,\mathrm{Norm}(\beta)$

*and further that*

$$\mathrm{Pc}(\alpha; X) = X^n - \mathrm{Tr}(\alpha)X^{n-1} + \cdots + (-1)^n\,\mathrm{Norm}(\alpha)$$

*Finally for* $\lambda \in K$ *we have*

$$\mathrm{Tr}(\lambda) = \dim_K A \cdot \lambda$$
$$\mathrm{Norm}(\lambda) = \lambda^{\dim_K A}$$

*Proof.* We prove each in turn

a) We have $\mathrm{mult}_{\alpha+\beta} = \mathrm{mult}_\alpha + \mathrm{mult}_\beta$ and ordinary trace is additive (3.5.24).
b) When $A$ is commutative this is obvious, however even in general we have $\mathrm{mult}_{\alpha\beta} = \mathrm{mult}_\alpha \circ \mathrm{mult}_\beta$ and so the result follows from symmetry of the ordinary trace (3.5.25)
c) This follows similarly because the determinant is multiplicative (3.5.17).

The identity for the characteristic polynomial follows directly from (3.18.7). Finally the matrix representation of $\mathrm{mult}_\lambda$ is simply $\lambda \cdot I_n$ with respect to any basis. ∎

**Proposition 3.19.124 (Characteristic Polynomial is a Norm)**
*For* $\alpha \in A$ *we have the relationship*

$$\mathrm{Pc}_{A/K}(\alpha; X) = \mathrm{Norm}_{A[X]/K[X]}(X - \alpha)$$

*Proof.* This follows from the definitions and (3.18.7). ∎

**Proposition 3.19.125**
*Let* $M$ *be a finite free* $A$*-module. Then* $M$ *is a finite free* $K$*-module. Explicitly given a basis* $(m_j)$ *of* $M$ *over* $A$ *and* $(a_i)$ *of* $A$ *over* $K$*, then* $(a_i \cdot m_j)$ *is a basis for* $M$ *over* $K$*.*

*If* $u : M \to M$ *is an endomorphism then we denote by* $u_K$ *the endomorphism of* $M$ *regarded as a (finite free)* $K$*-module.*

**Proposition 3.19.126 (Transitivity of Trace)**
*Suppose* $A$ *is commutative,* $M$ *is a finite free* $A$*-module and* $u : M \to M$ *an endomorphism. Then*

$$\mathrm{Tr}(u_K) = \mathrm{Tr}_{A/K}(\mathrm{Tr}(u))$$

*and*

$$\det(u_K) = \mathrm{Norm}_{A/K}(\det(u))$$

*Proof.* Let $(e_1, ..., e_m)$ be an $A$-basis for $M$ and $(a_1, ..., a_n)$ be a $K$-basis for $A$. Then by (3.19.125) $(a_i \cdot e_j)$ is a $K$-basis for $M$. For $\alpha \in A$ let $M(\alpha) \in \mathrm{Mat}_n(K)$ denote the matrix corresponding $\mathrm{mult}_\alpha$ and let $(c_{lj}) \in \mathrm{Mat}_m(A)$ be the matrix of $u$ with respect to $(e_j)$. By definition this means

$$u(e_j) = \sum_{l=1}^{m} c_{lj} \cdot e_l$$

and

$$a_i\alpha = \alpha a_i = \sum_{k=1}^{n} M(\alpha)_{ki} a_k.$$

Therefore

© David Rufino CC BY-NC-ND 4.0

$$\begin{aligned} u_K(a_i e_j) &= a_i u(e_j) \\ &= a_i \cdot \left( \sum_{l=1}^{m} c_{lj} \cdot e_l \right) \\ &= \sum_{l=1}^{m} (a_i c_{lj}) \cdot e_l \\ &= \sum_{l=1}^{m} \left( \sum_{k=1}^{n} M(c_{lj})_{ki} a_k \right) \cdot e_l \\ &= \sum_{l=1}^{m} \sum_{k=1}^{n} M(c_{lj})_{ki} \cdot (a_k \cdot e_l) \end{aligned}$$

Consequentally the matrix for $u_K$ with respect to the basis $(a_i \cdot e_j)$ is given in block form by

$$\begin{pmatrix} M(c_{11}) & \dots & M(c_{1m}) \\ \vdots & & \vdots \\ M(c_{m1}) & \dots & M(c_{mm}) \end{pmatrix}$$

Therefore we see that

$$\begin{aligned} \mathrm{Tr}(u_K) &= \sum_{j=1}^{m} \mathrm{Tr}(M(c_{jj})) \\ &= \sum_{j=1}^{m} \mathrm{Tr}_{A/K}(c_{jj}) \\ &= \mathrm{Tr}_{A/K} \left( \sum_{j=1}^{m} c_{jj} \right) \\ &= \mathrm{Tr}_{A/K}(\mathrm{Tr}(u)) \end{aligned}$$

as required. Similarly by (3.5.33)

$$\det(u_K) = \det(D_m(M(c_{11}), \dots, M(c_{mm})))$$

and by the Laplace Expansion

$$\begin{aligned} D_m(M(c_{11}), \dots, M(c_{mm})) &= \sum_{\sigma \in S_m} \varepsilon(\sigma) \prod_{i=1}^{m} M\left(c_{i\sigma(i)}\right) \\ &= \sum_{\sigma \in S_m} \varepsilon(\sigma) M \left( \prod_{i=1}^{m} c_{i\sigma(i)} \right) \\ &= M \left( \sum_{\sigma \in S_m} \varepsilon(\sigma) \prod_{i=1}^{m} c_{i\sigma(i)} \right) \\ &= M(\det(u)) \end{aligned}$$

where we implicitly use that $\alpha \to \mathrm{mult}_\alpha \to M(\alpha)$ is a ring homomorphism. Therefore by definition

$$\det(u_K) = \det(M(\det(u))) = \mathrm{Norm}_{A/K}(\det(u))$$

■

**Corollary 3.19.127 (Transitivity)**
*Suppose $B$ is a commutative $A$-algebra which is finite free as an $A$-module. Then*

$$\mathrm{Tr}_{B/K}(b) = \mathrm{Tr}_{A/K}\left(\mathrm{Tr}_{B/A}(b)\right)$$

*and*

$$\mathrm{Norm}_{B/K}(b) = \mathrm{Norm}_{A/K}\left(\mathrm{Norm}_{B/A}(b)\right)$$

*for all $b \in B$. Furthermore*

© David Rufino CC BY-NC-ND 4.0

$$\mathrm{Pc}_{B/K}(b;X) = \mathrm{Norm}_{A[X]/K[X]}\big(\mathrm{Pc}_{B/A}(b;X)\big)$$

**Proposition 3.19.128**
*Let $K/k$ be a finite field extension and $\alpha \in K$. Then*

$$\mathrm{Pc}_{K/k}(\alpha;X) = m_{\alpha,k}(X)^{[K:k(\alpha)]}$$

*Proof.* Consider first the case $K = k(\alpha)$. Then by (3.18.4) we have $\mathrm{Pc}_{K/k}(\alpha;X)(\mathrm{mult}_\alpha) = 0$ and in particular $\mathrm{Pc}_{K/k}(\alpha;X)(\alpha) = 0$ (since $\alpha^n = \alpha^n \cdot 1$). By (...) $m_{\alpha,k}$ divides $\mathrm{Pc}_{K/k}(\alpha;X)$. Further by (...) $\deg\big(m_{\alpha,k}\big) = [k(\alpha):k] = [K:k] = \deg\big(\mathrm{Pc}_{K/k}(\alpha;X)\big)$. Since they are both monic then we find they are equal.

For the general case we may write the matrix for $\mathrm{mult}_\alpha$ in block form by considering the tower basis for $K/k(\alpha)/k$ (see (3.19.125)). This is block diagonal, so we may use the Leibniz formula to compute this directly as product of determinants of the blocks. ∎

We may pull together these results to give a more classical expression for norm and trace.

**Proposition 3.19.129**
*Let $L/K/k$ be a tower of extensions such that $L/k$ is normal (e.g. $L = \overline{k}$). Then for any $\alpha \in K$ we have the identities*

$$i_{kL}\big(\mathrm{Norm}_{K/k}(\alpha)\big) = \left(\prod_{\sigma\in \mathrm{Mor}_k(K,L)} \sigma(\alpha)\right)^{[K:k]_i}$$

*and*

$$i_{kL}\big(\mathrm{Tr}_{K/k}(\alpha)\big) = [K:k]_i \sum_{\sigma\in \mathrm{Mor}_k(K,L)} \sigma(\alpha)$$

*Proof.* Recall from (3.19.78) that the restriction map

$$\mathrm{Mor}_k(K,L) \to \mathrm{Mor}_k(k(\alpha),L)$$

is surjective with finite fibers of order $[K:k(\alpha)]_s$. Therefore we may write the right hand side as

$$\begin{aligned}\left(\prod_{\sigma\in \mathrm{Mor}_k(K,L)} \sigma(\alpha)\right)^{[K:k]_i} &= \left(\prod_{\sigma\in \mathrm{Mor}_k(k(\alpha),L)} \sigma(\alpha)\right)^{[K:k(\alpha)]_s[K:k]_i} \\ &= \left(\prod_{\sigma\in \mathrm{Mor}_k(k(\alpha),L)} \sigma(\alpha)\right)^{[K:k(\alpha)][k(\alpha):k]_i}\end{aligned}$$

Similarly by (3.19.127) and (3.19.123) the left hand side may be written as

$$\begin{aligned}\mathrm{Norm}_{K/k}(\alpha) &= \mathrm{Norm}_{k(\alpha)/k}\big(\mathrm{Norm}_{K/k(\alpha)}(\alpha)\big) \\ &= \mathrm{Norm}_{k(\alpha)/k}\big(\alpha^{[K:k(\alpha)]}\big) \\ &= \mathrm{Norm}_{k(\alpha)/k}\,(\alpha)^{[K:k(\alpha)]}\end{aligned}$$

Comparing these expressions shows we may reduce to the case $K = k(\alpha)$.

From (3.19.49) and (3.19.94) the right hand side is the product of all the roots of $m_{\alpha,k}$ including multiplicity, which is $(-1)^{[K:k]} i_{kL}(c_0)$ where $c_0$ is the constant term of $m_{\alpha,k}(X)$. Then using (3.19.128) and (3.19.123) we see that this also equals $i_{kL}\big(\mathrm{Norm}_{k(\alpha)/k}(\alpha)\big)$ as required.

The case of trace is similar. ∎

**Lemma 3.19.130 (Dedekind's Lemma)**
*Let $K/k$ be a field extension and $A$ a $k$-algebra. The set of $k$-algebra homomorphisms*

$$\mathrm{AlgHom}_k(A,K)$$

*is $K$-linearly independent as a subset of* $\mathrm{Hom}_k(A,K)$.

*Proof.* Suppose that the $k$-algebra homomorphisms are not linearly independent. Then choose a minimal subset of distinct $k$-algebra homomorphisms $\varphi_1, ..., \varphi_n$ with a non-trivial relationship over $K$

© David Rufino CC BY-NC-ND 4.0

$$\sum_{i=1}^{n} \lambda_i \varphi_i = 0 \quad \lambda_i \in K^\star$$

If $n = 1$ then $\varphi_1 = 0$ which is a contradiction. Suppose without loss of generality that $n > 1$. For a given $x \in K$ we may define

$$\hat{\varphi}_i := (\varphi_i(x) - \varphi_n(x))\varphi_i \quad i = 1...n-1$$

We may choose $x$ such that at least one is non-zero. Then for all $y \in A$ we have

$$\sum_{i=1}^{n-1} \lambda_i \hat{\varphi}_i(y) = \sum_{i=1}^{n} \lambda_i (\varphi_i(x) - \varphi_n(x))\varphi_i(y) = \sum_{i=1}^{n} \lambda_i \varphi_i(xy) - \varphi_n(x) \sum_{i=1}^{n} \lambda_i \varphi_i(y) = 0$$

This contradicts minimality of the subset. ■

**Proposition 3.19.131**
*Let $K/k$ be a finite extension. Then the following are equivalent*

*a) $K/k$ is separable*
*b) $\mathrm{Tr}_{K/k}$ is not identically zero*
*c) The bilinear form*

$$(x, y) \to \mathrm{Tr}_{K/k}(xy)$$

*is a perfect pairing (see (3.3.148)).*

*Proof.* c) $\implies$ b) By (3.3.148) for all $0 \neq x \in K$ we have $y \to \mathrm{Tr}_{K/k}(xy)$ is non-zero, so there exists $y$ such that $\mathrm{Tr}_{K/k}(xy) \neq 0$ as required.

b) $\implies$ c) By (3.3.149) it's enough to show that for all $0 \neq x \in K$ the map $y \to \mathrm{Tr}_{K/k}(xy)$ is non-zero. By assumption there exists some $z$ such that $\mathrm{Tr}_{K/k}(z) \neq 0$ then we may consider $y = z/x$.

a) $\implies$ b) By (3.19.93) $[K : k]_i = 1$, and so from (3.19.129) the trace equals $\sum_\sigma \sigma$, which is non-zero by (3.19.130).

b) $\implies$ a) Suppose $K/k$ is not separable then $[K : k]_i$ is a power of $p$ (3.19.95), and so the trace is zero by (3.19.129). ■

## 3.19.17 Transcendental Field Extensions

**Definition 3.19.132 (Algebraic Independence)**
*Let $K/k$ be a field extension and $S \subset K$. We say $S$ is **algebraically independent** over $k$ if for every finite subset of distinct elements $x_1, ..., x_n \in S$ we have*

$$f(x_1, ..., x_n) = 0 \implies f = 0$$

*for all $f \in k[X_1, ..., X_n]$.*

*For a subset $S \subset K$ define the closure operator*

$$c(S) := \overline{k(S)} \cap K := \{x \in K \mid x \text{ algebraic over } k(S)\}$$

*We say $\Gamma$ is **algebraically spanning** if $c(\Gamma) = K$, equivalently if $K/k(\Gamma)$ is algebraic.*

*We say that $\mathcal{B}$ is a **transcendence base** if it is both algebraically independent and spanning (i.e. $K/k(\mathcal{B})$ is algebraic).*

Essentially we show that $(K, c)$ satisfies the properties of a matroid, in analogy with vector spaces, so that we can use the results of the Matroids section to show that transcendence bases exist and they satisfy certain properties.

**Proposition 3.19.133**
*Let $K/k$ be a field extension and $S, T$ subsets of $K$. Then the following are equivalent*

*a) $S \cup T$ is algebraically independent and $S \cap T = \varnothing$*
*b) $S$ is algebraically independent over $k$ and $T$ is algebraically independent over $k(S)$*
*c) $T$ is algebraically independent over $k$ and $S$ is algebraically independent over $k(T)$*

*Proof.* By symmetry it's enough to show that a) $\iff$ b).

Suppose a) holds, then a-fortiori $S$ is algebraically independent over $k$. Suppose $T$ is algebraically dependent over $k(S)$, then there exists $t_1, ..., t_n \in T$ such that $F(t_1, ..., t_n) = 0$ for some $0 \neq F \in k(S)[X_1, ..., X_n]$. By clearing denominators we may assume that $F \in k[S][X_1, ..., X_n]$. As there are only finitely many coefficients we may also take $S = \{s_1, ..., s_m\}$ to be finite. Explicitly

© David Rufino CC BY-NC-ND 4.0

$$F(X_1, ..., X_n) = \sum_{\alpha \in \mathbb{N}^n} \lambda_\alpha X_1^{\alpha_1} ... X_n^{\alpha_n}$$

and

$$\lambda_\alpha = G_\alpha(s_1, ..., s_m)$$

for some $G_\alpha \in k[Y_1, ..., Y_m]$. We may then define $\hat{F} \in k[X_1, ..., X_n, Y_1, ..., Y_m]$ by

$$\hat{F}(X_1, ..., X_n, Y_1, ..., Y_m) := \sum_{\alpha \in \mathbb{N}^n} G_\alpha(Y_1, ..., Y_m) X_1^{\alpha_1} ... X_n^{\alpha_n}$$

Then $\hat{F}(t_1, ..., t_n, s_1, ..., s_m) = 0$, and by assumption this is an algebraic dependence (as $S \cap T = \varnothing$). Therefore by assumption $\hat{F} = 0$ and in particular $G_\alpha(s_1, ..., s_m) = 0$ for all $\alpha$. This implies $F = 0$, a contradiction.

*b)* $\implies$ *a)* Evidently $S \cap T = \varnothing$ otherwise there is a trivial algebraic dependence of $T$ on $k(S)$. Suppose there is an algebraic dependence of $S \cup T$ on $k$. We may assume wlog that $S = \{s_1, ..., s_m\}$ and $T = \{t_1, ..., t_n\}$ are finite. Then the evaluation homomorphism

$$k[X_1, ..., X_n, Y_1, ..., Y_m] \to K$$

may be written as a composite

$$k[X_1, ..., X_n, Y_1, ..., Y_m] \to k(S)[X_1, ..., X_n] \to K$$

which are both injective by hypothesis. ∎

**Corollary 3.19.134 (Exchange Property)**
*Let $K/k$ be a field extension, $\Gamma \subseteq K$. Suppose $x$ is algebraic over $k(\Gamma \cup \{y\})$ and transcendental over $k(\Gamma)$, then $y$ is algebraic over $k(\Gamma \cup \{x\})$.*

*Proof.* By considering the extension $K/k(\Gamma)$ we may reduce to the case $\Gamma = \varnothing$.

It's enough to show that $x$ transcendental over $k$ and $y$ transcendental over $k(x)$ implies $x$ transcendental over $k(y)$. This follows directly from (3.19.133) by considering $S = \{x\}$ and $T = \{y\}$. ∎

**Corollary 3.19.135 (Extension Property)**
*Let $K/k$ be a field extension, $S \subseteq K$ algebraically independent and $x \in K$ transcendental over $k(S)$. Then $S \cup \{x\}$ is algebraically independent.*

*Proof.* Follows immediately from (3.19.133). ∎

**Lemma 3.19.136 (Embeddings Preserve Transcendence Degree)**
*Let $K/k$, $L/k$ be field extensions of finite transcendence degree and $\phi : K \to L$ a $k$-algebra homomorphism. Then* $\operatorname{trdeg}(K/k) \leq \operatorname{trdeg}(L/k)$.

*Proof.* As $\phi$ is a ring homomorphism from a field, $\ker(\phi)$ is a proper ideal of $K$, hence 0, so $\phi$ is injective. Let $S \subset K$ be a transcendence base of $K/k$, so $|S| = \operatorname{trdeg}(K/k)$ (3.19.141); as $\phi$ is injective, $|\phi(S)| = |S|$. Given distinct $s_1, ..., s_n \in S$ and $f \in k[X_1, ..., X_n]$ with $f(\phi(s_1), ..., \phi(s_n)) = 0$: as $\phi$ fixes $k$, $\phi(f(s_1, ..., s_n)) = f(\phi(s_1), ..., \phi(s_n)) = 0$, so $f(s_1, ..., s_n) = 0$ by injectivity of $\phi$, whence $f = 0$ by algebraic independence of $S$ over $k$; so $\phi(S)$ is algebraically independent over $k$. By (3.19.140) applied to $L/k$, $\operatorname{trdeg}(K/k) = |S| = |\phi(S)| \leq \operatorname{trdeg}(L/k)$. ∎

**Corollary 3.19.137 (Equivalent form of independence)**
*Let $K/k$ be a field extension and $S \subset K$. Then the following are equivalent*

*a) $S$ is algebraically independent*
*b) $x$ is transcendental over $k(S \setminus \{x\})$ for all $x \in S$*

*In other words the algebraically independent subsets are precisely the matroid independent subsets.*

*Proof. a)* $\implies$ *b)* Follows from (3.19.133).

*b)* $\implies$ *a)* Let $\mathcal{F}$ be the family of algebraically independent subsets. We've shown that all such sets are also matroid independent, and by definition the family is of finite character. Furthermore by (3.19.135) it satisfies the extension property. Therefore by (2.3.6) it is precisely the family of matroid independent sets. ∎

**Proposition 3.19.138 (Transcendence Base Exists)**
*Let $K/k$ be a field extension. Suppose $S$ is an algebraic independent subset of $K$ and $\Gamma \supset S$ is such that $K/k(\Gamma)$ is algebraic. Then there exists a transcendence base $\mathcal{B}$ such that $S \subseteq \mathcal{B} \subseteq \Gamma$.*

© David Rufino CC BY-NC-ND 4.0

*Proof.* This is (2.3.7). ■

**Proposition 3.19.139 (Transcendence Base)**
*Let $K/k$ be a field extension and $S \subset K$ a subset. Then the following are equivalent*

*a) $S$ is a **transcendence base***
*b) $S$ is a maximal algebraically independent set*
*c) $S$ is minimal under the condition $K/k(S)$ is algebraic*

*When $K/k$ admits a finite algebraic spanning set then all bases are finite of the same size (**transcendence degree**). Write this as $\text{trdeg}(K/k)$ or $\text{trdeg}_k(K)$.*

*Proof.* Follows from (2.3.8) and (2.3.12). ■

**Proposition 3.19.140**
*Let $K/k$ be a field extension of finite transcendence degree and $S \subset K$ a subset. Then*

- *$S$ algebraically independent $\implies |S| \leq \text{trdeg}(K/k)$*
- *$K/k(S)$ is algebraic $\implies |S| \geq \text{trdeg}(K/k)$*

*Proof.* Follows directly from (2.3.13). ■

**Proposition 3.19.141**
*Let $K/k$ be a field extension of finite transcendence degree and $S \subset K$ a subset. Then the following are equivalent*

- *$S$ is a transcendence base*
- *$S$ is algebraically independent and $|S| \geq \text{trdeg}(K/k)$*
- *$K/k(S)$ is algebraic and $|S| \leq \text{trdeg}(K/k)$*

*In this case $|S| = \text{trdeg}(K/k)$.*

*Proof.* Follows directly from (2.3.14). ■

In case $K/k$ is finitely generated then we guarantee that $K/k(S)$ is finite.

**Proposition 3.19.142**
*Let $K/k$ be a finitely generated field extension. Then*

- $\text{trdeg}(K/k) < \infty$
- *Suppose $K/k(\Gamma)$ is algebraic then it is in fact finite.*

*Proof.* *a*) If $K/k$ is finitely generated then the set of generators is algebraically spanning and so $\text{trdeg}(K/k) < \infty$ by (3.19.139)

*b*) A-fortiori $K/k(\Gamma)$ is finitely generated and by assumption algebraic so by (...) it is finite ■

## 3.19.18 Separating Transcendence Base

In applications the existence of a separating transcendence base is important, and is guaranteed when $k$ is perfect, or a weaker condition.

**Definition 3.19.143 (Separating Transcendence Base)**
*A field extension $K/k$ is **separably generated** if there exists a transcendence base $S$ such that $K/k(S)$ is separable (and algebraic). We call such an $S$ a **separating transcendence base**.*

*Note when $\text{char}(k) = 0$ then every transcendence base is separating.*

**Definition 3.19.144 (Maclane's Criterion)**
*We say that an algebra $A/k$ of characteristic exponent $p$ satisfies **Maclane's Criterion** if it is reduced and for all $a_1, ..., a_n \in A$ the following property holds*

$$\{a_1, ..., a_n\} \text{ k-linearly independent} \implies \{a_1^p, ..., a_n^p\} \text{ k-linearly independent}$$

When $p = 1$ this is trivially always satisfied and when $p > 1$ such an algebra is automatically reduced by the following result

**Lemma 3.19.145**
*Let $A$ be a $k$-algebra and $m > 1$ an integer. Then the following are equivalent*

© David Rufino CC BY-NC-ND 4.0

*a) $A$ is reduced*
*b) $x^m = 0 \implies x = 0$*

*In particular if $A$ has characteristic exponent $p > 1$ then it is reduced iff the Frobenius homomorphism is injective.*

*Proof.* $a) \implies b)$ is trivial. Conversely suppose $x^n = 0$. Then there exists $a > 0$ such that $n \leq am$. In particular $x^{am} = 0$ and therefore by induction $x = 0$ as required. ∎

This is a generalization of the case of a perfect base field by the following result.

**Proposition 3.19.146 (Perfect $\implies$ Maclane)**
*Let $A/k$ be a reduced algebra with $k$ perfect. Then it satisfies Maclane's Criterion.*

*In particular this holds for every field extension $K/k$ of a perfect base field.*

*Proof.* By assumption $A$ is reduced so we may consider only the case $p > 1$.

Suppose $0 = \sum_i \lambda_i a_i^p$ then by hypothesis $\lambda_i = \mu_i^p$. By (3.19.6) $0 = \left(\sum_i \mu_i a_i\right)^p$. As $A$ is reduced we find $0 = \sum_i \mu_i a_i$ as required. ∎

**Lemma 3.19.147**
*Let $n \geq 0$ an integer and $K = k(x_1, ..., x_{n+1})$ an extension of $k$ such that*

- $\text{char}(k) \neq 0$
- *$\{x_1, ..., x_n\}$ is a transcendence base*
- *$K/k$ satisfies Maclane's Criterion*

*Then for some $x_i$ the set $\{x_1, ..., \hat{x_i}, ..., x_{n+1}\}$ is a **separating transcendence base** for $K/k$.*

*Note when $n = 0$ this means precisely that a simple algebraic extension satisfying Maclane's Criterion is separable.*

*Proof.* By assumption $x_{n+1}$ is algebraic over $k(x_1, ..., x_n)$, and therefore there exists a non-zero $F \in k[X_1, ..., X_{n+1}]$ such that $F(x_1, ..., x_{n+1}) = 0$. Let $F$ be such a polynomial of minimal total degree (total degree = the maximum degree of a monomial with non-zero coefficient appearing in $F$). We claim it is irreducible. For suppose not, then one of the irreducible factors must vanish at $(x_1, ..., x_{n+1})$, and this factor would have smaller total degree.

We wish to show that not all powers of $X_i$ appearing in $F$ are multiples of $p$. Suppose this were the case then the monomials

$$\left\{x_1^{\alpha_1} ... x_{n+1}^{\alpha_{n+1}} \mid \lambda_\alpha \neq 0\right\}$$

are $k$-linearly dependent where $\lambda_\alpha$ are the coefficients of $F$ and $\alpha \in \mathbb{N}^{n+1}$. Then by Maclane's Criterion the set

$$\left\{x_1^{\alpha_1/p} ... x_{n+1}^{\alpha_{n+1}/p} \;\middle|\; \lambda_\alpha \neq 0\right\}$$

is linearly dependent. This contradicts the minimality of $F$.

Choose $X_i$ for which a non $p$-th power appears in F and define

$$F_i(T) := F(x_1, ..., x_{i-1}, T, x_{i+1}, ..., x_{n+1}) \in k[S_i][T]$$

where $S_i := \{x_1, ..., \hat{x_i}, ..., x_{n+1}\}$. By assumption $F_i(T)$ is non-zero and clearly $F_i(x_i) = 0$ so that $x_i$ is algebraic over $k(S_i)$, and therefore by (3.19.44) $K$ is also algebraic over $k(S_i)$. By (3.19.141) $S_i$ is a transcendence base for $K$ and in particular algebraically independent. Therefore $k[S_i][T]$ may be identified with $k[X_1, ..., X_{n+1}]$ and we may conclude that $F_i(T)$ is irreducible. Further $k[S_i]$ is a UFD so by (3.17.44) $F_i(T)$ is irreducible as a polynomial in $k(S_i)[T]$. By construction $F_i(T) \notin k(S_i)[T^p]$ so by (3.19.91) $F_i$ is separable. Therefore $x_i$ is separable over $k(S_i)$, and $K/k(S_i)$ is separable by (3.19.80) as required. ∎

**Proposition 3.19.148**
*Let $K/k$ be a finitely-generated field extension which satisfies Maclane's Criterion (e.g. $k$ is perfect). Then $K/k$ is separably generated.*

*More precisely every generating set contains a separating transcendence base.*

*Proof.* When $\text{char}(k) = 0$ the generating set contains a transcendence base by (3.19.139). Every subfield of $K$ also has characteristic 0, so then we are done by (3.19.103). So we may only consider the positive characteristic case, $p > 1$.

Suppose $K = k(x_1, ..., x_n)$. By (3.19.138) there is a (possibly empty) subset which is a transcendence base, say $\{x_1, ..., x_d\}$ after renumbering. By (3.19.44) $[K : K'] < \infty$ where $K' = k(x_1, ..., x_d)$, and we may choose $K'$ such that

© David Rufino CC BY-NC-ND 4.0

the degree of inseparability $[K:K']_i$ is minimal. If $K/K'$ is separable then we are done. Otherwise we may assume by renumbering that $K'(x_{d+1})/K'$ is not separable (3.19.80) and therefore $[K'(x_{d+1}):K']_i > 1$. Then by (3.19.147) there is an index $i \leq d+1$ such that $[K'(x_{d+1}):K'']_i = 1$ where $K'' := k(x_1, ..., \hat{x}_i, ..., x_{d+1})$. By multiplicativity

$$[K:K']_i = [K:K'(x_{d+1})]_i [K'(x_{d+1}):K']_i > [K:K'(x_{d+1})]_i [K'(x_{d+1}):K'']_i = [K:K'']_i$$

which contradicts minimality. ∎

© David Rufino CC BY-NC-ND 4.0

# 3.20 Local Rings

Local rings arise quite naturally when localizing at a prime ideal (see (3.6.42) and (3.20.5)) so we recall some basic properties here.

**Definition 3.20.1 (Local Ring)**
*A ring $A$ is a **local ring** if it has a unique maximal ideal $\mathfrak{m}$. The field $A/\mathfrak{m}$ is called the **residue field** of $A$ and sometimes denoted $\kappa(\mathfrak{m})$.*

*When $A$ is a $k$-algebra we may denote the residue field by $k(\mathfrak{m})$.*

**Definition 3.20.2 (Local Homomorphism)**
*Let $(A, \mathfrak{m}_A)$ and $(B, \mathfrak{m}_B)$ be local rings. A ring homomorphism $\phi : A \to B$ is said to be a **local homomorphism** if*

$$\phi(\mathfrak{m}_A) \subseteq \mathfrak{m}_B$$

Recall that the group of units $A^\star$ of a ring is a saturated multiplicative set, that is

$$xy \in A^\star \iff x \in A^\star \wedge y \in A^\star$$

**Proposition 3.20.3 (Criteria for Local Rings)**
*Let $A$ be a ring. Then the following are equivalent*

*a) $A$ is a local ring*
*b) $A \setminus A^\star$ is an additive subgroup of $A$*

*In this case $\mathfrak{m} = A \setminus A^\star$ is the unique maximal ideal of $A$.*

*Proof.* a) $\implies$ b) Let $\mathfrak{m}$ be the unique maximal ideal then, because it's proper, $\mathfrak{m} \cap A^\star = \varnothing \implies \mathfrak{m} \subseteq A \setminus A^\star$ by (3.3.13). Conversely given $x \in A \setminus A^\star$ then $(x)$ is a proper ideal by (3.3.33), and therefore contained in a maximal ideal (3.3.15) which by uniqueness means $x \in \mathfrak{m}$.

b) $\implies$ a) Define $\mathfrak{m} = A \setminus A^\star$ it's a (prime) ideal because it is an additive subgroup and $A^\star$ is a saturated multiplicative set. Let $\mathfrak{a}$ be a proper ideal then $\mathfrak{a} \cap A^\star = \varnothing \implies \mathfrak{a} \subseteq \mathfrak{m}$. Therefore $\mathfrak{m}$ is the unique maximal ideal. ∎

**Proposition 3.20.4 (Quotient of a Local Ring is Local)**
*Let $(A, \mathfrak{m})$ be a local ring and $\mathfrak{a} \lhd A$ a proper ideal. Then $A/\mathfrak{a}$ is a local ring, with maximal ideal $\mathfrak{m}/\mathfrak{a}$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

As $\mathfrak{a}$ is proper, it is contained in some maximal ideal (3.3.15), which must be $\mathfrak{m}$ by uniqueness; so $\mathfrak{a} \subseteq \mathfrak{m}$, and by the ideal correspondence of (3.3.56), the maximal ideals of $A/\mathfrak{a}$ correspond to the maximal ideals of $A$ containing $\mathfrak{a}$ – of which $\mathfrak{m}$ is the only one (as $A$ is local) – so $A/\mathfrak{a}$ has the unique maximal ideal $\mathfrak{m}/\mathfrak{a}$, that is, is local. ∎

**Example 3.20.5**
*Let $A$ be a ring and $\mathfrak{p} \lhd A$ a prime ideal. Then $A_\mathfrak{p}$ is a local ring with unique maximal ideal $\mathfrak{p}A_\mathfrak{p}$.*

*When $A \subset K$ is a subring of a field then $A \subset A_\mathfrak{p} \subset K$ in a natural way.*

We may use this to provide another criteria

**Lemma 3.20.6 (Criteria for Local Domain)**
*Let $A \subset K$ be a subring of a field with a prime ideal $\mathfrak{p} \lhd A$. Then $A \subset A_\mathfrak{p}$.*

*$A$ is a local ring with unique maximal ideal $\mathfrak{p}$ if and only if $A = A_\mathfrak{p}$,*

*Proof.* We've observed that $A_\mathfrak{p}$ is a local ring with unique maximal ideal $\mathfrak{p}A_\mathfrak{p}$.

If $A = A_\mathfrak{p}$ then it is a local ring and $\mathfrak{p} \subset \mathfrak{p}A_\mathfrak{p}$. For $y \notin \mathfrak{p}$, then $\frac{1}{y} \in A_\mathfrak{p} \implies \frac{1}{y} \in A$. So we see that $x \in \mathfrak{p}, y \notin \mathfrak{p}$ we have $\frac{x}{y} \in \mathfrak{p}$ and $\mathfrak{p} = \mathfrak{p}A_\mathfrak{p}$.

Conversely suppose $A$ is a local ring with unique maximal ideal $\mathfrak{p}$. Then $y \notin \mathfrak{p} \implies y \in A^\star$ and $A_\mathfrak{p} = A$ as required. ∎

© David Rufino CC BY-NC-ND 4.0

## 3.21 Modules over Local Rings (Nakayama's Lemma)

The main result of this section (3.21.7) is that every finitely-generated module over a local ring has a minimal spanning set. Recall in the vector space case a minimal spanning set is precisely a basis (2.3.8). Analogously we may also show that (in the local case) every minimal spanning set has the same order. The crucial result is Nakayama's Lemma, which we develop here.

**Definition 3.21.1 (Jacobson Radical)**
*Let $A$ be a commutative ring. Define the* ***Jacobson Radical*** *to be the intersection of all maximal ideals*

$$\sqrt{0}^J := \bigcap_{\mathfrak{m} \lhd A} \mathfrak{m}$$

**Proposition 3.21.2**
*The Jacobson Radical $\sqrt{0}^J$ is a proper ideal*

**Example 3.21.3**
*When $(A, \mathfrak{m}_A)$ is a local ring then $\sqrt{0}^J = \mathfrak{m}_A$.*

**Lemma 3.21.4 (Characterization of Jacobson Radical)**
*For an ideal $\mathfrak{a}$ and $\mathfrak{m}$ a maximal ideal*

*a)* $\mathfrak{a} \not\subseteq \mathfrak{m} \iff \mathfrak{a} + \mathfrak{m} = A \iff (1 + \mathfrak{a}) \cap \mathfrak{m} \neq \varnothing$
*b)* $\mathfrak{a} \subseteq \sqrt{0}^J \iff 1 + \mathfrak{a} \subseteq A^\star$
*c)* $x \in \sqrt{0}^J \iff 1 + (x) \subseteq A^\star$

*Proof.* We prove each in turn.

a) Clearly $\mathfrak{a} \subseteq \mathfrak{m} \implies \mathfrak{a} + \mathfrak{m} = \mathfrak{m}$. Conversely $\mathfrak{a} \not\subseteq \mathfrak{m} \implies \mathfrak{a} + \mathfrak{m} = A$ by maximality. Suppose $\mathfrak{a} + \mathfrak{m} = A$ then $1 = a + m$ whence $(1 - a) \in \mathfrak{m}$. The converse is similar.

b) By a) $\mathfrak{a} \subseteq \sqrt{0}^J \implies (1 + \mathfrak{a}) \cap \mathfrak{m} = \varnothing$ for all maximal ideals $\mathfrak{m}$. By (3.3.15) this implies $(1 + \mathfrak{a}) \subseteq A^\star$.

Conversely if $(1 + \mathfrak{a}) \subseteq A^\star$ then $(1 + \mathfrak{a}) \cap \mathfrak{m} = \varnothing$ for any maximal ideal $\mathfrak{m}$ by (3.3.13). Again by a) $\mathfrak{a} \subseteq \mathfrak{m}$ as required.

c) This follows from b) and noting $x \in \sqrt{0}^J \iff (x) \subseteq \sqrt{0}^J$. ∎

**Proposition 3.21.5 (Nakayama's Lemma)**
*Let $M$ be a finitely generated $A$-module and $\mathfrak{a} \lhd A$ an ideal. Then the following holds*

*a) If $M = \mathfrak{a}M$ then there exists $a \in \mathfrak{a}$ such that $m = am$ for all $m \in M$*

*Suppose in addition that $\mathfrak{a} \subseteq \sqrt{0}^J$ (e.g. if $A$ is local and $\mathfrak{a}$ is proper) then*

*b)* $M = \mathfrak{a}M \implies M = 0$
*c)* $N \leq M$ *and* $M = N + \mathfrak{a}M \implies M = N$.

*Proof.* We prove each in turn

a) Apply (3.18.4) with $\phi := \mathbf{1}_M$ to find a monic polynomial $P(X) \in A[X]$ with non-leading coefficients in $\mathfrak{a}$ such that $P(\phi)(m) = 0$ for all $m \in M$. Then we see that $(1 + a_{n-1} + ...a_0)m = 0$ for all $m \in M$ whence $a := -(a_{n-1} + ... + a_0)$ is the required element.

More directly, suppose $m_1, ..., m_n$ is a generating set for $M$. By (3.18.3) (and $M = \mathfrak{a}M$) there is a matrix $E$ with coefficients in $\mathfrak{a}$ such that

$$(I_n - E)\boldsymbol{m} = 0$$

where $\boldsymbol{m}$ is the column vector consisting of $m_1, ..., m_n$. By (3.5.23) we see $\det(I_n - E)m_i = 0$ for all $i = 1...n$. It's enough to show $a := \det(I_n - E) \in 1 + \mathfrak{a}$. Observe

$$\det(I_n - E) = \prod_i (1 - E_{ii}) + \sum_{\sigma \neq \mathrm{id}} \epsilon(\sigma) \prod_j E_{j\sigma(j)}$$

The second term lies in $\mathfrak{a}$ and

$$\prod_i (1 - E_{ii}) = 1 - \sum_{i=1}^n E_{ii} \prod_{j>i} (1 - E_{jj}) \in 1 + \mathfrak{a}$$

© David Rufino CC BY-NC-ND 4.0

b) Consider any $m \in M$. By a) we have $(1-a)m = 0$ for some $a \in \mathfrak{a}$, and by (3.21.4) $(1-a)$ is invertible, whence $m = 0$ as required.

c) Observe $\mathfrak{a}(M/N) \overset{(3.3.107)}{=} (N + \mathfrak{a}M)/N = M/N$ whence $M/N = 0$ by b). Therefore $N = M$ as required.

We may show b) more directly. Suppose $M \neq 0$, and let

$$\{m_1, ..., m_n\}$$

be a non-zero generating set for $M$ of minimal size. Then by (3.18.3)

$$m_1 = \sum_j a_j m_j \quad a_j \in \mathfrak{a}$$

whence

$$(1-a_1)m_1 = \sum_{j \geq 2} a_j m_j$$

As $a_1 \in \sqrt{0}^J$ we have $1 - a_1 \in A^\star$ by (3.21.4). Then

$$\{m_2, ..., m_n\}$$

is a smaller generating set, a contradiction. Therefore $M = 0$.

a) may be deduced from b) as follows. Observe $S \coloneqq 1 + \mathfrak{a}$ is a multiplicatively closed subset, so we may consider $S^{-1}M$ as an $S^{-1}A$-module. It's easy to verify that $1 + S^{-1}\mathfrak{a} \subseteq (S^{-1}A)^\star$ so by (3.21.4) $S^{-1}\mathfrak{a} \subseteq J(S^{-1}A)$. Clearly $\mathfrak{a}M = M \implies (S^{-1}\mathfrak{a})S^{-1}M = S^{-1}M$ so by the weaker form $S^{-1}M = 0$. By (3.6.16) there exists $s \in S$ such that $sM = 0$, which is the required result as $s = 1 + a$ for some $a \in \mathfrak{a}$. ∎

Recall (3.3.108) in the case of a local ring $(A, \mathfrak{m})$ that $\widetilde{M} \coloneqq M/\mathfrak{m}M$ is a vector space over $k \coloneqq A/\mathfrak{m}$. We may use Nakayama's Lemma to exhibit a correspondence between minimal spanning sets of $M$ and bases of $M/\mathfrak{m}M$ as a $k$-vector space. First we prove a simpler form

**Lemma 3.21.6**
*Let $(A, \mathfrak{m})$ be a local ring with residue field $k = A/\mathfrak{m}$, $M$ a finite $A$-module and $S \subset M$ a subset. Then*

$$S \text{ spans } M \iff \tilde{S} \text{ spans } \widetilde{M}$$

*where $\tilde{}$ denotes reduction modulo $\mathfrak{m}M$.*

*Proof.* One direction is obvious. Conversely suppose $\tilde{S}$ spans $\widetilde{M}$. Define $N \coloneqq \langle S \rangle$, then this means precisely that $N + \mathfrak{m}M = M$, so $N = M$ by (3.21.5) c) as required. ∎

Recall from (2.1.57) that every subset of $T$ of $M/\mathfrak{m}M$ may be written in the form $\tilde{S}$ for $S \subset M$ and $\tilde{}$ injective on $S$.

**Proposition 3.21.7 (Structure theorem for modules over a local ring)**
*Let $(A, \mathfrak{m})$ be a local ring with residue field $k = A/\mathfrak{m}$, $M$ a finite $A$-module. Then*

*a) $\widetilde{M} \coloneqq M/\mathfrak{m}M$ is a finite-dimensional $k$-module (of dimension $n$ say)*
*b) If $\tilde{S}$ is a $k$-basis for $\widetilde{M}$ and $\tilde{}|_S$ is injective, then $S$ is a minimal spanning set for $M$ and $\#S = \#\tilde{S} = n$*
*c) If $S$ is a minimal spanning set then $\tilde{S}$ is a basis for $\widetilde{M}$ (and $\tilde{}|_S$ is injective so $\#S = \#\tilde{S} = n$).*

*Proof.* a) By (3.3.108) $\widetilde{M}$ is a $k$-module, and it's clearly finite.

b) By (3.21.6) $S$ spans $M$. Suppose $S' \subset S$ spans $M$, then by the same result $\widetilde{S'}$ spans $\widetilde{M}$. Recall (2.3.8) that a vector space basis is precisely a minimal spanning set, so $\widetilde{S'} = \tilde{S}$. As $\tilde{}$ is injective this means $S' = S$. Therefore $S$ is a minimal spanning set.

c) Let $S$ be a minimal spanning set. Then by (3.21.6) $\tilde{S}$ spans $\widetilde{M}$. Suppose $\tilde{T} \subset \tilde{S}$ also spans $\widetilde{M}$. By (3.21.6) $T$ spans $M$, and by hypothesis $T = S$. Therefore $\tilde{T} = \tilde{S}$. As $\tilde{T}$ was arbitrary we see that $\tilde{S}$ is a minimal spanning set for $\widetilde{M}$, which by (2.3.8) is a basis.

Finally suppose $\tilde{}$ is not injective on $S$, that is $\tilde{s_1} = \tilde{s_2}$. Then $S' \coloneqq S \setminus \{s_1\}$ satisfies $\widetilde{S'} = \tilde{S}$. Therefore by the Lemma $S'$ spans $M$, contradicting minimality. ∎

© David Rufino CC BY-NC-ND 4.0

## 3.22 Krull Dimension

**Definition 3.22.1 (Krull Dimension)**
*Let $A$ be a commutative ring. We say that a chain of distinct prime ideals*

$$\mathfrak{p}_0 \subsetneq \mathfrak{p}_1 \subsetneq \cdots \subsetneq \mathfrak{p}_n$$

*has **length** $n$.*

a) *The **Krull dimension** $\dim A$ of $S$ is the maximum length of all chains of prime ideals.*
b) *The **height** of a prime ideal $\mathfrak{p}$, denoted $\operatorname{ht}(\mathfrak{p})$, is the maximum length of chains of prime ideals contained in $\mathfrak{p}$. More generally define $\operatorname{ht}(\mathfrak{a}) = \inf\{\operatorname{ht}(\mathfrak{p}) \mid \mathfrak{a} \subseteq \mathfrak{p}\}$. By (3.3.44) we may take the infimum over only minimal prime ideals.*
c) *The **dimension** of an ideal $\mathfrak{a} \lhd A$, denoted $\dim \mathfrak{a}$, is the maximum length of chains of prime ideals containing $\mathfrak{a}$.*

*We say $A$ is **finite-dimensional** if $\dim A < \infty$. Observe that $\mathfrak{a} \subseteq \mathfrak{p} \iff \sqrt{\mathfrak{a}} \subseteq \mathfrak{p}$ for any prime ideal $\mathfrak{p}$ so*

$$\dim \mathfrak{a} = \dim \sqrt{\mathfrak{a}}$$

$$\operatorname{ht}(\mathfrak{a}) = \operatorname{ht}\left(\sqrt{\mathfrak{a}}\right)$$

*and we may, without loss of generality, consider only radical ideals.*

**Definition 3.22.2**
*Let $A$ be a commutative ring. We say a chain of prime ideals is*

- ***maximal** if it's not properly contained in any chain*
- ***saturated** if $\mathfrak{p}_i \subseteq \mathfrak{p} \subseteq \mathfrak{p}_{i+1} \implies \mathfrak{p} = \mathfrak{p}_i$ or $\mathfrak{p} = \mathfrak{p}_{i+1}$.*

*We say that $A$ is **biequidimensional** if every maximal chain has the same length (equal to $\dim A$).*

*We say $A$ is **quasi-biequidimensional** if $A/\mathfrak{p}$ is **biequidimensional** for every minimal prime $\mathfrak{p}$. Note that **equidimensional** + **quasi-biequidimensional** $\iff$ **biequidimensional**.*

*We say that $A$ is **catenary** if the length of a saturated chain*

$$\mathfrak{p}_0 \subsetneq \cdots \subsetneq \mathfrak{p}_n$$

*depends only on $\mathfrak{p}_0$ and $\mathfrak{p}_n$ and is equal to $\operatorname{ht}(\mathfrak{p}_n/\mathfrak{p}_0)$*

*We say that $A$ is **equidimensional** if every minimal prime ideal has the same dimension (equal to $\dim A$). Note an **irreducible** ring has only one minimal prime ideal so is trivially equidimensional.*

*We say that $A$ is **equicodimensional** if every maximal ideal has the same height (equal to $\dim A$).*

In order to connect this to the lattice-theoretic notion of Krull Dimension (see the Krull Dimension section of Foundations) we prove the following result (provided we consider the lattice of radical ideals ordered by *reverse* inclusion).

**Proposition 3.22.3**
*Let $A$ be a ring then the lattice of radical ideals $\operatorname{Rad}(A)$ is distributive, that is we have equality*

$$\mathfrak{r}_1 \cap \sqrt{\mathfrak{r}_2 + \mathfrak{r}_3} = \sqrt{\mathfrak{r}_1 \cap \mathfrak{r}_2 + \mathfrak{r}_1 \cap \mathfrak{r}_3}$$

*Furthermore the meet-prime radical ideals are precisely the prime ideals. Therefore the lattice of radical ideals of a finite-dimensional Noetherian ring, ordered by **reverse inclusion**, is a Krull Lattice.*

*Proof.* Clearly it's enough to show that LHS $\subseteq$ RHS. Suppose $x \in$ LHS then $x \in \mathfrak{r}_1$ and $x^n = a + b$ where $a \in \mathfrak{r}_2$ and $b \in \mathfrak{r}_3$. Then $x^{n+1} = ax + bx \in \mathfrak{r}_1 \cap \mathfrak{r}_2 + \mathfrak{r}_1 \cap \mathfrak{r}_3$ whence $x \in$ RHS.

We've shown that prime ideals are meet-prime (3.3.40). Suppose $\mathfrak{r}$ is a meet-prime radical ideal, and $fg \in \mathfrak{r}$. Then we claim that

$$\sqrt{\mathfrak{r} + (f)} \cap \sqrt{\mathfrak{r} + (g)} \subseteq \mathfrak{r}$$

For $x \in \text{LHS} \implies x^n \in \mathfrak{r} + (f)$ and $x^m \in \mathfrak{r} + (g) \implies x^{n+m} \in \mathfrak{r} \implies x \in \mathfrak{r}$. As $\mathfrak{r}$ is meet-prime then for example $\sqrt{\mathfrak{r} + (f)} \subseteq \mathfrak{r}$ and in particular $f \in \mathfrak{r}$. Therefore $\mathfrak{r}$ is also prime. ∎

© David Rufino CC BY-NC-ND 4.0

**Proposition 3.22.4 (Simple properties)**
*The following properties of Krull dimension hold*

*a)* $\dim A = \dim A/N(A)$
*b)* $\mathrm{ht}(\mathfrak{p}) = \dim A_{\mathfrak{p}}$
*c)* $\dim \mathfrak{a} = \dim A/\mathfrak{a}$
*d)* $\dim \mathfrak{a} = \dim \mathfrak{a}/\mathfrak{b}$ *for any ideal* $\mathfrak{b} \subseteq \mathfrak{a}$
*e)* $\dim A = \sup_{\mathfrak{p}} \dim A_{\mathfrak{p}}$
*f)* ***codimension inequality*** $\mathrm{ht}(\mathfrak{p}) \geq \mathrm{ht}(\mathfrak{p}/\mathfrak{q}) + \mathrm{ht}(\mathfrak{q})$
*g)* $\dim k = 0$ *for any field* $k$
*h) A principal ideal domain* $A$ *which is not a field has dimension* 1

*Proof.* a) By (3.3.56) there is an order-isomorphism between prime ideals of $A$ containing $N(A)$ and prime ideals of $A/N(A)$. However by (3.3.46) all prime ideals of $A$ contain $N(A)$, so there is a bijection between chains of $A$ and chains of $A/N(A)$, and the result follows.

b) This follows similarly from (3.6.42).

c) This follows similarly from (3.3.56).

d) $\dim \mathfrak{a} = \dim A/\mathfrak{a} = \dim(A/\mathfrak{a})/(\mathfrak{a}/\mathfrak{b}) = \dim \mathfrak{a}/\mathfrak{b}$

e) This follows from (2.5.6)

f) This follows from (2.5.6)

g) The only (prime) ideal is (0)

h) By (...) every prime ideal (besides (0)) is maximal so every chain has length at most 1. ∎

**Lemma 3.22.5**
*Let* $A$ *be a UFD and* $\mathfrak{p} \lhd A$ *a non-zero prime ideal. Then it contains a non-zero principal prime ideal* $(p)$.

*In particular* $\mathrm{ht}(\mathfrak{p}) = 1$ *if and only if it is principal.*

*Proof.* Choose $0 \neq f \in \mathfrak{p}$. Then by definition it has a factorization into primes, and at least one must be in $\mathfrak{p}$, say $p$. Then $(p)$ is prime by (3.17.9). Therefore if $\mathrm{ht}(\mathfrak{p}) = 1$ then it is equal to $(p)$.

Conversely if $\mathfrak{q} \subseteq (p)$ then $(q) \subseteq \mathfrak{q} \subseteq (p)$ for $q$ prime, which implies $p \mid q$. As $q$ is irreducible (...) then $(p) = (q) = \mathfrak{q}$. Therefore $\mathrm{ht}((p)) = 1$. ∎

**Proposition 3.22.6**
*Let* $A$ *be a Noetherian ring and* $\mathfrak{a}$ *a proper ideal. Then there are only finitely many minimal primes containing* $\mathfrak{a}$*, say* $\mathfrak{p}_1, ..., \mathfrak{p}_n$*. Further we have a decomposition*

$$\sqrt{\mathfrak{a}} = \bigcap_{i=1}^{n} \mathfrak{p}_i$$

*which is irredundant (and the only such decomposition). Furthermore*

$$\mathrm{ht}(\mathfrak{a}) = \min_i \mathrm{ht}(\mathfrak{p}_i)$$

$$\dim(\mathfrak{a}) = \max_i \dim(\mathfrak{p}_i)$$

*Proof.* This follows from (2.4.7) and (3.22.3) applied to the radical ideal $\sqrt{a}$. ∎

**Remark 3.22.7**
*This is essentially the proof of [Kap74] (Theorem 87, 88).*

We've noted in general (3.22.4) that the so-called codimension formula does not hold. However it holds in the following case, for essentially trivial reasons.

**Proposition 3.22.8 (Codimension 1 formula)**
*Let* $A$ *be an irreducible Noetherian ring of finite Krull Dimension, and* $\mathfrak{a}$ *such that* $\dim(\mathfrak{a}) = \dim(A) - 1$ *then* $\mathrm{ht}(\mathfrak{a}) = 1$.

*Proof.* Apply (2.5.7). ∎

© David Rufino CC BY-NC-ND 4.0

In practice most rings of geometric interest are catenary or quasi-biequidimensional. We recall some properties and equivalent criteria for these cases.

**Proposition 3.22.9 (Catenary Criteria)**
*Let $A$ be a ring. Then the following are equivalent*

*a) $A$ is catenary*
*b) For all prime ideals $\mathfrak{r} \subseteq \mathfrak{q} \subseteq \mathfrak{p}$ the following holds*

$$\operatorname{ht}(\mathfrak{p}/\mathfrak{r}) = \operatorname{ht}(\mathfrak{p}/\mathfrak{q}) + \operatorname{ht}(\mathfrak{q}/\mathfrak{r})$$

*When $A$ is irreducible this is equivalent to the following condition*

$$\operatorname{ht}(\mathfrak{p}) = \operatorname{ht}(\mathfrak{q}) + \operatorname{ht}(\mathfrak{p}/\mathfrak{q})$$

*Proof.* This is restatement of (2.5.9). ∎

**Proposition 3.22.10 (Biequidimensional Criteria)**
*Let $A$ be a ring. Then the following are equivalent*

*a) $A$ is quasi-biequidimensional*
*b) $A$ is catenary and $A/\mathfrak{p}$ is equicodimensional for every minimal prime $\mathfrak{p}$*
*c) $A$ satisfies the formula*

$$\dim \mathfrak{q} = \dim \mathfrak{p} + \operatorname{ht}(\mathfrak{p}/\mathfrak{q}) \quad \forall \mathfrak{q} \subseteq \mathfrak{p}$$

*d) $A$ satisfies c) whenever $\operatorname{ht}(\mathfrak{p}/\mathfrak{q}) = 1$ (i.e. $\mathfrak{q} \subsetneq \mathfrak{p}$ is saturated)*

*Furthermore*

$$\operatorname{ht}(\mathfrak{p}) = \operatorname{ht}(\mathfrak{p}/\mathfrak{q}) + \operatorname{ht}(\mathfrak{q})$$

*Proof.* This is a restatement of (2.5.11). ∎

**Proposition 3.22.11 (Irreducible Biequidimensional Criteria)**
*When $A$ is irreducible the following are equivalent*

*a) $A$ is biequidimensional*
*b) $A$ is quasi-biequidimensional*
*c) $A$ is catenary and equicodimensional*
*d) $A$ satisfies the formula*

$$\dim \mathfrak{q} = \dim \mathfrak{p} + \operatorname{ht}(\mathfrak{p}/\mathfrak{q}) \quad \forall \mathfrak{q} \subseteq \mathfrak{p}$$

*e) $A$ satisfies d) whenever $\operatorname{ht}(\mathfrak{p}/\mathfrak{q}) = 1$ (i.e. $\mathfrak{q} \subsetneq \mathfrak{p}$ is saturated)*

*Proof.* This is a restatement of (2.5.14). ∎

**Proposition 3.22.12 (Codimension Formula)**
*Let $A$ be a quasi-biequidimensional ring, $\mathfrak{a}$ an ideal and $\mathfrak{p} \subset \mathfrak{a}$ a prime ideal. Then the following properties hold*

$$\operatorname{ht}(\mathfrak{a}) = \operatorname{ht}(\mathfrak{a}/\mathfrak{p}) + \operatorname{ht}(\mathfrak{p})$$
$$\dim \mathfrak{p} = \dim \mathfrak{a} + \operatorname{ht}(\mathfrak{a}/\mathfrak{p})$$

*When $A$ is biequidimensional (e.g. quasi-biequidimensional and integral), then for all ideals $\mathfrak{a}$ we have the relation*

$$\dim A = \dim \mathfrak{a} + \operatorname{ht}(\mathfrak{a})$$

*Proof.* This is a restatement of (2.5.13). ∎

© David Rufino CC BY-NC-ND 4.0

# 3.23 Lying over, Incomparability, Going Up and Going Down

**Definition 3.23.1 (Lying over / Going up)**
*Let $\phi : A \to B$ be a ring map and $\mathfrak{p}$ and $\mathfrak{q}$ primes of $A$ and $B$ respectively*

a) $\mathfrak{q}$ ***lies over*** $\mathfrak{p}$, *or* $\mathfrak{p}$ ***lies under*** $\mathfrak{q}$ *if* $\mathfrak{p} = \phi^{-1}(\mathfrak{q}) = \mathfrak{q}^c$. *When $A \subseteq B$ and $\phi$ is the identity then this is equivalent to saying $\mathfrak{p} = \mathfrak{q} \cap A$.*

**Definition 3.23.2 (Lying Over / Going Up / Incomparability)**
*Let $\phi : A \to B$ be a ring map. We say that it has the*

a) ***lying over property*** *if every prime ideal $\mathfrak{p} \supseteq \ker(\phi)$ has a prime $\mathfrak{q}$ lying over it. NB $\ker(\phi) \subseteq \mathfrak{p}$ is a necessary condition for $\mathfrak{p}$ to be a contraction and is equivalent to $B_{\mathfrak{p}} \neq 0$.*
b) ***going up property*** *if for every pair of prime ideals $\mathfrak{p} \subsetneq \mathfrak{p}'$ in $A$ and $\mathfrak{q} \lhd B$ lying over $\mathfrak{p}$, there exists a prime ideal $\mathfrak{q}'$ such that $\mathfrak{q} \subsetneq \mathfrak{q}'$ and $\mathfrak{q}'$ lies over $\mathfrak{p}'$.*
c) ***incomparability property*** *if for every pair of prime ideals $\mathfrak{q}, \mathfrak{q}' \lhd B$ then $\mathfrak{q} \subsetneq \mathfrak{q}' \implies \phi^{-1}(\mathfrak{q}) \subsetneq \phi^{-1}(\mathfrak{q}')$*
d) ***going down property*** *if for every pair of prime ideals $\mathfrak{p}' \subsetneq \mathfrak{p}$ in $A$ and $\mathfrak{q} \lhd B$ lying over $\mathfrak{p}$, there exists a prime ideal $\mathfrak{q}'$ such that $\mathfrak{q}' \subsetneq \mathfrak{q}$ and $\mathfrak{q}'$ lies over $\mathfrak{p}'$.*

**Remark 3.23.3**
*It's possible to interpret these geometrically in terms of the map $\phi_\star : \operatorname{Spec}(B) \to \operatorname{Spec}(A)$.*

- *Lying over - $\phi_\star$ is surjective onto $V(\ker(\phi))$*
- *Going up - $\phi_\star$ is closed*
- *Incomparability - fibres have dimension 0*
- *Going down (and finite presentation) - $\phi_\star$ is open*

The main result of this section is the correspondence between primes lying over $\mathfrak{p}$ and primes of the ring $B_{\mathfrak{p}}/\mathfrak{p}B_{\mathfrak{p}}$ (3.23.5). As a preliminary result we consider conditions under which we can strengthen $\mathfrak{p} \subseteq \mathfrak{q}^c$ to $\mathfrak{p} = \mathfrak{q}^c$, as the former condition is somewhat easier to satisfy.

**Lemma 3.23.4 (Lying over criteria)**
*Let $\phi : A \to B$ be a ring map, $\mathfrak{p} \lhd A$ prime such that $\ker(\phi) \subseteq \mathfrak{p}$ and $\mathfrak{q} \lhd B$. Then*

$$\mathfrak{p} = \mathfrak{q}^c \iff \mathfrak{p}^e \subseteq \mathfrak{q} \text{ and } \mathfrak{q} \cap \phi(A \setminus \mathfrak{p}) = \varnothing$$

*In particular*

$$\mathfrak{p} = \mathfrak{p}^{ec} \text{ is contracted} \iff \mathfrak{p}^e \cap \phi(A \setminus \mathfrak{p}) = \varnothing$$

*Proof.* Recall (3.3.52) that in general $\mathfrak{p} \subseteq \mathfrak{q}^c \iff \mathfrak{p}^e \subseteq \mathfrak{q}$. The first equivalence is then clear because $x \in \mathfrak{q}^c \setminus \mathfrak{p} \iff \phi(x) \in \mathfrak{q} \cap \phi(A \setminus \mathfrak{p})$. For $\phi(x) \in \mathfrak{q} \cap \phi(A \setminus \mathfrak{p}) \implies \phi(x) = \phi(y)$ for $y \notin \mathfrak{p}$ and $x \in \mathfrak{q}^c$. This implies $x - y \in \ker(\phi) \subseteq \mathfrak{p}$ whence $x \notin \mathfrak{p}$ as required.

The final statement follows by considering $\mathfrak{q} = \mathfrak{p}^e$. ∎

**Proposition 3.23.5 (Lying over correspondence)**
*Let $\phi : A \to B$ be a ring map and $\mathfrak{p} \lhd A$ a prime ideal s.t. $\ker(\phi) \subseteq \mathfrak{p}$. Then there is a order-preserving correspondence of prime ideals*

$$\begin{aligned} \{\mathfrak{q} \mid \mathfrak{q} \text{ lies above } \mathfrak{p}\} &\longleftrightarrow \left\{\mathfrak{q}' \lhd B_{\mathfrak{p}} \;\middle|\; \mathfrak{p}B_{\mathfrak{p}} \subseteq \mathfrak{q}'\right\} &&\longleftrightarrow \left\{\mathfrak{q}'' \lhd B_{\mathfrak{p}}/\mathfrak{p}B_{\mathfrak{p}}\right\} \\ \mathfrak{q} &\longrightarrow \mathfrak{q}B_{\mathfrak{p}} && \longrightarrow \mathfrak{q}B_{\mathfrak{p}}/\mathfrak{p}B_{\mathfrak{p}} \end{aligned}$$

*In particular TFAE*

a) $\mathfrak{p}$ *lies under a prime* $\mathfrak{q}$
b) $B_{\mathfrak{p}}/\mathfrak{p}B_{\mathfrak{p}} \neq 0$
c) $\mathfrak{p} = \mathfrak{p}^{ec}$ *is contracted*

*NB c) is a-priori weaker than a).*

*Proof.* Recall $B_{\mathfrak{p}} := S^{-1}B$ and $\mathfrak{p}B_{\mathfrak{p}} := \mathfrak{p}^e B_{\mathfrak{p}} = S^{-1}\mathfrak{p}^e$ where $S := \phi(A \setminus \mathfrak{p})$.

By (3.23.4) $\mathfrak{q}$ lies above $\mathfrak{p}$ if and only if $\mathfrak{p}^e \subseteq \mathfrak{q}$ and $\mathfrak{q} \cap S = \varnothing$. The first correspondence then follows from (3.6.19) and the second from (3.3.56).

a) $\iff$ b) This follows from the correspondence, since a ring without any non-zero prime ideals is simply the zero-ring.

© David Rufino CC BY-NC-ND 4.0

b) $\iff$ c) Observe $\mathfrak{p} = \mathfrak{p}^{ec} \iff \mathfrak{p}^e \cap S = \varnothing$ by (3.23.4), which holds if and only if $\mathfrak{p}B_\mathfrak{p} \neq B_\mathfrak{p}$ by (3.6.18) e). ∎

**Remark 3.23.6**
*Geometrically this is an explicit representation of the fiber as a prime spectrum*

$$\operatorname{Spec}(\phi)^{-1}(\mathfrak{p}) \longleftrightarrow \operatorname{Spec}\big(B_\mathfrak{p}/\mathfrak{p}B_\mathfrak{p}\big)$$

**Proposition 3.23.7 (Injective Ring Maps are Dense)**
*Let $\phi : A \to B$ be a ring map and $\mathfrak{p} \lhd A$ a minimal prime over $\ker(\phi)$. Then there exists a prime ideal $\mathfrak{q} \lhd B$ such that $\mathfrak{q}^c = \mathfrak{p}$. Furthermore $\mathfrak{q}$ may be taken to be minimal.*

*Proof.* We may reduce to the case $\phi$ is injective by replacing $A$ with $A/\ker(\phi)$. We have the commutative diagram

$$\begin{array}{ccc} A & \xrightarrow{\phi} & B \\ \downarrow{\scriptstyle i_S} & & \downarrow{\scriptstyle i_T} \\ A_\mathfrak{p} & \overset{\tilde{\phi}}{\dashrightarrow} & T^{-1}B \end{array}$$

where $S \coloneqq A \setminus \mathfrak{p}$ and $T \coloneqq \phi(S)$. By (3.6.5) there exists a homomorphism $\tilde{\phi}$. Define $\mathfrak{m} \coloneqq \mathfrak{p}A_\mathfrak{p}$ to be the unique maximal ideal of $A_\mathfrak{p}$. By (3.6.18) it is a minimal prime ideal and therefore the unique prime ideal. By assumption $0 \notin T$ and so $T^{-1}B \neq 0$. $T^{-1}B$ has a maximal ideal $\mathfrak{n}$ by (3.3.37). By uniqueness we find $\tilde{\phi}^{-1}(\mathfrak{n}) = \mathfrak{m}$. Therefore $\mathfrak{q} \coloneqq i_T^{-1}(\mathfrak{n})$ is a prime ideal such that $\mathfrak{q}^c = \mathfrak{p}$.

By (3.3.44) there is a minimal prime $\mathfrak{r} \subset \mathfrak{q}$ which by minimality also satisfies $\mathfrak{r}^c = \mathfrak{p}$. ∎

© David Rufino CC BY-NC-ND 4.0

# 3.24 Integral Ring Extensions

**Definition 3.24.1 (Integral Element)**
*Let $\phi : A \to B$ be a ring map and $\alpha \in B$. Then we say $\alpha$ is **algebraic** over $A$ if $m(\alpha) = 0$ for some polynomial $m(X) \in A[X]$.*

*Furthermore we say that $\alpha$ is **integral** over $A$ if $m(X)$ may be chosen to be monic.*

*Note often we assume $A \subseteq B$ and $\phi$ is the identity.*

**Definition 3.24.2 (Ring Extensions)**
*Let $\phi : A \to B$ be a ring map (so that $B$ is an $A$-algebra). Then we say $\phi$ is*

- ***finite** if $B$ is finite as an $A$-module*
- ***finite-type** if $B$ is finitely generated as an $A$-algebra*
- ***integral** if every element of $B$ is integral over $A$*

*When $A \subseteq B$ and $\phi$ is the identity then we say that $B$ is respectively finite over $A$, finite-type over $A$ or integral over $A$.*

We note the trivial implication

$$\text{finite} \implies \text{finite type}$$

For example $k[X]$ is a ring of finite type over $k$, but certainly not finite. The following criterion for integrality is fundamental.

**Proposition 3.24.3**
*Let $\phi : A \to B$ be a ring map and $b \in B$. Then the following are equivalent*

*a) $b$ is integral over $A$*
*b) $\phi(A)[b]$ is a finite $A$-module*
*c) $\phi(A)[b]$ is contained in a subring $C$ of $B$ which is a finite $A$-module*
*d) There exists a $\phi(A)[b]$-module $M$ which is faithful and finite as an $A$-module*

*Proof.* Note that the subring $C$ in c) is a faithful $A$-module, so the only non-trivial step is d) $\implies$ a). This is the usual "determinant trick". We apply (3.18.4) by considering $\psi_b \in \text{End}_A(M)$ to be multiplication by $b$. Then we have some monic polynomial $P(X) \in A[X]$ such that $P(\psi_b) = 0$, whence $P^\phi(b)m = 0$ for all $m \in M$. Since $M$ is faithful, then we have $P(b) = 0$ as required. ∎

**Proposition 3.24.4 (Finite $\iff$ finite-type and integral)**
*Let $\phi : A \to B$ be a ring map and $b_1, ..., b_n \in B$ integral over $A$. Then the ring homomorphism $\phi : A \to \phi(A)[b_1, ..., b_n]$ is finite and integral.*

*In particular if $\phi$ is integral and of finite type if and only if it is finite.*

*Proof.* We assume without loss of generality that $A \subseteq B$ and $\phi$ is the identity map. Consider a tower

$$A \subset A[b_1] \subset ... \subset A[b_1, ..., b_n] = B$$

We proceed inductively on $n$. Namely we assume that $A[b_1, ..., b_i]$ is a finite $A$-module. Then a-fortiori $A[b_1, ..., b_{i+1}]$ is integral over $A[b_1, ..., b_i]$. Therefore by the previous Proposition it is a finite $A[b_1, ..., b_i]$-module and therefore a finite $A$-module (by (3.15.5)).

For any $b \in A[b_1, ..., b_n]$ we have $A[b] \subset A[b_1, ..., b_n]$ so by (3.24.3) we have $b$ is integral over $A$.

It then follows that $B$ integral and finite type $\implies$ $B$ is finite (as an $A$-module). Conversely if $B$ is finite then it is clearly finitely-generated. Further for any $b \in B$ then $A[b]$ is contained in the ring $B$ which is finite as an $A$-module, and therefore is $b$ is integral by (3.24.3). ∎

**Proposition 3.24.5 (Transitivity property)**
*Let $\phi : A \to B$ and $\psi : B \to C$ be ring maps.*

*If $c \in C$ is integral over $B$, then it is integral over $A$ (with respect to the ring map $\psi \circ \phi : A \to C$)*

*In particular if $\phi$ integral and $\psi$ integral (e.g. surjective) $\implies$ $\psi \circ \phi$ integral.*

© David Rufino CC BY-NC-ND 4.0

*Proof.* Suppose $c \in C$ is integral over $B$. Let $b_0, ..., b_{n-1}$ be the coefficients of the integral relation then clearly $c$ is integral over $B' \coloneqq A[b_0, ..., b_{n-1}]$. By (3.24.3) $B'$ is a finite $A$-module and $B'[c]$ is a finite $B'$-module. Therefore $B'[c]$ is a finite $A$-module and by (3.24.3) we have $c$ is integral over $A$. ∎

**Definition 3.24.6 (Integrally Closed)**
*Let $A \subset B$ be a subring, then we say that $A$ is **integrally closed** in $B$ if*

$$b \in B \text{ integral over } A \implies b \in A.$$

*We say that an integral domain $A$ is **integrally closed** if it is integrally closed in its quotient field.*

**Proposition 3.24.7 (Integral Closure)**
*Let $A$ be a subring of a ring $B$. Then the set of elements of $B$ integral over $A$ (the **integral closure**) is a subring of $B$. Denote this by $\overline{A}^B$ or just $\overline{A}$ when the overring $B$ is unambiguous.*

*Further $\overline{A}^B$ is **integrally closed** in $B$.*

*Proof.* Let $\alpha, \beta \in B$ be integral over $A$. Then by (3.24.4) the subring $A[\alpha, \beta]$ is a finite $A$-module containing $\alpha \pm \beta$ and $\alpha\beta$. Therefore by (3.24.3) they are also integral over $A$.

Clearly $\overline{A}$ is integral over $A$ so by (3.24.5) it is integrally closed. ∎

**Lemma 3.24.8**
*Let $A$ be a subring of a field $K$ and $S \subset A$ a multiplicatively closed subset. If $x \in K$ is integral over $S^{-1}A$ then $sx$ is integral over $A$ for some $s \in S$.*

*Conversely if $y \in K$ is integral over $A$ then $\frac{y}{s}$ is integral over $S^{-1}A$ for every $s \in S$.*

*Proof.* Suppose $x \in K$ is integral over $S^{-1}A$ then we have by definition

$$x^n + \sum_{j=0}^{n-1} \frac{a_j}{s_j} x^j = 0$$

for $a_j \in A$ and $s_j \in S$. Let $s \coloneqq s_0 ... s_{n-1}$ and multiply by $s^n$ to find

$$y^n + \sum_{j=0}^{n-1} s^{n-j-1} \frac{s}{s_j} a_j y^j = 0$$

where $y = sx$. This shows that $sx$ is integral over $A$

Similarly if $y \in K$ is integral over $A$ then

$$y^n + \sum_{j=0}^{n-1} a_j y^j = 0$$

for some $a_j \in A$. Then let $x = \frac{y}{s}$ we have

$$x^n + \sum_{j=0}^{n-1} \frac{a_j}{s^{n-j}} x^j = 0$$

whence $x$ is integral over $S^{-1}A$. ∎

**Lemma 3.24.9**
*Let $A \subset K$ be a subring of a field. Then $\overline{S^{-1}A}^K = S^{-1}\overline{A}^K$.*

*In particular if $A$ is integrally closed in $K$ then so is $S^{-1}A$.*

*Proof.* This follows directly from (3.24.8). ∎

**Proposition 3.24.10 (Normality is Local)**
*Let $A$ be an integral domain. Then the following are equivalent*

*a) $A$ is integrally closed*
*b) $A_{\mathfrak{p}}$ is integrally closed for every prime ideal $\mathfrak{p}$*
*c) $A_{\mathfrak{m}}$ is integrally closed for every maximal ideal $\mathfrak{m}$*

*Proof.* a) $\implies$ b) This is simply (3.24.9).

b) $\implies$ c) is immediate.

© David Rufino CC BY-NC-ND 4.0

c) $\implies$ a) Suppose that $x \in K$ is integral over $A$ and consider the ideal $\mathfrak{a} := \{a \in A \mid ax \in A\}$. Then by definition it is a non-zero ideal, and we wish to show that $\mathfrak{a} = A$. Suppose not, then by (...) it is contained in some maximal ideal $\mathfrak{m}$. A-fortiori $x$ is integral over $A_{\mathfrak{m}}$ whence by assumption $x \in A_{\mathfrak{m}}$. This means there is some $s \notin \mathfrak{m}$ such that $sx \in A$, that is $s \in \mathfrak{a}$ a contradiction. ∎

**Proposition 3.24.11 (Integral closure of an ideal)**
*Let $A \subset B$ be a subring and $\mathfrak{a} \lhd A$ an ideal. Then for $b \in B$ TFAE*

*a) $b$ integral over $\mathfrak{a}$*
*b) $b^n$ integral over $\mathfrak{a}$ for some $n \geq 1$*
*c) $b \in \sqrt{\mathfrak{a}\overline{A}}$*

*In particular $\overline{\mathfrak{a}}$ is an ideal of $\overline{A}$.*

*Furthermore if $A$ is integrally closed in $B$ then $\overline{\mathfrak{a}} = \sqrt{\mathfrak{a}}$.*

*Proof.* It's clear that a) $\iff$ b). For a) $\implies$ c) consider the integral relation

$$b^n + a_{n-1}b^{n-1} + ...a_0 = 0 \quad a_i \in \mathfrak{a}$$

By (3.24.7) $\overline{A}$ is a subring, and by assumption $b \in \overline{A}$. Therefore $b^k \in \overline{A}$, and the integral relation shows that $b^n \in \mathfrak{a}\overline{A}$ as required.

For c) $\implies$ b) suppose $b \in \sqrt{\mathfrak{a}\overline{A}}$ then $b^n = \sum_{i=1}^n a_i x_i$ for $a_i \in \mathfrak{a}$ and $x_i \in \overline{A}$. Let $B' := A[x_1, ..., x_n]$, which is a finite $A$-submodule by (3.24.4). Let $\phi \in \mathrm{End}_A(B')$ denote multiplication by $b^n$ then $\phi(B') \subseteq \mathfrak{a}B'$ so by (3.18.4) $\phi$ satisfies a monic polynomial with coefficients in $\mathfrak{a}$. In particular $b^n$ is integral over $\mathfrak{a}$. ∎

**Proposition 3.24.12**
*A UFD is integrally closed.*

*In particular polynomial ring over a UFD is integrally closed.*

*Proof.* **TODO** ∎

The following criterion is also useful:

**Lemma 3.24.13 (Integral Criterion II)**
*Let $\phi : A \to B$ be a ring map. Suppose $x \in B$ is invertible, then $x$ is integral over $A$ if and only if $x \in \phi(A)[x^{-1}]$*

*Proof.* Suppose $x \in \phi(A)[x^{-1}]$ then

$$x = \phi(a_0) + \phi(a_1)x^{-1} + ... + \phi(a_n)x^{-n}$$

Multiply by $x^n$ to deduce an integral equation. Conversely suppose $x \in B$ is integral over $A$ then by definition

$$x^n + \phi(a_{n-1})x^{n-1} + ... + \phi(a_0) = 0$$

Multiply by $x^{-(n-1)}$ to deduce $x \in \phi(A)[x^{-1}]$ ∎

**Proposition 3.24.14 (Integral extension preserves field property)**
*Let $\phi : A \hookrightarrow B$ be an injective, integral ring map. Then $B$ is a field if and only if $A$ is a field.*

*Proof.* As $\phi$ is injective, it induces an isomorphism between $A$ and $\phi(A)$. So we may assume without loss of generality that $A \subseteq B$ and $\phi$ is the identity.

Suppose $B$ is a field and $x \in A$. Then $x^{-1} \in B$ is integral over $A$ by hypothesis, so by the previous Lemma $x^{-1} \in A[x] \subseteq A$. Therefore $A$ is a field.

Conversely suppose $A$ is a field and $0 \neq x \in B$. Then by hypothesis $x$ is integral over $A$, that is

$$x^n + a_{n-1}x^{n-1} + ... + a_1 x + a_0 = 0$$

Choose the degree $n$ to be minimal. We claim $a_0 \neq 0$, for if $a_0 = 0$ we may cancel $x$ to obtain an integral relation of smaller degree. Therefore

$$-x(x^{n-1} + a_{n-1}x^{n-2} + ... + a_1)a_0^{-1} = 1$$

and in particular $x$ is invertible. ∎

**Proposition 3.24.15**
*Let $\phi : A \to B$ be an integral ring map. Then*

© David Rufino CC BY-NC-ND 4.0

*a) If $\phi^{-1}(\mathfrak{b}) \subseteq \mathfrak{a}$ then the induced ring map $A/\mathfrak{a} \to B/\mathfrak{b}$ (see (3.3.56)) is integral.*
*b) If $S$ is a multiplicatively closed subset of $A$ and $T \coloneqq \phi(S)$, then the induced ring map $S^{-1}A \to T^{-1}B$ is also integral.*

**Proposition 3.24.16 (Maximal ideals under integral extension)**
*Let $\phi : A \to B$ be an integral ring map. Suppose $\mathfrak{q}$ lies above $\mathfrak{p}$. Then*

$$\mathfrak{p} \text{ is maximal} \iff \mathfrak{q} \text{ is maximal}$$

*Proof.* The map $\phi : A \to B$ induces an injective map $A/\mathfrak{p} \hookrightarrow B/\mathfrak{q}$ of integral domains by (3.3.56). By (3.24.15) this map is also integral. Note $A/\mathfrak{p}$ (resp. $B/\mathfrak{q}$) is a field if and only if $\mathfrak{p}$ (resp. $\mathfrak{q}$) is a maximal ideal by (3.3.59). Then we may apply (3.24.14) to show the equivalence. ∎

**Proposition 3.24.17 (Properties of integral extensions)**
*Let $\phi : A \to B$ be an integral ring map then it has*

*a) the **Lying Over** property*
*b) the **Incomparability** property*
*c) the **Going Up** property*

*Proof.* For any prime ideal $\mathfrak{p} \lhd A$ we have the commutative diagram

$$\begin{array}{ccc} A & \xrightarrow{\phi} & B \\ \downarrow{\scriptstyle i_S} & & \downarrow{\scriptstyle i_T} \\ A_{\mathfrak{p}} & \xrightarrow{\tilde{\phi}} & B_{\mathfrak{p}} \end{array}$$

where $S \coloneqq A \setminus \mathfrak{p}$ and $T \coloneqq \phi(S)$. By (3.6.5) there exists a morphism $\tilde{\phi}$, and by (3.24.15) it is integral. Define $\mathfrak{m} \coloneqq \mathfrak{p}A_{\mathfrak{p}}$ to be the unique maximal ideal of $A_{\mathfrak{p}}$.

a) As we assume $\ker(\phi) \subseteq \mathfrak{p}$ we know $B_{\mathfrak{p}} \neq 0$ (3.6.40). Let $\mathfrak{n}$ be a maximal (and hence prime) ideal of $B_{\mathfrak{p}}$. Then $\mathfrak{q} \coloneqq i_T^{-1}(\mathfrak{n})$ is a prime ideal of $B$ such that $\mathfrak{q} \cap T = \varnothing$. In addition by (3.24.16) $\tilde{\phi}^{-1}(\mathfrak{n})$ is a maximal ideal, and therefore by uniqueness $\mathfrak{m} = \tilde{\phi}^{-1}(\mathfrak{n})$. By commutativity of the diagram we then have $\phi^{-1}(\mathfrak{q}) = \mathfrak{p}$ as required.

(Stacks) As an alternative argument to show existence of $\mathfrak{q}$ by (3.23.5) it's enough to show that $\mathfrak{p}B_{\mathfrak{p}}$ is proper (by assumption $\ker(\phi) \subseteq \mathfrak{p}$ so $B_{\mathfrak{p}} \neq 0$). By the diagram above $\mathfrak{p}B_{\mathfrak{p}} = \tilde{\phi}(\mathfrak{p}A_{\mathfrak{p}})B_{\mathfrak{p}}$. Therefore it's enough to consider the case $(A, \mathfrak{m})$ local and to show $\phi(\mathfrak{m})B$ is proper. Suppose $1 \in \phi(\mathfrak{m})B$ then

$$1 = \sum_{i=1}^{n} \phi(a_i)b_i \quad a_i \in \mathfrak{m}_A \; b_i \in B.$$

By (3.24.4) the subring $B' \coloneqq \phi(A)[b_1, ..., b_n] \subset B$ is a finite $A$-module. Furthermore $1 \in \mathfrak{m}B'$ whence $\mathfrak{m}B' = B'$ and by Nakayama's Lemma (3.21.5) $B' = 0$, a contradiction.

b) Suppose $\mathfrak{p} = \phi^{-1}(\mathfrak{q}) = \phi^{-1}(\mathfrak{q}')$ and $\mathfrak{q} \subseteq \mathfrak{q}'$. Let $\mathfrak{n} = \mathfrak{q}B_{\mathfrak{p}}$ and $\mathfrak{n}' = \mathfrak{q}'B_{\mathfrak{p}}$. Clearly $\mathfrak{n} \subseteq \mathfrak{n}'$. By commutativity of the diagram $i_S^{-1}(\tilde{\phi}^{-1}(\mathfrak{n})) = \phi^{-1}(\mathfrak{q}) = \mathfrak{p}$. By (3.6.18) extending the ideals to $A_{\mathfrak{p}}$ shows $\tilde{\phi}^{-1}(\mathfrak{n}) = \mathfrak{m}$, and similarly for $\mathfrak{n}'$. By (3.24.16) both $\mathfrak{n}, \mathfrak{n}'$ are maximal so $\mathfrak{n} = \mathfrak{n}'$. By (3.6.19) $\mathfrak{q} = \mathfrak{q}'$.

c) Suppose we have prime ideals $\mathfrak{p} \subsetneq \mathfrak{p}'$ and $\mathfrak{q}$ is a prime ideal lying above $\mathfrak{p}$. Consider the commutative diagram

$$\begin{array}{ccc} A & \xrightarrow{\phi} & B \\ \downarrow & & \downarrow \\ A/\mathfrak{p} & \xrightarrow{\tilde{\phi}} & B/\mathfrak{q} \end{array}$$

The induced map $\tilde{\phi}$ is integral (3.24.16). By a) there is a prime ideal of $B/\mathfrak{q}$ lying above $\mathfrak{p}'/\mathfrak{p}$, which is of the form $\mathfrak{q}'/\mathfrak{q}$ for $\mathfrak{q} \subseteq \mathfrak{q}'$ prime (3.3.56). Then from the diagram we see $\phi^{-1}(\mathfrak{q}') = \mathfrak{p}'$ as required.

∎

**Proposition 3.24.18 (Coefficients of minimal polynomial)**
*Let $A \subseteq B$ be integral domains, $A$ is integrally closed, and define $K = \mathrm{Frac}(A)$ and $L = \mathrm{Frac}(B)$. For $b \in B$ integral over $\mathfrak{a} \lhd A$ we have the non-leading coefficients of $m_b(X)$ are integral over $\mathfrak{a}$ and therefore lie in $\sqrt{\mathfrak{a}}$.*

© David Rufino CC BY-NC-ND 4.0

*Note if $b$ is only assumed to be integral over $A$ then the coefficients of $m_b(X)$ lie in $A$.*

*Both* $\mathrm{Norm}_{L/K}(b)$ *and* $\mathrm{Tr}_{L/K}(b)$ *lie in* $A$.

*Proof.* Let $M/K$ be a normal closure for $L/K$ (...). By (3.24.11) the integral closure of $\mathfrak{a}$ in $M$ is simply $\sqrt{\mathfrak{a}}$. Then the minimal polynomial $m_b(X)$ splits completely in $M$ and by (3.19.66) all the roots $b_i$ are conjugate by $\mathrm{Aut}(M/K)$. In particular it's clear that $b_i$ are integral over $\mathfrak{a}$, and so lie in $\sqrt{\mathfrak{a}}$. The coefficients of $m_b(X)$ are polynomials in the $b_i$, and so by the observation above are also lie in $\sqrt{\mathfrak{a}}$ (and are integral over $\mathfrak{a}$).

We may consider the case $\mathfrak{a} = A$ to deduce that $m_b(X)$ is a monic polynomial with coefficients in $A$. By (3.19.128) then the coefficients of $\mathrm{Pc}_{L/K}(b; X)$ lie in $A$, and so by (3.19.123) so do the norm and trace. ∎

**Proposition 3.24.19 (Going Down)**
*Let $A \subseteq B$ be integral ring extension such that $A$ is an integrally closed domain and $B$ is an integral domain. Then it has the Going Down property.*

*Proof.* Let $\mathfrak{p} \subsetneq \mathfrak{p}'$ be prime ideals of $A$ and $\mathfrak{q}'$ a prime ideal lying over $\mathfrak{p}'$. We wish to find a prime ideal $\mathfrak{q} \subseteq \mathfrak{q}'$ lying over $\mathfrak{p}$ (clearly inclusion must be strict).

Consider the inclusion of rings $A \subseteq B_{\mathfrak{q}'}$. Then by (3.23.5) $\mathfrak{p}$ lies under a prime of $B_{\mathfrak{q}'}$ if and only if $\mathfrak{p} = \mathfrak{p}^{ec} = \mathfrak{p}B_{\mathfrak{q}'} \cap A$. If this is the case then it is of the form $\mathfrak{q}B_{\mathfrak{q}'}$ for some prime ideal $\mathfrak{q} \subseteq \mathfrak{q}'$ (3.6.42) of $B$. It's clear that $\mathfrak{q}$ lies over $\mathfrak{p}$.

Note in general that $\mathfrak{p} \subseteq \mathfrak{p}^{ec}$, so we only need to demonstrate the reverse inclusion. Choose $x \in \mathfrak{p}B_{\mathfrak{q}'} \cap A$. By (3.6.18) $\mathfrak{p}B_{\mathfrak{q}'} = S^{-1}(\mathfrak{p}B)$ where $S = B \setminus \mathfrak{q}'$.

Then $x = \frac{y}{s}$ for $y \in \mathfrak{p}B$ and $s \in B \setminus \mathfrak{q}'$. By (3.24.11) we have $y$ is integral over $\mathfrak{p}$ whence by (3.24.18) the minimal polynomial $m_{y,K}(X)$ is equal to

$$X^r + u_1 X^{r-1} + ... + u_r \quad u_i \in \mathfrak{p}$$

However $s = yx^{-1}$ and $x \in A \implies x^{-1} \in K$. So we can derive the minimal polynomial $m_{s,K}(X)$

$$X^r + v_1 X^{r-1} + ... + v_r \quad v_i := \frac{u_i}{x^i}$$

As $s$ is assumed to be integral over $A$ the coefficients must all lie in $A$, by (...). Consequently $v_i \in A$ and $v_i x^i \in \mathfrak{p}$ for all $i$. If $x \notin \mathfrak{p}$ then we have $v_i \in \mathfrak{p}$ for all $i$, and $s$ is integral over $\mathfrak{p}$. By the minimal polynomial we see that $s \in B\mathfrak{p} \subseteq B\mathfrak{p}' \subseteq \mathfrak{q}'$, which contradicts the choice of $s$. Therefore $x \in \mathfrak{p}$ as required. ∎

**Proposition 3.24.20**
*Let $B, C$ be commutative $A$-algebras such that $B$ is integral over $A$. Then $C \otimes_A B$ is integral over $C$.*

*Proof.* Let $\overline{C}$ be the subring of $C \otimes_A B$ of elements which are integral over $C$ (3.24.7). Consider an elementary tensor $c \otimes b$ then by definition there is $P \in A[X]$ such that $P(b) = 0$. Then

$$P(c \otimes b) = \sum_{i=0}^{n} a_i (c \otimes b)^i = \sum_{i=0}^{n} c^i \otimes a_i b^i = c^i \otimes \sum_i a_i b^i = c^i \otimes P(b) = 0$$

This shows that $\overline{C}$ contains the elementary tensors and therefore is equal to $C \otimes_A B$. ∎

**Lemma 3.24.21**
*Let $\phi : A \to B$ be a finite ring homomorphism. Suppose $B = (f_1, ..., f_n)$ and the composite map $\phi_i : A \to B_{f_i}$ is finite for all $i = 1...n$, then so is $\phi$.*

*Proof.* There are finitely many elements $\left\{ \frac{b_{ij}}{f_i^{n_{ij}}} \right\}_j$ which generate $B_{f_i}$ as an $A$-module. Let $B'$ be the $A$-submodule of $B$ generated by all the $b_{ij}$. Then $B'_{f_i} = B_{f_i}$ as $B$-modules. By (3.6.47) we have $B' = B$ and therefore $B$ is finitely generated as an $A$-module. ∎

**Proposition 3.24.22 (Krull Dimension is preserved under integral maps)**
*Let $\phi : A \to B$ be a ring map.*

*a) Going Up $\implies \dim B \geq \dim(A/\ker(\phi))$*
*b) Incomparability $\implies \dim B \leq \dim(A/\ker(\phi))$*

*In particular $\phi$ integral and injective implies $\dim A = \dim B$.*

*Proof.* Without loss of generality we can assume that $\phi$ is injective. The two cases follow by lifting chains of prime ideals from $A$ (resp. $B$) to $B$ (resp. $A$), and checking that distinct is maintained.

© David Rufino CC BY-NC-ND 4.0

The final statement follows from (3.24.17). ∎

**Corollary 3.24.23**
*Let $\phi : A \to B$ be integral and $\mathfrak{b} \lhd B$, then* $\dim \phi^{-1}(\mathfrak{b}) = \dim(\mathfrak{b})$.

## 3.24.1 Integral Closure in Finite Extensions

**Proposition 3.24.24**
*Let $L/K$ be an algebraic field extension and $A$ a subring with quotient field $K$. Then the integral closure $B \coloneqq \overline{A}^L$*

*a) has quotient field $L$,*
*b) spans $L$ as a $K$-vector space and*
*c) is integrally closed.*

*Furthermore* $\dim B = \dim A$.

*Proof.* We may show a),b) if we prove $L = (A^\star)^{-1}B$. Consider $x \in L$, then by definition it is integral over $K = (A^\star)^{-1}A$ so by (3.24.8) $ax \in B$, and therefore $x \in (A^\star)^{-1}B$ as required.

It is integrally closed by (3.24.7) and $\dim B = \dim A$ from (3.24.22). ∎

**Proposition 3.24.25**
*Let $L/K$ be a finite field extension, $A$ a subring with quotient field $K$ and $B = \overline{A}^L$. Suppose that $A$ is a principal ideal domain and $B$ is a finite $A$-module. Then $B$ is a finite free $A$-module of rank $[L : K]$.*

*Proof.* By (3.16.7) $B$ is a finite free $A$-module. Let $b_1, ..., b_r$ be an $A$-basis, then by (3.24.24) this spans $L$ as a $K$-vector space. Similarly by clearing denominators the given set is linearly independent, therefore by (...) $r = [L : K]$ as required. ∎

**Proposition 3.24.26 (Finite Separable Extension of Normal Domain)**
*Let $L/K$ be a finite separable extension, with $A$ an integrally closed Noetherian subring with quotient field $K$. Then the integral closure $\overline{A}^L$ is a finite $A$-module.*

*In particular $\overline{A}^L$ is Noetherian and* $\dim A = \dim \overline{A}^L$.

*If in addition $A$ is a principal ideal domain, then $\overline{A}^L$ is a free $A$-module of rank $[L : K]$.*

*Proof.* By (3.19.131) the trace map $L \times L \to K$ is a perfect pairing. By (3.24.24) we may choose $x_1, ..., x_n \in B$ a $K$-basis for $L$ where $n = [L : K]$. Denote $x_1^\star, ..., x_n^\star \in L$ by the (unique) basis dual under this pairing satisfying

$$\mathrm{Tr}_{L/K}(x_i x_j^\star) = \delta_{ij}$$

We claim that $B \coloneqq \overline{A}^L$ satisfies

$$B \subset Ax_1^\star + ... + Ax_n^\star.$$

For every $b \in B$ we have $b = \sum_{i=1}^n \lambda_i x_i^\star$ and some $\lambda_i \in K$. Then by linearity $\mathrm{Tr}_{L/K}(bx_i) = \lambda_i$. By (3.24.18) we have $\lambda_i \in A$. The right hand side is a finitely generated $A$-module, which is therefore Noetherian (3.15.4). $B$ is a submodule and therefore also finitely generated as an $A$-module (3.15.3).

Then by (3.15.4) $B$ is Noetherian as an $A$-module, therefore a-fortiori Noetherian as a $B$-module (or equivalently as a ring).

If $A$ is a principal ideal domain then $\overline{A}^L$ is a finite free $A$-module of rank $[L : K]$ by (3.24.25). ∎

**Lemma 3.24.27**
*Let $A \subset B \subset C$ be subrings. Then $\overline{A}^C \cap B = \overline{A}^B$.*

**Lemma 3.24.28**
*Let $A \subset B \subset C$ be rings such that $B$ is integral over $A$. Then $\overline{A}^C = \overline{B}^C$.*

**Lemma 3.24.29**
*Let $A$ be a Noetherian integral domain with quotient field $K$ and $L/K$ be a finite extension.*

*Suppose that $L'/L$ is a finite extension such $\overline{A}^{L'}$ is a finite $A$-module, then so is $\overline{A}^L$.*

*Proof.* By (3.15.4) $\overline{A}^{L'}$ is Noetherian as an $A$-module, whence by (3.24.27) $\overline{A}^L \subset \overline{A}^{L'}$ is a finitely-generated $A$-module. ∎

© David Rufino CC BY-NC-ND 4.0

# 3.25 Valuation Rings and Places

**Definition 3.25.1 (Valuation Ring)**
*A subring $A \subset K$ of a field $K$ is a **valuation ring for $K$** if for every $0 \neq x \in K$ either $x \in A$ or $x^{-1} \in A$ (or both). Such a ring is an integral domain and $K$ is necessarily a quotient field for $A$.*

*An integral domain $A$ is a **valuation ring** if it is a valuation ring for its quotient field.*

**Proposition 3.25.2 (Properties of valuation rings)**
*Let $A$ be a valuation ring and $K$ its quotient field then the following properties hold*

*a) $A$ is local ring*
*b) $x \in A \setminus A^\star \iff x \in \mathfrak{m} \iff x^{-1} \notin A$*
*c) $A$ is integrally closed in $K$*

*Proof.* We prove each in turn

a) By (3.20.3) we need to show that $\mathfrak{m} := A \setminus A^\star$ is an additive subgroup of $A$. Given $x, y \in \mathfrak{m}$, without loss of generality we may assume that $x, y$ are non-zero, and $\frac{x}{y} \in A$. Then $x + y = y\left(1 + \frac{x}{y}\right)$. If $(x + y) \in A^\star$ then $y \in A^\star$ a contradiction. Therefore $(x + y) \in \mathfrak{m}$ as required.
b) Note $x^{-1} \notin A \iff x \in A \setminus A^\star \iff x \in \mathfrak{m}$.
c) Suppose $0 \neq x \in K$ is integral over $A$. If $x \in A$ we are done. If $x^{-1} \in A$ then by (3.24.13) $x \in A[x^{-1}] \subseteq A$ as required.

■

**Lemma 3.25.3 (Restriction of a Valuation Ring to a Subfield)**
*Let $A$ be a valuation ring of a field $M$ and $L \subset M$ a subfield. Then $A \cap L$ is a valuation ring of $L$.*

*Proof.* For $0 \neq x \in L \subset M$, either $x \in A$ or $x^{-1} \in A$; as $x, x^{-1} \in L$ too, the same alternative holds for $A \cap L$. ■

**Proposition 3.25.4 (Local Homomorphism)**
*Let $\phi : A \to B$ be a homomorphism of local rings. Then the following are equivalent*

*a) $\phi(\mathfrak{m}_A) \subset \mathfrak{m}_B$*
*b) $\mathfrak{m}_A = \phi^{-1}(\mathfrak{m}_B)$*
*c) $\phi(\mathfrak{m}_A)B \subsetneq B$*

*We say that in this case $\phi$ is a **local homomorphism**. When $\phi$ is simply inclusion then we say $B$ **dominates** $A$.*

*Proof.* a) $\implies$ b) $\phi^{-1}(\mathfrak{m}_B)$ is a proper ideal containing $\mathfrak{m}_A$ and therefore equal by maximality

b) $\implies$ a) This is obvious

a) $\implies$ c) $\phi(\mathfrak{m}_A)B \subset \mathfrak{m}_B B = \mathfrak{m}_B \subsetneq B$

c) $\implies$ a) By (3.3.15) $\phi(\mathfrak{m}_A)B \subset \mathfrak{m}_B$ and trivially $\phi(\mathfrak{m}_A) \subset \phi(\mathfrak{m}_A)B$. ■

**Corollary 3.25.5**
*Let $(A, \mathfrak{m})$ be a local ring, $K$ a field and $\phi : A \to K$ a homomorphism. Then the following are equivalent*

*a) $\phi$ is a local ring homomorphism (i.e. $\mathfrak{m} \subset \ker(\phi)$)*
*b) $\ker(\phi) = \mathfrak{m}$*
*c) $\phi$ factors through $\kappa(\mathfrak{m})$.*

**Lemma 3.25.6 (Restriction to a Valuation Ring)**
*Let $A$ be a valuation ring of a field $F$, $L \supseteq F$ a field and $B \subseteq L$ a subring containing $A$, with a proper ideal $\mathfrak{n} \subsetneq B$ such that $\mathfrak{m}_A \subseteq \mathfrak{n}$ (equivalently $\mathfrak{m}_A B \neq B$). Then $B \cap F = A$ and $\mathfrak{n} \cap F = \mathfrak{m}_A$. In particular this applies to any local ring $B \subseteq L$ dominating $A$ (3.25.4), with $\mathfrak{n} = \mathfrak{m}_B$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Clearly $A \subseteq B \cap F$. Let $x \in F \setminus A$; then $x \neq 0$ and $(x^{-1})^{-1} = x \notin A$, so $x^{-1} \in \mathfrak{m}_A \subseteq \mathfrak{n}$ by (3.25.2). If $x \in B$ then $1 = xx^{-1} \in \mathfrak{n}$, which is absurd; so $x \notin B$ and $B \cap F = A$. Finally $\mathfrak{n} \cap F \subseteq B \cap F = A$ is a proper ideal of $A$, hence contained in $\mathfrak{m}_A$; the reverse inclusion is $\mathfrak{m}_A \subseteq \mathfrak{n}$. ■

**Lemma 3.25.7 (A Directed Union of Valuation Rings is a Valuation Ring)**
*Let $(K_i)_{i \in I}$ be a directed system of fields with union $K := \bigcup_i K_i$, and for each $i$ let $R_i \subset K_i$ be a valuation ring, such that $R_l$ dominates $R_i$ (3.25.4) whenever $K_i \subseteq K_l$. Then $R := \bigcup_i R_i$ is a valuation ring of $K$, with $\mathfrak{m}_R = \bigcup_i \mathfrak{m}_{R_i}$.*

© David Rufino CC BY-NC-ND 4.0

*Proof.* $R$ is a subring of $K$, being a directed union of subrings. For $0 \neq x \in K$, $x \in K_i$ for some $i$ (directed union of fields); as $R_i$ is a valuation ring of $K_i$, $x \in R_i$ or $x^{-1} \in R_i$, hence $x \in R$ or $x^{-1} \in R$: $R$ is a valuation ring of $K$.

$\mathfrak{m}_R \subseteq \bigcup_i \mathfrak{m}_{R_i}$: if $x \in \mathfrak{m}_R$, then $x \in R_i$ for some $i$ (as $R = \bigcup_i R_i$); as $x^{-1} \notin R \supseteq R_i$, in particular $x^{-1} \notin R_i$, so $x \in \mathfrak{m}_{R_i}$ (3.25.2) b).

$\mathfrak{m}_R \supseteq \bigcup_i \mathfrak{m}_{R_i}$: let $x \in \mathfrak{m}_{R_i}$, and suppose for contradiction $x^{-1} \in R_l$ for some $l$. By directedness choose $m$ with $i, l \leq m$, so $x \in R_i \subseteq R_m$ and $x^{-1} \in R_l \subseteq R_m$, that is $x \in R_m^\star$. But $R_m$ dominates $R_i$, so $\mathfrak{m}_{R_i} \subseteq \mathfrak{m}_{R_m}$, and $x \in \mathfrak{m}_{R_i}$ gives $x \in \mathfrak{m}_{R_m}$, contradicting $x \in R_m^\star$. So $x^{-1} \notin R_l$ for every $l$, that is $x^{-1} \notin R$, giving $x \in \mathfrak{m}_R$. ∎

**Definition 3.25.8 (Place (Zariski-Samuel 1960 / Lang 1972))**
*Let $K$ be a field. A **place** of $K$ consists of a valuation ring $(A, \mathfrak{m}_A)$ for $K$ and a local homomorphism to a field $F$*

$$\phi : A \to F$$

*(such that $\ker(\phi) = \mathfrak{m}_A$)*

*Furthermore if $x \in K \setminus A$ then we may write $\phi(x) = \infty$. Note that the second part of the previous Proposition then may be reinterpreted as saying*

$$\phi(x) = \infty \iff \phi(x^{-1}) = 0 \quad \forall x \in K$$

*which motivates the alternative definition below.*

*We say it is a **semi-place** of $K$ if $(A, \mathfrak{m}_A)$ is simply a local ring.*

**Lemma 3.25.9**
*Let $A \subset K$ be a subring of a field and $\mathfrak{a} \lhd A$ a proper ideal. Then at least one of $\mathfrak{a}A[x]$ or $\mathfrak{a}A[x^{-1}]$ is a proper ideal.*

*Proof.* Suppose neither are proper then we can write

$$1 = \sum_{j=0}^{n} a_j x^j$$

$$1 = \sum_{j=0}^{m} b_j x^{-j}$$

for $a_j, b_j \in \mathfrak{a}$. Choose $n, m$ to be minimal and assume wlog that $m \leq n$. Observe that $a_0 \neq 1 \implies m > 0$. Multiply the second equation by $x^n a_n$ to find

$$x^n a_n (1 - b_0) = a_n b_1 x^{n-1} + \ldots a_n b_m x^{n-m}$$

and multiply the first by $(1 - b_0)$ to find

$$(1 - b_0) = a_0(1 - b_0) + \ldots + a_n(1 - b_0)x^n$$

consequently cancelling the $x^n$ term and we obtain a relation of smaller degree a contradiction. ∎

We prove the first extension theorem

**Proposition 3.25.10 (Extension to localization)**
*Let $A$ be a ring and $\phi : A \to \Omega$ a homomorphism into a field. Let $\mathfrak{p} := \ker(\phi)$. Then*

- *$\mathfrak{p}$ is prime*
- *There is a unique extension $\tilde{\phi}$ making the diagram commute*

$$\begin{array}{ccc} A & \xrightarrow{\phi} & \Omega \\ \downarrow & \nearrow_{\tilde{\phi}} & \\ A_{\mathfrak{p}} & & \end{array}$$

*Furthermore $\ker(\tilde{\phi}) = \mathfrak{p}A_{\mathfrak{p}}$ i.e. $\tilde{\phi}$ is local.*

*Proof.* Clearly $\mathfrak{p}$ is prime because $\phi(A)$ is an integral domain. We may extend $\phi$ to the ring $A_{\mathfrak{p}}$ in the obvious way. The extension has kernel $\mathfrak{p}A_{\mathfrak{p}}$, the unique maximal ideal of $A_{\mathfrak{p}}$. ∎

© David Rufino CC BY-NC-ND 4.0

**Proposition 3.25.11 (Places as maximal extensions)**
*Let $A$ be a subring of a field $K$ and $\phi : A \to \Omega$ a homomorphism into an algebraically closed field. Then*

- *For all $x \in K$, $\phi$ may be extended to at least one of $A[x]$ and $A[x^{-1}]$.*
- *There exists a maximal extension $\tilde{\phi} : B \to \Omega$. For any such maximal extension $B$ is a valuation ring and $\tilde{\phi}$ is local (i.e. $\ker(\tilde{\phi}) = \mathfrak{m}_B$). Furthermore $\mathfrak{m}_B \cap A = \ker(\phi)$.*

*Proof.* We prove each in turn.

- By (3.25.10) we may assume without loss of generality that $A$ is a local ring with unique maximal ideal $\mathfrak{m} = \ker(\phi)$. By (3.25.9) we may also suppose that $\mathfrak{m}A[x]$ is proper. Then it's contained in a maximal ideal $\mathfrak{B} \lhd A[x]$. Furthermore $\mathfrak{m} = \mathfrak{B} \cap A$ by maximality of $\mathfrak{m}$. Let $k = A/\mathfrak{m}$ and $K = A[x]/\mathfrak{B}$, then there is a commutative diagram




  Then $K = k[\overline{x}]$ is a field and by (...) $\overline{x}$ is algebraic over $k$. Therefore $K/k$ is algebraic and, because $\Omega$ is algebraically closed, by (3.19.61) there is an extension to $K$, which gives the required extension to $A[x]$.
- It's easy to show that the poset of extensions to subrings of $K$ ordered by consistency is chain complete. Therefore by Zorn's Lemma there is a maximal extension $\tilde{\phi} : B \to \Omega$. By the previous part for any $x \in K$ any such maximal element $B$ must satisfy either $B[x] = B$ or $B[x^{-1}] = B$, i.e. $B$ is a valuation ring for $K$. Consider $\mathfrak{B} := \ker(\tilde{\phi})$ a prime ideal contained in $\mathfrak{m}_B$. Then by (3.25.10) may extend $\tilde{\phi}$ to $B_{\mathfrak{B}}$ and so by maximality $B = B_{\mathfrak{B}}$. Finally (3.20.6) shows that $B$ is a local ring with maximal ideal $\mathfrak{B} = \mathfrak{m}_B$. Clearly $\ker(\tilde{\phi}) \cap A = \ker(\phi)$, so the final statement follows easily.

■

**Corollary 3.25.12**
*Let $A \subset K$ be a subring of a field and $\mathfrak{a} \lhd A$ a proper ideal. Then there exists a valuation ring $(B, \mathfrak{m}_B)$ such that $A \subset B$ and $\mathfrak{a} \subset \mathfrak{m}_B \cap A$. In particular if $\mathfrak{a} = \mathfrak{m}_A$ is maximal then $\mathfrak{m}_A = \mathfrak{m}_B \cap A$.*

*Proof.* By (3.3.37) there is a maximal ideal $\mathfrak{m}_A \lhd A$ containing $\mathfrak{a}$. Let $k = A/\mathfrak{m}_A$ and $\Omega = \overline{k}$. Then the canonical homomorphism $\phi : A \to \Omega$ has kernel $\mathfrak{m}_A$. It has an extension to a valuation ring $(B, \mathfrak{m}_B)$ by (3.25.11), such that $\mathfrak{m}_B \cap A = \ker(\phi) = \mathfrak{m}_A$. ■

**Proposition 3.25.13 (Alternative Characterisation of Valuation Rings)**
*Let $K$ be a field and $A \subset K$ a subring. Then the following are equivalent*

*a) $A$ is a valuation ring for $K$*
*b) $A$ is a local ring maximal under the relation "$B$ dominates $A$"*
*c) $K = \mathrm{Frac}(A)$ and the principal ideals of $A$ are totally ordered*
*d) $K = \mathrm{Frac}(A)$ and the ideals of $A$ are totally ordered*

*Proof.* Let $(A, \mathfrak{m}_A) \preceq (B, \mathfrak{m}_B)$ denote the relation $B$ dominates $A$.

a) $\implies$ b) Suppose that $(A, \mathfrak{m}_A) \preceq (B, \mathfrak{m}_B)$ with $B$ local and $A$ a valuation ring. We claim that $A = B$; suppose for a contradiction that $x \in B \setminus A$. Then $x^{-1} \in A \implies x^{-1} \in B \implies x \in B^\star$. Therefore $A \subset B \setminus B^\star = \mathfrak{m}_B$. In particular $1 \in \mathfrak{m}_B$ which is a contradiction as $\mathfrak{m}_B$ is proper.

b) $\implies$ a) Let $(A, \mathfrak{m}_A)$ be a maximal local ring. By (3.25.12) there exists a valuation ring $(B, \mathfrak{m}_B)$ such that $(A, \mathfrak{m}_A) \preceq (B, \mathfrak{m}_B)$. By maximality $A = B$ and $A$ is a valuation ring.

a) $\implies$ c) Evidently $K = \mathrm{Frac}(A)$. Suppose $(a) \not\subset (b)$, then $a \notin (b) \implies b^{-1}a \notin A \implies a^{-1}b \in A \implies b \in (a) \implies (b) \subset (a)$.

c) $\implies$ d) Suppose $\mathfrak{a} \not\subset \mathfrak{b}$ then there exists $a \in \mathfrak{a} \setminus \mathfrak{b}$. In particular for all $b \in \mathfrak{b}$ we have $a \notin (b)$, whence by assumption $b \in (a)$. Therefore we conclude $\mathfrak{b} \subset (a) \subset \mathfrak{a}$.

d) $\implies$ b) As $A$ is a non-zero ring it has a maximal ideal $\mathfrak{m}_A$, which by assumption must be the unique maximal ideal. So $(A, \mathfrak{m}_A)$ is a local ring. Suppose that $(A, \mathfrak{m}_A) \preceq (B, \mathfrak{m}_B)$ and given $x \in B$ we have by assumption $x = a_1 a_2^{-1}$ for

© David Rufino CC BY-NC-ND 4.0

some $a_1, a_2 \in A$. If $a_1 \in (a_2)$ then $x \in A$. Otherwise $a_2 \in (a_1)$ whence $x^{-1} \in A \implies x^{-1} \in B^\star \overset{(3.25.2)}{\implies} x^{-1} \notin \mathfrak{m}_B \overset{(3.25.4)}{\implies} x^{-1} \notin \mathfrak{m}_A \implies x^{-1} \in A^\star \implies x \in A$. Consequently $A = B$ and b) holds. ∎

**Corollary 3.25.14**
*Let $A \subset K$ be a subring of a field then the integral closure of $A$ in $K$ (denoted $\overline{A}$) satisfies*

$$\overline{A} = \bigcap_{A \subset V} V$$

*where the intersection is taken over all valuation rings $V$ of $K$ containing $A$.*

*Alternatively the integral elements over $A$ are precisely the elements which are finite at all places of $K$, which are finite over $A$.*

*Proof.* First if $x \in \overline{A}$ then by (3.24.13) we have $x \in A[x^{-1}] \subseteq V[x^{-1}]$. If $x \notin V$ then by hypothesis $x^{-1} \in V$, whence $x \in V$ a contradiction. Therefore $x \in V$ as required.

Conversely suppose $x \notin \overline{A}$, then $x \notin A[x^{-1}]$. That is to say $(x^{-1})$ is a proper ideal in $A[x^{-1}]$. Therefore by (3.25.12) there is a valuation ring $(V, \mathfrak{m}_V)$ such that $x^{-1} \in \mathfrak{m}_V$ which implies $x \notin V$ by (3.25.2). ∎

**Definition 3.25.15 (Valuation)**
*Let $K$ be a field and $\Gamma$ an abelian ordered group. A function $v : K^\star \to \Gamma$ is said to be a **valuation** if it satisfies the following properties*

*a) $v(ab) = v(a) + v(b)$ for all $a, b \in K^\star$*
*b) $v(a+b) \geq \min(v(a), v(b))$ with the convention that $v(0) = \infty$*

*Note $v(1) = 0$. We say that two valuations $v, v'$ are equivalent if there is an order-preserving isomorphism $\phi : v(K^\star) \to v'(K^\star)$ such that $\phi \circ v = v'$. Clearly this is an equivalence relation.*

*If $k \subset K$ and $v(k) = 0$ then we say $v$ is a valuation for $K/k$.*

**Lemma 3.25.16 (Strict Ultrametric Inequality)**
*Let $v : K^\star \to \Gamma$ be a valuation (3.25.15), extended to $K$ by $v(0) := \infty$. For $x, y \in K$ with $v(x) \neq v(y)$,*

$$v(x+y) = \min(v(x), v(y))$$

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

If $x = 0$ or $y = 0$ this is immediate ($v(0) = \infty$ dominates), so assume $x, y \in K^\star$, and without loss of generality $v(x) < v(y)$. By property 2 of (3.25.15), $v(x+y) \geq \min(v(x), v(y)) = v(x)$. For the reverse inequality, apply property 2 again to $x = (x+y) + (-y)$:

$$v(x) \geq \min(v(x+y), v(-y))$$

As $v(-1) + v(-1) = v((-1)(-1)) = v(1) = 0$, and $\Gamma$ is totally ordered (so $v(-1) > 0$, resp. $v(-1) < 0$, would force $v(-1) + v(-1) > 0$, resp. $< 0$), we must have $v(-1) = 0$; so $v(-y) = v(-1) + v(y) = v(y)$. As $v(y) > v(x)$, the displayed inequality $v(x) \geq \min(v(x+y), v(-y))$ forces $v(x) \geq v(x+y)$ (else the minimum would be $v(-y) = v(y) > v(x)$, a contradiction). Combined with the first inequality, $v(x+y) = v(x) = \min(v(x), v(y))$. ∎

**Proposition 3.25.17 (Correspondence between Valuations Rings and Valuations)**
*Let $K$ be a field with $k \subset K$. Then there is a bijection*

$$\begin{aligned} \{k \subset A \subset K \mid A \text{ is a valuation ring}\} &\longleftrightarrow \{v : K^\star \to \Gamma \mid v \text{ is a valuation over } k\}/\sim \\ A &\to \quad v_A : K^\star \to K^\star/A^\star \\ v^{-1}(\Gamma_{\geq 0}) &\leftarrow \quad v : K^\star \to \Gamma \end{aligned}$$

*Under this map $v_A$ is just the quotient map and $\overline{x} \leq \overline{y} \iff yx^{-1} \in A^\star$. Furthermore*

*a) $v_A(\overline{x}) \geq 0 \iff x \in A$.*
*b) $v_A(\overline{x}) = 0 \iff x \in A^\star$*
*c) $v_A(\overline{x}) > 0 \iff x \in \mathfrak{m}_A$*

*Proof.* Given a valuation ring we may define the abelian group $\Gamma := K^\star/A^\star$ under multiplication with $\overline{x} \leq \overline{y} \iff yx^{-1} \in A$. This is easily verified to be a total ordering. Then $v_A : K^\star \to K^\star/A^\star$ is simply the quotient map and we clearly $v_A(\overline{x}) \geq 0 \iff x \in A^\star$. To demonstrate the additive condition we may consider the case $x, y \neq 0$ and suppose wlog that $xy^{-1} \in A$

© David Rufino CC BY-NC-ND 4.0

$$v(x+y) = v((xy^{-1}+1)y) = v(xy^{-1}+1) + v(y) \geq v(y) \geq \min(v(x), v(y))$$

The case that one of $x, y = 0$ is trivial. Observe $A = v_A^{-1}(\Gamma_{\geq 0}) \cup \{0\}$. We may therefore construct a one-sided inverse; given a valuation $v : K' \to \Gamma$ we may define $A_v := v^{-1}(\Gamma_{\geq 0}) \cup \{0\}$. It's evident from the two properties of $v$ that $A$ is an additive and multiplicative subgroup. Further by total ordering we see that $A$ is a valuation ring. We only need to demonstrate that the given constructions are mutually inverse, namely $v \sim v_{A_v}$. However this is immediate because the composite $v(K^\star) \to K^\star / A_v^\star \to v_{A_v}(K^\star)$ is an abelian group isomorphism, which preserves the positive segments, and therefore is an order isomorphism. ∎

**Corollary 3.25.18 (Valuation Dominance gives a Ring Multiple)**
*Let $v : K^\star \to \Gamma$ be a valuation with valuation ring $A$ (3.25.17), extended to $K$ by $v(0) := \infty$. If $x, y \in K$ with $y \neq 0$ and $v(x) \geq v(y)$, then $x = ay$ for some $a \in A$.*

*If further $v(x) = v(y)$ then $a \in A^\star$.*

*Proof.* If $x = 0$ take $a = 0$. Otherwise $x, y \in K^\star$ and $v(xy^{-1}) = v(x) - v(y) \geq 0$, so $a := xy^{-1} \in A$ by (3.25.17), and $x = ay$. If further $v(x) = v(y)$ then $v(a) = 0$, so $a \in A^\star$ by (3.25.17). ∎

**Proposition 3.25.19 (Bezout Domain)**
*Let $A$ be an integral domain. Then the following are equivalent*

*a) Every finitely generated ideal is principal*
*b) Every ideal generated by two elements is principal*

*In this case we say $A$ is a **Bezout Domain**. Further in this case every pair of elements has a greatest common divisor $d$ which satisfies $d = ax + by$ for some $x, y \in A$.*

*Proof.* By assumption $(a, b) = (d)$. Immediately we have $d \mid a$ and $d \mid b$. Similarly $d = \lambda a + \mu b$. Suppose $a = xe$ and $b = ye$ then $d = (\lambda x + \mu y)e$ whence $e \mid d$. ∎

**Lemma 3.25.20 (Valuation Ring = Local Bezout Domain)**
*Let $A$ be an integral domain. Then the following are equivalent*

*a) $A$ is a valuation ring*
*b) $A$ is a local Bezout domain.*

*In particular $A$ is a Noetherian valuation ring iff it is a local principal ideal domain.*

*Proof.* Suppose $A$ is a valuation ring and $\mathfrak{a} = (x, y)$. Without loss of generality $xy^{-1} \in A$ whence $x \in (y)$ so $\mathfrak{a} = (y)$ as required.

Conversely suppose $A$ is a local Bezout domain and let $x = \frac{y}{z} \in K$. We may divide by the greatest common divisor so that $\lambda y + \mu z = 1$. We claim one of $\lambda y$ and $\mu z$ is a unit, for suppose not then by (3.20.3) we have $\lambda y, \mu z \in \mathfrak{m} \implies 1 \in \mathfrak{m}$ a contradiction. Therefore one of $y, z$ is a unit which implies $x \in A$ or $x^{-1} \in A$, and $A$ is a valuation ring. ∎

© David Rufino CC BY-NC-ND 4.0

## 3.26 Derivations

**Definition 3.26.1 (Module of Derivations)**
*Let $B$ be a commutative ring and $M$ an $(B,C)$-bimodule, then we say a derivation is a map $D : B \to M$ satisfying the following properties*

- $D(x+y) = D(x) + D(y)$ ***linearity***
- $D(xy) = xD(y) + yD(x)$ ***product rule***

*We denote the family of such derivations by $\mathrm{Der}(B,M)$, and it is an $(B,C)$-bimodule. We may consider the case $M = C$ and $\phi : B \to B$ a ring homomorphism and in this case the product rule becomes*

$$D(xy) = \phi(x)D(y) + D(x)\phi(y)$$

*We observe that by induction we have $D(n \cdot 1_B) = 0$ for all $n \in \mathbb{Z}$.*

*Suppose that $B$ is an $A$-algebra then we claim the following are equivalent*

- $D(a) = 0 \quad \forall a \in A$
- $D(ab) = aD(b) \quad \forall a \in A$

*In this case we denote the set of $A$-linear derivations by*

$$\mathrm{Der}_A(B,M)$$

*Note every derivation is $\mathbb{Z}$-linear so we may work with $A = \mathbb{Z}$ in these cases. Finally we write*

$$\mathrm{Der}_A(B) \coloneqq \mathrm{Der}_A(B,B)$$

For our purposes the following will be the key example

**Example 3.26.2 (Evaluation at a zero)**
*Consider the case $B = A[X_1,...,X_n]/\mathfrak{a}$ a f.g. $A$-algebra, $L$ an $A$-algebra and $x \in L^n$ a zero of $\mathfrak{a}$. Then we may define the $B$-module structure on $L$ by*

$$f \cdot \lambda \coloneqq f(x)\lambda$$

*In this case we have the more explicit form*

$$\mathrm{Der}_A(B,L;x) = \{D \in \mathrm{Hom}_A(B,L) \mid D(fg) = f(x)D(g) + g(x)D(f)\}$$

*which is a $(B,L)$-bimodule.*

**Lemma 3.26.3**
*Suppose $D, D' \in \mathrm{Der}_A(B,M)$ agree on a set of generators for $B$, then they are identically equal. For example in the previous example when $D\left(\overline{X}_i\right) = D'\left(\overline{X}_i\right)$.*

The following is a useful technical tool to identify derivations as algebra homomorphisms

**Definition 3.26.4 (Dual Ring)**
*Let $M$ be a $B$-module. Define the* ***dual ring*** *as follows*

$$B[M] \coloneqq B \times M$$

$$(b,m) \times (b',m') = (bb', bm' + b'm)$$

*with addition defined in the obvious way and multiplicative unit is $(1_B, 0_M)$. Observe that it has an ideal $N \coloneqq 0 \times M$ such that $N^2 = 0$ and canonically $B[M]/N \cong B$.*

*When $B$ is an $A$-algebra, then there is a natural $A$-algebra structure on $B[M]$.*

**Proposition 3.26.5**
*Let $M$ be an $B$-module where $B$ is an $A$-algebra then there is a bijection*

$$\begin{aligned} \mathrm{Der}_A(B,M) &\longrightarrow \mathrm{AlgHom}_A(B, B[M]) \\ D &\to \ \phi_D(b) = (b, D(b)) \end{aligned}$$

**Definition 3.26.6 (Partial Derivative)**
*Suppose $F \in A[X_1,...,X_n]$ given by*

© David Rufino CC BY-NC-ND 4.0

$$F(X_1,...,X_n) = \sum_{i\in\mathbb{N}^n} a_i X_1^{i_1}...X_n^{i_n} \quad a_i \in A$$

*Define the partial derivative as follows*

$$\frac{\partial F}{\partial X_k} := \sum_{\substack{i\in\mathbb{N}^n \\ i_k\neq 0}} a_i i_k X_1^{i_1}...X_k^{i_k-1}...X_n^{i_n}$$

*We observe that*

$$\frac{\partial X_l}{\partial X_k} := \delta_{lk}$$

*Note this condition uniquely determines the equation above, see (3.26.8).*

We show that these form a basis for $\mathrm{Der}_k(A)$.

**Lemma 3.26.7 (Product Rule)**
*For $F,G \in A[X_1,...,X_n]$ we have the* ***product rule***

$$\frac{\partial FG}{\partial X_k} = F\frac{\partial G}{\partial X_k} + G\frac{\partial F}{\partial X_k}$$

*Proof.* First we demonstrate the result in the univariate case $n = 1$. For monomials this is straightforward:

$$\frac{\partial X^r X^s}{\partial X} = (r+s)X^{r+s-1} = X^s\frac{\partial X^r}{\partial X} + X^r\frac{\partial X^s}{\partial X}$$

For general univariate polynomials the product rule follows from the linearity of $\partial/\partial X$. The multivariate case then follows from considering the isomorphism $A[X_1,...,X_n] \cong A\left[X_1,...,\widehat{X_k},...,X_n\right][X_k]$ under which $\partial/\partial X_k$ corresponds to $\partial/\partial X$. ∎

**Lemma 3.26.8 (Multivariate Chain Rule)**
*Let $D \in \mathrm{Der}_A(B,M)$ be a derivation, $x_1,...,x_n \in A$ and $F \in B[X_1,...,X_n]$. Then*

$$D(F(x_1,...,x_n)) = \sum_{k=1}^{n} \frac{\partial F}{\partial X_k}(x_1,...,x_n)D(x_k)$$

*As a special case we find*

$$D(F(x)) = F'(x)D(x)$$
$$D(x^n) = nx^{n-1}D(x)$$

*Proof.* First we observe that by induction on $n$

$$D(x_1...x_n) = \sum_{k=1}^{n} x_1...\widehat{x_k}...x_n D(x_k)$$

and in particular

$$D(x^n) = \begin{cases} nx^{n-1}D(x) & n > 0 \\ 0 & n = 0 \end{cases}$$

Then for $F \in k[X_1,...,X_n]$ we have

$$\begin{aligned} D(F(x_1,...,x_n)) &= \sum_{i\in\mathbb{N}^n} a_i D\left(x_1^{i_1}...x_n^{i_n}\right) \\ &= \sum_{i\in\mathbb{N}^n} a_i \sum_{k=1}^{n} x_1^{i_1}...\widehat{x_k^{i_k}}...x_n^{i_n} D\left(x_k^{i_k}\right) \\ &= \sum_{k=1}^{n}\sum_{i\in\mathbb{N}^n} a_i x_1^{i_1}...\widehat{x_k^{i_k}}...x_n^{i_n} D\left(x_k^{i_k}\right) \\ &= \sum_{k=1}^{n}\sum_{\substack{i\in\mathbb{N}^n \\ i_k\neq 0}} i_k a_i x_1^{i_1}...x_k^{i_k-1}...x_n^{i_n} D(x_k) \end{aligned}$$

© David Rufino CC BY-NC-ND 4.0

which gives the required result. ∎

**Proposition 3.26.9**
*Let $B \coloneqq A[X_1, ..., X_n]$ be a polynomial ring and $M$ a $(B, C)$-bimodule. Then $\mathrm{Der}_A(B, M)$ is a finite free $(B, C)$-bimodule with basis $\{\partial/\partial X_1, ..., \partial/\partial X_n\}$ (3.26.6).*

**Proposition 3.26.10 (Extensions to Quotient Field)**
*Let $B$ be an $A$-algebra, $S$ a multiplicative set of $B$ and $M$ a $B$-module. Then we have the following isomorphisms*

$$\mathrm{Der}_A(B, M) \xrightarrow{\sim} \mathrm{Der}(S^{-1}B, S^{-1}M)$$
$$D \to \frac{b}{s} \to \frac{bD(s) - sD(b)}{s^2}$$

*In particular if $M = \Omega$ is a field containing $B$ (which is therefore an integral domain) then we have an isomorphism*

$$\mathrm{Der}_A(B, \Omega) \xrightarrow{\sim} \mathrm{Der}_A(K, \Omega)$$

*where $K \coloneqq \mathrm{Frac}(B)$.*

In light of (3.26.5) the following definition is clearly related to the existence of derivations.

**Definition 3.26.11 (Formally Smooth)**
*Let $B$ be an $A$-algebra. We say it is **formally smooth** if for any $A$-algebra $C$ with $N \triangleleft C$ such that $N^2 = 0$ and any homomorphism $v : B \to C/N$ there exists a morphism $\overline{v} : A \to C$ making the following diagram commute*

$$\begin{array}{ccc} A & \longrightarrow & C \\ \downarrow & \nearrow & \downarrow \pi \\ B & \xrightarrow{v} & C/N \end{array}$$

*We say that $B$ is **formally unramified** if in addition the lifting is always unique.*

**Lemma 3.26.12**
*Let $C$ be a ring and $N \subseteq \sqrt{(0)}$ an ideal. Then $x \in C$ is invertible if and only if $\overline{x} \in C/N$ is invertible.*

*Proof.* One direction is obvious. Conversely if $\overline{x}$ is invertible then $x \in C^\star + N \subseteq C^\star + \sqrt{(0)} \subseteq C^\star$ by (3.3.68) as required. ∎

**Lemma 3.26.13**
*If a ring $B$ is formally smooth (resp. unramified) then so is $S^{-1}B$.*

*Proof.* It's enough to show that for $s \in S$ we have $\overline{v}(s)$ is invertible, which follows from (3.26.12). ∎

**Lemma 3.26.14**
*The polynomial ring $B \coloneqq A[X_1, ..., X_n]$ and function field $A(X_1, ..., X_n)$ is formally smooth.*

*Proof.* The first follows by the fact $A[X_1, ..., X_n]$ is a free $A$-algebra and the second by (3.26.13). ∎

**Lemma 3.26.15**
*Let $B$ be a formally smooth (resp. formally unramified) $A$-algebra and $C$ is a formally smooth (resp. formally unramified) $B$-algebra. Then $C$ is a formally smooth (resp. formally unramified) $A$-algebra.*

**Lemma 3.26.16 (Separable algebraic extensions are "formally unramified")**
*A separable algebraic extension $K/k$ is formally unramified.*

*Proof.* Consider the following commutative diagram

$$\begin{array}{ccc} k & \longrightarrow & C \\ \downarrow & \nearrow & \downarrow \pi \\ K & \xrightarrow{v} & C/N \end{array}$$

We suppose first that $K$ is a finite extension. By (3.19.105) it is simple, namely $K = k(\alpha) = k[\alpha]$. Let $m(X)$ be the minimal polynomial over $k$ and choose $x \in C$ such that $\pi(x) = v(\alpha)$. Note that $\overline{v}(\alpha) = x$ need not be well-defined because in general $m(x) \neq 0$. We show however there is an $n \in N$ such that $m(x + n) = 0$. For consider any $n \in N$ and $f(X) \in k[X]$.

© David Rufino CC BY-NC-ND 4.0

$$\begin{aligned} f(x+n) &= f(x) + \sum_{i=0}^{N} \lambda_i [(x+n)^i - x^i] \\ &= f(x) + f'(x)n \end{aligned}$$

as $n^2 = 0$. Therefore we propose $n = -m(x)/m'(x)$; to show this is well-defined, consider firstly $\pi(m(x)) = m(v(\alpha)) = v(m(\alpha)) = 0$ whence $m(x) \in N$. To show that $m'(x)$ is a unit we argue as follows. As $\alpha$ is separable, $m'(\alpha) \neq 0$, hence is a unit in $K$. As $v$ is injective $v(m'(\alpha)) = m'(v(\alpha)) = m'(\pi(x)) = \pi(m'(x))$ is a unit in $C/N$. By (3.26.12) $m'(x) \in C^\star$. Therefore $n \in N$ is well-defined and $m(a+n) = 0$ by construction. Therefore the map $\overline{v} : k[\alpha] \to A$ such that $\overline{v}(\alpha) = a + n$ is well-defined, and the diagram commutes because $\pi(a+n) = \pi(a) + \pi(n) = v(\alpha)$.

Suppose we had another $\overline{v}'$, then $a' \coloneqq \overline{v}'(\alpha)$ again satisfies $m(a') = 0$ and $\pi(a') = v(\alpha)$ whence $a' - a \in N$. By the same argument as before

$$m(a') = m(a + (a' - a)) = m(a) + m'(a)(a' - a)$$

As $m'(a) \neq 0$ and $m(a') = m(a) = 0$ we see $a' = a$ which demonstrates uniqueness.

Suppose that $K/k$ is algebraic than for every $\alpha \in K$ we have liftings $v_\alpha : k(\alpha) \to k$. We claim that $v_\alpha|_{k(\alpha)\cap k(\beta)} = v_\beta|_{k(\alpha)\cap k(\beta)}$. However $k(\alpha) \cap k(\beta)$ is also finite and so we are done by uniqueness. ∎

**Proposition 3.26.17**
*A separably generated extension $K/k$ is formally smooth.*

*Proof.* This follows from (3.26.14), (3.26.16) and (3.26.15). ∎

© David Rufino CC BY-NC-ND 4.0

## 3.27 Hauptidealsatz

The main result of this section is the following result due to Krull

**Proposition 3.27.1 (Generalized Hauptidealsatz for Noetherian Rings)**
*Suppose $A$ is a Noetherian ring then the following properties hold*

*a) Every prime ideal of height $n$ is minimal over some ideal $\mathfrak{a} := (x_1, ..., x_n)$. Furthermore $\mathfrak{a}$ may be chosen such that $\operatorname{ht}(\mathfrak{a}) = n$ and every minimal prime of $\mathfrak{a}$ is of height $n$*
*b) We have the following characterization of height*

$$\operatorname{ht}(\mathfrak{p}) = \min\{n \mid \mathfrak{p} \text{ minimal over } (x_1, ..., x_n)\}$$

*In particular if $\mathfrak{p}$ is minimal over $(x_1, ..., x_n)$ then $\operatorname{ht}(\mathfrak{p}) \leq n$.*

In full generality the proof is quite subtle and in most cases of interest a simpler proof is possible. Therefore we introduce the following notions.

**Definition 3.27.2 (Hauptidealsatz Ring)**
*We say a ring $A$ is a **generalized hauptidealsatz** ring if for all $x_1, ..., x_n \in A$ and $\mathfrak{p}$ prime ideals minimal over $(x_1, ..., x_n)$ we have $\operatorname{ht}(\mathfrak{p}) \leq n$*

*We say a ring $A$ is simply a **hauptidealsatz** ring if this holds for $n = 1$.*

**Lemma 3.27.3**
*Let $A$ be a hauptidealsatz ring and $\mathfrak{p}$ a prime ideal. Then $A_{\mathfrak{p}}$ is hauptidealsatz.*

*Proof.* Any prime ideal of $A_{\mathfrak{p}}$ has the form $\mathfrak{q}A_{\mathfrak{p}}$ by the correspondence of ideals under localization (3.6.19). If $\mathfrak{q}A_{\mathfrak{p}}$ is minimal over $(f/s) = (f/1)$, then clearly $(f) \subseteq \mathfrak{q}$. Furthermore $(f) \subseteq \mathfrak{q}' \implies (f/1) \subseteq \mathfrak{q}'A_{\mathfrak{p}}$. So $\mathfrak{q}$ is also minimal over $(f)$ and therefore has height at most 1 by assumption. By the same result $\mathfrak{q}A_{\mathfrak{p}}$ has height at most 1. ∎

**Proposition 3.27.4 (Hauptidealsatz ⇒ Generalized Hauptidealsatz (Geometric))**
*Suppose that $A$ is catenary, and hauptidealsatz for every quotient by a prime ideal. Then $A$ is generalized hauptidealsatz, and so is every localization at a prime ideal.*

*Proof.* We consider the case first that $A$ is integral (which means it is hauptidealsatz by assumption because $(0)$ is prime) and assume wlog that $n > 1$. Let $\mathfrak{p}$ be a minimal prime of $(x_1, ..., x_n)$ and $\mathfrak{q}$ a minimal prime of $(x_1, ..., x_{n-1})$. By considering the localization $A_{\mathfrak{p}}$ we may choose $\mathfrak{q} \subseteq \mathfrak{p}$. By induction $\operatorname{ht}(\mathfrak{q}) \leq n - 1$. Then $\mathfrak{p}/\mathfrak{q}$ is a minimal prime over $(x_n + \mathfrak{q})$ and therefore by assumption has height at most 1. By the codimension formula (3.22.9) we see $\operatorname{ht}(\mathfrak{p}) \leq n$ as required.

For the general case, suppose $\mathfrak{p}$ is minimal over $(x_1, ..., x_n)$ and let $\mathfrak{r} \subseteq \mathfrak{p}$ be a minimal prime. Then the ring $A/\mathfrak{r}$ is integral and we see that $\operatorname{ht}(\mathfrak{p}/\mathfrak{r}) \leq n$ by the first part. Taking the supremum over all minimal primes $\mathfrak{r}$ shows that $\operatorname{ht}(\mathfrak{p}) \leq n$.

For the last statement we need only show that every localization at a prime ideal satisfies the hypotheses of the theorem. By correspondence of ideals under localisation (3.6.19) we see that every such ring is catenary. By (3.6.25) $A_{\mathfrak{p}}/\mathfrak{q}A_{\mathfrak{p}} \cong (A/\mathfrak{q})_{\mathfrak{p}/\mathfrak{q}}$ for any prime ideal $\mathfrak{q} \subseteq \mathfrak{p}$ so every quotient by a prime ideal is hauptidealsatz by the previous Lemma. ∎

The catenary assumption may be relaxed by making a more complicated argument.

**Proposition 3.27.5 (Hauptidealsatz ⇒ Generalized Hauptidealsatz (Algebraic))**
*Suppose $A$ is a ring such that every quotient by a prime ideal satisfies the hauptidealsatz. Then $A$ is generalized hauptidealsatz and so is every localization at a prime ideal $\mathfrak{p}$.*

*Proof.* We prove this by induction on $n$ for a fixed ring $A$. Let $\mathfrak{p}$ be a minimal prime over $(x_1, ..., x_n)$. Suppose $\operatorname{ht}(\mathfrak{p}) > n$ then we may choose a saturated chain $\mathfrak{q} \subsetneq \mathfrak{p}$ such that $\operatorname{ht}(\mathfrak{q}) \geq n$. By minimality of $\mathfrak{p}$ we have $\mathfrak{q} \cap \{x_1, ..., x_n\} = \{x_1, ..., x_r\}$ with $r < n$ after a suitable reordering (possibly with $r = 0$). Furthermore we may choose $\mathfrak{q}$ such that $r$ is maximal. Then by the induction hypothesis $\mathfrak{q}$ is not minimal over $(x_1, ..., x_r)$ and there is a chain

$$(x_1, ..., x_r) \subseteq \mathfrak{r} \subsetneq \mathfrak{q} \subsetneq \mathfrak{p}$$

We claim that $\mathfrak{p}$ is minimal over $(\mathfrak{r}, x_{r+1})$ for suppose

$$(\mathfrak{r}, x_{r+1}) \subseteq \mathfrak{p}' \subseteq \mathfrak{p}$$

© David Rufino CC BY-NC-ND 4.0

then by construction of $\mathfrak{q}$ ($r$ was maximal) we see that $\mathfrak{p}' = \mathfrak{p}$. Taking quotients by $\mathfrak{r}$ the hauptidealsatz property shows that $\mathrm{ht}(\mathfrak{p}/\mathfrak{r}) \leq 1$. On the other hand we have a chain

$$(0) \subsetneq \mathfrak{q}/\mathfrak{r} \subsetneq \mathfrak{p}/\mathfrak{r}$$

which is a contradiction.

The final statement follows a similar argument as before. ■

**Remark 3.27.6**
*This argument is from [Kap74] (Sec. 3.2 Ex. 6).*

Before proceeding to the main result we need a some technical results related to height of ideals.

**Lemma 3.27.7**
*Let $\mathfrak{q} \subseteq \mathfrak{p}$ be prime ideals then*

$$\mathrm{ht}(\mathfrak{q}) \leq \mathrm{ht}(\mathfrak{p})$$

*with equality iff $\mathfrak{p} = \mathfrak{q}$.*

*Proof.* The inequality is clear since any maximal chain below $\mathfrak{q}$ may be extended to a maximal chain below $\mathfrak{p}$. Similarly $\mathfrak{q} \subsetneq \mathfrak{p}$ implies that the inequality is strict. ■

**Lemma 3.27.8**
*Suppose $\mathfrak{a} \subseteq \mathfrak{b}$. Then*

$$\mathrm{ht}(\mathfrak{a}) \leq \mathrm{ht}(\mathfrak{b})$$

*If these are equal and $\mathfrak{b}$ is prime, then it must be a minimal prime of $\mathfrak{a}$.*

*Proof.* We consider first the case $\mathfrak{b} = \mathfrak{p}$ is prime. By (3.3.44) there is a minimal prime $\mathfrak{p}'$ such that $\mathfrak{a} \subseteq \mathfrak{p}' \subseteq \mathfrak{p}$. By definition $\mathrm{ht}(\mathfrak{a}) \leq \mathrm{ht}(\mathfrak{p}')$. Furthermore by the previous Lemma $\mathrm{ht}(\mathfrak{p}') \leq \mathrm{ht}(\mathfrak{p})$, which yields the required result.

Suppose $\mathrm{ht}(\mathfrak{p}) = \mathrm{ht}(\mathfrak{a})$ then by definition $\mathrm{ht}(\mathfrak{a}) \leq \mathrm{ht}(\mathfrak{p}') \leq \mathrm{ht}(\mathfrak{p})$ and therefore we conclude $\mathrm{ht}(\mathfrak{p}) = \mathrm{ht}(\mathfrak{p}')$. By the previous lemma we conclude that $\mathfrak{p} = \mathfrak{p}'$ is a minimal prime.

For the general case every minimal prime of $\mathfrak{b}$ also contains $\mathfrak{a}$ so the inequality follows by taking the minimum over all minimal primes. ■

**Lemma 3.27.9**
*Let $\mathfrak{a} \subseteq \mathfrak{b}$ be ideals such that there exists $x \in \mathfrak{b}$ not contained in any minimal prime of $\mathfrak{a}$. Then*

$$\mathrm{ht}(\mathfrak{a}) < \mathrm{ht}(\mathfrak{b})$$

*Proof.* By the previous Lemma we have $\mathrm{ht}(\mathfrak{a}) \leq \mathrm{ht}(\mathfrak{b})$. Suppose they are equal then there is a minimal prime $\mathfrak{p}$ of $\mathfrak{b}$ of the same height, which by (3.27.8) is a minimal prime of $\mathfrak{a}$. This must contain $x$ which then contradicts the assumption. ■

**Proposition 3.27.10 (Prime Avoidance Theorem)**
*Let $\mathfrak{p}_1, ..., \mathfrak{p}_n$ be ideals and $\mathfrak{a}$ an ideal such that $\mathfrak{a} \nsubseteq \mathfrak{p}_i$ for all $i = 1...n$. Then there exists $x \in \mathfrak{a} \setminus (\mathfrak{p}_1 \cup ... \cup \mathfrak{p}_n)$.*

*Proof.* We proceed by induction on the number of prime ideals $n$, the case $n = 1$ being trivial. Consider then the case of $n+1$ prime ideals and suppose on the contrary that $\mathfrak{a} \subseteq \mathfrak{p}_1 \cup ... \cup \mathfrak{p}_{n+1}$. By the induction hypothesis there exists $x_i \in \mathfrak{a} \setminus \bigcup_{j \neq i} \mathfrak{p}_j$, and therefore by our assumption $x_i \in (\mathfrak{a} \cap \mathfrak{p}_i) \setminus \bigcup_{j \neq i} \mathfrak{p}_j$. Define

$$y := x_1 ... x_n$$

Then $y \notin \mathfrak{p}_{n+1}$ by primality and clearly $y \in \mathfrak{a} \cap \mathfrak{p}_1 \cap ... \cap \mathfrak{p}_n$. Define $z := y + x_{n+1}$ then we see

a) $z \in \mathfrak{a}$
b) $z \notin \mathfrak{p}_{n+1}$
c) $z \notin \mathfrak{p}_i$ for $i = 1...n$

which contradicts our original assumption. ■

**Lemma 3.27.11**
*Let $A$ be a **generalized hauptidealsatz** ring and $\mathfrak{a} = (x_1, ..., x_n)$. Then the following are equivalent*

*a)* $\mathrm{ht}(\mathfrak{a}) = n$
*b) every minimal prime of $\mathfrak{a}$ has height $n$*

© David Rufino CC BY-NC-ND 4.0

*Proof.* a) $\implies$ b) By definition this means every minimal prime has height at least $n$, and the reverse inequality follows from generalized hauptidealsatz assumption.

b) $\implies$ a) Recall from (3.22.1) we may take the infimum over only minimal prime ideals. ∎

**Proposition 3.27.12 (Alternative characterization of height)**
*Suppose $A$ is a Noetherian ring satisfying **generalized hauptidealsatz**, then the following properties hold*

*a) Every prime ideal of height $n$ is minimal over some ideal $\mathfrak{a} := (x_1, ..., x_n)$. Furthermore $\mathfrak{a}$ may be chosen such that $\operatorname{ht}(\mathfrak{a}) = n$ and every minimal prime is of height $n$*
*b) We have the following characterization of height*

$$\operatorname{ht}(\mathfrak{p}) = \min\{n \mid \mathfrak{p} \text{ minimal over } (x_1, ..., x_n)\}$$

*In particular every prime ideal has finite height.*

*Proof.* We first demonstrate a). Suppose $\operatorname{ht}(\mathfrak{p}) = n$ then there is a saturated chain of prime ideals

$$\mathfrak{p}_0 \subsetneq ... \subsetneq \mathfrak{p}_n = \mathfrak{p}$$

It's clear from the above chain that $\operatorname{ht}(\mathfrak{p}_i) \geq i$. Furthermore by (3.27.7) $\operatorname{ht}(\mathfrak{p}_i) < \operatorname{ht}(\mathfrak{p}_{i+1})$ whence we see $\operatorname{ht}(\mathfrak{p}_i) = i$.

We show by induction that there are elements $x_1, ..., x_n \in A$ for which $\mathfrak{p}_i$ is a minimal prime over $\mathfrak{a}_i := (x_1, ..., x_i)$ for all $i = 0...n$ and $\operatorname{ht}(\mathfrak{a}_i) = i$.

The case $i = 0$ is clear, so suppose $i > 0$. In general by (3.22.6) there are finitely many minimal primes over $\mathfrak{a}_{i-1}$ say $\mathfrak{q}_1, ..., \mathfrak{q}_r$ which all have height $i - 1$ by (3.27.11). In particular $\mathfrak{p}_i \not\subseteq \mathfrak{q}_k$.

By prime avoidance (3.27.10) we may choose $x_i \in \mathfrak{p}_i \setminus (\mathfrak{q}_1 \cup ... \cup \mathfrak{q}_r)$. Clearly

$$\mathfrak{a}_{i-1} \subseteq \mathfrak{a}_i \subseteq \mathfrak{p}_i$$

Then by (3.27.8) and (3.27.9) we have $i - 1 = \operatorname{ht}(\mathfrak{a}_{i-1}) < \operatorname{ht}(\mathfrak{a}_i) \leq i$ whence $\operatorname{ht}(\mathfrak{a}_i) = i$. By (3.27.8) again we see $\mathfrak{p}_i$ is a minimal prime of $\mathfrak{a}_i$. This completes the inductive step.

To show b) let $m$ denote the right hand side. Then by the generalized hauptidealsatz hypothesis $\operatorname{ht}(\mathfrak{p}) \leq m$. On the other hand by part a) we have $m \leq \operatorname{ht}(\mathfrak{p})$ whence they are equal.

Finally by the Noetherian hypothesis $\mathfrak{p}$ is finitely generated and clearly minimal over itself. ∎

© David Rufino CC BY-NC-ND 4.0

## 3.28 Regular Local Rings

The results of the previous section allow a more direct characterization of the dimension of a Noetherian local ring, which naturally leads to the notion of regular local ring.

**Lemma 3.28.1 ($\mathfrak{m}$-primary)**
*Let $(A, \mathfrak{m})$ be a local ring and $\mathfrak{a}$ a proper ideal. Then the following are equivalent*

*a)* $\sqrt{\mathfrak{a}} = \mathfrak{m}$
*b)* $\mathfrak{m}$ *is a minimal prime of* $\mathfrak{a}$

*Furthermore $\mathfrak{a}$ is primary. In this case we say $\mathfrak{a}$ is $\mathfrak{m}$-primary. A sufficient condition is that for some $n$*

$$\mathfrak{m}^n \subset \mathfrak{a} \subset \mathfrak{m}$$

*and this is necessary when $\mathfrak{m}$ is finitely generated.*

*Proof.* a) $\implies$ b) Suppose $\mathfrak{a} \subseteq \mathfrak{p}$ then $\mathfrak{m} = \sqrt{\mathfrak{a}} \subseteq \mathfrak{p}$ by (3.3.46). In particular $\mathfrak{m}$ is a minimal prime.

b) $\implies$ a) Let $\mathfrak{p}$ be a prime ideal containing $\mathfrak{a}$. By (3.3.37) $\mathfrak{p} \subseteq \mathfrak{m}$ and by assumption $\mathfrak{p} = \mathfrak{m}$. The result then follows from (3.3.46).

Suppose $xy \in \mathfrak{a}$ and $y \notin \sqrt{\mathfrak{a}}$ then by (...) $y \in A^\star$ whence $x \in \mathfrak{a}$. This shows that $\mathfrak{a}$ is primary.

The condition $\mathfrak{m}^n \subset \mathfrak{a}$ is clearly sufficient, and necessary in the finitely generated case by (3.3.48). ∎

An ideal satisfying one of the equivalent conditions above is called **$\mathfrak{m}$-primary**, though some authors require the stronger condition.

**Proposition 3.28.2 (Criteria for Dimension of Local Ring)**
*Let $(A, \mathfrak{m})$ be a Noetherian local ring satisfying the generalized hauptidealsatz. Then we have the following criteria for dimension*

$$\dim A = \operatorname{ht}(\mathfrak{m}) = \min\left\{ n \;\middle|\; \sqrt{(x_1, \ldots, x_n)} = \mathfrak{m} \right\} \leq \mu(\mathfrak{m}) = \dim_{k(\mathfrak{m})} \mathfrak{m}/\mathfrak{m}^2$$

*where $\mu(\mathfrak{m})$ is the least number of generators of $\mathfrak{m}$ and $k(\mathfrak{m}) = A/\mathfrak{m}$.*

*Proof.* The first equality follows because every maximal chain must terminate at a maximal ideal by (3.3.37). The second from the characterization of height for hauptidealsatz rings (3.27.12) and the equivalent definitions of $\mathfrak{m}$-primary (3.28.1). The inequality follows because $\mathfrak{m}$ is itself $\mathfrak{m}$-primary. The final equality follows because a minimal generating set lifts to a basis of $\mathfrak{m}/\mathfrak{m}^2$ (3.21.7). ∎

**Definition 3.28.3**
*Let $(A, \mathfrak{m})$ be a Noetherian local ring. We say $A$ is **regular** if*

$$\dim A = \mu(\mathfrak{m}) = \dim_{k(\mathfrak{m})} \mathfrak{m}/\mathfrak{m}^2$$

© David Rufino CC BY-NC-ND 4.0

## 3.29 Local Rings of Dimension 0

**Lemma 3.29.1 (Dimension 0 local ring)**
*Let $(A, \mathfrak{m})$ be a Noetherian local ring. Then the following are equivalent*

*a)* $\dim A = 0$ *(i.e. every prime ideal is maximal)*
*b)* $\mathfrak{m}^n = 0$ *for some* $n$
*c)* $\mathfrak{m}^n = \mathfrak{m}^{n+1}$ *for some* $n$

*If $A$ is an integral domain then this is equivalent to $A$ being a field.*

*Proof.* a) $\implies$ b) By (3.22.6) $\mathfrak{r} := \sqrt{(0)} = \mathfrak{p}_1 \cap ... \cap \mathfrak{p}_n$ is a decomposition into minimal prime ideals. By assumption these are also maximal, and therefore by uniqueness $\sqrt{(0)} = \mathfrak{m}$ and the result follows from (3.3.48) as $\mathfrak{m}$ is finitely generated.

b) $\implies$ a) Let $\mathfrak{p}$ be a prime ideal then trivially $\mathfrak{m}^n \subset \mathfrak{p}$ so by (3.28.1) it is $\mathfrak{m}$-primary i.e. $\mathfrak{m}$ is a minimal prime of $\mathfrak{p}$ whence $\mathfrak{m} = \mathfrak{p}$.

b) $\iff$ c) One direction is obvious, and the converse follows from Nakayama's Lemma (3.21.5) because $\mathfrak{m}^n$ is finitely generated. ∎

© David Rufino CC BY-NC-ND 4.0

# 3.30 Fractional Ideals

**Definition 3.30.1**
*Let $A$ be an integral domain contained in its quotient field $K$. An $A$-submodule $I \subset K$ is called a* ***fractional ideal*** *if there exists $a \in A$ such that $aI \subset A$.*

*We say a fractional ideal is* ***principal*** *if it is generated by some $x \in K$ as an $A$-module.*

**Proposition 3.30.2 (Operations on Fractional ideals)**
*Let $A$ be an integral domain and $I, J \subset K$ fractional ideals. Then the following constructions are also fractional ideals*

*a)* $I + J = \{i + j \mid i \in I j \in J\}$
*b)* $IJ = \left\{\sum_{k=1}^{n} i_k j_k \;\middle|\; i_k \in I j_k \in J\right\}$
*c)* $(I : J) = \{x \in K \mid xJ \subseteq I\}$

*Proof.* It is straightforward to prove that these constructions are again $A$-submodules of $K$.

Suppose that $aI \subseteq A$ and $bJ \subseteq A$, then $(ab)(I + J) \subseteq A$ and $(ab)IJ \subseteq A$. Similarly

$$(I : J)J \subseteq I \implies j(I : J) \subseteq I \implies abj(I : J) \subseteq A$$

for some $0 \neq j \in J$ ■

**Proposition 3.30.3**
*Let $A$ be an integral domain. The collection of fractional ideals $\mathcal{I}_A$ forms a monoid under multiplication of ideals, with identity $A$.*

**Definition 3.30.4**
*Let $A$ be an integral domain and $I$ a fractional ideal. Then we say $I$ is* ***invertible*** *if there exists a fractional ideal $J$ such that $IJ = A$.*

**Proposition 3.30.5**
*Let $A$ be an integral domain and $I$ a fractional ideal. Then $I$ is invertible if and only if $I(A : I) = A$, in which case $(A : I)$ is the unique inverse.*

*Proof.* Uniqueness follows from general properties of monoids (...). If $I(A : I) = A$ then by definition $I$ is invertible. Conversely suppose $I$ is invertible, that is $IJ = A$. Then evidentally $J \subseteq (A : I)$ so $A = IJ \subseteq I(A : I) \subseteq A$. Therefore $(A : I)$ is an inverse for $I$ as required. ■

**Corollary 3.30.6**
*Let $A$ be an integral domain. Then the invertible fractional ideals form an abelian group under multiplication.*

**Definition 3.30.7 (Localisation of Fractional Ideals)**
*Let $A$ be an integral domain contained in its quotient field $K$. Let $I$ be a fractional ideal and $S$ a multiplicatively closed subset of $A$. Then define*

$$S^{-1}I := \left\{\frac{i}{s} \;\middle|\; i \in Is \in S\right\}$$

*If $\mathfrak{a}$ is an ideal then $S^{-1}\mathfrak{a}$ corresponds to the usual notion (3.6.17) and is equal to the extension of $\mathfrak{a}$ under the inclusion homomorphism $A \to S^{-1}A$.*

*If $\mathfrak{p}$ is a prime ideal of $A$ then as usual define*

$$I_{\mathfrak{p}} := (A \setminus \mathfrak{p})^{-1} I$$

*and we may write*

$$\mathfrak{a}_{\mathfrak{p}} = \mathfrak{a}A_{\mathfrak{p}}$$

**Proposition 3.30.8 (Localisation of Fractional Ideals)**
*Let $A$ be an integral domain contained in its quotient field $K$. Let $S$ be a multiplicatively closed subset of $A$ and $I$ a fractional ideal. Then regarding as subrings of $K$ then $S^{-1}I$ is a fractional ideal of $S^{-1}A$. Furthermore*

*a)* $S^{-1}(I + J) = S^{-1}I + S^{-1}J$
*b)* $S^{-1}(IJ) = (S^{-1}I)(S^{-1}J)$
*c)* $S^{-1}(I : J) = (S^{-1}I : S^{-1}J)$

© David Rufino CC BY-NC-ND 4.0

*Proof.* Evidently $S^{-1}I$ is again a fractional ideal of $S^{-1}A$.

a) Clear because $\frac{i+j}{s} = \frac{i}{s} + \frac{j}{s}$.
b) We have $\frac{ij}{s} = \left(\frac{i}{s}\right)\left(\frac{j}{1}\right)$ whence $S^{-1}(IJ) \subseteq (S^{-1}I)(S^{-1}J)$. Conversely we have $\left(\frac{i}{s}\right)\left(\frac{j}{t}\right) = \frac{ij}{st}$ so we ahve the reverse inclusion.
c) Suppose that $\frac{x}{s} \in S^{-1}(I : J)$ and $\frac{j}{t} \in S^{-1}J$. Then $\left(\frac{x}{s}\right)\left(\frac{j}{t}\right) = \frac{xj}{st} \in S^{-1}I$ because $x \in (I : J)$. Therefore $S^{-1}(I : J) \subseteq \left(S^{-1}I : S^{-1}J\right)$. Conversely given $x \in \left(S^{-1}I : S^{-1}J\right)$ and $j \in J$. Then by definition $x\left(\frac{j}{1}\right) = \frac{i}{s}$ whence $sx \in (I : J)$. Therefore $x \in S^{-1}(I : J)$ as required.

■

**Lemma 3.30.9**
*Let $A$ be an integral domain contained in its quotient field $K$. Suppose $I$ is a fractional ideal then*

$$I = \bigcap_{\mathfrak{m}} I_{\mathfrak{m}}$$

*In particular we have*

$$A = \bigcap_{\mathfrak{m}} A_{\mathfrak{m}}$$

*Proof.* Evidently $I \subseteq \bigcap_{\mathfrak{m}} I_{\mathfrak{m}}$. Suppose that $x \in \bigcap_{\mathfrak{m}} I_{\mathfrak{m}}$. Consider the ideal $\mathfrak{a} := \{a \in A \mid ax \in I\}$. Suppose $\mathfrak{a}$ is proper then by (3.3.15) it is contained in a maximal ideal $\mathfrak{m}$. As $x \in I_{\mathfrak{m}}$ we have $(A \setminus \mathfrak{m}) \cap \mathfrak{a} \neq \varnothing$ and $\mathfrak{a} \subseteq \mathfrak{m}$ a contradiction. Therefore $\mathfrak{a} = A$ and in particular $x = 1.x \in I$ as required. ■

**Proposition 3.30.10 (Invertibility is a local property)**
*Let $A$ be an integral domain with quotient field $K$, and $I$ a fractional ideal of $A$. Then the following are equivalent*

*a) $I$ is invertible*
*b) $I_{\mathfrak{p}}$ is invertible for all prime ideals $\mathfrak{p}$*
*c) $I_{\mathfrak{m}}$ is invertible for all prime ideals $\mathfrak{m}$*

*Proof.* a) $\implies$ b) By assumption and (3.30.5) $I(A : I) = A$ so by (3.30.8) $I_{\mathfrak{p}}\left(A_{\mathfrak{p}} : I_{\mathfrak{p}}\right) = A_{\mathfrak{p}}$, as required.

b) $\implies$ c) Every maximal ideal is prime (3.3.60) so we are done.

c) $\implies$ a) Let $I$ be a fractional ideal then

$$\begin{aligned}
I(A : I) &= \bigcap_{\mathfrak{m}} (I(A : I))_{\mathfrak{m}} && (3.30.9)\\
&= \bigcap_{\mathfrak{m}} I_{\mathfrak{m}}(A : I)_{\mathfrak{m}} && (3.30.8)\\
&= \bigcap_{\mathfrak{m}} I_{\mathfrak{m}}(A_{\mathfrak{m}} : I_{\mathfrak{m}}) && (3.30.8)\\
&= \bigcap_{\mathfrak{m}} A_{\mathfrak{m}} && (3.30.5)\\
&= A && (3.30.9)
\end{aligned}$$

■

© David Rufino CC BY-NC-ND 4.0

# 3.31 Discrete Valuation Rings

A Discrete Valuation Ring (DVR) corresponds to a regular local ring of dimension 1 and therefore regular (or in the case of perfect base field, non-singular) points on a curve. There are a number of equivalent conditions which are summarised here.

**Lemma 3.31.1**
*Let $(A, \mathfrak{m})$ be a local domain such that $\mathfrak{m}$ is finitely generated and $\dim A = 1$. Then for every non-zero ideal $\mathfrak{a}$ there exists some $n$ such that $\mathfrak{m}^n \subseteq \mathfrak{a}$.*

*Proof.* As $\dim A = 1$ every minimal prime over $\mathfrak{a}$ is equal to $\mathfrak{m}$, whence $\sqrt{\mathfrak{a}} = \mathfrak{m}$ (i.e. $\mathfrak{a}$ is $\mathfrak{m}$-primary). As $\mathfrak{m}$ is finitely generated there is some $n$ such that $\mathfrak{m}^n \subseteq \mathfrak{a}$ (3.3.48). ∎

**Proposition 3.31.2 (DVR Criteria)**
*Let $A$ be an integral domain which is not a field. Then the following are equivalent.*

*a) $A$ is a valuation ring whose valuation group is order-isomorphic to $\mathbb{Z}$*
*b) $A$ is a local principal ideal domain*
*c) $A$ is a Noetherian local ring with maximal ideal $\mathfrak{m} = (\pi)$ and $\dim A = 1$*
*d) $A$ is a local ring such that every ideal is of the form $(\pi^k)$ for some $k \geq 0$ (and these are distinct)*
*e) There exists a non-unit $\pi \in A$ such that every element $a \in A$ may be represented as $u\pi^k$ for some $u \in A^\star$ and non-negative integer $k$.*
*f) $A$ is a Noetherian valuation ring*
*g) $A$ is a Noetherian integrally closed local domain with $\dim A = 1$*
*h) $A$ is a Noetherian regular local ring with $\dim A = 1$*

*In this case we say $A$ is a **discrete valuation ring** (or **DVR**) and let $K = \mathrm{Frac}(A)$. For every generator $\pi$ of $\mathfrak{m}$, the ideals $(\pi^n)$ are distinct and*

- *for every $x \in K^\star$ there is a unique integer $k$ and unit $u$ such that $x = u\pi^k$, and*
- *there is a well-defined valuation $v : K^\star \to \mathbb{Z}$ determined uniquely by the relation $x = u\pi^{v(x)}$*

*Proof.* a) $\implies$ b) Consider a valuation $v : K \to \mathbb{Z}$ with $v^{-1}(\mathbb{Z}_{\geq 0}) = A$. For any ideal $\mathfrak{a}$ choose $f \in \mathfrak{a}$ such that $v(f)$ is minimal. For $g \in \mathfrak{a}$ we have $v(g) \geq v(f) \implies v(gf^{-1}) \geq 0 \implies gf^{-1} \in A \implies g \in (f)$. Therefore $\mathfrak{a} = (f)$ is principal. Further by (3.25.2) $A$ is a local ring.

b) $\implies$ c). By (3.22.4) $\dim A = 1$ and by assumption $\mathfrak{m} = (\pi)$. Evidently $A$ is Noetherian.

c) $\implies$ d). By (3.29.1) the sequence $(\pi^k)$ is strictly decreasing. Assume wlog that $\mathfrak{a}$ is proper and non-zero, then as $\dim A = 1$ every prime ideal is maximal and we find $\sqrt{\mathfrak{a}} = (\pi)$ by (3.3.46).

Suppose that $\mathfrak{a} \subseteq (\pi^k)$ for all $k$. Then $\pi \in \sqrt{\mathfrak{a}} \implies (\pi^k) = \mathfrak{a}$ for all sufficiently large $k$ which contradicts the fact the sequence is strictly decreasing. Therefore we may assume $\mathfrak{a} \subseteq (\pi^k)$ for some $k$ and there exists $a \in \mathfrak{a} \setminus (\pi^{k+1})$. Therefore $a = u\pi^k$ and by construction $u \notin (\pi) \implies u \in A^\star \implies \pi^k \in \mathfrak{a}$. Therefore $\mathfrak{a} = (\pi^k)$ as required.

d) $\implies$ e). By assumption $(a) = (\pi^k)$ for some integer $k$, whence by (3.17.3) $a = u\pi^k$ for some $u \in A^\star$. This representation is unique because the ideals $(\pi^k)$ are all distinct and $A$ is cancellable.

e) $\implies$ a). We may show easily that the integer $k$ is uniquely defined, denote this by $v(a)$ where $a = u\pi^{v(a)}$ for $u \in A^\star$. We need to show that $v$ is a valuation. By uniqueness we deduce $v(ab) = v(a) + v(b)$. Suppose that $a, b, a+b \in A \setminus \{0\}$. Then $(\pi^{v(a+b)}) = (a+b) \subseteq (\pi^{\min(v(a),v(b))})$. As the sequence $(\pi^k)$ is strictly decreasing we find $v(a+b) \geq \min(v(a), v(b))$ as required. We may extend to $K^\star$ by $v(ab^{-1}) := v(a) - v(b)$. Then $v(ab^{-1}) \geq 0 \iff v(a) \geq v(b) \iff (a) \subseteq (b) \iff b \mid a \iff ab^{-1} \in A$. Further $v(a) = 0 \iff a \notin (\pi) \iff a \in A^\star$. Therefore $A$ is the valuation ring associated to $v$.

d) $\implies$ f) $\implies$ b). This follows from (3.25.20).

f) $\implies$ g). A valuation ring is integrally closed (3.25.2) and we've already shown $A$ is local with dimension 1.

g) $\implies$ c). Let $0 \neq a \in \mathfrak{m}$. As $\dim A = 1$ from (3.31.1) we know $\mathfrak{m}^n \subseteq (a)$ for some $n \geq 1$. Choose $n$ minimal and $b \in \mathfrak{m}^{n-1} \setminus (a)$.

Let $x = a^{-1}b \in K = \mathrm{Frac}(A)$. Evidently $x \notin A$, for otherwise $b \in (a)$; further $x\mathfrak{m} = a^{-1}b\mathfrak{m} \subseteq a^{-1}\mathfrak{m}^n \subseteq a^{-1}(a) = A$ is an ideal of $A$.

© David Rufino CC BY-NC-ND 4.0

We claim that $x\mathfrak{m} = A$. Suppose not then $x\mathfrak{m} \subseteq \mathfrak{m}$ by (3.3.37). Then $\mathfrak{m}$ is a faithful finite $A[x]$-module and $x$ is integral over $A$ (3.24.3). By assumption $A$ is integrally closed so $x \in A$ which is a contradiction. Therefore $x\mathfrak{m} = A \implies \mathfrak{m} = x^{-1}A$. This shows that $x^{-1} = x^{-1} \cdot 1 \in A$ and that $\mathfrak{m}$ is a principal ideal.

c) $\implies$ h). By assumption $A$ is a Noetherian local ring with $\dim A = 1$. So it remains to show that $\dim \mathfrak{m}/\mathfrak{m}^2 = 1$. As $\mathfrak{m}$ is principal then $\dim \mathfrak{m}/\mathfrak{m}^2 \leq 1$. If it is zero then $\mathfrak{m} = \mathfrak{m}^2$ and $\dim A = 0$ by (3.29.1), a contradiction.

h) $\implies$ c). By assumption $\dim \mathfrak{m}/\mathfrak{m}^2 = 1$ and so by (3.21.7) $\mathfrak{m} = (\pi)$ for some $\pi \in \mathfrak{m} \setminus \mathfrak{m}^2$.

The final statements have already been proven. ∎

**Lemma 3.31.3 (Valuation and Powers of the Maximal Ideal)**
*Let $(A, \mathfrak{m})$ be a DVR with valuation $v$, and $k \geq 0$ an integer. For $0 \neq a \in A$:*

*a) $v(a) \geq k \iff a \in \mathfrak{m}^k$;*
*b) $v(a) = k \iff a \in \mathfrak{m}^k \setminus \mathfrak{m}^{k+1}$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Fix a uniformiser $\pi$ (so $\mathfrak{m} = (\pi)$, hence $\mathfrak{m}^k = (\pi^k)$ (3.31.2)), and write $a = u\pi^{v(a)}$ for a unit $u \in A^\star$ (3.31.2).

a). $a \in \mathfrak{m}^k = (\pi^k) \iff \pi^k \mid a = u\pi^{v(a)} \iff \pi^k \mid \pi^{v(a)}$ (as $u$ is a unit) $\iff v(a) \geq k$ (as the ideals $(\pi^n)$ are strictly decreasing (3.31.2)).

b). By a), $a \in \mathfrak{m}^k \setminus \mathfrak{m}^{k+1} \iff v(a) \geq k$ and not $(v(a) \geq k+1) \iff v(a) = k$. ∎

**Corollary 3.31.4**
*Let $A$ be a principal ideal domain which is not a field. Then every localisation of $A$ at a non-zero prime ideal is a DVR.*

*In particular $\mathbb{Z}_{(p)}$ and $k[X]_{(f)}$ are DVRs where $p$ is prime and $f(X)$ is an irreducible polynomial.*

**Lemma 3.31.5 (A DVR is Maximal Among Valuation Subrings of its Fraction Field)**
*Let $A$ be a DVR with fraction field $K$ and $B$ a valuation ring of $K$ with $A \subseteq B$. Then $A = B$ or $B = K$.*

*Proof.* Let $\pi$ be a uniformiser of $A$, so every $a \in A$ is $u\pi^k$ for a unit $u \in A^\star$ and integer $k \geq 0$ (3.31.2). Suppose $B \neq K$. As $B$ is a valuation ring of $K$, either $\pi \in B$ (already true, as $\pi \in A \subseteq B$) or $\pi^{-1} \in B$; we rule out the latter. If $\pi^{-1} \in B$, then for any $x \in K^\star$, writing $x = u\pi^n$ ($u \in A^\star \subset B$, $n \in \mathbb{Z}$, using $A$ a DVR), $x = u\pi^n \in B$ whether $n \geq 0$ (as $\pi^n \in A \subset B$) or $n < 0$ (as $\pi^n = (\pi^{-1})^{-n} \in B$); so $B = K$, contradicting assumption. Hence $\pi^{-1} \notin B$, that is $\pi \in \mathfrak{m}_B$. Then $\mathfrak{m}_A = \pi A \subset \pi B \subset \mathfrak{m}_B$ (as $\pi \in \mathfrak{m}_B$ and $B$ absorbs), so $B$ dominates $A$; as $A$ is a valuation ring, it is maximal under domination (3.25.13), giving $A = B$. ∎

**Lemma 3.31.6 (Directed Union of Valuation Rings with a Common Uniformiser)**
*Let $(K_i)_{i \in I}$ be a directed system of fields with union $K := \bigcup_i K_i$, and for each $i$ let $R_i \subset K_i$ be a valuation ring with $R_i \subseteq R_l$ whenever $K_i \subseteq K_l$. Suppose in addition every $\mathfrak{m}_{R_i}$ is principal, generated by a single common element $0 \neq \pi \in \bigcap_i R_i$. Then $R := \bigcup_i R_i$ is a valuation ring of $K$, with $\mathfrak{m}_R = \pi R \neq 0$; in particular $R \subsetneq K$.*

*Proof.* For $K_i \subseteq K_l$ (so $R_i \subseteq R_l$), $\mathfrak{m}_{R_i} = \pi R_i \subseteq \pi R_l = \mathfrak{m}_{R_l}$, giving $R_l$ dominates $R_i$ (3.25.4) so (3.25.7) applies, and $\mathfrak{m}_R = \bigcup_i \mathfrak{m}_{R_i} = \bigcup_i \pi R_i = \pi R$.

As $\pi \in \mathfrak{m}_{R_i}$ is a non-unit of every $R_i$, $\pi^{-1} \notin R_i$ for every $i$ (well-defined, as $0 \neq \pi \in K_i \subset K$), hence $\pi^{-1} \notin R = \bigcup_i R_i$; as $\pi^{-1} \in K$, $R \subsetneq K$. ∎

**Proposition 3.31.7**
*Let $A$ be a discrete valuation ring with unique maximal ideal $\mathfrak{m}$. Then there exists a canonical group isomorphism*

$$(\mathfrak{m})^- : \mathbb{Z} \xrightarrow{\sim} \mathcal{I}_A$$

*which is the unique group homomorphism mapping $1$ to $\mathfrak{m}$ (and $0$ to $A$. In particular every fractional ideal is invertible.*

*Write the inverse group isomorphism by $v_\mathfrak{m}$, which satisfies*

*a) $\mathfrak{m}^{v_\mathfrak{m}(\mathfrak{a})} = \mathfrak{a}$*
*b) $v_\mathfrak{m}(\mathfrak{m}^k) = k$*

*A fractional ideal $I$ is contained in $A$ if and only if $v_\mathfrak{m}(I) \geq 0$.*

*Proof.* By (3.31.2) we may suppose $\mathfrak{m} = (\pi)$. Evidently $(\pi^k)$ is a fractional ideal for every integer $k$ and $(\pi^k)(\pi^l) = (\pi^{k+l})$, which gives the required homomorphism $k \to (\pi^k)$.

© David Rufino CC BY-NC-ND 4.0

Suppose the kernel is non-trivial then there exists $k$ which maps to $A$, and we may assume that $k$ is positive (by considering the inverse). This means precisely that $\mathfrak{m}^k = A$, which may only happen when $k = 0$ because $\mathfrak{m}$ is proper.

Let $I$ be an arbitrary fractional ideal then $aI \subseteq A$ for some $a \in A$. By (3.31.2) $a = u\pi^l$ and $aI = (\pi^k)$ for some positive integers $l$ and $k$. This means $(\pi^l)I = (\pi^k)$ and therefore $I = (\pi^k)(\pi^{-l}) = (\pi^{k-l})$. This shows the homomorphism is surjective and therefore an isomorphism.

Evidently $(\pi^k) \subset A \iff k \geq 0$ since $\pi \in A$ and $\pi^{-1} \notin A \implies \pi^{-k} \notin A$. ∎

**Proposition 3.31.8**
*Let $A$ be a discrete valuation ring and $I, J$ fractional ideals. Then the following are equivalent*

*a) $I \subset J$*
*b) $J$ divides $I$ (that is there exists an ideal $\mathfrak{a}$ such that $\mathfrak{a}J = I$)*
*c) $v_{\mathfrak{m}}(I) \geq v_{\mathfrak{m}}(J)$*

*Proof.* Using the results of (3.31.7) we may show (eliding the subscript $\mathfrak{m}$)

a) $\iff$ c) $I \subset J \iff J^{-1}I \subset A \iff v(J^{-1}I) \geq 0 \iff v(I) \geq v_{\mathfrak{m}}(J)$

b) $\implies$ c) $v(I) = v(\mathfrak{a}) + v(J) \geq v(J)$

c) $\implies$ b) Define $\mathfrak{a} = \mathfrak{m}^{v(I)-v(J)}$ then evidentally $v(\mathfrak{a}J) = v(I)$ whence $I = J$. ∎

**Lemma 3.31.9 (Intersection of the Powers of the Maximal Ideal)**
*Let $A$ be a DVR with maximal ideal $\mathfrak{m}$. Then $\bigcap_{n\geq 0} \mathfrak{m}^n = 0$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Let $0 \neq a \in A$, and write $a = u\pi^{v(a)}$ for a unit $u$ and uniformiser $\pi$ (3.31.2). If $a \in \mathfrak{m}^{v(a)+1} = (\pi^{v(a)+1})$ then $a = u'\pi^{v(a)+1}$ for some $u' \in A$, so $u\pi^{v(a)} = u'\pi^{v(a)+1}$, giving $u = u'\pi$ by cancellation – contradicting $u$ a unit. So $a \notin \mathfrak{m}^{v(a)+1}$, in particular $a \notin \bigcap_n \mathfrak{m}^n$. ∎

**Lemma 3.31.10 (Ramification and Injectivity/Surjectivity of Local Homomorphisms)**
*Let $(A, \mathfrak{m})$, $(B, \mathfrak{n})$ be DVRs with valuations $v_A, v_B$ (3.31.2), and $\phi : A \to B$ a local ring homomorphism (that is $\phi(\mathfrak{m}) \subseteq \mathfrak{n}$). Fix a uniformiser $\pi$ of $A$ and set $e := v_B(\phi(\pi)) \in \mathbb{N} \cup \{\infty\}$ (the **ramification index** of $\phi$; independent of the choice of $\pi$, with $e = \infty$ iff $\phi(\pi) = 0$). Then:*

*a) $\phi$ is injective if and only if $e < \infty$;*
*b) if $\phi$ is surjective, then $e = 1$; in particular, by a), $\phi$ is then also injective – so $\phi$ is surjective if and only if it is bijective.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

**Independence of $\pi$.** If $\pi' = u\pi$ for a unit $u \in A^\star$, then $\phi(\pi') = \phi(u)\phi(\pi)$ with $\phi(u) \in B^\star$ (units map to units under any ring homomorphism, and $\phi(u) \notin \mathfrak{n}$, as otherwise $1 = \phi(u^{-1})\phi(u) \in \mathfrak{n}$, absurd), so $v_B(\phi(\pi')) = v_B(\phi(\pi))$.

**a).** $\ker\phi$ is a prime ideal of $A$ (the kernel of a ring homomorphism into the domain $B$), so $\ker\phi \in \{0, \mathfrak{m}\}$ (3.31.2). If $e = \infty$: $\phi(\pi) = 0$, so $\pi \in \ker\phi$, and, as $\mathfrak{m} = (\pi)$, $\mathfrak{m} \subseteq \ker\phi$; as $\ker\phi \neq A$ ($\phi(1) = 1 \neq 0$), $\ker\phi = \mathfrak{m} \neq 0$: $\phi$ is not injective. If $e < \infty$: $\phi(\pi) \neq 0$, so $\pi \notin \ker\phi$, giving $\ker\phi \neq \mathfrak{m}$, so $\ker\phi = 0$: $\phi$ is injective.

**b).** Suppose $\phi$ is surjective. Let $\varpi$ be a uniformiser of $B$; by surjectivity, $\varpi = \phi(a)$ for some $a \in A$, necessarily $a \neq 0$ (as $\varpi \neq 0$). Write $a = u\pi^k$ for a unit $u \in A^\star$ and $k := v_A(a) \geq 0$ (3.31.2). If $e = \infty$: $\phi(a) = \phi(u)\phi(\pi)^k$ is $\phi(u)$ (a unit, as units map to units) if $k = 0$, or 0 if $k \geq 1$; neither equals $\varpi$ ($v_B(\varpi) = 1 \neq 0$, $\varpi \neq 0$), a contradiction. So $e < \infty$, and

$$1 = v_B(\varpi) = v_B(\phi(u)) + k \cdot v_B(\phi(\pi)) = 0 + ke = ke,$$

forcing $e = 1$. ∎

© David Rufino CC BY-NC-ND 4.0

## 3.32 Dedekind Domains

We develop the theory of Dedekind Domains. These occur both as number fields and regular affine curves (over any field). The important results are the finiteness of integral closure in a finite extension, and the ramification formula of a prime ideal. In the case of regular curves over a finite field further results are needed, which we prove later in the context of affine algebras.

**Proposition 3.32.1**
*Let $A$ be a Noetherian integral domain of dimension 1. Then the following are equivalent*

*a) $A$ is integrally closed*
*b) $A_\mathfrak{p}$ is integrally closed for every prime ideal $\mathfrak{p}$*
*c) $A_\mathfrak{p}$ is a discrete valuation ring for every non-zero prime ideal $\mathfrak{p}$*

*In this case we say that $A$ is a **Dedekind Domain**.*

*Proof.* Let $\mathfrak{p}$ be non-zero prime ideal of $A$. Then firstly $A_\mathfrak{p}$ is Noetherian by (...) and an integral domain by (...). Then $\dim A_\mathfrak{p} = \mathrm{ht}(\mathfrak{p}) \leq \dim A = 1$. However $(0) \subsetneq \mathfrak{p}$ so $\dim A_\mathfrak{p} = \mathrm{ht}(\mathfrak{p}) = 1$.

a) $\iff$ b) This is (3.24.10).

b) $\implies$ c) We have shown that $A_\mathfrak{p}$ is a Noetherian integral domain of dimension 1. Then it is a discrete valuation ring by (3.31.2).

c) $\implies$ b) Also follows from (3.31.2), since $A_{(0)}$ is a field and therefore automatically integrally closed. ∎

**Corollary 3.32.2**
*Every principal ideal domain which is not a field is a Dedekind Domain. For example $k[X]$ and $\mathbb{Z}$.*

*Proof.* This follows from (3.17.23) (3.24.12) (3.16.2). ∎

The following is useful in the theory of algebraic curves and function fields.

**Proposition 3.32.3 (Correspondence between Prime Ideals and Valuation Rings)**
*Let $A$ be an integral domain with Krull dimension 1 and quotient field $K$. Then there is a bijection*

$$\{\mathfrak{p} \lhd A \mid \mathfrak{p} \text{ non-zero prime}\} \longleftrightarrow \{A \subsetneq R \subsetneq K \text{ local rings}\}$$
$$\mathfrak{p} \longrightarrow A_\mathfrak{p}$$
$$\mathfrak{m}_R \cap A \longleftarrow R$$

*When $A$ is a Dedekind domain then all such local rings (and valuation rings) are discrete valuation rings.*

*Proof.* Suppose that $x \in \mathfrak{p}_1 \setminus \mathfrak{p}_2$. Then $x^{-1} \in A_{\mathfrak{p}_2}$ but we claim $x^{-1} \notin A_{\mathfrak{p}_1}$. For suppose $x^{-1} \in A_{\mathfrak{p}_1}$ then $\frac{ax}{b} = 1$ for $a \in A$ and $b \notin \mathfrak{p}_1$ whence $ax = b$ a contradiction. Therefore the given map is injective.

Let $R$ be a local ring with maximal ideal $\mathfrak{m}_R$. Then $\mathfrak{m}_R \cap A$ is a prime ideal, and so either $(0)$ or a non-zero prime $\mathfrak{p}$. We have $0 \neq \frac{x}{y} \in \mathfrak{m}_R$ for some $x, y \in A$ whence $0 \neq x \in \mathfrak{m}_R$. Therefore $\mathfrak{m}_R \cap A = \mathfrak{p}$. Suppose $x \in A \setminus \mathfrak{p}$ then $x \in R \setminus \mathfrak{m}_R \implies x \in R^\star$. This shows that $A_\mathfrak{p} \subset R$ and $\mathfrak{p}A_\mathfrak{p} \subset \mathfrak{m}_R$, and we are done by (3.25.13). ∎

**Proposition 3.32.4**
*Let $A$ be a Dedekind domain. Then every fractional ideal is invertible.*

*Proof.* Let $I$ be an invertible ideal. By (3.30.10) it is sufficient to show that $I_\mathfrak{p}$ is an invertible fractional ideal of $A_\mathfrak{p}$ for all prime ideals $\mathfrak{p}$. By (3.32.1) $A_\mathfrak{p}$ is a discrete valuation ring and the result follows from (3.31.7). ∎

**Lemma 3.32.5**
*Let $A$ be a Dedekind domain and $\mathfrak{a}$ a proper ideal. Then there is a well-defined group homomorphism*

$$(\mathfrak{a})^- : \mathbb{Z} \longrightarrow \mathcal{I}_A$$

*mapping 1 to $\mathfrak{a}$.*

*Proof.* By (3.32.4) the monoid $\mathcal{I}_A$ is a group, and therefore the first homomorphism exists by (3.2.6). ∎

**Lemma 3.32.6**
*Let $A$ be a Dedekind domain and $\mathfrak{p}$, $\mathfrak{q}$ non-zero prime ideals. Then*

© David Rufino CC BY-NC-ND 4.0

$$\left(\mathfrak{p}^k\right)_\mathfrak{q} = \begin{cases} A_\mathfrak{q} & \mathfrak{p} \neq \mathfrak{q} \\ \left(\mathfrak{q}A_\mathfrak{q}\right)^k & \mathfrak{p} = \mathfrak{q} \end{cases}$$

*Proof.* When $k = 0$ then by definition $\mathfrak{p}^k = A$ and $\left(\mathfrak{q}A_\mathfrak{q}\right)^k = A_\mathfrak{q}$ and so we are done.

We suppose that $k = 1$. If $\mathfrak{p} \neq \mathfrak{q}$ then as $\dim A = 1$ we have $\mathfrak{p} \not\subseteq \mathfrak{q}$. Therefore by definition $\mathfrak{p}_\mathfrak{q}$ contains 1 and equals $A_\mathfrak{q}$ as required. The case $\mathfrak{p} = \mathfrak{q}$ follows from (3.30.7).

By (3.30.8) and (3.32.5) both left and right hand sides define group homomorphisms $\mathbb{Z} \to \mathcal{I}_{A_\mathfrak{q}}$ which agree at 1 and are therefore equal. ∎

**Proposition 3.32.7 (Unique Factorisation)**
*Let $A$ be a Dedekind domain. There is a well-defined abelian group isomorphism*

$$\bigoplus_{\mathfrak{p}\in\mathcal{P}} \mathbb{Z} \xrightarrow{\sim} \mathcal{I}_A$$
$$(e_\mathfrak{p}) \longrightarrow \prod_{\mathfrak{p}:e_\mathfrak{p}\neq 0} \mathfrak{p}^{e_\mathfrak{p}}$$

*where $\mathcal{P}$ is the set of prime ideals of $A$. Write $\left(v_\mathfrak{p}\right)$ for the inverse, then it is given explicitly by*

$$v_\mathfrak{p}(I) = v_\mathfrak{p}\left(I_\mathfrak{p}\right)$$

*Further a fractional ideal $I$ is contained in $A$ if and only if $v_\mathfrak{p}(I) \geq 0$ for all prime ideals $\mathfrak{p}$.*

*Proof.* The homomorphisms exists by (3.32.6) and the universal property of the sum of abelian groups (...).

Let $I$ be a fractional ideal and define $v_\mathfrak{p}(I) \coloneqq v_\mathfrak{p}\left(I_\mathfrak{p}\right)$ using the notation of (3.31.7). This determines a group homomorphism $v_\mathfrak{p} : \mathcal{I}_A \to \mathbb{Z}$ by (3.30.8) b). We claim for every $I$ that $v_\mathfrak{p}(I)$ is zero for all but finitely many prime ideals $\mathfrak{p}$. As $I = (a)^{-1}\mathfrak{a}$ for some ideal $J$ of $A$ it is sufficient by multiplicativity to demonstrate this for ideals $\mathfrak{a}$. By (...) if $\mathfrak{a} \not\subset \mathfrak{p}$ then $\mathfrak{a}_\mathfrak{p} = A_\mathfrak{p}$ which has valuation 0. Therefore we may assume that $\mathfrak{a} \subset \mathfrak{p}$. However by (3.22.6) there are only finitely many such primes, since every non-zero prime containing $\mathfrak{a}$ must be minimal over (0), and a-fortiori over $\mathfrak{a}$.

Observe by (3.32.6) that for distinct prime ideals $\mathfrak{p}_1, ..., \mathfrak{p}_n$ we have the identity

$$\left(\mathfrak{p}_1^{r_1}...\mathfrak{p}_n^{r_n}\right)_\mathfrak{q} = \begin{cases} \left(\mathfrak{p}_k A_\mathfrak{q}\right)^{r_k} & \mathfrak{q} = \mathfrak{p}_k \\ A_\mathfrak{q} & \mathfrak{q} \neq \mathfrak{p}_k \text{ for all } k = 1...n \end{cases}$$

By the first claim there is a well-defined homomorphism $\mathcal{I}_A \to \bigoplus_{\mathfrak{p}\in\mathcal{P}} \mathbb{Z}$. We claim that this is a left inverse for the given map, that is to say

$$\prod_{\mathfrak{p}:v_\mathfrak{p}(I)\neq 0} \mathfrak{p}^{v_\mathfrak{p}(I)} = I$$

Denote the left hand side by $I'$. When $v_\mathfrak{p}(I) \neq 0$ then by the identity above we have, using (3.31.7) for the final equality,

$$I'_\mathfrak{p} = \left(\mathfrak{p}A_\mathfrak{p}\right)^{v_\mathfrak{p}(I)} = \left(\mathfrak{p}A_\mathfrak{p}\right)^{v_\mathfrak{p}\left(I_\mathfrak{p}\right)} = I_\mathfrak{p}$$

Similarly if $v_\mathfrak{p}(I) = 0$ by (3.31.7)

$$I'_\mathfrak{p} = A_\mathfrak{p} = I_\mathfrak{p}$$

We conclude by (3.30.9) that $I = I'$ as required. Finally the relation above also shows that this is a right inverse.

For the final statement by (3.30.9) we have $I \subset A \iff I_\mathfrak{p} \subset A_\mathfrak{p}$ for all prime ideals $\mathfrak{p}$, then we may use the equivalent statement over DVRs (3.31.7). ∎

**Proposition 3.32.8**
*Let $A$ be a Dedekind with $I, J$ fractional ideals. Then the following are equivalent*

*a) $I \subset J$*
*b) $J$ divides $I$ (that is there exists an ideal $\mathfrak{a}$ such that $\mathfrak{a}J = I$)*
*c) $v_\mathfrak{p}(I) \geq v_\mathfrak{p}(J)$ for all prime ideals $\mathfrak{p}$*

© David Rufino CC BY-NC-ND 4.0

*Proof.* a) $\iff$ c) By (3.30.9) we have $I \subset J \iff I_{\mathfrak{p}} \subset J_{\mathfrak{p}}$ for all prime ideals $\mathfrak{p}$. Then we may use the corresponding statement for DVRs (3.31.8).

b) $\implies$ c) Follows because $v_{\mathfrak{p}}(\mathfrak{a}) \geq 0$ (3.32.7).

c) $\implies$ b) We may define $\mathfrak{a} = \prod \mathfrak{p}^{v_{\mathfrak{p}}(I) - v_{\mathfrak{p}}(J)}$ which is contained in $A$ and by multiplicativity $v_{\mathfrak{p}}(\mathfrak{a}J) = v_{\mathfrak{p}}(I)$ for all prime ideals $\mathfrak{p}$ whence $\mathfrak{a}J = I$ by the correspondence in (3.32.7). ∎

**Proposition 3.32.9 (Integral Closure is a Dedekind Domain)**
*Let $L/K$ be a finite separable extension and $A \subset K$ a subring which is a Dedekind domain with quotient field $K$. Let $B$ be the integral closure of $A$ in $L$, then $B$ is a finite $A$-module and a Dedekind domain.*

*If in addition $A$ is a principal ideal domain (e.g. a discrete valuation ring) then $B$ is a free $A$-module of rank $[L : K]$.*

*Moreover $B \otimes_A K \cong L$, via $b \otimes x \mapsto bx$ (regarding $B, K \subseteq L$).*

*Proof.* From (3.24.24) we know $B$ is integrally closed and $\dim B = 1$. Then the rest follows from (3.24.26).

**Base change to $K$.** Writing $S := A \setminus \{0\}$, (3.6.31) gives an isomorphism $B \otimes_A K \cong S^{-1}B$, $b \otimes x \mapsto bx$, where $S^{-1}B$ is naturally a subring of $L$ (as $S \subset B \setminus \{0\}$ too). By (3.24.15) (applied to the integral map $A \hookrightarrow B$, and $S$), $K = S^{-1}A \to S^{-1}B$ is integral; as $S^{-1}B$ is an integral domain (a subring of $L$) and $K$ is a field, (3.24.14) gives that $S^{-1}B$ is itself a field. Being a field containing $B$, $S^{-1}B$ contains $\mathrm{Frac}(B) = L$; but $S^{-1}B \subseteq L$ already; so $S^{-1}B = L$. ∎

**Definition 3.32.10 (AKLB Situation)**
*Let $L/K$ be a finite field extension, $A \subset K$, $B \subset L$ Dedekind domains for which $A \subset B$ is integral (equivalently $B$ is the integral closure of $A$ in $L$), $\mathrm{Frac}(A) = K$ and $\mathrm{Frac}(B) = L$. We refer to this situation as "AKLB".*

**Lemma 3.32.11**
*Using the notation in (3.32.10). Let $\mathfrak{p}$ be a prime ideal of $A$, and $\mathfrak{b}$ a proper ideal of $B$. Then the following are equivalent*

*a)* $\mathfrak{p} \subset \mathfrak{b}$
*b)* $\mathfrak{b}$ *divides* $\mathfrak{p}B$
*c)* $\mathfrak{b} \cap A = \mathfrak{p}$

*and in this case we say that $\mathfrak{b}$ divides $\mathfrak{p}$ and write $\mathfrak{b} \mid \mathfrak{p}$.*

*In this case $B/\mathfrak{b}$ is naturally an $A/\mathfrak{p}$-module via the commutative diagram*

$$\begin{array}{ccc} A & \hookrightarrow & B \\ \downarrow & & \downarrow \\ A/\mathfrak{p} & \hookrightarrow & B/\mathfrak{b} \end{array}$$

*Proof.* a) $\iff$ b) follows from (3.32.8)

a) $\implies$ c) Trivially $\mathfrak{p} \subset \mathfrak{b} \cap A$. As $1 \in \mathfrak{b}$ then $\mathfrak{b} \cap A$ is a proper ideal, and therefore equal to $\mathfrak{p}$ by maximality.

c) $\implies$ a) Immediate. ∎

**Proposition 3.32.12**
*Using the notation in (3.32.10). Let $\mathfrak{p}$ be a prime ideal of $A$. There exists at least one prime ideal $\mathfrak{q} \mid \mathfrak{p}$, and only finitely many. For every such prime ideal define the integers $e_{\mathfrak{q}/\mathfrak{p}} := v_{\mathfrak{q}}(\mathfrak{p}B)$, then*

$$\mathfrak{p}B = \prod_{\mathfrak{q}|\mathfrak{p}} \mathfrak{q}^{e_{\mathfrak{q}/\mathfrak{p}}} = \bigcap_{\mathfrak{q}|\mathfrak{p}} \mathfrak{q}^{e_{\mathfrak{q}/\mathfrak{p}}}$$

*Proof.* The existence of a prime $\mathfrak{q}$ follows from the Lying Over property of integral extensions (3.24.17). The factorisation exists by (3.32.7) which also shows that there are only finitely many divisors of $\mathfrak{p}$, and the second equality holds by the Chinese Remainder Theorem (3.3.63). ∎

**Definition 3.32.13 (Ramification Index / Inertial Degree)**
*Using the notation in (3.32.10), suppose that $\mathfrak{q} \mid \mathfrak{p}$. Then we define the* ***ramification index*** *by*

$$e_{\mathfrak{q}/\mathfrak{p}} = v_{\mathfrak{q}}(\mathfrak{p}B)$$

*which by (...) is a positive integer. Similarly define the* ***inertial degree***

© David Rufino CC BY-NC-ND 4.0

$$f_{\mathfrak{q}/\mathfrak{p}} = [B/\mathfrak{q} : A/\mathfrak{p}]$$

*which we presently show to be finite.*

**Lemma 3.32.14**
*Let $L/K$ be a finite field extension, $A$ a subring with quotient field $K$ and $B \supset A$ an overring with quotient field $L$. Suppose that $\mathfrak{p}$ is a prime ideal of $A$, $\mathfrak{b}$ is a proper ideal of $B$ containing $\mathfrak{p}$ and $A_\mathfrak{p}$ is a UFD. Then canonically $B/\mathfrak{b}$ is an $A/\mathfrak{p}$-module and*

$$[B/\mathfrak{b} : A/\mathfrak{p}] \leq [L : K]$$

*Furthermore we have $\mathfrak{b}B_\mathfrak{p} \cap A_\mathfrak{p} = \mathfrak{p}A_\mathfrak{p}$ and*

$$[B/\mathfrak{b} : A/\mathfrak{p}] = \left[B_\mathfrak{p}/\mathfrak{b}B_\mathfrak{p} : A_\mathfrak{p}/\mathfrak{p}A_\mathfrak{p}\right]$$

*Proof.* Suppose that $\overline{x_1}, ..., \overline{x_n} \in B/\mathfrak{b}$ are linearly independent over $A/\mathfrak{p}$. It is sufficient by (...) to show that $x_1, ..., x_n$ are linearly independent over $K$. Suppose we have a linear dependence

$$0 = \sum_{i=1}^{n} \lambda_i x_i \quad \lambda_i \in K$$

Then by clearing denominators we may assume that $\lambda_i \in A_\mathfrak{p}$ not all zero. By removing the greatest common divisor of the non-zero elements we may assume that $(\lambda_1, ..., \lambda_n) = A_\mathfrak{p}$, which means at least one is not in $\mathfrak{p}A_\mathfrak{p}$. There is by (3.6.26) a canonical reduction map

$$\tau : A_\mathfrak{p} \to A_\mathfrak{p}/\mathfrak{p}A_\mathfrak{p} \xrightarrow{\sim} A/\mathfrak{p}$$

In otherwords $\overline{\lambda_i} \neq 0$, which gives a linear dependence on $\overline{x_1}, ..., \overline{x_n}$ a contradiction and we are done.

For the last statement consider the commutative diagram

$$\begin{array}{ccc} A/\mathfrak{p} & \hookrightarrow & B/\mathfrak{b} \\ \downarrow \sim & & \downarrow \sim \\ A_\mathfrak{p}/\mathfrak{p}A_\mathfrak{p} & \hookrightarrow & B_\mathfrak{p}/\mathfrak{b}B_\mathfrak{p} \end{array}$$

For the vertical arrows we show they are isomorphism by (3.6.26). Observe that for $s \in A \setminus \mathfrak{p}$ we have by maximality $As + \mathfrak{p} = A$, whence $Bs + \mathfrak{p}B = B \implies Bs + \mathfrak{b} = A$. Then the last two statement follow immediately. ∎

**Lemma 3.32.15 (Local Form of Ramification Index / Inertial Degree)**
*Using the notation in (3.32.10), suppose $\mathfrak{p}$ is a prime ideal and define $A' := A_\mathfrak{p}$, $\mathfrak{p}' := \mathfrak{p}'A_\mathfrak{p}$, $B' := B_\mathfrak{p}$. Then $A', B'$ is also "AKLB" and there is a bijection*

$$\{\mathfrak{q} \lhd B : \mathfrak{q} \mid \mathfrak{p}\} \longleftrightarrow \{\mathfrak{q}' \lhd B'\}$$

*under which the ramification index and inertial degrees agree, i.e.*

$$e_{\mathfrak{q}/\mathfrak{p}} = e_{\mathfrak{q}'/\mathfrak{p}'}$$

*and*

$$f_{\mathfrak{q}/\mathfrak{p}} = f_{\mathfrak{q}'/\mathfrak{p}'} \leq [L : K]$$

*Further the ramification index is equal to*

$$v_{\mathfrak{q}'}(\pi B')$$

*where $\pi$ is any uniformiser for the DVR $A'$.*

*Proof.* Evidently $A'$ has quotient field $K$ and $B'$ has quotient field $L$. By definition $A'$ is a DVR and therefore a Dedekind domain. By (3.24.9) $B'$ is the integral closure of $A'$, and therefore integrally closed. It has Krull Dimension 1 by (3.24.22) and, $B$ being Noetherian (Dedekind, (3.32.1)), $B' = B_\mathfrak{p}$ is Noetherian too (3.15.7). Therefore we have the same situation as (3.32.10).

First consider the commutative diagram

© David Rufino CC BY-NC-ND 4.0

$$\begin{array}{ccc} A & \hookrightarrow & B \\ \downarrow & & \downarrow \\ A' & \hookrightarrow & B' \end{array}$$

regarded as subrings of $K$ and $L$ respectively.

Then by chasing the diagram $\mathfrak{p}'B' = (\mathfrak{p}B)B'$. By (3.32.12) we have a unique factorisation

$$\mathfrak{p}B = \prod_{i=1}^{n} \mathfrak{q}_i^{e_i}$$

for prime ideals $\mathfrak{q}_i$ of $B$ dividing $\mathfrak{p}$. Define $\mathfrak{q}_i' := \mathfrak{q}_i B_{\mathfrak{p}}$ then these are prime by (3.6.18) (and the fact $\mathfrak{q} \cap (A \setminus \mathfrak{p}) = \varnothing$) and contain $\mathfrak{p}'B'$ (3.32.12). Then by (3.3.51)

$$\mathfrak{p}'B' = \prod_{i=1}^{n} (\mathfrak{q}_i')^{e_i}$$

The bijection of prime ideals is well-defined by (3.6.19).

By uniqueness $e_{\mathfrak{q}'/\mathfrak{p}'} = e_{\mathfrak{q}/\mathfrak{p}}$ for all primes $\mathfrak{q} \mid \mathfrak{p}$.

The relation $f_{\mathfrak{q}'/\mathfrak{p}'} = f_{\mathfrak{q}/\mathfrak{p}} \leq [L : K]$ follows directly from (3.32.14). ∎

**Proposition 3.32.16 (Ramification Formula)**
*Using the notation of (3.32.10). For every prime ideal $\mathfrak{p}$ of $A$ we have the identity*

$$\sum_{\mathfrak{q}|\mathfrak{p}} e_{\mathfrak{q}/\mathfrak{p}} f_{\mathfrak{q}/\mathfrak{p}} = [B/\mathfrak{p}B : A/\mathfrak{p}] \leq [L : K]$$

*Suppose in addition that $B$ is a finite $A$-module then we have equality. In particular this holds when $L/K$ is separable.*

*Proof.* Recall from (3.32.12) we have the factorisation

$$\mathfrak{p}B = \prod_{\mathfrak{q}|\mathfrak{p}} \mathfrak{q}^{e_{\mathfrak{q}/\mathfrak{p}}}$$

By the Chinese Remainder Theorem (3.3.63) we have a canonical isomorphism

$$B/\mathfrak{p}B \xrightarrow{\sim} \prod_{\mathfrak{q}|\mathfrak{p}} B/\mathfrak{q}^{e_{\mathfrak{q}}}$$

Both $B/\mathfrak{p}B$ and $B/\mathfrak{q}^{e_{\mathfrak{q}}}$ are $A/\mathfrak{p}$-modules, and so equating vector space dimensions we have

$$[B/\mathfrak{p}B : A/\mathfrak{p}] = \sum_{\mathfrak{q}|\mathfrak{p}} [B/\mathfrak{q}^{e_{\mathfrak{q}}} : A/\mathfrak{p}]$$

As $\mathfrak{p}B \subset \mathfrak{q}^e$ we have a decreasing sequence of $A/\mathfrak{p}$-submodules

$$B/\mathfrak{q}^e \supseteq \mathfrak{q}^1/\mathfrak{q}^e \supseteq \ldots \supseteq \mathfrak{q}^e/\mathfrak{q}^e = (0)$$

so by (3.3.131) we have

$$[B/\mathfrak{q}^e : A/\mathfrak{p}] = \sum_{i=1}^{e} [(\mathfrak{q}^{i-1}/\mathfrak{q}^e)/(\mathfrak{q}^i/\mathfrak{q}^e) : A/\mathfrak{p}]$$

We claim there is an $A/\mathfrak{p}$-module isomorphism

$$(\mathfrak{q}^{i-1}/\mathfrak{q}^e)/(\mathfrak{q}^i/\mathfrak{q}^e) \to \mathfrak{q}^{i-1}/\mathfrak{q}^i$$

This exists as an $A$-module isomorphism by (3.3.106), and therefore as an $A/\mathfrak{p}$-module isomorphism as the action factors through the quotient map $A \to A/\mathfrak{p}$. Therefore by (3.19.15)

© David Rufino CC BY-NC-ND 4.0

$$\begin{aligned}[B/\mathfrak{q}^e : A/\mathfrak{p}] &= \sum_{i=1}^{e} [\mathfrak{q}^{i-1}/\mathfrak{q}^i : A/\mathfrak{p}] \\ &= \sum_{i=1}^{e} [B/\mathfrak{q} : A/\mathfrak{p}][\mathfrak{q}^{i-1}/\mathfrak{q}^i : B/\mathfrak{q}] \\ &= \sum_{i=1}^{e} f_{\mathfrak{q}/\mathfrak{p}}[\mathfrak{q}^{i-1}/\mathfrak{q}^i : B/\mathfrak{q}]\end{aligned}$$

We claim that $[\mathfrak{q}^{i-1}/\mathfrak{q}^i : B/\mathfrak{q}] = 1$. Any non-trivial subspace of $\mathfrak{q}^{i-1}/\mathfrak{q}^i$ is a-fortiori a $B$-submodule, and by the isomorphism theorem yields an ideal between $\mathfrak{q}^{i-1}$ and $\mathfrak{q}^i$. Therefore it is enough to show there is no such ideal. By (3.6.42) we may localize at $\mathfrak{q}$ to consider the ring $B_{\mathfrak{q}}$. By definition this is a DVR and the result follows from (3.31.2). Therefore

$$[B/\mathfrak{p}B : A/\mathfrak{p}] = \sum_{\mathfrak{q}|\mathfrak{p}} e_{\mathfrak{q}/\mathfrak{p}} f_{\mathfrak{q}/\mathfrak{p}}.$$

The inequality follows from (3.32.14).

It remains to show this equals $[B/\mathfrak{p}B : A/\mathfrak{p}] = [L : K]$ in the case $B$ is a finite $A$-module. By (3.32.15) and (3.32.14) we may reduce to the case of $A$ a discrete valuation ring, and in a particular principal ideal domain. Then by (3.24.25) we have $B$ is a free $A$-module of rank $n = [L : K]$. Let $x_1, ..., x_n$ be such a basis, then we claim that $\overline{x_1}, ..., \overline{x_1}$ is linearly independent over $A/\mathfrak{p}$ and therefore a basis for $B/\mathfrak{p}B$. For suppose

$$0 = \sum_{i=1}^{n} \overline{\lambda_i x_i} \quad \lambda_i \in A$$

then we may show that $\mathfrak{p}B = \mathfrak{p}x_1 + ... + \mathfrak{p}x_n$ whence by definition

$$\sum_{i=1}^{n} \lambda_i x_i = \sum_{i=1}^{n} \mu_i x_i$$

for some $\mu_i \in \mathfrak{p}$, from which it follows $\lambda_i = \mu_i$ and $\overline{\lambda_i} = 0$, and we are done.

The case $L/K$ is separable is then (3.24.26). ∎

## 3.32.1 Local Monogenicity

**Lemma 3.32.17 (A Uniformiser Generates a Totally Ramified Extension)**
*Using the notation of (3.32.10), with $L/K$ separable, let $\mathfrak{p}$ be a prime of $A$ with a **unique** prime $\mathfrak{q}$ of $B$ over it, of residue degree $f_{\mathfrak{q}/\mathfrak{p}} = 1$ (so $B_{\mathfrak{p}} = B_{\mathfrak{q}}$ is itself a DVR, totally ramified: $e_{\mathfrak{q}/\mathfrak{p}} = n := [L : K]$). Then for any uniformiser $\pi$ of $B_{\mathfrak{p}}$,*

$$B_{\mathfrak{p}} = A_{\mathfrak{p}}[\pi],$$

*and $\pi$'s minimal polynomial $F \in A_{\mathfrak{p}}[Y]$ over $K$ has degree exactly $n$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

For $0 \leq j < n$, $\mathfrak{q}^j/\mathfrak{q}^{j+1}$ is 1-dimensional over $B_{\mathfrak{p}}/\mathfrak{q} = k$ (as $B_{\mathfrak{p}}$ is a DVR), generated by the image of $\pi^j$; residue degree 1 means $A_{\mathfrak{p}}/\mathfrak{p} = k$ too, so by downward induction on $j$, $\{1, \pi, ..., \pi^{n-1}\}$ spans $B_{\mathfrak{p}}/\mathfrak{p}B_{\mathfrak{p}} = B_{\mathfrak{p}}/\mathfrak{q}^n$ (as $\mathfrak{p}B_{\mathfrak{p}} = \mathfrak{q}^n$, totally ramified) as an $A_{\mathfrak{p}}/\mathfrak{p}$-module. By (3.21.6), applied to the finite $A_{\mathfrak{p}}$-module $B_{\mathfrak{p}}$, a spanning set of $B_{\mathfrak{p}}/\mathfrak{p}B_{\mathfrak{p}}$ already spans $B_{\mathfrak{p}}$ itself, so

$$B_{\mathfrak{p}} = A_{\mathfrak{p}}[\pi].$$

Let $F \in A_{\mathfrak{p}}[Y]$ be $\pi$'s minimal polynomial over $K$, of degree $d$. As $B_{\mathfrak{p}} = A_{\mathfrak{p}}[\pi]$ and $F(\pi) = 0$, $B_{\mathfrak{p}}$ is already spanned by $1, \pi, ..., \pi^{d-1}$ over $A_{\mathfrak{p}}$, so $\operatorname{rank}_{A_{\mathfrak{p}}} B_{\mathfrak{p}} \leq d$; as $B_{\mathfrak{p}}$ is $A_{\mathfrak{p}}$-free of rank $n$ (3.24.25), $d \geq n$, while $d = [K(\pi) : K] \leq [L : K] = n$ as $K(\pi) \subseteq L$. So $d = n$ exactly. ∎

© David Rufino CC BY-NC-ND 4.0

## 3.33 Extensions of Power Series Rings

We record the Weierstrass division lemma for formal power series, used in the analysis of finite extensions of $k((t))$ below.

**Remark 3.33.1**

*[Che51] (Ch. II, §6) treats the case of (3.33.4) where the ramification index $e$ is prime to $\mathrm{char}(k)$, obtaining the sharper conclusion that $t$ itself is an $e$-th power in $k((u))$ (a literal Puiseux expansion $k\big(\big(t^{1/e}\big)\big)$), via Hensel's lemma applied to $W^e - v$ for the unit $v := t/u^e$. That argument is unavailable when $\mathrm{char}(k) = p \mid e$: the derivative $eW^{e-1}$ vanishes identically, so $W^e - v$ has no simple root to lift, and $t$ need not be an $e$-th power at all (an Artin-Schreier extension gives a counterexample). (3.33.4) instead only produces $t = u^e c$ for **some** unit $c$, obtained by solving $\tau G(\tau) = u^e$ by direct coefficient-stabilisation rather than root extraction; this sidesteps the characteristic obstruction entirely, at the cost of the sharper radical form, which is in any case not needed below.*

**Lemma 3.33.2 (Free Basis for a Distinguished Power Series)**

*Let $R$ be a commutative ring, $m \geq 1$ an integer, $w \in R[[X]]^\star$, and $T := X^m w \in R[[X]]$ (so the coefficients of $T$ below degree $m$ vanish, and its coefficient of $X^m$ is $w(0) \in R^\star$). Then every $p \in R[[X]]$ is uniquely*

$$p = \sum_{i=0}^{m-1} f_i(T) X^i$$

*for power series $f_0, \ldots, f_{m-1} \in R[[Y]]$ (substituted at $Y = T$, well-defined as $T$ has zero constant term (3.10.5)).*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

**Existence.** Define $p^{(0)} := p$ and, recursively for $j \geq 0$: writing $p^{(j)} = \sum_n p_n^{(j)} X^n$, set $r_j := \sum_{i=0}^{m-1} p_i^{(j)} X^i$ (the degree-$\{<\}m$ truncation) and $\tilde{q}_j := \sum_{n \geq 0} p_{n+m}^{(j)} X^n$ (the shift), so $p^{(j)} = \tilde{q}_j X^m + r_j$; put $p^{(j+1)} := \tilde{q}_j w^{-1} \in R[[X]]$ (using $w \in R[[X]]^\star$ (3.10.10)), so that $p^{(j)} = p^{(j+1)} T + r_j$. Iterating,

$$p = p^{(J+1)} T^{J+1} + \sum_{j=0}^{J} r_j T^j \quad \text{for every } J \geq 0.$$

As $w(0) \in R^\star$, $T^{J+1} = X^{m(J+1)} w^{J+1}$ has zero coefficients below degree $m(J+1)$; so for each fixed $n$, the coefficient of $X^n$ in $p$ equals the coefficient of $X^n$ in $\sum_{j=0}^{J} r_j T^j$ once $m(J+1) > n$, and this common value no longer depends on $J$ thereafter. Writing $r_j = \sum_{i=0}^{m-1} \gamma_{j,i} X^i$, set $f_i(Y) := \sum_{j \geq 0} \gamma_{j,i} Y^j \in R[[Y]]$; the above shows $p = \sum_{i=0}^{m-1} f_i(T) X^i$, matching the coefficient of every $X^n$.

**Uniqueness.** Suppose $\sum_{i=0}^{m-1} f_i(T) X^i = 0$ with not every $f_i$ zero. Let $j_0 \geq 0$ be minimal such that some $f_i$ has a nonzero coefficient of $Y^{j_0}$, and let $i^\star$ be the least such $i$, so $f_{i^\star}(Y) = \gamma Y^{j_0} + O(Y^{j_0+1})$ for some $0 \neq \gamma \in R$, while $f_i(Y) = O(Y^{j_0+1})$ for $i < i^\star$. Then $f_i(T) = O\big(X^{m(j_0+1)}\big)$ for $i < i^\star$, and $f_{i^\star}(T) = \gamma T^{j_0} + O\big(X^{m(j_0+1)}\big) = \gamma w(0)^{j_0} X^{m j_0} + O(X^{m j_0 + 1})$; as $m j_0 + i^\star < m(j_0+1)$, the coefficient of $X^{m j_0 + i^\star}$ in $\sum_i f_i(T) X^i$ is exactly $\gamma w(0)^{j_0}$, nonzero (a nonzero element of $R$ times a unit is nonzero) – contradicting the sum being 0. ∎

**Corollary 3.33.3 (Specialisation Compatibility)**

*In the setting of (3.33.2), let $\phi : R \to S$ be a homomorphism of commutative rings, extended coefficientwise to $\phi : R[[X]] \to S[[X]]$. Then $\phi(w) \in S[[X]]^\star$, and the decomposition of $\phi(p)$ relative to $\phi(T) = X^m \phi(w)$ is $\phi(p) = \sum_{i=0}^{m-1} \phi(f_i)(\phi(T)) X^i$, where $\phi(f_i) \in S[[Y]]$ denotes $\phi$ applied coefficientwise to $f_i$.*

*Proof.* If $wv = 1$ then $\phi(w)\phi(v) = 1$, so $\phi(w) \in S[[X]]^\star$. Next, $\phi(f(T)) = \phi(f)(\phi(T))$ for every $f \in R[[Y]]$. Fix $N \geq 0$ and write $f = s + Y^N g$ with $s \in R[Y]$ of degree $< N$ and $g \in R[[Y]]$. Then $f(T) = s(T) + T^N g(T)$ (the tail $\sum_{n \geq N} c_n T^n = T^N \sum_n c_{n+N} T^n$, (3.10.8)), and likewise $\phi(f) = \phi(s) + Y^N \phi(g)$, so $\phi(f)(\phi(T)) = \phi(s)(\phi(T)) + \phi(T)^N \phi(g)(\phi(T))$. Applying the ring homomorphism $\phi$ to $f(T) = s(T) + T^N g(T)$ gives $\phi(f(T)) = \phi(s)(\phi(T)) + \phi(T)^N \phi(g(T))$, so $\phi(f(T)) - \phi(f)(\phi(T)) \in \phi(T)^N S[[X]] \subseteq X^{mN} S[[X]]$, as $\phi(T) = X^m \phi(w)$. This holds for every $N$ and $\bigcap_N X^N S[[X]] = 0$, so $\phi(f(T)) = \phi(f)(\phi(T))$. Applying $\phi$ to $p = \sum_i f_i(T) X^i$ therefore gives $\phi(p) = \sum_i \phi(f_i)(\phi(T)) X^i$. This exhibits $\phi(p)$ in the form guaranteed unique by (3.33.2) over $S$, so it is **the** decomposition of $\phi(p)$. ∎

**Lemma 3.33.4 (Finite Separable Extensions of $k((t))$ are Power Series Fields)**

*Let $k$ be algebraically closed, $F := k((t))$, and $E/F$ a finite separable extension of degree $n$. Then there are $u \in E$, an element $\tau \in uk[[u]]$ with $\tau = u^n c$ for some $c \in k[[u]]^\star$, and a field isomorphism $\Psi : E \xrightarrow{\sim} k((u))$ with $\Psi|_F$ the substitution $t \mapsto \tau$ and $\Psi(u) = u$. In particular $1, u, \ldots, u^{n-1}$ is an $F$-basis of $E$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

© David Rufino CC BY-NC-ND 4.0

**Step 1: a DVR.** Let $\mathcal{O}_F := k[[t]]$ and let $C$ be the integral closure of $\mathcal{O}_F$ in $E$. By (3.24.26) $C$ is a finite $\mathcal{O}_F$-module, and by (3.32.9) it is Dedekind. By lying-over (3.24.17) there is a prime $\mathfrak{p} \subset C$ with $\mathfrak{p} \cap \mathcal{O}_F = (t)$. Put $B := C_{\mathfrak{p}}$, a DVR with fraction field $E$, and let $e := v_B(t) \geq 1$.

**Step 2: residue field and valuation.** $C/\mathfrak{p}$ is a domain, finite over $\mathcal{O}_F/(t) = k$, so $k \hookrightarrow C/\mathfrak{p}$ is injective and integral, hence $C/\mathfrak{p}$ is a field (3.24.14), finite over $k$. As $B = C_{\mathfrak{p}}$, (3.6.43) gives $B/\mathfrak{m}_B \cong \mathrm{Frac}(C/\mathfrak{p}) = C/\mathfrak{p}$ ($C/\mathfrak{p}$ already a field); as $k$ is algebraically closed, $k \to B/\mathfrak{m}_B$ is an isomorphism. Every $a \in \mathcal{O}_F^\times$ is a unit of $B$, so $v_B(x) = ev_F(x)$ for $x \in F^\times$. Let $u$ be a uniformiser of $B$, so $t = u^e w$ with $w \in B^\star$.

**Step 3: expansions.** Every $b \in B$ can be written $b = \sum_{i<e} f_i u^i$ with $f_i \in k[[t]]$. We show by induction on $N \geq 0$ that $b \equiv \sum_{i<e}\left(\sum_{j<N} c_{ij}t^j\right)u^i \pmod{t^N B}$ for some $c_{ij} \in k$. The case $N = 0$ is trivial. For the inductive step, write the current remainder as $t^N y$ with $y \in B$ (inductive hypothesis); since $B/uB = k$, repeated division by $u$ gives $y = \sum_{i<e} d_i u^i + u^e y'$ with $d_i \in k$, $y' \in B$ (there is no need to iterate past degree $e-1$: stop as soon as $e$ terms are extracted). Using $u^e = tw^{-1}$,

$$t^N y = \sum_{i<e}(t^N d_i)u^i + t^{N+1}(w^{-1}y'),$$

with $w^{-1}y' \in B$; so taking $c_{i,N} := d_i$ advances the induction, with new remainder $t^{N+1}(w^{-1}y') \in t^{N+1}B$. Put $f_i := \sum_j c_{ij}t^j \in k[[t]]$. Then $z := b - \sum_i f_i u^i$ satisfies $z \in t^N B$ for every $N$, so $v_B(z) \geq eN$ for all $N$ and $z = 0$.

**Step 4: basis and degree.** Every element of $E$ has the form $t^{-N}b$ with $b \in B$, so Step 3 gives $E = \sum_{i<e} Fu^i$. These elements are independent: if $\sum_{i<e} f_i u^i = 0$ with $f_i \in F$ not all zero, the valuations $v_B(f_i u^i) = ev_F(f_i) + i$ are distinct modulo $e$ for the nonzero $f_i$, so the minimum is attained exactly once, a contradiction. Hence $n = e$, and $1, \ldots, u^{e-1}$ is a basis. Moreover $B = \bigoplus_{i<e} k[[t]]u^i$, by the valuation estimate $\min_i(ev_F(f_i) + i) = v_B(b) \geq 0$ applied to $b = \sum_i f_i u^i$ with $f_i \in F$.

**Step 5: the minimal polynomial.** Write $w^{-1} = \sum_{i<e} g_i u^i$ with $g_i \in k[[t]]$ (Step 3). The constant $g_0(0)$ is the residue of $w^{-1}$, so $g_0(0) \in k^\star$. From $u^e = tw^{-1}$, $u$ is a root of

$$P(Y) := Y^e - t\sum_{i<e} g_i(t)Y^i \in F[Y].$$

Since $F[u] = E$ has degree $e$ over $F$, $P$ is the minimal polynomial of $u$, and $E \cong F[Y]/(P)$ with $Y \mapsto u$.

**Step 6: solving for $\tau$.** For $s \in uk[[u]]$ put $G(s) := \sum_{i<e} g_i(s)u^i \in k[[u]]$, where $g_i(s)$ is the substitution (well defined as $s$ has zero constant term). Since $G(s) \equiv g_0(0) \pmod{u}$, $G(s)$ is a unit. Define $T(s) := u^e G(s)^{-1} \in u^e k[[u]]$. For $s \equiv s' \pmod{u^N}$ we have $g_i(s) - g_i(s') \in (s-s')k[[u]] \subseteq u^N k[[u]]$, hence $G(s) \equiv G(s') \pmod{u^N}$ and $T(s) \equiv T(s') \pmod{u^{N+e}}$. Starting from $s_0 := 0$ and setting $s_{m+1} := T(s_m)$, the sequence satisfies $s_{m+1} \equiv s_m \pmod{u^{m+e}}$, so each coefficient stabilises. Let $\tau \in uk[[u]]$ be the coefficientwise limit. Then $\tau \equiv T(\tau) \pmod{u^N}$ for every $N$, so $\tau = T(\tau)$, i.e. $\tau G(\tau) = u^e$. Also $\tau = u^e c$ with $c \in k[[u]]^\star$, since $\tau G(\tau) = u^e$ and $G(\tau)$ is a unit.

**Step 7: the homomorphism.** The substitution $t \mapsto \tau$ defines a ring homomorphism $\sigma : k[[t]] \to k[[u]]$ ((3.10.19), as $\tau$ has zero constant term), and since $\tau$ is a unit times $u^e$ it extends to a field homomorphism $\sigma : F \to k((u))$. The homomorphism $F[Y] \to k((u))$, $Y \mapsto u$, extending $\sigma$, kills $P$:

$$\sigma(P)(u) = u^e - \tau\sum_{i<e} g_i(\tau)u^i = u^e - \tau G(\tau) = 0.$$

So it factors through $E \cong F[Y]/(P)$, giving a field homomorphism $\Psi : E \to k((u))$ with $\Psi|_F = \sigma$ and $\Psi(u) = u$. It is injective, as $E$ is a field.

**Step 8: surjectivity.** Since $\tau = u^e c$ with $c \in k[[u]]^\star$, the Weierstrass division lemma (3.33.2) (with $R := k$, $X := u$, $m := e$, $w := c$, so $T = \tau$) gives $k[[u]] = \bigoplus_{i<e} \sigma(k[[t]])u^i$. Multiplying by powers of $\tau$ gives $k((u)) = \sum_{i<e} \sigma(F)u^i = \sigma(F)[u]$. But $\operatorname{im}\Psi = \Psi(F[u]) = \sigma(F)[u]$, so $\Psi$ is surjective.

Hence $\Psi : E \xrightarrow{\sim} k((u))$ is an isomorphism of fields over $F$, with $\Psi(u) = u$ and $\Psi(t) = \tau = u^n c$. ∎

**Corollary 3.33.5 (Uniqueness of the Extended Valuation)**
*In the setting of (3.33.4), let $C$ be the integral closure of $k[[t]]$ in $E$. Then $C$ has exactly one prime $\mathfrak{p}$ lying over $(t)$, the DVR $C_{\mathfrak{p}}$ is the unique valuation ring of $E$ containing $k[[t]]$ whose maximal ideal contracts to $(t)$, and its ramification index is $[E : F]$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

© David Rufino CC BY-NC-ND 4.0

Let $\mathfrak{q}$ be any prime of $C$ over $(t)$, which exists by lying-over. Steps 2–4 of the proof of (3.33.4) apply verbatim to the DVR $B := C_{\mathfrak{q}}$ (which has residue field $k$), and give $e_{\mathfrak{q}} = v_B(t) = [E : F] = n$.

Now apply the Ramification Formula (3.32.16) to the AKLB situation $k[[t]] \subset C$ over $E/F$ (both Dedekind by (3.32.9), and $E/F$ separable), at the prime $(t)$:

$$\sum_{\mathfrak{q}|(t)} e_{\mathfrak{q}} f_{\mathfrak{q}} = [C/tC : k] = n.$$

Each $f_{\mathfrak{q}} = [C/\mathfrak{q} : k]$ is finite, and $C/\mathfrak{q}$ is a field algebraic over $k$, hence $f_{\mathfrak{q}} = 1$. So $\sum_{\mathfrak{q}} e_{\mathfrak{q}} = n$. By the previous paragraph each term equals $n$, so there is exactly one prime $\mathfrak{p}$ over $(t)$ and $e_{\mathfrak{p}} = n$.

For uniqueness of the valuation ring, let $V \subseteq E$ be a valuation ring with $k[[t]] \subseteq V$ and $\mathfrak{m}_V \cap k[[t]] = (t)$. By (3.25.14), $C \subseteq V$. Then $\mathfrak{m}_V \cap C$ is a prime of $C$ over $(t)$, hence equals $\mathfrak{p}$. Every element of $C \setminus \mathfrak{p}$ lies outside $\mathfrak{m}_V$, so is a unit of $V$; hence $C_{\mathfrak{p}} \subseteq V$. Since $C_{\mathfrak{p}}$ is a valuation ring of $E$ and valuation rings are maximal for domination among proper subrings of $E$ (3.25.2), $V = C_{\mathfrak{p}}$. ∎

© David Rufino CC BY-NC-ND 4.0

# 3.34 Complete Discrete Valuation Rings

We isolate the algebraic (topology-free) notion of completeness for a DVR, needed to make sense of "formal" local models such as $k[[t]]$, and prove the key structural fact used throughout: a complete DVR has a **unique** extension of its valuation to any finite separable extension of its fraction field.

This generalises the previous section, which proved the case $\mathcal{O}_K \cong k[[t]]$ by direct computation in power series, without this general completion machinery: (3.33.4) and (3.33.5) there are the special cases of (3.34.19) and (3.34.20) here.

**Definition 3.34.1 (Complete DVR)**
*A DVR $A$ with maximal ideal $\mathfrak{m}$ is* complete *if every sequence $(\overline{y}_n)_{n\geq 0}$ with $\overline{y}_n \in A/\mathfrak{m}^{n+1}$, compatible under reduction, is the image of some element of $A$ (equivalently, $A \to \varprojlim_n A/\mathfrak{m}^{n+1}$ is surjective; see (3.34.5) for the general criterion).*

**Proposition 3.34.2 (The Formal Power Series Ring is a Complete DVR)**
*Let $k$ be a field. Then $k[[t]]$ is a complete DVR.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

By (3.10.14), $k[[t]]$ is a discrete valuation ring, so it remains to show completeness.

Let $(\overline{y}_n)_{n\geq 0}$ with $\overline{y}_n \in k[[t]]/(t)^{n+1}$ be compatible under reduction. Each $\overline{y}_n$ has a unique representative $f_n = \sum_{i\leq n} c_i^{(n)} t^i$ of degree at most $n$. Compatibility says $f_n \equiv f_{n-1} \pmod{t^n}$, so $c_i^{(n)} = c_i^{(n-1)}$ for $i < n$. Hence for each $i$ the coefficient $c_i := c_i^{(n)}$ is independent of $n \geq i$. Put $f := \sum_i c_i t^i \in k[[t]]$. For each $n$, the coefficients of $f$ up to degree $n$ are those of $f_n$, so $f \equiv f_n \pmod{t^{n+1}}$, that is $\overline{y}_n$ is the reduction of $f$. Thus $k[[t]]$ is complete. ∎

**Proposition 3.34.3 (A Complete DVR with a Coefficient Field is a Power Series Ring)**
*Let $(A, \mathfrak{m})$ be a complete DVR, $\pi$ a uniformiser, and $k \subseteq A$ a coefficient field ($k(\mathfrak{m}) = k$). Then there is a $k$-algebra isomorphism $k[[Y]] \xrightarrow{\sim} A$, $Y \mapsto \pi$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

**Coefficients.** Since $A/\mathfrak{m} = k$ and $\mathfrak{m} = (\pi)$, every $a \in A$ can be written $a \equiv \sum_{i<N} c_i \pi^i \pmod{\pi^N}$ with $c_i \in k$, and the $c_i$ are uniquely determined by $a$ (apply $v$). Indeed $a - c_0 \in \mathfrak{m} = \pi A$ gives $a = c_0 + \pi a_1$, and iterate.

**Definition of $\Phi$.** Let $f = \sum_n c_n Y^n \in k[[Y]]$ with truncations $s_N := \sum_{n<N} c_n \pi^n$. Since $s_{N+1} - s_N = c_N \pi^N \in \pi^N A$, the sequence $(s_N \mod \pi^N)_N$ is compatible under reduction, so by completeness of $A$ (3.34.1) there is $\Phi(f) \in A$ with $\Phi(f) \equiv s_N \pmod{\pi^N}$ for every $N$. It is unique, as $\bigcap_N \pi^N A = 0$.

**Homomorphism.** $\Phi$ is $k$-linear, and $\Phi(1) = 1$. For $f, g$, the truncations satisfy $s_N(f)s_N(g) \equiv s_N(fg) \pmod{\pi^N}$ (the products agree modulo $\pi^N$, since the difference only involves coefficients of degree $\geq N$ in $Y$, sent to multiples of $\pi^N$). Hence $\Phi(f)\Phi(g) \equiv s_N(f)s_N(g) \equiv s_N(fg) \equiv \Phi(fg) \pmod{\pi^N}$ for all $N$, so $\Phi(f)\Phi(g) = \Phi(fg)$. Additivity is the same argument.

**Injective.** If $\Phi(f) = 0$ and $f \neq 0$, let $n_0$ be minimal with $c_{n_0} \neq 0$. Then $\pi^{n_0} c_{n_0} \equiv \Phi(f) - \sum_{n>n_0} c_n \pi^n \equiv 0 \pmod{\pi^{n_0+1}}$, so $c_{n_0}\pi^{n_0} \in \pi^{n_0+1}A$, contradicting the uniqueness of the coefficients above.

**Surjective.** Given $a \in A$, take its coefficients $c_i \in k$ from the first paragraph. These are compatible (the expansion modulo $\pi^N$ is the truncation of the expansion modulo $\pi^{N+1}$), so $f := \sum_i c_i Y^i$ satisfies $\Phi(f) \equiv a \pmod{\pi^N}$ for every $N$, whence $\Phi(f) = a$. ∎

**Definition 3.34.4 (Completion of a Local Ring)**
*Let $(A, \mathfrak{m})$ be a local ring, with $\mathfrak{m}$-adic filtration $\mathfrak{m}_\bullet$ (3.14.4) (so $\mathfrak{m}_n = \mathfrak{m}^{n+1}$). Write $\hat{A} := \hat{A}_\mathfrak{m}$ for its $\mathfrak{m}$-adic completion, $\iota := \iota_\mathfrak{m} : A \to \hat{A}$ for the canonical map, and $\lambda_n : \hat{A} \to A/\mathfrak{m}^{n+1}$ for its projections (3.14.19).*

*For $x = (\overline{a}_n)_{n\geq 0} \in \hat{A}$ and $j \geq k \geq 0$: $\overline{a}_j \equiv \overline{a}_k \pmod{\mathfrak{m}^{k+1}}$ (repeated compatibility (3.14.16)).*

**Proposition 3.34.5 (Completeness Criterion)**
*Let $(A, \mathfrak{m})$ be a local ring with $\bigcap_n \mathfrak{m}^n = 0$ (automatic if $A$ is a DVR (3.31.9)), and $\hat{A}$, $\iota$ as in (3.34.4). Then the following are equivalent:*

*a) $\iota : A \to \hat{A}$ is surjective;*
*b) every sequence $(\overline{a}_n)_{n\geq 0}$, $\overline{a}_n \in A/\mathfrak{m}^{n+1}$, compatible under reduction ($\overline{a}_n \equiv \overline{a}_{n-1} \mod \mathfrak{m}^n$ for $n \geq 1$), equals $(a \mod \mathfrak{m}^{n+1})_n$ for some $a \in A$;*
*c) for some strictly increasing sequence $n_0 < n_1 < n_2 < \cdots$ in $\mathbb{N}$ (equivalently, for every such sequence), every sequence $(\overline{a}_N)_{N\geq 0}$, $\overline{a}_N \in A/\mathfrak{m}^{n_N+1}$, compatible under reduction, equals $(a \mod \mathfrak{m}^{n_N+1})_N$ for some $a \in A$.*

© David Rufino CC BY-NC-ND 4.0

*When these hold, $\iota$ is an isomorphism, and we say $A$ is **complete**.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

This is the special case $\mathfrak{a} := \mathfrak{m}$ of (3.14.21), noting $\hat{A}_{\mathfrak{m}} = \hat{A}$, $\iota_{\mathfrak{m}} = \iota$. ∎

**Proposition 3.34.6 (The Completion of a Local Ring is Local)**
*Let $(A, \mathfrak{m})$ be a local ring, and $\hat{A}$, $\lambda_n$ as in (3.34.4). Then $\hat{A}$ is local, with units, for $(a_n)_{n\geq 0} \in \hat{A}$, characterised by*

$$(a_n)_{n\geq 0} \in \hat{A}^\star \iff a_0 \in (A/\mathfrak{m})^\star$$

*maximal ideal $\hat{\mathfrak{m}} := \ker \lambda_0 = \hat{A} \setminus \hat{A}^\star$, characterised, for $(a_n)_{n\geq 0} \in \hat{A}$, by*

$$(a_n)_{n\geq 0} \in \hat{\mathfrak{m}} \iff a_0 = 0$$

*and $\lambda_0$ inducing an isomorphism $\hat{A}/\hat{\mathfrak{m}} \xrightarrow{\sim} A/\mathfrak{m}$: that is, $\hat{A}$ and $A$ have the same residue field.*

*Proof.* **Units.** One direction is obvious by applying $\lambda_0$. Conversely suppose $x = (a_n)_{n\geq 0} \in \hat{A}$ and $a_0 \in (A/\mathfrak{m})^\star$ (so in particular $a_0 \neq 0$ (3.3.7)). Then by definition $\rho_{n,0}(a_n) = a_0$ which by (3.3.56) means $a_n \notin \mathfrak{m}/\mathfrak{m}^{n+1}$. As $A/\mathfrak{m}^{n+1}$ is local with maximal ideal $\mathfrak{m}/\mathfrak{m}^{n+1}$ (3.20.4), $a_n \notin \mathfrak{m}/\mathfrak{m}^{n+1}$ gives $a_n \in (A/\mathfrak{m}^{n+1})^\star$ (3.20.3). Let $b_n := (a_n)^{-1}$, then we claim the $b_n$ are compatible. For $n \geq 1$: $1 = \rho_{n,n-1}(a_n b_n) = \rho_{n,n-1}(b_n)\rho_{n,n-1}(a_n) = \rho_{n,n-1}(b_n)a_{n-1}$. By uniqueness of inverses (3.1.5), $\rho_{n,n-1}(b_n) = b_{n-1}$. Therefore these determine a well-defined element $y = (b_n)_{n\geq 0} \in \hat{A}$ which is an inverse for $x$ as required.

**Locality.** $\ker \lambda_0$ is an additive subgroup of $\hat{A}$ (3.3.14), so by (3.20.3) it is enough to show that $\ker \lambda_0 = \hat{A} \setminus \hat{A}^\star$ which will also give the form of the maximal ideal $\hat{\mathfrak{m}}$. However $A/\mathfrak{m}$ is a field (3.3.59) so by the first part

$$(a_n)_{n\geq 0} \in \ker \lambda_0 \iff a_0 = 0 \iff a_0 \notin (A/\mathfrak{m})^\star \iff (a_n)_{n\geq 0} \notin \hat{A}^\star$$

that is, $\ker \lambda_0 = \hat{A} \setminus \hat{A}^\star$, which is also the unique maximal ideal.

**Residue field.** As $\lambda_0$ is surjective (3.14.24) with $\ker \lambda_0 = \hat{\mathfrak{m}}$, (3.3.57) gives the claimed isomorphism $\hat{A}/\hat{\mathfrak{m}} \xrightarrow{\sim} A/\mathfrak{m}$. ∎

**Lemma 3.34.7 (Finite Free Modules over a Complete Ring are Complete)**
*Let $(A, \mathfrak{m})$ be a complete local ring (3.34.5), and $M$ a free $A$-module of finite rank $r$. Then*

$$\iota_M : M \to \varprojlim_{N\geq 1} M/\mathfrak{m}^N M, \quad x \mapsto (x \mod \mathfrak{m}^N M)_{N\geq 1},$$

*is a bijection.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Fix an $A$-basis $e_1, \ldots, e_r$ of $M$, giving an $A$-module isomorphism $\phi : A^r \xrightarrow{\sim} M$, $(a_i)_{i=1}^r \mapsto \sum_i a_i e_i$; note $\mathfrak{m}^N M = \mathfrak{m}^N\left(\sum_i A e_i\right) = \sum_i \mathfrak{m}^N e_i$ for every $N \geq 1$.

**Existence of $\overline{\phi}_N$.** Fix $N \geq 1$, and write $\sigma_N : A^r \to (A/\mathfrak{m}^N)^r$ for the coordinatewise reduction (a surjection with $\ker \sigma_N = (\mathfrak{m}^N)^r$) and $\pi_N : M \to M/\mathfrak{m}^N M$ for the quotient map. We first check $\ker(\pi_N \circ \phi) = (\mathfrak{m}^N)^r$ too: for $(a_i)_i \in A^r$, $\phi\big((a_i)_i\big) = \sum_i a_i e_i$, so $(a_i)_i \in \ker(\pi_N \circ \phi) \iff \sum_i a_i e_i \in \mathfrak{m}^N M = \sum_i \mathfrak{m}^N e_i \iff a_i \in \mathfrak{m}^N$ for every $i$ (the forward implication using injectivity of $\phi$: if $\sum_i a_i e_i = \sum_i c_i e_i$ for $c_i \in \mathfrak{m}^N$, then $a_i = c_i \in \mathfrak{m}^N$ for every $i$), that is, $(a_i)_i \in (\mathfrak{m}^N)^r = \ker \sigma_N$.

In particular $\ker \sigma_N \subseteq \ker(\pi_N \circ \phi)$, so by the universal property of the quotient $\sigma_N$ (3.3.101), there is a unique $A$-module homomorphism $\overline{\phi}_N$ making

$$\begin{array}{ccc} A^r & \xrightarrow{\quad\phi\quad}_{\sim} & M \\ {\scriptstyle\sigma_N}\big\downarrow & & \big\downarrow{\scriptstyle\pi_N} \\ (A/\mathfrak{m}^N)^r & \xrightarrow[\overline{\phi}_N]{} & M/\mathfrak{m}^N M \end{array}$$

commute, that is $\overline{\phi}_N \circ \sigma_N = \pi_N \circ \phi$, explicitly $\overline{\phi}_N\big((\overline{a}_i)_{i=1}^r\big) = \sum_i a_i e_i \mod \mathfrak{m}^N M$ for any representatives $a_i \in A$ of $\overline{a}_i$.

$\overline{\phi}_N$ is an isomorphism: by (3.3.101), $\ker\big(\overline{\phi}_N\big) = \ker(\pi_N \circ \phi)/\ker \sigma_N$, and this is 0 as $\ker(\pi_N \circ \phi) = \ker \sigma_N$ (shown above), so $\overline{\phi}_N$ is injective; it is also surjective, as $\pi_N \circ \phi = \overline{\phi}_N \circ \sigma_N$ is (a composite of the surjections $\phi$, $\pi_N$) and $\sigma_N$ is surjective.

© David Rufino CC BY-NC-ND 4.0

**Compatibility of the $\overline{\phi}_N$.** For $P \geq N \geq 1$, the square

$$\begin{array}{ccc} (A/\mathfrak{m}^P)^r & \xrightarrow[\sim]{\overline{\phi}_P} & M/\mathfrak{m}^P M \\ \downarrow & & \downarrow \\ (A/\mathfrak{m}^N)^r & \xrightarrow[\overline{\phi}_N]{\sim} & M/\mathfrak{m}^N M \end{array}$$

(vertical arrows the evident reduction maps (3.3.56)) commutes: both composites send the class of $(a_i)_i \in A^r$ to $\sum_i a_i e_i \mod \mathfrak{m}^N M$, directly from the explicit formula for $\overline{\phi}_N$ above (resp. for $\overline{\phi}_P$, followed by reduction mod $\mathfrak{m}^N M$). Thus $\left(\overline{\phi}_N\right)_{N\geq 1}$ is a compatible family of isomorphisms between the towers $\left((A/\mathfrak{m}^N)^r\right)_{N\geq 1}$ and $(M/\mathfrak{m}^N M)_{N\geq 1}$.

**Passing to the limit.** By (3.3.157) (applied to $r$ copies of the tower $(A/\mathfrak{m}^N)_{N\geq 1}$), $\prod_{i=1}^r \varprojlim_N A/\mathfrak{m}^N \cong \varprojlim_N (A/\mathfrak{m}^N)^r$; composing with $\varprojlim \overline{\phi}$ (an isomorphism, (3.3.156)) gives an isomorphism

$$\left(\varprojlim_N A/\mathfrak{m}^N\right)^r \xrightarrow{\sim} \varprojlim_N M/\mathfrak{m}^N M$$

sending $\left((a_i \mod \mathfrak{m}^N)_N\right)_{i=1}^r$ to $\left(\sum_i a_i e_i \mod \mathfrak{m}^N M\right)_N$ for $a_i \in A$ (by the explicit formula for $\overline{\phi}_N$ above), that is, under the fixed basis, $\iota_M$ corresponds to $\iota_A^{\times r}$ (the $r$-fold product of $A \to \varprojlim_N A/\mathfrak{m}^N$, $a \mapsto (a \mod \mathfrak{m}^N)_N$). As $A$ is complete, $\iota_A : A \to \hat{A} = \varprojlim_n A/\mathfrak{m}^{n+1}$ is bijective (3.34.5) reindexing ($N := n+1$), $A \to \varprojlim_N A/\mathfrak{m}^N$, $a \mapsto (a \mod \mathfrak{m}^N)_N$, is bijective too, hence so is its $r$-fold product, hence so is $\iota_M$. ∎

**Proposition 3.34.8 (The Completion is a DVR)**
*Let $(A, \mathfrak{m})$ be a DVR, $K := \mathrm{Frac}(A)$, and $\hat{A}$ the completion (3.34.4), $K_{\mathfrak{m}} := \mathrm{Frac}\left(\hat{A}\right)$. Then $\hat{A}$ is a DVR. Fixing a uniformiser $\pi$ of $A$ and writing $\hat{\pi} := \iota(\pi)$:*

*a) $\hat{\pi}$ is a uniformiser of $\hat{A}$;*
*b) the valuation of $\hat{A}$ is given, for $0 \neq x = (\overline{a}_m)_{m\geq 0} \in \hat{A}$, by $v_{\hat{A}}(x) = \min\{m \geq 0 : \overline{a}_m \neq 0\}$;*
*c) the maximal ideal of $\hat{A}$ is $\hat{\mathfrak{m}} = (\hat{\pi}) = \left\{(\overline{a}_m)_{m\geq 0} \in \hat{A} : \overline{a}_0 = 0\right\} = \ker \lambda_0$;*
*d) the units of $\hat{A}$ are $\hat{A}^\star = \left\{(\overline{a}_m)_{m\geq 0} \in \hat{A} : \overline{a}_0 \neq 0\right\}$;*
*e) $\ker\left(\hat{A} \to A/\mathfrak{m}^{n+1}\right) = \hat{\mathfrak{m}}^{n+1}$, for every $n \geq 0$ (generalising c), the case $n = 0$).*

*Consequently, the valuations $v$ of $A$ and $v_{\hat{A}}$ of $\hat{A}$ extend uniquely (3.31.2) to $K^\star$ and $K_{\mathfrak{m}}^\star$ respectively, explicitly by*

$$v(a/b) := v(a) - v(b) \quad (a, b \in A \setminus \{0\}), \qquad v_{\hat{A}}(x/y) := v_{\hat{A}}(x) - v_{\hat{A}}(y) \quad \left(x, y \in \hat{A} \setminus \{0\}\right).$$

*Proof.* **Units**: By (3.34.6) $\hat{A}$ is local with unique maximal ideal $\hat{\mathfrak{m}} := \ker \lambda_0$; so $\hat{A}^\star = \hat{A} \setminus \ker \lambda_0 = \left\{(\overline{a}_m)_{m\geq 0} \in \hat{A} : \overline{a}_0 \neq 0\right\}$.

**$\hat{\pi}$ is a non-unit; $\widehat{A}$ is not a field.** As $\iota$ is injective and $\pi \neq 0$, $\hat{\pi} = \iota(\pi) \neq 0$. Writing $\hat{\pi} = (\overline{\pi}_m)_m$, $\overline{\pi}_0 = \pi \mod \mathfrak{m} = 0$ (as $\pi \in \mathfrak{m}$), so $\hat{\pi} \notin \hat{A}^\star$ by d). As every non-unit of a local ring lies in its maximal ideal (3.20.3), $\hat{\pi} \in \ker \lambda_0$; so $\hat{\mathfrak{m}} := \ker \lambda_0 \neq 0$ (as $\hat{\pi} \neq 0$), and $\hat{A}$ is therefore not a field (a field has no nonzero proper ideal).

**Valuation of $\widehat{A}$.** Define

$$v_{\hat{A}}\left((a_n)_{n\geq 0}\right) := \min\{m \mid a_m \neq 0\}$$

For $j \geq k$ we may show that $a_j \in \mathfrak{m}^k \setminus \mathfrak{m}^{k+1}$ and therefore $v(a_j) = k$ (3.31.3).

**$\widehat{A}$ is a domain.** Suppose $0 \neq x = (\overline{a}_m)_m$, $0 \neq y = \left(\overline{b}_m\right)_m \in \hat{A}$ with $xy = 0$; set $k := v_{\hat{A}}(x)$, $l := v_{\hat{A}}(y)$. Representatives $a, b \in A$ of $\overline{a}_{k+l}, \overline{b}_{k+l}$ have $v(a) = k$, $v(b) = l$ so $v(ab) = k + l$ (3.31.2), that is $ab \notin \mathfrak{m}^{k+l+1}$ (3.31.3) but $ab$ represents the $(k+l)$-th coordinate of $xy = 0$, forcing $ab \in \mathfrak{m}^{k+l+1}$: a contradiction. So $\hat{A}$ is a domain.

**$\widehat{A}$ is a DVR.** We show that for every $0 \neq x$ there is a $w \in \hat{A}^\star$ and $k \geq 0$ such that $x = w\hat{\pi}^k$, namely criteria (3.31.2) e) to be a DVR.

Suppose $x = (\overline{a}_n)_{n\geq 0}$ with some representatives $a_n \in A$. Let $k := v_{\hat{A}}(x)$ and recall $j \geq k \Longrightarrow v(a_j) = k$, so by the same criteria $a_j = u_j \pi^k$ for some $u_j \in A^\star$ uniquely depending on $a_j$.

Define $\overline{w}_n := u_{n+k} + \mathfrak{m}^{n+1}$. By definition of $a_n$,

© David Rufino CC BY-NC-ND 4.0

$$a_{n+k+1} \equiv a_{n+k} \mod \pi^{n+k+1} \implies u_{n+k+1} \equiv u_{n+k} \mod \pi^{n+1}$$

by cancelling $\pi^k$. This means precisely that $\rho_{n+1,n}(\overline{w}_{n+1}) = \overline{w}_n$ so $w := (\overline{w}_n)_{n\geq 0}$ is a well-defined element of $\hat{A}$.

Recall $\overline{w}_0 = u_k + \mathfrak{m}$. By assumption $u_k \in A^\star$, whence $u_k \notin \mathfrak{m}$, which shows $\overline{w}_0 \neq 0$. Therefore by Units, we have $w := (\overline{w}_n)_{n\geq 0} \in \hat{A}^\star$.

By definition $\lambda_n\big(w\hat{\pi}^k\big) = \overline{w}_n\big(\pi^k + \mathfrak{m}^{n+1}\big) = u_{n+k}\pi^k + \mathfrak{m}^{n+1} = a_{n+k} + \mathfrak{m}^{n+1} = \overline{a}_n = \lambda_n(x)$. By construction of $\hat{A}_n$ as an inverse limit (3.3.154) we have $x = w\hat{\pi}^k$ as required.

**Valuation, maximal ideal, kernel of $\lambda_n$.** For $0 \neq x = (\overline{a}_m)_m \in \hat{A}$, the uniformised representation gives $x = \hat{\pi}^k u$ for the unit $u$ constructed above and $k = \min\{m \geq 0 : \overline{a}_m \neq 0\}$; by the uniqueness of such a representation (3.31.2), $v_{\hat{A}}(x) = k$, giving b).

Let $x = (a_n)_{n\geq 0}$ then $\lambda_n(x) = 0 \iff a_n = 0 \iff a_0 = \ldots = a_n = 0 \iff v_{\hat{A}}(x) \geq n+1$. Then using that $\hat{A}$ is a DVR with unique maximal ideal $\hat{\mathfrak{m}} := (\hat{\pi})$ this is equivalent to $x \in \hat{\mathfrak{m}}^{n+1}$ (3.31.3) giving e); of which c) is a special case. ∎

**Lemma 3.34.9 (Local is Continuous, DVR Case)**
*Let $(A, \mathfrak{m})$, $(B, \mathfrak{n})$ be DVRs, and $\phi : A \to B$ a ring homomorphism. Then $\phi$ is local (that is $\phi(\mathfrak{m}) \subseteq \mathfrak{n}$) if and only if $\phi$ is continuous (3.14.5) and then $\phi(\mathfrak{m}^{n+1}) \subseteq \mathfrak{n}^{n+1}$ for every $n \geq 0$ (same-index witness).*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

This is (3.14.15) with $\mathfrak{a} := \mathfrak{m}$, $\mathfrak{b} := \mathfrak{n}$ ($\mathfrak{n}$ maximal, hence radical (3.3.36)); the same-index witness $\phi(\mathfrak{m}^{n+1}) \subseteq \mathfrak{n}^{n+1}$ is exactly that lemma's forward-direction computation ($k := n+1$). ∎

**Proposition 3.34.10 (Taut Criterion, DVR Case)**
*Let $(A, \mathfrak{m})$, $(B, \mathfrak{n})$ be DVRs, $\phi : A \to B$ a local ring homomorphism, and $e$ its ramification index (3.31.10). Then $\phi$ is taut (3.14.10) if and only if $e = 1$ and $B = \phi(A) + \mathfrak{n}$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Fix a uniformiser $\pi$ of $A$, so $e = v_B(\phi(\pi))$ (3.31.10). For every $n \geq 0$: $\phi(\mathfrak{m}^{n+1}) = \phi((\pi^{n+1})) = (\phi(\pi)^{n+1})$, which is $\mathfrak{n}^{e(n+1)}$ if $e < \infty$ (as $v_B(\phi(\pi)^{n+1}) = e(n+1)$), and 0 if $e = \infty$.

**Case $e = \infty$.** For any strictly increasing $n_0 < n_1 < n_2 < \cdots$, the taut recursion at $N$ reads $\mathfrak{n}^{n_N+1} = \phi(\mathfrak{m}^{n_N+1}) + \mathfrak{n}^{n_{N+1}+1} = \mathfrak{n}^{n_{N+1}+1}$, forcing $n_N + 1 = n_{N+1} + 1$, that is $n_N = n_{N+1}$: impossible, as $n_N < n_{N+1}$. So $\phi$ is not taut.

**Case $e < \infty$.** As ideals of a DVR are totally ordered by inclusion, $\mathfrak{n}^{e(n_N+1)} + \mathfrak{n}^{n_{N+1}+1} = \mathfrak{n}^{\min(e(n_N+1), n_{N+1}+1)}$; so the taut recursion at $N$, $\mathfrak{n}^{n_N+1} = \mathfrak{n}^{\min(e(n_N+1), n_{N+1}+1)}$, holds iff $\min(e(n_N+1), n_{N+1}+1) = n_N + 1$. As $n_{N+1} > n_N$ gives $n_{N+1} + 1 > n_N + 1$ always, this holds iff $e(n_N+1) = n_N + 1$, that is (as $n_N + 1 \geq 1$) iff $e = 1$.

So if $e \neq 1$, the recursion fails at every $N$, for every choice of $(n_N)_N$: $\phi$ is not taut. If $e = 1$: $\phi(\mathfrak{m}^{n+1}) = \mathfrak{n}^{n+1}$ for every $n \geq 0$, so the recursion holds automatically, for every $N$ and every choice of $n_{N+1} > n_N$; so $\phi$ is taut iff $B = \phi(A) + \mathfrak{n}^{n_0+1}$ for some $n_0 \geq 0$ – equivalently (as $\mathfrak{n}^{n_0+1} \subseteq \mathfrak{n}$ makes this weakest, hence necessary and sufficient, at $n_0 := 0$) iff $B = \phi(A) + \mathfrak{n}$. ∎

**Lemma 3.34.11 (Functoriality of Completion, DVR Case)**
*Let $(A, \mathfrak{m})$, $(B, \mathfrak{n})$ be DVRs, with completions $\big(\hat{A}, \hat{\mathfrak{m}}\big)$, $\iota_A$, $\lambda_n^A$ and $\big(\hat{B}, \hat{\mathfrak{n}}\big)$, $\iota_B$, $\lambda_n^B$ (3.34.4), and $\phi : A \to B$ a local ring homomorphism with ramification index $e$ (3.31.10). This is (3.14.27) ($\phi$ being continuous (3.34.9)), giving: there is a unique continuous ring homomorphism $\hat{\phi} : \hat{A} \to \hat{B}$ with $\hat{\phi} \circ \iota_A = \iota_B \circ \phi$, which is moreover **local** (3.34.9).*

*Moreover this is functorial: $\widehat{\mathrm{id}_A} = \mathrm{id}_{\hat{A}}$; and, for a further DVR $(C, \mathfrak{a}_C)$ and a local ring homomorphism $\psi : B \to C$ (so $\psi \circ \phi : A \to C$ is again local), $\widehat{\psi \circ \phi} = \hat{\psi} \circ \hat{\phi} : \hat{A} \to \hat{C}$.*

*Moreover $\hat{\phi}$ has the same ramification index $e$; so by (3.31.10) a), $\hat{\phi}$ is injective if and only if $e < \infty$.*

*Moreover $\hat{\phi}$ is bijective if and only if it is surjective, if and only if $e = 1$ and $B = \phi(A) + \mathfrak{n}$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

By (3.34.9), $\phi$ is continuous; so (3.14.27) gives the stated $\hat{\phi}$ and functoriality directly.

As $\hat{A}, \hat{B}$ are DVRs (3.34.8) with maximal ideals $\hat{\mathfrak{m}}, \hat{\mathfrak{n}}$, and $\hat{\phi}$ is continuous for exactly their native filtrations (these being $\ker \lambda_\bullet^A, \ker \lambda_\bullet^B$), (3.34.9) (applied to $\hat{A}, \hat{B}, \hat{\phi}$) gives $\hat{\phi}$ local; and this is the unique local map, uniqueness among local maps coinciding with uniqueness among continuous ones (by that lemma).

© David Rufino CC BY-NC-ND 4.0

**Ramification index.** Fix a uniformiser $\pi$ of $A$, so $e = v_B(\phi(\pi))$, and $\iota_A(\pi)$ is a uniformiser of $\hat{A}$ (3.34.8) a). For $b \in B$, $b \neq 0$: $\iota_B(b)$ has $\lambda_m^B$-coordinate $b \mod \mathfrak{n}^{m+1}$ for every $m$ (3.14.16), so, by (3.34.8) b), $v_{\hat{B}}(\iota_B(b)) = \min\{m \geq 0 : b \notin \mathfrak{n}^{m+1}\} = v_B(b)$. So, by the commuting square,

$$v_{\hat{B}}\big(\hat{\phi}(\iota_A(\pi))\big) = v_{\hat{B}}(\iota_B(\phi(\pi))) = v_B(\phi(\pi)) = e$$

(both sides $\infty$ if $\phi(\pi) = 0$, by the same computation applied formally): $\hat{\phi}$'s own ramification index (with respect to the uniformiser $\iota_A(\pi)$) is $e$.

**Injectivity criterion.** This is (3.31.10) a), applied to $\hat{\phi}$ (with ramification index $e$, as above).

**Surjectivity criterion.** If $e = 1$ and $B = \phi(A) + \mathfrak{n}$: by (3.34.10), $\phi$ is taut, so (3.14.27)'s own surjectivity criterion gives $\hat{\phi}$ surjective. Conversely, suppose $\hat{\phi}$ is surjective; by (3.31.10) b) (applied to $\hat{\phi}$, ramification index $e$), $e = 1$, so $\phi(\pi)$ is a uniformiser of $B$ (3.31.2), giving $\phi(\mathfrak{m}^{n+1}) = (\phi(\pi)^{n+1}) = \mathfrak{n}^{n+1}$ for every $n \geq 0$; so (3.3.56) gives induced maps $\overline{\phi}_n : A/\mathfrak{m}^{n+1} \to B/\mathfrak{n}^{n+1}$ for every $n \geq 0$, and $\hat{\phi} = \varprojlim \overline{\phi}_n$ (3.3.156) (taking $n(m) := m$ throughout), that is $\lambda_n^B \circ \hat{\phi} = \overline{\phi}_n \circ \lambda_n^A$ for every $n \geq 0$. As $\lambda_n^A, \lambda_n^B$ are surjective (3.14.24) and $\hat{\phi}$ is surjective, $\overline{\phi}_n \circ \lambda_n^A$ is surjective, hence so is $\overline{\phi}_n$ (its image already equals that of $\overline{\phi}_n \circ \lambda_n^A$, $\lambda_n^A$ being surjective); taking $n := 0$, $\overline{\phi}_0$ surjective is exactly $B = \phi(A) + \mathfrak{n}$. Finally, bijective trivially gives surjective, and conversely (3.31.10) b) (applied to $\hat{\phi}$) already gives that $\hat{\phi}$ surjective implies $\hat{\phi}$ bijective: so $\hat{\phi}$ is bijective if and only if it is surjective. ∎

**Proposition 3.34.12 (Universal Property of Completion, DVR Case)**
*Let $(A, \mathfrak{m})$ be a DVR and $\big(\hat{A}, \hat{\mathfrak{m}}\big)$ its completion (3.34.4), $(B, \mathfrak{n})$ a complete DVR, with $\iota_B : B \xrightarrow{\sim} \hat{B}$ its own completion (3.34.5), and $\phi : A \to B$ a local homomorphism with ramification index $e$ (3.31.10). This is (3.14.28) ($\phi$ being continuous (3.34.9)), giving: there is a unique **local** ring homomorphism $\hat{\phi} : \hat{A} \to B$ such that $\hat{\phi} \circ \iota = \phi$ (local and continuous being equivalent here (3.34.9)).*

*Moreover $\hat{\phi}$ is injective if and only if $e < \infty$.*

*Further $\hat{\phi}$ is bijective if and only if it is surjective, if and only if $e = 1$ and $B = \phi(A) + \mathfrak{n}$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

By (3.34.9), $\phi$ is continuous, with $\phi(\mathfrak{m}^{n+1}) \subseteq \mathfrak{n}^{n+1}$ for every $n \geq 0$; and $B$ is complete ((3.14.20), $\iota_B$ being an isomorphism). So (3.14.28) gives the stated $\hat{\phi}$ (for the filtration $\ker \lambda_\bullet = \hat{\mathfrak{m}}^{\bullet+1}$ of $\hat{A}$, (3.34.8) e)), namely $\hat{\phi} = \iota_B^{-1} \circ \Psi$ where $\Psi : \hat{A} \to \hat{B}$ is the unique continuous ring homomorphism with $\Psi \circ \iota = \iota_B \circ \phi$ – exactly (3.34.11)'s own $\hat{\phi}$ for $\phi$. As $\iota_B$ is an isomorphism, $\hat{\phi}$ is injective (resp. surjective) if and only if $\Psi$ is.

As $\hat{A}, B$ are DVRs (3.34.8) (applied to $A$), with maximal ideals $\hat{\mathfrak{m}}, \mathfrak{n}$, and $\hat{\phi}$ is continuous for exactly their native filtrations $\hat{\mathfrak{m}}^{\bullet+1}$, $\mathfrak{n}^{\bullet+1}$, (3.34.9) (applied to $\hat{A}, B, \hat{\phi}$) gives $\hat{\phi}$ local; and this is the unique local map making the triangle commute, uniqueness among local maps coinciding with uniqueness among continuous ones (3.34.9).

**Injectivity criterion.** By (3.34.11)'s own injectivity criterion, $\Psi$ is injective if and only if $e < \infty$; so, by the above, so is $\hat{\phi}$.

**Surjectivity criterion.** By (3.34.11)'s own surjectivity criterion, $\Psi$ is bijective if and only if it is surjective, if and only if $e = 1$ and $B = \phi(A) + \mathfrak{n}$; so, by the above ($\hat{\phi} = \iota_B^{-1} \circ \Psi$, an isomorphism composed with $\Psi$), the same holds for $\hat{\phi}$. ∎

**Proposition 3.34.13 (The Completion Matches the Filtration)**
*Let $(A, \mathfrak{m})$ be a DVR and $\big(\hat{A}, \hat{\mathfrak{m}}\big)$ its completion (3.34.8). There is a unique ring homomorphism $\nu_n : A/\mathfrak{m}^{n+1} \to \hat{A}/\hat{\mathfrak{m}}^{n+1}$ such that the square*

$$\begin{array}{ccc} A & \xrightarrow{\ \iota\ } & \hat{A} \\ {\scriptstyle \overline{\pi}_n}\downarrow & & \downarrow{\scriptstyle \pi_n} \\ A/\mathfrak{m}^{n+1} & \xrightarrow[\ \nu_n\ ]{} & \hat{A}/\hat{\mathfrak{m}}^{n+1} \end{array}$$

*commutes. Further $\nu_n$ is an isomorphism, with inverse $\mu_n := \nu_n^{-1}$. Moreover, writing $\rho_n : A/\mathfrak{m}^{n+1} \to A/\mathfrak{m}^n$ and $\hat{\rho}_n : \hat{A}/\hat{\mathfrak{m}}^{n+1} \to \hat{A}/\hat{\mathfrak{m}}^n$ for the natural reductions (3.3.56) $(\nu_n)_{n\geq 0}$ is natural in $n$: for every $n \geq 1$, the square*

© David Rufino CC BY-NC-ND 4.0

$$\begin{array}{ccc} A/\mathfrak{m}^{n+1} & \xrightarrow[\sim]{\nu_n} & \hat{A}/\hat{\mathfrak{m}}^{n+1} \\ \rho_n \downarrow & & \downarrow \hat{\rho}_n \\ A/\mathfrak{m}^{n} & \xrightarrow[\sim]{\nu_{n-1}} & \hat{A}/\hat{\mathfrak{m}}^{n} \end{array}$$

*commutes, that is* $\hat{\rho}_n \circ \nu_n = \nu_{n-1} \circ \rho_n$ *(equivalently,* $\rho_n \circ \mu_n = \mu_{n-1} \circ \hat{\rho}_n$*).*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Writing $\lambda_n : \hat{A} \to A/\mathfrak{m}^{n+1}$ for the projections (3.34.4): $\ker \lambda_n = \hat{\mathfrak{m}}^{n+1}$ (3.34.8) e), so the statement is exactly (3.14.30) (with $\mathfrak{a} := \mathfrak{m}$), applied using this equality in place of the general inclusion $\mathfrak{m}^{n+1}\hat{A} \subseteq \ker \lambda_n$. ∎

**Corollary 3.34.14 (The Completion is Complete)**
*Let* $(A, \mathfrak{m})$ *be a DVR, and* $\hat{A}$ *as in (3.34.4). Then* $\hat{A}$ *is complete (3.34.5).*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

By (3.14.26) (applied to $(A, \mathfrak{a}_\bullet)$, $\mathfrak{a}_n := \mathfrak{m}^{n+1}$ (3.14.19)), $\left(\hat{A}, \ker \lambda_\bullet\right)$ is complete (3.14.20) as $\ker \lambda_n = \hat{\mathfrak{m}}^{n+1}$ for every $n \geq 0$ (3.34.8) e), this is exactly completeness of $\hat{A}$ in the sense of (3.34.5). ∎

**Lemma 3.34.15 (Completeness of the Fraction Field)**
*Let* $(A, \mathfrak{m})$ *be a complete DVR,* $\pi$ *a uniformiser, and* $K := \mathrm{Frac}(A)$*. Then* $(\pi^n A)_{n\in\mathbb{Z}}$ *is a filtration of* $K$ *as an* $A$*-module (3.14.2) (each* $\pi^n A$ *free of rank* 1*), with* $K = \bigcup_n \pi^n A$*; and the canonical map*

$$\jmath : K \to \varprojlim_{n\geq 0} K/\pi^n A, \quad x \mapsto (x \mod \pi^n A)_{n\geq 0},$$

*(3.14.17) is bijective.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

**Filtration, exhaustiveness.** $\pi^{n+1}A \subseteq \pi^n A$ for every $n \in \mathbb{Z}$ (as $\pi^{n+1} = \pi \cdot \pi^n$), and $x \mapsto \pi^n x$ is an $A$-module isomorphism $A \xrightarrow{\sim} \pi^n A$: a filtration of $K$ by finite free $A$-submodules (3.14.2). Every nonzero $x \in K$ is $x = u\pi^{v(x)}$ for a unit $u \in A^\star$ and its valuation $v(x) \in \mathbb{Z}$ (3.31.2), so $x \in \pi^{v(x)}A$: $K = \bigcup_n \pi^n A$.

**Surjectivity of $\jmath$.** Let $(\overline{x}_n)_{n\geq 0}$, $\overline{x}_n \in K/\pi^n A$, be compatible under reduction. Pick a representative $x_0 \in K$ of $\overline{x}_0$ and, by exhaustiveness, $N \geq 1$ with $x_0 \in \pi^{-N}A$. Choosing representatives $x_n \in K$ of $\overline{x}_n$ $(n \geq 0)$ inductively with $x_n - x_{n-1} \in \pi^{n-1}A$ for $n \geq 1$ (possible by compatibility): as $\pi^{n-1}A \subseteq \pi^0 A \subseteq \pi^{-N}A$ for every $n \geq 1$, every $x_n$ lies in $M := \pi^{-N}A$.

As $\mathfrak{m}^{n+N}M = \pi^{n+N}A \cdot \pi^{-N}A = \pi^n A$ for every $n \geq 0$, the sequence $(x_n \mod \pi^n A)_{n\geq 0}$, viewed inside $M$, matches $(x_n \mod \mathfrak{m}^{n+N}M)_{n\geq 0}$. Extend it to a compatible sequence $(\overline{y}_j)_{j\geq 1}$, $\overline{y}_j \in M/\mathfrak{m}^j M$, by setting $\overline{y}_{n+N} := x_n \mod \mathfrak{m}^{n+N}M$ for $n \geq 0$ (compatible among themselves, as the $x_n$ already are) and, for $1 \leq j < N$, $\overline{y}_j$ the image of $\overline{y}_N = x_0 \mod \mathfrak{m}^N M$ under the reduction map $M/\mathfrak{m}^N M \to M/\mathfrak{m}^j M$ (so compatible with the rest too). By (3.34.7) ($A$ complete, $M$ free of rank 1), $M \to \varprojlim_{j\geq 1} M/\mathfrak{m}^j M$ is bijective, so there is $x \in M$ with $x \equiv \overline{y}_j \pmod{\mathfrak{m}^j M}$ for every $j \geq 1$; in particular, at $j = n + N$ $(n \geq 0)$, $x \equiv x_n \equiv \overline{x}_n \pmod{\pi^n A}$. So $\jmath(x) = (\overline{x}_n)_n$: $\jmath$ is surjective.

**Bijectivity.** $\bigcap_n \pi^n A = \bigcap_{n\geq 0} \mathfrak{m}^n = 0$ (3.31.9), so $(K, \pi^\bullet A)$ is separated; by (3.14.20), surjectivity of $\jmath$ then gives bijectivity. ∎

**Proposition 3.34.16 (Completion of the Fraction Field)**
*Let* $(A, \mathfrak{m})$ *be a DVR,* $\pi$ *a uniformiser,* $K := \mathrm{Frac}(A)$*, and* $\hat{A}, \iota, \lambda_\bullet$ *its completion (3.34.8),* $K_{\mathfrak{m}} := \mathrm{Frac}\left(\hat{A}\right)$*. Write* $\hat{K}$*,* $\jmath$ *for the completion of* $K$ *as a filtered* $A$*-module for* $(\pi^n A)_{n\in\mathbb{Z}}$ *(3.14.17) (3.34.15). Then* $\iota$ *extends (uniquely, (3.6.5)) to an injective ring homomorphism* $\iota_K : K \to K_{\mathfrak{m}}$*, and there is a unique ring structure on* $\hat{K}$ *for which the canonical bijection* $\Psi : \hat{K} \xrightarrow{\sim} K_{\mathfrak{m}}$ *(constructed below) is a ring isomorphism, with* $\Psi \circ \jmath = \iota_K$*.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

As $\bigcap_n \mathfrak{m}^n = 0$ (3.31.9), $\iota$ is injective, so (3.6.5) extends it to an injective ring homomorphism $\iota_K : K \to K_{\mathfrak{m}}$, $\iota_K(a/s) = \iota(a)/\iota(s)$. Write $\hat{\pi} := \iota(\pi)$, a uniformiser of $\hat{A}$ (3.34.8).

**Exactness.** For $a \in A$, $j \geq 0$: $\ker \lambda_{j-1} = \hat{\mathfrak{m}}^j = \hat{\pi}^j \hat{A}$ (3.34.8) (reading $\ker \lambda_{-1} := \hat{A}$ for $j = 0$, trivially true), and $\lambda_{j-1} \circ \iota$ is the reduction map $A \to A/\mathfrak{m}^j$, so $\iota(a) \in \hat{\pi}^j \hat{A} \iff a \in \mathfrak{m}^j = \pi^j A$. For general $a/s \in K$ $(a, s \in A,\ s \neq 0)$, write $s = u\pi^{v(s)}$ for a unit $u \in A^\star$ (so $\iota(u) \in \hat{A}^\star$): fixing $n \geq 0$, $\hat{\pi}^n \iota(s)\hat{A} = \hat{\pi}^{n+v(s)}\hat{A}$ ($\iota(u)$ a unit), so

© David Rufino CC BY-NC-ND 4.0

$$\iota_K(a/s) \in \hat{\pi}^n \hat{A} \iff \iota(a) \in \hat{\pi}^{n+v(s)} \hat{A} \iff a \in \pi^{n+v(s)} A \iff a/s \in \pi^n A.$$

So $\iota_K^{-1}\left(\hat{\pi}^n \hat{A}\right) = \pi^n A$ for every $n \geq 0$.

**Density.** Given $y \in K_{\mathfrak{m}}$, $n \geq 0$: $y \in \hat{\pi}^{-M} \hat{A}$ for some $M \geq 0$ (every nonzero element of $K_{\mathfrak{m}}$ has a valuation, (3.34.8)), say $y = \hat{\pi}^{-M} \hat{a}$, $\hat{a} \in \hat{A}$; as $\lambda_{n+M} \circ \iota$ is the (surjective) reduction map $A \to A/\mathfrak{m}^{n+M+1}$, $\hat{a} = \iota(a) + c$ for some $a \in A$, $c \in \ker \lambda_{n+M} = \hat{\mathfrak{m}}^{n+M+1}$; so $y = \iota_K(\pi^{-M} a) + \hat{\pi}^{-M} c$, with $\hat{\pi}^{-M} c \in \hat{\pi}^{-M} \hat{\mathfrak{m}}^{n+M+1} = \hat{\pi}^{n+1} \hat{A} \subseteq \hat{\pi}^n \hat{A}$. So $K_{\mathfrak{m}} = \iota_K(K) + \hat{\pi}^n \hat{A}$.

**The bijection $\Psi$.** By Exactness and Density, $\iota_K$ induces, for every $n \geq 0$, a bijection $K/\pi^n A \xrightarrow{\sim} K_{\mathfrak{m}}/\hat{\pi}^n \hat{A}$, compatible with reduction (both induced by $\iota_K$); assembling over $n$, $\hat{K} = \varprojlim_n (K/\pi^n A) \xrightarrow{\sim} \varprojlim_n \left(K_{\mathfrak{m}}/\hat{\pi}^n \hat{A}\right)$. As $\hat{A}$ is complete (3.34.14), (3.34.15) (applied to the complete DVR $\hat{A}$, uniformiser $\hat{\pi}$) gives that $K_{\mathfrak{m}} \to \varprojlim_n \left(K_{\mathfrak{m}}/\hat{\pi}^n \hat{A}\right)$ is bijective; composing with its inverse gives $\Psi : \hat{K} \xrightarrow{\sim} K_{\mathfrak{m}}$. Unwinding the construction on $\jmath(x)$ for $x \in K$ gives $\Psi(\jmath(x)) = \iota_K(x)$: $\Psi \circ \jmath = \iota_K$.

**Transport.** Define $x \cdot_{\hat{K}} y := \Psi^{-1}(\Psi(x)\Psi(y))$ for $x, y \in \hat{K}$: this makes $\hat{K}$ a ring and $\Psi$ a ring isomorphism, by construction. As $\Psi \circ \jmath = \iota_K$ is then a ring homomorphism and $\Psi$ a ring isomorphism, $\jmath = \Psi^{-1} \circ \iota_K$ is a ring homomorphism too. Uniqueness of a ring structure on $\hat{K}$ making $\Psi$ an isomorphism is immediate, a ring structure transported along a bijection being unique. ∎

**Remark 3.34.17**
*So, up to the ring isomorphism $\Psi$, the completion of $K = \mathrm{Frac}(A)$ as a filtered $A$-module for the fractional-ideal filtration $(\pi^n A)_{n \in \mathbb{Z}}$ – **not** the filtration induced from $A$ (3.14.2), which is degenerate ($\mathfrak{a}_n K = K$ already for $n \geq 0$, $K$ being a field) – recovers exactly $K_{\mathfrak{m}} = \mathrm{Frac}\left(\hat{A}\right)$, with $\jmath$ corresponding to the canonical field extension $\iota_K$.*

**Corollary 3.34.18 (The Completion via any Exponent)**
*Let $(A, \mathfrak{m})$ be a DVR, and $\hat{A}$ as in (3.34.4). For every integer $e \geq 1$ there is a canonical isomorphism $\theta_{\mathfrak{m},e} : \hat{A} \xrightarrow{\sim} \varprojlim_N A/\mathfrak{m}^{Ne}$, $(\overline{a}_n)_{n \geq 0} \mapsto (\overline{a}_{Ne-1})_{N \geq 1}$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

This is (3.14.31) with $\mathfrak{a} := \mathfrak{m}$, noting $\hat{A}_{\mathfrak{m}} = \hat{A}$. ∎

**Proposition 3.34.19 (Uniqueness of the Prime Above a Complete Base)**
*Let $(\mathcal{O}_K, \mathfrak{m})$ be a complete DVR, $K := \mathrm{Frac}(\mathcal{O}_K)$, $L/K$ a finite separable extension, and $\mathcal{O}_L$ the integral closure of $\mathcal{O}_K$ in $L$. Then $\mathcal{O}_L$ has a unique prime over $\mathfrak{m}$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

As $\mathcal{O}_K$ is a PID and $L/K$ is finite separable, (3.32.9) gives that $\mathcal{O}_L$ is a Dedekind domain, finite free of rank $n := [L : K]$ over $\mathcal{O}_K$. Write $\mathfrak{q}_1, \ldots, \mathfrak{q}_s$ for the primes of $\mathcal{O}_L$ over $\mathfrak{m}$ (there is at least one, by Lying Over (3.24.17), and – $\mathcal{O}_L$ being Dedekind and integral over the DVR $\mathcal{O}_K$ – these are exactly the maximal ideals of $\mathcal{O}_L$), with ramification indices $e_l$; by the Ramification Formula (3.32.16), $\mathfrak{m}\mathcal{O}_L = \prod_l \mathfrak{q}_l^{e_l}$.

As $\mathcal{O}_K$ is complete and $\mathcal{O}_L$ is finite free over it, (3.34.7) gives that $\mathcal{O}_L \to \varprojlim_N \mathcal{O}_L/\mathfrak{m}^N \mathcal{O}_L$ is bijective: $\mathcal{O}_L$ is $\mathfrak{m}\mathcal{O}_L$-adically complete (3.14.21).

The ideals $\mathfrak{q}_l^{e_l}$ ($l = 1, \ldots, s$) are pairwise comaximal, so the Chinese Remainder Theorem (3.3.63) gives a ring isomorphism $\mathcal{O}_L/\mathfrak{m}\mathcal{O}_L \xrightarrow{\sim} \prod_{l=1}^s \mathcal{O}_L/\mathfrak{q}_l^{e_l}$. If $s \geq 2$, the right side has a nontrivial idempotent (e.g. $(1, 0, \ldots, 0)$, not 0 or 1 as $s \geq 2$); by (3.14.36) (applied to $\mathcal{O}_L$, $\mathfrak{m}\mathcal{O}_L$, complete above), this lifts to an idempotent $e \in \mathcal{O}_L$ with $e \not\equiv 0, 1 \pmod{\mathfrak{m}\mathcal{O}_L}$, so $e \neq 0, 1$ in $\mathcal{O}_L$. But $\mathcal{O}_L$, a subring of the field $L$, is a domain, so $e^2 = e$ forces $e(e-1) = 0$, hence $e = 0$ or $e = 1$: a contradiction. So $s = 1$. ∎

**Corollary 3.34.20 (Uniqueness of the Extended Valuation)**
*Let $(\mathcal{O}_K, \mathfrak{m})$ be a complete DVR, $K := \mathrm{Frac}(\mathcal{O}_K)$, $k := \mathcal{O}_K/\mathfrak{m}$, and $L/K$ a finite separable extension of degree $n$. Let $\mathcal{O}_L$ be the integral closure of $\mathcal{O}_K$ in $L$. Then $\mathcal{O}_L$ is itself a **complete** DVR, with a unique prime $\mathfrak{q}$ lying above $\mathfrak{m}$, of ramification index $e$ and residue degree $f$, with $ef = n$.*

*If moreover $f = 1$ (so $L/K$ is **totally ramified**: $e = n$), then $\mathcal{O}_L = \mathcal{O}_K[\varpi]$ for any uniformiser $\varpi$ of $\mathcal{O}_L$ (3.32.17) and, if $\mathcal{O}_K \cong k[[t]]$ for a uniformiser $t$, then $\mathcal{O}_L \cong k[[\varpi]]$ and $L \cong k((\varpi))$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

By (3.34.19), $\mathcal{O}_L$ has a unique prime $\mathfrak{q}$ over $\mathfrak{m}$; writing $e, f$ for its ramification index and residue degree, the Ramification Formula (3.32.16) gives $ef = n$, and $\mathcal{O}_L$, being Dedekind and local (its only nonzero prime being $\mathfrak{q}$), is itself a DVR (3.31.2) g) – that is, a place of $L$, the unique one over $\mathfrak{m}$.

© David Rufino CC BY-NC-ND 4.0

**$\mathcal{O}_L$ is complete.** As $\mathcal{O}_K$ is complete and $\mathcal{O}_L$ is finite free of rank $n$ over it (3.32.9), (3.34.7) gives that $\mathcal{O}_L \to \varprojlim_N \mathcal{O}_L/\mathfrak{m}^N\mathcal{O}_L$ is bijective. As $\mathfrak{m}\mathcal{O}_L = \mathfrak{q}^e$ (the unique prime, Ramification Formula), $\mathfrak{m}^N\mathcal{O}_L = \mathfrak{q}^{Ne}$ for every $N \geq 1$; so this says exactly that every sequence compatible under reduction at the levels $\mathfrak{q}^{Ne} = \mathfrak{q}^{n_N+1}$ ($n_N := Ne-1$, $N \geq 1$) arises from an element of $\mathcal{O}_L$. By (3.34.5) (applied to the DVR $\mathcal{O}_L$, this cofinal sequence, c) $\iff$ a)), the same holds for $\mathcal{O}_L$'s own $\mathfrak{q}$-adic tower: $\mathcal{O}_L$ is complete.

**If moreover $f = 1$:** $e = n$ by $ef = n$. This is exactly the hypothesis of (3.32.17) (a unique prime of $\mathcal{O}_L$ over $\mathfrak{m}$, of residue degree 1), giving $\mathcal{O}_L = \mathcal{O}_K[\varpi]$ for any uniformiser $\varpi$. Finally, if $\mathcal{O}_K \cong k[[t]]$: $\mathcal{O}_L/\mathfrak{q}$ is a degree-$f = 1$ extension of $k$, hence equal to $k$ (via the embedding $k \hookrightarrow \mathcal{O}_K \hookrightarrow \mathcal{O}_L \twoheadrightarrow \mathcal{O}_L/\mathfrak{q}$, an isomorphism as $f = 1$): $k \subseteq \mathcal{O}_L$ is a coefficient field. As $\mathcal{O}_L$ is already complete (above), (3.34.3) (applied to $\mathcal{O}_L$, $\varpi$, $k$) gives $\mathcal{O}_L \cong k[[\varpi]]$, and $L \cong k((\varpi))$. ∎

**Proposition 3.34.21 (Splitting via a Primitive Element)**
*Let $(\mathcal{O}_K, \mathfrak{m})$ be a DVR, $K := \mathrm{Frac}(\mathcal{O}_K)$, $L/K$ a finite separable extension of degree $n$, $\mathcal{O}_L$ the integral closure of $\mathcal{O}_K$ in $L$, with primes $\mathfrak{q}_1, \ldots, \mathfrak{q}_s$ over $\mathfrak{m}$ (Lying Over (3.24.17) these are exactly the maximal ideals of the Dedekind domain $\mathcal{O}_L$ (3.32.9), it being integral over the DVR $\mathcal{O}_K$). Write $\hat{\mathcal{O}}_K, \iota$ for the completion of $\mathcal{O}_K$ (3.34.8), $K_{\mathfrak{m}} := \mathrm{Frac}\big(\hat{\mathcal{O}}_K\big)$, and $\iota_K : K \to K_{\mathfrak{m}}$ for the extension of $\iota$ (3.6.5) ($\iota$ injective, (3.31.9)); similarly, for each $l$, write $\hat{\mathcal{O}}_{L,\mathfrak{q}_l} := \varprojlim_n \mathcal{O}_L/\mathfrak{q}_l^n$, $\iota_l$, for the completion of $\mathcal{O}_L$ at $\mathfrak{q}_l$ (3.14.16), $L_{\mathfrak{q}_l} := \mathrm{Frac}\big(\hat{\mathcal{O}}_{L,\mathfrak{q}_l}\big)$, and $\iota_{L,l} : L \to L_{\mathfrak{q}_l}$ for the extension of $\iota_l$ ($\iota_l$ injective, (3.31.9), applied to the DVR $(\mathcal{O}_L)_{\mathfrak{q}_l}$, using $\mathcal{O}_L \cap \mathfrak{q}_l^n(\mathcal{O}_L)_{\mathfrak{q}_l} = \mathfrak{q}_l^n$).*

*For each $l$, there is moreover a unique ring homomorphism $\hat{\phi}_l : \hat{\mathcal{O}}_K \to \hat{\mathcal{O}}_{L,\mathfrak{q}_l}$ with $\hat{\phi}_l \circ \iota = \iota_l|_{\mathcal{O}_K}$; it is injective, and extends (uniquely, still denoted $\hat{\phi}_l$) to an injective ring homomorphism $K_{\mathfrak{m}} \to L_{\mathfrak{q}_l}$, with $\hat{\phi}_l \circ \iota_K = \iota_{L,l}|_K$. These fit into a commutative cube*




*(the bottom face the rings, the top face their fraction fields, the depth edges the completion maps $\iota$, $\iota_l$, and the vertical edges the canonical inclusions of a domain into its fraction field).*

*Then there is a canonical isomorphism of $K_{\mathfrak{m}}$-algebras*

$$\Phi : L \otimes_K K_{\mathfrak{m}} \xrightarrow{\sim} \prod_{l=1}^{s} L_{\mathfrak{q}_l}, \quad x \otimes y \mapsto \big(\iota_{L,l}(x)\hat{\phi}_l(y)\big)_{l=1}^{s}.$$

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

**Structure maps $\hat{\phi}_l$, and the cube.** As $\mathfrak{q}_l$ lies over $\mathfrak{m}$, $\iota_l(\mathfrak{m}\mathcal{O}_L) \subseteq \iota_l(\mathfrak{q}_l) \subseteq \hat{\mathfrak{q}}_l$ (continuity of $\iota_l$ (3.14.16)): $\iota_l|_{\mathcal{O}_K} : \mathcal{O}_K \to \hat{\mathcal{O}}_{L,\mathfrak{q}_l}$ is local. As $\hat{\mathcal{O}}_{L,\mathfrak{q}_l}$ is a complete DVR, (3.34.12) (applied to $\phi := \iota_l|_{\mathcal{O}_K}$) gives a unique continuous ring homomorphism $\hat{\phi}_l : \hat{\mathcal{O}}_K \to \hat{\mathcal{O}}_{L,\mathfrak{q}_l}$ with $\hat{\phi}_l \circ \iota = \iota_l|_{\mathcal{O}_K}$; it is injective (ramification index $v_{\hat{\mathcal{O}}_{L,\mathfrak{q}_l}}(\iota_l(\pi))$, for a uniformiser $\pi$ of $\mathcal{O}_K$, finite as $\iota_l(\pi) \neq 0$), so (3.6.5) extends it to an injective ring homomorphism $\hat{\phi}_l : K_{\mathfrak{m}} \to L_{\mathfrak{q}_l}$; as $\hat{\phi}_l \circ \iota_K$ and $\iota_{L,l}|_K$ agree on $\mathcal{O}_K$ (both restricting to $\iota_l|_{\mathcal{O}_K}$) and $K = \mathrm{Frac}(\mathcal{O}_K)$, the uniqueness clause of (3.6.5) gives $\hat{\phi}_l \circ \iota_K = \iota_{L,l}|_K$ on all of $K$.

The bottom face of the cube commutes by $\hat{\phi}_l$'s defining property. The left and right faces commute because each diagonal edge ($K \to K_{\mathfrak{m}}$, $L \to L_{\mathfrak{q}_l}$) is, by construction, the extension along (3.6.5) of the corresponding ring-level map ($\mathcal{O}_K \to \hat{\mathcal{O}}_K$, $\mathcal{O}_L \to \hat{\mathcal{O}}_{L,\mathfrak{q}_l}$). The front face ($\mathcal{O}_K \to \mathcal{O}_L \to L$ versus $\mathcal{O}_K \to K \to L$) is immediate, both being the inclusion $\mathcal{O}_K \to L$. The top face is exactly $\hat{\phi}_l \circ \iota_K = \iota_{L,l}|_K$, just shown.

**Primitive element.** Fix a primitive element $\theta \in \mathcal{O}_L$ for $L/K$ (3.19.105), with minimal polynomial $g \in \mathcal{O}_K[Y]$ (coefficients integral, as $\mathcal{O}_K$ is integrally closed and $\theta$ integral over it), monic of degree $n$, separable. Viewing $g \in K_{\mathfrak{m}}[Y]$ via $\iota_K$ (still denoted $g$): as $K_{\mathfrak{m}}[Y]$ is a UFD, $g = g_1 \cdots g_r$ for monic irreducible $g_j \in K_{\mathfrak{m}}[Y]$; as $g$ is separable, so is each $g_j$, and the $g_j$ are pairwise coprime (distinct irreducibles in a PID). As $L \cong K[Y]/(g)$ via $\theta \mapsto Y$ (3.19.40),

© David Rufino CC BY-NC-ND 4.0

$$L \otimes_K K_{\mathfrak{m}} \cong K_{\mathfrak{m}}[Y]/(g) \cong \prod_{j=1}^{r} K_{\mathfrak{m}}[Y]/(g_j) =: \prod_{j=1}^{r} E_j$$

(the first isomorphism by (3.4.42), the second the Chinese Remainder Theorem (3.3.63) for the pairwise coprime $(g_j)$). Let $\iota_j : L \to E_j$ be reduction modulo $g_j$ ($\theta \mapsto Y \mod g_j$, well-defined as $g_j \mid g$), a $K$-algebra map, injective as $L$ is a field; $E_j$ is a field of degree $d_j := \deg g_j$ over $K_{\mathfrak{m}}$, with $\sum_j d_j = n$, and $\iota_j(\theta)$ a root of $g_j$.

**Each factor is a completion.** Write $\mathcal{O}_{E_j}$ for the integral closure of $\hat{\mathcal{O}}_K$ in $E_j$: as $\hat{\mathcal{O}}_K$ is complete and $E_j/K_{\mathfrak{m}}$ is finite separable, (3.34.20) gives that $\mathcal{O}_{E_j}$ is itself a complete DVR, with a unique prime over $\hat{\mathfrak{m}}$. As $E_j = K_{\mathfrak{m}}(\iota_j(\theta))$ is the $K_{\mathfrak{m}}$-span of $1, \iota_j(\theta), \dots, \iota_j(\theta)^{d_j-1}$, and $\iota_j(L) = \iota_j(K)[\iota_j(\theta)]$ the corresponding $\iota_j(K)$-span, while $\iota_j(K)$ is dense in $K_{\mathfrak{m}}$ ($\mathcal{O}_K$ dense in $\hat{\mathcal{O}}_K$ by construction of the completion (3.14.16), and $K, K_{\mathfrak{m}}$ their fraction fields), $\iota_j(L)$ is dense in $E_j$ (coefficientwise, $E_j$ finite-dimensional over $K_{\mathfrak{m}}$). So $\left(\mathcal{O}_{E_j}, \iota_j|_{\mathcal{O}_L}\right)$ is, up to this identification, the completion of $\mathcal{O}_L$ at the prime obtained by pulling back $\mathcal{O}_{E_j}$'s valuation along $\iota_j$: there is a (unique) field isomorphism $\Theta_j : E_j \xrightarrow{\sim} L_{\mathfrak{q}_{l(j)}}$, for some $l(j) \in \{1, \dots, s\}$, with $\Theta_j \circ \iota_j = \iota_{L,l(j)}$ and $\Theta_j|_{K_{\mathfrak{m}}} = \hat{\phi}_{l(j)}$ (both structure maps agreeing, by uniqueness of the extended valuation).

**Distinctness.** $j \neq j' \implies l(j) \neq l(j')$: equal primes would force $E_j \cong L_{\mathfrak{q}_{l(j)}} = L_{\mathfrak{q}_{l(j')}} \cong E_{j'}$ compatibly with $\iota_j, \iota_{j'}$, hence equal minimal polynomials $g_j = g_{j'}$ for $\iota_j(\theta), \iota_{j'}(\theta)$ over $K_{\mathfrak{m}}$ – contradicting $j \neq j'$ (distinct factors of $g$).

**Exhaustiveness, without a degree count.** For each $l \in \{1, \dots, s\}$: as $g$ has $\mathcal{O}_K$-coefficients and $\iota_{L,l}$ is a $K$-algebra homomorphism, $g\big(\iota_{L,l}(\theta)\big) = \iota_{L,l}(g(\theta)) = 0$; as $\hat{\phi}_l \circ \iota_K = \iota_{L,l}|_K$ (Structure maps, above), this is the same as evaluating the factored $g = g_1 \cdots g_r \in K_{\mathfrak{m}}[Y]$ at $\iota_{L,l}(\theta)$ via $\hat{\phi}_l$, giving $0 = \prod_j g_j\big(\iota_{L,l}(\theta)\big)$ (coefficients of each $g_j$ pushed via $\hat{\phi}_l$), a product of elements of the field $L_{\mathfrak{q}_l}$; so $g_j\big(\iota_{L,l}(\theta)\big) = 0$ for some $j$, necessarily unique (for $j \neq j'$, a Bézout relation $1 = ag_j + bg_{j'}$ in $K_{\mathfrak{m}}[Y]$, pushed via $\hat{\phi}_l$ and evaluated at $\iota_{L,l}(\theta)$, rules out both vanishing at once) – defining $j(l)$.

We check $j(l(j)) = j$ for every $j$: by construction of $\Theta_j$, $\Theta_j\big(\iota_j(\theta)\big) = \iota_{L,l(j)}(\theta)$ and $\Theta_j|_{K_{\mathfrak{m}}} = \hat{\phi}_{l(j)}$, so applying $\Theta_j$ to $g_j\big(\iota_j(\theta)\big) = 0$ (coefficients via $K_{\mathfrak{m}} \hookrightarrow E_j$) gives $g_j\Big(\iota_{L,l(j)}(\theta)\Big) = 0$ (coefficients via $\hat{\phi}_{l(j)}$); by uniqueness in the definition of $j(\cdot)$, $j(l(j)) = j$. Symmetrically (the roles of $\big(E_j, \iota_j, \hat{\phi}_{l(j)}\big)$ and $\big(L_{\mathfrak{q}_l}, \iota_{L,l}, \hat{\phi}_l\big)$ being interchangeable in this argument, both exhibiting $\theta$'s image as a root of the matching factor of $g$), $l(j(l)) = l$ for every $l$.

So $j \mapsto l(j)$ and $l \mapsto j(l)$ are mutually inverse bijections between $\{1, \dots, r\}$ and $\{1, \dots, s\}$; in particular $r = s$, and relabelling by $l(j)$,

$$\Phi : L \otimes_K K_{\mathfrak{m}} \xrightarrow{\sim} \prod_{j=1}^{r} E_j \xrightarrow[\prod_j \Theta_j]{\sim} \prod_{l=1}^{s} L_{\mathfrak{q}_l}$$

is an isomorphism of $K_{\mathfrak{m}}$-algebras, with $\Phi(x \otimes 1)$ having $l$-th coordinate $\Theta_{j(l)}\Big(\iota_{j(l)}(x)\Big) = \iota_{L,l(j(l))}(x) = \iota_{L,l}(x)$ (using $\Theta_{j(l)} \circ \iota_{j(l)} = \iota_{L,l(j(l))}$, above, and $l(j(l)) = l$, just shown) for every $x \in L$. As $\Phi$ is $K_{\mathfrak{m}}$-linear and $\Phi(1 \otimes y) = \Big(\hat{\phi}_l(y)\Big)_l$ for $y \in K_{\mathfrak{m}}$ (the same computation, with $\theta$ replaced by the trivial case $x := 1 \in L$, so $\iota_{L,l}(1) = 1$, composed with multiplication by $y$ – equivalently, $1 \otimes y = y \cdot (1 \otimes 1)$ and $\Phi(1 \otimes 1) = (1)_l$), $\Phi(x \otimes y) = \Phi((x \otimes 1)(1 \otimes y)) = \Phi(x \otimes 1)\Phi(1 \otimes y) = \Big(\iota_{L,l}(x)\hat{\phi}_l(y)\Big)_l$ for every $x \in L$, $y \in K_{\mathfrak{m}}$: the stated formula. ∎

**Proposition 3.34.22 (Coefficient Expansion of a DVR with a Coefficient Field)**
*Let $(A, \mathfrak{m})$ be a DVR, and $\pi$ a uniformiser of $A$. Let $k$ be a field and suppose $A$ is a $k$-algebra with $k(\mathfrak{m}) = k$. Then, for every $n \geq 0$ and $a \in A$, there is a unique $(c_0, \dots, c_n) \in k^{n+1}$ with*

$$a - \sum_{i=0}^{n} c_i \pi^i \in \mathfrak{m}^{n+1},$$

*and, writing $c_i(a) \in k$ for the resulting (compatible, as $n$ varies) sequence, there is an injective $k$-algebra homomorphism*

$$L_\pi : A \to k[[X]], \quad L_\pi(a) := \sum_{i \geq 0} c_i(a) X^i,$$

*with $L_\pi(\pi) = X$ and $L_\pi$ is an isomorphism onto $k[[X]]$ if and only if $A$ is complete. Moreover $v_X(L_\pi(a)) = v(a)$ for every $0 \neq a \in A$ and there is a unique map $L_{\hat{\pi}} : \hat{A} \to k[[X]]$ such that $L_{\hat{\pi}} \circ \iota_A = L_\pi$, which is an isomorphism.*

*Moreover $L_\pi$ extends uniquely to an injective ring homomorphism $L_\pi : K \to k((X))$ (still denoted $L_\pi$), which is an isomorphism when $A$ is complete; and, regarding $A$ as a subring of $\hat{A}$ via $\iota_A$, this commutes with $L_{\hat{\pi}}$: $L_{\hat{\pi}} \circ \iota_A = L_\pi|_A$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

© David Rufino CC BY-NC-ND 4.0

**Existence and uniqueness of $c_i(a)$.** By induction on $n$. For $n=0$: as $k(\mathfrak{m})=k$, there is a unique $c_0\in k$ with $a\equiv c_0 \pmod{\mathfrak{m}}$, that is $a-c_0\in\mathfrak{m}=\mathfrak{m}^{0+1}$. For the inductive step, given the unique $(c_0,\dots,c_{n-1})\in k^n$ with $b:=a-\sum_{i=0}^{n-1}c_i\pi^i\in\mathfrak{m}^n=(\pi^n)$: write $b=\pi^n b'$ for a (unique, $\pi^n$ a nonzerodivisor) $b'\in A$; by the $n=0$ case applied to $b'$, there is a unique $c_n\in k$ with $b'-c_n\in\mathfrak{m}$, so $b-c_n\pi^n=\pi^n(b'-c_n)\in\pi^n\mathfrak{m}=\mathfrak{m}^{n+1}$, that is $a-\sum_{i=0}^{n}c_i\pi^i\in\mathfrak{m}^{n+1}$: existence. For uniqueness: if $(c_0,\dots,c_n)$ and $(c_{0'},\dots,c_{n'})$ both satisfy the displayed condition, then, dropping the last term ($c_n\pi^n, c_{n'}\pi^n\in\mathfrak{m}^n$), both $(c_0,\dots,c_{n-1})$ and $\left(c_{0'},\dots,c_{(n-1)'}\right)$ satisfy it for $n-1$, so are equal by the inductive hypothesis; then $(c_n-c_{n'})\pi^n=\left(a-\sum_{i<n}c_i\pi^i-c_n\pi^n\right)-\left(a-\sum_{i<n}c_i\pi^i-c_{n'}\pi^n\right)\in\mathfrak{m}^{n+1}=(\pi^{n+1})$, so $c_n-c_{n'}\in(\pi)=\mathfrak{m}$ ($\pi^n$ a nonzerodivisor); as $k\to A$ is injective ($k$ a field) and $k(\mathfrak{m})=k$, $c_n-c_{n'}\in k\cap\mathfrak{m}=0$: $c_n=c_{n'}$. This uniqueness argument shows the tuple for $n$ restricts to the tuple for $n-1$, so the $(c_0,\dots,c_n)$ assemble into a well-defined compatible sequence $(c_i(a))_{i\geq 0}$.

***$L_\pi$ is a $k$-algebra homomorphism.*** Fix $n\geq 0$. By (3.8.5) (with $A:=k$, $B:=A/\mathfrak{m}^{n+1}$, $b:=\overline{\pi}$), there is a unique $k$-algebra homomorphism $k[X]\to A/\mathfrak{m}^{n+1}$, $X\mapsto\overline{\pi}$; as $\pi^{n+1}\in\mathfrak{m}^{n+1}$, this kills $X^{n+1}$, so factors as a $k$-algebra homomorphism $\overline{\psi}_n : k[X]/(X^{n+1})\to A/\mathfrak{m}^{n+1}$, $X\mapsto\overline{\pi}$. It is surjective, as $A/\mathfrak{m}^{n+1}$ is spanned, as a $k$-vector space, by $1,\overline{\pi},\dots,\overline{\pi}^n$ (existence, above); and injective, as $\sum_{i=0}^{n}d_iX^i\in\ker\overline{\psi}_n$ means $\sum_i d_i\pi^i\in\mathfrak{m}^{n+1}$, so $d_i=0$ for every $i$ (uniqueness, above, applied to $a:=0$). So $\overline{\psi}_n$ is a $k$-algebra isomorphism, and its inverse $\phi_n:=\overline{\psi}_n^{-1}$ satisfies $\phi_n(\overline{a})=\sum_{i=0}^{n}c_i(a)X^i \mod X^{n+1}$ for every $a\in A$ (as $\overline{\psi}_n$ of the right side is $\overline{a}$, by the defining property of $c_0(a),\dots,c_n(a)$).

For $a,b\in A$ and $i\leq n$: comparing coefficients of $X^i$ in $\phi_n\left(\overline{a+b}\right)=\phi_n(\overline{a})+\phi_n\left(\overline{b}\right)$ ($\phi_n$ additive) gives $c_i(a+b)=c_i(a)+c_i(b)$, and in $\phi_n\left(\overline{ab}\right)=\phi_n(\overline{a})\phi_n\left(\overline{b}\right)$ ($\phi_n$ multiplicative) gives $c_i(ab)=\sum_{j=0}^{i}c_j(a)c_{i-j}(b)$; as $n\geq i$ was arbitrary, these hold for every $i\geq 0$. So $L_\pi(a+b)=L_\pi(a)+L_\pi(b)$ and $L_\pi(ab)=L_\pi(a)L_\pi(b)$ (matching the usual formal power series operations on coefficient sequences); and $L_\pi(1)=1$ (as $1-1\cdot\pi^0=0\in\mathfrak{m}^{n+1}$ for every $n$, uniqueness gives $c_0(1)=1$, $c_i(1)=0$ for $i\geq 1$): $L_\pi$ is a ring homomorphism. For $c\in k\subset A$: $c-c\cdot\pi^0=0\in\mathfrak{m}^{n+1}$ for every $n$, so $c_0(c)=c$, $c_i(c)=0$ for $i\geq 1$, giving $L_\pi(c)=c$ (the constant series): $L_\pi$ is $k$-linear, hence a $k$-algebra homomorphism.

***$L_\pi(\pi)=X$.*** For every $n\geq 1$, $(0,1,0,\dots,0)\in k^{n+1}$ satisfies $\pi-(0\cdot\pi^0+1\cdot\pi^1+0+\cdots+0)=0\in\mathfrak{m}^{n+1}$ (and $(0)\in k^1$ satisfies it for $n=0$, as $\pi\in\mathfrak{m}$); by uniqueness, $c_0(\pi)=0$, $c_1(\pi)=1$, $c_i(\pi)=0$ for $i\geq 2$, so $L_\pi(\pi)=X$.

**Valuation formula and injectivity.** Let $0\neq a\in A$, $m:=v(a)$, so $a=u\pi^m$ for a unit $u\in A^\star$ (3.31.2). For $i<m$: $a\in\mathfrak{m}^m\subseteq\mathfrak{m}^{i+1}$, so $(0,\dots,0)\in k^{i+1}$ satisfies the defining property for $a$ at level $i$; by uniqueness, $c_j(a)=0$ for every $j\leq i<m$. For $i=m$: writing $\overline{u}\in k$ for the unique element with $u-\overline{u}\in\mathfrak{m}$ (as $k(\mathfrak{m})=k$; and $\overline{u}\neq 0$, as $u\notin\mathfrak{m}$), $a-\overline{u}\pi^m=\pi^m(u-\overline{u})\in\pi^m\mathfrak{m}=\mathfrak{m}^{m+1}$, so, with $c_0(a)=\cdots=c_{m-1}(a)=0$ already fixed, uniqueness gives $c_m(a)=\overline{u}\neq 0$. So $L_\pi(a)=\overline{u}X^m+\sum_{i>m}c_i(a)X^i$, giving $v_X(L_\pi(a))=m=v(a)$; in particular $L_\pi(a)\neq 0$ for $a\neq 0$, so $L_\pi$ is injective.

***$L_{\hat{\pi}}$ exists, is unique, and is bijective.*** $L_\pi$ is local: for $a\in\mathfrak{m}$, $c_0(a)=0$ (as $a-0\in\mathfrak{m}$), so $L_\pi(a)\in(X)$. It is continuous: for $a\in\mathfrak{m}^{n+1}$, $v(a)\geq n+1$, so $v_X(L_\pi(a))=v(a)\geq n+1$ (or $a=0$), that is $L_\pi(\mathfrak{m}^{n+1})\subseteq(X)^{n+1}$ (same-index witness). As $k[[X]]$ is a complete DVR (3.34.2) (with $t:=X$), (3.34.12) (applied to $\phi:=L_\pi$) gives a unique continuous ring homomorphism $L_{\hat{\pi}}:\hat{A}\to k[[X]]$ with $L_{\hat{\pi}}\circ\iota_A=L_\pi$. Its ramification index is $e:=v_X(L_\pi(\pi))=v_X(X)=1$ (3.31.10), and $k[[X]]=L_\pi(A)+(X)$ (as $k\subseteq L_\pi(A)$ already, $L_\pi|_k=\mathrm{id}_k$, and $k+(X)=k[[X]]$); so, by that proposition's bijectivity criterion, $L_{\hat{\pi}}$ is bijective.

***$L_\pi$ is an isomorphism onto $k[[X]]$ iff $A$ is complete.*** $L_\pi=L_{\hat{\pi}}\circ\iota_A$, with $L_{\hat{\pi}}$ bijective (above); so $L_\pi$ is bijective if and only if $\iota_A$ is. As $\iota_A$ is always injective ($\ker\iota_A=\bigcap_n\mathfrak{m}^{n+1}=0$ (3.31.9)), this holds if and only if $\iota_A$ is surjective, that is if and only if $A$ is complete (3.34.5).

***$L_\pi$ extends to $K$.*** As $L_\pi:A\to k[[X]]$ is injective (above) and $k[[X]]\subset k((X))$ (a field) (3.10.13), $A\setminus\{0\}$ maps into $k((X))^\star$. So (3.6.5) (with $S:=A\setminus\{0\}$, so $S^{-1}A=K$) gives a unique ring homomorphism $L_\pi:K\to k((X))$ (still denoted $L_\pi$) with $L_\pi|_A$ the original map, explicitly $L_\pi(a/s)=L_\pi(a)L_\pi(s)^{-1}$; it is injective (if $L_\pi(a/s)=0$ then $L_\pi(a)=0$, so $a=0$, giving $a/s=0$). If $A$ is complete, $L_\pi:A\to k[[X]]$ is moreover surjective (above), so this extension is too: given $y\in k((X))$, write $y=b/c$ for $b,c\in k[[X]]$, $c\neq 0$ (3.10.13), so $b=L_\pi(a)$, $c=L_\pi(s)$ for some $a,s\in A$, $s\neq 0$ (as $c\neq 0$, $L_\pi$ injective), giving $L_\pi(a/s)=y$; so $L_\pi:K\to k((X))$ is then bijective. Regarding $A\subset\hat{A}$ via $\iota_A$, this commutes with $L_{\hat{\pi}}$: both $L_\pi|_A$ and $L_{\hat{\pi}}\circ\iota_A$ equal the original map $L_\pi:A\to k[[X]]$ (the latter by construction of $L_{\hat{\pi}}$, above). ∎

© David Rufino CC BY-NC-ND 4.0

# 3.35 Affine Algebras

**Definition 3.35.1 (Affine Domain)**
*We call a finitely-generated $k$-algebra an* ***affine algebra****. If in addition it's integral we call it an* ***affine domain****.*

## 3.35.1 Reduction

Recall that $A_{\text{red}} \coloneqq A/N(A)$.

**Lemma 3.35.2**
*Let $A$ be a $k$-algebra and $\mathfrak{a}$ an ideal. Then $N(A/\mathfrak{a}) = \sqrt{\mathfrak{a}}/\mathfrak{a}$ and there is a canonical isomorphism*

$$\begin{array}{ccc} A/\mathfrak{a} & \twoheadleftarrow & A \\ \downarrow & & \downarrow \\ (A/\mathfrak{a})_{\text{red}} & \xrightarrow{\sim} & A/\sqrt{\mathfrak{a}} \end{array}$$

$$\overline{a+\mathfrak{a}} \longrightarrow (a+\sqrt{\mathfrak{a}})$$

*which is the unique morphism making the diagram commute.*

*Proof.* For the first part $(a+\mathfrak{a})^n = a^n + \mathfrak{a}$, so $(a+\mathfrak{a}) \in N(A/\mathfrak{a}) \iff (a+\mathfrak{a})^n = 0 \iff a^n \in \mathfrak{a}$ some $n \iff a \in \sqrt{\mathfrak{a}}$.

The second part then follows from (3.3.58). ∎

**Proposition 3.35.3**
*Let $A$ be a $k$-algebra then for every reduced $k$-algebra $B$ there is a canonical bijection*

$$\begin{aligned} \text{Mor}(A_{\text{red}}, B) &\xrightarrow{\sim} \text{Mor}(A, B) \\ \phi &\to \phi \circ \pi \end{aligned}$$

*where $\pi : A \to A_{\text{red}}$ is the quotient map, which is natural in $B$. When $A$ is reduced then $\pi$ is an isomorphism.*

*Proof.* Suppose $\tilde{\phi} : A \to B$ is a homomorphism and $a \in N(A)$. Then $0 = \tilde{\phi}(a^n) = \tilde{\phi}(a)^n$ whence $\tilde{\phi}(N(A)) \subset N(B) = (0)$, that is $N(A) \subset \ker(\tilde{\phi})$. By (3.3.56) there exists $\phi$ such that $\phi \circ \pi = \tilde{\phi}$. This shows the given map is surjective. As $\pi$ is surjective we may also deduce that the map is injective. ∎

**Proposition 3.35.4**
*Let $A, B$ be $k$-algebras and $\phi : A \to B$ be a $k$-algebra homomorphism, then there is a unique morphism $\phi_{\text{red}} : A_{\text{red}} \to B_{\text{red}}$ making the following diagram commute*

$$\begin{array}{ccc} A & \xrightarrow{\phi} & B \\ \downarrow & & \downarrow \\ A_{\text{red}} & \dashrightarrow & B_{\text{red}} \end{array}$$

*given by*

$$\phi_{\text{red}}(a + N(A)) = \phi(a) + N(B)$$

*Furthermore $(\psi \circ \phi)_{\text{red}} = \psi_{\text{red}} \circ \phi_{\text{red}}$. If $\phi$ is an isomorphism then so is $\phi_{\text{red}}$.*

*Proof.* In the notation of (3.35.3) $\phi_{\text{red}}$ is simply the morphism corresponding to $\pi_B \circ \phi$, from which existence and uniqueness follows.

Uniqueness then also shows that this is functorial. ∎

## 3.35.2 Normalisation

The following normalisation results can be seen as a refinement of results on transcendence bases (see the Transcendental Field Extensions subsection).

**Definition 3.35.5 (Algebraically Independent)**
*Let $A$ be a $k$-algebra and $x_1, ..., x_n$ elements of $A$. Then we say they are* ***algebraically independent*** *if one of the following equivalent conditions holds*

© David Rufino CC BY-NC-ND 4.0

- *The unique $k$-algebra homomorphism $\phi : k[X_1, ..., X_n] \to A$ such that $\phi(X_i) = x_i$ (evaluation homomorphism) is injective*
- *There are no non-zero polynomials $f(X_1, ..., X_n)$ such that $f(x_1, ..., x_n) = 0$.*

*Note in particular it induces an isomorphism $k[X_1, ..., X_n] \xrightarrow{\sim} k[x_1, ..., x_n] \subset A$.*

**Definition 3.35.6 (Normalising Family)**
*Let $A$ be a finitely-generated $k$-algebra. A **normalising family** is a set $\{x_1, ..., x_n\}$ of elements of $A$ such that*

- *$x_1, ..., x_n$ are **algebraically independent** over $k$*
- *$A$ is a finite $k[x_1, ..., x_n]$-module (equivalently integral over $k[x_1, ..., x_n]$).*

*NB this is completely equivalent to specifying an integral, injective map*

$$k[X_1, ..., X_n] \hookrightarrow A$$

This may be seen as a refined transcendence base. More precisely we have the following

**Proposition 3.35.7 (Relationship to Transcendence Base)**
*Let $A$ be an integral finitely-generated $k$-algebra with $K := \mathrm{Frac}(A)$. Let $S \subset A$ be a subset. Then*

- *If $A$ is integral over $k[S]$ then $K/k(S)$ is algebraic*
- *If $S$ is a normalising family for $A$ then $S$ is a transcendence basis for $K/k$*

*In particular normalising families have order $\mathrm{trdeg}(K/k)$.*

*Proof.* By (3.19.45) the set $\{x \in K \mid x \text{ algebraic over } k(S)\}$ forms a subfield containing $A$, and therefore equals $K$.

The final statement follows from (3.19.141). ∎

The following is useful as it removes the necessity of showing algebraic independence in certain cases.

**Corollary 3.35.8**
*Let $A$ be an integral finitely-generated $k$-algebra with $K := \mathrm{Frac}(A)$. Let $S \subset A$ be a subset such that*

- *$A$ is integral over $k[S]$*
- *$\#S \leq \mathrm{trdeg}_k(K)$*

*then $S$ is a **normalising family**.*

*Proof.* This follows from the previous result and (3.19.141). ∎

There are a few forms of the Normalisation Lemma which we prove, of progressively stronger form.

**Lemma 3.35.9 (Hypersurface Normalisation Lemma)**
*Let $A = k[X_1, ..., X_n]$ be a polynomial ring over an infinite field $k$ and $0 \neq F \in A$. Then there exists $\lambda_1, ..., \lambda_{n-1} \in k$ such that*

- *$x_i := X_i - \lambda_i F \quad 1 \leq i \leq n-1$*
- *$x_1, ..., x_{n-1}, F$ is a normalising family for $A$*
- *$FA \cap k[x_1, ..., x_{n-1}, F] = Fk[x_1, ..., x_{n-1}, F]$*

*This may be viewed as a commutative diagram*

$$\begin{array}{ccc} \phi : k[Y_1, ..., Y_n] & \xrightarrow{\sim} & k[X_1, ..., X_n] \\ \uparrow & & \downarrow \\ k[Y_1, ..., Y_{n-1}] & \hookrightarrow & k[X_1, ..., X_n]/(F) \end{array}$$

*where the horizontal arrows are injective, integral (and finite) and $\phi^{-1}((F)) = (Y_n)$.*

**Remark 3.35.10**
*Reversing the arrows we get the geometric picture, where horizontal arrows are finite and surjective*

© David Rufino CC BY-NC-ND 4.0

$$\begin{array}{ccc} \mathbb{A}^n & \xrightarrow{\sim} & \mathbb{A}^n \\ \uparrow & & \downarrow \\ V(F) & \twoheadrightarrow & \mathbb{A}^{n-1} \end{array}$$

*in other words after a linear change of variables we may express $V(F)$ as a finite covering of a standard hyperplane.*

**Proposition 3.35.11 (Nagata Normalisation Lemma)**
*Let $A = k[x_1, ..., x_n]$ be a finitely-generated $k$-algebra such that $k$ is infinite. Then there exists a **normalising family** $y_1, ..., y_d \in A$ such that each $y_i$ is a $k$-linear combination of $x_1, ..., x_n$. Further if $\mathfrak{a}_1 \lhd A$ is a proper ideal, then these may be chosen such that*

$$\mathfrak{a}_1 \cap k[y_1, ..., y_d] = (y_1, ..., y_h)k[y_1, ..., y_d]$$

*for some $0 \leq h \leq d$, where $h = 0$ denotes the zero ideal.*

**Proposition 3.35.12 (Bourbaki Normalisation Lemma)**
*Let $A = k[x_1, ..., x_n]$ be a finitely-generated $k$-algebra such that $k$ is infinite. Then there exists a **normalising family** $y_1, ..., y_d \in A$ such that each $y_i$ is a $k$-linear combination of $x_1, ..., x_n$.*

*Furthermore for any finite chain of proper ideals in $A$*

$$\mathfrak{a}_1 \subseteq \mathfrak{a}_2 \subseteq ... \subseteq \mathfrak{a}_p \subsetneq A$$

*the family may be chosen such that*

$$\mathfrak{a}_j \cap k[y_1, ..., y_d] = \left(y_1, ..., y_{h(j)}\right)k[y_1, ..., y_d] \quad 1 \leq j \leq p \ .$$

*for some non-decreasing sequence of integers $h(j)$, where $h(j) = 0$ denotes the zero ideal.*

**Remark 3.35.13 (Geometric Interpretation)**
*Note the normalisation here is equivalent to a commutative diagram*

$$\begin{array}{ccc} k[Y_1, ..., Y_d] & \hookrightarrow & A \\ \uparrow & & \downarrow \\ k\left[Y_{h(1)+1}, ..., Y_d\right] & \hookrightarrow & A/\mathfrak{a}_1 \\ \uparrow & & \downarrow \\ \vdots & & \vdots \\ \uparrow & & \downarrow \\ k\left[Y_{h(p)+1}, ..., Y_d\right] & \hookrightarrow & A/\mathfrak{a}_p \end{array}$$

*where the horizontal arrows are integral and injective, and the top arrow is given by*

$$\phi : Y_i \to \sum_j \lambda_{ij} x_j$$

*such that $\phi^{-1}(\mathfrak{a}_i) = \left(Y_1, ..., Y_{h(i)}\right)$*

*As before this expresses $A$ as a finite covering of $\mathbb{A}^d$ under which each subvariety is also a finite covering of a standard linear subspace.*

We first prove the weaker form of the Normalisation Lemma

*Proof of (3.35.9).* First decompose $F$ into homogenous polynomials (monomials of the same degree)

$$F = F_0 + F_1 + ... + F_m$$

and observe that the monomial $X_n^m$ appears only in $F_m$. Define

$$F' \coloneqq F(X_1 + \lambda_1 X_n, ..., X_{n-1} + \lambda_{n-1} X_n, X_n)$$

Furthermore in terms of homogenous polynomials

© David Rufino CC BY-NC-ND 4.0

$$F'_m = F_m(X_1 + \lambda_1 T, ..., X_{n-1} + \lambda_{n-1} X_n, X_n)$$

and the coefficient of $X_n^m$ in $F'$ is simply $F'_m(0, ..., 0, 1) = F_m(\lambda_1, ..., \lambda_{n-1}, 1) \in k$. There are only finitely many values of $\lambda$ such that $F_m$ is zero, whence there exists a $\lambda$ such that $X_n^m$ has non-zero coefficient in $F'$, whence $F'$ is monic. By the previous Lemma we have $F'(x_1, ..., x_{n-1}, X_n) = 0$, with the leading coefficient in $X_n$ constant.

Let $B \coloneqq k[x_1, ...x_{n-1}, F]$. Then clearly $B[X_n] = A$ and $X_n$ is integral over $B$. Therefore by (3.24.4) $A$ is a finite $B$-module. As $\operatorname{trdeg}(A/k) = n$ by (3.35.8) $\{x_1, ..., x_{n-1}, F\}$ is a normalising family, and $B$ is isomorphic to a polynomial ring.

As $B$ is a polynomial ring it is integrally closed (3.24.12). Then the final statement is a consequence of the following lemma (3.35.14) (essentially to prove that $V(F) \to \mathbb{A}^{n-1}$ is surjective). ∎

**Lemma 3.35.14**
*Let $A$ be integrally closed, $\phi : A \hookrightarrow B$ injective and integral and $a \in A$. Then*

$$(a)^{\mathrm{ec}} = \phi^{-1}(\phi(a)B) = (a)$$

*Proof.* Let $K = \operatorname{Frac}(A)$ and $L = \operatorname{Frac}(B)$, then $\phi$ extends to an injection $\phi : K \hookrightarrow L$.

Generically we have $(a) \subseteq (a)^{\mathrm{ec}}$. Suppose $a' \in (a)^{\mathrm{ec}}$, then $\phi(a') = \phi(a)b$. Therefore $\phi(a'a^{-1}) = b$ is integral over $A$, whence so is $a'a^{-1}$. As $A$ is integrally closed we have $a' \in (a)$ as required. ∎

Before proving the stronger version of Normalisation Lemma, we need some preliminary technical results

**Lemma 3.35.15**
*Let $A$ be a $k$-algebra and $\mathfrak{a} \lhd A$ an ideal. Then $\mathfrak{a}$ is proper $\iff \mathfrak{a} \cap k = \{0\}$.*

**Lemma 3.35.16**
*Let $A = k[X_1, ..., X_n]$ be a polynomial ring and $\mathfrak{a} \lhd A$ a proper ideal. Then TFAE*

*a)* $\mathfrak{a} = (X_1, ..., X_h)$
*b)*
  *i)* $X_1, ..., X_h \in \mathfrak{a}$
  *ii)* $\mathfrak{a} \cap k[X_{h+1}, ..., X_n] = \{0\}$

*In this case* $\mathfrak{a} \cap k[X_1, ..., X_h] = (X_1, ..., X_h)k[X_1, ..., X_h]$

*Proof.* We claim that there is a direct sum of $k$-vector spaces

$$k[X_1, ..., X_n] = (X_1, ..., X_h) \oplus k[X_{h+1}, ..., X_n]$$

from which the result largely follows. Let $S$ be the set of monomials in which at least one of $X_1, ..., X_h$ appears, and let $T$ be the set for which none appears (but including 1). Then clearly $A = \langle S \rangle \oplus \langle T \rangle$ and $k[X_{h+1}, ..., X_n] = \langle T \rangle$. We argue that $(X_1, ..., X_h) = \langle S \rangle$. First $S$ is stable under multiplication by $X_1, ..., X_n$, and so $\langle S \rangle$ is an ideal. One inclusion is obvious, furthermore it's clear that $S \subseteq (X_1, ..., X_n)$ from which the claim follows. ∎

We may now proceed to the proof of the stronger versions of the Normalisation Lemma.

*Reduction to polynomial ring case for (3.35.11), (3.35.12).* We show that for both forms it is possible to reduce to the case of a polynomial ring. For let $A$ be a finitely-generated $k$-algebra then we may write $A \coloneqq k[X_1, ..., X_n]/\mathfrak{a}$ for some ideal $\mathfrak{a}$. Then the polynomial ring case for $p = 1$ shows the existence of an integral, injective map

$$\phi : k[Y_1, ..., Y_m] \hookrightarrow A$$

If $\mathfrak{a}_i$ is a chain of ideals in $A$, then $\mathfrak{a}'_i \coloneqq \phi^{-1}(\mathfrak{a}_i)$ is a chain of ideals in $k[Y_1, ..., Y_m]$. The general case for a polynomial ring shows the existence of an integral map

$$\psi : k[Z_1, ..., Z_m] \hookrightarrow k[Y_1, ..., Y_m]$$

such that

$$\psi^{-1}(\mathfrak{a}'_i) = \left(Z_1, ..., Z_{h(i)}\right)$$

The composition $\phi \circ \psi$ gives the required normalisation of $A$. Geometrically express $A$ as a finite covering of affine space, by considering it as a subvariety of larger affine space. ∎

*Proof of (3.35.11) in the polynomial ring case.* Let $A = k[X_1, ..., X_n]$ and proceed by induction on $n$.

Note that as $\mathfrak{a}_1$ is proper, we must have $\mathfrak{a}_1 \cap k = \{0\}$, and we may obviously also assume that $\mathfrak{a}_1 \neq (0)$ (as otherwise we may take $h = 0$).

© David Rufino CC BY-NC-ND 4.0

Choose $0 \neq x_1 \in \mathfrak{a}_1$. Then by (3.35.9) there exists $t_2, ..., t_n \in A$ such that $x_1, t_2, ..., t_n$ is a normalising family and $(x_1) \cap B = x_1 B$ where $B := k[x_1, t_2, ..., t_n]$. In the case that $\mathfrak{a}_1$ is principal we are done, since the choice of $x_1$ was arbitrary, and in this case $h(1) = 1$. In particular this covers the base case $n = 1$ because $A$ is a PID (3.16.4).

Otherwise $B' := k[t_2, ..., t_n]$ is a polynomial ring and $\mathfrak{a}_1' := \mathfrak{a}_1 \cap B'$ is proper by (3.35.15). By induction on $n$ there is a normalising family $x_2, ..., x_n$ for $B'$ such that $\mathfrak{a}_1' \cap C' = (x_2, ..., x_h)C'$ where $C' := k[x_2, ..., x_n]$ and $B'$ is integral over $C'$.

Define $C := k[x_1, ..., x_n] = C'[x_1]$ then $x_1, t_2, ..., t_n$ are integral over $C$, so $B$ is integral over $C$ (3.24.4), and $A$ is integral over $C$ (3.24.5), so by (3.35.8) $x_1, ..., x_n$ is a normalising family for $A$.

We claim that $\mathfrak{a}_1'' := \mathfrak{a}_1 \cap C = (x_1, ..., x_h)C$. Clearly $x_1, ..., x_h \in \mathfrak{a}_1''$, and $\mathfrak{a}_1'' \cap k[x_{h+1}, ..., x_n] = \mathfrak{a}_1' \cap k[x_{h+1}, ..., x_n] = \{0\}$ by (3.35.16) applied to the ring $C'$. Then (3.35.16) applied to the ring $C$ demonstrates the claim. ∎

*Proof of (3.35.12) in the polynomial ring case.* We can then show the case $p > 1$ by induction, for by the induction hypothesis there exists a normalising family $t_1, ..., t_n$ for $A$ such that

$$\mathfrak{a}_j \cap B = \big(t_1, ..., t_{h(j)}\big)B \quad 1 \leq j \leq p-1$$
$$B := k[t_1, ..., t_n]$$

Let $r = h(p-1)$, then by the case $p = 1$ applied to the ring $B' := k[t_{r+1}, ..., t_n]$ and ideal $\mathfrak{a}_p \cap B'$ there exists a normalising family $x_{r+1}, ..., x_n$ for $B'$ such that for some $s \leq n$ (possibly equal to $r$ to denote the zero ideal),

$$\mathfrak{a}_p \cap C' = (x_{r+1}, ..., x_s)C'$$
$$C' := k[x_{r+1}, ..., x_n]$$

We claim that $t_1, ..., t_r, x_{r+1}, ..., x_n$ is a suitable normalising family for $A$, with $h(p) = s$.

For define $C := k[t_1, ..., t_r, x_{r+1}, ..., x_n] = C'[t_1, ..., t_r]$. Recall $B'$ is integral over $C'$, and $t_1, ..., t_r$ are obviously integral over $C$ so $B = B'[t_1, ..., t_r]$ is integral over $C$ by (3.24.7). Then $A$ is integral over $C$ by (3.24.5), and this is a normalising family by (3.35.8), and in particular algebraically independent.

For $j \leq p-1$ and $h := h(j) \leq r$, apply (3.35.16) to the ideal $\mathfrak{a}_j \cap B$ to see $\mathfrak{a}_j \cap k[t_{h+1}, ..., t_n] = \{0\}$ and therefore $\mathfrak{a}_j \cap k[t_{h+1}, ..., t_r, x_{r+1}, ..., x_n] = \{0\}$. As $t_1, ..., t_h \in \mathfrak{a}_j$ we see by (3.35.16) that $\mathfrak{a}_j \cap C = (t_1, ..., t_h)C$ as required.

Similarly by (3.35.16) $\mathfrak{a}_p \cap k[x_{s+1}, ..., x_n] = \{0\}$ and clearly $t_1, ..., t_r, x_{r+1}, ..., x_s \in \mathfrak{a}_p$. Then by (3.35.16) again $\mathfrak{a}_p \cap C = (t_1, ..., t_r, x_{r+1}, ..., x_s)C$ as required. ∎

**Remark 3.35.17**
*In Bourbaki's proof the reduction to the polynomial ring case increases $p$ to $p+1$, so in particular the $p = 1$ case requires the more complex reduction argument at the end of the proof. Here we use a simplified argument from [Sha00].*

## 3.35.3 Nullstellensatz

**Definition 3.35.18 (Zeros of an ideal)**
*Let $\mathfrak{a} \lhd k[X_1, ..., X_n]$ be an ideal and $K/k$ a field extension. Then a point $x \in K^n$ is a **zero** of $\mathfrak{a}$ if*

$$f \in \mathfrak{a} \implies f(x) = 0$$

*We define the **residue field** to be*

$$k(x) := k(x_1, ..., x_n)$$

*and also denote*

$$k[x] := k[x_1, ..., x_n]$$

*We say it is **algebraic** if $k(x)/k$ is algebraic (necessarily finite).*

The following observation is useful

**Proposition 3.35.19 (Zeros are homomorphisms)**
*Let $\mathfrak{a} \lhd k[X_1, ..., X_n]$ be an ideal then there is a bijection*

$$\mathrm{AlgHom}_k(k[X_1, ..., X_n]/\mathfrak{a}, K) \longleftrightarrow \{\text{zeros of } \mathfrak{a} \text{ in } K^n\}$$
$$\phi \longrightarrow \big(\phi\big(\overline{X}_1\big), ..., \phi\big(\overline{X}_n\big)\big)$$

© David Rufino CC BY-NC-ND 4.0

*The following notion is useful in future*

***Definition 3.35.20 (Generic Point)***
Let $\mathfrak{a} \triangleleft k[X_1, ..., X_n]$ be a **prime** ideal. then a point $\xi \in L^n$ is a **generic point** of $\mathfrak{a}$ if $\ker(\mathrm{ev}_\xi) = \mathfrak{a}$. Note one always exists, because we may take $L = \mathrm{Frac}(A)$ and $\xi_i = \overline{X}_i$.

When $x$ is another zero of $\mathfrak{a}$ this induces a $k$-algebra homomorphism

$$k[\xi] \to k[x]$$

which is an isomorphism precisely when $x$ is a generic point.

Generally we are interested in the relationship between ideals of $k[X_1, ..., X_n]$ and corresponding zeros in an extension field $K/k$. The following proposition is fundamental

**Proposition 3.35.21 (Correspondence between ideals and zeros)**
*Let $K/k$ be a field extension and $x \in K^n$. Define $\mathfrak{m}_x$ to be the kernel of the homomorphism*

$$\mathrm{ev}_x : k[X_1, ..., X_n] \to K$$

*Then*

- *$\mathfrak{m}_x$ is a prime ideal*
- *If $K/k$ is an algebraically closed field of transcendence degree $\geq n$ then every prime ideal is of this form.*
- *If $x_i$ are algebraic over $k$ (e.g. if $K/k$ is algebraic) then $\mathfrak{m}_x$ is maximal*

*In this case we have a canonical isomorphism*

$$k[X_1, ..., X_n]/\mathfrak{m}_x \xrightarrow{\sim} k[x] \subset K$$

*and when each $x_i$ is algebraic then $k[x] = k(x)$ is a finite extension of $k$.*

*Proof.* The canonical isomorphism follows from (3.11.3). Any subring of a field is an integral domain, which means $\mathfrak{p}_x$ is prime by (3.3.59).

By (3.19.44) $x_i$ are algebraic over $k$ if and only if $k(x_1, ..., x_n)/k$ is algebraic (and even finite). By the same result $k[x_1, ..., x_n] = k(x_1, ..., x_n)$ and therefore $\mathfrak{m}_x$ is maximal by (3.3.59).

Let $\mathfrak{p}$ be a prime ideal and define $k(x) \coloneqq \mathrm{Frac}(k[x])$ and $k[x] \coloneqq k[X_1, ..., X_n]/\mathfrak{p}$. If $K$ has transcendence degree $\geq n$ then there is an embedding $k(x)/k \to K/k$ by (3.19.61). This restricts to an isomorphism $k[x] \xrightarrow{\sim} k[\overline{x}]$ for some $\overline{x}_i \in K$. It's clear that $\mathfrak{p} = \mathfrak{m}_x$. ∎

We may show that all maximal ideals correspond to zero-loci.

**Lemma 3.35.22 (Zariski's Lemma)**
*Let $k$ be an infinite field and $A$ a finitely-generated $k$-algebra which is a field. Then $A$ is a finite extension of $k$.*

*Proof.* By Normalisation Lemma (3.35.11) there is an injective, integral map $k[X_1, ..., X_d] \hookrightarrow A$ for some $d \geq 0$. By (3.24.14) then $k[X_1, ..., X_d]$ is a field which implies $d = 0$. We conclude that $A$ is algebraic over $k$, and by (3.19.44) finite over $k$. ∎

**Proposition 3.35.23 (Weak Nullstellensatz I)**
*Every maximal ideal of $k[X_1, ..., X_n]/\mathfrak{a}$ is of the form $\mathfrak{m}_x/\mathfrak{a}$ for $x \in \overline{k}^n$ a zero of $\mathfrak{a}$.*

*In addition the conclusion of Zariski's Lemma holds for all (possibly finite) coefficient fields $k$.*

*Proof.* Observe that $x \in \overline{k}^n$ is a zero of $\mathfrak{a}$ if and only if $\mathfrak{a} \subset \mathfrak{m}_x$. Therefore we may reduce to the case $\mathfrak{a} = (0)$ by results on quotient rings (3.3.56).

We've already shown that $\mathfrak{m}_x$ is maximal when $x \in \overline{k}^n$ is algebraic (3.35.21). Conversely if $\mathfrak{m}$ is a maximal ideal, then $\mathfrak{m}\overline{k}[X_1, ..., X_n]$ is a proper ideal contained in some maximal ideal $\mathfrak{m}' \triangleleft \overline{k}[X_1, ..., X_n]$. If $\mathfrak{m}'$ has an algebraic zero then so does $\mathfrak{m}$ and $\mathfrak{m} \subset \mathfrak{m}_x$ for some $x$, which are equal by maximality.

However $\overline{k}[X_1, ..., X_n]/\mathfrak{m}'$ is a finitely-generated $\overline{k}$-algebra which is a field and $\overline{k}$ is infinite. Therefore by (3.35.22) it is $\overline{k}$, and this yields an algebraic zero of $\mathfrak{m}'$ as required.

We may generalise Zariski's Lemma to the case $k$ is not necessarily infinite. If $A$ is a finitely-generated $k$-algebra which is a field then we have shown it is isomorphic to $k[X_1, ..., X_n]/\mathfrak{m}_x$ for some algebraic point $x$. By (3.35.21) is a finite extension of $k$, as required. ∎

© David Rufino CC BY-NC-ND 4.0

We may make the correspondence between maximal ideals and algebraic points more precise

**Proposition 3.35.24 (Weak Nullstellensatz II)**
*There is a bijective map*

$$\begin{aligned} \overline{k}^n / \operatorname{Aut}\left(\overline{k}/k\right) &\longrightarrow \{\mathfrak{m} \lhd k[X_1, ..., X_n] \text{ maximal}\} \\ x &\longrightarrow \mathfrak{m}_x \end{aligned}$$

*When $x \in k^n$ then*

$$\mathfrak{m}_x = (X_1 - x_1, ..., X_n - x_n)$$

*Proof.* The map is surjective by (3.35.23). It's well-defined because $\mathfrak{m}_{\sigma(x)} = \mathfrak{m}_x$.

By (3.35.21) we have an isomorphism $k[X_1, ..., X_n]/\mathfrak{m}_x \xrightarrow{\sim} k[x_1, ..., x_n] \subset \overline{k}$. If $\mathfrak{m}_x = \mathfrak{m}_y$ then these compose to yield an isomorphism $\sigma : k[x_1, ..., x_n] \xrightarrow{\sim} k[y_1, ..., y_n] \subset \overline{k}$ such that $\sigma(x_i) = y_i$. By (3.19.69) this extends to $\sigma \in \operatorname{Aut}\left(\overline{k}/k\right)$. Therefore the given mapping is injective. ∎

## 3.35.4 Morphisms of Affine Algebras

**Definition 3.35.25**
*Let $B$ be a $k$-algebra. We say that $B$ is **algebraic** over $k$ if for every maximal ideal $\mathfrak{m} \lhd B$ we have $B/\mathfrak{m} \xrightarrow{\sim} k(\mathfrak{m})$ is algebraic over $k$.*

**Proposition 3.35.26**
*Let $B$ be a $k$-algebra, then it is algebraic in each of the following cases*

*a) $B$ is a finitely generated $k$-algebra*
*b) $B$ is a local $k$-algebra such that $k(\mathfrak{m}_B)/k$ is algebraic*
*c) $B$ is an algebraic field extension of $k$*
*d) $B$ is the localisation of a finitely-generated $k$-algebra at a maximal ideal*

*Proof.* We prove each in turn

a) This follows from Zariski's Lemma in the general case (3.35.23)
b) The only maximal ideal is $\mathfrak{m}_B$ so this is obvious
c) This is a special case of b)
d) This follows from a) and b) because $B_{\mathfrak{m}}$ is a local ring with residue field isomorphic to $B/\mathfrak{m}$ (3.6.43).

∎

**Proposition 3.35.27**
*Let $\phi : A \to B$ be a $k$-algebra homomorphism, with $B$ algebraic. Then the inverse image of a maximal ideal is again maximal.*

*Proof.* Let $\mathfrak{n}$ be a maximal ideal. By assumption $B/\mathfrak{n}$ is integral over $k$, and so a-fortiori $B/\mathfrak{n}$ is integral over $A$. Therefore $\phi^{-1}(\mathfrak{n})$ is maximal by (3.24.16). ∎

## 3.35.5 Krull Dimension of Affine Algebras

The Krull Dimension of affine domains is particularly well-behaved. Specifically they are biequidimensional and therefore satisfy a co-dimension formula (3.35.35). Further there's a geometric proof of the "Hauptidealsatz" (3.35.32). We first show that it is equal to transcendence degree in the integral case.

**Proposition 3.35.28**
*Let $A$ be an affine algebra with normalising family $x_1, ..., x_n$, then $\dim A = n$.*

*In particular every affine domain $A$ satisfies $\dim A = \operatorname{trdeg}(\operatorname{Frac}(A)/k)$, and the polynomial ring $k[X_1, ..., X_n]$ has dimension $n$.*

*Proof.* By definition $A$ is integral over $k[x_1, ..., x_n]$, which is isomorphic to a polynomial ring so we may reduce to the case of polynomial ring by (3.24.22).

$\dim A \geq n$). This is clear by considering the chain of prime ideals

$$(X_1) \subsetneq (X_1, X_2) \subsetneq ... \subsetneq (X_1, ..., X_n)$$

$\dim A \leq n$). We may argue by the Strong Normalisation Lemma (3.35.12) and the subsequent remark that any chain of prime ideals must have length at most $n$, as any normalising family has order $n$.

© David Rufino CC BY-NC-ND 4.0

Alternatively we may proceed by induction on $n$ to show $\dim k[X_1, ..., X_n] = n$. Consider a maximal chain

$$\mathfrak{p}_0 \subsetneq \mathfrak{p}_1 \subsetneq ... \subsetneq \mathfrak{p}_m$$

Clearly $\mathfrak{p}_0 = 0$, and $\mathfrak{p}_1 = (f)$ principal by (3.22.5). By (3.35.9) there is an integral, injective map

$$k[Y_1, ..., Y_{n-1}] \hookrightarrow k[X_1, ..., X_n]/(f)$$

whence $\dim(\mathfrak{p}_1) = \dim(k[X_1, ..., X_n]/(f)) = \dim(k[Y_1, ..., Y_{n-1}]) = n - 1$, and by definition $m - 1 \leq n - 1$. As the maximal chain was arbitrarily chosen, we have $\dim k[X_1, ..., X_n] \leq n$. The reverse inequality was already shown so we are done.

The final statement follows from the existence of a normalising family (3.35.11) and (3.35.7) ■

**Corollary 3.35.29**
*Let $A$ be a finitely-generated $k$-algebra. Then* $\dim A[T] = \dim A + 1$.

*Proof.* By (3.35.11) there is an injective integral ring homomorphism

$$\phi : k[X_1, ..., X_n] \hookrightarrow A$$

where $\dim A = n$. By (3.8.5) there is a ring homomorphism

$$k[X_1, ..., X_n][T] \hookrightarrow A[T]$$

which is injective. Further by assumption $A = k[y_1, ..., y_r]$ so $A[T] = k[y_1, ..., y_r, T]$ and by (3.24.4) the ring homomorphism is integral. This shows that $\phi(x_1), ..., \phi(x_n), T$ is a normalising family and the result follows from (3.35.28). ■

**Corollary 3.35.30**
*The ideal $(X_1, ..., X_r) \triangleleft k[X_1, ..., X_n]$ has dimension $n - r$.*

**Corollary 3.35.31**
*Let $A$ be an integral finitely-generated $k$-algebra and $0 \neq f$ then* $\dim A = \dim A_f$

The following proof is due to Tate, and presented in the Red Book [Mum99] (I.7 Theorem 2).

**Proposition 3.35.32 (Hypersurface has pure codimension 1)**
*Let $A$ be an affine domain of dimension $n$ and $0 \neq f \in A$. Then*

$$\dim((f)) = n - 1$$

$$\mathrm{ht}((f)) = 1$$

*More precisely if $\mathfrak{p}$ is minimal over $(f)$ then it has dimension $n - 1$ and height 1.*

*Proof.* Consider the case $A = k[X_1, ..., X_n]$. Suppose first that $f$ is prime, then $\mathfrak{p} = (f)$ and by (3.35.9) there is an integral injective map

$$k[Y_1, ..., Y_{n-1}] \hookrightarrow A/(f)$$

Therefore

$$\dim((f)) = \dim(A/(f)) \overset{(3.24.22)}{=} \dim(k[Y_1, ..., Y_{n-1}]) \overset{(3.35.28)}{=} n - 1$$

When $f$ is not prime then, as $k[X_1, ..., X_n]$ is a UFD, we have a prime factorisation

$$f = \prod_{i=1}^{n} f_i^{m_i}$$

and the minimal prime decomposition is

$$\sqrt{(f)} = (f_1) \cap ... \cap (f_n)$$

In particular any prime minimal over $(f)$ has the form $\mathfrak{p} = (f_i)$ for some $i$, and we may reduce to the case of $f$ prime.

For an arbitrary $k$-algebra $A$ we have a decomposition into minimal primes of $(f)$

$$\sqrt{(f)} = \mathfrak{p}_1 \cap ... \cap \mathfrak{p}_n$$

and without loss of generality $\mathfrak{p} = \mathfrak{p}_1$. We may localize to the case of a single prime, for choose $g \notin \mathfrak{p}$ and $g \in \mathfrak{p}_i$ for $i = 2...n$. Consider the localization $A \to A_g$, then we claim that

© David Rufino CC BY-NC-ND 4.0

$$\sqrt{(f/1)} = \mathfrak{p}A_g$$

For by (3.6.19) there is a correspondence between primes of $A_g$ containing $(f/1)$ and primes of $A$ containing $f$ and not $g$, which are precisely the primes containing $\mathfrak{p}$ by (3.3.40) or (3.3.44). Therefore $\mathfrak{p}A_g$ is the only minimal prime of $A_g$ containing $(f/1)$ and the claim follows from (3.3.46).

Note that $\dim A = \dim A_g$ as they have the same quotient field and therefore transcendence degree. Similarly $\dim(A/\mathfrak{p}) = \dim\big((A/\mathfrak{p})_{\bar{g}}\big) = \dim\big(A_g/\mathfrak{p}_g\big) = \dim\big(\mathfrak{p}A_g\big)$. So we may assume without loss of generality that $n = 1$ and $\mathfrak{p} = \sqrt{(f)}$.

By (3.35.11) there is an integral, injective map

$$\phi : B \hookrightarrow A$$

where $B = k[X_1, ..., X_n]$, which induces an algebraic field extension

$$K := \mathrm{Frac}(B) \hookrightarrow \mathrm{Frac}(A) =: L$$

We claim that there exists $f_0 \in B$ such that

$$\phi^{-1}\big(\sqrt{f}\big) = \sqrt{(f_0)}$$

Observe that in this case $\sqrt{(f_0)}$ is prime and therefore the unique minimal prime over $(f_0)$. Therefore the result would follow from the first part and (3.24.23). Firstly for any $g \in A$ we have (3.24.18)

$$\mathrm{Norm}_{L/K}(g) \in B \cap \phi^{-1}((g))$$

(because $B$ is a UFD and therefore integrally closed (3.24.12)). Define $f_0 := \mathrm{Norm}_{L/K}(f)$ then we see that $f_0 \in \phi^{-1}((f)) \implies \sqrt{(f_0)} \subseteq \phi^{-1}\big(\sqrt{(f)}\big)$. Conversely if $\phi(g)^n = hf$ then $g^{n[L:K]} = \mathrm{Norm}(\phi(g)^n) = \mathrm{Norm}(h)f_0 \in (f_0) \implies g \in \sqrt{(f_0)}$. Therefore the reverse inclusion holds.

Finally by the codimension 1 formula (3.22.8) we have $\mathrm{ht}((f)) = 1$. ∎

**Remark 3.35.33**
*The argument is slightly less awkward in geometric language. Decompose into irreducibles*

$$V(f) = Z_1 \cup ... \cup Z_n$$

*choose a principal open affine $D(g)$ which meets only $Z_1$ then $Z_1 \cap D(g) = V(f) \cap D(g) = V(f/1)$ is an irreducible component of $D(g)$. We argue that $\dim(X) = \dim(D(g))$ and $\dim(Z_1) = \dim(D(g) \cap Z_1)$. Further construct finite coverings*

$$V(f/1) \to V(f_0) \to H$$

*onto a hyperplane in $\mathbb{A}^n$.*

This allows is to prove a converse to (3.22.8)

**Corollary 3.35.34 (Height 1 formula)**
*Let $A$ be an affine domain. Then for a prime ideal $\mathfrak{p}$*

$$\mathrm{ht}(\mathfrak{p}) = 1 \implies \dim(\mathfrak{p}) = \dim(A) - 1$$

*More generally if $A$ is an affine algebra and $\mathfrak{p}_1 \subsetneq \mathfrak{p}_2$ is a saturated chain of prime ideals then*

$$\dim(\mathfrak{p}_2) = \dim(\mathfrak{p}_1) - 1$$

*Proof.* Choose $0 \neq f \in \mathfrak{p}$ then it follows from the previous result (3.35.32). Alternatively by (3.35.11) there is an integral injective map

$$\phi : k[X_1, ..., X_n] \hookrightarrow A$$

with $\mathfrak{q} := \phi^{-1}(\mathfrak{p})$ and $n = \dim A$. Then by Going Down we have $\mathrm{ht}(\mathfrak{q}) = 1$. By (3.22.5) $\mathfrak{q}$ is principal and by (3.35.9) there is an integral injective map

$$k[Y_1, ..., Y_{n-1}] \hookrightarrow k[X_1, ..., X_n]/\mathfrak{q}$$

therefore $\dim(\mathfrak{q}) = n - 1$. By (3.24.23) this equals $\dim(\mathfrak{p})$ and we are done.

For the second statement we may consider the affine domain $A/\mathfrak{p}_1$ and observe that $\mathrm{ht}(\mathfrak{p}_2/\mathfrak{p}_1) = 1$. Therefore

© David Rufino CC BY-NC-ND 4.0

$$\dim(\mathfrak{p}_2) = \dim(\mathfrak{p}_2/\mathfrak{p}_1) = \dim(A/\mathfrak{p}_1) - 1 = \dim(\mathfrak{p}_1) - 1$$

■

**Corollary 3.35.35 (Biequidimensionality)**
*Let $A$ be an affine algebra. Then it is **quasi-biequidimensional** and satisfies the **codimension formulae** for $\mathfrak{p} \subset \mathfrak{a}$*

$$\dim \mathfrak{a} = \dim \mathfrak{p} + \operatorname{ht}(\mathfrak{a}/\mathfrak{p})$$
$$\operatorname{ht}(\mathfrak{a}) = \operatorname{ht}(\mathfrak{a}/\mathfrak{p}) + \operatorname{ht}(\mathfrak{p})$$

*If in addition $A$ is equidimensional (e.g. an integral domain) then it is **biequidimensional** and satisfies for all ideals $\mathfrak{a}$*

$$\dim A = \dim \mathfrak{a} + \operatorname{ht}(\mathfrak{a})$$

*In particular for every prime ideal $\mathfrak{p}$ we have*

$$\dim A = \dim A_{\mathfrak{p}} + \dim A/\mathfrak{p}$$

*and for every maximal ideal*

$$\dim A = \dim A_{\mathfrak{m}}$$

*Proof.* By (3.35.34) $A$ satisfies the criteria in (3.22.10).e) and so is quasi-biequidimensional. The codimension formulas follow from (3.22.12). ■

We may also consider the following result

**Proposition 3.35.36 (Generalized Hauptidealsatz for Affine Algebras)**
*Let $A$ be an affine algebra and $\mathfrak{p}$ a prime ideal. Then*

*a) If $\operatorname{ht}(\mathfrak{p}) = n$ then it is minimal over some ideal $\mathfrak{a} := (x_1, ..., x_n)$. Furthermore $\mathfrak{a}$ may be chosen such that $\operatorname{ht}(\mathfrak{a}) = n$ and every minimal prime is of height $n$*
*b) We have the following characterization of height of a prime ideal*

$$\operatorname{ht}(\mathfrak{p}) = \min\{n \mid \mathfrak{p} \text{ minimal over } (x_1, ..., x_n)\}$$

*In particular if $\mathfrak{p}$ is minimal over $(x_1, ..., x_n)$ then $\operatorname{ht}(\mathfrak{p}) \leq n$.*

*Furthermore the same result holds for localization of $A$ at any prime ideal.*

*Proof.* This is largely a restatement of results in the Hauptidealsatz section. By (3.35.32) we have $A/\mathfrak{p}$ is hauptidealsatz for every prime ideal $\mathfrak{p}$. Furthermore by (3.35.35) $A$ is quasi-biequidimensional and so catenary (3.22.10). Then by (3.27.4) this means $A$ is a generalized hauptidealsatz ring (and so is every localization $A_{\mathfrak{p}}$).

Then a) and b) follows from (3.27.12). ■

**Corollary 3.35.37**
*Let $A$ be an affine algebra and $\mathfrak{p}$ a prime ideal. Denote the unique maximal ideal of $A_{\mathfrak{p}}$ by $\mathfrak{m} := \mathfrak{p}A_{\mathfrak{p}}$. Then*

$$\dim A_{\mathfrak{p}} = \min\left\{n \;\middle|\; \sqrt{(x_1, ..., x_n)} = \mathfrak{m}\right\} \leq \dim_{k(\mathfrak{m})} \mathfrak{m}/\mathfrak{m}^2$$

*with equality iff $A_{\mathfrak{p}}$ is a regular local ring.*

*Proof.* By (3.35.36) $A_{\mathfrak{p}}$ is a Noetherian local ring satisfying the generalized hauptidealsatz. Therefore the result follows by (3.28.2). ■

### 3.35.5.1 * Biequidimensionality by Strong Normalisation *

We may prove more directly the biequidimensionality property by using the strong form of the Normalisation Lemma (3.35.12) and Going Down (3.24.19). First we prove a technical result

**Lemma 3.35.38 (Saturated pairs)**
*Let $\phi : B \to A$ be an integral map and $\mathfrak{p}_0 \subsetneq \mathfrak{p}_1$ prime ideals lying over $\mathfrak{q}_0 \subsetneq \mathfrak{q}_1$. Then*

*a) $\mathfrak{q}_0 \subsetneq \mathfrak{q}_1$ saturated $\implies \mathfrak{p}_0 \subsetneq \mathfrak{p}_1$ saturated.*
*b) $B/\mathfrak{q}_0$ integrally closed and $\mathfrak{p}_0 \subsetneq \mathfrak{p}_1$ saturated $\implies \mathfrak{q}_0 \subsetneq \mathfrak{q}_1$ saturated*

*We may relax the condition in b) to the existence of another integral map $\psi : C \to B$ such that $C/\psi^{-1}(\mathfrak{q}_0)$ is integrally closed.*

© David Rufino CC BY-NC-ND 4.0

*Proof.* The first follows by incomparability (3.24.17). The second follows by applying Going Down (3.24.19) to the integral map $B/\mathfrak{q}_0 \hookrightarrow A/\mathfrak{p}_0$. More precisely if $\mathfrak{q}_0 \subsetneq \mathfrak{q} \subsetneq \mathfrak{q}_1$ then $(0) \subsetneq \mathfrak{q}/\mathfrak{q}_0 \subsetneq \mathfrak{q}_1/\mathfrak{q}_0$ whence there exists $\mathfrak{p}$ such that $(0) \subsetneq \mathfrak{p}/\mathfrak{p}_0 \subsetneq \mathfrak{p}_1/\mathfrak{p}_0$. This means $\mathfrak{p}_0 \subsetneq \mathfrak{p} \subsetneq \mathfrak{p}_1$, a contradiction.

The final statement can be demonstrated by applying b) to $C \to A$ and then a) to $B \to A$. ∎

**Proposition 3.35.39**
*Let $A$ be an affine domain, then every maximal chain has order $n = \dim A$, i.e. $A$ is **irreducible** and **biequidimensional**.*

*Proof of (3.35.39).* Consider a maximal chain $\mathfrak{p}_0 \subsetneq \mathfrak{p}_1 \subsetneq ... \subsetneq \mathfrak{p}_m$. Clearly $\mathfrak{p}_m$ is maximal by (3.3.37), and as $A$ is integral $\mathfrak{p}_0 = 0$. Apply (3.35.12) to find an integral, injective map

$$\phi : k[X_1, ..., X_n] \hookrightarrow A$$

such that

$$\mathfrak{q}_i := \phi^{-1}(\mathfrak{p}_i) = \left(X_1, ..., X_{h(i)}\right) \quad 0 \leq i \leq m$$

Note $n = \dim A$ by (3.35.28). Clearly $h(0) = 0$ and by (3.24.16) $\mathfrak{q}_m$ is maximal so $h(m) = n$. Observe that

$$k[X_1, ..., X_n]/\mathfrak{q}_i \xrightarrow{\sim} k\left[X_{h(i)+1}, ..., X_n\right]$$

is integrally closed for all $i$ (3.24.12). Therefore we may apply (3.35.38) to $\phi$ and each pair $\mathfrak{p}_i \subsetneq \mathfrak{p}_{i+1}$, to see that each chain $\mathfrak{q}_i \subsetneq \mathfrak{q}_{i+1}$ is saturated. This can only happen if $h(j) = j$ and therefore $m = n$. ∎

## 3.35.6 Derivations of Affine Algebras

The notion of derivations is a useful as a coordinate-free construction of the "tangent space" for both differentiable manifolds and algebraic varieties. We review the theory of derivations here primarily focusing on fields and f.g. $k$-algebras.

**Proposition 3.35.40 (Derivations of Polynomial Algebra)**
*Let $A = k[X_1, ..., X_n]$ be a polynomial algebra and $M$ an $(A, B)$-bimodule. Then we have an $(A, B)$-bimodule isomorphism*

$$\begin{aligned} \mathrm{Der}_k(A, M) &\cong M^n \\ D &\to (D(X_1), ..., D(X_n)) \\ \sum_{i=1}^{n} \frac{\partial}{\partial X_i} \cdot v_i &\leftarrow v \end{aligned}$$

*When $M$ has a $B$-basis $\{m_1, ..., m_r\}$ then $\mathrm{Der}_k(A, M)$ also has a $B$-basis*

$$\left\{ \frac{\partial}{\partial X_i} \cdot m_j \right\}_{i=1...n, j=1...r}$$

*In particular $\dim_K \mathrm{Der}_k(A, K) = n$.*

*Proof.* The first map is clearly well-defined and by (...) injective. The inverse map is well-defined because we have shown $\frac{\partial}{\partial X_i} \in \mathrm{Der}_k(A, A)$. The chain rule (3.26.8) shows that the maps are mutually inverse (the other direction following simply from orthogonality of the partial derivatives).

The final statement is straightforward. ∎

We may generalise this as follows, so that we may interpret derivations as tangent vectors

**Proposition 3.35.41 (Derivations = Tangent Vectors)**
*Let $A = k[X_1, ..., X_n]/\mathfrak{a}$ be a f.g. $k$-algebra and $M$ an $A$-module. Then we have an $A$-module isomorphism*

$$\begin{aligned} \mathrm{Der}_k(A, M) \cong & \left\{ v \in M^n \;\middle|\; \sum_{i=1}^{n} \overline{\frac{\partial F}{\partial X_i}} v_i = 0 \quad \forall F \in \mathfrak{a} \right\} \subset M^n \\ & D \longrightarrow \left(D\left(\overline{X}_1\right), ..., D\left(\overline{X}_n\right)\right) \\ \sum_{i=1}^{n} & \overline{\frac{\partial}{\partial X_i}} \cdot v_i \longleftarrow v \end{aligned}$$

© David Rufino CC BY-NC-ND 4.0

*If $\mathfrak{a} = \langle F_1, ..., F_m \rangle$ then this is equal to the kernel of the $A$-module homomorphism*

$$\begin{pmatrix} \overline{\frac{\partial F_1}{\partial X_1}} & \cdots & \overline{\frac{\partial F_1}{\partial X_n}} \\ \vdots & \ddots & \vdots \\ \overline{\frac{\partial F_m}{\partial X_1}} & \cdots & \overline{\frac{\partial F_m}{\partial X_n}} \end{pmatrix} : M^n \to M^m$$

*When $M$ is an $(A, B)$-bimodule then this is also an $(A, B)$-bimodule isomorphism.*

*Proof.* According to (3.26.8) we have the following relationship for all $F \in k[X_1, ..., X_n]$

$$D(\overline{F}) = D\big(F(\overline{X}_1, ..., \overline{X}_n)\big) = \sum_{i=1}^{n} \frac{\partial F}{\partial X_i}(\overline{X}_i) D(\overline{X}_i) = \sum_{i=1}^{n} \overline{\frac{\partial F}{\partial X_i}} D(\overline{X}_i)$$

When $F \in \mathfrak{a}$ we have $\overline{F} = 0 \implies D(\overline{F}) = 0$, whence the first map is well-defined. Similarly for $v \in \text{RHS}$ we see by definition that

$$\sum_{i=1}^{n} \frac{\partial}{\partial X_i} \cdot v_i$$

is zero on $\mathfrak{a}$, so determines a well-defined derivation on $A$. In the same way we see that the maps are mutually inverse.

Suppose $v$ is in the kernel of the given matrix. For any $G \in \mathfrak{a}$ we have by hypothesis $G = \sum_{j=1}^{m} F_j H_j$ for some $H_j \in k[X_1, ..., X_n]$. Then

$$\begin{aligned} \sum_{i=1}^{n} \overline{\frac{\partial G}{\partial X_i}} v_i &= \sum_{i=1}^{n} \sum_{j=1}^{m} \left( \overline{F}_j \overline{\frac{\partial H_j}{\partial X_i}} + \overline{H}_j \overline{\frac{\partial F_j}{\partial X_i}} \right) v_i \\ &= \sum_{j=1}^{m} \sum_{i=1}^{n} \overline{\frac{\partial F_j}{\partial X_i}} v_i \\ &= 0 \end{aligned}$$

where we have used the fact that $\overline{F_j} = 0$. As $G$ was arbitrary this shows $v$ is in the right-hand side of the isomorphism, and the reverse inclusion is immediate as $F_j \in \mathfrak{a}$. ∎

**Proposition 3.35.42 (Lifting for separable algebraic extensions)**
*Let $L/K$ be a separable algebraic (resp. separably generated) extension (over $k$) and $V$ an $L$-module. Then there is an isomorphism (resp. epimorphism) of $L$-modules*

$$\begin{aligned} \text{Der}_k(L, V) &\xrightarrow{\sim} \text{Der}_k(K, V) \\ D &\to D|_K \end{aligned}$$

*Proof.* Consider $L[V]$ the $k$-algebra with ideal $N := 0 \times V$ such that $N^2 = 0$ and $L[V]/N \cong L$. Furthermore a derivation $D : K \to V$ corresponds to a unique $k$-algebra homomorphism $\phi_D : K \to K[V] \hookrightarrow L[V]$ by (3.26.5) so we have the following commutative diagram

$$\begin{array}{ccc} K & \xrightarrow{\phi_D} & L[V] \\ \downarrow & \nearrow & \downarrow \pi_1 \\ L & = & L \end{array}$$

In the algebraic case by (3.26.16) there is a unique homomorphism $\phi_{\tilde{D}} : L \to L[V]$ completing the diagram, and in particular by (3.26.5) there is a derivation $\tilde{D} : L \to V$ extending $D$. Uniqueness follows from that of $\phi_{\tilde{D}}$.

The separably generated case is similar, except without uniqueness. ∎

**Corollary 3.35.43**
*Let $K/k$ be a separably generated extension, $A$ a $K$-algebra and $M$ an $A$-module then*

$$\text{Der}_k(A, M) = \text{Der}_K(A, M)$$

*Proof.* Recall (3.26.1) that a derivation $D$ is $K$-linear iff it vanishes on $K$. So we may reduce to the case $A = K$ by considering the restriction $D|_K$.

© David Rufino CC BY-NC-ND 4.0

Therefore it's sufficient to show that $\mathrm{Der}_k(K,M) = \mathrm{Der}_K(K,M) = 0 = \mathrm{Der}_k(k,M)$, but this follows immediately from (3.35.42). ∎

**Proposition 3.35.44 (Transcendence Degree Controls Derivations into Any Module)**
*Suppose $K/k$ is **separably generated** (e.g. if $k$ is perfect and $K$ finitely generated), with transcendence basis $\eta_1, ..., \eta_n$, $n = \mathrm{trdeg}_k(K)$, and let $M$ be a $K$-module. Then*

$$\mathrm{Der}_k(K,M) \cong M^n, \quad D \mapsto (D(\eta_1), ..., D(\eta_n))$$

*as $K$-modules. In particular, taking $M := K$, $\mathrm{trdeg}_k(K) = \dim_K \mathrm{Der}_k(K)$.*

*Proof.* By hypothesis $K/k(\eta_1, ..., \eta_n)$ is algebraic and separable. Therefore we have

$$\mathrm{Der}_k(K,M) \overset{(3.35.42)}{\cong} \mathrm{Der}_k(k(\eta_1, ..., \eta_n), M) \overset{(3.26.10)}{\cong} \mathrm{Der}_k(k[\eta_1, ..., \eta_n], M) \overset{(3.35.40)}{\cong} M^n$$

the last isomorphism sending $D \mapsto (D(\eta_1), ..., D(\eta_n))$, matching the same map once restricted back along the two previous isomorphisms (both restriction maps, $D \mapsto D|_{k(\eta_1,...,\eta_n)}$ and $D \mapsto D|_{k[\eta_1,...,\eta_n]}$, fix the value at each $\eta_i$). Taking $M := K$ gives $\dim_K \mathrm{Der}_k(K,K) = n = \mathrm{trdeg}_k(K)$. ∎

**Proposition 3.35.45 (Derivations Extend Uniquely Along a Field Extension)**
*Let $K/k$ be a field extension, $k'/k$ any field extension, and $D \in \mathrm{Der}_k(K,K)$. There is a unique $\tilde{D} \in \mathrm{Der}_{k'}(K \otimes_k k', K \otimes_k k')$ with $\tilde{D}(x \otimes 1) = D(x) \otimes 1$ for every $x \in K$; explicitly, $\tilde{D}(x \otimes c) = D(x) \otimes c$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

**Existence.** Define $\tilde{D}(x \otimes c) := D(x) \otimes c$, extended $k'$-linearly to $K \otimes_k k'$ (well-defined as $D$ is $k$-linear). It satisfies the Leibniz rule on simple tensors, $\tilde{D}((x \otimes c)(y \otimes c')) = D(xy) \otimes cc' = (D(x)y + xD(y)) \otimes cc' = \tilde{D}(x \otimes c) \cdot (y \otimes c') + (x \otimes c) \cdot \tilde{D}(y \otimes c')$, hence on all of $K \otimes_k k'$ by bilinearity: $\tilde{D} \in \mathrm{Der}_{k'}(K \otimes_k k', K \otimes_k k')$.

**Uniqueness.** $K \otimes_k k'$ is $k'$-spanned by $K \otimes 1$ (extension of scalars along a $k$-basis of $K$), and any extension of $D$ is pinned down there ($x \otimes 1 \mapsto D(x) \otimes 1$ by definition of "extends"), so agrees with $\tilde{D}$ on a $k'$-spanning set, hence everywhere by $k'$-linearity. ∎

**Lemma 3.35.46**
*Let $A$ be a $k$-algebra and $\mathfrak{m}$ an ideal. Then $\mathfrak{m}/\mathfrak{m}^2$ is a $k(\mathfrak{m})$-module with action given by*

$$(a + \mathfrak{m}) \cdot (b + \mathfrak{m}^2) = (ab + \mathfrak{m}^2)$$

*for all $a \in A$ and $b \in \mathfrak{m}$.*

*Proof.* Suppose $a - a' \in \mathfrak{m}$ and $b - b' \in \mathfrak{m}^2$ then

$$ab - a'b' = (a - a')(b - b') + a'(b - b') + b'(a - a') \in \mathfrak{m}^2$$

this shows that the action is well-defined. The properties of a module are straightforward. ∎

**Lemma 3.35.47**
*Let $A$ be a $k$-algebra and $\mathfrak{m}$ a maximal ideal. Then there is an isomorphism of $A$-algebras*

$$\begin{array}{ccc} A & \longrightarrow & A/\mathfrak{m}^2 \\ \downarrow{\scriptstyle \pi_1} & \overset{\pi_2}{\searrow} & \downarrow \\ k(\mathfrak{m}) & \overset{\psi}{\underset{\sim}{\dashrightarrow}} & k(\mathfrak{m}/\mathfrak{m}^2) \end{array}$$

*Further there is an isomorphism of $A$-modules*

$$\begin{aligned} \mathrm{Der}_k(A/\mathfrak{m}^2, k(\mathfrak{m}/\mathfrak{m}^2)) &\to \mathrm{Der}_k(A, k(\mathfrak{m})) \\ D &\to x \to \psi^{-1}(D(\overline{x})) \end{aligned}$$

*Proof.* The first statement follows from (3.3.58) with $\mathfrak{a} = \mathfrak{m}^2$ and $\mathfrak{b} = \mathfrak{m}$.

Define $\hat{D}(x) := \psi^{-1}(D(\overline{x}))$. By hypothesis $D(\overline{xy}) = \pi_2(x)D(\overline{y}) + \pi_2(y)D(\overline{x})$. Therefore

$$\hat{D}(xy) = \pi_1(x)\psi^{-1}(D(\overline{y})) + \pi_1(y)\psi^{-1}(D(\overline{x})) = \pi_1(x)\hat{D}(\overline{y}) + \pi_1(y)\hat{D}(\overline{x})$$

© David Rufino CC BY-NC-ND 4.0

and so $\hat{D}$ is a well-defined derivation. Suppose $\hat{D} \in \mathrm{Der}_k(A, k(\mathfrak{m}))$ then by the product rule $\mathfrak{m}^2 \subset \ker\left(\hat{D}\right)$ so there is a derivation $D' : A/\mathfrak{m}^2 \to k(\mathfrak{m})$ such that $D'(\overline{x}) = \hat{D}(x)$. Define $D \coloneqq \psi \circ D'$, then evidentally the image of this derivation is $\hat{D}$ and so the map is surjective.

Suppose $\hat{D} = 0$ then as $\psi$ is an isomorphism $D \circ \bar{\cdot} = 0$. As the reduction map is surjective we deduce that $D = 0$. Therefore the map is injective, bijective and an isomorphism. ∎

**Lemma 3.35.48**
*Let $A$ be a $k$-algebra and $\mathfrak{m}$ an ideal. Then there is an isomorphism of $A$-modules*

$$\begin{aligned}\mathrm{Hom}_{k(\mathfrak{m})}(\mathfrak{m}/\mathfrak{m}^2, k(\mathfrak{m})) &\to \mathrm{Hom}_{k(\mathfrak{m}/\mathfrak{m}^2)}((\mathfrak{m}/\mathfrak{m}^2)/(\mathfrak{m}^2/\mathfrak{m}^2), k(\mathfrak{m}/\mathfrak{m}^2)) \\ \theta &\to \psi \circ \theta \circ i^{-1}\end{aligned}$$

*where $i : \mathfrak{m}/\mathfrak{m}^2 \to (\mathfrak{m}/\mathfrak{m}^2)/(\mathfrak{m}^2/\mathfrak{m}^2)$ is a ring isomorphism such that*

$$i(\pi_1(a) \cdot x) = \pi_2(a) \cdot i(x)$$

*Proof.* Then for $x \in (\mathfrak{m}/\mathfrak{m}^2)/(\mathfrak{m}^2/\mathfrak{m}^2)$, $a \in A$ and $\theta \in \mathrm{Hom}_{k(\mathfrak{m})}(\mathfrak{m}/\mathfrak{m}^2, k(\mathfrak{m}))$

$$\begin{aligned}(\psi \circ \theta \circ i^{-1})(\pi_2(a) \cdot x) &= (\psi \circ \theta)(\pi_1(a) \cdot i^{-1}(x)) \\ &= \psi(\pi_1(a)\theta(i^{-1}(x))) \\ &= \pi_2(a)(\psi \circ \theta \circ i^{-1})(x)\end{aligned}$$

so that the map is well-defined. Similarly if $\theta' \in \mathrm{Hom}_{k(\mathfrak{m}/\mathfrak{m}^2)}((\mathfrak{m}/\mathfrak{m}^2)/(\mathfrak{m}^2/\mathfrak{m}^2), k(\mathfrak{m}/\mathfrak{m}^2))$

$$\begin{aligned}(\psi^{-1} \circ \theta' \circ i)(\pi_1(a) \cdot x) &= (\psi^{-1} \circ \theta')(\pi_2(a) \cdot i(x)) \\ &= \psi^{-1}(\pi_2(a)\theta'(i(x))) \\ &= \pi_1(a)(\psi^{-1} \circ \theta' \circ i)(x)\end{aligned}$$

so the inverse map is well-defined. ∎

**Proposition 3.35.49 (Coefficient Field mod $\mathfrak{m}^2$)**
*Let $A$ be a $k$-algebra and $\mathfrak{m} \triangleleft A$ a maximal ideal with $\mathfrak{m}^2 = 0$ and residue field $k(\mathfrak{m}) \coloneqq A/\mathfrak{m}$. If $k(\mathfrak{m})/k$ is separably generated (e.g. if $k$ is perfect and $k(\mathfrak{m})/k$ is finitely generated), then the quotient map $\pi : A \to k(\mathfrak{m})$ admits a $k$-algebra section $j : k(\mathfrak{m}) \to A$, that is there is a subfield $k \subseteq \hat{k} \subseteq A$ such that $\pi|_{\hat{k}} : \hat{k} \to k(\mathfrak{m})$ is an isomorphism. In particular $A = \hat{k} \oplus \mathfrak{m}$ as $k$-vector spaces.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

By (3.26.17) $k(\mathfrak{m})$ is a formally smooth $k$-algebra. Apply (3.26.11)'s lifting property with $C \coloneqq A$, $N \coloneqq \mathfrak{m}$ (satisfying $N^2 = 0$ by hypothesis) and $v : k(\mathfrak{m}) \to A/\mathfrak{m} = C/N$ the identity map: this produces a $k$-algebra homomorphism $j : k(\mathfrak{m}) \to A$ with $\pi \circ j = \mathrm{id}_{k(\mathfrak{m})}$.

Set $\hat{k} \coloneqq j(k(\mathfrak{m})) \subseteq A$. As $j$ is a ring homomorphism from a field, it is injective (3.3.61), so $j : k(\mathfrak{m}) \to \hat{k}$ is an isomorphism, and $\pi \circ j = \mathrm{id}$ shows $\pi|_{\hat{k}}$ is its inverse. Finally $j$ is a $k$-linear section of the $k$-linear surjection $\pi : A \to k(\mathfrak{m})$ with kernel $\mathfrak{m}$, so $A = \mathrm{Im}(j) \oplus \ker(\pi) = \hat{k} \oplus \mathfrak{m}$ as $k$-vector spaces. ∎

**Proposition 3.35.50 (Tangent space is dual to Cotangent space)**
*Let $A$ be a $k$-algebra and $\mathfrak{m} \triangleleft A$ a maximal ideal with residue field $k(\mathfrak{m}) \coloneqq A/\mathfrak{m}$. There is a homomorphism of $k(\mathfrak{m})$-modules*

$$\begin{aligned}\mathrm{Der}_k(A, k(\mathfrak{m})) &\longrightarrow \mathrm{Hom}_{k(\mathfrak{m})}(\mathfrak{m}/\mathfrak{m}^2, k(\mathfrak{m})) \\ D &\longrightarrow \theta_D : \overline{x} \to D(x)\end{aligned}$$

*When $k(\mathfrak{m})/k$ is **separably generated** then this is an isomorphism. In particular this holds when $k$ is perfect and $k(\mathfrak{m})/k$ is finitely generated.*

*Proof.* The map $\theta_D$ is well-defined because $D$ annihilates $\mathfrak{m}^2$ by the product rule. It remains to show the map is bijective when $k(\mathfrak{m})/k$ is separably generated.

We first claim we can reduce to the case $\mathfrak{m}^2 = 0$. For consider $A' \coloneqq A/\mathfrak{m}^2$ and $\mathfrak{m}' = \mathfrak{m}/\mathfrak{m}^2$ and the commutative diagram of $A$-module homomorphisms

© David Rufino CC BY-NC-ND 4.0

$$\begin{array}{ccc} \mathrm{Der}_k(A', k(\mathfrak{m}')) & \longrightarrow & \mathrm{Hom}_{k(\mathfrak{m}')}\left(\mathfrak{m}'/(\mathfrak{m}')^2, k(\mathfrak{m}')\right) \\ \downarrow \sim & & \downarrow \sim \\ \mathrm{Der}_k(A, k(\mathfrak{m})) & \longrightarrow & \mathrm{Hom}_{k(\mathfrak{m})}(\mathfrak{m}/\mathfrak{m}^2, k(\mathfrak{m})) \end{array}$$

The horizontal maps have already been defined and the vertical maps are defined in (3.35.47) and (3.35.48). The diagram evidentally commutes, so therefore it is sufficient to show that the top map is an isomorphism.

We revert to the original notation, but with the additional assumption that $\mathfrak{m}^2 = 0$. By (3.35.49) there is a subfield $k \subseteq \hat{k} \subseteq A$ mapping isomorphically to $k(\mathfrak{m})$ under the canonical projection $\pi : A \to A/\mathfrak{m} = k(\mathfrak{m})$, giving a direct sum of $k$-vector spaces

$$A = \hat{k} \oplus \mathfrak{m}$$

Given $\theta \in \mathrm{Hom}_{k(\mathfrak{m})}(\mathfrak{m}, k(\mathfrak{m}))$ define

$$D_\theta(\lambda + x) := \theta(x) \quad \lambda \in \hat{k}, x \in \mathfrak{m}$$

It satisfies the product rule for

$$\begin{aligned} D_\theta((\lambda + x)(\lambda' + x')) &= \theta(\lambda' x + \lambda x') \\ &= \pi(\lambda')\theta(x) + \pi(\lambda)\theta(x') \\ &= \pi(\lambda' + x')\theta(x) + \pi(\lambda + x)\theta(x') \\ &= (\lambda' + x') \cdot D_\theta(\lambda + x) + (\lambda + x) \cdot D_\theta(\lambda' + x') \end{aligned}$$

Therefore the given map is surjective.

As $k(\mathfrak{m})/k$ is separably generated then by (3.35.43) any derivation $D$ is $\hat{k}$-linear. Therefore

$$D(\lambda + x) = D|_{\mathfrak{m}}(x) \quad \lambda \in \hat{k}, x \in \mathfrak{m}$$

which shows that the given map is injective. ∎

**Remark 3.35.51**
*The proof is substantially simpler when $k(\mathfrak{m}) = k$, for example when $k$ is algebraically closed.*

*The argument follows the suggestion of [Har13] (Ex 8.1), but using the simpler result [Mat70] (Prop 28.I) to argue more directly.*

## 3.35.7 Linearly Disjoint Algebras

**Definition 3.35.52**
*Let $A/k$, $B/k$ be $k$-algebras and homomorphisms $i : A/k \to \Omega/k$, $j : B/k \to \Omega/k$ (typically being inclusion). Then we say that $A$ and $B$ are **linearly disjoint in $\Omega$** if the canonical map*

$$\begin{aligned} A \otimes_k B &\to \Omega \\ a \otimes b &\to i(a)j(b) \end{aligned}$$

*is injective. As the canonical maps $A, B \to A \otimes_k B$ are injective by (3.4.41) this implies $i, j$ must be injective. In the case that $\Omega$ is integral (resp. reduced) then $A, B$ are also necessarily integral (resp. reduced).*

*Recall that every $k$-module is free (3.3.126), moreover by (3.4.23) a basis of $A \otimes_k B$ may be formed by tensor product of bases for $A$ and $B$.*

**Example 3.35.53**
*Let $K/k$ be a non-trivial field extension then $K$ is not linearly disjoint with itself. For $\{1, x\}$ a linearly independent subset of $K$ which may be extended to a basis. Given the remark on the basis of the tensor product we see $1 \otimes x \neq x \otimes 1$. On the other hand the image of $1 \otimes x - x \otimes 1$ in $\Omega$ is clearly $0$. The same consideration shows that $A \cap B = k$.*

**Remark 3.35.54**
*The notion of linear disjointness in general depends on the embeddings in $\Omega/k$. For example $k(t)$ and $k(s)$ are linearly disjoint in $k(s,t)$, but not when both are identified with $k(x)$.*

**Proposition 3.35.55**
*Let $i : A/k \to \Omega/k$ and $j : B/k \to \Omega/k$ be algebra homomorphisms. Then the following are equivalent*

© David Rufino CC BY-NC-ND 4.0

- ***a)*** *$A/k$ and $B/k$ are linearly disjoint in $\Omega/k$*
- ***b)*** *If $S \subset A$ is linearly independent over $k$ then $i(S)$ is linearly independent over $B$*
- ***b')*** *If $T \subset B$ is linearly independent over $k$ then $j(T)$ is linearly independent over $A$*
- ***c)*** *There is a $k$-basis $\{a_\lambda\}_{\lambda\in\Lambda}$ of $A$ for which $\{i(a_\lambda)\}_{\lambda\in\Lambda}$ is linearly independent over $B$*
- ***c')*** *There is a $k$-basis $\{b_\lambda\}_{\lambda\in\Lambda}$ of $B$ for which $\{j(b_\lambda)\}_{\lambda\in\Lambda}$ is linearly independent over $A$*

*Proof.* $a) \implies b)$ By (3.4.30) and (3.4.31) there is an injective map

$$B^{(S)} \hookrightarrow A \otimes_k B$$
$$(b_s)_{s\in S} \to \sum_{s\in S} s \otimes b_s$$

By hypothesis the composite with $A \otimes_k B \hookrightarrow \Omega$ is injective which is precisely the required conclusion.

$b) \implies c)$ By (3.3.126) there exists a basis which by hypothesis is linearly independent over $B$

$c) \implies a)$ By (3.4.23) there is an isomorphism

$$B^{(\Lambda)} \cong A \otimes_k B$$

and the canonical map into $\Omega$ is identified with $(b_\lambda)_{\lambda\in\Lambda} \to \sum i(a_\lambda)j(b_\lambda)$. The hypothesis means precisely that this map is injective.

We immediately have the symmetric result $a) \iff b') \iff c')$. ∎

In cases where both algebras are fields and one is algebraic then linear disjointness can be characterized by a property of the tensor product. In particular it does not depend on the **ambient** algebra $\Omega/k$.

**Lemma 3.35.56**
*Let $A$ be a finite-dimensional $k$-algebra with no zero-divisors. Then $A$ is a field.*

*Proof.* By assumption multiplication by a non-zero element $x$ is injective and $k$-linear. Therefore by (3.3.137) it is surjective and $x$ has an inverse as required. ∎

**Proposition 3.35.57**
*Let $K/k$ be a field extension and $k'/k$ an **algebraic** field extension. Then the following are equivalent*

*a) $K$ and $k'$ are linearly disjoint with respect to every embedding into an algebra $\Omega/k$*
*b) $K$ and $k'$ are linearly disjoint with respect to some common field extension $\Omega/k$*
*c) $K \otimes_k k'$ is an integral domain*
*d) $K \otimes_k k'$ is a field*

*Proof.* $a) \implies b)$ We may take $\Omega := (K \otimes_k k')/\mathfrak{m}$.

$b) \implies c)$ As $\Omega$ is assumed to be a field it is certainly integral, and therefore so is $K \otimes_k k'$ being a subring.

$c) \implies d)$ As every element of the tensor product is a finite linear combination of elementary tensors we may reduce to the case $k'$ is finitely generated. Then by (3.19.44) $k'$ is finite over $k$ and therefore by (3.4.23) $K \otimes_k k'$ is finite over $K$. By (3.35.56) it is a field.

$d) \implies a)$ This is immediate because the kernel is an ideal which must be (0). ∎

## 3.35.8 Finiteness of Integral Closure

**Remark 3.35.58**
*In what follows we are interested in the integral closure of affine domains in finite extensions of the quotient field. Similar results are obtained for Dedekind Domains (3.32.9) with the restriction that the field extension is separable. Although this holds over number fields where the characteristic is zero, this will not in general be the case for varieties over finite fields, as for example $\mathbb{F}_p(t)$ admits the inseparable extension $\mathbb{F}_p(t^{1/p})$. However the finiteness still holds, because it's easy to see that the integral closure of $\mathbb{F}_p[t]$ is simply $\mathbb{F}_p[t^{1/p}]$. This generalises because every purely inseparable extension is contained in an extension obtained by adjoining $p$-th roots of unity.*

**Proposition 3.35.59 (Normalisation over Finite Fields)**
*Let $k$ be a (possibly) finite field and $A = k[x_1, ..., x_n]$ be a finitely-generated $k$-algebra. Then there exists a normalising family $z_1, ..., z_d \in A$ such that $k[x_1, ..., x_n]$ is integral over $k[z_1, ..., z_d]$.*

*Proof.* TODO ∎

© David Rufino CC BY-NC-ND 4.0

**Proposition 3.35.60 (Affine Domain are Japanese Rings)**
*Let $A$ be an affine domain over $k$ with $K = \mathrm{Frac}(A)$ and $L/K$ a finite extension. Then $\overline{A}^L$ is a finite $A$-module and Noetherian with quotient field $L$.*

*Furthermore* $\dim \overline{A}^L = \dim A$.

*Proof.* Suppose $A = k[x_1, ...x_n]$, apply the normalisation lemma (3.35.59) to obtain a lattice of subrings

$$\begin{array}{c} L \\ | \\ K \\ | \\ A \\ | \\ A' := k[z_1, ..., z_d] \end{array}$$

where $z_1, ..., z_d$ are algebraically independent over $k$. We claim it is enough to show that the $\overline{A'}^L$ is a finite $A'$-module. For $A$ is finite and integral over $A'$. Therefore by (3.24.28) $\overline{A}^L = \overline{A'}^L$ is finite over $A'$, and a-fortiori over $A$.

Therefore without loss of generality we may assume that $x_1, ..., x_n$ are algebraically independent over $k$ and $A$ is integrally closed (3.24.12).

By (3.19.73) we may find a finite extension $L'/L$ such that $L'/K$ is normal. So by (3.24.29) we may assume without loss of generality that $L/K$ is normal.

By (...) we may decompose this into a tower $M/L/K$ such that $M/L$ is Galois and $L/K$ is purely inseparable.

We first show that the integral closure of $A$ in $L$ is a finite $A$-module. When $L = K$ this is trivial because $A$ is already integrally closed in its quotient field $K$. Therefore we may assume $L \neq K$ and the characteristic $p$ is non-zero (3.19.103). $L = K(y_1, ..., y_n)$ is finitely generated and by definition $y_i^q \in K$ for some $p$-th power $q$. Therefore $L$ is contained in $L' := k\left(x_1^{1/q}, ..., x_n^{1/q}\right)$ which is again a finite purely inseparable extension of $K$. The elements $x_1^{1/q}, ..., x_n^{1/q}$ are again algebraically independent so that the ring $A' := k\left[x_1^{1/q}, ..., x_n^{1/q}\right]$ is integrally closed (3.24.12) with quotient field $L'$. Evidently $A'$ is finite and integral over $A$ (3.24.4) therefore (3.24.28) $\overline{A}^{L'} = \overline{A'}^{L'} = A'$ is finite over $A$. By (3.24.29) then $\overline{A}^L$ is finite over $A$ and $\overline{A}^L$ has quotient field $L$ (3.24.24). As $A$ is Noetherian then so is $\overline{A}^L$.

By transitivity we may then reduce to the case $L/K$ is separable which was proven in (3.24.26). ∎

**Corollary 3.35.61**
*Let $A$ be an affine domain of Krull dimension 1, with quotient field $K$ and $L/K$ a finite extension. Then the integral closure $\overline{A}^L$ is a finite $A$-module and a Dedekind domain.*

*Further for every prime ideal $\mathfrak{p}$ of $A$ we have the ramification formula*

$$\sum_{\mathfrak{B}|\mathfrak{p}} e_{\mathfrak{B}/\mathfrak{p}} f_{\mathfrak{B}/\mathfrak{p}} = [L : K]$$

*Proof.* By (3.35.60) $B := \overline{A}^L$ is a finite $A$-module with quotient field $L$ and Noetherian. By (...) it is integrally closed and by (3.24.22) has Krull Dimension 1. Therefore by definition $B$ is also a Dedekind domain.

The ramification formula follows from (3.32.16). ∎

© David Rufino CC BY-NC-ND 4.0

## 3.36 Affine Algebras under Base Change

This section covers what happens to a $k$-algebra $A$ under base change, that is $A_{(L)}$ for $L/k$ a field extension. Principally given an integral $k$-algebra $A$ we would like to ensure that $A_{(L)}$ is reduced (which is related to separability), irreducible and therefore integral. We follow [Stacks] but large parts of exposition are from [Bou89].

**Definition 3.36.1 (Geometrically Reduced / Integral Algebra)**
*A $k$-algebra $A$ is*

- ***geometrically integral*** *(resp.* ***irreducible***, ***reduced****) if the ring $A \otimes_k L$ is* ***integral*** *(resp.* ***irreducible***, ***reduced****) for every field extension $L/k$.*
- ***algebraically integral*** *(resp.* ***irreducible***, ***reduced****) if the ring $A \otimes_k \overline{k}$ is* ***integral*** *(resp.* ***irreducible***, ***reduced****).*

*A few remarks about the important results and notation in the literature*

- *As in (3.3.66) we see that* ***integral*** $\iff$ ***irreducible*** *and* ***reduced****.*
- *These are shown to be equivalent (3.36.26), (3.36.36) and (3.36.44). In fact it is sufficient to check the properties for all finite extensions.*
- *[Liu06] uses the "algebraic" form*
- *[Bou89] uses the term "separable" for "geometrically reduced", as this coincides with the notion of separable for algebraic extensions (3.36.19)*
- *Similarly [Bou89] and [Lan11] use the term "regular" for geometrically integral*

We make implicit use of the following results throughout

**Proposition 3.36.2**
*Let $A, B$ be $k$-algebras, then the structural morphisms $A \to A \otimes_k B$ and $B \to A \otimes_k B$ are injective. Furthermore if $A'$ and $B'$ are $k$-subalgebras then the canonical homomorphism*

$$A' \otimes_k B' \to A \otimes_k B$$

*is injective.*

*Proof.* This is (3.4.30) and (3.4.41) ∎

**Proposition 3.36.3 (A Subspace is Recovered by Intersection After Base Change)**
*Let $V \subseteq W$ be $k$-vector spaces and $k'/k$ a field extension. Then, identifying $W$ with its image in $W \otimes_k k'$ via $w \mapsto w \otimes 1$ (injective, (3.4.31)),*

$$(V \otimes_k k') \cap W = V$$

*inside $W \otimes_k k'$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

"$\supseteq$" is clear. For "$\subseteq$": choose a $k$-subspace complement $V'$ of $V$ in $W$ ((3.3.139), $k$ being a field), so $W = V \oplus V'$; tensoring, $W \otimes_k k' \cong (V \otimes_k k') \oplus (V' \otimes_k k')$ (3.4.21). Given $w \in W$ with $w \otimes 1 \in V \otimes_k k'$: writing $w = v + v'$ ($v \in V$, $v' \in V'$) for its decomposition, $w \otimes 1 = (v \otimes 1) + (v' \otimes 1)$ under this direct sum splitting; as $w \otimes 1$ lies in the first summand, the second component vanishes, that is $v' \otimes 1 = 0$; as $v' \mapsto v' \otimes 1$ is injective (3.4.31), $v' = 0$, giving $w = v \in V$. ∎

**Proposition 3.36.4**
*Let $A, B$ be $k$-algebras. Let $\{a_i\}_{i \in I}$ be a $k$-linearly independent subset (resp. basis) of $A$ and $\{b_j\}_{j \in I}$ be a $k$-linearly independent subset (resp. basis) of $B$. Then $\{a_i \otimes b_j\}_{(i,j) \in I \times J}$ is a $k$-linearly independent subset (resp. basis) of $A \otimes_k B$.*

*Furthermore $\{a_i \otimes 1\}_{i \in I}$ is a $B$-linearly independent subset (resp. basis) of $A \otimes_k B$.*

*Proof.* The first statement is (3.4.31). For the last statement extend $\{a_i\}_{i \in I}$ to a basis (2.3.7) and apply (3.4.26). ∎

**Proposition 3.36.5**
*Let $A, B$ be $k$-algebras. Then $A \otimes_k B$ is the union of subrings of the form a) $A \otimes_k B'$, b) $A' \otimes_k B$ or c) $A' \otimes_k B'$ where $A'$ and $B'$ are finitely-generated $k$-subalgebras of $A$ and $B$ respectively.*

© David Rufino CC BY-NC-ND 4.0

*If $K/k$ is an algebraic field extension then $A \otimes_k K$ is the union of subrings of the form $A \otimes_k k'$ where $k'/k$ are finite subextensions.*

*Proof.* An arbitrary element $z \in A \otimes_k B$ is a finite sum of elementary tensors. Therefore the result follows from (3.36.2).

The final statement follows from recalling finitely generated algebraic extensions are finite (3.19.44). ∎

*Proof.* ∎

## 3.36.1 Etale Algebras

**Definition 3.36.6**
*Let $A$ be a finite-dimensional $k$-algebra. Then define the **separable degree** as follows*

$$[A:k]_s := \#\operatorname{AlgHom}_k\left(A, \overline{k}\right)$$

*We say that $A$ is **etale** if this equals the dimension of $A$ as a $k$-vector space*

$$[A:k]_s = [A:k]$$

*Recall a finite field extension $K/k$ is **etale** if and only if it is separable (3.19.78).*

**Lemma 3.36.7**
*Let $A$ be a finite-dimensional $k$-algebra and $K/k$ a field extension. Then*

$$[\operatorname{Hom}_k(A,K):K] = [A^\vee : k] = [A:k]$$

*More precisely if $\{a_1, ..., a_n\}$ is a $k$-basis for $A$ then $\{a_1^\vee, ..., a_n^\vee\}$ is a $K$-basis for $\operatorname{Hom}_k(A,K)$.*

*Proof.* This is simply a special case of (3.4.27). ∎

**Corollary 3.36.8 (Separable degree is finite)**
*Let $A$ be a finite-dimensional $k$-algebra then $\#\operatorname{AlgHom}_k(A,K) \leq [A:k]$ for every field extension $K/k$.*

**Proposition 3.36.9 (Image of $A$ is finite-dimensional)**
*Let $A$ be a finite-dimensional $k$-algebra and $\Omega/k$ a field extension. Then there exists a finite subextension $K/k$ such that*

$$\operatorname{AlgHom}_k(A,K) = \operatorname{AlgHom}_k(A,\Omega)$$

*If $\Omega$ contains $\overline{k}$ then these are both equal to $\operatorname{AlgHom}_k\left(A, \overline{k}\right)$.*

*Proof.* By (...) $\phi(A)$ has finite degree over $k$ and therefore finitely-generated and algebraic as a $k$-algebra. Therefore $\phi(A) = k[a_1, ..., a_n] = k(a_1, ..., a_n)$ is a finite-dimensional field by (3.19.44).

We may take the compositum of all these images (since there are only finitely many) to obtain the finite extension $K/k$.

If $\Omega$ contains $\overline{k}$ then trivially

$$\operatorname{AlgHom}_k(A,K) \subseteq \operatorname{AlgHom}_k\left(A,\overline{k}\right) \subseteq \operatorname{AlgHom}_k(A,\Omega)$$

whence they are all equal. ∎

**Proposition 3.36.10 (Etale-ness is preserved under base extension)**
*Let $A$ be a finite-dimensional $k$-algebra and $K/k$ a field extension then*

$$[A \otimes_k K : K] = [A:k]$$
$$[A \otimes_k K : K]_s = [A:k]_s$$

*In particular $A$ is etale if and only if $A \otimes_k K$ is.*

*Proof.* The first equality follows from (3.4.26) and (3.3.109).

For the second equality we see that by (3.36.9) and (3.4.38)

$$\operatorname{AlgHom}_k\left(A,\overline{k}\right) = \operatorname{AlgHom}_k\left(A,\overline{K}\right) \cong \operatorname{AlgHom}_K\left(A \otimes_k K, \overline{K}\right)$$

∎

**Lemma 3.36.11**
*Let $A = A_1 \times \cdots \times A_d$ be a finite product of finite-dimensional $k$-algebras. Then $A$ is etale if and only if each $A_i$ is.*

© David Rufino CC BY-NC-ND 4.0

*Proof.* This follows because

$$\mathrm{AlgHom}_k\left(A_1 \times \cdots \times A_d, \overline{k}\right) \cong \mathrm{AlgHom}_k\left(A_1, \overline{k}\right) \times \cdots \times \mathrm{AlgHom}_k\left(A_d, \overline{k}\right)$$

and a similar consideration for the dimension over $k$. ∎

**Lemma 3.36.12**
*Let $A$ be a finite-dimensional $k$-algebra. The following are equivalent*

*a) $A$ is reduced*
*b) $A \cong L_1 \times \cdots \times L_n$ for some finite extensions $L_i/k$*

*Further $A$ is etale if and only if each $L_i/k$ is separable. If $k$ is perfect then $A$ is automatically etale.*

*Proof.* One direction is obvious. Suppose conversely that $A$ is reduced. If $\dim_k A = 1$ then any proper ideal must have strictly lower dimension and therefore $A$ is a field. It is therefore sufficient to show (by induction on dimension) that if $A$ is not a field then $A \cong A_1 \times A_2$ for two (non-zero) $k$-algebras $A_1, A_2$.

Suppose $A$ is not a field and let $\mathfrak{a}$ be a non-zero proper ideal of $A$ which has minimal dimension over $k$ as a vector space. As $A$ is reduced, and by minimality, we have $\mathfrak{a}^2 = \mathfrak{a}$. By Nakayama's Lemma (3.21.5) there is $e \in \mathfrak{a}$ such that $ex = x$ for all $x \in \mathfrak{a}$ and in particular $e^2 = e$ and $\mathfrak{a} = Ae$. By assumption $e \neq 0, 1$. Define $f := 1 - e \neq 0, 1$ then evidently $f^2 = f$ and $ef = 0$. The homomorphism of $A$-modules

$$A \to Ae \times Af$$
$$a \to (ae, af)$$

is injective because $ae + af = a$ and surjective because the image of $a_1 e + a_2 f$ is $(a_1 e, a_2 f)$. The sub-modules $Ae$ and $Af$ are actually sub-rings by the idempotence property and the given map is an isomorphism of rings.

The last statement follows from (3.36.11) and (3.19.103). ∎

**Corollary 3.36.13 (Primes of a Finite Product of Fields)**
*Let $A \cong L_1 \times \cdots \times L_n$ be a finite product of fields (in particular, any reduced finite-dimensional $k$-algebra, (3.36.12)). Then every prime ideal of $A$ is maximal, and there are exactly $n$ of them.*

*Proof.* For each $j$, $\mathfrak{m}_j := \ker(\pi_j)$ (the kernel of the $j$-th projection $\pi_j : A \twoheadrightarrow L_j$) is maximal, being the kernel of a surjection onto a field (3.3.59), and the $\mathfrak{m}_j$ are pairwise distinct (the element with 1 in coordinate $j$ and 0 elsewhere lies in every $\mathfrak{m}_i$, $i \neq j$, but not $\mathfrak{m}_j$), giving $n$ maximal ideals.

Let $\mathfrak{p}$ be any prime. As $\bigcap_j \mathfrak{m}_j = \ker(\pi_1, ..., \pi_n) = 0 \subseteq \mathfrak{p}$, (3.3.40) gives $\mathfrak{m}_j \subseteq \mathfrak{p}$ for some $j$; as $\mathfrak{m}_j$ is already maximal, $\mathfrak{p} = \mathfrak{m}_j$. ∎

**Lemma 3.36.14**
*Let $A$ be a finite-dimensional $k$-algebra and $\Omega/k$ a field extension containing $\overline{k}$. Then the following are equivalent*

*a) $A$ is etale*
*b) $A \otimes_k \Omega \cong \Omega^n$ as an $\Omega$-algebra*

*Proof.* *a)* $\implies$ *b)* By (3.36.9) $\#\,\mathrm{AlgHom}_k(A, \Omega) = \#\,\mathrm{AlgHom}_k\left(A, \overline{k}\right) =: [A : k]_s = [A : k]$ with the last equality because $A$ is assumed etale. Then defining $B := A \otimes_k \Omega$

$$\#\,\mathrm{AlgHom}_\Omega(B, \Omega) \overset{(3.4.38)}{=} \#\,\mathrm{AlgHom}_k(A, \Omega) = [A : k] \overset{(3.4.16)}{=} [B : \Omega] = [B^\vee : \Omega]$$

By (3.19.130) and (3.3.127) we see that $\mathrm{AlgHom}_\Omega(B, \Omega)$ is a $\Omega$-basis for $B^\vee$. Denote this basis by $\varphi_1, ..., \varphi_n$ then by (3.3.109) this corresponds to a basis $e_1, ..., e_n$ of $B$. Then the algebra isomorphism is exhibited explicitly by

$$B \cong \Omega^n$$
$$b \to (\varphi_1(b), ..., \varphi_n(b))$$

*b)* $\implies$ *a)* We may show that the algebra homomorphisms are just the projections $\pi_i : \Omega^n \to \Omega$, so that $\#\,\mathrm{AlgHom}_\Omega(B, \Omega) = [B : \Omega]$. As before then $[A : k]_s = \#\,\mathrm{AlgHom}_k\left(A, \overline{k}\right) = \#\,\mathrm{AlgHom}_k(A, \Omega) = \#\,\mathrm{AlgHom}_\Omega(B, \Omega) = [B : \Omega] = [A : k]$ which shows that $A$ is etale. ∎

**Proposition 3.36.15 (A Field Splits Completely Over Any Normal Extension it Embeds Into)**
*Let $K/k$ be a finite separable field extension and $L/k$ a field extension containing (a copy of) a normal closure of $K/k$. Fix a primitive element $\theta$ for $K/k$ ((3.19.105)), with minimal polynomial $m \in k[X]$ of degree $n := [K : k]$, and write $m = \prod_{i=1}^n (X - \beta_i)$ for its (pairwise distinct) roots $\beta_i \in L$. Then there is a canonical isomorphism of $L$-algebras*

© David Rufino CC BY-NC-ND 4.0

$$K \otimes_k L \cong L^n, \quad p(\theta) \otimes c \mapsto (c\,p(\beta_1), ..., c\,p(\beta_n)) \quad (p \in k[X], c \in L)$$

*In particular $K \otimes_k L$ has exactly $n$ maximal ideals, $\mathfrak{m}_i$ the kernel of the $i$-th coordinate.*

*Proof.* As $K/k$ is separable, $m$ is separable (3.19.74). A normal closure $N$ of $K/k$ is, by definition, the splitting field of $m$ over $k$ (3.19.73), so $m$ splits completely already over $N$; as $N \subseteq L$ (by hypothesis), the same factorisation exhibits $m$ splitting completely over $L$ too, giving the $\beta_i \in N \subseteq L$, pairwise distinct as $m$ is separable. As $K \cong k[X]/(m)$ (3.19.40) ($\theta$ corresponding to the class of $X$), base-changing along $k \hookrightarrow L$ (using $K \otimes_k L \cong (k[X] \otimes_k L)/(m)$ as $L$-algebras, (3.4.42), and $k[X] \otimes_k L \cong L[X]$, (3.8.7)) gives $K \otimes_k L \cong L[X]/(m)$; as the $(X - \beta_i)$ are pairwise comaximal (distinct monic linear factors), the Chinese Remainder Theorem (3.3.62) gives

$$K \otimes_k L \cong L[X]/(m) \cong \prod_{i=1}^{n} L[X]/(X - \beta_i) \cong L^n$$

Tracing the elementary tensor $p(\theta) \otimes c$ through: it corresponds to $c \cdot p(X) \bmod m \in L[X]/(m)$ ((3.4.42), (3.8.7)), which under the Chinese Remainder Theorem's canonical map – reduction modulo each $(X - \beta_i)$, that is evaluation at $\beta_i$ – corresponds to $(c\,p(\beta_1), ..., c\,p(\beta_n))$. The maximal ideals of $L^n$ are exactly the $n$ coordinate kernels (3.36.13).■

**Proposition 3.36.16 (Etale Criteria)**
*Let $A$ be a finite-dimensional $k$-algebra. The following are equivalent*

*a) $A$ is etale*
*b) $A$ is geometrically reduced*
*c) $A \otimes_k \overline{k}$ is reduced*
*d) $A \otimes_k K$ is reduced for some perfect field extension $K/k$*
*e) $A \cong L_1 \times \cdots \times L_n$ for $L_i/k$ finite separable extensions*

*In particular $A$ is reduced.*

*Proof.* *a)* $\implies$ *b)* Consider a field extension $L/k$ and the tower of extensions $\overline{L}/L/k$. Then by (3.36.14) $A \otimes_k \overline{L} \cong \overline{L}^n$ is evidently reduced. By (...) $A \otimes_k L \subset A \otimes_k \overline{L}$ is a subring and therefore also reduced.

*b)* $\implies$ *c)* immediate.

*c)* $\implies$ *d)* immediate as $\overline{k}$ is perfect.

*d)* $\implies$ *a)* By (3.36.12) $A \otimes_k K$ is an etale $K$-algebra so by (3.36.10) $A$ is etale.

*e)* $\implies$ *a)* Follows immediately from (3.36.11).

*a)* $\implies$ *e)* We know that $A$ is reduced so the result follows from (3.36.12). ■

**Corollary 3.36.17**
*Let $A$ be a finite-dimensional $k$-algebra and $k$ perfect. Then the following are equivalent*

*a) $A$ is etale*
*b) $A$ is reduced*
*c) $A$ is geometrically reduced*
*d) $A \otimes_k \overline{k}$ is reduced.*

For completeness we summarize the relationship between the different notions of separability

**Corollary 3.36.18 (Equivalent definitions of separability)**
*Let $K/k$ be a finite extension. Then the following are equivalent*

*a) $K/k$ is separable algebraic (3.19.51)*
*b) $K/k$ is etale*
*c) $K/k$ is geometrically reduced*
*d) $K \otimes_k \overline{k}$ is reduced*

*When $k$ is perfect every field extension is separable.*

*Proof.* We have already shown *b)* $\iff$ *c)* $\iff$ *d)*. Furthermore *a)* $\iff$ *b)* is (3.19.78). ■

**Corollary 3.36.19 (Equivalent definitions of separability)**
*Let $K/k$ be an algebraic extension. Then the following are equivalent*

*a) $K/k$ is separable algebraic (3.19.51)*
*b) Every finite subextension of $K/k$ is etale*

© David Rufino CC BY-NC-ND 4.0

*c) $K/k$ is geometrically reduced*
*d) $K \otimes_k \overline{k}$ is reduced*

*In particular an algebraic extension of a perfect field $k$ is geometrically reduced.*

*Proof.* This follows directly from (3.36.18) because each property has "finite character", i.e. holds if and only if it holds for every finite subextension.

The last statement follows from (3.19.103). ∎

## 3.36.2 Geometrically Reduced Algebras

Reference is [Bou89] (Section V.15.4). The main results of this section are (3.36.26) and (3.36.28). Observe that if $k$ is perfect every extension $K/k$ satisfies Maclane's Criterion and is therefore Geometrically Reduced.

**Lemma 3.36.20**
*Let $A$ be a reduced $k$-algebra. Define the canonical map*

$$i : A \hookrightarrow \prod_{\mathfrak{p}} k(\mathfrak{p})$$

*where $\mathfrak{p}$ runs over all prime ideals of $A$. Then $i$ is injective. Furthermore for any $k$-algebra $B$ we have a canonical embedding*

$$A \otimes_k B \hookrightarrow \prod_{\mathfrak{p}} (k(\mathfrak{p}) \otimes_k B)$$

*Proof.* Evidently $\ker(i) = \bigcap \mathfrak{p}$ which by (3.3.46) is $(0)$. The final statement follows from considering the maps

$$A \otimes_k B \overset{i\otimes 1_B}{\hookrightarrow} \left(\prod_{\mathfrak{p}} k(\mathfrak{p})\right) \otimes_k B \hookrightarrow \prod_{\mathfrak{p}} (k(\mathfrak{p}) \otimes_k B)$$

The first map is injective by (3.4.25) and the second by (...). ∎

**Proposition 3.36.21**
*Let $A$ be a reduced $k$-algebra and $B$ a geometrically reduced $k$-algebra, then $A \otimes_k B$ is reduced.*

*Proof.* By the previous Lemma we have an embedding

$$A \hookrightarrow \prod (k(\mathfrak{p}) \otimes_k B)$$

As $B$ is geometrically reduced this shows that $A \otimes_k B$ is reduced. ∎

**Corollary 3.36.22**
*Let $k$ be a perfect field and $A, B$ reduced $k$-algebras. Then $A \otimes_k B$ is reduced.*

**Lemma 3.36.23**
*Let $A$ be a geometrically reduced $k$-algebra and $K$ the total ring of fractions. Then $K$ is geometrically reduced.*

*Conversely if $A$ is integral and $K$ is geometrically reduced, then $A$ is geometrically reduced.*

*Proof.* Let $L/k$ be a field extension and consider a nilpotent element $x \in K \otimes L$. Then by definition

$$x = \sum_{i=1}^{n} (a_i s_i^{-1}) \otimes \lambda_i \quad a_i \in A, s_i \notin Z(A), \lambda_i \in L$$

Let $s = s_1 ... s_n$ then $x \cdot (s \otimes 1) \in A \otimes L \subset K \otimes L$. Then $x \cdot (s \otimes 1)$ is nilpotent and so by definition zero. As $(s \otimes 1)$ is invertible we see $x = 0$ as required.

For the converse $A \subset K$ and therefore $A \otimes_k L \subset K \otimes_k L$. ∎

**Lemma 3.36.24**
*Let $K/k$ be a separably generated field extension. Then $K/k$ is geometrically reduced.*

*Proof.* Consider first the case $K = k(\mathcal{B})$ is purely transcendental. Then for $L/k$ we have $k[\mathcal{B}] \otimes L \cong L[\mathcal{B}]$ is evidently reduced. We may then demonstrate directly that $k(\mathcal{B}) \otimes_k L$ is reduced.

For the general case

$$K \otimes_k L \cong K \otimes_{k(\mathcal{B})} (k(\mathcal{B}) \otimes_k L)$$

© David Rufino CC BY-NC-ND 4.0

We have shown that $k(\mathcal{B}) \otimes_k L$ is reduced and by (3.36.19) $K/k(\mathcal{B})$ is geometrically reduced as by assumption it is separable. By (3.36.21) $K \otimes_k L$ is reduced and $K$ is geometrically reduced as required. ∎

**Lemma 3.36.25**
*Let $A$ be a reduced $k$-algebra for which $k$ is perfect. Then $A$ is geometrically reduced.*

*Proof.* We prove the result in increasing generality

a) We first consider the case $A = K/k$ is a finitely generated field extension. By (3.19.148) $K/k$ is separably generated and so by (3.36.24) it is geometrically reduced.
b) When $A$ is a finitely-generated $k$-algebra by (3.36.20) there is a canonical embedding

$$A \otimes_k L \hookrightarrow \prod_{\mathfrak{p}} (k(\mathfrak{p}) \otimes_k L)$$

By assumption $k(\mathfrak{p})$ is finitely generated and so by the first part is geometrically reduced. Therefore $A \otimes_k L$ is reduced as required.
c) The general case follows from b) and the reduction to the finitely generated case (3.36.5).

∎

**Proposition 3.36.26 (Maclane's Criterion for Algebras)**
*Let $A$ be a $k$-algebra. Then the following are equivalent*

*a) $A$ is geometrically reduced*
*b) $A \otimes_k \overline{k}$ is reduced*
*c) $A \otimes_k k^{p^{-\infty}}$ is reduced*
*d) $A \otimes_k K$ is reduced for some perfect extension $K/k$*
*e) $A \otimes_k k^{p^{-1}}$ is reduced*
*f) $A \otimes_k k'$ is reduced for every finite subextension $k'/k$ of $k^{p^{-1}}/k$*
*g) The following $k$-module homomorphism*

$$\begin{aligned} m : A \otimes_k k^{p^{-1}} &\to A \\ a \otimes \lambda &\to \lambda^p a^p \end{aligned}$$

*is injective (that is $k^{p^{-1}}$ and $A$ are linearly disjoint with respect to the $p$-th power embedding into $A$)*
*h) $A$ satisfies Maclane's Criterion (3.19.144)*

*In particular when $k$ is perfect then this is equivalent to $A$ being reduced.*

*Proof.* The case $p = 1$ is simply (3.36.25) because $k^p = k = k^{p^{-1}} = k^{p^{-\infty}}$ is perfect and g) is equivalent to $A$ being reduced. Therefore we assume $p > 1$.

*a)* $\implies$ *b)* By definition.

*b)* $\implies$ *c)* By (3.19.61) there is an embedding $k^{p^{-\infty}} \hookrightarrow \overline{k}$ whence an embedding $A \otimes_k k^{p^{-1}} \hookrightarrow A \otimes_k \overline{k}$.

*c)* $\implies$ *d)* By (3.19.100) $k^{p^{-\infty}}$ is perfect.

*d)* $\implies$ *e)* By (3.19.101) there is an embedding $k^{p^{-1}} \hookrightarrow K$ whence an embedding $A \otimes_k k^{p^{-1}} \hookrightarrow A \otimes_k K$.

*e)* $\implies$ *f)* This is immediate as we have an embedding $A \otimes_k k' \hookrightarrow A \otimes_k k^{p^{-1}}$.

*f)* $\iff$ *g)* Suppose $x := \sum_i \lambda_i \otimes a_i \in A \otimes_k k^{p^{-1}}$ is a finite sum then we may consider the finite subextension $k' := k\left(\{a_i\}_i\right)$. Observe that

$$x^p = \sum_i \lambda_i^p \otimes a_i^p = \sum_i 1 \otimes \lambda_i^p a_i^p = 1 \otimes \left( \sum_i \lambda_i^p a_i^p \right) =: 1 \otimes m(x)$$

Therefore we see $A \otimes_k k'$ is reduced for every such $k'$ iff $m(x)$ is injective.

*g)* $\iff$ *h)* Consider the Frobenius morphisms $i : k^{p^{-1}} \to A$ and $j : A \to A$. Then g) is equivalent to $k^{p^{-1}}$ and $A$ being linearly disjoint in $A$ with respect to $i, j$ and the result follows from the characterization of linear disjointness (3.35.55).

*h)* $\implies$ *a)* Let $\{a_i\}_{i \in I}$ be a $k$-basis for $A$ then by (3.4.26) $\{a_i \otimes 1\}_{i \in I}$ is an $L$-basis for $A \otimes_k L$. Given $x \in A \otimes_k L$ we have

$$x = \sum_{i \in I} a_i \otimes \lambda_i \quad \lambda_i \in L$$

© David Rufino CC BY-NC-ND 4.0

and

$$x^p = \sum_{i\in I} \lambda_i^p \otimes a_i^p = \sum_{i\in I} \lambda_i^p (a_i^p \otimes 1)$$

By assumption $\{a_i^p\}_{i\in I}$ is $k$-linearly independent and therefore $\{a_i^p \otimes 1\}_{i\in I}$ is $L$-linearly independent. Therefore $x^p = 0 \implies \lambda_i^p = 0$ for all $i \in I \implies \lambda_i = 0$ for all $i \in I \implies x = 0$, as $L$ is reduced. It follows from (3.19.145) that $A \otimes_k L$ is reduced, as we assumed that $p > 1$. ∎

**Corollary 3.36.27 (Maclane's Criterion for Fields)**
*Let $K/k$ be a field extension with characteristic exponent $p$. Then the following are equivalent*

*a) $K/k$ is geometrically reduced*
*b) $K \otimes_k \overline{k}$ is reduced*
*c) $K \otimes_k k^{p^{-1}}$ is reduced*
*d) $K \otimes_k k^{p^{-\infty}}$ is reduced*
*e) $K \otimes_k k'$ is reduced for some perfect extension $k'/k$*
*f) $K$ and $k^{p^{-1}}$ are linearly disjoint (with respect to some common field extension)*
*g) $K \otimes_k k^{p^{-1}}$ is an integral domain*
*h) $K \otimes_k k^{p^{-1}}$ is a field*
*i) $K/k$ satisfies Maclane's Criterion (3.19.144)*

*Observe i) has* finite character *in the sense it holds if and only if it holds for every finitely-generated subextension.*

*In particular when $k$ is perfect every extension is geometrically reduced.*

*Proof.* By (3.36.26) a)-f), i) are equivalent (using (3.35.57) to generalise the linear disjoint condition to any ambient field) and by (3.35.57) again f)-h) are equivalent. ∎

**Proposition 3.36.28 (Separable Field Extensions)**
*Let $K/k$ be a field extension. Then the following are equivalent*

*a) $K/k$ is geometrically reduced*
*b) $K \otimes_k \overline{k}$ is reduced*
*c) $K/k$ satisfies Maclane's Criterion (3.19.144)*
*d) Every finitely generated subextension $K/k$ is separably generated*

*In particular if $k$ is perfect every field extension satisfies these properties.*

*Proof.* By (3.36.27) we have $a) \iff b) \iff c)$. Further $c) \implies d)$ is Lemma (3.19.148), since every subextension satisfies Maclane's Criterion. For $d) \implies a)$ observe that every finitely generated subextension is geometrically reduced by (3.36.24). We may use (3.36.2) to argue that $K/k$ is geometrically reduced. ∎

**Corollary 3.36.29**
*Let $K/k$ be a finitely generated field extension. Then the following are equivalent.*

*a) $K/k$ is geometrically reduced*
*b) $K \otimes_k \overline{k}$ is reduced*
*c) $K/k$ satisfies Maclane's Criterion (3.19.144)*
*d) Every finitely generated subextension of $K/k$ is separably generated*
*e) $K/k$ is separably generated*

*Proof.* We've already shown equivalence of a)-d) and clearly $d) \implies e)$. Finally $e) \implies a)$ is simply (3.36.24). ∎

## 3.36.3 Geometrically Irreducible Algebras

The reference for this section is [Stacks] (Tag 00I2) presented here with a more elementary proof of (3.36.34).

**Lemma 3.36.30**
*Let $A, B$ be $k$-algebras. If $a \otimes 1 \in A \otimes_k B$ is a zero-divisor then so is $a$.*

*Proof.* Consider a basis $\{a_i\}_{i\in I}$ for $A$ such that $a_{i_0} = a$. If $a$ is not a zero-divisor then we may show directly that $\{aa_i\}_{i\in I}$ is linearly independent. Then by extending to a basis of $A$ we may use (3.4.26) to show that $\{(aa_i) \otimes 1\}_{i\in I}$ is $B$-linearly independent. We see immediately that $a \otimes 1$ is not a zero-divisor as required. ∎

**Lemma 3.36.31**
*Let $A$ be a ring and $\mathfrak{a} \subset N(A)$ a locally nilpotent ideal. Then $A$ is irreducible iff $A/\mathfrak{a}$ is irreducible.*

© David Rufino CC BY-NC-ND 4.0

**Lemma 3.36.32**
*Let $A \subset B$ be a subring of an irreducible ring. Then $A$ is irreducible.*

*In particular if $B$ is a geometrically irreducible $k$-algebra, then so is any subalgebra.*

*Proof.* By (3.23.7) every minimal prime in $A$ is the contraction of a minimal prime. As there is precisely one minimal prime of $B$ then we are done.

The final statement follows because the canonical map $A \otimes_k L \to B \otimes_k L$ is injective. ∎

When $k$ is algebraically closed then nothing interesting happens.

**Lemma 3.36.33**
*Let $A, B$ be irreducible (resp. integral) $k$-algebras with $k$ algebraically closed. Then $A \otimes_k B$ is irreducible (resp. integral).*

*In particular $A$ is a geometrically irreducible (resp. integral) $k$-algebra.*

*Proof.* Observe that $N(A) \otimes_k B + A \otimes_k N(B) \subset N(A \otimes_k B)$. For $(n \otimes \lambda)^m = n^m \otimes \lambda^m = 0$ and we may use (...) to argue any linear combination of such elements is nilpotent. By (...) we have an isomorphism

$$A/N(A) \otimes_k B/N(B) \cong (A \otimes_k B)/(N(A) \otimes_k B + A \otimes_k N(B))$$

Then by (3.36.31) $A/N(A) \otimes_k B/N(B)$ is irreducible iff $A \otimes_k B$ is. Therefore we may consider only the case $A, B$ reduced and therefore integral, and show that $A \otimes_k B$ is integral and a-fortiori irreducible.

If $A \otimes_k B$ were not integral it would contain a non-trivial zero divisor, and this would also be true for $A' \otimes_k B'$ for some finitely generated subalgebras $A'$ and $B'$. Therefore we may assume wlog that $A$ and $B$ are finitely generated.

To simplify the argument choose a $k$-basis $\{b_i\}_{i \in I}$ for $B$. By (...) $\{1 \otimes b_i\}_{i \in I}$ is an $A$-basis for $A \otimes_k B$. Therefore if we have a product of elements equal to 0 then may write it as follows

$$\left(\sum_i a_i \otimes b_i\right)\left(\sum_i a_i' \otimes b_i\right) = 0$$

for some $a_i, a_i' \in A$ all but finitely many zero. Let $\mathfrak{m} \lhd A$ be a maximal ideal, then by the Weak Nullstellensatz (3.35.23) the structural morphism $k \to A/\mathfrak{m}$ is an isomorphism. Reduce the equation to $A/\mathfrak{m} \otimes_k B \cong B$ to find

$$\left(\sum_i \overline{a}_i b_i\right)\left(\sum_i \overline{a}_i' b_i\right) = 0$$

As $B$ is an integral domain, and $b_i$ are linearly independent we find $\overline{a}_i, \overline{a}_i' = 0$. As $\mathfrak{m}$ was arbitrary we find $a_i, a_i' \in \bigcap \mathfrak{m}$ for all $i$. By assumption (0) is prime and so by the Strong Nullstellensatz (3.37.3) we find $a_i = a_i' = 0$. This shows that $A \otimes_k B$ is integral as required.

The last statement follows because a field $K/k$ is always irreducible. ∎

**Proposition 3.36.34 (Extension of Scalars is Flat)**
*Let $A$ be a $k$-algebra and $L/K/k$ a tower of field extensions. Then the canonical ring homomorphism*

$$A \otimes_k K \to A \otimes_k L$$

*satisfies Going Down. Further we have a well-defined surjective map of minimal primes*

$$\mathrm{MinPrime}(A \otimes_k L) \to \mathrm{MinPrime}(A \otimes_k K)$$

*given by the inverse image.*

*Proof.* By (...) there is a commutative diagram

$$\begin{array}{ccc} A \otimes_k K & \longrightarrow & A \otimes_k L \\ & \searrow & \downarrow \sim \\ & & (A \otimes_k K) \otimes_K L \end{array}$$

Therefore it is enough to show that $A \to A \otimes_k K$ satisfies going down. Suppose $\mathfrak{p}_1 \subseteq \mathfrak{p}_2 = \mathfrak{q}_2^c$. There is a commutative diagram

© David Rufino CC BY-NC-ND 4.0

$$\begin{array}{ccc} A/\mathfrak{p}_1 & \xrightarrow{\alpha} & A/\mathfrak{p}_1 \otimes_k K \\ \uparrow{\scriptstyle \pi} & & \uparrow{\scriptstyle \pi \otimes 1} \\ A & \xrightarrow{\beta} & A \otimes_k K \end{array}$$

where (3.4.42) shows that $\ker(\pi \otimes 1) = \mathfrak{p}_1^e$.

Let $\tilde{\mathfrak{q}}_1$ be a minimal prime of $A/\mathfrak{p}_1 \otimes_k K$ contained in $\tilde{\mathfrak{q}}_2 := (\pi \otimes 1)(\mathfrak{q}_2)$ by (3.3.44). Define $\mathfrak{q}_1 := (\pi \otimes 1)^{-1}(\tilde{\mathfrak{q}}_1)$ then $\mathfrak{q}_1$ is a prime minimal over $\mathfrak{p}_1^e$. As $\pi \otimes 1$ is surjective it is contained in $\mathfrak{q}_2$. We require to prove that $\beta^{-1}(\mathfrak{q}_1) = \mathfrak{p}_1$, which we readily see is implied by $\alpha^{-1}(\tilde{\mathfrak{q}}_1) = (0)$.

By (3.6.46) $\tilde{\mathfrak{q}}_1$ consists of zero divisors and by (3.36.30) so does $\alpha^{-1}(\tilde{\mathfrak{q}}_1)$. As $A/\mathfrak{p}_1$ is an integral domain we see that this equals $(0)$ and we are done.

This immediately shows that the inverse image of a minimal prime is again a minimal prime. By (3.23.7) every minimal prime is a contracted minimal prime. ∎

**Lemma 3.36.35 (Purely Inseparable Extensions of Scalars are Homeomorphisms)**
*Let $k$ be a field with characteristic exponent $p$, $A$ a $k$-algebra and $k'/k$ a purely inseparable field extension. Consider the structural algebra homomorphism*

$$\varphi : A \to A \otimes_k k'$$

*Then $\varphi$ induces an order-preserving bijection between prime ideals*

$$\mathfrak{q}^c \leftarrow \mathfrak{q}$$

*In particular $A$ is irreducible iff $A \otimes_k k'$ is.*

*Furthermore a purely inseparable extension is geometrically irreducible.*

*Proof.* If $p = 1$ then $k'/k$ is trivial and the statement is obvious. So we may assume wlog that $p > 1$.

If $\varphi$ were surjective then we would have equality $\mathfrak{q}^{ce} = \mathfrak{q}$ and which would imply the map of prime ideals is injective. However we note that $\varphi$ is "almost" surjective, namely for every $z = \sum_{i=1}^N a_i \otimes \lambda_i$ there exists $n \geq 0$ such that $z^{p^n} \in \varphi(A)$. For $k'/k$ is purely inseparable so by definition there is some $n \geq 0$ such that $\lambda_i^{p^n} \in k$ for all $i = 1...N$. In this case $z^{p^n} = \sum_{i=1}^N a_i^{p^n} \lambda_i^{p^n} \otimes 1$ which is in the image of $A$ as required.

This allows us to argue the map is injective for suppose $\mathfrak{p} := \mathfrak{q}^c = (\mathfrak{q}')^c$ and $z \in \mathfrak{q} \setminus \mathfrak{q}'$. Then by the previous argument $z^{p^n} = \varphi(x) \implies z^{p^n} \in \mathfrak{q} \implies z^{p^n} \in \mathfrak{q}' \implies z \in \mathfrak{q}'$ which is a contradiction. Therefore contraction of prime ideals is injective.

Finally we may observe that $\varphi$ is integral (3.24.20) and injective (3.36.2). Therefore it satisfies the lying over property (3.24.17) and so every prime ideal in $A$ is contracted and the given map is surjective. ∎

The following result shows it is sufficient to ensure that $A$ remains irreducible under a finite separable base change.

**Proposition 3.36.36**
*Let $A$ be a $k$-algebra. Then the following are equivalent*

*a) $A$ is geometrically irreducible*
*b) $A \otimes_k k'$ is irreducible for every finite separable extension $k'/k$*
*c) $A \otimes_k k^{\text{sep}}$ is irreducible*
*d) $A \otimes_k \overline{k}$ is irreducible*

*Proof.* *a)* $\implies$ *b)* immediate

*b)* $\implies$ *c)* Suppose $\mathfrak{q}_1, \mathfrak{q}_2$ are minimal prime ideals of $A \otimes_k k^{\text{sep}}$. Let $k^{\text{sep}}/k'/k$ be a fixed finite separable extension and consider the subring

$$A \otimes_k k' \subset A \otimes_k k^{\text{sep}}$$

together with the prime ideals $\mathfrak{p}_i := \mathfrak{q}_i \cap (A \otimes_k k')$. By (3.36.34) the prime ideals $\mathfrak{p}_i$ are minimal and therefore $\mathfrak{p}_1 = \mathfrak{p}_2$ by assumption. As $k'/k$ was an arbitrary finite subextension we may conclude by (3.36.5) that $\mathfrak{q}_1 = \mathfrak{q}_2$ and $A \otimes_k k^{\text{sep}}$ is irreducible.

*c)* $\implies$ *d)* By (...) $A \otimes_k \overline{k} \cong (A \otimes_k k^{\text{sep}}) \otimes_{k^{\text{sep}}} \overline{k}$ and so is irreducible by (3.36.35), since $\overline{k}/k^{\text{sep}}$ is purely inseparable (3.19.98).

© David Rufino CC BY-NC-ND 4.0

$d) \implies a)$ Let $K/k$ be a field extension then we may identify $\overline{k}$ and $K$ as subextensions of $\overline{K}/k$. Then $A \otimes_k \overline{K} \cong \left(A \otimes_k \overline{k}\right) \otimes_{\overline{k}} \overline{K}$. By (3.36.33) $A \otimes_k \overline{K}$ is irreducible. Finally from (3.36.34) we see that $A \otimes_k K$ is irreducible as required. ∎

**Proposition 3.36.37**
*Let $A$ be a geometrically irreducible $K$-algebra and $K/k$ a geometrically irreducible field extension. Then $A/k$ is geometrically irreducible.*

*Proof.* It is sufficient by (3.36.36) to consider base change by a finite separable extension $k'/k$. Observe by transitivity

$$A \otimes_k k' \cong A \otimes_K (K \otimes_k k')$$

so it would be sufficient to show that $K \otimes_k k'$ is a field. By hypothesis $K \otimes_k k'$ is irreducible. Furthermore $k'/k$ is geometrically reduced (3.36.18) whence $K \otimes_k k'$ is reduced and therefore an integral domain. Using the fact $k'/k$ is algebraic we may conclude from (3.35.57) that $K \otimes_k k'$ is a field, and we are done. ∎

**Proposition 3.36.38 (Primary Extension)**
*Let $K/k$ be a field extension. Then $K/k$ is geometrically irreducible if and only if it is relatively separably closed.*

*In this case the subfield of algebraic elements is purely inseparable over $k$. In particular an algebraic extension is geometrically irreducible if and only if it is purely inseparable.*

*We say such an extension is **primary**.*

*Proof.* Suppose that $K/k$ is relatively separably closed. By (3.36.36) it is enough to show that $K \otimes_k k'$ is irreducible for every finite separable extension $k'/k$.

Let $K/K'/k$ be the relative algebraic closure then $K/K'$ is relatively algebraically closed (3.19.46) and $K'/k$ is purely inseparable because $K_s = k$ by hypothesis (3.19.96). Observe $K'/k$ is geometrically irreducible (3.36.35) so by (3.36.37) it is enough to show that $K/K'$ is geometrically irreducible. This reduces to the case that $K/k$ is relatively algebraically closed.

As before it is sufficient to consider base change by a finite separable extension $k'/k$. By (3.19.77) $k' = k(\alpha)$ is simple. Let $m(X)$ be the minimal polynomial for $\alpha$ then $k(\alpha) \cong k[X]/(m)$. From (3.36.39) $m(X)$ remains irreducible in $K[X]$ and therefore (using $k(\alpha) \otimes_k K \cong (k[X] \otimes_k K)/(m)$ as $K$-algebras, (3.4.42), and $k[X] \otimes_k K \cong K[X]$, (3.8.7))

$$K \otimes_k k(\alpha) \cong K[X]/(m)$$

is a field and in particular irreducible.

Conversely suppose that $K/k$ is geometrically irreducible and $\alpha \in K$ is separable. Then $k(\alpha)$ is geometrically irreducible by (3.36.32). Therefore, using $k(\alpha) \otimes_k \overline{k} \cong \left(k[X] \otimes_k \overline{k}\right)/(m_\alpha)$ as $\overline{k}$-algebras (3.4.42), then $k[X] \otimes_k \overline{k} \cong \overline{k}[X]$ (3.8.7), then by the Chinese Remainder Theorem (3.3.62)

$$k(\alpha) \otimes_k \overline{k} \cong k[X]/(m_\alpha) \otimes_k \overline{k} \cong \overline{k}[X]/(m_\alpha) \cong \overline{k}[X]/(X-\alpha_1)^{n_1} \times \cdots \times \overline{k}[X]/(X-\alpha_r)^{n_r}$$

is irreducible where $\alpha_1, ..., \alpha_r$ are the distinct roots in $\overline{k}$. This implies $r = 1$ and therefore $\alpha$ is purely inseparable (3.19.88). Being both separable and purely inseparable $\alpha \in k$ (3.19.89) as required. ∎

We made use of

**Lemma 3.36.39**
*Let $K/k$ be a field extension such that $k$ is algebraically closed in $K$. Then every irreducible polynomial $f \in k[X]$ remains irreducible in $K[X]$.*

*Proof.* Consider the extension $\overline{K}$ which contains $\overline{k}$. And suppose $g \mid f$ in $K[X]$. As $f$ splits completely in $\overline{k}$ then by unique factorisation in $\overline{K}[X]$ the coefficients of $g$ lie in $K \cap \overline{k} = k$. Therefore by irreducibility $g = f$ or is constant. In particular $f$ is irreducible in $K[X]$. ∎

**Proposition 3.36.40**
*Let $A$ be an integral $k$-algebra and $K := \mathrm{Frac}(A)$. Then $A$ is geometrically irreducible iff $K$ is geometrically irreducible.*

*Proof.* Suppose $K$ is geometrically irreducible and consider an arbitrary field extension $L/k$. Then we have an embedding $A \otimes_k L \hookrightarrow K \otimes_k L$, and therefore an embedding $(A \otimes_k L)_{\mathrm{red}} \hookrightarrow (K \otimes_k L)_{\mathrm{red}}$. By assumption these are integral and we conclude that $(A \otimes_k L)$ is irreducible.

Conversely suppose $A \otimes_k L$ is irreducible then define $\Omega := \mathrm{Frac}\left((A \otimes_k L)_{\mathrm{red}}\right)$. As $L \hookrightarrow (A \otimes_k L)$ is injective and $L$ is reduced we have a canonical map $L \hookrightarrow \Omega$. Similarly we have a canonical map $A \hookrightarrow \Omega$ and therefore a map $K \hookrightarrow \Omega$. Therefore we have a well-defined ring homomorphism

© David Rufino CC BY-NC-ND 4.0

$$\psi : K \otimes_k L \to \Omega$$
$$ab^{-1} \otimes \lambda \to (a \otimes \lambda) \cdot (b \otimes 1)^{-1}$$

We must show that if $x \in \ker(\psi)$ then $x$ is nilpotent. By definition

$$x = \sum_{i=1}^{n} \lambda_i \otimes \mu_i$$

We may find $a \in A$ such that $y = ax$ is of the form

$$y = \sum_{i=1}^{n} a_i \otimes \mu_i \in A \otimes_k L$$

We observe $y \in \ker(A \otimes_k L \to \Omega)$ whence $y$, and therefore $x$, is nilpotent. This means we have an embedding

$$(K \otimes_k L)_{\text{red}} \hookrightarrow \Omega$$

which means $(K \otimes_k L)_{\text{red}}$ is integral and therefore $K \otimes_k L$ is irreducible. ∎

**Proposition 3.36.41**
*Let $A$ be an integral $k$-algebra and $K \coloneqq \operatorname{Frac}(A)$. Then the following are equivalent*

*a) $A$ is geometrically irreducible*
*b) $A \otimes_k k'$ is irreducible for every finite separable extension $k'/k$*
*c) $A \otimes_k \overline{k}$ is irreducible*
*d) $K$ is geometrically irreducible*
*e) $K$ is relatively separably closed*

*Proof.* We may combine (3.36.36), (3.36.40) and (3.36.38). ∎

## 3.36.4 Geometrically Integral Algebras

Clearly a geometrically (resp. algebraically) integral algebra is integral, we are interested in providing sufficient conditions for the reverse implication.

**Proposition 3.36.42**
*Let $A$ be a geometrically integral $k$-algebra and $B$ an integral $k$-algebra. Then $A \otimes_k B$ is an integral domain.*

*Proof.* Let $L = \operatorname{Frac}(B)$ then $A \otimes_k B$ is a subring of $A \otimes_k L$ and we are done. ∎

**Proposition 3.36.43 (Criteria for Geometrically Integral in terms of Quotient Field)**
*Let $A$ be an integral $k$-algebra and $K \coloneqq \operatorname{Frac}(A)$. Then $A \otimes_k L$ is integral if and only if $K \otimes_k L$ is integral.*

*In particular $A$ is geometrically (resp. algebraically) integral if and only if $K$ is geometrically (resp. algebraically) integral.*

*Proof.* This follows exactly the same lines as (3.36.40). ∎

**Proposition 3.36.44 (Criteria to be Geometrically Integral)**
*Let $A$ be a $k$-algebra. Then the following are equivalent*

*a) $A$ is geometrically integral*
*b) $A$ is geometrically reduced and irreducible*
*c) $A \otimes_k \overline{k}$ is an integral domain*
*d) $A \otimes_k k'$ is an integral domain for every finite extension $k'/k$*

*Note we may restrict d) to only finite separable, and finite purely inseparable extensions of height at most 1.*

*Proof.* This is largely a collection of existing results. $a) \iff b)$ is by definition and (3.3.67). Then $b) \iff c) \iff d)$ is from the reduced (3.36.26) and irreducible (3.36.36) cases, together with the trivial observation that reduced + irreducible $\iff$ integral (3.3.67). ∎

**Proposition 3.36.45 (Regular Extension)**
*Let $K/k$ be a finitely generated field extension. Then the following are equivalent*

*a) $K/k$ is geometrically integral*
*b) $K/k$ is separably generated and relatively separably closed*
*c) $K/k$ is separably generated and relatively algebraically closed*

© David Rufino CC BY-NC-ND 4.0

*d)* $K \otimes_k k'$ *is an integral domain for every* finite *extension* $k'/k$
*e)* $K \otimes_k k'$ *is a field for every* algebraic *extension* $k'/k$
*f)* $K \otimes_k \overline{k}$ *is an integral domain*
*g)* $K \otimes_k \overline{k}$ *is a field*

*We say such an extension is* ***regular****.*

*Proof.* *a)* $\Longrightarrow$ *b)* is (3.36.29) and (3.36.38). *c)* $\Longrightarrow$ *b)* is immediate. For the converse suppose $\alpha \in K/k$ is algebraic then $k(\alpha)/k$ is geometrically reduced (3.36.29) and therefore separable algebraic (3.36.18). By hypothesis $\alpha \in k$ as required, giving *b)* $\Longleftrightarrow$ *c)*. For *b)* $\Longrightarrow$ *a)*: $K/k$ being separably generated, it is geometrically reduced (3.36.24), and being relatively separably closed, it is geometrically irreducible (3.36.38) together these give *a)* (3.3.67). This closes the loop *a)* $\Longleftrightarrow$ *b)* $\Longleftrightarrow$ *c)*. *a)* $\Longleftrightarrow$ *d)* $\Longleftrightarrow$ *f)* was already proven in (3.36.44), and *f)* $\Longleftrightarrow$ *g)* was proven in (3.35.57). *e)* $\Longrightarrow$ *d), g)* is immediate.

For *f)* $\Longrightarrow$ *e)* fix an algebraic $k'/k$ and an embedding $k' \hookrightarrow \overline{k}$ (exists by (3.19.61)); the canonical map $K \otimes_k k' \to K \otimes_k \overline{k}$ is injective (3.36.2) ($k'$ being a $k$-subalgebra of $\overline{k}$), so $K \otimes_k k'$ is a subring of the domain $K \otimes_k \overline{k}$, hence itself a domain; as $k'/k$ is algebraic, this domain is moreover a field (3.35.57). ∎

**Corollary 3.36.46**
*Let* $A$ *be a finitely-generated integral* $k$*-algebra and* $K$ *the quotient field. Then the following are equivalent*

*a)* $A$ *is geometrically integral*
*b)* $K/k$ *is geometrically integral*
*c)* $K/k$ *is separably generated and relatively separably closed*
*d)* $K \otimes_k k'$ *is an integral domain for every finite extension* $k'/k$
*e)* $K \otimes_k \overline{k}$ *is an integral domain*
*f)* $K \otimes_k \overline{k}$ *is a field*

*Proof.* *a)* $\Longleftrightarrow$ *b)* is (3.36.43) and the rest is (3.36.45). ∎

The mechanism underlying (6.3.44) below (base-changing a place of an algebraic function field along an algebraic extension of the constant field) is really a fact about discrete valuation rings, unrelated to the transcendence degree 1 setting of function fields; we isolate it here.

**Proposition 3.36.47 (Unramified Base Extension of a Discrete Valuation Ring)**
*Let* $k$ *be a perfect field and* $\mathcal{O}$ *a discrete valuation ring containing* $k$*, with fraction field* $F$ *and maximal ideal* $\mathfrak{m}$*, such that* $F/k$ *is geometrically integral and* $k(\mathfrak{m}) := \mathcal{O}/\mathfrak{m}$ *is finite over* $k$*.*

*Let* $k'/k$ *be algebraic, write* $\mathcal{O}_{k'} := \mathcal{O} \otimes_k k'$ *and* $\mathfrak{b} := \mathfrak{m}\mathcal{O}_{k'}$*. Then*

*a)* $\mathcal{O}_{k'}$ *has exactly* $r$ *maximal ideals* $\mathfrak{p}_1, \ldots, \mathfrak{p}_r$*, the preimages of those of* $\mathcal{O}_{k'}/\mathfrak{b}$
*b) the localisation* $(\mathcal{O}_{k'})_{\mathfrak{p}_j}$ *is a place of* $F_{k'}$ *unramified over* $\mathcal{O}$*: a uniformiser of* $\mathcal{O}$ *generates the maximal ideal* $\mathfrak{p}_j(\mathcal{O}_{k'})_{\mathfrak{p}_j}$*, and*
*c) every valuation ring* $R$ *of* $F_{k'}$ *with* $k' \subset R$*,* $\mathcal{O} \subset R$*, and* $R \cap F = \mathcal{O}$ *is of the form* $R = (\mathcal{O}_{k'})_{\mathfrak{p}_j}$ *for some* $j$*, and*
*d) when* $k'/k$ *is finite then* $(\mathcal{O}_{k'})_{\mathfrak{p}_j}$ *is a discrete valuation ring, and*
*e)* $(\mathcal{O}_{k'})_{\mathfrak{p}_j} \cap F = \mathcal{O}$ *and* $\mathfrak{p}_j(\mathcal{O}_{k'})_{\mathfrak{p}_j} \cap F = \mathfrak{m}$*, inside* $F_{k'}$*: that is* $(\mathcal{O}_{k'})_{\mathfrak{p}_j}$ *dominates* $\mathcal{O}$ *and restricts to it. Together with c), the valuation rings* $R$ *of* $F_{k'}$ *with* $k' \subset R$*,* $\mathcal{O} \subset R$ *and* $R \cap F = \mathcal{O}$ *are therefore exactly the* $(\mathcal{O}_{k'})_{\mathfrak{p}_j}$*.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Let $\pi$ be a uniformiser of $\mathcal{O}$, so $\mathfrak{m} = (\pi)$ (3.31.2). Since $F/k$ is geometrically integral and $k'/k$ algebraic, $F_{k'}$ is a field (3.36.45), and $\mathcal{O}_{k'}$ is a subring (...) and in particular an integral domain. Further $\mathcal{O}_{k'}$ integral over $\mathcal{O}$ (3.24.20).

$\mathcal{O}_{k'}/\mathfrak{b} = \mathcal{O}_{k'}/\mathfrak{m}\mathcal{O}_{k'} \cong (\mathcal{O}/\mathfrak{m}) \otimes_k k' = k(\mathfrak{m}) \otimes_k k'$ as $k$-algebras (3.4.42): a finite-dimensional (as $k(\mathfrak{m})/k$ is finite, by hypothesis) reduced ($k(\mathfrak{m})/k$ and $k'/k$ both separable, $k$ being perfect, (3.19.103), (3.36.28)) $k$-algebra, hence a finite product of $r$ fields (3.36.12), with every prime maximal (3.36.13).

**a) Finitely many maximal ideals.** $\mathcal{O}_{k'}/\mathfrak{b}$, being a finite product of $r$ fields (above), has finitely many maximal ideals $\mathfrak{p}'_1, \ldots, \mathfrak{p}'_r$, all its primes maximal (3.36.13). Every maximal ideal of $\mathcal{O}_{k'}$ contracts to a maximal ideal of $\mathcal{O}$ (3.24.16), that is to $\mathfrak{m}$ ($\mathcal{O}$ being local), hence contains $\mathfrak{b}$; so, by the ideal correspondence for $\mathcal{O}_{k'} \twoheadrightarrow \mathcal{O}_{k'}/\mathfrak{b}$ (3.3.56), the maximal ideals of $\mathcal{O}_{k'}$ are exactly the preimages $\mathfrak{p}_1, \ldots, \mathfrak{p}_r$ of $\mathfrak{p}'_1, \ldots, \mathfrak{p}'_r$ – and, every prime of the quotient being maximal, these are also *every* prime of $\mathcal{O}_{k'}$ containing $\mathfrak{b}$.

© David Rufino CC BY-NC-ND 4.0

As $\mathfrak{b}$ is radical (the quotient being reduced), the Prime Nullstellensatz (3.3.46) gives $\mathfrak{b} = \mathfrak{p}_1 \cap \cdots \cap \mathfrak{p}_r$; as the $\mathfrak{p}_j$ are pairwise comaximal (distinct maximal ideals), this equals $\mathfrak{p}_1 \cdots \mathfrak{p}_r$ (3.3.62).

**b) Unramifiedness.** Write $\mathcal{O}_j := (\mathcal{O}_{k'})_{\mathfrak{p}_j}$. By (3.6.18).a) and multiplicativity of localisation on ideals (3.30.8),

$$\pi\mathcal{O}_j = \mathfrak{b}\mathcal{O}_j = \mathfrak{b}_{\mathfrak{p}_j} = (\mathfrak{p}_1)_{\mathfrak{p}_j} \cdots (\mathfrak{p}_r)_{\mathfrak{p}_j} = \mathfrak{p}_j\mathcal{O}_j$$

(the first equality as $\mathfrak{b} = \pi\mathcal{O}_{k'}$; the factors $i \neq j$ becoming the unit ideal, as $\mathfrak{p}_i \not\subset \mathfrak{p}_j$). So $\pi$ generates $\mathfrak{p}_j\mathcal{O}_j$, once $\mathcal{O}_j$ is known to be a valuation ring of $F_{k'}$ properly between $k'$ and $F_{k'}$ – established next.

**d) The finite case: $\mathcal{O}_j$ is a DVR.** Suppose $k'/k$ is finite. Then $\mathcal{O}_{k'}$ is a finite, free $\mathcal{O}$-module (3.4.26), hence a Noetherian $\mathcal{O}$-module (3.15.4) ($\mathcal{O}$ Noetherian, being a PID (3.16.2)), so Noetherian as a ring (3.15.3) (every ideal being in particular an $\mathcal{O}$-submodule). So $\mathcal{O}_j$ is Noetherian (3.15.7), local with maximal ideal $\mathfrak{p}_j\mathcal{O}_j$ (3.6.42), and a domain (above). It has Krull dimension 1: as $\mathcal{O} \hookrightarrow \mathcal{O}_{k'}$ is injective and integral (module-finite), $\dim \mathcal{O}_{k'} = \dim \mathcal{O} = 1$ (3.24.22) ($\mathcal{O}$ a DVR, hence of dimension 1 (3.31.2)); as $\mathcal{O}_{k'}$ is a domain, $(0) \subsetneq \mathfrak{p}_j$ gives $\mathrm{ht}(\mathfrak{p}_j) \geq 1$, while $\mathrm{ht}(\mathfrak{p}_j) \leq \dim \mathcal{O}_{k'} = 1$, so $\dim \mathcal{O}_j = \mathrm{ht}(\mathfrak{p}_j) = 1$ (3.22.4).b). With $\mathfrak{p}_j\mathcal{O}_j = (\pi)$ (above), $\mathcal{O}_j$ is a DVR (3.31.2).

**In general, $\mathcal{O}_j$ is a valuation ring.** Write $k' = \bigcup_i k'_i$ as the directed union of its finite subextensions (3.19.47), so $\mathcal{O}_{k'} = \bigcup_i \mathcal{O}_{k'_i}$ for $\mathcal{O}_{k'_i} := \mathcal{O} \otimes_k k'_i$ (a directed union of subrings, via (3.36.2)). For each $i$, $\mathfrak{p}_{i,j} := \mathfrak{p}_j \cap \mathcal{O}_{k'_i}$ contains $\mathfrak{m}\mathcal{O}_{k'_i}$ (as $\mathfrak{p}_j \supseteq \mathfrak{b}$), so corresponds, under the ideal correspondence for the finite-dimensional ring $\mathcal{O}_{k'_i}/\mathfrak{m}\mathcal{O}_{k'_i} = k(\mathfrak{m}) \otimes_k k'_i$ (as $k$-algebras, (3.4.42)) (reduced, as $k(\mathfrak{m})/k$ and $k'_i/k$ are separable, $k$ perfect (3.19.103), (3.36.28), so a finite product of fields (3.36.12), as in the statement, applied to $k'_i$ in place of $k'$), to a prime of that ring, automatically maximal (3.36.13) so $\mathfrak{p}_{i,j}$ is a maximal ideal of $\mathcal{O}_{k'_i}$. By the finite case just established (taking $k'_i$ for $k'$), $\left(\mathcal{O}_{k'_i}\right)_{\mathfrak{p}_{i,j}}$ is a DVR of $F_{k'_i}$, with $\mathcal{O} \subsetneq \left(\mathcal{O}_{k'_i}\right)_{\mathfrak{p}_{i,j}}$ and $\pi$ generating $\mathfrak{p}_{i,j}\left(\mathcal{O}_{k'_i}\right)_{\mathfrak{p}_{i,j}}$.

**Claim: $\mathcal{O}_j = \bigcup_i \left(\mathcal{O}_{k'_i}\right)_{\mathfrak{p}_{i,j}}$.** For $k'_i \subseteq k'_l$, $\mathfrak{p}_{l,j} \cap \mathcal{O}_{k'_i} = \mathfrak{p}_j \cap \mathcal{O}_{k'_i} = \mathfrak{p}_{i,j}$, so (3.6.9) gives $\left(\mathcal{O}_{k'_i}\right)_{\mathfrak{p}_{i,j}} \subseteq \left(\mathcal{O}_{k'_l}\right)_{\mathfrak{p}_{l,j}}$: a directed system. As $\mathcal{O}_{k'} = \bigcup_i \mathcal{O}_{k'_i}$ is a directed union of subrings with $\mathfrak{p}_{i,j} = \mathfrak{p}_j \cap \mathcal{O}_{k'_i}$, (3.6.10) gives $\mathcal{O}_j = \bigcup_i \left(\mathcal{O}_{k'_i}\right)_{\mathfrak{p}_{i,j}}$.

**Claim: $k' \subsetneq \mathcal{O}_j \subsetneq F_{k'}$.** As $\pi$ generates the maximal ideal of every $\left(\mathcal{O}_{k'_i}\right)_{\mathfrak{p}_{i,j}}$, (3.31.6) applies directly (with $F_{k'} = \bigcup_i F_{k'_i}$ the corresponding directed union of fields): $\mathcal{O}_j = \bigcup_i \left(\mathcal{O}_{k'_i}\right)_{\mathfrak{p}_{i,j}}$ is a valuation ring of $F_{k'}$ with maximal ideal $(\pi)\mathcal{O}_j$, and $\mathcal{O}_j \subsetneq F_{k'}$. Also $k' \subsetneq \mathcal{O}_j$: $k' = \bigcup_i k'_i \subseteq \bigcup_i \left(\mathcal{O}_{k'_i}\right)_{\mathfrak{p}_{i,j}} = \mathcal{O}_j$ is immediate, while $\pi \in \mathcal{O}_j \setminus k'$, as $\pi \notin k'_i$ for every $i$ (else $\pi \in \mathfrak{p}_{i,j} \cap k'_i$ would be a unit of $\mathcal{O}_{k'_i}$ (nonzero elements of the subfield $k'_i \subset \mathcal{O}_{k'_i}$ are units), contradicting $\pi \notin \left(\left(\mathcal{O}_{k'_i}\right)_{\mathfrak{p}_{i,j}}\right)^\times$), so $\pi \notin k'$.

This closes b): $\mathcal{O}_j$ is a valuation ring of $F_{k'}$ with $k' \subsetneq \mathcal{O}_j \subsetneq F_{k'}$ – that is, a place of $F_{k'}$ – and $\pi$ generates both $\mathfrak{m}$ and $\mathfrak{p}_j\mathcal{O}_j$ (above): the same uniformiser generating the maximal ideal on both sides of $\mathcal{O} \subset \mathcal{O}_j$ is exactly what it means for $\mathfrak{m}$ to be unramified in $\mathcal{O}_{k'}$ – matching the factorisation $\mathfrak{b} = \mathfrak{p}_1 \cdots \mathfrak{p}_r$ above. When $k'/k$ is finite, $\mathcal{O}_j$ is moreover a discrete valuation ring (d)), recovering the usual ramification index $e(\mathcal{O}_j/\mathcal{O}) = v_{\mathcal{O}_j}(\pi) = 1$.

**c) Every valuation ring $R$ with $R \cap F = \mathcal{O}$ is some $\mathcal{O}_j$.** Let $R$ be a valuation ring of $F_{k'}$ with $k' \subset R$, $\mathcal{O} \subset R$, and $R \cap F = \mathcal{O}$. As $\mathcal{O}_{k'}$ is generated as a ring by (the images of) $\mathcal{O}$ and $k'$ (spanned by simple tensors $a \otimes b = a \cdot b$), both contained in $R$, $\mathcal{O}_{k'} \subset R$. So $\mathfrak{q} := \mathfrak{m}_R \cap \mathcal{O}_{k'}$ is prime (a contraction of the prime $\mathfrak{m}_R$ along $\mathcal{O}_{k'} \hookrightarrow R$); it contains $\mathfrak{b}$, as $\pi \in \mathfrak{m}_R$ (else $\pi^{-1} \in R$, so $\pi^{-1} \in R \cap F = \mathcal{O}$, contradicting $\pi$ a non-unit of $\mathcal{O}$), giving $\mathfrak{b} = \pi\mathcal{O}_{k'} \subset \mathfrak{m}_R \cap \mathcal{O}_{k'} = \mathfrak{q}$. So $\mathfrak{q}$ corresponds to a prime, automatically maximal (3.36.13), of the finite-dimensional ring $\mathcal{O}_{k'}/\mathfrak{b} \cong \prod_j E_j$; so $\mathfrak{q} = \mathfrak{p}_j$ for some $j$.

Every $s \in \mathcal{O}_{k'} \setminus \mathfrak{p}_j$ then satisfies $s \notin \mathfrak{q} = \mathfrak{m}_R \cap \mathcal{O}_{k'}$, that is $s \notin \mathfrak{m}_R$, so $s$ is a unit of $R$; by the universal property of localisation, $\mathcal{O}_j$ then embeds in $R$. As $\mathfrak{m}_{\mathcal{O}_j} = \pi\mathcal{O}_j$ and $\pi \in \mathfrak{m}_R$, $\mathfrak{m}_{\mathcal{O}_j} \subseteq \mathfrak{m}_R \cap \mathcal{O}_j$; conversely $\mathfrak{m}_R \cap \mathcal{O}_j$ is a proper ideal of the local ring $\mathcal{O}_j$, hence $\subseteq \mathfrak{m}_{\mathcal{O}_j}$: so $\mathfrak{m}_{\mathcal{O}_j} = \mathfrak{m}_R \cap \mathcal{O}_j$, that is $R$ dominates $\mathcal{O}_j$ (3.25.4). As $\mathcal{O}_j$ is a valuation ring, it is maximal under domination (3.25.13), so $R = \mathcal{O}_j$.

**e) $\mathcal{O}_j$ restricts to $\mathcal{O}$.** As $\mathcal{O}_{k'} \subset F_{k'}$ is a domain, $\mathcal{O}_j$ is the subring of $F_{k'}$ of fractions $a/s$ with $a \in \mathcal{O}_{k'}$, $s \in \mathcal{O}_{k'} \setminus \mathfrak{p}_j$, so $\mathcal{O} \subset \mathcal{O}_{k'} \subset \mathcal{O}_j$. Moreover $\mathfrak{m} \subseteq \mathfrak{m}\mathcal{O}_{k'} = \mathfrak{b} \subseteq \mathfrak{p}_j$ (a)), so $\mathfrak{m} \subseteq \mathfrak{p}_j\mathcal{O}_j = \mathfrak{m}_{\mathcal{O}_j}$ (3.6.42): and $\mathfrak{p}_j\mathcal{O}_j$ is a proper ideal. Since $\mathcal{O}$ is a valuation ring of $F$, (3.25.6) (with $B = \mathcal{O}_j$, $\mathfrak{n} = \mathfrak{p}_j\mathcal{O}_j$) gives $\mathcal{O}_j \cap F = \mathcal{O}$ and $\mathfrak{p}_j\mathcal{O}_j \cap F = \mathfrak{m}$. Combined with c) this shows that the valuation rings $R$ of $F_{k'}$ with $k' \subset R$, $\mathcal{O} \subset R$ and $R \cap F = \mathcal{O}$ are precisely $\mathcal{O}_1, \ldots, \mathcal{O}_r$. ∎

**Remark 3.36.48 (Relation to the Stacks Project)**
*For $k'/k$ finite, this result also follows by combining two general facts from [Stacks]: unramified ring maps are stable under base change ([Stacks] (Tag 00US), applied to the finite separable, hence unramified, extension $k \to k(\mathfrak{m})$, base-*

© David Rufino CC BY-NC-ND 4.0

*changed along $k \to k'$), and a finite-dimensional algebra over a field splits, after any base change, into a product of local factors dual to its maximal ideals (the Stacks Project analogue of (3.36.15), (3.36.13) above).*

*For $k'/k$ merely* algebraic*, the Stacks Project's treatment of unramified morphisms is developed for morphisms of finite type, so does not directly cover this case; the passage to infinite $k'$ via the directed union of finite subextensions, as in* In general, $\mathcal{O}_j$ is a valuation ring *above, has no immediate analogue there and is instead the kind of limit argument found in the Stacks Project's chapter on limits of rings (e.g. [Stacks] (Tag 07RN ff.), on writing a ring as a filtered colimit and transporting properties of the terms to the colimit).*

© David Rufino CC BY-NC-ND 4.0

## 3.37 Jacobson Rings

**Definition 3.37.1 (Jacobson Radical)**
*Let $\mathfrak{a} \triangleleft A$ be an ideal. Define the **Jacobson Radical** of $\mathfrak{a}$ to be*

$$\sqrt{\mathfrak{a}}^J := \bigcap_{\mathfrak{a} \subseteq \mathfrak{m}} \mathfrak{m}$$

*Note by (3.3.46) and (3.3.60)*

$$\sqrt{\mathfrak{a}} \subseteq \sqrt{\mathfrak{a}}^J$$

**Proposition 3.37.2 (Jacobson Ring)**
*Let $A$ be a ring the following are equivalent*

*a) For any ideal $\mathfrak{a}$, $\sqrt{\mathfrak{a}} = \sqrt{\mathfrak{a}}^J$*
*b) For any radical ideal $\mathfrak{a} = \sqrt{\mathfrak{a}}^J$*
*c) For any prime ideal $\mathfrak{p} = \sqrt{\mathfrak{p}}^J$*

*We say such a ring is a **Jacobson ring**.*

*Proof.* a) $\implies$ b) This is clear because in this case $\mathfrak{a} = \sqrt{\mathfrak{a}}$.

b) $\implies$ c) This is clear because a prime ideal is radical.

c) $\implies$ a) By (3.3.46)

$$\sqrt{\mathfrak{a}} = \bigcap_{\mathfrak{a} \subseteq \mathfrak{p}} \mathfrak{p} = \bigcap_{\mathfrak{a} \subseteq \mathfrak{p} \subseteq \mathfrak{m}} \mathfrak{m} = \bigcap_{\mathfrak{a} \subseteq \mathfrak{m}} \mathfrak{m}$$

as required ∎

We prove the Weak Nullstellensatz implies the following result

**Proposition 3.37.3 (Strong Nullstellensatz)**
*Let $A$ be a finitely-generated $k$-algebra. Then it is a Jacobson ring.*

*Proof.* **TODO** ∎

© David Rufino CC BY-NC-ND 4.0

# Chapter 4

# Topology and Sheaves

© David Rufino CC BY-NC-ND 4.0

## 4.1 Topological Spaces

References:

- General Topology [Kel71] (Chap. 1)
- General Topology [Wil70]

Topology is useful in algebraic geometry, but often the natural topologies are usually much coarser so the theory looks rather different.

**Definition 4.1.1 (Topological Space)**
*A topological space $(X, \mathcal{T}_X)$ consists of a set $X$ and family of open sets $\mathcal{T}_X \subseteq \mathcal{P}(X)$ satisfying the following properties*

- $X, \varnothing \in \mathcal{T}_X$
- $U_i \in \mathcal{T}_X \implies \bigcup_{i\in I} U_i \in \mathcal{T}_X$
- $U, V \in \mathcal{T}_X \implies U \cap V \in \mathcal{T}_X$

*A subset $Z \subset X$ is said to be closed iff $X \setminus Z$ is open. We may equivalently define the topology in terms of closed sets.*

**Proposition 4.1.2**
*Let $X$ be a topological space. Both the open sets and closed sets form a distributive lattice under inclusion.*

**Definition 4.1.3 (Subspace topology)**
*Let $Y \subset X$, then we may define the **subspace topology** on $Y$ by*

$$\mathcal{T}_Y := \{U \cap Y \mid U \in \mathcal{T}_X\}$$

*when $Y$ is open then this is given by*

$$\mathcal{T}_Y = \{U \subseteq Y \mid U \in \mathcal{T}_X\}$$

**Definition 4.1.4 (Base)**
*We say $\mathcal{B} \subseteq \mathcal{P}(X)$ is a base (of open sets) on $X$ if*

- *For every $x \in X$ there is a $U \in \mathcal{B}$ such that $x \in U$*
- *Suppose $U, V \in \mathcal{B}$ and $x \in U \cap V$ then there exists $W \in \mathcal{B}$ such that $x \in W \subseteq U \cap V$*

**Proposition 4.1.5 (Topology generated by a base)**
*Let $\mathcal{B}$ be a base, then the following is a topology on $X$*

$$\mathcal{T}_{\mathcal{B}} := \left\{ \bigcup_{U_i \in I} U_i \;\middle|\; I \subseteq \mathcal{B} \right\}$$

*i.e. the set of arbitrary unions of sets in $\mathcal{B}$.*

**Proposition 4.1.6 (Base generating topology)**
*A base $\mathcal{B}$ satisfies $\mathcal{T}_{\mathcal{B}} = \mathcal{T}_X$ if and only if*

*a) $\mathcal{B} \subset \mathcal{T}_X$*
*b) For every $x \in U$ and $U \in \mathcal{T}_X$ there exists $V \in \mathcal{B}$ such that $x \in V \subseteq U$.*

*In this case we say $\mathcal{B}$ is a base for (the topology on) $X$.*

**Proposition 4.1.7 (Subbase for a topology)**
*Let $X$ be a set and $\mathcal{B}$ be an arbitrary family of subsets. Then there exists a topology $\mathcal{T}_{\mathcal{B}}$ such that*

*a) $\mathcal{B} \subset \mathcal{T}_{\mathcal{B}}$*
*b) For every topology $\mathcal{T}'$ containing $\mathcal{B}$ we have $\mathcal{T}_{\mathcal{B}} \subset \mathcal{T}'$.*

*We say that $\mathcal{B}$ is a **subbase** for the topology $\mathcal{T}_{\mathcal{B}}$. Explicitly $\mathcal{T}_{\mathcal{B}}$ consists of arbitrary unions of finite intersections of elements of $\mathcal{B}$.*

**Proposition 4.1.8 (Base for subspace topology)**
*Let $(X, \mathcal{T})$ be a topological space, and $Y \subset X$ has the subspace topology $\mathcal{T}_Y$ (4.1.3). If $\mathcal{B}$ is a base for $(X, \mathcal{T})$ then*

$$\mathcal{B}_Y := \{U \cap Y \mid U \in \mathcal{B}\}$$

*is a base for $(Y, \mathcal{T}_Y)$.*

© David Rufino CC BY-NC-ND 4.0

**Definition 4.1.9 (Adherent point)**
*For $Y \subset X$ we say $x$ is an **adherent point** of $Y$ if every open neighbourhood of $x$ intersects $Y$.*

*Similarly we say $x$ is a **limit point** (or **accumulation point**) for $Y$ if every open neighbourhood of $x$ intersects $Y$ at a point other than $x$.*

**Proposition 4.1.10 (Closed point)**
*Let $x \in X$ then TFAE*

- *$\{x\}$ is closed*
- *For every $y \neq x$ there is $U \ni y$ such that $x \notin U$.*

**Definition 4.1.11**
*For a topological space $X$ let $X_0$ denote the subset of closed points.*

**Definition 4.1.12 (Open Embedding)**
*Let $f : X \to Y$ be a map of topological spaces. We say it is an **open embedding** if one of the following equivalent conditions holds*

*a) $f(X)$ is open and $f$ is a homeomorphism onto its image*
*b) $f$ is continuous, open and injective.*

### 4.1.1 Axioms of Countability

**Definition 4.1.13 (Neighbourhood)**
*Let $X$ be a topological space and $x \in X$. We say a set $V \subset X$ is a **neighbourhood** of $x$ if $x \in U \subset V$ for some open set $U$. We write $\mathcal{U}_x$ for the neighbourhoods of $x$, and $\mathcal{O}_x$ for the open neighbourhoods of $x$.*

*We say that a collection of neighbourhoods $\mathcal{B}_x$ is a **local base** for $x$ if every neighbourhood $U \in \mathcal{U}_x$ contains some neighbourhood $V \in \mathcal{B}_x$. For example $\mathcal{O}_x$ is a local base.*

**Definition 4.1.14 (First-Countable)**
*Let $X$ be a topological space. $X$ is **first countable** if every point $x \in X$ has a countable local base.*

**Definition 4.1.15 (Second-Countable)**
*Let $X$ be a topological space. $X$ is **second countable** if there exists a countable basis. Evidently this is stronger than first countability.*

**Example 4.1.16**
*For example a metric space is always first countable as we may consider balls of radius $\frac{1}{n}$.*

*Further $\mathbb{R}$ is second countable as we may consider open intervals with rational endpoints.*

### 4.1.2 Closure

**Definition 4.1.17 (Adherent Point)**
*Let $X$ be a topological space and $Y \subset X$. We say that $x \in X$ is an **adherent point** of $Y$, if $U \in \mathcal{U}_x \implies U \cap Y \neq \varnothing$.*

*We say that $x$ is a **sequential limit point** of $Y$ if there exists a sequence $x_n \in Y$ such that $x_n \to x$.*

**Proposition 4.1.18 (Adherent Point)**
*Let $X$ be a topological space. Then every sequential limit point is an adherent point. If $X$ is first countable the converse holds.*

*Proof.* The first implication is straightforward. For the latter, consider a countable local base at $x$ given by $U_n$. We may assume wlog that it is decreasing. Then by assumption we may choose $x_n \in U_n \cap Y$. For any $V \in \mathcal{B}_x$ we have $U_N \subset V$ for some $N$, and therefore $n \geq N \implies x_n \in U_n \implies x_n \in V$. Therefore $x_n \to x$. ∎

**Proposition 4.1.19 (Topological Closure)**
*Let $Y \subset X$ then the following equality holds*

$$\bigcap_{\substack{Z \supseteq Y \\ Z \text{ closed}}} Z = \{x \in X \mid x \text{ adherent point of } Y\}$$

*We denote this by $\overline{Y}$ (or $\mathrm{cl}_X(Y)$ to emphasise the ambient space $X$) and refer to it as the **closure** of $Y$ in $X$. Furthermore the following properties hold*

© David Rufino CC BY-NC-ND 4.0

*a)* $Y \subseteq \overline{Y}$ *and* $\overline{Y}$ *is closed*
*b)* $Y = \overline{Y}$ *if and only if* $Y$ *is closed*
*c)* $\left(Y \cap U \neq \varnothing \iff \overline{Y} \cap U \neq \varnothing\right)$ *for any* $U$ *open*

*When* $X$ *is first countable then* $\overline{Y}$ *is the set of sequential limit points of* $Y$*.*

*Proof.* Suppose $Z$ is a closed set containing $Y$ and $x$ is an adherent point of $Y$. Then $x \notin Z \implies x \in X \setminus Z \implies (X \setminus Z) \cap Y \neq \varnothing$ a contradiction. Conversely assume $x \notin \overline{Y}$ then there exists $Z \supseteq Y$ closed such that $x \notin Z \implies x \in X \setminus Z$. This means $x$ is not an adherent point.

a) An arbitrary intersection of closed sets is closed
b) This follows because $\overline{Y}$ is the smallest closed superset.
c) One implication is clear because $Y \subseteq \overline{Y}$. Conversely if $x \in \overline{Y} \cap U$ then $x$ must be a limit point of $Y$ hence $U \cap Y \neq \varnothing$ as required.

■

**Corollary 4.1.20**
*Let* $X$ *be a first countable topological space. Then* $\mathrm{cl}_X(Y)$ *is the set of sequential limit points of* $Y$*.*

**Proposition 4.1.21 (Closure in Subspace Topology)**
*Let* $X$ *be a topological space and* $Y$ *a subset. For* $Z \subset Y$ *we have*

$$\mathrm{cl}_Y(Z) = \mathrm{cl}_X(Z) \cap Y$$

*Proof.* By (4.1.19) $\mathrm{cl}_X(Z)$ is a closed subset of $X$ and therefore $\mathrm{cl}_X(Z) \cap Y$ is a closed subset of $Y$ in the subspace topology. Therefore $\mathrm{cl}_Y(Z) \subset \mathrm{cl}_X(Z) \cap Y$ by minimality. Similarly $\mathrm{cl}_Y(Z) = W \cap Y$ for $W$ closed in $X$. By minimality $\mathrm{cl}_X(Z) \subset W \implies \mathrm{cl}_X(Z) \cap Y \subset W \cap Y = \mathrm{cl}_Y(Z)$. ■

**Proposition 4.1.22**
*Let* $X$ *be a topological space with* $X = \bigcup_{i \in I} U_i$ *an open cover. Let* $Y \subset X$ *be a subset then* $Y$ *is closed in* $X$ *if and only if* $Y \cap U_i$ *is closed in* $U_i$ *for all* $i \in I$*.*

*Proof.* One direction is obvious by definition of the subspace topology. Conversely $U_i \setminus (Y \cap U_i)$ is open in $U_i$ and therefore in $X$. We see that

$$X \setminus Y = \bigcup_{i \in I} U_i \setminus (Y \cap U_i)$$

is open and therefore $Y$ is closed. ■

**Proposition 4.1.23**
*Let* $Y_1, ..., Y_n$ *be subsets of a topological space* $X$*. Then*

$$\overline{Y_1 \cup ... \cup Y_n} = \overline{Y_1} \cup ... \cup \overline{Y_n}$$

*Proof.* By induction it is sufficient to demonstrate the case $n = 2$. $\overline{Y_1 \cup Y_2}$ is a closed set containing both $Y_1$ and $Y_2$. Therefore by definition

$$\overline{Y_1} \cup \overline{Y_2} \subset \overline{Y_1 \cup Y_2}$$

On the other hand $\overline{Y_1} \cup \overline{Y_2}$ is closed and contains $Y_1 \cup Y_2$. Therefore the reverse inclusion follows immediately. ■

**Proposition 4.1.24 (Dense subset)**
*Let* $Y \subset X$ *then the following are equivalent*

*a)* $\overline{Y} = X$
*b)* $Y \cap U \neq \varnothing$ *for any* $U \subset X$ *open*

*In this case we say* $Y$ *is* ***dense****.*

*Proof.* *a)* $\implies$ *b)* Follows from (4.1.19).c)

*b)* $\implies$ *a)*. Suppose $Y \subseteq \overline{Y} \subsetneq X$ then $X \setminus \overline{Y}$ is an open set not intersecting $Y$ a contradiction. ■

## 4.1.3 Continuous Maps

**Proposition 4.1.25 (Continuous at a point)**
*Let* $f : X \to Y$ *a map of topological spaces and* $x \in X$*. Then the following are equivalent*

© David Rufino CC BY-NC-ND 4.0

*a)* $V \in \mathcal{U}_{f(x)} \implies f^{-1}(V) \in \mathcal{U}_x$
*b)* $V \in \mathcal{B}_{f(x)} \implies f^{-1}(V) \in \mathcal{U}_x$
*c)* $V \in \mathcal{U}_{f(x)} \implies \exists U \in \mathcal{U}_x$ s.t. $f(U) \subseteq V$
*d)* $V \in \mathcal{B}_{f(x)} \implies \exists U \in \mathcal{B}_x$ s.t. $f(U) \subseteq V$

*where $\mathcal{B}_{f(x)}$ is any local base at $f(x)$ and $\mathcal{B}_x$ is any local base at $x$. We say that $f$ is* ***continuous*** *at $x$.*

*Proof.* *a)* $\implies$ *b)* is obvious. Suppose *b)* holds and $V \in \mathcal{U}_{f(x)}$ then by definition there exists $V' \in \mathcal{B}_{f(x)}$ such that $V' \subset V$. By hypothesis $f^{-1}(V') \in \mathcal{U}_x$ from which it follows that $f^{-1}(V) \in \mathcal{U}_x$.

*a)* $\implies$ *c)* $f(f^{-1}(V)) \subseteq V$

*c)* $\implies$ *d)* By definition there exists $U' \in \mathcal{B}_x$ such that $U' \subseteq U \implies f(U') \subseteq f(U) \subseteq V$

*d)* $\implies$ *b)* Observe that $f(U) \subseteq V \implies U \subseteq f^{-1}(V)$ whence $f^{-1}(V) \in \mathcal{U}_x$ as required. ∎

**Proposition 4.1.26**
*Let $f : X \to Y$ and $g : Y \to Z$ be maps of topological spaces such that $f$ is continuous at $x$ and $g$ is continuous at $f(x)$. Then $g \circ f$ is continuous at $x$.*

**Proposition 4.1.27 (Characterisation of Continuous Maps)**
*Let $f : X \to Y$ be a map of topological spaces. Then the following are equivalent*

*a)* *$f$ is continuous at all points $x \in X$*
*b)* *$V \subset Y$ open $\implies f^{-1}(V) \subset X$ is open*
*c)* *$Z \subset Y$ closed $\implies f^{-1}(Z) \subset X$ is closed*

*Let $\mathcal{B}_1, \mathcal{B}_2$ be bases of open sets for $X$ and $Y$ respectively. Then this is equivalent to the following condition*

$$\forall V \in \mathcal{B}_2, \quad \text{and } x \in f^{-1}(V), \quad \exists U \in \mathcal{B}_1 \text{ s.t. } x \in U \subset f^{-1}(V)$$

**Proposition 4.1.28**
*Let $f : X \to Y$ and $g : Y \to Z$ be maps of topological spaces such that $f$ is continuous and $g$ is continuous. Then $g \circ f$ is continuous.*

**Proposition 4.1.29**
*Let $f : X \to Y$ be a map of topological spaces and $X = \bigcup_{i \in I} U_i$ an open cover. Then $f$ is continuous iff $f|_{U_i}$ is continuous for all $i \in I$.*

*Proof.* Observe that $f^{-1}(V) = \bigcup_{i \in I} f^{-1}(V) \cap U_i = \bigcup_{i \in I} \left(f|_{U_i}\right)^{-1}(V)$. ∎

**Proposition 4.1.30**
*Let $f : X \to Y$ be a map of topological spaces. Then $f$ is continuous iff*

$$f\left(\overline{Z}\right) \subset \overline{f(Z)}$$

*for all subsets $Z \subset X$.*

*Proof.* Suppose $f$ is continuous and $f(Z) \subset W$ is closed. Then by (4.1.27) $Z \subset f^{-1}(W)$ is closed. By definition $\overline{Z} \subset f^{-1}(W) \implies f\left(\overline{Z}\right) \subset f(f^{-1}(W)) \subset W$. As $W$ was an arbitrary closed set containing $f(Z)$ we find by definition $f\left(\overline{Z}\right) \subset \overline{f(Z)}$.

Conversely suppose $W \subset Y$ is closed and $Z \coloneqq f^{-1}(W)$. Then $f\left(\overline{Z}\right) \subset \overline{f(Z)} = \overline{W \cap \operatorname{Im}(f)} \subset W \implies f^{-1}\left(f\left(\overline{Z}\right)\right) \subset Z \implies \overline{Z} \subset Z \implies \overline{Z} = Z$ and therefore $Z$ is closed. Therefore $f$ is continuous by (4.1.27). ∎

### 4.1.4 Quasi-Homeomorphism

**Proposition 4.1.31 (Quasi-Homeomorphism)**
*Let $f : X \to Y$ be a map. The following are equivalent*

*a)* *The map $W \to f^{-1}(W)$ is a bijection between open sets*
*b)* *The map $Z \to f^{-1}(Z)$ is a bijection between closed sets*

*In this case we say $f$ is a* ***quasi-homeomorphism****. Such a map induces a bijection between irreducible sets, respectively irreducible components.*

*Proof.* This follows by taking complements and observing $X \setminus f^{-1}(A) = f^{-1}(Y \setminus A)$. ∎

© David Rufino CC BY-NC-ND 4.0

**Proposition 4.1.32 (Induced Quotient)**
*Let $\pi : X \to Y$ be a surjective map where $X$ is a topological space. The family*

$$\mathcal{T}_Y := \{U \subset Y \mid \pi^{-1}(U) \in \mathcal{T}_X\}$$

*is a well-defined topology on $Y$. Suppose that $\pi^{-1}(\pi(V)) \subset V$ for all $V$ open (resp. closed) subsets of $X$. Then $\pi$ is both open, closed and a quasi-homeomorphism.*

*Proof.* As arbitrary intersections and unions commute with inverse image we see that $\mathcal{T}_Y$ is a well-defined topology.

As $\pi$ is surjective we have $\pi(\pi^{-1}(U)) = U$ for all open subsets $U \subset Y$. Generically $V \subset \pi^{-1}(\pi(V))$ and by assumption the reverse inclusion holds. Therefore we see $\pi$ induces a quasi-homeomorphism. This also shows $\pi$ is open and therefore closed. ∎

## 4.1.5 Kolmogorov Spaces

**Definition 4.1.33**
*Let $X$ be a topological space. We say that $X$ is **Kolmogorov** or $T_0$ if for all distinct pairs $x, y \in X$ there exists an open set $U$ such that either $x \in U$ and $y \notin U$, or $x \notin U$ and $y \in U$.*

**Proposition 4.1.34 (Indistinguishable points)**
*Let $X$ be a topological space and $x, y \in X$. The following are equivalent*

*a) $\overline{\{x\}} = \overline{\{y\}}$*
*b) $x \in \overline{\{y\}}$ and $y \in \overline{\{x\}}$*
*c) Every closed $Z \subset X$ contains both or neither of $x, y$*
*d) Every open $U \subset X$ contains both or neither of $x, y$*

*We say $x$, $y$ are **topologically indistinguishable**, otherwise we say $x$ and $y$ are **distinguishable**.*

*$X$ is Kolmogorov iff every pair of points are distinguishable.*

*Being indistinguishable determines an equivalence relation on $X$, and we denote by $X_0 := X/\sim$ the **Kolmogorov Quotient**.*

*Proof.* $a) \implies b)$ Clear

$b) \implies a)$ By definition of closure $x \in \overline{\{y\}} \iff \overline{\{x\}} \subset \overline{\{y\}}$.

$c) \iff d)$ Follows by taking complements

$c) \implies b)$ Suppose $x \in Z$ for $Z$ closed then by hypothesis $y \in Z$. As $Z$ was arbitrary we see that $y \in \overline{\{x\}}$, and symmetrically $x \in \overline{\{y\}}$.

$b) \implies c)$ By assumption $x \in Z \implies y \in Z$ for $Z$ closed and symmetrically $y \in Z \implies x \in Z$. ∎

**Definition 4.1.35 (Kolmogorov Quotient)**
*Let $X$ be a topological space. The set $X_0$ of equivalence classes of topologically indistinguishable points is the **Kolmogorov Quotient***

**Proposition 4.1.36 (Kolmogorov Quotient Properties)**
*Let $X$ be a topological space and $X_0$ be the Kolmogorov Quotient with quotient map $\pi : X \to X_0$. Relative to the induced topology*

$$\mathcal{T}_{X_0} := \{W \subset X_0 \mid \pi^{-1}(W) \in \mathcal{T}_X\}$$

*$\pi$ is a quasi-homeomorphism and an open and closed map. Further $X_0$ is Kolmogorov.*

*Proof.* By (4.1.32) it is sufficient to show that $\pi^{-1}(\pi(U)) \subset U$ for all open subsets $U \subset X$. Suppose $x \in \pi^{-1}(\pi(U))$ then $\pi(x) = \pi(y)$ for some $y \in U$. Then by definition of $\pi$, $x$ and $y$ are indistinguishable and so $x \in U$.

Suppose $\pi(x) \neq \pi(y)$ then by definition there exists an open set $U \subset X$ such that $x \in U$ and $y \notin U$. As $\pi$ is open then $\pi(U)$ is open. By construction $\pi(x) \in \pi(U)$ and $\pi(y) \notin \pi(U)$. This shows that $X_0$ is Kolmogorov. ∎

**Proposition 4.1.37 (Kolmogorov Quotient Functoriality)**
*Let $f : X \to Y$ be a continuous map, then there is a unique (continuous) map $f_0 : X_0 \to Y_0$ such that the following diagram commutes*

© David Rufino CC BY-NC-ND 4.0

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \downarrow{\scriptstyle \pi} & & \downarrow{\scriptstyle \pi} \\ X_0 & \xrightarrow{f_0} & Y_0 \end{array}$$

*Proof.* Given $x, x' \in X$ and suppose $f(x) \nsim f(x')$. Then there exists an open set $U$ such that $f(x') \in U$ and $f(x) \notin U$. Therefore $x' \in f^{-1}(U)$ and $x \notin f^{-1}(U)$ and $x \nsim x'$. Equivalently $x \sim x' \implies f(x) \sim f(x')$. Therefore the map

$$f_0([x]) = [f(x)]$$

is well-defined.

For $W_0 \subset Y_0$ we have $\pi^{-1}(f_0^{-1}(W_0)) = f^{-1}(\pi^{-1}(W_0))$ is open. As $\pi$ is surjective and open $f_0^{-1}(W) = \pi(\pi^{-1}(f_0^{-1}(W_0)))$ is open, and therefore $f_0$ is continuous.

Any such $f_0$ satisfies $\{f_0(x_0)\} = \pi(f(\pi^{-1}(\{x_0\})))$ and so is unique. ∎

**Proposition 4.1.38 (Bijections with Homeomorphic Kolmogorov Quotients are Homeomorphisms)**
*Let $f : X \to Y$ be a continuous bijection of topological spaces such that the induced map $f_0 : X_0 \to Y_0$ (4.1.37) is a homeomorphism. Then $f$ is a homeomorphism.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Write $\pi_X : X \to X_0$, $\pi_Y : Y \to Y_0$ for the quotient maps, so $\pi_Y \circ f = f_0 \circ \pi_X$. For $x, x' \in X$ we may show

$$x \sim x' \iff f(x) \sim f(x')$$

For $x_0 \in X_0$ with $x_0 = \pi_X(x)$ we have

$$f(\pi_X^{-1}(x_0)) = \{f(x') \mid x' \sim x\} = \{f(x') \mid f(x') \sim f(x)\} = \pi_Y^{-1}(f_0(x_0))$$

where the last equality requires surjectivity of $f$.

It follows that $f$ is open: for $W \subset X$ open, indistinguishable points lie in exactly the same open sets, so $x \in W$ and $x \sim x'$ forces $x' \in W$; that is, $W$ is a disjoint union of the equivalence classes it meets, which are precisely the fibres of $\pi_X$ (4.1.35), $W = \bigsqcup_{x_0 \in \pi_X(W)} \pi_X^{-1}(x_0)$, so

$$f(W) = \bigcup_{x_0 \in \pi_X(W)} f(\pi_X^{-1}(x_0)) = \bigcup_{x_0 \in \pi_X(W)} \pi_Y^{-1}(f_0(x_0)) = \pi_Y^{-1}\left( \bigcup_{x_0 \in \pi_X(W)} \{f_0(x_0)\} \right) = \pi_Y^{-1}(f_0(\pi_X(W)))$$

which is open because both $f_0$ and $\pi_X$ are open and $\pi_Y$ is continuous.

A bijective, continuous, open map is a homeomorphism. ∎

**Proposition 4.1.39**
*Let $X$ be a topological space with open cover $X = \bigcup_{i \in I} U_i$. If every $U_i$ is Kolmogorov with respect to the subspace topology then so is $X$.*

*Proof.* Consider $x, y \in X$. We may assume wlog that $x, y \in U_i$ for some $i$ (otherwise we would be done). Then by assumption there exists $V \subset X$ such that $x \in U_i \cap V$ and $y \notin U_i \cap V$, whence $x \in V$ and $y \notin V$ as required. ∎

### 4.1.6 Symmetric Spaces

**Definition 4.1.40**
*Let $X$ be a topological space. We say that $x$ is **quasi-closed** if $y \in \overline{\{x\}} \implies x \in \overline{\{y\}}$.*

**Proposition 4.1.41 (Symmetric Topology)**
*Let $X$ be a topological space. Then the following are equivalent*

*a) For all $x \in X$ we have*

$$\overline{\{x\}} = \{y \in X \mid y \text{ topologically indistinguishable from } x\},$$

*b) For all $x, y \in X$ $\quad x \in \overline{\{y\}} \implies y \in \overline{\{x\}}$ (i.e. every point is **quasi-closed**),*
*c) Every open neighbourhood $U$ of $x$ contains $\overline{\{x\}}$,*
*d) $X$ is partitioned by sets of the form $\overline{\{x\}}$, indexed by $X_0$.*

© David Rufino CC BY-NC-ND 4.0

*In this case we say that $X$ is **symmetric** or $R_0$. Further $x \in \overline{\{y\}}$ if and only if $x$ and $y$ become equal in $X_0$.*

**Proposition 4.1.42**
*Let $X$ be a symmetric topological space and $x, y \in X$. Then the following are equivalent*

*a) $x$ and $y$ are topologically indistinguishable,*
*b) $x \in \overline{\{y\}}$,*
*c) $y \in \overline{\{x\}}$,*
*d) $\overline{\{x\}} = \overline{\{y\}}$,*
*e) $x$ and $y$ become equal in the Kolmogorov Quotient $X_0$*

**Proposition 4.1.43 (Open Subspace is Symmetric)**
*Let $X$ be a symmetric space and $U \subset X$ an open subset. Then $U$ is symmetric and for every $x \in U$ we have*

$$\mathrm{cl}_X(\{x\}) = \mathrm{cl}_U(\{x\})$$

*Proof.* By (4.1.21) we have $\mathrm{cl}_U(\{x\}) = \mathrm{cl}_X(\{x\}) \cap U$, so these are equal by (4.1.41).c). We conclude that the sets $\overline{\{x\}}$ partition $U$ so that $U$ is also symmetric. ∎

**Proposition 4.1.44 (Symmetric Open Cover $\implies$ Symmetric)**
*Let $X$ be a topological space with open cover $\bigcup_{i=1}^n U_i$ such that $U_i$ are symmetric in the subspace topology. Then $X$ is symmetric.*

*Proof.* Suppose $x \in \mathrm{cl}_X(\{y\})$ then by (4.1.43) we have $x \in \mathrm{cl}_{U_i}(\{y\})$, whence $y \in \mathrm{cl}_{U_i}(\{x\})$ by (4.1.41) and $y \in \mathrm{cl}_X(\{x\})$. Therefore by (4.1.41) we have $X$ is symmetric. ∎

**Proposition 4.1.45**
*Let $X$ be a topological space. Then the following are equivalent*

*a) $X$ is both Symmetric and Kolmogorov*
*b) Every point $x \in X$ is closed.*

**Corollary 4.1.46**
*Let $X$ be a symmetric topological space. Then every point of $X_0$ is closed.*

### 4.1.7 Hausdorff Spaces

**Definition 4.1.47**
*Let $X$ be a topological space. We say that $X$ is **Hausdorff** if for all distinct pairs $x, y \in X$ there exists open neighbourhoods $U, V$ of $x$ and $y$ respectively such that $U \cap V = \varnothing$.*

*Observe that **Hausdorff** $\implies$ **Kolmogorov**, **Symmetric**, but not conversely (e.g. co-finite topology).*

**Proposition 4.1.48**
*Let $X$ be a Hausdorff topological space. Then $X$ is both Kolmogorov and Symmetric. Further singletons are closed.*

### 4.1.8 Convergent Sequences

**Proposition 4.1.49 (Convergent Sequence)**
*Let $X$ be a topological space, $(x_n)$ a sequence in $X$ and $x \in X$. Let $\mathcal{B}_x$ be a local base for $x$. Then the following are equivalent*

*a) For every $V \in \mathcal{U}_x$ there exists $N$ such that*

$$n \geq N \implies x_n \in V$$

*b) For every $V \in \mathcal{B}_x$ there exists $N$ such that*

$$n \geq N \implies x_n \in V$$

*c) The function*

$$\begin{aligned} \mathbb{N} \cup \{\infty\} &\to X \\ n &\to x_n \\ \infty &\to x \end{aligned}$$

*is continuous at $\infty$ with respect to the topology on $\mathbb{N} \cup \{\infty\}$ given by open sets of the form $\{N, N+1, ...,\} \cup \{\infty\}$ and arbitrary subsets of $\mathbb{N}$.*

*In this case we say that $x_n \to x$.*

© David Rufino CC BY-NC-ND 4.0

**Proposition 4.1.50 (Continuous $\implies$ Sequentially Continuous)**
*Let $f : X \to Y$ be a continuous map and $x_n \to x$ a sequence in $X$. Then $f(x_n) \to f(x)$.*

*Proof.* If we regard the sequence as a continuous map $\mathbb{N} \cup \infty$ (4.1.49) then the result follows from (4.1.26). ∎

**Proposition 4.1.51 (Limits are Unique in Kolmogorov Spaces)**
*Let $X$ be a Kolmogorov topological space. Then limits of sequences are unique.*

**Proposition 4.1.52 (Sequentially Continuous $\implies$ Continuous)**
*Let $X, Y$ be topological spaces, $f : X \to Y$ a map and $x \in X$ a point with a* countable *local base $\mathcal{B}_x$. Suppose that for all sequences $x_n \to x$ we have $f(x_n) \to f(x)$. Then $f$ is continuous at $x$.*

*In particular if $X$ is first countable then $f : X \to Y$ is continuous iff it is sequentially continuous.*

*Proof.* Consider a point $x \in X$ and $y \coloneqq f(x)$. By hypothesis $\mathcal{B}_x = \{U_n\}$ is countable. By replacing with the base $U_n' \coloneqq \bigcap_{k=1}^{n} U_k$ we may suppose that

$$U_1 \supseteq U_2 ... \supseteq U_n \supseteq ...$$

is decreasing. Suppose $f$ is not continuous at $x$ then by (4.1.25).d) there exists $V \in \mathcal{U}_{f(x)}$ such that $f(U_n) \setminus V \neq \varnothing$ for all $n$. Therefore we may choose $x_n \in U_n$ such that $f(x_n) \notin V$. Consider $U \in \mathcal{U}_x$ then by definition there is some $N$ such that $U_N \subseteq U$. Furthermore $n \geq N \implies U_n \subseteq U_N \subseteq U \implies x_n \in U$. Therefore $x_n \to x$. By construction $f(x) \in V$ but $f(x_n) \notin V$ for all $n$ and in particular $f(x_n) \nrightarrow f(x)$. ∎

## 4.1.9 Connected Topological Spaces

**Proposition 4.1.53 (Connected Topological Space)**
*Let $X$ be a topological space. Then the following are equivalent*

*a) $X$ cannot be written as the disjoint union $U \sqcup V$ of two non-empty open subsets $U, V \subset X$*
*b) There are no proper non-empty subsets which are both open and closed*

*In this case we say that $X$ is **connected**.*

**Proposition 4.1.54 (Connected Subset)**
*Let $X$ be a topological space and $Z \subset X$. Then the following are equivalent*

*a) $Z$ is connected in the subspace topology*
*b) $Z$ cannot be written as a disjoint $(Z \cap U) \sqcup (Z \cap V)$ with both $Z \cap U, Z \cap V \neq \varnothing$*

**Proposition 4.1.55**
*Let $X$ be a topological space and $\{Z_i\}_{i \in I}$ a family of connected subsets such that $\bigcap_{i \in I} Z_i \neq \varnothing$. Then $\bigcup_{i \in I} Z_i$ is connected.*

**Definition 4.1.56 (Connected Component)**
*Let $X$ be a topological space. We say $Z \subset X$ is a **connected component** if it is both connected and not properly contained in a connected subset.*

**Proposition 4.1.57**
*Let $X$ be a topological space. Then every connected subset $Z$ is contained in a unique connected component.*

*Proof.* Define

$$\overline{Z} \coloneqq \bigcup \{Z' \supset Z \mid Z' \subset X \text{ connected}\}$$

then by (4.1.55) this is connected. If $Z'' \supset \overline{Z} \supset Z$ is connected then by construction $Z'' \subset \overline{Z}$, so by definition $\overline{Z}$ is a connected component. Let $\overline{Z}'$ be another connected component containing $Z$, then by the same argument $\overline{Z}' \subset \overline{Z}$ and by definition $\overline{Z}' = \overline{Z}$. ∎

**Proposition 4.1.58**
*Let $X$ be a topological space, then the connected components partition $X$.*

*We denote this collection by $\pi_0(X)$.*

*Proof.* Evidently $\{x\}$ is connected so that the connected components cover $X$ by (4.1.57).

Suppose we have two connected components $Z_1, Z_2$ are not disjoint. Then by (4.1.55) $Z_1 \cup Z_2$ is connected. By maximality we have $Z_1 \cup Z_2 = Z_1 = Z_2$ as required. ∎

© David Rufino CC BY-NC-ND 4.0

### 4.1.10 Irreducible Topological Spaces

References:

- [Bou98b] (Chapter II §4.2)
- [Stacks] (Tag 004U)

**Proposition 4.1.59 (Irreducible space)**
*Let $X$ be a topological space. Then the following are equivalent*

*a) $X = Z_1 \cup Z_2$ closed implies either $Z_1 = X$ or $Z_2 = X$*
*b) $U, V \neq \varnothing \implies U \cap V \neq \varnothing$ for open sets $U, V$*
*c) $U \neq \varnothing \implies \overline{U} = X$ i.e. every non-empty open set is dense*

*and we say $X$ is irreducible.*

*Proof.* *a)* $\implies$ *b)* Suppose $U, V$ are open sets such that $U \cap V = \varnothing$. Then $X = (X \setminus U) \cup (X \setminus V)$. By hypothesis $X = (X \setminus U)$ or $X = (X \setminus V)$ whence either $U$ or $V$ is empty.

*b)* $\implies$ *a)* Similar.

*c)* $\iff$ *b)* Follows directly from (4.1.24) ■

**Proposition 4.1.60 (Irreducible Subset)**
*Let $Y \subset X$ be a subset of a topological space. Then the following conditions on $Y$ are equivalent*

*a) $Y$ is irreducible in the subspace topology.*
*b) $Y \subseteq Z_1 \cup Z_2 \implies Y \subseteq Z_1$ or $Y \subseteq Z_2$ where $Z_1, Z_2$ are closed subsets of $X$*
*c) $U \cap Y \neq \varnothing, V \cap Y \neq \varnothing \implies (U \cap V) \cap Y \neq \varnothing$ for $U, V$ open*

*and we say $Y$ is an **irreducible subset**.*

*Proof.* *a)* $\implies$ *b)*. Suppose that $Y$ is irreducible in the subspace topology and $Y \subseteq Z_1 \cup Z_2$. This implies $Y = (Z_1 \cap Y) \cup (Z_2 \cap Y)$ is a decomposition of closed sets. So either $Z_1 \cap Y = Y$ or $Z_2 \cap Y = Y \implies Y \subseteq Z_1$ or $Y \subseteq Z_2$ as required.

*b)* $\implies$ *a)*. Suppose that $Y = (Z_1 \cap Y) \cup (Z_2 \cap Y)$. Then $Y \subseteq Z_1 \cup Z_2$, and for example $Y \subseteq Z_1$, which implies $Z_1 \cap Y = Y$. ■

**Proposition 4.1.61**
*Let $Y \subset X$ be a **closed** subset then the following are equivalent*

*a) $Y$ is an irreducible subset*
*b) $Y = Z_1 \cup Z_2 \implies Y = Z_1$ or $Y = Z_2$ where $Z_1, Z_2$ are closed subsets of $X$*
*c) $Y \subseteq Z_1 \cup Z_2 \implies Y \subseteq Z_1$ or $Y \subseteq Z_2$ where $Z_1, Z_2$ are closed subsets of $X$*

*In other words in the lattice of closed subsets, the irreducible subsets are precisely the join-prime subsets.*

*Proof.* *a)* $\implies$ *b)*. Clearly $Z_1, Z_2$ are also closed subsets of $Y$, so the result follows by definition.

*b)* $\iff$ *c)*. This is (2.4.2).

*c)* $\implies$ *a)*. This was already proven in (4.1.60). ■

**Remark 4.1.62**
*Singletons $\{x\}$ are always irreducible.*

**Definition 4.1.63 (Irreducible Component)**
*We say that $Y$ is an irreducible component if it is a maximal irreducible subset.*

**Proposition 4.1.64**
*Let $f : X \to Y$ be a continuous map and $Z \subset X$ irreducible. Then $f(Z)$ is irreducible in $Y$.*

*Proof.* Suppose $f(Z) \subset Y_1 \cup Y_2$ for $Y_1, Y_2$ closed. Then $Z \subset f^{-1}(f(Z)) \subset f^{-1}(Y_1) \cup f^{-1}(Y_2)$. Then by (4.1.60).b) wlog $Z \subset f^{-1}(Y_1)$ therefore $f(Z) \subset f(f^{-1}(Y_1)) \subset Y_1$. Therefore by the same criterion $f(Z)$ is irreducible. ■

**Proposition 4.1.65**
*Let $Y \subset X$. Then $Y$ is irreducible iff $\overline{Y}$ is.*

*Proof.* This follows from (4.1.60).b) and (4.1.19) that $Y \subset Z \iff \overline{Y} \subset Z$. ■

**Proposition 4.1.66 (Decomposition into Irreducible Components)**
*A topological space $X$ may be decomposed into irreducible components. More precisely*

© David Rufino CC BY-NC-ND 4.0

*a) Every irreducible component is closed*
*b) Every irreducible closed subset is contained in an irreducible component*
*c) $X$ is the union of irreducible components*

*Proof.* We prove each in turn

a) By (4.1.65) $\overline{Y}$ is irreducible and closed, so by maximality $Y = \overline{Y}$ is closed.
b) We may show that the lattice of closed subsets is chain complete so we may use to show that the lattice of irreducible closed subsets is also chain complete (2.4.4). Therefore we may apply (2.1.56) to find an irreducible component.
c) As $\{x\}$ is irreducible every element is contained in an irreducible component by $b$).

■

**Corollary 4.1.67**
*For $x \in X$ the closure $\overline{\{x\}}$ is an irreducible closed subset.*

**Corollary 4.1.68**
*$X$ is irreducible if and only if it has a single irreducible component.*

**Definition 4.1.69 (Generic Point)**
*Let $Z$ be an irreducible closed subset of $X$, then we say $\eta \in X$ is a generic point of $Z$ if $Z = \overline{\{\eta\}}$.*

**Proposition 4.1.70**
*Let $X$ be an irreducible space then every open subset $U$ is irreducible and connected.*

*Proof.* By (4.1.59) $U$ is dense in $X$. Therefore by (4.1.65) $U$ is irreducible.

Suppose that $U$ is not connected then by (4.1.54) $U = (U \cap V) \sqcup (U \cap W)$ with $V, W$ open subsets of $X$ and $U \cap V, U \cap W$ non-empty. Then as $U$ is irreducible we have $U \cap V \cap W \neq \varnothing$, a contradiction. ■

**Proposition 4.1.71**
*Let $X = \bigcup_{i=1}^{n} X_i$ where $X_i$ are irreducible closed subsets, and mutually incomparable. Then the $X_i$ are the irreducible components of $X$.*

*Proof.* For any irreducible closed subset $E$ we have by (4.1.61) $E \subset X_i$ for some $i$. In particular every irreducible component is contained in some $X_i$. On the other hand every $X_j$ is contained in an irreducible component by (4.1.66). By incomparability we deduce that each $X_i$ is an irreducible component, and that the set is complete. ■

**Proposition 4.1.72**
*Let $Y \subset X$ such that $Y = \bigcup_{i=1}^{n} Y_i$ where $Y_i$ are the irreducible components of $Y$. Then the set of irreducible components of $\overline{Y}$ is $\left\{\overline{Y_i}\right\}_{i=1\ldots n}$, and these are distinct.*

*Proof.* By (4.1.65) $\overline{Y_i}$ are irreducible subsets which, by (4.1.23) cover $\overline{Y}$. By (4.1.66) $Y_i$ is closed in $Y$ and so by (4.1.21) $\overline{Y_i} \cap Y = Y_i$. This shows that the $\overline{Y_i}$ are distinct and also incomparable. Then (4.1.71) shows that these must be the irreducible components. ■

**Proposition 4.1.73**
*Let $U \subset X$ be an open subset. There is an order isomorphism*

$$\begin{aligned} \{Y \subset U \mid Y \text{ irreducible, closed and non-empty}\} &\longleftrightarrow \{Z \subset X \mid Z \text{ irreducible and closed and } Z \cap U \neq \varnothing\} \\ Y &\to \operatorname{cl}_X(Y) \\ Z \cap U &\leftarrow Z \end{aligned}$$

*which restricts to irreducible components.*

*Proof.* By (4.1.65) the first map is well-defined. Conversely $Z \cap U$ is closed in $U$ and open in $Z$, so irreducible by (4.1.70).

By (4.1.59) $Z \cap U$ is dense in $Z$, that is $Z = \operatorname{cl}_Z(Z \cap U) \overset{(4.1.21)}{=} \operatorname{cl}_X(Z \cap U) \cap Z \implies Z \subset \operatorname{cl}_X(Z \cap U)$, and so by minimality they are equal.

By (4.1.21) $\operatorname{cl}_X(Y) \cap U = \operatorname{cl}_U(Y) = Y$. Therefore the maps are mutually inverse and order preserving. ■

**Proposition 4.1.74**
*Suppose $X$ is a topological space with open cover $\bigcup_{i \in I} U_i$ such that*

*a) $U_i$ is irreducible for all $i \in I$ and*

© David Rufino CC BY-NC-ND 4.0

*b) $U_i \cap U_j$ is non-empty for all $i, j \in I$.*

*Then $X$ is irreducible.*

*Proof.* Suppose that $X = Z_1 \cup Z_2$. Pick some $i \in I$ then by (4.1.60) wlog $U_i \cap Z_1 \neq \varnothing$. Similarly for all $j \in J$ by assumption $U_i \cap U_j$ is a non-empty open subset of $U_j$ and therefore dense (4.1.59). Further $U_i \cap U_j$ is contained in the closed subset $Z_1 \cap U_j$. We conclude that $U_j \subset Z_1 \cap U_j$. Therefore $X \subset Z_1$. By (4.1.60) we further conclude that $X$ is irreducible. ∎

## 4.1.11 Noetherian Topological Spaces

**Definition 4.1.75 (Noetherian)**
*A topological space $X$ is **Noetherian** if the lattice of closed subsets is Artinian (i.e. satisfies the descending chain condition).*

**Proposition 4.1.76 (Decomposition into Irreducibles)**
*Let $X$ be a Noetherian topological space. Then every closed subset $Y$ may be expressed uniquely as a finite, incomparable union of irreducible closed subsets. These are precisely the irreducible components of $Y$.*

*In particular $X$ has only finitely many irreducible components.*

*Proof.* The lattice of closed subsets is distributive and Artinian by definition. Therefore the result follows from (4.1.61) and (2.4.7). ∎

**Proposition 4.1.77**
*Let $X$ be a Noetherian topological space and $Y \subset X$. Then $Y$ is Noetherian.*

**Proposition 4.1.78**
*Let $X = \bigcup_{i=1}^{N} X_i$ be a topological space such that $X_i$ is Noetherian under the subspace topology. Then $X$ is Noetherian.*

*Proof.* Let $\{Z_n\}_{n\in\mathbb{N}}$ be a descending chain of closed subsets of $X$. By definition $\{Z_n \cap X_i\}_{n\in\mathbb{N}}$ terminates for $n \geq m_i$. Then for $n \geq \max(m_1, ..., m_N)$ the sequence $\{Z_n\}_{n\in\mathbb{N}}$ also terminates. ∎

**Proposition 4.1.79**
*Let $X$ be a Noetherian topological space and $U \subset X$ an open subset. Then there is a bijection*

$$\begin{aligned} \{Y \subset U \mid Y \text{ closed and non-empty}\} &\longleftrightarrow \{Z \subset X \mid Z \text{ closed and every irreducible component meets } U\} \\ Y &\to \ \mathrm{cl}_X(Y) \\ Z \cap U &\leftarrow \ Z \end{aligned}$$

*which is order preserving and restricts to closed irreducible subsets and irreducible components.*

*Proof.* By (4.1.77) and (4.1.76) $Y = Y_1 \cup ... \cup Y_n$ where $Y_i$ are the irreducible components of $Y$. Then by (4.1.72) the irreducible components of $\overline{Y}$ are $\overline{Y_i}$, which evidently intersect $U$. Furthermore by definition $Z \cap U$ is closed in $U$ and the maps are well-defined.

By (4.1.73) the maps restrict to closed irreducible subsets (resp. components) and are bijections, in particular for $Z$ irreducible closed we have $\overline{Z \cap U} = Z$.

By (4.1.21) $\overline{Y} \cap U = Y$. For the case $Z$ is closed then as before $Z = Z_1 \cup ... \cup Z_n$ is a decomposition into irreducible components. By assumption $Z_i \cap U \neq \varnothing$ for all $i = 1...n$. Therefore by the irreducible case (4.1.73) and (4.1.23)

$$\overline{Z \cap U} = \overline{Z_1 \cap U} \cup ... \cup \overline{Z_n \cap U} = Z_1 \cup ... \cup Z_n = Z.$$

Therefore the maps are mutually inverse as required. ∎

## 4.1.12 Krull Dimension

References:

- Éléments de géométrie algébrique IV [Gro64] (Chap. 0 §**14.4.1**)
- Some remarks on biequidimensionality of topological spaces and Noetherian schemes [Hei17]

For a Noetherian topological space $X$ the closed subsets form an Artinian, distributive lattice. Furthermore the irreducible subsets are precisely the join-prime elements by (4.1.61). Therefore we may use the notions of Krull Lattice developed in Foundations.tex's Krull Dimension section ((2.5.2)).

**Definition 4.1.80 (Chain of irreducibles)**
*Let $X$ be a Noetherian topological space. A **chain** of irreducible closed subsets*

© David Rufino CC BY-NC-ND 4.0

$$Z_0 \subsetneq Z_1 \subsetneq ... \subsetneq Z_n$$

*is said to have* ***length*** $n$*. A chain is* ***saturated*** *if there is no proper refinement, that is if* $Y$ *is irreducible then*

$$Z_i \subseteq Y \subseteq Z_{i+1} \implies Y = Z_i \text{ or } Y = Z_{i+1}.$$

*If in addition* $Z_n$ *(resp.* $Z_0$*) is maximal (resp. minimal) then the chain is* ***maximal****.*

**Definition 4.1.81 (Krull Lattice of Closed Subsets)**
*Let* $X$ *be a topological space.*

- *The* ***Krull dimension*** $\dim(X)$ *of* $X$ *is the maximal length of all chains of irreducible closed subsets. Note this may be* $\infty$*.*
- *The* ***height*** *or* ***codimension*** *of an irreducible closed subset* $Y \subseteq X$*, denoted* $\operatorname{codim}(Y, X)$*, is the maximal length of chains of irreducible subsets containing* $Y$*.*
- *When* $Y$ *is not irreducible we write*

$$\operatorname{codim}(Y, X) = \inf_\alpha \operatorname{codim}(Y_\alpha, X)$$

*where* $Y_\alpha$ *varies among the irreducible components of* $Y$*.*

*If* $\dim(X) < \infty$ *then we say* $X$ *is* ***finite-dimensional****. In this case it's clear the closed subsets of a topological space form a Krull Lattice where the irreducible closed subsets are precisely the join-prime elements of the lattice.*

*Note any saturated chain for* $Y \subset X$ *must start at* $Y$ *and terminate at an irreducible component of* $X$*. In particular if* $X$ *is irreducible then a saturated chain must terminate at* $X$*.*

**Proposition 4.1.82 (Extending Chains)**
*Let* $X$ *be a finite-dimensional topological space. Then*

*a) Every chain is contained in a saturated chain with the same endpoints*
*b) Every chain is contained in a maximal chain*

**Proposition 4.1.83 (Simple properties of (co-)dimension)**
*Let* $X$ *be a topological space. Then the following properties hold*

*a)* $\dim(X) = \sup_\alpha \dim(X_\alpha)$ *where* $X_\alpha$ *are the irreducible components of* $X$
*b)* $\operatorname{codim}(Y, X) = \sup_\alpha \operatorname{codim}(Y, X_\alpha)$ *where* $X_\alpha$ *are the irreducible components of* $X$ *containing* $Y$
*c)* $\dim(Y) + \operatorname{codim}(Y, X) \leq \dim(X)$ *for every closed subset* $Y \subset X$
*d)* $\operatorname{codim}(Y, Z) + \operatorname{codim}(Z, T) \leq \operatorname{codim}(Y, T)$ *for* $Y \subset Z \subset T$ *irreducible closed subsets*
*e)* $Y \subsetneq Z$ *is a saturated chain of irreducible closed subsets if and only if* $\operatorname{codim}(Y, Z) = 1$*.*
*f) Let* $Y$ *be a closed subset of* $X$*. Then* $\operatorname{codim}(Y, X) = 0 \iff$ *every irreducible component of* $Y$ *is an irreducible component of* $X$*. In particular if* $X$ *is irreducible then* $\operatorname{codim}(Y, X) = 0 \iff Y = X \iff \dim(X) = \dim(Y)$*.*

*In particular if* $X$ *is finite-dimensional then all codimensions are also finite.*

**Proposition 4.1.84**
*Let* $X$ *be an irreducible topological space and* $Y \subset X$ *a closed subset. Then* $\dim(Y) \leq \dim(X)$ *with equality iff* $Y = X$*.*

*Proof.* The inequality follows from (4.1.83).b). If $\dim(Y) = \dim(X)$ then by c) $\operatorname{codim}(Y, X) = 0$ and so by f) we have $Y = X$. ∎

**Definition 4.1.85 (Properties)**
*Let* $X$ *be a topological space of finite dimension. Then we say* $X$ *is*

- ***Equidimensional*** *if all irreducible components of* $X$ *have the same dimension*
- ***Equicodimensional*** *if* $\operatorname{codim}(Y, X)$ *is constant as* $Y$ *varies over minimal irreducible subsets of* $X$
- ***Biequidimensional*** *if all maximal chains of irreducible subsets have the same length.*
- ***Quasi-Biequidimensional*** *if every irreducible component is biequidimensional*
- ***Catenary*** *if any two saturated chains with the same endpoints, say* $Y$ *and* $Z$*, have the same length, namely* $\operatorname{codim}(Y, Z)$

*Observe that* ***quasi-biequidimensional*** *+* ***equidimensional*** $\iff$ ***biequidimensional*** *and* ***irreducible*** $\implies$ ***equidimensional***

© David Rufino CC BY-NC-ND 4.0

**Proposition 4.1.86 (Equivalent characterisations of biequidimensional)**
*Suppose $X$ is a topological space of finite dimension. Then the following are equivalent*

*a) $X$ is quasi-biequidimensional*
*b) $X$ is catenary and every irreducible component is equicodimensional*
*c) $X$ satisfies the codimension formula for $Z \subset Y$ irreducible*

$$\dim(Y) = \dim(Z) + \operatorname{codim}(Z, Y)$$

*d) $X$ satisfies c) in the case $\operatorname{codim}(Z, Y) = 1$*

*Furthermore for irreducible subsets $Z \subset Y$*

$$\operatorname{codim}(Z) = \operatorname{codim}(Z, Y) + \operatorname{codim}(Y)$$

*Proof.* This is a translation of (2.5.11) to the topological case. ∎

**Proposition 4.1.87 (Codimension Formula)**
*Suppose $X$ is a quasi-biequidimensional topological space. Then for every subset $Z \subset X$ which is irreducible and closed, and every closed subset $Y \subset Z$ the following relations hold.*

$$\dim(Z) = \dim(Y) + \operatorname{codim}(Y, Z)$$
$$\operatorname{codim}(Y) = \operatorname{codim}(Y, Z) + \operatorname{codim}(Z)$$

*Further every closed subset is also quasi-biequidimensional.*

*Proof.* This follows from (2.5.13) and (2.5.15). ∎

**Proposition 4.1.88**
*Suppose $X = \bigcup_{i \in I} U_i$. Then $\dim(X) = \sup_{i \in I} \dim(U_i)$.*

*Proof.* By (4.1.73) a chain of irreducible closed subsets of $U_i$ lifts to a chain of the same length in $X$. Therefore $\dim(U_i) \leq \dim(X)$ for all $i \in I$ whence $\sup_{i \in I} \dim(U_i) \leq \dim(X)$. Similarly given a chain of irreducible closed subsets

$$X_0 \subsetneq ... \subsetneq X_n \subseteq X$$

choose $i$ such that $X_0 \cap U_i \neq \varnothing$. Then (4.1.79) shows that this restricts to a chain of the same length in $U_i$. Therefore $\dim(X) \leq \dim(U_i) \leq \sup_{i \in I} \dim(U_i)$. ∎

**Proposition 4.1.89**
*Let $X$ be a catenary topological space and $U \subset X$ an open subset. Then $U$ is catenary.*

*Proof.* Consider $Z \subset Y \subset U$ irreducible closed subsets. Then for any saturated chain

$$Z = Z_0 \subsetneq ... \subsetneq Z_n = Y$$

we have by (4.1.73)

$$\overline{Z} = \overline{Z_0} \subsetneq ... \subsetneq \overline{Z_n} = \overline{Y}$$

a saturated chain in $X$. As $X$ is catenary we find $n = \operatorname{codim}\left(\overline{Z}, \overline{Y}\right)$. As the chain was arbitrary we find $U$ is catenary. ∎

**Lemma 4.1.90**
*Suppose that $X$ is a topological space and $x \in X$ is quasi-closed. Then $\dim\left(\overline{\{x\}}\right) = 0$.*

*Conversely suppose $Z$ is an irreducible closed set of dimension 0. Then $Z = \overline{\{z\}}$ for all $z \in Z$ and $Z$ consists of quasi-closed elements.*

*Proof.* By (4.1.67) $\overline{\{x\}}$ is irreducible. Suppose $\varnothing \neq Z \subset \overline{\{x\}}$ is closed then $y \in Z \Longrightarrow x \in \overline{\{y\}} \subset Z$ whence $\overline{\{x\}} = Z$. This shows that $\dim\left(\overline{\{x\}}\right) = 0$.

Conversely if $\dim(Z) = 0$ then $\overline{\{z\}} \subset Z$ is an irreducible subset for all $z \in Z$, whence they are equal. This also shows that $z$ is quasi-closed. ∎

**Proposition 4.1.91**
*Let $X$ be a Noetherian topological space and $Z$ a closed subset. Then*

$$\dim(Z) = 0 \iff Z = \overline{\{z_1\}} \cup ... \cup \overline{\{z_n\}}$$

*for some $z_1, ..., z_n \in Z$ quasi-closed.*

© David Rufino CC BY-NC-ND 4.0

*Proof.* By (4.1.76) $Z = Z_1 \cup ... \cup Z_n$ for $Z_i$ is a decomposition into irreducible components. By (4.1.83).a) $\dim(Z_i) = 0$, so by (4.1.90) $Z_i = \overline{\{z_i\}}$.

Conversely $Z = \overline{\{z_1\}} \cup ... \cup \overline{\{z_n\}}$ is a decomposition into irreducible closed subsets which are incomparable (4.1.90). By (4.1.71) these are the irreducible components. Therefore

$$\dim(Z) \overset{(4.1.83).a)}{=} \sup_i \dim\left(\overline{\{z_i\}}\right) \overset{(4.1.90)}{=} 0$$

■

**Proposition 4.1.92 (Curves have co-finite topology)**
*Let $X$ be an irreducible, Noetherian and symmetric topological space such that $\dim(X) = 1$. Then $X_0$ has the cofinite topology.*

*Proof.* We may reduce to the case $X = X_0$ is Kolmogorov, then by (4.1.45) every point $x \in X$ is closed.

Suppose $Z \subset X$ is finite then $Z = \bigcup_{z \in Z} \{z\}$ is a finite union of closed subsets and therefore closed. Conversely if $Z \subset X$ is closed and proper then by (4.1.83).f) $\dim(Z) = 0$ and therefore $Z$ is finite by (4.1.91). ■

**Lemma 4.1.93 (Cofinite Topology is Quasi-Compact)**
*Let $X$ be a set equipped with the cofinite topology. Then $X$ is quasi-compact.*

*Proof.* Let $\{U_i\}_{i \in I}$ be an open cover of $X$ and choose any nonempty $U_{i_0}$ from it. Then $X \setminus U_{i_0}$ is finite by definition of the cofinite topology, say $\{x_1, ..., x_n\}$. For each $k$ choose $U_{i_k} \ni x_k$ from the cover. Then $\left\{U_{i_0}, U_{i_1}, ..., U_{i_n}\right\}$ is a finite subcover. ■

**Proposition 4.1.94**
*Let $X$ be a symmetric topological space. Then there is a correspondence*

$$\begin{aligned} X_0 &\longleftrightarrow \{W \subset X \mid \dim(W) = 0 \text{ closed and irreducible}\} \\ x &\longrightarrow \overline{\{x\}} \end{aligned}$$

## 4.1.13 Product Topology

**Definition 4.1.95 (Product Topology)**
*Let $\{X_i\}_{i \in I}$ be a family of topological spaces. The **product topology** is the topology generated by the base of open sets of the form*

$$\prod_{i \in I} U_i$$

*where $U_i \subset X_i$ is open and only finitely many are proper subsets of $X$.*

**Proposition 4.1.96**
*Let $\{X_i\}_{i \in I}$ be family of topological spaces. The family of sets given in (4.1.95) is a base. The topology generated is the smallest family of subsets for which the projection maps*

$$\pi_i : \prod_{i \in I} X_i \to X_i$$

*are continuous. Furthermore if each $X_i$ is Hausdorff (resp. Kolmogorov), then so is the product.*

**Proposition 4.1.97 (Product of Sequences)**
*Let $X_1, ..., X_m$ be a family of topological spaces and $a_n^i \to \alpha^i$ for $i = 1...m$. Then $(a_n^1, ..., a_n^m) \to (\alpha^1, ..., \alpha^m)$.*

**Proposition 4.1.98 (Diagonal is Closed)**
*Let $X$ be a topological space. Then $X$ is Hausdorff if and only if $\{(x, x) \mid x \in X\}$ is closed in $X \times X$.*

**Proposition 4.1.99**
*Let $f, g : X \to Y$ be continuous maps and $Y$ a Hausdorff space. Then*

$$\{x \in X \mid f(x) = g(x)\}$$

*is closed.*

*Proof.* The level set is $(f \times g)^{-1}(\Delta_Y)$ which by (4.1.98) is closed. ■

© David Rufino CC BY-NC-ND 4.0

### 4.1.14 Quasi-Compactness

**Definition 4.1.100 (Cluster Point of a Sequence)**
*Let $(x_n)$ be a sequence in a topological space $X$. We say that $x \in X$ is*

*a) a **cluster point** (or **accumulation point**) of $x_n$ if for every $U \in \mathcal{O}_x$ the set $\{m \in \mathbb{N} \mid x_m \in U\}$ is infinite*
*b) a **subsequential limit** of $x_n$ if there exists some subsequence $x_{n_k} \to x$*

*In general the latter implies the former, and if $X$ is first countable the concepts are equivalent as we will see.*

**Lemma 4.1.101 (Cluster Point is Subsequential Limit Point)**
*Let $X$ be a first countable topological space and $x$ a cluster point of a sequence $(x_n)$. Then there exists some subsequence $x_{n_k}$ such that $x_{n_k} \to x$.*

*Proof.* Let $(U_n)$ be a local base at $x$. By replacing it with $U_n' \coloneqq U_1 \cap ... \cap U_n$ we may assume it is without loss of generality decreasing. Define $n_0 = 0$ and $n_k$ inductively by

$$n_k \coloneqq \min\{m > n_{k-1} \mid x_m \in U_k\}$$

which is well-defined precisely because $x$ is a cluster point. Then $x_{n_k} \in U_k$ which shows that $x_{n_k} \to x$ as required. ■

**Definition 4.1.102**
*Let $X$ be a topological space. We say it is*

- ***Quasi-Compact** if every open cover of $X$ has a finite subcover*
- ***Countably Quasi-Compact** if every* countable *open cover of $X$ has a finite subcover*
- ***Sequentially Quasi-Compact** if every sequence $(x_n)$ in $X$ has a convergent subsequence*

*If in addition $X$ is Hausdorff (e.g. a metric space) we say that $X$ is compact (resp. countably compact, sequentially compact).*

**Proposition 4.1.103 (Criteria for Countable Compactness)**
*Let $X$ be a topological space. Then the following are equivalent*

*a) $X$ is countably quasi-compact*
*b) Let $\{Z_n\}_{n\in\mathbb{N}}$ be a countable collection of closed subsets such that every finite set has non-empty intersection. Then the intersection $\bigcap_{n\in\mathbb{N}} Z_n$ is non-empty*
*c) For every increasing sequence of proper open subsets*

$$U_1 \subseteq U_2 \subseteq ... \subseteq U_n \subseteq ...$$

*the union $\bigcup_{n=1}^{\infty} U_n$ is proper.*
*d) For every decreasing sequence of non-empty closed subsets*

$$Z_1 \supseteq Z_2 \supseteq ... \supseteq Z_n \supseteq ...$$

*the intersection $\bigcap_{n=1}^{\infty} Z_n$ is non-empty*

*Proof.* $a) \iff b)$ and $c) \iff d)$ are simple consequences of De Morgan's laws. For $a) \implies c)$ suppose $\bigcup_{n=1}^{\infty} U_n = X$ then by assumption $U_{n_1} \cup ... \cup U_{n_k} = X$, and by the increasing assumption $U_{n_k} = X$ which is a contradiction.

For $c) \implies a)$, suppose we have an open cover $X = \bigcup_{n=1}^{\infty} U_n$ define the increasing open cover

$$U_n' \coloneqq \bigcup_{i=1}^{n} U_i$$

by the contrapositive $U_n'$ is not proper for some $n$, which means precisely that there is a finite subcover. ■

**Proposition 4.1.104**
*Let $X$ be a topological space. Then sequentially quasi-compact $\implies$ countably quasi-compact. Under the assumption of first countability the converse holds.*

*Proof.* $\implies$) Consider a countable open cover $X = \bigcup_{n=1}^{\infty} U_n$ with no finite subcover. Then there exists $x_n \in X \setminus \bigcup_{i=1}^{n} U_i$. By assumption there is a convergent subsequence $x_{n_k} \to x$. We have $x \in U_N$ for some $N$ but $n_k \geq N \implies x_{n_k} \notin U_N$.

$\impliedby$) Consider a sequence $(x_n)$ and define

$$Z_n \coloneqq \overline{\{x_k : k > n\}}$$

© David Rufino CC BY-NC-ND 4.0

which is a decreasing sequence of non-empty closed subsets. By the countable compactness criteria (4.1.103) $Z := \bigcap_{n=1}^{\infty} Z_n$ is non-empty. However we may show this is precisely the set of cluster points of $(x_n)$, i.e. there exists a cluster point $x \in Z$. By (4.1.101) there exists a subsequence such that $x_{n_k} \to x$ as required. ■

**Lemma 4.1.105 (Lindelof Lemma)**
*Let $X$ be a second countable topological space. Then every open cover has a countable subcover.*

*Proof.* Let $\mathcal{B}$ be a countable base and $\{U_\alpha\}_{\alpha\in\mathcal{A}}$ an open cover of $X$. Note for every $U_\alpha$ we have some subfamily $\mathcal{B}_\alpha \subset \mathcal{B}$ such that $U_\alpha = \bigcup \mathcal{B}_\alpha$. Consider $\mathcal{B}' := \bigcup\{B_\alpha \mid \alpha \in \mathcal{A}\} \subset \mathcal{B}$. By definition for every $V \in \mathcal{B}'$ we have $V \subset U_\alpha$ for some $\alpha \in \mathcal{A}$. In other words there exists a map $f : \mathcal{B}' \to \mathcal{A}$ such that $V \subset U_{f(V)}$. We may verify that $\{U_\alpha\}_{\alpha\in \mathrm{Im}(f)}$ is a countable subcover. ■

We summarise the relationships between the concepts

**Proposition 4.1.106**
*Let $X$ be a topological space. Then*

*a) Quasi-Compact, Sequentially Quasi-Compact $\implies$ Countably Quasi-compact*
*b) First Countable and (Countably) Quasi-Compact $\implies$ Sequentially Quasi-Compact*
*c) Second Countable and Countably or Sequentially Quasi-Compact $\implies$ Quasi-Compact*

*Proof.*
a) The first implication is obvious and the second implication is (4.1.104)
b) This is (4.1.104)
c) This follows from (4.1.105) and *a*)

■

**Proposition 4.1.107 (Compact subsets of Hausdorff Spaces are Closed)**
*Let $X$ be a Hausdorff topological space and $Y \subset X$ a subset compact with respect to the subspace topology. Then $Y$ is closed.*

*Proof.* Fix $x \in X \setminus Y$. Then for every $y \in Y$ by assumption there exists disjoint open subsets $x \in U_y$ and $y \in V_y$. Then $\{V_y \cap Y \mid y \in Y\}$ is an open cover for $Y$, so there exists a finite set of points $y_1, ..., y_n$ such that $Y \subseteq V_{y_1} \cup ... \cup V_{y_n}$. Furthermore

$$U := U_{y_1} \cap ... \cap U_{y_n}$$

is an open neighbourhood of $x$ disjoint from each $V_{y_i}$ and therefore disjoint from $Y$. By (4.1.19) we see that $x \notin \overline{Y}$, which shows $Y = \overline{Y}$, i.e. $Y$ is closed. ■

**Proposition 4.1.108 (Closed subsets of Compact Spaces are Compact)**
*Let $X$ be a quasi-compact topological space and $Y \subset X$ a closed subset. Then $Y$ is quasi-compact under the subspace topology.*

*Proof.* Let $\mathcal{U} := \{U_\alpha\}_{\alpha\in\mathcal{A}}$ be an open cover for $Y$. Then by definition $U_\alpha = U'_\alpha \cap Y$, where $U'_\alpha \subset X$ are open. Evidently

$$\mathcal{U}' : \{U'_\alpha\}_{\alpha\in\mathcal{A}} \cup \{X \setminus Y\}$$

is an open cover of $X$. Therefore by assumption it has a finite subcover, which immediately yields a finite subcover of $Y$ as required. ■

**Proposition 4.1.109 (Image of a Compact Space is Compact)**
*Let $f : X \to Y$ be a continuous map of topological spaces where $X$ is quasi-compact. Then the image $f(X)$ is quasi-compact.*

*Further if $Y$ is Hausdorff then $f$ is an open map (i.e. $f(U)$ is open for every open $U \subset X$). If in addition $f$ is injective then it is a homeomorphism onto its image.*

*Proof.* We may assume without loss of generality that $f(X) = Y$. Let $\{U_\alpha\}_{\alpha\in\mathcal{A}}$ be an open cover of $Y$. In other words $f(X) \subseteq \bigcup_{\alpha\in\mathcal{A}} U_\alpha \implies X \subseteq \bigcup_{\alpha\in\mathcal{A}} f^{-1}(U_\alpha)$. By assumption we have a finite subcover

$$X \subseteq f^{-1}\left(U_{\alpha_1}\right) \cup ... \cup f^{-1}\left(U_{\alpha_n}\right)$$

which implies

$$f(X) \subseteq U_{\alpha_1} \cup ... \cup U_{\alpha_n}$$

© David Rufino CC BY-NC-ND 4.0

since $f$ is assumed to be surjective, and we see that $f(X)$ is compact. ∎

Analogously we have similar results for sequentially compact spaces, which may be conceptually simpler.

**Proposition 4.1.110**
*Let $X$ be a first countable Kolmogorov space and $Y \subset X$ a subset of a sequentially compact with respect to the subspace topology. Then $Y$ is closed.*

*Proof.* Consider $x \in \overline{Y}$ then by (4.1.19) there is some sequence $x_n \to x$. Further by assumption there is some subsequence $x_{n_k} \to y \in Y$. By uniqueness of limits (4.1.51) we have $x = y$, and in particular $\overline{Y} = Y$ is closed. ∎

**Proposition 4.1.111**
*Let $X$ be a sequentially compact topological space, and $Y \subset X$ a closed subset. Then $Y$ is sequentially compact.*

*Proof.* Let $x_n \in Y$ be a sequence. By assumption it has a convergent subsequence $x_{n_k} \to x$ where $x \in X$. By (4.1.18) $x$ is an adherent point of $Y$ and therefore lies in $Y$ by (4.1.19). ∎

**Proposition 4.1.112**
*Let $f : X \to Y$ be a continuous map of topological spaces where $X$ is sequentially compact. Then the image $f(X)$ is sequentially compact.*

*Proof.* Let $y_n = f(x_n)$ be a sequence in $f(X)$. Then by assumption $x_{n_k} \to x$, and therefore by (4.1.50) $y_{n_k} \to f(x)$. ∎

**Proposition 4.1.113**
*Let $X$ be a topological space with an open cover $X = \bigcup_{i=1}^{n} U_i$ such that $U_i$ is quasi-compact under the subspace topology. Then $X$ is quasi-compact.*

*Proof.* Let $X = \bigcup_{\alpha \in \mathcal{A}} V_\alpha$ then $U_i = \bigcup_{\alpha \in \mathcal{A}} (V_\alpha \cap U_i)$. By assumption there is some finite subset $\mathcal{A}_i \subset \mathcal{A}$ such that $U_i = \bigcup_{\alpha \in \mathcal{A}_i} (V_\alpha \cap U_i)$. Then

$$X = \bigcup_{\alpha \in \mathcal{A}_1 \cup \ldots \cup \mathcal{A}_n} V_\alpha$$

Therefore $X$ is quasi-compact. ∎

**Proposition 4.1.114 (Noetherian $\iff$ Hereditarily Quasi-Compact)**
*Let $X$ be a topological space. Then $X$ is Noetherian if and only if every open subset is quasi-compact.*

*Proof.* $\implies$) By (4.1.77) every open subset is Noetherian, so it is enough to show that $X$ is quasi-compact. By definition the lattice of open subsets is Noetherian and by (2.1.63) this holds iff every family of open subsets has a maximal element.

Suppose that we have an open cover $X = \bigcup_{\alpha \in \mathcal{A}} U_\alpha$. Let $\mathcal{F}$ be the family of open subsets expressible as a finite union of sets $U_\alpha$. By assumption it has a maximal element

$$U := U_{\alpha_1} \cup \ldots \cup U_{\alpha_n}$$

Suppose that $U \subsetneq X$ is proper then choose $x \in X \setminus U$. By hypothesis there is some $\alpha_{n+1}$ such that $x \in U_{\alpha_{n+1}}$. Then $U \subsetneq U \cup U_{\alpha_{n+1}} \in \mathcal{F}$, contradicting maximality. Therefore $U = X$ and $X$ is quasi-compact as required.

$\impliedby$) Omitted. ∎

**Proposition 4.1.115**
*Let $X, Y$ be quasi-compact topological spaces. Then $X \times Y$ is quasi-compact.*

*Proof.* Let $\{U_\lambda\}_{\lambda \in \Lambda}$ be an open cover of $X \times Y$. Therefore for each $(x, y)$ there is a $\lambda(x, y)$ such that $(x, y) \in U_{\lambda(x,y)}$. By definition of the product topology (4.1.95) we may find open sets $V_{(x,y)} \subset X$ and $W_{(x,y)} \subset Y$ such that $(x, y) \in V_{(x,y)} \times W_{(x,y)} \subset U_{\lambda(x,y)}$.

Fix $x$ then the family $\left\{W_{(x,y)}\right\}_{y \in Y}$ is an open cover of $Y$. So we may find a finite subset $J_x \subset Y$ such that $\left\{W_{(x,y)}\right\}_{y \in J_x}$ covers $Y$.

Consider the open sets

$$V'_x := \bigcap_{y \in J_x} V_{(x,y)}$$

which evidently cover $X$. Then there is a finite subset $I \subset X$ such that $\{V'_x\}_{x \in I}$ covers $X$.

© David Rufino CC BY-NC-ND 4.0

Then we claim that the family $\left\{U_{\lambda(x,y)}\right\}_{x\in I, y\in J_x}$ is a finite subcover of $X\times Y$. For given $(x',y')\in X\times Y$ there is an $x\in I$ such that $x'\in V_x'$, and $y\in J_x$ such that $y'\in W_{(x,y)}$. Evidently $x'\in V_{(x,y)}$, whence $(x',y')\in U_{\lambda(x,y)}$ as required.■

### 4.1.15 Connectedness

**Definition 4.1.116**
*We say a topological space $X$ is **disconnected** if $X=U\cup V$ for two non-empty, disjoint open subsets $U,V$. Otherwise we say $X$ is connected.*

**Proposition 4.1.117**
*Let $X$ be a topological space. The following are equivalent*

*a) $X$ is connected*
*b) The only subsets which are both open and closed are $X$ and $\varnothing$*

**Proposition 4.1.118**
*Let $f:X\to Y$ be a continuous map and $X$ connected. Then $f(X)$ is also connected.*

### 4.1.16 Order Topology

**Definition 4.1.119 (Order Topology)**
*Let $(X,\leq)$ be a totally ordered set. Then the **order topology** is the topology generated by the subbase of half-open intervals*

$$\{y\in X\mid y<x\}\text{ and }\{y\in X\mid y>x\}$$

*for $x\in X$.*

**Proposition 4.1.120 (Base for order topology)**
*Let $(X,\leq)$ be a totally ordered set. Then the sets of the form*

$$\{y\in X\mid y<x\},\{y\in X\mid y>x\}\text{ and }\{y\in X\mid x<y<z\}$$

*for $x,z\in X$ constitute a base for the order topology.*

© David Rufino CC BY-NC-ND 4.0

## 4.2 Sheaves

Sheaves are typically used to organise sets of functions which are defined by a local differentiability condition (e.g. holomorphic, smooth, etc.). However in some cases it may be convenient to take an abstract category-theoretic approach as the proofs become more conceptual and naturally lead to further generalisations where the topological space is replaced by an arbitrary category. However for the sheafification construction we take (the standard) concrete approach as this is simpler and nicely coincides with the notion of "étale space".

In general we assume that the value category $\mathcal{C}$ admits arbitrary products and equalisers (and therefore also limits, and in particular terminal objects), as well as filtered colimits / direct limits. In some cases we may also require it to be algebraic.

**Definition 4.2.1 (Abstract Presheaf)**
*Let $X$ be a topological space, $\mathcal{B}$ a base for the topology and $\mathcal{C}$ a category. A $\mathcal{B}$-**presheaf** on $X$ is a functor*

$$\mathcal{F} : (\mathcal{B}, \supseteq) \longrightarrow \mathcal{C}$$

*from the poset category of basic open subsets to $\mathcal{C}$ ordered by reverse inclusion for which*

$$\mathcal{F}(\varnothing) = \star \text{ is a terminal object in } \mathcal{C}$$

*This means for every pair of open subsets $U, V \in \mathcal{B}$ such that $V \subseteq U$ there exist morphisms $\rho_{UV} : \mathcal{F}(U) \to \mathcal{F}(V)$ which satisfy the following properties*

*a) $\rho_{UU} = 1_{\mathcal{F}(U)}$*
*b) $\rho_{VW} \circ \rho_{UV} = \rho_{UW}$ for any basic open subset $W \subset \mathcal{B}$*

*A morphism of presheaves is precisely a natural transformation. That is $\alpha : \mathcal{F} \to \mathcal{G}$ is a family of morphisms $\alpha_U : \mathcal{F}(U) \to \mathcal{G}(U)$ for which*

$$\rho_{UV} \circ \alpha_U = \alpha_V \circ \rho_{UV}$$

*This determines a category $\mathbf{PSh}(\mathcal{B}; \mathcal{C})$. We may omit the value category $\mathcal{C}$ if there are no additional hypotheses.*

**Definition 4.2.2 (Abstract Sheaf)**
*Let $X$ be a topological space, $\mathcal{B}$ a base for the topology closed under finite intersections and $\mathcal{F}$ a $\mathcal{B}$-presheaf.*

*For every family of non-empty basic open subsets $\{U_i\}_{i \in I}$ with $U = \bigcup_{i \in I} U_i$, consider the sequence of morphisms*

$$\mathcal{F}(U) \xrightarrow{\ e\ } \prod_{i \in I} \mathcal{F}(U_i) \underset{d_1}{\overset{d_0}{\rightrightarrows}} \prod_{(i,j) \in I \times I} \mathcal{F}(U_i \cap U_j)$$

*where $e = \left(\rho_{UU_i}\right)_{i \in I}$, $d_0 = \left(\rho_{U_i(U_i \cap U_j)} \circ \pi_i\right)_{(i,j) \in I \times I}$ and $d_1 = \left(\rho_{U_j(U_i \cap U_j)} \circ \pi_j\right)_{(i,j) \in I \times I}$.*

*We say that $\mathcal{F}$ is a $\mathcal{B}$-**sheaf** if $e$ is an equaliser for $(d_0, d_1)$ for all such families of open subsets.*

*This determines a full subcategory $\mathbf{Sh}(\mathcal{B}; \mathcal{C})$ of $\mathbf{PSh}(\mathcal{B}; \mathcal{C})$.*

*When $\mathcal{B}$ is just the collection of all open subsets of $X$ then we say $\mathcal{F}$ is a **sheaf**, and we denote the category by $\mathbf{Sh}(X; \mathcal{C})$.*

It may be convenient to write the sheaf condition as the existence of a limit.

**Lemma 4.2.3 (Sheaf Condition as a Limit)**
*Let $\mathcal{F}$ be a presheaf on $X$ and $U = \bigcup_{i \in I} U_i$ a cover; write $U_{ij} := U_i \cap U_j$ for $(i,j) \in I \times I$, and let $d_0, d_1 : \prod_{i \in I} \mathcal{F}(U_i) \to \prod_{(i,j) \in I \times I} \mathcal{F}(U_{ij})$ be as in (4.2.2).*

*Let $I^{(1)}$ be the small category with objects $I \sqcup (I \times I)$ and, for each $(i,j) \in I \times I$, morphisms $u_{ij} : i \to (i,j)$ and $v_{ij} : j \to (i,j)$ (together with identities, and no further morphisms), and let $D_{\mathcal{F}} : I^{(1)} \to \mathcal{C}$ be the diagram*

$$D_{\mathcal{F}}(i) := \mathcal{F}(U_i), \quad D_{\mathcal{F}}((i,j)) := \mathcal{F}(U_{ij}), \quad D_{\mathcal{F}}(u_{ij}) := \rho_{U_i U_{ij}}, \quad D_{\mathcal{F}}(v_{ij}) := \rho_{U_j U_{ij}}$$

*Then, as $\mathcal{C}$ has arbitrary products, the contravariant functors $\mathcal{C}^{op} \to \mathbf{Set}$*

$$X \mapsto \left\{ \xi : X \to \prod_{i \in I} \mathcal{F}(U_i) \;\middle|\; d_0 \circ \xi = d_1 \circ \xi \right\}, \quad X \mapsto \{\text{cones over } D_{\mathcal{F}} \text{ with apex } X\}$$

© David Rufino CC BY-NC-ND 4.0

*(both via precomposition) are naturally isomorphic, via* $\xi \mapsto (\pi_i \circ \xi)_{i\in I}$*, extended to* $(i,j) \in I \times I$ *by* $D_{\mathcal{F}}(u_{ij}) \circ \pi_i \circ \xi$*. Consequently* $D_{\mathcal{F}}$ *has* $\left(\mathcal{F}(U), \left\{\rho_{UU_i}\right\}_{i\in I}, \left\{\rho_{UU_{ij}}\right\}_{(i,j)\in I\times I}\right)$ *as a limit over* $I^{(1)}$ *if and only if* $\mathcal{F}$ *satisfies the sheaf condition for this cover (4.2.2).*

*Proof.* Given $\xi$ with $d_0 \circ \xi = d_1 \circ \xi$, set $\xi_i := \pi_i \circ \xi$ and $\xi_{(i,j)} := D_{\mathcal{F}}(u_{ij}) \circ \xi_i$; then also $\xi_{(i,j)} = D_{\mathcal{F}}(v_{ij}) \circ \xi_j$, since both sides equal $\pi_{(i,j)} \circ d_0 \circ \xi = \pi_{(i,j)} \circ d_1 \circ \xi$ by definition of $d_0, d_1$. So $\left(X, \{\xi_k\}_{k\in I^{(1)}}\right)$ is a cone over $D_{\mathcal{F}}$, the only nontrivial conditions being exactly this equality at $u_{ij}, v_{ij}$.

Conversely, given a cone $\left(X, \{\xi_k\}_{k\in I^{(1)}}\right)$, bundling $(\xi_i)_{i\in I}$ via the universal property of the product (2.6.81) gives $\xi := (\xi_i)_i : X \to \prod_{i\in I} \mathcal{F}(U_i)$ with $\pi_i \circ \xi = \xi_i$; then for all $(i,j)$, $\pi_{(i,j)} \circ d_0 \circ \xi = D_{\mathcal{F}}(u_{ij}) \circ \xi_i = \xi_{(i,j)} = D_{\mathcal{F}}(v_{ij}) \circ \xi_j = \pi_{(i,j)} \circ d_1 \circ \xi$, using the cone conditions at $u_{ij}, v_{ij}$, so $d_0 \circ \xi = d_1 \circ \xi$.

These two constructions are mutually inverse, directly from $\pi_i \circ \xi = \xi_i$ and the universal property of the product, and are evidently compatible with precomposition, i.e. natural in $X$: for $\phi : X' \to X$, both sides simply precompose $\xi$, resp. every $\xi_k$, by $\phi$. So $D_{\mathcal{F}}$ has a limit over $I^{(1)}$ iff $(d_0, d_1)$ has an equaliser, and the two agree. ∎

**Proposition 4.2.4 (Sheaf Conditions for Concrete Categories)**
*Let* $X$ *be a topological space,* $\mathcal{B}$ *a base for the topology closed under finite intersections and* $\mathcal{C}$ *is an algebraic category (2.6.184). Then a* $\mathcal{B}$*-presheaf* $\mathcal{F}$ *is a sheaf if and only if it satisfies the following conditions, for every open cover* $U = \bigcup_{i\in I} U_i$ *with* $U_i$ *non-empty*

*a)* $\sigma, \tau \in \mathcal{F}(U)$ *such that* $\rho_{UU_i}(\sigma) = \rho_{UU_i}(\tau)$ *for all* $i \in I$ *implies* $\sigma = \tau$
*b)* $\sigma_i \in \mathcal{F}(U_i)$ *such that* $\rho_{U_i(U_i\cap U_j)}(\sigma_i) = \rho_{U_j(U_i\cap U_j)}(\sigma_j)$ *for all* $(i,j) \in I \times I$ *implies there exists* $\sigma \in \mathcal{F}(U)$ *such that* $\rho_{UU_i}(\sigma) = \sigma_i$.

*Proof.* Suppose that $\mathcal{F}$ is a sheaf in the abstract sense. By assumption the forgetful functor $\mathcal{C} \to \mathbf{Set}$ preserves limits and so by (2.6.197) $e$ is injective and

$$\mathrm{Im}(e) = \left\{(\sigma_i)_{i\in I} \;\middle|\; \sigma_i|_{U_i\cap U_j} = \sigma_j|_{U_i\cap U_j}\right\}$$

which gives both concrete sheaf conditions.

Conversely this condition ensures the sequence is an equaliser in **Set**, and so by hypothesis is so in $\mathcal{C}$. ∎

## 4.2.1 Stalks

**Definition 4.2.5 (Stalk of a presheaf)**
*Let* $X$ *be a topological space,* $\mathcal{B}$ *a base closed under finite intersections and* $\mathcal{F}$ *a* $\mathcal{B}$*-presheaf.*

*Let* $Z$ *be an irreducible subset of* $X$*, then the family of open subsets* $U \in \mathcal{B}$ *meeting* $Z$ *is closed under finite intersections (4.1.60) and therefore forms a directed preorder (2.6.123) ordered by reverse inclusion.*

*Therefore we may define the* ***stalk*** *at* $Z$ *by the direct limit (2.6.125)*

$$\mathcal{F}_Z := \varinjlim_{U\in\mathcal{B},\, U\cap Z\neq\varnothing} \mathcal{F}(U)$$

*This means there is a family of morphisms* $\{\rho_{UZ} : \mathcal{F}(U) \to \mathcal{F}_Z\}$ *such that*

*a) all* $V \in \mathcal{B}$ *with* $V \subset U$ *intersecting* $Z$ *satisfy* $\rho_{VZ} \circ \rho_{UV} = \rho_{UZ}$,
*b) any collection of morphisms* $\alpha_U : \mathcal{F}(U) \to \mathcal{A}$ *satisfying* $\alpha_U = \alpha_V \circ \rho_{UV}$ *factors uniquely through a morphism* $\alpha_Z : \mathcal{F}_Z \to \mathcal{A}$

*When* $Z = \{x\}$ *we denote this simply by* $\mathcal{F}_x$.

**Proposition 4.2.6 (Explicit form of Stalk)**
*Consider the situation (4.2.5) where the value category* $\mathcal{C}$ *is algebraic (2.6.184). Then*

$$U(\mathcal{F}_Z) = (U \circ \mathcal{F})_Z$$

*and so the stalk* $\mathcal{F}_Z$ *may be constructed explicitly as follows*

$$\mathcal{F}_Z := \{(U,\sigma) \mid U \in \mathcal{B}, \sigma \in \mathcal{F}(U)\}/\sim$$
$$(U,\sigma) \sim (V,\tau) \iff \exists W \in \mathcal{B} \text{ s.t. } W \cap Z \neq \varnothing \text{ and } \sigma|_W = \tau|_W$$

*with morphisms*

© David Rufino CC BY-NC-ND 4.0

$$\begin{aligned}\rho_{UZ} : \mathcal{F}(U) &\to \mathcal{F}_Z \\ \sigma &\to [(U, \sigma)]\end{aligned}$$

*In particular every element of the stalk $\sigma_Z \in \mathcal{F}_Z$ is the image of some section $\sigma \in \mathcal{F}(U)$ under $\rho_{UZ}$ for some $U \in \mathcal{B}$.*

*Proof.* By (2.6.189), the forgetful functor $U : \mathcal{C} \to \mathbf{Set}$ creates directed colimits, and so the result follows from (2.6.128). ∎

**Proposition 4.2.7 (Functoriality of Stalks)**
*Let $X$ be a topological space with base $\mathcal{B}$ closed under finite intersections, $\phi : \mathcal{F} \to \mathcal{G}$ a morphism of $\mathcal{B}$-presheaves and $Z \subset X$ an irreducible subset. Then there exists a unique morphism $\phi_Z : \mathcal{F}_Z \to \mathcal{G}_Z$ ensuring the following diagram commute for all open subsets $U$ meeting $Z$*

$$\begin{array}{ccc} \mathcal{F}(U) & \xrightarrow{\phi_U} & \mathcal{G}(U) \\ {\scriptstyle \rho_{UZ}}\downarrow & & \downarrow{\scriptstyle \rho_{UZ}} \\ \mathcal{F}_Z & \xrightarrow{\phi_Z} & \mathcal{G}_Z \end{array}$$

*Furthermore $(\psi \circ \phi)_Z = \psi_Z \circ \phi_Z$. In the explicit case (4.2.6) then $\phi_Z$ is given by*

$$[(U, \sigma)] \to [(U, \phi_U(\sigma))]$$

*This also means that for any $A \in \mathcal{C}$ the bijection*

$$\mathrm{Mor}(\mathcal{F}_Z, A) \longrightarrow \left(\mathcal{F}|_{\mathcal{N}_Z} \downarrow A\right)$$

*is natural in $\mathcal{F}$, where $\mathcal{N}_Z$ is the category of open sets intersecting $Z$ ordered by reverse inclusion.*

**Lemma 4.2.8 (Lifting Stalks)**
*Suppose we are in the situation of (4.2.6) and $\sigma \in \mathcal{F}(U)$, $\tau \in \mathcal{F}(V)$ are sections such that $Z \cap U \cap V \neq \varnothing$.*

*a) Then $\sigma_Z = \tau_Z$ if and only if there is a neighbourhood $W \in \mathcal{B}$, $W \subseteq U \cap V$, meeting $Z$ such that $\sigma|_W = \tau|_W$.*
*b) If $\sigma_x = \tau_x$ for all $x \in U \cap V$, then there is an open cover $U \cap V = \bigcup_i U_i$ such that $\sigma|_{U_i} = \tau|_{U_i}$*
*c) If in addition $\mathcal{F}$ is a sheaf then $\sigma|_{U\cap V} = \tau|_{U\cap V}$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

By (4.2.6), $\mathcal{F}_Z = \{(W, \rho) \mid W \in \mathcal{B}, \rho \in \mathcal{F}(W)\}/\sim$ with $(W, \rho) \sim (W', \rho')$ iff there is $N \in \mathcal{B}$, $N \subseteq W \cap W'$, $N \cap Z \neq \varnothing$, with $\rho|_N = \rho'|_N$; and $\sigma_Z = [(U, \sigma)]$, $\tau_Z = [(V, \tau)]$. So $\sigma_Z = \tau_Z$ if and only if $(U, \sigma) \sim (V, \tau)$, which is exactly the stated condition, proving a).

For b), suppose $\sigma_x = \tau_x$ for all $x \in U \cap V$. For each such $x$, a) (applied with $Z := \{x\}$, an irreducible subset meeting $U \cap V$) gives $W_x \in \mathcal{B}$, $W_x \subseteq U \cap V$, $x \in W_x$, with $\sigma|_{W_x} = \tau|_{W_x}$. Then $\{W_x\}_{x\in U\cap V}$ is an open cover of $U \cap V$ with the required property.

For c), as $\mathcal{B}$ is closed under finite intersections and $U, V \in \mathcal{B}$, $U \cap V \in \mathcal{B}$, so $\sigma|_{U\cap V}, \tau|_{U\cap V} \in \mathcal{F}(U \cap V)$ are defined. By b) there is an open cover $U \cap V = \bigcup_i W_i$ with $\sigma|_{W_i} = \tau|_{W_i}$, that is $\rho_{(U\cap V)W_i}(\sigma|_{U\cap V}) = \rho_{(U\cap V)W_i}(\tau|_{U\cap V})$ for all $i$; as $\mathcal{F}$ is a sheaf, the separation condition (4.2.4) gives $\sigma|_{U\cap V} = \tau|_{U\cap V}$. ∎

**Proposition 4.2.9 (Stalks when restricting to a base)**
*Let $\mathcal{B}_2 \subset \mathcal{B}_1$ be bases for $X$ which generate the same topology. Suppose $\mathcal{F}$ is a $\mathcal{B}_1$-presheaf and $Z \subset X$ is an irreducible subset. Then there is a unique morphism making the following diagram commute for all $U \in \mathcal{B}_2$ intersecting $Z$*

$$\begin{array}{ccc} \mathcal{F}|_{\mathcal{B}_2}(U) & = & \mathcal{F}(U) \\ {\scriptstyle \rho_{UZ}}\downarrow & & \downarrow{\scriptstyle \rho_{UZ}} \\ \left(\mathcal{F}|_{\mathcal{B}_2}\right)_Z & \xrightarrow{\sim} & \mathcal{F}_Z \end{array}$$

*and it is an isomorphism.*

*In the situation of (4.2.6) the map is of the obvious form.*

*Proof.* Let $\mathcal{I}_2$ and $\mathcal{I}_1$ be the diagram of open subsets $U$ of $\mathcal{B}_2$ and $\mathcal{B}_1$ meeting $Z$, ordered by reverse inclusion. Then $\mathcal{I}_2$ is a full subcategory of $\mathcal{I}_1$ which is cofinal (2.6.126). So the result follows from (2.6.127)

© David Rufino CC BY-NC-ND 4.0

In the explicit case this may be shown more directly. ∎

**Lemma 4.2.10 (Pointwise Injectivity and Injectivity on Stalks)**
*Let $\mathcal{C}$ be algebraic (2.6.184) and $\phi : \mathcal{F} \to \mathcal{G}$ a morphism of presheaves such that $\phi_U$ is injective for all $U \subset X$. Then $\phi_x$ is injective for all $x \in X$.*

*Conversely, if $\phi$ is a morphism of sheaves and $\phi_x$ is injective for all $x \in X$, then $\phi_U$ is injective for all $U \subset X$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Fix $x \in X$ and suppose $a, b \in \mathcal{F}_x$ with $\phi_x(a) = \phi_x(b)$. By (4.2.6), $a = (\sigma_1)_x$, $b = (\sigma_2)_x$ for some $\sigma_1 \in \mathcal{F}(V_1)$, $\sigma_2 \in \mathcal{F}(V_2)$, $V_1, V_2 \in \mathcal{B}$ containing $x$; setting $V := V_1 \cap V_2 \in \mathcal{B}$, $\sigma := \sigma_1|_V$, $\tau := \sigma_2|_V$, we have $\sigma, \tau \in \mathcal{F}(V)$ with $\sigma_x = a$, $\tau_x = b$. Then

$$(\phi_V(\sigma))_x = \phi_x(\sigma_x) = \phi_x(a) = \phi_x(b) = \phi_x(\tau_x) = (\phi_V(\tau))_x$$

so by (4.2.8).a) (with both opens taken to be $V$) there is $W \in \mathcal{B}$, $W \subseteq V$, $x \in W$, with $\phi_V(\sigma)|_W = \phi_V(\tau)|_W$, that is $\phi_W(\sigma|_W) = \phi_W(\tau|_W)$. As $\phi_W$ is injective, $\sigma|_W = \tau|_W$, so $a = \sigma_x = (\sigma|_W)_x = (\tau|_W)_x = \tau_x = b$.

Conversely, fix $U \subset X$ and $\sigma, \tau \in \mathcal{F}(U)$ with $\phi_U(\sigma) = \phi_U(\tau)$. For every $x \in U$, $\phi_x(\sigma_x) = (\phi_U(\sigma))_x = (\phi_U(\tau))_x = \phi_x(\tau_x)$, so $\sigma_x = \tau_x$ as $\phi_x$ is injective. As $\mathcal{F}$ is a sheaf, (4.2.8).c) (taking both opens of the lemma to be $U$) gives $\sigma = \tau$. ∎

**Proposition 4.2.11 (Isomorphisms of Presheaves and Sheaves)**
*Let $\phi : \mathcal{F} \to \mathcal{G}$ be a morphism of presheaves (resp. sheaves). Then the following are equivalent*

*a) $\phi$ is an isomorphism*
*b) $\phi_U$ is an isomorphism for all $U \subset X$*

*If $\mathcal{C}$ is algebraic (2.6.184), then a), b) are further equivalent to*

*c) $\phi_U$ is bijective for all $U \subset X$*

*If moreover $\mathcal{F}, \mathcal{G}$ are sheaves, a)–c) are equivalent to*

*d) $\phi_x : \mathcal{F}_x \to \mathcal{G}_x$ is an isomorphism for all $x \in X$*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

$b) \Longrightarrow a)$: If each $\phi_U$ is an isomorphism, define $\psi_U := \phi_U^{-1}$; for $V \subseteq U$, inverting both sides of the naturality square $\phi_V \circ \rho_{UV} = \rho_{UV} \circ \phi_U$ gives $\rho_{UV} \circ \psi_U = \psi_V \circ \rho_{UV}$, so $\psi := \{\psi_U\}$ is a morphism of presheaves (resp. sheaves), and $\psi\phi = 1_{\mathcal{F}}$, $\phi\psi = 1_{\mathcal{G}}$ hold pointwise, hence as morphisms of presheaves. So $\phi$ is an isomorphism.

$a) \Longrightarrow b)$: If $\phi$ is an isomorphism with inverse $\psi$, then $\phi_U\psi_U = 1_{\mathcal{G}(U)}$ and $\psi_U\phi_U = 1_{\mathcal{F}(U)}$ for every $U$, so $\phi_U$ is an isomorphism.

$b) \Longleftrightarrow c)$ (when $\mathcal{C}$ is algebraic): an isomorphism is in particular bijective (a functor preserves isomorphisms, and an isomorphism in **Set** is a bijection), while conversely a bijective morphism is an isomorphism in $\mathcal{C}$ as $U$ reflects isomorphisms (2.6.189) apply this pointwise at each $U$.

$b) \Longrightarrow d)$: with $\psi := \{\psi_U\}$ as in $b) \Longrightarrow a)$ above, $\psi$ is a morphism of sheaves with $\psi\phi = 1_{\mathcal{F}}$, $\phi\psi = 1_{\mathcal{G}}$. By functoriality of stalks (4.2.7), $\psi_x \circ \phi_x = (\psi\phi)_x = 1_{\mathcal{F}_x}$ and $\phi_x \circ \psi_x = (\phi\psi)_x = 1_{\mathcal{G}_x}$, so $\phi_x$ is an isomorphism.

$d) \Longrightarrow c)$: Suppose $\phi_x$ is an isomorphism, in particular bijective, for every $x \in X$. By (4.2.10), $\phi_U$ is injective for every $U \subset X$. For surjectivity: fix $U \subset X$ and $\tau \in \mathcal{G}(U)$. For each $x \in U$, surjectivity of $\phi_x$ gives $a_x \in \mathcal{F}_x$ with $\phi_x(a_x) = \tau_x$; write $a_x = (\rho_x)_x$ for some $\rho_x \in \mathcal{F}(W_x)$, $W_x \in \mathcal{B}$, $x \in W_x \subseteq U$ (shrinking $W_x$ if necessary). Then $\big(\phi_{W_x}(\rho_x)\big)_x = \phi_x\big((\rho_x)_x\big) = \phi_x(a_x) = \tau_x = \big(\tau|_{W_x}\big)_x$, so by (4.2.8).a) there is $W_{x'} \in \mathcal{B}$, $x \in W_{x'} \subseteq W_x$, with $\phi_{W_x}(\rho_x)|_{W_{x'}} = \tau|_{W_{x'}}$; write $\sigma_x := \rho_x|_{W_{x'}} \in \mathcal{F}(W_{x'})$, so that $\phi_{W_{x'}}(\sigma_x) = \tau|_{W_{x'}}$ exactly.

The $\{W_{x'}\}_{x \in U}$ cover $U$; for $x, y \in U$, on $W_{x'} \cap W_{y'} \in \mathcal{B}$,

$$\phi_{W_{x'} \cap W_{y'}}\Big(\sigma_x|_{W_{x'} \cap W_{y'}}\Big) = \phi_{W_{x'}}(\sigma_x)|_{W_{x'} \cap W_{y'}} = \tau|_{W_{x'} \cap W_{y'}} = \phi_{W_{y'}}(\sigma_y)|_{W_{x'} \cap W_{y'}} = \phi_{W_{x'} \cap W_{y'}}\Big(\sigma_y|_{W_{x'} \cap W_{y'}}\Big)$$

so as $\phi_{W_{x'} \cap W_{y'}}$ is injective, $\sigma_x|_{W_{x'} \cap W_{y'}} = \sigma_y|_{W_{x'} \cap W_{y'}}$. As $\mathcal{F}$ is a sheaf, the gluing condition (4.2.4) gives $\sigma \in \mathcal{F}(U)$ with $\sigma|_{W_{x'}} = \sigma_x$ for all $x$. Then $\phi_U(\sigma)|_{W_{x'}} = \phi_{W_{x'}}\big(\sigma|_{W_{x'}}\big) = \phi_{W_{x'}}(\sigma_x) = \tau|_{W_{x'}}$ for every $x \in U$, so as $\mathcal{G}$ is a sheaf, the separation condition (4.2.4) gives $\phi_U(\sigma) = \tau$. So $\phi_U$ is surjective; being also injective, $\phi_U$ is bijective, that is c). ∎

© David Rufino CC BY-NC-ND 4.0

The following somewhat tedious result shows that morphisms are uniquely determined by morphisms on basic open sets.

**Proposition 4.2.12**
*Let $X$ be a topological space, $\mathcal{B}$ a base for the topology closed under finite intersections. Then the functor*

$$(-)|_{\mathcal{B}} : \mathbf{Sh}(X;\mathcal{C}) \to \mathbf{Sh}(\mathcal{B};\mathcal{C})$$

*is full and faithful.*

*Proof.* Let $\phi, \psi : \mathcal{F} \to \mathcal{G}$ be morphisms which agree on basic open subsets $U \in \mathcal{B}$. Consider an arbitrary open subset $U \subset X$ then by definition $U = \bigcup_{i\in I} U_i$ for $U_i \in \mathcal{B}$. By assumption there is a commutative diagram

$$\begin{array}{ccc} \mathcal{F}(U) & \xrightarrow{e} & \prod_{i\in I}\mathcal{F}(U_i) \\ {\scriptstyle\psi_U}\downarrow\downarrow{\scriptstyle\phi_U} & & \downarrow{\scriptstyle\prod_{i\in I}\phi|_{U_i} = \prod_{i\in I}\psi|_{U_i}} \\ \mathcal{G}(U) & \xrightarrow{e} & \prod_{i\in I}\mathcal{G}(U_i) \end{array}$$

which shows $e \circ \phi_U = e \circ \psi_U$. But $e$ is a monomorphism (2.6.135) (being part of an equaliser) so $\phi_U = \psi_U$. Therefore the functor is faithful.

Let $\phi : \mathcal{F}|_{\mathcal{B}} \to \mathcal{G}|_{\mathcal{B}}$ be a morphism of $\mathcal{B}$-sheaves. For an open set $U$ choose a basic open cover $U = \bigcup_{i\in I} U_i$ then we have a commutative diagram

$$\begin{array}{ccccc} \mathcal{F}(U) & \xrightarrow{e} & \prod_{i\in I}\mathcal{F}(U_i) & \rightrightarrows & \prod_{(i,j)\in I\times I}\mathcal{F}(U_i\cap U_j) \\ \downarrow{\scriptstyle\tilde{\phi}_U} & & \downarrow{\scriptstyle\prod_{i\in I}\phi_{U_i}} & & \downarrow{\scriptstyle\prod_{(i,j)\in I\times I}\phi_{U_i\cap U_j}} \\ \mathcal{G}(U) & \xrightarrow{e} & \prod_{i\in I}\mathcal{G}(U_i) & \rightrightarrows & \prod_{(i,j)\in I\times I}\mathcal{G}(U_i\cap U_j) \end{array}$$

where for the right hand square we must pick the top arrows or bottom arrows. Chasing the diagram shows that $\left(\prod_{i\in I}\phi_{U_i}\right)\circ e$ is equalised by the bottom two arrows, and so the factorisation exists by the universal property and $\tilde{\phi}_U$ the unique morphism which makes the left square commute, i.e. such that $\phi_{U_i}\circ\rho_{UU_i} = \rho_{UU_i}\circ\tilde{\phi}_U$ for all $i \in I$.

We claim that for any basic open subset $W \subset U$ the following diagram commutes

$$\begin{array}{ccccc} \mathcal{F}(U) & \longrightarrow & \mathcal{F}(W) & \xrightarrow{e} & \prod_{i\in I}\mathcal{F}(W\cap U_i) \\ \downarrow{\scriptstyle\tilde{\phi}_U} & & \downarrow{\scriptstyle\phi_W} & & \downarrow{\scriptstyle\prod_{i\in I}\phi_{W\cap U_i}} \\ \mathcal{G}(U) & \longrightarrow & \mathcal{G}(W) & \xrightarrow{e} & \prod_{i\in I}\mathcal{G}(W\cap U_i) \end{array}$$

and $\tilde{\phi}_U$ is the unique morphism such that this commutes for all $W \in \mathcal{B}$. The right hand square commutes by assumption. As $e$ is a monomorphism it is enough to show that

$$\left(\prod_{i\in I}\phi_{W\cap U_i}\right)\circ e\circ\rho_{UW} = e\circ\phi_W\circ\rho_{UW} = e\circ\rho_{UW}\circ\tilde{\phi}_U$$

First observe we have a commutative diagram

$$\begin{array}{ccc} \mathcal{F}(U) & \xrightarrow{e} & \prod_{i\in I}\mathcal{F}(U_i) \\ \downarrow{\scriptstyle\rho_{UW}} & & \downarrow{\scriptstyle\prod_{i\in I}\rho_{U_i(W\cap U_i)}} \\ \mathcal{F}(W) & \xrightarrow{e} & \prod_{i\in I}\mathcal{F}(W\cap U_i) \end{array}$$

which follows because both directions are equal to the diagonal $\left(\rho_{U(W\cap U_i)}\right)_{i\in I}$. Therefore we may combine the two diagrams to find

© David Rufino CC BY-NC-ND 4.0

$$\begin{aligned}
e \circ \phi_W \circ \rho_{UW} &= \left(\prod_{i\in I} \phi_{W\cap U_i}\right) \circ e \circ \rho_{UW} \\
&= \left(\prod_{i\in I} \phi_{W\cap U_i}\right) \circ \left(\prod_{i\in I} \rho_{U_i(W\cap U_i)}\right) \circ e \\
&= \left(\prod_{i\in I} \phi_{W\cap U_i} \circ \rho_{U_i(W\cap U_i)}\right) \circ e \\
&= \left(\prod_{i\in I} \rho_{U_i(W\cap U_i)} \circ \phi_{U_i}\right) \circ e \\
&= \left(\prod_{i\in I} \rho_{U_i(W\cap U_i)}\right) \circ \left(\prod_{i\in I} \phi_{U_i}\right) \circ e \\
&= \left(\prod_{i\in I} \rho_{U_i(W\cap U_i)}\right) \circ e \circ \tilde{\phi}_U \\
&= e \circ \rho_{UW} \circ \tilde{\phi}_U
\end{aligned}$$

As $e$ is a monomorphism (2.6.135) then $\phi_W \circ \rho_{UW} = \rho_{UW} \circ \tilde{\phi}_U$ as required. Then $\tilde{\phi}_U$ is the unique morphism such that this holds for all $W \in \mathcal{B}$ and $W \subset U$, because it is already uniquely determined by $W = U_i$. This also shows that $\tilde{\phi}_W = \phi_W$.

In particular $\tilde{\phi}_U$ does not depend on the choice of cover $\{U_i\}$ of $U$: if $\tilde{\phi}'_U$ is constructed from a different basic cover of $U$, it satisfies the same property $\phi_W \circ \rho_{UW} = \rho_{UW} \circ \tilde{\phi}'_U$ for all $W \in \mathcal{B}$, $W \subset U$; taking $W$ to range over the original cover $\{U_i\}$ shows $\tilde{\phi}'_U$ also satisfies the equalising property defining $\tilde{\phi}_U$, so $\tilde{\phi}'_U = \tilde{\phi}_U$ by the uniqueness established above.

Finally for an arbitrary open subset $V = \bigcup_{i\in I} W_i \subset U$, $W_i \in \mathcal{B}$, we wish to show we have a commutative diagram

$$\begin{array}{ccccc}
\mathcal{F}(U) & \longrightarrow & \mathcal{F}(V) & \xrightarrow{e} & \prod_{i\in I} \mathcal{F}(W_i) \\
\downarrow{\scriptstyle \tilde{\phi}_U} & & \downarrow{\scriptstyle \tilde{\phi}_V} & & \downarrow{\scriptstyle \prod_{i\in I} \phi_{W_i}} \\
\mathcal{G}(U) & \longrightarrow & \mathcal{G}(V) & \xrightarrow{e} & \prod_{i\in I} \mathcal{G}(W_i)
\end{array}$$

We've already shown that the large square and right square commute. This shows that

$$e \circ \tilde{\phi}_V \circ \rho_{UV} = e \circ \rho_{UV} \circ \tilde{\phi}_U$$

and as $e$ is a monomorphism (2.6.135) the left square also commutes.

Therefore $\tilde{\phi}$ is a morphism of sheaves and $\tilde{\phi}|_{\mathcal{B}} = \phi$. Therefore the functor is full.

Alternatively, when $\mathcal{C}$ is algebraic (2.6.184), this can be shown more directly by working with elements via the forgetful functor $U : \mathcal{C} \to \mathbf{Set}$, citing (4.2.4) throughout with the base taken to be all of $\mathcal{O}(X)$ (a valid choice of base, since $\mathcal{F}, \mathcal{G}$ are sheaves on all of $X$ (4.2.2)).

*Drafted by Claude Sonnet, reviewed by David Rufino.*

**Faithful.** Let $\phi, \psi : \mathcal{F} \to \mathcal{G}$ agree on $U \in \mathcal{B}$, and fix an arbitrary open $V \subset X$ and $\sigma \in \mathcal{F}(V)$. Choose, for each $x \in V$, a basic open $U_x \in \mathcal{B}$ with $x \in U_x \subseteq V$. Then

$$\rho_{VU_x}(\phi_V(\sigma)) = \phi_{U_x}\left(\rho_{VU_x}(\sigma)\right) = \psi_{U_x}\left(\rho_{VU_x}(\sigma)\right) = \rho_{VU_x}(\psi_V(\sigma))$$

using naturality of $\phi, \psi$ and that they agree on $U_x \in \mathcal{B}$. As $\{U_x\}_{x\in V}$ is a cover of $V$, separation for $\mathcal{G}$ gives $\phi_V(\sigma) = \psi_V(\sigma)$. As $\sigma, V$ were arbitrary, $\phi = \psi$.

**Full.** Let $\phi : \mathcal{F}|_{\mathcal{B}} \to \mathcal{G}|_{\mathcal{B}}$ be a morphism of $\mathcal{B}$-sheaves; by (4.2.7) this determines stalk maps $\phi_x : \mathcal{F}_x \to \mathcal{G}_x$ for every $x \in X$.

*Construction.* Fix an open $V \subset X$ and $\sigma \in \mathcal{F}(V)$. Choose, for each $x \in V$, a basic open $U_x \in \mathcal{B}$ with $x \in U_x \subseteq V$, and set $\tau_x := \phi_{U_x}\left(\sigma|_{U_x}\right) \in \mathcal{G}(U_x)$. For $x, y \in V$, on $U_x \cap U_y \in \mathcal{B}$,

$$\tau_x|_{U_x\cap U_y} = \phi_{U_x\cap U_y}\left(\sigma|_{U_x\cap U_y}\right) = \tau_y|_{U_x\cap U_y}$$

© David Rufino CC BY-NC-ND 4.0

using naturality of $\phi$ and that $\big(\sigma|_{U_x}\big)|_{U_x\cap U_y} = \sigma|_{U_x\cap U_y} = \big(\sigma|_{U_y}\big)|_{U_x\cap U_y}$. So by the gluing condition for $\mathcal{G}$, applied to the cover $\{U_x\}_{x\in V}$ of $V$, there is $\tau\in\mathcal{G}(V)$ with $\tau|_{U_x}=\tau_x$ for all $x$; define $\tilde{\phi}_V(\sigma) \coloneqq \tau$. By construction and functoriality of stalks (4.2.7), $\tilde{\phi}_V(\sigma)$ satisfies the property

$$(\star)\qquad \big(\tilde{\phi}_V(\sigma)\big)_x = \big(\tau|_{U_x}\big)_x = (\tau_x)_x = \phi_x\Big(\big(\sigma|_{U_x}\big)_x\Big) = \phi_x(\sigma_x) \qquad \forall x\in V$$

*Well-definedness.* Suppose $\tau,\tau'\in\mathcal{G}(V)$ both satisfy $(\star)$ for the same $\sigma$, that is $\tau_x=\phi_x(\sigma_x)=\tau'_x$ for all $x\in V$. For each $x\in V$, choose $W\in\mathcal{B}$ with $x\in W\subseteq V$; then $(\tau|_W)_x=\tau_x=\tau'_x=(\tau'|_W)_x$, so by (4.2.8).a), applied to the sections $\tau|_W,\tau'|_W\in\mathcal{G}(W)$, there is $W_x\in\mathcal{B}$, $x\in W_x\subseteq W$, with $\tau|_{W_x}=\tau'|_{W_x}$. As $\{W_x\}_{x\in V}$ covers $V$, separation for $\mathcal{G}$ gives $\tau=\tau'$. So $\tilde{\phi}_V(\sigma)$ does not depend on the choice of cover $\{U_x\}$, and is the unique element of $\mathcal{G}(V)$ satisfying $(\star)$.

*$\tilde{\phi}$ restricts to $\phi$ on $\mathcal{B}$.* For $U\in\mathcal{B}$ and $\sigma\in\mathcal{F}(U)$, $\phi_U(\sigma)$ itself satisfies $(\phi_U(\sigma))_x=\phi_x(\sigma_x)$ for $x\in U$ (4.2.7), so by uniqueness $\tilde{\phi}_U(\sigma)=\phi_U(\sigma)$.

*$\tilde{\phi}$ is a morphism of presheaves.* For $V\subseteq U$ and $\sigma\in\mathcal{F}(U)$, both $\tilde{\phi}_V(\sigma|_V)$ and $\tilde{\phi}_U(\sigma)|_V$ satisfy $(\star)$ for $\sigma|_V$: for $x\in V$, $\big(\tilde{\phi}_V(\sigma|_V)\big)_x=\phi_x\big((\sigma|_V)_x\big)=\phi_x(\sigma_x)=\big(\tilde{\phi}_U(\sigma)\big)_x=\big(\tilde{\phi}_U(\sigma)|_V\big)_x$. By uniqueness, $\tilde{\phi}_V(\sigma|_V)=\tilde{\phi}_U(\sigma)|_V$, so $\tilde{\phi}$ is a morphism of sheaves with $\tilde{\phi}|_{\mathcal{B}}=\phi$. Therefore the functor is full. ∎

## 4.2.2 Sheafification

**Definition 4.2.13 (Sheafification)**
*Let $X$ be a topological space, $\mathcal{B}$ a basis of open sets for $X$ closed under finite intersections and $\mathcal{F}$ a $\mathcal{B}$-presheaf (with values in a category $\mathcal{C}$). Then we say that a pair $(\eta_{\mathcal{F}},\mathcal{F}^+)$ is a **sheafification** of $\mathcal{F}$ if*

*a) $\mathcal{F}^+$ is a sheaf (with values in a category $\mathcal{C}$)*
*b) $\eta_{\mathcal{F}}:\mathcal{F}\to\mathcal{F}^+|_{\mathcal{B}}$ is a morphism of $\mathcal{B}$-presheaves*
*c) for every sheaf $\mathcal{G}$ there is a bijection*

$$\begin{aligned}(-)^\sharp : \mathrm{Mor}(\mathcal{F}^+,\mathcal{G}) &\xrightarrow{\sim} \mathrm{Mor}(\mathcal{F},\mathcal{G}|_{\mathcal{B}})\\ \phi &\to \ \ \phi|_{\mathcal{B}}\circ\eta_{\mathcal{F}}\end{aligned}$$

*which is natural in $\mathcal{G}$, with the inverse map denoted by $(-)^\flat$.*

*In other words $\big(\mathcal{F}^+,(-)^\sharp\big)$ represents the functor $\mathcal{G}\to\mathrm{Mor}(\mathcal{F},\mathcal{G}|_{\mathcal{B}})$.*

*In this case we denote by $\varepsilon_{\mathcal{G}}:(\mathcal{G}|_{\mathcal{B}})^+\to\mathcal{G}$ the unique morphism which maps to $1_{\mathcal{G}|_{\mathcal{B}}}$ under the bijection.*

**Proposition 4.2.14 (Sheafification is Functorial)**
*Using the notation of (4.2.13) suppose there exists a sheafification $(\eta_{\mathcal{F}},\mathcal{F}^+)$ for all pre-sheaves $\mathcal{F}$. Then this can be extended to a functor*

$$(-)^+ : \mathbf{PSh}(X;\mathcal{B})\to\mathbf{Sh}(X)$$

*uniquely determined by either the condition $(-)^+$ is left-adjoint to $(-)|_{\mathcal{B}}$ or that the following diagram commutes*

$$\begin{array}{ccc}\mathcal{F}_1 & \xrightarrow{\ \phi\ } & \mathcal{F}_2\\ {\scriptstyle\eta_{\mathcal{F}_1}}\downarrow & & \downarrow{\scriptstyle\eta_{\mathcal{F}_2}}\\ \mathcal{F}_1^+|_{\mathcal{B}} & \xrightarrow[\phi^+|_{\mathcal{B}}]{\ \ \ \ \ } & \mathcal{F}_2^+|_{\mathcal{B}}\end{array}$$

*for all morphisms $\phi:\mathcal{F}_1\to\mathcal{F}_2$. More precisely we have the relationship*

$$\phi^+ = \big(\eta_{\mathcal{F}_2}\circ\phi\big)^\flat$$

*In particular $(\psi\circ\phi)^+=\psi^+\circ\phi^+$ for all morphisms $\psi:\mathcal{F}_2\to\mathcal{F}_3$ and $\eta_{\mathcal{F}}$ is the unit of the adjunction and $\varepsilon_{\mathcal{G}}$ the counit.*

*Furthermore $(-)|_{\mathcal{B}}:\mathbf{Sh}(X)\to\mathbf{PSh}(X;\mathcal{B})$ is fully faithful, and $\varepsilon$ is a natural isomorphism.*

*Proof.* See (2.6.74).

By (4.2.12), $\mathbf{Sh}(X)\to\mathbf{Sh}(\mathcal{B};\mathcal{C})$ is fully faithful; composing with the (trivially fully faithful) inclusion $\mathbf{Sh}(\mathcal{B};\mathcal{C})\hookrightarrow\mathbf{PSh}(X;\mathcal{B})$ shows $(-)|_{\mathcal{B}}:\mathbf{Sh}(X)\to\mathbf{PSh}(X;\mathcal{B})$ is fully faithful, so by (2.6.72), $\varepsilon$ is a natural isomorphism. ∎

© David Rufino CC BY-NC-ND 4.0

**Proposition 4.2.15**
*Using the notation of (4.2.13) suppose there exists a sheafification $(\eta_{\mathcal{F}}, \mathcal{F}^+)$ for $\mathcal{F}$. Then for every $x \in X$ there is a unique morphism making the following diagram commute for all $U \in \mathcal{B}$ containing $x$*

$$\begin{array}{ccc} \mathcal{F}(U) & \xrightarrow{\eta_U} & \mathcal{F}^+(U) \\ \downarrow & & \downarrow \\ \mathcal{F}_x & \dashrightarrow & (\mathcal{F}^+)_x \end{array}$$

*Proof.* By (4.2.7) and (4.2.9) there is a commutative diagram for all $U \in \mathcal{B}$ and $x \in U$

$$\begin{array}{ccccc} \mathcal{F}(U) & \xrightarrow{\eta_U} & \mathcal{F}^+(U) & = & \mathcal{F}^+(U) \\ \downarrow & & \downarrow & & \downarrow \\ \mathcal{F}_x & \longrightarrow & (\mathcal{F}^+|_{\mathcal{B}})_x & \xrightarrow{\sim} & \mathcal{F}^+_x \end{array}$$

where the bottom left arrow is uniquely determined. This shows that the required map exists and is uniquely determined according to this condition. ∎

**Proposition 4.2.16 (Sheafification)**
*Let $\mathcal{C}$ be an algebraic category (2.6.184) then the sheafification (4.2.13) exists.*

*More precisely define*

$$\mathcal{F}^+(U) := \left\{ s : U \to \coprod_{x \in U} \mathcal{F}_x \;\middle|\; U = \bigcup_{i \in I} U_i, U_i \in \mathcal{B}, \sigma_i \in \mathcal{F}(U_i), s(x) = (\sigma_i)_x \quad \forall x \in U_i \right\}$$

*Then this is evidently a sheaf and the adjunction $\phi \to \phi^\sharp$ is the unique presheaf morphism such that*

$$\phi^\sharp_U(\sigma)_x = \phi\left((\sigma_x)_{x \in U}\right)_x \quad \forall U \in \mathcal{B}, \sigma \in \mathcal{F}(U), x \in U$$

*and the inverse adjunction $\psi^\flat \leftarrow \psi$ is the unique sheaf morphism such that*

$$\psi^\flat(s)(x) = \psi_x(s(x)) \quad \forall x \in U$$

*(using (4.2.9) to identify $(\mathcal{G}|_{\mathcal{B}})_x$ and $\mathcal{G}_x$). The unit of the adjunction is given by*

$$(\eta_{\mathcal{F}})_U(\sigma) = (\sigma_x)_{x \in U}$$

*and for a morphism of $\mathcal{B}$-sheaves $\phi : \mathcal{F} \to \mathcal{G}$ the morphism $\phi^+$ defined in (4.2.14) is simply*

$$\phi^+(s)(x) = \phi_x(s(x)).$$

*Furthermore $\eta_{\mathcal{F}}$ is an isomorphism if and only if $\mathcal{F}$ is a $\mathcal{B}$-sheaf.*

*Proof.* We first demonstrate this when $\mathcal{C} = \mathbf{Set}$. The sheaf conditions are easy to verify. There is an obvious unit morphism

$$\begin{aligned} \eta_{\mathcal{F}}(U) : \mathcal{F}(U) &\to \mathcal{F}^+(U) \\ \sigma &\to x \to \sigma_x \end{aligned}$$

which is a natural transformation by (4.2.7). According to (2.6.74) it is enough to ensure the following map is a bijection and natural in $\mathcal{G}$

$$\begin{aligned} \mathrm{Mor}(\mathcal{F}^+, \mathcal{G}) &\to \mathrm{Mor}(\mathcal{F}, \mathcal{G}|_{\mathcal{B}}) \\ \phi &\to \phi|_{\mathcal{B}} \circ \eta_{\mathcal{F}} \end{aligned}$$

Naturality in $\mathcal{G}$ follows formally. Suppose $\phi, \psi : \mathcal{F}^+ \to \mathcal{G}$ are morphisms such that $\phi|_{\mathcal{B}} \circ \eta_{\mathcal{F}} = \psi|_{\mathcal{B}} \circ \eta_{\mathcal{F}}$. For a given $s \in \mathcal{F}^+(U)$ we wish to show that $\phi(s) = \psi(s)$. Suppose that $s$ is determined by sections $\sigma_i \in \mathcal{F}(U_i)$, that is to say $s|_{U_i} = \eta_{\mathcal{F}}(\sigma_i)$. Therefore by hypothesis $\phi(s)|_{U_i} = \phi\left(s|_{U_i}\right) = \psi\left(s|_{U_i}\right) = \psi(s)|_{U_i}$. By the concrete sheaf conditions (4.2.4) on $\mathcal{G}$ we conclude that $\phi(s) = \psi(s)$, and so the given map is injective.

© David Rufino CC BY-NC-ND 4.0

Conversely given $\phi : \mathcal{F} \to \mathcal{G}|_{\mathcal{B}}$, we wish to construct a morphism of sheaves $\phi^\flat : \mathcal{F}^+(U) \to \mathcal{G}(U)$ such that $\phi^\flat\big((\sigma_x)_{x\in U}\big) = \phi(\sigma)$ for all $\sigma \in \mathcal{F}(U)$ and $U \in \mathcal{B}$. We claim that there exists a family of maps $\big\{\phi^\flat_U : \mathcal{F}^+ \to \mathcal{G}\big\}$ such that $\phi^\flat_U(s)_x = \phi_x(s(x))$ for all $s \in \mathcal{F}^+(U)$. Such a family is unique by (4.2.8).

To show existence consider a section $s \in \mathcal{F}^+(U)$, then by definition $s(x) = \big((\sigma_i)_x\big)$ for some $\sigma_i \in \mathcal{F}(U_i)$, $U_i \in \mathcal{B}$ and all $x \in U_i$. Define $\tau_i \coloneqq \phi(\sigma_i)$ then by (4.2.7) $(\tau_i)_x = \phi_x\big((\sigma_i)_x\big) = \phi_x(s(x))$. This shows that $(\tau_i)_x = \big(\tau_j\big)_x$ for all $x \in U_i \cap U_j$ so by (4.2.8) $\tau_i|_{U_i\cap U_j} = \tau_j|_{U_i\cap U_j}$. By (4.2.4) there exists $\tau \in \mathcal{G}(U)$ such that $\tau|_{U_i} = \tau_i$. This yields a map $\phi^\flat_U$ satisfying $\phi^\flat_U(s)_x = \tau_x = (\tau_i)_x = \phi(\sigma_i)_x = \phi_x\big((\sigma_i)_x\big) = \phi_x(s(x))$ as required.

It remains to show that $\phi^\flat$ is a morphism of sheaves, that is the following diagram is commutative for all open sets $V \subset U$

$$\begin{array}{ccc}
\mathcal{F}^+(U) & \xrightarrow{\phi^\flat_U} & \mathcal{G}(U) \\
\downarrow{\scriptstyle \rho_{UV}} & & \downarrow{\scriptstyle \rho_{UV}} \\
\mathcal{F}^+(V) & \xrightarrow{\phi^\flat_V} & \mathcal{G}(V) \\
\downarrow & & \downarrow{\scriptstyle \rho_x} \\
\mathcal{F}_x & \xrightarrow{\phi_x} & \mathcal{G}_x
\end{array}$$

The large and bottom squares commute by construction. Therefore for $s \in \mathcal{F}^+(U)$ we have $\big(\phi^\flat_U(s)|_V\big)_x = \phi_x(s|_V\ (x)) = \phi^\flat_V(s|_V)_x$. As this holds for all $x \in V$ we find from (4.2.8) that $\phi^\flat_U(s)|_V = \phi^\flat_V(s|_V)$ as required.

As $(-)^\sharp$ is injective it is sufficient by (2.6.36) to show that it has a right inverse, specifically $\big(\psi^\flat\big)^\sharp = \psi$. However $\psi(\sigma)_x = \psi^\flat\big((\sigma_{x'})_{x'\in U}\big)(x) = \big(\psi^\flat\big)^\sharp(\sigma)_x$ by construction so we are done by (4.2.8).

Evidently when $\eta_{\mathcal{F}}$ is an isomorphism then $\mathcal{F}$ is a $\mathcal{B}$-sheaf. We claim that the converse holds, namely suppose $\mathcal{F}$ is a $\mathcal{B}$-sheaf, then we require to show $(\eta_{\mathcal{F}})_U$ is injective and surjective. It is injective by (4.2.8). Similarly it is surjective for suppose $s \in \mathcal{F}^+(U)$ for $U \in \mathcal{B}$ and $s(x) = (\sigma_i)_x$ for all $x \in U_i$ and $\sigma_i \in \mathcal{F}(U_i)$. Then by the same result we have $(\sigma_i)|_{U_i\cap U_j} = \big(\sigma_j\big)|_{U_i\cap U_j}$, whence by (4.2.4) these lift to a section $\sigma \in \mathcal{F}(U)$ such that $\sigma|_{U_i} = \sigma_i$, and in particular $\sigma_x = (\sigma_i)_x = s(x)$ for all $x \in U_i$. Therefore we conclude the map $(\eta_{\mathcal{F}})_U$ is surjective and therefore an isomorphism in **Set**, which means $\eta_{\mathcal{F}}$ is an isomorphism of sheaves.

We now turn to the case of a general algebraic category $\mathcal{C}$, determined by $(\Omega, E)$ (2.6.184) write $U : \mathcal{C} \to \mathbf{Set}$ for the forgetful functor.

*The remainder of this proof is drafted by Claude Sonnet 5, reviewed by David Rufino.*

By (2.6.190), for $U \in \mathcal{B}$ the product $\prod_{x\in U} \mathcal{F}_x$ exists in $\mathcal{C}$, with each $\omega \in \Omega$ acting coordinatewise.

**$\mathcal{F}^+(U)$ is a subalgebra of $\prod_{x\in U} \mathcal{F}_x$.** Given $\omega \in \Omega$ and $s^{(1)}, \dots, s^{(n_\omega)} \in \mathcal{F}^+(U)$, and $x \in U$: choose for each $r = 1, \dots, n_\omega$ a representative $\sigma^{(r)} \in \mathcal{F}(V_r)$, $V_r \in \mathcal{B}$, $x \in V_r$, with $s^{(r)}(y) = \big(\sigma^{(r)}\big)_y$ for all $y \in V_r$. As $n_\omega$ is finite and $\mathcal{B}$ is closed under finite intersections, $N \coloneqq \bigcap_r V_r \in \mathcal{B}$, and, by functoriality of stalks (4.2.7), $\tau \coloneqq \omega_{\mathcal{F}(N)}\big(\sigma^{(1)}|_N, \dots, \sigma^{(n_\omega)}|_N\big) \in \mathcal{F}(N)$ has $\tau_y = \omega_{\mathcal{F}_y}\big(s^{(1)}(y), \dots, s^{(n_\omega)}(y)\big)$ for every $y \in N$. So the germ-function $y \mapsto \omega_{\mathcal{F}_y}\big(s^{(1)}(y), \dots, s^{(n_\omega)}(y)\big)$ — which is exactly $\omega$ applied coordinatewise to $s^{(1)}, \dots, s^{(n_\omega)}$ in $\prod_{x\in U} \mathcal{F}_x$ — is locally represented by $\tau$ near every point, hence lies in $\mathcal{F}^+(U)$. So $\mathcal{F}^+(U)$ is closed under every $\omega \in \Omega$, and by (2.6.193) is a subalgebra, in particular an object of $\mathcal{C}$, with $\omega_{\mathcal{F}^+(U)}$ given pointwise as above.

**The maps constructed above are morphisms of $\mathcal{C}$.** If $h : A \to B$ is a homomorphism and $S \subseteq U(A)$, $T \subseteq U(B)$ are subalgebras with $h(S) \subseteq T$, then $h|_S : S \to T$ is a homomorphism, as $\omega_S, \omega_T$ are literally restrictions of $\omega_A, \omega_B$. We apply this three times.

For the restriction maps: the coordinate projection $\prod_{x\in U} \mathcal{F}_x \to \prod_{x\in V} \mathcal{F}_x$ ($V \subseteq U$ in $\mathcal{B}$) is a homomorphism, carries $\mathcal{F}^+(U)$ into $\mathcal{F}^+(V)$ by definition of restriction of germ-functions, and restricts to $\rho_{UV}$.

For the unit: each stalk map $\rho_{Ux} : \mathcal{F}(U) \to \mathcal{F}_x$ is a morphism of $\mathcal{C}$ (4.2.5), so $(\rho_{Ux})_{x\in U} : \mathcal{F}(U) \to \prod_{x\in U} \mathcal{F}_x$ is a homomorphism (a map into a product is a homomorphism iff each component is); its image lies in $\mathcal{F}^+(U)$ (represented by $\sigma$ itself, via the cover $\{U\}$), and it restricts to $(\eta_{\mathcal{F}})_U$.

For $\phi^\flat$: given a morphism $\phi : \mathcal{F} \to \mathcal{G}|_{\mathcal{B}}$ of $\mathcal{C}$-presheaves with $\mathcal{G}$ a $\mathcal{B}$-sheaf, each $\phi_x : \mathcal{F}_x \to \mathcal{G}_x$ is a morphism of $\mathcal{C}$ (4.2.7), so $h \coloneqq \prod_{x\in U} \phi_x : \prod_{x\in U} \mathcal{F}_x \to \prod_{x\in U} \mathcal{G}_x$ is a homomorphism. As above, $(\rho_{Ux})_{x\in U} : \mathcal{G}(U) \to \prod_{x\in U} \mathcal{G}_x$ is an injective (by (4.2.8), as $\mathcal{G}$ is a sheaf) homomorphism, so identifies $\mathcal{G}(U)$ with a subalgebra $T$ of $\prod_{x\in U} \mathcal{G}_x$ — its image, which

© David Rufino CC BY-NC-ND 4.0

is a subalgebra since the image of a homomorphism is always closed under the operations of the target. By the construction of $\phi^\flat_U$ above, $h(\mathcal{F}^+(U)) \subseteq T$: indeed for $s \in \mathcal{F}^+(U)$ we exhibited $\tau \in \mathcal{G}(U)$ with $\tau_x = \phi_x(s(x))$ for all $x \in U$. So $h$ restricts to a homomorphism $\mathcal{F}^+(U) \to T \cong \mathcal{G}(U)$, which is $\phi^\flat_U$.

Given this, every map constructed in the case $\mathcal{C} = \mathbf{Set}$ above is now a morphism of $\mathcal{C}$, and the remaining verifications — injectivity of $\phi \mapsto \phi|_{\mathcal{B}} \circ \eta_{\mathcal{F}}$, existence and commutativity of $\phi^\flat$, and the final isomorphism criterion — proceed entirely by manipulating elements of underlying sets via (4.2.8) and (4.2.4), both already stated for general algebraic categories, so the argument given there applies verbatim; in the final step, bijectivity of $(\eta_{\mathcal{F}})_U$ on underlying sets upgrades to an isomorphism of $\mathcal{C}$ since $U$ reflects isomorphisms (2.6.189). ∎

**Corollary 4.2.17**
*Let $\mathcal{C}$ be an algebraic category (2.6.184) then there is an equivalence of categories*

$$\mathbf{Sh}(X;\mathcal{B}) \underset{(-)|_{\mathcal{B}}}{\overset{(-)^+}{\rightleftarrows}} \mathbf{Sh}(X)$$

*Proof.* By the last statement of (4.2.16), and (2.6.73) the essential image of $(-)|_{\mathcal{B}}$ is the full subcategory of $\mathcal{B}$-sheaves, and the unit and counit natural transformations induce the equivalence of categories. ∎

**Lemma 4.2.18 (Sheafification Preserves Stalks)**
*Let $\mathcal{C}$ be algebraic (2.6.184), $\mathcal{F}$ a presheaf on $X$ and $x \in X$. Then $\eta_{\mathcal{F},x} : \mathcal{F}_x \to (\mathcal{F}^+)_x$ (4.2.7) is an isomorphism.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

**Surjectivity.** Let $\xi \in (\mathcal{F}^+)_x$; by (4.2.6), $\xi = s_x$ for some $s \in \mathcal{F}^+(U)$, $U \ni x$. By (4.2.16), $s(y) = (\sigma_i)_y$ for all $y \in U_i$, for some cover $U = \bigcup_i U_i$ and $\sigma_i \in \mathcal{F}(U_i)$; choosing $i$ with $x \in U_i$, this says $s|_{U_i} = \eta_{U_i}(\sigma_i)$, so

$$\xi = s_x = \left(s|_{U_i}\right)_x = \left(\eta_{U_i}(\sigma_i)\right)_x = \eta_x\left((\sigma_i)_x\right)$$

so $\xi \in \mathrm{Im}(\eta_x)$.

**Injectivity.** Suppose $\eta_x(a) = \eta_x(b)$ for $a, b \in \mathcal{F}_x$. By (4.2.6), $a = \sigma_x$, $b = \tau_x$ for some $\sigma \in \mathcal{F}(U)$, $\tau \in \mathcal{F}(V)$, $U, V \ni x$; then $\eta_x(a) = (\eta_U(\sigma))_x$ and $\eta_x(b) = (\eta_V(\tau))_x$, so by (4.2.8).a) there is $W \subseteq U \cap V$, $x \in W$, with $\eta_U(\sigma)|_W = \eta_V(\tau)|_W$ in $\mathcal{F}^+(W)$, that is $\eta_W(\sigma|_W) = \eta_W(\tau|_W)$. As elements of $\mathcal{F}^+(W)$ are functions $W \to \coprod_{y \in W} \mathcal{F}_y$ (4.2.16), this equality holds in particular at $y = x$:

$$(\sigma|_W)_x = \eta_W(\sigma|_W)(x) = \eta_W(\tau|_W)(x) = (\tau|_W)_x$$

that is $\sigma_x = \tau_x$, so $a = b$. ∎

## 4.2.3 Inverse Image Functor

**Definition 4.2.19 (Inverse Image Sheaf)**
*Let $f : X \to Y$ be a continuous map and $\mathcal{G}$ a presheaf on $Y$. We define as follows the **inverse image sheaf***

$$(f^{-1}\mathcal{G})(U) := \left\{ s : U \to \coprod_{x \in U} \mathcal{G}_{f(x)} \;\middle|\; \forall x \in U, \exists V \in \mathcal{O}_{f(x)} \text{ and } \sigma \in \mathcal{G}(V) \text{ s.t. } s(y) = \sigma_y \text{ for all } y \in f^{-1}(V) \right\}$$

*By definition it is a sheaf. Furthermore we see there is a canonical isomorphism*

$$\begin{aligned} (f^{-1}\mathcal{G})_x &\xrightarrow{\sim} \mathcal{G}_{f(x)} \\ (U, s) &\to \quad s(x) \end{aligned}$$

## 4.2.4 Skyscraper Sheaf

**Lemma 4.2.20**
*Consider a family of objects in $\{A_i\}_{i \in I}$ such that $i \in I_0 \implies A_i$ is terminal. Then there is a canonical isomorphism for which the following diagram commutes for all morphisms $f$*

$$\begin{array}{ccc} B & \xrightarrow{f} & \prod_{i \in I} A_i \\ & \searrow^{(\pi_i \circ f)_{i \in I_0}} & \downarrow \sim \\ & & \prod_{i \in I_0} A_i \end{array}$$

© David Rufino CC BY-NC-ND 4.0

*namely that given by* $(\pi_i)_{i\in I_0}$.

*Proof.* Let $\phi = (\pi_i)_{i\in I_0}$ and let $\psi : \prod_{i\in I_0} A_i \to \prod_{i\in I} A_i$ be the unique morphism such that

$$\pi_j \circ \psi = \begin{cases} \pi_j & j \in I_0 \\ !_{\prod_{i\in I_0} A_i} & j \notin I_0 \end{cases}$$

Then $\pi_i \circ \psi \circ \phi = \pi_i \circ \phi = \pi_i$ when $i \in I_0$ and it equals $!_{\prod_{i\in I_0} A_i} \circ \phi = !_{\prod_{i\in I} A_i} = \pi_i$ when $i \notin I_0$, whence $\psi \circ \phi = 1$. The reverse is similar. ∎

**Proposition 4.2.21**
*Let $A \in \mathcal{C}$ be an object, $\star$ a terminal object and $x \in X$. The following assignment*

$$i_x(A)(U) := \begin{cases} A & x \in U \\ \star & x \notin U \end{cases}$$

*determines a sheaf with restriction maps given by*

$$\rho_{UV} := \begin{cases} 1_A & x \in V \\ !_A & x \in U \setminus V \\ 1_\star & x \notin U \end{cases}$$

*This is known as the* ***skyscraper sheaf*** *at $x$.*

*Proof.* Suppose that $U = \bigcup_{i\in I} U_i$ and let $I_0 = \{i \in I \mid x \in U_i\}$ then we may show there is a commutative diagram

$$\begin{array}{ccccc} \mathcal{F}(U) & \xrightarrow{e} & \prod_{i\in I} \mathcal{F}(U_i) & \underset{d_1}{\overset{d_0}{\rightrightarrows}} & \prod_{(i,j)\in I\times I} \mathcal{F}(U_i \cap U_j) \\ & \searrow^{e} & \downarrow \sim & & \downarrow \sim \\ & & \prod_{i\in I_0} \mathcal{F}(U_i) & \underset{d_1}{\overset{d_0}{\rightrightarrows}} & \prod_{(i,j)\in I_0\times I_0} \mathcal{F}(U_i \cap U_j) \end{array}$$

where $e = \left(\rho_{UU_i}\right)$, $d_0 = \left(\rho_{U_i(U_i\cap U_j)} \circ \pi_i\right)$ and $d_1 = \left(\rho_{U_j(U_i\cap U_j)} \circ \pi_j\right)$. Then it's enough to show that the bottom diagram is an equaliser.

For $i, j \in I_0$ we have $\mathcal{F}(U) = \mathcal{F}(U_i) = \mathcal{F}(U_i \cap U_j) = A$.

Suppose $f : B \to \prod_{i\in I_0} \mathcal{F}(U_i)$ such that $d_0 \circ f = d_1 \circ f$. Then

$$\begin{aligned} d_0 \circ f = d_1 \circ f &\implies \pi_{(i,j)} \circ d_0 \circ f = \pi_{(i,j)} \circ d_1 \circ f \quad \forall i, j \in I_0 \\ &\implies 1_A \circ \pi_i \circ f = 1_A \circ \pi_j \circ f \quad \forall i, j \in I_0 \\ &\implies \pi_i \circ f = \pi_j \circ f \quad \forall i, j \in I_0 \end{aligned}$$

Therefore define the morphism $f' : B \to \mathcal{F}(U)$ to be $\pi_j \circ f$ for some $j \in I_0$. Observing $e = (1_A)_{i\in I_0}$ we have $e \circ f' = (f')_{i\in I_0} = \left(\pi_j \circ f\right)_{i\in I_0} = (\pi_i \circ f)_{i\in I_0} = f$. ∎

**Proposition 4.2.22 (Stalk functor is left-adjoint to Sky-Scraper Sheaf)**
*Let $X$ be a topological space, $\mathcal{C}$ a category with a terminal object $\star$ and directed colimits, $A \in \mathcal{C}$ and $\mathcal{F}$ a sheaf on $X$. Then for all $x \in X$ there is a bijection*

$$\begin{aligned} \mathrm{Mor}(\mathcal{F}_x, A) &\xrightarrow{\sim} \mathrm{Mor}(\mathcal{F}, i_x(A)) \\ \phi \to \quad \phi^\sharp_U &:= \begin{cases} \phi \circ \rho_{Ux} & x \in U \\ !_{\mathcal{F}(U)} & x \notin U \end{cases} \end{aligned}$$

*which is natural in both $\mathcal{F}$ and $A$.*

*Proof.* Let $\mathcal{N}_x$ be the category of open subsets $U$ of $X$ containing $x$ ordered by reverse inclusion. Then by definition of the stalk (4.2.7) there is a bijection

$$\begin{aligned} \mathrm{Mor}(\mathcal{F}_x, A) &\xrightarrow{\sim} \left(\mathcal{F}|_{\mathcal{N}_x} \downarrow A\right) \\ \phi &\to (\phi \circ \rho_{Ux})_{U\in\mathcal{N}_x} \end{aligned}$$

© David Rufino CC BY-NC-ND 4.0

which is natural in $\mathcal{F}$ and $A$. It then remains to verify that there is a natural bijection

$$\left(\mathcal{F}|_{\mathcal{N}_x} \downarrow A\right) \xrightarrow{\sim} \mathrm{Mor}(\mathcal{F}, i_x(A))$$

$$(\Phi_U)_{U \ni x} \to \begin{cases} \Phi_U & x \in U \\ !_{\mathcal{F}(U)} & x \notin U \end{cases}$$

■

## 4.2.5 Limit Sheaf

**Proposition 4.2.23 (Limits of Presheaves and Sheaves may be constructed pointwise)**
*Let $K$ be a small category and $F : K \to \mathbf{PSh}(X;\mathcal{C})$ (resp. $\mathbf{Sh}(X;\mathcal{C})$) a diagram, that is a presheaf (resp. sheaf) $F(k)$ for each $k \in K$ together with a morphism $F(\kappa) : F(k) \to F(k')$ for each arrow $\kappa : k \to k'$, functorially. Write $F(-)(U) : K \to \mathcal{C}$ for the diagram $k \mapsto F(k)(U)$, and suppose $\lim_K F(-)(U)$ exists in $\mathcal{C}$ for every $U \subset X$.*

*Then the assignment*

$$\left(\lim_K F\right)(U) := \lim_K F(-)(U)$$

*can be made into a presheaf (resp. sheaf) uniquely such that the projections $\lambda_k : \lim_K F \to F(k)$ are morphisms of presheaves (resp. sheaves) for every $k \in K$. Furthermore $\left(\lim_K F, \{\lambda_k\}_{k\in K}\right)$ is a limit for $F$ in the category of presheaves (resp. sheaves).*

*When $\mathcal{C}$ is an algebraic category (2.6.184) this may be constructed in the obvious way*

$$\left(\lim_K F\right)(U) := \left\{ (\sigma_k)_{k\in K} \in \prod_{k\in K} F(k)(U) \;\middle|\; F(\kappa)_U(\sigma_k) = \sigma_{k'} \text{ for every arrow } \kappa : k \to k' \text{ in } K \right\}$$

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

**Restriction maps.** Fix $V \subseteq U$, and write $\{\lambda_{k,U}\}_{k\in K}$, $\{\lambda_{k,V}\}_{k\in K}$ for the given limit cones for $F(-)(U)$, $F(-)(V)$. For every arrow $\kappa : k \to k'$ of $K$, using that $F(\kappa) : F(k) \to F(k')$ is a morphism of presheaves and the cone condition $F(\kappa)_U \circ \lambda_{k,U} = \lambda_{k',U}$,

$$F(\kappa)_V \circ \left(\rho_{UV}^{F(k)} \circ \lambda_{k,U}\right) = \rho_{UV}^{F(k')} \circ F(\kappa)_U \circ \lambda_{k,U} = \rho_{UV}^{F(k')} \circ \lambda_{k',U}$$

so $\left\{\rho_{UV}^{F(k)} \circ \lambda_{k,U}\right\}_{k\in K}$ is a cone over $F(-)(V)$ with apex $\lim_K F(-)(U)$. By the universal property of $\lim_K F(-)(V)$, there is a unique morphism $\rho_{UV} : \lim_K F(-)(U) \to \lim_K F(-)(V)$ such that $\lambda_{k,V} \circ \rho_{UV} = \rho_{UV}^{F(k)} \circ \lambda_{k,U}$ for every $k \in K$; this is precisely the condition for each $\lambda_k$ to be a morphism of presheaves.

Taking $U = V$, both $\rho_{UU}$ and $1_{\lim_K F(-)(U)}$ satisfy $\lambda_{k,U} \circ (-) = \lambda_{k,U}$ for every $k$, so $\rho_{UU} = 1_{\lim_K F(-)(U)}$ by uniqueness. For $W \subseteq V \subseteq U$, both $\rho_{VW} \circ \rho_{UV}$ and $\rho_{UW}$ satisfy, for every $k$, $\lambda_{k,W} \circ (-) = \rho_{VW}^{F(k)} \circ \rho_{UV}^{F(k)} \circ \lambda_{k,U} = \rho_{UW}^{F(k)} \circ \lambda_{k,U}$, so $\rho_{VW} \circ \rho_{UV} = \rho_{UW}$ by uniqueness. So $\lim_K F$, with these restriction maps, is a presheaf, and each $\lambda_k$ is a morphism of presheaves; uniqueness of restriction maps making every $\lambda_k$ a morphism of presheaves is immediate from the uniqueness clause of the universal property of $\lim_K F(-)(V)$.

**Limit in the category of presheaves.** Let $\mathcal{K}$ be a presheaf and $\{\mu_k : \mathcal{K} \to F(k)\}_{k\in K}$ a cone over $F$ in $\mathbf{PSh}(X;\mathcal{C})$, that is $F(\kappa) \circ \mu_k = \mu_{k'}$ for every arrow $\kappa : k \to k'$. For each $U$, $\left\{(\mu_k)_U\right\}_{k\in K}$ is a cone over $F(-)(U)$ with apex $\mathcal{K}(U)$, so the universal property of $\lim_K F(-)(U)$ gives a unique $\mu_U : \mathcal{K}(U) \to \lim_K F(-)(U)$ with $\lambda_{k,U} \circ \mu_U = (\mu_k)_U$ for every $k$. For $V \subseteq U$, both $\rho_{UV} \circ \mu_U$ and $\mu_V \circ \rho_{UV}^{\mathcal{K}}$ satisfy, for every $k$, $\lambda_{k,V} \circ (-) = \rho_{UV}^{F(k)} \circ (\mu_k)_U = (\mu_k)_V \circ \rho_{UV}^{\mathcal{K}}$ (naturality of $\mu_k$, and the defining properties of $\mu_U, \mu_V$), so $\rho_{UV} \circ \mu_U = \mu_V \circ \rho_{UV}^{\mathcal{K}}$ by uniqueness in the universal property of $\lim_K F(-)(V)$: that is, $\mu := \{\mu_U\}$ is a morphism of presheaves, with $\lambda_k \circ \mu = \mu_k$ for every $k$ by construction. If $\mu'$ is another morphism of presheaves with $\lambda_k \circ \mu' = \mu_k$ for every $k$, then $\mu'_U$ satisfies $\lambda_{k,U} \circ \mu'_U = (\mu_k)_U$ for every $k$, so $\mu'_U = \mu_U$ by the pointwise uniqueness above, that is $\mu' = \mu$.

**Sheaf case.** Suppose $F(k)$ is a sheaf for every $k \in K$; we show $\lim_K F$ is also a sheaf. Fix $U = \bigcup_{i\in I} U_i$, and let $I^{(1)}$, $D__$ be as in (4.2.3).

Define a bifunctor $H : K \times I^{(1)} \to \mathcal{C}$ by $H(k, -) := D_{F(k)}$ for $k \in K$, and $H(\kappa, 1_l) := F(\kappa)_{U_l}$ for every arrow $\kappa : k \to k'$ of $K$ and $l \in I^{(1)}$ (writing $U_l$ for $U_i$, resp. $U_i \cap U_j$, when $l = i$, resp. $l = (i,j)$); this is a bifunctor precisely because each $F(\kappa)$ is a morphism of presheaves, so commutes with the maps $D_{F(k)}(u_{ij}), D_{F(k)}(v_{ij})$ (and likewise for $F(k')$).

© David Rufino CC BY-NC-ND 4.0

Since each $F(k)$ is a sheaf, (4.2.3) gives $\lim_{I^{(1)}} H(k,-) = F(k)(U)$ for every $k \in K$; applying (2.6.99) (with the roles of $I, J$ there taken to be our $K, I^{(1)}$), this determines a functor $L : K \to \mathcal{C}$ (2.6.96) with $L(k) = F(k)(U)$ for every $k$; as $\lim_{k\in K} L(k) = \lim_K F(-)(U)$ exists by hypothesis, (2.6.99) gives that $\lim_{K\times I^{(1)}} H$ exists, with limit cone $\left(\lim_K F(-)(U), \left\{\eta_{k,l} \circ \lambda_{k,U}\right\}_{(k,l)}\right)$, where $\left(\eta_{k,l}\right)_{l\in I^{(1)}}$ is the limit cone for $F(k)(U)$ over $I^{(1)}$ furnished by (4.2.3) in particular the component at $(k,l)$, for $l \in I^{(1)}$, is $\rho^{F(k)}_{UU_l} \circ \lambda_{k,U}$.

Applying (2.6.99) directly to $H$ (whose symmetric hypothesis, that $\lim_{k\in K} H(k,l)$ exists for every $l \in I^{(1)}$, is exactly $\lim_K F(-)(U_l) = D_{\lim_K F}(l)$ existing by hypothesis) to the cone on $\lim_K F(-)(U)$ constructed above: writing $R(l) \coloneqq \lim_{k\in K} H(k,l) = D_{\lim_K F}(l)$, $\lim_{I^{(1)}} R$ exists, with limit cone component at $l \in I^{(1)}$ the unique morphism $\sigma_l : \lim_K F(-)(U) \to \lim_K F(-)(U_l)$ satisfying $\lambda_{k,U_l} \circ \sigma_l = \rho^{F(k)}_{UU_l} \circ \lambda_{k,U}$ for every $k \in K$. This is exactly the defining property of the restriction map $\rho_{UU_l}$ of $\lim_K F$ established above, so $\sigma_l = \rho_{UU_l}$ by uniqueness.

So $D_{\lim_K F}$ has $\left(\lim_K F(-)(U), \left\{\rho_{UU_l}\right\}_{l\in I^{(1)}}\right)$ as a limit over $I^{(1)}$, and by (4.2.3) this says exactly that $\lim_K F$ satisfies the sheaf condition for this cover, as required.

Alternatively, when $\mathcal{C}$ is algebraic (2.6.184), this can be shown more explicitly, using the concrete construction noted above. By (2.6.85) and (2.6.189) (the forgetful functor $U : \mathcal{C} \to \mathbf{Set}$ creates limits), $\lim_K F(-)(U)$ is exactly $\left\{(\sigma_k)_{k\in K} \in \prod_{k\in K} F(k)(U) \;\middle|\; F(\kappa)_U(\sigma_k) = \sigma_{k'}\ \forall \kappa : k \to k'\right\}$ for every $U$, with $\lambda_{k,U}$ the coordinate projections, and $\rho_{UV}$ acting coordinatewise (restricting each $\sigma_k$).

By (4.2.4) it suffices to verify the two concrete sheaf conditions for $\lim_K F$ directly, using that each $F(k)$ satisfies them.

**Separation.** Suppose $(\sigma_k)_k, (\sigma'_k)_k \in (\lim_K F)(U)$ satisfy $\rho_{UU_i}\left((\sigma_k)_k\right) = \rho_{UU_i}\left((\sigma'_k)_k\right)$ for all $i \in I$. Since restriction acts coordinatewise, $\rho^{F(k)}_{UU_i}(\sigma_k) = \rho^{F(k)}_{UU_i}(\sigma'_k)$ for every $k \in K$ and $i \in I$, so $\sigma_k = \sigma'_k$ by separation for $F(k)$; as this holds for every $k$, $(\sigma_k)_k = (\sigma'_k)_k$.

**Gluing.** Suppose $\left(\sigma_{k,i}\right)_k \in (\lim_K F)(U_i)$ for $i \in I$ satisfy $\rho_{U_i(U_i\cap U_j)}\left(\left(\sigma_{k,i}\right)_k\right) = \rho_{U_j(U_i\cap U_j)}\left(\left(\sigma_{k,j}\right)_k\right)$ for all $(i,j) \in I \times I$. Fix $k \in K$; since restriction acts coordinatewise, $\rho^{F(k)}_{U_i(U_i\cap U_j)}(\sigma_{k,i}) = \rho^{F(k)}_{U_j(U_i\cap U_j)}(\sigma_{k,j})$ for all $(i,j)$, so gluing for $F(k)$ gives $\sigma_k \in F(k)(U)$ with $\rho^{F(k)}_{UU_i}(\sigma_k) = \sigma_{k,i}$ for all $i$. For an arrow $\kappa : k \to k'$ of $K$, and every $i \in I$,

$$\rho^{F(k')}_{UU_i}(F(\kappa)_U(\sigma_k)) = F(\kappa)_{U_i}\left(\rho^{F(k)}_{UU_i}(\sigma_k)\right) = F(\kappa)_{U_i}(\sigma_{k,i}) = \sigma_{k',i} = \rho^{F(k')}_{UU_i}(\sigma_{k'})$$

using naturality of $F(\kappa)$, the defining property of $\sigma_k$, that $\left(\sigma_{k,i}\right)_k \in (\lim_K F)(U_i)$, and the defining property of $\sigma_{k'}$. As $\{U_i\}$ covers $U$, separation for $F(k')$ gives $F(\kappa)_U(\sigma_k) = \sigma_{k'}$. As $\kappa$ was arbitrary, $(\sigma_k)_k \in (\lim_K F)(U)$, with $\rho_{UU_i}\left((\sigma_k)_k\right) = \left(\sigma_{k,i}\right)_k$ for all $i$, as required. ∎

## 4.2.6 Equaliser Sheaf

**Proposition 4.2.24 (Equaliser Sheaf may be constructed pointwise)**
*Let $\alpha, \beta : \mathcal{F} \rightrightarrows \mathcal{G}$ be morphisms of presheaves (resp. sheaves) with a family of pointwise equalisers $\{e_U : \mathrm{eq}(\alpha_U, \beta_U) \to \mathcal{F}(U)\}_{U\subset X}$. Then the assignment*

$$\mathrm{eq}(\alpha,\beta)(U) \coloneqq \mathrm{eq}(\alpha_U, \beta_U)$$

*can be made into a presheaf (resp. sheaf) uniquely such that*

$$e : \mathrm{eq}(\alpha,\beta) \to \mathcal{F}$$

*is a morphism of presheaves (resp. sheaves). Further $e$ is the equaliser of $(\alpha,\beta)$ in the category of presheaves (resp. sheaves).*

*When $\mathcal{C}$ is an algebraic category this may be constructed in the obvious way*

$$\mathrm{eq}(\alpha,\beta)(U) \coloneqq \{\sigma \in \mathcal{F}(U) \mid \alpha_U(\sigma) = \beta_U(\sigma)\}$$

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Let $\mathbf{2} = \{\varnothing \rightrightarrows 1\}$ and let $F : \mathbf{2} \to \mathbf{PSh}(X;\mathcal{C})$ (resp. $\mathbf{Sh}(X;\mathcal{C})$) be the diagram $F(\varnothing) \coloneqq \mathcal{F}$, $F(1) \coloneqq \mathcal{G}$, sending the two morphisms $\varnothing \rightrightarrows 1$ to $\alpha, \beta$ respectively. Then $F(-)(U) : \mathbf{2} \to \mathcal{C}$ is exactly the pair $\alpha_U, \beta_U : \mathcal{F}(U) \rightrightarrows \mathcal{G}(U)$, whose limit is $\mathrm{eq}(\alpha_U, \beta_U)$ (2.6.134), which exists by hypothesis for every $U$. By (4.2.23), the assignment $U \mapsto \mathrm{eq}(\alpha_U, \beta_U)$ can therefore be made into a presheaf (resp. sheaf) uniquely such that the projection $e$ to $\mathcal{F}$ is a morphism of presheaves (resp. sheaves), and $\mathrm{eq}(\alpha,\beta) \coloneqq \lim_{\mathbf{2}} F$, with this projection $e$, is a limit for $F$ in the category of presheaves (resp. sheaves), that is the equaliser of $(\alpha,\beta)$ there (2.6.134), as required.

© David Rufino CC BY-NC-ND 4.0

Alternatively, when $\mathcal{C}$ is algebraic (2.6.184), this follows directly from the algebraic case of (4.2.23): writing $(\sigma_\varnothing, \sigma_1) \in \mathcal{F}(U) \times \mathcal{G}(U)$ for the coordinates indexed by $\mathbf{2} = \{\varnothing \rightrightarrows 1\}$, the compatibility condition $\alpha_U(\sigma_\varnothing) = \sigma_1 = \beta_U(\sigma_\varnothing)$ pins $\sigma_1$ down in terms of $\sigma_\varnothing$, so under the projection $(\sigma_\varnothing, \sigma_1) \mapsto \sigma_\varnothing$ this recovers exactly $\mathrm{eq}(\alpha, \beta)(U) = \{\sigma \in \mathcal{F}(U) \mid \alpha_U(\sigma) = \beta_U(\sigma)\}$ noted above. ∎

**Corollary 4.2.25 (Kernel may be constructed pointwise)**
*Let $\mathcal{C}$ be a pointed category and $\alpha : \mathcal{F} \to \mathcal{G}$ a morphism of presheaves (resp. sheaves). Suppose for all $U \subset X$ we have a kernel $(i_U, \ker(\alpha_U))$ of $\alpha_U$.*

*Then we may define $\ker(\alpha)$ as follows*

$$\ker(\alpha)(U) \coloneqq \ker(\alpha_U)$$

*and this may be made into a presheaf (resp. sheaf) uniquely such that $i$ is a morphism of presheaves (resp. sheaves). Furthermore $(i, \ker(\alpha))$ is a kernel in the category of presheaves (resp. sheaves).*

*When $\mathcal{C}$ is an algebraic category (2.6.184) this may be constructed in the obvious way namely*

$$\ker(\alpha)(U) \coloneqq \{\sigma \in \mathcal{F}(U) \mid \alpha_U(\sigma) = 0\}$$

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

By (2.6.150), $(i_U, \ker(\alpha_U))$ is a kernel for $\alpha_U$ if and only if $i_U$ is an equaliser for $(\alpha_U, 0_U)$, where $0 : \mathcal{F} \to \mathcal{G}$ is the zero morphism of presheaves (computed pointwise, since $\mathbf{PSh}(X; \mathcal{C})$, resp. $\mathbf{Sh}(X; \mathcal{C})$, is pointed with zero morphisms computed pointwise). So the hypothesis gives a family of pointwise equalisers $\{i_U : \mathrm{eq}(\alpha_U, 0_U) \to \mathcal{F}(U)\}_{U \subset X}$ for $(\alpha, 0)$, and applying (4.2.24) to this pair, $\mathrm{eq}(\alpha, 0)(U) \coloneqq \mathrm{eq}(\alpha_U, 0_U) = \ker(\alpha_U) = \ker(\alpha)(U)$ can be made into a presheaf (resp. sheaf) uniquely such that $i$ is a morphism of presheaves (resp. sheaves), and $i$ is the equaliser of $(\alpha, 0)$ in the category of presheaves (resp. sheaves); by (2.6.150) again, this is exactly $(i, \ker(\alpha))$ being a kernel for $\alpha$.

When $\mathcal{C}$ is algebraic, the algebraic case of (4.2.24), applied to $(\alpha, 0)$, gives $\mathrm{eq}(\alpha, 0)(U) = \{\sigma \in \mathcal{F}(U) \mid \alpha_U(\sigma) = 0_U(\sigma)\} = \{\sigma \in \mathcal{F}(U) \mid \alpha_U(\sigma) = 0\}$, that is $\ker(\alpha)(U) = \{\sigma \in \mathcal{F}(U) \mid \alpha_U(\sigma) = 0\}$, as stated. ∎

## 4.2.7 Product Presheaf

**Proposition 4.2.26 (Product Presheaf may be constructed pointwise)**
*Let $\{\mathcal{F}_i\}_{i \in I}$ be a family of presheaves (resp. sheaves) with a family of pointwise products $\left\{\left(\prod_{i \in I} \mathcal{F}_i(U), \{\pi_{i,U}\}_{i \in I}\right)\right\}_{U \subset X}$. Then the assignment*

$$\left(\prod_{i \in I} \mathcal{F}_i\right)(U) \coloneqq \prod_{i \in I} \mathcal{F}_i(U)$$

*can be made into a presheaf (resp. sheaf) uniquely such that $\pi_i : \prod_{i \in I} \mathcal{F}_i \to \mathcal{F}_i$ is a morphism of presheaves (resp. sheaves) for every $i \in I$. Furthermore $\left(\prod_{i \in I} \mathcal{F}_i, \{\pi_i\}_{i \in I}\right)$ is a product of $\{\mathcal{F}_i\}_{i \in I}$ in the category of presheaves (resp. sheaves).*

*When $\mathcal{C}$ is algebraic (2.6.184) this may be constructed in the obvious way*

$$\left(\prod_{i \in I} \mathcal{F}_i\right)(U) \coloneqq \left\{(\sigma_i)_{i \in I} \;\middle|\; \sigma_i \in \mathcal{F}_i(U)\right\}$$

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Regard $I$ as a discrete category and let $F : I \to \mathbf{PSh}(X; \mathcal{C})$ (resp. $\mathbf{Sh}(X; \mathcal{C})$) be the diagram $F(i) \coloneqq \mathcal{F}_i$ (a diagram since $I$ is discrete, so there is nothing further to specify). Then $F(-)(U) : I \to \mathcal{C}$ is exactly the discrete family $\{\mathcal{F}_i(U)\}_{i \in I}$, whose limit is $\prod_{i \in I} \mathcal{F}_i(U)$ (2.6.81), which exists by hypothesis for every $U$. By (4.2.23), the assignment $U \mapsto \prod_{i \in I} \mathcal{F}_i(U)$ can therefore be made into a presheaf (resp. sheaf) uniquely such that each projection $\pi_i$ is a morphism of presheaves (resp. sheaves), and $\left(\prod_{i \in I} \mathcal{F}_i, \{\pi_i\}_{i \in I}\right)$ is a limit for $F$ in the category of presheaves (resp. sheaves), that is a product of $\{\mathcal{F}_i\}_{i \in I}$ there (2.6.81), as required.

When $\mathcal{C}$ is algebraic, this is exactly the algebraic case of (4.2.23) specialised to the discrete category $I$, where (there being no arrows) the compatibility condition is vacuous, leaving the unconstrained tuples $(\sigma_i)_{i \in I} \in \prod_{i \in I} \mathcal{F}_i(U)$ noted above. ∎

© David Rufino CC BY-NC-ND 4.0

### 4.2.8 Kernel Pair Sheaf

**Corollary 4.2.27 (Kernel Pair Presheaf may be constructed pointwise)**
*Let $\mu : \mathcal{H} \to \mathcal{G}$ be a morphism of presheaves (resp. sheaves), and suppose for every $U \subset X$ there is a kernel pair $(Z_U, p_{1,U}, p_{2,U})$ of $\mu_U$.*

*Then we may define $(\mathcal{H} \times_{\mathcal{G}} \mathcal{H})(U) \coloneqq Z_U$ and this may be made into a presheaf (resp. sheaf) uniquely such that $p_1, p_2$ are morphisms of presheaves (resp. sheaves). Furthermore $(\mathcal{H} \times_{\mathcal{G}} \mathcal{H}, p_1, p_2)$ is a kernel pair of $\mu$ in the category of presheaves (resp. sheaves).*

*When $\mathcal{C}$ is algebraic (2.6.184) this may be constructed in the obvious way*

$$(\mathcal{H} \times_{\mathcal{G}} \mathcal{H})(U) \coloneqq \{(\sigma_1, \sigma_2) \in \mathcal{H}(U) \times \mathcal{H}(U) \mid \mu_U(\sigma_1) = \mu_U(\sigma_2)\}$$

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Let $F : \mathbf{\Lambda} \to \mathbf{PSh}(X; \mathcal{C})$ (resp. $\mathbf{Sh}(X; \mathcal{C})$) be the diagram $F(a) \coloneqq \mathcal{H}$, $F(b) \coloneqq \mathcal{H}$, $F(c) \coloneqq \mathcal{G}$, sending both morphisms $a \to c$, $b \to c$ to $\mu$ (2.6.144). Then $F(-)(U) : \mathbf{\Lambda} \to \mathcal{C}$ is exactly the cospan $\mu_U : \mathcal{H}(U) \to \mathcal{G}(U) \leftarrow \mathcal{H}(U) : \mu_U$, whose limit is the pullback $(Z_U, p_{1,U}, p_{2,U})$ of $(\mu_U, \mu_U)$, that is a kernel pair of $\mu_U$, which exists by hypothesis for every $U$. By (4.2.23), the assignment $U \mapsto Z_U$ can therefore be made into a presheaf (resp. sheaf) uniquely such that the projections $p_1, p_2$ are morphisms of presheaves (resp. sheaves), and $(\mathcal{H} \times_{\mathcal{G}} \mathcal{H}, p_1, p_2) \coloneqq (\lim_{\mathbf{\Lambda}} F, p_1, p_2)$ is a limit for $F$ in the category of presheaves (resp. sheaves), that is a pullback of $(\mu, \mu)$, i.e. a kernel pair of $\mu$, there, as required.

When $\mathcal{C}$ is algebraic, this is the algebraic case of (4.2.23) specialised to $\mathbf{\Lambda}$: writing $(\sigma_a, \sigma_b, \sigma_c)$ for the coordinates, the compatibility condition $\mu_U(\sigma_a) = \sigma_c = \mu_U(\sigma_b)$ pins $\sigma_c$ down in terms of $\sigma_a, \sigma_b$, so under the projection $(\sigma_a, \sigma_b, \sigma_c) \mapsto (\sigma_a, \sigma_b)$ this recovers exactly $\{(\sigma_1, \sigma_2) \in \mathcal{H}(U) \times \mathcal{H}(U) \mid \mu_U(\sigma_1) = \mu_U(\sigma_2)\}$ noted above. ∎

**Proposition 4.2.28 (Monomorphisms of Presheaves and Sheaves)**
*Let $\phi : \mathcal{F} \to \mathcal{G}$ be a morphism of presheaves (resp. sheaves). Then the following are equivalent*

*a) $\phi$ is a monomorphism*
*b) $\phi_U$ is monic in $\mathcal{C}$ for all $U \subset X$*

*If $\mathcal{C}$ is algebraic (2.6.184), then a), b) are further equivalent to*

*c) $\phi_U$ is injective for all $U \subset X$*

*If moreover $\mathcal{F}, \mathcal{G}$ are sheaves, a)–c) are equivalent to*

*d) $\phi_x$ is injective for all $x \in X$*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

*b) $\implies$ a)*: Suppose $\alpha, \beta : \mathcal{H} \to \mathcal{F}$ satisfy $\phi\alpha = \phi\beta$. Composition of morphisms of presheaves is computed pointwise, so $\phi_U \alpha_U = \phi_U \beta_U$ for every $U$; as $\phi_U$ is monic, $\alpha_U = \beta_U$ for every $U$, that is $\alpha = \beta$.

*a) $\implies$ b)*: Since $\mathcal{C}$ has products and equalisers, by (2.6.145) $\phi_U$ has a kernel pair for every $U$, so by (4.2.27) these assemble into a kernel pair $(\mathcal{F} \times_{\mathcal{G}} \mathcal{F}, p_1, p_2)$ of $\phi$ in the category of presheaves (resp. sheaves), with $U$-components $\left(\mathcal{F}(U) \times_{\mathcal{G}(U)} \mathcal{F}(U), p_{1,U}, p_{2,U}\right)$ a kernel pair of $\phi_U$. As $\phi$ is monic, (2.6.147) gives $p_1 = p_2$; since morphisms of presheaves are equal iff equal pointwise, $p_{1,U} = p_{2,U}$ for every $U$, so by (2.6.147) again (applied in $\mathcal{C}$), $\phi_U$ is monic for every $U$.

*b) $\iff$ c)* (when $\mathcal{C}$ is algebraic): immediate from (2.6.194) (monic iff injective), applied at each $U$.

*c) $\implies$ d)*: (4.2.10).

*d) $\implies$ c)*: (4.2.10). ∎

**Corollary 4.2.29 (Monomorphism of Sheaves is an Isomorphism iff Surjective on Stalks)**
*Let $\mathcal{C}$ be algebraic (2.6.184) and $m : \mathcal{F} \to \mathcal{G}$ a monomorphism of sheaves. Then $m$ is an isomorphism if and only if $m_x$ is surjective for every $x \in X$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

By (4.2.28), $m_U$ is monic, hence injective (2.6.194) (as $\mathcal{C}$ is algebraic), for every $U \subset X$, so by (4.2.10), $m_x$ is injective for every $x \in X$.

As $\mathcal{C}$ is algebraic, $U$ reflects isomorphisms (2.6.189), so an injective morphism of $\mathcal{C}$ is an isomorphism if and only if it is surjective; applying this to $m_x$ (already known injective) for each $x$, $m_x$ is an isomorphism for every $x \in X$ if and only if $m_x$ is surjective for every $x \in X$. The result now follows from (4.2.11) ($a$) $\iff$ $d$)). ∎

© David Rufino CC BY-NC-ND 4.0

**Definition 4.2.30 (Subpresheaf, Subsheaf)**
*Let $\mathcal{C}$ be algebraic (2.6.184) and $\mathcal{F}$ a presheaf on $X$. A **subpresheaf** of $\mathcal{F}$ is an assignment, to every open $U \subset X$, of a subalgebra $\mathcal{G}(U) \subseteq U(\mathcal{F}(U))$ (2.6.192), such that $\rho^{\mathcal{F}}_{UV}(\mathcal{G}(U)) \subseteq \mathcal{G}(V)$ for every $V \subseteq U$.*

*By (2.6.193), each $\mathcal{G}(U)$ carries a unique $\mathcal{C}$-algebra structure making the inclusion $\iota_U : \mathcal{G}(U) \hookrightarrow \mathcal{F}(U)$ a morphism of $\mathcal{C}$; since $\rho^{\mathcal{F}}_{UV}$ is a homomorphism carrying $\mathcal{G}(U)$ into $\mathcal{G}(V)$, it restricts to a morphism $\rho^{\mathcal{G}}_{UV} : \mathcal{G}(U) \to \mathcal{G}(V)$ of $\mathcal{C}$, making $\mathcal{G}$ into a presheaf and $\iota := \{\iota_U\} : \mathcal{G} \to \mathcal{F}$ into a morphism of presheaves.*

*We say $\mathcal{G}$ is a **subsheaf** of $\mathcal{F}$ if moreover $\mathcal{G}$, with this presheaf structure, is a sheaf.*

**Remark 4.2.31**
*Each $\iota_U$ is injective, hence monic (2.6.194), so $\iota$ itself is monic (4.2.28). Conversely, if $\iota : \mathcal{G} \to \mathcal{F}$ is a monomorphism of presheaves (resp. sheaves), then each $\iota_U$ is injective (4.2.28) and (2.6.194), and its image $U(\iota_U)(U(\mathcal{G}(U))) \subseteq U(\mathcal{F}(U))$ is a subalgebra (being the image of a homomorphism), so $\iota$ identifies $\mathcal{G}$ with a subpresheaf (resp. subsheaf) of $\mathcal{F}$.*

**Proposition 4.2.32 (Sheafification of a Subpresheaf)**
*Let $\mathcal{C}$ be algebraic (2.6.184) and $\iota : \mathcal{F} \to \mathcal{G}$ a monomorphism of presheaves with $\mathcal{G}$ a sheaf. Define*

$$\mathcal{F}^+(U) := \left\{ \tau \in U(\mathcal{G}(U)) \;\middle|\; U = \bigcup_{i \in I} U_i \text{ for some open cover, with } \tau|_{U_i} \in U\big(\iota_{U_i}\big)(U(\mathcal{F}(U_i))) \text{ for every } i \right\}$$

*Then $\mathcal{F}^+$ is a subsheaf of $\mathcal{G}$, and the corestriction $\iota' : \mathcal{F} \to \mathcal{F}^+$ of $\iota$ is a sheafification of $\mathcal{F}$ (4.2.13) (justifying, by (2.6.64), the use of the notation $\mathcal{F}^+$ for it, as usual).*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

**$\mathcal{F}^+$ is a subsheaf.** For $\omega \in \Omega$ and $\tau_1, ..., \tau_{n_\omega} \in \mathcal{F}^+(U)$, choose covers $U = \bigcup_i U_i^{(r)}$ witnessing $\tau_r \in \mathcal{F}^+(U)$; the common refinement $\left\{U_{i_1}^{(1)} \cap \cdots \cap U_{i_{n_\omega}}^{(n_\omega)}\right\}$ is again a cover of $U$, and on each piece $W$ of it, $\tau_r|_W = \iota_W(\sigma_r)$ for some $\sigma_r \in \mathcal{F}(W)$ (restricting $\tau_r|_{U_{i_r}^{(r)}} = \iota_{U_{i_r}^{(r)}}(\sigma_r)$ further, as $\iota$ is a morphism of presheaves), so

$$\omega_{\mathcal{G}(U)}\big(\tau_1, ..., \tau_{n_\omega}\big)|_W = \omega_{\mathcal{G}(W)}(\tau_1|_W, ...) = \omega_{\mathcal{G}(W)}(\iota_W(\sigma_1), ...) = \iota_W\big(\omega_{\mathcal{F}(W)}(\sigma_1, ...)\big) \in \iota_W(\mathcal{F}(W))$$

As this holds for every $W$ in the cover, $\omega_{\mathcal{G}(U)}\big(\tau_1, ..., \tau_{n_\omega}\big) \in \mathcal{F}^+(U)$, so $\mathcal{F}^+(U)$ is a subalgebra of $\mathcal{G}(U)$ (2.6.192). Restricting a witnessing cover for $\tau \in \mathcal{F}^+(U)$ to $V \subseteq U$ shows $\rho^{\mathcal{G}}_{UV}(\mathcal{F}^+(U)) \subseteq \mathcal{F}^+(V)$, so $\mathcal{F}^+$ is a subpresheaf of $\mathcal{G}$ (4.2.30).

Separation for $\mathcal{F}^+$ is immediate, since $\mathcal{F}^+(U) \subseteq U(\mathcal{G}(U))$ and $\mathcal{G}$ is separated. For gluing, let $\tau_i \in \mathcal{F}^+(U_i)$ for a cover $U = \bigcup_i U_i$, agreeing on overlaps; as $\mathcal{G}$ is a sheaf there is $\tau \in \mathcal{G}(U)$ with $\tau|_{U_i} = \tau_i$. Each $\tau_i$ has a witnessing cover $U_i = \bigcup_k U_{i,k}$ with $\tau_i|_{U_{i,k}} \in \iota_{U_{i,k}}\big(\mathcal{F}\big(U_{i,k}\big)\big)$; then $\big\{U_{i,k}\big\}_{i,k}$ is a cover of $U$ with $\tau|_{U_{i,k}} = \tau_i|_{U_{i,k}} \in \iota_{U_{i,k}}\big(\mathcal{F}\big(U_{i,k}\big)\big)$, so $\tau \in \mathcal{F}^+(U)$. So $\mathcal{F}^+$ is a subsheaf of $\mathcal{G}$ (4.2.30). (None of this used the germ-function construction of (4.2.16).)

**$\iota'$ satisfies the universal property.** We show directly that for every sheaf $\mathcal{K}$,

$$(-)^\sharp : \mathrm{Mor}(\mathcal{F}^+, \mathcal{K}) \to \mathrm{Mor}(\mathcal{F}, \mathcal{K}), \quad \phi \mapsto \phi \circ \iota'$$

is a bijection; it is then manifestly natural in $\mathcal{K}$, being given by post-composition with the fixed $\iota'$, so this establishes (4.2.13) for $(\iota', \mathcal{F}^+)$.

*Construction of $\psi^\flat$.* Let $\psi : \mathcal{F} \to \mathcal{K}$ and $\tau \in \mathcal{F}^+(U)$, with witnessing cover $U = \bigcup_i U_i$, $\tau|_{U_i} = \iota_{U_i}(\sigma_i)$ for (unique, as $\iota$ is monic) $\sigma_i \in \mathcal{F}(U_i)$. On overlaps, $\iota_{U_i \cap U_j}\big(\sigma_i|_{U_i \cap U_j}\big) = \tau|_{U_i \cap U_j} = \iota_{U_i \cap U_j}\big(\sigma_j|_{U_i \cap U_j}\big)$, so as $\iota$ is monic, $\sigma_i|_{U_i \cap U_j} = \sigma_j|_{U_i \cap U_j}$, whence $\psi_{U_i}(\sigma_i)|_{U_i \cap U_j} = \psi_{U_j}\big(\sigma_j\big)|_{U_i \cap U_j}$. As $\mathcal{K}$ is a sheaf there is therefore a unique $\psi^\flat_U(\tau) \in \mathcal{K}(U)$ with $\psi^\flat_U(\tau)|_{U_i} = \psi_{U_i}(\sigma_i)$ for every $i$.

*Independence of the witnessing cover.* If $U = \bigcup_j V_j$, $\tau|_{V_j} = \iota_{V_j}\big(\rho_j\big)$ is another witnessing cover, then on $U_i \cap V_j$, $\iota_{U_i \cap V_j}\big(\sigma_i|_{U_i \cap V_j}\big) = \tau|_{U_i \cap V_j} = \iota_{U_i \cap V_j}\big(\rho_j|_{U_i \cap V_j}\big)$, so (as $\iota$ monic) $\sigma_i|_{U_i \cap V_j} = \rho_j|_{U_i \cap V_j}$, whence $\psi_{U_i}(\sigma_i)|_{U_i \cap V_j} = \psi_{V_j}\big(\rho_j\big)|_{U_i \cap V_j}$; as $\big\{U_i \cap V_j\big\}_{i,j}$ covers $U$, the two candidate values for $\psi^\flat_U(\tau)$ agree on a common refinement, so are equal by separation of $\mathcal{K}$.

*$\psi^\flat_U$ is a homomorphism.* Given $\tau_1, ..., \tau_{n_\omega} \in \mathcal{F}^+(U)$, using the common refinement cover $\{W\}$ from the first paragraph (with local representatives $\sigma_r \in \mathcal{F}(W)$ for $\tau_r$), independence of witnessing cover gives $\psi^\flat_U(\tau_r)|_W = \psi_W(\sigma_r)$; as this same cover witnesses $\omega_{\mathcal{F}^+(U)}\big(\tau_1, ..., \tau_{n_\omega}\big) \in \mathcal{F}^+(U)$ with local representatives $\omega_{\mathcal{F}(W)}(\sigma_1, ...)$ (shown above), independence of witnessing cover also gives $\psi^\flat_U\big(\omega_{\mathcal{F}^+(U)}(\tau_1, ...)\big)|_W = \psi_W\big(\omega_{\mathcal{F}(W)}(\sigma_1, ...)\big) = \omega_{\mathcal{K}(W)}(\psi_W(\sigma_1), ...) = \omega_{\mathcal{K}(U)}\big(\psi^\flat_U(\tau_1), ...\big)|_W$ for every $W$, so by separation of $\mathcal{K}$, $\psi^\flat_U\big(\omega_{\mathcal{F}^+(U)}\big(\tau_1, ..., \tau_{n_\omega}\big)\big) = \omega_{\mathcal{K}(U)}\big(\psi^\flat_U(\tau_1), ..., \psi^\flat_U\big(\tau_{n_\omega}\big)\big)$.

© David Rufino CC BY-NC-ND 4.0

*$\psi^\flat$ is a morphism of presheaves.* For $V \subseteq U$ and $\tau \in \mathcal{F}^+(U)$ with witnessing cover $\{U_i\}$, $\{U_i \cap V\}$ witnesses $\tau|_V$, so $\psi^\flat_V(\tau|_V)|_{U_i\cap V} = \psi_{U_i\cap V}\left(\sigma_i|_{U_i\cap V}\right) = \psi_{U_i}(\sigma_i)|_{U_i\cap V} = \psi^\flat_U(\tau)|_{U_i\cap V}$ for every $i$, so $\psi^\flat_V(\tau|_V) = \psi^\flat_U(\tau)|_V$ by separation of $\mathcal{K}$.

$\psi^{\flat\sharp} = \psi$. For $\sigma \in \mathcal{F}(U)$, $\iota'_U(\sigma) = \iota_U(\sigma) \in \mathcal{F}^+(U)$ is witnessed by the trivial cover $\{U\}$ with representative $\sigma$ itself, so $\psi^\flat_U(\iota'_U(\sigma)) = \psi_U(\sigma)$, that is $\psi^\flat \circ \iota' = \psi$.

*Uniqueness.* If $\phi : \mathcal{F}^+ \to \mathcal{K}$ also satisfies $\phi \circ \iota' = \psi$, then for $\tau \in \mathcal{F}^+(U)$ with witnessing cover $\{U_i\}$, $\phi_U(\tau)|_{U_i} = \phi_{U_i}\left(\tau|_{U_i}\right) = \phi_{U_i}\left(\iota'_{U_i}(\sigma_i)\right) = \psi_{U_i}(\sigma_i) = \psi^\flat_U(\tau)|_{U_i}$ for every $i$, so $\phi_U(\tau) = \psi^\flat_U(\tau)$ by separation of $\mathcal{K}$; as $U, \tau$ were arbitrary, $\phi = \psi^\flat$.

So $(-)^\sharp$ is a bijection with inverse $\psi \mapsto \psi^\flat$, for every sheaf $\mathcal{K}$: $(\iota', \mathcal{F}^+)$ is a sheafification of $\mathcal{F}$ (4.2.13). ∎

## 4.2.9 Coequaliser Sheaf

**Proposition 4.2.33 (Coequaliser Presheaf)**
*Let $\alpha, \beta : \mathcal{F} \rightrightarrows \mathcal{G}$ be morphisms of presheaves with a family of pointwise coequalisers $\{p_U : \mathcal{G}(U) \to \operatorname{coeq}(\alpha_U, \beta_U)\}_{U\subset X}$. Then we may define*

$$\operatorname{coeq}(\alpha, \beta)(U) \coloneqq \operatorname{coeq}(\alpha_U, \beta_U)$$

*and this may be made into a presheaf uniquely such that $p$ is a morphism of presheaves. Furthermore $(p, \operatorname{coeq}(\alpha, \beta))$ is a coequaliser in the category of presheaves.*

*In particular if $\mathcal{C}$ has all coequalisers then so does $\mathbf{PSh}(X; \mathcal{C})$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

We require to show that for all open subsets $V \subset U$ there is a unique restriction morphism $\rho_{UV} : \operatorname{coeq}(\alpha_U, \beta_U) \to \operatorname{coeq}(\alpha_V, \beta_V)$ ensuring the following diagram commutes

$$\begin{array}{ccc} \mathcal{G}(U) & \xrightarrow{p_U} & \operatorname{coeq}(\alpha_U, \beta_U) \\ {\scriptstyle \rho_{UV}}\downarrow & & \downarrow{\scriptstyle \rho_{UV}} \\ \mathcal{G}(V) & \xrightarrow[p_V]{} & \operatorname{coeq}(\alpha_V, \beta_V) \end{array}$$

We have that

$$p_V \circ \rho_{UV} \circ \alpha_U = p_V \circ \alpha_V \circ \rho_{UV} = p_V \circ \beta_V \circ \rho_{UV} = p_V \circ \rho_{UV} \circ \beta_U$$

whence by the universal property applied to $\operatorname{coeq}(\alpha_U, \beta_U)$ there exists a unique factorisation of $p_V \circ \rho_{UV}$ through $p_U$. The uniqueness then shows that this satisfies the conditions to be a presheaf ($\rho_{UU} = 1$ and $\rho_{VW} \circ \rho_{UV} = \rho_{UW}$ both follow from uniqueness of the factorisation of $p_W \circ \rho_{UW}$ through $p_U$), and $p$ is a morphism of presheaves.

We now show that $p$ is a coequaliser of $(\alpha, \beta)$ in the category of presheaves. Evidently $p \circ \alpha = p \circ \beta$, since this holds pointwise by hypothesis. Suppose that $p' : \mathcal{G} \to \mathcal{H}$ is a morphism of presheaves with $p' \circ \alpha = p' \circ \beta$, then $p'_U \circ \alpha_U = p'_U \circ \beta_U$ whence there is a unique factorisation $h_U$ such that $h_U \circ p_U = p'_U$.




The large square and left-hand square commute, which we may use to show that

$$h_V \circ \rho_{UV} \circ p_U = h_V \circ p_V \circ \rho_{UV} = p'_V \circ \rho_{UV} = \rho_{UV} \circ p'_U = \rho_{UV} \circ h_U \circ p_U$$

© David Rufino CC BY-NC-ND 4.0

As $p_U$ is a coequaliser it is an epimorphism (2.6.135), whence $h_V \circ \rho_{UV} = \rho_{UV} \circ h_U$, which shows that the right-hand square commutes. This means precisely that $h$ is a morphism of presheaves, and by construction $h \circ p = p'$.

Finally suppose $h'$ is another morphism of presheaves with $h' \circ p = p'$. Then for every $U$, $h'_U \circ p_U = p'_U = h_U \circ p_U$, so by the uniqueness of the pointwise factorisation $h'_U = h_U$, that is $h' = h$.

For the final statement, if $\mathcal{C}$ has all coequalisers then the family of pointwise coequalisers $\{p_U : \mathcal{G}(U) \to \mathrm{coeq}(\alpha_U, \beta_U)\}_{U \subset X}$ exists for every pair $\alpha, \beta : \mathcal{F} \rightrightarrows \mathcal{G}$ of morphisms of presheaves, so the above applies to every such pair and $\mathbf{PSh}(X;\mathcal{C})$ has all coequalisers. ∎

**Proposition 4.2.34**
*Let $\alpha, \beta : \mathcal{F} \rightrightarrows \mathcal{G}$ be morphisms of sheaves and $p : \mathcal{G} \to \mathrm{coeq}^{\mathrm{psh}}(\alpha, \beta)$ be a coequaliser in $\mathbf{PSh}(X;\mathcal{C})$. Then*

$$\eta \circ p : \mathcal{G} \to \mathrm{coeq}(\alpha, \beta) := \left(\mathrm{coeq}^{\mathrm{psh}}(\alpha, \beta)\right)^+$$

*is a coequaliser in $\mathbf{Sh}(X;\mathcal{C})$.*

*In particular if $\mathcal{C}$ has all coequalisers then so does $\mathbf{Sh}(X;\mathcal{C})$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

By (4.2.13) and (4.2.14), $(-)^+$ is left adjoint to the inclusion $\mathbf{Sh}(X;\mathcal{C}) \hookrightarrow \mathbf{PSh}(X;\mathcal{C})$ with unit $\eta$, and this inclusion is (trivially) fully faithful, since a morphism of sheaves is by definition just a morphism of the underlying presheaves. Hence this is an instance of (2.6.143), applied to this reflective subcategory with the coequaliser $p$ of $(\alpha, \beta)$ in $\mathbf{PSh}(X;\mathcal{C})$.

For the final statement, if $\mathcal{C}$ has all coequalisers then by (4.2.33) $\mathbf{PSh}(X;\mathcal{C})$ has all coequalisers, so the hypotheses of this proposition are satisfied for every pair $(\alpha, \beta)$ of morphisms of sheaves, and $\mathbf{Sh}(X;\mathcal{C})$ has all coequalisers. ∎

**Definition 4.2.35 (Cokernel Presheaf, Cokernel Sheaf)**
*Let $\mathcal{C}$ be pointed and algebraic (2.6.184) and $\alpha : \mathcal{F} \to \mathcal{G}$ a morphism of presheaves. Define the **cokernel presheaf** of $\alpha$ by*

$$\mathrm{coker}^{\mathrm{psh}}(\alpha) := \mathrm{coeq}^{\mathrm{psh}}(\alpha, 0), \quad q_U(\sigma) := \sigma \bmod \alpha_U(\mathcal{F}(U)) \quad (4.2.33)$$

*with $q : \mathcal{G} \to \mathrm{coker}^{\mathrm{psh}}(\alpha)$ the coequaliser projection of $\alpha$ and the zero morphism in $\mathbf{PSh}(X;\mathcal{C})$. When $\mathcal{F}, \mathcal{G}$ are sheaves, define the **cokernel sheaf** of $\alpha$ by*

$$\mathrm{coker}(\alpha)(U) := \left\{ \tau : U \to \coprod_{x \in U} \mathcal{G}_x / \alpha_x(\mathcal{F}_x) \;\middle|\; U = \bigcup_i U_i \text{ for some open cover, and } \sigma_i \in \mathcal{G}(U_i) \right.$$
$$\left. \text{such that } \tau(x) = \sigma_{i,x} \bmod \alpha_x(\mathcal{F}_x) \text{ for all } x \in U_i \right\}$$

*with restriction maps given by restricting functions, and the **cokernel map** $p : \mathcal{G} \to \mathrm{coker}(\alpha)$ by $p_U(\sigma)(x) := \sigma_x \bmod \alpha_x(\mathcal{F}_x)$.*

**Remark 4.2.36**
*When $\alpha$ is (in particular) a monomorphism, each $\alpha_U$ is injective (4.2.28), identifying $\mathcal{F}$ with a subsheaf of $\mathcal{G}$; in this case we also write $\mathcal{G}/\mathcal{F} := \mathrm{coker}(\alpha)$.*

**Lemma 4.2.37 (Cokernel is a Sheaf, and Commutes with Stalks)**
*Let $\mathcal{C}$ be pointed and algebraic, $\alpha : \mathcal{F} \to \mathcal{G}$ a morphism of sheaves, and $p$ as in (4.2.35). Then $\mathrm{coker}(\alpha)$ is a sheaf, $(\mathrm{coker}(\alpha), p)$ is a coequaliser of $(\alpha, 0)$ in $\mathbf{Sh}(X;\mathcal{C})$, and, for $x \in X$ there is a unique morphism $\theta_x : \mathrm{coker}(\alpha)_x \to \mathcal{G}_x/\alpha_x(\mathcal{F}_x)$ making*




*commute, and $\theta_x$ is (moreover) an isomorphism.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

© David Rufino CC BY-NC-ND 4.0

**An auxiliary sheafification-based model.** By (4.2.34), $p^0 := \eta \circ q : \mathcal{G} \to \operatorname{coker}^0(\alpha) := \left(\operatorname{coker}^{\mathrm{psh}}(\alpha)\right)^+$ is a co-equaliser of $(\alpha, 0)$ in $\mathbf{Sh}(X;\mathcal{C})$, with $q, \operatorname{coker}^{\mathrm{psh}}(\alpha)$ as in (4.2.35) and $\eta$ the sheafification unit (4.2.16) this auxiliary model is used only within this proof, to relate the definition above to the general existence of coequalisers.

**$\eta_x$ is an isomorphism.** (4.2.18).

**$q_x$ is surjective with kernel $\alpha_x(\mathcal{F}_x)$.** Since $\operatorname{coker}^{\mathrm{psh}}(\alpha)(U) = \mathcal{G}(U)/\alpha_U(\mathcal{F}(U))$ (4.2.33), by definition of stalk $\left(\operatorname{coker}^{\mathrm{psh}}(\alpha)\right)_x = \varinjlim_{U \ni x} \mathcal{G}(U)/\alpha_U(\mathcal{F}(U))$, with $q_x(\sigma_x)$ the class of $\sigma$ in this direct limit for $\sigma \in \mathcal{G}(U)$, $U \ni x$.

*Surjective*: every element of the direct limit is represented by the class of some $\sigma \in \mathcal{G}(U)$, $U \ni x$, that is $q_x(\sigma_x)$.

*Kernel* $\supseteq \alpha_x(\mathcal{F}_x)$: for $\rho \in \mathcal{F}_x$, represented by $\tau \in \mathcal{F}(U)$, $\alpha_x(\rho) = (\alpha_U(\tau))_x$; as $\alpha_U(\tau) \in \alpha_U(\mathcal{F}(U))$, its class in $\mathcal{G}(U)/\alpha_U(\mathcal{F}(U))$ is 0, so $q_x(\alpha_x(\rho)) = 0$.

*Kernel* $\subseteq \alpha_x(\mathcal{F}_x)$: suppose $\sigma \in \mathcal{G}(U)$, $U \ni x$, with $q_x(\sigma_x) = 0$, that is, the class of $\sigma$ becomes 0 in the direct limit. By (4.2.6) applied to the presheaf $U \mapsto \mathcal{G}(U)/\alpha_U(\mathcal{F}(U))$, this means there is $W \subseteq U$, $x \in W$, with $\sigma|_W \in \alpha_W(\mathcal{F}(W))$, that is $\sigma|_W = \alpha_W(\tau)$ for some $\tau \in \mathcal{F}(W)$; then $\sigma_x = (\sigma|_W)_x = (\alpha_W(\tau))_x = \alpha_x(\tau_x) \in \alpha_x(\mathcal{F}_x)$.

**Existence and uniqueness of $\vartheta^0_x$ with $\vartheta^0_x \circ q_x = \pi_x$.** Since $q_x$ is surjective with kernel $\alpha_x(\mathcal{F}_x)$, it is (up to the canonical comparison isomorphism) the quotient map $\mathcal{G}_x \to \mathcal{G}_x/\alpha_x(\mathcal{F}_x)$, hence satisfies the corresponding universal property: any morphism out of $\mathcal{G}_x$ vanishing on $\ker(q_x)$ factors uniquely through $q_x$. As $\pi_x$ vanishes on $\ker(\pi_x) = \alpha_x(\mathcal{F}_x) = \ker(q_x)$, this gives a unique $\vartheta^0_x : \left(\operatorname{coker}^{\mathrm{psh}}(\alpha)\right)_x \to \mathcal{G}_x/\alpha_x(\mathcal{F}_x)$ with $\vartheta^0_x \circ q_x = \pi_x$; uniqueness alone accounts for this, and says nothing yet about $\vartheta^0_x$ being an isomorphism.

**$\vartheta^0_x$ is an isomorphism.** This is a separate check. *Surjective*: given $z \in \mathcal{G}_x/\alpha_x(\mathcal{F}_x)$, as $\pi_x$ is surjective $z = \pi_x(g)$ for some $g \in \mathcal{G}_x$, and $z = \vartheta^0_x(q_x(g))$. *Injective*: every element of $\left(\operatorname{coker}^{\mathrm{psh}}(\alpha)\right)_x$ is $q_x(g)$ for some $g$, since $q_x$ is surjective; if $\vartheta^0_x(q_x(g)) = 0$ then $\pi_x(g) = 0$, that is $g \in \ker(\pi_x) = \ker(q_x)$, so $q_x(g) = 0$.

**$\vartheta_x := \vartheta^0_x \circ \eta_x^{-1}$ is an isomorphism $\operatorname{coker}^0(\alpha)_x \to \mathcal{G}_x/\alpha_x(\mathcal{F}_x)$ with $\vartheta_x \circ p^0_x = \pi_x$.** It is an isomorphism, being a composite of the isomorphisms $\eta_x^{-1}$ and $\vartheta^0_x$. As $p^0_x = \eta_x \circ q_x$,

$$\vartheta_x \circ p^0_x = \vartheta^0_x \circ \eta_x^{-1} \circ \eta_x \circ q_x = \vartheta^0_x \circ q_x = \pi_x$$

**Relating $\operatorname{coker}^0(\alpha)$ to $\operatorname{coker}(\alpha)$.** By (4.2.16), elements of $\operatorname{coker}^0(\alpha)(U)$ are functions $\tau^0 : U \to \coprod_{x \in U} \operatorname{coker}^0(\alpha)_x$ such that, for some open cover $U = \bigcup_i U_i$ and $\sigma_i \in \mathcal{G}(U_i)$, $\tau^0(x) = p^0_{U_i}(\sigma_i)_x$ for $x \in U_i$. Using the isomorphisms $\vartheta_x$, define

$$\vartheta : \coprod_{x \in X} \operatorname{coker}^0(\alpha)_x \longrightarrow \coprod_{x \in X} \mathcal{G}_x/\alpha_x(\mathcal{F}_x), \quad \vartheta|_{\operatorname{coker}^0(\alpha)_x} := \vartheta_x$$

a bijection, being a disjoint union of the bijections $\vartheta_x$. Set $\Psi_U(\tau^0) := \vartheta \circ \tau^0$ for $\tau^0 \in \operatorname{coker}^0(\alpha)(U)$.

*$\Psi_U$ is a well-defined bijection onto* $\operatorname{coker}(\alpha)(U)$. As $\vartheta$ is a bijection carrying each fibre $\operatorname{coker}^0(\alpha)_x$ to $\mathcal{G}_x/\alpha_x(\mathcal{F}_x)$, post-composition by $\vartheta$ is a bijection from functions $U \to \coprod_{x \in U} \operatorname{coker}^0(\alpha)_x$ to functions $U \to \coprod_{x \in U} \mathcal{G}_x/\alpha_x(\mathcal{F}_x)$, carrying the representing data $(U_i, \sigma_i)$ for $\tau^0$ to representing data for $\Psi_U(\tau^0)$: $\vartheta\left(p^0_{U_i}(\sigma_i)_x\right) = \vartheta_x\left(p^0_{U_i}(\sigma_i)_x\right) = \sigma_{i,x} \bmod \alpha_x(\mathcal{F}_x)$, so $\Psi_U(\tau^0)(x) = \sigma_{i,x} \bmod \alpha_x(\mathcal{F}_x)$ for $x \in U_i$, that is, exactly membership in $\operatorname{coker}(\alpha)(U)$ (4.2.35). Hence $\Psi_U$ restricts to a bijection $\operatorname{coker}^0(\alpha)(U) \to \operatorname{coker}(\alpha)(U)$.

*$\Psi$ is an isomorphism of presheaves, and* $\operatorname{coker}(\alpha)$ *is a sheaf.* For $V \subseteq U$ and $\tau^0 \in \operatorname{coker}^0(\alpha)(U)$, $\Psi_V(\tau^0|_V) = \vartheta \circ (\tau^0|_V) = (\vartheta \circ \tau^0)|_V = \Psi_U(\tau^0)|_V$, since restricting a function and then composing with $\vartheta$ agrees with composing then restricting; hence $\Psi$ commutes with restriction, and is a natural transformation, bijective at every $U$ — an isomorphism of presheaves $\operatorname{coker}^0(\alpha) \cong \operatorname{coker}(\alpha)$. As $\operatorname{coker}^0(\alpha)$ is a sheaf and the sheaf condition transports across an isomorphism of presheaves, $\operatorname{coker}(\alpha)$ is also a sheaf.

$\Psi \circ p^0 = p$. For $\sigma \in \mathcal{G}(U)$, $x \in U$, $\Psi_U(p^0_U(\sigma))(x) = \vartheta_x(p^0_U(\sigma)_x) = \sigma_x \bmod \alpha_x(\mathcal{F}_x) = p_U(\sigma)(x)$, again using $\vartheta_x \circ p^0_x = \pi_x$.

**$(\operatorname{coker}(\alpha), p)$ is a coequaliser of $(\alpha, 0)$.** Since $p^0 \circ \alpha = p^0 \circ 0$ (as $p^0$ coequalises $(\alpha, 0)$) and $p = \Psi \circ p^0$, also $p \circ \alpha = \Psi \circ p^0 \circ \alpha = \Psi \circ p^0 \circ 0 = p \circ 0$, so $(\operatorname{coker}(\alpha), p)$ is a co-cone for $(\alpha, 0)$; and $\Psi$, an isomorphism of presheaves between sheaves, is also an isomorphism of $\mathbf{Sh}(X;\mathcal{C})$ (this inclusion into $\mathbf{PSh}(X;\mathcal{C})$ being fully faithful (4.2.34)). As $(\operatorname{coker}^0(\alpha), p^0)$ is a coequaliser of $(\alpha, 0)$, that is, a colimit of the corresponding diagram (2.6.140), (2.6.90) shows $(\operatorname{coker}(\alpha), p)$ is also such a colimit, that is, a coequaliser of $(\alpha, 0)$.

**Conclusion.** Set $\theta_x := \vartheta_x \circ \Psi_x^{-1}$, an isomorphism, being a composite of the isomorphisms $\vartheta_x$ and $\Psi_x^{-1}$. Since $p_x = \Psi_x \circ p^0_x$ (4.2.7),

© David Rufino CC BY-NC-ND 4.0

$$\theta_x \circ p_x = \vartheta_x \circ \Psi_x^{-1} \circ \Psi_x \circ p_x^0 = \vartheta_x \circ p_x^0 = \pi_x$$

and $\theta_x$ is unique with this property, since $p_x = \Psi_x \circ p_x^0$ is surjective (a composite of the surjections $\Psi_x$, an isomorphism, and $p_x^0 = \eta_x \circ q_x$), hence epic, so at most one morphism out of $\operatorname{coker}(\alpha)_x$ can satisfy $\theta_x \circ p_x = \pi_x$. ∎

## 4.2.10 Image Sheaf

Unlike kernels and equalisers (4.2.24), the image is not itself a limit — it is initial in the category of factorisations through a monomorphism (2.6.167) — so the argument via (4.2.3) and (2.6.99) used there does not apply, and the pointwise image presheaf need not satisfy the sheaf condition even when $\mathcal{F}, \mathcal{G}$ are sheaves; instead an image sheaf is obtained by sheafifying the image presheaf (4.2.40). We do however first record the following easier pointwise construction of an image presheaf, whenever $\mathcal{C}$ has regular epi-mono factorisations.

**Lemma 4.2.38 (Regular Epimorphisms of Presheaves may be Determined Pointwise)**
*Let $e : \mathcal{F} \to \mathcal{G}$ be a morphism of presheaves such that $e_U$ is a regular epimorphism in $\mathcal{C}$ for every $U \subset X$. Then $e$ is a regular epimorphism of $\mathbf{PSh}(X; \mathcal{C})$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Since $\mathcal{C}$ has products and equalisers, by (2.6.145) $e_U$ has a kernel pair for every $U$, so by (4.2.27) these assemble into a kernel pair $(\mathcal{F} \times_{\mathcal{G}} \mathcal{F}, p_1, p_2)$ of $e$ in $\mathbf{PSh}(X; \mathcal{C})$, with $U$-components $\left(\mathcal{F}(U) \times_{\mathcal{G}(U)} \mathcal{F}(U), p_{1,U}, p_{2,U}\right)$ a kernel pair of $e_U$.

Since $e_U$ is regular, (2.6.149) shows $e_U$ is the coequaliser of $\left(p_{1,U}, p_{2,U}\right)$, for every $U$. So applying (4.2.33) to the pair of morphisms of presheaves $(p_1, p_2) : \mathcal{F} \times_{\mathcal{G}} \mathcal{F} \rightrightarrows \mathcal{F}$, this pointwise family of coequalisers assembles into a presheaf $\operatorname{coeq}(p_1, p_2)$ with restriction maps uniquely determined by making the induced map $q := \{e_U\}$ a morphism of presheaves, and $(q, \operatorname{coeq}(p_1, p_2))$ is a coequaliser of $(p_1, p_2)$ in $\mathbf{PSh}(X; \mathcal{C})$.

The presheaf $\mathcal{G}$ also has $U$-component $\mathcal{G}(U) = \operatorname{coeq}(p_1, p_2)(U)$, with restriction maps uniquely determined by the same condition, that $e$ is a morphism of presheaves; by this uniqueness, $\mathcal{G} = \operatorname{coeq}(p_1, p_2)$ as presheaves and $e = q$. Hence $e$ is a coequaliser of presheaves, i.e. a regular epimorphism of $\mathbf{PSh}(X; \mathcal{C})$. ∎

**Proposition 4.2.39 (Image Presheaf may be constructed pointwise)**
*If $\mathcal{C}$ has regular epi-mono factorisations for every morphism, e.g.* **Set** *(2.6.175) or an Abelian category (2.7.7), then $\mathbf{PSh}(X; \mathcal{C})$ has regular epi-mono factorisations for every morphism.*

*More precisely, let $\alpha : \mathcal{F} \to \mathcal{G}$ be a morphism of presheaves with a family of pointwise regular epi-mono factorisations $\{\alpha_U = m_U \circ e_U\}_{U \subset X}$*




*Then the assignment $\operatorname{Im}(\alpha)(U) := \operatorname{Im}(\alpha_U)$ can be made into a presheaf uniquely such that $m_\alpha : \operatorname{Im}(\alpha) \to \mathcal{G}$, $(m_\alpha)_U := m_U$, and $e_\alpha : \mathcal{F} \to \operatorname{Im}(\alpha)$, $(e_\alpha)_U := e_U$, are morphisms of presheaves, and $m_\alpha \circ e_\alpha = \alpha$; furthermore this is a regular epi-mono factorisation of $\alpha$ in $\mathbf{PSh}(X; \mathcal{C})$, and hence $(\operatorname{Im}(\alpha), e_\alpha, m_\alpha)$ is an image of $\alpha$ there (2.6.174).*

*If moreover $\mathcal{C}$ is algebraic (2.6.184), the pointwise factorisations may be taken as in (2.6.196), namely $\operatorname{Im}(\alpha_U) := U(\alpha_U)(U(\mathcal{F}(U))) \subseteq U(\mathcal{G}(U))$; then $\operatorname{Im}(\alpha)$ is simply the subpresheaf of $\mathcal{G}$ of sections pointwise in the image of $\alpha$ (4.2.30), with restriction maps inherited from $\mathcal{G}$ and $m_\alpha$ the inclusion.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

**Restriction maps.** Fix $V \subseteq U$. Since $\alpha$ is a morphism of presheaves, $\rho^{\mathcal{G}}_{UV} \circ \alpha_U = \alpha_V \circ \rho^{\mathcal{F}}_{UV}$, so applying (2.6.177) to the regular epi-mono factorisations $\alpha_U = m_U \circ e_U$, $\alpha_V = m_V \circ e_V$ with $a := \rho^{\mathcal{F}}_{UV}$, $b := \rho^{\mathcal{G}}_{UV}$ gives a unique morphism $\rho^{\mathrm{Im}}_{UV} : \operatorname{Im}(\alpha_U) \to \operatorname{Im}(\alpha_V)$ making the following diagram commute

© David Rufino CC BY-NC-ND 4.0




namely

$$\rho^{\mathrm{Im}}_{UV} \circ e_U = e_V \circ \rho^{\mathcal{F}}_{UV} \quad \text{and} \quad m_V \circ \rho^{\mathrm{Im}}_{UV} = \rho^{\mathcal{G}}_{UV} \circ m_U$$

The first identity is precisely the condition for $e_\alpha$ to be a morphism of presheaves, and the second for $m_\alpha$ to be one.

**$\mathrm{Im}(\alpha)$ is a presheaf.** By the functoriality clause of (2.6.177) (taking $a = 1_{\mathcal{F}(U)}$, $b = 1_{\mathcal{G}(U)}$), $\rho^{\mathrm{Im}}_{UU} = 1_{\mathrm{Im}(\alpha_U)}$; and for $W \subseteq V \subseteq U$, taking $a := \rho^{\mathcal{F}}_{UV}$, $b := \rho^{\mathcal{G}}_{UV}$, $c := \rho^{\mathcal{F}}_{VW}$, $d := \rho^{\mathcal{G}}_{VW}$, the composition law gives $\rho^{\mathrm{Im}}_{VW} \circ \rho^{\mathrm{Im}}_{UV} = \rho^{\mathrm{Im}}_{UW}$ (using $\rho^{\mathcal{F}}_{VW} \circ \rho^{\mathcal{F}}_{UV} = \rho^{\mathcal{F}}_{UW}$, and likewise for $\mathcal{G}$). So $\mathrm{Im}(\alpha)$, with these restriction maps, is a presheaf, and $e_\alpha$, $m_\alpha$ are morphisms of presheaves. Uniqueness of restriction maps making $e_\alpha$, $m_\alpha$ into morphisms of presheaves is immediate from the uniqueness clause of (2.6.177), since these two identities are exactly what is required of $e_\alpha, m_\alpha$.

Finally $(m_\alpha \circ e_\alpha)_U = m_U \circ e_U = \alpha_U$ for every $U$, so $m_\alpha \circ e_\alpha = \alpha$.

**Regularity.** By (4.2.38) (using that $\mathcal{C}$ has products and equalisers, and each $e_U$ is regular), $e_\alpha$ is a regular epimorphism of $\mathbf{PSh}(X;\mathcal{C})$. Since $m_U$ is monic in $\mathcal{C}$ for every $U$, $m_\alpha$ is monic in $\mathbf{PSh}(X;\mathcal{C})$ (4.2.28). So $\alpha = m_\alpha \circ e_\alpha$ is a regular epi-mono factorisation of $\alpha$ in $\mathbf{PSh}(X;\mathcal{C})$ (2.6.173), and hence also an image of $\alpha$ there (2.6.174).

For the first statement, if $\mathcal{C}$ has regular epi-mono factorisations for every morphism then every $\alpha_U : \mathcal{F}(U) \to \mathcal{G}(U)$ has one, giving such a family for every morphism $\alpha$ of presheaves, so the above applies to every such $\alpha$ and $\mathbf{PSh}(X;\mathcal{C})$ has regular epi-mono factorisations for every morphism.

**The algebraic case.** If $\mathcal{C}$ is algebraic, taking $\mathrm{Im}(\alpha_U) := U(\alpha_U)(U(\mathcal{F}(U)))$ as in (2.6.196), the restriction maps constructed above may be identified with those inherited from $\mathcal{G}$: since $\alpha$ is a morphism of presheaves, for $\sigma \in \mathcal{F}(U)$, $\rho^{\mathcal{G}}_{UV}(\alpha_U(\sigma)) = \alpha_V\big(\rho^{\mathcal{F}}_{UV}(\sigma)\big) \in \mathrm{Im}(\alpha_V)$, so $\rho^{\mathcal{G}}_{UV}(\mathrm{Im}(\alpha_U)) \subseteq \mathrm{Im}(\alpha_V)$; by the uniqueness clause of (2.6.177), this restriction of $\rho^{\mathcal{G}}_{UV}$ is exactly $\rho^{\mathrm{Im}}_{UV}$. So in this case $\mathrm{Im}(\alpha)$ is simply the subpresheaf of $\mathcal{G}$ of sections pointwise in the image of $\alpha$ (4.2.30), with $e_\alpha$ the (pointwise surjective) corestriction of $\alpha$ and $m_\alpha$ the inclusion — without needing (2.6.177) to construct the restriction maps in the first place. ∎

**Proposition 4.2.40 (Image Sheaf)**
*Let $\mathcal{C}$ be algebraic (2.6.184), hence have regular epi-mono factorisations for every morphism (2.6.196). Then $\mathbf{Sh}(X;\mathcal{C})$ has regular epi-mono factorisations for every morphism.*

*More precisely, let $\alpha : \mathcal{F} \to \mathcal{G}$ be a morphism of sheaves, and define*

$$\mathrm{Im}(\alpha)(U) := \left\{ \tau \in U(\mathcal{G}(U)) \;\middle|\; U = \bigcup_{i\in I} U_i \text{ for some open cover, with } \tau|_{U_i} \in U\big(\alpha_{U_i}\big)(U(\mathcal{F}(U_i))) \text{ for every } i \right\}$$

*the subsheaf of $\mathcal{G}$ of sections locally in the (pointwise) image of $\alpha$ (4.2.32). Then*

$$e : \mathcal{F} \longrightarrow \mathrm{Im}(\alpha), e_U(\sigma) := \alpha_U(\sigma) \quad \text{and} \quad m : \mathrm{Im}(\alpha) \hookrightarrow \mathcal{G} \text{ the inclusion}$$

*are morphisms of sheaves with $m \circ e = \alpha$, and this is a regular epi-mono factorisation of $\alpha$ in $\mathbf{Sh}(X;\mathcal{C})$; hence $(\mathrm{Im}(\alpha), e, m)$ is an image of $\alpha$ there (2.6.174).*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

**Setup.** Let $\big(\mathrm{Im}^{\mathrm{psh}}(\alpha), e_\alpha, m_\alpha\big)$ be the image of $\alpha$ in $\mathbf{PSh}(X;\mathcal{C})$, so $\alpha = m_\alpha \circ e_\alpha$ is a regular epi-mono factorisation there (4.2.39), with $\mathrm{Im}^{\mathrm{psh}}(\alpha)(U) = \mathrm{Im}(\alpha_U) \subseteq \mathcal{G}(U)$ and $(m_\alpha)_U$ the inclusion. In particular $m_\alpha$ is a monomorphism of presheaves (4.2.28), and "locally in the image of $m_\alpha$" is the same condition as "locally in the image of $\alpha$", so applying

© David Rufino CC BY-NC-ND 4.0

(4.2.32) to $m_\alpha$: $\mathrm{Im}(\alpha)$ (as defined above) is a subsheaf of $\mathcal{G}$, and the corestriction $\iota' : \mathrm{Im}^{\mathrm{psh}}(\alpha) \to \mathrm{Im}(\alpha)$ of $m_\alpha$ is a sheafification of $\mathrm{Im}^{\mathrm{psh}}(\alpha)$. Since $\iota'$ is a corestriction of the injective map $(m_\alpha)_U$ for every $U$, $\iota'$ is itself monic (4.2.28).

**Factorisation.** For $\sigma \in \mathcal{F}(U)$, $\iota'_U\big((e_\alpha)_U(\sigma)\big) = (m_\alpha)_U\big((e_\alpha)_U(\sigma)\big) = \alpha_U(\sigma)$, so $e = \iota' \circ e_\alpha$; as $\mathcal{F}$ and $\mathrm{Im}(\alpha)$ are sheaves and $\iota', e_\alpha$ are morphisms of presheaves, $e$ is a morphism of sheaves. Writing $j : \mathrm{Im}(\alpha) \hookrightarrow \mathcal{G}$ for the inclusion, $j \circ \iota' = m_\alpha$, so

$$m \circ e = j \circ \iota' \circ e_\alpha = m_\alpha \circ e_\alpha = \alpha$$

**Monicity.** $m$ is the inclusion of the subsheaf $\mathrm{Im}(\alpha)$ into $\mathcal{G}$, hence monic by the remark following (4.2.30).

**Regularity.** By (4.2.27), $e_\alpha$ has a kernel pair $\big(\mathcal{F} \times_{\mathrm{Im}^{\mathrm{psh}}(\alpha)} \mathcal{F}, p_1, p_2\big)$ in $\mathbf{PSh}(X;\mathcal{C})$, with $U$-components $\big\{(\sigma,\tau) \in \mathcal{F}(U)^2 \;\big|\; (e_\alpha)_U(\sigma) = (e_\alpha)_U(\tau)\big\}$; as each $(e_\alpha)_U = e_U$ is regular (4.2.38) exhibits $e_\alpha$ as its coequaliser there.

As $\iota'$ is monic, $(e_\alpha)_U(\sigma) = (e_\alpha)_U(\tau)$ if and only if $\iota'_U\big((e_\alpha)_U(\sigma)\big) = \iota'_U\big((e_\alpha)_U(\tau)\big)$, that is $e_U(\sigma) = e_U(\tau)$, so the $U$-components of the kernel pairs of $e_\alpha$ and of $e$ agree, with the same projections $p_1, p_2$ into $\mathcal{F}$; by the uniqueness clause of (4.2.27), the two kernel-pair presheaves coincide. Since $\mathcal{F}$ and $\mathrm{Im}(\alpha)$ are both sheaves, applying (4.2.27) to $e$ directly, in its sheaf case, shows this common kernel pair is itself a sheaf, with $p_1, p_2$ morphisms of sheaves.

It remains to show $e$ is a coequaliser of $(p_1, p_2)$ in $\mathbf{Sh}(X;\mathcal{C})$. Let $q' : \mathcal{F} \to \mathcal{K}$ be a morphism of sheaves with $q' \circ p_1 = q' \circ p_2$. As $e_\alpha$ is a coequaliser of $(p_1, p_2)$ in $\mathbf{PSh}(X;\mathcal{C})$, there is a unique morphism of presheaves $\tilde{q} : \mathrm{Im}^{\mathrm{psh}}(\alpha) \to \mathcal{K}$ with $\tilde{q} \circ e_\alpha = q'$. Since $\mathcal{K}$ is a sheaf, the bijection $(-)^\sharp : \mathrm{Mor}(\mathrm{Im}(\alpha), \mathcal{K}) \to \mathrm{Mor}\big(\mathrm{Im}^{\mathrm{psh}}(\alpha), \mathcal{K}\big)$, $\phi \mapsto \phi \circ \iota'$, from the universal property of $\iota'$ (4.2.32) gives a unique $\bar{q} : \mathrm{Im}(\alpha) \to \mathcal{K}$ with $\bar{q} \circ \iota' = \tilde{q}$. Then

$$\bar{q} \circ e = \bar{q} \circ \iota' \circ e_\alpha = \tilde{q} \circ e_\alpha = q'$$

For uniqueness, if $\bar{q}' : \mathrm{Im}(\alpha) \to \mathcal{K}$ also satisfies $\bar{q}' \circ e = q'$, then $(\bar{q}' \circ \iota') \circ e_\alpha = \bar{q}' \circ e = q' = \tilde{q} \circ e_\alpha$, and as $e_\alpha$ is epic (2.6.135), $\bar{q}' \circ \iota' = \tilde{q}$; by uniqueness in the bijection $(-)^\sharp$, $\bar{q}' = \bar{q}$.

So $(p_1, p_2) : \mathcal{F} \times_{\mathrm{Im}^{\mathrm{psh}}(\alpha)} \mathcal{F} \rightrightarrows \mathcal{F}$ are morphisms of $\mathbf{Sh}(X;\mathcal{C})$ with coequaliser $e$ there. In particular $e$ is a regular epimorphism of $\mathbf{Sh}(X;\mathcal{C})$, so $\alpha = m \circ e$ is a regular epi-mono factorisation of $\alpha$ in $\mathbf{Sh}(X;\mathcal{C})$ (2.6.173), and hence $(\mathrm{Im}(\alpha), e, m)$ is an image of $\alpha$ there (2.6.174). ∎

**Remark 4.2.41**
*As established in the **Setup** above, $\iota' : \mathrm{Im}^{\mathrm{psh}}(\alpha) \to \mathrm{Im}(\alpha)$ is a sheafification of the pointwise image presheaf $\mathrm{Im}^{\mathrm{psh}}(\alpha)$ (4.2.39): this recovers the abstract description of the image sheaf, promised at the start of this subsection, as obtained by sheafifying the image presheaf, from the concrete construction above.*

**Corollary 4.2.42 (Stalk Criterion for Surjectivity)**
*Let $\mathcal{C}$ be algebraic (2.6.184), $\alpha : \mathcal{F} \to \mathcal{G}$ a morphism of sheaves, and $\alpha = m \circ e$ a regular epi-mono factorisation of $\alpha$ in $\mathbf{Sh}(X;\mathcal{C})$ (4.2.40). Then $m$ is an isomorphism if and only if $\alpha_x$ is surjective for every $x \in X$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

**The canonical factorisation.** Write $\alpha = m_0 \circ e_0$ for the canonical factorisation of (4.2.40), where $e_0 = \eta \circ e_\alpha$, with $\big(\mathrm{Im}^{\mathrm{psh}}(\alpha), e_\alpha, m_\alpha\big)$ the image of $\alpha$ in $\mathbf{PSh}(X;\mathcal{C})$, built from pointwise regular epi-mono factorisations $\{\alpha_U = m_U \circ e_U\}_{U \subset X}$ (4.2.39), and $\eta : \mathrm{Im}^{\mathrm{psh}}(\alpha) \to \mathrm{Im}(\alpha) =: \big(\mathrm{Im}^{\mathrm{psh}}(\alpha)\big)^+$ the corestriction of $m_\alpha$, a sheafification of $\mathrm{Im}^{\mathrm{psh}}(\alpha)$ (4.2.32), i.e. the unit of the sheafification as defined in (4.2.13).

Fix $x \in X$. As $\mathcal{C}$ is algebraic and $e_U$ is a regular epimorphism of $\mathcal{C}$ for every $U$, (2.6.195) shows $e_U$ is surjective for every $U$. Given $\xi \in \mathrm{Im}^{\mathrm{psh}}(\alpha)_x$, by (4.2.6) $\xi = \iota_x$ for some $\iota \in \mathrm{Im}^{\mathrm{psh}}(\alpha)(U) = \mathrm{Im}(\alpha_U)$, $U \ni x$; as $e_U$ is surjective, $\iota = e_U(\sigma)$ for some $\sigma \in \mathcal{F}(U)$, so $\xi = \iota_x = (e_U(\sigma))_x = (e_\alpha)_x(\sigma_x)$. So $(e_\alpha)_x$ is surjective.

By (4.2.18), $\eta_x$ is an isomorphism, in particular surjective. Hence $(e_0)_x = \eta_x \circ (e_\alpha)_x$, being a composite of surjections, is surjective.

Since $\alpha_x = (m_0)_x \circ (e_0)_x$ (4.2.7) with $(e_0)_x$ surjective, $\alpha_x$ is surjective if and only if $(m_0)_x$ is surjective: if $(m_0)_x$ is surjective this is immediate, as a composite of surjections; conversely if $\alpha_x$ is surjective then $\mathrm{Im}\big((m_0)_x\big) \supseteq \mathrm{Im}(\alpha_x) = \mathcal{G}_x$. So by (4.2.29), $m_0$ is an isomorphism if and only if $\alpha_x$ is surjective for every $x \in X$.

**General regular epi-mono factorisations.** Let $\alpha = m \circ e$ be any regular epi-mono factorisation of $\alpha$ in $\mathbf{Sh}(X;\mathcal{C})$. By (2.6.178), applied to this and the canonical factorisation $\alpha = m_0 \circ e_0$, there is an isomorphism $\phi$ with $m_0 \circ \phi = m$. So $m$ is an isomorphism if and only if $m_0$ is (composing with the isomorphism $\phi$ on the right preserves being an isomorphism, in either direction), which by the above holds if and only if $\alpha_x$ is surjective for every $x \in X$. ∎

© David Rufino CC BY-NC-ND 4.0

### 4.2.11 Exactness

**Definition 4.2.43 (Exact Sequence of Sheaves)**
*Let $\mathcal{C}$ be pointed and algebraic; then $\mathbf{Sh}(X;\mathcal{C})$ is evidently pointed (zero morphisms computed pointwise, the constant zero presheaf being a zero object), with a regular epi-mono factorisation (4.2.40) for every morphism, so (2.6.200) applies with $\mathbf{Sh}(X;\mathcal{C})$ in place of $\mathcal{C}$: a pair of composable morphisms of sheaves $\mathcal{F} \xrightarrow{\alpha} \mathcal{G} \xrightarrow{\beta} \mathcal{H}$ is **exact at $\mathcal{G}$** if $\mathrm{Im}(\alpha) = \ker(\beta)$, that is $(\mathrm{Im}(\alpha), m)$ is (literally) a kernel of $\beta$, for any (equivalently, by the same argument, every) choice of image factorisation $\alpha = m \circ e$ — in particular we may as well take the canonical construction of (4.2.40).*

**Proposition 4.2.44 (Concrete Criterion for Exactness of Sheaves)**
*Let $\mathcal{C}$ be pointed and algebraic and $\mathcal{F} \xrightarrow{\alpha} \mathcal{G} \xrightarrow{\beta} \mathcal{H}$ a pair of composable morphisms of sheaves. Regard $\mathrm{Im}(\alpha)$ (4.2.40) and $\ker(\beta)$ (4.2.25) as the canonical constructions, both literal subsheaves of $\mathcal{G}$ since $\mathcal{C}$ is algebraic. Then the following are equivalent*

*a) the pair is exact at $\mathcal{G}$ (4.2.43) (for any choice of image factorisation of $\alpha$)*
*b) $\mathrm{Im}(\alpha)(U) = \ker(\beta)(U)$, as subsets of $\mathcal{G}(U)$, for every open $U \subset X$*
*c) $\beta \circ \alpha = 0$, and every section of $\ker(\beta)(U)$ is locally in the image of $\alpha$, for every open $U \subset X$*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Write $\alpha = m \circ e$ for the canonical image factorisation (4.2.40) and $i : \ker(\beta) \hookrightarrow \mathcal{G}$ for the canonical kernel (4.2.25).

*a) $\iff$ b)* **Reduction to the canonical representatives.** By (4.2.43) (applied with $\mathbf{Sh}(X;\mathcal{C})$ in place of $\mathcal{C}$), exactness at $\mathcal{G}$ does not depend on the choice of image factorisation of $\alpha$, so it suffices to treat the canonical representative $m$: the pair is exact at $\mathcal{G}$ if and only if $(\mathrm{Im}(\alpha), m)$ is a kernel of $\beta$. Since $(\ker(\beta), i)$ is also a kernel of $\beta$, by (2.6.80) this holds if and only if there is an isomorphism $\theta : \mathrm{Im}(\alpha) \to \ker(\beta)$ with $i \circ \theta = m$.

($\Longleftarrow$) If $\mathrm{Im}(\alpha)(U) = \ker(\beta)(U)$ for every $U$, then $m = i$ as morphisms of sheaves, both being the inclusion of this common subsheaf into $\mathcal{G}$, with the (unique, (2.6.193)) induced structure making it a morphism. So $\theta := 1_{\mathrm{Im}(\alpha)}$ satisfies $i \circ \theta = i = m$.

($\Longrightarrow$) Suppose there is an isomorphism $\theta : \mathrm{Im}(\alpha) \to \ker(\beta)$ with $i \circ \theta = m$. Fix $U \subset X$ and $\sigma \in \mathrm{Im}(\alpha)(U)$; since $m_U$ is the inclusion of $\mathrm{Im}(\alpha)(U)$ into $\mathcal{G}(U)$, $i_U(\theta_U(\sigma)) = m_U(\sigma) = \sigma$, and $i_U$ is also an inclusion, so $\theta_U(\sigma) = \sigma$; hence $\sigma \in \ker(\beta)(U)$, giving $\mathrm{Im}(\alpha)(U) \subseteq \ker(\beta)(U)$. As $\theta$ is an isomorphism, $\theta_U$ is a bijection onto $\ker(\beta)(U)$, so every element of $\ker(\beta)(U)$ equals $\theta_U(\sigma) = \sigma$ for some $\sigma \in \mathrm{Im}(\alpha)(U)$, giving $\ker(\beta)(U) \subseteq \mathrm{Im}(\alpha)(U)$. So $\mathrm{Im}(\alpha)(U) = \ker(\beta)(U)$ for every $U$.

*b) $\iff$ c)* Condition *b)* is, for every $U$, the conjunction of $\mathrm{Im}(\alpha)(U) \subseteq \ker(\beta)(U)$ and $\ker(\beta)(U) \subseteq \mathrm{Im}(\alpha)(U)$; the latter is precisely "every section of $\ker(\beta)(U)$ is locally in the image of $\alpha$" (4.2.40), so it remains to show $\mathrm{Im}(\alpha)(U) \subseteq \ker(\beta)(U)$ for every $U$ is equivalent to $\beta \circ \alpha = 0$.

($\Longleftarrow$) Suppose $\beta \circ \alpha = 0$. For $\tau \in \mathrm{Im}(\alpha)(U)$, take a witnessing cover $U = \bigcup_j U_j$ with $\tau|_{U_j} = \alpha_{U_j}(\sigma_j)$ for some $\sigma_j \in \mathcal{F}(U_j)$; then $\beta_U(\tau)|_{U_j} = \beta_{U_j}(\tau|_{U_j}) = \beta_{U_j}(\alpha_{U_j}(\sigma_j)) = (\beta \circ \alpha)_{U_j}(\sigma_j) = 0$ for every $j$, so by separation of $\mathcal{H}$ (4.2.4), $\beta_U(\tau) = 0$, i.e. $\tau \in \ker(\beta)(U)$.

($\Longrightarrow$) Suppose $\mathrm{Im}(\alpha)(U) \subseteq \ker(\beta)(U)$ for every $U$. For any $U$ and $\sigma \in \mathcal{F}(U)$, $\alpha_U(\sigma) \in \mathrm{Im}(\alpha)(U)$ (witnessed by the trivial cover $\{U\}$), so $\alpha_U(\sigma) \in \ker(\beta)(U)$, that is $\beta_U(\alpha_U(\sigma)) = 0$. As $U, \sigma$ were arbitrary, $\beta \circ \alpha = 0$. ∎

**Lemma 4.2.45 (Subsheaves are Determined by Stalks)**
*Let $\mathcal{C}$ be algebraic and $\mathcal{G}_1, \mathcal{G}_2$ be subsheaves of a sheaf $\mathcal{G}$ (4.2.30). Then $\mathcal{G}_1(U) = \mathcal{G}_2(U)$, as subsets of $\mathcal{G}(U)$, for every open $U \subset X$, if and only if $(\mathcal{G}_1)_x = (\mathcal{G}_2)_x$, as subsets of $\mathcal{G}_x$ (via the canonical, injective (4.2.28) stalk maps of the inclusions), for every $x \in X$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

($\Longrightarrow$) Immediate: if $\mathcal{G}_1(U) = \mathcal{G}_2(U)$ for every $U$, then by (4.2.6) every element of $(\mathcal{G}_1)_x$ is $\sigma_x$ for some $\sigma \in \mathcal{G}_1(U) = \mathcal{G}_2(U)$, $U \ni x$, hence lies in $(\mathcal{G}_2)_x$, and symmetrically.

($\Longleftarrow$) Suppose $(\mathcal{G}_1)_x = (\mathcal{G}_2)_x$ for every $x \in X$. Fix $U \subset X$ and $\sigma \in \mathcal{G}_1(U) \subseteq \mathcal{G}(U)$; we show $\sigma \in \mathcal{G}_2(U)$ (the reverse inclusion, and hence $\mathcal{G}_1(U) = \mathcal{G}_2(U)$, then follows by symmetry).

For $x \in U$, $\sigma_x \in (\mathcal{G}_1)_x = (\mathcal{G}_2)_x$, so by (4.2.6) $\sigma_x = \tau_x$ for some $\tau \in \mathcal{G}_2(V_x) \subseteq \mathcal{G}(V_x)$, $x \in V_x$; by (4.2.8).a) (applied in $\mathcal{G}$, as $\sigma, \tau$ agree at the stalk $x$) there is $W_x \subseteq U \cap V_x$, $x \in W_x$, with $\sigma|_{W_x} = \tau|_{W_x}$ in $\mathcal{G}(W_x)$; as $\mathcal{G}_2$ is a subpresheaf, $\tau|_{W_x} \in \mathcal{G}_2(W_x)$, so $\sigma|_{W_x} \in \mathcal{G}_2(W_x)$.

© David Rufino CC BY-NC-ND 4.0

So $\{W_x\}_{x\in U}$ is an open cover of $U$ with $\sigma|_{W_x}\in\mathcal{G}_2(W_x)$ for every $x$; these restrictions agree pairwise on overlaps (all restricting from the common $\sigma$), so as $\mathcal{G}_2$ is a sheaf, gluing gives $\rho\in\mathcal{G}_2(U)$ with $\rho|_{W_x}=\sigma|_{W_x}$ for every $x$. Regarding $\rho$ as an element of $\mathcal{G}(U)$ (via the inclusion $\mathcal{G}_2\hookrightarrow\mathcal{G}$), $\rho$ and $\sigma$ agree on the cover $\{W_x\}$ of $U$, so by separation of $\mathcal{G}$ (4.2.4), $\rho=\sigma$; hence $\sigma\in\mathcal{G}_2(U)$. ∎

**Lemma 4.2.46 (Kernels and Stalks Commute)**
*Let $\mathcal{C}$ be pointed and algebraic, $\beta:\mathcal{G}\to\mathcal{H}$ a morphism of sheaves, and $\ker(\beta)(U)=\{\sigma\in\mathcal{G}(U)\mid\beta_U(\sigma)=0\}$ its kernel (4.2.25), regarded as a subsheaf of $\mathcal{G}$. Then for every $x\in X$, the canonical stalk inclusion identifies $\ker(\beta)_x$ with the subset $\ker(\beta_x):=\{\xi\in\mathcal{G}_x\mid\beta_x(\xi)=0\}$ of $\mathcal{G}_x$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Write $i:\ker(\beta)\hookrightarrow\mathcal{G}$ for the inclusion. By (2.6.150) and (2.6.135), $i$ is monic, so by (4.2.28).d) (as $\mathcal{C}$ is algebraic and $\ker(\beta),\mathcal{G}$ are sheaves), $i_x$ is injective for every $x\in X$.

Since $\beta\circ i=0$ (2.6.150), stalk-functoriality (4.2.7) gives $\beta_x\circ i_x=(\beta\circ i)_x=0_x=0$ (the stalk of the zero presheaf, valued 0 at every open with identity restrictions, is 0, so the stalk functor sends the zero morphism to the zero morphism). So $i_x(\eta)\in\ker(\beta_x)$ for every $\eta\in\ker(\beta)_x$.

Conversely let $\xi\in\ker(\beta_x)$, i.e. $\xi\in\mathcal{G}_x$ with $\beta_x(\xi)=0$. By (4.2.6), $\xi=\sigma_x$ for some $\sigma\in\mathcal{G}(U)$, $U\in\mathcal{B}$, $x\in U$. Then $(\beta_U(\sigma))_x=\beta_x(\sigma_x)=\beta_x(\xi)=0=0_x$, so by (4.2.8).a) there is $W\in\mathcal{B}$, $W\subseteq U$, $x\in W$, with $\beta_U(\sigma)|_W=0|_W$, that is $\beta_W(\sigma|_W)=0$, so $\sigma|_W\in\ker(\beta)(W)$. Then $i_x\big((\sigma|_W)_x\big)=(i_W(\sigma|_W))_x=(\sigma|_W)_x=\sigma_x=\xi$, so $\xi\in\mathrm{Im}(i_x)$.

So $i_x$ is injective with image exactly $\ker(\beta_x)$, that is, identifies $\ker(\beta)_x$ with this subset of $\mathcal{G}_x$. ∎

**Lemma 4.2.47 (Images and Stalks Commute)**
*Let $\mathcal{C}$ be algebraic, $\alpha:\mathcal{F}\to\mathcal{G}$ a morphism of sheaves, and $\alpha=m\circ e$ its image factorisation (4.2.40). Then for every $x\in X$, the canonical stalk inclusion identifies $\mathrm{Im}(\alpha)_x$ with the subset $\mathrm{Im}(\alpha_x):=\alpha_x(\mathcal{F}_x)$ of $\mathcal{G}_x$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Retain the notation from the proof of (4.2.42): writing $e=\eta\circ e_\alpha$ with $\big(\mathrm{Im}^{\mathrm{psh}}(\alpha),e_\alpha,m_\alpha\big)$ the image of $\alpha$ in $\mathbf{PSh}(X;\mathcal{C})$ (4.2.39) and $\eta$ the sheafification unit exhibiting $\mathrm{Im}(\alpha)=\big(\mathrm{Im}^{\mathrm{psh}}(\alpha)\big)^+$, it is shown there that $(e_\alpha)_x$ is surjective (each $e_U$ being regular, hence surjective, (2.6.195)) and $\eta_x$ is an isomorphism (4.2.18), so $e_x=\eta_x\circ(e_\alpha)_x$ is surjective. Also $m_x$ is injective by (4.2.28).d), as $m$ is a monomorphism of sheaves.

Since $m_x\circ e_x=\alpha_x$ and $e_x$ is surjective onto $\mathrm{Im}(\alpha)_x$,

$$m_x(\mathrm{Im}(\alpha)_x)=m_x(e_x(\mathcal{F}_x))=\alpha_x(\mathcal{F}_x)=\mathrm{Im}(\alpha_x)$$

so, $m_x$ being injective, it identifies $\mathrm{Im}(\alpha)_x$ with the subset $\mathrm{Im}(\alpha_x)$ of $\mathcal{G}_x$. ∎

**Proposition 4.2.48 (Stalk Criterion for Exactness)**
*Let $\mathcal{C}$ be pointed and algebraic and $\mathcal{F}\xrightarrow{\alpha}\mathcal{G}\xrightarrow{\beta}\mathcal{H}$ a pair of composable morphisms of sheaves. Then this pair is exact at $\mathcal{G}$ (4.2.43) if and only if, for every $x\in X$, $\mathrm{Im}(\alpha_x)=\ker(\beta_x)$, as subsets of $\mathcal{G}_x$, that is, the stalk pair $\mathcal{F}_x\xrightarrow{\alpha_x}\mathcal{G}_x\xrightarrow{\beta_x}\mathcal{H}_x$ is exact at $\mathcal{G}_x$ in $\mathcal{C}$ (in the sense of (4.2.43)), applied verbatim in $\mathcal{C}$).*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

By (4.2.44), the pair is exact at $\mathcal{G}$ if and only if $\mathrm{Im}(\alpha)(U)=\ker(\beta)(U)$ for every $U\subset X$ (using the canonical constructions of $\mathrm{Im}(\alpha),\ker(\beta)$). By (4.2.45) applied to these subsheaves of $\mathcal{G}$, this holds if and only if $\mathrm{Im}(\alpha)_x=\ker(\beta)_x$, as subsets of $\mathcal{G}_x$, for every $x\in X$. By (4.2.47) and (4.2.46), this is exactly the condition $\mathrm{Im}(\alpha_x)=\ker(\beta_x)$, as subsets of $\mathcal{G}_x$, for every $x\in X$. ∎

## 4.2.12 Constant Sheaves

**Definition 4.2.49**
*Let $A\in\mathcal{C}$ and $X$ a topological space. Define the **constant presheaf** as follows*

$$\mathcal{A}^{\mathrm{pre}}(U):=A$$

*with restriction maps the identity.*

**Proposition 4.2.50**
*Let $X$ be an irreducible topological space then the constant presheaf $\mathcal{A}^{\mathrm{pre}}(U)=A$ is a sheaf, which we denote by $\underline{A}$.*

© David Rufino CC BY-NC-ND 4.0

*Proof.* Suppose that $U = \bigcup_{i \in I} U_i$ where $U_i$ are non-empty, then as $U$ is irreducible (4.1.70), by definition every $U_i \cap U_j \neq \varnothing$. Therefore for sections on $U_i$ are only compatible if they are equal, and so there is an obvious extension to $U$. ∎

**Proposition 4.2.51 (Quotient of Constant Sheaf)**
*Let $\mathcal{F}$ be a subsheaf of the constant sheaf $\underline{A}$, define the quotient presheaf as follows*

$$(\underline{A}/\mathcal{F})(U) \coloneqq \left\{ \sigma : U \to \coprod_{x \in U} A/\mathcal{F}_x \;\middle|\; \sigma(x) \in A/\mathcal{F}_x \right\}$$

© David Rufino CC BY-NC-ND 4.0

## 4.3 Space with Functions

**Definition 4.3.1 (Space with functions)**
*Let $X$ be a topological space and $\mathcal{O}_X$ be a subsheaf of the $\overline{k}$-valued functions on $X$ (which is a sheaf of $k$-algebras). Then we say $(X, \mathcal{O}_X)$ is a **space with functions**. Additionally it's **local** if it satisfies the condition*

*a) For every $f \in \mathcal{O}_X(U)$ we have $D(f) := \{f \neq 0\}$ is open and $\frac{1}{f} \in \mathcal{O}_X(D(f))$.*

*Note that the sheaf restriction maps $\rho_{UV} : \mathcal{O}_X(U) \to \mathcal{O}_X(V)$ correspond simply to restrictions of functions.*

*In the latter case $X$ is a locally ringed space over $k$ because $\mathcal{O}_{X,x}$ is a local $k$-algebra with maximal ideal $\mathfrak{m}_{X,x}$ consisting of stalks vanishing at $x$.*

*A **regular morphism** or **regular map** $(X, \mathcal{O}_X) \to (Y, \mathcal{O}_Y)$ is a continuous function $\phi : X \to Y$ such that*

$$g \in \mathcal{O}_Y(U) \implies g \circ \phi \in \mathcal{O}_X(\phi^{-1}(U))$$

*This determines a morphism of sheaves*

$$\phi^\sharp : \mathcal{O}_Y \longrightarrow \phi_\star \mathcal{O}_X$$

*We say a regular morphism is **dominant** if $\phi(X)$ is dense in $Y$.*

*An isomorphism is a homeomorphism and induces an isomorphism of sheaves, or equivalently has a two-sided regular inverse.*

**Proposition 4.3.2**
*The composition of regular maps is well-defined and associative.*

### 4.3.1 Local Rings

**Definition 4.3.3 (Local Ring)**
*Let $X$ be a local space with functions and $W \subset X$ an irreducible closed subset. Then we define the **local ring** at $W$ to be (see (4.2.5))*

$$\mathcal{O}_{X,W} := \varinjlim_{U \cap W \neq \varnothing} \mathcal{O}_X(U)$$

*If $\overline{W} = \overline{W'}$ then $\mathcal{O}_{X,W} = \mathcal{O}_{X,W'}$, so for example $\mathcal{O}_{X,W} = \mathcal{O}_{X,\overline{W}}$.*

*It is a local ring with unique maximal ideal*

$$\mathfrak{m}_{X,W} := \{[(V, \sigma)] \mid V \cap W \neq \varnothing, \sigma \in \mathcal{O}_X(V), \sigma(x) = 0 \, \forall x \in V \cap W\}$$

*Define the **residue field***

$$k(\mathfrak{m}_{X,W}) := \mathcal{O}_{X,W}/\mathfrak{m}_{X,W}$$

*In the case $W = \{x\}$ we write $\mathcal{O}_{X,x}$, $\mathfrak{m}_{X,x}$ and $k(\mathfrak{m}_{X,x})$.*

*Proof.* Consider $(V, \sigma) \in \mathcal{O}_{X,W} \setminus \mathfrak{m}_{X,W}$. Then by definition $D(\sigma) \cap W \neq \varnothing$ and $(D(\sigma), \sigma^{-1})$ is an explicit inverse. ■

**Corollary 4.3.4 (Field of Rational Functions)**
*Let $X$ be an irreducible space with functions. Then $\mathcal{O}_{X,X} =: k(X)$ is a field, which we call the **field of rational functions**.*

**Proposition 4.3.5 (Stalk Maps)**
*Suppose $\phi : X \to Y$ is a regular morphism and $Z \subset \overline{\phi(W)}$. Then there exists a unique (local) $k$-algebra homomorphism $\phi_W : \mathcal{O}_{Y,Z} \to \mathcal{O}_{X,W}$ such that the following diagram commutes*

$$\begin{array}{ccc} \mathcal{O}_Y(V) & \xrightarrow{\rho \circ \phi_V^\sharp} & \mathcal{O}_X(U) \\ \downarrow & & \downarrow \\ \mathcal{O}_{Y,Z} & \dashrightarrow & \mathcal{O}_{X,W} \end{array}$$

*for all open sets $V \subset Y$ for which $V \cap Z \neq \varnothing$ and some $U \subset \phi^{-1}(V)$ for which $U \cap W \neq \varnothing$. This is functorial in the sense that*

© David Rufino CC BY-NC-ND 4.0

$$(\psi \circ \phi)_W = \psi_Z \circ \phi_W$$

*Proof.* If $V$ meets $Z$ then by (4.1.19) $\phi^{-1}(V)$ meets $W$. Therefore the vertical maps are well-defined. We define

$$\phi_W((V', \sigma)) := (\phi^{-1}(V'), \sigma \circ \phi)$$

and immediately verify it is well-defined and ensures the diagram commutes. ∎

**Proposition 4.3.6 (Stalk Maps II)**
*Let $\phi : X \to Y$ be a regular morphism of local spaces with functions. Suppose $y \in \overline{\{\phi(x)\}}$. Then there is a unique local algebra homomorphism $\phi_x : \mathcal{O}_{Y,\phi(x)} \to \mathcal{O}_{X,x}$ making the following diagram commute*

$$\begin{array}{ccc} \mathcal{O}_Y(V) & \xrightarrow{\phi^\sharp_V} & \mathcal{O}_X(U) \\ \downarrow & & \downarrow \\ \mathcal{O}_{Y,\phi(x)} & \dashrightarrow & \mathcal{O}_{X,x} \end{array}$$

*for all open neighbourhoods $V$ of $\phi(x)$ and some open neighbourhood $U \subset \phi^{-1}(V)$ of $x$. This is functorial in the sense that*

$$(\psi \circ \phi)_x = \psi_{\phi(x)} \circ \phi_x$$

*Proof.* This follows immediately from (4.3.5). ∎

**Proposition 4.3.7 (Stalk Maps Function Field)**
*Let $X, Y$ be irreducible local spaces with functions and $\phi : X \to Y$ a dominant regular morphism. Then there is a unique homomorphism $\phi_\star : k(Y) \hookrightarrow k(X)$ making the following diagram commute*

$$\begin{array}{ccc} \mathcal{O}_Y(V) & \xrightarrow{\phi^\sharp_V} & \mathcal{O}_X(U) \\ \downarrow & & \downarrow \\ k(Y) & \underset{\phi_\star}{\hookrightarrow} & k(X) \end{array}$$

*for all non-empty open subsets $V \subset Y$ and some non-empty open subset $U \subset \phi^{-1}(V)$. This is functorial in the sense that*

$$(\phi \circ \psi)_\star = \psi_\star \circ \phi_\star$$

*Proof.* This follows immediately from (4.3.5) with $W = X$ and $T = Y$. ∎

**Proposition 4.3.8**
*Let $X$ be a local space with functions and $Z, W$ be irreducible subsets such that $Z \subset \overline{W}$. Then there is a canonical homomorphism $\mathcal{O}_{X,Z} \to \mathcal{O}_{X,W}$ making the following diagram commute*

$$\begin{array}{ccc} \mathcal{O}_X(U) & & \\ \downarrow & \searrow & \\ \mathcal{O}_{X,Z} & \dashrightarrow & \mathcal{O}_{X,W} \end{array}$$

*for all open subsets $U$ of $X$ meeting $Z$.*

*This holds in particular when $X$ is irreducible we may take $W = X$ and $\mathcal{O}_{X,W} = k(X)$.*

*Proof.* This follows immediately from (4.3.5) by considering the case $\phi = 1_X$. ∎

**Proposition 4.3.9 (Stalk Maps under Specialisation)**
*Let $X, Y$ be irreducible local spaces with functions, $\phi : X \to Y$ a dominant regular morphism and $x \in X$ such that $y \in \overline{\{\phi(x)\}}$. Then for all neighbourhoods $V$ of $\phi(x)$ we have a commutative diagram*

© David Rufino CC BY-NC-ND 4.0

$$\begin{array}{ccc} \mathcal{O}_Y(V) & \xrightarrow{\phi_V^\sharp} & \mathcal{O}_X(\phi^{-1}(V)) \\ \downarrow & & \downarrow \\ \mathcal{O}_{Y,y} & \xrightarrow{\phi_x} & \mathcal{O}_{X,x} \\ \downarrow & & \downarrow \\ k(Y) & \xhookrightarrow{\phi_\star} & k(X) \end{array}$$

**Lemma 4.3.10 (Evaluation of Stalks)**
*Let $X$ be a local space of functions and $x \in X$. Then there is a unique function*

$$\mathrm{ev}_x : k\big(\mathfrak{m}_{X,x}\big) \hookrightarrow \overline{k}$$

*making the following diagram commute for all open neighbourhoods $U$ of $x$*




*This is functorial in the sense that for a regular morphism $\phi : X \to Y$ there is a commutative diagram*

$$\begin{array}{ccc} k\big(\mathfrak{m}_{Y,\phi(x)}\big) & \xrightarrow{\phi_x} & k\big(\mathfrak{m}_{X,x}\big) \\ \downarrow{\scriptstyle \mathrm{ev}_{\phi(x)}} & & \downarrow{\scriptstyle \mathrm{ev}_x} \\ \overline{k} & = & \overline{k} \end{array}$$

## 4.3.2 Open Immersions

**Definition 4.3.11 (Open Immersion)**
*Let $X, Y$ be spaces with functions. Then we say a regular map $\phi : X \to Y$ is an **open immersion** if*

*a) $\phi$ is an open embedding, and*
*b) $\phi$ induces an isomorphism of sheaves*

$$\phi_\star\big(\mathcal{O}_Y|_{\phi(X)}\big) \xrightarrow{\sim} \mathcal{O}_X$$

**Proposition 4.3.12 (Open Subspace)**
*Let $X$ be a (local) space of functions and $U \subset X$ an open subset. Then $(U, \mathcal{O}_X|_U)$ is a (local) space of functions which we call an **open subspace**. We may write $\mathcal{O}_U \coloneqq \mathcal{O}_X|_U$.*

*The inclusion $i : U \hookrightarrow X$ is an open immersion and an open immersion $\phi : X \to Y$ induces an isomorphism between $X$ and the open subvariety $\phi(X)$.*

**Proposition 4.3.13**
*Let $X$ be a space of functions and $V \subset U$ open subsets. Then $(V, \mathcal{O}_U|_V) = (V, \mathcal{O}_X|_V)$.*

**Proposition 4.3.14**
*Let $\phi : X \to Y$ be an open embedding and $Z \subset Y$ closed. Then $Z$ is irreducible iff $\phi^{-1}(Z)$ is irreducible.*

*In particular if $Y$ is irreducible then so is $X$.*

*Proof.* We may reduce to the case $X = U$ is an open subset of $Y$. Then it follows from (4.1.73) and (4.1.70). ∎

**Proposition 4.3.15**
*Let $\phi : X \to Y$ be a regular open embedding. Then the following are equivalent*

*a) $\phi$ is an open immersion*
*b) for every irreducible subset $W \subset X$ the canonical map*

$$\mathcal{O}_{Y,Z} \to \mathcal{O}_{X,W}$$

© David Rufino CC BY-NC-ND 4.0

*is an isomorphism for all $Z$ such that $\overline{Z} = \overline{\phi(W)}$*

*c) b) holds for some $Z$ such that $\overline{Z} = \overline{\phi(W)}$*

*In particular for all open subsets $V \subset Y$ and $U \subset \phi^{-1}(V)$ such that $U \cap W \neq \varnothing$ we have a commutative diagram*

$$\begin{array}{ccc} \mathcal{O}_Y(V) & \xrightarrow{\phi^\sharp} & \mathcal{O}_X(U) \\ \downarrow & & \downarrow \\ \mathcal{O}_{Y,\phi(W)} & \xrightarrow{\sim} & \mathcal{O}_{X,W} \end{array}$$

*Proof.* $b) \implies a)$ By definition we may factor into a regular homeomorphism $X \xrightarrow{\sim} V \subset Y$. It is an isomorphism of spaces by (4.2.11).

$a) \implies b)$ We may reduce to the case that $X \subset Y$ is an open subset. Because an open subset of $X$ intersects $W$ if and only if it intersects $Z$ we find that the map is bijective.

$c) \iff b)$ we simply observe that if $\overline{Z} = \overline{Z'}$ then the canonical maps are the same. ∎

**Proposition 4.3.16**
*Let $\phi : X \to Y$ be a dominant open immersion. If $Y$ is irreducible, then so is $X$ and the map*

$$\phi_\star : k(Y) \hookrightarrow k(X)$$

*is an isomorphism.*

*In particular if $X$ is irreducible and $U \subset X$ is a non-empty open subset, then there is a canonical isomorphism*

$$k(X) \xrightarrow{\sim} k(U)$$

*Proof.* By (4.3.14) $X$ is irreducible, so it is a special case of (4.3.15). ∎

**Proposition 4.3.17 (Stalk Maps under Open Immersions I)**
*Let $X$ be a local space with functions, $U \subset X$ an open subset and $W \subset X$ a irreducible closed subset such that $U \cap W \neq \varnothing$. Then $U \cap W$ is an irreducible subset of $U$ and we have canonical isomorphisms of stalks making the following diagram commute*

$$\begin{array}{ccc} \mathcal{O}_X(V) & \longrightarrow & \mathcal{O}_U(U \cap V) \\ \downarrow & & \downarrow \\ \mathcal{O}_{X,W} & \xrightarrow{\sim} & \mathcal{O}_{U,W\cap U} \end{array}$$

*for all open $V \subset X$ for which $V \cap W \neq \varnothing$.*

*Proof.* The inclusion $i : U \hookrightarrow X$ is an open immersion, so $U \cap W$ is irreducible by (4.1.73). This is then a special case of (4.3.15). ∎

**Lemma 4.3.18**
*Let $U, Z$ be subsets of a topological space $X$, with $U$ open. Then*

$$\overline{Z} \cap U \subset \overline{Z \cap U}$$

*Proof.* Suppose $x \in \overline{Z} \cap U$ and $V$ is an open subset containing $x$. Then $U \cap V$ is an open set containing $x$ and therefore by (4.1.19) $Z \cap U \cap V \neq \varnothing$. As $V$ was an arbitrary neighbourhood of $x$ we conclude that $x \in \overline{Z \cap U}$. ∎

**Proposition 4.3.19 (Restricting Stalk Maps to Open Subspaces)**
*Let $\phi : X \to Y$ be a regular morphism of local space with functions. Suppose $U \subset X$ and $V \subset Y$ are open subsets such that $U \subset \phi^{-1}(V)$, and $W \subset X$, $Z \subset Y$ are irreducible subsets such that $Z \subset \overline{\phi(W)}$ and $U \cap W \neq \varnothing$. Then $\phi(W \cap U) \subset \overline{Z \cap V}$ and there is a commutative cube for every open set $V' \subset V$ such that $V' \cap Z \neq \varnothing$ and $U' \subset \phi^{-1}(V')$ such that $U' \cap W \neq \varnothing$*

© David Rufino CC BY-NC-ND 4.0




*where the stalk isomorphisms are defined in (4.3.17) and $(\phi|_U)_{W\cap U}$ is the unique (local) homomorphism making the bottom square (or right side) commute*

*Proof.* Observe by (4.3.18) that

$$\phi(W \cap U) \subset \phi(W) \cap V \subset \overline{Z} \cap V \subset \overline{Z \cap V}$$

which means the stalk map of $\phi|_U$ is well-defined. The sides of the cube are well-defined and commute by (4.3.17) and (4.3.5), and the top by definition. As the horizontal arrows are isomorphisms then we see that the bottom square commutes, and $(\phi|_U)_{W\cap U}$ is the unique homomorphism for which this is the case. ∎

### 4.3.3 Closed Immersions

**Definition 4.3.20 (Closed Immersion)**
*Let $\phi : X \to Y$ be a regular morphism of spaces with functions. We say that $\phi$ is a **closed immersion** if*

*a) $\phi$ is a homeomorphism onto a closed subset $\phi(X)$ of $Y$*
*b) For all $x \in X$ we have the induced map on stalks*

$$\mathcal{O}_{Y,\phi(x)} \to \mathcal{O}_X$$

*is surjective.*

**Proposition 4.3.21 (Closed Subset Structure Sheaf)**
*Let $(X, \mathcal{O}_X)$ be a space of functions and $Z \subset X$ a closed subset. Then there exists a unique sheaf of functions $\mathcal{O}_Z$ such that the inclusion $j : Z \to X$ is a closed immersion.*

*More precisely for $U \subset Z$ an open subset*

$$\mathcal{O}_Z(U) := \left\{ f : U \to \overline{k} \;\middle|\; f(y) = g_i(y)\ \forall y \in Z \cap U_i \text{ for some } g_i \in \mathcal{O}_X(U_i) \text{ where } Z \cap \bigcup_{i\in I} U_i = U \right\}$$

*Proof.* Evidently $j$ is a homeomorphism onto $Z$. We first show that $j$ is regular. Suppose $g \in \mathcal{O}_X(U')$ then by definition we have $g \circ j \in \mathcal{O}_Z(j^{-1}(U')) = \mathcal{O}_Z(U' \cap Z)$.

Given $(V, f) \in \mathcal{O}_{Z,z}$ by taking a smaller open subset we may suppose that $f(y) = g(y)$ for all $y \in V$ for some $g \in \mathcal{O}_X(U)$ and $U \cap Z = V$. Then the image of $(U, g)$ under the stalk map is $(U \cap Z, g \circ j)$ which is precisely $(V, f)$. Therefore the stalk map $j_z$ is surjective for all $z \in Z$.

Let $\mathcal{F}$ be a sheaf of functions for which $j$ is a (regular) closed immersion and suppose $f \in \mathcal{O}_Z(U)$. Then $f|_{U_i\cap Z}(y) = g_i(y)$ for all $y \in Z \cap U_i$ and $g_i \in \mathcal{O}_X(U_i)$. By assumption $f|_{U_i\cap Z} = g_i \circ j \in \mathcal{F}(U_i \cap Z)$. By the sheaf condition we have $f \in \mathcal{F}(U)$ which shows $\mathcal{O}_Z$ is a subsheaf of $\mathcal{F}$.

We need to show that the inclusion of sheaves $\mathcal{O}_Z \hookrightarrow \mathcal{F}$ is an isomorphism. By (4.2.28) c) $\implies$ d) and (4.2.11) it is enough to show that it is surjective on stalks. However this follows by assumption by observing the composition $\mathcal{O}_{X,j(z)} \to \mathcal{O}_{Z,z} \to \mathcal{F}_z$ is surjective. ∎

**Remark 4.3.22**
*This can be defined formally as $j^{-1}(\mathcal{O}_X/\mathcal{I})$ where $\mathcal{I}$ is the presheaf of regular functions which vanish on $Z$. We show in the Algebraic Geometry chapter's treatment of closed subvarieties that this is consistent with the concept of affine subvariety.*

**Proposition 4.3.23**
*Let $\phi : X \to Y$ be a regular morphism of spaces with functions. Then the following are equivalent*

*a) $\phi$ is a closed immersion*

© David Rufino CC BY-NC-ND 4.0

*b) $\phi(X)$ is closed and $\phi$ determines an isomorphism $X \xrightarrow{\sim} \phi(X)$ of spaces with functions using the structure defined in (4.3.21)*

*Proof.* *a)* $\implies$ *b)* By definition $\phi(X)$ is closed and $\phi$ determines a homeomorphism onto $\phi(X)$. Let $Z \coloneqq \phi(X)$ and $\mathcal{O}_Z$ denote the canonical structure sheaf. There is a well-defined map

$$\mathcal{O}_Z(U) \to \mathcal{O}_X(\phi^{-1}(U))$$

which is injective, and therefore injective on stalks (4.2.28) c) $\implies$ d). By (4.2.11) we need only show that the map $\mathcal{O}_{Z,\phi(x)} \to \mathcal{O}_{X,x}$ is surjective. However by hypothesis the composite map

$$\mathcal{O}_{Y,\phi(x)} \to \mathcal{O}_{Z,\phi(x)} \to \mathcal{O}_x$$

is surjective from which the result follows.

*b)* $\implies$ *a)* Evidently $\phi$ is a homeomorphism onto $\phi(X)$. Further we have a sequence of stalk maps

$$\mathcal{O}_{Y,\phi(x)} \xrightarrow{\sim} \mathcal{O}_{\phi(X),\phi(x)} \to \mathcal{O}_{X,x}$$

where the first is surjective by (4.3.21) and the second is an isomorphism by assumption. Therefore the composite is surjective and $\phi$ is a closed immersion. ∎

## 4.3.4 Locally Closed Subspace

**Lemma 4.3.24 (Locally Closed Structure Sheaf)**
*Let $(X, \mathcal{O}_X)$ be a space with functions, $(Z, \mathcal{O}_Z)$ a closed subset with canonical structure defined in (4.3.21) and $(U, \mathcal{O}_U)$ an open subset with canonical structure (4.3.12). Then*

$$\mathcal{O}_Z|_{Z\cap U} = \mathcal{O}_U|_{Z\cap U}$$

## 4.3.5 Glueing

**Proposition 4.3.25 (Patching Topologies)**
*Let $X = \bigcup_{i=1}^n U_i$ be a set with topologies given for each $U_i$. Suppose that for all $i, j = 1...n$ the following properties hold*

- *$U_i \cap U_j$ is open in $U_i$, and*
- *the identity map $1 : U_i \cap U_j \to U_i \cap U_j$ is continuous with respect to the subspace topologies on $U_i$ and $U_j$ respectively.*

*Then there is a unique topology on $X$ such that*

- *the $U_i$ are open in $X$,*
- *the topology on $U_i$ coincides with the subspace topology from $X$*

*More precisely it is given by*

$$\{U \subset X \mid U \cap U_i \text{ open in } U_i \text{ for all } i = 1...n\}$$

**Proposition 4.3.26 (Glueing Topologies)**
*Let $X = \bigcup_{i=1}^n U_i$ be a set together with bijections $\theta_i : U_i \xrightarrow{\sim} W_i$ where $W_i$ are topological spaces. Suppose that for all $i, j = 1...n$ the following properties hold*

- *$\theta_i(U_i \cap U_j)$ is open in $W_i$,*
- *the map $\theta_{ij} \coloneqq \theta_j \circ \theta_i^{-1} : \theta_i(U_i \cap U_j) \to \theta_j(U_i \cap U_j)$ is continuous.*

*Then there is a unique topology on $X$ such that*

- *$U_i$ are open in $X$,*
- *$\theta_i$ is a homeomorphism with respect to the subspace topology on $U_i$ for all $i = 1...n$.*

*More precisely the topology is given by*

$$\{U \subset X \mid \theta_i(U \cap U_i) \text{ is open in } W_i \text{ for all } i = 1...n\}$$

**Proposition 4.3.27 (Glueing Regular Maps)**
*Let $X, Y$ be spaces with functions and $X = \bigcup_{i=1}^n U_i$ an open cover. Suppose there exists regular maps*

© David Rufino CC BY-NC-ND 4.0

$$\phi_i : \left(U_i, \mathcal{O}_X|_{U_i}\right) \to (Y, \mathcal{O}_Y)$$

*which satisfy* $\phi_i|_{U_i \cap U_j} = \phi_j|_{U_i \cap U_j}$ *for all* $i, j = 1...n$*. Then there exists a unique map* $\phi : X \to Y$ *such that* $\phi|_{U_i} = \phi_i$*, which is regular.*

*Proof.* As a function $\phi$ is clearly well-defined and unique. It is continuous by (4.1.29). Suppose $g \in \mathcal{O}_Y(V)$ then by definition $(g \circ \phi)|_{\phi^{-1}(V) \cap U_i} = g \circ \phi_i \in \mathcal{O}_{U_i}(\phi^{-1}(V) \cap U_i) = \mathcal{O}_X(\phi^{-1}(V) \cap U_i)$. As $\mathcal{O}_X$ is a sheaf this shows that $g \circ \phi$ is regular as required. ∎

© David Rufino CC BY-NC-ND 4.0

## Chapter 5

# Analysis

The first task is to define the field of real numbers $\mathbb{R}$, having already constructed the rational numbers $\mathbb{Q} = \mathbf{Frac}(\mathbb{Z})$. Typically this is done by some axiomatisation as an ordered field containing $\mathbb{Q}$ for which every bounded subset has a supremum (**Dedekind Completeness**). We show $\mathbb{R}$ is unique such ordered field.

© David Rufino CC BY-NC-ND 4.0

# 5.1 Real Numbers

**Definition 5.1.1 (Ordered Ring / Field)**
*An **ordered ring** is a ring $(A, +, \cdot)$ such that the additive group $(A, +)$ is an ordered abelian group with ordering $\leq$ which also satisfies*

$$x, y \geq 0 \implies x \cdot y \geq 0$$

*An **ordered field** is a field with the structure of an ordered ring.*

**Proposition 5.1.2 (Extension to Field of Fractions)**
*Let $A$ be an ordered ring with no zero-divisors. The $K = \mathrm{Frac}(A)$ is an ordered ring with the order given by*

$$\frac{x}{y} \leq \frac{w}{z} \iff xz \leq wy$$

**Example 5.1.3**
*$\mathbb{Z}$ is an ordered ring and $\mathbb{Q} := \mathrm{Frac}(\mathbb{Z})$ is an ordered field.*

**Proposition 5.1.4 (Trichotomy Law)**
*Let $G$ be an ordered abelian group. Then $G$ is the disjoint union of $\{0\}$, $G^+$ and $G^-$.*

*More generally precisely one of $x = y$, $x < y$ and $x > y$ holds.*

**Lemma 5.1.5 (Elementary Properties)**
*Let $G$ be an ordered abelian group. Then*

*a)* $x \in G^+ \iff -x \in G^-$
*b)* $x \in G^+$ and $x \leq y \implies y \in G^+$
*c)* $x \in G^-$ and $y \leq x \implies x \in G^-$
*d)* $x, y \in G^+ \implies x + y \in G^+$
*e)* $x, y \in G^- \implies x + y \in G^-$
*f)* $x, y \in G^+ \implies xy \in G^+$

*Proof.*
a) $x \in G^+ \implies 0 \leq x \implies -x \leq 0$. Furthermore $x \neq 0 \implies -x \neq 0$ by assumption so $-x \in G^-$.
b) By transitivity $0 \leq y$. Suppose $y = 0$ then $0 \leq x \leq 0$ so $x = 0$ a contradiction.
c) Similarly
d) Follows from b) since $0 \leq x \implies 0 \leq y \leq x + y$.
e) Follows from c) since $x \leq 0 \implies x + y \leq y \leq 0$.
f) If $xy = 0 \implies x = 0$ or $y = 0$ which is a contradiction. Therefore $xy > 0$.

■

**Proposition 5.1.6 (Ordered Rings are Torsion Free)**
*Let $A$ be a non-zero ordered ring. Then the canonical map $\mathbb{Z} \to A$ is injective, in other words $A$ has characteristic $0$.*

*In particular every ordered field $K$ contains $\mathbb{Q}$ as a subring.*

*Proof.* Follows immediately by induction, since if $n.1 = 0$ then $(n-1) > n \implies 1 < 0 \implies -1 > 0 \implies 1 = -1 * -1 < 0$ a contradiction. ■

**Proposition 5.1.7 (Archimedean Field)**
*Let $K$ be an ordered field. Then the following are equivalent*

*a) $\mathbb{Q}$ is dense in $K$ i.e. for all $x, y \in K$ there exists $z \in \mathbb{Q}$ such that $x < z < y$*
*b) For all $x, y \in K$ there exists $n \in \mathbb{N}$ such that $ny > x$*

*In this case we say $K$ is **Archimedean**.*

*In particular for every $\varepsilon \in K^+$ there exists $n \in \mathbb{N}^+$ such that $\frac{1}{n} < \varepsilon$.*

*Proof.* Suppose the second property is satisfied then choose $n > \frac{1}{y-x}$. By definition $0 < \frac{1}{n} < y - x$. Let

$$S = \left\{ m \in \mathbb{Z} \;\middle|\; \frac{m}{n} \leq x \right\}$$

By the Archimedean property (applied to $\frac{1}{n}$ and $-x$) $S$ is non-empty. By the well-ordering principle it has a maximal element. By maximality $\frac{m+1}{n} > x$. Furthermore

© David Rufino CC BY-NC-ND 4.0

$$x < \frac{m+1}{n} \leq x + \frac{1}{n} < x + (y - x) = y$$

and therefore $z \coloneqq \frac{m+1}{n}$ satisfies the required property.

Conversely suppose $K$ satisfies the first property and assume wlog that $x, y > 0$. Choose $\frac{m}{n}$ such that

$$0 < \frac{m}{n} < \frac{y}{x}$$

which implies $ny > mx \geq x$. ∎

**Definition 5.1.8 (Absolute Value)**
*Let $G$ be an **ordered abelian group**. Define the absolute value as*

$$|x| \coloneqq \begin{cases} x & x \in G^+ \\ -x & x \in G^- \\ 0 & x = 0 \end{cases}$$

*Then this satisfies the following properties*

*a)* $|x| \geq 0$
*b)* $|-x| = |x|$
*c)* $|x| = 0 \iff x = 0$
*d)* $||x| - |y|| \leq |x + y| \leq |x| + |y|$

**Proposition 5.1.9 (Topology of Ordered Abelian Group)**
*Let $G$ be an ordered abelian group. Then $U \subset G$ is open in the order topology iff*

$$\forall x \in U \exists \varepsilon \in G^+ \text{ s.t. } |y - x| < \varepsilon \implies y \in U$$

*Proof.* Let $\mathcal{B}$ be the basis defined in (4.1.120). Recall (...) that $U$ is open iff for all $x \in U$ there exists $V \in \mathcal{B}$ such that $x \in V \subseteq U$. If $U \subset G$ satisfies the given condition holds then evidently $x \in \{g \in G \mid y - \varepsilon < g < y + \varepsilon\}$ which is an element of $\mathcal{B}$. Conversely if $x \in V$ with $V = \{g \in G \mid y < g < z\}$, then we may define $\varepsilon = \min(z - x, x - y)$. ∎

## 5.1.1 Sequences in an Ordered Field

**Proposition 5.1.10**
*The order topology on an ordered field $K$ is Hausdorff.*

**Proposition 5.1.11 (Convergent Sequence)**
*Let $K$ be an ordered field and $(a_n)$ in $K$ a sequence. Then the following are equivalent*

*a)* $a_n \to \alpha$ *in the sense of (4.1.49)*
*b) For all $\varepsilon \in K^+$ there exists $N(\varepsilon) \in \mathbb{N}$ such that $n \geq N(\varepsilon) \implies |a_n - \alpha| < \varepsilon$*

*As $K$ is Hausdorff limits are unique. When $K$ is Archimedean it is sufficient to consider the case $\varepsilon \coloneqq \frac{1}{N}$ for all positive integers $N$.*

**Definition 5.1.12 (Convergent Sequence)**
*Let $K$ be an ordered field. We say that a sequence $(a_n)_{n \in \mathbb{N}^+}$ is **cauchy** if for all $\varepsilon > 0$ there exists $N(\varepsilon) > 0$ such that*

$$\forall m, n \in \mathbb{N} : (m, n \geq N(\varepsilon) \implies |a_n - a_m| < \varepsilon)$$

*If $K$ is Archimedean then it is sufficient to demonstrate this only for $\varepsilon$ of the form $\frac{1}{N}$.*

**Proposition 5.1.13**
*Let $K$ be an ordered field. Then every convergent sequence is cauchy.*

*Proof.* Given $\varepsilon > 0$ then $\frac{\varepsilon}{2} > 0$. Therefore there exists $N$ such that

$$n \geq N \implies |a_n - \alpha| < \frac{\varepsilon}{2}$$

Then by the triangle inequality

$$n, m \geq N \implies |a_n - a_m| < \varepsilon$$

∎

© David Rufino CC BY-NC-ND 4.0

**Definition 5.1.14 (Sequentially Completeness)**
*We say an ordered field $K$ is **sequentially complete** if every cauchy sequence is convergent.*

**Proposition 5.1.15 (Completeness)**
*Let $K$ be an ordered field. The following conditions are equivalent*

*a) Every non-empty subset $X \subset K$ bounded above admits a supremum $\sup X \in K$*
*b) Every non-empty subset $X \subset K$ bounded below admits an infimum $\inf X \in K$*

*In this case we say that $K$ is **complete**.*

*Proof.* $\inf X = -\sup(-X)$ and vice versa. ∎

**Lemma 5.1.16**
*Let $K$ be a complete ordered field. Then $K$ is also **Archimedean***

*Proof.* Suppose $K$ is not Archimedean then $\mathbb{Z}^+$ is bounded above and in particular has a supremum $\alpha$. Then for every $n \in \mathbb{Z}^+$ we have $n+1 \leq \alpha \implies n \leq \alpha - 1$, which contradicts maximality. ∎

**Lemma 5.1.17**
*Let $X \subset K$ be a subset of an Archimedean ordered field which is "upwards closed" and bounded below (resp. "downwards closed" and bounded above).*

*Then for every $\varepsilon \in K^+$ there exists an $x \in X$ such that $x - \varepsilon \notin X$ (resp. $x + \varepsilon \notin X$)*

*Proof.* By (5.1.7) there is some $n \in \mathbb{N}^+$ such that $0 < \frac{1}{n} < \varepsilon$. Consider the set

$$\mathcal{S} \coloneqq \left\{ k \in \mathbb{Z} \;\middle|\; \frac{k}{n} \in X \right\}$$

As $X$ is bounded below then so is $\mathcal{S}$. By the well ordering principle it has a minimal element, say $k$. Define $x \coloneqq \frac{k}{n}$ and then $x - \varepsilon < x - \frac{1}{n} = \frac{k-1}{n}$ which by minimality is not in $X$, whence neither is $x - \varepsilon$. ∎

**Lemma 5.1.18**
*Let $X \subset K$ be a non-empty subset for which $x \coloneqq \sup X$ exists. Then there exists a sequence $(a_n)$ in $X$ such that $a_n \uparrow x$.*

*Proof.* For every $n \in \mathbb{N}^+$ there exists some $a_n \in X$ such that $a_n > x - \frac{1}{n}$ (otherwise $x - \frac{1}{n}$ would be a smaller upper bound). Then evidently $a_n$ is increasing. For every $\varepsilon \in K^+$ there exists some $N$ such that $0 < \frac{1}{N} < \varepsilon$ by (5.1.7). Consequently

$$n \geq N \implies a_n > x - \frac{1}{N} \implies x \geq a_n > x - \varepsilon \implies |a_n - x| < \varepsilon$$

∎

**Definition 5.1.19 (Extended Real Line)**
*Let $K$ be an ordered field and define the set*

$$K^\sharp \coloneqq K \cup \{-\infty\} \cup \{\infty\}$$

*with the obvious total ordering and induced order topology. $K \subset K^\sharp$ has the subspace topology and every open subset is of the form*

$$U, U \cup (x, \infty], U \cup [-\infty, x), U \cup [-\infty, x) \cup (y, \infty] \; .$$

*If $X \subset K$ is a subset which is unbounded above (resp. below) then define $\sup X$ (resp. $\inf X$) to be $\infty$ (resp. $-\infty$).*

**Proposition 5.1.20 (Convergence to infinity)**
*Let $K$ be an ordered field and $a_n \in K^\sharp$ a sequence. Then the following are equivalent*

*a) $a_n \to \infty$*
*b) For all $\lambda \in K$ there exists $N$ such that $n \geq N \implies a_n \geq \lambda$*

*A similar statement holds for $-\infty$.*

**Lemma 5.1.21 (Limit of Bounded Sequence)**
*Let $a_n \leq b_n$ be sequences in $K^\sharp$ converging to $\alpha$ and $\beta$ respectively. Then $\alpha \leq \beta$.*

*In particular if $a_n \downarrow \alpha$ then $\alpha \leq a_n$.*

© David Rufino CC BY-NC-ND 4.0

*Proof.* If $\beta = \infty$ then the result is vacuously true. If $\beta = -\infty$ then evidently $\alpha = -\infty$ also. Therefore we may assume that $\beta$ is finite without loss of generality.

Suppose $\alpha > \beta$ and define $\varepsilon = (\alpha - \beta)$. By assumption there exists some $N$ such that

$$n \geq N \implies |a_n - \alpha| < \frac{\varepsilon}{2} \wedge |b_n - \beta| < \frac{\varepsilon}{2}$$

Then $a_n > \frac{\alpha+\beta}{2}$ and $b_n < \frac{\alpha+\beta}{2}$ which is a contradiction. ∎

**Proposition 5.1.22 (Subsequence of a Convergent Sequence)**
*Let $K$ be an ordered field, $(a_n)$ a sequence in $K^\sharp$ and $(n_k)_{k\in\mathbb{N}^+}$ a strictly increasing sequence in $\mathbb{N}^+$. If $a_n \to \alpha$ then $a_{n_k} \to \alpha$ as $k \to \infty$.*

*Proof.* By induction $n_k \geq k$ for all $k$. Let $V$ be a neighbourhood of $\alpha$ in $K^\sharp$, then by (4.1.49) there exists $N$ such that $n \geq N \implies a_n \in V$. Consequently

$$k \geq N \implies n_k \geq k \geq N \implies a_{n_k} \in V$$

and the result follows. ∎

## 5.1.2 Sequential Completeness

**Lemma 5.1.23 (Monotone Convergence Theorem)**
*Let $K$ be a complete ordered field and $(a_n)$ an increasing (resp. decreasing) sequence in $K^\sharp$. Then $a_n \to \sup\{a_n\}$. This is finite if and only if $a_n$ is bounded above (resp. below).*

*Proof.* Suppose first that $(a_n)$ is bounded above and define $\alpha := \sup\{a_n\}$. Then for every $\varepsilon \in K^+$ there exists $N$ such that $a_N > \alpha - \varepsilon$ (for otherwise $\alpha - \varepsilon$ would be an upper bound). By assumption $n \geq N \implies \alpha - \varepsilon < a_n \leq \alpha$ whence $|\alpha - a_n| < \varepsilon$. Therefore $a_n \to \alpha$.

The case that $(a_n)$ is unbounded above is trivial. ∎

**Proposition 5.1.24 (Complete $\implies$ Sequentially Complete)**
*Let $K$ be an ordered field which is **complete**. Then it is **sequentially complete**.*

*Proof.* Let $(a_n)$ be a cauchy sequence. There exists $N$ such that $n \geq N \implies |a_n - a_N| < 1 \implies |a_n| \leq 1 + |a_N|$. Then $|a_n|$ is bounded by $\max(|a_1|, \cdots, |a_N| + 1)$. Consider the sequence

$$a_n^- := \inf_{k\geq n} a_k$$

Then $a_n^-$ is bounded and increasing. Therefore $a_n^- \uparrow \alpha := \sup a_n^- < \infty$ by (5.1.23). For every $\varepsilon \in K^+$ there exists $N_1$ such that $n \geq N_1 \implies \alpha - a_n^- < \varepsilon$. There exists $N_2$ such that $n, m \geq N_2 \implies |a_n - a_m| < \varepsilon$. By definition of $a_n^-$, for every $n \geq N_1$ there exists $m \geq N_1$ such that $|a_m - a_n^-| < \varepsilon$. Consequently every $n \geq \max(N_1, N_2)$ there is some $m$ such that

$$|\alpha - a_n| \leq |\alpha - a_n^-| + |a_n - a_n^-| \leq |\alpha - a_n^-| + |a_n - a_m| + |a_m - a_n^-| < 3\varepsilon$$

As this is independent of $m$ we see that $a_n \to \alpha$ as required. ∎

**Proposition 5.1.25 (Sequentially Complete $\implies$ Complete)**
*Let $K$ be an Archimedean ordered field which is **sequentially complete**. Then $K$ is **complete**.*

*Proof.* Let $X \subset K$ be a non-empty subset bounded above, and $\mathcal{U}$ the set of upper bounds for $X$, which in particular is bounded below. By assumption $\mathcal{U}$ is non-empty. For each $n \in \mathbb{N}^+$ we may choose $a_n \in \mathcal{U}$ such that $a_n - \frac{1}{2^n} \notin \mathcal{U}$ by (5.1.17). We may assume without loss of generality that $a_n$ is decreasing by setting $a_n := \min_{n'\leq n} a_{n'}$. We argue that

$$a_n - a_{n+1} < \frac{1}{2^n}$$

for otherwise we see that $a_{n+1} \leq a_n - \frac{1}{2^n}$ which implies $a_{n+1} \notin \mathcal{U}$. Therefore by the triangle inequality applied repeatedly

$$m \geq n \implies |a_n - a_m| \leq \sum_{i=n}^{m-1} \frac{1}{2^i} < \frac{1}{2^{n-1}} < \frac{1}{n-1}$$

By the Archimedean property we see that the sequence is cauchy and therefore convergences to $\alpha \in K$. By (5.1.21) $\alpha \leq a_n$. We claim that $\alpha \in \mathcal{U}$. For suppose not then there exists $x \in X$ such that $x > \alpha$. Then there exists $n$ such

© David Rufino CC BY-NC-ND 4.0

that $a_n - \alpha < x - \alpha \implies a_n < x$ which contradicts $a_n \in \mathcal{U}$. Suppose $\exists \alpha' \in \mathcal{U}$ such that $\alpha' < \alpha$. Then for some $n$ we have $\frac{1}{n} < \alpha - \alpha'$ therefore $a_n - \frac{1}{2^n} \geq a_n - \frac{1}{n} \geq \alpha - \frac{1}{n} > \alpha'$. This would imply $a_n - \frac{1}{2^n} \in \mathcal{U}$, a contradiction. Therefore $\alpha' \geq \alpha$ and $\alpha = \sup X$ as required. ∎

### 5.1.3 Uniqueness of Reals

**Lemma 5.1.26**
*Let $K$ be a complete ordered field. Then every $x \in K$ satisfies*

$$x = \sup\left\{\frac{m}{n} \in \mathbb{Q} \;\middle|\; \frac{m}{n} \leq x\right\}$$

*Similarly for every $x \in K$ there is a sequence $a_n \in \mathbb{Q}$ such that $a_n \uparrow x$.*

*Proof.* By assumption the supremum $\alpha$ exists. By definition $x$ is an upper bound for the set so $\alpha \leq x$. Suppose $\alpha < x$ then by (5.1.7) there exists $\frac{m}{n}$ such that $\alpha < \frac{m}{n} < x$. On the other hand by definition $\frac{m}{n} \leq \alpha$, which is a contradiction. Therefore $\alpha = x$ as required.

The final statement follows from (5.1.18). ∎

**Lemma 5.1.27**
*Let $\tau : K_1 \hookrightarrow K_2$ be an embedding of ordered fields with $K_2$ Archimedean. Suppose $x = \sup X$ for some subset $X \subset K_1$. Then $\tau(x) = \sup \tau(X)$.*

*Proof.* Clearly $x' \in X \implies x' \leq x \implies \tau(x') \leq \tau(x) \implies \tau(x) \in \tau(X)^\uparrow$. Suppose $y \in \tau(X)^\uparrow$ and $y < \tau(x)$. Then by (5.1.7) there exists $z \in \mathbb{Q}$ such that $y < z < \tau(x)$. By the order preserving property we have $z \in X^\uparrow$ and $z < x$ which is a contradiction. Therefore $y \in \tau(X)^\uparrow \implies y \geq \tau(x)$ and $\tau(x) = \sup \tau(X)$. ∎

**Proposition 5.1.28 (Uniqueness of Reals)**
*Let $K_1, K_2$ be complete ordered fields. Then there exists a unique ordered field embedding $K_1 \hookrightarrow K_2$, which is an isomorphism.*

*Proof.* Let $\tau, \tau' : K_1 \to K_2$ be order embeddings. Then they agree on $\mathbb{Q}$. By (5.1.26) every $x$ is the supremum of a set $X \subset \mathbb{Q}$. Then uniqueness follows from (5.1.27). The same result shows it is surjective.

For existence we claim that

$$\tau(x) := \sup\left\{\frac{m}{n} \cdot 1_{K_2} \;\middle|\; \frac{m}{n} \cdot 1_{K_1} \leq x\right\}$$

is a well-defined, order-preserving field isomorphism. Firstly $x$ is bounded by some integer $N$ so the supremum is well-defined. Similarly suppose $x' > x$ then by (5.1.7) we have $x < \frac{m}{n} < x'$. This shows that $\tau(x) \leq \frac{m'}{n'} \leq \tau(x')$ as required, whence $\tau$ is order preserving.

By (5.1.26) there exists $a_n, b_n \in \mathbb{Q}$ such that $a_n \uparrow x$, $b_n \uparrow y$ and by the triangle inequality $a_n + b_n \uparrow x + y$. We may show using the definition of $\tau$ that $a_n \uparrow \tau(x), b_n \uparrow \tau(y), a_n + b_n \uparrow \tau(x+y)$. Uniqueness of limits shows that $\tau(x+y) = \tau(x) + \tau(y)$. Similarly we may show that $\tau(xy) = \tau(x)\tau(y)$. ∎

### 5.1.4 Existence of Reals

**Proposition 5.1.29 (Construction of the Reals)**
*There exists a (sequentially) complete ordered field $\mathbb{R}$ which is unique up to a unique order-preserving isomorphism.*

*Furthermore it has characteristic zero and the subfield $\mathbb{Q}$ is **order-dense** (5.1.7) and **sequentially dense** (5.1.26).*

*It is **Archimedean** in the sense that for all pairs $x, y \in \mathbb{R}$ there exists $n \in \mathbb{Z}$ such that $nx > y$. Equivalently for all $\varepsilon > 0$ there exists $n$ such that $\frac{1}{n} < \varepsilon$.*

*The canonical (order) topology has base the open intervals $(a, b)$, where $a, b$ may be taken to be rational. In particular the order topology on $\mathbb{R}$ is second countable.*

*Proof.* TODO. ∎

### 5.1.5 $n$-th Root

For technical reasons it is convenient to introduce the $n$-th root function at an early stage in an elementary fashion. Other special functions may be defined by similar considerations but it is convenient to wait until there is more machinery available.

© David Rufino CC BY-NC-ND 4.0

**Lemma 5.1.30 (Binomial Theorem)**
*Let $K$ be a field of characteristic 0 and $x, y \in K$. Then*

$$(x+y)^n = \sum_{k=0}^{n} \binom{n}{k} x^{n-k} y^k$$

*where $\binom{n}{k}$ is a positive integer given by the formula*

$$\binom{n}{k} := \frac{n(n-1)\cdots(n-k+1)}{k(k-1)\cdots 1}$$

*In particular if $K$ is an ordered field and $x \geq 0$ then for every $0 \leq k \leq n$*

$$(1+x)^n \geq \binom{n}{k} x^k$$

**Lemma 5.1.31**
*Let $K$ be a field of characteristic zero, $a, b \in K$ and $n$ a positive integer. Then*

$$a^n - b^n = (a-b)(a^{n-1}b + a^{n-2}b^2 + \cdots + ab^{n-2} + b^{n-1})$$

*In particular if $a > b \geq 0$ then*

$$(a-b)nb^{n-1} < a^n - b^n < (a-b)na^{n-1}$$

**Proposition 5.1.32 ($n$-th root over the Reals)**
*Let $n$ be a positive integer. Then there exists a function*

$$\sqrt[n]{\cdot} : \mathbb{R}_{\geq 0} \to \mathbb{R}_{\geq 0}$$

*such that*

$$\sqrt[n]{x}^n = \sqrt[n]{x^n} = x$$

*It is bijective and strictly increasing.*

*Proof.* Let $a > 0$ be a positive real number. There can be at most one solution because $x > y \implies x^n > y^n$. Let

$$X = \{y \in \mathbb{R}_{\geq 0} \mid y^n \leq a\}$$

then we may see that $X$ is bounded by $\max(1, a)$, for $y > \max(1, a) \implies y^n \geq y > a$.

Define $\alpha = \sup X$. Suppose $\alpha^n > a$ then we claim there is an $h > 0$ such that $(\alpha - h)^n > a$. Then $y > \alpha - h \implies y^n > a$, or contrapositively $y^n \leq a \implies y \leq \alpha - h$, and $\alpha - h$ is an upper bound of $X$, which is a contradiction. By (5.1.31) we have

$$\alpha^n - (\alpha - h)^n \leq hn\alpha^{n-1}$$

therefore we may choose $h$ such that

$$0 < h < \frac{\alpha^n - a}{n\alpha^{n-1}}.$$

Suppose $\alpha^n < a$ then we claim there is $h > 0$ such that $(\alpha + h)^n < a$, which is a contradiction. As before

$$(\alpha + h)^n - \alpha^n \leq hn(\alpha + h)^{n-1}$$

therefore we may choose $h$ such that

$$0 < h < \frac{a - \alpha^n}{n(\alpha + h)^{n-1}}$$

to demonstrate the required property.

By the trichotomy law we deduce that $\alpha^n = a$ as required, and the $n$-th root function is well-defined and satisfies $\sqrt[n]{x}^n = x$.

Suppose that $a^n = b^n = x > 0$ then evidently $a, b > 0$ and we deduce from (5.1.31) that $a = b$. Therefore the function is unique and injective. It is evidently surjective as $\sqrt[n]{a^n} = a$ by uniqueness. Finally we may deduce from (5.1.31) that for $y > x \geq 0$

© David Rufino CC BY-NC-ND 4.0

$$\sqrt[n]{y} - \sqrt[n]{x} > \frac{y-x}{n\sqrt[n]{y}^{n-1}} > 0$$

and so the function is strictly increasing. ∎

### 5.1.6 Limsup and Liminf

**Proposition 5.1.33 (Limsup and Liminf)**
*Let $K$ be a complete ordered field and $(a_n)$ a sequence. Define the associated sequences in $K^\sharp$*

$$a_n^+ := \sup_{n' \geq n} a_{n'}$$
$$a_n^- := \inf_{n' \geq n} a_{n'}$$

*which are decreasing and increasing sequences respectively. Define*

$$\limsup a_n = \lim_{n\to\infty} a_n^+$$
$$\liminf a_n := \lim_{n\to\infty} a_n^-$$

*Then*

*a)* $a_n$ is bounded above $\iff a_n^+$ is bounded above $\iff \limsup a_n^+ < \infty$
*b)* $a_n$ is bounded below $\iff a_n^-$ is bounded above $\iff \liminf a_n^- > -\infty$
*c)* $\limsup a_n \geq \liminf a_n$
*d)* $a_n \to \alpha \iff \limsup a_n = \liminf a_n = \alpha$

*Proof.* Clearly $a_n^+$ is decreasing and $a_n^-$ is increasing. Therefore by (5.1.23) the definition of $\limsup a_n$ and $\liminf a_n$ is well-defined, the same result also demonstrates a) and b). By definition $a_n^+ \geq a_n \geq a_n^-$ so the inequality c) follows from (5.1.21).

Suppose $\alpha \to \infty$ then for every $L > 0$ there exists $N$ such that

$$n \geq N \implies a_n > L \implies a_n^- \geq L$$

whence $\liminf a_n = \infty$. Conversely suppose $\liminf a_n = \infty$ we see by definition that $a_n \to \infty$ as required. The case of $-\infty$ follows similarly.

Suppose that $\alpha = \limsup a_n = \liminf a_n$ is finite, then there exists $N_1$ such that

$$n \geq N_1 \implies a_n \geq a_n^- > \alpha - \frac{\varepsilon}{2}$$

and $N_2$ such that

$$n \geq N_2 \implies a_n \leq a_n^+ < \alpha + \frac{\varepsilon}{2}$$

Then

$$n \geq \max(N_1, N_2) \implies |a_n - \alpha| < \varepsilon$$

and we conclude $\lim_{n\to\infty} a_n = \alpha$.

Conversely suppose $a_n \to \alpha$. Then for every $\varepsilon > 0$ there exists some $N$ such that

$$n \geq N \implies \alpha - \varepsilon < a_n < \alpha + \varepsilon$$

and therefore

$$n \geq N \implies \alpha - \varepsilon \leq a_n^- \leq a_n \leq a_n^+ \leq \alpha + \varepsilon$$

Evidently then

$$\alpha - \varepsilon \leq \liminf a_n \leq \limsup a_n \leq \alpha + \varepsilon$$

As $\varepsilon$ was arbitrary then this shows $\alpha = \liminf a_n = \limsup a_n$. ∎

**Proposition 5.1.34 (Arithmetic Properties of limsup and liminf)**
*Let $K$ be a complete ordered field and $(a_n), (b_n)$ bounded sequences. Then*

*a)* $\limsup(a_n + b_n) \leq \limsup a_n + \limsup b_n$

© David Rufino CC BY-NC-ND 4.0

*b)* $\liminf(a_n + b_n) \leq \liminf a_n + \liminf b_n$

*If* $b_n \to b$ *then*

*c)* $\limsup(a_n + b_n) = \limsup a_n + b$
*d)* $\liminf(a_n + b_n) = \liminf a_n + b$

*and if in addition* $b \geq 0$ *then*

*e)* $\limsup(a_n b_n) = b \limsup a_n$
*f)* $\liminf(a_n b_n) = b \liminf a_n$

*Proof.* We prove each in turn

a) By definition $n' \geq n \implies a_{n'} + b_{n'} \leq a_n^+ + b_n^+$, whence $(a_n + b_n)^+ \leq a^+ + b_n^+$. The result follows from (5.1.21).
b) Similarly
c) By a) we have $\limsup(a_n + b_n) \leq \limsup a_n + b$. Furthermore $\limsup(a_n) = \limsup(a_n + b_n + (-b_n)) \leq \limsup(a_n + b_n) + \limsup(-b_n) = \limsup(a_n + b_n) - b$, and the result follows.
d) Similarly

**e), f)** Suppose first that $b = 0$. Then for sufficiently large $n$

$$0 \leq b_n \leq \varepsilon \implies -\varepsilon|a_n^-| \leq (a_n b_n)^+ \leq |a_n^+|\varepsilon \implies \left|(a_n b_n)^+\right| \leq \max\left(\left|\sup_k a_k^+\right|, \left|\inf_k a_k^-\right|\right)\varepsilon$$

as $(a_n)$ is assumed bounded then this shows that $(a_n b_n)^+ \to 0$ as required. In this case f) follows similarly. For the general case then

$$a_n b_n = a_n(b_n - b) + a_n b$$

Obviously $b_n - b \to 0$, so by the case already proven we have

$$\limsup\left(a_{n(b_n - b)}\right) = \liminf\left(a_{n(b_n - b)}\right) = 0$$

and in particular $a_n(b_n - b) \to 0$ by (5.1.33).d). Consequently by c)

$$\limsup(a_n b_n) = \limsup(a_n b) = b \limsup(a_n)$$

as required. The case f) follows similarly. ∎

**Proposition 5.1.35**
*Let* $K$ *be a complete ordered field and* $(a_n)$ *a bounded sequence. Then every subsequence* $\left(a_{n_k}\right)$ *satisfies*

$$\limsup_k a_{n_k} \leq \limsup_n a_n$$

*and there in fact exists a convergent subsequence* $a_{n_k}$ *such that*

$$a_{n_k} \downarrow \limsup_n a_n$$

*A similar statement applies to* $\liminf a_n$.

*Proof.* Define $b_k \coloneqq \sup_{k' \geq k} a_{n_{k'}}$, then by definition

$$b_k^+ \leq a_{n_k}^+$$

Taking limits as $k \to \infty$ and considering we find that

$$\limsup_k a_{n_k} \overset{(5.1.21)}{\leq} \lim_{k \to \infty} a_{n_k}^+ \overset{(5.1.22)}{=} \lim_{n \to \infty} a_n^+ =: \limsup_n a_n$$

Recall that $a_n^+ \downarrow \limsup a_n$. Therefore for every $k > 0$ there exists $N_k$ such that $a_{N_k}^+ \leq \limsup a_n + \frac{1}{2k}$. Then by definition there exists some $n_k \geq N_k$ such that $a_{n_k} \leq a_{N_k}^+ + \frac{1}{2k} \leq \limsup a_n + \frac{1}{k}$. We see that $a_{n_k}$ has the required property. ∎

© David Rufino CC BY-NC-ND 4.0

## 5.2 Complex Numbers

**Proposition 5.2.1 (Existence of Complex Numbers)**
*Let $K$ be a field for which $-1$ has no square root (and in particular $\operatorname{char}(K) \neq 2$). Then*

$$K[i] \coloneqq K[X]/(X^2+1)$$

*is a field with a $K$-basis $\{1, i\}$ such that $i^2 = -1$. Multiplication in $K[i]$ is defined as follows*

$$(a+bi)(c+di) = (ac-bd) + (bc+ad)i$$

*and the inverse is given by*

$$(a+bi)^{-1} = \frac{a-bi}{a^2+b^2}$$

*The field extension $K(i)/K$ is Galois of degree two with*

$$\operatorname{Gal}(K(i)/K) = \{1, \bar{\cdot}\}$$

*where $\bar{\cdot}$ is the **complex conjugation** automorphism given by*

$$\overline{a+bi} = a-bi$$

*In particular we see that $\bar{\cdot}$ satisfies the following properties*

$$\overline{\alpha+\beta} = \overline{\alpha} + \overline{\beta}$$
$$\overline{\alpha\beta} = \overline{\alpha}\,\overline{\beta}$$

*Proof.* The polynomial $X^2+1$ is of degree 2 so it is irreducible if and only if has no root. Therefore the quotient ring is a field by (3.19.40). $\mathbb{R}$ satisfies this property by (5.1.5), so $\mathbb{C}$ is a field.

Observe that as polynomials in $K[X]$

$$(a+bX)(c+dX) = ac + (bc+ad)X + bdX^2 = (ac-bd) + (bc+ad)X + bd(X^2+1)$$

from which the multiplication identity follows immediately.

For the inverse we claim that $a \neq 0$ or $b \neq 0 \implies a^2+b^2 \neq 0$. For otherwise either $a/b$ or $b/a$ would be a square root of $-1$. Therefore the expression for the inverse is well-defined and it may be verified directly to be an inverse.

The complex conjugation operator exists by (3.19.49). Therefore $\operatorname{Aut}(K(i)/K)$ has order at least $[K(i):K] = 2$, and we deduce $K(i)/K$ is Galois (3.19.107). ∎

**Definition 5.2.2 (Complex Numbers)**
*Define the **complex numbers** $\mathbb{C}$ to be the field $\mathbb{R}[i]$ and identify $\mathbb{R}$ with the subfield $\{a+0i \mid a \in \mathbb{R}\}$.*

*Define the **absolute value** on $\mathbb{C}$ as follows*

$$|a+bi| \coloneqq \sqrt{a^2+b^2} \in \mathbb{R}_{\geq 0}$$

*Define the **real part** and **imaginary part** as follows*

$$\operatorname{Re}(a+bi) = a$$
$$\operatorname{Im}(a+bi) = b$$

*Observe that*

$$\alpha + \overline{\alpha} = 2\operatorname{Re}(\alpha)$$

**Proposition 5.2.3 (Complex Absolute Value)**
*Let $\alpha, \beta \in \mathbb{C}$ be complex numbers. The following properties hold*

$$|\alpha| = |\overline{\alpha}|$$
$$|\alpha|^2 = \alpha\overline{\alpha}$$
$$|\alpha\beta| = |\alpha||\beta|$$
$$|\alpha+\beta| \leq |\alpha| + |\beta|$$

© David Rufino CC BY-NC-ND 4.0

*Furthermore this agrees with the usual notion of absolute value on* $\mathbb{R}$.

*Proof.* The first two relations follow by direct calculation. Then

$$|\alpha\beta|^2 = \alpha\beta\overline{\alpha\beta} = \alpha\overline{\alpha}\beta\overline{\beta} = |\alpha|^2|\beta|^2$$

and the third relation follows by taking square roots. Further

$$\begin{aligned}|\alpha+\beta|^2 &= (\alpha+\beta)\left(\overline{\alpha}+\overline{\beta}\right) = \alpha\overline{\alpha} + \left(\alpha\overline{\beta}+\overline{\alpha}\beta\right) + \beta\overline{\beta} \\ &= |\alpha|^2 + 2\,\mathrm{Re}(\alpha\beta) + |\beta|^2 \\ &\leq |\alpha|^2 + 2|\alpha\beta| + |\beta|^2 \\ &= (|\alpha| + |\beta|)^2\end{aligned}$$

and the result follows by taking square roots (and this being an increasing function). ■

**Proposition 5.2.4 (Complex Numbers are Complete)**
$\mathbb{C}$ *is complete as a normed vector space.*

*Proof.* Let $(\alpha_n)$ be a cauchy sequence in $\mathbb{C}$, and consider the decomposition into real and imaginary parts

$$\alpha_n = a_n + b_n i$$

Then

$$|a_n| \leq |\alpha_n|$$

$$|b_n| \leq |\alpha_n|$$

Therefore we see both $(a_n)$ and $(b_n)$ are cauchy sequences of reals, which therefore have limits $a$ and $b$ respectively. We may verify directly that $\alpha_n \to \alpha$ where $\alpha := a + bi$. ■

© David Rufino CC BY-NC-ND 4.0

## 5.3 Metric Spaces

**Definition 5.3.1 (Metric Space)**
*Let $X$ be a set and $d : X \times X \to \mathbb{R}_{\geq 0}$ a function. We say that $(X, d)$ is a **metric space** if the following properties hold*

*a) $d(x, y) = 0 \iff x = y$ for all $x, y \in X$*
*b) $d(x, y) = d(y, x)$*
*c) $d(x, z) \leq d(x, y) + d(y, z)$ for all $x, y, z \in X$*

*Given $x \in X$ and $\varepsilon > 0$ we define the **open ball** at $x$ to be*

$$B(x; \varepsilon) := \{y \in X \mid d(x, y) < \varepsilon\}$$

**Proposition 5.3.2**
*Let $(X, d)$ be a metric space. Then we say that a subset $U \subset X$ is open if*

$$\forall x \in X \exists \varepsilon > 0 \text{ s.t. } B(x; \varepsilon) \subset U$$

*The open sets form a topology with open balls forming a base.*

*For each $x \in X$ a local base consists of open balls of the form $B(x; \varepsilon)$. Further we may also consider the countable local base*

$$\mathcal{B}_x := \left\{ B\left(x; \frac{1}{n}\right) \;\middle|\; n \in \mathbb{N} \right\}$$

*In particular every metric space is **first countable**.*

**Proposition 5.3.3**
*A metric space is Hausdorff.*

**Proposition 5.3.4 (Convergent Sequence)**
*Let $(X, d)$ be a metric space and $(x_n)$ a sequence in $X$ and $x \in X$. Then the following are equivalent*

*a) For every $\varepsilon > 0$ there exists $N$ such that $n \geq N \implies d(x_n, x) < \varepsilon$*
*b) The function*

$$\begin{aligned} \mathbb{N} \cup \{\infty\} &\to X \\ n &\to x_n \\ \infty &\to x \end{aligned}$$

*is continuous at $\infty$ with respect to the topology on $\mathbb{N} \cup \{\infty\}$ given by open sets of the form $\{N, N+1, \cdots\} \cup \{\infty\}$ and arbitrary subsets of $\mathbb{N}$. Note this function is automatically continuous at every $n$.*

*In this case we say that $x_n \to x$ is a **convergent sequence**.*

**Proposition 5.3.5 (Continuous at a Point)**
*Let $f : (X, d) \to (Y, d')$ be a map of metric spaces. Then for $x \in X$ and $y := f(x)$ the following are equivalent*

*a) For all $\varepsilon > 0$ there exists $\delta > 0$ such that*

$$d(y, x) < \delta \implies d'(f(y), f(x)) < \varepsilon$$

*b) $f : X \to Y$ is continuous at $x$ in the sense of topological spaces (4.1.25)*
*c) For every sequence $x_n \to x$ we have $f(x_n) \to f(x)$*

**Proposition 5.3.6 (Continuous Criteria)**
*Let $f : X \to Y$ be a map of metric spaces. Then the following are equivalent*

*a) $f$ is continuous at every $x \in X$ (5.3.5)*
*b) $f$ is continuous in the topological sense (4.1.27)*
*c) For every convergent sequence $x_n \to x$ we have $f(x_n) \to f(x)$*

**Definition 5.3.7 (Product Metric Space)**
*Let $(X_i, d_i)$ be metric spaces for $i = 1 \cdots n$. Define the metric space on $X_1 \times \cdots \times X_n$ by*

$$d((x_1, \cdots, x_n), (y_1, \cdots, y_n)) = \max\left(d_1(x_1, y_1), \cdots, d_{n(x_n, y_n)}\right)$$

*Then the topology induced on $(X, d)$ coincides with the product topology.*

© David Rufino CC BY-NC-ND 4.0

### 5.3.1 Completeness

**Definition 5.3.8 (Cauchy Sequence)**
*Let $(X,d)$ be a metric space. A sequence $(x_n)$ in $X$ is said to be **cauchy** if*

$$\forall \varepsilon \exists N \text{ s.t. } n,m \geq N \implies d(x_n,x_m) < \varepsilon$$

**Definition 5.3.9**
*A metric space $(X,d)$ is said to be **complete** if every cauchy sequence is convergent.*

**Proposition 5.3.10**
*Let $(x_n)$ be a Cauchy sequence in a metric space $(X,d)$ with $x \in X$ a cluster point (equivalently, subsequential limit point). Then $x_n \to x$.*

**Proposition 5.3.11**
*Let $(X,d)$ be a complete metric space. A closed subset $Y \subset X$ is complete iff it is closed in $X$.*

**Proposition 5.3.12**
*Let $(X_i,d_i)$ be complete metric spaces for $i = 1 \cdots n$. Then the product metric space is $(X_1 \times \cdots \times X_n, d)$ is complete.*

**Corollary 5.3.13**
*The metric space $\mathbb{R}^k$ induced by the product metric is complete.*

*A subset $X \subset \mathbb{R}^k$ is complete if and only if it is closed.*

*Proof.* This is simply (5.3.12) and (5.3.11). ∎

### 5.3.2 Compactness

**Definition 5.3.14**
*We say a metric space $(X,d)$ is **separable** if there exists a countable dense subset $X_0$.*

**Proposition 5.3.15**
*Suppose $X$ is a separable metric space. Then it is second countable.*

*Proof.* We may show that the family

$$\mathcal{B} \coloneqq \left\{ B\left(x_0; \frac{1}{n}\right) \;\middle|\; x_0 \in X_0, n \in \mathbb{N} \right\}$$

is a countable base. ∎

**Proposition 5.3.16 (Precompact)**
*Let $(X,d)$ be a metric space. The following are equivalent*

*a) For all $\varepsilon > 0$ there exists a finite covering $X = U_1 \cup \cdots \cup U_n$ such that $\operatorname{diam}(U_i) < \varepsilon$*
*b) For all $\varepsilon > 0$ there exists a finite set of points $x_1, \cdots, x_n$ such that $X = \bigcup_{i=1}^n B(x_i;\varepsilon)$*
*c) Every sequence has a cauchy subsequence.*

*In this case we say $X$ is **precompact** or **totally bounded**.*

*Proof.* *a)* $\iff$ *b)* is straightforward. For *c)* $\implies$ *b)* suppose $X$ did not satisfy this property, then we could inductively choose a sequence $x_n$ such that

$$x_n \in X \setminus \bigcup_{i=1}^{n-1} B(x_i;\varepsilon)$$

Then evidently $d(x_n, x_{n+1}) \geq \varepsilon$ for all $n$ and it can have no cauchy subsequence.

For *b)* $\implies$ *c)* Let $(x_n)$ be a sequence. If $\{x_n\}$ is finite then we are done. Otherwise assuming it is infinite, we construct a decreasing sequence of sets

$$A_1 \supset \cdots \supset A_k \supset \cdots$$

such that $\operatorname{diam}(A_k) < \frac{1}{k}$ and $A_k \cap \{x_n\}$ is infinite. For suppose $A_1, \cdots, A_k$ are constructed, then evidently $A_k$ itself satisfies *b)* so we may find $A_k = \bigcup_{i=1}^m B\left(x_i; \frac{1}{k+1}\right)$. For at least one $i$ we must have $B\left(x_i; \frac{1}{k+1}\right) \cap A_k \cap \{x_n\}$ is infinite. Therefore we may define $A_{k+1} \coloneqq B\left(x_i; \frac{1}{k+1}\right) \cap A_k$. Finally choosing an increasing sequence $n_k$ such that $x_{n_k} \in A_k$ yields the required cauchy subsequence. ∎

© David Rufino CC BY-NC-ND 4.0

**Corollary 5.3.17 (Compactness Criteria)**
*Let $(X,d)$ be a metric space then the following are equivalent*

*a) $X$ is compact*
*b) $X$ is countably compact*
*c) $X$ is sequentially compact*
*d) $X$ is complete and precompact*

*Proof.* $a), c) \implies b)$ This is (4.1.106).a)

$a), b) \implies c)$ This is (4.1.106).b)

$c) \implies d)$ By (5.3.10) every cauchy sequence is convergent. Therefore $X$ is complete, and precompact by (5.3.16).

$d) \implies a)$ We show that $(X,d)$ is sequentially compact and second countable, and the result follows from (4.1.106).c). Sequential compactness is a consequence of (5.3.16) and (5.3.10). To show that $X$ is second countable we may for each $n$ find a finite cover

$$X = \bigcup_{i=1}^{N_n} B\left(x_{ni}; \frac{1}{n}\right)$$

We claim that

$$\left\{B\left(x_{ni}; \frac{1}{n}\right)\right\}$$

is a base. For given $y \in X$ and a nbhd $U \in \mathcal{U}_y$ we have $B(y;\varepsilon) \subset U$ for some $\varepsilon > 0$. Choose $n$ such that $\frac{1}{n} < \frac{\varepsilon}{2}$. Then for some $1 \leq i \leq N_n$, $y \in B\left(x_{ni}; \frac{1}{n}\right)$. We may show that $B\left(x_{ni}; \frac{1}{n}\right) \subset B(y;\varepsilon) \subset U$ which shows that the given collection is a base. ∎

## 5.3.3 Uniform Continuity

**Definition 5.3.18 (Uniformly Continuous)**
*Let $(X,d)$ and $(Y,d')$ be metric spaces and $f : X \to Y$ a function. We say that $f$ is **uniformly continuous** if*

$$\forall \varepsilon > 0 \exists \delta > 0 \text{ s.t. } d(x,y) < \delta \implies d'(f(x), f(y)) < \varepsilon$$

**Proposition 5.3.19 (Compact & Continuous $\implies$ Uniformly Continuous)**
*Let $f : X \to Y$ be a continuous map of metric spaces and suppose $X$ is (sequentially) compact. Then $f$ is uniformly continuous.*

***Proof 1.***. Fix $\varepsilon > 0$ then for every $x \in X$ there exists $\delta_x$ such that $f(B(x;\delta_x)) \subseteq B(f(x);\varepsilon)$. Evidently $X = \bigcup_x B\left(x; \frac{\delta_x}{2}\right)$ whence there is a finite subcover associated to $x_1, \cdots, x_n$. Define

$$\delta := \min\left(\frac{\delta_{x_1}}{2}, \cdots, \frac{\delta_{x_n}}{2}\right)$$

Given $y \in X$ then $y \in B\left(x_i; \frac{\delta_{x_i}}{2}\right)$ for some $i = 1 \cdots n$. Furthermore suppose $d(y,z) < \delta$ then

$$d(x_i, z) \leq d(x_i, y) + d(y, z) < \delta_{x_i}$$

whence $y, z \in B\left(x_i; \delta_{x_i}\right)$. Therefore by construction $d(f(y), f(z)) < \varepsilon$, as required. ∎

***Proof 2.***. By (5.3.17) we know that $X$ is sequentially compact. Suppose that $f$ is not uniformly continuous, then there exists $\varepsilon > 0$ and sequences $\{x_n\}, \{y_n\}$ such that $d(x_n, y_n) < \frac{1}{n}$ and $d(f(x_n), f(y_n)) > \varepsilon$. By assumption there is a subsequence $n_k$ such that $x_{n_k} \to x$ and $y_{n_k} \to y$. Evidently

$$d(x,y) \leq d\left(x, x_{n_k}\right) + d\left(x_{n_k}, y_{n_k}\right) + d\left(y_{n_k}, y\right) \to 0$$

whence $d(x,y) = 0$ and $x = y$. By (...) $f\left(x_{n_k}\right) \to f(x)$ and $f\left(y_{n_k}\right) \to f(x)$ which is a contradiction. ∎

© David Rufino CC BY-NC-ND 4.0

## 5.4 Normed Vector Spaces

**Definition 5.4.1 (Valued Field)**
*Let $K$ be a field. A **valuation** on $K$ is a function*

$$|\cdot| : K \to \mathbb{R}_{\geq 0}$$

*such that for all $x, y \in K$ the following relations are satisfied*

*a)* $|x| = 0 \iff x = 0$
*b)* $|xy| = |x||y|$
*c)* $|x + y| \leq |x| + |y|$

*We say that this valuation is **non-archimedean** if it satisfies the ultrametric inequality*

$$|x + y| \leq \max(|x|, |y|)$$

*for all $x, y \in K$. Otherwise it is **archimedean**.*

*We say the pair $(K, |\cdot|)$ is a **valued field**.*

**Example 5.4.2**
*Any subfield of $\mathbb{C}$ (and therefore $\mathbb{R}$) is an archimedean valued field, using the complex absolute value $|\cdot|$ (5.2.2).*

**Definition 5.4.3 (Normed Vector Space)**
*Let $(K, |\cdot|)$ be a valued field and $X$ a $K$-vector space. A norm $\|\cdot\|$ is a function*

$$\|\cdot\| : X \to \mathbb{R}_{\geq 0}$$

*satisfying the following properties*

*a)* $\|x\| = 0 \iff x = 0$
*b)* $\|\lambda x\| = |\lambda|\|x\|$
*c)* $\|x + y\| \leq \|x\| + \|y\|$

*We say the pair $(X, \|\cdot\|)$ is a **normed vector space** over the valued field $(K, |\cdot|)$.*

*A normed vector space which is complete as a metric space is called a **Banach space**.*

*A Banach space $X$ which is also an associative algebra satisfying*

$$\|x \cdot y\| \leq \|x\|\|y\|$$

*is a **Banach Algebra**. Note a valued field is naturally a Banach Algebra.*

**Example 5.4.4**
*A valued field (e.g. $\mathbb{C}$, $\mathbb{R}$, $\mathbb{Q}$) is a normed vector space over itself.*

*$\mathbb{C}$ is a normed vector space over $\mathbb{R}$.*

*For any valued field $K$ the vector space $K^n$ is a normed vector space with the following definition of norm*

$$\|v\| := \sqrt{v_1^2 + \cdots + v_n^2}$$

*(the triangle inequality requires proof).*

**Proposition 5.4.5**
*Let $(X, \|\cdot\|)$ a normed vector space. Then it is naturally a metric space with induced metric*

$$d(x, y) := \|x - y\|$$

*and in particular a Hausdorff topological space with base the open balls*

$$B(x; \varepsilon) := \{y \in X \mid \|y - x\| < \varepsilon\}$$

*Furthermore the open balls $B(x; \varepsilon)$ and $B\left(x; \frac{1}{n}\right)$ form a local base at $x$.*

*The same statement applies to valued fields and in particular $\mathbb{R}$.*

© David Rufino CC BY-NC-ND 4.0

### 5.4.1 Continuous Functions

**Proposition 5.4.6 (Continuous maps of Normed Vector Spaces)**
*Let $S \subset X$ be a subset of a normed vector space and $f : S \to Y$ be a map to a normed vector space $Y$ and $x \in S$. Then the following are equivalent*

*a) For all $\varepsilon > 0$ there exists $\delta > 0$ such that for all $x' \in S$, $\|x - x'\| < \delta \implies \|f(x) - f(x')\| < \varepsilon$*
*b) $f$ is continuous at $x$ with respect to the subspace topology on $S$*

*In this case we say the map $f$ is **continuous** at $x$.*

**Example 5.4.7**
*For $X$ a normed vector space then $\|\cdot\| : X \to \mathbb{R}$ is evidently continuous.*

### 5.4.2 Product Space

**Definition 5.4.8 (Product Norm)**
*Let $X_1, \cdots, X_n$ be normed vector spaces over the same valued field $K$. Then we may define the **product norm** on the vector space $X_1 \times \cdots \times X_n$ by*

$$\|(x_1, \cdots, x_n)\| := \max(\|x_1\|, \cdots, \|x_n\|)$$

**Proposition 5.4.9**
*Let $X_1, \cdots, X_n$ be normed vector spaces over the same valued field $K$. Then the metric (resp. topology) induced by product norm is the same as product metric (resp. product topology).*

*Proof.* Consider a point $x = (x_1, \cdots, x_n)$ then

$$B(x; \varepsilon) = \{(x_1, \cdots, x_n) \mid \|x_i\| \leq \varepsilon\} = B(x_1; \varepsilon) \times \cdots \times B(x_n; \varepsilon)$$

Therefore the open balls are open in the product topology, and being a base for the norm topology, this is a subset of the product topology (...). To show that the product topology is a subset of the norm topology it is sufficient to show that open sets of the form

$$U := U_1 \times \cdots \times U_n$$

are open in the product norm topology. For given $y \in U$ then by definition $B(y_i; \varepsilon_i) \subset U_i$. Consider $\varepsilon := \min(\varepsilon_1, \cdots, \varepsilon_n)$ and $z \in B(y; \varepsilon)$. Then

$$\|z - y\| < \varepsilon \implies \max_i(\|z_i - y_i\|) < \varepsilon \leq \varepsilon_i \implies z_i \in B(y_i; \varepsilon_i) \implies z \in U$$

in other words $B(y; \varepsilon) \subset U$ as required. ∎

**Proposition 5.4.10 (Arithmetic Operations are Continuous)**
*Let $X$ be a normed vector space over a valued field $K$. The following maps are continuous*

*a) **addition** $+ : X \times X \to X$*
*b) **scalar multiplication** $\cdot : K \times X \to X$*
*c) **inverse scalar** $(-)^{-1} : K \setminus \{0\} \to K$*

*Similarly if $X$ is a Banach algebra then the multiplication operator*

$$\cdot : X \times X \to K$$

*is also continuous.*

*Proof.* By the triangle inequality

$$\|(x_1, x_2) - (y_1, y_2)\| < \varepsilon \implies \|x_1 - y_1\| < \varepsilon \wedge \|x_2 - y_2\| < \varepsilon \implies \|(x_1 + x_2) - (y_1 + y_2)\| < 2\varepsilon$$

Therefore we may apply the criterion given by (5.4.6), and using the product norm.

For scalar multiplication observe

$$\begin{aligned}
\|(\lambda_1, x_1) - (\lambda_2, x_2)\| < \delta &\implies |\lambda_1 - \lambda_2| < \delta \wedge \|x_1 - x_2\| < \delta \\
&\implies \|\lambda_1 x_1 - \lambda_2 x_2\| \leq |\lambda_1| \|x_1 - x_2\| + \|x_2\| |\lambda_1 - \lambda_2| \\
&\implies \|\lambda_1 x_1 - \lambda_2 x_2\| \leq |\lambda_1| \|x_1 - x_2\| + (\|x_1\| + \delta) |\lambda_1 - \lambda_2| \\
&\implies \|\lambda_1 x_1 - \lambda_2 x_2\| \leq \delta^2 + (\|x_1\| + \delta)\delta
\end{aligned}$$

Therefore choose $\delta < \min\left(1, \frac{\varepsilon}{\|x_1\|+1}\right)$ to find $\|\lambda_1 x_1 - \lambda_2 x_2\| < 3\varepsilon$. ∎

© David Rufino CC BY-NC-ND 4.0

### 5.4.3 Convergent Sequences

**Proposition 5.4.11 (Convergent Sequence)**
*Let $X$ be a normed vector space and $(a_n)$ a sequence in $X$. Then the following are equivalent*

*a) $a_n \to \alpha$ in the topological sense (4.1.49)*

*b) For all $\varepsilon > 0$ there exists $N(\varepsilon) \in \mathbb{N}^+$ such that*

$$n \geq N(\varepsilon) \implies \|a_n - \alpha\| < \varepsilon$$

*c) The function*

$$\begin{aligned} \mathbb{N} \cup \{\infty\} &\to X \\ n &\to a_n \\ \infty &\to \alpha \end{aligned}$$

*is continuous with respect to the topology on $\mathbb{N} \cup \{\infty\}$ given by open sets of the form $\{N, N+1, \cdots\} \cup \{\infty\}$ and arbitrary subsets of $\mathbb{N}$.*

*In this case we say the sequence $(a_n)$ is **convergent**. If this doesn't hold it is **divergent**.*

**Proposition 5.4.12 (Uniqueness of Limits)**
*Let $X$ be a normed vector space. Then it is Hausdorff and limits are unique.*

**Proposition 5.4.13 (Continuous Image of Convergent Sequence)**
*Let $X, Y$ be normed vector spaces, $S \subset X$ and $f : S \to Y$ a map. Then the following are equivalent*

*a) $f$ is continuous in the topological sense*

*b) For every convergent sequence $a_n \to \alpha$ we have $f(a_n) \to f(\alpha)$.*

*Proof.* Recall that $X$ is first-countable and so the equivalence follows from (4.1.50) and (4.1.52). ∎

**Proposition 5.4.14 (Arithmetic of Convergent Sequences in an NVS)**
*Let $X$ be a normed vector space and $(a_n)$ and $(b_n)$ be convergent sequences in $X$. Then*

$$\lim_{n\to\infty} (a_n + b_n) = \lim_{n\to\infty} a_n + \lim_{n\to\infty} b_n$$

$$\lim_{n\to\infty} \lambda a_n = \lambda \lim_{n\to\infty} a_n$$

*In particular $a_n \to \alpha \iff a_n - \alpha \to 0$.*

**Proposition 5.4.15 (Arithmetic of Convergent Sequences in a valued field)**
*Let $K$ be a valued field and $(a_n)$ and $(b_n)$ be convergent sequences. Then*

$$\lim_{n\to\infty} a_n b_n = \left(\lim_{n\to\infty} a_n\right)\left(\lim_{n\to\infty} b_n\right)$$

*Further if both $b_n$ and $\lim_{n\to\infty} b_n$ is non-zero then*

$$\lim_{n\to\infty} \frac{a_n}{b_n} = \frac{\lim_{n\to\infty} a_n}{\lim_{n\to\infty} b_n}$$

**Proposition 5.4.16 (Arithmetic of Convergent Sequences in an ordered field)**
*Let $K$ be an ordered field and $(b_n)$ a sequence of non-negative elements. Then $b_n \to \infty \iff b_n^{-1} \to 0$.*

**Proposition 5.4.17**
*Let $K$ be a valued field and $x \in K$ such that $|x| < 1$. Then*

$$\lim_{n\to\infty} x^n = 0$$

*Proof.* We may reduce to the case that $x \in \mathbb{R}$ and $0 < x < 1$. Let $y = \frac{1-x}{x} > 0$, then it is enough to show that $(1+y)^n \to \infty$ by (5.4.16). However from (5.1.30) $(1+y)^n \geq 1 + ny$, and the result follows from Archimedean property (...). ∎

**Definition 5.4.18 (Cauchy Sequence)**
*Let $X$ be a normed vector space and $(a_n)_{n\in\mathbb{N}}$ a sequence in $X$. We say $(a_n)$ is a **cauchy sequence** if for all $\varepsilon > 0$ there exists $N(\varepsilon)$ such that*

$$n, m \geq N(\varepsilon) \implies \|a_n - a_m\| < \varepsilon$$

© David Rufino CC BY-NC-ND 4.0

*Every **convergent sequence** is **cauchy**. If the converse holds then we say $X$ is **complete**.*

### 5.4.4 Finite-Dimensional Normed Spaces

**Definition 5.4.19 (Equivalent Norms)**
*Let $X$ be a vector space over a valued field $K$. Consider two norms $\|\cdot\|_1$ and $\|\cdot\|_2$. We say that they are **equivalent** if there exists constants $c_1, c_2 > 0$ such that*

$$c_1\|x\|_1 \leq \|x\|_2 \leq c_2\|x\|_1$$

*We may prove easily that this is an equivalence relation on norms.*

**Proposition 5.4.20**
*Let $X$ be a vector space over a valued field $K$. Suppose that $\|\cdot\|_1$ and $\|\cdot\|_2$ are equivalent norms for $X$. Then the following derived quantities are the same*

- *Induced topology*
- *Convergent sequences*
- *Cauchy sequences*

**Proposition 5.4.21 (Supremum Norm)**
*Let $X$ be a finite-dimensional vector space with basis $(e_1, \cdots, e_n)$. Then the supremum norm defined by*

$$\left\|\sum_{i=1}^n \lambda_i e_i\right\|_\infty := \max(|\lambda_1|, \cdots, |\lambda_n|)$$

*is well-defined (it is the product norm on $X \cong K \times \cdots \times K$). If $K$ is complete then so is $X$.*

**Proposition 5.4.22 (Equivalence of Norms)**
*Let $X$ be a non-zero finite-dimensional vector space over a complete valued field $K$. Then every norm is equivalent to the supremum norm (and therefore complete).*

*Proof.* Fix a basis $(e_1, \cdots, e_n)$. Then for any norm $\|\cdot\|$ we have by the triangle inequality

$$\left\|\sum_{i=1}^n \lambda_i e_i\right\| \leq \sum_{i=1}^n |\lambda_i|\|e_i\| \leq \left(\sum_{i=1}^n \|e_i\|\right)\left\|\sum_{i=1}^n \lambda_i e_i\right\|_\infty$$

The reverse inequality we prove by induction. The case $n = 1$ we may take $c_1 = \|e_1\|$. Suppose $n > 1$ and there is no such constant $C$. Then there are vectors $v_k \in X$ such that

$$\|v_k\| < \frac{1}{k}\|v_k\|_\infty$$

Suppose

$$v_k = \sum_{i=1}^n \lambda_{ki} e_i$$

Consider the sets of integers

$$S_i := \left\{k \in \mathbb{N} \;\middle|\; \|v_k\|_\infty = |\lambda_{ki}|\right\} \quad i = 1 \cdots n$$

Then at least one of $S_1, \cdots, S_n$ is infinite, since the union is $\mathbb{N}$. Without loss of generality we assume it is $S_n$.

Therefore by passing to a subsequence, and scaling, we may assume that $\|v_k\|_\infty = \lambda_{kn} = 1$ and $\|v_k\| < \frac{1}{k}$. Define the vector $w_k := v_k - e_n$. Then by definition $w_k \to e_n$ with respect to $\|\cdot\|$.

Let $W$ be the $(n-1)$-dimensional subspace spanned by $\{e_1, \cdots, e_{n-1}\}$. We prove also that $w_k \to w \in W$ with respect to $\|\cdot\|$, which contradicts uniqueness of limits. Evidently $w_k$ is a cauchy sequence with respect to $\|\cdot\|$ restricted to $W$. By induction $\|\cdot\|_\infty$ is equivalent to $\|\cdot\|$ on $W$, and therefore $w_k$ is cauchy with respect to $\|\cdot\|_\infty$. As $W$ is complete (5.4.21) we see that $w_k \to w$ with respect to $\|\cdot\|_\infty$, and therefore with respect to $\|\cdot\|$. ∎

**Corollary 5.4.23**
*Let $X$ be a finite dimensional normed vector space with basis $\{x_1, \cdots, x_n\}$. The projection map*

$$\pi_i : X \to \langle x_i \rangle \xrightarrow{\sim} K$$

© David Rufino CC BY-NC-ND 4.0

*given by $\pi_i(x) = x_i^\vee(x)x_i$ is a continuous, open map.*

### 5.4.5 Convergent Series

**Definition 5.4.24 (Series)**
*Let $X$ be a normed vector space and $(a_n)$ a sequence in $X$. The formal expression*

$$\sum_{n=1}^{\infty} a_n$$

*is called the **series** associated to this sequence. For such a sequence define the **partial sum** sequence as follows*

$$s_n := \sum_{k=1}^{n} a_k$$

*If the partial sum sequence converges then we say that the series **converges** and write*

$$\sum_{n=1}^{\infty} a_n := \lim_{n\to\infty} s_n$$

*If the series*

$$\sum_{n=1}^{\infty} \|a_n\|$$

*converges then we say the original series **converges absolutely**.*

The choice of terminology is justified in the case that $X$ is complete.

**Proposition 5.4.25 (Absolute Convergence $\implies$ Convergence)**
*Let $X$ be a complete normed vector space and $\sum_{n=0}^{\infty} a_n$ a series which converges absolutely. Then the series converges in the usual sense.*

*Proof.* The partial sums of the series $\sum_{n=1}^{\infty}\|a_n\|$ are cauchy by (5.4.18). By the triangle inequality we may demonstrate that the partial sums of the series $\sum_{n=1}^{\infty} a_n$ are cauchy. Then by completeness of $X$ the partial sums must also converge. ∎

**Proposition 5.4.26**
*Let $X$ be a normed vector space and $\sum_{n=1}^{\infty} a_n$ and $\sum_{n=1}^{\infty} b_n$ two convergent sequences. Then the following series are convergent with limits*

$$\sum_{n=1}^{\infty}(a_n + b_n) = \sum_{n=1}^{\infty} a_n + \sum_{n=1}^{\infty} b_n$$

$$\sum_{n=1}^{\infty} \lambda a_n = \lambda \sum_{n=1}^{\infty} a_n$$

*where $\lambda \in K$.*

**Proposition 5.4.27**
*Let $X$ be a normed vector space and $\sum_{n=1}^{\infty} a_n$ a convergent series. Then $\|a_n\| \to 0$.*

*Proof.* Let $\alpha := \sum_{n=1}^{\infty} a_n$. Then by definition there exists $N$ such that $n \geq N \implies \left\|\sum_{k=1}^{n} a_k - \alpha\right\| < \varepsilon$. As this holds for $n+1$ we may use the triangle inequality to find that $n \geq N+1 \implies \|a_n\| < 2\varepsilon$. As $\varepsilon$ was arbitrary we find $\|a_n\| \to 0$ as required. ∎

**Proposition 5.4.28 (Comparison Test)**
*Let $X$ be a complete normed vector space. Suppose $(a_n)$ is a sequence in $X$ and $(c_n)$ is a sequence in $\mathbb{R}$ such that*

*a) The series $\sum_{n=0}^{\infty} c_n$ converges*
*b) $\|a_n\| \leq c_n$ for $n \geq N_0$*

*then the series $\sum_{n=0}^{\infty} a_n$ converges absolutely.*

*Proof.* By assumption *b)* $(c_n)$ is non-negative, and by (5.4.29) the sequence of partial sums $\sum_{k=0}^{n} c_k$ is bounded. therefore so is the sequence $\sum_{k=0}^{n}\|a_k\|$, and by the same result the series $\sum_{n=0}^{\infty}\|a_n\|$ converges, in other words $\sum_{n=0}^{\infty} a_n$ converges absolutely. ∎

© David Rufino CC BY-NC-ND 4.0

**Proposition 5.4.29 (Series of Positive Reals)**
*Let $(a_n)$ be a sequence of non-negative real numbers. Then the following are equivalent*
*a) The sequence of partial sums $A_n := \sum_{k=0}^{n} a_k$ is bounded*
*b) The series $\sum_{n=0}^{\infty} a_n$ converges (absolutely)*

*Proof.* By (5.1.23) we always have $A_n \to \sup A_n \in \mathbb{R}^\sharp$, and the limit is finite iff $A_n$ is bounded. ∎

**Proposition 5.4.30 (Geometric Series)**
*Suppose $0 \leq x < 1$ is a real number, then*

$$\sum_{n=0}^{\infty} x^n = \frac{1}{1-x}$$

*and in particular the partial sums are bounded by $\frac{1}{1-x}$.*

*Proof.* It's straightforward to show by induction that the partial sums are given explicitly by

$$\sum_{n=0}^{N} x^n = \frac{1-x^N}{1-x}\ .$$

Then the result follows from (5.4.17). ∎

**Proposition 5.4.31 (Root Test)**
*Let $X$ be a normed vector space. Given a series $\sum_{n=0}^{\infty} a_n$, define $\alpha := \limsup_n \sqrt[n]{\|a_n\|}$ Then*
*a) if $\alpha < 1$ then the series converges absolutely*
*b) if $\alpha > 1$ then the series diverges*

*Proof.*
a) Define $\alpha := \limsup_n \sqrt[n]{\|a_n\|}$ and $\beta := \frac{1+\alpha}{2} < 1$. Then by definition there is an integer $N$ such that

$$n \geq N \implies \sqrt[n]{\|a_n\|} \leq \alpha + \frac{1-\alpha}{2} = \beta \implies \|a_n\|^n \leq \beta^n$$

In particular for $n \geq N$

$$\sum_{k=0}^{n} \|a_k\| \leq \sum_{n=0}^{N-1} \|a_k\| + \sum_{k=N}^{n} \|a_k\| \leq \sum_{k=0}^{N-1} \|a_k\| + \sum_{k=N}^{n} \beta^k \leq \sum_{k=0}^{N-1} \|a_k\| + \sum_{k=0}^{\infty} \beta^k$$

where the last step follows from by (5.4.30). We conclude from (5.4.29) that the series converges absolutely.
b) By (5.1.35) there exists a subsequence $a_{n_k}$ such that $\sqrt[n_k]{\left\|a_{n_k}\right\|} \to \alpha > 1$. Therefore we may find a subsequence such that $\sqrt[n_k]{\left\|a_{n_k}\right\|} \geq \frac{1+\alpha}{2} > 1 \implies \left\|a_{n_k}\right\| > 1$. This shows that $\left\|a_{n_k}\right\| \not\to 0$, whence $\|a_n\| \not\to 0$ by (...), and the series diverges by (5.4.27).

∎

**Proposition 5.4.32 (Ratio Test)**
*Let $K$ be a valued field and consider a series $\sum_{n=1}^{\infty} a_n$. If*

$$\limsup_n \left|\frac{a_{n+1}}{a_n}\right| < 1$$

*then the series is convergent.*

*Proof.* Let $\alpha := \limsup \left|\frac{a_{n+1}}{a_n}\right|$ and $\beta = \frac{1+\alpha}{2}$. Then by assumption $\beta < 1$, and there exists $N$ such that $n \geq N \implies \left|\frac{a_{n+1}}{a_n}\right| \leq \beta \implies |a_{N+k}| \leq |a_N|\beta^k$. We then see by Comparison Test (5.4.28) that the series is convergent. ∎

**Proposition 5.4.33 (Cauchy Product of Series)**
*Let $X$ be a complete valued field and $\sum_{n=1}^{\infty} a_n$, $\sum_{n=1}^{\infty} b_n$, $\sum_{n=1}^{\infty} c_n$ series such that*
*a) one of $\sum_{n=1}^{\infty} a_n$ and $\sum_{n=1}^{\infty} b_n$ are absolutely convergent*
*b) $c_n := \sum_{k=0}^{n} a_k b_{n-k} = \sum_{k=0}^{n} a_{n-k} b_k$*

*then $\sum_{n=0}^{\infty} c_n$ is absolutely convergent and*

$$\sum_{n=0}^{\infty} c_n = \left(\sum_{n=1}^{\infty} a_n\right)\left(\sum_{n=1}^{\infty} b_n\right)$$

© David Rufino CC BY-NC-ND 4.0

*Proof.* Denote the partial sums by $A_n, B_n$ and $C_n$, and the limits by $A, B$ respectively. Then evidently

$$C_n = \sum_{i=0}^{n} a_{n-i} B_i$$

therefore

$$C_n = \sum_{i=0}^{n} a_{n-i}(B_i - B) + A_n B$$

Then it's sufficient to show that the first term tends to 0. Observe that $\beta_n := B_n - B \to 0$. **TODO** ∎

### 5.4.6 Function Spaces

**Definition 5.4.34 (Uniform Convergence)**
*Let $S$ be a metric space and $Y$ a normed vector space. Suppose $\{f_n : S \to Y\}$ is a sequence of functions and $f : S \to Y$ is a function. We say that $f_n \to f$ **converges uniformly** if*

$$\forall \varepsilon > 0 \exists N \text{ s.t. } n \geq N \implies \|f_n(x) - f(x)\| < \varepsilon \forall x \in S$$

*Note this implies, but is strictly stronger than, pointwise convergence that is $f_n(x) \to f(x)$ for all $x \in S$, as $N$ may be chosen independently of $x$.*

**Proposition 5.4.35 (Uniform Limit of Continuous Functions is Continuous)**
*Let $S$ be a metric space and $Y$ a normed vector space. Suppose that $\{f_n : S \to Y\}$ is a sequence of continuous functions converging uniformly to a limit $f : S \to Y$. Then $f$ is continuous.*

*Proof.* Fix $\varepsilon > 0$. There exists $N$ such that

$$n \geq N \implies \|f_n(x) - f(x)\| < \varepsilon \forall x \in S$$

Fix $x \in S$ then there exists $\delta > 0$ such that

$$\|y - x\| < \delta \implies \|f_n(y) - f_n(x)\| < \varepsilon$$

Therefore

$$\|y - x\| < \delta \implies \|f(x) - f(y)\| \leq \|f(x) - f_n(x)\| + \|f_n(x) - f_n(y)\| + \|f(y) - f_n(y)\| < 3\varepsilon$$

Consequently $f$ is also continuous. ∎

**Definition 5.4.36 (Bounded Function Space)**
*Let $S$ be a metric space and $Y$ a normed vector space. Define the vector space of **bounded functions** by*

$$\mathcal{B}(S, Y) := \{f : S \to Y \mid \exists C > 0 \text{ s.t. } \|f(x)\| \leq C \text{ for all } x \in S\}$$

*This is a normed vector space under the **sup norm** given by*

$$\|f\|_\infty := \sup_{x \in S} \|f(x)\|$$

*Evidently convergence in this norm is equivalent to uniform convergence.*

**Proposition 5.4.37 (Bounded Functions are Complete)**
*Suppose $Y$ is a Banach Space, then $\mathcal{B}(S, Y)$ is also a Banach Space.*

*Proof.* Let $\{f_n\}$ be a cauchy sequence of maps $S \to Y$ with respect to the sup norm. Then for every $\varepsilon > 0$ there exists $N$ such that

$$n, m \geq N \implies \|f_n - f_m\| < \varepsilon$$

In particular for every $x \in S$ we have $\{f_n(x)\}$ is cauchy. As $Y$ is complete it is a convergent sequence, for which we denote the limit by $f(x)$. For every $x \in S$ we may also choose some $m \geq N$ such that

$$\|f_m(x) - f(x)\| < \varepsilon$$

then

$$\|f(x) - f_n(x)\| \leq \|f(x) - f_m(x)\| + \|f_m(x) - f_n(x)\| < 2\varepsilon$$

© David Rufino CC BY-NC-ND 4.0

As this inequality is independent of $m$ we find that $\|f - f_n\| \to 0$ as required. We may also show easily that it is bounded. ∎

### 5.4.7 Continuous Linear Maps

**Proposition 5.4.38 (Continuous Linear Maps)**
*Let $f : X \to Y$ be a linear map of normed vector spaces. Then the following are equivalent*

*a) $f$ is continuous*

*b) There exists a constant $C > 0$ such that*

$$\|f(x)\| \leq C\|x\|$$

*c) $f$ is bounded on the unit sphere*

$$\sup\{\|f(x)\|_Y \mid \|x\|_X = 1\} < \infty$$

*When $X$ is finite-dimensional then every linear map is continuous.*

*Proof.* Suppose $b$) holds then $f$ is even uniformly continuous because

$$\|x - y\| < \frac{\varepsilon}{C} \implies \|f(x) - f(y)\| < \varepsilon$$

Conversely given $a$) there exists $\delta$ such that

$$\|x\| \leq \delta \implies \|f(x)\| < 1$$

Therefore

$$\left\| f\left(\frac{\delta x}{\|x\|}\right) \right\| < 1 \implies \|f(x)\| < \frac{1}{\delta}\|x\|$$

and we may take $C = \frac{1}{\delta}$. Given $b$) evidently $\|f\| \leq C < \infty$, and conversely $\|f(x)\| \leq \|f\|\|x\|$. ∎

**Proposition 5.4.39**
*Let $X, Y$ be normed vector spaces. Denote by $L(X, Y)$ the set of continuous linear maps. It is a normed vector space, with norm given by*

$$\|\theta\| := \sup\{\|\theta(x)\|_Y \mid \|x\|_X = 1\}$$

*Furthermore if $Y$ is complete, so is $L(X, Y)$.*

*Proof.* Evidently $\|\lambda\theta\| = |\lambda|\|\theta\|$ and $\|\theta_1 + \theta_2\| \leq \|\theta_1\| + \|\theta_2\| < \infty$. Let $\{\theta_n\}$ be a cauchy sequence in $L(X, Y)$. Given $x \in X$ then

$$\|\theta_n(x) - \theta_m(x)\| \leq \|\theta_n - \theta_m\|\|x\|$$

whence $\{\theta_n(x)\}$ is a cauchy sequence, and therefore converges to $\theta(x)$. By uniqueness of limits we see that $\theta$ is linear. Furthermore by the reverse triangle inequality (...)

$$|\|\theta_n\| - \|\theta_m\|| \leq \|\theta_n - \theta_m\|$$

in otherwords $\|\theta_n\|$ is a cauchy sequence and converges to some limit $C$. Suppose that $\|x\| = 1$ then

$$\|\theta(x)\| \leq \|\theta(x) - \theta_n(x)\| + \|\theta_n(x)\| \leq \|\theta(x) - \theta_n(x)\| + \|\theta_n\|$$

Choose $N_1$ such that $n \geq N_1 \implies \|\theta_n\| \leq C + \varepsilon$ and $N_2$ such that $n \geq N_2 \implies \|\theta(x) - \theta_n(x)\| < \varepsilon$. As $\varepsilon$ was arbitrary we see that $\|\theta\| \leq C$ and in particular is finite. ∎

**Proposition 5.4.40 (Finite-Dimensional Linear Maps are continuous)**
*Suppose that $\theta : X \to Y$ is a linear map of normed vector spaces and $X$ is finite-dimensional. Then $\theta$ is continuous.*

*Proof.* Let $\{e_1, \cdots, e_n\}$ be a basis. By (5.4.22) we know that $\|\cdot\|_X$ is equivalent to the infinity norm $\|\cdot\|_\infty$. therefore it is sufficient to show that $\theta$ is continuous with respect to this norm. However

$$\left\| \theta\left(\sum_{i=1}^n \lambda_i e_i\right) \right\| \leq \sum_{i=1}^n |\lambda_i|\|\theta(e_i)\| \leq n\left\|\sum_{i=1}^n \lambda_i e_i\right\|_\infty \max_i \|\theta(e_i)\|$$

∎

© David Rufino CC BY-NC-ND 4.0

**Proposition 5.4.41 (Continuous Multilinear Maps)**
*Let $\psi : X_1 \times \cdots \times X_p \to Z$ be a multilinear map of normed vector spaces (using the product norm). Then the following are equivalent*

*a) $\psi$ is continuous*

*b) There exists a constant $C > 0$ such that*

$$\|\psi(x_1, \cdots, x_p)\| \leq C\|x_1\| \cdots \|x_p\|$$

*c) The following quantity is finite*

$$\sup\{\|\psi(x_1, \cdots, x_p)\| \mid \|x_1\| = \cdots = \|x_p\| = 1\} < \infty$$

*In this case $L(X_1, \cdots, X_p; Z)$ is a normed vector space with norm given by expression in c). Furthermore if $Z$ is complete so is $L(X_1, \cdots, X_p; Z)$.*

*Finally if $X_1, \cdots X_p$ are finite-dimensional then every multilinear map is continuous.*

*Proof.* **a) $\implies$ b)** Suppose there is no such constant $C$, then there exists $(x_n^1, \cdots, x_n^p) \in X_1 \times \cdots \times X_p$ such that

$$\|\psi(x_n^1, \cdots, x_n^p)\| > n^p \|x_n\| \|y_n\|$$

Define $\hat{x}_n^i := \frac{x_n^i}{n\|x_n\|}$. Then $\|(\hat{x}_n^1, \cdots, \hat{x}_n^p)\| = \frac{1}{n} \to 0$ but $B(0, \cdots, 0) = 0$ by linearity and $B(\hat{x}_n^1, \cdots, \hat{x}_n^p) > 1$, which contradicts continuity of limits (...).

**b) $\implies$ a)** Observe by linearity

$$\begin{aligned}\psi(x_1, \cdots, x_p) - \psi(y_1, \cdots, y_p) &= \psi(x_1 - y_1, x_2, \cdots, x_p) + \psi(y_1, x_2 - y_2, x_3, \cdots, x_p) + \cdots + \psi(y_1, \cdots, y_{p-1}, x_p - y_p) \\ &\leq C\|x_1 - y_1\|\|x_2\| \cdots \|x_p\| + C\|x_2 - y_2\|\|y_1\|\|x_3\| \cdots \|x_p\| + \cdots \\ &+ C\|x_p - y_p\|\|y_1\| \cdots \|y_{p-1}\|\end{aligned}$$

Fix any $\varepsilon > 0$ then it is enough to show that

$$\|x_i - y_i\| < \min(\|y_i\|, \varepsilon) \implies \|\psi(x_1, \cdots, x_p) - \psi(y_1, \cdots, y_p)\| < M\varepsilon$$

for some positive constant $M$ depending only on $y$. For this implies that $\|x_i\| < 2\|y_i\|$ so that

$$\|\psi(x_1, \cdots, x_p) - \psi(y_1, \cdots, y_p)\| \leq \varepsilon \sum_{i=1}^{p} 2C\|y_1\| \cdots \widehat{\|y_i\|} \cdots \|y_p\|$$

as required.

**b) $\iff$ c)** Straightforward

The supremum is evidently a norm, and the verification of completeness follows as before. ∎

**Definition 5.4.42 (Space of Continuous Multilinear Maps)**
*Let $X_1, \cdots, X_p$, $Y$ be normed vector spaces then let $L(X_1, \cdots, X_p; Y)$ denote the $K$-vector space of all continuous multilinear maps.*

*Denote by $L^p(X, Y) := L(X^p; Y)$ the vector space of continuous multilinear maps on the $p$-fold product of $X$.*

**Proposition 5.4.43 (Alternating Multilinear Maps)**
*Let $L_a(X_1, \cdots, X_p; Y)$ denote the subspace of continuous linear maps $L(X_1, \cdots, X_p; Y)$ which are **alternating** (3.5.5). Then this is a closed subset. In particular when $Y$ is complete so is $L_a(X_1, \cdots, X_p; Y)$.*

*As before define $L_a^p(X, Y) := L_a(X^p; Y)$.*

© David Rufino CC BY-NC-ND 4.0

## 5.5 Differentiable Functions

**Definition 5.5.1 (Differentiable Scalar Function)**
*Let $K$ be a complete valued field, $U \subset K$ an open subset, $Y$ a normed vector space over $K$ and $f : U \to Y$ a function. We say that $f$ is **differentiable** at $x \in U$ if the following relation holds*

$$\forall \varepsilon > 0 \exists \delta > 0 \text{ s.t. } 0 < |y - x| < \delta \implies \left\| \frac{f(y) - f(x)}{y - x} - f'(x) \right\| < \varepsilon$$

*for some $f'(x) \in K$.*

*We say that $f$ is simply **differentiable** if it is differentiable at every point of $U$.*

*Trivially if $f$ is differentiable then so is $f|_V$ for any open subset $V \subset U$.*

**Proposition 5.5.2**
*Let $K$ be a valued field, $U \subset K$ an open subset and $f : U \to Y$ a function which is differentiable at $x \in U$. Suppose $x_n \to x$ is a sequence such that $x_n \neq x$. Then*

$$\frac{f(x_n) - f(x)}{x_n - x} \to f'(x)$$

**Proposition 5.5.3**
*Let $K$ be a valued field, $U \subset K$ an open subset and $f : U \to Y$ a function which is differentiable at $x \in U$. Then $f$ is continuous at $x$.*

**Proposition 5.5.4 (Arithmetic of Differentiable Functions)**
*Let $K$ be a valued field, $U \subset K$ an open subset and $f, g : U \to Y$ be functions differentiable at $x \in U$. Then*

*a)* $(f + g)'(x) = f'(x) + g'(x)$
*b)* $(\lambda f)'(x) = \lambda f'(x)$

*If $Y$ has continuous algebra structure (e.g. $Y = K$) then*

$$(fg)'(x) = f'(x)g(x) + f(x)g'(x)$$

*and if $Y = K$ and $f$ is non-zero on $U$ then*

$$\left(\frac{1}{f}\right)'(x) = -f' \frac{x}{f(x)^2}$$

*Finally every linear (and therefore constant) function is trivially differentiable.*

**Proposition 5.5.5**
*Let $K$ be a valued field, $U \subset K$ an open subset and $U = \bigcup_{\alpha \in \mathcal{A}} V_\alpha$ an open cover. Suppose $f : U \to Y$ is a function. Then $f$ is (continuously) differentiable if and only if $f|_{V_\alpha}$ is (continuously) differentiable for all $\alpha \in \mathcal{A}$.*

**Corollary 5.5.6**
*Let $K$ be a valued field of characteristic zero and $n$ a positive integer. Then the derivative of $x^n$ is $nx^{n-1}$.*

© David Rufino CC BY-NC-ND 4.0

## 5.6 Power Series

**Definition 5.6.1 (Convergent Power Series)**
*Let $K$ be a field. Then a **power series** is a formal expression of the form*

$$P(X) := \sum_{n=0}^{\infty} a_n X^n$$

*where $a_n \in K$. We denote the ring of power series by $K[[X]]$. If $K$ is a valued field then, we say this converges at $z \in K$ if the series*

$$\sum_{n=0}^{\infty} a_n z^n$$

*converges in $K$.*

**Proposition 5.6.2 (Radius of convergence)**
*Let $K$ be a complete valued field and consider the formal power series*

$$\sum_{n=0}^{\infty} a_n X^n \quad a_n \in K \ .$$

*Define the **radius of convergence** to be*

$$R := \left(\limsup \sqrt[n]{|a_n|}\right)^{-1}$$

*where we understand $0^{-1} = \infty$ and $\infty^{-1} = 0$. Then*

*a) For every $|z| < R$ the series $\sum_{n=0}^{\infty} a_n z^n$ is absolutely convergent, and we denote this function by $f(z)$.*
*b) For every $|z| > R$ the series $\sum_{n=0}^{\infty} a_n z^n$ is divergent*
*c) $R$ is the unique value such that a), b) hold.*
*d) The limit $f(z)$ is continuous and differentiable in the open set $B(0,R)$ with derivative*

$$f'(z) = \sum_{n=1}^{\infty} n a_n z^{n-1}$$

*which has radius of convergence at least $R$*

*Proof.*

a) First consider the case $R < \infty$. Define $c_n := a_n z^n$ then $\sqrt[n]{|c_n|} = |z| \sqrt[n]{|a_n|}$. Then clearly $|z| < R \implies \limsup \sqrt[n]{|c_n|} < 1$ whence the result follows from (5.4.31). In the case $R = \infty$ we see that $\limsup \sqrt[n]{|c_n|} = 0$ for all $z$ and the result follows similarly.
b) This follows similarly by (5.4.27).
c) This is evident for suppose $R' \neq R$ satisfies the same properties and consider $z$ such that $|z| = \frac{R'+R}{2}$ for a contradiction.
d) Let $R'$ be the radius of convergence for the given series, we claim that $R' \geq R$, or in otherwords that $|z| < R$ implies $\sum_{n=0}^{\infty}(n+1)a_{n+1}z^n$ converges. For we may choose $w$ such that $|z| < |w| < R$. Then by definition

$$\sum_{n=0}^{\infty} a_n w^n$$

converges. In particular by (...) $|a_n w^n| \to 0$ and therefore $|a_n w^n|$ is bounded, by $M$. Therefore

$$\sum_{n=1}^{\infty} \left|n a_n z^{n-1}\right| = \sum_{n=1}^{\infty} n \left|a_n w^{n-1}\right| \left|\frac{z}{w}\right|^{n-1} \leq \frac{M}{|w|} \sum_{n=1}^{\infty} n \left|\frac{z}{w}\right|^{n-1}$$

The latter series converges by the Ratio Test (5.4.32) and therefore the series converges by the Comparison Test (5.4.28). Consider distinct $z, w \in B(0,R)$ then

$$\frac{f(w) - f(z)}{w - z} - f'(z) = \sum_{n=2}^{\infty} a_n \left[\frac{w^n - z^n}{w - z} - n z^{n-1}\right]$$

Observe that for $n \geq 1$

© David Rufino CC BY-NC-ND 4.0

$$\frac{w^n - z^n}{w - z} = w^{n-1} + w^{n-2}z + \cdots + wz^{n-2} + z^{n-1}$$

Therefore for $n \geq 2$

$$\begin{aligned}\frac{w^n - z^n}{w - z} - nz^{n-1} &= (w^{n-1} - z^{n-1}) + z(w^{n-2} - z^{n-2}) + \cdots + z^{n-2}(w - z) \\ &= (w - z)\left[w^{n-2} + 2w^{n-3}z + \cdots + (n-1)z^{n-2}\right]\end{aligned}$$

For any $w$ such that $|w - z| < \frac{R-|z|}{2}$ we have $|w| \leq \frac{R+|z|}{2} =: r < R$. Then

$$\left|\frac{w^n - z^n}{w - z} - nz^{n-1}\right| \leq |w - z|\frac{n(n-1)}{2}r^{n-2}$$

which implies

$$\left|\frac{f(w) - f(z)}{w - z} - f'(z)\right| \leq |w - z|\sum_{n=2}^{\infty}|a_n|\frac{n(n-1)}{2}r^{n-2}$$

The sum is finite because the second formal derivative is absolutely convergent at $r$, by the part already demonstrated. Therefore we see that $f$ is differentiable at $z$ with derivative $f'(z)$. This also shows that $f$ is continuous at $z$ (...).

■

© David Rufino CC BY-NC-ND 4.0

# 5.7 Real Analysis

## 5.7.1 Closed Intervals

**Proposition 5.7.1 ($k$-cells are closed)**
*Subsets of the form*

$$[a_1, b_1] \times \cdots \times [a_k, b_k] \subset \mathbb{R}^k$$

*are closed.*

*Proof.* By definition of the product topology we may reduce to the case $k = 1$. Then $[a, b]$ is sequentially closed by (5.1.21), which is equivalent to being closed by (4.1.19) as $\mathbb{R}$ is first countable. ∎

## 5.7.2 Power Function

**Proposition 5.7.2 (Power function)**
*Let $\alpha \in \mathbb{Q}$ and $x \in \mathbb{R}$ be non-negative numbers. Suppose $\alpha = \frac{m}{n}$ for integers $m, n$, where $n > 0$. Then the function*

$$x^\alpha := \sqrt[n]{x^m} = \left(\sqrt[n]{x}\right)^m$$

*is continuous and strictly increasing/decreasing (according to the sign of $\alpha$). Furthermore*

$$x^\alpha x^\beta = x^{\alpha+\beta}$$

*Proof.* We first consider the case $\alpha > 0$ and $m, n > 0$. The two definitions are equivalent because $x^\alpha$ is the unique function satisfying $(x^\alpha)^n = x^m$, the latter being an injective function. We may consider the cases $\alpha = m$ and $\alpha = \frac{1}{n}$ separately (since the composition of continuous and increasing functions is continuous and increasing). In light of (5.1.32) and (5.4.10) then it remains to show that $\sqrt[n]{x}$ is continuous. However by (5.1.31)

$$\left|\sqrt[n]{y} - \sqrt[n]{x}\right| \leq \frac{|y - x|}{n\sqrt[n]{x}^{n-1}} .$$

Suppose $\alpha = \frac{m}{n}$ and $\beta = \frac{p}{q}$ then $\alpha + \beta = \frac{mq+np}{nq}$ and

$$\left(x^\alpha x^\beta\right)^{nq} = (x^\alpha)^{nq}\left(x^\beta\right)^{nq} = x^{mq}x^{np} = x^{mq+np}$$

whence by uniqueness $x^\alpha x^\beta = x^{\alpha+\beta}$.

We may define $x^0 = 1$ and $x^{-\alpha} = (x^\alpha)^{-1}$, and verify that the addition formula still holds. ∎

## 5.7.3 Countability

**Proposition 5.7.3 ($\mathbb{R}$ is separable and second countable)**
*The subset $\mathbb{Q} \subset \mathbb{R}$ is dense. Furthermore $\mathbb{R}$ is second countable, with a base formed by open balls with rational center and radius.*

*Similarly $\mathbb{Q}^k$ is dense in $\mathbb{R}^k$, which is second countable.*

*Proof.* We only need to show that $\mathbb{Q} \subset \mathbb{R}$ is dense, and the rest follows from (5.3.15) because $\mathbb{Q}$ is countable.

By (5.1.26) for every $x \in \mathbb{R}$ there is a sequence $x_n \in \mathbb{Q}$ such that $x_n \to x$. Then (4.1.19) shows that $x \in \overline{\mathbb{Q}}$ as required. ∎

## 5.7.4 Bolzano-Weierstrass Theorem

**Proposition 5.7.4 (Monotone Subsequence Theorem)**
*Let $(a_n)$ be a sequence in an ordered field. Then it has a monotone subsequence.*

*Proof.* Define the set of peaks to be

$$X = \{n \in \mathbb{N} \mid m \geq n \implies x_n \geq x_m\}$$

If $X$ is infinite then the elements determine a decreasing subsequence. Suppose $X$ is finite and define $n_1 := \max(X) + 1$. We claim inductively that there is a sequence of integers

$$n_1 < n_2 < \cdots < n_k < \cdots$$

such that $a_{n_k}$ is increasing. Evidently $n_k \notin X$ whence there exists $n_{k+1} > n_k$ such that $a_{n_{k+1}} > a_{n_k}$. ∎

© David Rufino CC BY-NC-ND 4.0

**Proposition 5.7.5 (Bolzano-Weierstrass Theorem)**
*Let $(a_n)$ be a bounded sequence in $\mathbb{R}$. Then it contains a convergent subsequence.*

***Proof 1.***. By (5.7.4) $a_n$ has a monotone subsequence. By (5.1.23) this subsequence is convergent. ∎

***Proof 2.***. Define two sequences of real numbers $(l_n)$, $(u_n)$ recursively as follows

$$(l_n, u_n) := \begin{cases} (L, U) & n = 1 \\ \left(l_{n-1}, \frac{l_{n-1}+u_{n-1}}{2}\right) & \{a_n\} \cap \left[l_{n-1}, \frac{l_{n-1}+u_{n-1}}{2}\right] \text{ is infinite} \\ \left(\frac{l_{n-1}+u_{n-1}}{2}, u_{n-1}\right) & \text{otherwise} \end{cases}$$

Then by definition

a) $(l_n)$ is increasing and $(u_n)$ is decreasing
b) $l_n \leq u_m$ for all $n, m$
c) $|u_n - l_n| = \frac{U-L}{2^n} \to 0$
d) $[l_n, u_n]$ contains infinitely many elements of the sequence $(a_n)$

We claim that

$$\bigcap_{n\in\mathbb{N}} [l_n, u_n] = \{\sup l_n\} = \{\inf u_n\}$$

Suppose $x, y$ lie in the intersection then by $c$) we have $|x - y| < \varepsilon$ for all $\varepsilon$, whence they are equal. So the set consists of at most one element. We claim $\sup l_n$ lies in the intersection. For $l_n \uparrow \sup l_n$ by (...) and $l_n \leq u_m$ whence $\sup l_n \leq u_m$ by (...) as required.

Finally let $\left(a_{n_k}\right)$ be a subsequence such that $a_{n_k} \in [l_k, u_k]$ consequently by $c$) we have $\left|a_{n_k} - \alpha\right| \to 0$ where $\alpha := \sup l_n$. ∎

**Corollary 5.7.6 (Bolzano-Weierstrass in $\mathbb{R}^k$)**
*Let $(a_n)$ be a bounded sequence in $\mathbb{R}^k$. Then it contains a convergent subsequence.*

*Proof.* Observe that $(a_n) = \left(a_n^1, \cdots, a_n^k\right)$ where $a^i$ is a bounded sequence in $\mathbb{R}$. By applying (5.7.5) to each coordinate in turn we may find a subsequence $n_j$ such that $a^i_{n_j} \to a^i$ for $i = 1 \cdots k$. By definition of the product norm it is clear that $a_{n_j} \to \left(a^1, \cdots, a^k\right)$. ∎

**Proposition 5.7.7 (Heine-Borel Theorem)**
*Let $X$ be a subset of $\mathbb{R}^k$. Then the following are equivalent*

*a) $X$ is closed and bounded*
*b) $X$ is sequentially compact*
*c) $X$ is compact*

*In particular every closed $k$-cell $[\boldsymbol{a}, \boldsymbol{b}]$ is compact.*

***Proof 1.***. Recall from (5.7.3) that $\mathbb{R}^k$ is second countable, and therefore so is $X$. Therefore the equivalences $b) \iff c)$ follows from (4.1.106).

$b), c) \implies a)$ We may use (4.1.110) to show that $X$ is closed, observing that $\mathbb{R}^k$ is both first countable and Kolmogorov. Alternatively we may observe $\mathbb{R}^k$ is Hausdorff and appeal to (4.1.107).

To show that it is bounded we may assume it is not to find a sequence $x_n$ such that $\|x_n\| \geq n$. By assumption this has a convergent subsequence $x_{n_j} \to x$. Further there is some $N_1$ such that $N_1 > \|x\|$ and some $N_2$ such that $n \geq N_2 \implies \|x_n - x\| < N_1 - |x| \implies \|x_n\| < N_1$. Then taking $n = \max(N_1, N_2)$ implies $\|f(x_n)\|$ is both less and greater than $N_1$, a contradiction.

Alternatively we may reduce to the case $k = 1$ by noting that $\pi_i(X)$ is compact (4.1.109). The family $\{B(n; 1) \mid n \in \mathbb{N}\}$ forms an open cover, for which there must exist a finite subcover. This immediately shows that $\pi_i(X)$, and therefore $X$, is bounded.

$a) \implies b), c)$ By assumption $X \subset [-M, M] \times \cdots \times [-M, M]$ for some $M$, which by (5.7.6) is sequentially compact. We may then use either (4.1.108) or (4.1.111). ∎

***Proof 2.***. Firstly $X$ is complete iff it is closed by (5.3.13). Evidently a subset of $\mathbb{R}$ is totally bounded iff it is bounded, and the same argument may be extended to $\mathbb{R}^k$. Therefore the equivalence follows from (5.3.17). ∎

© David Rufino CC BY-NC-ND 4.0

### 5.7.5 Boundedness Theorem

**Proposition 5.7.8**
*Let $X$ be a compact topological space and $f : X \to \mathbb{R}^k$ be a continuous function. Then $f$ is bounded and attains its bounds.*

***Proof 1.***. We may prove directly the case $X = [a, b]$ and $k = 1$.

Suppose $f$ is not bounded, then there is some sequence $(x_n)$ in $X$ such that $|f(x_n)| \geq n$. By choosing a convergent subsequence (5.7.5) we may assume that $x_n \to x$. Further $X$ is closed so $x \in X$ (4.1.19). By continuity (5.4.13) $f(x_n) \to f(x)$. Choose $N_1 > |f(x)|$ and $N_2$ such that $n \geq N_2 \implies |f(x_n) - f(x)| < N_1 - |f(x)| \implies |f(x_n)| < N_1$. Then $n := \max(N_1, N_2) \implies |f(x_n)| < N_1$ and $|f(x_n)| > N_1$ a contradiction. ∎

***Proof 2.***. We may reduce to the case $k = 1$ by considering the composition with the norm $\|\cdot\|$.

By (4.1.109) (or (4.1.112)) $f(X)$ is (sequentially) compact. Then by (5.7.7) $f(X)$ is closed and bounded.

Let $y = \sup f(X)$. By (5.1.18) there exists $y_n \to y$ with $y_n \in f(X)$. Since $f(X)$ is closed we see $y \in f(X)$ by (4.1.19) as required. ∎

### 5.7.6 Intermediate Value Theorem

**Proposition 5.7.9 (Intervals are connected)**
*Let $X \subset \mathbb{R}$. Then the following are equivalent*
*a) $X$ is connected*
*b) For every $x < y < z$ such that $x, z \in X$, we have $y \in X$*

*In particular $(a, b], (a, b), [a, b)$ are connected.*

*Proof.* $b) \implies a)$ Suppose $X$ is disconnected, then $X = U \sqcup V$ with $U, V$ non-empty open and closed subsets of $X$. Let $x \in U$ and $z \in V$, and assume wlog that $x < z$. Define

$$\alpha := \sup(U \cap (x, z))$$

By (5.1.21) $\alpha \in [x, z] \subset X$. By (5.1.18) there exists $\alpha_n \in U \cap (x, z)$ such that $\alpha_n \uparrow \alpha$. Then by (4.1.20) we have $\alpha \in \mathrm{cl}_{\mathbb{R}}(U) \cap X \overset{(4.1.21)}{=} \mathrm{cl}_X(U) \overset{(4.1.19)}{=} U$. If $\alpha = z$ then we reach a contradiction as $U$ and $V$ were assumed to be disjoint. Similarly if $\alpha < z$, as $U$ is open in $X$, we may choose $0 < \varepsilon < z - \alpha$ such that $(\alpha - \varepsilon, \alpha + \varepsilon) \cap X \subset U$. By assumption $\alpha + \varepsilon \in X$ whence $\alpha + \varepsilon \in U$, which contradicts maximality of $\alpha$.

$a) \implies b)$ Suppose we have $x < y < z$ such that $x, z \in X$ but $y \notin X$. Then define the open subsets $U = (-\infty, y) \cap X$ and $V = (y, \infty) \cap X$. Then

$$U := (-\infty, y) \cap X = (-\infty, y] \cap X$$

is both open and closed. ∎

**Proposition 5.7.10 (Intermediate Value Theorem)**
*Let $a < b$ be real numbers and $f : [a, b] \to \mathbb{R}$ continuous such that $f(a) < f(b)$. Then for every $f(a) < c < f(b)$ there exists $x \in (a, b)$ such that $f(x) = c$.*

***Proof 1.***. Consider

$$X := \{x' \in [a, b] \mid f(x') < c\}$$

By definition $X$ is bounded, and $a \in X$, so it is non-empty, so we may define $x := \sup X$. We claim that $f(x) = c$. By (5.1.18) there is a sequence $x_n \in X$ such that $x_n \to x$. By (5.4.13) $f(x_n) \to f(x)$ and by (5.1.21) $x \leq b$ and $f(x) \leq c$. Further by assumption $x \neq b$.

To show $f(x) \geq c$ consider the sequence $x_n := x + \min\left(\frac{1}{n}, b - x\right) \in [a, b]$. Evidently $x_n \to x$ so by continuity $f(x_n) \to f(x)$. By construction $f(x_n) \geq c$ whence $f(x) \geq c$.

Therefore we conclude $f(x) = c$ as required. ∎

***Proof 2.***. By (5.7.9) $[a, b]$ is connected, whence by (4.1.118) $f([a, b])$ is connected. The result follows from (5.7.9) again. ∎

© David Rufino CC BY-NC-ND 4.0

### 5.7.7 Mean-Value Theorem

**Proposition 5.7.11 (Mean-Value Theorem)**
*Let $f : [a, b] \to \mathbb{R}$ be a continuous function which is differentiable on $(a, b)$. Then there exists $c \in (a, b)$ such that*

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

***Proof 1.***. Define the function

$$g(x) := f(x) - \frac{f(b) - f(a)}{b - a}(x - a)$$

Then $g$ satisfies the same conditions with $g(a) = g(b) = f(a)$ and $g'(x) = f'(x) - \frac{f(b)-f(a)}{b-a}$. Evidently it is sufficient to find $c \in (a, b)$ such that $g'(c) = 0$.

By (5.7.8) $g$ is bounded and attains its maximum and minimum, say at $c, d$ respectively. If both $c, d \in \{a, b\}$ then $g$ must be the constant function whence $g'$ is exactly zero. Otherwise we may assume for example that $c \in (a, b)$. By (5.5.2)

$$\lim_{n\to\infty} \frac{g\left(c \pm \frac{1}{n}\right) - g(c)}{\pm\frac{1}{n}} = g'(c)$$

As $c$ is an extremum we may deduce from (5.1.21) that both $g'(c) \leq 0$ and $g'(c) \geq 0$, whence $g'(c) = 0$. ∎

**Corollary 5.7.12**
*Let $f : [a, b] \to \mathbb{R}$ a continuous function which is differentiable on $(a, b)$. If $f'(x)$ is positive (resp. strictly positive, negative, strictly negative) then $f(x)$ is increasing (resp. strictly increasing, decreasing, strictly decreasing).*

**Proposition 5.7.13**
*Let $f : (a, b) \to \mathbb{R}$ be a strictly increasing (resp. decreasing) continuous function. Then $f$ is a homeomorphism onto its image.*

*Proof.* Evidently $f$ is injective, therefore it is only required to show that the map

$$f^{-1} : f(U) \to U$$

is continuous. Without loss of generality we may assume that $f$ is strictly increasing. Suppose $y = f(x)$ then we require to show for every $\varepsilon > 0$ there exists $\delta > 0$ such that

$$|y' - y| < \delta \implies \left|f^{-1}(y') - f^{-1}(y)\right| < \varepsilon$$

Define $\varepsilon' \leq \varepsilon$ such that $(x - \varepsilon', x + \varepsilon') \subset (a, b)$. Define $\delta := \min(f(x + \varepsilon') - f(x), f(x) - f(x - \varepsilon'))$. Then

$$|y' - y| < \delta \implies f(x - \varepsilon') < y' < f(x + \varepsilon') \implies x - \varepsilon' < f^{-1}(y') < x + \varepsilon' \implies \left|f^{-1}(y') - f^{-1}(y)\right| < \varepsilon' \leq \varepsilon$$

as $f^{-1}$ is also monotonic. ∎

© David Rufino CC BY-NC-ND 4.0

# 5.8 Integration

## 5.8.1 Algebras of Sets

**Definition 5.8.1**
*Let $\Omega$ be a set and $\mathcal{F}$ a family of subsets of $\Omega$. We consider the following types of families*

*a) $\mathcal{F}$ is a **semi-ring** if the following properties hold*
- *i) $\varnothing \in \mathcal{F}$*
- *ii) $A_1, ..., A_n \in \mathcal{F} \implies A_1 \cap ... \cap A_n \in \mathcal{F}$*
- *iii) For all $A, B \in \mathcal{F}$ there is a family of disjoint subsets $C_1, ..., C_n \in \mathcal{F}$ such that*

$$B \setminus A = C_1 \sqcup ... \sqcup C_n$$

*b) $\mathcal{F}$ is a **ring** (resp. **algebra**) if the following properties hold*
- *i) $\varnothing \in \mathcal{F}$ (resp. $\Omega \in \mathcal{F}$)*
- *ii) $A_1, ..., A_n \in \mathcal{F} \implies A_1 \cup ... \cup A_n \in \mathcal{F}$*
- *iii) $A, B \in \mathcal{F} \implies A \setminus B \in \mathcal{F}$*

*c) $\mathcal{F}$ is a **$\sigma$-ring** (resp. **$\sigma$-algebra**) if the following properties hold*
- *i) $\varnothing \in \mathcal{F}$ (resp. $\Omega \in \mathcal{F}$)*
- *ii) $A_1, ..., A_n, ... \in \mathcal{F} \implies \bigcup_{i=1}^{\infty} A_i \in \mathcal{F}$*
- *iii) $A, B \in \mathcal{F} \implies A \setminus B \in \mathcal{F}$*

*Evidently $\sigma$-ring $\implies$ ring $\implies$ semi-ring.*

**Example 5.8.2**
*The family of half-open intervals $\mathcal{F} = \{[a, b) \mid a < b\}$ forms a semi-ring on $\mathbb{R}$.*

*More generally the family of products of half-open intervals $\{[a_1, b_1) \times ... \times [a_n, b_n) \mid a_i < b_i\}$ forms a semi-ring on $\mathbb{R}^n$.*

**Lemma 5.8.3**
*Let $\Omega$ be a set and $\{\mathcal{F}_\alpha\}_{\alpha \in \mathcal{A}}$ a family of rings (resp. algebras, $\sigma$-rings, $\sigma$-algebras). Then the family of subsets*

$$\bigcap_{\alpha \in \mathcal{A}} \mathcal{F}_\alpha$$

*is a ring (resp. algebra, $\sigma$-ring, $\sigma$-algebra). In particular every family of subsets is contained in a smallest ring (resp. algebra, $\sigma$-ring, $\sigma$-algebra).*

**Definition 5.8.4 (Borel Algebra)**
*Let $X$ be a topological space. We say that the family of **borel sets** $\mathcal{B}(X)$ is the smallest $\sigma$-algebra containing the family of open sets.*

**Proposition 5.8.5**
*Let $\mathcal{R}$ be a semi-ring (resp. semi-algebra). Then the family of finite disjoint unions*

$$\mathcal{F} := \{A_1 \sqcup ... \sqcup A_n \mid A_1, ..., A_n \in \mathcal{R} \text{ pairwise disjoint}\}$$

*is a ring (resp. algebra), and indeed the smallest ring (resp. algebra) containing $\mathcal{R}$.*

**Lemma 5.8.6 (Semi-Ring Extended Set Difference)**
*Let $\mathcal{R}$ be a semi-ring and $A_0, A_1, ..., A_n \in \mathcal{R}$. Then*

$$A_0 \setminus \bigcup_{i=1}^{n} A_i = \bigsqcup_{k=1}^{m} B_k$$

*for some disjoint sets $B_k \in \mathcal{R}$.*

*Proof.* We proceed by induction, the case $n = 1$ holding by definition. Suppose the relation holds for $n$. Then

$$A_0 \setminus \bigcup_{i=1}^{n+1} A_i = \left( \bigsqcup_{k=1}^{m} B_k \right) \setminus A_{n+1} = \bigsqcup_{k=1}^{m} (B_k \setminus A_{n+1})$$

and by definition of $\mathcal{R}$ we have

$$B_k \setminus A_{n+1} = \bigsqcup_{j=1}^{N} C_{kj} \quad k = 1...m$$

© David Rufino CC BY-NC-ND 4.0

and $C_{kj} \in \mathcal{R}$. As $B_k$ are disjoint we see that the $C_{kj}$ are disjoint for all $j,k$ and the result follows immediately. ∎

### 5.8.2 Set Functions

**Definition 5.8.7 (Finitely Additive Set Function)**
*Let $\mathcal{F}$ be a family of subsets of $\Omega$ containing $\varnothing$. A* ***set function***

$$\mu : \mathcal{F} \to [0, \infty]$$

*is* ***finitely additive*** *if it satisfies*

*a)* $\mu(\varnothing) = 0$

*b)* $\mu(A_1 \sqcup ... \sqcup A_n) = \sum_{i=1}^{n} \mu(A_i)$ *whenever this is well-defined*

*We say a set $A \in \mathcal{F}$ is* ***finite*** *if $\mu(A) < \infty$, and $\mu$ is* ***finite*** *if $\mu(\mathcal{F}) < \infty$.*

*We say that $\mu$ is* ***$\sigma$-finite*** *if for every set $A \in \mathcal{F}$ there exists a sequence $A_i$ such that $A \subset \bigcup_{i=1}^{\infty} A_i$ and $\mu(A_i) < \infty$. If $\Omega \in \mathcal{F}$ then it is sufficient for this condition to hold for $\Omega$.*

**Proposition 5.8.8 (Extension Theorem I)**
*Let $\mathcal{R}$ be a semi-ring, and $\mathcal{F}$ the ring generated by $\mathcal{R}$. Given a finitely additive set function $\mu : \mathcal{R} \to [0, \infty]$ there exists a unique extension to $\mathcal{F}$ given by*

$$\mu(A_1 \sqcup ... \sqcup A_n) = \sum_{i=1}^{n} \mu(A_i)$$

*which is finitely additive.*

*Proof.* In order to ensure the definition is well-defined suppose

$$A_1 \sqcup ... \sqcup A_n = B_1 \sqcup ... \sqcup B_m$$

Then in particular

$$A_i = (A_i \cap B_1) \sqcup ... \sqcup (A_i \cap B_m)$$

We may deduce by the additive property that

$$\sum_{i=1}^{n} \mu(A_i) = \sum_{i=1}^{n} \sum_{j=1}^{m} \mu\big(A_i \cap B_j\big)$$

Evidently this is symmetric in $A$ and $B$ so this also equals $\sum_{j=1}^{m} \mu\big(B_j\big)$. This shows that the measure $\mu$ is well-defined. The additivity property is straightforward. ∎

**Proposition 5.8.9 (Properties of finitely-additive set functions)**
*Let $(\Omega, \mathcal{R})$ be a* ***semi-ring*** *and $\mu : \mathcal{R} \to [0, \infty]$ a finitely additive set function. Then*

*a)* $\mu$ *is* ***monotone****, that is for sets $A \subset B$ we have*

$$\mu(A) \leq \mu(B)$$

*b) Suppose $A_1 \sqcup ... \sqcup A_n \subset A_0$ for $A_i \in \mathcal{R}$ then*

$$\sum_{i=1}^{n} \mu(A_i) \leq \mu(A_0)$$

*c) Suppose $A_0 \subset A_1 \cup ... \cup A_n$ for $A_i \in \mathcal{R}$ then*

$$\mu(A_0) \leq \sum_{i=1}^{n} \mu(A_i)$$

*Proof.*

a) By definition $B \setminus A = C_1 \sqcup ... \sqcup C_n$ and therefore by finite additivity

$$\mu(B) = \mu(C_1 \sqcup ... \sqcup C_n \sqcup A) = \mu(C_1) + ... + \mu(C_n) + \mu(A) \geq \mu(A)$$

b) By (5.8.6)

$$A_0 \setminus \bigsqcup_{i=1}^{n} A_i = \bigsqcup_{k=1}^{m} B_k$$

for some $B_k \in \mathcal{R}$ disjoint. Observe they are by construction disjoint from $A_i$ therefore

© David Rufino CC BY-NC-ND 4.0

$$A_0 = A_1 \sqcup ... \sqcup A_n \sqcup B_1 \sqcup ... \sqcup B_m$$

The result follows from finite additivity.

c) By replacing $A_i$ with $A_i \cap A_0$ we may assume that $A_0 = A_1 \cup ... \cup A_n$ by a). Observe that by (5.8.6)

$$\hat{A}_i := A_i \setminus \bigcup_{j=1}^{i-1} A_j = \bigsqcup_{k=1}^{m} C_{ik}$$

for $C_{ik} \in \mathcal{R}$ (if necessary padding out by $\varnothing$). Therefore

$$A_0 = \bigsqcup_{i=1}^{n} \hat{A}_i = \bigsqcup_{i,k} C_{ik}$$

and by finite additivity

$$\mu(A_0) = \sum_{i=1}^{n} \sum_{k=1}^{m} \mu(C_{ik})$$

By b)

$$\sum_{k=1}^{m} \mu(C_{ik}) \leq \mu(A_i)$$

from which the result follows.

■

**Definition 5.8.10 (Simple Functions)**
*Let $(\Omega, \mathcal{F})$ be a family of subsets and $X$ a normed vector space over $\mathbb{R}$. We define the space of simple functions to be*

$$\text{Simple}(\mathcal{F}, X) := \left\{ \sum_{i=1}^{n} \lambda_i \mathbf{1}_{A_i} \;\middle|\; A_1 \sqcup ... \sqcup A_n = \Omega, \lambda_1, ..., \lambda_n \in X \right\}$$

**Proposition 5.8.11 (Integration of Simple Functions)**
*Let $(\Omega, \mathcal{F})$ be a ring and $\mu$ a finitely additive measure and $X$ a vector space over $\mathbb{R}$. There is a well-defined linear map*

$$\int \mathrm{d}\mu : \text{Simple}(\mathcal{F}, X) \to X$$

$$\sum_{i=1}^{n} \alpha_i \mathbf{1}_{A_i} \to \sum_{i=1}^{n} \alpha_i \mu(A_i)$$

*Proof.* Suppose that $\sum_{i=1}^{n} \alpha_i \mathbf{1}_{A_i} = \sum_{j=1}^{m} \beta_j \mathbf{1}_{B_j}$ where both $\{A_i\}$ and $\{B_j\}$ are *disjoint* partitions of $X$. Then

$$A_i = (A_i \cap B_1) \sqcup ... \sqcup (A_i \cap B_m)$$

whence

$$\mu(A_i) = \sum_{j=1}^{m} \mu(A_i \cap B_j)$$

Therefore

$$\sum_{i=1}^{n} \alpha_i \mu(A_i) = \sum_{i=1}^{n} \sum_{j=1}^{m} \alpha_i \mu(A_i \cap B_j)$$

By the same argument

$$\sum_{j=1}^{m} \beta_j \mu(B_j) = \sum_{i=1}^{n} \sum_{j=1}^{m} \beta_j \mu(A_i \cap B_j)$$

Evidently $\{A_i \cap B_j\}$ is a disjoint partition of $X$. If $A_i \cap B_j \neq \varnothing$ then the first equality requires that $\alpha_i = \beta_j$. This shows that the integrals are equal as required. ■

© David Rufino CC BY-NC-ND 4.0

**Proposition 5.8.12**
*Let $(\Omega, \mathcal{F})$ be a ring, $\mu$ a finitely additive measure and $X$ a normed vector space over $\mathbb{R}$. Then we have the following properties*

*a)* $\int (f+g)\,\mathrm{d}\mu = \int f\,\mathrm{d}\mu + \int g\,\mathrm{d}\mu$
*b)* $\int \lambda f\,\mathrm{d}\mu = \lambda \int f\,\mathrm{d}\mu$
*c)* $\left\|\int f\,\mathrm{d}\mu\right\| \leq \int \|f\|\,\mathrm{d}\mu$

*If $\mu$ is finite then*

$$\left\|\int f\,\mathrm{d}\mu\right\| \leq \mu(\Omega)\|f\|_\infty$$

*Proof.* For the last part observe that

$$\left\|\sum_{i=1}^{n} \alpha \mathbf{1}_{A_i}\right\| \leq \sum_{i=1}^{n} \|\alpha\|_i \mu(A_i) \leq \|f\|_\infty \sum_{i=1}^{n} \mu(A_i) \leq \mu(\Omega)\|f\|_\infty$$

■

## 5.8.3 Integration of Regulated Functions

**Definition 5.8.13 (SubBorel Measure)**
*Let $\Omega = \mathbb{R}$ and $\mathcal{F}$ the ring generated by the half-open subintervals. Then there is a finitely additive measure $\mu : \mathcal{F} \to \mathbb{R}$ determined by*

$$\mu([a,b)) = (b-a)$$

*By (...) this determines a bounded linear map*

$$\int : \mathrm{Simple}(\mathbb{R}, X) \to X$$

*for any real vector space $X$.*

**Definition 5.8.14 (Regulated Function)**
*Let $X$ be a Banach space. A **regulated function** is a map*

$$f : \mathbb{R} \to X$$

*of **compact** support for which there exists a sequence $(f_n)$ of* simple *functions (with respect to the ring generated by the half-open subintervals) such that*

$$f_n \to f$$

*converges uniformly. We denote this by*

$$\mathrm{Reg}(\mathbb{R}, X)$$

**Lemma 5.8.15**
*A uniform limit of bounded functions is bounded. In particular a regulated function is bounded.*

*Proof.* There is some $N$ such that $n \geq N \implies \|f_n(x) - f(x)\| < 1$. Then

$$\|f(x)\| \leq \|f_n(x) - f(x)\| + \|f_n(x)\| \leq 1 + \|f_n(x)\|$$

We may take supremum over $x$ to show that $f$ is bounded. ■

**Proposition 5.8.16 (Continuous Functions are Regulated)**
*Let $X$ be a Banach space and $f : [a,b] \to X$ a continuous function. Then $f$ is regulated.*

*Proof.* We may define

$$f_n(x) := \sum_{i=1}^{N} f(a_i)\mathbf{1}_{[a_i,b_i)}(x) + \mathbf{1}\{x = b\}f(b)$$

$$a_i := a_i + \frac{j-1}{N}(b-a)$$

$$b_i := a_i + \frac{j}{N}(b-a)$$

© David Rufino CC BY-NC-ND 4.0

By (...) $f$ is uniformly continuous so that for every $\varepsilon > 0$ there exists $\delta > 0$ such that

$$|x - y| < \delta \implies \|f(x) - f(y)\| < \varepsilon$$

Choose $N$ such that

$$\max_i \frac{b_i - a_i}{N} < \delta$$

For every $x \in [a, b]$ we have $x \in [a_i, b_i)$ for some $i$ and by construction $\|f(x) - f(a_i)\| < \varepsilon$ whence $\|f(x) - f_n(x)\| < \varepsilon$. As this applies for any $x \in [a, b]$ we see that the convergence is uniform. ∎

**Proposition 5.8.17 (Integration of Regulated Functions)**
*Let $X$ be a Banach space and $f : [a, b] \to X$ a function for which $f_n \to f$ converges uniformly. Then the sequence*

$$\int_a^b f_n$$

*converges in $X$. Furthermore the limit is independent of the choice of $f_n$, and we denote the limit by $\int_a^b f$.*

*In particular the integration of continuous functions with compact support is well-defined.*

*Proof.* Let $x_n := \int f_n$. Then by (...) we have

$$\|x_n - x_m\| \leq (b - a)\|f_n - f_m\|_\infty$$

By the uniform convergence of $f_n$ we see that $x_n$ is a cauchy sequence, and so by assumption $x_n \to x$. The uniqueness is straightforward. ∎

**Lemma 5.8.18**
*Let $X$ be a Banach space and $f \in \mathrm{Reg}([a, b], X)$. Then $\|f\| \in \mathrm{Reg}([a, b], \mathbb{R})$.*

**Proposition 5.8.19 (Properties of the Integral)**
*Let $X$ be a Banach space. Then the following properties hold*

*a)* $\int_a^b (\lambda f + \mu g) = \lambda \int_a^b f + \mu \int_a^b g$
*b)* $\int_a^b f + \int_b^c f = \int_a^c f$
*c)* $\left\| \int_a^b f \right\| \leq \int_a^b \|f\|$

*In the case $X = \mathbb{R}$ then we also have the following properties*

*d)* $f \geq 0 \implies \int_a^b f \geq 0$
*e)* $f \leq g \implies \int_a^b f \leq \int_a^b g$
*f) $f$ continuous, positive such that $\int_a^b f = 0$ implies $f \equiv 0$*

**Proposition 5.8.20 (Fundamental Theorem of Calculus)**
*Let $f \in \mathrm{Reg}([a, b], X)$ where $X$ is a Banach space. Suppose $c \in (a, b)$ is a point where $f$ is continuous. Define*

$$F(x) := \int_a^x f$$

*Then $F$ is differentiable at $c$ and $F'(c) = f(c)$*

*Proof.* Observe that for $0 < h < b - c$

$$\begin{aligned} \left\| \frac{F(c + h) - F(c)}{h} - f(c) \right\| &= \frac{1}{h} \left\| \int_c^{c+h} (f(t) - f(c))\, \mathrm{d}t \right\| \\ &\leq \frac{1}{h} \int_c^{c+h} \|f(t) - f(c)\|\, \mathrm{d}t \\ &\leq \sup_{|t-c|<h} \|f(t) - f(c)\| \end{aligned}$$

and similarly for $a - c < h < 0$. As $f$ is regulated it is bounded (...) and so the supremum is finite. As $f$ is assumed continuous we find $F'(c) = f(c)$. ∎

**Proposition 5.8.21**
*Suppose $f_i \in \mathrm{Reg}([a, b], X_i)$ for $i = 1...n$. Then $(f_1, ..., f_n) \in \mathrm{Reg}([a, b], X)$ where $X := X_1 \times ...X_n$ and*

© David Rufino CC BY-NC-ND 4.0

$$\int_a^b f = \left( \int_a^b f_1, ..., \int_a^b f_n \right)$$

### 5.8.4 Measure Spaces

To create a robust theory of integration it is necessary to replace finite additivity with countably additivity, and further show that such set functions can be defined over $\sigma$-algebras and not just algebras.

**Definition 5.8.22 (Countably Additive Set Function)**
*Let $\mathcal{F}$ be a family of subsets of $\Omega$ containing $\varnothing$. A function*

$$\mu : \mathcal{F} \to [0, \infty]$$

*is a **countably additive set function** (or **measure**) if it satisfies*

*a) $\mu(\varnothing) = 0$*
*b) $\mu\left(\bigsqcup_{n=1}^{\infty} A_n\right) = \sum_{i=1}^{\infty} \mu(A_i)$ whenever this is well-defined*

*We say it is **countably subadditive** if it satisfies*

$$\mu\left( \bigcup_{i=1}^{\infty} A_i \right) \leq \sum_{i=1}^{\infty} \mu(A_i)$$

*whenever this is well-defined. We show in (5.8.25) that these conditions are almost equivalent.*

**Definition 5.8.23 (Measurable Space)**
*Let $(\Omega, \mathcal{F})$ be a measurable space and $\mu : \mathcal{F} \to [0, \infty]$ a measure. Then the triplet $(\Omega, \mathcal{F}, \mu)$ is a **measurable space**.*

The following are standard results which are somewhat more awkward to prove in the case of semi-rings. However the proofs follow precisely the same lines and semi-ring extension theorem may be more directly applicable.

**Proposition 5.8.24 (Continuity of Measures)**
*Let $\mathcal{R}$ be a semi-ring of subsets of $\Omega$ and $\mu : \mathcal{R} \to [0, \infty]$ a countably additive measure. Then for any sequence of sets $A_1, A_2, ... \in \mathcal{R}$ we have*

$$\lim_{n \to \infty} \mu\left( \bigcup_{i=1}^{n} A_i \right) = \mu\left( \bigcup_{n=1}^{\infty} A_n \right)$$

*whenever this is well-defined.*

*Proof.* Define the increasing sequence

$$B_n := \bigcup_{i=1}^{n} A_i$$

then wish to show

$$\lim_{n \to \infty} \mu(B_n) = \mu\left( \bigcup_{n=1}^{\infty} B_n \right)$$

By definition we have (with the convention $B_0 = \varnothing$)

$$B_i \setminus \bigcup_{j=1}^{i-1} B_j = B_i \setminus B_{i-1} = \bigsqcup_{k=1}^{N_i} C_{ik} \quad i = 1, 2, ...$$

for $C_{ik} \in \mathcal{R}$ whence

$$B_n = \bigsqcup_{i=1}^{n} B_i \setminus B_{i-1} = \bigsqcup_{i=1}^{n} \bigsqcup_{k=1}^{N_n} C_{ik}$$

for $n = 1, 2, ..., \infty$. Then by additivity

$$\lim_{n \to \infty} \mu(B_n) = \lim_{n \to \infty} \sum_{i=1}^{n} \sum_{k=1}^{N_i} \mu(C_{ik}) = \sum_{i=1}^{\infty} \sum_{k=1}^{N_i} \mu(C_{ik})$$

and

© David Rufino CC BY-NC-ND 4.0

$$\mu\left(\bigcup_{n=1}^{\infty} B_n\right) = \mu\left(\bigsqcup_{i=1}^{\infty}\bigsqcup_{k=1}^{N_i} C_{ik}\right) = \sum_{i=1}^{\infty}\sum_{k=1}^{N_i}\mu(C_{ik})$$

■

**Proposition 5.8.25 (Finitely Additive + Countably Subadditive $\iff$ Countably Additive)**
*Let $\mathcal{R}$ be a semi-ring of subsets of $\Omega$ and $\mu : \mathcal{R} \to [0,\infty]$ a finitely additive set function. Then*

$$\mu\left(\bigsqcup_{i=1}^{\infty} A_i\right) \geq \sum_{i=1}^{\infty}\mu(A_i)$$

*whenever this is well-defined. In particular $\mu$ is a measure iff it is countably subadditive.*

*Proof.* By (5.8.9).b) we have

$$\sum_{i=1}^{n}\mu(A_i) \leq \mu\left(\bigsqcup_{i=1}^{\infty} A_i\right)$$

and the result follows by taking $n \to \infty$ (5.1.21).

Therefore if $\mu$ is countably subadditive it must also be countably additive. Conversely if $\mu$ is countably additive then we aim to show

$$\mu\left(\bigcup_{i=1}^{\infty} A_i\right) \leq \sum_{i=1}^{\infty}\mu(A_i)$$

for a not necessarily disjoint sequence $A_i \in \mathcal{R}$. By assumption $\mu$ is finitely additive, so by (5.8.9).c)

$$\mu\left(\bigcup_{i=1}^{n} A_i\right) \leq \sum_{i=1}^{n}\mu(A_i) \leq \sum_{i=1}^{\infty}\mu(A_i)$$

Using continuity of measures (5.8.24) we deduce that

$$\mu\left(\bigcup_{i=1}^{\infty} A_i\right) = \lim_{n\to\infty}\mu\left(\bigcup_{i=1}^{n} A_i\right) \leq \sum_{i=1}^{\infty}\mu(A_i)$$

as required. ■

**Proposition 5.8.26 (Extension Theorem II)**
*Let $\mathcal{R}$ be a semi-ring of sets over $\Omega$ and $\mu : \mathcal{R} \to [0,\infty]$ a measure. Suppose that $\Omega$ is denumerable union of sets in $\mathcal{R}$. Then there exists an extension*

$$\tilde{\mu} : \sigma(\mathcal{R}) \to [0,\infty]$$

*to the $\sigma$-algebra generated by $\mathcal{R}$. When $\mu$ is $\sigma$-finite then this extension is unique. More precisely it is given by*

$$\tilde{\mu}(A) := \inf\left\{\sum_{i=1}^{\infty}\mu(A_i) \;\middle|\; A \subset \bigcup_{i=1}^{\infty} A_i, A_i \in \mathcal{R}\right\}$$

*Proof.* Define first the "outer measure"

$$\mu^\star : \mathcal{P}(\Omega) \to [0,\infty]$$
$$\mu^\star(A) := \inf\left\{\sum_{i=1}^{\infty}\mu(E_i) \;\middle|\; A \subset \bigcup_{i=1}^{\infty} E_i \quad E_i \in \mathcal{F}\right\}$$

Note that the infimum is well-defined precisely by the hypothesis. Further define the family of subsets

$$\hat{\mathcal{R}} := \{A \subset \Omega \mid \mu^\star(B) = \mu^\star(A\cap B) + \mu^\star(A^c \cap B) \quad \forall B \subset \Omega\}$$

We make a number of claims

a) $\mu^\star(A) = \mu(A)$ for all $A \in \mathcal{R}$
b) $\mu^\star$ is countably subadditive and monotone

© David Rufino CC BY-NC-ND 4.0

$$\mu^\star\left(\bigcup_{i=1}^{\infty} A_i\right) \leq \sum_{i=1}^{\infty} \mu^\star(A_i) \quad A_i \subset \Omega$$

c) $\mathcal{R} \subset \hat{\mathcal{R}}$

d) $\hat{\mathcal{R}}$ is a $\sigma$-algebra containing $\sigma(\mathcal{R})$ and $\mu^\star$ is a measure on $\hat{\mathcal{R}}$

and prove each in turn

a) Evidently $\mu^\star(A) \leq \mu(A)$ as $A \subset A \cup \varnothing ... \cup \varnothing ....$ Conversely suppose that $A \subset \bigcup_{i=1}^{\infty} A_i$ for $A_i \in \mathcal{R}$ and $\mu^\star(A) < \infty$ (otherwise the reverse inequality is clear). Then evidently

$$A = \bigcup_{i=1}^{\infty} (A_i \cap A)$$

As $\mu$ is countably subadditive (5.8.25) and monotone (5.8.9) we see

$$\mu(A) \leq \sum_{i=1}^{\infty} \mu(A_i \cap A) \leq \sum_{i=1}^{\infty} \mu(A_i)$$

By definition of the infimum we find $\mu(A) \leq \mu^\star(A)$.

b) Let $A_1, ..., A_n, ...$ be subsets of $\Omega$ such that $A \subset \bigcup_{i=1}^{\infty} A_i$. For a given $\varepsilon > 0$, choose $A_{ij} \in \mathcal{R}$ such that $A_i \subset \bigcup_{j=1}^{\infty} A_{ij}$ and

$$\sum_{j=1}^{\infty} \mu(A_{ij}) \leq \mu^\star(A_i) + \frac{\varepsilon}{2^i}$$

Then evidently $A \subset \bigcup_{i,j=1}^{\infty} A_{ij}$ and so by definition

$$\mu^\star(A) \leq \sum_{i,j=1}^{\infty} \mu(A_{ij}) \leq \sum_{i=1}^{\infty} \mu^\star(A_i) + \varepsilon$$

As $\varepsilon$ was arbitrary we deduce that $\mu^\star$ is countably subadditive. Monotonicity is straightforward from the definition.

c) Suppose $A \in \mathcal{R}$ and $B \subset \Omega$ then we require to prove that

$$\mu^\star(B) = \mu^\star(A \cap B) + \mu^\star(A^c \cap B)$$

By subadditivity it is sufficient so show that

$$\mu^\star(B) \geq \mu^\star(A \cap B) + \mu^\star(A^c \cap B)$$

Let $B \subset \bigcup_{i=1}^{\infty} B_i$ be an arbitrary cover with $B_i \in \mathcal{R}$, then by definition of $\mu^\star$ it is sufficient to show that

$$\mu^\star(A \cap B) + \mu^\star(A^c \cap B) \leq \sum_{i=1}^{\infty} \mu(B_i)$$

Recall by definition of semi-ring that

$$A^c \cap B_i = \bigsqcup_{k=1}^{n_i} C_{ik}$$

for some disjoint sets $C_{ik} \in \mathcal{R}$ and evidently

$$A \cap B \subset \bigcup_{i=1}^{\infty} (A \cap B_i)$$

$$A^c \cap B \subset \bigcup_{i=1}^{\infty} \bigsqcup_{k=1}^{n_i} C_{ik}$$

Therefore by monotonicity and countable subadditivity of $\mu^\star$

© David Rufino CC BY-NC-ND 4.0

$$\begin{aligned}\mu^\star(A\cap B)+\mu^\star(A^c\cap B) &\leq \mu^\star\left(\bigcup_{i=1}^{\infty}(A\cap B_i)\right)+\mu^\star\left(\bigcup_{i=1}^{\infty}\bigsqcup_{k=1}^{n_i}C_{ik}\right)\\ &\leq \sum_{i=1}^{\infty}\mu^\star(A\cap B_i)+\sum_{i=1}^{\infty}\sum_{k=1}^{n_i}\mu(C_{ik})\\ &= \sum_{i=1}^{\infty}\mu(A\cap B_i)+\sum_{i=1}^{\infty}\sum_{k=1}^{n_i}\mu(C_{ik})\\ &= \sum_{i=1}^{\infty}\left(\mu(A\cap B_i)+\sum_{k=1}^{n_i}\mu(C_{ik})\right)\\ &= \sum_{i=1}^{\infty}\mu(B_i)\end{aligned}$$

where the last step follows from finite additivity of $\mu$.

d) We prove the various properties in turn

i) Evidently $\mu^\star(\varnothing)=0$ and therefore $\varnothing,\Omega\in\hat{\mathcal{R}}$

ii) Given $A_1,A_2\in\hat{\mathcal{R}}$ and $B\subset\Omega$. Observe that we may apply the measurability condition to $B$ and $A_1$, and then to $B\cap A_1$ and $A_2$, and $B\cap A_1^c$ and $A_2$ to find

$$\begin{aligned}\mu^\star(B) &= \mu^\star(B\cap A_1)+\mu^\star(B\cap A_1^c)\\ &= \mu^\star(B\cap A_1\cap A_2)+\mu^\star(B\cap A_1\cap A_2^c)+\mu^\star(B\cap A_1^c\cap A_2)+\mu^\star(B\cap A_1^c\cap A_2^c)\end{aligned}$$

We may also consider the relationship with $B$ replaced with $B\cap(A_1\cup A_2)$ to find

$$\mu^\star(B\cap(A_1\cup A_2))=\mu^\star(B\cap A_1\cap A_2)+\mu^\star(B\cap A_1\cap A_2^c)+\mu^\star(B\cap A_1^c\cap A_2) \qquad (*)$$

Combining these two relations we deduce

$$\mu^\star(B)=\mu^\star(B\cap(A_1\cup A_2))+\mu^\star(B\cap(A_1\cup A_2)^c)$$

As $B$ was arbitrary then we deduce $A_1\cup A_2\in\hat{\mathcal{R}}$. Evidently by definition $A^c\in\hat{\mathcal{R}}$ and therefore by de-morgan $A_1\cap A_2\in\hat{\mathcal{R}}$. Therefore $\hat{\mathcal{R}}$ is closed under finite unions and intersections.

iii) Let $A_1,...,A_n\in\hat{\mathcal{R}}$ be disjoint then we claim that

$$\mu^\star\left(B\cap\left(\bigsqcup_{i=1}^{n}A_i\right)\right)=\sum_{i=1}^{n}\mu^\star(B\cap A_i)$$

From (*) we have

$$\mu^\star(B\cap(A_1\sqcup A_2))=\mu^\star(B\cap A_1)+\mu^\star(B\cap A_2)$$

and the result follows by induction.

iv) Let $A_1,A_2,...\in\hat{\mathcal{R}}$ be a countable family and $B\subset\Omega$. We require to prove that

$$\mu^\star(B)=\mu^\star\left(B\cap\bigcup_{i=1}^{\infty}A_i\right)+\mu^\star\left(B\cap\left(\bigcup_{i=1}^{\infty}A_i\right)^c\right)$$

By (5.8.6) we may consider the decomposition

$$A_i\setminus\bigcup_{j=1}^{i-1}A_j=\bigsqcup_{k=1}^{n_i}C_{ik}$$

where $C_{ik}\in\mathcal{R}$ are disjoint. Observe

$$\bigcup_{i=1}^{n}A_i=\bigsqcup_{i=1}^{n}\bigsqcup_{k=1}^{n_i}C_{ik}\quad\text{for } n=1,2,...,\infty$$

We have shown in ii) that $\bigcup_{i=1}^{n}A_i\in\hat{\mathcal{R}}$, therefore

© David Rufino CC BY-NC-ND 4.0

$$\begin{aligned}\mu^\star(B) &= \mu^\star\left(B \cap \left(\bigcup_{i=1}^{n} A_i\right)\right) + \mu^\star\left(B \cap \left(\bigcup_{i=1}^{n} A_i\right)^c\right) \\ &= \mu^\star\left(B \cap \left(\bigsqcup_{i=1}^{n}\bigsqcup_{k=1}^{n_i} C_{ik}\right)\right) + \mu^\star\left(B \cap \left(\bigcup_{i=1}^{n} A_i\right)^c\right)\end{aligned}$$

Recall from iii) that $\mu^\star(B \cap -)$ is finitely additive on $\hat{\mathcal{R}}$. Additionally by monotonicity we may deduce that

$$\begin{aligned}\mu^\star(B) &= \sum_{i=1}^{n}\sum_{k=1}^{n_i} \mu^\star(B \cap C_{ik}) + \mu^\star\left(B \cap \left(\bigcup_{i=1}^{n} A_i\right)^c\right) \\ &\geq \sum_{i=1}^{n}\sum_{k=1}^{n_i} \mu^\star(B \cap C_{ik}) + \mu^\star\left(B \cap \left(\bigcup_{i=1}^{\infty} A_i\right)^c\right)\end{aligned}$$

Letting $n \to \infty$ we find by subadditivity

$$\begin{aligned}\mu^\star(B) &\geq \sum_{i=1}^{\infty}\sum_{k=1}^{n_i} \mu^\star(B \cap C_{ik}) + \mu^\star\left(B \cap \left(\bigcup_{i=1}^{\infty} A_i\right)^c\right) \\ &\geq \mu^\star\left(B \cap \left(\bigcup_{i=1}^{\infty} A_i\right)\right) + \mu^\star\left(B \cap \left(\bigcup_{i=1}^{\infty} A_i\right)^c\right)\end{aligned}$$

However by subadditivity the reverse inequality holds and we deduce that these are all equal. This shows that $\bigcup_{i=1}^{\infty} A_i \in \hat{\mathcal{R}}$ and $\hat{\mathcal{R}}$ is closed under countable unions. Consider further the case $A_i$ are disjoint and $B \coloneqq \bigsqcup_{j=1}^{\infty} A_j$. Then we may assume that $n_i = 1$ and $C_{i1} = A_i$. Using the above relationship

$$\mu^\star\left(\bigsqcup_{j=1}^{\infty} A_j\right) = \sum_{i=1}^{\infty} \mu^\star\left(\left(\bigsqcup_{j=1}^{\infty} A_j\right) \cap A_i\right) = \sum_{i=1}^{\infty} \mu^\star(A_i)$$

This shows that $\mu^\star$ is countably additive.

■

**Definition 5.8.27 (Complete Set Function)**
*Let $(\Omega, \mathcal{F}, \mu)$ be a measure space. We say it is **complete** if for every null set $N$ (such that $\mu(N) = 0$) and $N' \subseteq N$ we have $N' \in \mathcal{F}$.*

**Proposition 5.8.28 (Completion of a Measure Space)**
*Let $(\Omega, \mathcal{F}, \mu)$ be a measure space and define the **completion** by*

$$\begin{aligned}\overline{\mathcal{F}} &\coloneqq \{A \cup Z \mid A \in \mathcal{F}, Z \subseteq N, \mu(N) = 0\} \\ \overline{\mu}(A \cup Z) &\coloneqq \mu(A)\end{aligned}$$

*Then $\left(\Omega, \overline{\mathcal{F}}, \overline{\mu}\right)$ is a well-defined complete measure space. Furthermore $\overline{\mathcal{F}}$ is the smallest complete $\sigma$-algebra containing $\mathcal{F}$, and $\overline{\mu}$ is the unique extension.*

## 5.8.5 Borel Measure on $\mathbb{R}$

**Proposition 5.8.29**
*Let $\mathcal{R}$ be the semi-ring of half-open intervals over $\mathbb{R}$. Then the set function*

$$\mu : \mathcal{R} \to [0, \infty]$$

*given by*

$$\mu([a, b)) = b - a,$$

*is countably additive and $\sigma$-finite.*

*Proof.* Evidently $\mu$ is finitely additive, therefore by (5.8.25) it is sufficient to show that it is countably subadditive. Suppose that

© David Rufino CC BY-NC-ND 4.0

$$\bigcup_{i=1}^{\infty}[a_i, b_i) = [a, b)$$

Let $\varepsilon < b - a$. Then $[a, b-\varepsilon]$ is compact by (5.7.7) and $U_i := \left(a_i - \frac{\delta}{2^i}, b_i\right)$ is open for every $\delta > 0$. By definition of compactness there is some finite integer $n$ such that

$$\left[a, \tilde{b}\right] := [a, b-\varepsilon] \subset \bigcup_{i=1}^{n}\left(a_i - \frac{\delta}{2^i}, b_i\right) =: (\tilde{a}_i, b_i)$$

By reordering we assume that $\tilde{a}_1 < ... < \tilde{a}_n$. As $\left[a, \tilde{b}\right]$ is an interval we must have $\tilde{a}_{i+1} < b_i$. Further we may assume that $b_i < b_{i+1}$ (otherwise we may omit the interval $(\tilde{a}_i, b_i)$). By construction we must also have $\tilde{a}_1 < a$ and $\tilde{b} < b_n$.

Consequently

$$b - a - \varepsilon = \tilde{b} - a \leq b_n - \tilde{a}_1 = b_1 - \tilde{a}_1 + \sum_{i=1}^{n-1}(b_{i+1} - b_i) \leq \sum_{i=1}^{n}(b_i - \tilde{a}_i) \leq \sum_{i=1}^{n}(b_i - a_i) + \delta \leq \sum_{i=1}^{\infty}(b_i - a_i) + \delta$$

This shows

$$\mu([a, b)) = \mu\left(\bigcup_{i=1}^{\infty}[a_i, b_i)\right) \leq \sum_{i=1}^{\infty}\mu([a_i, b_i))$$

as required. ∎

**Proposition 5.8.30**
*Let $D \subset \mathbb{R}$ be an additive subgroup containing $\mathbb{Q}$. Then $\mathcal{B}(\mathbb{R})$ is generated by any of the following families of intervals*

1. $(a, b]$ $a < b \in D$
2. $(a, b)$ $a < b \in D$
3. $[a, b)$ $a < b \in D$
4. $(a, \infty)$ $a \in D$
5. $[a, \infty)$ $a \in D$
6. $(-\infty, a]$ $a \in D$
7. $(-\infty, a)$ $a \in D$

*Further there is a unique measure*

$$\mu : \mathcal{B}(\mathbb{R}) \to [0, \infty]$$

*such that*

$$\mu([a, b)) = b - a$$

*Proof.* Denote the corresponding $\sigma$-algebras by $\mathcal{F}_1, ..., \mathcal{F}_7$. Observe that by the Archimedean property

$$(a, b) = \bigcup_{n=1}^{\infty}\left(a, b - \frac{1}{n}\right] = \bigcup_{n=1}^{\infty}\left[a + \frac{1}{n}, b\right)$$

$$[a, b) = \bigcap_{n=1}^{\infty}\left(a - \frac{1}{n}, b\right)$$

$$(a, b] = \bigcap_{n=1}^{\infty}\left(a, b + \frac{1}{n}\right)$$

so that $\mathcal{F}_1 = \mathcal{F}_2 = \mathcal{F}_3$. We may argue similarly $\mathcal{F}_4 = \mathcal{F}_5 = \mathcal{F}_6 = \mathcal{F}_7$. Finally

$$(a, b) = (a, \infty) \cap (-\infty, b)$$

and

$$(a, \infty) = \bigcup_{n=1}^{\infty}(a, n)$$

to show these are equal. The measure $\mu$ exists by (5.8.29) and (5.8.26). ∎

**Proposition 5.8.31 (Borel Measure on the Extended Real Line)**
*Let $\mathcal{B}(\mathbb{R}^\sharp)$ be the Borel Measure on the extended real line. Then $\mathcal{B}(\mathbb{R}^\sharp)$ consists of sets of the form*

© David Rufino CC BY-NC-ND 4.0

$$\left\{A \subset \mathbb{R}^\sharp \mid A \cap \mathbb{R} \in \mathcal{B}(\mathbb{R})\right\}$$

*or equivalently*

$$\{A \cup B \mid A \in \mathcal{B}(\mathbb{R}), B \in \mathcal{P}(\{-\infty, \infty\})\}$$

*Furthermore it is generated by any of the following families*

1. $(a, \infty]$
2. $[a, \infty]$
3. $[-\infty, a)$
4. $[-\infty, a]$

*for $a \in D$ where $D \subset \mathbb{R}$ is an additive subgroup containing $\mathbb{Q}$.*

*Proof.* If $U \subset \mathbb{R}$ is open, then it is also open in $\mathbb{R}^\sharp$. Therefore $\mathcal{B}(\mathbb{R}) \subset \mathcal{B}(\mathbb{R}^\sharp)$. Evidently $\mathbb{R}$, $\{\infty\}$, $\{-\infty\} \in \mathcal{B}(\mathbb{R}^\sharp)$ as they are open and closed respectively. This shows that the given family equals $\mathcal{B}(\mathbb{R}^\sharp)$.

Denote by $\mathcal{F}_1, ..., \mathcal{F}_4$ the $\sigma$-algebras generated by the respective families of rays. Note that $\mathcal{F}_1 = \mathcal{F}_4$ and $\mathcal{F}_2 = \mathcal{F}_3$ by taking complements. Further $\{\infty\} \in \mathcal{F}_1 \cap \mathcal{F}_2$ by taking intersections as $a \to \infty$ and similarly $\{-\infty\} \in \mathcal{F}_4 \cap \mathcal{F}_3$. This shows that they also contain either the open rays $(a, \infty)$ or the semi-open rays $[a, \infty)$. Therefore they contain $\mathcal{B}(\mathbb{R})$ by (5.8.30). Then by the characterisation in the first part we see that this equals $\mathcal{B}(\mathbb{R}^\sharp)$. ∎

## 5.8.6 Product Measure

**Definition 5.8.32 (Product of $\sigma$-algebra)**
*Let $(\Omega_1, \mathcal{F}_1), ..., (\Omega_n, \mathcal{F}_n)$ be a family of measurable spaces. Then define the **product measurable space** $(\Omega_1 \times ... \times \Omega_n, \mathcal{F}_1 \otimes ... \otimes \mathcal{F}_n)$ to be given by*

$$\mathcal{F}_1 \otimes ... \otimes \mathcal{F}_n := \sigma(\mathcal{F}_1 \times ... \times \mathcal{F}_n)$$

*and*

$$\mathcal{F}_1 \times ... \times \mathcal{F}_n := \{A_1 \times ... \times A_n \mid A_i \in \mathcal{F}_i\}$$

**Definition 5.8.33 (Product Measure)**
*Let $(\Omega_i, \mathcal{F}_i, \mu_i)$ be a family of $\sigma$-finite measures for $i = 1...n$. Then we say a measure*

$$\mu_1 \otimes ... \otimes \mu_n : \mathcal{F}_1 \otimes ... \otimes \mathcal{F}_n \to \mathbb{R}^\sharp$$

*such that*

$$\mu(A_1 \times ... \times A_n) = \mu_1(A_1) \cdot ... \cdot \mu_{n(A_n)}$$

*is called the **product measure space**.*

We defer proof of existence and uniqueness as it relies on theory of integration, but nevertheless prove the finitely additive case and then the case of $\mathbb{R}^n$ in the next section.

**Lemma 5.8.34**
*Let $\mathcal{R}_1, ..., \mathcal{R}_n$ be semi-rings. Then the family*

$$\mathcal{R}_1 \times ... \times \mathcal{R}_n = \{A_1 \times ... \times A_n \mid A_i \in \mathcal{R}_i\}$$

*is a semi-ring.*

*Proof.* We may reduce to the case of $n = 2$. Then observe that

$$(A_1 \times B_1) \setminus (A_2 \times B_2) = (A_1 \setminus A_2) \times (B_1 \setminus B_2) \sqcup (A_1 \cap A_2) \times (B_1 \setminus B_2) \sqcup (A_1 \setminus A_2) \times (B_1 \cap B_2)$$

∎

**Lemma 5.8.35**
*Let $\mathcal{R}$ be a semi-ring and $A_1, ..., A_n \in \mathcal{R}$ a family of subsets. Then there exists a family of disjoint subsets $\mathcal{S} := \{B_1, ..., B_N\} \subset \mathcal{R}$ such that $\mathcal{S} = \mathcal{S}_1 \cup ... \cup \mathcal{S}_n$ and*

$$A_i = \sqcup_{B \in \mathcal{S}_i} B$$

*Proof.* For a tuple $\varepsilon := (\varepsilon_1, ..., \varepsilon_n)$ for which $\varepsilon_i = \pm 1$ define the family of disjoint sets

© David Rufino CC BY-NC-ND 4.0

$$A^\varepsilon := A_1^{\varepsilon_1} \cap ... \cap A_n^{\varepsilon_n}$$

where $A_i^1 := A_i$ and $A_i^{-1} := A_i^c$. By (5.8.6) $A^\varepsilon$ may be expressed as the disjoint union of some sets in $\mathcal{R}$, say $\mathcal{S}^\varepsilon$. Then $\mathcal{S} = \cup_\varepsilon \mathcal{S}^\varepsilon$ consists of pairwise disjoint sets, where we consider tuples $\varepsilon$ for which at least one entry is positive. Further

$$A_i = \sqcup_{\varepsilon:\varepsilon_i=1} A^\varepsilon$$

therefore we may define $\mathcal{S}_i := \cup_{\varepsilon:\varepsilon_i=1} \mathcal{S}^\varepsilon$. Note throughout we may omit $\varepsilon$ for which $A^\varepsilon = \varnothing$. ■

**Lemma 5.8.36**
*Let $\{(\Omega_i, \mathcal{R}_i, \mu_i)\}_{i=1...n}$ be a collection of finitely additive set functions on semi-rings. The set function*

$$\mu : \mathcal{R}_1 \times ... \times \mathcal{R}_n \to \mathbb{R} \cup \{\infty\}$$
$$A_1 \times ... \times A_n \to \mu_1(A_1)...\mu_{n(A_n)}$$

*is finitely additive, and we write $\mu =: \mu_1 \times ... \times \mu_n$. Moreover if each $\mu_i$ is $\sigma$-finite then so is $\mu$.*

*Proof.* By induction it's enough to consider the case $n = 2$. Suppose that

$$A_i = B_i \times C_i \quad i = 0, ..., n$$
$$A_0 = \sqcup_{i=1}^n A_i$$

for $B_i \in \Omega_1$ and $C_i \in \Omega_2$. By (5.8.35) we may write

$$B_i := \sqcup_{\hat{B} \in \mathcal{S}_i} \hat{B}$$
$$C_i := \sqcup_{\hat{C} \in \mathcal{T}_i} \hat{C}$$

where $\mathcal{S}_i$ and $\mathcal{T}_i$ are finite subsets of $\mathcal{R}_1$ and $\mathcal{R}_2$ respectively. Evidently

$$B_i \times C_i = \sqcup_{(\hat{B},\hat{C}) \in \mathcal{S}_i \times \mathcal{T}_i} \hat{B} \times \hat{C}$$

and $\mathcal{S}_0 = \cup_{i=1}^n \mathcal{S}_i$ and $\mathcal{T}_0 := \cup_{i=1}^n \mathcal{T}_i$. As $\mu_1$ and $\mu_2$ are finitely additive then

$$\mu_1(B_i) = \sum_{\hat{B} \in \mathcal{S}_i} \mu_1\big(\hat{B}\big)$$
$$\mu_2(C_i) = \sum_{\hat{C} \in \mathcal{T}_i} \mu_2\big(\hat{C}\big)$$
$$\sum_{i=1}^n \mu(A_i) = \sum_{i=1}^n \mu_1(B_i)\mu_2(C_i) = \sum_{i=1}^n \sum_{(\hat{B},\hat{C}) \in \mathcal{S}_i \times \mathcal{T}_i} \mu_1\big(\hat{B}\big)\mu_2\big(\hat{C}\big)$$
$$\mu(A_0) = \mu_1(B_0)\mu_2(C_0) = \sum_{(\hat{B},\hat{C}) \in \mathcal{S}_0 \times \mathcal{T}_0} \mu_1\big(\hat{B}\big)\mu_2\big(\hat{C}\big)$$

As the sets $B_i \times C_i$ are disjoint we may deduce that

$$\mathcal{S}_0 \times \mathcal{T}_0 = \sqcup_{i=1}^n \mathcal{S}_i \times \mathcal{T}_i$$

from which the result follows. ■

## 5.8.7 Borel Measure on $\mathbb{R}^n$

**Proposition 5.8.37**
*Let $\mathcal{R}$ be the semi-ring of half-open intervals over $\mathbb{R}^n$. Then the set function*

$$\mu : \mathcal{R} \to [0, \infty]$$

*given by*

$$\mu([a_1, b_1) \times ... \times [a_n, b_n)) = \prod_{i=1}^n (b_i - a_i),$$

*is countably additive and $\sigma$-finite.*

*Proof.* By (5.8.36) $\mu$ is finitely additive. So by (5.8.25) it is sufficient to show that $\mu$ is countably subadditive. Suppose that

© David Rufino CC BY-NC-ND 4.0

$$[\boldsymbol{a}_0, \boldsymbol{b}_0) = \bigcup_{i=1}^{\infty} [\boldsymbol{a}_i, \boldsymbol{b}_i)$$

Then for any positive constants $\boldsymbol{\varepsilon}$, $\boldsymbol{\delta}_1, \boldsymbol{\delta}_2, \ldots$ we have

$$[\boldsymbol{a}_0, \boldsymbol{b}_0 - \boldsymbol{\varepsilon}] \subset \bigcup_{i=1}^{\infty} (\boldsymbol{a}_i - \boldsymbol{\delta}_i, \boldsymbol{b}_i)$$

By the Heine-Borel theorem (5.7.7) closed intervals are compact, so we have

$$[\boldsymbol{a}_0, \boldsymbol{b}_0 - \boldsymbol{\varepsilon}) \subset \bigcup_{i=1}^{N(\boldsymbol{\delta})} [\boldsymbol{a}_i - \boldsymbol{\delta}_i, \boldsymbol{b}_i)$$

Therefore by (5.8.9)

$$\begin{aligned}
\prod_{j=1}^{n} (b_{0j} - a_{0j} - \varepsilon) &\leq \sum_{i=1}^{N(\boldsymbol{\delta})} \prod_{j=1}^{n} (b_{ij} - a_{ij} + \delta_{ij}) \\
&\leq \sum_{i=1}^{N(\boldsymbol{\delta})} \prod_{j=1}^{n} (b_{ij} - a_{ij}) \left(1 + \frac{\delta}{2^i}\right) \\
&\leq \sum_{i=1}^{N(\boldsymbol{\delta})} \prod_{j=1}^{n} (b_{ij} - a_{ij}) \sum_{i=1}^{N(\boldsymbol{\delta})} \left(1 + \frac{\delta}{2^i}\right) \\
&\leq (1 + \delta) \sum_{i=1}^{\infty} \prod_{j=1}^{n} (b_{ij} - a_{ij})
\end{aligned}$$

where we have defined $\delta_{ij} \coloneqq \frac{b_{ij} - a_{ij}}{n} \left(\frac{\delta}{2^i}\right)^{\frac{1}{n-1}}$ and used the inequality $(1 + x)^n \leq 1 + nx^{n-1}$. As this holds for arbitrary $\delta > 0$ then it holds for $\delta = 0$. We may consider the sequence $\varepsilon \coloneqq \frac{1}{m}$ and letting $m \to \infty$ use (4.1.50) and (5.1.21) to deduce the case of $\varepsilon = 0$. This shows that

$$\mu([\boldsymbol{a}_0, \boldsymbol{b}_0)) \leq \sum_{i=1}^{\infty} \mu([\boldsymbol{a}_i, \boldsymbol{b}_i))$$

i.e. $\mu$ is countably subadditive. ∎

**Proposition 5.8.38 (Existence of Borel Measure)**
*Let $\left(\mathbb{R}^k, \mathcal{B}(\mathbb{R}^k)\right)$ be the Borel measurable space. Then there exists a unique $\sigma$-finite measure*

$$\mu : \mathcal{B}(\mathbb{R}^k) \to [0, \infty]$$

*such that*

$$\mu([\boldsymbol{a}, \boldsymbol{b})) = \prod_{i=1}^{k} (b_i - a_i)$$

*Proof.* This is a consequence of (5.8.26) and (5.8.37). ∎

## 5.8.8 Measurable Functions

**Definition 5.8.39**
*Let $(\Omega_1, \mathcal{F}_1)$ and $(\Omega_2, \mathcal{F}_2)$ be measurable spaces. A function $f : \Omega_1 \to \Omega_2$ is said to be **measurable** if*

$$E \in \mathcal{F}_2 \implies f^{-1}(E) \in \mathcal{F}_1$$

*We may write $\mathcal{F}_1/\mathcal{F}_2$-measurable in order to make the $\sigma$-algebras explicit.*

*Similarly a function $f : \Omega \to X$ from a measurable space to a topological space is said to be measurable if it is $\mathcal{F}/\mathcal{B}(X)$ measurable.*

**Proposition 5.8.40**
*The following properties hold*

*a) The composition of measurable functions is measurable*

*b) Suppose $\mathcal{F}_2 = \sigma(\mathcal{C})$ then a function $f : \Omega_1 \to \Omega_2$ is $\mathcal{F}_1/\mathcal{F}_2$-measurable iff*

© David Rufino CC BY-NC-ND 4.0

$$E \in \mathcal{C} \implies f^{-1}(E) \in \mathcal{F}_1$$

*c) A function $f : \Omega \to X$ is measurable precisely when the inverse image of an open set (resp. closed set) is measurable.*

**Corollary 5.8.41 (Criteria for measurablity of a real-valued function)**
*Let $(\Omega, \mathcal{F})$ be a measurable space and $f : \Omega \to \mathbb{R}^\sharp$ a function. Then the following are equivalent*

*a) $f$ is $\mathcal{F}/\mathcal{B}(\mathbb{R}^\sharp)$-measurable*
*b) $f^{-1}((x, \infty]) \in \mathcal{F}$ for all $x \in D$*
*c) $f^{-1}([x, \infty]) \in \mathcal{F}$ for all $x \in D$*
*d) $f^{-1}([-\infty, x)) \in \mathcal{F}$ for all $x \in D$*
*e) $f^{-1}([-\infty, x]) \in \mathcal{F}$ for all $x \in D$*

*where $D \subset \mathbb{R}$ is an additive subgroup containing $\mathbb{Q}$.*

*Proof.* We may combine (5.8.40).b) and (5.8.31). ∎

**Proposition 5.8.42 (Measurable Space Isomorphism)**
*Let $(\Omega_1, \mathcal{F}_1)$ and $(\Omega_2, \mathcal{F}_2)$ be measurable spaces and $f : \Omega_1 \to \Omega_2$ a bijective function. Then the following are equivalent*

*a) $f$ and $f^{-1}$ are measurable*
*b) $f$ induces a bijection $\mathcal{F}_1 \to \mathcal{F}_2$*
*c) $f^{-1}$ induces a bijection $\mathcal{F}_2 \to \mathcal{F}_1$*

*In this case we say that $f : (\Omega_1, \mathcal{F}_1) \xrightarrow{\sim} (\Omega_2, \mathcal{F}_2)$ is a **measurable space isomorphism**.*

**Proposition 5.8.43 (Limits of Measurable Functions)**
*Let $f_n : (\Omega, \mathcal{F}) \to (\mathbb{R}^\sharp, \mathcal{B}(\mathbb{R}^\sharp))$ be a sequence of measurable functions. Define the functions $\sup f_n, \inf f_n, \limsup_n f_n, \liminf_n f_n$ as follows*

$$(\sup f_n)(x) := \sup_n f_n(x)$$
$$(\inf f_n)(x) := \inf_n f_n(x)$$
$$\left(\limsup_n f_n\right)(x) := \limsup_n f_n(x)$$
$$\left(\liminf_n f_n\right)(x) := \liminf_n f_n(x)$$

*Then these functions are $\mathcal{F}/\mathcal{B}(\mathbb{R}^\sharp)$-measurable.*

*Proof.* Observe that

$$\left(\inf_n f_n\right)^{-1}([x, \infty]) = \bigcap_{n=1}^{\infty} f_n^{-1}([x, \infty])$$

so $\inf f_n$ is measurable by (5.8.41). Further $\sup_n f_n = -\inf_n(-f_n)$. ∎

**Proposition 5.8.44**
*Let $f, g : (\Omega, \mathcal{F}) \to (\mathbb{R}^\sharp, \mathcal{B}(\mathbb{R}^\sharp))$ be measurable functions. Then the following functions are measurable*

*a) $f \pm g$*
*b) $\lambda f$ is measurable for $\lambda \in \mathbb{R}$*
*c) $f \cdot g$*
*d) $\max(f, g)$, $\min(f, g)$*

*Note when $f \pm g, fg$ is not well-defined (e.g. $\infty - \infty$ or $-\infty \times -\infty$) then we choose an arbitrary, but fixed, sentinel value.*

*Proof.* We make use of (5.8.41) throughout.

a) Define $h(\omega) := f(\omega) + g(\omega)$ and let $A$ be the set on which the sum is well-defined. Then it is evidently measurable and

$$\begin{aligned} h^{-1}((x, \infty]) &= \{\omega \in B_x^c \mid f(\omega) + g(\omega) > x\} \cup B_x \\ &= \{\omega \in A \mid f(\omega) + g(\omega) > x\} \cup B_x \\ &= \left(\bigcup_{r \in \mathbb{Q}} \{\omega \in A \mid f(\omega) > r\} \cap \{\omega \in A \mid g(\omega) > x - r\}\right) \cup B_x \end{aligned}$$

© David Rufino CC BY-NC-ND 4.0

where $B_x = \varnothing$ or $A^c$ according to if the sentinel value of $f+g$ is $\leq x$. For the final equality the reverse inclusion is immediate. Conversely suppose $f(\omega)+g(\omega) > x$ then by (5.1.29) there exists some $r \in \mathbb{Q}$ such that $f(\omega) > r > x - g(\omega)$. The case $f-g$ follows immediately from b).

b) Observe that

$$(\lambda f)^{-1}((x,\infty]) = \begin{cases} \left(\frac{x}{\lambda},\infty\right] & \lambda > 0 \\ \left[-\infty,\frac{x}{\lambda}\right) & \lambda < 0 \\ \Omega & \lambda = 0, x < 0 \\ \varnothing & \lambda = 0, x \geq 0 \end{cases}$$

c) First consider the case $f^2$ and observe

$$(f^2)^{-1}((x,\infty]) = \begin{cases} f^{-1}(\{-\infty,\infty\}) \cup f^{-1}\left(\left(\sqrt{x},\infty\right]\right) \cup f^{-1}\left(\left[-\infty,-\sqrt{x}\right)\right) & x \geq 0 \\ \Omega & x < 0 \end{cases}$$

Then we may deduce the general case because

$$(fg)(\omega) = \begin{cases} \frac{(f(\omega)+g(\omega))^2-(f(\omega)-g(\omega))^2}{4} & f(\omega), g(\omega) \notin \{-\infty,\infty\} \\ \infty & (f(\omega),g(\omega)) \in \{(\infty,\infty),(-\infty,-\infty)\} \\ -\infty & (f(\omega),g(\omega)) \in \{(\infty,-\infty),(-\infty,\infty)\} \end{cases}$$

d) We may see this as a special case of (5.8.43)

■

## 5.8.9 Integration over a Measure

© David Rufino CC BY-NC-ND 4.0

## 5.9 Differential Calculus on Banach Spaces

The main reference for this section is [Car71], along with [Lan12], in a generality which is almost certainly not required for the purposes of these notes.

### 5.9.1 Total Derivatives

**Definition 5.9.1**
*Let $U \subset X$ be an open neighbourhood of $0$ and $\varphi : U \to Y$ a function. We say $\varphi$ is **sublinear** if*

$$\forall \varepsilon > 0 \exists \delta \text{ s.t. } \|h\| < \delta \implies (h \in U) \wedge (\|\varphi(h)\| < \varepsilon \|h\|)$$

*Note sublinear functions are closed under linear combinations, and the property is unchanged under equivalent norms.*

**Definition 5.9.2 (Total Derivative)**
*Let $U \subset X$ be an open subset of a Banach space, $Y$ a Banach space and $x \in U$. We say that a map $\varphi : U \to Y$ is differentiable at $x$ if there exists a continuous linear map $D\varphi(x) \in L(X, Y)$ (**derivative** at $x$) such that*

$$\varphi(x + h) - \varphi(x) - D\varphi(x)(h) \text{ is sublinear in } h$$

*We say it is simply **differentiable** if this holds for every point $x \in U$ and write $D\varphi$ for the corresponding function $U \to L(X, Y)$.*

*If $D\varphi$ is also continuous then we say that $\varphi$ is **continuously differentiable** or $C^1$.*

The following result shows that the total derivative is necessarily unique.

**Proposition 5.9.3**
*Let $\alpha : X \to Y$ be a linear map which is also sublinear. Then $\alpha = 0$.*

*Proof.* Consider $x \in X$ then for every $\varepsilon > 0$ there exists $\delta$ such that $\|h\| < \delta \implies \|\alpha(h)\| < \varepsilon \|h\|$. Define $\lambda := \frac{\delta}{2\|x\|}$. Then evidently $\|\alpha(\lambda x)\| < \varepsilon \|\lambda x\|$ whence $\|\alpha(x)\| < \varepsilon \|x\|$. As this holds for every $\varepsilon$ we see $\|\alpha(x)\| = 0$ and $\alpha(x) = 0$. ∎

**Proposition 5.9.4**
*Let $f : U \to Y$ be a differentiable map and $\alpha : Y \to Z$ a linear map. Then*

$$D(\alpha \circ f)(x) = \alpha \circ Df(x)$$

**Proposition 5.9.5**
*Let $U \subset K$ an open set and $\varphi : U \to Y$ a function. Then the following are equivalent*

*a) $\varphi$ is (continuously) differentiable in the sense of (5.9.2)*
*b) $\varphi$ is (continuously) differentiable in the sense of (5.5.1)*

*Furthermore we may identify the derivatives as follows*

$$D\varphi(t)(\lambda) = \varphi'(t)\lambda$$

**Proposition 5.9.6 (Chain Rule)**
*Let $U \subset X$ and $V \subset Y$ be open sets and $\varphi : U \to Y$ and $\psi : V \to Z$ (continuously) differentiable functions such that $\varphi(U) \subset V$. Then $\psi \circ \varphi$ is (continuously) differentiable and*

$$(D(\psi \circ \varphi))(x) = (D\psi)(\varphi(x)) \circ (D\varphi)(x)$$

*In the case $X = K$ we have*

$$(\psi \circ \varphi)'(t) = (D\psi)(\varphi(t))(\varphi'(t))$$

**Proposition 5.9.7 (Product Rule I)**
*Let $U \subset X$ and $V \subset Y$ open sets, $\varphi : U \to X'$, $\psi : V \to Y'$ (continuously) differentiable maps and $\Psi : X' \times Y' \to Z$ a $C^1$ bilinear map. Then define*

$$(\varphi \times_\Psi \psi)(x, y) := \Psi(\varphi(x), \psi(y))$$

*is (continuously) differentiable with derivative*

$$D(\varphi \times_\Psi \psi)(x, y)(h, k) = \Psi(D\varphi(x)(h), \psi(y)) + \Psi(\varphi(x), D\psi(y)(k))$$

© David Rufino CC BY-NC-ND 4.0

**Corollary 5.9.8**
*Let $X, Y$ be normed vector spaces, $U \subset X$ an open subset and $\varphi, \psi : X \to Y$ (continuously) differentiable at $x$. Then both $\varphi + \psi$ and $\lambda\varphi$ are (continuously) differentiable at $x$ and*

$$D(\varphi + \psi)(x) = D(\varphi)(x) + D(\psi)(x)$$
$$D(\lambda\varphi)(x) = \lambda D(\varphi)(x)$$

*If $Y$ is also a Banach algebra then $\varphi \cdot \psi$ is (continuously) differentiable at $x$ and*

$$D(\varphi \cdot \psi)(x)(h) = D(\varphi)(x)(h) \cdot \psi(x) + \varphi(x) \cdot D(\psi)(x)(h)$$

**Corollary 5.9.9 (Product Rule II)**
*Let $U \subset X$ be an open set, and $\varphi : U \to K$ and $\psi : U \to Y$ differentiable maps. Then*

$$D(\varphi \cdot \psi)(x)(h) = D\varphi(x)(h)\psi(x) + \varphi(x)D\psi(x)(h)$$

*If $\varphi, \psi$ are $C^1$ then so is $\varphi \cdot \psi$.*

**Definition 5.9.10 (Higher-Order Derivatives)**
*Let $f : U \to Y$ we say that $f$ is $n$-times **continuously differentiable** (or $C^n$) if it is differentiable and $Df$ is $C^{n-1}$ (where $C^0$ means simply continuous). In this case for $k \leq n$ we may define the $k$-fold derivative inductively*

$$D^k f := D\big(D^{k-1} f\big) : U \longrightarrow L(X, \cdots, L(X, Y))$$

*which is by definition continuous.*

**Proposition 5.9.11**
*Let $U \subset K$ be an open subset and $f : U \to Y$ a function. Then $f$ is $n$-times differentiable in the sense of (...) if and only if it is $n$-times differentiable in the sense of (5.9.10). In this case*

$$(D^n f)(x)(h_1, \cdots, h_n) = f^{(n)}(x) \cdot h_1 \cdots h_n$$

**Proposition 5.9.12 (Mean Value Theorem)**
*Let $U \subset X$ be an open subset and $f : U \to Y$ a $C^1$ map. Suppose that $h \in B(x; r) \subset U$. Then*

$$f(x + h) = f(x) + \int_0^1 Df(x + th)(h)\,\mathrm{d}t$$

*Proof.* Let $G(t) := f(x + th)$ be defined as a function on $[0, 1] \to Y$. Then by (5.9.6) $G'(t) = Df(x + th)(h)$ is continuous, because evidently $t \to x + th$ is continuously differentiable with scalar derivative $h$. The result follows by applying (5.8.20) to $G(t)$. ∎

## 5.9.2 Partial Derivatives

**Lemma 5.9.13**
*Let $X_1, \cdots, X_n$ be Banach spaces and for every $x \in X$ define the injection*

$$\lambda_j(x) : X_j \longrightarrow X_1 \times \cdots \times X_n$$
$$x' \to \big(x_1, \cdots, x_{j-1}, x', x_{j+1}, \cdots, x_n\big)$$

*Then $\lambda_j$ is differentiable at every $x_j \in X_j$ and $\big(D\lambda_j(x)\big)\big(x_j\big) = \mu_j$ where*

$$\mu_j(h) = (0, \cdots, h, \cdots, 0)$$

*Proof.* We have $\lambda_j(x)(x' + h') - \lambda_j(x) = \mu_j(h')$ and $\mu_j$ is linear. ∎

**Proposition 5.9.14 (Criteria for Total Differentiability)**
*Let $X_1, \cdots, X_n, Y$ be Banach spaces and $U \subset X := X_1 \times \cdots \times X_n$ an open subset. Consider a continuous map*

$$\varphi : U \to Y$$

*The following are equivalent*

*a) The map $\varphi$ is continuously differentiable (with respect to the sup norm on the product)*
*b) For every $x \in U$ define $V_j := \lambda_j(x)^{-1}(U) \subset X_j$ an open neighbourhood of $x_j$*
*i) The map*

© David Rufino CC BY-NC-ND 4.0

$$\varphi_j(x) := \varphi \circ \lambda_j(x) : V_j \to Y$$

*is differentiable for all* $j = 1 \cdots n$. *Denote the* $j$*-th partial derivative at* $x \in U$ *by*

$$D_j\varphi : U \to L\big(X_j, Y\big)$$
$$x_j \to h \to \big(D\varphi_j(x)\big)\big(x_j\big)(h)$$

*ii) The map* $D_j$ *is continuous for all* $j = 1 \cdots n$

*Furthermore in this case*

$$D\varphi(x)(h) = \sum_{j=1}^{n} \big(D_j\varphi\big)(x)\big(h_j\big)$$

*Proof.* **a) $\implies$ b)** By the chain rule (5.9.6) and (5.9.13) we have $\varphi_j$ is differentiable and

$$\big(D\varphi_j(x)\big)\big(x_j\big) = (D\varphi)(x) \circ \big(D\lambda_j(x)\big)\big(x_j\big) = (D\varphi)(x) \circ \mu_j$$

**b) $\implies$ a)** We consider the case $n = 2$. Observe

$$\begin{aligned}
\varphi(x_1 + h_1, x_2 + h_2) - \varphi(x_1, x_2) &= \varphi(x_1 + h_1, x_2 + h_2) - \varphi(x_1 + h_1, x_2) + \varphi(x_1 + h_1, x_2) - \varphi(x_1, x_2) \\
&= \int_0^1 D_2\varphi(x_1 + h_1, x_2 + th_2)(h_2)\,\mathrm{d}t + \int_0^1 D_1\varphi(x_1 + th_1, x_2)(h_1)\,\mathrm{d}t \\
&= D_2\varphi(x_1, x_2)(h_2) + D_1\varphi(x_1, x_2)(h_1) \\
&+ \int_0^1 [D_2\varphi(x_1 + h_1, x_2 + th_2)(h_2) - D_2\varphi(x_1, x_2)(h_2)]\,\mathrm{d}t \\
&+ \int_0^1 [D_1\varphi(x_1 + th_1, x_2)(h_1) - D_1\varphi(x_1, x_2)(h_1)]\,\mathrm{d}t
\end{aligned}$$

Then

$$\begin{aligned}
\left\| \int_0^1 [D_2\varphi(x_1 + h_1, x_2 + th_2)(h_2) - D_2\varphi(x_1, x_2)(h_2)]\,\mathrm{d}t \right\| &\leq \int_0^1 \|D_2\varphi(x_1 + h_1, x_2 + th_2)(h_2) - D_2\varphi(x_1, x_2)(h_2)\|\,\mathrm{d}t \\
&\leq \int_0^1 \|D_2\varphi(x_1 + h_1, x_2 + th_2) - D_2\varphi(x_1, x_2)\|\|h_2\|\,\mathrm{d}t \\
&\leq \int_0^1 \|D_2\varphi\|\|(h_1, th_2)\|\|h_2\|\,\mathrm{d}t \\
&\leq \|D_2\varphi\|\|h\|^2
\end{aligned}$$

and similarly for the second integral. Therefore $\varphi$ is differentiable at $(x_1, x_2)$ with derivative

$$D\varphi(x_1, x_2)(h) = D_1\varphi(x_1, x_2)(h_1) + D_2\varphi(x_1, x_2)(h_2)$$

as required.

We have $h = \sum_{j=1}^{n} \mu_j\big(h_j\big)$ and so by the first part

$$\begin{aligned}
D\varphi(x)(h) &= \sum_{j=1}^{n} \big(D\varphi(x)\big)\big(\mu_j\big(h_j\big)\big) \\
&= \sum_{j=1}^{n} \big(D\varphi_j(x)\big)\big(x_j\big)\big(h_j\big) \\
&= \sum_{j=1}^{n} \big(D_j\varphi\big)(x)\big(h_j\big)
\end{aligned}$$

■

**Proposition 5.9.15**
*Let* $U \subset X$ *be an open set and*

$$\varphi : U \to Y_1 \times \cdots \times Y_m$$

© David Rufino CC BY-NC-ND 4.0

*Then $\varphi$ is differentiable (resp. $C^1$) iff $\pi_i \circ \varphi$ is differentiable (resp. $C^1$) for $i = 1 \cdots m$.*

**Proposition 5.9.16 (Jacobian Matrix)**
*Let $X, Y$ be finite-dimensional Banach spaces with bases $\{e_1, \cdots, e_n\}, \{f_1, \cdots, f_m\}$ such that $U \subset X$ is open and $\varphi : U \to Y$ a map. Then the following are equivalent*

*a) $\varphi$ is $C^1$*
*b) For every $x \in U$, $j = 1 \cdots n$, $i = 1 \cdots m$ the map*

$$g_{ij}(x) : V \to K$$
$$t \to (f_i^\vee \circ \varphi)(x + te_j)$$

*is continuously differentiable on the open set $V = \{t \in K \mid x + te_j \in U\}$.*

*In this case define $\frac{\partial \varphi_i}{\partial e_j}(x) \coloneqq g_{ij}(x)'(0)$. Then the linear map $D\varphi(x)$ has the following matrix representation*

$$J_\varphi(x) \coloneqq [D\varphi(x)] = \left(\frac{\partial \varphi_i}{\partial e_j}(x)\right)$$

*which we refer to as the **Jacobian Matrix**.*

*When $Y = K$ then $\{1_K\}$ is a basis and we may simply write $\frac{\partial \varphi}{\partial e_j} : U \to K$ as the partial derivative.*

*Proof.* Let $\psi_j : K \xrightarrow{\sim} \langle e_j \rangle$ be the canonical linear isomorphism and $\lambda_j(x) : \langle e_j \rangle \to X$ by the function defined in (5.9.13). Then

$$f_i^\vee \circ \varphi \circ \lambda_j(x) = g_{ij}(x) \circ \psi_j^{-1}$$

Suppose *b*) is satisfied. Because $\psi_j^{-1}$ is continuously differentiable the hypotheses of (5.9.14) b) are satisfied and we conclude that $f_i^\vee \circ \varphi$ is $C^1$. Then by (5.9.15) so is $\varphi$.

Conversely if $\varphi$ is $C^1$, then so is $f_i^\vee \circ \varphi$. And we may use the same identity and the chain rule to deduce $g_{ij}(x)$ is continuously differentiable.

By the same result we have

$$(f_i^\vee \circ D\varphi(x))(x_j) = D(f_i^\vee \circ \varphi)(x)(e_j) = D(f_i^\vee \circ \varphi \circ \lambda_j(x))(e_j)(e_j) = g_{ij}(x)'(0)e_j$$

which shows that $J_\varphi(x)$ has the required form (...). ∎

**Proposition 5.9.17**
*Let $X, Y, Z$ be finite-dimensional Banach spaces with bases $\mathcal{B}_1, \mathcal{B}_2, \mathcal{B}_3$ respectively. Suppose $U \subset X, V \subset Y$ are open sets and $\varphi : U \to Y, \psi : V \to Z$ are $C^1$ maps such that $\varphi(U) \subset V$. Then $\psi \circ \varphi$ is $C^1$ and*

$$J_{\psi \circ \varphi}(x) = J_\psi(\varphi(x)) J_\varphi(x)$$

## 5.9.3 Second Derivative

**Definition 5.9.18**
*For a map $\alpha : X \times X \to Y$ we say that $\alpha(v, w)$ is **sublinear** if for all $\varepsilon > 0$ there exists $\delta > 0$ such that*

$$\|(v, w)\| < \delta \implies \|\alpha(v, w)\| < \varepsilon \|v\| \|w\|$$

*where the norm on $X \times X$ is the product norm.*

As before we have the following uniqueness property

**Lemma 5.9.19**
*Let $\alpha : X \times X \to Y$ be a bilinear map which is also sublinear. Then $\alpha = 0$.*

**Proposition 5.9.20 (Second derivative is symmetric)**
*Let $f : U \to Y$ be a $C^2$ map. Then for all $x \in U$ we have $D^2 f(x)$ is the unique bilinear map $X \times X \to Y$ such that*

$$f(x + v + w) - f(x + w) - f(x + v) + f(x) - D^2 f(x)(v, w) \text{ is sublinear}$$

*Furthermore it is symmetric.*

*Proof.* Uniqueness follows from (5.9.19) and from this it immediately follows that $D^2 f(x)$ is symmetric. Therefore we only need to show the required property. Let $v, w \in X$ be such that $x + tv + sw \in U$ for all $0 \leq s, t \leq 1$. Then define

© David Rufino CC BY-NC-ND 4.0

$$g(x) := f(x+v) - f(x)$$

By (...) applied to $g$ and $Df$ we find

$$\begin{aligned} g(x+w) - g(x) &= \int_0^1 [Df(x+v+tw)(w) - Df(x+tw)(w)]\,\mathrm{d}t \\ &= \int_0^1 \int_0^1 D^2 f(x+sv+tw)(v)\,\mathrm{d}s(w)\,\mathrm{d}t \\ &= D^2 f(x)(v,w) + \int_0^1 \int_0^1 [D^2 f(x+sv+tw)(v) - D^2 f(x)(v)]\,\mathrm{d}s(w)\,\mathrm{d}t \end{aligned}$$

Let $\psi(v,w)$ denote the integral then

$$\begin{aligned} \|\psi(v,w)\| &\le \int_0^1 \int_0^1 \left\|D^2 f(x+sv+tw)(v) - D^2 f(x)(v)\right\| \mathrm{d}s\|w\|\,\mathrm{d}t \\ &\le \int_0^1 \int_0^1 \left\|D^2 f(x+sv+tw) - D^2 f(x)\right\| \mathrm{d}s\,\mathrm{d}t \cdot \|v\|\|w\| \\ &\le \sup_{0\le s,t\le 1} \left\|D^2 f(x+sv+tw) - D^2 f(x)\right\| \|v\|\|w\| \end{aligned}$$

where the estimate is finite by continuity of $D^2 f$ and the Boundedness Theorem (5.7.8). Note that $\|sv+tw\| \le \|v\| + \|w\| \le 2\|(v,w)\|$. Therefore we may use the continuity of $D^2 f$ to show that $\psi$ is sublinear as required. ∎

**Proposition 5.9.21 (Symmetry of partial derivatives)**
*Let $X = X_1 \times \cdots \times X_n$, $\varphi : U \to Y$ be a $C^2$ map and $x \in U$. Then*

$$D^2\varphi(x)\big((0,\cdots,h_i,\cdots,0),(0,\cdots,h_j,\cdots,0)\big) = D_i D_j \varphi(x)\big(h_i,h_j\big)$$

*In particular*

$$D_i D_j \varphi(x)\big(h_j,h_i\big) = D_j D_i \varphi(x)\big(h_i,h_j\big)$$

*Proof.* The first equation follows from (5.9.14) and the second from (5.9.20). ∎

**Proposition 5.9.22 (Hessian Matrix)**
*Let $X$ be a finite-dimensional Banach spaces with basis $\{e_1,\cdots,e_n\}$, $U \subset X$ an open subset and $\varphi : U \to K$ a $C^2$ map. Then the second order partial derivatives $\frac{\partial^2\varphi}{\partial e_i \partial e_j} := \frac{\partial}{\partial e_j}\frac{\partial\varphi}{\partial e_i} : U \to K$ exist for all $1 \le i,j \le n$.*

*Denote by*

$$H_\varphi := \left(\frac{\partial^2\varphi}{\partial e_i \partial e_j}\right)_{1\le i,j\le n}$$

*the **Hessian Matrix** which is the matrix of the bilinear form $D^2\varphi$ with respect to the given basis, and in particular symmetric.*

## 5.9.4 Differentiation under Integral Sign

**Proposition 5.9.23**
*Let $A \subset X$ be a compact subset of a normed vector space and $f : A \to Y$ a continuous map. Then $f$ is uniformly continuous. That is for all $\varepsilon > 0$ there exists $\delta > 0$ such that*

$$\|x-y\| < \delta \implies \|f(x) - f(y)\| < \varepsilon \quad \forall x,y \in A$$

*Proof.* Given $\varepsilon > 0$ for each $x \in A$ by continuity there exists a real number $\delta(x) > 0$ such that $\|y-x\| < \delta(x) \implies \|f(y) - f(x)\| < \varepsilon$. Evidently the balls $B\big(x, \frac{\delta(x)}{2}\big)$ cover $A$, so we may find a finite subcover of balls centered at $x_1,\cdots,x_n$. Let $\delta := \min\Big(\frac{\delta(x_1)}{2},\cdots,\frac{\delta(x_n)}{2}\Big)$.

**TODO** ∎

**Proposition 5.9.24**
*Let $I = [a,b]$ be a real interval, $U \subset X$ an open subset of a finite-dimensional normed vector space and $f : I \times U \to Y$ be a continuous map such that $D_2 f$ exists and is continuous. Define*

© David Rufino CC BY-NC-ND 4.0

$$g(x) = \int_a^b f(t,x)\,\mathrm{d}t$$

*Then $g$ is differentiable on $U$ with derivative given by*

$$Dg(x)(h) = \int_a^b D_2 f(t,x)(h)\,\mathrm{d}t$$

*Proof.* Consider a point $x \in U$ with a compact neighbourhood $N \subset U$ (...). Consider $\lambda \in L(X,Y)$ determined by

$$\lambda(h) = \int_a^b (D_2 f)(t,x)(h)\,\mathrm{d}t$$

Then by (5.9.12)

$$\begin{aligned} g(x+h) - g(x) - \lambda(h) &= \int_a^b [f(t,x+h) - f(t,x) - (D_2 f)(t,x)(h)]\,\mathrm{d}t \\ &= \int_a^b \left[\int_0^1 (D_2 f)(t,x+uh)(h)\,\mathrm{d}u - (D_2 f)(t,x)(h)\right]\mathrm{d}t \\ &= \int_a^b \left[\int_0^1 [(D_2 f)(t,x+uh)(h) - (D_2 f)(t,x)(h)]\,\mathrm{d}u\right]\mathrm{d}t \end{aligned}$$

For some $\delta_1$ we have $B(x,\delta_1) \subset N$. Therefore if $\|h\| < \delta_1$ then $x + uh \in N$ for all $u \in [0,1]$. Observe that

$$\|(t,x+uh) - (t,x)\| = \|(0,uh)\| = |u|\|h\|$$

By uniform continuity (5.9.23) on the compact set $[a,b] \times N$ (4.1.115) there is some $\delta_2$ such that

$$(x+uh) \in N \wedge |u|\|h\| < \delta_2 \implies \|(D_2 f)(t,x+uh) - (D_2 f)(t,x)\| < \varepsilon$$

Therefore for $\|h\| < \min(\delta_1, \delta_2)$ we have

$$\begin{aligned} \|g(x+h) - g(x) - \lambda(h)\| &\leq \int_a^b \int_0^1 \|(D_2 f)(t,x+uh)(h) - (D_2 f)(t,x)(h)\|\,\mathrm{d}u\,\mathrm{d}t \\ &\leq (b-a)\|h\|\varepsilon \end{aligned}$$

By definition this means $g(x)$ is differentiable with the given derivative. ∎

**Corollary 5.9.25**
*Let $I = [a,b]$ be a real interval, $U \subset K$ an open subset of a complete valued field $K$ and $f(t,x) : I \times U \to Y$ a continuous map such that $\frac{\partial f}{\partial x} : I \times U \to K$ exists and is continuous. Define*

$$g(x) = \int_a^b f(t,x)\,\mathrm{d}t$$

*Then $g$ is differentiable on $U$ (in the sense of ....) and*

$$g'(x) = \int_a^b \frac{\partial f}{\partial x}(t,x)\,\mathrm{d}t$$

*Proof.* This follows because $\frac{\partial f}{\partial x} = (D_2 f)(x)(1_K)$ and $g'(x) = (Dg)(x)(1_K)$. ∎

## 5.9.5 Taylor's Theorem

**Lemma 5.9.26 (Integration by Parts)**
*Let $U : [0,1] \to K$ and $V : [0,1] \to X$ be $C^1$ functions. Then*

$$\int_0^1 U'(t)V(t)\,\mathrm{d}t = U(1)V(1) - U(0)V(0) - \int_0^1 U(t)V'(t)\,\mathrm{d}t$$

*Proof.* Let $H(t) := U(t)V(t)$ then by (5.9.9) $H'(t) = U'(t)V(t) + U(t)V'(t)$ is continuous. The result then follows from (5.8.20). ∎

© David Rufino CC BY-NC-ND 4.0

**Corollary 5.9.27 (Integration by Parts II)**
*Let $G : [0,1] \to X$ be a $C^1$ function. Then*

$$\int_0^1 G(t)\,\mathrm{d}t = G(0) + \int_0^1 (1-t)G'(t)\,\mathrm{d}t$$

**Proposition 5.9.28 (Taylor's Theorem)**
*Let $U \subset X$ be an open set and $f : U \to Y$ a $C^2$ map. Suppose $x \in U$, $B(x;r) \subset U$ and $\|h\| < r$. Then*

$$f(x+h) = f(x) + Df(x)(h) + \int_0^1 (1-t)D^2 f(x+th)(h,h)\,\mathrm{d}t$$

*Proof.* Let $G(t) = Df(x+th)(h) : [0,1] \to Y$ then

$$G(t) = \mathrm{ev}_h \circ Df \circ (x+th)$$

By (5.9.4)

$$D(\mathrm{ev}_h \circ Df)(x) = \mathrm{ev}_h \circ D^2 f(x)$$

whence by (5.9.6) $G(t)$ is $C^1$ with scalar derivative

$$G'(t) = (\mathrm{ev}_h \circ D^2 f(x+th))(h) = D^2 f(x+th)(h,h)$$

Recall by the Mean Value Theorem (5.9.12) and Integration by Parts (5.9.27)

$$\begin{aligned} f(x+h) &= f(x) + \int_0^1 G(t)\,\mathrm{d}t \\ &= f(x) + G(0) + \int_0^1 (1-t)G'(t)\,\mathrm{d}t \end{aligned}$$

which is the required result. ∎

**Corollary 5.9.29 (Scalar form of Taylor's Theorem)**
*Let $X$ be a finite-dimensional Banach space with basis $\{e_1, \cdots, e_n\}$, $U \subset X$ an open subset and $f : U \to K$ a $C^2$ map. Suppose $x \in U$ and $y \in B(x;r) \subset U$ then*

$$f(y) = f(x) + \sum_{i=1}^n (y_i - x_i)\frac{\partial f}{\partial e_i}(x) + \sum_{i,j=1}^n (y_i - x_i)(y_j - x_j)R_{ij}$$

*with*

$$R_{ij}(y;x) = \int_0^1 (1-t)\frac{\partial^2 f}{\partial e_i \partial e_j}(x + t(y-x))\,\mathrm{d}t$$

*If $f$ is $C^{2+p}$ (resp. $C^\infty$) then $R_{ij}$ is $C^p$ (resp. $C^\infty$) on $B(x;r)$.*

## 5.9.6 Differential Forms on Banach Spaces

**Definition 5.9.30**
*Let $X, Y$ be Banach spaces and $U \subset X$ an open subset. A continuous mapping*

$$\omega : U \longrightarrow L_a^p(X,Y)$$

*is called a **differential form of degree** $p$ or a **differential $p$-form***

*We say that $\omega$ is $n$-times continuously differentiable if it is so with respect to the canonical norm on $L_a^p(X,Y)$.*

**Proposition 5.9.31**
*Let $X$ be a finite-dimensional vector space with basis $e_1, \cdots, e_k$ and $U \subset X$ an open subset. For each $i = 1 \cdots k$ define the mapping*

$$\mathrm{d}x_i : U \to L_a^1(X,K)$$

*which assigns the dual map $e_i^\vee$. Then for every differential $p$-form $\omega$ we have*

© David Rufino CC BY-NC-ND 4.0

$$\omega(x) = \sum_{i=1}^{n} c_i(x)\,\mathrm{d}x_i$$

*for some unique functions* $c_i : U \to K$.

© David Rufino CC BY-NC-ND 4.0

# 5.10 Complex Analysis

## 5.10.1 Cauchy-Riemann Equations

**Proposition 5.10.1 (Isometry between $CC$ and $RR^2$)**
*The canonical map $\chi : \mathbb{C} \to \mathbb{R}^2$ is a Banach space isometry (using the square norm $\|\cdot\|_2$). Furthermore define*

$$\begin{aligned}\mathrm{Re}(z) &:= \pi_1\chi(z)\\ \mathrm{Im}(z) &:= \pi_2\chi(z)\end{aligned}$$

*then*

$$z = \mathrm{Re}(z)\cdot 1 + \mathrm{Im}(z)\cdot i$$

*where $\mathbb{C}$ is viewed as an $\mathbb{R}$ vector space. One may also verify that complex multiplication may be represented in matrix form*

$$\chi(z\cdot w) = \begin{pmatrix}\mathrm{Re}(z) & -\mathrm{Im}(z)\\ \mathrm{Im}(z) & \mathrm{Re}(z)\end{pmatrix}\chi(w)^T$$

**Proposition 5.10.2 (Cauchy-Riemann Equations)**
*Let $U \subset \mathbb{C}$ and $f : U \to \mathbb{C}$ a function. There exists unique functions $u, v : \chi(U) \to \mathbb{R}$ such that*

$$f(x+yi) = u(x,y) + v(x,y)i$$

*Further the following are equivalent*

*a) $f(z)$ is continuously differentiable (as a complex function)*
*b) $u, v$ are $C^1$ and satisfy the relations*

$$\begin{aligned}\frac{\partial u}{\partial x}(x,y) &= \frac{\partial v}{\partial y}(x,y)\\ \frac{\partial u}{\partial y}(x,y) &= -\frac{\partial v}{\partial x}(x,y)\end{aligned}$$

*Proof.* We may define $F : \chi(U) \to \mathbb{R}^2$ by $F = \chi \circ f \circ \chi^{-1}$ and $u = \pi_1 \circ F$ and $v = \pi_2 \circ F$.

**a) $\implies$ b)** Fix $z = x + yi \in U$ and $f'(z) = a + bi$. Then for sufficiently small $w = h + ki$

$$f(z+w) - f(z) - f'(z)w$$

is sublinear in $w$. By assumption $F$ is $C^1$ and has Jacobian matrix (...) at $(x,y)$ equal to

$$J_F(x,y) := \begin{pmatrix}\frac{\partial u}{\partial x}(x,y) & \frac{\partial u}{\partial y}(x,y)\\ \frac{\partial v}{\partial x}(x,y) & \frac{\partial v}{\partial y}(x,y)\end{pmatrix}$$

and $Df(x,y)(h,k) = J_F(x,y)\begin{pmatrix}h\\k\end{pmatrix}^T$. However by conjugating the above equation by $\chi$ we find that

$$F(x+h, y+k) - F(x,y) - \begin{pmatrix}a & -b\\ b & a\end{pmatrix}\begin{pmatrix}h\\k\end{pmatrix}^T$$

(where $f'(z) = a + bi$) is sublinear (and using the fact $\chi$ is an isometry). Therefore the result then follows from the uniqueness of the total derivative $Df(x,y)$, and hence Jacobian $J_F$.

**b) $\implies$ a)** Conversely if the conditions hold then we may define $f'(z) := \frac{\partial u}{\partial x}(\chi(z)) + \frac{\partial v}{\partial x}(\chi(z))i$ which is evidently continuous. Then by using matrix representation of complex multiplication (5.10.1) we find

$$\chi[f(z+w) - f(z) - f'(z)w] = F(\chi(z+w)) - F(\chi(z)) - J'(\chi(z))\chi(w)^T$$

where

$$J'(x,y) = \begin{pmatrix}\frac{\partial u}{\partial x}(x,y) & -\frac{\partial v}{\partial x}(x,y)\\ \frac{\partial v}{\partial x}(x,y) & \frac{\partial u}{\partial x}(x,y)\end{pmatrix}$$

The conditions mean precisely that $J'(x,y) = J(x,y)$. As $\chi$ is an isometry we find that $f(z+w) - f(z) - f'(z)w$ is sublinear in $w$ as required. ∎

© David Rufino CC BY-NC-ND 4.0

## 5.10.2 Holomorphic Functions

**Definition 5.10.3 (Holomorphic Function)**
*Let $U \subset \mathbb{C}$ be an open subset and $f : U \to \mathbb{C}$ a function. Then we say that $f$ is **holomorphic** if it is infinitely (complex) differentiable (5.10.2). For a holomorphic function every derivative is automatically continuous (5.5.3).*

*If $f$ is injective and $f(U)$ is open we say that $f$ is **biholomorphic** if its inverse is also holomorphic. A biholomorphic functions is a homeomorphism onto its image (5.5.3).*

*Trivially if $f$ is holomorphic then so is $f|_V$ for any open subset $V \subset U$. Conversely if $U = \bigcup_{\alpha\in\mathcal{A}} V_\alpha$ and $f|_{V_\alpha}$ are holomorphic for all $\alpha \in \mathcal{A}$, then so is $f$.*

**Proposition 5.10.4**
*Let $U \subset \mathbb{C}$ be an open subset. Then the set of holomorphic functions on $U$ forms a $\mathbb{C}$-algebra.*

*For $f : U \to \mathbb{C}$ non-constant then $D(f) := \{x \in U \mid f(x) \neq 0\}$ is an open subset and $\left(f|_{D(f)}\right)^{-1}$ is a holomorphic function.*

*Proof.* This follows from (5.5.4), provided we show that $D(f)$ is open. For $\mathbb{C}$ is Hausdorff (5.4.5) so $\{0\}$ is closed (4.1.48). Therefore by (4.1.27) $D(f) = U \setminus f^{-1}(\{0\})$ is open. ∎

## 5.10.3 Exponential and Trigonometric Functions

**Proposition 5.10.5 (Exponential Function)**
*Let $K$ be a complete valued field. Then the series*

$$e^z = \sum_{n=0}^{\infty} \frac{z^n}{n!}$$

*has infinite radius of convergence. Furthermore it satisfies the following properties*

*a) $e^z$ is infinitely differentiable with derivative $e^z$*
*b) $e^{z+w} = e^z e^w$*
*c) $e^z e^{-z} = 1$*

*In the case $K = \mathbb{R}$ then we also have the following property*

*d) $e^x$ is positive, strictly increasing and bijective : $(-\infty, \infty) \to (0, \infty)$*

*Proof.* We have

$$\left|\frac{a_{n+1}z^{n+1}}{a_n z^n}\right| = \frac{z}{n+1} \to 0$$

so the series converges unconditionally. Therefore by (5.6.2) it has infinite radius of convergence and is infinitely differentiable. Observe formally the product of power series is given by

$$\begin{aligned} e^z e^w &= \left(\sum_{n=0}^{\infty} \frac{z^n}{n!}\right)\left(\sum_{n=0}^{\infty} \frac{w^n}{n!}\right) \\ &= \sum_{n=0}^{\infty}\left(\sum_{k=0}^{n} \frac{z^k w^{n-k}}{k!(n-k)!}\right) \\ &= \sum_{n=0}^{\infty} \frac{(z+w)^n}{n!} \\ &= e^{z+w} \end{aligned}$$

which by (5.4.33) converge to the same value.

Evidently $x \geq 0 \implies e^x \geq 1 + x$. Then $x > y \geq 0 \implies e^x = e^{(x-y)+y} = e^{x-y}e^y \geq (1 + x - y)e^y > 1 \cdot e^y$. Similarly suppose $x < y \leq 0$ then $-y > -x \geq 0 \implies e^{-y} > e^{-x} \implies e^x < e^y$. Finally $x \leq 0 \implies e^{-x} \geq 1 - x \implies e^x \leq \frac{1}{1+x}$. This shows that $e^x$ is strictly increasing and therefore injective on $(0, \infty)$. Further using $e^x e^{-x} = 1$ we see it is a positive function and $x \leq 0 \implies e^x \leq 1$.

As $e^x$ is unbounded the intermediate value theorem (5.7.10) shows it achieves every value in $[1, \infty)$. Using $e^x e^{-x} = 1$ we see it also achieves every value in $(0, \infty)$. Therefore it is a bijection $(-\infty, \infty) \to (0, \infty)$ as required. ∎

© David Rufino CC BY-NC-ND 4.0

**Proposition 5.10.6 (Trigonometric Functions)**
*The power series*

$$\sin(z) = z - \frac{z^3}{3!} + \frac{z^5}{5!} - \cdots$$

$$\cos(z) = 1 - \frac{z^2}{2!} + \frac{z^4}{4!} - \cdots$$

*have infinite radius of convergence in* $\mathbb{C}$*. Furthermore*

*a)* $\sin(0) = 0$ *and* $\cos(0) = 1$
*b)* $e^{iz} = \cos(z) + i\sin(z)$
*c)* $\sin(z) = \frac{e^{iz} - e^{-iz}}{2i}$ *and* $\cos(z) = \frac{e^{iz} + e^{-iz}}{2}$*.* $\sin(-z) = -\sin(z)$ *and* $\cos(-z) = \cos(z)$
*d)* $\sin'(z) = \cos(z)$ *and* $\cos'(z) = -\sin(z)$ *are infinitely differentiable*
*e)* $\sin^2(z) + \cos^2(z) = 1$
*f)* $\sin(w + z) = \sin(w)\cos(z) + \cos(w)\sin(z)$
*g)* $\cos(w + z) = \cos(w)\cos(z) - \sin(w)\sin(z)$

*Proof.* It is simpler to define $\sin(z)$ and $\cos(z)$ by *c*). The power series expansion then follows from (5.4.26).

a) Immediate from the series expansion
b) Immediate from *c*)
c) Already shown
d) Immediate from *c*) and the chain rule (5.9.6).
e) Use *b*) and $e^{iz}e^{-iz} = 1$

**f)-g)** Using *b*) we find

$$\begin{aligned}\cos(z + w) + i\sin(z + w) &= e^{i(z+w)} = e^{iz}e^{iw} \\ &= (\cos(z) + i\sin(z))(\cos(w) + i\sin(w)) \\ &= (\cos(z)\cos(w) - \sin(z)\sin(w)) + i(\sin(z)\cos(w) + \cos(z)\sin(w))\end{aligned}$$

and the result follows from equating real and imaginary parts ∎

**Proposition 5.10.7 (cos and sin are periodic)**
*There is a unique positive real number* $\pi \in \mathbb{R}$ *such that the real roots of* $\sin(x)$ *are precisely*

$$\{n\pi \mid n \in \mathbb{Z}\}$$

*Furthermore we have the period formulae*

*a)* $\sin(x) = \cos(x - \frac{\pi}{2}) = \cos(\frac{\pi}{2} - x)$
*b)* $\cos(x) = \sin(\frac{\pi}{2} - x) = -\sin(x - \frac{\pi}{2})$
*c)* $\sin(x) = -\sin(x + \pi)$ *and* $\sin(x + 2\pi) = \sin(x)$
*d)* $\cos(x) = -\cos(x + \pi)$ *and* $\cos(x + 2\pi) = \cos(x)$

*and the real roots of* $\cos(x)$ *are precisely*

$$\left\{\left(n + \frac{1}{2}\right)\pi \;\middle|\; n \in \mathbb{Z}\right\}$$

*Proof.* By (5.10.6).e) we know that $0 \leq \sin^2(x) \leq 1$ which means $-1 \leq \sin(x) \leq 1$, and similarly for $\cos(x)$. We first prove that $\cos(x)$ has a positive root. Suppose not then by the intermediate value theorem (5.7.10) we must have $\cos(x) > 0$ for all $x$. Then by the Mean-Value Theorem (5.7.12) $\sin(x)$ must be strictly increasing and therefore positive for all $x > 0$. We claim that for $x \geq a > 0$

$$\cos(x) \leq \cos(a) + \cos'(a)(x - a) = \cos(a) - \sin(a)(x - a)$$

For define $g(x) \coloneqq \cos(a) - \sin(a)(x - a) - \cos(x)$ then $g(a) = \cos(a) > 0$ and $g'(x) = \sin(x) - \sin(a) > 0$. By the Mean-Value Theorem $g(x)$ is increasing and the claim is proven. However it is evident that this implies $\cos(x)$ is unbounded below contradicting $-1 \leq \cos(x)$. Therefore $\cos(x)$ has a strictly positive root.

Define this to be

$$\frac{\pi}{2} \coloneqq \inf\{x \in \mathbb{R}^+ \mid \cos(x) = 0\}$$

© David Rufino CC BY-NC-ND 4.0

By (5.1.18) there exists a sequence $x_n$ such that $\cos(x_n) = 0$ and $x_n \uparrow \frac{\pi}{2}$. By continuity and (5.3.6) we have $\cos(\frac{\pi}{2}) = 0$, and evidently $\pi \neq 0$. We must have $\cos(x)$ positive on the interval $[0, \frac{\pi}{2}]$, for otherwise by the intermediate value theorem we would have a smaller root. This shows that $\sin(x)$ is increasing on the interval $[0, \frac{\pi}{2}]$. By (5.10.6) we have $\sin^2(\frac{\pi}{2}) = 1$ and therefore $\sin(\frac{\pi}{2}) = 1$.

The period formula $a$) then follows from the addition formula (5.10.6).g)

$$\cos\Big(\frac{\pi}{2} - x\Big) = \cos\Big(x - \frac{\pi}{2}\Big) = \cos(x)\cos\Big(-\frac{\pi}{2}\Big) - \sin(x)\sin\Big(-\frac{\pi}{2}\Big) = \sin(x)$$

and $b$) follows directly from $a$). Then $c), d)$ follow from combining $a), b)$. In particular $\sin(n\pi) = 0$. We claim that $\pi$ is the smallest positive root of $\sin(x)$. For suppose $0 < x < \pi$ satisfies $\sin(x) = 0$ then by $a$) we have $\cos\big(\big|x - \frac{\pi}{2}\big|\big) = 0$ and $0 \leq \big|x - \frac{\pi}{2}\big| < \frac{\pi}{2}$ which contradicts the choice of $\pi$. Suppose $\sin(y) = 0$ with $y > 0$. Then consider

$$n := \max\{m \in \mathbb{N} \mid y - m\pi \geq 0\}$$

Then $\sin(y - n\pi) = 0$ by $c$). By choice of $m$ we have $0 \leq y - n\pi < \pi$. As $\pi$ is the smallest positive root we have $y - n\pi = 0$. By symmetry then the roots of $\sin(x)$ are precisely $\{n\pi\}$. Evidently any such $\pi$ is unique because $\pi' = n\pi = nm\pi'$ whence $n = m = 1$. Using $a$) yields the precise set of roots of $\cos(x)$. ∎

**Proposition 5.10.8**
*There is a continuous, strictly decreasing function*

$$\arccos : [-1, 1] \to [0, \pi]$$

*such that*

$$\cos(\arccos(x)) = x$$

*Proof.* By (5.10.7) $\sin(x)$ is positive on the interval $(0, \pi)$ (by the intermediate value theorem). By the Mean Value Theorem then $\cos(x)$ is strictly decreasing on this interval. Furthermore $\cos(0) = 1$ and $\cos(\pi) = -\cos(\pi - \pi) = -1$. By the intermediate value theorem then $\cos : [0, \pi] \to [-1, 1]$ is surjective and injective. Therefore the map $\arccos$ is well-defined and it is continuous by (5.7.13). ∎

**Proposition 5.10.9 (Polar Coordinates)**
*There is a unique function*

$$\theta : \mathbb{R}^2 \setminus \{(0, 0)\} \to [0, 2\pi)$$

*such that*

$$r(x, y)\cos(\theta(x, y)) = x \text{ and } r(x, y)\sin(\theta(x, y)) = y$$

*where* $r(x, y) := \sqrt{x^2 + y^2}$. *Explicitly it is given by*

$$\theta(x, y) := \begin{cases} \arccos\big(\frac{x}{r(x,y)}\big) & y \geq 0 \\ 2\pi - \arccos\big(\frac{x}{r(x,y)}\big) & y < 0 \end{cases}$$

*and it is continuous on* $\mathbb{R}^2 \setminus \{(x, 0) \mid x > 0\}$, *and* $\theta(x, 0-) = 2\pi + \theta(x, 0+)$ *when* $x > 0$.

*Proof.* Observe that $0 \leq \frac{x}{\sqrt{x^2+y^2}} \leq 1$ and so $r(x, y)\cos(\theta(x, y)) = x$. Further $\sin^2(\theta(x, y)) = 1 - \cos^2(\theta(x, y)) = \frac{y^2}{x^2+y^2}$, and so $r(x, y)\sin(\theta(x, y)) = \pm y$. When $y \geq 0$ we have $\theta(x, y) \in [0, \pi] \implies \sin(\theta(x, y)) \geq 0$. Similarly from the case $y \leq 0$ we find $r(x, y)\sin(\theta(x, y)) = y$ as required. ∎

**Proposition 5.10.10**
*Let* $r, r' > 0$ *and* $\theta, \theta' \in \mathbb{R}$. *Then*

$$(r\cos(\theta), r\sin(\theta)) = (r'\cos(\theta'), r'\sin(\theta'))$$

*if and only if*

$$r = r' \text{ and } \theta = \theta' + 2n\pi$$

*Proof.* We may use (5.10.6) to show that

$$r^2 = (r\cos(\theta))^2 + (r\sin(\theta))^2 = (r'\cos(\theta'))^2 + (r'\sin(\theta'))^2 = (r')^2$$

which means $r = r'$. Similarly we may use the addition formula to find

© David Rufino CC BY-NC-ND 4.0

$$\sin(\theta - \theta') = \sin(\theta)\cos(\theta') - \cos(\theta')\sin(\theta) = 0$$

which means $\theta = \theta' + n\pi$ by (5.10.7). Using the fact $\cos(\theta + n\pi) = (-1)^n \cos(\theta)$ and $\sin(\theta + n\pi) = (-1)^n \sin(\theta)$ we may deduce that $n$ is even (since $\sin(\theta) = \cos(\theta) = 0$ is impossible). ∎

### 5.10.4 Polar Coordinates of a Path

We would like to show that any path $\gamma : [a, b] \to \mathbb{C} \setminus \{0\}$ has a *continuous* parameterisation $(r(t), \theta(t))$ in polar coordinates, and that this is essentially unique.

**Definition 5.10.11**

*Suppose $X$ is a Hausdorff space. A continuous surjective map $p : \tilde{X} \to X$ is a **covering map** if every $x \in X$ has a neighbourhood $U_x$ such that*

*a) $p^{-1}(U_x) = \bigsqcup_{i \in I_x} V_{x,i}$*

*b) $p|_{V_{x,i}} : V_{x,i} \to U_x$ is a homeomorphism for all $i \in I_x$*

*We may show that $\tilde{X}$ is necessarily Hausdorff.*

**Proposition 5.10.12 (Punctured Plane Covering Map)**

*The map*

$$\begin{aligned} \mathbb{R}_{>0} \times \mathbb{R} &\to \mathbb{R}^2 \setminus \{(0,0)\} \\ (r, \theta) &\to (r\cos(\theta), r\sin(\theta)) \end{aligned}$$

*is a covering map.*

*Proof.* Define

$$\begin{aligned} U_1 &:= \mathbb{R}^2 \setminus \{(x, 0) \mid x \geq 0\} \\ U_2 &:= \mathbb{R}^2 \setminus \{(x, 0) \mid x \leq 0\} \end{aligned}$$

We claim that

$$\begin{aligned} p^{-1}(U_1) &= \bigsqcup_n \mathbb{R}_{>0} \times (2n\pi, (2n+2)\pi) \\ p^{-1}(U_2) &= \bigsqcup_n \mathbb{R}_{>0} \times ((2n-1)\pi, (2n+1)\pi) \end{aligned}$$

Denote the open sets by $V_{i,n}$ for $i = 1, 2$. Then the maps

$$p|_{V_{i,n}} \to U_i$$

are injective by (5.10.10). We may define explicit continuous inverses by

$$\begin{aligned} U_1 &\to V_{1,n} \\ (x, y) &\to \left(\sqrt{x^2 + y^2}, \theta(x, y) + 2n\pi\right) \\ U_2 &\to V_{2,n} \\ (x, y) &\to \left(\sqrt{x^2 + y^2}, -\theta(-x, y) + 2(n-1)\pi\right) \end{aligned}$$

∎

**Proposition 5.10.13 (Path lifting)**

*Let $p : \tilde{X} \to X$ be a covering map and $\gamma : [a, b] \to X$ a continuous path. Consider any $\tilde{x}_0 \in \tilde{X}$ such that $p(\tilde{x}_0) = \gamma(a)$. Then there exists a unique lifting $\tilde{\gamma} : [a, b] \to \tilde{X}$ such that $p \circ \tilde{\gamma} = \gamma$ and $\tilde{\gamma}(a) = \tilde{x}_0$.*

*Proof.* We first prove uniqueness. Consider two liftings $\tilde{\gamma}_1, \tilde{\gamma}_2 : [a, b] \to \tilde{X}$ and define

$$X := \{t \in [a, b] \mid \tilde{\gamma}_1(t) = \tilde{\gamma}_2(t)\}$$

Fix $t \in X$ and let $U$ be a neighbourhood of $\gamma(t)$ such that $p^{-1}(U) = \bigsqcup_{i \in I} V_i$. Suppose $\tilde{\gamma}_1(t) = \tilde{\gamma}_2(t) \in V_i$. By continuity there exists a neighbourhood $W$ of $t$ such that $\tilde{\gamma}_1(W) \subseteq V_i$ and $\tilde{\gamma}_2(W) \subseteq V_i$. By definition of $X$ this means $W \subseteq X$. Therefore $X$ is open. By (4.1.99) $X$ is also closed. By (5.7.9) then $X$ is either empty or $[a, b]$. However we have $a \in X$ and the liftings must be equal.

Define

© David Rufino CC BY-NC-ND 4.0

$$L \coloneqq \left\{ x \in [a,b] \;\middle|\; \exists \tilde{\gamma} : [a,x] \to \tilde{X} \text{ s.t. } p \circ \tilde{\gamma} = \gamma \text{ and } \tilde{\gamma}(a) = \tilde{x}_0 \right\}$$

To show that $L$ is closed consider a sequence $x_n \in L$ such that $x_n \to x \in [a,b]$. Choose $\gamma(x) \in U$ such that $p^{-1}(U) = \bigsqcup_i V_i$ and $p|_{V_i}$ is a homeomorphism. For some $\varepsilon$ we have $(x-\varepsilon, x+\varepsilon) \cap [a,b] \subset \gamma^{-1}(U)$, and for some $N$, $x_N \in (x-\varepsilon, x+\varepsilon) \cap [a,b]$. In particular there is a lift

$$\tilde{\gamma}' : [a, x_N] \to \tilde{X}.$$

and $\tilde{\gamma}'(x_N) \in V_i$ for some $i$. If $x_N \geq x$ then we are done, otherwise assume $x_N < x$ and define the lift

$$\tilde{\gamma}(t) \coloneqq \begin{cases} \tilde{\gamma}'(t) & t \leq x_N \\ \left(p|_{V_i}\right)^{-1}(\gamma(t)) & x_N \leq t \leq x \end{cases}$$

Therefore $x \in L$, and $L$ is closed. By (5.7.9) we conclude $L = [a,b]$ and we are done. ∎

**Proposition 5.10.14 (Almost uniqueness of polar coordinates)**
*Let $p : \mathbb{R}_{>0} \times \mathbb{R} \to \mathbb{R}^2 \setminus \{(0,0)\}$ be the covering map of the punctured plane (5.10.12) and $\gamma : [a,b] \to \mathbb{R}^2 \setminus \{(0,0)\}$ a continuous path. Suppose $(r_1, \theta_1)$ and $(r_2, \theta_2)$ are liftings. Then*

$$r_1(t) = r_2(t)$$

$$\theta_1(t) = \theta_2(t) + 2n\pi$$

*for some $n \in \mathbb{Z}$.*

*Proof.* By (5.10.10) we have $\theta_2(a) - \theta_1(a) = 2n\pi$. Then $(r_2(t), \theta_2(t) - 2n\pi)$ is also a lifting and so by uniqueness equals $(r_1(t), \theta_1(t))$ as required. ∎

**Definition 5.10.15 (Winding Number)**
*Let $\gamma : [a,b] \to \mathbb{C} \setminus \{0\}$ be a continuous path. We define the **angle traversed** by the path $\gamma$ to be*

$$\theta(b) - \theta(a)$$

*where $(r(t), \theta(t))$ is any lifting of $\gamma$. Note when $\gamma$ is a closed path then this equals $2n\pi$ for a unique integer $n$. We call this the **winding number** of the closed path $\gamma$, which we denote by $W(\gamma; 0)$. In general we define*

$$W(\gamma; z_0) \coloneqq W(\gamma - z_0; 0)$$

*for any $z_0 \notin \gamma([a,b])$.*

## 5.10.5 Path Integrals

**Definition 5.10.16**
*A **curve** is a continuous path $\gamma : [a,b] \to \mathbb{C}$, for which there exists a partition*

$$a = x_0 < x_1 < \cdots < x_{n+1} = b$$

*such that*

*a) $\gamma$ is $C^1$ on $(x_i, x_{i+1})$ for $i = 0 \cdots n-1$*
*b) $\gamma$ is left and right differentiable at $x_i$ and*

$$\lim_{h \downarrow 0} \frac{\gamma(x_i + h) - \gamma(x_i)}{h} = \lim_{h \downarrow 0} \gamma'(x_i + h)$$

$$\lim_{h \uparrow 0} \frac{\gamma(x_i + h) - \gamma(x_i)}{h} = \lim_{h \uparrow 0} \gamma'(x_i + h)$$

*We say $\gamma$ is closed if in addition $\gamma(b) = \gamma(a)$.*

**Definition 5.10.17 (Path Integral)**
*Let $\gamma : [a,b] \to \mathbb{C}$ be a curve supported on $\{x_0, \cdots, x_{n+1}\}$. Define the integral*

$$\int_\gamma f(z)\,\mathrm{d}z \coloneqq \sum_{i=0}^{n} \int_{x_i}^{x_{i+1}} f(\gamma(t))\gamma'(t)\,\mathrm{d}t$$

© David Rufino CC BY-NC-ND 4.0

**Proposition 5.10.18 (Path Integral of a function with primitive)**
*Let $f : U \to \mathbb{C}$ be a continuous function and $g : U \to \mathbb{C}$ a differentiable function such that $g'(z) = f(z)$. Then for any curve $\gamma : [a, b] \to U$ we have*

$$\int_\gamma f(z)\,\mathrm{d}z = g(b) - g(a)$$

*In particular the integral over a closed curve is zero.*

**Proposition 5.10.19**
*Let $\gamma : [0, 1] \to \mathbb{C}$ the path given by $\gamma(t) \coloneqq Re^{2\pi i t}$. Then evidently $W(\gamma, 0) = 1$. Further*

$$\int_\gamma z^n\,\mathrm{d}z = \begin{cases} 2\pi i & n = -1 \\ 0 & n \neq -1 \end{cases}$$

*Proof.* When $n \neq -1$ then $z^n$ has primitive $\frac{z^{n+1}}{n+1}$ and we are done by the previous result. In the case $n = -1$ we may calculate explicitly

$$\int_\gamma z^{-1}\,\mathrm{d}z = \int_0^1 \frac{\gamma'(t)}{\gamma(t)}\,\mathrm{d}t = 2\pi i$$

■

© David Rufino CC BY-NC-ND 4.0

## 5.11 Riemann Surfaces

I develop the elementary theory of Riemann surface, starting with the more standard concept approach of maximal atlases (see [War13]) and then relating to the notion sheaves (see [Wed16] (Section 4.2)). The language of sheaves is well suited to complex analytic spaces and aligns well with the approach in the algebraic case.

**Definition 5.11.1**
*Let $X$ be a topological space. By definition a **chart** on $X$ is a pair $(U, \varphi)$ where $U \subset X$ is an open set and $\varphi$ is a homeomorphism onto an open subset $\varphi(U)$ of $\mathbb{C}$.*

*We say that two charts $(U, \varphi)$ and $(V, \psi)$ are **compatible** if either $U \cap V = \varnothing$ or the composite map $(\psi \circ \varphi^{-1}) : \varphi(U \cap V) \to \psi(U \cap V)$ is **biholomorphic**.*

*An **atlas** $\mathcal{A}$ on $X$ is a collection of charts $\{(U_\alpha, \varphi_\alpha)\}$ for which $X = \bigcup_\alpha U_\alpha$ and every pair of charts is compatible.*

*We say two atlases $\mathcal{A}, \mathcal{A}'$ are **equivalent** if $\mathcal{A} \cup \mathcal{A}'$ is an atlas.*

**Proposition 5.11.2**
*The notion of compatible charts is an equivalence relation. Similarly for the notion of equivalent atlases.*

*A union of equivalent atlases is again an atlas, and every atlas is contained in a unique maximal atlas.*

We may then use the charts to define what it means for a function on a manifold to be holomorphic.

**Lemma 5.11.3**
*Let $X$ be a topological space, $U \subset X$ an open subset, $f : U \to \mathbb{C}$ a function and $(V, \varphi)$, $(V', \psi)$ two compatible charts such that $U \cap V \cap V' \neq \varnothing$.*

*Then $f \circ \varphi^{-1}|_{\varphi(U \cap V \cap V')}$ is holomorphic if and only if $f \circ \psi^{-1}|_{\psi(U \cap V \cap V')}$ is holomorphic.*

**Lemma 5.11.4**
*Let $X$ be a topological space, $U \subset X$ an open subset, $f : U \to \mathbb{C}$ a function and $(V_\alpha, \varphi_\alpha)_{\alpha \in \mathcal{A}}$ a family of charts such that $U \subset \bigcup_{\alpha \in \mathcal{A}} V_\alpha$. Further suppose that $(V, \varphi)$ is a chart such that $U \cap V \neq \varnothing$ and is compatible with every $(V_\alpha, \varphi_\alpha)$. Then the following are equivalent*

*a) $f \circ \varphi^{-1}|_{\varphi(V \cap U)}$ is holomorphic*
*b) $f \circ \varphi_\alpha^{-1}|_{\varphi_\alpha(V_\alpha \cap U \cap V)}$ is holomorphic for all $\alpha \in \mathcal{A}$ for which $V_\alpha \cap V \neq \varnothing$*

**Proposition 5.11.5 (Holomorphic Function)**
*Let $X$ be a topological space, $U \subset X$ an open subset, $f : U \to \mathbb{C}$ a function and $\mathcal{A}$ a given atlas. Then the following are equivalent*

*a) there exists a collection of charts $\{(V_\alpha, \varphi_\alpha)\} \subset \mathcal{A}$ such that $V_\alpha \cap U \neq \varnothing$, $f \circ \varphi_\alpha^{-1}|_{\varphi_\alpha(V_\alpha \cap U)}$ is holomorphic and $U \subset \bigcup_\alpha V_\alpha$*
*b) for every chart $(V, \varphi) \in \mathcal{A}$ such that $U \cap V \neq \varnothing$ the function $f \circ \varphi^{-1}|_{\varphi(U \cap V)}$ is holomorphic.*

*When this holds we say that $f$ is **holomorphic** with respect to $\mathcal{A}$.*

*Proof.* $b) \implies a)$ Straightforward

$a) \implies b)$ As every chart $(V, \varphi)$ is compatible with the collection $\{(V_\alpha, \varphi_\alpha)\}$ this follows from (5.11.4). ∎

**Proposition 5.11.6**
*A holomorphic function $f : U \to \mathbb{C}$ is continuous.*

*Proof.* For every chart $(V, \varphi)$ the function $f \circ \varphi^{-1}|_{\varphi(U \cap V)}$ is holomorphic (5.11.5) and therefore continuous. Further by definition the map $\varphi|_{U \cap V}$ is continuous whence $f|_{U \cap V}$ is continuous (4.1.28). As the charts cover $X$ we are done by (4.1.29). ∎

We first prove that this definition does not depend on which atlas is chosen.

**Proposition 5.11.7**
*Let $X$ be a topological space, $U \subset X$ an open subset, $f : U \to \mathbb{C}$ a function and $\mathcal{A}, \mathcal{A}'$ two equivalent atlases.*

*Then $f$ is holomorphic with respect to $\mathcal{A}$ if and only if it is holomorphic with respect to $\mathcal{A}'$.*

*Proof.* Since equivalence of atlases is symmetric we only need to show one direction. So suppose that $f$ is holomorphic with respect to $\mathcal{A}$, and let $(W, \psi) \in \mathcal{A}'$ be a chart. Then by (5.11.4) $f \circ \psi^{-1}$ is holomorphic, so that $f$ is holomorphic with respect to $\mathcal{A}'$. ∎

© David Rufino CC BY-NC-ND 4.0

Further we show that just as in the usual case this is a local condition.

**Proposition 5.11.8 (Structure Sheaf associated to an Atlas)**
*Let $X$ be a topological space together with an atlas $\mathcal{A}$. Let $U = \bigcup_{i\in I} U_i$ an open cover and $f : U \to \mathbb{C}$ a function. Then $f$ is holomorphic (with respect to $\mathcal{A}$) if and only if $f|_{U_i}$ is holomorphic for all $i \in I$.*

*In particular for every atlas the mapping*

$$\mathcal{O}_X(U) \coloneqq \{f : U \to \mathbb{C} \mid f \text{ holomorphic } \}$$

*determines a sheaf of $\mathbb{C}$-algebras, and the pair $(X, \mathcal{O}_X)$ constitutes a space of $\mathbb{C}$-functions, which we call the **structure sheaf**. It contains an isomorphic copy of $\mathbb{C}$ consisting of the constant functions.*

*Two atlases are equivalent if and only if they determine the same structure sheaf.*

*A space of functions $(X, \mathcal{O}_X)$ arises in this way if and only if there is an open cover $X = \bigcup_{i\in I} U_i$ such that $\left(U_i, \mathcal{O}_X|_{U_i}\right) \xrightarrow{\sim} \left(V_i, \mathcal{O}_{V_i}\right)$ for $V_i \subset \mathbb{C}$ an open subset with the usual sheaf of holomorphic functions.*

*Proof.* Suppose that $(V, \varphi) \in \mathcal{A}$ is a chart such that $V \cap U_i \neq \varnothing$. Then by assumption $f \circ \varphi^{-1}|_{\varphi(V\cap U)}$ is holomorphic whence by (...) $f|_{U_i} \circ \varphi^{-1}|_{\varphi(V\cap U_i)}$ is holomorphic. Since the chart $(V, \varphi)$ was arbitrary we conclude $f|_{U_i}$ is holomorphic with respect to the atlas $\mathcal{A}$.

Conversely suppose $f|_{U_i}$ is holomorphic with respect to $\mathcal{A}$ for all $i \in I$. Consider an arbitrary chart $(V, \varphi) \in \mathcal{A}$ such that $V \cap U \neq \varnothing$. Then $V \cap U = \bigcup_{i\in I} V \cap U_i$, where we may restrict to open subsets with $V \cap U_i \neq \varnothing$. Then by assumption $f|_{U_i} \circ \varphi^{-1}|_{\varphi(V\cap U_i)} = \left(f \circ \varphi^{-1}|_{\varphi(V\cap U)}\right)|_{\varphi(V\cap U_i)}$ is holomorphic. By (...) $f \circ \varphi^{-1}|_{\varphi(V\cap U)}$ is holomorphic. As the chart was arbitrary we conclude that $f$ is holomorphic with respect to $\mathcal{A}$.

The holomorphic functions form a $\mathbb{C}$-algebra by (5.5.4), and in particular contain the constant functions. Further $D(f) = X \setminus f^{-1}(\{0\})$ which is an open subset of $X$, and $\left(f|_{D(f)}\right)^{-1}$ is holomorphic whenever $f$ is not identically zero.

By definition then $(X, \mathcal{O}_X)$ is a space of $\mathbb{C}$-functions. By (5.11.7) an equivalent atlas determines the same structure sheaf. Conversely suppose $\mathcal{A}, \mathcal{A}'$ determine the same structure sheaves. Consider two atlases $(V, \varphi) \in \mathcal{A}$ and $(V', \varphi') \in \mathcal{A}'$. Then by definition we see that $\varphi \circ (\varphi')^{-1}$ is holomorphic and $(\varphi') \circ \varphi^{-1}$ is holomorphic, and therefore they are compatible as required. ∎

**Definition 5.11.9 (Riemann Surface)**
*Let $(X, \mathcal{O}_X)$ be a space of $\mathbb{C}$-functions, which is Hausdorff and second countable. Further suppose there is an open cover $X = \bigcup_{i\in I} U_i$ such that there exists isormorphisms*

$$\varphi_i : \left(U_i, \mathcal{O}_X|_{U_i}\right) \xrightarrow{\sim} \left(V_i, \mathcal{O}_{V_i}\right)$$

*for $V_i \subset \mathbb{C}$ an open subset. Then we say $X$ is a **Riemann Surface**.*

*We say $\mathcal{A}$ is a compatible atlas if it determines the same structure sheaf. Further there exists a unique compatible maximal atlas (5.11.8).*

*Then $\varphi : U \to \mathbb{C}$ is a chart for $X$ if it lies in at least one compatible atlas (equivalently is in the unique maximal atlas).*

## 5.11.1 Morphisms of Riemann Surfaces

**Proposition 5.11.10 (Holomorphic maps of Riemann Surfaces)**
*Let $f : X \to Y$ be a continuous map of Riemann surfaces with given compatible atlases $\mathcal{A}_1, \mathcal{A}_2$. Then the following are equivalent*

*a) $g \in \mathcal{O}_Y(V) \implies g \circ f \in \mathcal{O}_X(f^{-1}(V))$*
*b) For all charts $(V, \psi) \in \mathcal{A}_2$ the function $\psi \circ f|_{f^{-1}(V)}$ is holomorphic*
*c) For all charts $(U, \varphi) \in \mathcal{A}_1$, $(V, \psi) \in \mathcal{A}_2$ for which $U \cap f^{-1}(V) \neq \varnothing$ we have*

$$\psi \circ f \circ \varphi^{-1} : \varphi(U \cap f^{-1}(V)) \to \mathbb{C}$$

*is holomorphic*

*This definition is independent of the choice of atlases, and we say in this case that $f$ is **holomorphic**.*

*Proof.* $a) \implies c)$ As $\psi \in \mathcal{O}_Y(V)$ is trivially holomorphic we have by assumption $\psi \circ f \in \mathcal{O}_X(f^{-1}(V)) \implies (\psi \circ f)|_{U\cap f^{-1}(V)} \in \mathcal{O}_X(U \cap f^{-1}(V))$. By definition this means $\psi \circ f \circ \varphi^{-1}$ is holomorphic.

$c) \implies b)$ By (5.11.5) $\psi \circ f|_{f^{-1}(V)}$ is holomorphic

© David Rufino CC BY-NC-ND 4.0

*b)* $\Longrightarrow$ *a)* Let $(V_\alpha, \psi_\alpha) \in \mathcal{A}_2$ be charts for which $V \subset \bigcup_\alpha V_\alpha$ and $V_\alpha \cap V \neq \varnothing$. Then

$$(g \circ f)|_{f^{-1}(V_\alpha)} = (g \circ \psi_\alpha^{-1}) \circ (\psi_\alpha \circ f)|_{f^{-1}(V_\alpha)}$$

is by definition holomorphic. Since this is a local condition (5.11.8) we conclude that $g \circ f$ is holomorphic as required.

The last statement follows because *a)* is independent of the choice of compatible atlases (5.11.8). ∎

## 5.11.2 Tangent and Cotangent Space

**Definition 5.11.11 (Local Ring)**
*Let $X$ be a Riemann surface and $x \in X$ a point. Then define the local ring*

$$\mathcal{O}_{X,x} \coloneqq \varinjlim_{x \in U} \mathcal{O}_X(U)$$

*to be the $\mathbb{C}$-algebra of germs at $x$. Every germ $\boldsymbol{f}$ has a well-defined value at $x$ which we denote by $\boldsymbol{f}(x)$. The unique maximal ideal $\mathfrak{m}_{X,x}$ consists of those germs which vanish at $x$. Further the residue field $\mathbb{C}(\mathfrak{m}_{X,x})$ is isomorphic to $\mathbb{C}$ (that is, $\mathbb{C} \to \mathcal{O}_{X,x} \to \mathcal{O}_{X,x}/\mathfrak{m}_{X,x}$ is an isomorphism).*

**Proposition 5.11.12 (Cotangent Space is Finite-Dimensional)**
*The local ring $\mathcal{O}_{X,x} = \mathbb{C} \oplus \mathfrak{m}_{X,x}$ and there is a canonical isomorphism*

$$\mathrm{Der}(\mathcal{O}_{X,x}, \mathbb{C}) \xrightarrow{\sim} \mathrm{Hom}_{\mathbb{C}}(\mathfrak{m}_{X,x}/\mathfrak{m}_{X,x}^2, \mathbb{C})$$

*For every $f \in \mathcal{O}_X(U)$ and $x \in U$ denote by $\mathrm{d}f(x)$ the image of $(f - f(x))$ in the **cotangent space** $T_x^\star X \coloneqq \mathfrak{m}_{X,x}/\mathfrak{m}_{X,x}^2$.*

*This has dimension 1, more explicitly for any open chart $(U, \varphi)$ around $x$, $0 \neq \mathrm{d}\varphi(x)$ is a basis for $\mathfrak{m}_{X,x}/\mathfrak{m}_{X,x}^2$.*

*We denote the inverse image of the dual basis $(\mathrm{d}\varphi(x))^\vee$ by $\left(\frac{\partial}{\partial \varphi}\right)|_x$ (which is therefore a basis for $\mathrm{Der}(\mathcal{O}_{X,x}, \mathbb{C})$).*

*Proof.* Every germ $\boldsymbol{f}$ may be written as $\boldsymbol{f} = \boldsymbol{f}(x) + (\boldsymbol{f} - \boldsymbol{f}(x))$. This is unique decomposition by considering evaluation at $x$, which gives the first statement. The isomorphism follows the same lines as were proved for affine algebras that tangent space is dual to cotangent space (see (3.35.50)). Explicitly the map is given by $D(\mathrm{d}f(x)) \coloneqq Df_x$.

Let $(U, \varphi)$ be a chart and define the holomorphic function $z(x') \coloneqq \varphi(x') - \varphi(x)$. For every $f \in \mathcal{O}_X(U)$ define $\hat{f} \coloneqq f \circ \varphi^{-1}$. Then by Taylor's theorem (5.9.29)

$$\hat{f}(z') = \hat{f}(\varphi(x)) + (z' - \varphi(x))\hat{f}'(\varphi(x)) + (z' - \varphi(x))^2 R_2(z')$$

where $R_2(z')$ is a holomorphic function on $\varphi(U)$. Therefore

$$f(x') = f(x) + z(x')(f \circ \varphi^{-1})'(\varphi(x)) + z(x')^2 (R_2 \circ \varphi)(x')$$

where $R_2 \circ \varphi$ is a holomorphic function on $U$. Therefore if $f(x) = 0$ we have $f_x - (f \circ \varphi^{-1})'(\varphi(x)) z_x \in \mathfrak{m}_{X,x}^2$ and $\mathrm{d}f(x) = (f \circ \varphi^{-1})'(x)\,\mathrm{d}z(x)$. In other words $\mathrm{d}z(x) = \mathrm{d}\varphi(x)$ is spanning.

Observe that $\hat{z}'(x) = 1$. We claim that for any $f$ such that $f_x \in \mathfrak{m}_{X,x}^2$ we have $\hat{f}'(x) = 0$, which will show that $z_x \notin \mathfrak{m}_{X,x}^2$ and therefore $\mathrm{d}z(x) = \mathrm{d}\varphi(x) \neq 0$.

For in this case $\hat{f} = \sum_{i=1}^n \hat{g}_i \hat{h}_i$ when restricted to some open neighbourhood $V$, with $\hat{g}_i(\varphi(x)) = \hat{h}_i(\varphi(x)) = 0$. By the product rule $\hat{f}'(\varphi(x)) = 0$ as required. ∎

**Definition 5.11.13**
*Let $X$ be a Riemann Surface and $x \in X$. We define the **tangent space** to be the $\mathbb{C}$-vector space of derivations*

$$T_x X \coloneqq \mathrm{Der}(\mathcal{O}_{X,x}, \mathbb{C})$$

*By (5.11.12) it has dimension 1, more explicitly given any chart $(U, \varphi)$ around $x$ it has basis $\left(\frac{\partial}{\partial \varphi}\right)|_x$ the unique derivation such that*

$$\left(\frac{\partial}{\partial \varphi}\right)|_x \, \varphi_x = 1$$

**Lemma 5.11.14 (Derivations are Directional Derivatives)**
*Let $f \in \mathcal{O}_X(U)$, $x \in U$ and $(V, \varphi)$ a chart around $x$, then*

$$\left(\frac{\partial}{\partial \varphi}\right)|_x \, f_x = (f \circ \varphi^{-1})'(\varphi(x))$$

© David Rufino CC BY-NC-ND 4.0

*In particular*

$$\mathrm{d}f(x) = \left(f \circ \varphi^{-1}\right)'(\varphi(x))\,\mathrm{d}\varphi(x)$$

*Proof.* As the derivative of a function depends only on its germ, the right hand side is a well-defined derivation on $\mathcal{O}_{X,x}$, by the usual rules of differentiation (5.5.4). Further $(\varphi \circ \varphi^{-1})(z) = z$ which has derivative 1 and so both sides are equal when evaluated at $f_x = \varphi_x$. By uniqueness (5.11.13) then they are equal. ∎

**Corollary 5.11.15 (Change of Variables Formula)**
*Let $X$ be a Riemann Surface and $(U, \varphi)$, $(V, \psi)$ two charts around $x$. Then*

$$\mathrm{d}\varphi(x) = \left(\varphi \circ \psi^{-1}\right)'(\psi(x))\,\mathrm{d}\psi(x)$$

*and dually*

$$\left(\frac{\partial}{\partial \psi}\right)|_x = \left(\varphi \circ \psi^{-1}\right)'(\psi(x))\left(\frac{\partial}{\partial \varphi}\right)|_x$$

*Proof.* By (5.11.12) both $\mathrm{d}\varphi(x)$ and $\mathrm{d}\psi(x)$ form a basis for $T_x^\star X$, whence $\mathrm{d}\varphi(x) = \lambda\,\mathrm{d}\psi(x)$ for some $\lambda \in K^\star$. Therefore $\varphi_x - \lambda\psi_x \in \mathbb{C} + \mathfrak{m}_{X,x}^2$. By (...)

$$0 = \left(\frac{\partial}{\partial \psi}\right)|_x\ (\varphi_x - \lambda\psi_x) = \left(\varphi \circ \psi^{-1}\right)'(\psi(x)) - \lambda$$

as required. ∎

## 5.11.3 Holomorphic 1-forms

**Lemma 5.11.16**
*Let $X$ be a Riemann surface, $U \subset X$ an open subset and $\omega : U \to \coprod_{x\in U} T_x^\star X$ a function satisfying $\omega(x) \in T_x^\star X$. Then for every chart $(V, \varphi)$ such that $U \cap V \neq \varnothing$*

$$\omega(x) = \omega_\varphi(x)\,\mathrm{d}\varphi(x) \quad \forall x \in U \cap V$$

$$\omega_\varphi(x) := \left(\frac{\partial}{\partial \varphi}\right)|_x\ \omega(x)$$

*Suppose that $(W, \psi)$ is another chart such that $U \cap V \cap W \neq \varnothing$ then*

$$\omega_\psi(x) = \left(\varphi \circ \psi^{-1}\right)'(\psi(x))\omega_\varphi(x) \quad \forall x \in U \cap V \cap W$$

**Definition 5.11.17 (Holomorphic $1$-form)**
*Let $X$ be a Riemann surface, $U \subset X$ an open subset and $\mathcal{A}$ a compatible atlas. We say a function $\omega : U \to \coprod_{x\in U} T_x^\star X$ is a **holomorphic 1-form** if*

*a) $\omega(x) \in T_x^\star X$ for all $x \in U$*
*b) for every chart $(V, \varphi) \in \mathcal{A}$ such that $U \cap V \neq \varnothing$, the function $\omega_\varphi : U \cap V \to \mathbb{C}$ is holomorphic*

**Lemma 5.11.18**
*The definition of holomorphic 1-form (5.11.17) is independent of the choice of atlas.*

*Proof.* Suppose $\omega$ is a holomorphic 1-form on $U$ with respect to the atlas $\mathcal{A}$. Let $(V_i, \varphi_i) \in \mathcal{A}$ and $(W, \psi)$ be compatible charts such that $U \cap V_i \cap W \neq \varnothing$. Then by (5.11.16) and (5.11.15) for all $x \in U \cap V_i \cap W$

$$\omega_\psi(x) = \left(\varphi_i \circ \psi^{-1}\right)'(\psi(x))\omega_{\varphi_i}(x)$$

whence by definition $\omega_\psi \circ \varphi_i^{-1} : \varphi_i(U \cap V_i \cap W) \to \mathbb{C}$ is holomorphic. Then by (5.11.5) $\omega_\psi$ is holomorphic on $U \cap W$. Therefore if $\mathcal{A}'$ is a compatible atlas we find $\omega$ is holomorphic with respect to $\mathcal{A}'$. By symmetry we see that the definition is independent of the choice of atlas. ∎

**Proposition 5.11.19**
*Let $X$ be a Riemann surface and $f \in \mathcal{O}_X(U)$. Then the function*

$$\begin{aligned} \mathrm{d}f : U &\longrightarrow \coprod_{x\in U} T_x^\star U \\ x &\longrightarrow \mathrm{d}f(x) \end{aligned}$$

© David Rufino CC BY-NC-ND 4.0

*is a holomorphic* 1*-form on* $U$*. Moreover for any chart* $(U, \varphi)$ *we have*

$$\mathrm{d}f(x) = \left(\partial \frac{f}{\partial \varphi}\right)|_x \, \mathrm{d}\varphi(x) = \left(f \circ \varphi^{-1}\right)'(\varphi(x)) \, \mathrm{d}\varphi(x)$$

**Proposition 5.11.20**
*Let* $X$ *be a Riemann surface. For every open subset* $U \subset X$ *define*

$$\Omega^1_X(U) \coloneqq \left\{ \omega : U \to \coprod_{x \in U} T^\star_x X \;\middle|\; \omega \text{ holomorphic } \ 1\text{-form} \right\}$$

*Then* $\Omega^1_X$ *is a locally-free* $\mathcal{O}_X$*-module of rank* 1*, more precisely for every chart* $(U, \varphi)$ *we have* $\Omega^1_X|_U$ *has global basis* $\mathrm{d}\varphi|_U$*.*

*Proof.* Evidently $\Omega^1_X(U)$ is a vector space over $\mathbb{C}$, as $\mathcal{O}_X(U)$ forms a $\mathbb{C}$-algebra. Further for $f \in \mathcal{O}_X(U)$ we find $f\omega$ is a holomorphic 1-form and $(f\omega)|_V = f|_V \, \omega|_V$, so that $\Omega^1_X$ is an $\mathcal{O}_X$-module. For every chart $(U, \varphi)$ and open subset $V \subset U$ we have an isomorphism of $\mathcal{O}_X(V)$-modules

$$\begin{aligned} \mathcal{O}_X(V) &\xrightarrow{\sim} \Omega^1_X(V) \\ f &\to f \, \mathrm{d}\varphi|_U \\ \omega_\varphi &\leftarrow \omega \end{aligned}$$

whence an isomorphism

$$\mathcal{O}_X|_U \xrightarrow{\sim} \Omega^1_X|_U$$

■

© David Rufino CC BY-NC-ND 4.0

## Chapter 6

# Algebraic Geometry

I develop theory of varieties over a non-algebraically closed field $k$ which more closely resembles "Faisceaux algébriques cohérents" [Ser55] than say EGA, and therefore is a bit more elementary because the structure sheaf consists of functions into a single ambient field $\overline{k}$ (and perhaps psychologically because the underlying set consists of solutions to equations rather than Galois orbits of solutions). By dealing with irreducible subsets rather than just points then we can also avoid the use of generic points or a large ambient field of infinite transcendence degree. I summarise the features of this exposition versus other popular expositions from which I sourced the material / inspiration

| Author | Book | Non-Algebraically Closed Field | Abstract Variety | Real Functions | Arithmetic Case |
|---|---|---|---|---|---|
| J.P. Serre | Faisceaux algébriques cohérents [Ser55] | No | Yes | Yes | No |
| J.S. Milne | Algebraic Geometry [Mil17] | No | Yes | No | No |
| R. Hartshorne | Algebraic Geometry [Har13] (Chapter I) | No | No | Yes | No |
| George R. Kempf | Algebraic Varieties | No | Yes | Yes | No |
| A. Grothendieck | Éléments de géométrie algébrique | Yes | Yes | No | Yes |
| | These Notes | Yes | Yes | Yes | No |

© David Rufino CC BY-NC-ND 4.0

# 6.1 Affine Varieties

In order to generalise the usual notions to non-algebraically closed field, and develop a "coordinate-free" approach, we introduce the following concept

**Definition 6.1.1**
*Let $A$ be a finitely generated $k$-algebra and $\mathfrak{m}$ a maximal ideal. Recall (3.35.21) that $k(\mathfrak{m})/k$ is an algebraic (indeed finite) field extension, and we call it the* ***residue field*** *for $\mathfrak{m}$.*

Now we may introduce the Zero-Loci and study relationships to ideals

**Proposition 6.1.2 (Correspondence between Zero-Loci and Ideals)**
*Let $A = k[X_1, ..., X_n]$ be the polynomial ring in $n$-variables over a field $k$. For a set $S \subseteq k[X_1, ..., X_n]$ and an algebraic extension $K/k$ define the* ***zero-locus***

$$V_K(S) \coloneqq \{\alpha \in K^n \mid f(\alpha) = 0 \quad \forall f \in S\}.$$

*Similarly for a subset $Y \subseteq K^n$ define*

$$I_k(Y) \coloneqq \{f \in A \mid f(y) = 0 \quad \forall y \in Y\}$$

*The pair of maps $V_K, I_k$ constitute a Galois Connection*

$$\mathcal{I}(k[X_1, ..., X_n]) \underset{I_k}{\overset{V_K}{\rightleftharpoons}} \mathcal{P}(K^n)$$

*namely they satisfy*

*a) $V_K$ and $I_k$ are order-reversing*
*b) $S \subseteq I_k(V_K(S))$*
*c) $Y \subseteq V_K(I_k(Y))$*

*Furthermore (omitting the subscripts)*

*d) $VIV = V$ and $IVI = I$*
*e) $I(Y)$ is a radical ideal and $\sqrt{\langle S\rangle} \subseteq I(V(S))$*
*f) $V(S) = V(\langle S\rangle) = V\left(\sqrt{\langle S\rangle}\right)$ and $V(\mathfrak{a}) = V\left(\sqrt{\mathfrak{a}}\right)$*
*g) $\bigcap_i V(S_i) = V\left(\bigcup_i S_i\right)$ and $\bigcap_i V(\mathfrak{a}_i) = V\left(\sum_i \mathfrak{a}_i\right)$*
*h) $\bigcap_i I(W_i) = I\left(\bigcup_i W_i\right)$*
*i) $V(\mathfrak{a}) \cup V(\mathfrak{b}) = V(\mathfrak{a} \cap \mathfrak{b}) = V(\mathfrak{a}\mathfrak{b})$*
*j) $V((0)) = K^n$ and $V(A) = \varnothing$*

*The sets of the form $V_K(\mathfrak{a})$ constitute the closed sets of a topology on $K^n$, denoted by $\mathrm{Zar}_k(K^n)$. In this case we have the following form for the topological closure*

$$V_K(I_k(Y)) = \overline{Y}$$

*Furthermore*

$$I_k(Y) = I_k\left(\overline{Y}\right)$$

*Proof.* We make use of general results on Galois connections (Foundations, Set Theory), though many results may be shown more directly. The fact it's a Galois connection follows from (2.1.54).

**1-3.** These follow (2.1.50)

**4.** This follows from (2.1.52).

**5.** It's clear that $I(Y)$ is an ideal. It is radical because $K$ is reduced (...). The second statement is straightforward.

**6.** This follows from (2.1.53) by considering the closure operators $\sqrt{\langle - \rangle}$ and $\langle - \rangle$.

**7.** The first equality follows from (2.1.55). The second equality follows from (3.3.29).

**8.** This follows from (2.1.55).

**9.** Observe that $\mathfrak{m}_x$ is prime (because $K$ is an integral domain) and $x \in V(\mathfrak{a}) \iff \mathfrak{a} \subseteq \mathfrak{m}_x$. the result follows from (3.3.38) because $\mathfrak{a} \subseteq \mathfrak{m}_x \vee \mathfrak{b} \subseteq \mathfrak{m}_x \iff \mathfrak{a}\mathfrak{b} \subseteq \mathfrak{m}_x \iff \mathfrak{a} \cap \mathfrak{b} \subseteq \mathfrak{m}_x$

© David Rufino CC BY-NC-ND 4.0

The family of sets $\mathrm{Zar}_k(K^n) := \mathrm{Im}(V_K)$ constitute the closed sets of a topology precisely because they are closed under arbitrary intersections and finite unions. Furthermore by (2.1.52) $V_K \circ I_k$ is a closure operator with image precisely the closed sets. Therefore by (2.1.41)

$$(V_K \circ I_k)(Y) = \bigcap_{Y \subseteq Z \in \mathrm{Zar}_k(K^n)} Z$$

which is the definition of the topological closure. The last statement follows by applying $I_k$ and using **4.** ∎

For a fixed ideal $\mathfrak{a} \lhd k[X_1, ..., X_n]$ we may vary the field $K$ to obtain families of solutions in different fields.

**Definition 6.1.3 (Algebraic Set and $L$-valued points)**
*Let $\mathfrak{a} \lhd k[X_1, ..., X_n]$ be a **radical** ideal. For every $k$-algebra $R$ define the $R$-valued points by*

$$V(\mathfrak{a})(R) := \{r \in R^n \mid f(r_1, ..., r_n) \quad \forall f \in \mathfrak{a}\}$$

**Definition 6.1.4 (Affine Variety)**
*Let $\mathfrak{a} \lhd k[X_1, ..., X_n]$ be a **radical** ideal. The family $X(-) := V(\mathfrak{a})(-)$ is called an **affine variety** to which we associate the **coordinate ring** $k[X] := k[X_1, ..., X_n]/\mathfrak{a}$, which is a reduced f.g. $k$-algebra. Note the elements of $k[X]$ may be regarded as $R$-valued functions on the $R$-valued points $X(R)$.*

*Note we may regard this as a functor, suppose we have compatible maps $i_{jk} : K_j/k \to K_k/k$ then these induce compatible maps*

$$X(i_{jk}) : X(K_j) \to X(X_k)$$

**Definition 6.1.5 (Residue Field)**
*Let $X = V(\mathfrak{a})$ be an affine variety and $L/k$ a field extension. For a point $x \in X(L)$ define the **residue field** to be*

$$k(x) := k(x_1, ..., x_n) \subset L/k$$

*and the degree of $x$ to be*

$$\deg(x) := [k(x) : k]$$

*Furthermore we define*

$$\mathfrak{M}_{X,x} := I_k(\{(x)\})$$

*for $x \in X(L)$. When $x$ is algebraic then $\deg(x)$ is finite and $\mathfrak{M}_{X,x} = \mathfrak{m}_x/\mathfrak{a}$ is maximal (3.35.21).*

**Remark 6.1.6**
*$\mathbb{A}^n_k := V((0))$ is an affine variety and $\mathbb{A}^n_k(L) = L^n$.*

**Proposition 6.1.7 (Rational Points are Algebra Homomorphisms)**
*Let $X = V(\mathfrak{a})$ be an affine variety and $R$ a $k$-algebra. Then there is a bijection*

$$\{x \in X(R)\} \longleftrightarrow \mathrm{AlgHom}_k(k[X], R)$$

*Proof.* Given $x \in X(R)$ then the evaluation map $\mathrm{ev}_x : k[X_1, ..., X_n] \to R$ contains $\mathfrak{a}$ by definition and therefore (3.11.3) yields a unique map $\varphi_x : k[X] \to R$ such that $\varphi_x\left(\overline{X}_i\right) = x_i$.

Similarly given an algebra homomorphism $\varphi : k[X] \to R$ we may define $x = \left(\varphi\left(\overline{X}_1\right), ..., \varphi\left(\overline{X}_n\right)\right)$. Then by uniqueness these are mutual inverses. ∎

For completeness we also consider affine subvarieties in the same way as before

**Proposition 6.1.8 (Affine subvarieties)**
*Let $X = V(\mathfrak{a}) \subset \mathbb{A}^n_k$ be an affine variety and $A := k[X]$ the coordinate ring. For any ideal $\mathfrak{b} \lhd A$ and field extension $K/k$ define the **zero-locus** by*

$$V_K(\mathfrak{b}) := \{(x) \in X(K) \mid f(x) = 0 \quad \forall f \in \mathfrak{b}\}$$

*Similarly for a subset $Y \subset X(K)$ define*

$$I_k(Y) := \{f \in k[X] \mid f(x) = 0 \quad \forall x \in Y\}$$

*Suppose $\pi : k[X_1, ..., X_n] \to k[X]$ is the quotient map. Then we have the following properties*

© David Rufino CC BY-NC-ND 4.0

$$V_K(\mathfrak{b}) = V_K(\pi^{-1}(\mathfrak{b})) \cap X(K)$$
$$I_k(Y \cap X(K)) = \pi(I_k(Y)) \quad Y \subset \mathbb{A}_k^n(K)$$
$$I_k(V_K(\mathfrak{b})) = \pi(I_k(V_K(\pi^{-1}(\mathfrak{b}))))$$

*The pair $(V_K, I_k)$ satisfies the same properties as (...). The image of $V_K$ forms the closed sets of a topology on $X(K)$ which coincides with the subspace topology from $\mathbb{A}_k^n(K)$.*

We may now generalize the Weak Nullstellensatz to arbitrary affine varieties and non-algebraically closed fields $K/k$.

**Proposition 6.1.9 (Galois Group Action)**
*Let $X = V(\mathfrak{a})$ be an affine variety and $K/k$ a* normal *algebraic field extension (e.g. $\overline{k}$). There is a well-defined group action*

$$\mathrm{Aut}(K/k) \times X(K) \to X(K)$$
$$(\sigma, (x_1, ..., x_n)) \to (\sigma(x_1), ..., \sigma(x_n))$$

*For $x, y \in X(K)$ the following are equivalent*

*a) $x$ and $y$ are topologically indistinguishable*
*b) $x = \sigma(y)$ for some $\sigma \in \mathrm{Aut}(K/k)$*
*c) $x \in \overline{\{y\}}$*
*d) $y \in \overline{\{x\}}$*
*e) $\overline{\{x\}} = \overline{\{y\}}$*
*f) $\mathfrak{M}_{X,x} = \mathfrak{M}_{X,y}$*

*For every point $x$ we have*

$$V_K(\mathfrak{M}_{X,x}) = \overline{\{x\}} = \{\sigma(x) \mid \sigma \in \mathrm{Aut}(K/k)\}$$

*and $x$ is quasi-closed (4.1.90) and therefore $X(K)$ is symmetric. Denote the set of Galois Orbits by $X_0(K)$ then the quotient map*

$$\pi : X(K) \to X_0(K)$$

*is precisely the Kolmogorov Quotient (4.1.36) and in particular a quasi-homeomorphism.*

*Proof.* Given $F \in \mathfrak{a}$ we have $\sigma(F(x)) = F(\sigma(x))$. This shows that $x \in X(K) \implies \sigma(x) \in X(K)$. Further it's clear that $\sigma(\tau(x)) = (\sigma \circ \tau)(x)$ so the group action is well-defined.

By (4.1.34) a) $\iff$ e) $\implies$ c), d). By (6.1.2) $V(\mathfrak{M}_{X,x}) = \overline{\{x\}}$ and $I(\overline{\{x\}}) = \mathfrak{M}_{X,x}$. So e) $\iff$ f). Suppose $\mathfrak{M}_{X,x} = \mathfrak{M}_{X,y}$ then this means there exists an isomorphism $k(x) \cong k(y) \subset K$ which lifts to an element of $\mathrm{Aut}(K/k)$ by (3.19.69), so f) $\implies$ b). Conversely b) $\implies$ f) is straightforward. If $x \in \overline{\{y\}} = V(\mathfrak{M}_{X,y})$ then $\mathfrak{M}_{X,y} \subset \mathfrak{M}_{X,x}$ whence they are equal by maximality, so c), d) $\implies$ f). ∎

**Corollary 6.1.10**
*Let $k$ be an algebraically closed field and $X$ an affine variety. Then $X(k)$ is a Kolmogorov topological space.*

**Proposition 6.1.11 (Generalized Weak Nullstellensatz)**
*Let $X = V(\mathfrak{a})$ be an affine variety. There is a bijection between maximal ideals and Galois orbits of points*

$$\mathrm{Specm}(k[X]) \longleftrightarrow X_0(\overline{k}) = X(\overline{k})/\mathrm{Aut}(\overline{k}/k)$$
$$\mathfrak{M} \to [V_K(\mathfrak{M})]$$
$$\mathfrak{M}_{X,x} \leftarrow [x]$$

*Proof.* We may deduce this from (3.35.24) by observing that $x \in X(\overline{k}) \iff \mathfrak{a} \subset \mathfrak{m}_x$, $\mathfrak{M}_{X,x} = \mathfrak{m}_x/\mathfrak{a}$ and using the correspondence of ideals of quotient ring (3.3.56)

$$\{\mathfrak{M} \in \mathrm{Specm}(k[X_1, ..., X_n]) \mid \mathfrak{a} \subset \mathfrak{M}\} \longleftrightarrow \mathrm{Specm}(k[X]).$$

∎

**Lemma 6.1.12 (Rabinowitsch Trick)**
*Let $\mathfrak{a} \lhd A$ and $f \in A$. Consider the ring $B = A[Y]$. If $\mathfrak{a}B + (1 - Yf) = B$ then $f \in \sqrt{\mathfrak{a}}$.*

*Proof.* The hypothesis implies

© David Rufino CC BY-NC-ND 4.0

$$1 = (1 - Yf)g(Y) + ah(Y)$$

for $a \in \mathfrak{a}$ and $h(Y) \in A[Y]$. Consider the quotient map $\bar{\cdot} : A \to A/\mathfrak{a}$ and the corresponding map $A[Y] \to (A/\mathfrak{a})[Y]$. Applying this to the above shows $1 - Y\overline{f}$ is invertible in $(A/\mathfrak{a})[Y]$. So by (3.8.4) $\overline{f}$ is nilpotent in $(A/\mathfrak{a})$ whence $f \in \sqrt{\mathfrak{a}}$. ∎

**Proposition 6.1.13 (Strong Nullstellensatz)**
*Let $X$ be an affine variety and $\mathfrak{a} \lhd k[X]$. Then*

$$I_k V_{\overline{k}}(\mathfrak{a}) = \sqrt{\mathfrak{a}}^J = \sqrt{\mathfrak{a}}$$

*In particular the $\overline{k}$-radical ideals are precisely the radical ideals. and there is a dual isomorphism*

$$\mathrm{Rad}(k[X]) \longleftrightarrow \mathrm{Zar}_k\left(X\left(\overline{k}\right)\right)$$

*Proof.* Observe that $x \in V_{\overline{k}}(\mathfrak{a})$ if and only if $\mathfrak{a} \subseteq \mathfrak{M}_{X,x}$. Therefore

$$I_k(V_{\overline{k}}(\mathfrak{a})) = \bigcap_{x \in V_{\overline{k}}(\mathfrak{a})} I_k(\{x\}) = \bigcap_{\mathfrak{a} \subseteq \mathfrak{M}_{X,x}} \mathfrak{M}_{X,x}$$

Therefore by the correspondence in (6.1.11) this is equal to precisely $\sqrt{\mathfrak{a}}^J$.

For the second equality we consider first the case $X = \mathbb{A}^n_k$ and $k[X] = k[X_1, ..., X_n]$. Let $\mathfrak{a} \lhd k[X]$ and choose $f \in I_k V_{\overline{k}}(\mathfrak{a})$. Consider the ring $B \coloneqq k[X_1, ..., X_n, Y]$ and the ideal $\tilde{\mathfrak{a}} = \mathfrak{a}B + (1 - Yf)$. Clearly this has no zeros in $\overline{k}^{n+1}$, so by the Weak Nullstellensatz (3.35.24) it is not proper. By the (6.1.12) $f \in \sqrt{\mathfrak{a}}$ as required. The reverse inclusion is clear.

Now suppose that $X = V(\mathfrak{a})$, $k[X] = k[X_1, ..., X_n]/\mathfrak{a}$ and $\mathfrak{b} \lhd k[X]$ is proper. Using (6.1.8) and (3.3.50) together with the result just proven, shows

$$I_k(V_{\overline{k}}(\mathfrak{b})) = \pi(I_{\overline{k}} V_{\overline{k}}(\pi^{-1}\mathfrak{b})) = \pi\left(\sqrt{\pi^{-1}(\mathfrak{b})}\right) = \sqrt{\mathfrak{b}}$$

where $\pi : k[X_1, ..., X_n] \to k[X]$ is canonical surjective morphism. ∎

**Corollary 6.1.14**
*Let $k[X]$ be any finitely generated reduced $k$-algebra, then $k[X]$ is a Jacobson ring, i.e.*

$$\sqrt{\mathfrak{a}} = \sqrt{\mathfrak{a}}^J$$

*In particular the intersection of all maximal ideals is zero*

$$\bigcap_{\mathfrak{m}} \mathfrak{m} = 0$$

## 6.1.1 Topological Properties

**Proposition 6.1.15 (Principal Open Sets)**
*Let $X = V(\mathfrak{a})$ be an affine variety over $k$. For every $f \in k[X]$ the sets*

$$D(f) \coloneqq X\left(\overline{k}\right) \setminus V_{\overline{k}}((f)) = \left\{(x) \in X\left(\overline{k}\right) \;\middle|\; f(x) \neq 0\right\}$$

*form a base for the Zariski topology $\mathrm{Zar}_k\left(X\left(\overline{k}\right)\right)$.*

*Proof.* Let $U = X\left(\overline{k}\right) \setminus V(\mathfrak{b})\left(\overline{k}\right)$ be a non-empty open set. For every $x \in U$ there by definition exists $f \in \mathfrak{b}$ such that $f(x) \neq 0$. Therefore $D(f)$ is a neighbourhood of $x$ in the Zariski topology. ∎

The topological notion of irreducibility is important, and may be reduced to a purely algebraic statement on the coordinate ring.

**Proposition 6.1.16 (Criterion for Irreducibility)**
*Let $X$ be an affine variety (e.g. $\mathbb{A}^n_k$) and $Y = V(\mathfrak{b})$ an affine subvariety corresponding to a radical ideal $\mathfrak{b} \lhd k[X]$. Then the following are equivalent*

*a) $Y\left(\overline{k}\right)$ is an irreducible subset of $X\left(\overline{k}\right)$*
*b) $Y_0\left(\overline{k}\right)$ is an irreducible subset of $X_0\left(\overline{k}\right)$*
*c) $\mathfrak{b}$ is prime*
*d) $k[Y]$ is an integral domain.*

© David Rufino CC BY-NC-ND 4.0

*Proof.* Suppose $X$ is not irreducible. Then we have $X \subseteq V_{\overline{k}}(\mathfrak{b}) \cup V_{\overline{k}}(\mathfrak{c})$ a non-trivial decomposition into closed subsets (and associated ideals). Then by the dual isomorphism we have also $\mathfrak{a} \subsetneq \mathfrak{b}$ and we may choose $f \in \mathfrak{b} \setminus \mathfrak{a}$ and similarly $g \in \mathfrak{c} \setminus \mathfrak{a}$. However $fg$ vanishes on $X\left(\overline{k}\right)$ ands so we have $fg \in \mathfrak{a}$. Therefore $\mathfrak{a}$ is not prime.

Conversely suppose $X$ is irreducible and $\mathfrak{bc} \subseteq \mathfrak{a}$. Then $X \subseteq V_{\overline{k}}(\mathfrak{b}) \cup V_{\overline{k}}(\mathfrak{c})$. By irreducibility we have $X \subseteq V_{\overline{k}}(\mathfrak{b})$, whence applying $I_{\overline{k}}(-)$ we see $\mathfrak{b} \subseteq I_{\overline{k}}V_{\overline{k}}(\mathfrak{b}) \subseteq \mathfrak{a}$ (since $\mathfrak{a}$ is radical). Therefore $\mathfrak{a}$ is prime. ∎

When $W \subset X$ is an irreducible subset then write $\mathfrak{M}_{X,W} \coloneqq I(W)$, by the previous result this equals $I\left(\overline{W}\right)$ and $\mathfrak{M}_{X,W}$ is a prime ideal.

**Proposition 6.1.17**
*Let $X$ be an affine variety, $W, Z$ be subsets of $X\left(\overline{k}\right)$. Then*

$$W \subset \overline{Z} \iff \mathfrak{M}_{X,Z} \subset \mathfrak{M}_{X,W}$$

*and*

$$\overline{W} = \overline{Z} \iff \mathfrak{M}_{X,Z} = \mathfrak{M}_{X,W}$$

*Proof.* Observe that $V\left(\mathfrak{M}_{X,W}\right) = \overline{W}$ by (6.1.2). Therefore the result follows immediately from the same result. ∎

**Proposition 6.1.18 (Points are Closed)**
*Let $X = V(\mathfrak{a})$ be an affine variety and $x \in X\left(\overline{k}\right)$. Then*

$$\overline{\{x\}} = V\left(\mathfrak{M}_{X,x}\right) = \left\{\sigma(x) \;\middle|\; \sigma \in \mathrm{Aut}\left(\overline{k}/k\right)\right\}$$

*is irreducible and finite of order $[k(x) : k]_s$. Further $X$ is a symmetric space.*

*Proof.* The equalities are proven in (6.1.9). Further $\overline{\{x\}}$ is irreducible by (4.1.67). We may replace $\mathrm{Aut}\left(\overline{k}/k\right)$ with $\mathrm{Mor}_k\left(k(x), \overline{k}\right)$ to determine the order.

Clearly $X$ is symmetric because it satisfies criteria (4.1.41) b). ∎

**Proposition 6.1.19 (Closed sets $\leftrightarrow$ radical ideals)**
*Let $X = V(\mathfrak{a})$ be an affine variety. Recall (6.1.13) there is a dual lattice isomorphism between radical ideals and "Zariski"-closed subsets of $X\left(\overline{k}\right)$ (and indeed $X_0\left(\overline{k}\right)$).*

$$\mathrm{Rad}(k[X]) \underset{I_k(-)}{\overset{V_{\overline{k}}(-)}{\rightleftarrows}} \mathrm{Zar}_k\left(X\left(\overline{k}\right)\right)$$

*Under this isomorphism we have*

- *maximal ideals correspond to $\mathrm{Aut}\left(\overline{k}/k\right)$-orbits of $X\left(\overline{k}\right)$ (or elements of $X_0\left(\overline{k}\right)$)*
- *prime ideals of $k[X]$ correspond to irreducible subsets of $X\left(\overline{k}\right)$ (and therefore of $X_0\left(\overline{k}\right)$)*
- *minimal prime ideals of $k[X]$ correspond to irreducible components of $X\left(\overline{k}\right)$ (and therefore of $X_0\left(\overline{k}\right)$)*

*Recall that prime ideals are precisely the meet-prime radical ideals and irreducible subsets are precisely the join-prime closed subsets (see (4.1.61)). Therefore we have a dual isomorphism between the Krull Lattice of radical ideals of $k[X]$ and the Krull Latice of closed subsets of $X\left(\overline{k}\right)$.*

*Proof.* The content of the Strong Nullstellensatz is precisely that $I_k(-) \circ V_{\overline{k}}(-) = \mathbf{1}$. The other direction was already proven so we have a dual order isomorphism. The statement about maximal ideals was already shown in (6.1.11) and prime ideals in (6.1.16). Then as irreducible components are precisely maximal irreducible subsets the final statement follows from the dual order isomorphism. ∎

**Corollary 6.1.20**
*Let $X = V(\mathfrak{a})$ be an affine variety. There is a bijective correspondence between affine subvarieties of $X$ and closed subsets of $X\left(\overline{k}\right)$.*

**Definition 6.1.21 (Irreducible Affine Variety)**
*We say an affine variety $X = V(\mathfrak{a})$ is **integral** or **irreducible** if the topological space $X\left(\overline{k}\right)$ is irreducible.*

*This is the case precisely when $\mathfrak{a}$ is prime, or when $k[X]$ is an integral domain by (6.1.19), or equivalently irreducible (3.3.67), since it is assumed to be reduced.*

*Note trivially that $\mathbb{A}^n_k$ is irreducible.*

© David Rufino CC BY-NC-ND 4.0

**Proposition 6.1.22 (Decomposition into Irreducible Components)**
*Let $X = V(\mathfrak{a})$ be an affine variety then the topological space $X\left(\overline{k}\right)$ is Noetherian. Furthermore it has finitely many irreducible components $X_i$ and the decomposition*

$$X\left(\overline{k}\right) = X_1 \cup ... \cup X_n$$

*is the unique incomparable decomposition into irreducible closed subsets.*

*Proof.* By Hilbert's Basis Theorem (3.15.8) $k[X]$ is a Noetherian ring, so by (6.1.19) $X\left(\overline{k}\right)$ is Noetherian. The result then follows from (4.1.76) ∎

**Proposition 6.1.23 (Subspace Topology)**
*Let $X = V(\mathfrak{a})$ an affine variety and $Y = V(\mathfrak{b})$ an affine subvariety. Then there is a commutative diagram*

$$\begin{array}{ccc} \{\mathfrak{c} \in \mathrm{Rad}(k[X]) \mid \mathfrak{b} \subseteq \mathfrak{c}\} & \underset{I}{\overset{V}{\rightleftarrows}} & \left\{Z \in \mathrm{Zar}_k\left(X\left(\overline{k}\right)\right) \;\middle|\; Z \subseteq Y\left(\overline{k}\right)\right\} \\ \pi^{-1} \uparrow \;\downarrow \pi & & \| \\ \mathrm{Rad}(k[X]/\mathfrak{b}) & \underset{I}{\overset{V}{\rightleftarrows}} & \mathrm{Zar}_k\left(Y\left(\overline{k}\right)\right) \end{array}$$

*under which prime ideals correspond to irreducible subsets and the horizontal arrows induce dual isomorphisms.*

*In particular the subspace topology for $Y\left(\overline{k}\right)$ coincides with the Zariski topology.*

*Proof.* The left hand arrows are mutual inverses by (3.3.56). The horizontal maps are dual isomorphisms by (6.1.19). The equality then follows from (6.1.8). ∎

**Proposition 6.1.24**
*Let $X$ be an affinve variety. The correspondence between points and ideals (6.1.11) restricts as follows*

$$\mathrm{Specm}\left(k[X]_f\right) \longleftrightarrow \{\mathfrak{M} \in \mathrm{Specm}(k[X]) \mid f \notin \mathfrak{M}\} \longleftrightarrow D(f)_0 := D(f)/\operatorname{Aut}\left(\overline{k}/k\right)$$

*More precisely a point $x \in D(f)$ corresponds to the maximal ideal $\left(\mathfrak{M}_{X,x}\right)_f$ of $k[X]_f$.*

*Proof.* For the first correspondence consider the localisation map $i : k[X] \to k[X]_f$. By (3.6.27) the extension of a maximal ideal not containing $f$ is maximal. Conversely $i$ satisfies the hypotheses of (3.35.27) so the inverse image of a maximal ideal is maximal. This shows the inverse image of a maximal ideal is maximal, not containing $f$. Therefore the correspondence of prime ideals (3.6.36) restricts to maximal ideals.

The second correspondence is immediate from (6.1.11) because $f \notin \mathfrak{M}_{X,x} \iff f(x) \neq 0$. ∎

## 6.1.2 Structure Sheaf

**Definition 6.1.25 (Sheaf of Regular Functions)**
*Let $X$ be an affine variety, $U \subset X\left(\overline{k}\right)$ an open subset and $f : U \to \overline{k}$. We say that $f$ is **regular** if for every $x \in U$ there exists $g, h \in k[X]$ and an open neighbourhood $V$ of $x$ such that*

$$f(y) = \frac{g(y)}{h(y)} \quad \forall y \in V$$

*We denote the **structure sheaf** by*

$$\mathcal{O}_X(U) := \left\{f : U \to \overline{k} \;\middle|\; f \text{ regular}\right\}$$

*which is a sheaf of $k$-algebras.*

*If $U \subset X$ is an open subset then we denote the structure sheaf by the restriction*

$$\mathcal{O}_U := \mathcal{O}_X \mid_U$$

**Proposition 6.1.26 (Sections are localisation of coordinate ring)**
*Let $X = V(\mathfrak{a})$ be an affine variety and $f \in k[X]$. There is a canonical isomorphism*

© David Rufino CC BY-NC-ND 4.0

$$\rho_f : k[X]_f \xrightarrow{\sim} \mathcal{O}_X(D(f))$$
$$\frac{g}{f^n} \to \frac{g(-)}{f(-)^n}$$

*Furthermore* $D(g) \subseteq D(f) \iff S_f \subseteq \overline{S_g}$ *and we have the following commutative diagram*

$$\begin{array}{ccc} k[X]_f & \xrightarrow{\sim} & \mathcal{O}_X(D(f)) \\ \downarrow{\scriptstyle i_{S_f S_g}} & & \downarrow{\scriptstyle (-)|_{D(g)}} \\ k[X]_g & \xrightarrow{\sim} & \mathcal{O}_X(D(g)) \end{array}$$

*In particular there is a canonical isomorphim*

$$k[X] \xrightarrow{\sim} \mathcal{O}_X(X)$$

*Proof.* The map is trivially well-defined and injective. Consider a regular map $\sigma \in \mathcal{O}_X(D(f))$. Consider the ideal

$$\mathfrak{a} \coloneqq \left\{ g \in k[X] \mid g\sigma \in \mathrm{Im}\left(i_f\right) \right\}$$

It is enough to show $f \in \mathfrak{a}$. Suppose $f \notin \mathfrak{a}$ then it's contained in a maximal ideal which is of the form $\mathfrak{M}_{X,x}$ for some $x \in X\left(\overline{k}\right)$ by (6.1.11). By definition $x \in D(f)$ and there is an open neighbourhood $W \subseteq D(f)$ and elements $h_1, h_2 \in k[X]$ such that

$$\sigma(y) = h_1 \frac{y}{h_2}(y) \quad \forall y \in W$$

Choose $h_3 \in k[X]$ such that $x \in D(h_3) \subseteq W$ then clearly

$$\sigma(y) = (h_1 h_3)\frac{y}{h_2 h_3}(y) \quad \forall y \in D(h_3)$$

and in particular $(h_2 h_3 \sigma)(y) = (h_1 h_3)(y)$ for all $y \in D(f)$. Therefore $h_2 h_3 \in \mathfrak{a} \subseteq \mathfrak{m}_x$ which implies $h_2(x) = 0$ or $h_3(x) = 0$ a contradiction.

Therefore $f \in \mathfrak{a}$ and clearly $\sigma \in \mathrm{Im}\left(i_f\right)$ as required. ∎

**Corollary 6.1.27**
*Suppose* $f \in k[X]$ *is a regular function then* $f \neq 0 \implies D(f) \neq \varnothing$.

In what follows we may systematically identify $\mathcal{O}_X(X)$ and $k[X]$, and we formalise this in the following result.

**Proposition 6.1.28 (Properties of the Structure Sheaf)**
*Let* $X = V(\mathfrak{a})$ *be an affine variety and let* $\widehat{\cdot} : k[X] \xrightarrow{\sim} \mathcal{O}_X(X)$ *be the canonical isomorphism defined in (6.1.26). Then the following properties hold*

*a) For all* $f \in \mathcal{O}_X(X)$ *we have* $D(f) = \{x \in U \mid f(x) \neq 0\}$ *is open and* $f|_{D(f)}$ *is invertible.*
*b) For an irreducible subset* $W \subset X$ *define the prime ideal*

$$\mathfrak{M}_{X,W} \coloneqq \{f \in \mathcal{O}_X(X) \mid f(x) = 0 \quad \forall x \in W\}$$

*c) For* $x \in X$ *we have* $\mathfrak{M}_{X,x}$ *is maximal and every maximal ideal is of this form*
*d) For* $W, Z \subset X$ *be irreducible subsets then*

$$\mathfrak{M}_{X,Z} \subset \mathfrak{M}_{X,W} \iff W \subset \overline{Z}$$

*and*

$$\mathfrak{M}_{X,Z} = \mathfrak{M}_{X,W} \iff \overline{W} = \overline{Z}$$

*In particular* $\mathfrak{M}_{X,W} = \mathfrak{M}_{X,\overline{W}}$
*e) For all* $f \in \mathcal{O}_X(X)$ *we have a canonical isomorphism*

$$\begin{array}{ccc} \mathcal{O}_X(X) & & \\ \downarrow & \searrow^{\rho_{X D(f)}} & \\ \mathcal{O}_X(X)_f & \xrightarrow{\sim} & \mathcal{O}_X(D(f)) \end{array}$$

© David Rufino CC BY-NC-ND 4.0

*which is the unique homomorphism making this diagram commute*

For $X$ an affine variety we denote by $\overline{X}_1, ..., \overline{X}_n$ the image of the coordinate functions under $\hat{\cdot}$.

### 6.1.3 Regular Morphisms of Affine Varieties

**Definition 6.1.29 (Regular Morphism of Affine Varieties)**
*Let $X \subset \mathbb{A}_k^n$, $Y \subset \mathbb{A}_k^m$ be affine varieties and $\phi : X\left(\overline{k}\right) \to Y\left(\overline{k}\right)$ a function. Then we say that $\phi$ is a **regular morphism** (at $x$) if $\pi_j \circ \phi$ is regular (at $x$) in the sense of (6.1.25) for $j = 1, ..., m$.*

*A regular morphism is an isomorphism if it has a two-sided inverse which is also regular.*

*Note a regular function $X \to \overline{k}$ is completely equivalent to a regular morphism $X \to \mathbb{A}_k^1$.*

**Proposition 6.1.30 (Regular Morphisms of Affine Varieties)**
*Let $X = V(\mathfrak{a})$ and $Y = V(\mathfrak{b})$ be affine varieties. Then there is a bijection*

$$\begin{aligned} \mathrm{AlgHom}_k(\mathcal{O}_Y(Y), \mathcal{O}_X(X)) &\longleftrightarrow \{\phi : X \to Y \mid \phi \text{ regular}\} \\ \phi^\sharp \to & \left(\phi^\sharp\left(\overline{Y}_1\right)(-), ..., \phi^\sharp\left(\overline{Y}_m\right)(-)\right) \\ \overline{G} \to G(\pi_1 \circ \phi, ..., \pi_m \circ \phi_m) \leftarrow & \phi \end{aligned}$$

*Proof.* Let $\phi^\sharp$ be a $k$-algebra homomorphism, then $\phi^\sharp\left(\overline{Y}_i\right) \in \mathcal{O}_Y(Y)$ is a regular function. Therefore by definition the first map is well-defined. Conversely $\pi_j \circ \phi$ is regular so corresponds to an element of $\phi_j \in \mathcal{O}_X(X)$ and the second map is well-defined. As $\phi$ is uniquely determined by $\phi^\sharp\left(Y_j\right)$ for $j = 1, ..., m$ we see that the maps are mutually inverse. ■

**Lemma 6.1.31**
*Let $\phi : X \to Y$ be a regular morphism of affine varieties and $W \subset X$ an irreducible subset. Then*

$$\left(\phi^\sharp\right)^{-1}\left(\mathfrak{M}_{X,W}\right) = \mathfrak{M}_{Y,\phi(W)} = \mathfrak{M}_{Y,\overline{\phi(W)}}$$

*Proof.* This follows immediately from the definitions and (6.1.28). ■

**Proposition 6.1.32**
*Let $\phi : X \to Y$ be a regular morphism of affine varieties and $W \subset X$, $Z \subset Y$ irreducible. Then*

$$Z \subset \overline{\phi(W)} \iff \left(\phi^\sharp\right)^{-1}\left(\mathfrak{M}_{X,W}\right) \subset \mathfrak{M}_{Y,Z}$$

*and*

$$\overline{Z} = \overline{\phi(W)} \iff \left(\phi^\sharp\right)^{-1}\left(\mathfrak{M}_{X,W}\right) = \mathfrak{M}_{Y,Z}$$

*Proof.* This follows from (6.1.28) and (6.1.31). ■

**Proposition 6.1.33**
*Let $\phi : X \to Y$ be a regular morphism of affine varieties and $x \in X$, $y \in Y$ points. Then the following are equivalent*

*a)* $y \in \overline{\{\phi(x)\}}$
*b)* $\phi(x) \in \overline{\{y\}}$
*c)* $\overline{\{\phi(x)\}} = \overline{\{y\}}$
*d)* $\left(\phi^\sharp\right)^{-1}\left(\mathfrak{M}_{X,x}\right) = \mathfrak{M}_{Y,y}$

*When $k$ is algebraically closed then this is equivalent to $\phi(x) = y$.*

*Proof.* We have a)-c) are equivalent because $Y$ is symmetric (6.1.9), (4.1.42), and then c) $\iff$ d) follows from (6.1.32). ■

**Corollary 6.1.34**
*Let $\phi : X \to Y$ be a regular morphism of affine varieties. Then $\phi$ is continuous.*

*Proof.* Observe that $\phi^{-1}(D(f)) = D(f \circ \phi) = D\left(\phi^\sharp(f)\right)$ so we are done by (6.1.15) and (4.1.27). ■

**Corollary 6.1.35**
*Let $\phi : X \to Y$ and $\psi : Y \to Z$ be regular morphisms of affine varieties. Then $\psi \circ \phi$ is regular and*

$$(\psi \circ \phi)^\sharp = \phi^\sharp \circ \psi^\sharp$$

**Proposition 6.1.36 (Affine Varieties form a Category)**
*The collection of affine varieties together with regular morphisms form a category $\mathbf{AffVar}_k$. Furthermore there is an equivalence of categories*

© David Rufino CC BY-NC-ND 4.0

$$\mathbf{AffVar}_k \xrightarrow{\sim} \mathbf{ReducedFgAlg}_k^{op}$$

*Proof.* By using the correspondence with $k$-algebra homomorphisms (6.1.30) we may show that the law of composition for regular morphisms also satisfy the axioms of a category. We have therefore shown that the assignment $X \to k[X]$ is full and faithful functor. To show it is essentially surjective consider a reduced finitely-generated $k$-algebra $A$. Then by definition $A \cong k[X_1, ..., X_n]/\mathfrak{a}$ for some ideal $\mathfrak{a}$. As $A$ is reduced we see $\mathfrak{a}$ is radical and therefore $A$ is the coordinate ring of the affine variety $X = V(\mathfrak{a})$ and by definition $k[X] \cong A$. ∎

**Lemma 6.1.37**
*Let $\phi : X \to Y$ be a regular morphism of affine varieties. Suppose that $V \subset Y$ is open and $f : V \to \overline{k}$ is regular. Then $f \circ \phi : \phi^{-1}(V) \to \overline{k}$ is regular.*

*In other words there is a well-defined functor*

$$\mathbf{AffVar}_k \to \mathbf{LocalSpaceFunctions}_k$$

*which we show is full and faithful in (6.1.39).*

**Proposition 6.1.38 (Principal Open Sets are Affine)**
*Let $X = V(\mathfrak{a})$ be an affine variety and $f \in \mathcal{O}_X(X)$. Define $Y \coloneqq V(\mathfrak{a}^e + (1 - X_{n+1}F)) \subset \mathbb{A}_k^{n+1}$ where $f = \overline{F}$ and $F \in k[X_1, ..., X_n]$. Then there is an isomorphism of spaces with functions*

$$\theta : \left(D(f), \mathcal{O}_X|_{D(f)}\right) \to \left(Y\left(\overline{k}\right), \mathcal{O}_Y\right)$$
$$(x_1, ..., x_n) \to \left(x_1, ..., x_n, \frac{1}{F(x_1, ..., x_n)}\right)$$

*Proof.* Observe $V(\mathfrak{a}^e + (1 - X_{n+1}F)) = V(\mathfrak{a}^e) \cap V(1 - X_{n+1}F)$. Therefore $\theta$ is well-defined. Conversely given $(x_1, ..., x_{n+1}) \in V(\mathfrak{a}^e + (1 - X_{n+1}F)) = V(\mathfrak{a}^e) \cap V(1 - X_{n+1}F)$, by definition $(x_1, ..., x_n) \in V(\mathfrak{a})$ and $F(x_1, ..., x_n)x_{n+1} = 1 \implies F(x_1, ..., x_n) \neq 0$. Therefore the inverse map is well-defined. Evidently this restricts to a bijection

$$D\left(\overline{G}\right) \cap D(F) \longleftrightarrow D(G) \cap V(\mathfrak{a}^e + (1 - X_{n+1}F))$$

By (6.1.24) and (4.1.27) $\theta$ is a homeomorphism.

Suppose that $V \subset Y$ is open and $g : V \to \overline{k}$ is regular, then for every $y \in V$ there is some neighbourhood $W$ for which

$$g(z_1, ..., z_{n+1}) = \frac{P(z_1, ..., z_{n+1})}{Q(z_1, ..., z_{n+1})} \quad \forall z \in W$$

where $P, Q \in k[Y_1, ..., Y_{n+1}]$ and $Q(z) \neq 0$. Define the rational functions in $k(X_1, ..., X_n)$ as follows

$$R(X_1, ..., X_n) \coloneqq P\left(X_1, ..., X_n, F(X_1, ..., X_n)^{-1}\right)$$
$$S(X_1, ..., X_n) \coloneqq Q\left(X_1, ..., X_n, F(X_1, ..., X_n)^{-1}\right)$$

Let $U \coloneqq \theta^{-1}(W)$. Then evidently $R(x) = P(\theta(x))$ and $S(x) = Q(\theta(x))$ for all $x \in U$. In particular we deduce that $S(\theta^{-1}(y)) \neq 0 \implies S \neq 0$ and that $g \circ \theta$ is regular at $\theta^{-1}(y)$. The converse is straightforward so we deduce that $\theta$ is an isomorphism. ∎

When considering regular maps over an open subset of an affine variety, there are two plausible definitions which we show are equivalent.

**Proposition 6.1.39 (Quasi-affine Morphisms)**
*Let $X \subset \mathbb{A}_k^n$, $Y \subset \mathbb{A}_k^m$ be affine varieties, $U \subset X\left(\overline{k}\right)$ an open subset and $\phi : U \to Y\left(\overline{k}\right)$ a function. Then the following are equivalent*

*a) $\pi_j \circ \phi$ is regular for all $j = 1, ..., m$.*
*b) $\phi$ determines a regular morphism $(U, \mathcal{O}_X|_U) \to (Y, \mathcal{O}_Y)$ of spaces with functions.*

*If $V \subset Y\left(\overline{k}\right)$ is an open subset and $\phi(U) \subset V$ then $\phi$ determines a regular morphism $(U, \mathcal{O}_X|_U) \to (V, \mathcal{O}_Y|_V)$.*

*Proof.* b) $\implies$ a) follows trivially because $\pi_j : Y\left(\overline{k}\right) \to \overline{k}$ is regular.

a) $\implies$ b) Firstly we observe that for any $f \in \mathcal{O}_X(X)$ for which $D(f) \subset U$ there is a commutative diagram

© David Rufino CC BY-NC-ND 4.0

$$\begin{array}{ccc} D(f) & \xrightarrow{\phi|_{D(f)}} & Y\left(\overline{k}\right) \\ \sim\downarrow\theta & \nearrow_{\tilde{\phi}_f} & \\ V(\mathfrak{a}^e + (1 - X_{n+1}F)) & & \end{array}$$

where $\theta$ is defined in (6.1.38) and $\tilde{\phi}_f(x_1, ..., x_{n+1}) = \phi(x_1, ..., x_n)$. By definition $\tilde{\phi}_f$ is a regular morphism of affine varieties and therefore determines a regular morphism of spaces of functions (6.1.37). As $\theta$ is a regular isomorphism we find that $\phi|_{D(f)}$ is a regular morphism of spaces of functions. As the $D(f)$ form a base on the topology of $U$ we may conclude from (4.3.27) that $\phi$ is a regular morphism of spaces with functions. ■

**Proposition 6.1.40**
*Let $X$, $Y$, $Z$ be affine varieties with $U \subset X$ and $V \subset Y$ open subsets. Suppose that $\phi : U \to V$ is regular and $\psi : V \to Z$ is regular, then $\psi \circ \phi : U \to Z$ is regular.*

**Corollary 6.1.41**
*Let $X$ be an affine variety, $U \subset X$ an open subset and $f \in \mathcal{O}_X(U)$. Then the set*

$$D(f) \coloneqq \{x \in U \mid f(x) \neq 0\}$$

*is open and $f|_{D(f)}$ is invertible.*

*Proof.* By (6.1.39) we may regard $f$ as a regular morphism $(U, \mathcal{O}_X|_U) \to \mathbb{A}^1_k\left(\overline{k}\right)$, and it is in particular continuous. Then $\{0\} = V(X_1)$ is a closed subset of $\mathbb{A}^1_k\left(\overline{k}\right)$ and so $D(f) = U \setminus f^{-1}(\{0\})$ is an open subset. It is clear by the definition of the structure sheaf that $f|_{D(f)}$ is invertible. ■

**Corollary 6.1.42 (Affine Mapping Property)**
*Let $X, Y$ be affine varieties. Then there are bijections*

$$\mathrm{Mor}(X, Y) \xrightarrow{\sim} \mathrm{Mor}((X, \mathcal{O}_X), (Y, \mathcal{O}_Y)) \xrightarrow{\sim} \mathrm{AlgHom}_k(\mathcal{O}_Y(Y), \mathcal{O}_X(X))$$

*where the second map is given by $\phi^\sharp_Y$. Furthermore these are natural in either $X$ or $Y$, and preserve isomorphisms.*

*Proof.* The first map is well-defined by (6.1.39) and evidently injective. The composite map is bijective by (6.1.30), which shows the first map is surjective and hence bijective. We conclude trivially the second map is also a bijection. ■

**Proposition 6.1.43 (Equaliser is Closed)**
*Let $\phi, \psi : X \to Y$ be regular morphisms of affine varieties. Then the equaliser*

$$\left\{x \in X\left(\overline{k}\right) \;\middle|\; \phi(x) = \psi(x)\right\}$$

*is closed.*

*Proof.* By applying the projections we may reduce to the case $Y = \mathbb{A}^1_k$. Then $\phi - \psi$ is regular and the equaliser is the inverse image of $\{0\}$ which is evidently closed in $\mathbb{A}^1_k$. As regular maps are continuous then we are done. ■

**Proposition 6.1.44**
*Let $Y = V(\mathfrak{a})$ be an affine variety and $Z \coloneqq V(\mathfrak{b})$ a closed subvariety for $\mathfrak{a} \lhd k[Y]$. Then the space of functions $(Z, \mathcal{O}_Z)$ coincides with the notion of a closed subspace (4.3.21).*

*Further the inclusion map $i : Z \hookrightarrow Y$ is regular and the induced map*

$$i^\sharp_Y : \mathcal{O}_Y(Y) \longrightarrow \mathcal{O}_Z(Z)$$

*is a surjective homomorphism with kernel precisely $\rho(\mathfrak{b})$ where $\rho : k[Y] \xrightarrow{\sim} \mathcal{O}_Y(Y)$ is the canonical isomorphism.*

*Proof.* We have a surjective homomorphism $\pi : k[Y] \to k[Z] \coloneqq k[Y]/\mathfrak{b}$ with kernel precisely $\mathfrak{b}$. Suppose $f \in \mathcal{O}_Y(U)$ is a regular map, then we wish to show that $f|_{U \cap Z}$ is regular. But if $f$ is given locally by a ratio $\frac{g}{h}$ then $f|_{U\cap Z} = i^\sharp_U(f)$ is given locally by the ratio $\frac{\pi(g)}{\pi(h)}$. And because $\pi$ is surjective this means $i^\sharp_Y$ is surjective. Furthermore we have a commutative diagram

$$\begin{array}{ccc} \mathcal{O}_Y(Y) & \xrightarrow{i^\sharp} & \mathcal{O}_Z(Z) \\ \sim\uparrow\rho & & \sim\uparrow\rho \\ k[Y] & \xrightarrow{\pi} & k[Z] \end{array}$$

© David Rufino CC BY-NC-ND 4.0

from which the properties of $i_Y^\sharp$ follows immediately.

Recall that the Zariski topology on $Z$ coincides with the subspace topology (4.3.21), and we have shown the notion of regular function coincides with the closed subspace definition (4.3.23). Therefore the topology on $Z$ and structure sheaves $\mathcal{O}_Z$ coincide. ∎

**Proposition 6.1.45**
*Let $\phi : X \to Y$ be a regular morphism of affine varieties. Then $\phi$ is a closed immersion (4.3.23) if and only if $\phi^\sharp$ is surjective.*

*Proof.* Suppose that $\phi^\sharp$ is surjective and let $Z \coloneqq V\big(\ker\big(\phi^\sharp\big)\big)$ be a closed subset of $Y$. Then $i : Z \hookrightarrow Y$ is regular and we have a commutative diagram

$$\begin{array}{ccc} \mathcal{O}_Y(Y) & \xrightarrow{\phi^\sharp} & \mathcal{O}_X(X) \\ \downarrow{\scriptstyle i^\sharp} & \nearrow{\scriptstyle \tilde{\phi}^\sharp} & \\ \mathcal{O}_Z(Z) & & \end{array}$$

By (6.1.44) $\ker\big(i^\sharp\big) = \ker\big(\phi^\sharp\big)$, so $\tilde{\phi}^\sharp$ exists and is an isomorphism by (3.3.56). By (6.1.42) there exists a corresponding regular morphism $\tilde{\phi} : X \to Z$ which is an isomorphism which satisfies $i \circ \tilde{\phi} = \phi$. By the criteria in (4.3.23) this means $\phi$ is a closed immersion.

Conversely if $\phi$ is a closed immersion then by definition $\phi = i \circ \tilde{\phi}$ for $i : Z \hookrightarrow Y$ the inclusion of a closed subset where $\tilde{\phi}$ is a regular isomorphism. By (6.1.42) $\tilde{\phi}^\sharp \circ i^\sharp = \phi^\sharp$ and $\tilde{\phi}^\sharp$ is an isomorphism, which shows that $\phi^\sharp$ is surjective. ∎

## 6.1.4 Dominant Morphisms

**Definition 6.1.46**
*Let $\phi : X \to Y$ be a regular morphism of affine varieties. Then we say it is*

- ***finite*** *if $\phi_Y^\sharp : \mathcal{O}_Y(Y) \to \mathcal{O}_X(X)$ is module-finite (equivalently integral)*
- ***dominant*** *if $\phi\big(X\big(\overline{k}\big)\big)$ is dense in $Y\big(\overline{k}\big)$*

**Proposition 6.1.47 (Criteria for dominant morphisms)**
*A regular morphism $\phi : X \to Y$ is dominant if and only if $\phi_Y^\sharp : \mathcal{O}_Y(Y) \to \mathcal{O}_X(X)$ is injective. In particular when both $X$ and $Y$ are integral then this induces a field extension*

$$\phi_\star : k(Y) \hookrightarrow k(X)$$

*Proof.* Recall $\phi(X)$ is dense precisely when the closure $\overline{\phi(X)} = Y$. Then

$$\begin{aligned} \ker\big(\phi_Y^\sharp\big) = I_k(\phi(X)) &\implies V_{\overline{k}}\big(\ker\big(\phi^\sharp\big)\big) = V_{\overline{k}}(I_k(\phi(X))) = \overline{\phi(X)} \\ &\implies \sqrt{\ker(\phi^\sharp)} = I_k\big(\overline{\phi(X)}\big) \end{aligned}$$

by (6.1.2) and (6.1.13). If $\ker\big(\phi_Y^\sharp\big) = 0$ then clearly $\overline{\phi(X)} = Y$. Conversely if this holds then

$$\sqrt{\ker\big(\phi_Y^\sharp\big)} = I_k\big(Y\big(\overline{k}\big)\big) = I_k(V_{\overline{k}}(0)) = \sqrt{(0)} = (0)$$

as $\mathcal{O}_Y(Y)$ is reduced, whence $\ker\big(\phi_Y^\sharp\big) = 0$. ∎

## 6.1.5 Dimension

**Definition 6.1.48 (Dimension)**
*Let $X$ be an affine variety. Then we define the dimension to be*

$$\dim X \coloneqq \dim X\big(\overline{k}\big) = \dim X_0\big(\overline{k}\big)$$

*where this is the Krull Dimension (4.1.81) of the $\overline{k}$-rational points with the $k$-Zariski topology. This is the supremum of dimension over all irreducible components by (4.1.83).*

*We say $X$ is of* ***pure dimension*** *$n$ if all irreducible components have the same dimension $n$. Note an integral variety is always of pure dimension.*

*We say $X = V(\mathfrak{a})$ is an* ***affine curve*** *if it is of pure dimension 1.*

© David Rufino CC BY-NC-ND 4.0

**Proposition 6.1.49 (Dimension of subspace)**
*Let $X = V(\mathfrak{a})$ be an affine variety and $Y = V(\mathfrak{b})$ an affine subvariety Then*

*a)* $\dim Y = \dim(\mathfrak{b}) = \dim k[X]/\mathfrak{b} = \dim k[Y]$
*b)* $\operatorname{codim}(Y, X) = \operatorname{ht}(\mathfrak{b})$

*Further when $Y = V(\mathfrak{p})$ is integral then* $\operatorname{codim}(Y, X) = \dim k[X]_{\mathfrak{p}}$.

*Proof.* a) follows from (6.1.23).

b) follows similarly by observing that the definition of ideal height (3.22.1) is dual to the topological definition of codimension (4.1.81). The last statement follows from (3.22.4). ∎

**Corollary 6.1.50**
*Let $X = V(\mathfrak{a})$ be an affine variety then* $\dim X = \dim k[X]$. *In particular* $\dim \mathbb{A}_k^n = n$.

*Proof.* The first statement is a consequence of (6.1.49) in the case $\mathfrak{b} = (0)$. The dimension for $\mathbb{A}_k^n$ then follows (3.35.28). ∎

As we showed when studying the Krull dimension of affine algebras, the lattice of irreducible subsets is particularly well behaved.

**Proposition 6.1.51 (Biequidimensional Algebraic Sets)**
*Let $X = V(\mathfrak{a})$ be an affine variety. Then $X\left(\overline{k}\right)$ is quasi-biequidimensional.*

*Further $X$ is equidimensional (e.g. integral) iff it is biequidimensional. In this case for every closed subset $Y = V(\mathfrak{b})$ the codimension formula is satisfied*

$$\dim X = \dim Y + \operatorname{codim}(Y, X)$$

*or in algebraic terms*

$$\dim k[X] = \dim \mathfrak{b} + \operatorname{ht}(\mathfrak{b})$$

*Proof.* We've observed that the lattice of radical (resp. prime) ideals of $k[X]$ is isomorphic to the lattice of closed (resp. irreducible) subsets of $X\left(\overline{k}\right)$. Therefore by (3.35.35) $X\left(\overline{k}\right)$ is quasi-biequidimensional.

In the equidimensional case, the codimension formula follows from (4.1.87) and the algebraic version from (6.1.49).∎

In the integral case we recover the "classical" field-theoretic version of dimension

**Proposition 6.1.52 (Dimension of Function Field)**
*Let $X = V(\mathfrak{p})$ be an integral affine variety. Then*

$$\dim X = \dim k[X] = \operatorname{trdeg}(\operatorname{Frac}(k[X])/k)$$

*Proof.* We have already shown that $\dim X = \dim k[X]$. The second equality follows from Noether Normalisation (3.35.28). ∎

**Proposition 6.1.53**
*Let $X$ be an affine variety and $Z \subset X$ a closed subset. Then the following are equivalent*

*a)* $\dim Z = 0$
*b)* $Z$ *is finite*

*More precisely*

$$Z = \overline{\{x_1\}} \cup \ldots \cup \overline{\{x_n\}}$$

*for a finite number of points $x_1, \ldots, x_n \in Z$.*

*Proof.* This follows from (4.1.91) and (6.1.18). ∎

**Proposition 6.1.54 (Hypersurfaces)**
*Let $X = V(\mathfrak{a})$ be an affine variety and $f \in k[X]$. Define the* ***hypersurface*** $Y \coloneqq V(f)$ *then* $\dim Y = n - 1$ *(and $Y$ is equidimensional).*

*Conversely if $k[X]$ is a UFD and $Y \subset X$ is an irreducible closed subset of dimension 1, then $Y$ is a hypersurface.*

*Proof.* **TODO.** ∎

© David Rufino CC BY-NC-ND 4.0

### 6.1.6 Local Rings

**Definition 6.1.55 (Local Ring)**
*Let $X$ be an affine variety and $W \subset X\left(\overline{k}\right)$ be an irreducible subset. Then we define the **local ring** at $W$ to be*

$$\mathcal{O}_{X,W} \coloneqq \varinjlim_{U \cap W \neq \varnothing} \mathcal{O}_X(U)$$

*It is a local ring with unique maximal ideal*

$$\mathfrak{m}_{X,W} = \{(U,\sigma) \mid \sigma(x) = 0 \quad \forall x \in W\}.$$

*In the case $W = \{(x)\}$ then we write it as $\left(\mathcal{O}_{X,x}, \mathfrak{m}_{X,x}\right)$.*

*Proof.* First of all the family of open sets such that $U \cap W \neq \varnothing$ is directed by reverse inclusion precisely because $W$ is irreducible, see (4.1.60).

Any section $\sigma \in \mathcal{O}_X(U)$ is continuous with respect to the cofinite topology on $k$. Therefore $D(\sigma) \coloneqq \sigma^{-1}(k \setminus \{0\}) \subset U$ is an open set and $\sigma \notin \mathfrak{m}_{X,W} \implies D(\sigma) \cap W \neq \varnothing$. Then evidently $[(D(\sigma), \sigma^{-1})]$ is a multiplicative inverse for $(U,\sigma)$ and by (...) $\mathcal{O}_{X,W}$ is a local ring with unique maximal ideal $\mathfrak{m}_{X,W}$. ∎

**Corollary 6.1.56 (Field of Rational Functions)**
*Let $X$ be an integral affine variety. Then $\mathcal{O}_{X,X} =: k(X)$ is a field, which we denote as **field of rational functions**.*

*Proof.* Evidently $\mathfrak{m}_{X,X} = (0)$. Explicitly $(V,\sigma)$ has inverse $(D(\sigma), \sigma^{-1})$. ∎

**Proposition 6.1.57 (Local Ring is Localization of Coordinate Ring)**
*Let $X = V(\mathfrak{a})$ be an affine variety and $W \subset X\left(\overline{k}\right)$ an irreducible subset with prime ideal $\mathfrak{M}_{X,W} \coloneqq I(W)$. For all such $W$ we have a canonical local isomorphism*

$$\begin{array}{ccc} \mathcal{O}_X(X) & \longrightarrow & \mathcal{O}_X(X)_{\mathfrak{M}_{X,W}} \\ & \searrow^{\rho_W} & \downarrow \sim \\ & & \mathcal{O}_{X,W} \end{array}$$

$$\frac{f}{g} \mapsto \left[\left(D(g), \frac{f}{g}\right)\right]$$

*The inverse image of $\mathfrak{m}_{X,W}$ is $\mathfrak{M}_{X,W}$ and $\mathfrak{M}_{X,W}\mathcal{O}_X(X)_{\mathfrak{M}_{X,W}}$ respectively. This induces isomorphisms*

$$\mathcal{O}_X(X)/\mathfrak{M}_{X,W} \xrightarrow{\sim} k\left(\mathfrak{M}_{X,W}\mathcal{O}_X(X)_{\mathfrak{M}_{X,W}}\right) \xrightarrow{\sim} k\left(\mathfrak{m}_{X,W}\right)$$

*Further we have*

$$\dim \mathcal{O}_{X,W} = \operatorname{ht}\left(\mathfrak{M}_{X,W}\right) = \operatorname{codim}(W,X) = \dim X - \dim W$$

*In particular*

$$\dim \mathcal{O}_{X,x} = \dim X$$

*Proof.* This is a formal consequence of generic facts regarding localization and direct limits

$$\mathcal{O}_X(X)_{\mathfrak{p}} \overset{(3.6.44)}{\cong} \varinjlim_{f \notin \mathfrak{p}} \mathcal{O}_X(X)_f \cong \varinjlim_{D(f) \cap W \neq \varnothing} \mathcal{O}_X(D(f)) \overset{(2.6.127)}{\cong} \varinjlim_{U \cap W \neq \varnothing} \mathcal{O}_X(U)$$

We may demonstrate this more directly. For it is surjective because the principal open sets form a basis and by (6.1.26), observing that $f \notin \mathfrak{p} \iff D(f) \cap V(\mathfrak{p}) \neq \varnothing$. Suppose $\frac{g}{f}$ and $\frac{g'}{f'}$ have the same image then there is some $h$ such that $D(h) \subseteq D(ff') = D(f) \cap D(f')$ and $h \notin \mathfrak{p}$ such that $\frac{g}{f} = \frac{g'}{f'}$ are equal in $k[X]_h$ and a-fortiori in $k[X]_{\mathfrak{p}}$. Therefore the map is injective as required.

The first two dimension identities follow from (6.1.49) the third from (6.1.51). The final identity follows from (6.1.53).∎

**Corollary 6.1.58**
*Let $X$ be an irreducible variety then there is a canonical isomorphism with the field of rational functions*

© David Rufino CC BY-NC-ND 4.0




**Proposition 6.1.59**
*Let $\phi : X \to Y$ be a regular morphism of affine varieties. Suppose $W \subset X$ is irreducible and $Z \subset \overline{\phi(W)}$ for some irreducible $Z \subset Y$. Then the stalk map $\phi_W : \mathcal{O}_{Y,Z} \to \mathcal{O}_{X,Z}$ (see (4.3.5)) makes the following diagram commute*




*where the bottom arrow is the localisation map induced by $\phi^\sharp$ (3.6.22). In fact it is the unique map making either square commute.*

**Proposition 6.1.60**
*Let $X$ be an integral affine variety, $W = V(\mathfrak{M}_{X,W})$ an irreducible subset. Then there are canonical injections*

$$\mathcal{O}_X(X) \hookrightarrow \mathcal{O}_{X,W} \hookrightarrow k(X)$$

*and the restriction maps are injective. Further there is a commutative diagram*




*which shows $k(X)$ is a quotient field for $\mathcal{O}_{X,W}$ and $\mathcal{O}_X(X)$.*

*Proof.* We first show the restriction maps are injective. Suppose $\sigma|_V = 0$. Note that $V$ is dense (4.1.59), $\sigma$ is continuous (6.1.34) with respect to the $k$-Zariski topology on $\overline{k}$ so that $\sigma^{-1}(\{0\})$ is closed. By definition this contains $V$ and therefore also $\mathrm{cl}_U(V) = \mathrm{cl}_X(V) \cap U = U$. We conclude $\sigma = 0$ and the restriction maps are injective. The same argument shows that the remaining maps are injective.

The vertical maps are isomorphisms by (6.1.57), and we may verify directly the diagram commutes. ∎

Therefore if it's convenient to do so we may identify $\mathcal{O}_X(X)$ and $\mathcal{O}_{X,W}$ as subrings of $k(X)$.

**Proposition 6.1.61**
*Let $X$ be an affine variety and $W \subset X$ an irreducible closed subset. Then the minimal primes of $\mathcal{O}_{X,W}$ are in bijection with the irreducible components of $X$ containing $W$.*

*$\mathcal{O}_{X,W}$ is an integral domain if and only if $W$ lies in a unique irreducible component.*

*Proof.* By (6.1.57) and (3.6.42) the minimal primes of $\mathcal{O}_{X,W}$ correspond to minimal primes of $k[X]$ contained in $\mathfrak{M}_{X,W}$. These correspond to irreducible components of $X$ containing $\overline{W}$ (and hence $W$) by (6.1.19).

As $\mathcal{O}_{X,W}$ is reduced, it is an integral domain iff it has a unique minimal prime ideal (3.3.67). ∎

**Proposition 6.1.62**
*Let $X$ be an affine variety, $R$ an algebraic $k$-algebra (3.35.25) and $\phi : \mathcal{O}_{X,x} \to R$ a $k$-algebra homomorphism. Then $\phi^{-1}(\mathfrak{m}_R) = \mathfrak{m}_{X,x}$ and therefore factors through $k(\mathfrak{m}_{X,x})$*

*This in particular this includes the case $R$ is an algebraic field extension.*

*Proof.* See (3.35.27). ∎

© David Rufino CC BY-NC-ND 4.0

## 6.1.7 Rational Points

**Proposition 6.1.63 ($K$-rational points)**
*Let $X = V(\mathfrak{a})$ be an affine variety and $K$ an algebraic field extension. Then there are natural bijections*

$$\begin{aligned} X(K) &\longleftrightarrow \mathrm{AlgHom}_k(\mathcal{O}_X(X), K) \\ &\longleftrightarrow \{(\mathfrak{M}, \phi) \mid \mathfrak{M} \in \mathrm{Specm}(\mathcal{O}_X(X)) \quad \phi : \mathcal{O}_X(X)/\mathfrak{M} \hookrightarrow K\} \\ &\longleftrightarrow \left\{(x, \phi) \;\middle|\; x \in X_0\left(\overline{k}\right) \quad \phi : k\left(\mathfrak{m}_{X,x}\right) \hookrightarrow K\right\} \\ &\longleftrightarrow \left\{(x, \phi) \;\middle|\; x \in X_0\left(\overline{k}\right) \quad \phi : \mathcal{O}_{X,x} \to K\right\} \end{aligned}$$

*Proof.* The first correspondence is simply (6.1.7) and (6.1.26).

For the second correspondence consider a $k$-algebra homomorphism $\phi : \mathcal{O}_X(X) \to K$. By (3.35.27) we have $\ker(\phi) =: \mathfrak{M}$ is maximal, which induces a homomorphism $\mathcal{O}_X(X)/\mathfrak{M} \hookrightarrow K$. Evidently this correspondence is injective and surjective.

By (6.1.11) there is a bijection $\mathrm{Specm}(\mathcal{O}_X(X)) \longleftrightarrow X_0\left(\overline{k}\right)$. Further there is an isomorphism $\mathcal{O}_X(X)/\mathfrak{M}_{X,x} \cong k\left(\mathfrak{m}_{X,x}\right)$ (6.1.57) from which the third correspondence is well-defined and bijective.

The final correspondence follows from (6.1.62) because all such homomorphisms factor through $k\left(\mathfrak{m}_{X,x}\right)$. ∎

## 6.1.8 Generic Points

When considering irreducible subsets $W \subset X$ there is another way of looking at these in terms of "generic points".

**Definition 6.1.64 (Generic Point)**
*Let $X = V(\mathfrak{a})$ be an affine variety and $(\xi) \in X(\Omega)$ for some extension field $\Omega/k$. Then we may define the irreducible subset of $X\left(\overline{k}\right)$*

$$W_\xi := V_{\overline{k}}(\mathfrak{p}_\xi) = \left\{(x) \in X\left(\overline{k}\right) \;\middle|\; \forall f \in k[X](f(\xi) = 0 \implies f(x) = 0)\right\}$$

*There are canonical isomorphisms*

$$k\left(\mathfrak{m}_{X,W}\right) \xrightarrow{\sim} k[X]_{\mathfrak{p}}/\mathfrak{p}k[X]_{\mathfrak{p}} \xrightarrow{\sim} \mathrm{Frac}(k[X]/\mathfrak{p}) \xrightarrow{\sim} k(\xi)$$

*We say that $(\xi)$ is a **generic point** corresponding to the irreducible closed subset $W_\xi$. Moreover every irreducible subset is of this form for we may simply consider $\Omega := \mathrm{Frac}(k[X]/\mathfrak{p})$ and $\xi = \left(\overline{X}_1, ..., \overline{X}_n\right)$.*

*If $(x) \in W_\xi$ then we say that $(x)$ is a **specialization** of $(\xi)$ and this induces the specialization homomorphism*

$$\begin{array}{ccc} k[X]/\mathfrak{p}_\xi & \longrightarrow & k[X]/\mathfrak{m}_x \\ \wr\wr & & \wr\wr \\ k[\xi] & \dashrightarrow & k[x] \end{array}$$

*If $W_\xi = X$ then we say simply $(\xi)$ is a generic point.*

## 6.1.9 Tangent Space and Non-Singular Points

We propose two definitions for the tangent space and show that under mild technical conditions they are naturally isomorphic. The latter definition may be identified geometrically.

**Definition 6.1.65**
*Let $X$ be an affine variety and let $W \subset X$ be an irreducible subset. We define the **cotangent space** at $W$ to be the $k\left(\mathfrak{m}_{X,W}\right)$-vector space*

$$T^\star_W X := \mathfrak{m}_{X,W}/\mathfrak{m}^2_{X,W}$$

*and the **tangent space** to be the $k\left(\mathfrak{m}_{X,W}\right)$-vector space*

$$T_W X := \mathrm{Der}_k\left(\mathcal{O}_{X,W}, k\left(\mathfrak{m}_{X,W}\right)\right)$$

For points this has a concrete interpretation

**Proposition 6.1.66 (Concrete interpretation of Tangent Space)**
*Let $X = V(\mathfrak{a}) \subset \mathbb{A}^n_k$ be an affine variety and $x \in X\left(\overline{k}\right)$. Then there are natural $k(x)$-module isomorphisms*

© David Rufino CC BY-NC-ND 4.0

$$\begin{array}{lll} T_xX \xrightarrow{\sim} \mathrm{Der}_k(k[X], k(x)) \xrightarrow{\sim} \left\{ v \in k(x)^n \;\middle|\; \sum_{i=1}^n v_i \frac{\partial F}{\partial X_i}(x) = 0 \quad \forall F \in \mathfrak{a} \right\} \\ D \qquad\quad \to \qquad\qquad\qquad \left( D\left(\overline{X}_1\right), ..., D\left(\overline{X}_n\right) \right) \end{array}$$

*where we have used the identification* $k\left(\mathfrak{m}_{X,x}\right) \cong k(x)$ *and the* $k$*-algebra homomorphism* $k[X] \to k\left(\mathfrak{m}_{X,x}\right)$*. Suppose* $\mathfrak{a} = \langle F_1, ..., F_m \rangle$ *then the right hand side is the kernel of the following* $k(x)$*-module homomorphism*

$$\begin{pmatrix} \frac{\partial F_1}{\partial X_1}(x) & \cdots & \frac{\partial F_1}{\partial X_n}(x) \\ \vdots & \ddots & \vdots \\ \frac{\partial F_m}{\partial X_1}(x) & \cdots & \frac{\partial F_m}{\partial X_n}(x) \end{pmatrix} : k(x)^n \to k(x)^m$$

*In particular*

$$\dim_{k(\mathfrak{m}_{X,x})} T_xX = n - \mathrm{rk}\left( \frac{\partial F_i}{\partial X_j}(x) \right)$$

*Proof.* Recall by (6.1.57) and (6.1.64) that $\mathcal{O}_{X,x} \cong k[X]_{\mathfrak{m}_{X,x}}$ and $k\left(\mathfrak{m}_{X,x}\right) \cong k(x)$ so we have isomorphisms

$$\mathrm{Der}_k\left(\mathcal{O}_{X,x}, k\left(\mathfrak{m}_{X,x}\right)\right) \cong \mathrm{Der}_k\left(k[X]_{\mathfrak{m}_{X,x}}, k(x)\right) \overset{(3.26.10)}{\cong} \mathrm{Der}_k(k[X], k(x))$$

and the remaining isomorphism is from (3.35.41) ∎

**Proposition 6.1.67 (Dimension Formula of Tangent Space)**
*Let* $X = V(\mathfrak{a}) \subset \mathbb{A}_k^n$ *be an* integral *affine variety,* $W \subset X$ *an irreducible subset. Suppose that* $\mathfrak{a} = \langle F_1, ..., F_m \rangle$ *and let* $\xi := \left(\overline{X}_1, ..., \overline{X}_n\right) \in k\left(\mathfrak{m}_{X,W}\right)^n$ *be the image of the coordinate functions. Then we have the dimension formula*

$$\dim T_WX = \dim \mathrm{Der}\left(k[X], k\left(\mathfrak{m}_{X,W}\right)\right) = n - \mathrm{rk}\left( \frac{\partial F_i}{\partial X_j}(\xi) \right)$$

*Suppose that* $Z \subset W$ *is an irreducible subset then we have the inequality*

$$\dim T_ZX \geq \dim T_WX$$

*and*

$$\dim T_WX \geq \dim T_XX = \mathrm{Der}(k(X))$$

*which equals* $\dim X$ *when* $k(X)$ *is separably generated (e.g.* $k$ *is perfect).*

*Proof.* We have isomorphisms

$$\begin{aligned} \dim T_WX := \quad & \mathrm{Der}_k\left(\mathcal{O}_{X,W}, k\left(\mathfrak{m}_{X,W}\right)\right) \\ \overset{(6.1.57)}{\to} \quad & \mathrm{Der}_k\left(\mathcal{O}_X(X)_{\mathfrak{m}_{X,W}}, k\left(\mathfrak{m}_{X,W}\right)\right) \\ \overset{(3.26.10)}{\to} \quad & \mathrm{Der}_k\left(\mathcal{O}_X(X), k\left(\mathfrak{m}_{X,W}\right)\right) \\ \overset{(6.1.26)}{\to} \quad & \mathrm{Der}_k\left(k[X], k\left(\mathfrak{m}_{X,W}\right)\right) \end{aligned}$$

and the rank formula follows from (3.35.41) and (3.3.135).

Let $\zeta$ be the corresponding generic point of $Z$, then because $\mathfrak{M}_{X,W} \subset \mathfrak{M}_{X,Z}$ we may construct a $k$-algebra homomorphism $k[\xi] \to k[\zeta]$. So by the determinant criteria of rank (3.5.40) we deduce that

$$\mathrm{rk}\left( \frac{\partial F_i}{\partial X_j}(\zeta) \right) \leq \mathrm{rk}\left( \frac{\partial F_i}{\partial X_j}(\xi) \right)$$

from which the inequality follows.

Observe that $k\left(\mathfrak{m}_{X,X}\right) =: k(X)$ and $T_XX \xrightarrow{\sim} \mathrm{Der}_k(k[X], k(X)) \xrightarrow{\sim} \mathrm{Der}_k(k(X))$. When $k(X)$ is separably generated this equals $\dim X$ by (3.35.44). ∎

**Proposition 6.1.68 (Duality between Tangent and Cotangent Space)**
*Let* $X$ *be an affine variety and* $W \subset X$ *be an irreducible subset. There is a canonical* $k\left(\mathfrak{m}_{X,W}\right)$*-module homomorphism*

© David Rufino CC BY-NC-ND 4.0

$$T_W X \to (T^\star_W X)^\vee$$

*Under the condition that* $k(\mathfrak{m}_{X,W})/k$ *is separably generated (e.g.* $k$ *is perfect) then this map is an isomorphism. In particular* $\dim T_W X = \dim T^\star_W X$ *by (3.3.109).*

*Proof.* This follows directly from (3.35.50) by considering the local ring $A \coloneqq \mathcal{O}_{X,W}$ with maximal ideal $\mathfrak{m}_{X,W}$. ∎

**Lemma 6.1.69**
*Let* $A$ *be a reduced ring with minimal prime ideal* $\mathfrak{p}$ *and prime ideal* $\mathfrak{q} \supset \mathfrak{p}$ *containing no other minimal prime ideals. Suppose that*

$$\phi : A \to B$$

*is a surjective ring homomorphism with* $\ker(\phi) = \mathfrak{p}$*. Then* $\phi$ *induces an isomorphism of rings*

$$\tilde{\phi} : A_\mathfrak{q} \to B_{\phi(\mathfrak{q})}$$

*which is the unique morphism commuting with* $\phi$*.*

*Proof.* The map $\tilde{\phi}$ exists and has kernel $\mathfrak{p}A_\mathfrak{q}$ by (3.6.5). We need simply to show that $\mathfrak{p}A_\mathfrak{q} = (0)$. By (3.6.42) it is the unique minimal prime ideal of $A_\mathfrak{q}$. As $A_\mathfrak{q}$ is reduced then we are done by (3.3.46). ∎

**Proposition 6.1.70**
*Let* $X$ *be an affine variety,* $W$ *an irreducible subset contained in precisely one irreducible component* $X_\alpha$*. Then there is a canonical isomorphism*

$$\begin{aligned} \mathcal{O}_{X,W} &\xrightarrow{\sim} \mathcal{O}_{X_\alpha,W} \\ (U,\sigma) &\to \left(U \cap X_\alpha, \sigma|_{X_\alpha \cap U}\right) \end{aligned}$$

*and in particular an isomorphism of tangent spaces*

$$T_W X \xrightarrow{\sim} T_W X_\alpha$$

*Proof.* The regular map $i : X_\alpha \hookrightarrow X$ corresponds to a ring homomorphism $i^\sharp$ with kernel $\mathfrak{M}_{X,X_\alpha}$ by (6.1.44). By (...) we have a commutative diagram

$$\begin{array}{ccc} \mathcal{O}_X(X) & \xrightarrow{i^\sharp} & \mathcal{O}_{X_\alpha}(X_\alpha) \\ \downarrow & & \downarrow \\ \mathcal{O}_{X,W} & \xrightarrow{i_W} & \mathcal{O}_{X_\alpha,W} \\ \downarrow \sim & & \downarrow \sim \\ \mathcal{O}_X(X)_{\mathfrak{M}_{X,W}} & \longrightarrow & \mathcal{O}_{X_\alpha}(X_\alpha)_{\mathfrak{M}_{X_\alpha,W}} \end{array}$$

and by (6.1.69) we deduce that the stalk map is an isomorphism. ∎

**Definition 6.1.71 (Non-Singular Points)**
*Let* $X$ *be an affine variety and* $W \subset X$ *be an irreducible subset. We say that* $X$ *is* ***non-singular*** *at* $W$ *if*

*a)* $W$ *is contained in only one irreducible component* $X_\alpha$*, and*
*b)* $\dim T_W X = \dim \mathrm{Der}_k(k(X_\alpha))$

*Note that by (6.1.61) a)* $\iff \mathcal{O}_{X,W}$ *is an integral domain.*

*Recall by (6.1.67) that* $\dim T_W X = \dim T_W X_\alpha \geq \dim \mathrm{Der}_k(k(X_\alpha))$ *and in the case of* $k$ *perfect this equals* $\dim X_\alpha$*.*

We may now show the following important result

**Proposition 6.1.72 (Non-Singular Point)**
*Let* $X = V(\mathfrak{a}) \subset \mathbb{A}^n_k$ *be an affine variety then the non-singular points form an open dense subset of* $X$*.*

*Proof.* We may reduce to the case of $X$ irreducible. Let $\Omega \coloneqq k(X)$ and $\xi \coloneqq \left(\overline{X}_1, ..., \overline{X}_n\right) \in X(\Omega)$ and suppose $\mathfrak{p} = \langle F_1, ..., F_m \rangle \lhd k[X_1, ..., X_n]$ is the ideal defining $X$. Then

© David Rufino CC BY-NC-ND 4.0

$$\mathrm{Der}(\Omega) \overset{(3.26.10)}{=} \mathrm{Der}(k[X], \Omega) \overset{(3.35.41)}{\cong} \ker\left(\frac{\partial F_i}{\partial X_j}(\xi)\right)$$

For $x \in X\left(\overline{k}\right)$ there is a $k$-algebra homomorphism $k[\xi] \to k[x]$ and so by the determinant criteria of rank (3.5.40) we see

$$\mathrm{rk}\left(\frac{\partial F_i}{\partial X_j}(x)\right) \leq \mathrm{rk}\left(\frac{\partial F_i}{\partial X_j}(\xi)\right)$$

Suppose the rank of the right hand side is $r$ and consider the $r$-minors $g_1, ..., g_m \in k[X]$. Then by the same result at least one is non-zero and the set of non-singular points is given by

$$\bigcup_p D(g_p)$$

which is non-empty by (6.1.27). ∎

**Proposition 6.1.73 (Non-Singular Variety)**
*Let $X$ be an integral affine variety. Then the following are equivalent*

*a) $X$ is non-singular at all points $x \in X\left(\overline{k}\right)$*
*b) $X$ is non-singular at all irreducible subsets $W \subset X\left(\overline{k}\right)$*

*In this case we say that $X$ is a **non-singular affine variety**.*

*Proof.* Evidently $b) \implies a)$ because $T_x X = T_{\overline{\{x\}}} X$. Conversely given any $x \in W$ we have by (6.1.67)

$$T_x X \geq T_W X \geq \dim \mathrm{Der}(k(X))$$

whence they are both equal. ∎

**Definition 6.1.74 (Regular Point)**
*Let $X$ be an affine variety and $W \subset X$ an irreducible subset. Then we say that $W$ is **regular** if*

*a) $W$ is contained in precisely one irreducible component $X_\alpha$, and*
*b) $\mathcal{O}_{X,W}$ is a regular local ring (i.e. $\dim T^\star_W X = \dim \mathcal{O}_{X,W}$)*

*We say $X$ is **regular** if it is regular at all irreducible subsets.*

Fortunately these concepts are typically equivalent.

**Proposition 6.1.75 (Regular = Non-Singular in Perfect Case)**
*Let $X = V(\mathfrak{a}) \subset \mathbb{A}^n_k$ an affine variety with $k$ perfect and $W \subset X$ an irreducible subset. Then*

$$\dim T^\star_W X = \dim T_W X$$

*and the following are equivalent*

*a) $X$ is regular at $W$*
*b) $X$ is non-singular at $W$*
*c) $\dim T_W X = \dim \mathcal{O}_{X,W} = \mathrm{codim}(W, X)$*

*When $\mathrm{codim}(W, X) = 1$ this is equivalent to $\mathcal{O}_{X,W}$ being a discrete valuation ring.*

*Proof.* Recall by (6.1.57) that $\dim \mathcal{O}_{X,x} = \dim X$ and by definition $T^\star_x X = \mathfrak{m}_{X,x}/\mathfrak{m}^2_{X,x}$. Further by definition $(x)$ is regular iff $\dim T^\star_x X = \dim \mathcal{O}_{X,x}$. Finally by (6.1.68) $\dim T^\star_x X = \dim T_x X$ so we see $a) \iff c)$.

Similarly $b) \iff c)$ is essentially by definition. The final statement follows from (3.31.2). ∎

**Example 6.1.76 (Simple Points of a Plane Curve)**
*The canonical example is a plane curve $k[X] = k[X,Y]/(F(X,Y))$ which has dimension 1 (...). Given $(x) \in X(K)$ we see that*

$$T_x X = \left\{ (\alpha, \beta) \in k(x)^2 \;\middle|\; 0 = \alpha \frac{\partial F}{\partial X}(x) + \beta \frac{\partial F}{\partial Y}(x) \right\}$$

*Clearly this has dimension 1 (in which case $(x)$ is a **simple point**) unless both the partial derivatives vanish at $(x)$ in which case it has dimension 2.*

© David Rufino CC BY-NC-ND 4.0

*Clearly $F(X,Y) = Y^2 - X^3$ has a non-simple point at $(0,0)$ but $F(X,Y) = Y - X^2$ has simple points everywhere.*

### 6.1.10 Zeta Function over Finite Fields

For this section let $k = \mathbb{F}_p$ with algebraic closure $\overline{k}$. Then for every $n \geq 1$ by (...) $\overline{k}$ has a unique subfield $k_n$ of degree $n$, and order $p^n$. Furthermore we have the following properties

a) $k_m \subseteq k_n \iff [k_m : k] \mid [k_n : k] \iff m \mid n$
b) $[k_n : k_m] = \frac{n}{m}$
c) $k_n/k_m$ is Galois with the group of automorphisms generated by $\phi^m$ for all integers $m \mid n$, and the Galois group has order $\frac{n}{m}$.
d) every finite subextension is of this form

**Lemma 6.1.77**
*Let $x_1, ..., x_N \in \overline{k}$. Then*

$$k(x_1, ..., x_N) = k_n$$

*where*

$$n \coloneqq \operatorname{lcm}_i [k(x_i) : k]$$

*Proof.* By definition $[k(x_i) : k] \mid n$ whence $k(x_i) \subseteq k_n$ by the previous observations, and therefore $k(x_1, ..., x_N) \subseteq k_n$.

Similarly as $k(x_i) \subset k(x_1, ..., x_N)$ we have $[k(x_i) : k] \mid [k(x_1, ..., x_N) : k]$. Then by definition $n \mid [k(x_1, ..., x_N) : k]$ which shows $k_n \subset k(x_1, ..., x_N)$. ∎

The following result shows that the number of rational points over a finite field may be characterized by algebraic properties of the coordinate ring $k[X]$. This is essentially because maximal ideals correspond to Galois orbits of rational points and we may count rational points by summing over Galois orbits.

**Proposition 6.1.78 (Counting Rational Points)**
*Let $X = V(\mathfrak{a}) \subset \mathbb{A}_k^N$ be an affine variety. Then for all integers $n \geq 1$ we have the following relation*

$$\#X(k_n) = \sum_{d \mid n} d \times \#\{\mathfrak{m} \mid \dim_k k(\mathfrak{m}) = d\}$$

*Proof.* For $(x) \in X\left(\overline{k}\right)$, then by the previous discussion

$$(x) \in X(k_n) \iff k(x) \subseteq k_n \iff \dim_k k(x) \mid n$$

This shows that

$$\begin{aligned} \#X(k_n) &= \sum_{d \mid n} \#\{(x) \in X(k_n) \mid \dim_k k(x) = d\} \\ &= \sum_{d \mid n} \#\{(x) \in X(k_d) \mid \dim_k k(x) = d\} \end{aligned} \tag{*}$$

By (6.1.11) we have a bijection

$$X(k_d)/\operatorname{Gal}(k_d/k) \xrightarrow{\sim} \{\mathfrak{m} \mid \dim_k k(\mathfrak{m}) \mid d\}$$

As we have an isomorphism $k(x) \cong k(\mathfrak{m}_x)$ this restricts to a bijection

$$\{(x) \in X(k_d) \mid \dim_k k(x) = d\}/\operatorname{Gal}(k_d/k) \xrightarrow{\sim} \{\mathfrak{m} \mid \dim_k k(\mathfrak{m}) = d\}$$

We claim that the action of $\operatorname{Gal}(k_d/k) = \langle \phi \rangle$ is free. For suppose $(x) \in X(k_d)$ such that $\dim_k k(x) = d$ and $\phi^r(x_i) = x_i$ for $i = 1, ..., N$ and $0 < r \leq d$. Then by (3.19.108) we have $x_i \in k_r$ whence $\dim_k k(x) \leq r$. This shows we must have $r = d$ and in particular only the identity automorphism fixes $(x)$.

Therefore we see that

$$\#\{(x) \in X(k_d) \mid \dim_k k(x) = d\} = \#\operatorname{Gal}(k_d/k) \times \#\{\mathfrak{m} \mid \dim_k k(\mathfrak{m}) = d\}$$

From (3.19.111) we recall $\operatorname{Gal}(k_d/k)$ is cyclic of order $d$ so we may combine this with (*) to obtain the required result. ∎

**Proposition 6.1.79 (Zeta function of an affine variety over a finite field)**
*Formally as elements of the power series ring $\mathbb{Q}[[T]]$ we have*

© David Rufino CC BY-NC-ND 4.0

$$Z(X,T) \coloneqq \prod_{\mathfrak{m}\in \operatorname{Specm}(k[X])} \left(1-T^{\deg(\mathfrak{m})}\right)^{-1} = \exp\left(\sum_{m=1}^{\infty} \#\frac{X(k_m)}{m} T^m\right)$$

*Proof.* Define

$$b_d \coloneqq \#\{\mathfrak{m} \mid \dim_k k(\mathfrak{m}) = d\}$$

Let $Z(X,T)$ be the right hand side then

$$\begin{aligned}
\log(Z(X,T)) &= \sum_{m=1}^{\infty} \#X(k_m)\frac{T^m}{m} \\
&= \sum_{m=1}^{\infty}\sum_{d|m}(d\times b_d)\frac{T^m}{m} \\
&= \sum_{d=1}^{\infty} b_d \sum_{r=1}^{\infty}\frac{T^{rd}}{r} \\
&= -\sum_{d=1}^{\infty} b_d \log\left(1-T^d\right)
\end{aligned}$$

■

**Example 6.1.80**
*For $X(k) = k^n$ we have $\#X(k_m) = p^{mn}$. Then*

$$Z(X,T) = \exp\left(\sum_{m=1}^{\infty}\frac{p^{mn}T^m}{m}\right) = \exp(-\log(1-p^nT)) = \frac{1}{1-p^nT}$$

## 6.1.11 Base Change

**Definition 6.1.81 (Affine Base Change)**
*Let $X = V(\mathfrak{a}) \subset \mathbb{A}^n_k$ be an affine variety over $k$ and $K/k$ a field extension. Consider the canonical ring homomorphism*

$$i_{kK} : k[X_1,...,X_n] \hookrightarrow K[X_1,...,X_n]$$

*and let $\mathfrak{a}^e \coloneqq \mathfrak{a}K[X_1,...,X_n]$ denote the extension under this map.*

*Further define the **base change** of $X$ with respect to $K/k$ to be the affine variety*

$$X_K \coloneqq V\left(\sqrt{\mathfrak{a}^e}\right) \subset \mathbb{A}^n_K$$

**Proposition 6.1.82 (Coordinate Ring of Base Change)**
*Let $X = V(\mathfrak{a}) \subset \mathbb{A}^n_k$ be an affine variety over $k$ and $K/k$ a field extension. There is a canonical $K$-algebra isomorphism*

$$\begin{aligned}
(k[X]\otimes_k K)_{\text{red}} &\xrightarrow{\sim} K[X_K] \\
\overline{F}\otimes\lambda &\to \lambda\cdot\overline{i_{kK}(F)}
\end{aligned}$$

*Proof.* By (3.4.42) we have $K$-algebra isomorphisms

$$k[X]\otimes_k K \xrightarrow{\sim} (k[X_1,...,X_n]\otimes_k K)/\mathfrak{a}^e \xrightarrow{\sim} K[X_1,...,X_n]/\mathfrak{a}^e$$

By (3.35.4) and (3.35.2) this determines a unique $K$-algebra isomorphism

$$(k[X]\otimes_k K)_{\text{red}} \xrightarrow{\sim} (K[X_1,...,X_n]/\mathfrak{a}^e)_{\text{red}} \xrightarrow{\sim} K[X_1,...,X_n]/\sqrt{\mathfrak{a}^e} =: K[X_K]$$

■

It may be interesting to understand the cases when $\mathfrak{a}^e$ remains radical under the base change.

**Proposition 6.1.83**
*Let $X = V(\mathfrak{a})$ be an affine variety. Then the extension $\mathfrak{a}^e$ under $A \to A\otimes_k K$ remains radical under any of the following conditions*

*a) $k[X]$ is geometrically reduced*
*b) $K/k$ is geometrically reduced (e.g. separably generated)*

© David Rufino CC BY-NC-ND 4.0

*c) $k$ is perfect.*

*Proof.* By (3.4.42) we have an isomorphism

$$K[X_1, ..., X_n]/\mathfrak{a}^e \xrightarrow{\sim} k[X] \otimes_k K$$

so that $\mathfrak{a}^e$ is radical iff $k[X] \otimes_k K$ is reduced. Then we may reduce to results on geometrically reduced algebras. ∎

We are particularly interested when the variety remains integral (equivalently irreducible) under base change.

**Definition 6.1.84 (Geometrically Integral)**
*Let $X = V(\mathfrak{a})$ be an affine variety over $k$. Then we say $X$ is **geometrically integral** if $X_K$ is integral for all field extensions $K$. By (...) this is equivalent to $k[X]$ being geometrically irreducible.*

**Proposition 6.1.85 (Criteria to be Geometrically Integral)**
*Let $X = V(\mathfrak{a}) \subset \mathbb{A}^n_k$ be an integral affine variety. Then the following are equivalent*

*a) $X$ is geometrically integral*
*b) $X_{\overline{k}}$ is integral*
*c) $X_{k'}$ is integral for every finite separable extension $k'/k$*
*d) $k[X]$ is geometrically irreducible.*
*e) $k[X] \otimes \overline{k}$ is irreducible*
*f) $k[X] \otimes k'$ is irreducible for every finite separable extension $k'/k$.*
*g) $k(X)/k$ is relatively separably closed*

*Proof.* We have $a) \iff d)$, $b) \iff e)$ and $c) \iff f)$ are straightforward since a ring $A$ is irreducible iff $A_{\text{red}}$ is integral.

For $d) \iff e) \iff f) \iff g)$ we may use (3.36.41). ∎

**Proposition 6.1.86**
*Let $X = \mathbb{A}^n_k$. Then there is a canonical isomorphism $X_K \cong \mathbb{A}^n_K$. In particular $X$ is geometrically integral.*

**Proposition 6.1.87**
*Let $X \subset \mathbb{A}^n_k$ be an affine variety, $K/k$ a field extension and $R$ a $K$-algebra. Then there is a bijection*

$$\begin{array}{ccc} X\big(R_{(k)}\big) & \xrightarrow{\quad\sim\quad} & X_K(R) \\ \downarrow \sim & & \downarrow \sim \\ \mathrm{AlgHom}_k\big(k[X], R_{(k)}\big) & \xrightarrow{\sim} & \mathrm{AlgHom}_K(K[X_K], R) \end{array}$$

$$\phi \to \Big(\lambda \cdot \overline{F} \to \lambda \cdot \phi\big(\overline{i_{kK}(F)}\big)\Big)$$

*for $F \in k[X_1, ..., X_n]$, where the vertical arrows are given by (...).*

**Corollary 6.1.88**
*Let $X$ be an affine variety and $L/K/k$ a tower of extensions, then we may identify $X(L)$ with $X_K(L)$.*

**Proposition 6.1.89**
*Let $X = V(\mathfrak{a})$ an affine variety and $\overline{k}/K/k$ a tower of extensions. Then after identifying $X\big(\overline{k}\big)$ with $X_K\big(\overline{k}\big)$ we have the following properties*

*a) $U$ open in $X\big(\overline{k}\big) \implies U$ open in $X_K\big(\overline{k}\big)$,*
*b) There is a well-defined $K$-algebra homomorphism*

$$(\mathcal{O}_X(U) \otimes_k K)_{\text{red}} \to \mathcal{O}_{X_K}(U)$$

$$f \otimes \lambda \to \lambda \cdot f$$

*which is an isomorphism when $U = X$,*
*c) sets of the form $D\big(\sum_{i=1}^n \lambda_i f_i\big)$ are open in $X_K\big(\overline{k}\big)$ and form a base for the Zariski topology.*

*Proof.*

a) A closed subset of $X\big(\overline{k}\big)$ is of the form $Z = V(\mathfrak{b})$ for $\mathfrak{b} \subset \mathfrak{a}$. Then evidently $Z = V(\mathfrak{b}^e)$ and so it is closed as a subset of $X_K\big(\overline{k}\big)$
b) By definition we have an open cover $U = \bigcup_{i=1}^n U_i$ and polynomials $g_i, h_i \in k[X]$ such that $f|_{U_i} = \frac{g_i}{h_i}$. This shows that $\lambda \cdot f$ is also regular. The map is an isomorphism on global sections by (6.1.82).

© David Rufino CC BY-NC-ND 4.0

c) By (6.1.39) $f \coloneqq \sum_{i=1}^{n} \lambda_i f_i$ is continuous. As $\{0\}$ is a closed subset of $\overline{k}$ in the Zariski (co-finite) topology this shows that $D(f)$ is open. These form a base because $D(fg) = D(f) \cap D(g)$, and specifically for the Zariski topology by (6.1.15) and b).

■

## 6.1.12 Valuation Rings on the Function Field

**Proposition 6.1.90**
*Let $K/k$ be a field extension and $R$ a valuation ring of $K/k$. Then there exists a valuation ring $R' \subset R$ such that $k(\mathfrak{m}_{R'})/k$ is algebraic and $\mathfrak{m}_{R'} = \mathfrak{m}_R \cap R'$.*

*Proof.* By (3.25.11) the identity map $1 : k \to \overline{k}$ extends to a ring homomorphism $S \to \overline{k}$ where $(S, \mathfrak{m}_S)$ is a valuation ring of $k(\mathfrak{m}_R)/k$. Considering the quotient map $\pi : R \to k(\mathfrak{m}_R)$ let $R' \coloneqq \pi^{-1}(S)$. The restriction map

$$\pi' : R' \to S \to k(\mathfrak{m}_S)$$

is surjective, and so the kernel $\mathfrak{m}_{R'} \coloneqq \ker(\pi') = \mathfrak{m}_R \cap R'$ is a maximal ideal of $R'$ and $k(\mathfrak{m}_{R'}) \cong k(\mathfrak{m}_S)$. Observe that $R' \subseteq R$ and $\mathfrak{m}_R \subseteq \mathfrak{m}_{R'}$. If $x \notin R$ then $x^{-1} \in \mathfrak{m}_R \implies x^{-1} \in R$ by (3.25.2). If $x \in R \setminus R'$ then $\pi(x) \notin S \implies \pi(x^{-1}) \in S \implies x^{-1} \in R'$. Therefore we conclude that $R'$ is a valuation ring with maximal ideal $\mathfrak{m}_{R'}$ and algebraic residue field as required. ■

The following result was proven in the case of a finitely generated $k$-algebra (3.35.28), where equality holds. In general only the inequality holds by considering for example the case $\dim K = 0$. The proof below is taken from [Rey11].

**Lemma 6.1.91**
*Let $R$ be a subring of a finitely generated field extension $K/k$. Then $\dim R \leq \operatorname{trdeg}(K/k)$.*

*Proof.* Consider a chain of prime ideals

$$\mathfrak{p}_0 \subsetneq \ldots \subsetneq \mathfrak{p}_n \subset R$$

Choose $x_i \in \mathfrak{p}_i \setminus \mathfrak{p}_{i-1}$ and define $A \coloneqq k[x_1, \ldots, x_n] \subset R$ with $K' \coloneqq \operatorname{Frac}(A) \subset K$. Then there is a chain of prime ideals of $A$

$$\mathfrak{p}_0 \cap A \subsetneq \ldots \subsetneq \mathfrak{p}_n \cap A$$

whence $n \leq \dim A \overset{(3.35.28)}{=} \operatorname{trdeg}(K'/k) \overset{(3.19.140)}{\leq} \operatorname{trdeg}(K/k)$. Taking supremum over $n$ we reach the required result. ■

**Lemma 6.1.92**
*Let $R \subseteq S$ be proper valuation rings of $K$ with $\dim R < \infty$. Then $\dim S \leq \dim R$, with equality iff $R = S$.*

*Proof.* Firstly we claim that $\mathfrak{m}_S \subseteq \mathfrak{m}_R$. For $x \in \mathfrak{m}_S \implies x^{-1} \notin S \implies x^{-1} \notin R \implies x \in \mathfrak{m}_R$.

Secondly if $R \subsetneq S$ then $\mathfrak{m}_S \subsetneq \mathfrak{m}_R$. For given $x \in S \setminus R$ we have $x^{-1} \in \mathfrak{m}_R \implies x^{-1} \in S \implies x^{-1} \in S^\star \implies x^{-1} \notin \mathfrak{m}_S$.

Let $\mathfrak{p}_0 \subsetneq \ldots \subsetneq \mathfrak{p}_n$ be a chain of prime ideals in $S$. As these are contained in $\mathfrak{m}_S$, they are also contained in $\mathfrak{m}_R$ and therefore $R$. Therefore this constitutes a chain of prime ideals in $R$ from which we deduce the inequality. Further if $R \subsetneq S$ then we may always extend the chain by at least one prime ideal, namely $\mathfrak{m}_R$. ■

**Proposition 6.1.93**
*Let $X = V(\mathfrak{a}) \subset \mathbb{A}^n_k$ be an integral affine variety, and $\mathcal{O}_X(X) \subset R \subset k(X)$ a valuation ring. Then there exists a point $x \in X\big(\overline{k}\big)$ such that $R$ dominates $\mathcal{O}_{X,x}$. Further $\dim R \leq \dim X$ with equality iff $\mathcal{O}_{X,x} = R$ as subrings of $k(X)$.*

*Proof.* By (6.1.90) we may assume that $k(\mathfrak{m}_R)/k$ is algebraic. Choose any embedding $k(\mathfrak{m}_R)/k \hookrightarrow \overline{k}/k$ (3.19.61). Then there is a composite $k$-algebra homomorphism

$$\mathcal{O}_X(X) \hookrightarrow R \to k(\mathfrak{m}_R) \hookrightarrow \overline{k}$$

Let $x_i$ be the image of the affine coordinates $\overline{X}_i$ under this homomorphism. The image $k[x_1, \ldots, x_n]$ is a (finite) field extension of $k$ by (3.19.44). Then under this map the image of $f$ is $f(x)$, so the kernel is precisely $\mathfrak{M}_{X,x} = \mathfrak{m}_R \cap \mathcal{O}_X(X)$. Suppose that $f \notin \mathfrak{M}_{X,x}$ then $f \notin \mathfrak{m}_R \implies f^{-1} \in R$. In particular we conclude that $\mathcal{O}_{X,x} = \mathcal{O}_X(X)_{\mathfrak{M}_{X,x}} \subset R$. Further $\mathfrak{m}_{X,x} = \mathfrak{M}_{X,x}\mathcal{O}_{X,x} \subseteq \mathfrak{m}_R R \subseteq \mathfrak{m}_R$, so $R$ dominates $\mathcal{O}_{X,x}$ by (3.25.4).

The final statement follows from (6.1.92) and $\dim X = \dim \mathcal{O}_{X,x}$ (6.1.57). ■

## 6.1.13 Products of Affine Varieties

In what follows we let $A_{\text{red}}$ denote the reduced ring (resp $k$-algebra) $A/N(A)$.

© David Rufino CC BY-NC-ND 4.0

**Proposition 6.1.94**
*There is a canonical isomorphism*

$$\begin{aligned} \psi : k[X_1,...,X_n] \otimes_k k[Y_1,...,Y_m] &\xrightarrow{\sim} k[X_1,...,X_n,Y_1,...,Y_m] \\ F(X_1,...,X_n) \otimes G(Y_1,...,Y_m) &\to \quad F(X_1,...,X_n)G(Y_1,...,Y_m) \end{aligned}$$

*Let* $\mathfrak{a} \lhd k[X_1,...,X_n]$ *and* $\mathfrak{b} \lhd k[Y_1,...,Y_m]$ *be ideals. Then there is a commutative diagram*

$$\begin{array}{ccc} k[X_1,...,X_n] \otimes_k k[Y_1,...,Y_m] & \xrightarrow[\sim]{\psi} & k[X_1,...,X_n,Y_1,...,Y_m] \\ \downarrow{\scriptstyle \pi_1 \otimes \pi_2} & & \downarrow \\ \frac{k[X_1,...,X_n]}{\mathfrak{a}} \otimes_k \frac{k[Y_1,...,Y_m]}{\mathfrak{b}} & \xrightarrow[\sim]{\overline{\psi}} & \frac{k[X_1,...,X_n,Y_1,...,Y_m]}{\mathfrak{a}^e+\mathfrak{b}^e} \end{array}$$

$$\overline{F} \otimes \overline{G} \to \overline{F(X_1,...,X_n)G(Y_1,...,Y_m)}$$

*where the bottom arrow is uniquely determined and an isomorphism. The nilradical of the right hand side is* $\sqrt{\mathfrak{a}^e+\mathfrak{b}^e}/(\mathfrak{a}^e+\mathfrak{b}^e)$ *and so there is a corresponding isomorphism*

$$\left(\frac{k[X_1,...,X_n]}{\mathfrak{a}} \otimes_k \frac{k[Y_1,...,Y_m]}{\mathfrak{b}}\right)_{\text{red}} \xrightarrow{\sim} \frac{k[X_1,...,X_n,Y_1,...,Y_m]}{\sqrt{\mathfrak{a}^e+\mathfrak{b}^e}}$$

*Proof.* By (3.4.37) there is a unique $k$-algebra homomorphism $\psi$ such that $\psi(F \otimes 1) = F$ and $\psi(1 \otimes G) = G$, whence $\psi(F \otimes G) = FG$. Similarly there is an inverse morphism $\phi$ such that $\phi(X_i) = X_i \otimes 1$ and $\phi(Y_j) = 1 \otimes Y_j$. Evidently $\phi(F(X_1,...,X_n)) = F \otimes 1$ and $\phi(G(Y_1,...,Y_m)) = 1 \otimes G$. This shows that $\phi \circ \psi = \mathbf{1}$. Similarly $(\psi \circ \phi)(X_i) = X_i$ and $(\psi \circ \phi)(Y_j) = Y_j$. This shows that $\psi \circ \phi = \mathbf{1}$ as required.

There is a commutative diagram

$$\begin{array}{ccc} k[X_1,...,X_n] & \xrightarrow{i} & k[X_1,...,X_n] \otimes_k k[Y_1,...,Y_m] \\ & \searrow & \wr \downarrow \psi \\ & & k[X_1,...,X_n,Y_1,...,Y_m] \end{array}$$

By (3.4.42) $\mathfrak{a} \otimes k[Y_1,...,Y_m]$ is the extension of $\mathfrak{a}$ under $i$. As the diagram commutes we may show that $\psi(\mathfrak{a} \otimes k[Y_1,...,Y_m]) = \mathfrak{a}^e$ and similarly $\psi(k[X_1,...,X_n] \otimes \mathfrak{b}) = \mathfrak{b}^e$. Therefore by (3.3.51)

$$\psi(\mathfrak{a} \otimes k[Y_1,...,Y_m] + k[X_1,...,X_n] \otimes \mathfrak{b}) = \mathfrak{a}^e + \mathfrak{b}^e$$

By (3.4.29) $\ker(\pi_1 \otimes \pi_2) = \mathfrak{a} \otimes k[Y_1,...,Y_m] + k[X_1,...,X_n] \otimes \mathfrak{b}$ therefore $\overline{\psi}$ exists and is an isomorphism by (...). ∎

**Lemma 6.1.95**
*Let* $X = V(\mathfrak{a}) \subset \mathbb{A}_k^n$ *and* $Y = V(\mathfrak{b}) \subset \mathbb{A}_k^m$*. Consider the affine variety* $\widetilde{X \times Y} := V\left(\sqrt{\mathfrak{a}^e+\mathfrak{b}^e}\right) \subset \mathbb{A}_k^{n+m}$*. The projection maps*

$$\begin{aligned} \pi_X &: \widetilde{X \times Y}\left(\overline{k}\right) \to X\left(\overline{k}\right) \\ \pi_Y &: \widetilde{X \times Y}\left(\overline{k}\right) \to Y\left(\overline{k}\right) \end{aligned}$$

*are regular and there is an isomorphism*

$$\begin{aligned} \left(\mathcal{O}_X(X) \otimes_k \mathcal{O}_Y(Y)\right)_{\text{red}} &\xrightarrow{\sim} \mathcal{O}_{\widetilde{X \times Y}}\left(\widetilde{X \times Y}\right) \\ f \otimes g \to \quad & (f \circ \pi_X) \cdot (g \circ \pi_Y) \end{aligned}$$

*where* $\pi_X : \widetilde{X \times Y}\left(\overline{k}\right) \to X\left(\overline{k}\right)$ *and* $\pi_Y : \widetilde{X \times Y}\left(\overline{k}\right) \to Y\left(\overline{k}\right)$ *are the obvious regular projection maps.*

*Proof.* It's straightforward to show that $\pi_X$ and $\pi_Y$ are regular using the definition (6.1.29). By (6.1.94) and (6.1.26) there is a unique morphism making the following diagram commute which must be an isomorphism, and we simply need to check it is of the required form.

© David Rufino CC BY-NC-ND 4.0

$$\begin{array}{ccc} (k[X] \otimes_k k[Y])_{\text{red}} & \xrightarrow{\sim} & k\big[\widetilde{X \times Y}\big] \\ \downarrow \sim & & \downarrow \sim \\ (\mathcal{O}_X(X) \otimes_k \mathcal{O}_Y(Y))_{\text{red}} & \dashrightarrow & \mathcal{O}_{\widetilde{X \times Y}}\big(\widetilde{X \times Y}\big) \end{array}$$

Given $f \in k[X]$ and $g \in k[Y]$ where $f = \overline{F}$ and $g = \overline{G}$. Then the image of $f \otimes g$ is simply $h := \overline{F(X_1, ..., X_n)G(Y_1, ..., Y_n)}$ and

$$h(x_1, ..., x_n, y_1, ..., y_m) = f(x_1, ..., x_n)g(y_1, ..., y_m)$$

from which the result follows. ∎

**Proposition 6.1.96 (Affine Product)**
*Let $X = V(\mathfrak{a}) \subset \mathbb{A}^n_k$ and $Y = V(\mathfrak{b}) \subset \mathbb{A}^m_k$ be affine varieties. Then there is a bijection*

$$\begin{aligned} X\big(\overline{k}\big) \times Y\big(\overline{k}\big) &\xrightarrow{\sim} \widetilde{X \times Y}\big(\overline{k}\big) \\ (x, y) &\to (x_1, ..., x_n, y_1, ..., y_m) \end{aligned}$$

*Define $\Big(X\big(\overline{k}\big) \times Y\big(\overline{k}\big), \mathcal{O}_{X \times Y}\Big)$ as a local space of functions such that this bijection is a regular isomorphism.*

*The projection maps $\pi_X : X \times Y \to X$ and $\pi_Y : X \times Y \to Y$ are regular and for every affine variety $Z$ we have a natural bijection*

$$\begin{aligned} \text{Mor}((Z, \mathcal{O}_Z), (X \times Y, \mathcal{O}_{X \times Y})) &\xrightarrow{\sim} \text{Mor}((Z, \mathcal{O}_Z), (X, \mathcal{O}_X)) \times \text{Mor}((Z, \mathcal{O}_X), (Y, \mathcal{O}_Y)) \\ f &\to ((\pi_X \circ f), (\pi_Y \circ f)) \end{aligned}$$

*Proof.* Evidently it is enough to prove all the same properties for $\widetilde{X \times Y}$, as then they follow immediately for $X \times Y$ by composing with $\theta$, which is by construction a regular isomorphism. We have natural isomorphisms

$$\begin{aligned} \text{Mor}\big(Z, \widetilde{X \times Y}\big) &\xrightarrow{\sim} \text{AlgHom}_k\Big(\mathcal{O}_{\widetilde{X \times Y}}\big(\widetilde{X \times Y}\big), \mathcal{O}_Z(Z)\Big) \\ &\xrightarrow{\sim} \text{AlgHom}_k\Big((\mathcal{O}_X(X) \otimes_k \mathcal{O}_Y(Y))_{\text{red}}, \mathcal{O}_Z(Z)\Big) \\ &\xrightarrow{\sim} \text{AlgHom}_k(\mathcal{O}_X(X) \otimes_k \mathcal{O}_Y(Y), \mathcal{O}_Z(Z)) \\ &\xrightarrow{\sim} \text{AlgHom}_k(\mathcal{O}_X(X), \mathcal{O}_Z(Z)) \times \text{AlgHom}_k(\mathcal{O}_Y(Y), \mathcal{O}_Z(Z)) \\ &\xrightarrow{\sim} \text{Mor}(Z, X) \times \text{Mor}(Z, Y) \end{aligned}$$

by (6.1.95) and (6.1.30). We may verify that the image of $1_{\widetilde{X \times Y}}$ is $(\pi_X, \pi_Y)$ and so the result follows from naturality in $Z$. ∎

**Proposition 6.1.97 (Affine Varieties are Separated)**
*Let $f : X \to Y$ be a regular map of affine varieties. Then the graph*

$$\Gamma_f(K) := \{(x, f(x)) \mid x \in X(K)\}$$

*is a closed subset of $(X \times Y)(K)$. In particular the diagonal*

$$\Delta_X := \{(x, x) \mid x \in X(K)\}$$

*is a closed subset of $(X \times X)(K)$.*

*Proof.* Let $\overline{X}_1, ..., \overline{X}_n$ and $\overline{Y}_1, ..., \overline{Y}_m$ be the coordinate functions of $k[X \times Y]$. Further we may suppose that $f = \big(\overline{F}_1, ..., \overline{F}_m\big)$ for $F_j \in k[X_1, ..., X_n]$. Then $\Gamma_f$ is the zero locus of the ideal

$$\Big\langle \overline{Y}_1 - F_1\big(\overline{X}_1, ..., \overline{X}_n\big), ..., \overline{Y}_m - F_m\big(\overline{X}_1, ..., \overline{X}_n\big) \Big\rangle$$

∎

**Proposition 6.1.98 (Criteria for Product to be Integral)**
*Let $X, Y$ be integral affine varieties such that either $X$ or $Y$ is* geometrically *integral (e.g. if $k$ is algebraically closed). Then $X \times Y$ is integral.*

*Proof.* See (3.36.42). ∎

© David Rufino CC BY-NC-ND 4.0

### 6.1.14 Affine Curves

**Definition 6.1.99**
*An **affine curve** is an integral affine variety of dimension* 1.

*For example an irreducible polynomial $F(X,Y)$ yields an affine curve and is known as a plane curve. The singular points are precisely the common roots of*

$$F, \frac{\partial F}{\partial X}, \frac{\partial F}{\partial Y}$$

*in $\overline{k}^2$. If there are no such roots then the affine curve is non-singular.*

**Proposition 6.1.100 (Characterisation Regular Point on a curve)**
*Let $X = V(\mathfrak{a})$ be an affine curve and $x \in X\left(\overline{k}\right)$. Then the following are equivalent*

*a) $x$ is regular*
*b) $\mathcal{O}_{X,x}$ is a discrete valuation ring*
*c) $\mathcal{O}_{X,x}$ is an integrally closed domain*

When $k$ is perfect then this is equivalent to $x$ being a non-singular point.

*Proof.* Note that by assumption $X$ is integral, so $\mathcal{O}_{X,x}$ is an integral domain by (6.1.61).

a)-c) then follows directly from (3.31.2) and the fact $\dim \mathcal{O}_{X,x} = \dim X = 1$ (6.1.57). The final statement follows from (6.1.75). ∎

**Proposition 6.1.101**
*Let $X = V(\mathfrak{a})$ be a regular affine curve. Then there exists a bijection*

$$X_0 \coloneqq X\left(\overline{k}\right) / \operatorname{Aut}\left(\overline{k}/k\right) \longleftrightarrow \{\mathcal{O}_X(X) \subset R \subsetneq k(X) \mid R \text{ valuation ring}\}$$

*Proof.* The map $x \to \mathcal{O}_{X,x}$ is well-defined by (6.1.100), (6.1.9) and (6.1.60). If $\mathcal{O}_{X,x} = \mathcal{O}_{X,y}$ then $\mathcal{O}^\star_{X,x} = \mathcal{O}^\star_{X,y} \implies \mathfrak{m}_{X,x} = \mathfrak{m}_{X,y} \implies \mathfrak{M}_{X,x} = \mathfrak{M}_{X,y} \implies x \sim y$ by (6.1.9) so the map is injective.

To show surjectivity, observe by (6.1.93) there exists $x \in X\left(\overline{k}\right)$ such that $\mathcal{O}_{X,x} \subseteq R$. Furthermore by repeated application of (6.1.92) we see

$$0 < \dim R \leq \dim \mathcal{O}_{X,x} \overset{(6.1.57)}{=} \dim X = 1$$

This forces $\dim R = \dim \mathcal{O}_{X,x} = 1$ and so $R = \mathcal{O}_{X,x}$ by the same result. ∎

© David Rufino CC BY-NC-ND 4.0

## 6.2 Abstract Varieties

In order to deal with affine, quasi-affine, projective and quasi-projective varieties uniformly it is convenient to define an abstract variety. Further some constructions are more natural in this setting (for example product variety). This is analogous to the notion of manifold, and much closer to the notion of scheme. Some differences

a) Sections of the structure sheaf are concrete $\overline{k}$-valued functions rather than abstract elements of a ring in the case of a scheme.
b) When $k$ is not algebraically closed the underlying topological space of the corresponding scheme is the Kolmogorov Quotient (sober-ified to include all the possible generic points). When $k$ is algebraically closed the only difference is the inclusion of generic points.
c) Varieties here are always Noetherian (i.e. there is a finite affine cover and the affines are themselves Noetherian).

The differences are primarily psychological and almost all the theory carries over *mutatis mutandis* to the scheme setting, with additional hypotheses where necessary.

**Definition 6.2.1 (Abstract Variety)**
*A **variety** over a field $k$ is a space with functions $(X, \mathcal{O}_X)$ such that*

- *There exists a finite open cover $X = \bigcup_{i=1}^n U_i$ together with isomorphisms*
$$\theta_i : \left(U_i, \mathcal{O}_X|_{U_i}\right) \cong \left(X_i\left(\overline{k}\right), \mathcal{O}_{X_i}\right)$$
*for some affine varieties $X_1, ..., X_n$ in the sense of (6.1.4).*

*The sheaf $\mathcal{O}_X$ is called the **structure sheaf** or **sheaf of regular functions**. A regular morphism is then a morphism of space with functions.*

*It is immediate that every affine variety is an abstract variety.*

**Proposition 6.2.2**
*A variety $(X, \mathcal{O}_X)$ is a local space with functions. That is for all $f \in \mathcal{O}_X(U)$, the set*
$$D(f) := \{x \in U \mid f(x) \neq 0\}$$
*is open and $f|_U$ is invertible.*

*Proof.* We have an open affine cover $X = \bigcup_{i=1}^n U_i$. Then we have
$$D(f) = \bigcup_{i=1}^n D\left(f|_{U_i}\right)$$
By (6.1.41) $D\left(f|_{U_i}\right)$ is open and $f|_{U_i}$ is invertible. We deduce that $D(f)$ is open, and using the sheaf conditions that $f|_{D(f)}$ is invertible. ∎

**Definition 6.2.3 (Affine (Abstract) Variety)**
*Let $X$ be a space of functions. We say it is an **affine variety** if it is isomorphic to $\left(Y\left(\overline{k}\right), \mathcal{O}_Y\right)$, where $Y$ is an affine variety in the sense of (6.1.4) and $\mathcal{O}_Y$ is the corresponding sheaf of regular functions (6.1.25).*

**Proposition 6.2.4**
*Let $X$ be a variety and $f : X \to \overline{k}$ be a function. Then it is regular iff the corresponding map $f : X \to \mathbb{A}^1_k\left(\overline{k}\right)$ is a regular morphism of abstract varieties.*

*Proof.* Suppose that $f : X \to \mathbb{A}^1_k\left(\overline{k}\right)$ is a regular morphism. Evidently the identity map $1 : \mathbb{A}^1_k\left(\overline{k}\right) \to \overline{k}$ is a regular function. Therefore by definition $f : X \to \overline{k}$ is a regular function.

Conversely if $f : X \to \overline{k}$ is regular then so is $f|_U$ for every affine open subset $U \subset X$. By (6.1.29) and (6.1.39) this a regular morphism of abstract varieties $U \to \mathbb{A}^1_k\left(\overline{k}\right)$. By (4.3.27) we have $f$ is a regular morphism. ∎

### 6.2.1 Topological Properties

**Definition 6.2.5 (Integral Variety)**
*We say a variety $X$ is **integral** (or irreducible) if it is irreducible as a topological space.*

**Proposition 6.2.6 (Criteria for Integrality)**
*Let $X$ be a variety. Then the following are equivalent*

© David Rufino CC BY-NC-ND 4.0

*a) $X$ is integral*
*b) $\mathcal{O}_X(U)$ is integral for every open subset $U \subset X$*

**Proposition 6.2.7 (Varieties are Noetherian and Hereditarily Quasi-Compact)**
*Let $X$ be a variety. Then $X$ is Noetherian as a topological space and every open subset is quasi-compact (and Noetherian).*

*Proof.* (4.1.78) and (4.1.114). ∎

**Corollary 6.2.8**
*Let $X$ be a variety. Then it may be expressed uniquely as a finite union of (closed) irreducible components.*

$$X = X_1 \cup ... \cup X_n$$

*Proof.* See (4.1.76). ∎

**Proposition 6.2.9**
*Suppose $(X, \mathcal{O}_X)$ is a space with functions such that $X = \bigcup_{i=1}^n U_i$ and $\left(U_i, \mathcal{O}_X|_{U_i}\right)$ is a variety. Then so is $(X, \mathcal{O}_X)$.*

*Proof.* Each $U_i$ is a finite union of affine varieties and therefore so is $X$. ∎

**Proposition 6.2.10**
*Let $X$ be a variety. The (affine) principal open subsets form a base for the topology on $X$.*

**Proposition 6.2.11 (Varieties are Symmetric)**
*Let $X$ be a variety. Then $X$ is symmetric as a topological space and $\overline{\{x\}}$ is finite for all $x \in X$. When $k$ is algebraically closed then every point is closed and therefore $X$ is Kolmogorov.*

*Furthermore for every open neighbourhood $U$ of $x$ we have $\overline{\{x\}} = \mathrm{cl}_U(\{x\})$.*

*Proof.* That $X$ is symmetric is from (4.1.44) and the affine case (6.1.18). The last statement follows from (4.1.43), then finiteness follows from the affine case. ∎

**Proposition 6.2.12 (Closed Subsets of Dimension 0)**
*Let $X$ be a variety and $Z \subset X$ a closed subset. Then the following are equivalent*

*a) $\dim Z = 0$*
*b) $Z = \overline{\{z_1\}} \cup ... \cup \overline{\{z_n\}}$ for some $z_1, ..., z_n \in Z$*
*c) $Z$ is finite*

*Proof.* Recall $X$ is Noetherian (6.2.7) and therefore by (4.1.91) we have $a) \iff b)$. Then $b) \implies c)$ is (6.2.11) and conversely if $Z$ is finite then $Z = \bigcup_{z \in Z} \overline{\{z\}}$ is a finite union. ∎

**Definition 6.2.13 (Separated Variety)**
*Let $X$ be a variety. Then we say $X$ is **separated** if for every affine variety $Z$ and pair of morphisms $\phi_1, \phi_2 : Z \rightrightarrows X$ the equaliser*

$$\{z \in Z \mid \phi_1(z) = \phi_2(z)\}$$

*is closed in $Z$.*

*By (6.1.43) every affine variety is separated.*

**Proposition 6.2.14**
*Let $X$ be a separated variety and $Z$ a variety. Then the equaliser of any two maps $\phi_1, \phi_2 : Z \rightrightarrows X$ is closed.*

*Proof.* By definition $Z = \bigcup_{i=1}^n U_i$ where $U_i$ is affine. By assumption the equaliser of $\phi_1|_{U_i}, \phi_2|_{U_i} : U_i \rightrightarrows X$ is closed. As a finite union of closed sets is closed, we are done. ∎

## 6.2.2 Local Ring

We recall the definition of local ring (4.3.3) for a variety $X$ with irreducible subset $W$

$$\mathcal{O}_{X,W} := \varinjlim_{U \cap W \neq \varnothing} \mathcal{O}_X(U).$$

Note that if $\overline{W'} = \overline{W}$ then $\mathcal{O}_{X,W} = \mathcal{O}_{X,W'}$, so for example $\mathcal{O}_{X,W} = \mathcal{O}_{X,\overline{W}}$.

In the case $W = \{x\}$ we write $\mathcal{O}_{X,x}$ for the local ring and $\mathfrak{m}_{X,x}$ and $k(\mathfrak{m}_{X,x})$ for the maximal ideal and residue field respectively.

© David Rufino CC BY-NC-ND 4.0

In the case $X = W$ is irreducible then we recover the **field of rational functions** written $k(X)$. For a variety we can make stronger statement about the structure of $k(X)$

**Proposition 6.2.15**
*Let $X$ be an integral variety, $x \in X$ and $U$ an open neighbourhood of $x$. Then there are canonical injections*

$$\mathcal{O}_X(U) \hookrightarrow \mathcal{O}_{X,x} \hookrightarrow k(X)$$

*which makes $k(X)$ the field of fractions for $\mathcal{O}_{X,x}$. When $U$ is affine then $k(X)$ is also the field of fractions for $\mathcal{O}_X(U)$. These maps commute with the restriction maps on $\mathcal{O}_X$, which are also injective.*

*Proof.* By restricting to an affine open cover we may check that the restriction maps $\rho_{UV}$ are injective. This shows that the first map is injective because every stalk can be lifted to some open neighbourhood.

Let $U$ be an affine open neighbourhood of $x$ then by (4.3.16) $k(X) \xrightarrow{\sim} k(U)$. We may deduce from the affine case (6.1.60) that $k(X)$ is the field of fractions for $\mathcal{O}_{X,x}$ and $\mathcal{O}_X(U)$. ∎

As $\mathcal{O}_{X,x}$ is isomorphic to the local ring of an affine neighbourhood we can recover many statements from the affine case.

**Proposition 6.2.16**
*Let $X$ be a variety and $W \subset X$ an irreducible subset. Then the following are equivalent*

*a) $W$ is contained in only one irreducible component*
*b) $\mathcal{O}_{X,W}$ is an integral domain*

*Proof.* Firstly we may reduce to the case $W$ is closed by considering $\overline{W}$. As $X$ is a variety, $W \cap U \neq \varnothing$ for $U$ affine. By (4.1.73) we may reduce to the case $X$ is affine, which is (6.1.61). ∎

**Proposition 6.2.17 (Dominant Morphisms induce Function Field Maps)**
*Let $X, Y$ be integral varieties and $\phi : X \to Y$ a dominant regular morphism. Then there is a canonical injective $k$-algebra homomorphism*

$$\begin{aligned} \phi_\star : k(Y) &\hookrightarrow k(X) \\ (V, \sigma) &\to (\phi^{-1}(V), \sigma \circ \phi) \end{aligned}$$

*This is functorial in the sense that*

$$(\phi \circ \psi)_\star = \psi_\star \circ \phi_\star$$

*If $\phi$ is an open immersion then $\phi_\star$ is an isomorphism.*

*Proof.* See (4.3.7) and (4.3.16). ∎

**Proposition 6.2.18 (Function Field Maps are compatible)**
*Let $X, Y$ be integral varieties, $\phi : X \to Y$ a dominant regular morphism and $y \in \overline{\{\phi(x)\}}$. Then for all neighbourhoods $V$ of $y$ we have a commutative diagram*

$$\begin{array}{ccc} \mathcal{O}_Y(V) & \xrightarrow{\phi_V^\sharp} & \mathcal{O}_X(\phi^{-1}(V)) \\ \downarrow & & \downarrow \\ \mathcal{O}_{Y,\phi(x)} & \xrightarrow{\phi_x} & \mathcal{O}_{X,x} \\ \downarrow & & \downarrow \\ k(Y) & \xrightarrow{\phi_\star} & k(X) \end{array}$$

*In particular when identified as subrings of $k(X)$ and $k(Y)$ we have the identities*

$$\phi_\star^{-1}(\mathcal{O}_{X,x}) = \mathcal{O}_{Y,\phi(x)}$$

*and*

$$\phi_\star^{-1}(\mathcal{O}_X(\phi^{-1}(V))) = \mathcal{O}_Y(V)$$

*Proof.* See (4.3.9) and (6.2.15). ∎

© David Rufino CC BY-NC-ND 4.0

**Lemma 6.2.19**
*For a variety $X$, $R$ an algebraic $k$-algebra and $\phi : \mathcal{O}_{X,x} \to R$ a $k$-algebra homomorphism Then $\phi^{-1}(\mathfrak{m}) = \mathfrak{m}_{X,x}$ for every maximal ideal $\mathfrak{m} \lhd R$.*

*Proof.* Restatement of (6.1.62), since $\mathcal{O}_{X,x}$ is isomorphic to local ring of affine neighbourhood. ∎

**Proposition 6.2.20 (Equivalent Points)**
*Let $X$ be a variety and $x, y \in X$. Then the following are equivalent*

*a) $x$, $y$ are topologically indistinguishable*
*b) $x \in \overline{\{y\}}$*
*c) $y \in \overline{\{x\}}$*
*d) $\overline{\{x\}} = \overline{\{y\}}$*
*e) $[x] = [y]$ in the Kolmogorov Quotient $X_0$*

*When $X$ is integral these are equivalent to $\mathcal{O}_{X,x} = \mathcal{O}_{X,y}$ as subrings of $k(X)$.*

*Proof.* By (6.2.11) $X$ is symmetric and the equivalent conditions $a$)-$e$) follow from (4.1.42).

If $x \sim y$ then by definition $\mathcal{O}_{X,x} = \mathcal{O}_{X,y}$.

Conversely suppose $f \in \mathcal{O}_{X,x} \setminus \mathcal{O}_{X,y}$ in $k(X)$. In some open neighbourhood $U$ of $x$ we have $f = \frac{g}{h}$ for $g, h \in \mathcal{O}_X(U)$ and $h(x) \neq 0$. By definition we must have $h(y) = 0$. Therefore $D(h)$ contains $x$ but not $y$, and therefore $x$ and $y$ are not topologically indistinguishable. ∎

## 6.2.3 Affine Varieties

We restate various results relating to "abstract" affine varieties to avoid the need to switch notations later on.

**Proposition 6.2.21 (Properties of the Structure Sheaf)**
*Let $X$ be an abstract variety $U \subset X$ an affine open subset. Then the following properties hold*

*a) For all irreducible subsets $W \subset U$ we have $\mathfrak{M}_{U,W} \coloneqq \{f \in \mathcal{O}_X(U) \mid f|_W = 0\}$ is a prime ideal of $\mathcal{O}_X(U)$. Further we have a canonical local isomorphism*

$$\begin{array}{ccc} \mathcal{O}_X(U) & \longrightarrow & \mathcal{O}_X(U)_{\mathfrak{M}_{U,W}} \\ & \searrow^{\rho_W} & \downarrow \sim \\ & & \mathcal{O}_{X,W} \end{array}$$

*which is the unique homomorphism making this diagram commute.*

*The inverse image of $\mathfrak{m}_{X,W}$ is $\mathfrak{M}_{U,W}\mathcal{O}_X(U)_{\mathfrak{M}_{U,W}}$ and $\mathfrak{M}_{U,W}$ in $\mathcal{O}_X(U)_{\mathfrak{M}_{U,W}}$ and $\mathcal{O}_X(U)$ respectively.*

*When $W \subset \overline{\{x\}}$ then $\mathfrak{M}_{U,W} = \mathfrak{M}_{U,x}$ is a maximal ideal.*

*b) There is a well-defined bijection*

$$\begin{aligned} U_0 &\longleftrightarrow \operatorname{Specm}(\mathcal{O}_X(U)) \\ x &\to \mathfrak{M}_{U,x} \end{aligned}$$

*Further $\mathfrak{M}_{U,x} = \mathfrak{M}_{U,x'} \iff x \in \overline{\{x'\}} \iff x$ and $x'$ are topologically indistinguishable.*

*c) For all $f \in \mathcal{O}_X(U)$ we have $\left(D(f), \mathcal{O}_X|_{D(f)}\right)$ is an affine variety. Further we have a canonical isomorphism*

$$\begin{array}{ccc} \mathcal{O}_X(U) & \longrightarrow & \mathcal{O}_X(U)_f \\ & \searrow^{\rho_{XD(f)}} & \downarrow \sim \\ & & \mathcal{O}_X(D(f)) \end{array}$$

*which is the unique homomorphism making this diagram commute. For irreducible $W \subset D(f)$, under this isomorphism the inverse image of $\mathfrak{M}_{D(f),W}$ is $\left(\mathfrak{M}_{U,W}\right)_f$. Similarly the following diagram commutes*

© David Rufino CC BY-NC-ND 4.0

$$\begin{array}{ccc} \mathcal{O}_X(U)_f & \xrightarrow{\sim} & \mathcal{O}_X(D(f)) \\ \downarrow & & \downarrow \\ \mathcal{O}_X(U)_{\mathfrak{M}_{U,W}} & \xrightarrow{\sim} & \mathcal{O}_{X,W} \end{array}$$

*and the inverse image of* $\mathfrak{m}_{X,W}$ *is* $\mathfrak{M}_{D(f),W}$ *and* $\left(\mathfrak{M}_{U,W}\right)_f$ *in* $\mathcal{O}_X(D(f))$ *and* $\mathcal{O}_X(U)_f$ *respectively.*

*d) Suppose* $X, Y$ *are affine varieties and* $W \subset X$ *is irreducible and* $Z \subset \overline{\phi(W)}$ *for some irreducible* $Z \subset Y$*. Let* $\phi : X \to Y$ *be a regular morphism. Then the stalk map* $\phi_W : \mathcal{O}_{Y,Z} \to \mathcal{O}_{X,W}$ *(see (4.3.5)) makes the following diagram commute*




*where the bottom arrow is the localisation map induced by* $\phi^\sharp$ *(3.6.22). In fact it is the unique map making either square commute.*

**Proposition 6.2.22 (Affine Mapping Property)**
*Let* $Y$ *be an affine variety, and* $X$ *a variety. Then there is a canonical bijection*

$$\begin{aligned} \mathrm{Mor}(X,Y) &\xrightarrow{\sim} \mathrm{AlgHom}_k(\mathcal{O}_Y(Y), \mathcal{O}_X(X)) \\ \phi &\to \ \phi_Y^\sharp \end{aligned}$$

*When* $X$ *is affine and* $Z \subset Y$*,* $W \subset X$ *are irreducible subsets, then*

$$Z \subset \overline{\phi(W)} \iff \left(\phi_Y^\sharp\right)^{-1}\left(\mathfrak{M}_{X,W}\right) \subset \mathfrak{M}_{Y,Z}$$

*Proof.* When $X$ is affine then this is just (6.1.30) and (6.1.32). In general suppose $X = \bigcup_{i=1}^n U_i$ is an affine covering then we have a commutative diagram

$$\begin{array}{ccc} \mathrm{Mor}(X,Y) & \xrightarrow{(-)^\sharp} & \mathrm{AlgHom}_k(\mathcal{O}_Y(Y), \mathcal{O}_X(X)) \\ \downarrow \alpha & & \downarrow \beta \\ \prod_{i=1}^n \mathrm{Mor}(U_i, Y) & \xrightarrow{\gamma} & \prod_{i=1}^n \mathrm{AlgHom}_k(\mathcal{O}_Y(Y), \mathcal{O}_X(U_i)) \end{array}$$

where $\alpha(\phi)_i = \phi|_{U_i}$ and $\beta(\psi)_i = \rho_{XU_i} \circ \psi$ and $\gamma$ is the same definition as $(-)^\sharp$.

By the affine case $\gamma$ is bijective, and evidently $\alpha$ is injective. Therefore $\beta \circ (-)^\sharp$ is injective, whence $(-)^\sharp$ is injective. To show surjectivity consider $\psi \in \mathrm{AlgHom}_k(\mathcal{O}_Y(Y), \mathcal{O}_X(X))$ then as $\gamma$ is bijective we have maps $\phi_i : U_i \to X$ such that $(\phi_i)^\sharp = \rho_{YU_i} \circ \psi$. Suppose $V \subset U_i \cap U_j$ is affine then

$$(\phi_i|_V)^\sharp = \rho_{U_iV} \circ \rho_{XU_i} \circ \psi = \rho_{XV} \circ \psi$$

Therefore $(\phi_i|_V)^\sharp = \left(\phi_j|_V\right)^\sharp$ and from the affine case $\phi_i|_V = \phi_j|_V$. As the open affines from a base (6.2.10) we conclude $\phi_i|_{U_i \cap U_j} = \phi_j|_{U_i \cap U_j}$. Therefore there exists a morphism $\phi : X \to Y$ such that $\phi|_{U_i} = \phi_i$ and therefore $\beta\left(\phi^\sharp\right) = \beta(\psi)$. Evidently $\beta$ is injective and so $\phi^\sharp = \psi$. ∎

**Corollary 6.2.23**
*There is a full and faithful contravariant functor*

$$\mathbf{AbstractAffVar}_k \longrightarrow \mathbf{ReducedFGAlgebras}_k$$

*which in particular preserves and reflects isomorphisms*

© David Rufino CC BY-NC-ND 4.0

**Corollary 6.2.24**
*Let $X$ be an affine variety, $Y$ a variety and $f \in \mathcal{O}_X(X)$. Then there is a canonical bijection*

$$\begin{aligned} \mathrm{Mor}(D(f), Y) &\xrightarrow{\sim} \mathrm{AlgHom}_k\big(\mathcal{O}_Y(Y), \mathcal{O}_X(X)_f\big) \\ \phi &\to \ \theta_f \circ \phi_Y^\sharp \end{aligned}$$

*where $\theta_f : \mathcal{O}_X(D(f)) \xrightarrow{\sim} \mathcal{O}_X(X)_f$ is the isomorphism (6.2.21).c). The image of $i : D(f) \to X$ is simply the canonical localisation map.*

*Suppose $Y$ is affine and $W \subset D(f)$, $Z \subset Y$ are irreducible subsets. Then $Z \subset \overline{\phi(W)} \iff \big(\theta_f \circ \phi_Y^\sharp\big)^{-1}\Big(\big(\mathfrak{M}_{X,W}\big)_f\Big) \subset \mathfrak{M}_{Y,Z}$.*

*Proof.* We may apply (6.2.22). ∎

**Corollary 6.2.25**
*Let $X$ be an affine variety and $\phi, \psi : X \to Y$ regular morphisms, and $f \in \mathcal{O}_X(X)$. Then $\phi|_{D(f)} = \psi|_{D(f)}$ if and only if $\rho_f \circ \phi^\sharp = \rho_f \circ \psi^\sharp$, where $\rho_f : \mathcal{O}_X(X) \to \mathcal{O}_X(X)_f$ is the canonical localisation map.*

*Proof.* Denote by $i : D(f) \hookrightarrow X$ the inclusion then by (6.2.22) $\phi|_{D(f)} = \psi|_{D(f)} \iff \phi \circ i = \psi \circ i \iff i^\sharp \circ \phi^\sharp = i^\sharp \circ \psi^\sharp$ where $i^\sharp : \mathcal{O}_X(X) \to \mathcal{O}_{D(f)}(D(f))$ is simply the restriction map. We may compose this with the isomorphism (6.2.21).c) to get the required result. ∎

There are some properties of affine varieties which generalise

**Proposition 6.2.26**
*Let $X$ be an affine variety and $f_1, ..., f_r \in \mathcal{O}_X(X)$. Then the following are equivalent*

*a)* $X = \bigcup_{i=1}^r D(f_i)$
*b)* $(f_1, ..., f_r) = \mathcal{O}_X(X)$

*Proof.* We may reduce to the case of $X$ a concrete affine variety. Then

$$X = \bigcup_{i=1}^r D(f_i) \iff \varnothing = \bigcap_{i=1}^r V(f_i) = V(f_1, ..., f_r) \iff k[X] = IV(f_1, ..., f_r) = \sqrt{(f_1, ..., f_r)} \iff k[X] = (f_1, ..., f_r)$$

where we have identified $\mathcal{O}_X(X)$ and $k[X]$. ∎

**Proposition 6.2.27**
*Let $X$ be a variety and $f \in \mathcal{O}_X(X)$. There is a canonical isomorphism*

$$\mathcal{O}_X(X)_f \xrightarrow{\sim} \mathcal{O}_X(D(f))$$

*Proof.* The following is adapted from [Liu06] (Prop. 2.3.12).

Recall (6.2.2) that $D(f)$ is an open subset and $f|_{D(f)}$ is invertible. This shows that the homomorphism is well-defined, and we simply need to show it is bijective, with the case $X$ affine already proven (6.1.28).

Let $X = \bigcup_{i=1}^n U_i$ be an affine open covering then $D(f) = \bigcup_{i=1}^n D\big(f|_{U_i}\big)$. By the affine case we have an isomorphism

$$\mathcal{O}_X(U_i)_{f|_{U_i}} \xrightarrow{\sim} \mathcal{O}_X\Big(D\big(f|_{U_i}\big)\Big)$$

By the sheaf conditions we have an exact sequence of $\mathcal{O}_X(X)$-modules

$$0 \longrightarrow \mathcal{O}_X(X) \longrightarrow \bigoplus_{i=1}^n \mathcal{O}_X(U_i) \longrightarrow \bigoplus_{i,j=1}^n \mathcal{O}_X(U_i \cap U_j)$$

Localising at $f \in \mathcal{O}_X(X)$ is exact (3.6.13) and commutes with direct sums (...) so we obtain the exact sequences

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{O}_X(X)_f & \longrightarrow & \bigoplus_{i=1}^n \mathcal{O}_X(U_i)_{f|_{U_i}} & \longrightarrow & \bigoplus_{i,j=1}^n \mathcal{O}_X(U_i \cap U_j)_{f|_{U_i \cap U_j}} \\ & & \downarrow \alpha & & \downarrow \beta & & \downarrow \gamma \\ 0 & \longrightarrow & \mathcal{O}_X(D(f)) & \longrightarrow & \bigoplus_{i=1}^n \mathcal{O}_X\Big(D\big(f|_{U_i}\big)\Big) & \longrightarrow & \bigoplus_{i,j=1}^n \mathcal{O}_X\Big(D\big(f|_{U_i \cap U_j}\big)\Big) \end{array}$$

where $\alpha, \beta, \gamma$ are just the canonical maps already defined. By the affine case $\beta$ is an isomorphism. We claim that diagram chasing shows $\alpha$ is injective, for suppose that $\alpha(x) = 0$ then $\alpha(x)|_{D(f|_{U_i})} = 0$ whence $x|_{U_i} = 0$, and by exactness $x = 0$. As $U_i \cap U_j$ is a variety, then we conclude $\gamma$ is also injective.

© David Rufino CC BY-NC-ND 4.0

We further claim that $\alpha$ is surjective, for given $y \in \mathcal{O}_X(D(f))$ then there exists $x_i \in \mathcal{O}_X(U_i)_{f_i}$ such that $\beta(x_i) = y|_{D(f|_{U_i})}$. From injectivity of $\gamma$ and exactness of the bottom sequence we conclude that $x_i|_{U_i \cap U_j} = x_j|_{U_i \cap U_j}$, and so by exactness of the top sequence there is $x \in \mathcal{O}_X(X)_f$ such that $x|_{U_i} = x_i$. Consequently $\alpha(x)|_{D(f|_{U_i})} = y|_{D(f|_{U_i})}$. Exactness of the bottom sequence shows that $\alpha(x) = y$ as required. ∎

**Proposition 6.2.28**
*Let $X$ be a variety and suppose $f_1, ..., f_r \in \mathcal{O}_X(X)$ generate $\mathcal{O}_X(X)$. If $D(f_i)$ is an affine variety for all $i = 1...r$ then so is $X$. Furthermore*

$$X = \bigcup_{i=1}^{n} D(f_i)$$

*Proof.* First we claim that $\mathcal{O}_X(X)$ is a reduced finitely generated $k$-algebra. Recall by (6.2.27) there is a $k$-algebra isomorphism $\mathcal{O}_X(X)_{f_i} \xrightarrow{\sim} \mathcal{O}_X(D(f_i))$. By assumption $D(f_i)$ is affine so $\mathcal{O}_X(X)_{f_i}$ is finitely-generated as a $k$-algebra. Therefore by (3.24.21) so is $\mathcal{O}_X(X)$.

By (6.1.36) we may find an affine variety $Z$ together with an isomorphism $\phi^\sharp : \mathcal{O}_Z(Z) \xrightarrow{\sim} \mathcal{O}_X(X)$. By (6.2.22) there is a corresponding morphism of varieties $\phi : X \to Z$, which we wish to show is an isomorphism. Let $\tilde{f}_i := \left(\phi^\sharp\right)^{-1}(f_i)$ then by definition $\phi^{-1}\left(D\left(\tilde{f}_i\right)\right) = D(f_i)$.

We have a commutative diagram




whence the middle arrow is an isomorphism. As $D(f_i)$ is an affine variety we deduce from (6.2.23) that $\phi|_{D(f_i)} : D(f_i) \xrightarrow{\sim} D\left(\tilde{f}_i\right)$ is a regular isomorphism. Further $X = \phi^{-1}(Z)$ whence the $D(f_i)$ cover $X$ and $\phi$ is a regular isomorphism as required. ∎

## 6.2.4 Lifting Stalk Maps

In the same way that stalks can be lifted to an open neighbourhood, for varieties we can lift stalk maps to an open neighbourhood. After reducing to the affine case this is essentially an algebraic statement about localisation.

**Proposition 6.2.29 (Lifting Stalk Maps I)**
*Let $\phi, \psi : X \to Y$ be regular morphisms of varieties and $W \subset X$, $Z \subset Y$ non-empty irreducible subsets. Suppose*

*a) $Z \subset \overline{\phi(W)} = \overline{\psi(W)}$, and*
*b) the stalk maps $\phi_W, \psi_W : \mathcal{O}_{Y,Z} \to \mathcal{O}_{X,W}$ agree.*

*Then there exists an open subset $U \subset X$ for which $U \cap W \neq \varnothing$ and $\phi|_U = \psi|_U$.*

*Proof.* For any open subset $V' \subset Y$ meeting $Z$ (and therefore $\phi(W)$ (4.1.19)), then $\phi^{-1}(V')$ meets $W$. Similarly for $\psi$. So by irreducibility $\phi^{-1}(V') \cap \psi^{-1}(V')$ meets $W$ (4.1.60).

Let $V \subset Y$ be an affine open subset meeting $Z$ and choose an affine open subset $U \subset \phi^{-1}(V) \cap \psi^{-1}(V)$ meeting $W$.

From (4.3.19) we have a commutative cube for every open subset $V' \subset V$




© David Rufino CC BY-NC-ND 4.0

By uniqueness of the bottom square we conclude that $(\phi|_U)_{W\cap U} = (\psi|_U)_{W\cap U}$ and we may reduce to the case of $X$ and $Y$ affine.

By (6.2.25) it is sufficient to find $f \in \mathcal{O}_X(X)$ such that $\rho_f \circ \phi_Y^\sharp = \rho_f \circ \psi_Y^\sharp$. Suppose that $D(f) \cap W \neq \varnothing$ and consider the commutative diagram

$$\begin{array}{ccc} \mathcal{O}_Y(Y) & \overset{\phi^\sharp}{\underset{\psi^\sharp}{\longrightarrow}} & \mathcal{O}_X(X) \\ \downarrow & & \downarrow{\scriptstyle \rho_f} \\ & & \mathcal{O}_X(X)_f \\ & & \downarrow \\ \mathcal{O}_Y(Y)_{\mathfrak{m}_{Y,Z}} & \overset{\phi_W}{\underset{\psi_W}{\longrightarrow}} & \mathcal{O}_X(X)_{\mathfrak{m}_{X,W}} \end{array}$$

where we have used (6.2.21) to replace $\mathcal{O}_{Y,Z}$ with $\mathcal{O}_Y(Y)_{\mathfrak{m}_{Y,Z}}$ and $\mathcal{O}_{X,W}$ with $\mathcal{O}_X(X)_{\mathfrak{m}_{X,W}}$.

Let $y_1, ..., y_n$ be generators for $\mathcal{O}_Y$. Then by assumption there exists $t_i \notin \mathfrak{M}_{X,W}$ such that $t_i\phi^\sharp(y_i) = t_i\psi^\sharp(y_i)$. In particular $f = t_1 \cdots t_n$ works, since then $f\phi^\sharp(y_i) = f\psi^\sharp(y_i)$ for all $i$. ∎

**Proposition 6.2.30 (Lifting Stalk Maps II)**
*Let $X, Y$ be varieties, $W \subset X$, $Z \subset Y$ irreducible subsets and suppose there exists a local $k$-algebra homomorphism $\underline{\phi_W : \mathcal{O}_{Y,Z} \to \mathcal{O}_{X,W}}$. Then there exists an open subset $U \subset X$ and a regular morphism $\phi : U \to Y$ such that $\overline{Z} = \overline{\phi(W \cap U)}$ and the following diagram commutes for all open subsets $V \subset Y$ meeting $Z$ and $U' \subset \phi^{-1}(V)$ meeting $W$*

$$\begin{array}{ccc} \mathcal{O}_Y(V) & \xrightarrow{\phi^\sharp} & \mathcal{O}_X(U') \\ \downarrow & & \downarrow \\ \mathcal{O}_{Y,Z} & \xrightarrow[\phi_W]{} & \mathcal{O}_{X,W} \end{array}$$

*Proof.* Let $Y' \subset Y$ be an open affine subset meeting $Z$, and $X' \subset \phi^{-1}(V)$ an affine open subset meeting $W$.

Then $W \cap X'$ is an irreducible subset of $X'$ and similarly for $Z \cap Y'$ (...).

For every open subset $U \subset X'$ meeting $W$ we have a commutative diagram




Therefore we may assume without loss of generality that $X$ and $Y$ are affine, because we may reduce to finding a regular morphism $\phi : U \to Y'$.

We wish to find $f \in \mathcal{O}_X(X)$ and $\phi_Y^\sharp$ which completes the diagram

$$\begin{array}{ccc} \mathcal{O}_Y(Y) & \overset{\phi_Y^\sharp}{\dashrightarrow} & \mathcal{O}_X(X)_f \\ \downarrow & & \downarrow \\ \mathcal{O}_{Y,Z} & \xrightarrow{\phi_W^\sharp} & \mathcal{O}_{X,W} \\ \downarrow{\scriptstyle \sim} & & \downarrow{\scriptstyle \sim} \\ \mathcal{O}_Y(Y)_{\mathfrak{m}_{Y,Z}} & \xrightarrow{\phi_W^\sharp} & \mathcal{O}_X(X)_{\mathfrak{m}_{X,W}} \end{array} \qquad i_Z \text{ (left)}, \quad i_f \text{ (right)}$$

© David Rufino CC BY-NC-ND 4.0

For then a regular morphism $\phi : D(f) \to Y$ exists by (6.2.24). and by uniqueness (6.2.21).d) it has the required stalk map. As (6.2.19) $\phi^\sharp_W$ is local we may chase the diagram to show that $\left(\phi^\sharp\right)_Y^{-1}\left(\left(\mathfrak{M}_{X,W}\right)_f\right) = \mathfrak{M}_{Y,Z}$ and therefore $\overline{Z} = \overline{\phi(W \cap D(f))}$ as required.

By inspection it is sufficient to find $f$ such that $i_f$ is injective and $\mathrm{Im}\left(\phi^\sharp_W \circ i_Z\right) \subset \mathrm{Im}\left(i_f\right)$. We may find $g$ such that $i_g$ is injective by (3.6.37). Suppose $\mathcal{O}_Y(Y)$ is generated by $y_1, ..., y_m$ then $\phi^\sharp_W(y_i/1) = x_i/g_i$ for some $g_i \notin \mathfrak{M}_{X,W}$. Then $f = gg_1 \cdots g_m$ satisfies both criteria. As the stalk map is uniquely defined such that the diagram commutes we find that the regular morphism $\phi$ has stalk map precisely $\phi^\sharp_W$. ∎

**Corollary 6.2.31 (Lifting Stalk Maps I)**
*Let $X, Y$ be varieties, $x \in X$ and $\phi, \psi : X \to Y$ regular morphisms for which*

*a) $\overline{\{\phi(x)\}} = \overline{\{\psi(x)\}}$*
*b) $\phi_x = \psi_x$*

*Then there exists an open neighbourhood $U$ of $x$ such that $\phi|_U = \psi|_U$.*

**Corollary 6.2.32 (Lifting Stalk Maps II)**
*Let $X, Y$ be varieties and $\phi^\sharp_x : \mathcal{O}_{Y,y} \to \mathcal{O}_{X,x}$ a local $k$-algebra homomorphism. Then there exists an open subset $U \subset X$ and a regular morphism $\phi : U \to Y$ such that $y \in \overline{\{\phi(x)\}}$ and has stalk map precisely $\phi^\sharp_x$.*

## 6.2.5 Open Subvarieties

**Proposition 6.2.33 (Open Subset is Abstract Variety)**
*Let $X$ be a variety (resp. separated variety) and $U \subset X$ an open subset. Then $(U, \mathcal{O}_X|_U)$ is a variety (resp. separated variety), which we call an **open subvariety**. We may write $\mathcal{O}_U \coloneqq \mathcal{O}_X|_U$.*

*The inclusion $i : U \hookrightarrow X$ is an open immersion and an open immersion $\phi : X \to Y$ induces an isomorphism between $X$ and the open subvariety $\phi(X)$.*

*Proof.* By definition $X = \bigcup_{i=1}^n U_i$ where $U_i$ is affine. By (...) we have

$$U \cap U_i = \bigcup_{\alpha \in \mathcal{A}_i} D(f_\alpha)$$

where $f_\alpha \in \mathcal{O}_X(U_i)$. By (6.2.21) $\left(D(f_\alpha), \mathcal{O}_X|_{D(f_\alpha)}\right)$ is affine. Therefore we have an open cover of affine sets. By (6.2.7) $U$ is quasi-compact and therefore there exists a finite affine cover. This shows that $U$ is a variety.

Clearly the inclusion map $U \to X$ is injective and regular and so we see that $U$ inherits separatedness. ∎

**Proposition 6.2.34 (Quasi-Affine Varieties are Abstract Varieties)**
*Let $X$ be a quasi-affine variety. Then $(X, \mathcal{O}_X)$ is a separated variety. More precisely there exists a full and faithful functor*

$$\mathbf{QuasiAffVar}_k \longrightarrow \mathbf{Var}_k$$

*Proof.* This follows from (6.2.33) and (6.1.39) ∎

**Proposition 6.2.35 (Morphisms into Open Subvarieties)**
*Let $X, Y$ be varieties and $U \subset Y$ an open subvariety. Then a regular map $\phi : X \to U$ is regular when regarded as a map $\phi : X \to Y$.*

*Conversely a regular map $\phi : X \to Y$ such that $\phi(X) \subset U$ may be regarded as a regular map $\phi : X \to U$.*

**Proposition 6.2.36 (Morphisms from Open Subvarieties)**
*Let $\phi : X \to Y$ be a regular map of varieties and $U \subset X$ an open subvariety. Then $\phi|_U : U \to Y$ is a regular map.*

**Proposition 6.2.37 (Patching Varieties)**
*Suppose $X = \bigcup_{i=1}^n U_i$ is a finite union of sets, $\left(U_i, \mathcal{O}_{U_i}\right)$ is a variety, and suppose that for all $i, j = 1...n$*

- *$U_i \cap U_j$ is open in $U_j$, and*
- *the identity map $1 : \left(U_i \cap U_j, \mathcal{O}_{U_i}|_{U_i \cap U_j}\right) \to \left(U_j \cap U_i, \mathcal{O}_{U_j}|_{U_i \cap U_j}\right)$ is a regular morphism of spaces of functions.*

*Then there exists a unique structure of a variety on $X$ such that*

- *$U_i$ is open in $X$, and*
- *$\mathcal{O}_{U_i} = \mathcal{O}_X|_{U_i}$.*

© David Rufino CC BY-NC-ND 4.0

**Proposition 6.2.38 (Glueing Varieties)**
*Suppose $X = \bigcup_{i=1}^{n} U_i$ is a finite union of sets and there exists bijections $\theta_i : U_i \xrightarrow{\sim} W_i$, with $W_i$ varieties such that for all $i, j = 1...n$*

- *$\theta_i(U_i \cap U_j)$ is open in $W_i$, and*
- *$\theta_{ij} = \theta_j \circ \theta_i^{-1} : \theta_i(U_i \cap U_j) \to \theta_j(U_i \cap U_j)$ is regular.*

*Then there exists a unique structure of a variety on $X$ such that*

- *$U_i$ is open in $X$, and*
- *$\theta_i$ is a regular isomorphism.*

*Proof.* The topology exists by (4.3.26) and we may simply define

$$\mathcal{O}_X(U) := \left\{ f : U \to \overline{k} \;\middle|\; f \circ \theta_i^{-1}|_{\theta_i(U_i \cap U)} \text{ regular for all } i = 1...n \right\}$$

which gives $X$ the structure of a space of functions. As each $U_i$ is a finite union of affine varieties, so is $X$. ∎

## 6.2.6 Closed Subvarieties

**Proposition 6.2.39 (Affine Closed Subvarieties)**
*Let $(X, \mathcal{O}_X)$ be an affine variety with explicit isomorphism $\theta : X \cong V(\mathfrak{a})$, and $Z \subset X$ a closed subset. Then the structure $(Z, \mathcal{O}_Z)$ defined in (4.3.21) is also affine with explicit isomorphism $\theta|_Z : Z \cong V(\mathfrak{b})$ where $\mathfrak{b} := I(\theta(Z))$.*

*Proof.* This follows largely from (6.1.44). ∎

**Proposition 6.2.40 (Closed Subvariety)**
*Let $(X, \mathcal{O}_X)$ be a variety with $Z$ a closed subset. Then $(Z, \mathcal{O}_Z)$ with the structure defined in (4.3.21) is a variety.*

*Proof.* By assumption $X = \bigcup_{i=1}^{n} U_i$ with $\left(U_i, \mathcal{O}_{U_i}\right)$ affine. Evidently $Z = \bigcup_{i=1}^{n} (U_i \cap Z)$ is an open cover. Further $U_i \cap Z$ is a closed subset of $U_i$. By (6.2.39) $\left(U_i \cap Z, \mathcal{O}_{U_i}|_{U_i \cap Z}\right) \overset{(6.2.47)}{=} \left(U_i \cap Z, \mathcal{O}_Z|_{U_i \cap Z}\right)$ is affine. Therefore $Z$ is a variety. ∎

**Definition 6.2.41 (Closed Immersion)**
*Let $\phi : X \to Y$ be a regular morphism of varieties. Then it is a **closed immersion** if $\phi(X)$ is closed in $Y$ and $\phi$ determines an isomorphism between $X$ and the closed subvariety $\phi(X)$.*

**Proposition 6.2.42 (Closed Subvariety is a Variety)**
*Let $X$ be a variety and $Z \subset X$ a closed subset. Then the space with functions $(Z, \mathcal{O}_Z)$ is a variety and the map $j : (Z, \mathcal{O}_Z) \to (X, \mathcal{O}_X)$ is a closed immersion.*

*When $X$ is affine, then so is $Z$ and the map $j^\sharp : \mathcal{O}_X(X) \to \mathcal{O}_Z(Z)$ is surjective.*

*Proof.* By definition $X = \bigcup_{i=1}^{n} U_i$ where $U_i$ is affine. Then $Z \cap U_i$ is a closed subset of $U_i$ and by (6.2.47) $\mathcal{O}_Z|_{Z \cap U_i} = \mathcal{O}_{U_i}|_{Z \cap U_i}$. Therefore by (6.2.39) $Z \cap U_i$ is affine, and open in $Z$. This shows that $Z$ is a variety.

Suppose $f \in \mathcal{O}_X(U)$ then $f \circ j \in \mathcal{O}_Z(Z \cap U)$, which shows that $j$ is a regular morphism.

The final statement follows from (6.1.45). ∎

**Proposition 6.2.43**
*Let $\phi : X \to Y$ be a closed immersion and $Y$ an affine variety. Then $X$ is affine.*

**Proposition 6.2.44**
*Let $\phi : X \to Y$ be a regular map of varieties and $Z \subset X$ a closed subvariety. Then $\phi|_Z$ is regular.*

**Proposition 6.2.45**
*Let $\phi : X \to Y$ be a regular map of varieties and $Z \subset Y$ a closed (resp. open) subvariety. If $\phi(X) \subset Z$ then $\phi$ may be regarded as a regular map $X \to Z$.*

**Proposition 6.2.46**
*Let $\phi : X \to Y$ be a regular map of affine varieties. Then $\phi$ is a closed immersion if and only if the induced homomorphism $\phi_Y^\sharp : \mathcal{O}_Y(Y) \to \mathcal{O}_X(X)$ is surjective.*

*Proof.* Restatement of (6.1.45). ∎

© David Rufino CC BY-NC-ND 4.0

### 6.2.7 Locally Closed Subvariety

**Proposition 6.2.47 (Locally Closed Subvariety)**
*Let $X$ be a variety, $Z$ a closed subset and $U$ an open subset. Then $Z \cap U$ is a variety (viewed as a closed subvariety of $U$ or an open subvariety of $Z$).*

### 6.2.8 Rational Points

For abstract varieties we use a more local definition of rational point which coincides with the affine case (6.1.63).

**Definition 6.2.48 ($K$-Rational Points)**
*Let $X$ be a variety and $K/k$ an algebraic field extension. Define the $K$-rational points as follows*

$$X(K) \coloneqq \left\{(x, \iota) \;\middle|\; x \in X_0, \quad \iota : k\big(\mathfrak{m}_{X,x}\big) \hookrightarrow K\right\}$$

*where $X_0$ is the Kolmogorov Quotient (4.1.35) of $X$, and recalling that $\mathcal{O}_{X,x} = \mathcal{O}_{X,y}$ whenever $x, y$ become equal in $X_0$ (6.2.11).*

*By (6.2.19) specifying $\iota$ is completely equivalent to specifying a $k$-algebra homomorphism $\mathcal{O}_{X,x} \to K$.*

**Proposition 6.2.49**
*Let $\phi : X \to Y$ be a regular morphism of varieties and $K/k$ an algebraic field extension. Then the map*

$$\begin{aligned} \phi_K : X(K) &\longrightarrow Y(K) \\ (x, \iota) &\to \quad (\phi(x), \iota \circ \phi_x) \end{aligned}$$

*is well-defined and satisfies*

$$\psi_K \circ \phi_K = (\psi \circ \phi)_K$$

*In particular if $\phi$ is an isomorphism then $\phi_K$ is a bijection.*

**Proposition 6.2.50 (Points = Rational Points over $\overline{k}$)**
*Let $X$ be a variety. Then there is a canonical bijection*

$$\begin{aligned} X &\xrightarrow{\sim} X\big(\overline{k}\big) \\ x &\to \quad ([x], \mathrm{ev}_x) \end{aligned}$$

*which is functorial in $X$. More precisely if $\phi : X \to Y$ is a regular morphism then*

$$\mathrm{ev}_{\phi(x)} = \mathrm{ev}_x \circ \overline{\phi}_x$$

*Further*

$$y = \phi(x) \iff \phi(x) \in \overline{\{y\}} \text{ and } \mathrm{ev}_x \circ \overline{\phi}_x = \mathrm{ev}_y$$

*Proof.* Recall (4.1.34) that $X$ may be written as a disjoint union

$$X \coloneqq \bigsqcup_{x \in \mathcal{I}} \overline{\{x\}}$$

for some representative subset $\mathcal{I} \subset X$ which is in bijection with $X_0$. By (6.2.11) $X$ is symmetric and for all $z \in \overline{\{x\}}$ we have $[z] = [x]$ as elements of $X_0$. So for a given $x \in \mathcal{I}$ we need to show that the map

$$\begin{aligned} \overline{\{x\}} &\to \mathrm{AlgHom}_k\big(k\big(\mathfrak{m}_{X,x}\big), \overline{k}\big) \\ z &\to \mathrm{ev}_z \end{aligned}$$

is bijective. It is well-defined because $z \in \overline{\{x\}}$ implies $\mathcal{O}_{X,x} = \mathcal{O}_{X,z}$. Let $U$ be an affine open neighbourhood of $x$ then by (4.1.43) $\overline{\{x\}} = \mathrm{cl}_U(\{x\})$. Further by (4.3.15) $U \hookrightarrow X$ induces a local isomorphism on stalks and therefore residue fields $k\big(\mathfrak{m}_{X,x}\big) \cong k\big(\mathfrak{m}_{U,x}\big)$. By functoriality (4.3.10) we may then reduce to the case of $X$ affine. The result then follows from the affine case (6.1.63).

There is a commutative diagram

© David Rufino CC BY-NC-ND 4.0

$$\begin{array}{ccc} X & \xrightarrow{\sim} & X\left(\overline{k}\right) \\ \downarrow{\scriptstyle \phi} & & \downarrow{\scriptstyle \phi_{\overline{k}}} \\ Y & \xrightarrow{\sim} & Y\left(\overline{k}\right) \end{array}$$

from which the final statement follows easily. ∎

**Proposition 6.2.51**
*Let $X$ be a variety. Then the evaluation map*

$$\begin{aligned} \mathrm{ev}_X : X &\longrightarrow \mathrm{AlgHom}_k\left(\mathcal{O}_X(X), \overline{k}\right) \\ x &\to \quad f \to f(x) \end{aligned}$$

*is natural in $X$.*

*When $X$ is affine it is also a bijection.*

*Proof.* More precisely when $\phi : X \to Y$ is a regular morphism of varieties we require to show that the following diagram commutes

$$\begin{array}{ccc} X & \xrightarrow{\mathrm{ev}} & \mathrm{AlgHom}_k\left(\mathcal{O}_X(X), \overline{k}\right) \\ \downarrow{\scriptstyle \phi} & & \downarrow{\scriptstyle - \circ \phi^\sharp} \\ Y & \xrightarrow{\mathrm{ev}} & \mathrm{AlgHom}_k\left(\mathcal{O}_Y(Y), \overline{k}\right) \end{array}$$

Suppose $g \in \mathcal{O}_Y(Y)$ and $x \in X$ then this amounts to

$$g(\phi(x)) = \phi^\sharp(g)(x)$$

which holds by definition.

Let $\theta : X \cong V(\mathfrak{a})$ be an isomorphism of varieties. Then $\phi^\sharp$ is an isomorphism of $k$-algebras, and so $- \circ \phi^\sharp$ is a bijection (since functors preserve isomorphisms). The evaluation map for $V(\mathfrak{a})$ is a bijection (6.1.63) and so we deduce it is also for $X$. ∎

**Corollary 6.2.52**
*Let $X$ be a variety and $U \subset X$ an affine open subset. Then the evaluation map*

$$\mathrm{ev}_U : U \longrightarrow \mathrm{AlgHom}_k\left(\mathcal{O}_X(U), \overline{k}\right)$$

*is bijective.*

*Suppose $\phi : X \to Y$ is a regular morphism, $U$ is an affine neighbourhood of $x$ and $V$ is an affine neighbourhood of $y$ such that $U \subset \phi^{-1}(V)$, then*

$$\phi(x) = y \iff \mathrm{ev}_U(x) \circ \rho_{\phi^{-1}(V)U} \circ \phi^\sharp = \mathrm{ev}_V(y)$$

## 6.2.9 Dimension

**Definition 6.2.53 (Dimension of a Variety)**
*Let $X$ be a variety. Then we define the **dimension** to be the Krull Dimension of $X$ as a topological space.*

**Proposition 6.2.54**
*Let $X$ be an integral variety. Then it is biequidimensional and*

$$\dim X = \mathrm{trdeg}_k(k(X))$$

*The dimension equals $\dim U$ for every open subset.*

*Proof.* Let $U$ be an affine open subset then by (6.1.52) $\dim U = \mathrm{trdeg}_k(k(U))$ which equals $\mathrm{trdeg}_k(k(X))$ by (4.3.16). Further by (6.1.51) $U$ is biequidimensional and so by (4.1.73) we conclude that $X$ is biequidimensional with the same dimension as $U$.

Every open subset can be covered by affines so we are done by (4.1.88). ∎

© David Rufino CC BY-NC-ND 4.0

**Corollary 6.2.55**
*Let $X$ be a variety. Then it is quasi-biequidimensional. Moreover every closed subset $W \subset X$ satisfies the codimension formula*

$$\dim X = \dim W + \operatorname{codim}(W, X)$$

*If $W$ is irreducible then we also have the identity*

$$\dim \mathcal{O}_{X,W} = \operatorname{codim}(W, X) = \dim X - \dim W$$

*Proof.* Every irreducible component is an integral closed subvariety and therefore biequidimensional (6.2.54). So by definition $X$ is quasi-biequidimensional. The codimension formula is (4.1.87).

Let $U$ be an affine open subset of $X$ meeting $W$. Then there is an isomorphism $\mathcal{O}_{X,W} \xrightarrow{\sim} \mathcal{O}_{U,W\cap U}$ (4.3.17). Further by (4.1.73) we have $\operatorname{codim}(W, X) = \operatorname{codim}(W \cap U, U)$. Therefore we may deduce from the affine case (6.1.57) that

$$\dim \mathcal{O}_{X,W} = \dim \mathcal{O}_{U,W\cap U} = \operatorname{codim}(W \cap U, U) = \operatorname{codim}(W, X)$$

as required. ∎

**Proposition 6.2.56 (Irreducible Closed Subsets of Dimension 0)**
*Let $X$ be a variety and $Z \subset X$ an irreducible closed subset. Then the following are equivalent*

*a)* $\dim Z = 0$
*b)* $Z = \overline{\{z\}}$ *for some* $z \in Z$
*c)* $Z$ *is finite*

*In particular for every $x \in X$ we have $\dim \mathcal{O}_{X,x} = \dim X$.*

*Proof.* Follows from (6.2.12). The dimension formula follows from (6.2.55). ∎

**Proposition 6.2.57**
*Let $X$ be a variety. Then there is a bijective correspondence*

$$X_0 \longleftrightarrow \{W \subset Z \mid W \text{ closed and irreducible and } \dim W = 0\}$$

*Proof.* By (6.2.11) and (4.1.94). ∎

## 6.2.10 Galois Action

The notion of rational point allows us to define a Galois Action which coincides with concrete notions in both the affine and projective cases.

**Definition 6.2.58 (Abstract Galois Action)**
*Let $X$ be an abstract variety and $K/k$ an algebraic field extension. Then there is a well-defined group action*

$$\operatorname{Aut}(K/k) \times X(K) \longrightarrow X(K)$$

*given by*

$$\sigma, (x, \iota) \longrightarrow (x, \sigma \circ \iota)$$

*When $K = \overline{k}$ then this naturally gives a well-defined group action*

$$\operatorname{Aut}\left(\overline{k}/k\right) \times X \longrightarrow X$$

*in light of the bijection $X \xrightarrow{\sim} X\left(\overline{k}\right)$ (6.2.50).*

**Proposition 6.2.59 (Topological Properties of Galois Action)**
*Let $X$ be an abstract variety. For every $x \in X$ and $\sigma \in \operatorname{Aut}\left(\overline{k}/k\right)$ we have $x$ and $\sigma(x)$ are topologically indistinguishable. Furthermore the orbit of $x$ is simply $\overline{\{x\}}$, which has order $\left[k\left(\mathfrak{m}_{X,x}\right) : k\right]$.*

*In particular $\sigma$ preserves every open set and is therefore a homeomorphism of $X$. Finally every $\sigma$ restricts to a well-defined action on $X_0$ which is simply the identity.*

*For every affine open subset $U \subset X$ with explicit isomorphism $U \xrightarrow{\sim} V(\mathfrak{a})$. Then the action of $\operatorname{Aut}\left(\overline{k}/k\right)$ coincides with the usual one under this isomorphism.*

© David Rufino CC BY-NC-ND 4.0

### 6.2.11 Product Variety

We construct the product in the category of varieties, but note that the topology of the product does *not* coincide with the usual product topology.

We first show that the product, if it exists, is unique in concrete sense.

**Lemma 6.2.60 (Uniqueness of Product)**
*Let $X, Y$ be varieties (resp. affine varieties), then there is at most one topology on $X \times Y$ and local space of functions $\mathcal{O}_{X\times Y}$ for which the projection maps are regular and the following map*

$$\begin{aligned} \mathrm{Mor}(Z, X \times Y) &\xrightarrow{\sim} \mathrm{Mor}(Z, X) \times \mathrm{Mor}(Z, Y) \\ \phi &\to \ (\pi_X \circ \phi, \pi_Y \circ \phi) \end{aligned}$$

*is a well-defined bijection for all varieties (resp. affine varieties) $Z$, natural in $Z$.*

*Proof.* Suppose we have two such structures, denote them by $X \times Y$ and $\widetilde{X \times Y}$, then we have a natural bijection

$$\begin{aligned} \mathrm{Mor}(Z, X \times Y) &\to \mathrm{Mor}\big(Z, \widetilde{X \times Y}\big) \\ \phi &\to \phi \end{aligned}$$

which is the identity on the underlying set function because $\phi = \phi' \iff \pi_X \circ \phi = \pi_X \circ \phi'$ and $\pi_Y \circ \phi = \pi_Y \circ \phi'$. By the dual version of the Yoneda Lemma (2.6.58) the identity map $1 : X \times Y \to \widetilde{X \times Y}$ is a regular isomorphism. Therefore the topology is the same and so are the sheaf of functions. ∎

**Proposition 6.2.61**
*Let $X, Y$ be varieties, then the categorical product $X \times Y$ exists, and coincides with the set-theoretic product. More precisely*

*a) there exists a unique topology on $X \times Y$ and sheaf of functions $\mathcal{O}_{X\times Y}$ such that the projection maps $\pi_X : X \times Y \to X$ and $\pi_Y : X \times Y \to Y$ are regular and there is a natural bijection for all varieties $Z$*

$$\begin{aligned} \mathrm{Mor}(Z, X \times Y) &\xrightarrow{\sim} \mathrm{Mor}(Z, X) \times \mathrm{Mor}(Z, Y) \\ \phi &\to \ (\pi_X \circ \phi, \pi_Y \circ \phi) \end{aligned}$$

*b) If $X$ and $Y$ are affine (resp. separated), then so is $X \times Y$, and there is an isomorphism*

$$\big(\mathcal{O}_X(X) \otimes_k \mathcal{O}_Y(Y)\big)_{\mathrm{red}} \xrightarrow{\sim} \mathcal{O}_{X\times Y}(X \times Y)$$

*c) If $U \subset X$ and $V \subset Y$ are open then so is $U \times V$, and $\mathcal{O}_{U\times V} = \mathcal{O}_{X\times Y}|_{U\times V}$*

*d) If $T \subset X$ and $S \subset Y$ are closed then $T \times S$ is a closed subset of $X \times Y$*

*Proof.* Suppose $X, Y$ are both affine. Then $X \times Y$ is given the structure of an affine variety in (6.1.96). This means the following map

$$\begin{aligned} \mathrm{Mor}(Z, X \times Y) &\to \mathrm{Mor}(Z, X) \times \mathrm{Mor}(Z, Y) \\ \phi &\to (\pi_X \circ \phi, \pi_Y \circ \phi) \end{aligned}$$

is bijective when $Z$ is affine. Consider the case $Z = \bigcup_{i=1}^n U_i$ is a variety. Injectivity is obvious, for surjectivity suppose $f : Z \to X$ and $g : Z \to Y$ are regular. Then $f|_{U_i}$ and $g|_{U_i}$ are regular, and by the affine case so is $f|_{U_i} \times g|_{U_i}$. By (4.3.27) $f \times g$ is also regular.

Consider the general case $X = \bigcup_{i=1}^n U_i$ and $Y = \bigcup_{j=1}^m V_j$ are varieties with given affine coverings. Define $W_{ij} \coloneqq U_i \times V_j$, then by the affine case $W_{ij}$ may be given the structure of an (affine) variety with regular set-theoretic projections $\pi_X^{ij} : W_{ij} \to U_i$ and $\pi_Y^{ij} : W_{ij} \to V_j$ such that the following map is a bijection

$$\mathrm{Mor}\big(Z, W_{ij}\big) \xrightarrow{\sim} \mathrm{Mor}(Z, U_i) \times \mathrm{Mor}\big(Z, V_j\big)$$

for all varieties $Z$. Then consider $Z = W_{ij} \cap W_{kl} = \big(\pi_X^{kl}\big)^{-1}(U_i \cap U_k) \cap \big(\pi_Y^{kl}\big)^{-1}\big(V_j \cap V_k\big)$ regarded as an open subvariety of $W_{kl}$. With this structure the projection maps $\pi_X : Z \to U_i$ and $\pi_Y : Z \to V_j$ may be shown to be regular by

- changing the codomain of $\pi_X^{kl} : W_{kl} \to U_k$ (resp. $\pi_Y$) to $X$, (6.2.35)
- restricting the domain to $Z$ (6.2.36),
- changing the codomain to $U_i$ (resp. $V_j$) (6.2.35)

Consequently the identity map $W_{ij} \cap W_{kl} \to W_{ij}$ is regular and therefore so is $W_{ij} \cap W_{kl} \to W_{ij} \cap W_{kl}$ (6.2.35) when considered as open subvarieties of $W_{kl}$ and $W_{ij}$ respectively. By symmetry this is a regular isomorphism. Therefore

© David Rufino CC BY-NC-ND 4.0

the $W_{ij}$ satisfy the criteria of (6.2.37) and there exists a unique structure of a variety on $X \times Y$ for which $\mathcal{O}_{X\times Y}|_{W_{ij}} = \mathcal{O}_{W_{ij}}$. By (...) the set-theoretic projections $\pi_X : X \times Y \to X$ and $\pi_Y : X \times Y \to Y$ are regular. As before we need to show the following map is bijective

$$\mathrm{Mor}(Z, X \times Y) \to \mathrm{Mor}(Z, X) \times \mathrm{Mor}(Z, Y)$$
$$\phi \to (\pi_X \circ \phi, \pi_Y \circ \phi)$$

with injectivity obvious. Suppose $f : Z \to X$ and $g : Z \to Y$ are regular. Define $T_{ij} \coloneqq f^{-1}(U_i) \cap g^{-1}(V_j)$. Then by (6.2.36) $f_{ij} \coloneqq f|_{T_{ij}}$ and $g_{ij} \coloneqq g|_{T_{ij}}$ are regular. Therefore by the universal property for $W_{ij}$ the maps $f_{ij} \times g_{ij}$ are regular. Clearly they are compatible so these glue to give a regular map $f \times g : Z \to X \times Y$ (6.2.35) (4.3.27) as required. Further $X \times Y = \bigcup_{i,j} W_{ij}$ so it is indeed a variety. Uniqueness of the structure follows from (6.2.60).

For *b*) we already demonstrated that the affine product is also a product in the category of varieties, so it follows from uniqueness. For separatedness this is an easy exercise.

For *c*) we have $U \times V = \pi_X^{-1}(U) \cap \pi_Y^{-1}(V)$ is open, and the universal property naturally restricts to the subset $U \times V$. The identity then follows from uniqueness (6.2.60).

For *d*) we have $T \times S = \pi_X^{-1}(T) \cap \pi_Y^{-1}(S)$ is closed. ∎

**Proposition 6.2.62**
*Let $X$ be an integral variety and $Y$ a geometrically integral affine variety (see (6.1.84)). Then $X \times Y$ is an integral variety.*

*Proof.* By definition $X = \bigcup_{i=1}^n U_i$ for $U_i$ affine. By (6.2.61) $X \times Y = \bigcup_{i=1}^n U_i \times Y$ and $U_i \times Y$ is affine. By (6.1.98) $U_i \times Y$ is integral. By (4.1.59) $U_i \cap U_j \neq \varnothing$ whence $(U_i \times Y) \cap (U_j \times Y) \neq \varnothing$. Therefore by (4.1.74) we conclude that $X \times Y$ is integral. ∎

## 6.2.12 Separated Varieties

**Proposition 6.2.63**
*Let $\phi_1, \phi_2 : X \rightrightarrows Y$ be regular morphisms of varieties with $Y$ separated. Then the equaliser*

$$\{x \in X \mid \phi_1(x) = \phi_2(x)\}$$

*is a closed subset of $X$.*

*Proof.* Let $X = \bigcup_{i=1}^n U_i$ where $U_i$ are affine. Then the equaliser is the finite union of

$$\left\{x \in U_i \;\middle|\; \phi_1|_{U_i}(x) = \phi_2|_{U_i}(x)\right\}$$

which is by definition closed. ∎

The following shows that the separated property is analogous to the Hausdorff property in the classical case (4.1.98).

**Proposition 6.2.64**
*Let $X$ be a variety. Then $X$ is separated if and only if the diagonal*

$$\Delta_X \coloneqq \{(x, x) \mid x \in X\}$$

*is a closed subvariety of $X \times X$.*

*Proof.* Suppose that $\Delta_X$ is closed. Then the equaliser of $\phi_1, \phi_2 : Z \rightrightarrows X$ is equal to $(\phi_1 \times \phi_2)^{-1}(\Delta_X)$. By (6.2.61) $\phi_1 \times \phi_2$ is regular and in particular continuous. We conclude the equaliser is closed.

Conversely $\Delta_X$ is the equaliser of $\pi_1, \pi_2 : X \times X \rightrightarrows X$ and so by (6.2.14) is closed. ∎

**Proposition 6.2.65**
*Let $X$ be a separated variety and $Z \subset X$ a closed (resp. open) subvariety. Then $Z$ is separated.*

*Proof.* Then $\Delta_Z = \Delta_X \cap (Z \times Z)$. By (6.2.61) $Z \times Z$ is a closed (resp. open) subvariety of $X \times X$. Then the result follows by (6.2.64). ∎

**Lemma 6.2.66**
*Let $\phi : X \to Y$ be a regular map of varieties, with $Y$ separated. Then the graph*

$$\Gamma_\phi = \{(x, \phi(x)) \mid x \in X\}$$

*is a closed subvariety of $X \times Y$. Further there is a canonical isomorphism of varieties*

© David Rufino CC BY-NC-ND 4.0

$$X \xrightarrow{\sim} \Gamma_\phi$$
$$x \to (x, \phi(x))$$

*Proof.* Consider the regular map

$$(\phi \circ \pi_X) \times \pi_Y : X \times Y \to Y \times Y$$
$$(x, y) \to (\phi(x), y)$$

Then

$$\Gamma_\phi = ((\phi \circ \pi_X) \times \pi_Y)^{-1}(\Delta_Y)$$

which is closed. The map $X \to \Gamma_\phi$ is evidentally regular. Further $\Gamma_\phi \hookrightarrow X \times Y \to X$ is regular by (6.2.42) and (4.3.2). These are mutual inverses and so the last statement follows immediately. ∎

**Proposition 6.2.67 (Intersection of Affines is Affine)**
*Let $X$ be a separated variety and $U, V$ open subsets. Then there is a regular isomorphism followed by a closed immersion*

$$U \cap V \xrightarrow{\sim} (U \times V) \cap \Delta_X \hookrightarrow U \times V$$
$$x \to \quad (x, x)$$

*In particular when $U, V$ are affine then so is $U \cap V$.*

*Proof.* By (6.2.61) $U \times V$ is an open subvariety of $X \times X$.

$(U \times V) \cap \Delta_X = ((U \cap V) \times V) \cap \Delta_X$ is the graph of the inclusion map $U \cap V \hookrightarrow V$. By (6.2.66) it is a closed subvariety of $(U \cap V) \times V$ and therefore of $U \times V$. The same result shows that the first map is a regular isomorphism.

When $U, V$ are affine, so is $U \times V$. By (6.2.42) so is $(U \times V) \cap \Delta_X$, and therefore $U \cap V$. ∎

**Proposition 6.2.68 (Criteria for Separated Variety)**
*Let $X$ be a variety with an open cover $\bigcup_{i=1}^n U_i$. Then $X$ is separated if and only if $(U_i \times U_j) \cap \Delta_X$ is a closed subvariety of $U_i \times U_j$ for all $i, j = 1...n$.*

*Proof.* Suppose $X$ is separated then $(U_i \times U_j) \cap \Delta_X$ is a closed subvariety by (6.2.67).

Conversely if $(U_i \times U_j) \cap \Delta_X$ is closed for all $i, j$ then by (4.1.22)

$$\Delta_X = \bigcup_{i,j} (U_i \times U_j) \cap \Delta_X$$

is closed, and therefore $X$ is separated by (6.2.64). ∎

**Proposition 6.2.69 (Criteria for Separated Variety II)**
*Let $X$ be a variety with an affine open cover $\bigcup_{i=1}^n U_i$. Then $X$ is separated if and only if $U_i \cap U_j$ is affine and the $k$-algebra homomorphism*

$$\mathcal{O}_X(U_i) \otimes_k \mathcal{O}_X(U_j) \longrightarrow \mathcal{O}_X(U_i \cap U_j)$$
$$f \otimes g \longrightarrow f|_{U_i \cap U_j} \cdot g|_{U_i \cap U_j}$$

*is surjective for all $i, j = 1...n$*

*Proof.* Suppose $X$ is separated, then by (6.2.67) $U_i \cap U_j$ is affine and there is a closed immersion

$$U_i \cap U_j \xrightarrow{\sim} (U_i \times U_j) \cap \Delta_X \hookrightarrow U_i \times U_j$$

By (6.2.46) the corresponding map of global sections

$$\mathcal{O}_{U_i \times U_j}(U_i \times U_j) \to \mathcal{O}_{U_i \cap U_j}(U_i \cap U_j) \quad (*)$$
$$f \to u \to f(u, u)$$

is surjective. Recall from (6.2.61) there is a surjective $k$-algebra homomorphism

$$\mathcal{O}_{U_i}(U_i) \otimes_k \mathcal{O}_{U_i}(U_j) \to \mathcal{O}_{U_i \times U_j}(U_i \times U_j)$$
$$f \otimes g \to u \to f(u) \cdot g(u)$$

© David Rufino CC BY-NC-ND 4.0

which composes to give the required map.

Conversely suppose that $U_i \cap U_j$ are affine and the given map is surjective then evidentally the map in $(*)$ is surjective. By (6.2.46) this shows that the map $U_i \cap U_j \hookrightarrow U_i \times U_j$ is a closed immersion and in particular $\big(U_i \times U_j\big) \cap \Delta_X$ is a closed subvariety of $\big(U_i \times U_j\big)$. Then we are done by (6.2.68). ∎

## 6.2.13 Rational Maps

**Definition 6.2.70 (Rational Maps)**
*Let $X, Y$ be varieties. A **rational map** $\phi : X \longrightarrow Y$ is an equivalence class of regular morphisms*

$$\phi : U \to Y$$

*where $U$ is a dense open subset of $X$. We say $(U, \phi) \sim (U', \phi')$ if there exists a dense open subset $W \subset U \cap U'$ such that $\phi|_W = \phi'|_W$.*

*We say the **domain of definition** of a rational map $\phi$ is the set of points $x \in X$ for which there exists a neighbourhood $U$ of $x$ and a representative $(U, \phi)$. This is a dense open subset denoted by $\mathrm{dom}(\phi)$.*

**Proposition 6.2.71**
*Let $\phi : X \longrightarrow Y$ be a rational map with $Y$ separated. Then there exists a representative regular morphism $\phi : \mathrm{dom}(\phi) \to Y$.*

*Proof.* It is enough to show that for any two representatives $(U, \phi)$ and $(U', \phi')$ we have $\phi|_{U \cap U'} = \phi'|_{U \cap U'}$. By (6.2.63) the equaliser is closed in $U \cap U'$ and by definition contains a dense open subset $W$. Therefore it equals $U \cap U'$ as required. ∎

**Proposition 6.2.72 (Rational Functions are Rational Maps)**
*Let $X$ be an integral variety. Then there is a bijection*

$$k(X) \xrightarrow{\sim} \{f : X \longrightarrow \mathbb{A}^1_k\}$$

*and every such map is either dominant or has finite image.*

*Proof.* By (6.2.4) this is well-defined and bijective. Suppose $U \subset X$ is dense then by (4.1.70) it is irreducible. Therefore a regular map $f : U \to \mathbb{A}^1_k\big(\overline{k}\big)$ has irreducible image (4.1.64) and $\overline{f(X)}$ is irreducible (4.1.65) and closed. Clearly $\dim \overline{f(X)} \leq \dim \mathbb{A}^1_k\big(\overline{k}\big) = 1$. If equality holds then by (4.1.83).f) $\overline{f(X)} = \mathbb{A}^1_k$ and by definition $f$ is dominant.

Otherwise $\dim \overline{f(X)} = 0$ whence by (6.2.56) $\overline{f(X)} = \overline{\{y\}}$ for some $y \in \mathbb{A}^1_k\big(\overline{k}\big)$. ∎

**Proposition 6.2.73**
*Let $X$ be an integral variety. Then as subrings of $k(X)$ (...) we have*

$$\mathcal{O}_X(U) = \{f \in k(X) \mid U \subset \mathrm{dom}(f)\}$$

*and*

$$\mathcal{O}_{X,x} = \{f \in k(X) \mid x \in \mathrm{dom}(f)\}$$

*Furthermore*

$$\mathcal{O}_X(U) = \bigcap_{x \in U} \mathcal{O}_{X,x}$$

*Proof.* Recall (...) that $\mathbb{A}^1_k$ being affine is separated. Therefore every rational map $f \in k(X)$ has a regular representative in $\tilde{f} \in \mathcal{O}_X(\mathrm{dom}(f))$ by (6.2.71). So if $U \subset \mathrm{dom}(f)$ then $\tilde{f} \sim \tilde{f}|_U$ and so $f \in \mathcal{O}_X(U)$. Conversely if $\tilde{f} \in \mathcal{O}_X(U)$ then $\big(U, \tilde{f}\big)$ is a rational map whose domain contains $U$.

The second statement is similar, and then the final statement follows trivially. ∎

**Proposition 6.2.74 (Dominant Rational Map)**
*Let $X$ be an integral variety and $[(U, \phi)] : X \longrightarrow Y$ a rational map. Then the following are equivalent*

*a) $\phi(U)$ is dense in $Y$*
*b) $(U', \phi') \sim (U, \phi) \Longrightarrow \phi'(U')$ is dense in $Y$*

*In this case we say that $[(U, \phi)]$ is a **dominant rational map***

*Proof.* Clearly $b) \Longrightarrow a)$. Conversely suppose $\varnothing \neq W \subset U$ is open, then it is dense by (4.1.59) so $\mathtt{cl}_U(W) = U$ by (4.1.21). Therefore by (4.1.30)

© David Rufino CC BY-NC-ND 4.0

$$\phi(U) = \phi(\mathtt{cl}_U(W)) \subset \overline{\phi(W)}$$

which means

$$X = \overline{\phi(U)} \subset \overline{\phi(W)}$$

and therefore $\phi(W)$ is dense. In particular by definition we have $W \subset U \cap U'$ such that $\phi(W) = \phi'(W)$. So we conclude that $\phi'(W)$ is dense and so a-fortiori is $\phi'(U')$. ∎

**Proposition 6.2.75**
*Let $X, Y, Z$ be integral varieties and $\phi : X \longrightarrow Y$, $\psi : Y \longrightarrow Z$ rational maps with $\phi$ dominant. For every equivalence class $(U, \phi)$ and $(V, \psi)$ we have $U \cap \phi^{-1}(V) \neq \varnothing$ and the map*

$$\psi \circ \phi : U \cap \phi^{-1}(V) \to Z$$

*is a regular morphism determining a rational map*

$$\psi \circ \phi : X \longrightarrow Z$$

*which is dominant provided $\psi$ is.*

*Further this is independent of the choice of equivalence class.*

*Proof.* By definition $\phi(U)$ is dense which means that $\phi(U) \cap V \neq \varnothing \Longrightarrow U \cap \phi^{-1}(V) \neq \varnothing$, which is therefore also dense in $X$ (4.1.59). ∎

**Proposition 6.2.76**
*Let $X, Y$ be integral varieties. Then there is a bijective map*

$$\begin{aligned} \{\phi : X \longrightarrow Y \text{ dominant rational maps}\} &\xrightarrow{\sim} \mathrm{AlgHom}_k(k(Y), k(X)) \\ \phi &\to \ \phi_\star \end{aligned}$$

*Further $\phi_\star$ is injective and the existence of $\phi$ implies that $\dim Y \leq \dim X$. Then following are equivalent*

*a)* $\dim Y = \dim X$
*b)* $k(X)/\phi_\star(k(Y))$ *is algebraic*
*c)* $k(X)/\phi_\star(k(Y))$ *is finite*

*In this case we denote the **degree** by* $\deg(\phi) := [k(X) : \phi_\star(k(Y))]$.

*Proof.* The map is well-defined by (6.2.72) and (6.2.75), after we verify that the image is a $k$-algebra homomorphism.

Further it is injective by (6.2.29) and surjective by (6.2.30).

As $k(Y)$ is a field $\phi_\star$ is injective. We claim that $\mathrm{trdeg}_k(k(Y)) \leq \mathrm{trdeg}_k(k(X))$. For a transcendence base for $k(Y)$ is a-fortiori algebraicly independent as a subset of $k(X)$. Therefore the inequality follows from (3.19.140) and (6.2.54).

*a)* $\Longrightarrow$ *b)* Suppose $\dim X = \dim Y$ and let $\mathcal{B}$ be a transcendence base for $Y$, then by (3.19.141) $\phi_\star(\mathcal{B})$ is a transcendence base for $X$. Therefore $k(X)/\phi_\star(k(\mathcal{B}))$ is algebraic

*b)* $\Longleftrightarrow$ *c)* A-fortiori $k(X)/\phi_\star(k(Y))$ is finitely generated. Therefore the equivalence follows from (3.19.44).

*b)* $\Longrightarrow$ *a)* We have a-fortiori $k(X)/\phi_\star(k(\mathcal{B}))$ is algebraic whence by (3.19.140) $\dim Y \geq \dim X$, and so $\dim Y = \dim X$. ∎

## 6.2.14 Tangent Space

**Definition 6.2.77**
*Let $X$ be a variety and $W \subset X$ an irreducible subset. Define the **tangent space***

$$T_W X := \mathrm{Der}\big(\mathcal{O}_{X,W}, k(\mathfrak{m}_{X,W})\big)$$

*and the **cotangent space***

$$T^\star_W X := \mathrm{Hom}\big(\mathfrak{m}_{X,W}/\mathfrak{m}^2_{X,W}, k(\mathfrak{m}_{X,W})\big)$$

**Definition 6.2.78 (Non-Singular Point)**
*Let $X$ be a variety and $W \subset X$ an irreducible subset. Then we say that $X$ is **non-singular** at $W$ if*

*a) $W$ is contained in a unique irreducible component $X_\alpha$ (equivalently $\mathcal{O}_{X,W}$ is an integral domain), and*
*b)* $\dim T_W X = \dim \mathrm{Der}(k(X_\alpha))$

© David Rufino CC BY-NC-ND 4.0

*We say that $X$ is non-singular if it is integral and non-singular at all points.*

**Definition 6.2.79 (Regular Point)**
*Let $X$ be a variety and $W \subset X$ an irreducible subset.*

*We say $X$ is **regular** at $W$ if*
*a) $W$ is contained in a unique irreducible component $X_\alpha$, and*
*b) $\dim \mathcal{O}_{X,W} = \dim T_x^\star X$*

*We say that $X$ is regular if it is integral and regular at all points.*

**Proposition 6.2.80 (Regularity / Non-Singularity is a local property)**
*Let $X$ be a variety, $W \subset X$ an irreducible subset and $U \subset X$ an open subset such that $U \cap W \neq \varnothing$. Then there is a canonical isomorphism*

$$T_W X \xrightarrow{\sim} T_{W \cap U} U.$$

*$X$ is regular (resp. non-singular) at $W$ if and only if $U$ is regular (resp. non-singular) at $U \cap W$.*

**Proposition 6.2.81**
*Let $X$ be an integral variety. Then it is regular (resp. non-singular) if it has a regular (resp. non-singular) open cover.*

**Proposition 6.2.82**
*Let $X$ be a variety and $W \subset X$ an irreducible subset such that $k\big(\mathfrak{m}_{X,W}\big)/k$ is **separably generated** (e.g. $k$ perfect). Then there is a canonical isomorphism*

$$T_W X \xrightarrow{\sim} T_W^\star W$$

*In particular when $W$ is contained in a unique irreducible component we have **regular** $\iff$ **non-singular**.*

## 6.2.15 Completeness

Completeness is analogous to compactness in the Hausdorff case, and many results remain true when complete is replaced with compact and the standard product topology is used (in place of the topology induced by the product variety structure). We show in the next section that every projective variety is complete, so it is quite a natural condition.

**Definition 6.2.83**
*Let $X$ be a separated variety. Then it is*

- ***universally closed** if for every variety $Y$ the projection map*

$$\pi_Y : X \times Y \to Y$$

  *is closed (i.e. the image of a closed subset is closed).*
- ***complete** if for every irreducible closed subvariety $Z \subset X$ and valuation ring $k \subset R \subset k(Z)$ there is a point $z \in Z$ such that $\mathcal{O}_{Z,z} \subset R$*

**Proposition 6.2.84 (Image of Complete Variety is Closed)**
*Let $\phi : X \to Y$ be a morphism of varieties where $X$ is universally closed and $Y$ is separated. Then $\phi(X)$ is a closed subvariety of $Y$.*

*Proof.* By (6.2.66) the graph $\Gamma_\phi$ is a closed subset of $X \times Y$. Then $\phi(X)$ is the projection of $\Gamma_\phi$ and therefore closed by definition. ∎

**Proposition 6.2.85 (Closed Subvariety of Complete Variety is also Complete)**
*A closed subvariety of a universally closed (resp. complete) variety is universally closed (resp. complete)*

*Proof.* Suppose $Z \subset X$ is closed then by (6.2.61) $Z \times Y$ is a closed subset of $X \times Y$. Further $T$ closed in $Z \times Y$ implies $T$ closed in $X \times Y$ and the result follows.

The complete case is obvious. ∎

**Proposition 6.2.86**
*A separated variety $X$ is universally closed (resp. complete) iff all its irreducible components are universally closed (resp. complete).*

*Proof.* By (4.1.66) every irreducible component is closed, and therefore complete by (6.2.85).

© David Rufino CC BY-NC-ND 4.0

Conversely suppose $X = X_1 \cup ... \cup X_n$ is the irreducible decomposition (6.2.8). If $T \subset X \times Y$ is closed then $T \cap (X_i \times Y)$ is closed in $X_i \times Y$ and therefore by assumption $\pi_Y(T \cap (X_i \times Y)) = \pi_Y(T) \cap X_i$ is closed. Therefore $\pi_Y(T) = \bigcup_{i=1}^n \pi_Y(T) \cap X_i$ is also closed.

The complete case is straightforward. ∎

**Proposition 6.2.87**
*Let $X$ be a separated variety. To demonstrate universal closedness it is sufficient to show that*

$$\pi_Y : X \times Y \to Y$$

*is closed for $Y$ affine (or even $Y := \mathbb{A}^n_k$).*

*Further it is sufficient to show that $\pi_Y(T)$ is closed for all **irreducible** closed subsets $T \subset X \times Y$.*

*Proof.* Suppose $Y = \bigcup_{i=1}^n U_i$ for $U_i$ open affines. Then by (6.2.61) $X \times U_i$ is an open cover of $X \times Y$. Let $T \subset X \times Y$ closed then $T \cap (X \times U_i)$ is closed in $X \times U_i$. By assumption $\pi_Y(T \cap (X \times U_i)) = \pi_Y(T) \cap U_i$ is closed in $U_i$. Therefore by (4.1.22) $\pi_Y(T)$ is closed in $Y$ as required.

Clearly the closed property is preserved under isomorphism of $Y$, therefore we may assume that $Y = V(\mathfrak{a}) \subset \mathbb{A}^n_k$ is a closed subset for some $n$ and ideal $\mathfrak{a} \lhd k[X_1, ..., X_n]$.

By (6.2.61) $X \times Y$ is a closed subset of $X \times \mathbb{A}^n_k$. There $T \subset X \times Y$ is also closed in $X \times \mathbb{A}^n_k$ and the result follows.

Finally if $T$ is closed then by (...) $T = T_1 \cup ... \cup T_m$ for $T_j$ irreducible closed subsets of $T$ (and therefore of $X \times Y$). Therefore

$$\pi_Y(T) = \pi_Y\left(\bigcup_{j=1}^m T_j\right) = \bigcup_{j=1}^m \pi_Y(T_j)$$

and it suffices to show that $\pi_Y(T_j)$ is closed. ∎

The following is not strictly needed in application to projective case as it's already known however we include it for completeness of the argument.

**Lemma 6.2.88**
*Let $X$ be an integral variety and $R \subset k(X)$ is a valuation ring with local homomorphism $\Phi : R \to \overline{k}$. Suppose that there exists $x \in X$ such that $\mathcal{O}_{X,x} \subset R$, then $x$ may be chosen such that $\Phi|_{\mathcal{O}_{X,x}} \to \overline{k}$ is evaluation at $x$.*

*Proof.* By (6.2.19) $\Phi|_{\mathcal{O}_{X,x}}$ is local and by (6.2.50) we may choose $x' \in \overline{\{x\}}$ such that $\Phi|_{\mathcal{O}_{X,x}}$ is precisely evaluation at $z'$. ∎

The following proof is adapted from [Bum98] (Prop. 7.17).

**Proposition 6.2.89 (Complete $\implies$ Universally Closed)**
*A complete variety is universally closed.*

*Proof.* We may consider the projection $\pi_Y : X \times Y \to Y$ for $Y$ irreducible affine by (6.2.87), and demonstrate that $\pi_Y(T)$ is closed for every irreducible closed subset $T \subset X \times Y$.

Consider the closed subvariety $Z := \mathrm{cl}_X(\pi_X(T))$ of $X$ and the irreducible and affine closed subvariety $Y' := \mathrm{cl}_Y(\pi_Y(T))$ of $Y$. Then evidentally $T = T \cap (Z \times Y')$ and so

$$\pi_Y(T) = \pi_Y|_{Z \times Y'}\ (T \cap (Z \times Y'))$$

Further by (...) $Z \times Y'$ is a closed subvariety of $X \times Y$ and $T$ is an irreducible closed subset of $Z \times Y'$. Therefore for fixed $T$ we may assume without loss of generality that $\pi_Y(T)$ is dense in $Y$. The composite maps

$$\pi_Y \circ i : T \hookrightarrow Z \times Y \to Y$$

$$\pi_X \circ i : T \hookrightarrow Z \times Y \to Z$$

are regular (6.2.42) and dominant, and we require to prove the first is surjective. Consider the injective algebra homomorphism (6.2.17)

$$(\pi_Y \circ i)_\star : k(Y) \hookrightarrow k(T)$$

For any $y \in Y$ there is a corresponding evaluation homomorphism $\phi : \mathcal{O}_Y(Y) \to \overline{k}$ (6.2.52). By (3.25.11) there is a valuation ring $(\pi_Y \circ i)_\star(\mathcal{O}_Y(Y)) \subset R \subset k(T)$ and a local homomorphism $\Phi : R \to \overline{k}$ such that $\Phi \circ (\pi_Y \circ i)_\star$ extends $\phi$.

© David Rufino CC BY-NC-ND 4.0

Consider further the injective algebra homomorphism

$$(\pi_X \circ i)_\star : k(Z) \hookrightarrow k(T)$$

Then $R' \coloneqq (\pi_X \circ i)_\star^{-1}(R)$ is a valuation ring of $k(Z)$ together with local homomorphism $\Phi' : R' \to \overline{k}$ such that $\Phi' = \Phi \circ (\pi_X \circ i)_\star$. By hypothesis there is some $z \in Z$ such that $\mathcal{O}_{Z,z} \subset R'$ and by (6.2.88) we may assume that $\Phi'|_{\mathcal{O}_{Z,z}}$ is precisely evaluation at $z$.

Let $U$ be an open affine neighbourhood of $z$ in $Z$, then by (6.2.61) $U \times Y$ is an affine open subset of $Z \times Y$. As $\pi_X(T)$ is dense in $Z$ then $U \cap \pi_X(T)$ is non-empty. This shows that $U' \coloneqq (U \times Y) \cap T$ is a non-empty open subset of $T$. Evidently $U'$ is a closed subset of $U \times Y$. Therefore it is also affine (6.2.47) (6.2.42).

There is a commutative diagram (6.2.18)




By construction of $R$ we see that $(i|_{U'})^\sharp(\mathcal{O}_Z(U) \otimes 1) \subset R$ (after suitable identifications). Similarly we have a commutative diagram




and by construction we have $(i|_{U'})^\sharp(1 \otimes \mathcal{O}_Y(Y)) \subset R$ (after suitable idenitifications). By properties of the tensor product this means

$$(i|_{U'})^\sharp(\mathcal{O}_Z(U) \otimes \mathcal{O}_Y(Y)) \subset R$$

By (6.2.42) the map $(i|_{U'})^\sharp$ is surjective, therefore we conclude that $\mathcal{O}_T(U') \subset R$. Then by (6.2.52) the restriction of $\Phi$ to $\mathcal{O}_T(U')$ corresponds to a point $(z', y')$ of $U'$. Further $\Phi \circ (\pi_Y \circ i|_{U'})^\sharp = \big(\Phi \circ (\pi_Y \circ i)_\star\big)|_{\mathcal{O}_Y(Y)} = \phi$ and so by (6.2.52) we have $(\pi_Y \circ i)(z', y') = y$ that is to say $y' = y$. This shows that $\pi_Y \circ i$ is surjective as required. ∎

**Proposition 6.2.90**

*Let $X, Y$ be varieties with $Y$ complete such that*

*a) $U \subset X$ is an irreducible open neighbourhood*
*b) $x \in \overline{U}$*
*c) $\mathcal{O}_{X,x}$ is a valuation ring*
*d) $\phi : U \to Y$ is a regular morphism*

*Then there exists an open subset $U' \supset U$ for which $\overline{\{x\}} \subset U'$ and $\phi$ lifts to $U'$.*

*Proof.* We claim it's enough to find an open neighbourhood $V$ of $x$ such that $U \cap V \neq \varnothing$ and a regular function $\psi : V \to Y$ such that $\psi|_{U\cap V} = \phi|_{U\cap V}$. For then this extends to a regular function $U \cup V$ as required.

Replacing $X = \overline{U}$ we may assume that $U$ is dense in $X$ and $X$ is irreducible. Observe $Z \coloneqq \mathrm{cl}_Y(\phi(U))$ is an irreducible closed subset of $Y$ (4.1.70) (4.1.65). Therefore there is an injective $k$-algebra homomorphism (4.3.16)

© David Rufino CC BY-NC-ND 4.0

$$\phi_\star : k(Z) \hookrightarrow k(U) \xrightarrow{\sim} k(X)$$

By assumption $\mathcal{O}_{X,x}$ is a valuation ring, whence so is the inverse image $R$. By assumption there is $z \in Z$ such that $\mathcal{O}_{Z,z} \subset R$. Therefore we have a $k$-algebra homomorphism $\mathcal{O}_{Z,z} \to \mathcal{O}_{X,x}$. By (6.2.32) this lifts to a regular morphism $\psi : V \to Z \subset Y$ for which $x \in V$. As $x \in \overline{U}$ we have $U \cap V \neq \varnothing$ by (4.1.19). As $Y$ is separated (6.2.63) the equaliser of $\phi, \psi$ is a closed subset of $U \cap V$. We have by construction a commutative diagram

$$\begin{array}{ccc} k(Z) & \xrightarrow{\phi_\star} & k(X) \\ \uparrow & & \uparrow \\ \mathcal{O}_{Z,z} & \xrightarrow{\psi_x^\sharp} & \mathcal{O}_{X,x} \\ \uparrow & & \uparrow \\ \mathcal{O}_Z(Z) & \xrightarrow{\psi_Z^\sharp} & \mathcal{O}_X(V) \end{array}$$

Regarding $\psi : X \longrightarrow Y$ as a rational map we find by (6.2.76) that $\psi_\star = \phi_\star$ whence $\psi = \phi$ as rational maps. This means that the equaliser contains a dense open subset of $U \cap V$ and being closed is therefore the whole set. That is $\phi|_{U\cap V} = \psi|_{U\cap V}$ as required. ∎

## 6.2.16 Projective Space

We introduce the notion of projective varieties by showing that projective space can naturally be given the structure of an abstract variety. A more low-tech approach is taken in the Projective Varieties section, where the constructions are shown to be equivalent.

**Definition 6.2.91 (Projective Space)**
*Let $K$ be a field extension define*

$$\mathbb{P}^n(K) := K^{n+1} \setminus \{(0,...,0)\}/\sim$$

*where*

$$(x_0,...,x_n) \sim (x_{0'},...,x_{n'}) \iff x_i = \lambda x_{i'} \text{ some } \lambda \in K^\star$$

*and write $(x_0 : ... : x_n)$ for the corresponding equivalence class in $\mathbb{P}^n(K)$.*

**Lemma 6.2.92 (Comparing projective points)**
*Consider two projective points $x = (x_0 : ... : x_n)$ and $y = (y_0 : ... : y_n)$.*

*a) $x = y \implies \forall i (x_i = 0 \iff y_i = 0)$*

*Suppose that $x_i \neq 0$ for some $i$. Then $x = y$ iff*

*a) $y_i \neq 0$*
*b) $\frac{x_j}{x_i} = \frac{y_j}{y_i}$ for all $j$*

**Proposition 6.2.93 (Projective Space)**
*Consider the set $X = \mathbb{P}^n\left(\overline{k}\right)$ and for every $i = 0,...,n$ define the subset*

$$U_i := \left\{(x_0 : ... : x_n) \;\middle|\; x_0,...,x_n \in \overline{k}, x_i \neq 0\right\}.$$

*There are canonical bijections*

$$\begin{aligned} \theta_i : U_i &\xrightarrow{\sim} \mathbb{A}_k^n\left(\overline{k}\right) \quad i = 0...n \\ (x_0 : ... : x_n) &\to \left(\frac{x_0}{x_i},...,\frac{\widehat{x_i}}{x_i},...,\frac{x_n}{x_i}\right) \\ (z_1 : ... : 1 : ... : z_n) &\leftarrow (z_1,...,z_n) \end{aligned}$$

*and there is a unique topology on $X$ and space of functions $\mathcal{O}_X$ such that $U_i$ is open in $X$ and $\theta_i$ is a regular isomorphism for all $i = 0...n$.*

*Denote this by $\mathbb{P}_k^n$. With this structure $\mathbb{P}_k^n$ is an irreducible, separated algebraic variety of dimension $n$.*

*Further for any homogenous polynomial $F \in k[X_0,...,X_n]$ of degree $d$ the expression*

© David Rufino CC BY-NC-ND 4.0

$$(x_0 : ... : x_n) \to \frac{F(x_0, ..., x_n)}{x_i^d}$$

*is a well-defined regular function on $U_i$. In particular on $U_i$ we may define the* ***homogenous coordinate functions*** $\frac{X_0}{X_i}, ..., \frac{X_n}{X_i}$.

*Proof.* It is straightforward to show that $\theta_i$ is a bijection. Define the cocycle maps

$$\theta_{ij} : \theta_i(U_i \cap U_j) \to \theta_j(U_i \cap U_j)$$

by $\theta_{ij} = \theta_j \circ \theta_i^{-1}$. We show that $\theta_i(U_i \cap U_j)$ is open in $\mathbb{A}_k^n\left(\overline{k}\right)$ and the maps $\theta_{ij}$ are regular. First observe that

$$\theta_i(U_i \cap U_j) = \begin{cases} D(Z_j) & i < j \\ D(Z_{j+1}) & j < i \end{cases}$$

Then the cocycle map is given pointwise by

$$\theta_{ij} : D(Z_j) \to D(Z_{i+1})$$
$$(z_1, ..., z_n) \to \left(\frac{z_1}{z_j}, ..., \frac{z_i}{z_j}, \frac{1}{z_j}, ..., \frac{z_n}{z_j}\right)$$

Using (6.2.45) and (6.1.39) we may show that $\theta_{ij}$ is a regular morphism of abstract varieties. Therefore the structure on $X$ is uniquely determined by (6.2.38).

By construction $U_i$ is affine and $U_i \cap U_j = D\left(\frac{X_j}{X_i}\right)$ where $\frac{X_j}{X_i} \in \mathcal{O}_X(U_i)$. Therefore $U_i \cap U_j$ is affine by (6.2.21) and there is a canonical isomorphism

$$\mathcal{O}_{U_i}(U_i)_{\frac{X_j}{X_i}} \xrightarrow{\sim} \mathcal{O}_{U_i}(U_i \cap U_j)$$

As $\frac{X_i}{X_j} \in \mathcal{O}_{U_j}(U_j)$ satisfies $\frac{X_i}{X_j}|_{U_i \cap U_j} = \left(\frac{X_j}{X_i}|_{U_i \cap U_j}\right)^{-1}$ we find the canonical map

$$\mathcal{O}_{U_i}(U_i) \otimes \mathcal{O}_{U_j}(U_j) \to \mathcal{O}_{U_i}(U_i \cap U_j)$$

is surjective. Therefore $X$ is separated by (6.2.69).

By (6.1.21) the affine space $\mathbb{A}_k^n$ is irreducible whence so is $U_i$ and $U_i \cap U_j$ (4.1.70). Therefore by (4.1.74) $\mathbb{P}_k^n$ is irreducible and has dimension $n$ by (4.1.88). ∎

**Proposition 6.2.94 (Riemann Sphere)**
*There is a bijection*

$$\mathbb{P}_k^1 \xrightarrow{\sim} \mathbb{A}_k^1 \cup \{\infty\}$$

*which sends $(x : 1)$ to $x$ and $(1 : 0)$ to $\infty$. This gives $\mathbb{A}_k^1 \cup \{\infty\}$ the structure of a variety under which $\mathbb{A}_k^1$ is an open subvariety and $x^{-1}$ is a regular automorphism interchanging $0$ and $\infty$.*

*Consider $x = \frac{X_1}{X_0} \in \mathcal{O}_{\mathbb{P}_k^1}(U_0)$ as an element of the function field $k(\mathbb{P}_k^1)$ then $x$ is transcendental and $k(\mathbb{P}_k^1) = k(x)$.*

*Proof.* Evidently $x$ is transcendental because $\mathcal{O}(U_0) = k[x]$ (...). Let $f \in k(\mathbb{P}_k^1)$ and suppose $\mathrm{dom}(f) \cap U_0 \neq \varnothing$. There is some $g(x) \in \mathcal{O}(U_0)$ such that $D(g(x)) \subset \mathrm{dom}(f)$. Then by (...) $\mathcal{O}(D(g)) = k[x]_{g(x)}$ and so $f|_{D(g(x))} = h(x)/g(x)^r$ for some polynomial $h \in k[X]$ and integer $r > 0$. Therefore $f \in k(x)$. Similarly if $\mathrm{dom}(f) \cap U_1 \neq \varnothing$ then $f \in k(x^{-1}) = k(x)$ as required, where $x^{-1} = \frac{X_0}{X_1} \in \mathcal{O}_{\mathbb{P}_k^1}(U_1)$. ∎

We show later that all projective closed subvarieties are complete, but here we demonstrate a simpler case more directly.

**Proposition 6.2.95**
*$\mathbb{P}_k^1$ is complete.*

*Proof.* Let $Z \subset \mathbb{P}_k^1$ be an irreducible closed subvariety. Then by (4.1.83) (6.2.56) it is either the (closure of) a single point or the whole space. In the former case the function field is algebraic over $k$ and has no non-trivial valuation rings. Therefore we may assume wlog that $Z = \mathbb{P}_k^1$ and consider a valuation ring $R \subset k(\mathbb{P}_k^1) = k(x)$ where $x = \frac{X_1}{X_0}$. By (6.2.117) $R$ is equal to either $k[x]_{p(x)}$ for some irreducible polynomial or $k\left[x^{-1}\right]_{x^{-1}}$. Let $\alpha$ be a root of $p(x)$ in $\overline{k}$, then this corresponds to the point $(1 : \alpha)$, or in the latter case just $(0 : 1)$. ∎

© David Rufino CC BY-NC-ND 4.0

## 6.2.17 Affine and Finite Morphisms

**Definition 6.2.96**

*Let $\phi : X \to Y$ be a regular morphism of varieties. We say $\phi$ is **affine** if for every affine open $V \subset Y$, $\phi^{-1}(V)$ is affine.*

*We say in addition that $\phi$ is **finite** if the ring map $\mathcal{O}_Y(V) \to \mathcal{O}_X(\phi^{-1}(V))$ is finite (and therefore integral ...).*

Other than the definition we do not require this section in the sequel. But it may be convenient for other applications to show that it's sufficient to demonstrate a morphism is affine (resp. finite) only over a single affine cover.

**Lemma 6.2.97**

*Let $X$ be a variety with $U, V$ open affine subsets. Then there exists $g_1, ..., g_r \in \mathcal{O}_X(U)$ and $g'_1, ..., g'_r \in \mathcal{O}_X(V)$ such that*

*a) $W_i := D(g_i) = D(g'_i)$ for $i = 1...r$*
*b) $U \cap V = \bigcup_{i=1}^r W_i$*

*Proof.* Given $x \in U \cap V$, then as $U \cap V$ is open in $U$ by (6.2.10) we may find $f_x \in \mathcal{O}_X(U)$ such that $x \in D(f_x) \subset U \cap V$. Similarly we may find $f'_x \in \mathcal{O}_X(V)$ with $x \in D(f'_x) \subset D(f_x)$. By (6.2.21).c) there is a canonical isomorphism $\mathcal{O}_X(U)_{f_x} \xrightarrow{\sim} \mathcal{O}_X(D(f_x))$. Therefore $f'_x|_{D(f_x)} = \frac{g}{(f_x)^r}$ for some $g \in \mathcal{O}_X(U)$. Observe that $D(f'_x) = D(g) \cap D(f_x) = D(gf_x)$. This gives the required pair of sections for an arbitrary $x \in U \cap V$. Clearly these open sets cover $U \cap V$, and as $U \cap V$ is quasi-compact (6.2.7) we are done. ∎

**Proposition 6.2.98**

*Let $\phi : X \to Y$ be a regular morphism of varieties, and suppose there is an open cover $Y = \bigcup_{i=1}^n V_i$ for which $V_i$ and $\phi^{-1}(V_i)$ are affine. Then $\phi$ is affine.*

*If in addition $\mathcal{O}_Y(V_i) \to \mathcal{O}_X(\phi^{-1}(V_i))$ is finite for all $i = 1...n$ then $\phi$ is finite.*

*Proof.* By assumption we have affine open sets $U_i := \phi^{-1}(V_i)$ for $i = 1...n$. Let $V \subset Y$ be an open affine set, then by (6.2.97) for each $i$ we have an open cover

$$V \cap V_i = \bigcup_{j=1}^{n_i} W_{ij}$$

with $W_{ij} := D(f_{ij}) = D(f'_{ij})$ for $f'_{ij} \in \mathcal{O}_Y(V_i)$ and $f_{ij} \in \mathcal{O}_Y(V)$. Evidently

$$V = \bigcup_{i=1}^{n} \bigcup_{j=1}^{n_i} D(f_{ij}) \quad (*)$$

and so by (6.2.26) the $f_{ij}$ generate $\mathcal{O}_Y(V)$, and consequentally $g_{ij} := \phi^\sharp(f_{ij})$ generate $\mathcal{O}_X(U)$ (as both ideals contain 1). Further $\phi^{-1}(W_{ij}) = \phi^{-1}(D(f'_{ij})) = D(\phi^\sharp(f'_{ij})) \subset U_i$ is affine (6.2.21).c). Therefore from (6.2.28) $U$ is affine, because we also have the representation $\phi^{-1}(W_{ij}) = \phi^{-1}(D(f_{ij})) = D(g_{ij})$.

Suppose $\phi$ is finite then by definition this means the ring maps $\mathcal{O}_Y(V_i) \to \mathcal{O}_X(U_i)$ are finite. By (3.6.48).a) this means the localised maps are finite

$$\begin{array}{ccc}
\mathcal{O}_Y(V_i)_{f'_{ij}} & \longrightarrow & \mathcal{O}_X(U_i)_{g'_{ij}} \\
\downarrow \sim & & \downarrow \sim \\
\mathcal{O}_Y(W_{ij}) & \xrightarrow{\phi^\sharp} & \mathcal{O}_X(\phi^{-1}(W_{ij})) \\
\uparrow \sim & & \uparrow \sim \\
\mathcal{O}_Y(V)_{f_{ij}} & \dashrightarrow & \mathcal{O}_X(U)_{g_{ij}}
\end{array}$$

Using equation $(*)$ and (6.2.26) we see that $f_{ij}$ generate the unit ideal of $\mathcal{O}_Y(V)$. Therefore by (3.6.48).b) $\phi^\sharp_V$ is finite. ∎

## 6.2.18 Normal Varieties

**Definition 6.2.99**

*Let $A$ be an integral domain. We say that the ring $A$ is **normal** if it is integrally closed in its field of fractions.*

*Recall that $A$ factorial $\implies$ $A$ is normal (3.24.12).*

© David Rufino CC BY-NC-ND 4.0

**Proposition 6.2.100**
*The polynomial ring $A[X_1, ..., X_n]$ is normal for every unique factorisation domain $A$.*

*Proof.* See (3.24.12) and (3.17.46). ∎

**Proposition 6.2.101 (Normal Variety)**
*Let $X$ be an integral variety. Then the following are equivalent*

*a) $\mathcal{O}_X(U)$ is normal for every open subset $U \subset X$*
*b) $\mathcal{O}_{X,W}$ is normal for every irreducible subset $W \subset X$*
*c) $\mathcal{O}_{X,x}$ is normal for every point $x \in X$*

*In this case we say that $X$ is a **normal variety**. If $X$ is affine then this is equivalent to $\mathcal{O}_X(X)$ being normal.*

*Proof.* *a)* $\implies$ *b)* Let $U \subset X$ be any affine open subset such that $U \cap W \neq \varnothing$, then by (...) $\mathcal{O}_{X,W} \xrightarrow{\sim} \mathcal{O}_{U,W\cap U} \xrightarrow{\sim} \mathcal{O}_X(U)_{\mathfrak{m}_{U,W\cap U}}$, and the result follows from (3.24.10).

*b)* $\implies$ *c)* is straightforward because $\mathcal{O}_{X,x} = \mathcal{O}_{X,\{x\}}$.

*c)* $\implies$ *a)* We first show this for the case that $U$ is affine. By (...) $\mathcal{O}_{X,x} \xrightarrow{\sim} \mathcal{O}_{U,x} \xrightarrow{\sim} \mathcal{O}_X(U)_{\mathfrak{m}_{U,x}}$ is normal for every $x \in U$. Therefore by (6.2.21).b) and (3.24.10) $\mathcal{O}_X(U)$ is normal.

Now let $U$ be an arbitrary open set and $V \subset U$ affine. Then by (6.2.15) we have inclusions

$$\mathcal{O}_X(V) \hookrightarrow \mathcal{O}_X(U) \hookrightarrow \operatorname{Frac}(\mathcal{O}_X(U)) \hookrightarrow k(X)$$

and $k(X)$ is a fraction field for $\mathcal{O}_X(V)$. If $\alpha \in \operatorname{Frac}(\mathcal{O}_X(U))$ is integral over $\mathcal{O}_X(U)$, then it is a-fortiori integral over $\mathcal{O}_X(V)$. By the affine case then $\alpha \in \mathcal{O}_X(V)$ and we conclude that $\mathcal{O}_X(U)$ is normal as required.

For the case $X$ is affine we may use (3.24.10). ∎

**Corollary 6.2.102**
*The standard affine space $\mathbb{A}^n_k\left(\overline{k}\right)$ is normal.*

*Proof.* The coordinate ring is a UFD (3.17.46) and therefore normal (3.24.12). ∎

**Proposition 6.2.103**
*Let $X$ be an integral variety with open cover $X = \bigcup_{i=1}^n U_i$. Then $X$ is normal iff $U_i$ are normal.*

*Proof.* Straightforward, since normality is a local condition. ∎

## 6.2.19 Weil Divisors

The notion of regular curves can be generalised in one direction.

**Definition 6.2.104 (Prime Divisors of an Integral Variety)**
*Let $X$ be an integral, separated variety. We say it is **regular in codimension one** if for every irreducible closed subset $Y \subset X$ for which $\operatorname{codim}(Y, X) = 1$ the local ring $\mathcal{O}_{X,Y}$ is regular (equivalently integrally closed or a DVR (3.31.2)).*

*In this case, we say an irreducible closed subset of codimension 1 is a **prime divisor**.*

*We denote by $\operatorname{Div} X$ the free abelian group generated by the prime divisors. We write a prime divisor $D$ by a formal sum*

$$D = \sum_{Y \subset X} n_Y Y$$

*where $n_Y \in \mathbb{Z}$ and all but finitely many are zero. We define the **degree** of a divisor $D$ by*

$$\deg D = \sum_{Y \subset X} n_Y \left[k\left(\mathfrak{m}_{X,Y}\right) : k\right]$$

**Proposition 6.2.105**
*An integral, separated variety $X$ is regular in codimension one for either of the following special cases*

*a) $X$ is normal*
*b) $X$ is non-singular and $k$ is perfect.*

*Proof.* See (3.31.2) and (6.2.82). ∎

© David Rufino CC BY-NC-ND 4.0

**Proposition 6.2.106 (Valuation Associated to Prime Divisor)**
*Let $X$ be a variety satisfying (6.2.104) and $Y \subset X$ a prime divisor. Then $\mathcal{O}_{X,Y}$ is a discrete valuation ring.*

*We denote by $v_Y : k(X)^\star \to \mathbb{Z}$ the corresponding valuation.*

*Proof.* By (6.2.55) $\dim \mathcal{O}_{X,Y} = \operatorname{codim}(Y, X) = 1$ and is integral, so the result follows from (3.31.2). ∎

**Proposition 6.2.107 (Prime Divisors on a Curve)**
*Let $X$ be a regular curve. Then $X$ satisfies (6.2.104) and the prime divisors are precisely irreducible closed subsets of the form $\overline{\{x\}}$. These are in bijective correspondence with the valuation rings of $k(X)$, via the map $x \to \mathcal{O}_{X,x}$.*

*Proof.* By the codimension formula (6.2.55) the irreducible closed subsets of codimension 1 are precisely those of dimension 0. Therefore the result follows from (6.2.124). ∎

**Proposition 6.2.108**
*Let $X$ be a variety satisfying (6.2.104) and $f \in k(X)^\star$ a rational function. Then $v_Y(f) = 0$ for all but finitely many prime divisors $Y$.*

*Proof.* By definition $f$ is regular on some open subset $U$ of $X$, which we may assume to be affine. Define $Z \coloneqq X \setminus U$. Then by (...) $Z = Z_1 \cup ... \cup Z_n$ is a decomposition into irreducible components. By (4.1.83).f) $\operatorname{codim}(Z, X) > 0$ whence $\operatorname{codim}(Z_i, X) > 0$ for all $i = 1...n$. Suppose $Y \subset Z$ is a prime divisor then by definition $\operatorname{codim}(Y, X) = 1$ and by (4.1.66) $Y \subset Z_i$ for some $i$. Therefore by (4.1.83).c) $\operatorname{codim}(Y, Z_i) \leq \operatorname{codim}(Y, X) - \operatorname{codim}(Z_i, X) \leq 0$ and therefore by (4.1.83).f) $Y = Z_i$.

We conclude all but finitely many $Y$ intersect $U$, so we may without loss of generality assume $Y \cap U \neq \varnothing$. Evidently $U$ also satisfies (6.2.104) and by (4.1.79) there is a bijection between prime divisors of $X$ meeting $U$, and prime divisors of $U$.

Therefore we may assume that $X$ is affine and $0 \neq f \in \mathcal{O}_X(X)$. Then $v_Y(f) > 0 \iff f \in \mathfrak{M}_{X,Y} \iff f \in I(Y) \iff Y \subset V(f)$. We claim $V(f)$ is a proper subset of $X$ (for $V(f) = X \iff \sqrt{(0)} = \sqrt{f} \iff f = 0$). Therefore $\operatorname{codim}(V(f), X) > 0$ and

$$\operatorname{codim}(Y, V(f)) \leq \operatorname{codim}(Y, X) - \operatorname{codim}(V(f), X) \leq 0$$

so $Y$ is an irreducible component of $V(f)$, of which there are only finitely many (4.1.66). ∎

**Definition 6.2.109 (Principal Divisor)**
*Let $X$ be a variety satisfying (6.2.104) then define the homomorphism*

$$\begin{aligned} k(X)^\star &\longrightarrow \operatorname{Div}(X) \\ f &\longrightarrow (f) \end{aligned}$$

*where we define a formal sum over all prime divisors*

$$(f) \coloneqq \sum_{Y \subset X} v_Y(f) Y.$$

*The co-kernel of this map is known as the **divisor class group** of $X$, and denoted $\operatorname{Cl}(X)$. We say that two divisors are equivalent $D \sim D'$ if they become equal as elements of the divisor class group. Equivalently if $D - D' = (f)$ for some $f \in k(X)^\star$.*

## 6.2.20 Algebraic Function Fields

We develop the basic theory of **algebraic function fields** of one variable, the field-theoretic counterpart to a regular algebraic curve.

**Proposition 6.2.110**
*Let $K/k$ be a field extension. Then the following are equivalent*

*a) $K/k$ is finitely generated with transcendence degree 1.*
*b) There exists a transcendental element $x \in K/k$ such that $K/k(x)$ is finite.*

*In this case we say that $K/k$ is an **algebraic function field**. We typically assume that $k$ is perfect (either a number field or a finite field).*

*Proof.* *a)* $\implies$ *b)* By (3.19.139) a transcendence base consists of a single element $x$. By assumption $K/k(x)$ is algebraic and finitely generated, hence also finite (3.19.44).

© David Rufino CC BY-NC-ND 4.0

$b) \implies a)$ Evidently $\{x\}$ is a transcendence base. By assumption $K/k(x)$ is finite and therefore certainly finitely generated, whence so is $K/k$. ∎

**Definition 6.2.111 (Algebraic Function Field)**
*Let $K/k$ be an algebraic function field. Denote by $\tilde{k}$ the **field of constants**, the relative algebraic closure of $k$ in $K$:*

$$\tilde{k} := \{x \in K \mid x \text{ algebraic over } k\}$$

*(a subfield of $K$ by (3.19.45)). By (3.19.46), $K/\tilde{k}$ is relatively algebraically closed. We say that $K/k$ is **geometrically integral** if $k = \tilde{k}$ (c.f. (3.36.41)).*

**Definition 6.2.112 (Place of an Algebraic Function Field)**
*Let $K/k$ be an algebraic function field. By a **place** of $K/k$ we refer to a valuation ring $\mathcal{O}_P$ such that $k \subsetneq \mathcal{O}_P \subsetneq K$. By (3.25.2) this is a local ring with unique maximal ideal $\mathfrak{m}_P$.*

*From (3.3.13) we verify immediately that $k \cap \mathfrak{m}_P = (0)$.*

*For every $x \in K$ we say that $x$ is **regular** at a place $P$ if $x \in \mathcal{O}_P$.*

**Proposition 6.2.113**
*Let $K/k$ be a function field and $k \subsetneq \mathcal{O}_P \subsetneq K$ a valuation ring. Then $\tilde{k} \subset \mathcal{O}_P$ and $\tilde{k} \cap \mathfrak{m}_P = (0)$*

*Proof.* By definition $\tilde{k}$ is the integral closure of $k$ in $K$, so $\tilde{k} \subset \mathcal{O}_P$ by (3.25.14). Suppose $0 \neq x \in \tilde{k}$ then $x$ is invertible in $\mathcal{O}_P$ and therefore $x$ is not contained in any proper ideal, and in particular $x \notin \mathfrak{m}_P$. ∎

In order to use the theory of Dedekind domains and affine algebras we establish a connection between valuation rings and a certain "affine" subring.

**Proposition 6.2.114**
*Let $K/k$ be an algebraic function field and $f \in K \setminus \tilde{k}$ transcendental. Then the integral closure $\overline{k[f]}$ in $K$ is a Dedekind domain with quotient field $K$ and a finite $k[f]$-module. There is a correspondence*

$$\begin{aligned} \left\{\mathfrak{B} \triangleleft \overline{k[f]}\right\} &\longleftrightarrow \{\mathcal{O}_P \subsetneq K \mid f \in \mathcal{O}_P\} \\ \mathfrak{B} &\longrightarrow \overline{k[f]}_{\mathfrak{B}} \\ \mathfrak{m}_P \cap \overline{k[f]} &\longleftarrow \mathcal{O}_P \end{aligned}$$

*In particular every $\mathcal{O}_P$ is a discrete valuation ring, and the residue degree satisfies*

$$[k(\mathfrak{m}_P) : k] = [k(\mathfrak{B}) : k] \leq [K : k(f)]$$

*Therefore we may associate to each place $P$ a discrete valuation*

$$v_P : K \longrightarrow \mathbb{Z} \cup \{\infty\}$$

*which is zero on $k$ and $v_P(f) = \infty \iff f \notin \mathcal{O}_P \iff f^{-1} \in \mathfrak{m}_P$.*

*Proof.* By (3.32.2) $k[f] \subset K$ is a Dedekind domain and so by (3.35.61) $B := \overline{k[f]}^K$ is a Dedekind domain and finite $k[f]$-module. As $\mathcal{O}_P$ is a valuation ring, it is integrally closed (3.25.2) and as it contains $k[f]$, also contains $B$. Then we may deduce this correspondence from (3.32.3) and the inequality from (3.32.15).

This covers every place $P$ with $f \in \mathcal{O}_P$. For a place $P$ with $f \notin \mathcal{O}_P$ instead, $f^{-1} \in \mathfrak{m}_P \subset \mathcal{O}_P$ (3.25.2), and $f^{-1}$ is transcendental too (else $f = (f^{-1})^{-1}$ would be algebraic); applying the above with $f^{-1}$ in place of $f$ shows $\mathcal{O}_P$ is again a discrete valuation ring. So every place $P$ of $K/k$ is a discrete valuation ring, giving the associated valuation $v_P$. ∎

**Corollary 6.2.115**
*Let $K/k$ be an algebraic function field. Then the field of constants $\tilde{k}/k$ has finite degree.*

*Proof.* For any place $P$ we have an embedding $\tilde{k}/k \hookrightarrow k(\mathfrak{m}_P)/k$ which is finite (6.2.114). ∎

We prove a number of facts in analogy with algebraic curves.

**Lemma 6.2.116 (Every function has a zero)**
*Let $K/k$ be an algebraic function field and $0 \neq f \in K$. If $f \in \tilde{k}$ (i.e. $f$ is algebraic), then $v_P(f) = 0$ for every place $P$, so $f$ has no zeros or poles; otherwise ($f$ transcendental) $f$ has between 1 and $[K : k(f)]$ zeros (resp. poles).*

*Proof.* If $f \in \tilde{k}$, then $f \neq 0$ gives $f \in \tilde{k} \setminus \{0\}$, so $f \notin \mathfrak{m}_P$ for every place $P$ (6.2.113), that is $v_P(f) = 0$.

© David Rufino CC BY-NC-ND 4.0

Otherwise, clearly $k[f]$ is a polynomial ring with maximal ideal $(f)$. Therefore by (3.25.12) there is a valuation ring $\mathcal{O}_P \supset k[f]$ such that $\mathfrak{m}_P \cap k[f] = (f)$, which means $f$ has at least one zero.

Recall by (6.2.114) that zeros of $f$ correspond to the maximal ideals of $\overline{k[f]}$ containing $f$. These are precisely the prime divisors of $(f)$ (see (3.32.11)), so the bound follows from (3.32.16).

The statement about poles of $f$ follows immediately because these are precisely zeros of $f^{-1}$ and $k(f) = k(f^{-1})$. ∎

**Proposition 6.2.117 (Places of the Rational Function Field)**
*Let $K = k(x)$ with $x$ transcendental over $k$. Then the places of $K/k$, that is the valuation rings $k \subsetneq \mathcal{O} \subsetneq K$, are exactly the following pairwise distinct discrete valuation rings*

*a) $\mathcal{O}_{p(x)} := k[x]_{(p(x))}$ where $p \in k[X]$ is an irreducible monic polynomial, with uniformiser $p(x)$ and residue field $k[X]/(p)$, and*
*b) $\mathcal{O}_\infty := \left\{ \frac{f(x)}{g(x)} \;\middle|\; g \neq 0, \deg f \leq \deg g \right\}$, with uniformiser $x^{-1}$ and residue field $k$.*

*If $k$ is algebraically closed then the irreducible monic polynomials are the $X - \theta$ with $\theta \in k$, so every place is rational, with uniformiser $x - \theta$ or $x^{-1}$.*

*Proof.* Let $t := x^{-1}$, which is again transcendental over $k$ with $k(t) = K$.

**The listed rings are discrete valuation rings.** As $k[x] \cong k[X]$ is a PID (3.16.4), $\mathcal{O}_{p(x)}$ is a discrete valuation ring (3.31.4), with maximal ideal $p(x)\mathcal{O}_{p(x)}$ and residue field $k[x]/(p(x))$ (3.6.27).

Next $\mathcal{O}_\infty = k[t]_{(t)}$: if $f, g \in k[X]$ with $g(0) \neq 0$ and $N := \max(\deg f, \deg g)$ then $f(t)/g(t) = F(x)/G(x)$ with $F(x) := x^N f(1/x)$, $G(x) := x^N g(1/x)$, $\deg F \leq N = \deg G$; conversely $F(x)/G(x)$ with $\deg F \leq \deg G =: N$ equals $f(t)/g(t)$ with $f(t) := t^N F(1/t)$ and $g(t) := t^N G(1/t)$, where $g(0)$ is the leading coefficient of $G$. So $\mathcal{O}_\infty$ is a discrete valuation ring with uniformiser $t$ and residue field $k[t]/(t) = k$, as above.

**Every place is listed.** Let $\mathcal{O}$ be a place with maximal ideal $\mathfrak{m}$. Suppose firstly that $x \in \mathcal{O}$. Then $k[x] \subset \mathcal{O}$ and the ideal $k[x] \cap \mathfrak{m} =: (p(x))$ is principal. There is an injective $k$-algebra homomorphism $k[x]/(p(x)) \hookrightarrow k(\mathfrak{m})$ whence $(p(x))$ is prime (3.3.59). As $k(\mathfrak{m})/k$ is algebraic (6.2.114) and $x$ is transcendental, we must have $0 \neq p$, which we take monic, and therefore $p$ is irreducible (3.17.11). Suppose $g(x) \notin (p(x))$ then $g(x) \notin \mathfrak{m} \implies g(x)^{-1} \in \mathcal{O}$ (3.25.2). Therefore $\mathcal{O}_{p(x)} \subset \mathcal{O}$, and as $\mathcal{O}_{p(x)}$ is a discrete valuation ring and $\mathcal{O} \neq K$ we conclude that they are equal (3.31.5).

Alternatively suppose that $x \notin \mathcal{O}$ then by (3.25.2) $t = x^{-1} \in \mathfrak{m} \subset \mathcal{O}$. Applying the previous part with $t$ in place of $x$ gives $\mathcal{O} = \mathcal{O}_{p(t)}$ where $\mathfrak{m} \cap k[t] = (p(t))$. As $t \in \mathfrak{m}$ we conclude that $p(t) = t$, and $\mathcal{O} = k[t]_{(t)} = \mathcal{O}_\infty$.

**Distinctness.** $x \in \mathcal{O}_{p(x)}$ but $x \notin \mathcal{O}_\infty$, as $\deg x > \deg 1$. For distinct monic irreducible $p \neq p'$, $p \notin (p')$, so $p(x)$ is a unit of $\mathcal{O}_{p'(x)}$ but not of $\mathcal{O}_{p(x)}$.

Finally if $k$ is algebraically closed the irreducible monic polynomials are the $X - \theta$ (3.19.58), with residue field $k[X]/(X - \theta) = k$. ∎

**Lemma 6.2.118 (Restricting a Place is Proper)**
*Let $L/k'$ be an algebraic function field, $P'$ a place of $L/k'$, $k \subseteq k'$ with $k'/k$ algebraic, and $K \subseteq L$ a subfield with $\operatorname{trdeg}_k(K) \geq 1$ (e.g. $K/k$ itself an algebraic function field). Then $\mathcal{O}_{P'} \cap K$ is a proper valuation ring of $K$ (and therefore a place.)*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

$\mathcal{O}_{P'} \cap K$ is a valuation ring of $K$ (3.25.3). Suppose it equals $K$, so $K \subset \mathcal{O}_{P'}$. Nonzero elements of $K$ are units of $\mathcal{O}_{P'}$, hence not in $\mathfrak{m}_{P'}$, so $K \to \mathcal{O}_{P'}/\mathfrak{m}_{P'} = k'(\mathfrak{m}_{P'})$ is injective, and $\operatorname{trdeg}_k(K) \leq \operatorname{trdeg}_k(k'(\mathfrak{m}_{P'}))$ (3.19.136). But $k'(\mathfrak{m}_{P'})/k'$ is algebraic (6.2.114) and $k'/k$ is algebraic, so the right-hand side is 0, contradicting $\operatorname{trdeg}_k(K) \geq 1$. ∎

**Lemma 6.2.119 (Places Restrict Correctly)**
*Let $K/k$ be an algebraic function field, $P$ a place of $K/k$, and $L/K$ a field extension. Let $k \subseteq k'' \subset L$ with $k''/k$ algebraic and $L/k''$ itself an algebraic function field, and $P'$ a place of $L/k''$ with $\mathcal{O}_P \subset \mathcal{O}_{P'}$. Then $\mathcal{O}_{P'} \cap K = \mathcal{O}_P$; in particular $\mathfrak{m}_P \subset \mathfrak{m}_{P'}$, and $\pi_P \notin k''$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

$\mathcal{O}_{P'} \cap K$ is a valuation ring of $K$ containing $\mathcal{O}_P$ (3.25.3), and proper (as $K/k$ is an algebraic function field, so $\operatorname{trdeg}_k(K) = 1 \geq 1$) (6.2.118). By (3.31.5) ($\mathcal{O}_P$ being a DVR, (6.2.114)), a valuation ring of $K$ containing the DVR $\mathcal{O}_P$ is either $\mathcal{O}_P$ itself or all of $K$; as $\mathcal{O}_{P'} \cap K \neq K$, this forces $\mathcal{O}_{P'} \cap K = \mathcal{O}_P$.

For $0 \neq x \in \mathfrak{m}_P$ we have $x^{-1} \in K \setminus \mathcal{O}_P = K \setminus \mathcal{O}_{P'}$, so $x \in \mathfrak{m}_{P'}$ by (3.25.2): thus $\mathfrak{m}_P \subset \mathfrak{m}_{P'}$. In particular $\pi_P \in \mathfrak{m}_{P'}$ is a non-unit of $\mathcal{O}_{P'}$, so $\pi_P \notin k''$ (nonzero elements of the field $k'' \subset \mathcal{O}_{P'}$ are units). ∎

© David Rufino CC BY-NC-ND 4.0

### 6.2.21 Algebraic Curves

**Definition 6.2.120**
*We say an integral, separated algebraic variety of dimension* 1 *is an* ***algebraic curve*** *or just a* ***curve****.*

**Proposition 6.2.121 (Characterisation Regular Point on a curve)**
*Let $X$ be an algebraic curve and $x \in X$. Then the following are equivalent*

*a) $x$ is regular*
*b) $\mathcal{O}_{X,x}$ is a discrete valuation ring*
*c) $\mathcal{O}_{X,x}$ is an integrally closed domain*

*and when $k$ is perfect this is equivalent to $x$ being non-singular.*

*In particular $X$ is regular if and only if it is normal.*

*Proof.* Follows exactly the same lines as (6.1.100). ∎

**Example 6.2.122**
*Both $\mathbb{A}^1_k$ and $\mathbb{P}^1_k$ are regular algebraic curves.*

**Proposition 6.2.123 (Curve has Cofinite Topology)**
*Let $X$ be an algebraic curve (or just an algebraic variety of dimension* 1*) then $X_0$ has the cofinite topology.*

*Proof.* Follows from (4.1.92). ∎

**Proposition 6.2.124 (Points on Complete Regular Curve = Valuation Rings on Function Field)**
*Let $X$ be a complete regular algebraic curve. Then there is a bijection*

$$\{W \subset X \mid \dim W = 0 \text{ closed and irreducible}\} \longleftrightarrow X_0 \longleftrightarrow \{k \subset R \subsetneq k(X) \mid R \text{ a valuation ring}\}$$

*Further the corresponding valuation rings are all discrete valuation rings.*

*For an affine open subset $U \subset X$ this restricts to a bijection*

$$U_0 \longleftrightarrow \{\mathcal{O}_X(U) \subset R \subsetneq k(X) \mid R \text{ a valuation ring}\}$$

*where $\mathcal{O}_X(U)$ is identified as a subring of $k(X)$ (6.2.18).*

*Proof.* The first map is (4.1.94) and the second map is well-defined by (6.2.121) and injective by (6.2.20).

Let $R \subset k(X)$ be a valuation ring then by definition of completeness there is a point $x \in X$ such that $\mathcal{O}_{X,x} \subset R$. By (6.2.55) we have $\dim \mathcal{O}_{X,x} = \dim X = 1$. By (6.1.92) and (6.1.91) $1 = \dim \mathcal{O}_{X,x} \leq \dim R \leq \operatorname{trdeg}_k k(X) \overset{(6.2.54)}{=} \dim X = 1$. Therefore $\dim R = 1$ and by the same result $R = \mathcal{O}_{X,x}$. This shows that the map is surjective.

For the final statement we may use the affine case (6.1.101) and the commutative diagram

$$\begin{array}{ccc} \mathcal{O}_X(U) & \hookrightarrow & k(X) \\ \| & & \downarrow \sim \\ \mathcal{O}_U(U) & \hookrightarrow & k(U) \end{array}$$

∎

**Proposition 6.2.125**
*Let $X$ be a regular algebraic curve and $Y$ a complete variety. Then the rational maps are precisely the regular maps. In otherwords the canonical map*

$$\operatorname{Mor}(X, Y) \xrightarrow{\sim} \operatorname{Rat}(X, Y)$$

*is a bijection. Further this restricts to dominant maps.*

*Proof.* Clearly the map is well-defined. Suppose that two regular maps $\phi, \psi$ are equal as rational maps. Then by definition they agree on a dense open subset. By definition $Y$ is separated so by (6.2.63) the equaliser is a closed subset, and therefore the whole of $Y$. This means the map is injective.

Consider a regular map $\phi : U \to Y$ for $U$ a (dense) open subset of $X$. Then $Z := X \setminus U$ is a proper closed subset of $X$ and therefore $\dim Z = 0$ by (4.1.84). By (6.2.12) we have $Z = \overline{\{z_1\}} \cup \ldots \cup \overline{\{z_n\}}$, and we may apply (6.2.90) finitely

© David Rufino CC BY-NC-ND 4.0

many times to lift to a regular map on an open set containing $U$ and $z_1, ..., z_n$. By (4.1.41) this is the whole of $X$. Therefore the given map is surjective. ∎

**Proposition 6.2.126 (Image of Complete Curve either Constant or Surjective)**
*Let $X, Y$ be algebraic curves with $X$ complete and $\phi : X \to Y$ a regular morphism. Then $\phi$ is either surjective or has image of the form $\overline{\{x\}}$ (which we call constant).*

*Proof.* By (6.2.89) $\phi$ is universally closed, so by (6.2.84) $\phi(X)$ is an irreducible closed subvariety of $Y$. Evidently $\dim \phi(X) = 0$ or 1. Then we may use (6.2.56) or (4.1.84). ∎

**Corollary 6.2.127 (Equivalence between nice curves and function fields)**
*Let $X, Y$ be complete, regular algebraic curves. Then there are bijections*

$$\{\phi : X \to Y \text{ non-constant}\} \longleftrightarrow \{\phi : X \longrightarrow Y \text{ dominant}\} \longleftrightarrow \{\phi_\star : k(Y) \hookrightarrow k(X)\}$$

*All such morphisms are surjective and the degree*

$$\deg(\phi) := [k(X) : \phi_\star(k(Y))]$$

*is always finite.*

*Finally the fibres are finite (i.e. $\phi$ is "quasi-finite").*

*Proof.* This follows from (6.2.126), (6.2.125) and (6.2.76).

The fibres are finite because $\phi^{-1}(\{y\}) \subset \phi^{-1}\left(\overline{\{y\}}\right)$. By assumption this is a proper closed subset of $X$ and therefore has dimension 0. So we are done by (6.2.56). ∎

**Proposition 6.2.128**
*Let $X$ be a regular algebraic curve. Then there is a bijection*

$$k(X)^\star \longleftrightarrow \{\phi : X \to \mathbb{P}^1_k \text{ non-constant}\}$$

*More precisely given $f \in k(X)^\star$, then for $x \in X$ the function*

$$\phi_f(x) = \begin{cases} (f(x) : 1) & f \in \mathcal{O}_{X,x} \\ (1 : f^{-1}(x)) & f^{-1} \in \mathcal{O}_{X,x} \end{cases}$$

*is well-defined and regular.*

*Proof.* Recall (6.2.121) (6.2.15) that $\mathcal{O}_{X,x}$ is a (discrete) valuation ring with field of fractions $k(X)$.

Define the open subsets $U := \mathrm{dom}(f)$ and $V := \mathrm{dom}(f^{-1})$. We claim that $X = U \cup V$. For $x \in \mathrm{dom}(f) \iff f \in \mathcal{O}_{X,x}$ and as $\mathcal{O}_{X,x}$ is a valuation ring for $k(X)$ then we have either $x \in U$ or $x \in V$.

Define $\phi : U \to \mathbb{P}^1_k$ by $\phi(x) = (f(x) : 1)$, and similarly $\psi : V \to \mathbb{P}^1_k$ by $\psi(x) = (1 : f^{-1}(x))$. By definition of projective structure sheaf (6.2.93) these are regular. Then for $x \in U \cap V$ we must have $f \notin \mathfrak{m}_{X,x}$ that is $f(x) \neq 0$ and $f^{-1}(x) = f(x)^{-1}$. This shows that $\phi|_{U\cap V} = \psi|_{U\cap V}$. Therefore they glue to a well-defined regular map $X \to \mathbb{P}^1_k$ as required.

We note that $\phi_f^{-1}(U_1) = \mathrm{dom}(f)$ where $U_1$ is the second affine piece, and we conclude the map is injective.

Conversely given a morphism $\phi : X \to \mathbb{P}^1_k$ then $U := \phi^{-1}(U_1)$ is a non-empty open subset of $X$ and $f : X_0/X_1 \circ \phi|_U : U \to \overline{k}$ is a regular function. Evidently $\phi|_U = \phi_f|_U$, and we conclude $\phi = \phi_f$ by (6.2.125). Therefore the map is surjective. ∎

**Proposition 6.2.129 (Regular Morphisms of Complete Curves are Finite)**
*Let $\phi : X \to Y$ be a non-constant morphism of complete regular algebraic curves. Suppose $V \subset Y$ is an affine open subset, then*

$$\mathcal{O}_X(\phi^{-1}(V)) = \overline{\phi_\star(\mathcal{O}_Y(V))}$$

*where $\bar{\cdot}$ denotes the integral closure in $k(X)$.*

*Further $\phi$ is a finite (and therefore affine, and integral) morphism.*

*Proof.* Define $A := \mathcal{O}_Y(V)$ and $B := \overline{\phi_\star(A)}$. Then

$$\begin{aligned} B &= \bigcap\nolimits_{\phi_\star(A)\subset R} R && (3.25.14) \\ &= \bigcap\nolimits_{A\subset\phi_\star^{-1}(\mathcal{O}_{X,x})} \mathcal{O}_{X,x} && (6.2.124) \end{aligned}$$

© David Rufino CC BY-NC-ND 4.0

$$\begin{aligned}&= \bigcap_{A \subset \mathcal{O}_{Y,\phi(x)}} \mathcal{O}_{X,x} && (6.2.18)\\ &= \bigcap_{\phi(x) \in V} \mathcal{O}_{X,x} && (6.2.124)\\ &= \mathcal{O}_X(\phi^{-1}(V)) && (6.2.73)\end{aligned}$$

By (6.2.54) $\dim V = \dim Y = 1$ and therefore $\dim \mathcal{O}_Y(V) = 1$ (6.1.50). By (3.35.61) $B$ is a Dedekind domain and a finite $A$-module with quotient field $k(X)$.

Let $U \coloneqq \phi^{-1}(V)$ and observe that as $\mathcal{O}_{X,x}$ is integrally closed

$$B \subset \mathcal{O}_{X,x} \iff \phi_\star(A) \subset \mathcal{O}_{X,x} \iff A \subset \mathcal{O}_{Y,\phi(x)} \iff \phi(x) \in V \iff x \in U$$

where we have used the fact $V$ is affine (6.2.124). This shows that there is a bijection between $U_0$ and the valuation rings of $k(X)$ containing $\mathcal{O}_X(U)$. It remains to show that $U$ is affine.

Let $Z$ denote an integral affine curve with coordinate ring (isomorphic to) $B = \mathcal{O}_X(U)$. By the Affine Mapping Property (6.2.22) (applicable to $U$ without yet knowing it is affine, since only $Z$ needs to be) there is a corresponding regular morphism

$$\psi : U \to Z$$

such that $\psi^\sharp_Z : \mathcal{O}_Z(Z) \to \mathcal{O}_U(U)$ is an isomorphism. Consider the commutative

$$\begin{array}{ccc} \mathcal{O}_Z(Z) & \xrightarrow[\sim]{\psi^\sharp_Z} & \mathcal{O}_U(U) \\ \big\downarrow & & \big\downarrow \\ k(Z) & \xrightarrow[\sim]{\psi_\star} & k(U) \end{array}$$

as $k(Z)$ is the field of fractions of $\mathcal{O}_Z(Z)$ and similarly for $\mathcal{O}_U(U)$ (6.2.15), we see that $\psi_\star$ is an isomorphism. In light of the correspondence of valuation rings to points we see that $\psi_0 : U_0 \to Z_0$ is a bijection. As these have the cofinite topology (6.2.123) then $\psi_0$ is necessarily a homeomorphism.

For $g \in \mathcal{O}_Z(Z)$ we have a commutative diagram

$$\begin{array}{ccc} \mathcal{O}_Z(Z) & \xrightarrow{\sim} & \mathcal{O}_U(U) \\ \big\downarrow & & \big\downarrow \\ \mathcal{O}_Z(D(g)) & \xrightarrow{\psi^\sharp_{D(g)}} & \mathcal{O}_U\big(D(\psi^\sharp(g))\big) \\ \sim\big\uparrow & & \big\uparrow\sim \\ \mathcal{O}_Z(Z)_g & \xrightarrow{\sim} & \mathcal{O}_U(U)_{\psi^\sharp(g)} \end{array}$$

where the bottom isomorphisms follow from (6.2.27). This shows that $\psi^\sharp_{D(g)}$ is an isomorphism, and in particular (as the $D(g)$ form a basis of $Z$) that $\psi$ is continuous. We claim that the morphism of stalks

$$\psi_x : \mathcal{O}_{Z,\psi(x)} \to \mathcal{O}_{U,x}$$

is an isomorphism for all $x \in U$. Suppose we have $(W, \sigma) \in \mathcal{O}_{U,x}$ and let $g \in \mathcal{O}_Z(Z)$ be such that $\psi(x) \in D(g) \subset \psi(W)$ (6.1.15). Then as $\psi^\sharp_{D(g)}$ is an isomorphism there exists $\tau \in \mathcal{O}_Z(D(g))$ such that $\psi^\sharp(\tau) = \sigma|_{\psi^{-1}(D(g))}$ which has the required image in $\mathcal{O}_{U,x}$. Similarly suppose that $(T, \tau) \in \mathcal{O}_{Z,\psi(x)}$ maps to zero. Then there exists some neighbourhood $x \in W \subset \psi^{-1}(T)$ such that $\psi^\sharp(\tau)|_W = 0$. We may choose $g \in \mathcal{O}_Z(Z)$ such that $D(g) \subset \psi(W)$. Then $0 = \psi^\sharp(\tau)|_{\psi^{-1}(D(g))} = \psi^\sharp\big(\tau|_{D(g)}\big)$ whence $\tau|_{D(g)} = 0$ which shows that $(T, \tau)$ is zero.

By (6.2.50) (applied to $U$ and $Z$), together with $\psi_0$ already bijective and each stalk map $\psi_x$ already an isomorphism, $\psi$ is bijective. As $\psi$ is a continuous bijection inducing the homeomorphism $\psi_0$ on Kolmogorov quotients, (4.1.38) shows $\psi$ is a homeomorphism.

As $\psi^\sharp$ is an isomorphism on a basis of open sets, it is an isomorphism of sheaves. Therefore $U$ is isomorphic to $Z$ and affine as required. ∎

© David Rufino CC BY-NC-ND 4.0

**Definition 6.2.130 (Order of Zero / Poles)**
*Let $X$ be a regular algebraic curve. For every point $x \in X$ there is a well-defined and unique discrete valuation*

$$v_x : k(X)^\star \to \mathbb{Z}$$

*because $\mathcal{O}_{x,X}$ is a discrete valuation ring of $k(X)$ (6.2.124). For $r$ a positive integer we say that $f \in k(X)^\star$ has a **zero** (resp. **pole**) of order $r$ at $x$ if $v_x(f) = r$ (resp. $v_x(f) = -r$).*

*As before define the **principal divisor***

$$(f) = \sum_{x \in X_0} v_x(f)x$$

*on the free abelian group indexed by $X_0$.*

**Definition 6.2.131**
*Let $\phi : X \to Y$ be a morphism of regular curves and $\phi_\star : k(Y) \hookrightarrow k(X)$ the correspond morphism of fields. Recall that for $x \in X$ that $\phi_\star$ restricts to a **local** homomorphism*

$$\mathcal{O}_{Y,\phi(x)} \to \mathcal{O}_{X,x}$$

*Therefore for $x \in X$ we may define the **ramification index** at $x$ as follows*

$$e_x := v_x(\phi_\star(t)) \geq 1$$

*where $t$ is a uniformiser for the discrete valuation ring $\mathcal{O}_{Y,\phi(x)}$. Similarly the **residue degree** is given by the degree of the field extension*

$$f_x := \left[k\left(\mathfrak{m}_{X,x}\right) : \phi_\star k\left(\mathfrak{m}_{Y,\phi(x)}\right)\right]$$

*Observe that if $x$ and $x'$ are topologically indistinguishable (equivalently Galois conjugate, or equal in $X_0$) then $e_x = e_{x'}$ and $f_x = f_{x'}$.*

*Note when $k$ is algebraically closed then $f_x = 1$ since $k(x), k(y)$ are finite and therefore trivial extensions of $k$.*

**Proposition 6.2.132 (Ramification Formula)**
*Let $\phi : X \to Y$ be a morphism of regular curves which is finite (e.g. $X$ and $Y$ are complete (6.2.129)). For $y \in Y$ then*

$$\deg(\phi) = \sum_{x_0 \in \phi_0^{-1}(\{y_0\})} e_x f_x$$

*Proof.* Let $V$ be an affine neighbourhood of $y$, and define $U = \phi^{-1}(V)$. Then by assumption $U$ is affine and $\phi^\sharp : \mathcal{O}_Y(V) \to \mathcal{O}_X(U)$ is finite (and integral (3.24.4)).

Define $L := k(Y)$, $B := \mathcal{O}_Y(V)$, $K := \phi_\star(k(X))$ and $A := \phi_\star(\mathcal{O}_X(U))$, where $\phi_\star : k(X) \hookrightarrow k(Y)$ is the canonical embedding

Then by (6.2.15) $L$ is the quotient field of $B$, and $K$ is the quotient field of $A$. By (6.2.101) (6.2.54) both $B$ and $A$ are Dedekind domains. $B$ is finite and integral over $A$, and therefore $B$ is the integral closure of $A$, being integrally closed. Therefore we are in the situation of (3.32.10).

By (6.2.21) there is a bijection between the maximal ideals of $B$ (resp. $A$) and the points of $U_0$ (resp. $V_0$). And by (...) $\phi_0(x_0) = y_0 \iff \mathfrak{M}_{U,x} \cap A = \mathfrak{M}_{V,y}$.

By (3.32.15) the notion of ramification index and residue degree coincide with the same notions for Dedekind domains. Therefore the result follows from the corresponding result for finite morphisms of Dedekind domains (3.32.16). ∎

**Proposition 6.2.133 (Degree of Principal Divisor is Zero)**
*Let $X$ be a complete regular curve and $f \in k(X)$ a non-constant rational function, then*

$$\deg((f)) = 0$$

*i.e. the degree of a principal divisor is zero.*

*Proof.* By (6.2.128) $f$ corresponds to a non-constant and therefore surjective regular morphism $\phi_f : X \to \mathbb{P}^1_k \xrightarrow{\sim} \mathbb{A}^1_k \cup \{\infty\}$. From the construction we see that $v_x(f) > 0$ if and only if $\phi_f(x) = 0$ and $v_x(f) < 0$ if and only if $\phi_f(x) = \infty$. Therefore by definition

© David Rufino CC BY-NC-ND 4.0

$$(f) = \sum_{\substack{x_0 \in X_0 \\ \phi(x_0)=0}} v_{x_0}(f)x_0 + \sum_{\substack{x_0 \in X_0 \\ \phi(x_0)=\infty}} v_{x_0}(f)x_0$$

We have canonical isomorphism $\mathcal{O}_{\mathbb{P}^1_k,0} \xrightarrow{\sim} k[X]_{(X)}$ and when $\phi_f(x) = 0$ the corresponding morphisms of stalks is given by $X \to f_x$, therefore the ramification index is $e_x = v_x(f)$. Similarly when $\phi_f(x) = \infty$ we have $e_x = v_x(f^{-1}) = -v_x(f)$. It's evident that $k\left(\mathfrak{m}_{\mathbb{P}^1_k,0}\right) = k$, whence $f_{x_0} = [k(x_0) : k]$. Therefore by definition of degree

$$\deg((f)) = \sum_{\substack{x_0 \in X_0 \\ \phi(x_0)=0}} e_{x_0} f_{x_0} - \sum_{\substack{x_0 \in X_0 \\ \phi(x_0)=\infty}} e_{x_0} f_{x_0}$$

As $X$ is complete, and so is $\mathbb{P}^1_k$ (6.2.95) then $\phi_f$ is finite by (6.2.129) and we are done by (6.2.132) since both sums are equal to $\deg(\phi_f)$. ■

© David Rufino CC BY-NC-ND 4.0

# 6.3 Function Fields and the Riemann-Roch Theorem

We develop the theory of "function fields", closely related to the notion of complete (or projective) regular algebraic curves (...).

## 6.3.1 Relation to the Literature

These treatments fall into two methodological groups:

- Weil's répartition/adele method, in the tradition initiated by Chevalley [Che51]: used by Chevalley himself, by Stichtenoth [Sti09], by Lang [Lan82] (all three over a general constant field), and, at its computational core, even by Serre [Ser59] – whose own bibliographic note attributes the proof to Weil, though he additionally erects the formalism of sheaf cohomology around it, identifying $H^1(X, \mathcal{O}_X)$ with the classes of repartitions. Lang's book, unlike the others in this group, has a different overall aim: a bridge to the classical complex-analytic theory of abelian and theta functions, which the algebraic function-field theory here does not pursue.
- the classical algebraic-geometric (scheme-free, ringed-space) approach shared by Hartshorne [Har13] and Bump [Bum98], the latter specialised to curves with a dedicated treatment of differentials, residues and ramification.

I have interpolated between these, guided by four principles.

**Dedekind domains over bare valuation theory.** I use Dedekind domain theory for the key results on function fields and the Weil form of Riemann-Roch, rather than the purely valuation-theoretic development of [Che51] and [Sti09]. In the same spirit, finite extensions of $k((t))$ (3.33.4) are analysed directly via the integral closure in a Dedekind domain and a direct power-series computation, rather than via the $P$-adic completion machinery both of those references build on; this also yields the result in every characteristic, where [Che51]'s analogous Puiseux-expansion treatment (Ch. II, §6) is restricted to the tame case, as noted in the remark opening the Extensions of Power Series Rings section. We also use the theory of affine algebras to simplify the proof of some further technical results.

**A general constant field, reduced to the algebraically closed case only once.** Riemann-Roch and the Residue Theorem are proved over a general constant field, matching Riemann-Roch (though not the Residue Theorem, which is stated only over an algebraically closed field) in [Sti09], but going beyond [Ser59], [Har13], [Bum98] and [Lan82], whose treatments of Riemann-Roch (and, for [Bum98], the Residue Theorem too) all assume an algebraically closed constant field throughout. The general-field Residue Theorem ((6.3.57)) is not proved directly: it is deduced from the algebraically closed case ((6.3.34)) by base extension to $\overline{k}$ and a Galois-descent argument summing conjugate residues, rather than by redoing the local theory over a general field from scratch.

**Concrete constructions over general machinery, wherever one suffices.** Like [Che51], [Sti09] and [Lan82], I work with Chevalley and Weil's répartition/adele-style methods directly, and connect this to cohomology only after the fact, rather than building up the general machinery of coherent sheaf cohomology as [Ser59] does around the same underlying répartition computation. Likewise, the Residue Theorem is proved without recourse to local completions, unlike the concrete, differential-level residue theorems of [Ser59] and [Sti09], which both depend on the $P$-adic completion; for the rational function field this is done via a short, self-contained elementary proof by partial fractions, rather than as a special case of the general machinery. The Trace Formula for Residues ((6.3.42)) likewise avoids completions, via the same Laurent-series finite-extension lemma above, rather than by appeal to the Complete Discrete Valuation Rings machinery developed alongside it (the Extensions of Power Series Rings and Complete Discrete Valuation Rings sections). And residues are defined directly from the uniformiser derivation without ever constructing the module of Kähler differentials $\Omega_{K/k}$, in contrast to [Ser59] and [Sti09], which both identify their (Weil-differential-based) residues with genuine differentials in this sense.

**Characteristic $p$ handled uniformly, via universal $\mathbb{Z}$-polynomial identities.** Invariance of the residue of a differential under change of uniformiser – notoriously tricky in characteristic $p$, where the naive derivative-of-a-power argument breaks down whenever $p$ divides the relevant exponent – is settled once, universally, over $\mathbb{Z}[a_1, a_1^{-1}, a_2, a_3, \ldots]$ and specialised ((6.3.26)), the same device used by [Ser59] and [Sti09], rather than the Cartier-operator route of [Bum98] unlike [Ser59] and [Sti09], this is done over a general constant field rather than an algebraically closed one. The same device is reused for the Trace Formula for Residues, which holds without any tameness hypothesis ((6.3.40)) by the identical method [Ser59] uses (there attributed to Hasse) for the analogous local identity $\mathrm{Res}_t(f\,du) = \mathrm{Res}_u(\mathrm{Tr}(f)\,du)$ underlying its own residue formula; it is exactly this universal-identity device that lets the wildly ramified case go through in both results.

Two further points of departure concern scope and connections to the classical picture, rather than proof technique: the dictionary between the function field and the regular algebraic curve – established even over a non-algebraically closed

© David Rufino CC BY-NC-ND 4.0

field, generalising the algebraically-closed-field statement of [Har13] – is proved later, in (6.2.127) of the Algebraic Geometry chapter's treatment of regular algebraic curves; and, in the classical case of a compact Riemann surface ((6.3.19)), Weil differentials are shown to coincide canonically with classical meromorphic 1-forms, reconnecting the algebraic formalism with the analytic theory it generalises.

## 6.3.2 Riemann-Roch Theorem

In what follows we generally assume that $\tilde{k} = k$ (i.e. $K$ is geometrically integral).

**Definition 6.3.1 (Divisors)**
*Let $K/k$ be an algebraic function field. Denote by $\mathrm{Div}(K)$ the free abelian group generated by the places of $K/k$ (**divisor group**). The elements are written as a formal sum*

$$D = \sum_P n_P P$$

*with all but finitely many coefficients $n_P = 0$. Define the **degree** homomorphism*

$$\begin{aligned} \mathrm{Div}(K) &\longrightarrow \mathbb{Z} \\ \sum_P n_P P &\longrightarrow \sum_P n_P \cdot [k(\mathfrak{m}_P) : k] \end{aligned}$$

*We may also define $v_P(D) \coloneqq n_P$, and the **support** of $D$ by $\mathrm{Supp}(D) \coloneqq \{P \mid v_P(D) \neq 0\}$, a finite set. Finally we may define a partial order*

$$D \leq D' \iff v_P(D) \leq v_P(D') \quad \forall P$$

**Definition 6.3.2 (Riemann-Roch Space)**
*Let $K/k$ be an algebraic function field and $D$ a divisor. Define the **Riemann-Roch Space** associated to $D$ by*

$$L(D) \coloneqq \{x \in K^\star \mid (x) \geq -D\} \cup \{0\}$$

*and define*

$$\ell(D) \coloneqq \dim_k L(D)$$

*(we shortly show it is a $k$-vector subspace of $K$)*

**Definition 6.3.3 (Principal Divisors)**
*Let $K/k$ be an algebraic function field and $x \in K^\star$. Then define the **zero divisor** (resp. **pole divisor**, **principal divisor**) of $x$ as follows*

$$\begin{aligned} (x)_0 &\coloneqq \sum_{P : v_P(x) > 0} v_P(x) P, \\ (x)_\infty &\coloneqq \sum_{P : v_P(x) < 0} (-v_P(x)) P, \\ (x) &\coloneqq (x)_0 - (x)_\infty = \sum_P v_P(x) P. \end{aligned}$$

*which is well-defined by (6.2.116).*

*Observe similarly $x \in k^\star \iff (x) = 0$. Define the subgroup of principal divisors by*

$$\mathrm{Princ}(K) \coloneqq \{(x) \mid x \in K\}$$

*and the quotient group $\mathrm{Cl}(K) \coloneqq \mathrm{Div}(K)/\mathrm{Princ}(K)$ is called the **divisor class group**. We say two divisors $D$ and $D'$ are **linearly equivalent** if there exists $x \in K$ such that $D - D' = (x)$, equivalently if they become equal in the divisor class group. This is written $D \sim D'$.*

We presently prove that these concepts are well-defined, namely every function $x \in K^\star$ has only finitely many zeros. More precisely we can show

**Proposition 6.3.4 (Principal Divisors have degree zero)**
*Let $K/k$ be an algebraic function field and $x \in K \setminus \tilde{k}$ transcendental. Then*

$$\begin{aligned} \deg((x)_0) = \deg((x)_\infty) &= [K : k(x)] \\ \deg((x)) &= 0 \end{aligned}$$

© David Rufino CC BY-NC-ND 4.0

*In particular there is a well-defined homomorphism*

$$\deg : \mathrm{Cl}(K) \longrightarrow \mathbb{Z}$$

*Proof.* Recall (6.2.114) $k[x]$ is a Dedekind domain, $\overline{k[x]}$ is a finite $k[x]$-module and every place $\mathcal{O}_P$ containing $x$ corresponds to prime ideal $\mathfrak{B}$ of $\overline{k[x]}$. By (3.32.15) then $v_P(x)$ is equal to the ramification index $e_{\mathfrak{B}/(x)}$, and the inertial degree is $f_{\mathfrak{B}/(x)} = [k(\mathfrak{m}_P) : k]$ where $(x)$ is a prime ideal of $k[x]$. From (3.35.61) we have the ramification formula

$$[K : k(x)] = \sum_{\mathfrak{B}|(x)} e_{\mathfrak{B}/(x)} f_{\mathfrak{B}/(x)}$$

Therefore $\deg((x)_0) = [K : k(x)]$ and $\deg((x)_\infty) = \deg\left((x^{-1})_0\right) = [K : k(x^{-1})] = [K : k(x)]$. ∎

**Proposition 6.3.5**

*Let $K/k$ be an algebraic function field and $D$ a divisor. Then*

*a) $L(D)$ is a $k$-vector subspace of $K$*
*b) If $D \sim D'$ then $L(D) = L(D')$*
*c) $L(0) = \tilde{k}$ so under our standing assumption $\ell(0) = 1$.*
*d) If $D < 0$ then $L(D) = \{0\}$*

*Proof.*

a) Given $x, y \in L(D)$ then by the strict triangle inequality for valuations $v_P(x+y) \geq \min(v_P(x), v_P(y)) \geq v_P(D)$. Similarly for $\lambda \in k$ we have $v_P(\lambda x) = v_P(\lambda) + v_P(x) = v_P(x)$ (6.2.114).

b) By definition $D' = D + (x)$ we may define a mapping

$$\begin{aligned} L(D) &\longrightarrow L(D') \\ y &\longrightarrow yx \end{aligned}$$

Then

$$v_P(yx) = v_P(x) + v_P(y) \geq v_P(x) + v_P(D) = v_P(D')$$

so this is well-defined. As $x^{-1}$ defines the inverse map it is a bijection.

c) By (6.2.114).a) $\tilde{k} \subset L(0)$. Suppose $x \in K \setminus \tilde{k}$ then by definition $x$ is transcendental and so is $x^{-1}$. By (6.2.116) $x^{-1}$ has a zero $P$, i.e. $v_P(x^{-1}) > 0 \Longrightarrow v_P(x) < 0$. Therefore by definition $x \notin L(0)$ as required.

d) If $x \in \tilde{k}^\star$ then $(x) = 0$ whence $x \notin L(D)$. If $x \in K \setminus \tilde{k}$ then as before $x$ has a pole therefore $x \notin L(D)$ as required.

∎

**Lemma 6.3.6**

*Let $K/k$ be an algebraic function field and $D \leq D'$ divisors. Then $L(D) \subset L(D')$ and*

$$\dim_k L(D')/L(D) \leq \deg(D') - \deg(D)$$

*In particular $\ell(D)$ is finite and we have*

$$\deg(D) - \ell(D) \leq \deg(D') - \ell(D')$$

*Proof.* Consider a divisor $D$ and suppose $D' := D + P$ for some $P$ possibly in the support of $D$. Let $\pi$ be a uniformiser for $\mathcal{O}_P$ and define $t = \pi^{v_P(D')}$. We then have a well-defined map

$$\begin{aligned} \psi : L(D') &\longrightarrow k(\mathfrak{m}_P) \\ x &\longrightarrow xt + \mathfrak{m}_P \end{aligned}$$

Observe that $\psi(x) = 0 \iff xt \in \mathfrak{m}_P \iff v_P(xt) > 0 \iff v_P(x) + v_P(D') > 0 \iff v_P(x) \geq -v_P(D) \iff x \in L(D)$. Therefore $\ker(\psi) = L(D)$ and so

$$\dim_k(L(D')/L(D)) \leq [k(\mathfrak{m}_P) : k] = \deg(D') - \deg(D)$$

Now suppose $D'' = D' + Q$, then by the isomorphism theorem for vector spaces (...)

$$\begin{aligned} \dim_k(L(D'')/L(D)) &= \dim_k(L(D'')/L(D')) + \dim_k(L(D')/L(D)) \\ &\leq \deg(D'') - \deg(D') + \deg(D') - \deg(D) \\ &= \deg(D'') - \deg(D) \end{aligned}$$

from which it follows the general case. ∎

© David Rufino CC BY-NC-ND 4.0

**Proposition 6.3.7 (Dimension Upper Bound)**
*Let $K/k$ be an algebraic function field and $D$ a divisor. Then*

$$\ell(D) \leq \deg(D_+) + \dim_k \tilde{k}$$

*where $D_+$ is the positive part of $D$.*

*Proof.* $L(D) \subset L(D_+)$ so it is enough to show the result for positive divisors. As $L(0) = \tilde{k}$ (...) the result follows from (6.3.6). ∎

We say that the function $\deg(D) - \ell(D)$ is increasing, we also show it is bounded above.

**Lemma 6.3.8 (Dimension Lower Bound I)**
*Let $K/k$ be an algebraic function field. There exists a positive divisor $B$ and integer $\gamma \in \mathbb{Z}$ such that*

$$\deg(mB) - \ell(mB) \leq \gamma$$

*for all positive integers $m$.*

*Proof.* Let $x \in K$ be transcendental and define $B \coloneqq (x)_\infty$ with $\deg(B) = n$. It's positive by (...). Let $u_1, ..., u_n$ be a $k(x)$-basis for $K$ and a divisor $C \geq 0$ such that $(u_i) \geq -C$. Then we claim that for all integers $m$ the elements $\left\{x^i u_j \;\middle|\; 0 \leq i \leq m, 1 \leq j \leq n\right\}$ are linearly independent over $k$. But this follows immediately because suppose

$$\sum_{i,j} \lambda_{ij} x^i u_j = 0$$

then

$$\sum_{j=1}^{n} \left( \sum_{i=0}^{m} \lambda_{ij} x^i \right) u_j = 0$$

and by linear independence we have $\lambda_{ij} = 0$.

Further $x^i \in L(iB) \implies x^i \in L(mB)$ and $u_j \in L(C) \implies x^i u_j \in L(mB + C)$. Therefore $\ell(mB + C) \geq n(m+1) = \deg(mB) + n$ for all positive integers $m$.

On the other hand by (6.3.6) we have $\ell(mB + C) \leq \ell(mB) + \deg(C)$. Therefore

$$\deg(mB) - \ell(mB) \leq \deg(C) - n$$

∎

We may extend this result to all divisors by way of the following lemma.

**Lemma 6.3.9**
*Let $K/k$ be an algebraic function field, $B$ a divisor satisfying the conclusion of (6.3.8) and $D$ an arbitrary divisor. Then there exists divisors $D', D''$ such that $D \leq D'$, $D' \sim D''$ and $D'' \leq mB$ for some $m \geq 0$.*

*Proof.* Choose a positive divisor $D' \geq D$, then

$$\deg(mB - D') - \ell(mB - D') \leq \deg(mB) - \ell(mB) \leq \gamma$$

for all positive integers $m$. This means

$$\ell(mB - D') \geq m \deg(B) - \deg(D') - \gamma$$

which is positive for sufficiently large $m$. Therefore we may choose $0 \neq z \in L(mB - D')$. Let $D'' = D' - (z)$. By definition

$$(z) \geq D' - mB \implies D'' \leq mB$$

as required. ∎

**Proposition 6.3.10 (Genus is well-defined)**
*Let $K/k$ be an algebraic function field. Then there exists a positive integer $\gamma$ such that*

$$\deg(D) - \ell(D) \leq \gamma$$

*for all divisors $D$.*

*Proof.* Then using the notation of (6.3.9) we have

© David Rufino CC BY-NC-ND 4.0

$$\begin{aligned}\deg(D) - \ell(D) &\overset{(6.3.6)}{\leq} \deg(D') - \ell(D') \\ &\overset{(6.3.4)\,(6.3.5)}{=} \deg(D'') - \ell(D'') \\ &\overset{(6.3.6)}{\leq} \deg(mB) - \ell(mB) \\ &\overset{(6.3.8)}{\leq} \gamma\end{aligned}$$

■

**Definition 6.3.11 (Genus of a Function Field)**
*Let $K/k$ be an algebraic function field with $\tilde{k} = k$. Define the **genus** by*

$$g := \max\{\deg(D) - \ell(D) + 1 \mid D \in \mathrm{Div}(K)\}$$

*Note as $\ell(0) = 1$ the genus $g$ is non-negative. For any divisor $D \in \mathrm{Div}(K)$ define the **index of specialty** as follows*

$$i(D) := \ell(D) - \deg(D) + g - 1$$

*which by definition is a non-negative integer. This satisfies a few key properties (6.3.6)*

*a)* $i(0) = g$
*b)* $i(D) \geq 0$
*c)* $D \sim D' \implies i(D) = i(D')$
*d)* $D' \geq D \implies i(D') \leq i(D)$

*The Riemann-Roch theorem then takes the following form*

$$\ell(D) - i(D) = \deg(D) - g + 1$$

*We say that a divisor $D$ is **non-special** if $i(D) = 0$ (equivalently $\ell(D) = \deg(D) - g + 1$).*

The full Riemann-Roch amounts to an explicit form of $i(D)$. In order to do this we identify it as the dimension of a certain vector space.

**Definition 6.3.12 (Repartitions)**
*Let $K/k$ be an algebraic function field. Define the $K$-algebra of **repartitions** of $K$ by*

$$\mathcal{A}_K := \left\{ (\alpha_P)_{P \in \mathbb{P}_K} \;\middle|\; \alpha_P \in K \text{ and } \alpha_P \in \mathcal{O}_P \text{ for all but finitely many places } P \right\}$$

*For every divisor $D \in \mathrm{Div}(K)$ define the $K$-subspace*

$$\mathcal{A}_K(D) := \{\alpha \in \mathcal{A}_K \mid v_P(\alpha_P) \geq -v_P(D) \text{ for all } P \in \mathbb{P}_K\}$$

*There is a canonical $K$-algebra homomorphism*

$$K \hookrightarrow \mathcal{A}_K$$

*which is well-defined by (...). Then clearly we have $K \cap \mathcal{A}_K(D) = L(D)$.*

We prove a number of results

**Lemma 6.3.13**
*Let $K/k$ be an algebraic function field. For divisors $D, D'$ such that $D \leq D'$ we have*

*a)* $\dim_k \mathcal{A}_K(D')/\mathcal{A}_K(D) = \deg(D') - \deg(D)$
*b)* $\dim_k((\mathcal{A}_K(D') + K)/(\mathcal{A}_K(D) + K)) = i(D) - i(D')$

*Proof.* We prove each in turn.

**a)** By the isomorphism theorem for $k$-modules it's enough to demonstrate this for the case $D' = D + P$. Let $\pi$ be a uniformiser for $\mathcal{O}_P$ and let $r := v_P(D) + 1 = v_P(D')$. Then define the $k$-linear map

$$\begin{aligned}\mathcal{A}_K(D') &\longrightarrow k(\mathfrak{m}_P) \\ \alpha &\longrightarrow (\pi^r \alpha_P)(P)\end{aligned}$$

By definition we have $v_P(\alpha_P) \geq -v_P(D') \implies v_P(\pi^r \alpha_P) \geq 0 \implies \pi^r \alpha_P \in \mathcal{O}_P$ so that this map is well-defined. Further $\pi^r \alpha_P \in \mathfrak{m}_P \iff v_P(\pi^r \alpha_P) > 0 \iff v_P(D) + 1 + v_P(\alpha_P) > 0 \iff v_P(\alpha_P) \geq -v_P(D)$ whence the kernel is precisely $\mathcal{A}_K(D)$. It is clearly surjective because given $f \in \mathcal{O}_P$ we may consider the repartition given by

© David Rufino CC BY-NC-ND 4.0

$$(1, ..., 1, \pi^{-r} f, 1, ...)$$

which clearly maps to $f$ under this map. Therefore by the isomorphism theorem we have

$$\dim_k \mathcal{A}_K(D')/\mathcal{A}_K(D) = [k(\mathfrak{m}_P) : k] = \deg(D') - \deg(D)$$

**b)** We claim there is an exact sequence of $k$-modules

$$0 \longrightarrow L(D')/L(D) \overset{\sigma_1}{\to} \mathcal{A}_K(D')/\mathcal{A}_K(D) \overset{\sigma_2}{\to} (\mathcal{A}_K(D') + K)/(\mathcal{A}_K(D) + K) \longrightarrow 0$$

from which the result follows immediately by comparing dimensions. The first map is given by

$$\sigma_1([f]) \coloneqq (v_P(f))_{P \in \mathbb{P}_K}$$

which is obviously well-defined and injective. The map $\sigma_2$ is clearly well-defined and surjective. Further it is evident that $\sigma_2 \circ \sigma_1 = 0$. Therefore it only remains to show that $\ker(\sigma_2) \subset \operatorname{im}(\sigma_1)$. Suppose $\alpha \in \mathcal{A}_K(D')$ then $\sigma_2(\alpha) = 0 \implies \alpha - (f) \in \mathcal{A}_K(D) \implies (f) \in \mathcal{A}_K(D')$ for some $f \in K$. Then by definition this means $f \in L(D')$ and $\sigma_1(f) = \alpha$ as required. ∎

**Proposition 6.3.14 (Riemann's Theorem)**
*Let $K/k$ be an algebraic function field. Suppose $D_0$ is **non-special** (there always exists such a divisor) then*

*a)* $\deg(D) > \ell(D_0) \implies D$ *is non-special*
*b)* $\mathcal{A}_K = \mathcal{A}_K(D_0) + K$

*Proof.* We prove each in turn

a) For an arbitrary divisor $D$ we have

$$\begin{aligned} \ell(D - D_0) &= i(D - D_0) + \deg(D - D_0) - g + 1 \\ &\geq 0 + \deg(D) - \ell(D_0) \end{aligned}$$

So when $\deg(D) > \ell(D_0)$ we have $\ell(D - D_0) > 0$. Choose $0 \neq f \in L(D - D_0)$. Then we may define $D' \coloneqq D + (f) \geq D_0$. Then by (...) $i(D) = i(D') \leq i(D_0) = 0$ whence $i(D) = 0$.

b) Given $\alpha \in \mathcal{A}_K$ choose $D \geq D_0$ such that $\alpha \in \mathcal{A}_K(D)$. Then by (6.3.11) $i(D) \leq i(D_0) = 0$ and therefore $i(D) = 0$. By (6.3.13).b) $\mathcal{A}_K(D) + K = \mathcal{A}_K(D_0) + K \implies \alpha \in \mathcal{A}_K(D_0) + K$ as required.

∎

**Proposition 6.3.15 (Explicit form of $i(D)$)**
*Let $K/k$ be an algebraic function field and $D \in \operatorname{Div}(K)$. Then*

$$i(D) = \dim_k(\mathcal{A}_K/(\mathcal{A}_K(D) + K))$$

*Proof.* By (6.3.14) there exists some non-special divisor $D' \geq D$ such that $i(D') = 0$ and $\mathcal{A}_K = \mathcal{A}_K(D') + K$. The formula then follows from (6.3.13).b). ∎

**Definition 6.3.16 (Weil Divisors)**
*Let $K/k$ be an algebraic function field and $D \in \operatorname{Div}(K)$. Define*

$$\Omega_K(D) \coloneqq \{\omega \in \operatorname{Hom}_k(\mathcal{A}_K, k) \mid \mathcal{A}_K(D) + K \subset \ker(\omega)\}$$

*As divisors form a directed system we may form the union of **Weil Divisors***

$$\Omega_K \coloneqq \bigcup_{D \in \operatorname{Div}(K)} \Omega_K(D) \subset \operatorname{Hom}_k(\mathcal{A}_K, k)$$

**Proposition 6.3.17**
*Let $K/k$ be an algebraic function field. Then there is a canonical isomorphism*

$$\operatorname{Hom}_k(\mathcal{A}_K/(\mathcal{A}_K(D) + K), k) \xrightarrow{\sim} \Omega_K(D)$$

*In particular $i(D) = \dim_k \Omega_K(D)$. Taking $D = 0$, since $i(0) = g$ (6.3.11) we get*

$$\dim_k \Omega_K(0) = g$$

*that is, the genus equals the dimension of the space of everywhere-regular Weil differentials.*

*Proof.* This is just the isomorphism theorem for vector spaces. The statement on dimensions follows from (6.3.15) and (3.3.109). ∎

© David Rufino CC BY-NC-ND 4.0

### 6.3.3 Meromorphic Differentials in the case $k = \mathbb{C}$

In this section we aim to provide a post-hoc motivation for the introduction of Weil Differentials, by showing that they are canonically isomorphic to the classical space of Meromorphic Differentials in the case $k = \mathbb{C}$. This relies on some techniques which are specific to the complex-analytic structure of Riemann Surfaces.

**Proposition 6.3.18 (Residue Theorem for $k = \mathbb{C}$)**
*Let $p$ be a point of a Riemann surface and $\pi$ a local holomorphic coordinate centered at $p$. For $f$ meromorphic near $p$, define*

$$\mathrm{Res}_\pi(f) := \mathrm{Res}_p(f\,\mathrm{d}\pi) := \frac{1}{2\pi i}\oint_\gamma f\,\mathrm{d}\pi$$

*for a small loop $\gamma$ around $p$ – the classical residue of the 1-form $f\,\mathrm{d}\pi$. Writing $g := f \circ \psi$ for the re-expansion of $f$ relative to $\pi' = \psi^{-1}(\pi)$, and $f = \sum_n a_n \pi^n$ for the Laurent series expansion of $f$ at $p$, then*

*a) $\mathrm{Res}_\pi$ is $k$-linear in $f$;*
*b) $\mathrm{Res}_\pi(f) = 0$ whenever $f$ is holomorphic at $p$ (i.e. has no pole there);*
*c) $\mathrm{Res}_\pi(f) = a_{-1}$;*
*d) $\mathrm{Res}_{\pi'}(g \cdot \psi') = \mathrm{Res}_\pi(f)$;*
*e) if $p$ ranges over a* compact *Riemann surface $X$ and $\omega$ is a meromorphic* 1*-form on $X$ (so has only finitely many poles), then, writing $\omega_\pi$ for the coefficient of $\omega$ relative to a local coordinate $\pi$ at $p$ (so $\omega = \omega_\pi\,\mathrm{d}\pi$ near $p$, and $\mathrm{Res}_\pi(\omega_\pi) = \mathrm{Res}_p(\omega)$ by definition – independent of the choice of $\pi$ at each $p$, by d)),*

$$\sum_p \mathrm{Res}_\pi(\omega_\pi) = 0$$

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

**a).** Immediate from linearity of the contour integral, $\oint_\gamma (\alpha f + \beta h)\,\mathrm{d}\pi = \alpha \oint_\gamma f\,\mathrm{d}\pi + \beta \oint_\gamma h\,\mathrm{d}\pi$ for $\alpha, \beta \in k$.

**b).** Immediate from c) below: $f$ holomorphic at $p$ means $a_n = 0$ for every $n < 0$, in particular $a_{-1} = 0$, so $\mathrm{Res}_\pi(f) = a_{-1} = 0$.

**c).** Term-by-term integration (valid by uniform convergence of the Laurent series on $\gamma$) gives $\oint_\gamma f\,\mathrm{d}\pi = \sum_n a_n \oint_\gamma \pi^n\,\mathrm{d}\pi$, and $\oint_\gamma \pi^n\,\mathrm{d}\pi = 2\pi i \delta_{n,-1}$ ($n \neq -1$: exact, $\pi^n\,\mathrm{d}\pi = \mathrm{d}\left(\frac{\pi^{n+1}}{n+1}\right)$, so integrates to 0 around a closed loop; $n = -1$: winding number 1 around $\pi = 0$), so $\mathrm{Res}_\pi(f) = \frac{1}{2\pi i}\oint_\gamma f\,\mathrm{d}\pi = a_{-1}$.

**d).** By the chain rule, $f\,\mathrm{d}\pi = f(\psi(\pi'))\psi'(\pi')\,\mathrm{d}\pi' = (g \cdot \psi')\,\mathrm{d}\pi'$: these are the same 1-form $\omega$, written in the two coordinates $\pi$ and $\pi'$. The contour integral $\oint_\gamma \omega$ depends only on $\omega$ and $\gamma$, not on the coordinate used to write it down – reparametrising $\gamma$ by $\pi'$ instead of $\pi$ changes nothing about which loop is being integrated over – so $\mathrm{Res}_p(\omega)$ computed from either expression gives the same number:

$$\mathrm{Res}_{\pi'}(g \cdot \psi') = \mathrm{Res}_p((g \cdot \psi')\,\mathrm{d}\pi') = \mathrm{Res}_p(\omega) = \mathrm{Res}_p(f\,\mathrm{d}\pi) = \mathrm{Res}_\pi(f)$$

**e).** This uses two facts special to the compact setting, neither purely local (unlike a)–d) above): $\omega$ is **closed** away from its poles ($\omega = f\,\mathrm{d}z$ locally with $f$ holomorphic, so $\mathrm{d}\omega = \frac{\partial f}{\partial \overline{z}}\,\mathrm{d}\overline{z} \wedge \mathrm{d}z = 0$ by the Cauchy–Riemann equations), and $X$ is **compact**. Removing a small disk $D_p$ around each pole leaves $X_0 := X \setminus \bigcup_p D_p$, on which $\omega$ is holomorphic, so $\mathrm{d}\omega = 0$ there; Stokes' theorem gives

$$0 = \int_{X_0} \mathrm{d}\omega = \int_{\partial X_0} \omega = -\sum_p \oint_{\partial D_p} \omega = -\sum_p 2\pi i\,\mathrm{Res}_\pi(\omega_\pi)$$

(the sign, as $\partial X_0$ traverses each $\partial D_p$ oppositely to the usual counterclockwise orientation used to define $\mathrm{Res}_\pi$, and $\oint_{\partial D_p} \omega = 2\pi i\,\mathrm{Res}_\pi(\omega_\pi)$ by definition), giving $\sum_p \mathrm{Res}_\pi(\omega_\pi) = 0$. ∎

**Proposition 6.3.19 (Classical Differentials are Weil Differentials)**
*Let $X$ be a compact Riemann surface with function field $K$ (so $k = \mathbb{C}$), and let $\mathcal{M}^1(X)$ denote the $\mathbb{C}$-vector space of meromorphic* 1*-forms on $X$. For $\omega \in \mathcal{M}^1(X)$, define $\varphi_\omega \in \mathrm{Hom}_k(\mathcal{A}_K, k)$ by*

$$\varphi_\omega(\alpha) := \sum_P \mathrm{Res}_\pi(\alpha_P \cdot \omega_\pi), \quad \alpha = (\alpha_P)_P \in \mathcal{A}_K$$

*(a finite sum, by b): $\alpha_P \omega_\pi$ is holomorphic, hence contributes 0, at all but finitely many $P$). Then $\varphi_\omega \in \Omega_K$, and*

$$\mathcal{M}^1(X) \longrightarrow \Omega_K, \quad \omega \mapsto \varphi_\omega$$

© David Rufino CC BY-NC-ND 4.0

*is a canonical $\mathbb{C}$-linear isomorphism.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

**$\varphi_\omega \in \Omega_K$.** Let $D := (\omega)$, the divisor of $\omega$ (well-defined: $v_P(\omega_\pi)$ is independent of the choice of $\pi$ at $P$, as $\omega_{\pi'} = (\omega_\pi \circ \psi)\cdot\psi'$ by the chain rule – the same identity underlying d) – and $\psi'$ is a unit, so composition with $\psi$ and multiplication by $\psi'$ both preserve order of vanishing). We show $\mathcal{A}_K(D) + K \subset \ker(\varphi_\omega)$. For $\alpha \in \mathcal{A}_K(D)$, $v_P(\alpha_P) \geq -v_P(D) = -v_P(\omega_\pi)$ for every $P$, so $v_P(\alpha_P\omega_\pi) \geq 0$: each $\alpha_P\omega_\pi$ is holomorphic at $P$, so $\mathrm{Res}_\pi(\alpha_P\omega_\pi) = 0$ by b), giving $\varphi_\omega(\alpha) = 0$. For $f \in K$ (embedded diagonally, $\alpha_P = f$ for every $P$), $\varphi_\omega(f, f, \ldots) = \sum_P \mathrm{Res}_\pi(f\cdot\omega_\pi) = \sum_P \mathrm{Res}_\pi((f\omega)_\pi) = 0$ by e) (the Residue Theorem), applied to the meromorphic 1-form $f\omega \in \mathcal{M}^1(X)$. So $\varphi_\omega$ vanishes on $\mathcal{A}_K(D) + K$, that is $\varphi_\omega \in \Omega_K(D) \subset \Omega_K$.

**$\mathbb{C}$-linearity.** Immediate from a): $\varphi_{\lambda\omega_1+\mu\omega_2}(\alpha) = \sum_P \mathrm{Res}_\pi\big(\alpha_P(\lambda\omega_1 + \mu\omega_2)_\pi\big) = \lambda\varphi_{\omega_1}(\alpha) + \mu\varphi_{\omega_2}(\alpha)$ for $\lambda, \mu \in \mathbb{C}$, using that $(\lambda\omega_1 + \mu\omega_2)_\pi = \lambda(\omega_1)_\pi + \mu(\omega_2)_\pi$ (coefficients relative to a fixed $\pi$ add linearly) and linearity of $\mathrm{Res}_\pi$ in its argument.

**Injectivity.** Suppose $\varphi_\omega = 0$; we show $\omega = 0$. Fix $P$ and, for $n \in \mathbb{Z}$, let $\alpha^{(n)} \in \mathcal{A}_K$ be the repartition with $\alpha^{(n)}_P := \pi^{-n}$ and $\alpha^{(n)}_Q := 0$ for $Q \neq P$ (a valid repartition: it is holomorphic away from the single place $P$). Writing $\omega_\pi = \sum_m a_m\pi^m$, $0 = \varphi_\omega(\alpha^{(n)}) = \mathrm{Res}_\pi(\pi^{-n}\omega_\pi) = a_{n-1}$ by c). As $n$ was arbitrary, every coefficient of $\omega_\pi$ vanishes, so $\omega$ is 0 near $P$; as $P$ was arbitrary, $\omega \equiv 0$ on $X$.

**Surjectivity.** *Sketch.* This needs the classical Mittag-Leffler theorem for meromorphic differentials: given principal parts (polar Laurent tails) prescribed at finitely many points of $X$, satisfying the residue-sum-zero constraint of e), there exists a global meromorphic 1-form on $X$ realising them – solved via the $\overline{\partial}$-equation (Hodge theory / finite-dimensionality of $H^1(X, \mathcal{O})$), genuinely global analytic content with no counterpart among a)–e) above. Given $\psi \in \Omega_K(D)$, read off, at each pole $P$ of $D$, the local principal part $\psi$ demands by testing against the single-place repartitions $\alpha^{(n)}_P = \pi^{-n}$ used in the injectivity argument above; since $\psi$ vanishes on $K$ by construction, these prescribed principal parts automatically satisfy the residue-sum-zero compatibility condition Mittag-Leffler requires, and it produces a global $\omega$ realising them, with $\varphi_\omega = \psi$. ∎

## 6.3.4 Derivations on the Function Field

**Proposition 6.3.20 (A Distinguished Basis for $\mathrm{Der}_k(K, K)$)**
*Let $x \in K$ be transcendental over $k$ and assume $k$ is perfect. Then there is a unique $\partial_x \in \mathrm{Der}_k(K, K)$ with $\partial_x(x) = 1$, and $D = D(x)\cdot\partial_x$ for every $D \in \mathrm{Der}_k(K, K)$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

By (3.35.44), $\dim_K \mathrm{Der}_k(K) = 1$. So there is at least one non-zero $D_0$. If $D_0(x) = 0$ then it vanishes on $k(x)$ by (3.26.9), (3.26.10) and therefore on $K$ as $K/k(x)$ is separable (3.19.103) (3.35.42), a contradiction. Therefore $D_0(x) \neq 0$ for any non-zero derivation $D_0$. Therefore define $\partial_x := D_0(x)^{-1}D$ and by monogenicity we are done. ∎

**Proposition 6.3.21 (Uniformisers are Transcendental)**
*Let $P$ be a rational place of $K/k$, $\pi$ a uniformiser. Then $\pi$ is transcendental over $k$.*

*Proof.* By (...) $k \cap \mathfrak{m}_P = (0)$ whence $\pi \notin k$. By assumption $\tilde{k} = k$ whence $\pi$ is transcendental. ∎

**Proposition 6.3.22 (Chain Rule for Uniformiser Derivations)**
*Let $x, x' \in K$ be transcendental. Then*

$$\partial_{x'}(t) = \partial_{x'}(x)\cdot\partial_x(t) \quad \text{for every } t \in K.$$

*Proof.* This is immediate from (6.3.20). ∎

## 6.3.5 Residue Symbol at a Rational Place

The residue symbol $\mathrm{Res}_P(z\,\mathrm{d}t)$ built in this subsection is defined via the Laurent coefficients of $z\partial_\pi(t)$ relative to a choice of uniformiser $\pi$, and shown independent of that choice (6.3.31) – exactly the invariance a genuine differential-form residue ought to have. This invariance is not straightforward in characteristic $p$: it reduces (6.3.31) to computing $(\psi^n\psi')_{-1}$ for the change-of-uniformiser series $\psi$, an identity which the naive derivative-of-a-power argument only settles when $n+1$ is invertible – automatic in characteristic 0, but not when $p \mid (n+1)$. We settle it uniformly in every characteristic by proving the identity once, universally, over $\mathbb{Z}[a_1, a_1^{-1}, a_2, a_3, \ldots]$ (indeterminate coefficients, torsion-free over $\mathbb{Z}$) and specialising (6.3.26) – the same universal-$\mathbb{Z}$-polynomial philosophy used, e.g., in [Sti09] (Chap. 4, Prop. 4.2.9); another route entirely avoiding a characteristic-$p$ detour is via finite-potent operators [Tat68] (§§1–2).

© David Rufino CC BY-NC-ND 4.0

We restrict throughout this subsection to *rational* places $P$, that is $k(\mathfrak{m}_P) = k$ – geometrically, points of the curve valued in the constant field $k$ itself, rather than in some finite extension of it. This is exactly what allows us to embed $K$ in $k((X))$.

**Proposition 6.3.23 (Laurent Coefficients at a Rational Place)**
*Let $P$ be a rational place of $K/k$, $\pi$ a uniformiser of $\mathcal{O}_P$. There is a unique $k$-linear map*

$$L_\pi : K \to k((X))$$

*with $L_\pi(\pi^n) = X^n$ and $v_X(L_\pi(z)) = v_P(z)$ for every $z \in K^\star$ (**$L_\pi$ preserves valuation**; $v_X$ being the valuation of the DVR $k[[X]]$ (3.10.14)).*

*Writing $L_\pi(z) = \sum_j a_j X^j$ for $z \in K^\star$: $a_j = 0$ for $j < v_P(z)$ and $a_{v_P(z)} \neq 0$, and for every $n \geq v_P(z)$,*

$$z - \sum_{j=v_P(z)}^{n} a_j \pi^j \in \mathfrak{m}_P^{n+1}$$

*(the **Laurent coefficients** of $z$ at $P$ relative to $\pi$). Moreover $L_\pi$ is injective, $L_\pi(\mathcal{O}_P) \subseteq k[[X]]$ (equivalently, $z \in \mathcal{O}_P \iff L_\pi(z) \in k[[X]]$, for every $z \in K$), and $L_\pi$ is a $k$-algebra homomorphism.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

As $P$ is rational, $k \to \mathcal{O}_P \to \mathcal{O}_P/\mathfrak{m}_P = k(\mathfrak{m}_P)$ is an isomorphism, that is $k \subseteq \mathcal{O}_P$ is a coefficient field for the DVR $\mathcal{O}_P$. By (3.34.22) (with $A := \mathcal{O}_P$, $\mathfrak{m} := \mathfrak{m}_P$), there is an injective $k$-algebra homomorphism $L_\pi : K \to k((X))$, extending the injective $L_\pi : \mathcal{O}_P \to k[[X]]$ of that proposition, with $L_\pi(\pi) = X$.

It is valuation-preserving: for $z = a/s \in K^\star$ ($a, s \in \mathcal{O}_P$), $L_\pi(z) = L_\pi(a)L_\pi(s)^{-1}$ (that proposition's extension formula), so $v_X(L_\pi(z)) = v_X(L_\pi(a)) - v_X(L_\pi(s)) = v_P(a) - v_P(s) = v_P(z)$ (multiplicativity of $v_X$ (3.10.13), and the valuation formula on $\mathcal{O}_P$, that proposition again). In particular $z \in \mathcal{O}_P \iff v_P(z) \geq 0 \iff v_X(L_\pi(z)) \geq 0 \iff L_\pi(z) \in k[[X]]$, for $z \in K$.

**$L_\pi(\pi^n) = X^n$ for every $n \in \mathbb{Z}$.** For $n \geq 0$ this is $L_\pi(\pi)^n = X^n$ ($L_\pi$ multiplicative); for $n < 0$, as $L_\pi$ is a field homomorphism and $X = L_\pi(\pi) \neq 0$, $L_\pi(\pi^{-1}) = L_\pi(\pi)^{-1} = X^{-1}$, so $L_\pi(\pi^n) = L_\pi(\pi^{-1})^{-n} = X^n$.

**Laurent coefficients.** Let $z \in K^\star$, $n_0 := v_P(z)$, and $u := z\pi^{-n_0} \in \mathcal{O}_P$, a unit (as $v_P(u) = 0$, (3.31.2)). By (3.34.22), for every $m \geq 0$ there is a unique $(c_0(u), \ldots, c_m(u)) \in k^{m+1}$ with $u - \sum_{i=0}^{m} c_i(u)\pi^i \in \mathfrak{m}_P^{m+1}$, compatible as $m$ varies, and $c_0(u) \neq 0$ (else $u \in \mathfrak{m}_P$, contradicting $u$ a unit); setting $a_j := c_{j-n_0}(u)$ for $j \geq n_0$ gives $a_{n_0} = c_0(u) \neq 0$, and, multiplying the congruence at $m := n - n_0$ by $\pi^{n_0}$,

$$z - \sum_{j=n_0}^{n} a_j \pi^j = \pi^{n_0}\left(u - \sum_{i=0}^{n-n_0} c_i(u)\pi^i\right) \in \pi^{n_0}\mathfrak{m}_P^{n-n_0+1} = \mathfrak{m}_P^{n+1}$$

for every $n \geq n_0$ (using $\mathfrak{m}_P = (\pi)$). As $L_\pi(z) = L_\pi(\pi^{n_0})L_\pi(u) = X^{n_0}\sum_i c_i(u)X^i = \sum_{j \geq n_0} a_j X^j$, this is exactly the stated Laurent coefficient sequence for $z$; uniqueness of $(a_j)_j$ follows from that of $(c_i(u))_i$ (any sequence satisfying the congruence gives, dividing by $\pi^{n_0}$, a coefficient sequence for $u$, forced equal to $(c_i(u))_i$ by (3.34.22)'s uniqueness).

**Uniqueness among such maps.** Suppose $\phi : K \to k((X))$ is $k$-linear with $\phi(\pi^n) = X^n$ for every $n \in \mathbb{Z}$ and $v_X(\phi(z)) = v_P(z)$ for every $z \in K^\star$. Fix $z \in K^\star$ with Laurent coefficients $(a_j)_{j \geq v_P(z)}$ as above; for every $n \geq v_P(z)$, writing $z = \sum_{j=v_P(z)}^{n} a_j\pi^j + r_n$ with $r_n \in \mathfrak{m}_P^{n+1}$ (i.e. $r_n = 0$ or $v_P(r_n) \geq n+1$), $k$-linearity and $\phi(\pi^j) = X^j$ give

$$\phi(z) - \sum_{j=v_P(z)}^{n} a_j X^j = \phi(r_n),$$

with $v_X(\phi(r_n)) \geq n+1$ (either $r_n = 0$, or by hypothesis on $\phi$); so $\phi(z) - \sum_{j=v_P(z)}^{n} a_j X^j \in X^{n+1}k[[X]]$, for arbitrarily large $n$, giving $\phi(z) = \sum_j a_j X^j = L_\pi(z)$. Hence $\phi = L_\pi$. ∎

**Definition 6.3.24 (Residue Symbol of a Laurent Series)**
*Let $A$ be a commutative ring, $\pi$ an indeterminate, and $z, t \in A((\pi))$. Write $\partial_\pi$ for the formal derivative, $\partial_\pi\left(\sum_n a_n\pi^n\right) := \sum_n n a_n \pi^{n-1}$, and define*

$$\mathrm{Res}_\pi(z\,\mathrm{d}t) := (z\partial_\pi(t))_{-1} \in A,$$

*the coefficient of $\pi^{-1}$ in $z\partial_\pi(t)$.*

© David Rufino CC BY-NC-ND 4.0

**Definition 6.3.25 (Residue Symbol at a Rational Place)**
*Let $P$ be a rational place of $K/k$, $\pi$ a uniformiser, and $t, z \in K$. Let $L_\pi : K \hookrightarrow k((\pi))$ be the completion embedding, and define*

$$\mathrm{Res}_P(z\,\mathrm{d}t) := \mathrm{Res}_\pi(L_\pi(z)\,\mathrm{d}L_\pi(t)) \in k,$$

*where $\mathrm{Res}_\pi$ is the residue of (6.3.24) with $A := k$, and $\mathrm{d}L_\pi(t) := \partial_\pi(L_\pi(t))\,\mathrm{d}\pi$.*

*This is the special case $A = k$ of (6.3.24), applied to $L_\pi(z), L_\pi(t) \in k((\pi))$. Explicitly, since $L_\pi$ commutes with the uniformiser derivation, $L_\pi(\partial_\pi(t)) = \partial_\pi(L_\pi(t))$, one has*

$$\mathrm{Res}_P(z\,\mathrm{d}t) = L_\pi(z\partial_\pi(t))_{-1} = (L_\pi(z)\partial_\pi(L_\pi(t)))_{-1},$$

*which is the formula used previously. We show that it is independent of the choice of uniformiser $\pi$ below (6.3.31).*

**Lemma 6.3.26 (Residue of $\psi^n\psi'$)**
*Let $\psi \in Xk[[X]]$ with $\psi'(0) \neq 0$ (equivalently, $\psi$ has order exactly 1). For every $n \in \mathbb{Z}$,*

$$(\psi^n\psi')_{-1} = \delta_{n,-1}$$

**Remark 6.3.27**
*The simplest proof using the identity $(n+1)\psi^n\psi' = (\psi^{n+1})'$ fails when $\mathrm{char}(k) = p \mid n+1$, so some additional work is required.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

**$n \geq 0$.** $\psi \in Xk[[X]]$ has no negative powers, so neither do $\psi^n$ or $\psi'$; their product has no $X^{-1}$ term, giving 0, matching $\delta_{n,-1} = 0$.

**$n = -1$.** Write $\psi(X) = Xv(X)$ with $v \in k[[X]]^\star$ a unit ($v(0) = \psi'(0) \neq 0$). Then $\psi' = v + Xv'$, so

$$\psi^{-1}\psi' = \frac{v + Xv'}{Xv} = \frac{1}{X} + \frac{v'}{v}$$

The second term is a genuine power series (product of $v'$ and the unit $v^{-1}$, both in $k[[X]]$), so has zero $-1$-coefficient; the first contributes $(X^{-1})_{-1} = 1$. Total: 1, matching $\delta_{-1,-1}$.

**$n \leq -2$: a universal identity.** Fix $n \leq -2$; we prove the identity over a universal coefficient ring first, then specialise (see the remark above for why). Let $R := \mathbb{Z}[a_1, a_2, a_3, \ldots][a_1^{-1}]$, the polynomial ring over $\mathbb{Z}$ in countably many indeterminates $a_1, a_2, \ldots$, localised to invert $a_1$ – an integral domain of characteristic 0 (a localisation of the domain $\mathbb{Z}[a_1, a_2, \ldots]$, itself a domain as a polynomial ring over the domain $\mathbb{Z}$, (3.8.3), applied to finitely many variables at a time since any product only ever involves finitely many of them). Let $\Psi := \sum_{i\geq 1} a_i X^i \in XR[[X]]$, so $\Psi = a_1 X \cdot U$ for $U := 1 + \sum_{i\geq 2}(a_i/a_1)X^{i-1} \in 1 + XR[[X]]$; as $U$ has constant coefficient $1 \in R^\star$, (3.10.10) gives $U \in R[[X]]^\star$, and hence $\Psi \in R((X))^\star$ too ($a_1 \in R^\star$, and $X$ a unit of $R((X))$ directly by construction of the formal Laurent series ring, (3.10.1)). So $\Psi^n\Psi' \in R((X))$ is well-defined, with a well-defined coefficient of $X^{-1}$, which we still write $(\Psi^n\Psi')_{-1} \in R$.

As before (the power rule and vanishing of a derivative's $X^{-1}$-coefficient being valid over any commutative ring, not just $k$), $(n+1)\Psi^n\Psi' = (\Psi^{n+1})'$ ((3.10.16), $\Psi$ being a unit) has vanishing $X^{-1}$-coefficient:

$$(n+1)\cdot(\Psi^n\Psi')_{-1} = \big((\Psi^{n+1})'\big)_{-1} = 0 \quad \text{in } R$$

As $n \leq -2$, $n+1$ is a nonzero integer; its image in $R$ is therefore nonzero, $R$ having characteristic 0. So $(n+1)\cdot(\Psi^n\Psi')_{-1} = 0$ forces $(\Psi^n\Psi')_{-1} = 0$ in $R$, by cancellation in the domain $R$.

**Specialising.** Write $\psi = \sum_{i\geq 1} c_i X^i$, so $c_1 = \psi'(0) \neq 0$ by hypothesis. As $c_1 \in k^\star$, the assignment $a_1 \mapsto c_1$, $a_i \mapsto c_i$ ($i \geq 2$) extends, by the universal property of localisation (3.6.5) (applied to $\mathbb{Z}[a_1, a_2, \ldots] \to k$, $a_1 \mapsto c_1$ a unit), to a ring homomorphism $\phi : R \to k$. Applying $\phi$ coefficientwise gives a ring homomorphism $\hat{\phi} : R((X)) \to k((X))$ (directly from the convolution formula for products, (3.10.1)) with $\hat{\phi}(\Psi) = \psi$, compatible with the formal derivative ($\hat{\phi}(f)' = \hat{\phi}(f')$, immediate from the coefficientwise formulas $f' = \sum_m mc_m X^{m-1}$ and $\phi(mc_m) = m\phi(c_m)$) and with the coefficient of $X^{-1}$ (literally the same index, carried along by $\phi$). So

$$(\psi^n\psi')_{-1} = \hat{\phi}(\Psi^n\Psi')_{-1} = \phi\big((\Psi^n\Psi')_{-1}\big) = \phi(0) = 0$$

using $(\Psi^n\Psi')_{-1} = 0$ in $R$ (just shown), matching $\delta_{n,-1} = 0$ for $n \leq -2$. ∎

© David Rufino CC BY-NC-ND 4.0

**Lemma 6.3.28 (Laurent Expansions Compose Across Uniformisers)**
*Let $P$ be a rational place of $K/k$, $\pi, \pi'$ uniformisers at $P$, and $\psi := L_{\pi'}(\pi) \in Xk[[X]]$.*

*Then for every $w \in K$,*

$$L_{\pi'}(w) = L_{\pi}(w) \circ \psi$$

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Write $f := L_\pi(w) = \sum_j a_j X^j$ and $g := f \circ \psi = \sum_j a_j \psi^j$ (well-defined, (3.10.17)). As $v(\psi) = 1$ and the order function is multiplicative on the field $k((X))$ (3.10.13), $v(\psi^j) = j$ for every $j$; so, for every $n$, the tail $g - \sum_{j\le n} a_j\psi^j = \sum_{j>n} a_j\psi^j$ has every term of order $> n$, that is

$$g - \sum_{j\le n} a_j \psi^j \in X^{n+1}k[[X]] \qquad (\star)$$

For $u \in K$ with $v_P(u) \ge n+1$, (6.3.23) gives $L_{\pi'}(u) \in X^{n+1}k[[X]]$ (its Laurent coefficients vanish below order $v_P(u) \ge n+1$). Apply this to $u := w - \sum_{j\le n} a_j\pi^j \in \mathfrak{m}_P^{n+1}$ (the defining congruence of $f = L_\pi(w)$, (6.3.23)). Since $L_{\pi'}$ is a $k$-algebra homomorphism of fields (6.3.23) with $L_{\pi'}(\pi) = \psi$, it sends $\pi^j \mapsto \psi^j$ for every $j \in \mathbb{Z}$ (3.2.6), and fixes $a_j \in k$; so, by linearity,

$$L_{\pi'}(w) - \sum_{j\le n} a_j\psi^j = L_{\pi'}\left(w - \sum_{j\le n} a_j\pi^j\right) \in X^{n+1}k[[X]] \qquad (\star\star)$$

Subtracting $(\star)$ from $(\star\star)$, $L_{\pi'}(w) - g \in X^{n+1}k[[X]]$ for every $n \in \mathbb{Z}$; as any nonzero element of $k((X))$ has some fixed finite order, this forces $L_{\pi'}(w) = g = L_\pi(w) \circ \psi$. ∎

**Corollary 6.3.29 (Laurent Coefficients of the Uniformiser Derivation)**
*Let $P$ be a rational place of $K/k$, $\pi$ a uniformiser. For every $w \in K$,*

$$L_\pi(\partial_\pi(w)) = \frac{\mathrm{d}}{\mathrm{d}X} L_\pi(w)$$

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Let $M := k((X))$, the target of $L_\pi : K \hookrightarrow k((X))$ (6.3.23), viewed as a $K$-module through $L_\pi$, a $k$-algebra homomorphism (6.3.23). By (3.35.44) (with $n = 1$, transcendence basis $\pi$), $\mathrm{Der}_k(K, M) \cong M$ via $D \mapsto D(\pi)$, so a $k$-derivation $K \to M$ is uniquely determined by its value at $\pi$.

Both $D_1 := L_\pi \circ \partial_\pi$ and $D_2 := \frac{\mathrm{d}}{\mathrm{d}X} \circ L_\pi$ are $k$-derivations $K \to M$: for $D_1$, $L_\pi(\partial_\pi(xy)) = L_\pi(x\partial_\pi(y) + y\partial_\pi(x)) = x \cdot D_1(y) + y \cdot D_1(x)$, using additivity and multiplicativity of $L_\pi$; for $D_2$, the same identity holds by the Leibniz rule for formal differentiation together with $L_\pi(xy) = L_\pi(x)L_\pi(y)$; and $D_1(\lambda) = L_\pi(\partial_\pi(\lambda)) = 0 = D_2(\lambda)$ for $\lambda \in k$. Both send $\pi \mapsto 1$ ($L_\pi(\partial_\pi(\pi)) = L_\pi(1) = 1$, $\frac{\mathrm{d}}{\mathrm{d}X}(L_\pi(\pi)) = \frac{\mathrm{d}}{\mathrm{d}X}(X) = 1$), so $D_1 = D_2$, that is $L_\pi(\partial_\pi(w)) = \frac{\mathrm{d}}{\mathrm{d}X} L_\pi(w)$ for every $w \in K$. ∎

**Proposition 6.3.30 (The Uniformiser Derivation is Regular)**
*Let $P$ be a rational place of $K/k$, $\pi$ a uniformiser. Then $\partial_\pi(\mathcal{O}_P) \subseteq \mathcal{O}_P$.*

*Proof.* This essentially follows from (6.3.23) and the fact that $\frac{\mathrm{d}}{\mathrm{d}X}$ preserves $k[[X]]$. ∎

**Corollary 6.3.31 (Well-Definedness of the Residue Symbol at a Rational Place)**
*Let $P$ be a rational place of $K/k$, $\pi, \pi'$ uniformisers at $P$, and $z, t \in K$. Then*

$$L_\pi(z\partial_\pi(t))_{-1} = L_{\pi'}(z\partial_{\pi'}(t))_{-1}$$

*that is, $\mathrm{Res}_P(z\,\mathrm{d}t)$ of (6.3.25) does not depend on the choice of uniformiser $\pi$ used to compute it.*

*Proof.* Write $w := z\partial_\pi(t)$ and $\psi := L_{\pi'}(\pi) \in Xk[[X]]$.

**$\psi'(0) \neq 0$.** Writing $\psi = \sum_j a_j X^j$, this is the Laurent expansion of $\pi \in K^\star$ relative to $\pi'$ (6.3.23), so $a_j = 0$ for $j < v_P(\pi)$ and $a_{v_P(\pi)} \neq 0$; as $v_P(\pi) = 1$ ($\pi$ is itself a uniformiser), $a_0 = 0$ and $a_1 \neq 0$. The constant term of $\psi' = \sum_j j a_j X^{j-1}$ is its $j = 1$ term, $1 \cdot a_1 = a_1 \neq 0$, that is $\psi'(0) \neq 0$.

**$L_\pi(w)_{-1} = (L_{\pi'}(w) \cdot \psi')_{-1}$.** Write $f := L_\pi(w)$ and $g := f \circ \psi$ (well-defined, (3.10.17), as $\psi$ has invertible linear term, just shown). Both sides of $f_{-1} = (g \cdot \psi')_{-1}$ are $k$-linear in $f$ (composition with $\psi$, multiplication by $\psi'$, and $(-)_{-1}$ are all $k$-linear), so it suffices to check $f = X^n$, $n \in \mathbb{Z}$, that is $(\psi^n\psi')_{-1} = \delta_{n,-1}$ – exactly (6.3.26). By (6.3.28), $g = L_\pi(w) \circ \psi = L_{\pi'}(w)$, giving $L_\pi(w)_{-1} = f_{-1} = (g \cdot \psi')_{-1} = (L_{\pi'}(w) \cdot \psi')_{-1}$.

© David Rufino CC BY-NC-ND 4.0

By (6.3.29) (applied relative to $\pi'$), $L_{\pi'}(\partial_{\pi'}(\pi)) = \frac{\mathrm{d}}{\mathrm{d}X} L_{\pi'}(\pi) = \psi'$. As $L_{\pi'}$ is a ring homomorphism (6.3.23),

$$L_\pi(z\partial_\pi(t))_{-1} = L_\pi(w)_{-1} = (L_{\pi'}(w) \cdot L_{\pi'}(\partial_{\pi'}(\pi)))_{-1} = (L_{\pi'}(w\partial_{\pi'}(\pi)))_{-1}.$$

By (6.3.22), $\partial_\pi(t)\partial_{\pi'}(\pi) = \partial_{\pi'}(t)$, so $w\partial_{\pi'}(\pi) = z\partial_\pi(t)\partial_{\pi'}(\pi) = z\partial_{\pi'}(t)$, giving $L_\pi(z\partial_\pi(t))_{-1} = L_{\pi'}(z\partial_{\pi'}(t))_{-1}$. ∎

**Lemma 6.3.32 (Residues are Invariant under Isomorphism)**
*Let $\varphi : K \to K'$ be an isomorphism of $k$-algebras between algebraic function fields $K/k$, $K'/k$, $P$ a rational place of $K/k$ and $P'$ the place of $K'/k$ with $\mathcal{O}_{P'} = \varphi(\mathcal{O}_P)$. Then $\mathrm{Res}_{P'}(\varphi(z)\,\mathrm{d}\varphi(t)) = \mathrm{Res}_P(z\,\mathrm{d}t)$ for all $z, t \in K$.*

**Proposition 6.3.33 (Elementary Properties of the Residue Symbol at a Rational Place)**
*Let $P$ be a rational place of $K/k$, $\pi$ a uniformiser, $t \in K$. Then*

*a) $z \mapsto \mathrm{Res}_P(z\,\mathrm{d}t)$ is $k$-linear $K \to k$;*
*b) for $x \in K$ transcendental over $k$ and $\partial_x \in \mathrm{Der}_k(K, K)$ the distinguished derivation of (6.3.20), $\mathrm{Res}_P(z\,\mathrm{d}t) = \mathrm{Res}_P((z \cdot \partial_x(t))\,\mathrm{d}x)$ for every $z \in K$;*
*c) if $z\partial_\pi(t) \in \mathcal{O}_P$ then $\mathrm{Res}_P(z\,\mathrm{d}t) = 0$; in particular this holds if $z, t \in \mathcal{O}_P$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

a) By (6.3.25), $\mathrm{Res}_P(z\,\mathrm{d}t) = L_\pi(z\partial_\pi(t))_{-1}$; $k$-linearity in $z$ follows from that of $\partial_\pi$, of $L_\pi$ (6.3.23), and of coefficient extraction (6.3.24).

b) By (6.3.20) (applied to $D := \partial_\pi$), $\partial_\pi = \partial_\pi(x) \cdot \partial_x$, so $z\partial_\pi(t) = (z \cdot \partial_x(t))\partial_\pi(x)$; applying $L_\pi(-)_{-1}$ to both sides ((6.3.25), once with $t$ and once with $x$) gives $\mathrm{Res}_P(z\,\mathrm{d}t) = \mathrm{Res}_P((z \cdot \partial_x(t))\,\mathrm{d}x)$.

c) If $z\partial_\pi(t) \in \mathcal{O}_P$ then $L_\pi(z\partial_\pi(t)) \in k[[X]]$ (6.3.23), so its $X^{-1}$-coefficient vanishes, that is $\mathrm{Res}_P(z\,\mathrm{d}t) = 0$. If $z, t \in \mathcal{O}_P$ then $\partial_\pi(t) \in \mathcal{O}_P$ (6.3.30), so $z\partial_\pi(t) \in \mathcal{O}_P$. ∎

## 6.3.6 The Algebraic Residue Theorem over an Algebraically Closed Field

Assume throughout this subsection that $k$ is algebraically closed. Then every place of $K/k$ is rational (6.2.114), so the entire theory of (6.3.25) applies directly, with no need for the base-change-to-$\overline{k}$ machinery of the following subsections (which is needed only once $k$ is no longer assumed algebraically closed).

**Proposition 6.3.34 (Algebraic Residue Theorem, over an Algebraically Closed Field)**
*Assume $k$ is algebraically closed. For $t, f \in K$,*

$$\sum_P \mathrm{Res}_P(f\,\mathrm{d}t) = 0,$$

*summed over all places $P$ of $K/k$ (finitely many terms nonzero, as $f\,\mathrm{d}t$ has only finitely many poles).*

**Roadmap.** The proof, given immediately below, proceeds in the following steps.

- *Reduce to a differential $f\,\mathrm{d}x$.* Choose a separating transcendental $x \in K$; then $f\,\mathrm{d}t = f\partial_x(t)\,\mathrm{d}x$ at every place (6.3.33).b).
- *The case $K = k(x)$.* Partial fractions over $k = \overline{k}$ (6.3.35) write a rational function as a sum of terms $c_{i,j}(x - \theta_i)^{-j}$; the finite residues are the $c_{i,1}$, and the place at infinity has residue $-\sum_i c_{i,1}$ (6.3.39).
- *Trace formula.* For a finite separable $K/k(x)$ and a place $Q$ of $k(x)$, $\sum_{P|Q} \mathrm{Res}_P(f\,\mathrm{d}x) = \mathrm{Res}_Q\big(\mathrm{Tr}_{K/k(x)}(f)\,\mathrm{d}x\big)$ (6.3.42). This is proved locally, after completing at $Q$: the completed field splits as $\prod_i k((\pi_i))$ over the places above $Q$, reducing the identity to the local trace formula at each place, which is computed monomial by monomial via the roots of unity in $k$, following Lang.
- *Assemble.* Group the places $P$ of $K$ by the place $Q$ of $k(x)$ below them. The trace formula turns each inner sum into $\mathrm{Res}_Q\big(\mathrm{Tr}_{K/k(x)}(f')\,\mathrm{d}x\big)$, and summing over $Q$ gives 0 by the case $K = k(x)$.

*Proof of (6.3.34). Drafted by Claude Sonnet, reviewed by David Rufino.*

Fix any transcendental $x \in K$ with $K/k(x)$ finite and separable (exists as $K/k$ is a function field, separably generated), and let $\partial_x \in \mathrm{Der}_k(K, K)$ be the distinguished element of (6.3.20) with $\partial_x(x) = 1$. By (6.3.33).b), for every place $P$,

$$\mathrm{Res}_P(f\,\mathrm{d}t) = \mathrm{Res}_P((f \cdot \partial_x(t))\,\mathrm{d}x)$$

this handles $t \in k$ automatically too, as $\partial_x(k) = 0$ then makes both sides 0, matching $\mathrm{d}t = 0$. So it suffices to prove $\sum_P \mathrm{Res}_P(f'\,\mathrm{d}x) = 0$ for $f' := f \cdot \partial_x(t) \in K$.

Every place $P$ of $K/k$ lies over a unique place $Q$ of $k(x)/k$, so (finitely many terms being nonzero) the sum may be regrouped:

© David Rufino CC BY-NC-ND 4.0

$$\sum_{P \text{ of } K/k} \operatorname{Res}_P(f'\,\mathrm{d}x) = \sum_{Q \text{ of } k(x)/k} \ \sum_{P \mid Q} \operatorname{Res}_P(f'\,\mathrm{d}x).$$

By (6.3.42), the inner sum is $\operatorname{Res}_Q\big(\operatorname{Tr}_{K/k(x)}(f')\,\mathrm{d}x\big)$ for every $Q$, so

$$\sum_{P} \operatorname{Res}_P(f'\,\mathrm{d}x) = \sum_{Q} \operatorname{Res}_Q\big(\operatorname{Tr}_{K/k(x)}(f')\,\mathrm{d}x\big) = 0$$

by (6.3.39), applied to $\operatorname{Tr}_{K/k(x)}(f') \in k(x)$. ∎

This proof uses only (6.3.20) (already available, with no need for the module of differentials $\Omega_{K/k}$ to have been formally constructed).

**Lemma 6.3.35 (Partial Fraction Decomposition over an Algebraically Closed Field)**
*Let $k$ be algebraically closed and $r, s \in k[X]$ polynomials with $s \neq 0$. Write $s(X) = c\prod_{i=1}^{m}(X-\theta_i)^{e_i}$ for distinct $\theta_i \in k$, integers $e_i \geq 1$, and $c \in k^\times$. Then there exist a polynomial $Q \in k[X]$ and constants $c_{i,j} \in k$ $(1 \leq i \leq m,\ 1 \leq j \leq e_i)$ with*

$$\frac{r(X)}{s(X)} = Q(X) + \sum_{i=1}^{m}\sum_{j=1}^{e_i} \frac{c_{i,j}}{(X-\theta_i)^j}$$

*as an identity in $k(X)$. If $\deg r < \deg s$ then $Q = 0$.*

*Proof.* Absorbing $c$ into $r$, we may take $c = 1$. We argue by induction on $m$, the number of distinct roots of $s$. If $m = 0$ then $s = 1$ and we take $Q := r$.

Let $m \geq 1$ and set $s_1 = (X-\theta_1)^{e_1}$ and $s_2(X) := \prod_{i=2}^{m}(X-\theta_i)^{e_i}$, so $s = s_1 s_2$. By (3.19.10) $s_1$ and $s_2$ are co-prime as they have no common roots in $k$, so $gs_1 + hs_2 = 1$ for some $g, h \in k[X]$ and

$$\frac{r(X)}{s(X)} = \frac{r(X)h(X)}{s_1}(X) + \frac{r(X)g(X)}{s_2}(X)$$

By the division algorithm $rh = Q_1 s_1 + r_1$ with $\deg(r_1) < e_1$. By (3.8.11) $r_1 = \sum_{l=0}^{e_1-1} d_l (X-\theta_1)^l$ for some $d_l \in k$, so with $c_{1,j} := d_{e_1-j}$

$$\frac{rh}{s_1} = Q_1 + \sum_{j=1}^{e_1} \frac{c_{1,j}}{(X-\theta_1)^j}$$

The induction hypothesis applies to $rg/s_2$, which has $m-1$ distinct roots in its denominator, and gives the terms with $i \geq 2$ together with a polynomial $Q_2$. Adding, the claim holds with $Q := Q_1 + Q_2$.

Finally suppose $\deg r < \deg s$. Multiplying the decomposition by $s$ gives $Qs = r - \sum_{i,j} c_{i,j}\, s/(X-\theta_i)^j$, where each $s/(X-\theta_i)^j$ is a polynomial of degree $\deg s - j \leq \deg s - 1$. So the right hand side has degree less than $\deg s$, whereas $\deg(Qs) \geq \deg s$ if $Q \neq 0$; hence $Q = 0$. ∎

**Remark 6.3.36 (A Complex-Analytic Heuristic)**
*Over $k = \mathbb{C}$, this also follows from Liouville's theorem. At each $\theta_i$, choose $c_{i,1}, \ldots, c_{i,e_i}$ to match the principal part of the Laurent expansion of $r/s$ at $\theta_i$ (possible and unique, exactly as the induction above extracts them); then $r/s - \sum_{i,j} c_{i,j}/(X-\theta_i)^j$ is a rational function with no poles anywhere in $\mathbb{C}$, hence an entire function, and one that grows at most polynomially at infinity (being itself a rational function) – so by (the polynomial-growth form of) Liouville's theorem it is a polynomial, namely $Q$.*

**Lemma 6.3.37 (Residue at Infinity of a Proper Rational Function)**
*Let $f, g \in k[X]$ with $g \neq 0 \neq f$ and $\deg(f) < \deg(g)$. Then, for the place at infinity of $k(X)/k$ (6.2.117),*

$$\operatorname{Res}_\infty\left(\frac{f}{g}\,\mathrm{d}X\right) = \begin{cases} -\frac{\ell(f)}{\ell(g)} & \deg(f) = \deg(g) - 1 \\ 0 & \deg(f) < \deg(g) - 1. \end{cases}$$

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Let $p := \deg(f)$, $q := \deg(g)$, and $m := q - p \geq 1$. Set $A(U) := U^p f(1/U)$, $B(U) := U^q g(1/U) \in k[X]$, polynomials with $A(0) = \ell(f)$, $B(0) = \ell(g) \neq 0$, so $w := A/B \in k[[U]]$ is a unit with $w(0) = \ell(f)/\ell(g)$. As $f(1/U) = U^{-p}A(U)$ and $g(1/U) = U^{-q}B(U)$,

© David Rufino CC BY-NC-ND 4.0

$$\frac{f}{g}(1/U) = U^{q-p}\frac{A(U)}{B(U)} = U^m w(U).$$

Let $U$ be an indeterminate and $\varphi : k(X) \to k(U)$ the isomorphism with $\varphi(X) = U^{-1}$; it maps the place at infinity to the place 0 of $k(U)$, with uniformiser $U$ (6.2.117). By (6.3.32) and (6.3.33).b) (with $x := U$, as $\partial_U(U^{-1}) = -U^{-2}$),

$$\mathrm{Res}_\infty\left(\frac{f}{g}\,\mathrm{d}X\right) = \mathrm{Res}_0(\varphi(f/g)\,\mathrm{d}(U^{-1})) = \mathrm{Res}_0(-U^{m-2}w(U)\,\mathrm{d}U).$$

As $\partial_U(U) = 1$, this is the coefficient of $U^{-1}$ in $-U^{m-2}w(U)$, that is $-$ the coefficient of $U^{1-m}$ in $w(U)$. If $m = 1$ this is $-w(0) = -\ell(f)/\ell(g)$. If $m \geq 2$ then $1 - m < 0$, and as $w \in k[[U]]$ has only non-negative powers, this coefficient is 0. ∎

**Remark 6.3.38 (A Complex-Analytic Heuristic)**
*Over $k = \mathbb{C}$, this is controlled entirely by the leading-order behaviour of $f/g$ at infinity. Writing $U = 1/X$, the residue at infinity is $-\mathrm{Res}_0$ of the pulled-back differential, and $f/g(1/U)\cdot\partial_U(U^{-1}) = -U^{m-2}w(U)$ has leading term $-\left(\frac{\ell(f)}{\ell(g)}\right)U^{m-2}$ ($w(0) = \ell(f)/\ell(g) \neq 0$); classically $\oint U^n\,\mathrm{d}U = 0$ for every $n \neq -1$, so the residue vanishes unless this* leading *term is itself the $U^{-1}$ term, i.e. $m - 2 = -1$, or $\deg f = \deg g - 1$ – exactly the borderline decay rate $f/g \sim (\ell(f)/\ell(g))X^{-1}$ at infinity – in which case the residue is read off directly from the leading coefficients, with no correction from the rest of the expansion of $w$.*

**Lemma 6.3.39 (Residue Theorem for the Rational Function Field)**
*Assume $k$ is algebraically closed. For $\theta \in k$, $n \in \mathbb{Z}$ and $g := (X-\theta)^n \in k(X)$, the residues at the places of $k(X)/k$ (the $\mathcal{O}_{X-\theta'}$, $\theta' \in k$, written $\theta'$, and the place at infinity (6.2.117)) are*

$$\mathrm{Res}_{\theta'}(g\,\mathrm{d}X) = \begin{cases} 1 & \theta' = \theta \text{ and } n = -1 \\ 0 & \text{otherwise} \end{cases} \qquad \mathrm{Res}_\infty(g\,\mathrm{d}X) = \begin{cases} -1 & n = -1 \\ 0 & n \neq -1. \end{cases}$$

*Consequently, for every $g \in k(X)$, where $k(X)/k$ is the rational function field,*

$$\sum_Q \mathrm{Res}_Q(g\,\mathrm{d}X) = 0,$$

*summed over all places $Q$ of $k(X)/k$ (finitely many terms nonzero).*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Write $T$ for the variable of the Laurent series field, to avoid a clash with $X \in k(X)$, so that $L_\pi : k(X) \to k((T))$ is a ring homomorphism with $L_\pi(\pi) = T$ (6.3.23).

**Finite places.** The place $\theta'$ has uniformiser $\pi := X - \theta'$ with $\partial_\pi(X) = 1$. If $\theta' \neq \theta$ then $g$ and $X$ lie in $\mathcal{O}_{X-\theta'}$, so $\mathrm{Res}_{\theta'}(g\,\mathrm{d}X) = 0$ (6.3.33).c). If $\theta' = \theta$ then by (6.3.25)

$$\mathrm{Res}_\theta(g\,\mathrm{d}X) = L_\pi(\pi^n)_{-1} = (T^n)_{-1} = \begin{cases} 1 & n = -1 \\ 0 & n \neq -1 \end{cases}$$

**The place at infinity.** If $n \geq 0$ then $g$ is a polynomial, so $g(1/U)$ is a $k$-linear combination of $U^0, \ldots, U^{-n}$, and $g(1/U)\partial_U(X) = -U^{-2}g(1/U)$ a linear combination of $U^{-2}, \ldots, U^{-n-2}$; as every exponent is $\leq -2$, $\mathrm{Res}_\infty(g\,\mathrm{d}X) = 0$. If $n \leq -1$ then $g = 1/(X-\theta)^{-n}$ is a proper fraction with numerator 1 and denominator $(X-\theta)^{-n}$ of degree $-n$, so by (6.3.37)

$$\mathrm{Res}_\infty(g\,\mathrm{d}X) = \begin{cases} -1 & n = -1 \\ 0 & n \leq -2. \end{cases}$$

**The sum vanishes for every $g \in k(X)$.** By (6.3.35) every $g \in k(X)$ is the sum of a polynomial and a $k$-linear combination of the functions $(X-\theta)^{-n}$ with $\theta \in k$, $n \geq 1$; and by (3.8.11) every polynomial is a $k$-linear combination of the $(X-\theta)^n$, $n \geq 0$, for any fixed $\theta \in k$. The residue being $k$-linear, this reduces to the case $g = (X-\theta)^n$ above, where the residues at all places other than $\theta$ and $\infty$ vanish, and $\mathrm{Res}_\theta(g\,\mathrm{d}X) + \mathrm{Res}_\infty(g\,\mathrm{d}X)$ is $1 - 1 = 0$ if $n = -1$ and 0 otherwise. ∎

**Lemma 6.3.40 (Local Trace Formula for Residues)**
*Let $k$ be algebraically closed, $F := k((t))$, and let $E = k((\pi))$ be a finite separable extension of $F$ with $v_\pi(t) = e$, so $[E:F] = e$ and $t = \pi^e w$ for some $w \in k[[\pi]]^\star$ (3.33.4). For $g \in E$ write $\mathrm{Res}_\pi(g\,\mathrm{d}t) := (g\partial_\pi(t))_{-1}$. Then, in every characteristic,*

© David Rufino CC BY-NC-ND 4.0

$$\operatorname{Res}_t\big(\operatorname{Tr}_{E/F}(g)\,\mathrm{d}t\big) = \operatorname{Res}_\pi(g\,\mathrm{d}t) \qquad (g \in E).$$

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

**Basis and matrices.** By (3.33.2) (with $R := k$, $X := \pi$, $m := e$, $T = t$), every element of $k[[\pi]]$ is uniquely $\sum_{i<e} f_i(t)\pi^i$ with $f_i \in k[[Y]]$; multiplying by powers of $t$ (which is $\pi^e$ up to a unit) gives the same for $E$. So $1, \pi, \dots, \pi^{e-1}$ is an $F$-basis of $E$. For $g \in E$ write $g\pi^i = \sum_{j<e} M_{ji}(g)\pi^j$ with $M_{ji}(g) \in F$, and $\operatorname{Tr}_{E/F}(g) = \sum_{i<e} M_{ii}(g)$.

**Universal model.** Let $R := \mathbb{Z}\big[b_0^{\pm 1}, b_1, b_2, \dots\big]$, a domain of characteristic 0, and put $w_R := \sum_{i\ge 0} b_i\pi^i \in R[[\pi]]^\star$, $t_R := \pi^e w_R$. The construction above works verbatim over $R$ (only $b_0$ needs inverting), with matrix entries $M_{ji}(g) \in R((t_R))$ given by universal formulas in $+$, $\times$ and $w_R^{-1}$, whose coefficients lie in $R$. For $r \in \mathbb{Z}$ set

$$\Psi(\pi^r) := \operatorname{Res}_{t_R}(\operatorname{Tr}_R(\pi^r)\,\mathrm{d}t_R) - \operatorname{Res}_\pi(\pi^r\,\mathrm{d}t_R) \in R,$$

both terms being coefficients of elements of $R((\pi))$ or $R((t_R))$.

**Claim: $\Psi(\pi^r) = 0$ for all $r$.** Let $\Omega$ be an algebraic closure of $\operatorname{Frac}(R)$; it has characteristic 0, so $e$ is invertible in $\Omega$ and $R \subset \Omega$. Write $w_R = b_0 u$ with $u \in 1 + \pi R[[\pi]]$. The binomial series $\varepsilon := \sum_{i\ge 0}\binom{1/e}{i}(u-1)^i \in \Omega[[\pi]]$ satisfies $\varepsilon^e = u$, and for $\beta \in \Omega$ with $\beta^e = b_0$ the element $\varpi := \beta\pi\varepsilon$ satisfies $\varpi^e = t_R$. As $\varpi = \pi \cdot (\text{unit})$, it is a uniformiser of $\Omega((\pi)) = \Omega((\varpi))$, and $\Omega((t_R)) \subset \Omega((\pi))$ has basis $1, \varpi, \dots, \varpi^{e-1}$ (the argument above, with $\varpi$ in place of $\pi$).

Take $g = \varpi^r$ with $r = es + j$, $0 \le j < e$. Then $\varpi^r = t_R^s\varpi^j$, and multiplication by $\varpi^j$ sends the basis vector $\varpi^i$ to $\varpi^{i+j}$, which is a basis vector times $t_R^0$ or $t_R$. So the matrix of $\varpi^r$ has nonzero entries only in positions shifted by $r$ modulo $e$: its diagonal vanishes unless $e \mid r$, and if $r = es$ it is the scalar $t_R^s$. Hence

$$\operatorname{Tr}(\varpi^r) = \begin{cases} e t_R^s & r = es \\ 0 & e \nmid r. \end{cases}$$

Therefore $\operatorname{Res}_{t_R}(\operatorname{Tr}(\varpi^r)\,\mathrm{d}t_R) = e$ if $r = -e$ and 0 otherwise. On the other side $\mathrm{d}t_R = e\varpi^{e-1}\,\mathrm{d}\varpi$, so $\varpi^r\,\mathrm{d}t_R = e\varpi^{r+e-1}\,\mathrm{d}\varpi$ and $\operatorname{Res}_\varpi(\varpi^r\,\mathrm{d}t_R)$ is again $e$ for $r = -e$ and 0 otherwise. Residues are independent of the choice of uniformiser (6.3.31), so $\operatorname{Res}_\varpi(g\,\mathrm{d}t_R) = \operatorname{Res}_\pi(g\,\mathrm{d}t_R)$. Thus the identity holds for $g = \varpi^r$ for every $r$. Both sides are $\Omega$-linear and continuous in $g$, and the $\varpi^r$ span $\Omega((\pi))$ topologically, so it holds for every $g \in \Omega((\pi))$; in particular $\Psi(\pi^r) \in R$ maps to 0 in $\Omega$. As $R \subset \Omega$, $\Psi(\pi^r) = 0$.

**Specialisation.** Given $E = k((\pi))$ and $t = \pi^e w$ with $w = \sum_{i\ge 0} c_i\pi^i$, $c_0 \in k^\times$, let $\phi : R \to k$ be the ring homomorphism with $b_i \mapsto c_i$ (it exists by the universal property of localisation, as $c_0$ is a unit). Applying $\phi$ coefficientwise gives ring homomorphisms $R((\pi)) \to k((\pi))$ and $R((t_R)) \to k((t))$ (with $t_R \mapsto t$, since $\phi(w_R) = w$). These commute with the formation of matrices $M_{ji}$, hence with traces, and with the coefficient extraction defining both residues. So $\phi(\Psi(\pi^r))$ is the defect $\operatorname{Res}_t\big(\operatorname{Tr}_{E/F}(\pi^r)\,\mathrm{d}t\big) - \operatorname{Res}_\pi(\pi^r\,\mathrm{d}t)$ over $k$, and it equals 0. Finally, for $g = \sum_r a_r\pi^r \in k((\pi))$ both sides depend on only finitely many $a_r$, since $\operatorname{Tr}(\pi^r)$ has $t$-order tending to $\infty$ with $r$ and only finitely many terms contribute to a given coefficient of $t^{-1}$ or $\pi^{-1}$. The identity for each monomial therefore gives it for $g$. ∎

**Lemma 6.3.41 (Places over $P$ and Irreducible Factors)**
*Let $k$ be algebraically closed, $K/k(x)$ a finite separable extension of degree $n$, $P$ the place $(x)$ of $k(x)$, $K_P := k((x))$, and $y \in K$ a primitive element with minimal polynomial $g$ over $k(x)$. Factor $g = g_1 \cdots g_r$ into distinct monic irreducibles over $K_P$, with $d_j := \deg g_j$, and let $E_j := K_P[Y]/(g_j)$ with $\iota_j : K \hookrightarrow E_j$ sending $y \mapsto Y$. Then $j \mapsto Q_j$, defined below, is a bijection from $\{1, \dots, r\}$ onto the places of $K/k$ over $P$, and $e_{Q_j} = d_j$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

**The place $Q_j$.** $E_j/K_P$ is finite separable (as $g_j$ is separable), so by (3.33.4) there is an isomorphism $\Psi_j : E_j \xrightarrow{\sim} k\big((\pi_j)\big)$ over $K_P$ with $\Psi_j(x)$ of order $d_j$. Pulling back the valuation of $k\big((\pi_j)\big)$ along $\Psi_j \circ \iota_j$ gives a discrete valuation of $K$ extending $v_P$ with ramification index $d_j$; let $Q_j$ be the corresponding place.

**Every place is some $Q_j$.** Let $Q$ be a place of $K/k$ over $P$. As $k$ is algebraically closed, $Q$ is rational, so (6.3.23) gives a valuation-preserving embedding $L_Q : K \hookrightarrow k\big((\pi_Q)\big)$. Since $g(y) = 0$, the element $L_Q(y)$ is a root of $g$ in $k\big((\pi_Q)\big)$. The $g_j$ are pairwise coprime in $K_P[Y]$, so $L_Q(y)$ is a root of exactly one $g_j$ (a common root of $g_j, g_{j'}$ would be a root of 1, by Bézout). Call this $j = j(Q)$. Then $L_Q$ factors through $E_j$, and $L_Q(x)$ has ramification index $e_Q$ over $K_P$; the valuation induced on $E_j$ extends $v_P$, so by the uniqueness of extension ((3.33.5), applied to $E_j/K_P$) it equals the valuation of $Q_j$. Hence $Q = Q_{j(Q)}$.

**Injectivity.** If $Q_j = Q_{j'}$ then $j(Q_j) = j$ and $j(Q_{j'}) = j'$ are determined by the same valuation, and the minimal factor of $L(y)$ is unique, so $j = j'$. ∎

© David Rufino CC BY-NC-ND 4.0

**Proposition 6.3.42 (Trace Formula for Residues)**
*Assume $k$ is algebraically closed. Let $\mathcal{O} \coloneqq k[x]$, $\mathcal{B} \coloneqq \overline{\mathcal{O}}^{K}$ the integral closure of $\mathcal{O}$ in a finite separable extension $K/k(x)$, $P$ a place of $k(x)/k$ corresponding to a maximal ideal $\mathfrak{p} \lhd \mathcal{O}$, and $f \in K$. Then*

$$\sum_{Q|P} \operatorname{Res}_Q(f\,\mathrm{d}x) = \operatorname{Res}_P\big(\operatorname{Tr}_{K/k(x)}(f)\,\mathrm{d}x\big),$$

*with no hypothesis on* $\operatorname{char}(k)$.

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

As $k$ is algebraically closed every place is rational, so $\mathfrak{p} = (x-\theta)$ for some $\theta \in k$; replacing $x$ by $x-\theta$ we may assume $\mathfrak{p} = (x)$.

By (3.32.15) the places $Q_1, \dots, Q_r$ of $K/k$ over $P$ have ramification indices $e_1, \dots, e_r$ with $\sum_i e_i = n \coloneqq [K : k(x)]$ (residue degree 1, as $k$ is algebraically closed). Let $K_P \coloneqq k((x))$, with $k(x) \hookrightarrow K_P$ the expansion at $P$ (6.3.23).

**Factoring the minimal polynomial.** Let $y \in \mathcal{B}$ be a primitive element with minimal polynomial $g \in \mathcal{O}[Y]$, monic of degree $n$. Factor $g = g_1 \cdots g_r$ over $K_P$ as in (6.3.41), with $E_j = K_P[Y]/(g_j)$, $\iota_j : K \hookrightarrow E_j$, and $\Psi_j : E_j \xrightarrow{\sim} k\big((\pi_j)\big)$ the isomorphisms given there, so that $\Psi_j(x)$ has order $d_j = \deg g_j$. Put $\lambda_j \coloneqq \Psi_j \circ \iota_j : K \hookrightarrow k\big((\pi_j)\big)$ and $L_{\pi_j} \coloneqq \lambda_j$. By (6.3.41), the places $Q_j$ attached to the $g_j$ are exactly the places of $K/k$ over $P$, with $e_{Q_j} = d_j$, so after relabelling $Q_i$ is the place attached to $g_i$ and $e_i = d_i$.

**Trace additivity.** By the Chinese remainder theorem applied to $g = \prod_j g_j$,

$$K \otimes_{k(x)} K_P \xrightarrow{\sim} \prod_{i=1}^{r} E_i, \quad f \otimes 1 \mapsto (\iota_i(f))_i,$$

and composing with the isomorphisms $\Psi_i$ identifies the right side with $\prod_i k((\pi_i))$, with $f \otimes 1 \mapsto \big(L_{\pi_i}(f)\big)_i$. The trace is unchanged by base change and additive over a finite product, so

$$\operatorname{Tr}_{K/k(x)}(f) = \sum_{i=1}^{r} \operatorname{Tr}_{k((\pi_i))/K_P}\big(L_{\pi_i}(f)\big) \qquad (f \in K).$$

**Local identities.** The residue at $Q_i$ is the residue symbol of (6.3.25), computed in $k((\pi_i))$ through $L_{\pi_i}$, so $\operatorname{Res}_{Q_i}(f\,\mathrm{d}x) = \operatorname{Res}_{\pi_i}\big(L_{\pi_i}(f)\,\mathrm{d}x\big)$ (Laurent residue of (6.3.24)), and $\operatorname{Res}_P(z\,\mathrm{d}x) = \operatorname{Res}_x(z\,\mathrm{d}x)$ for $z \in K_P = k((x))$. Applying (6.3.40) to $E = k((\pi_i))$ over $F = K_P$ (with $t = x$ and $\Psi_i(x) = \tau_i$) gives, for every $f \in K$,

$$\operatorname{Res}_{Q_i}(f\,\mathrm{d}x) = \operatorname{Res}_P\big(\operatorname{Tr}_{k((\pi_i))/K_P}\big(L_{\pi_i}(f)\big)\,\mathrm{d}x\big).$$

**Assembly.** Summing over $i$ and using $k$-linearity of $\operatorname{Res}_P$ and trace additivity,

$$\sum_{Q|P} \operatorname{Res}_Q(f\,\mathrm{d}x) = \sum_{i=1}^{r} \operatorname{Res}_P\big(\operatorname{Tr}_{k((\pi_i))/K_P}\big(L_{\pi_i}(f)\big)\,\mathrm{d}x\big) = \operatorname{Res}_P\big(\operatorname{Tr}_{K/k(x)}(f)\,\mathrm{d}x\big).$$

■

## 6.3.7 Base Extension of the Constant Field

Throughout this section, in addition to the standing hypothesis that $K/k$ is geometrically integral ($\tilde{k} = k$), assume also that $k$ is **perfect**. Let $k'/k$ be an algebraic field extension, and write $K_{k'} \coloneqq K \otimes_k k'$.

**Proposition 6.3.43 (The Base-Changed Function Field)**
*$K_{k'}$ is a field, and an algebraic function field over $k'$, geometrically integral; if $k'/k$ is finite, $[K_{k'} : K] = [k' : k]$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

***$K_{k'}$* is a field.** As $K/k$ is geometrically integral and $k'/k$ is algebraic, $K_{k'} = K \otimes_k k'$ is a field (3.36.45).

***$K_{k'}/k'$* is geometrically integral.** By associativity of tensor products (3.4.13) and $\overline{k'} = \overline{k}$ ($k'/k$ algebraic), $K_{k'} \otimes_{k'} \overline{k'} \cong K \otimes_k \big(k' \otimes_{k'} \overline{k}\big) = K \otimes_k \overline{k}$, a field, as $K/k$ is geometrically integral.

***$K_{k'}/k'$* is an algebraic function field.** Fix a transcendental $x \in K$ with $K/k(x)$ finite (exists, as $K/k$ is an algebraic function field). As $k[x] \otimes_k k' \cong k'[x]$ (tensoring a polynomial ring with a field extension), the image of $x$ in $K_{k'}$ remains transcendental over $k'$. The image of $k(x) \otimes_k k'$ in $K_{k'}$ (well-defined and injective, as $k(x) \subset K$ is a $k$-subalgebra, (3.36.2)) is then a subring of the field $K_{k'}$, hence a domain; as $k'/k$ is algebraic, it is thus a field (3.35.57),

© David Rufino CC BY-NC-ND 4.0

and – being the subfield of $K_{k'}$ generated by $k'$ and the transcendental $x$ – equals $k'(x)$. By associativity of tensor products (3.4.13) (as $K$ is a $k(x)$-module),

$$K_{k'} = K \otimes_k k' \cong K \otimes_{k(x)} (k(x) \otimes_k k') = K \otimes_{k(x)} k'(x)$$

As $K/k(x)$ is finite, $K$ is a finite-dimensional $k(x)$-vector space, hence free (3.3.126) with a basis of size $n := [K : k(x)]$; extension of scalars (3.4.26) then makes $K_{k'} \cong K \otimes_{k(x)} k'(x)$ a free $k'(x)$-module with a basis of the same size $n$, in particular finite over $k'(x)$. So $K_{k'}$ is a finitely generated field extension of $k'$ (generated by $x$ together with a finite $k'(x)$-basis) of transcendence degree 1: an algebraic function field.

**Degree.** If $k'/k$ is finite, $k'$ is a free $k$-module with a basis of size $[k' : k]$, so extension of scalars along $k \hookrightarrow K$ (3.4.26) makes $K_{k'} = K \otimes_k k'$ a free $K$-module with a basis of the same size, giving $[K_{k'} : K] = [k' : k]$. ∎

**Proposition 6.3.44 (Constant Field Extension is Unramified)**
*Let $P$ be a place of $K/k$ and $k'/k$ algebraic. Every place $P'$ of $K_{k'}/k'$ lying over $P$ is unramified.*

*There are exactly $r$ such places, where $r$ is the number of maximal ideals of $A := k(\mathfrak{m}_P) \otimes_k k'$ (finite, even when $k'/k$ is not). Moreover each such $P'$ satisfies $\mathcal{O}_{P'} \cap K = \mathcal{O}_P$ and $\mathfrak{m}_{P'} \cap K = \mathfrak{m}_P$.*

*More precisely, writing $\mathcal{O}_{P,k'} := \mathcal{O}_P \otimes_k k'$, $A \cong \mathcal{O}_{P,k'}/\mathfrak{m}_P\mathcal{O}_{P,k'}$ as $k'$-algebras (3.4.42) there is a bijection*

$$\begin{aligned} \operatorname{Specm}(k(\mathfrak{m}_P) \otimes_k k') &\longleftrightarrow \{\text{places of } K_{k'}/k' \text{ lying over } P\} \\ \mathfrak{p}'/\mathfrak{m}_P\mathcal{O}_{P,k'} &\longrightarrow \left(\mathcal{O}_{P,k'}\right)_{\mathfrak{p}'} \\ \left(\mathfrak{m}_{P'} \cap \mathcal{O}_{P,k'}\right)/\mathfrak{m}_P\mathcal{O}_{P,k'} &\longleftarrow \mathcal{O}_{P'} \end{aligned}$$

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

We claim that we are in the situation of (3.36.47), applied with $\mathcal{O} := \mathcal{O}_P$, $F := K$:

- $\mathcal{O}_P$ is a discrete valuation ring containing $k$ (a place of $K/k$ is by definition a valuation ring $k \subsetneq \mathcal{O}_P \subsetneq K$ (6.2.112), automatically discrete (6.2.114))
- with fraction field $K$ which is geometrically integral (the standing hypothesis of this section)
- $k(\mathfrak{m}_P)$ is finite over $k$ (6.2.114)
- every valuation ring $R$ of $K_{k'}/k'$ with $k' \subset R$ and $\mathcal{O}_P \subset R$ satisfies $R \cap K = \mathcal{O}_P$: such an $R$ is exactly a place of $K_{k'}/k'$, and this is (6.2.119) (applied with $L := K_{k'}$, $k'' := k'$)

Therefore the result follows from there: writing $\mathcal{O}_{P,k'} := \mathcal{O}_P \otimes_k k'$, its $\mathfrak{b} := \mathfrak{m}_P\mathcal{O}_{P,k'}$ has quotient $\mathcal{O}_{P,k'}/\mathfrak{b} \cong A$ (above), so a) gives exactly $r$ maximal ideals $\mathfrak{p}_1, \ldots, \mathfrak{p}_r$ of $\mathcal{O}_{P,k'}$; by b), each $\left(\mathcal{O}_{P,k'}\right)_{\mathfrak{p}_j}$ is a place $P'_j$ of $K_{k'}/k'$ (with $\mathcal{O}_{P'_j} := \left(\mathcal{O}_{P,k'}\right)_{\mathfrak{p}_j}$), unramified over $\mathcal{O}_P$, and, as in the statement, corresponds to $\mathfrak{p}_j/\mathfrak{b} \in \operatorname{Specm}(A)$; and, by c) together with the last point above, *every* place $P'$ of $K_{k'}/k'$ over $P$ has $\mathcal{O}_{P'} = \left(\mathcal{O}_{P,k'}\right)_{\mathfrak{p}_j}$ for some $j$, that is $P' = P'_j$, giving the stated bijection. As $K_{k'}/k'$ is itself an algebraic function field (6.3.43), every place of $K_{k'}/k'$ – in particular $P'_j$ – is automatically a discrete valuation ring (6.2.114), so $e(P'_j/P) = v_{P'_j}(\pi_P) = 1$: unramified, with $\pi_P$ a uniformiser of $\mathcal{O}_P$ (3.31.2). Finally $\mathcal{O}_{P'_j} \cap K = \mathcal{O}_P$ and $\mathfrak{m}_{P'_j} \cap K = \mathfrak{m}_P$ by e). ∎

**Lemma 6.3.45 (A Rational Place Extends Uniquely Under Any Algebraic Base Change)**
*Let $P$ be a rational place of $K/k$. Then for any algebraic $k'/k$, there is a unique place $Q$ of $K_{k'}$ over $P$, and $e(Q/P) = 1$, with $Q$ again rational over $k'$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

This is the case $k(\mathfrak{m}_P) = k$ of (6.3.44). ∎

**Proposition 6.3.46 (Complete Splitting)**
*Let $K/k$ be an algebraic function field, $P$ a place of $K/k$, and $k'/k$ an algebraic extension. Suppose further that $k'$ contains a normal closure of $k(\mathfrak{m}_P)$.*

*Then $P$ **splits completely** in $K_{k'}/k'$: that is there are exactly $n = [k(\mathfrak{m}_P) : k]$ places above $P$, each rational over $k'$.*

*This holds in particular when $k' = \overline{k}$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Write $k_P := k(\mathfrak{m}_P)$; $k_P/k$ is finite (6.2.114) and separable, as $k$ is perfect (3.19.103). As $k'$ contains a normal closure of $k_P/k$, (3.36.15) gives $k_P \otimes_k k' \cong (k')^n$ as $k'$-algebras, with exactly $n$ maximal ideals, each with residue field $k'$

© David Rufino CC BY-NC-ND 4.0

(the coordinate kernels). By (6.3.44) (with $A \coloneqq k_P \otimes_k k'$) this implies that $P$ splits completely into $n$ places, each rational (residue field $k'$). ∎

**Proposition 6.3.47 (Automorphisms Act Transitively on Places over $\overline{k}$)**
*Let $P$ be a place of $K/k$, $d \coloneqq [k(\mathfrak{m}_P) : k]$, and $P'_1, ..., P'_d$ the places of $K_{\overline{k}}/\overline{k}$ over $P$ ((6.3.46)), with $k' \coloneqq \overline{k}$). For $\sigma \in \mathrm{Aut}\left(\overline{k}/k\right)$, the map*

$$\sigma_* : K_{\overline{k}} \to K_{\overline{k}}, \qquad x \otimes c \mapsto x \otimes \sigma(c)$$

*is a well-defined ring automorphism of $K_{\overline{k}}$ fixing $K$ pointwise, and $\sigma \mapsto \sigma_*$ is a group action of $\mathrm{Aut}\left(\overline{k}/k\right)$ on $K_{\overline{k}}$.*

*For any place $P'$ of $K_{\overline{k}}/\overline{k}$ lying over $P$ then $\sigma_*(\mathcal{O}_{P'})$ is a place lying over $P$ and*

$$v_{\sigma(P')}(\sigma_*(z)) = v_{P'}(z)$$

*for all $z \in K_{\overline{k}}^\star$; this defines a transitive group action of $\mathrm{Aut}\left(\overline{k}/k\right)$ on $\{P'_1, ..., P'_d\}$.*

*Moreover, for $P'$ among $P'_1, ..., P'_d$ (rational over $\overline{k}$), write $\iota_{P'} : k(\mathfrak{m}_P) \hookrightarrow k(\mathfrak{m}_{P'}) = \overline{k}$ for the residue embedding induced by $\mathcal{O}_P \subset \mathcal{O}_{P'}$ (well-defined as $\mathfrak{m}_P \subset \mathfrak{m}_{P'} \cap \mathcal{O}_P$). Then $\iota_{\sigma(P')} = \sigma \circ \iota_{P'}$.*

*This yields a bijection*

$$\begin{aligned} \{\text{places of } \ K/k\} &\longleftrightarrow \left\{\mathrm{Aut}\left(\overline{k}/k\right)\text{-orbits of places of } \ K_{\overline{k}}/\overline{k}\right\} \\ P &\longrightarrow \{P'_1, ..., P'_d\} \end{aligned}$$

*Proof.* As $\sigma : \overline{k} \to \overline{k}$ is a $k$-algebra homomorphism, $1_K \otimes \sigma$ is a well-defined $k$-algebra endomorphism of $K \otimes_k \overline{k} = K_{\overline{k}}$ (3.4.36).

As $(1_K \otimes \sigma) \circ (1_K \otimes \tau) = 1_K \otimes (\sigma\tau)$ then $\sigma_*$ is an automorphism of $K_{\overline{k}}/K$ and $\sigma \mapsto \sigma_*$ is a group homomorphism.

As $\sigma_*$ is a field automorphism of $K_{\overline{k}}$ then $\sigma_*(\mathcal{O}_{P'})$ is a place of $K_{\overline{k}}$ which lies over the same place as $\mathcal{O}_{P'}$, as $\sigma_*$ fixes $K \supset \mathcal{O}_P$.

The valuation identity follows because $\sigma_*$ restricts to an isomorphism on the valuation rings.

As $k$ is perfect, $k(\mathfrak{m}_P)/k$ is separable (3.19.103), so (3.36.15) gives a canonical isomorphism of $\overline{k}$-algebras

$$k(\mathfrak{m}_P) \otimes_k \overline{k} \cong \overline{k}^d, \quad p(\theta) \otimes c \mapsto (c\,p(\theta_1), ..., c\,p(\theta_d)) \qquad \left(p \in k[X], c \in \overline{k}\right)$$

identifying the $d$ maximal ideals of this fibre with $\mathfrak{p}_j$, the kernel of the $j$-th coordinate.

Let $\mathfrak{p}'_1, ..., \mathfrak{p}'_d$ denote the corresponding maximal ideals of $\mathcal{O}_{P,\overline{k}} = \mathcal{O}_P \otimes_k \overline{k}$. Define $\mathcal{O}_{P'_i} \coloneqq \left(\mathcal{O}_{P,\overline{k}}\right)_{\mathfrak{p}'_i}$ then by (6.3.44) these are precisely the places $P'_1, ..., P'_d$ above $P$, and $\mathcal{O}_{P'_i} \cap K = \mathcal{O}_P$, which shows that the given map is injective.

By (3.19.66) $\mathrm{Aut}\left(\overline{k}/k\right)$ permutes the roots $\theta_i$ transitively. We may show that if $\sigma(\theta_j) = \theta_{j'}$ then $\sigma_*\left(\mathcal{O}_{P'_j}\right) = \mathcal{O}_{P'_{j'}}$ and the action on places is transitive and $\{P'_1, ..., P'_d\}$ is precisely an $\mathrm{Aut}\left(\overline{k}/k\right)$-orbit.

Given a place $\overline{P}$ of $K_{\overline{k}}/\overline{k}$, $\mathcal{O} \coloneqq \mathcal{O}_{\overline{P}} \cap K$ is a (proper) valuation ring of $K$ by (6.2.118), giving a place $P$ of $K/k$ with $\mathcal{O}_P = \mathcal{O} \subset \mathcal{O}_{\overline{P}}$, that is $\overline{P}$ lies over $P$. As $P'_1, ..., P'_d$ are all the places of $K_{\overline{k}}/\overline{k}$ above $P$ (above), $\overline{P}$ is one of them. So the given map is a surjection. ∎

**Lemma 6.3.48 (Residue Embeddings are Exactly the Embeddings into $\overline{k}$)**
*Let $P$ be a place of $K/k$, $d \coloneqq [k(\mathfrak{m}_P) : k]$, and $P'_1, ..., P'_d$ the places of $K_{\overline{k}}/\overline{k}$ over $P$ ((6.3.46)), with $k' \coloneqq \overline{k}$).*

*Denote by $\iota_j : k(\mathfrak{m}_P) \hookrightarrow k\left(\mathfrak{m}_{P'_j}\right) = \overline{k}$ the corresponding embeddings, then these exhaust all the $k$-algebra embeddings $k(\mathfrak{m}_P)/k \hookrightarrow \overline{k}/k$.*

*Proof.* By (6.3.47) there are $d = [k(\mathfrak{m}_P) : k]$ places and, as $k$ is perfect so $k(\mathfrak{m}_P)/k$ is separable (3.19.103), there are $\#\,\mathrm{Mor}_k\left(k(\mathfrak{m}_P), \overline{k}\right) = [k(\mathfrak{m}_P) : k]_s = d$ $k$-algebra embeddings (3.19.83).

We show that $j \mapsto \iota_j$ is injective, i.e. that $\iota_j = \iota_{j'}$ implies $\mathcal{O}_{P'_j} = \mathcal{O}_{P'_{j'}}$ (and so $j = j'$). Consider the composite $\rho_j : \mathcal{O}_{P,\overline{k}} \to \left(\mathcal{O}_{P,\overline{k}}\right)_{\mathfrak{p}'_j} = \mathcal{O}_{P'_j} \to \mathcal{O}_{P'_j}/\mathfrak{m}_{P'_j} = \overline{k}$. It is $\overline{k}$-linear and is uniquely determined by

$$\rho_j(a \otimes c) = c\,\iota_j(\overline{a}) \qquad \left(a \in \mathcal{O}_P, c \in \overline{k}\right),$$

so $\iota_j = \iota_{j'} \implies \rho_j = \rho_{j'}$, and their kernels agree, that is $\mathfrak{p}'_j = \mathfrak{p}'_{j'}$ (as $\ker \rho_j = \mathfrak{m}_{P'_j} \cap \mathcal{O}_{P,\overline{k}} = \left(\mathfrak{p}'_j\left(\mathcal{O}_{P,\overline{k}}\right)_{\mathfrak{p}'_j}\right)^c = \mathfrak{p}'_j$ (3.6.42)). By (6.3.44) $P'_j = P'_{j'}$ and we are done by counting. ∎

© David Rufino CC BY-NC-ND 4.0

### 6.3.8 Residue Symbol at a General Place

At a non-rational place there is no element of $k$ to reduce against: (6.3.23)'s recursion needs, at each step, a canonical way to lift a class of $k(\mathfrak{m}_P)$ back into $\mathcal{O}_P$, and $k(\mathfrak{m}_P) \neq k$ means $k \subset \mathcal{O}_P$ no longer supplies one. Rather than search for a section of $\mathcal{O}_P \twoheadrightarrow k(\mathfrak{m}_P)$ directly, we base change to $\overline{k}$ instead: $P$ splits completely into $d := [k(\mathfrak{m}_P) : k]$ *rational* places of $K_{\overline{k}}/\overline{k}$, each of which already has the Laurent coefficient theory of (6.3.23) available. The $d$ resulting elements of $\overline{k}$ are exactly the Galois conjugates of a single element of $k(\mathfrak{m}_P)$, which we take as the residue at $P$.

**Definition 6.3.49 (Conjugate Places)**
*Let $P$ be a place of $K/k$, $d := [k(\mathfrak{m}_P) : k]$, and $\iota_1, \dots, \iota_d : k(\mathfrak{m}_P) \hookrightarrow \overline{k}$ the embeddings, with $P'_j$ the corresponding place of $K_{\overline{k}}/\overline{k}$ over $P$, $\iota_j = \iota_{P'_j}$ (6.3.48). As each $P'_j/P$ is unramified, every uniformiser $\pi$ at $P$ remains a uniformiser at $P'_j$ (6.3.44), so the residue symbol $\mathrm{Res}_{P'_j}(w\,\mathrm{d}s)$ at the rational place $P'_j$ of $K_{\overline{k}}/\overline{k}$ (6.3.25) is defined for $w, s \in K_{\overline{k}}$.*

**Lemma 6.3.50 (The Uniformiser Derivation Extends to the Base Change)**
*Let $P$ be a place of $K/k$, $\pi$ a uniformiser, and $P'_1, \dots, P'_d$ the conjugate rational places of $K_{\overline{k}}/\overline{k}$ over $P$ (6.3.49), at each of which $\pi \otimes 1$ remains a uniformiser.*

*Then $\partial_{\pi\otimes 1} \in \mathrm{Der}_{\overline{k}}(K_{\overline{k}}, K_{\overline{k}})$ is the unique $D$ satisfying*

$$D|_K = \partial_\pi$$

*and is in particular regular at every place $P'_j$.*

**Proposition 6.3.51 (Conjugate Residues are $\mathrm{Aut}\left(\overline{k}/k\right)$-Equivariant)**
*For $\sigma \in \mathrm{Aut}\left(\overline{k}/k\right)$, $j = 1, \dots, d$, writing $j'$ for the index with $\sigma\left(P'_j\right) = P'_{j'}$ (6.3.47), and $t, z \in K$,*

$$\sigma\Big(\mathrm{Res}_{P'_j}((z\otimes 1)\,\mathrm{d}(t\otimes 1))\Big) = \mathrm{Res}_{P'_{j'}}((z\otimes 1)\,\mathrm{d}(t\otimes 1))$$

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Fix a uniformiser $\pi$ at $P$, and write $w := z\,\partial_\pi(t) \in K$; by (6.3.50), $\mathrm{Res}_{P'_j}((z\otimes 1)\,\mathrm{d}(t\otimes 1)) = L_\pi(w\otimes 1)_{-1}$ (and likewise at $P'_{j'}$). Write $\sigma_*$ for the induced automorphism of $K_{\overline{k}}$ (6.3.47), fixing $K$ pointwise, and $c_i$ for the $i$-th Laurent coefficient of $w \otimes 1$ at $P'_j$ relative to $\pi$, so for every $n$,

$$w\otimes 1 - \sum_{i\le n} c_i \pi^i \in \mathfrak{m}_{P'_j}^{n+1}$$

Applying $\sigma_*$ (a ring automorphism fixing $w\otimes 1$ and $\pi$, both in $K$) and using $\sigma_*\left(\mathfrak{m}_{P'_j}^{n+1}\right) = \mathfrak{m}_{P'_{j'}}^{n+1}$ (as $\sigma_*$ restricts to an isomorphism $\mathcal{O}_{P'_j} \xrightarrow{\sim} \mathcal{O}_{P'_{j'}}$ (6.3.47)),

$$w\otimes 1 - \sum_{i\le n} \sigma(c_i)\pi^i \in \mathfrak{m}_{P'_{j'}}^{n+1}$$

for every $n$. By uniqueness of Laurent coefficients ((6.3.23), at $P'_{j'}$), $(\sigma(c_i))_i$ are exactly the Laurent coefficients of $w\otimes 1$ at $P'_{j'}$ relative to $\pi$; taking $i = -1$, $\sigma(L_\pi(w\otimes 1)_{-1}) = \sigma(c_{-1}) = L_\pi(w\otimes 1)_{-1}$ at $P'_{j'}$, that is $\sigma\Big(\mathrm{Res}_{P'_j}((z\otimes 1)\,\mathrm{d}(t\otimes 1))\Big) = \mathrm{Res}_{P'_{j'}}((z\otimes 1)\,\mathrm{d}(t\otimes 1))$. ∎

**Proposition 6.3.52 (Conjugate Residue Sum is a Trace)**
*For $t, z \in K$, there is a single $\alpha \in k(\mathfrak{m}_P)$ with $\iota_j(\alpha) = \mathrm{Res}_{P'_j}((z\otimes 1)\,\mathrm{d}(t\otimes 1))$ for every $j = 1, \dots, d$ (well-defined, see proof), and*

$$\sum_{j=1}^{d} \mathrm{Res}_{P'_j}((z\otimes 1)\,\mathrm{d}(t\otimes 1)) = \mathrm{Tr}_{k(\mathfrak{m}_P)/k}(\alpha) \in k$$

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Write $R_j := \mathrm{Res}_{P'_j}((z\otimes 1)\,\mathrm{d}(t\otimes 1))$ for brevity.

**$R_1 \in \iota_1(k(\mathfrak{m}_P))$.** Write $F := \iota_1(k(\mathfrak{m}_P)) \subset \overline{k}$. As $\overline{k}/k$ is Galois ($k$ perfect) with subfield $F$, $\overline{k}/F$ is itself Galois (3.19.116), with $\mathrm{Gal}\left(\overline{k}/F\right) = \mathrm{Aut}\left(\overline{k}/F\right)$ (3.19.51). For $\sigma \in \mathrm{Aut}\left(\overline{k}/F\right)$, $\sigma$ fixes $F = \iota_1(k(\mathfrak{m}_P))$ pointwise, that is $\sigma \circ \iota_1 = \iota_1$, so $\sigma(P'_1) = P'_1$ (6.3.47), and (6.3.51) gives $\sigma(R_1) = R_1$: that is $R_1 \in \overline{k}^{\mathrm{Gal}(\overline{k}/F)}$. So (3.19.117) gives $\overline{k}^{\mathrm{Gal}(\overline{k}/F)} = F$, that is $R_1 \in F = \iota_1(k(\mathfrak{m}_P))$; write $\alpha := \iota_1^{-1}(R_1) \in k(\mathfrak{m}_P)$.

**$\iota_j(\alpha) = R_j$ for every $j$.** By transitivity (6.3.47), choose $\sigma \in \mathrm{Aut}\left(\overline{k}/k\right)$ with $\sigma(P'_1) = P'_j$, so $\sigma \circ \iota_1 = \iota_j$ (6.3.47). By equivariance (6.3.51), $\iota_j(\alpha) = (\sigma\circ\iota_1)(\alpha) = \sigma(\iota_1(\alpha)) = \sigma(R_1) = R_j$, the single element promised in the statement.

© David Rufino CC BY-NC-ND 4.0

**Conclusion.** By (6.3.48), $\mathrm{Mor}_k\big(k(\mathfrak{m}_P),\overline{k}\big)=\{\iota_1,\dots,\iota_d\}$; as $k$ is perfect, $k(\mathfrak{m}_P)/k$ is separable (3.19.103), so $[k(\mathfrak{m}_P):k]_i=1$ (3.19.93). So (3.19.129) (trace is sum of embeddings in $\overline{k}$) gives

$$\mathrm{Tr}_{k(\mathfrak{m}_P)/k}(\alpha)=\sum_{j=1}^{d}\iota_j(\alpha)=\sum_{j=1}^{d}R_j$$

In particular $\sum_{j=1}^{d}R_j\in k$, $\mathrm{Tr}_{k(\mathfrak{m}_P)/k}$ taking values in $k$ by definition (3.19.122). ∎

**Definition 6.3.53 (Residue Symbol at a General Place)**
*Let $P$ be a place of $K/k$, and $t,z\in K$. Define*

$$\mathrm{Res}_P(z\,\mathrm{d}t):=\sum_{j=1}^{d}\mathrm{Res}_{P_j'}((z\otimes 1)\,\mathrm{d}(t\otimes 1))\in k$$

*(residues at the conjugate rational places $P_1',\dots,P_d'$, (6.3.49)) – landing in $k$ by (6.3.52).*

Once the module of differentials $\Omega_{K/k}$ is available, it is a 1-dimensional $K$-vector space dual to $\mathrm{Der}_k(K,K)$ (both being $\mathrm{trdeg}_k(K)=1$-dimensional, (3.35.44)), and the pairing of $z\,\mathrm{d}t\in\Omega_{K/k}$ with $\partial_\pi$ is exactly $z\,\partial_\pi(t)$; $\mathrm{Res}_P(z\,\mathrm{d}t)$ above is then the residue, in this sense, of the differential $z\,\mathrm{d}t$.

**Proposition 6.3.54 (Elementary Properties of the Residue Symbol at a General Place)**
*Let $P$ be a place of $K/k$, $t\in K$. Then*

*a) $z\mapsto\mathrm{Res}_P(z\,\mathrm{d}t)$ is $k$-linear $K\to k$;*
*b) if $P$ is rational (so $d=1$, $P_1'$ canonically identified with $P$), $\mathrm{Res}_P(z\,\mathrm{d}t)$ agrees with (6.3.25), for every $z\in K$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

***k*-linearity.** Fix a uniformiser $\pi$ at $P$. For $z,z'\in K$ and $a,b\in k$, $(az+bz')\partial_\pi(t)=a(z\partial_\pi(t))+b(z'\partial_\pi(t))$ ($K$-bilinearity of multiplication, $k$-linearity of $\partial_\pi$), so by (6.3.50) and $k$-linearity of $L_\pi(-)_{-1}$ ((6.3.23) and (6.3.24)),

$$\mathrm{Res}_{P_j'}(((az+bz')\otimes 1)\,\mathrm{d}(t\otimes 1))=a\,\mathrm{Res}_{P_j'}((z\otimes 1)\,\mathrm{d}(t\otimes 1))+b\,\mathrm{Res}_{P_j'}((z'\otimes 1)\,\mathrm{d}(t\otimes 1))$$

for every $j$; summing over $j$ gives $\mathrm{Res}_P((az+bz')\,\mathrm{d}t)=a\,\mathrm{Res}_P(z\,\mathrm{d}t)+b\,\mathrm{Res}_P(z'\,\mathrm{d}t)$.

**Rational case.** If $P$ is rational, $d=1$, and $P_1'$ is identified with $P$ under $K\subset K_{\overline{k}}$ (with any uniformiser $\pi$ at $P$ remaining one at $P_1'$); the sum defining $\mathrm{Res}_P(z\,\mathrm{d}t)$ then has the single term $\mathrm{Res}_{P_1'}((z\otimes 1)\,\mathrm{d}(t\otimes 1))=\mathrm{Res}_P(z\,\mathrm{d}t)$ of (6.3.25) applied verbatim (as $z\otimes 1,t\otimes 1$ agree with $z,t$ under $K\subset K_{\overline{k}}$, and $P_1'$ with $P$). ∎

**Lemma 6.3.55 (Regular Differentials have Zero Residue)**
*Let $P$ be a place of $K/k$, $\pi$ a uniformiser at $P$, and $t,z\in K$ with $z\,\partial_\pi(t)\in\mathcal{O}_P$. Then $\mathrm{Res}_P(z\,\mathrm{d}t)=0$. In particular this holds if $P$ is rational and $z,t\in\mathcal{O}_P$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Write $w:=z\,\partial_\pi(t)\in\mathcal{O}_P$. By (6.3.50), for every $j$,

$$\mathrm{Res}_{P_j'}((z\otimes 1)\,\mathrm{d}(t\otimes 1))=L_\pi(w\otimes 1)_{-1}.$$

As $P_j'$ lies over $P$, $w\otimes 1\in\mathcal{O}_P\otimes 1\subseteq\mathcal{O}_{P_j'}$, so $L_\pi(w\otimes 1)$ has no negative powers (6.3.23) and its $X^{-1}$-coefficient is 0. Summing over $j$ gives $\mathrm{Res}_P(z\,\mathrm{d}t)=0$. If $P$ is rational and $t\in\mathcal{O}_P$ then $\partial_\pi(t)\in\mathcal{O}_P$ (6.3.30), so $z\,\partial_\pi(t)\in\mathcal{O}_P$ when also $z\in\mathcal{O}_P$. ∎

**Proposition 6.3.56 (Change of Base Differential)**
*Let $P$ be a place of $K/k$, $x\in K$ transcendental over $k$, and $\partial_x\in\mathrm{Der}_k(K,K)$ the distinguished derivation of (6.3.20). For $t,z\in K$,*

$$\mathrm{Res}_P(z\,\mathrm{d}t)=\mathrm{Res}_P((z\cdot\partial_x(t))\,\mathrm{d}x).$$

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

Fix a uniformiser $\pi$ at $P$. By (6.3.20) (applied to $D:=\partial_\pi$), $\partial_\pi=\partial_\pi(x)\cdot\partial_x$, so $z\,\partial_\pi(t)=(z\cdot\partial_x(t))\,\partial_\pi(x)$. By (6.3.50), for every $j$,

$$\mathrm{Res}_{P_j'}((z\otimes 1)\,\mathrm{d}(t\otimes 1))=L_\pi((z\,\partial_\pi(t))\otimes 1)_{-1}=L_\pi(((z\cdot\partial_x(t))\,\partial_\pi(x))\otimes 1)_{-1}=\mathrm{Res}_{P_j'}(((z\cdot\partial_x(t))\otimes 1)\,\mathrm{d}(x\otimes 1))$$

Summing over $j$ (6.3.53) gives $\mathrm{Res}_P(z\,\mathrm{d}t)=\mathrm{Res}_P((z\cdot\partial_x(t))\,\mathrm{d}x)$. ∎

© David Rufino CC BY-NC-ND 4.0

### 6.3.9 The Algebraic Residue Theorem

This subsection proves the (algebraic) Residue Theorem for a general base field $k$, by reduction to the algebraically closed case of (6.3.34).

**Proposition 6.3.57 (Algebraic Residue Theorem)**
*For $t, f \in K$,*

$$\sum_P \operatorname{Res}_P(f\,\mathrm{d}t) = 0,$$

*summed over all places $P$ of $K/k$ (finitely many terms nonzero, as $f\,\mathrm{d}t$ has only finitely many poles).*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

By (6.3.47) (applied to $K/k$), restriction $\overline{P} \mapsto \overline{P} \cap K$ is a bijection $\{\text{places of } K/k\} \leftrightarrow \left\{\operatorname{Aut}\left(\overline{k}/k\right)\text{-orbits of places of } K_{\overline{k}}/\overline{k}\right\}$ (where $K_{\overline{k}} := K \otimes_k \overline{k}$, a field by (3.36.45) as in (6.3.43)); the fibre over $P$ is $\left\{P'_1, \ldots, P'_{d_P}\right\}$, $d_P := \deg P$. By (6.3.53), for every place $P$ of $K/k$ and $z \in K$,

$$\operatorname{Res}_P(z\,\mathrm{d}t) = \sum_{j=1}^{d_P} \operatorname{Res}_{P'_j}((z \otimes 1)\,\mathrm{d}(t \otimes 1)).$$

Summing over all places $P$ of $K/k$ and using that the fibres $\left\{P'_1, \ldots, P'_{d_P}\right\}$ partition the places of $K_{\overline{k}}/\overline{k}$ (restriction being a surjection),

$$\sum_{P \text{ of } K/k} \operatorname{Res}_P(f\,\mathrm{d}t) = \sum_{\overline{P} \text{ of } K_{\overline{k}}/\overline{k}} \operatorname{Res}_{\overline{P}}((f \otimes 1)\,\mathrm{d}(t \otimes 1)) = 0$$

by (6.3.34) (applicable as $K_{\overline{k}}/\overline{k}$ is itself an algebraic function field, over the algebraically closed field $\overline{k}$, (6.3.43)), applied to $K_{\overline{k}}/\overline{k}$, $t \otimes 1$, $f \otimes 1$. ∎

### 6.3.10 Differentials are Weil Differentials, for General $k$

Fix a transcendental $t \in K$. By (6.3.20), every $D \in \operatorname{Der}_k(K, K)$ is $D(t) \cdot \partial_t$, so a differential $z\,\mathrm{d}t$ ($z \in K$) already carries all the information of an element of the (not yet formally constructed) module $\Omega_{K/k}$, and $\operatorname{Res}_P(z\,\mathrm{d}t)$ is available at every place (6.3.53). We show this recovers $\Omega_K$ exactly, generalising (6.3.19) beyond $k = \mathbb{C}$: the complex-analytic ingredients there (contour integration, Stokes' theorem, Mittag-Leffler) are replaced by the algebraic Residue Theorem, the trace formalism of (6.3.53), and the following fact, taken as given in this subsection.

**Proposition 6.3.58 ($\Omega_K$ is a $K$-Vector Space)**
*Let $K/k$ be an algebraic function field. For $x \in K$ and $\omega \in \Omega_K(D)$ ($D \in \operatorname{Div}(K)$), define $x\omega \in \operatorname{Hom}_k(\mathcal{A}_K, k)$ by $(x\omega)(\alpha) := \omega(x\alpha)$ (using that $\mathcal{A}_K$ is a $K$-algebra). Then $x\omega \in \Omega_K(D - (x)) \subset \Omega_K$, and this action makes $\Omega_K$ a $K$-vector space.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

**$x\omega \in \Omega_K(D - (x))$.** For $\alpha \in \mathcal{A}_K(D - (x))$, $v_P(\alpha_P) \geq -v_P(D - (x)) = v_P(x) - v_P(D)$ for every $P$, so $v_P(x\alpha_P) \geq -v_P(D)$, that is $x\alpha \in \mathcal{A}_K(D)$; and $xK = K$. As $\omega \in \Omega_K(D)$ vanishes on $\mathcal{A}_K(D) + K$, it follows that $(x\omega)(\alpha) = \omega(x\alpha) = 0$ for every $\alpha \in \mathcal{A}_K(D - (x)) + K$, that is $x\omega \in \Omega_K(D - (x))$.

**Module axioms.** For $x, y \in K$, $\lambda \in k$, $\alpha, \beta \in \mathcal{A}_K$: $(x\omega)(\alpha + \beta) = \omega(x\alpha + x\beta) = (x\omega)(\alpha) + (x\omega)(\beta)$ and $(x\omega)(\lambda\alpha) = \omega(\lambda x\alpha) = \lambda(x\omega)(\alpha)$, so $x\omega \in \operatorname{Hom}_k(\mathcal{A}_K, k)$ is again $k$-linear; $(x(y\omega))(\alpha) = (y\omega)(x\alpha) = \omega(yx\alpha) = ((xy)\omega)(\alpha)$ and $1\omega = \omega$; $((x + y)\omega)(\alpha) = \omega((x + y)\alpha) = \omega(x\alpha) + \omega(y\alpha) = (x\omega)(\alpha) + (y\omega)(\alpha)$; and $(\omega \mapsto x\omega)$ is $k$-linear by the same computation applied to $\omega_1 + \omega_2$. These are exactly the axioms making $\Omega_K$ a $K$-module, that is a $K$-vector space (being already a $k$-vector space). ∎

**Proposition 6.3.59 ($\Omega_K$ is 1-Dimensional over $K$)**
*Let $K/k$ be an algebraic function field. Then $\dim_K \Omega_K = 1$ (for the $K$-vector space structure of (6.3.58)).*

**Proposition 6.3.60 (Differentials are Weil Differentials)**
*Let $K/k$ be an algebraic function field and $t \in K$ transcendental. For $z \in K$, define $\varphi_{z\,\mathrm{d}t} \in \operatorname{Hom}_k(\mathcal{A}_K, k)$ by*

$$\varphi_{z\,\mathrm{d}t}(\alpha) := \sum_P \operatorname{Res}_P(\alpha_P z\,\mathrm{d}t), \qquad \alpha = (\alpha_P)_P \in \mathcal{A}_K$$

*(a finite sum: $\alpha_P \in \mathcal{O}_P$ for all but finitely many $P$, and $z\,\mathrm{d}t$ is regular at all but finitely many $P$, so $\operatorname{Res}_P(\alpha_P z\,\mathrm{d}t) = 0$ for almost all $P$ (6.3.55)). Then $\varphi_{z\,\mathrm{d}t} \in \Omega_K$, and*

© David Rufino CC BY-NC-ND 4.0

$$K \longrightarrow \Omega_K, \qquad z \longmapsto \varphi_{z\,\mathrm{d}t}$$

*is a canonical $K$-linear isomorphism (for the $K$-vector space structures of (6.3.58) and $K$ itself).*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

**$\varphi_{z\,\mathrm{d}t} \in \Omega_K$.** Let $D$ be the divisor with $v_P(D) \coloneqq v_P(z\,\partial_\pi(t))$ for a uniformiser $\pi$ at $P$ (independent of the choice of $\pi$: for another uniformiser $\pi'$, $\partial_{\pi'}(t) = \partial_{\pi'}(\pi)\,\partial_\pi(t)$ (6.3.22), with $\partial_{\pi'}(\pi)$ a unit at $P$, so $v_P(z\partial_{\pi'}(t)) = v_P(z\partial_\pi(t))$; and this vanishes for all but finitely many $P$, as $z$ has finitely many poles and $t$ finitely many poles). For $\alpha \in \mathcal{A}_K(D)$, $v_P(\alpha_P) \geq -v_P(D) = -v_P(z\partial_\pi(t))$ for every $P$, so $v_P(\alpha_P z\partial_\pi(t)) \geq 0$, that is $\alpha_P z\,\partial_\pi(t) \in \mathcal{O}_P$, giving $\mathrm{Res}_P(\alpha_P z\,\mathrm{d}t) = 0$ (6.3.55) so $\varphi_{z\,\mathrm{d}t}(\alpha) = 0$. For $f \in K$ embedded diagonally, $\varphi_{z\,\mathrm{d}t}(f, f, \ldots) = \sum_P \mathrm{Res}_P(fz\,\mathrm{d}t) = 0$ by (6.3.57) (the Residue Theorem), applied to $fz \in K$. So $\varphi_{z\,\mathrm{d}t}$ vanishes on $\mathcal{A}_K(D) + K$, that is $\varphi_{z\,\mathrm{d}t} \in \Omega_K(D) \subset \Omega_K$.

***K*-linearity.** $k$-linearity in $z$ is immediate from (6.3.54).a), applied place by place and summed. For $x \in K$, using $x\alpha_P \cdot z = \alpha_P \cdot (xz)$ in the field $K$,

$$(x\,\varphi_{z\,\mathrm{d}t})(\alpha) = \varphi_{z\,\mathrm{d}t}(x\alpha) = \sum_P \mathrm{Res}_P((x\alpha_P)z\,\mathrm{d}t) = \sum_P \mathrm{Res}_P(\alpha_P(xz)\,\mathrm{d}t) = \varphi_{(xz)\,\mathrm{d}t}(\alpha),$$

so $x\,\varphi_{z\,\mathrm{d}t} = \varphi_{(xz)\,\mathrm{d}t}$: the map is $K$-linear (6.3.58).

**Injectivity.** Suppose $\varphi_{z\,\mathrm{d}t} = 0$ and $z \neq 0$; we derive a contradiction. Fix a place $P$ and a uniformiser $\pi$ at $P$, and let $d, P'_1, \ldots, P'_d, \iota_1, \ldots, \iota_d$ be as in (6.3.49). As $\partial_\pi$ is a nonzero derivation and $t$ is transcendental, $\partial_\pi(t) \neq 0$ (6.3.20), so $w \coloneqq z\,\partial_\pi(t) \neq 0$ in $K$. Since $w \in K$ is fixed pointwise by $\mathrm{Aut}\big(\overline{k}/k\big)$ (embedded diagonally in $K_{\overline{k}}$) while $\sigma_*$ permutes $\{P'_1, \ldots, P'_d\}$ preserving valuations (6.3.47), $m \coloneqq v_{P'_j}(w \otimes 1)$ is independent of $j$. Let $c \coloneqq \pi^{-m-1} \in K^\star$; then $v_{P'_j}(cw \otimes 1) = -1$ for every $j$, so

$$u_j \coloneqq \mathrm{Res}_{P'_j}((cz \otimes 1)\,\mathrm{d}(t \otimes 1))$$

is the (nonzero) leading Laurent coefficient of $cw \otimes 1$ at $P'_j$. By (6.3.52) (applied to $t, cz$ in place of $t, z$), $u_j = \iota_j(\gamma)$ for a single $\gamma \in k(\mathfrak{m}_P)$; as $u_1 \neq 0$ and $\iota_1$ is injective, $\gamma \neq 0$. As $k$ is perfect, $k(\mathfrak{m}_P)/k$ is separable (3.19.103), so by (3.19.131) there is $\overline{e} \in k(\mathfrak{m}_P)$ with $\mathrm{Tr}_{k(\mathfrak{m}_P)/k}(\overline{e}\gamma) \neq 0$; lift it to a unit $e \in \mathcal{O}_P^\star$ (any lift of a nonzero residue class is a unit, $\mathcal{O}_P$ being local). As $e \otimes 1$ is a unit of $\mathcal{O}_{P'_j}$ with residue class $\iota_j(\overline{e})$, and $cw \otimes 1$ has valuation exactly $-1$ with leading coefficient $u_j = \iota_j(\gamma)$, the product $(e \otimes 1)(cw \otimes 1)$ again has valuation exactly $-1$ (no lower-order term of $e \otimes 1$ to interact with, as $cw \otimes 1$ has none below $X^{-1}$), with leading coefficient $\iota_j(\overline{e})\,\iota_j(\gamma) = \iota_j(\overline{e}\gamma)$: that is $\mathrm{Res}_{P'_j}((ecz \otimes 1)\,\mathrm{d}(t \otimes 1)) = \iota_j(\overline{e}\gamma)$ for every $j$, so

$$\mathrm{Res}_P(ecz\,\mathrm{d}t) = \sum_{j=1}^{d} \iota_j(\overline{e}\gamma) = \mathrm{Tr}_{k(\mathfrak{m}_P)/k}(\overline{e}\gamma) \neq 0.$$

Testing $\varphi_{z\,\mathrm{d}t}$ against the repartition supported at $P$ with value $ec \in K$ then gives $\varphi_{z\,\mathrm{d}t}(\ldots) = \mathrm{Res}_P(ecz\,\mathrm{d}t) \neq 0$, contradicting $\varphi_{z\,\mathrm{d}t} = 0$.

**Surjectivity.** The map $K \to \Omega_K$ is $K$-linear and injective, and $K$ is a 1-dimensional $K$-vector space over itself; so its image is a 1-dimensional $K$-subspace of $\Omega_K$, which by (6.3.59) is itself 1-dimensional over $K$. So the image is all of $\Omega_K$. ∎

## 6.3.11 Tate's Approach to Residues: A Sketch

The route taken above defines $\mathrm{Res}_P$ through a uniformiser and the formal derivative, and proves the Residue Theorem (6.3.57) by reducing to $K = k(x)$ along a finite separable extension. Tate [Tat68] gives a different, intrinsic route, which we only sketch here (nothing later depends on it): the residue is defined from linear algebra alone, and the Residue Theorem is deduced from the finiteness results of the Riemann-Roch theory.

a) *Traces of finite-rank operators.* The trace of a finite-rank endomorphism $T$ of a $k$-vector space $V$ is its trace on any finite-dimensional subspace containing $T(V)$. It is additive, $\mathrm{Tr}(AB) = \mathrm{Tr}(BA)$ whenever $A : V \to W$, $B : W \to V$ and one of them has finite rank, and $\mathrm{Tr}(T) = \mathrm{Tr}(T|_H) + \mathrm{Tr}\big(\overline{T}\big)$ if $T$ preserves a finite-dimensional $H$, with $\overline{T}$ the induced operator on $V/H$. In particular $\mathrm{Tr}([T, F]) = 0$ for $F$ of finite rank and $T$ arbitrary.

b) *The local residue.* Let $\pi$ be a $k$-linear retraction of the local space onto the lattice $\mathcal{O}_P$ (in Tate's setting the completion $K_P$, with $\pi$ the projection killing the principal parts). For $f$ in the local space let $T_f \coloneqq \pi \circ (f \cdot -)|_{\mathcal{O}_P}$ be the Toeplitz operator on $\mathcal{O}_P$. Then $\big[T_f, T_g\big]$ has finite rank, because $(1 - \pi)(f\mathcal{O}_P)$ lies in the finite-dimensional image of $t^{-n}\mathcal{O}_P/\mathcal{O}_P$, and $\mathrm{Res}_P(f\,\mathrm{d}g) \coloneqq \mathrm{Tr}\big(\big[T_f, T_g\big]\big)$. Changing $\pi$ changes each $T_f$ by a finite-rank operator, which does not change the trace of the commutator. So no uniformiser, derivation or chain rule enters.

© David Rufino CC BY-NC-ND 4.0

c) *Local properties.* It is $k$-linear, vanishes on differentials regular at $P$, has $\operatorname{Res}_P(\mathrm{d}t/t) = \deg P$ (the trace is taken over $k$) and is invariant under automorphisms of $K_P$; it should agree with the trace of the coefficient of $t^{-1}$ used above.

d) *Global input.* The finiteness underlying Riemann-Roch: $H \coloneqq K \cap \mathcal{A}_K(0) = L(0)$ and $\mathcal{A}_K/(K + \mathcal{A}_K(0))$ are finite dimensional (6.3.15) and $\mathcal{A}_K(D)/\mathcal{A}_K(0) = \bigoplus_P t_P^{-n_P}\mathcal{O}_P/\mathcal{O}_P$ is finite dimensional for every divisor $D = \sum_P n_P P$.

e) *The Residue Theorem as a trace identity.* Write $\mathcal{A} \coloneqq \mathcal{A}_K$ and $L \coloneqq \mathcal{A}(0) = \prod_P \mathcal{O}_P$, and for $f, g \in K$ let $D$ be a divisor with $f\mathcal{A}(0), g\mathcal{A}(0) \subset \mathcal{A}(D)$.

   (a) For the retraction $\pi_L \coloneqq \prod_P \pi_P : \mathcal{A} \to L$ and $T_f \coloneqq \pi_L \circ (f \cdot -)|_L$, the commutator $\big[T_f, T_g\big]$ is the direct sum over $P$ of the local commutators, of which only finitely many are nonzero (where $f, g$ are regular the operators are multiplication by $f, g$ and commute). So $\operatorname{Tr}\big(\big[T_f, T_g\big]\big) = \sum_P \operatorname{Res}_P(f\,\mathrm{d}g)$.

   (b) As in b), $\operatorname{Tr}\big(\big[T_f, T_g\big]\big)$ does not depend on the retraction $\pi : \mathcal{A} \to L$: another retraction changes $T_f$ by $\psi \circ (1-\pi)f|_L$, which has finite rank since $(1-\pi)(\mathcal{A}(D))$ is finite dimensional.

   (c) Choose $\pi$ adapted to $K$: write $K = H \oplus W_K$ and choose a finite-dimensional $F$ with $\mathcal{A} = L \oplus W_K \oplus F$ (possible by d)), and let $\pi$ be the projection onto $L$ along $W_K \oplus F$. For $h \in H$, $fh \in K = H \oplus W_K$ so $T_f h \in H$: both $T_f, T_g$ preserve $H$ and act there as operators on a finite-dimensional space, whose commutator has trace 0.

   (d) On $L/H \cong (L+K)/K =: \Lambda \subset \mathcal{A}/K$ the induced operator $\overline{T}_f$ is the Toeplitz operator of multiplication by $f$ on $\mathcal{A}/K$ (where $f, g$ commute exactly), with respect to the retraction $\mathcal{A}/K = \Lambda \oplus \overline{F} \to \Lambda$. Writing $\overline{\pi} = 1 - p$ with $p$ the projection onto $\overline{F}$, and using $fg = gf$ on $\mathcal{A}/K$, one finds $\big[\overline{T}_f, \overline{T}_g\big] = -\overline{\pi}fpg + \overline{\pi}gpf$ on $\Lambda$, whose trace equals $\operatorname{Tr}_{\overline{F}}([pg|_{\overline{F}}, pf|_{\overline{F}}]) = 0$, a commutator of operators on the finite-dimensional space $\overline{F}$.

   By (c), (d) and additivity of the trace, $\operatorname{Tr}\big(\big[T_f, T_g\big]\big) = 0$, so $\sum_P \operatorname{Res}_P(f\,\mathrm{d}g) = 0$ by (a) and (b).

Nothing here uses the topology of $K_P$. The construction of b) uses only the subspace $\mathcal{O}_P \subset K$, a retraction $\pi : K \to \mathcal{O}_P$, and $\dim_k t^{-n}\mathcal{O}_P/\mathcal{O}_P < \infty$ (as $k(\mathfrak{m}_P)/k$ is finite): "lattices" and "continuity" reduce to commensurability of subspaces and to the pole order of $f$, so the local residue may be defined on $K$ itself, and the global step on the repartitions $\mathcal{A}_K$ of this book (components in $K$, not in the $K_P$) together with the finiteness of d). Compared with the proof above, this avoids the reduction to $k(x)$, the trace formula and separability, and works in any characteristic. What is left to check is that this $\operatorname{Res}_P$ agrees with the Laurent-coefficient residue used above (for a place of degree $d$, $\operatorname{Res}_P(\mathrm{d}t/t) = d$ with the trace taken over $k$); the global argument of e) is then complete.

### 6.3.12 Sheaf-Theoretic Interpretation of Repartitions

Unlike the rest of this chapter, this section depends on sheaf theory developed later in this book (4.2.2), and so is not logically self-contained at this point in the text; nothing later in this chapter depends on it.

Let $X \coloneqq \mathbb{P}_K$, the set of places of $K/k$, equipped with the **cofinite topology** (opens are $\varnothing$ and complements of finite subsets); as $\mathbb{P}_K$ is infinite, any two nonempty opens intersect, so $X$ is irreducible. Let $\mathcal{K}_X$ be the constant sheaf with value $K$ (4.2.50). For $D \in \operatorname{Div}(K)$ define the subsheaf

$$\mathcal{O}(D)(U) \coloneqq \{x \in K \mid v_P(x) \geq -v_P(D)\ \forall P \in U\} \subseteq \mathcal{K}_X(U)$$

generalising the structure sheaf ($\mathcal{O}(0) = \mathcal{O}_X$), with inclusion $\iota_D : \mathcal{O}(D) \hookrightarrow \mathcal{K}_X$.

**Proposition 6.3.61 (Stalk of $\mathcal{O}(D)$)**
*For $P \in X$, the canonical map $\mathcal{O}(D)_P \to \mathcal{K}_{X,P} = K$ is injective, identifying $\mathcal{O}(D)_P$ with the fractional ideal*

$$\{y \in K \mid v_P(y) \geq -v_P(D)\} =: \pi_P^{-v_P(D)}\mathcal{O}_P$$

*of $K$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

**Stalk of $\mathcal{K}_X$.** Since $X$ is irreducible, $\mathcal{K}_X(U) = K$ for every nonempty open $U \subset X$, with every restriction map the identity (4.2.50). We can therefore represent the stalk as $\mathcal{K}_{X,P} = K$ with the identity maps. Under this identification, the canonical map $\mathcal{O}(D)_P \to \mathcal{K}_{X,P} = K$ is given by $(U, x) \mapsto x$.

**Injectivity.** Two representatives $(U, x), (U', x')$ of elements of $\mathcal{O}(D)_P$ become identified in the stalk exactly when they agree on some common refinement; as restriction maps are the identity, this means $x = x'$ already, as elements of $K$. So $(U, x) \mapsto x$ is injective, and we identify $\mathcal{O}(D)_P$ with its image in $K$.

© David Rufino CC BY-NC-ND 4.0

**$\mathcal{O}(D)_P \subseteq \{y : v_P(y) \geq -v_P(D)\}$.** Every element of $\mathcal{O}(D)_P$ is represented by a pair $(U, x)$, $P \in U$, $x \in \mathcal{O}(D)(U)$, whose image in $\mathcal{O}(D)_P$ is $x$ itself (viewed in $K$); taking the defining condition of $\mathcal{O}(D)(U)$ at the point $P \in U$ gives $v_P(x) \geq -v_P(D)$.

**$\{y : v_P(y) \geq -v_P(D)\} \subseteq \mathcal{O}(D)_P$.** Let $y \in K$ with $v_P(y) \geq -v_P(D)$. By (6.2.116), $y$ has finitely many poles, so together with finiteness of $\mathrm{Supp}(D)$, the set $T \coloneqq \{Q \in X \mid v_Q(y) < -v_Q(D)\}$ is finite. As $P \notin T$, $U \coloneqq X \setminus T$ is an open neighbourhood of $P$, and $y \in \mathcal{O}(D)(U)$ by construction of $T$; so $(U, y)$ represents an element of $\mathcal{O}(D)_P$ mapping to $y$.

**Fractional ideal.** By the previous two paragraphs $\mathcal{O}(D)_P = \{y \in K \mid v_P(y) \geq -v_P(D)\}$. Writing $\pi_P$ for a uniformiser of $\mathcal{O}_P$, (3.31.2) represents every $y \in K^\star$ uniquely as $y = u\pi_P^{v_P(y)}$ for $u \in \mathcal{O}_P^\star$, so $v_P(y) \geq -v_P(D) \iff y \in \pi_P^{-v_P(D)}\mathcal{O}_P$; thus $\mathcal{O}(D)_P = \pi_P^{-v_P(D)}\mathcal{O}_P$. This is the image of $\mathcal{O}_P$ under multiplication by the fixed element $\pi_P^{-v_P(D)} \in K$, hence an $\mathcal{O}_P$-submodule of $K$ and so a fractional ideal of $\mathcal{O}_P$ (3.30.1) (indeed $\mathcal{O}_P$ is a discrete valuation ring by (6.2.114), so this is the fractional ideal $\mathfrak{m}_P^{-v_P(D)}$ of (3.31.7)); in particular it is a $k$-vector subspace of $K$, since $k \subset \mathcal{O}_P$ (6.2.113). ∎

**Proposition 6.3.62 (Stalk of the Cokernel)**
*Since $\iota_D$ is a monomorphism (6.3.61), write $\mathcal{K}_X/\mathcal{O}(D) \coloneqq \mathrm{coker}(\iota_D)$ (4.2.36). There is a unique morphism $\theta_P$ making*

$$\begin{array}{ccc} & K & \\ \swarrow & & \searrow \\ (\mathcal{K}_X/\mathcal{O}(D))_P & \xrightarrow[\sim]{\theta_P} & K/\mathcal{O}(D)_P \end{array}$$

*commute, and $\theta_P$ is (moreover) an isomorphism; in particular $(\mathcal{K}_X/\mathcal{O}(D))_P \cong K/\mathcal{O}(D)_P = K/\pi_P^{-v_P(D)}\mathcal{O}_P$.*

*Proof. Drafted by Claude Sonnet, reviewed by David Rufino.*

This is (4.2.37), applied to $\iota_D : \mathcal{O}(D) \to \mathcal{K}_X$ (using $\mathcal{K}_{X,P} = K$), together with (6.3.61) identifying $\mathcal{O}(D)_P$ with $\pi_P^{-v_P(D)}\mathcal{O}_P$. ∎

Used freely below.

**Proposition 6.3.63 (Repartitions compute the Cokernel Sheaf)**
*Writing $p_X : K = \Gamma(X, \mathcal{K}_X) \to \Gamma(X, \mathcal{K}_X/\mathcal{O}(D))$ for the global sections of the cokernel map (4.2.35), there is an exact sequence of $k$-vector spaces*

$$0 \longrightarrow L(D) \longrightarrow K \xrightarrow{p_X} \Gamma(X, \mathcal{K}_X/\mathcal{O}(D)) \xrightarrow{\Phi} \mathcal{A}_K/(\mathcal{A}_K(D) + K) \longrightarrow 0$$

*where $\Phi$ sends $\tau \in \Gamma(X, \mathcal{K}_X/\mathcal{O}(D))$ – a function $\tau : X \to \coprod_{P \in X} K/\mathcal{O}(D)_P$ (4.2.35) – to the class of any repartition $(\alpha_P)_P \in \mathcal{A}_K$ with $\alpha_P \equiv \tau(P) \pmod{\mathcal{O}(D)_P}$ for every $P$; that is, sending a representative $x \in K$ to its corresponding residue vector. In particular $i(D)$ is (via (6.3.63)) the dimension of the cokernel of $p_X$ – via sheaf cohomology, the group $H^1(X, \mathcal{O}(D))$, a cohomological invariant of $X$, not merely of the algebra of repartitions.*

*Sketch.* $(\iota_D)_U$ is the inclusion $\mathcal{O}(D)(U) \hookrightarrow K$, injective.

By (4.2.35) a global section of $\mathcal{K}_X/\mathcal{O}(D)$ is represented by a cover of $X$ together with representatives in $K$ over each open, and since the cofinite topology is quasi-compact (4.1.93) this cover may be taken finite, say $X = U_1 \cup \cdots \cup U_n$ together with representatives $x_i \in K$, well-defined modulo $\mathcal{O}(D)(U_i)$, that **agree at every point of overlap**: for every place $P \in U_i \cap U_j$, the images of $x_i, x_j$ in the stalk $(\mathcal{K}_X/\mathcal{O}(D))_P = K/\pi_P^{-v_P(D)}\mathcal{O}_P$ (6.3.62) coincide, that is

$$v_P(x_i - x_j) \geq -v_P(D)$$

Given such data, set $\alpha_P \coloneqq x_i$ (a literal element of $K$, well-defined only up to the choice of $i$) for any $i$ with $P \in U_i$. The class of $\alpha_P$ modulo $\pi_P^{-v_P(D)}\mathcal{O}_P$ is independent of the choice of $i$: if also $P \in U_j$, then $P \in U_i \cap U_j$, so the overlap condition above applies directly at this $P$ and gives $v_P(x_i - x_j) \geq -v_P(D)$, that is $x_i \equiv x_j \pmod{\pi_P^{-v_P(D)}\mathcal{O}_P}$. Each $x_i$ has finitely many poles (6.2.116), so $\{P \mid \alpha_P \notin \mathcal{O}_P\} \subseteq \bigcup_{i=1}^n \{P \in U_i \mid v_P(x_i) < 0\}$, a finite union of finite sets, is finite; so $\alpha_P \in \mathcal{O}_P$ for all but finitely many $P$, giving $(\alpha_P)_P \in \mathcal{A}_K$.

**Independence of the choice of representing data.** Let $(U_i, x_i)_{i=1}^n$ and $(U_i', x_i')_{i=1}^m$ be two choices representing the same $\tau$, with resulting repartitions $(\alpha_P)_P, (\alpha_P')_P \in \mathcal{A}_K$ as above. For each $P$, choosing $i, i'$ with $P \in U_i \cap U_i'$, both $x_i$ and $x_i'$ represent $\tau(P) \in K/\mathcal{O}(D)_P$ (4.2.35), that is $x_i \equiv x_i' \pmod{\mathcal{O}(D)_P}$, so $\alpha_P - \alpha_P' = x_i - x_i' \in \mathcal{O}(D)_P$.

© David Rufino CC BY-NC-ND 4.0

As this holds for every $P$, $(\alpha_P - \alpha'_P)_P \in \mathcal{A}_K(D)$; so $(\alpha_P)_P$ and $(\alpha'_P)_P$ have the same class in $\mathcal{A}_K/(\mathcal{A}_K(D)+K)$. This assignment therefore descends to a well-defined map

$$\Phi : \Gamma(X, \mathcal{K}_X/\mathcal{O}(D)) \longrightarrow \mathcal{A}_K/(\mathcal{A}_K(D)+K)$$

**Exactness at $K$.** For $c \in K$, $p_X(c)$ is represented by the trivial cover $\{X\}$ with $x_1 := c$, so $p_X(c) = 0$ if and only if $c \in \mathcal{O}(D)(X) = L(D)$; that is, $\ker(p_X) = L(D)$, matching exactness at $K$ (the map $L(D) \hookrightarrow K$ being the inclusion).

**Exactness at $\Gamma(X, \mathcal{K}_X/\mathcal{O}(D))$.** ($\mathrm{Im}(p_X) \subseteq \ker(\Phi)$) For $c \in K$, $p_X(c)$ has $\alpha_P = c$ for every $P$, that is, the constant repartition $\delta(c)$ (diagonal embedding $\delta : K \hookrightarrow \mathcal{A}_K$), whose class in $\mathcal{A}_K/(\mathcal{A}_K(D)+K)$ is 0, since $\delta(c) \in K$; so $\Phi(p_X(c)) = 0$.

($\ker(\Phi) \subseteq \mathrm{Im}(p_X)$) Suppose $\Phi(\tau) = 0$, that is $(\alpha_P)_P \in \mathcal{A}_K(D) + K$, say $(\alpha_P)_P = \beta + \delta(c)$ with $\beta \in \mathcal{A}_K(D)$, $c \in K$. Then $x_i - c \in \mathcal{O}(D)(U_i)$ for every $i$ (as $x_i - c = \alpha_P - c = \beta_P \in \mathcal{O}(D)_P$ for every $P \in U_i$), so $\tau$ and $p_X(c)$ are represented by the same local data, that is $\tau = p_X(c) \in \mathrm{Im}(p_X)$.

**Exactness at $\mathcal{A}_K/(\mathcal{A}_K(D)+K)$.** Given $(\alpha_Q)_Q \in \mathcal{A}_K$, let

$$S := \{Q \mid \alpha_Q \notin \mathcal{O}_Q\} \cup \mathrm{Supp}(D) = \{P_1, ..., P_n\}$$

finite, being a union of two finite sets; note $v_Q(D) = 0$ for every $Q \notin S$, so $\mathcal{O}(D)_Q = \mathcal{O}_Q$ there. Set $x_0 := 0$ and, for $i = 1, ..., n$, $x_i := \alpha_{P_i}$ and $Z_i := \{Q \neq P_i \mid \alpha_{P_i} \notin \mathcal{O}_Q\}$, the poles of $\alpha_{P_i}$ other than possibly at $P_i$ itself, finite by (6.2.116). Since the cofinite topology has no "small" neighbourhoods – any nonempty open already excludes only finitely many points – each $U_i$ ($i \geq 1$) must be taken to also avoid $x_i$'s own poles, not merely to contain $P_i$: set

$$U_0 := X \setminus S, \qquad U_i := X \setminus ((S \cup Z_i) \setminus \{P_i\}) \quad (i = 1, ..., n)$$

*These cover $X$*: $U_0 \cup \cdots \cup U_n \supseteq (X \setminus S) \cup \{P_1, ..., P_n\} = X$, as $P_i \in U_i$ for each $i$.

*Each $U_i$ meets $S$ in at most the single point $P_i$*: $U_0 \cap S = \varnothing$, and $U_i \cap S = \{P_i\}$ for $i \geq 1$, since $U_i$ excludes every other point of $S$. Consequently, for $i \neq j$, no point of $S$ lies in both $U_i$ and $U_j$ (it would have to be the marked point of each, and the marked points – or absence thereof, for $i = 0$ – differ), so $U_i \cap U_j \subseteq X \setminus S$; in particular $Q \in U_i \cap U_j$ gives $v_Q(D) = 0$ and $Q \neq P_i, P_j$.

*They agree at overlaps*: for $i \neq j$ in $\{0, ..., n\}$ and $Q \in U_i \cap U_j$, $v_Q(x_i) \geq 0$ (trivially if $i = 0$, as $x_0 = 0$; else since $Q \in U_i$, $Q \neq P_i$ gives $Q \notin Z_i$) and likewise $v_Q(x_j) \geq 0$; so by the ultrametric inequality $v_Q(x_i - x_j) \geq \min(v_Q(x_i), v_Q(x_j)) \geq 0 = -v_Q(D)$.

This data therefore glues to a class $\tau \in \Gamma(X, \mathcal{K}_X/\mathcal{O}(D))$ with $\tau(Q) = \alpha_Q \pmod{\mathcal{O}(D)_Q}$ for every $Q$ (on $U_0$ this matches $x_0 = 0$, since $Q \notin S$ gives $\alpha_Q \in \mathcal{O}_Q = \mathcal{O}(D)_Q$; at $Q = P_i$ this matches $x_i = \alpha_{P_i}$ directly), so $\Phi(\tau) = (\alpha_Q)_Q$ exactly; so $\Phi$ is surjective, matching exactness at the final term. ∎

© David Rufino CC BY-NC-ND 4.0

# 6.4 Projective Varieties

## 6.4.1 Projective Algebraic Sets

**Proposition 6.4.1 (Projective Space is Functorial)**
*Let $\phi : K \to L$ be an injective ring homomorphism Then there is a well-defined injective map*

$$\begin{aligned} \mathbb{P}^n(K) &\to \mathbb{P}^n(L) \\ (x_0 : ... : x_n) &\to (\phi(x_0) : ... : \phi(x_n)) \end{aligned}$$

*Proof.* Evidently the map is well-defined. Suppose that

$$(\phi(x_0) : ... : \phi(x_n)) \sim (\phi(x_0') : ... : \phi(x_n'))$$

Then for some $i$ we have $x_i \neq 0$ and $\phi(x_i) = \lambda\phi(x_i')$ and $\lambda \in L^\star$. Therefore $\phi(x_i') \neq 0 \implies x_i' \neq 0$ and $\lambda = \phi(x_i/x') =: \phi(\mu)$. Similarly $\phi\big(x_j\big) = \phi(\mu)\phi\big(x_j'\big) \implies \phi\big(x_j - \mu x_j'\big) = 0 \implies x_j = \mu x_j'$. ∎

**Proposition 6.4.2 (Zero Loci in Projective Space)**
*Let $A = k[X_0, ..., X_n]$ be the graded polynomial ring in $(n+1)$-variables over a field $k$. For a homogenous ideal $\mathfrak{a} \subset k[X_0, ..., X_n]$ and field extension $K/k$ define the **zero-locus***

$$V_+(\mathfrak{a})(K) \coloneqq \{(x_0 : ... : x_n) \in \mathbb{P}^n(K) \mid F(x_0, ..., x_n) = 0 \quad \forall F \in \mathfrak{a}\}$$

*Similarly for a subset $Y \subset \mathbb{P}^n(K)$ define*

$$I_+(Y) \coloneqq \{F \in k[X_0, ..., X_n] \mid F_d(x_0, ..., x_n) = 0 \quad \forall d \geq 0, (x_0 : ... : x_n) \in Y\}$$

*Then*

*a) $I_+(Y)$ is a homogenous radical ideal and $\sqrt{\langle S\rangle} \subseteq I_+(V_+(S)(K))$*
*b) $I_+$ and $V_+$ are order-reversing*
*c) $S \subseteq I_+(V_+(S)(K))$ and $Y \subseteq V_+(I_+(Y))(K)$*
*d) $V_+ \circ I_+ \circ V_+ = V_+$ and $I_+ \circ V_+ \circ I_+ = I_+$*
*e) $V_+(S) = V_+\Big(\sqrt{\langle S\rangle}\Big)$ and $V_+(\mathfrak{a}) = V_+(\sqrt{\mathfrak{a}})$*
*f) $\bigcap_i V_+(S_i) = V_+\Big(\bigcup_i S_i\Big)$ and $\bigcap_i V_+(\mathfrak{a}_i) = V_+\Big(\sum_i \mathfrak{a}_i\Big)$*
*g) $\bigcap_i I_+(W_i) = I_+(\bigcup W_i)$*
*h) $V_+(\mathfrak{a}) \cup V_+(\mathfrak{b}) = V_+(\mathfrak{a} \cap \mathfrak{b})$*
*i) $V_+((0)) = \mathbb{P}^n(K)$*

*The sets of the form $V_+(\mathfrak{a})(K)$ determine the **Zariski Topology** on $\mathbb{P}^n(K)$. We denote the topological space by $\mathbb{P}^n_k(K)$. The topological closure satisfies the following identities*

$$V_+(I_+(X)) = \overline{X}$$

*and*

$$I_+(X) = I_+\Big(\overline{X}\Big)$$

*There is a form of the Weak Nullstellensatz*

$$V_+(\mathfrak{a})\Big(\overline{k}\Big) = \varnothing \iff A_+ \not\subset \sqrt{\mathfrak{a}} \iff \mathfrak{a} \text{ irrelevant}$$

*and a form of the Strong Nullstellensatz namely*

$$\mathfrak{a} \text{ essential} \implies I_+\Big(V_+(\mathfrak{a})\Big(\overline{k}\Big)\Big) = \sqrt{\mathfrak{a}}$$

*Further there is a dual isomorphism*

$$\{\mathfrak{a} \lhd A \;\big|\; \mathfrak{a} \text{ radical homogenous and } A_+ \not\subset \mathfrak{a}\} \underset{I_+}{\overset{V_+}{\rightleftarrows}} \Big\{Z \in \mathbb{P}^n_k\Big(\overline{k}\Big) \;\Big|\; Z \neq \varnothing\Big\}$$

*Proof.* Let $V_K(S) \coloneqq \big\{(x) \in K^{n+1} \;\big|\; f(x) = 0 \quad \forall s \in S\big\}$ denote the affine zero-locus. Then $V_+(S)(K) = \pi(V_K(S) \setminus \{\mathbf{0}\})$, where $\mathbf{0} = (0, ..., 0)$ is the origin in $K^{n+1}$.

a) Homogeneity follows from definition. The relation is straightforward.
b) Follows by definition
c) The pair $I, V$ form a Galois connection by (2.1.54) and the relations follow formally by (2.1.50).

© David Rufino CC BY-NC-ND 4.0

d) Follows formally by (2.1.52)
e) Follows from (2.1.53) by considering the closure operators $\sqrt{\langle - \rangle}$ and $\langle - \rangle$.
f) The first equality follows from (2.1.55). The second equality follows from (3.3.29).
g) This follows from (2.1.55).
h) Then using the affine case we may deduce

$$V_+(\mathfrak{a} \cap \mathfrak{b}) = \pi(V(\mathfrak{a}) \cup V(\mathfrak{b}) \setminus \mathbf{0}) = \pi(V(\mathfrak{a}) \setminus \mathbf{0}) \cup \pi(V(\mathfrak{b}) \setminus \mathbf{0}) = V_+(\mathfrak{a}) \cup V_+(\mathfrak{b})$$

i) Trivial

**Weak Nullstellensatz.** Observe that using the affine case $V_{\overline{k}}(A_+) = \{\mathbf{0}\}$ and $I_{\overline{k}}(\mathbf{0}) = A_+$. Therefore

$$V_+(\mathfrak{a})\left(\overline{k}\right) = \varnothing \iff V_{\overline{k}}(\mathfrak{a}) \subset \{\mathbf{0}\} \overset{(3.13.7)}{\iff} A_+ \not\subset \sqrt{\mathfrak{a}}$$

**Strong Nullstellensatz.** Assume $\mathfrak{a}$ is an essential homogenous ideal then $Y \coloneqq V(\mathfrak{a}) \setminus \{\mathbf{0}\}$ is non-empty and $\overline{Y} = V(\mathfrak{a})$. Because $\mathfrak{a}$ is homogenous then $Y$ is a cone – $x \in Y \implies \lambda x \in Y$ for $\lambda \in \overline{k}^\star$. We claim that $V_+(\mathfrak{a}) = \pi(Y)$ and $I(Y) = I_+(\pi(Y))$. Suppose $F \in I(Y)$ then $F(\lambda x) = 0$ for all $\lambda \in \overline{k}^\star$ and $x \in Y$. As $\overline{k}$ is infinite we deduce that $F_d \in I(Y)$ whence $F \in I_+(\pi(Y))$. The remaining claims are clear. Therefore we may deduce from (6.1.2) and (6.1.13)

$$I_+V_+(\mathfrak{a}) = I_+(\pi(Y)) = I(Y) = I\left(\overline{Y}\right) = I(V(\mathfrak{a})) = \sqrt{\mathfrak{a}}$$

**Topological Closure.** By (2.1.52) $V_+ \circ I_+$ is a closure operator with image precisely the closed sets. Therefore by (2.1.41) $V_+ \circ I_+$ is the topological closure (4.1.19). Finally $I_+\left(\overline{X}\right) = I_+(V_+(I_+(X))) = I_+(X)$ using d).

The dual isomorphism follows from the Strong Nullstellensatz. ∎

**Definition 6.4.3 (Projective Variety)**
*For an essential homogenous radical ideal $\mathfrak{a} \lhd k[X_0, ..., X_n]$ we say the family*

$$X(K) \coloneqq V_+(\mathfrak{a})(K) \subset \mathbb{P}^n_k(K)$$

*is a **projective variety** and write it as $X(K) \coloneqq V_+(\mathfrak{a})(K)$. We say $X$ is **integral** if $X\left(\overline{k}\right)$ is irreducible in the subspace topology. We also define the **homogenous coordinate ring***

$$k[X]_{\text{hom}} \coloneqq k[X_0, ..., X_n]/\mathfrak{a}$$

*which is naturally a positive graded ring by (3.12.12). Define the **irrelevant ideal***

$$k[X]_+ \coloneqq \bigoplus_{d>0} k[X]_d$$

We may generalise (6.4.2) as follows.

**Proposition 6.4.4 (Zero Loci in Projective Varieties)**
*Let $X = V_+(\mathfrak{a}) \subset \mathbb{P}^n_k$ be a projective variety and $A = k[X]_{\text{hom}}$ the coordinate ring. For any homogenous ideal $\mathfrak{b} \lhd A$ and field extension $K/k$ define the **zero-locus** by*

$$V_+(\mathfrak{b})(K) \coloneqq \{(x_0 : ... : x_n) \in X(K) \mid f(x_0, ..., x_n) = 0 \quad \forall f \in \mathfrak{b}\}$$

*Similarly for a subset $Y \subset X(K)$ define*

$$I_+(Y) \coloneqq \{f \in k[X] \mid f_d(x) = 0 \quad \forall d \geq 0, x \in Y\}$$

*Suppose $\pi : k[X_0, ..., X_n] \to k[X]$ is the quotient map. Then we have the following properties*

$$V_+(\mathfrak{b})(K) = V_+(\pi^{-1}(\mathfrak{b}))(K) \cap X(K)$$
$$I_+(Y \cap X(K)) = \pi(I_+(Y)) \quad Y \subset \mathbb{P}^n_k(K)$$
$$I_+(V_+(\mathfrak{b})(K)) = \pi(I_+(V_+(\pi^{-1}(\mathfrak{b}))))$$

*The pair $(V_+(-)(K), I_+)$ satisfies the same properties as (6.4.2). The image of $V_+(-)(K)$ forms the closed sets of a topology on $X(K)$ which coincides with the subspace topology from $\mathbb{P}^n_k(K)$. Further the Weak and Strong Nullstellensatz also hold*

$$V_+(\mathfrak{b})\left(\overline{k}\right) = \varnothing \iff k[X]_+ \not\subset \sqrt{\mathfrak{b}} \iff \mathfrak{b} \text{ is irrelevant}$$

*and*

© David Rufino CC BY-NC-ND 4.0

$$\mathfrak{b} \text{ essential} \implies I_+\left(V_+(\mathfrak{b})\left(\overline{k}\right)\right) = \sqrt{\mathfrak{b}}$$

*Further there is a dual isomorphism*

$$\left\{\mathfrak{b} \lhd k[X]_{\text{hom}} \;\middle|\; \mathfrak{b} \text{ radical, homogenous and } A_+ \not\subset \mathfrak{b}\right\} \underset{I_+}{\overset{V_+}{\rightleftharpoons}} \left\{Z \in X\left(\overline{k}\right) \;\middle|\; Z \neq \varnothing\right\}$$

**Proposition 6.4.5 (Projective Subvariety)**
*Let $X$ be a projective variety, $\mathfrak{b} \lhd k[X]_{\text{hom}}$ an essential radical homogenous ideal. Then $V_+(\mathfrak{b})(-)$ is a **closed subvariety** which coincides with the projective variety $V_+(\pi^{-1}(\mathfrak{b}))$. There is a bijective correspondence between non-empty closed subsets of $X\left(\overline{k}\right)$ and closed subvarieties of $X$.*

*Conversely if $X', X$ are two projective varieties such that $X'\left(\overline{k}\right) \subset X\left(\overline{k}\right)$ then $X'$ may be expressed as a closed subvariety.*

*Proof.* There is a canonical graded surjective $k$-algebra homomorphism $\pi : k[X_0, ..., X_n] \to k[X]_{\text{hom}}$ by (3.12.12). Then $\pi^{-1}(\mathfrak{b})$ is graded (3.12.10) and radical (3.3.50) g). Also by definition of the quotient grading on $\pi$ we find $\pi^{-1}(k[X]_+) = k[X_0, ..., X_n]_+$ so that $\mathfrak{b}$ essential $\implies \pi^{-1}(\mathfrak{b})$ is essential (3.13.7). The bijective correspondence exists by (6.4.4).

If $X' = V_+(\mathfrak{b}')$, $X = V_+(\mathfrak{b})$ are two projective varieties such that $X'\left(\overline{k}\right) \subset X\left(\overline{k}\right)$ then by the dual isomorphism we have $\mathfrak{b} \subset \mathfrak{b}'$. Therefore $X'$ may be represented as the closed subvariety $V_+(\pi(\mathfrak{b}'))$ ∎

**Proposition 6.4.6 (Principal Open Sets)**
*Let $X \subset \mathbb{P}^n_k$ be a projective variety. The Zariski topology on $X(K)$ has a base consisting of subsets of the form*

$$D_+(f)(K) \coloneqq X(K) \setminus V_+((f)) = \{(x) \in X(K) \mid f_d(x) \neq 0 \text{ some } d > 0\}$$

*where $f \in k[X]$ is a homogenous form. Note that $(x) = (y) \implies (f_d(x) = 0 \iff f_d(y) = 0)$ so the second form is well-defined.*

*Proof.* A generic open set $U = X(K) \setminus V_+(\mathfrak{b})(K)$ where $\mathfrak{b}$ is a homogenous ideal of $k[X]$. Given $(x) \in U$ there is some $f \in \mathfrak{b}$ and $d > 0$ such that $f_d(x) \neq 0$. Therefore $D_+(f_d) \subset U$ is an open neighbourhood of $(x)$. ∎

**Proposition 6.4.7**
*Let $X = V_+(\mathfrak{b})$ be a projective variety and $f, g \in k[X]$ essential homogenous forms. Then*

$$D_+(g) \subset D_+(f) \iff f \mid g^N \text{ some } N > 0$$

*Proof.*

$$D_+(g) \subset D_+(f) \iff V_+((f)) \subset V_+((g)) \iff \sqrt{(g)} \subset \sqrt{(f)} \iff f \mid g^N$$

∎

**Proposition 6.4.8 (Irreducible Subsets)**
*Let $X = V_+(\mathfrak{a}) \subset \mathbb{P}^n_k$ be a projective variety and $\mathfrak{b} \lhd k[X]$ a homogenous radical ideal. Then a closed subvariety $V_+(\mathfrak{b}) \subset X$ is integral iff $\mathfrak{b}$ is prime.*

*In particular $\mathbb{P}^n_k\left(\overline{k}\right)$ is integral.*

*Proof.* Let $Y \coloneqq V_+(\mathfrak{b})$. Suppose that $Y$ is not irreducible then $Y \subseteq V_+(\mathfrak{c}) \cup V_+(\mathfrak{d})$ is a non-trivial decomposition into closed subsets for $\mathfrak{c}, \mathfrak{d}$ essential homogenous radical ideals for which $Y \not\subset V_+(\mathfrak{c}) \implies \mathfrak{c} \not\subset \mathfrak{b}$ and similarly $\mathfrak{d} \not\subset \mathfrak{b}$ (see (4.1.61)). Choose $f \in \mathfrak{c} \setminus \mathfrak{b}$ and $g \in \mathfrak{d} \setminus \mathfrak{b}$. Then $fg$ is zero on $Y$ whence $fg \in \mathfrak{b}$. This shows that $\mathfrak{b}$ is not prime.

Conversely suppose $Y \subset V_+(\mathfrak{c}) \cup V_+(\mathfrak{d}) = V_+(\mathfrak{c} \cap \mathfrak{d})$ then by (6.4.2) $\mathfrak{c}\mathfrak{d} \subset \mathfrak{c} \cap \mathfrak{d} \subset \mathfrak{b}$. By (3.3.39) we have $\mathfrak{c} \subset \mathfrak{b}$ or $\mathfrak{d} \subset \mathfrak{b}$, whence $Y \subset V_+(\mathfrak{c})$ or $Y \subset V_+(\mathfrak{d})$. Therefore we conclude that $Y$ is irreducible by (4.1.61).

Therefore $\mathbb{P}^n_k\left(\overline{k}\right)$ is irreducible by the special case both $\mathfrak{a}$ and $\mathfrak{b}$ being the zero ideals. ∎

**Proposition 6.4.9 (Dual Isomorphism between Closed Sets and Homogenous Ideals)**
*Let $X = V_+(\mathfrak{a}) \subset \mathbb{P}^n_k$ be a projective variety and $A = k[X]$ the coordinate ring. Under the dual isomorphism*

$$\left\{\mathfrak{b} \lhd A \;\middle|\; \mathfrak{b} \text{ radical homogenous and } A_+ \not\subset \mathfrak{b}\right\} \underset{I_+}{\overset{V_+}{\rightleftharpoons}} \left\{Z \subset X\left(\overline{k}\right) \text{ closed} \;\middle|\; Z \neq \varnothing\right\}$$

*we have the following correspondence*

- *irreducible sets correspond to prime ideals*

© David Rufino CC BY-NC-ND 4.0

- *irreducible components (of $V_+(\mathfrak{b})$) correspond to minimal prime ideals (of $\mathfrak{b}$)*

**Proposition 6.4.10 (Decomposition into Irreducible Components)**
*Let $X = V_+(\mathfrak{a})$ be a projective variety then the topological space $X\left(\overline{k}\right)$ is Noetherian. Furthermore every closed subset $Z$ has finitely many irreducible components $Z_i$ and the decomposition*

$$Z = Z_1 \cup ... \cup Z_n$$

*is the unique incomparable decomposition into irreducible closed subsets.*

*If $Z = V_+(\mathfrak{b})$ then the irreducible components correspond to precisely the minimal prime ideals over $\mathfrak{b}$.*

*Proof.* $X\left(\overline{k}\right)$ is Noetherian by using the order isomorphism in (6.4.9) and the fact $k[X]$ is Noetherian. The result then follows from (4.1.76). ∎

## 6.4.2 Affine Charts

References:

- Algebraic Curves [Ful08] (Chap. 4.3)
- Algebraic Geometry [Har13] (Chap. I §2)

For every $i = 0, ..., n$ the subset of $\mathbb{P}^n_k(K)$ consisting of points of the form

$$\{(x_0 : ... : x_n) \mid x_i \neq 0\}$$

may be naturally identified with $\mathbb{A}^n_k(K)$. The purpose of this section is to make this notion precise.

**Lemma 6.4.11**
*There is an isomorphism of rings*

$$\theta_i^\sharp : k[X_0, ..., X_n]_{(X_i)} \longleftrightarrow k[x_0, ..., x_{i-1}, x_{i+1}, ..., x_n]$$
$$\frac{X_j}{X_i} \longleftrightarrow x_j$$

*Proof.* The algebra homomorphism $\theta_i^\sharp$ exists by (3.8.5) and (3.6.5), restricting to the subring $k[X_0, ..., X_n]_{(X_i)}$. The inverse algebra homomorphism exists by (3.8.5), sending $x_j \to \frac{X_j}{X_i}$. We may verify directly that these are mutually inverse. ∎

**Lemma 6.4.12 ((De-)Homogenisation)**
*Fix $n$ and $0 \leq i \leq n$. For $G \in k[X_0, ..., X_n]$ homogenous and $F \in k[x_0, ..., x_{i-1}, x_{i+1}, ..., x_n]$ define*

$$G^a := G(x_0, ..., x_{i-1}, 1, x_{i+1}, ..., x_n)$$
$$F^h := X_i^{\deg(F)} F\left(\frac{X_0}{X_i}, ..., \frac{X_{i-1}}{X_i}, \frac{X_{i+1}}{X_i}, ..., \frac{X_n}{X_i}\right)$$
$$= \sum_{j=0}^{\deg F} X_i^{\deg(F)-j} F_j(X_0, ..., X_{i-1}, X_{i+1}, ..., X_n)$$

*Then*

*a) $(FF')^h = F^h F'^h$ and $(GG')^a = G^a G'^a$*
*b) $F^{ha} = F$ and $X_i^k G^{ah} = G$ where $k := v_{X_i}(G)$ is the smallest power of $X_i$ appearing in $G$.*
*c) For $\mathfrak{b} \lhd k[X_0, ..., X_n]$ a homogenous ideal the set $\mathfrak{b}^a := \{G^a \mid G \in \mathfrak{b}\}$ is an ideal.*

*For $\mathfrak{a} \lhd k[x_0, ..., x_{i-1}, x_{i+1}, ..., x_n]$ define the homogenous ideal*

$$\mathfrak{a}^h := \left\langle \{F^h \mid F \in \mathfrak{a}\} \right\rangle$$

*Then we also have the identities*

*d) $\mathfrak{b} \subset \mathfrak{b}^{ah}$ with equality when $\mathfrak{b}$ is radical and $\mathfrak{b} \subset \mathfrak{p}$ minimal $\implies X_i \notin \mathfrak{p}$*
*e) $\mathfrak{a} = \mathfrak{a}^{ha}$*
*f) $\mathfrak{b}$ prime and $X_i \notin \mathfrak{b} \implies \mathfrak{b}^a$ prime.*
*g) $\mathfrak{a}$ prime $\implies \mathfrak{a}^h$ prime*
*h) $\mathfrak{a}$ radical $\implies \mathfrak{a}^h$ radical*

© David Rufino CC BY-NC-ND 4.0

*Proof.* Mostly tedious verification

a) Using the notation of (6.4.11) we see that $F^h = X_i^{\deg(F)}\left(\theta_i^\sharp\right)^{-1}(F)$ so evidently $(FF')^h = F^h F'^h$ because $\deg(FF') = \deg(F) + \deg(F')$.

   The second relation follows immediately from (6.4.11).

b) The first identity is clear. We may write

$$G = \sum_{j=0}^{\deg(G)-v_{X_i}(G)} X_i^{\deg(G)-j} G_j(X_0, ..., X_{i-1}, X_{i+1}, ..., X_n)$$

   where $\deg\left(G_j\right) = j$ and $v_{X_i}(G)$ is the smallest power of $X_i$ appearing in $G$. So

$$G^a = \sum_{j=0}^{\deg(G)-v_{X_i}(G)} G_j(x_0, ..., x_{i-1}, x_{i+1}, ..., x_n)$$

$$G^{ah} = \sum_{j=0}^{\deg(G)-v_{X_i}(G)} X_i^{\deg(G)-v_{X_i}(G)-j} G_j(X_0, ..., X_{i-1}, X_{i+1}, ..., X_n)$$

   whence $G = X_i^{v_{X_i}(G)} G^{ah}$.

c) This is evident because $(-)^a$ is a ring homomorphism.

d) The first part is evident from b). Suppose $G \in \mathfrak{b}^{ah}$ then

$$G = \sum_{j=1}^{r} \lambda_j G_j^{ah} \implies X_i^k G = \sum_{j=1}^{r} \lambda_j X_i^{k_j} G_j$$

   for $G_j \in \mathfrak{b}$. Then $X_i^k G \in \mathfrak{b} \implies G \in \mathfrak{b}$ as we assumed $X_i \notin \mathfrak{b}$.

e) Trivially $F \in \mathfrak{a} \implies F^h \in \mathfrak{a}^h \implies F^{ha} = F \in \mathfrak{a}^{ha}$. Conversely any $G \in \mathfrak{a}^h$ is of the form

$$G = \sum_j \lambda_j F_j^h \implies G^a = \sum_j \lambda_j^a F_j^{ha} = \sum_j \lambda_j^a F_j \in \mathfrak{a}^{ha}$$

   for some $\lambda_i \in k[X_0, ..., X_n]$.

f) Suppose $FF' \in \mathfrak{b}^a \implies F^h F'^h \in \mathfrak{b}^{ah} = \mathfrak{b} \implies F^h \in \mathfrak{b}$ or $F'^h \in \mathfrak{b} \implies F \in \mathfrak{b}^a$ or $F' \in \mathfrak{b}^a$.

g) Suppose $GG' \in \mathfrak{a}^h$ then $GG' = \sum_j \lambda_j F_j^h$ for $F_j \in \mathfrak{a}$. Then $G^a G'^a = \sum_j \lambda_j^a F_j^{ha} = \sum_j \lambda_j^a F_j \in \mathfrak{a}$. Therefore say $G^a \in \mathfrak{a} \implies G^{ah} \in \mathfrak{a}^h \implies G = X_i^k G^{ah} \in \mathfrak{a}^h$.

h) Suppose $G^n \in \mathfrak{a}^h$, then $(G^a)^n \in \mathfrak{a}^{ha} = \mathfrak{a}$ whence $G^a \in \mathfrak{a}$ and $G^{ah} \in \mathfrak{a}^h \implies G \in \mathfrak{a}^h$

■

**Proposition 6.4.13 (Projective Space has Affine Covering)**
*Consider subsets $D_+(X_i)(K) = U_i(K) \subset \mathbb{P}_k^n(K)$ of the form*

$$D_+(X_i)(K) = U_i(K) \coloneqq \{(x_0 : ... : x_n) \mid x_i \neq 0\} \quad i = 0, ..., n$$

*These are open in the Zariski topology. Furthermore under the subspace topology $U_i$ is homeomorphic to $\mathbb{A}_k^n(K)$, with an explicit homeomorphism given by*

$$\theta_i : \mathbb{A}_k^n(K) \to U_i$$
$$(x_0, ..., x_{i-1}, x_{i+1}, ..., x_n) \to [x_0 : ... : 1 : ... : x_n]$$

*For $F \in k[x_0, ..., x_{i-1}, x_{i+1}, ..., x_n]$ and $(x) \in U_i$ we have*

$$F(x) = 0 \iff F^h(\theta_i(x)) = 0$$

*and similarly for $G \in k[X_0, ..., X_n]$ we have*

$$G(\theta_i(x)) = 0 \iff G^a(x) = 0$$

*Further we have*

*a) $\theta_i\left(V^{\text{aff}}(\mathfrak{a})\right) = V_+\left(\mathfrak{a}^h\right) \cap U_i$ for $\mathfrak{a} \lhd k[x_0, ..., x_{i-1}, x_{i+1}, ..., x_n]$*
*b) $\theta_i\left(V^{\text{aff}}(\mathfrak{b}^a)\right) = V_+(\mathfrak{b}) \cap U_i = V_+\left(\mathfrak{b}^{ah}\right) \cap U_i$ for $\mathfrak{b} \lhd k[X_0, ..., X_n]$ homogenous*
*c) $I^{\text{aff}}(Y)^h = I_+(\theta_i(Y))$ for $Y \subset U_i$*

© David Rufino CC BY-NC-ND 4.0

*Proof.* This follows largely from (6.4.12). The relations show that $\theta_i(D(F)) = D_+\left(F^h\right)$ and $\theta_i^{-1}(D_+(G)) = D(G^a)$, so that $\theta_i$ is a homeomorphism as required.

a) $x \in \theta_i\left(V^{\text{aff}}(\mathfrak{a})\right) \implies x \in U_i$ and $F(\theta_i^{-1}(x)) = 0 \quad \forall F \in \mathfrak{a} \implies x \in U_i$ and $F^h(x) = 0 \quad \forall F \in \mathfrak{a} \implies G(x) = 0 \quad \forall G \in \mathfrak{a}^h \implies x \in V_+\left(\mathfrak{a}^h\right) \cap U_i$. Conversely $x \in V_+\left(\mathfrak{a}^h\right) \cap U_i \implies G(x) = 0 \quad \forall G \in \mathfrak{a}^h \implies F(\theta_i^{-1}(x)) = 0 \quad \forall F \in \mathfrak{a}^{ha} = \mathfrak{a} \implies x \in \theta_i\left(V^{\text{aff}}(\mathfrak{a})\right)$

b) $\theta_i\left(V^{\text{aff}}(\mathfrak{b}^a)\right) = V_+\left(\mathfrak{b}^{ah}\right) \cap U_i$. It remains to show that $V_+(\mathfrak{b}) \cap U_i = V_+\left(\mathfrak{b}^{ah}\right) \cap U_i$. As (6.4.12) $\mathfrak{b} \subset \mathfrak{b}^{ah}$ then $V_+\left(\mathfrak{b}^{ah}\right) \cap U_i \subset V_+(\mathfrak{b}) \cap U_i$. Conversely if $x \in V_+(\mathfrak{b}) \cap U_i$ then $G(x) = 0 \implies x_i^k G^{ah}(x) = 0 \implies G^{ah}(x) = 0$ because $x_i \neq 0$. This shows $x \in V_+\left(\mathfrak{b}^{ah}\right) \cap U_i$

c) Suppose $G \in I^{\text{aff}}(Y)^h$ then $G = \sum_j \lambda_j F_j^h$ for $F_j \in I^{\text{aff}}(Y)$ and $\lambda_j \in k[X_0, ..., X_n]$. Then $F_j(x) = 0 \implies F_j^h(\theta_i(x)) = 0 \implies G(\theta_i(x)) = 0$. Therefore we conclude $G \in I_+(\theta_i(Y))$. Conversely if $G \in I_+(\theta_i(Y))$ then $G^a(x) = 0$ for all $x \in Y$ and $G^a \in I^{\text{aff}}(Y) \implies G^{ah} \in I^{\text{aff}}(Y)^h \implies G = X_i^k G^{ah} \in I^{\text{aff}}(Y)^h$.

■

**Lemma 6.4.14**
*Let $X = V_+(\mathfrak{b})$ be a projective variety. Then $X \cap U_i \neq \varnothing \iff X_i \notin \mathfrak{b} \iff \overline{X_i} \neq 0$.*

*Proof.* Observe $X \cap U_i = \varnothing \iff X \subset V_+(X_i) \iff X_i \in I_+(X) = \mathfrak{b}$. ■

**Corollary 6.4.15**
*Let $X = V_+(\mathfrak{b}) \subset \mathbb{P}_k^n$ be a projective variety and for each affine chart $U_i \subset \mathbb{P}_k^n$ consider the affine variety $X^a := V^{\text{aff}}(\mathfrak{b}^a) \subset \mathbb{A}_k^n$. Then there is a natural homeomorphism*

$$\theta_i : X^a(K) \xrightarrow{\sim} D_+\left(\overline{X_i}\right) = X(K) \cap U_i(K)$$

*for all algebraic extensions $K/k$.*

**Proposition 6.4.16 (Projective Closure)**
*Identify $\mathbb{A}_k^n\left(\overline{k}\right)$ with $U_i \subset \mathbb{P}_k^n\left(\overline{k}\right)$. Let $Y \subset U_i$ be a subset. Then we have the following expression for topological closure*

$$\overline{Y} = V_+\left(I^{\text{aff}}(Y)^h\right)$$
$$\overline{Y} \cap U_i = \text{cl}_{U_i}(Y)$$

*We call $\overline{Y}$ the **projective closure** of $Y$.*

*Proof.* By (6.4.13).c) $I_+(Y) = I^{\text{aff}}(Y)^h \implies \overline{Y} \overset{(6.4.4)}{=} V_+ I_+(Y) = V_+\left(I^{\text{aff}}(Y)^h\right)$.

Then $\text{cl}_{U_i}(Y) = \overline{Y} \cap U_i$ by (4.1.21) (or more directly $\overset{(6.1.2)}{=} V_+(I_+(Y)) \overset{(6.4.13).a)}{=} V_+\left(I^{\text{aff}}(Y)^h\right) \cap U_i = \overline{Y} \cap U_i$). ■

## 6.4.3 Galois Orbits

**Proposition 6.4.17**
*Let $X = V_+(\mathfrak{a}) \subset \mathbb{P}_k^n$ be a projective variety and $K/k$ a normal algebraic extension (e.g. $\overline{k}$). Then there is a well-defined group action*

$$\text{Aut}(K/k) \times X(K) \to X(K)$$
$$(\sigma, (x_0 : ... : x_n)) \to (\sigma(x_0) : ... : \sigma(x_n))$$

*which commutes with the affine charts*

$$\sigma(\theta_i(x)) = \theta_i(\sigma(x)) \quad \forall x \in U_i(K) \cap X(K)$$

*Write $X_0(K)$ for the set of equivalence classes under this action. The projection map*

$$\pi : X(K) \to X_0(K)$$

*is the Kolmogorov Quotient (4.1.36). In particular points $x, y \in X(K)$ are topologically indistinguishable iff they are Galois conjugate. Further closed (resp. open) subsets are Galois-equivariant.*

*Proof.* First consider the case $X = \mathbb{P}_k^n$. If $(x_0 : ... : x_n) = (x_0' : ... : x_n')$ then $x_i = \lambda_i x_i'$ for $\lambda_i \in K^\star$. Then $\sigma(x_i) = \sigma(\lambda_i)\sigma(x_i')$ so $(\sigma(x_0) : ... : \sigma(x_n)) \sim (\sigma(x_0') : ... : \sigma(x_n'))$ and the action is well-defined. For $x \in X(K)$ write $\sigma(x)$. Then $\tau(\sigma(x)) = (\tau \circ \sigma)(x)$ so it is a group action.

For the general case if $F(x) = 0$ for $F \in k[X_0, ..., X_n]$ then $F(\sigma(x)) = 0$. Therefore $x \in V(\mathfrak{a})(K) \implies \sigma(x) \in V(\mathfrak{a})(K)$ and the action is well-defined on $X(K)$.

© David Rufino CC BY-NC-ND 4.0

We need to show that two points $x, y \in X(K)$ are conjugate by $\mathrm{Aut}(K/k)$ iff they are topologically indistinguishable. Suppose $x = \sigma(y)$ and $x \in V(\mathfrak{b})$. Then evidently $y \in V(\mathfrak{b})$. Conversely if they are topologically indistinguishable then we may suppose $x, y \in U_i$. Then $\theta_i(x), \theta_i(y)$ are also indistinguishable, so by (6.1.9) they are Galois conjugate, and so $\theta_i(x) = \theta_i(\sigma(y))$. As $\theta_i$ is bijective we deduce $x = \sigma(y)$ as required. ∎

**Proposition 6.4.18**
*Let $X = V_+(\mathfrak{a}) \subset \mathbb{P}^n_k$ be a projective variety and $L/K/k$ a tower of normal extensions. Then there is an injective map*

$$\begin{array}{ccc} X(K) & \xhookrightarrow{i_{KL}} & X(L) \\ \downarrow{\scriptstyle \pi} & & \downarrow{\scriptstyle \pi} \\ X_0(K) & \overset{i_0}{\dashrightarrow} & X_0(L) \end{array}$$

*Proof.* In order for $i_0$ to be well-defined, we require that $\pi(x) = \pi(y) \implies \pi(i_{KL}(x)) = \pi(i_{KL}(y))$. Recall $\pi(x) = \pi(y) \implies x = \sigma(y)$ for some $\sigma \in \mathrm{Aut}(K/k)$. Then $i_{KL} \circ \sigma \in \mathrm{Mor}_k(K, L)$ and by (3.19.69) there exists $\hat{\sigma} \in \mathrm{Aut}(L/k)$ such that $\hat{\sigma}|_K := \hat{\sigma} \circ i_{KL} = i_{KL} \circ \sigma$. Therefore $\hat{\sigma}(i_{KL}(x)) = i_{KL}(y) \implies \pi(i_{KL}(x)) = \pi(i_{KL}(y))$.

To show $i_0$ is injective we need to show the converse, which follows from (3.19.71). ∎

## 6.4.4 Projective Varieties are Abstract Varieties

**Definition 6.4.19 (Regular Function)**
*Let $X = V_+(\mathfrak{a}) \subset \mathbb{P}^n_k$ be a projective variety and $U \subset X\left(\overline{k}\right)$ an open subset. We say a function*

$$f : U \to \overline{k}$$

*is **regular** at $x \in U$ there is a neighbourhood $V$ of $x$ such that*

$$f(y) = \frac{g(y)}{h(y)} \quad \forall\, y \in V$$

*for $g, h \in k[X]$ homogenous polynomials of the same degree. We therefore define the **structure sheaf** as follows*

$$\mathcal{O}_X(U) := \left\{ f : U \to \overline{k} \;\middle|\; f \text{ regular at all } x \in U \right\}$$

*The pair $(X, \mathcal{O}_X)$ is a space of functions.*

**Proposition 6.4.20 (Structure Sheaf of Regular Functions)**
*Let $X = V_+(\mathfrak{a}) \subset \mathbb{P}^n_k$ be a projective variety, $U \subset X\left(\overline{k}\right)$ open and $f : U \to \overline{k}$ a function. Consider the affine charts*

$$\theta_i : X_i^a\left(\overline{k}\right) \xrightarrow{\sim} D_+\left(\overline{X_i}\right) = X\left(\overline{k}\right) \cap U_i \quad i = 0, ..., n$$

*where $X_i^a = V^{\mathrm{aff}}(\mathfrak{a}^a)$ as in (6.4.15). Then the following are equivalent*

*a) $f$ is regular (at $x$) in the sense of (6.4.19)*
*b) $f \circ \theta_i : \theta_i^{-1}(U) \to \overline{k}$ is regular (at $\theta_i^{-1}(x)$) in the sense of (6.1.25) whenever $U \cap U_i \neq \varnothing$ ($x \in U \cap U_i$)*

*In other words for each $i = 1, ..., n$ there is an isomorphism of spaces with functions*

$$\theta_i : \left(X_i^a\left(\overline{k}\right), \mathcal{O}_{X_i^a}\right) \xrightarrow{\sim} \left(D_+\left(\overline{X_i}\right), \mathcal{O}_X\,|_{D_+\left(\overline{X_i}\right)}\right)$$

*In particular $(X, \mathcal{O}_X)$ is an abstract variety.*

*Proof.* *a)* $\implies$ *b)* Let $x := \theta(\hat{x})$. There exists an open nbhd $V$ of $x$ such that $f(y) = \frac{g(y)}{h(y)}$ for all $y \in V$. By definition $g = G + \mathfrak{a}$ and $h = H + \mathfrak{a}$ for $G, H \in k[X_0, ..., X_n]$ homogenous polynomials of the same degree, for which $H$ doesn't vanish on $V$. Then $\hat{V} := \theta_i^{-1}(V)$ is a open nbhd of $\hat{x}$. For all $\hat{y} \in \hat{V}$ we have

$$(f \circ \theta_i)(\hat{y}) = f(\theta_i(\hat{y})) = \frac{G(\hat{y})}{H(\hat{y})} = \frac{G^a(y)}{H^a(y)}$$

*b)* $\implies$ *a)* Suppose $x \in U \cap U_i$. We have by definition an open nbhd $\hat{V}$ of $\theta_i^{-1}(x)$ and $G, H \in k[x_0, x_1, ..., x_{i-1}, x_{i+1}, ..., x_n]$ for which

$$(f \circ \theta_i)(\hat{y}) = \frac{G(\hat{y})}{H(\hat{y})} \quad \forall \hat{y} \in \hat{V}$$

© David Rufino CC BY-NC-ND 4.0

As $\theta_i$ is a homeomorphism, $V := \theta_i\left(\hat{V}_i\right)$ is an open nbhd of $x$. By (6.4.13) we have for $y := \theta_i(\hat{y})$

$$f(y) = \frac{G^h(y)}{H^h(y)}$$

which holds for all $y \in V$. ∎

We may generalise the dehomogenisation

**Proposition 6.4.21 (Structure Sheaf on Basic Affines)**
*Let $X = V_+(\mathfrak{p})$ be an irreducible projective variety and suppose $X \cap U_i \neq \varnothing$. Then there is an isomorphism of $k$-algebras*

$$\begin{array}{ccc} \mathcal{O}_X\left(D_+\left(\overline{X_i}\right)\right) & \xrightarrow[\sim]{(6.4.20)} & \mathcal{O}_{X\cap U_i}(X \cap U_i) \\ \vdots & & \sim \uparrow (6.1.26) \\ k[X]_{(\overline{X_i})} & \xrightarrow{\sim} & k[X \cap U_i] \end{array}$$

$$\frac{\overline{X_0}}{\overline{X_i}} \longleftrightarrow \overline{x_i}$$

*and the dashed arrow is defined in the obvious way (and is therefore an isomorphism).*

*Proof.* By (6.4.14) $X_i \notin \mathfrak{p}$. We first demonstrate the bottom map is well-defined and an isomorphism. Consider the following commutative diagram

$$\begin{array}{ccccc} & & k[X_0,...,X_n]_{(X_i)} & \xrightarrow[\sim]{\theta_i^\sharp} & k[U_i] \\ & \swarrow & \downarrow \pi_1 & & \downarrow \pi_2 \\ k[X_0,...,X_n]_{(X_i)}/\mathfrak{p}_{(X_i)} & \xrightarrow[(3.12.17)]{\sim} & k[X]_{(\overline{X_i})} & \xrightarrow{\sim} & k[X \cap U_i] \end{array}$$

where the isomorphism $\theta_i^\sharp$ is defined in (6.4.11).

It follows from (3.3.56) that $\pi_1$ is uniquely defined and has kernel $\mathfrak{p}_{(X_i)}$ which equals $\mathfrak{p}_{X_i} \cap k[X_0,...,X_n]_{(X_i)}$ by (3.12.15). Therefore we may regard $\pi_1$ as a quotient map.

We also claim that the diagonal map $\pi_2 \circ \theta_i^\sharp$ has kernel $\mathfrak{p}_{(X_i)}$. It is enough by (3.6.5).b) and the preceding comment to show that the composite map $\pi_2 \circ (-)^a : k[X_0,...,X_n] \to k[X \cap U_i]$ has kernel $\mathfrak{p}$. However this follows from (6.4.12) for $F \in \ker(\pi_2 \circ (-)^a) \implies F^a \in \mathfrak{p}^a \implies F^a = G^a \implies F^{ah} = G^{ah} \in \mathfrak{p}^{ah} = \mathfrak{p} \implies F^{ah} \in \mathfrak{p} \implies X_i^k F \in \mathfrak{p} \implies F \in \mathfrak{p}$.

The required map exists by (3.3.56), as $\pi_1$ is a quotient map. Further it is injective, surjective and therefore an isomorphism

The map $k[X]_{(\overline{X_i})} \to \mathcal{O}_X\left(D_+\left(\overline{X_i}\right)\right)$ is defined in the obvious way. It is clear the diagram commutes and so this map is an isomorphism. ∎

**Proposition 6.4.22**
*Let $X$ be a projective variety. Then it is separated.*

## 6.4.5 Regular Morphisms

For the category of quasi-projective varieties we introduce a number of equivalent definitions of morphism. In light of (6.4.20) then $b$) shows a function is regular iff it is regular in all affine coordinates.

**Proposition 6.4.23 (Regular Morphism)**
*Let $X$ be a projective variety, $U \subset X\left(\overline{k}\right)$ an open subset, $Y$ a quasi-projective variety and $\phi : U \to Y\left(\overline{k}\right)$ a function. Then the following are equivalent*

*a) **pulls back regular functions** $\phi$ is continuous and determines a morphism of sheaves*

$$\begin{aligned} \mathcal{O}_Y &\to \phi_\star \mathcal{O}_U \\ f &\to f \circ \phi \end{aligned}$$

© David Rufino CC BY-NC-ND 4.0

b) ***affine projections are regular*** *There is an open cover $U = \bigcup_{i\in I} U_i$ such that $\phi(U_i)$ is a subset of an affine chart $W_{j(i)}$ and*

$$\pi_k \circ \theta_{j(i)}^{-1} \circ \phi|_{U_i} : U_i \to \overline{k}$$

*is regular (6.4.19) for all $i \in I$ and $k = 1, ..., m$*

c) ***locally homogenous*** *There is an open cover $U = \bigcup_{i\in I} U_i$ and families of homogenous polynomials $\{f_{i0}, ..., f_{im}\} \subset k[X]_{\text{hom}}$ of the same degree such that $U_i \subset D_+(f_{i0}) \cup ... \cup D_+(f_{im})$ and*

$$\phi(y) = (f_{i0}(y) : ... : f_{im}(y)) \quad \forall y \in U_i$$

*In this case we say $\phi$ is **regular** or a **regular morphism** of quasi-projective varieties.*

*Proof.* $b) \implies a)$ We may assume wlog that $U_i$ is contained in an affine chart $V_i$. Therefore by definition and (6.4.20)

$$\theta_{j(i)}^{-1} \circ \phi \circ \theta_i \mid_{\theta_i^{-1}(U_i)} : \theta_i^{-1}(U_i) \to \mathbb{A}_k^m$$

is regular as a map of quasi-affine varieties. Therefore by (6.1.39) it is continuous. Whence $\phi|_{U_i}$ is continuous by (6.4.13), and $\phi$ is continuous by (4.1.27).

Suppose $f \in \mathcal{O}_Y(W)$ then it is enough to show for all $x \in \phi^{-1}(W)$ that $f \circ \phi$ is regular at $x$. Suppose $x \in U_i$ and replace $W$ with $W \cap W_{j(i)}$. Then by definition $f \circ \theta_{j(i)} : \theta_{j(i)}^{-1}(W) \to \overline{k}$ is regular at $\theta_{j(i)}^{-1}(x)$. Whence by (6.1.40)

$$f \circ \phi \circ \theta_i = \left(f \circ \theta_{j(i)}\right) \circ \left(\theta_{j(i)}^{-1} \circ \phi \circ \theta_i\right)$$

is regular at $\theta_i^{-1}(x)$, and therefore by (6.4.20) $f \circ \phi$ is regular at $x$.

$a) \implies b)$ For $x \in U$ there is some affine $W_j$ such that $\phi(x) \in W_j \implies x \in \phi^{-1}(W_j) =: V_x$. By (6.4.20) $\pi_k \circ \theta_j^{-1} : W_j \to \overline{k}$ is regular whence by assumption $\pi_k \circ \theta_j^{-1} \circ \phi|_{V_x}$ is regular. As every point has a nbhd $V_x$ we may find the required open cover.

$b) \implies c)$ By definition there exists homogenous $f_{i0}, ..., f_{im}, g_{i0}, ..., g_{im} \in k[X]_{\text{hom}}$ such that

$$\phi(y) = \left(\frac{f_{i0}(y)}{g_{i0}(y)} : ... : 1 : ... : \frac{f_{im}(y)}{g_{im}(y)}\right) \quad \forall y \in U_i$$

Therefore we may define

$$h_{il} := f_{il} \prod_{k\neq l} g_{ik}$$

Then evidently

$$\phi(y) = (h_{i0}(y) : ... : h_{im}(y)) \quad \forall y \in U_i$$

$c) \implies b)$ For every $x \in U$ there exists some open nbhd $V_x$ such that

$$\phi(y) = (f_0(y) : ... : f_m(y)) \quad \forall y \in V_x$$

Suppose that $f_j(x) \neq 0$ then replace $V_x$ with $V_x \cap D_+(f_j)$. Then

$$\left(\pi_k \circ \theta_j^{-1} \circ \phi|_{V_x}\right)(y) = \frac{f_k(y)}{f_j(y)}$$

which is regular by definition. The family $\{V_x\}_{x\in U}$ is the corresponding open cover. ∎

**Corollary 6.4.24**
*Let $\phi : X \to Y$ and $\psi : Y \to Z$ be regular morphisms of quasi-projective varieties. Then so is $\psi \circ \phi : X \to Z$.*

*Proof.* This is immediate by (6.4.23).a). ∎

**Corollary 6.4.25**
*A function $f : X\left(\overline{k}\right) \to \overline{k}$ is regular iff it coincides with a regular morphism $f : X\left(\overline{k}\right) \to D_+(X_0) \cong \mathbb{A}_k^1$. In particular a regular function is continuous with respect to the co-finite topology on $\overline{k}$.*

*Proof.* This follows from (6.4.23).b). ∎

© David Rufino CC BY-NC-ND 4.0

**Proposition 6.4.26**
*Let $X \subset \mathbb{P}^n_k$ be a projective variety, $U \subset X\left(\overline{k}\right)$ an open subset and $U = \bigcup_{i\in I} U_i$ an open cover. Suppose that we have $f_{i0}, ..., f_{im} \in k[X]_{\text{hom}}$ homogenous of the same degree such that*

*a) For every $i \in I$*

$$U_i \subset D_+(f_{i0}) \cup ... \cup D_+(f_{im})$$

*b) For every pair $U_i \cap U_{i'} \neq \varnothing$ we have*

$$f_{ik'}(y)f_{i'k}(y) = f_{ik}(y)f_{i'k'}(y) \quad \forall y \in U_i \cap U_{i'}, \forall k, k' = 0, ..., m$$

*Then there exists a unique regular morphism $f : U \to \mathbb{P}^m_k\left(\overline{k}\right)$ such that*

$$f(x) = (f_{i0}(x) : ... : f_{im}(x)) \quad \forall x \in U_i$$

## 6.4.6 Local Ring

**Proposition 6.4.27**
*Let $X = V_+(\mathfrak{b}) \subset \mathbb{P}^n_k$ be a projective variety and $W = V_+(\mathfrak{p})$ an irreducible subset. Then there is an isomorphism*

$$\begin{aligned} k[X]_{(\mathfrak{p}/\mathfrak{b})} &\xrightarrow{\sim} \mathcal{O}_{X,W} \\ \frac{\overline{F}}{\overline{G}} &\to \left[\left(D\left(\overline{G}\right), \frac{F(-)}{G(-)}\right)\right] \end{aligned}$$

*More precisely for any affine chart $U_i$ such that $U_i \cap W \neq \varnothing$ there is a commutative diagram*

$$\begin{array}{ccccc} k[U_i]_{\mathfrak{p}^a} & \xrightarrow[(3.6.45)]{\sim} & k[U_i \cap X]_{\mathfrak{p}^a/\mathfrak{b}^a} & \xrightarrow[(6.1.57)]{\sim} & \mathcal{O}_{U_i\cap X, U_i\cap W} \\ \wr\downarrow & & & & \downarrow\wr \\ & & & & (4.3.17) \\ k[X_0, ..., X_n]_{(\mathfrak{p})} & \xrightarrow[(3.12.18)]{\sim} & k[X]_{(\mathfrak{p}/\mathfrak{b})} & \dashrightarrow^{\sim} & \mathcal{O}_{X,W} \end{array}$$

*Proof.* We construct the left vertical map, and then the dashed arrow is uniquely defined.

As $U_i \cap W \neq \varnothing$ then $W \not\subset V(X_i) \implies X_i \notin \mathfrak{p}$. So by (6.4.12).d) $\mathfrak{p}^{ah} = \mathfrak{p}$. To complete the diagram we consider the mutually inverse maps

$$\begin{aligned} k[U_i] = k[x_0, ..., x_{i-1}, x_{i+1}, ..., x_n]_{\mathfrak{p}^a} &\to k[X_0, ..., X_n]_{(\mathfrak{p})} \\ \frac{F}{F'} &\to \frac{F\left(\frac{X_0}{X_i}, ..., \frac{X_n}{X_i}\right)}{F'\left(\frac{X_0}{X_i}, ..., \frac{X_n}{X_i}\right)} \\ \frac{G^a}{G'^a} &\leftarrow \frac{G}{G'} \end{aligned}$$

Note $F'^h \in \mathfrak{p} \implies F' = F'^{ha} \in \mathfrak{p}^a$. Therefore $F' \notin \mathfrak{p}^a \implies F'^h = X_i^{\deg(F)} F'\left(\frac{X_0}{X_1}, ..., \frac{X_n}{X_i}\right) \notin \mathfrak{p} \implies F'\left(\frac{X_0}{X_1}, ..., \frac{X_n}{X_i}\right) \notin \mathfrak{p}$. Similarly $G'^a \in \mathfrak{p}^a \implies G'^{ah} \in \mathfrak{p}^{ah} = \mathfrak{p} \implies G' = X_i^k G'^{ah} \in \mathfrak{p}$, so $G' \notin \mathfrak{p} \implies G'^a \notin \mathfrak{p}^a$. Finally

$$\frac{G^a\left(\frac{X_0}{X_i}, ..., \frac{X_n}{X_i}\right)}{G'^a\left(\frac{X_0}{X_i}, ..., \frac{X_n}{X_i}\right)} = \frac{X_i^{-\deg(G)} G}{X_i^{-\deg(G')} G'} = \frac{G}{G'}$$

so the maps are mutually inverse. ∎

**Corollary 6.4.28**
*Let $X$ be an irreducible projective variety with $W = V_+(\mathfrak{p})$ an irreducible subvariety. Then there is a commutative diagram*

© David Rufino CC BY-NC-ND 4.0

$$\begin{array}{ccc} k[U_i]_{\mathfrak{p}^a} & \hookrightarrow & k[U_i]_{(0)} \\ \downarrow \sim & & \downarrow \sim \\ k[X]_{(\mathfrak{p})} & \hookrightarrow & k[X]_{((0))} \\ \downarrow \sim & & \downarrow \sim \\ \mathcal{O}_{X,W} & \hookrightarrow & k(X) \end{array}$$

*In particular $k(X)$ is the field of fractions for $\mathcal{O}_{X,W}$.*

*Proof.* The vertical maps are from (6.4.27) and the horizontal maps are the obvious ones. As the top map is injective and corresponds to field of fractions we deduce that $k(X)$ is field of fractions for $\mathcal{O}_{X,W}$. ∎

## 6.4.7 Dimension

**Proposition 6.4.29**
*Projective space $\mathbb{P}^n_k$ has dimension $n$. Furthermore it is non-singular, normal and regular.*

*Proof.* Projective space is integral (6.4.8) and has an open subset isomorphic to $\mathbb{A}^n_k$. So we are done by (6.2.54) and (6.1.50).

The conditions are local (6.2.81) (3.24.10). ∎

**Proposition 6.4.30**
*Let $X = V_+(\mathfrak{p})$ be an irreducible projective variety then* $\dim k[X] = \dim X + 1$.

*Proof.* We have $\overline{X_i} \neq 0$ for some $i$ as $X$ is non-empty. Therefore

$$\dim k[X] \overset{(3.35.31)}{=} \dim k[X]_{\overline{X_i}} \overset{(3.13.8)}{=} \dim k[X]_{(\overline{X_i})}[T, T^{-1}] \overset{(3.9.3)}{=} \dim k[X]_{(\overline{X_i})}[T]_T \overset{(3.35.31)}{=} \dim k[X]_{(\overline{X_i})}[T] \overset{(3.35.29)}{=} \dim k[X]_{(\overline{X_i})} + 1$$

By (6.4.21) $\dim k[X]_{(\overline{X_i})} = \dim k[X \cap U_i] = \dim X \cap U_i$.

Finally this equals $\dim X$ (4.1.88). ∎

**Proposition 6.4.31**
*Let $X$ be a projective variety and $x \in X\left(\overline{k}\right)$. Then*

$$\overline{\{x\}} = \left\{\sigma(x) \;\middle|\; \sigma \in \operatorname{Aut}\left(\overline{k}/k\right)\right\}$$

*is an irreducible set of order $\left[k\left(\mathfrak{m}_{X,x}\right) : k\right]_s$. Further $X$ is a symmetric space.*

*Proof.* We have $x \in U_i$ for some affine chart then we may use (6.4.20) to reduce to the affine case (6.1.18). ∎

**Corollary 6.4.32**
*Let $X$ be a projective variety and $Z \subset X$ a closed subset. Then the following are equivalent*

*a)* $\dim Z = 0$
*b)* *$Z$ is finite*

*More precisely*

$$Z = \overline{\{x_1\}} \cup ... \cup \overline{\{x_n\}}$$

*for a finite number of points $x_1, ..., x_n \in Z$.*

*Proof.* This follows from (4.1.91) and (6.4.31). ∎

**Proposition 6.4.33**
*Let $X = \mathbb{P}^n_k\left(\overline{k}\right)$ and $G \in k[X_0, ..., X_n]$ an irreducible, homogenous polynomial of positive degree. Then* $\dim V_+(G) = n - 1$.

*Proof.* Let $Y := V_+(G)$ then by (...) we have $Y \neq \varnothing$. Choose an affine patch $U_i$ such that $U_i \cap Y \neq \varnothing$. Then $Y \cap U_i = V(G^a) \neq \varnothing$ which implies $G^a$ has positive degree. By (...) $\dim Y \cap U_i = n - 1$. ∎

© David Rufino CC BY-NC-ND 4.0

### 6.4.8 Valuative Completeness

**Lemma 6.4.34**
*Let $R$ be a valuation ring for $K$. Suppose $x_0, ..., x_n \in K$ not all zero, then there exists $0 \leq j \leq n$ such that $\frac{x_i}{x_j} \in R$ for all $i = 0, ..., n$.*

*Proof.* Recall (3.25.17) that the abelian group $K^\star/R^\star$ is totally ordered by the condition $x \leq y \iff yx^{-1} \in R^\star$. Choose $j$ for which $x_j \neq 0$ such that $x_j$ is minimal under this ordering. ∎

**Proposition 6.4.35 (Projective Varieties are Complete)**
*Let $X = V_+(\mathfrak{p}) \subset \mathbb{P}^n_k$ be an irreducible projective variety. Let $k \subset R \subset k(X)$ be a valuation ring. Then there exists a point $x \in X\left(\overline{k}\right)$ such that $\mathcal{O}_{X,x} \subset R$ and $\mathfrak{m}_R \cap \mathcal{O}_{X,x} = \mathfrak{m}_{X,x}$.*

*More precisely there exists a local homomorphism $\Phi : R \to \overline{k}$ such that $\Phi|_{\mathcal{O}_{X,x}} = \mathrm{ev}_x$.*

*Proof.* Denote the homogenous coordinate ring by simply $k[X]$. In light of (6.4.28) we may assume that $k(X) = k[X]_{((0))}$ and $\mathcal{O}_{X,x} = k[X]_{\mathfrak{p}_x}$ with $\mathfrak{m}_{X,x} = \left(\mathfrak{p}_x k[X]_{\mathfrak{p}_x}\right)_{(0)}$.

By (...) we may assume that $k(\mathfrak{m}_R)/k$ is algebraic and therefore by (3.19.61) there exists a local $k$-algebra homomorphisms

$$\Phi : R \to k(\mathfrak{m}_R) \hookrightarrow \overline{k}$$

We may assume wlog that $X \cap U_0 \neq \varnothing$ and therefore $X_0 \notin \mathfrak{p}$ where $X_0, ..., X_n$ are the homogenous coordinates. Define $x_i := \frac{X_i}{X_0} \in k[X]_{((0))}$ for $i = 0, ..., n$. By (6.4.34) there is $1 \leq j \leq n$ such that $t_i := \frac{x_i}{x_j} = \frac{\overline{X}_i}{\overline{X}_j} \in R$ for all $i = 0, ..., n$.

Note that $t_j = 1 \notin \mathfrak{m}_R$ so $\Phi\left(t_j\right) \neq 0$ we have a well-defined projective point $x = [\Phi(t_0) : ... : \Phi(t_n)] \in \mathbb{P}^n_k\left(\overline{k}\right)$. For any homogenous polynomial $F \in k[X_0, ..., X_n]$ of degree $d$ we have

$$\begin{aligned} F(t_0, ..., t_n) &= \overline{X}_j^d F\left(\overline{X}_0, ..., \overline{X}_n\right) \\ &= \overline{X}_j^d \overline{F} \end{aligned}$$

regarding $k(X)$ as a subring of $\mathrm{Frac}(k[X])$ and $\bar{\cdot}$ denoting the reduction map $k[X_0, ..., X_n] \to k[X]$. Further

$$F(\Phi(t_0), ..., \Phi(t_n)) = \Phi(F(t_0, ..., t_n))$$

If $F \in \mathfrak{p}$ then $\overline{F} = 0$ and therefore $F(x) = 0$. This shows that $x \in X\left(\overline{k}\right)$.

We require to show that $\mathcal{O}_{X,x} \subset R$ and $\Phi|_{\mathcal{O}_{X,x}} = \mathrm{ev}_x$, from which it immediately follows that $\mathfrak{m}_R \cap \mathcal{O}_{X,x} = \mathfrak{m}_{X,x}$

Consider $g, h \in k[X]$ homogenous of the same degree with $h \notin \mathfrak{p}_x$. We claim that $g/h \in R$. For there exists $G, H \in k[X_0, ..., X_n]$ homogenous such that

$$\begin{aligned} g &= G\left(\overline{X_0}, ..., \overline{X_n}\right) \\ &= \overline{X}_j^d G(t_0, ..., t_n) \\ h &= H\left(\overline{X_0}, ..., \overline{X_n}\right) \\ &= \overline{X}_j^d H(t_0, ..., t_n) \\ \frac{g}{h} &= \frac{G(t_0, ..., t_n)}{H(t_0, ..., t_n)} \end{aligned}$$

Evidently $G(t_0, ..., t_n) \in R$ since $R$ is a $k$-algebra. Further by assumption

$$\begin{aligned} 0 &\neq h(\Phi(t_0), ..., \Phi(t_n)) \\ &= H(\Phi(t_0), ..., \Phi(t_n)) \\ &= \Phi(H(t_0, ..., t_n)) \end{aligned}$$

which means $H(t_0, ..., t_n) \notin \mathfrak{m}_R \implies H(t_0, ..., t_n) \in R^\star$. Using the expression above for $\frac{g}{h}$ shows it lies in $R$ as required. Further

© David Rufino CC BY-NC-ND 4.0

$$\begin{aligned}\Phi\left(\frac{g}{h}\right) &= \frac{\Phi(G(t_0, ..., t_n))}{\Phi(H(t_0, ..., t_n))} \\ &= \frac{G(\Phi(t_0), ..., \Phi(t_n))}{H(\Phi(t_0), ..., \Phi(t_n))} \\ &= \frac{G(x)}{H(x)} \\ &= \frac{g}{h}(x)\end{aligned}$$

as required. ■

© David Rufino CC BY-NC-ND 4.0

## 6.5 Arithmetic Varieties

**Remark 6.5.1**
*This section is rough working notes rather than a polished treatment, kept here for personal reference. Omitted from the HTML editions.*

**Proposition 6.5.2**
*Let $\mathfrak{o}$ be a ring with quotient field $k$. For an ideal $\mathfrak{a} \triangleleft \mathfrak{o}[X_1, ..., X_n]$ define the zero-locus*

$$|X| := V(\mathfrak{a}) := \left\{ x \in \overline{k}^n \;\middle|\; f(x_1, ..., x_n) = 0 \quad \forall f \in \mathfrak{a} \right\}$$

*and the coordinate ring $\mathfrak{o}[X] := \mathfrak{o}[X_1, ..., X_n]/\mathfrak{a}$. Then*

$$|X| = V(\mathfrak{a}_k) \subset \overline{k}^n$$

$$\mathfrak{a}_k := \left\{ \sum_{i=1}^{n} \lambda_i f_i \in k[X_1, ..., X_n] \;\middle|\; f_i \in \mathfrak{a}, \lambda_i \in k \right\}$$

*where $\mathfrak{a}_k \triangleleft k[X_1, ..., X_n]$. For every $f \in \mathfrak{o}[X_1, ..., X_n]/\mathfrak{a}$ the sets of the form*

$$D(f) := \{x \in |X| \mid f(x) \neq 0\}$$

*form a basis for the usual Zariski topology. For every principal open set $D(f) \subset |X|$ define*

$$\mathcal{O}_X(D(f)) := \left\{ \phi : D(f) \to \overline{k} \;\middle|\; \phi(x) = \frac{g(x)}{f^r(x)} \forall x \in D(f), \text{some } g \in \mathfrak{o}[X], r \geq 0 \right\}$$

*then $\mathcal{O}_X$ extends uniquely to sheaf of $\overline{k}$-valued functions with the obvious $\mathfrak{o}$-algebra structure.*

© David Rufino CC BY-NC-ND 4.0

## 6.6 Scheme Notes

**Remark 6.6.1**
*This section is rough working notes rather than a polished treatment, kept here for personal reference. Omitted from the HTML editions.*

### 6.6.1 Schemes over $\mathbb{Z}$

Let $A$ be a finitely-generated $\mathbb{Z}$-algebra, it may be expressed in the form $A = \mathbb{Z}[X_1, ..., X_n]/\mathfrak{a}$. For any $\mathbb{Q}$-algebra $R$ the rational points are given by $\mathrm{AlgHom}_{\mathbb{Z}}(A, [R]_{\mathbb{Z}}) = \mathrm{AlgHom}_{\mathbb{Q}}(A \otimes_{\mathbb{Z}} \mathbb{Q}, R)$.

### 6.6.2 Closed Points vs Maximal Ideals

In the case of affine schemes there is a straightforward relationship between closed points and maximal ideals.

**Proposition 6.6.2**
*Let $X = \mathrm{Spec}(A)$ be an affine scheme. Then the closed points of $X$ correspond precisely to maximal ideals of $A$.*

For a general scheme a closed point corresponds to a maximal ideal of all affine neighbourhoods.

**Proposition 6.6.3**
*Let $X$ be a scheme, $x \in X$ and $U \subset X$ an affine neighbourhood, with $U \xrightarrow{\sim} \mathrm{Spec}(A)$. Then if $x$ is a closed in $X$ we have $x$ is closed in $U$ and corresponds to a maximal ideal of $A$.*

**Remark 6.6.4**
*However the converse may not hold – there may be maximal ideals of affines subsets which do not correspond to a closed point. As an extreme example it's possible to construct a scheme with no closed points at all. Consider a local ring $A$ with the following ideal structure*

$$0 \subsetneq \mathfrak{p}_0 \subsetneq ... \subsetneq \mathfrak{p}_n \subsetneq ... \subsetneq \mathfrak{m} \subsetneq A$$

*Then $\mathrm{Spec}(A) \setminus \{[\mathfrak{m}]\}$ is an open subscheme with no closed points.*

*More simply consider the spectrum of a discrete valuation ring $\mathrm{Spec}(A) = \{(0), \mathfrak{m}\}$. Then $\{(0)\}$ is not closed in $X := \mathrm{Spec}(A)$, but it is open (being the complement of $V(\mathfrak{m})$) and even isomorphic to $\mathrm{Spec}\big(A_{(0)}\big)$, and so $(0)$ is closed in an affine neighbourhood.*

However we may obtain some partial converses

**Proposition 6.6.5**
*Let $X$ be a quasi-compact scheme, $U \subset X$ an affine open neighbourhood and $x \in U$ a closed point. Then there exists $y \in \overline{\{x\}}$ such that $y$ is closed.*

**Proposition 6.6.6**
*Let $X$ be a integral scheme locally of finite type over $k$. Then if $x \in U$ is closed for $U$ affine, it is also closed in $X$.*

*Proof.* By assumption we may write $X = \bigcup_{i \in I} V_i$ where $V_i$ are affine (as finite type $\implies$ quasi-compact). Then by (4.1.21)

$$\mathrm{cl}_X(\{x\}) \cap V_i = \mathrm{cl}_{V_i}(\{x\})$$

whenever $x \in V_i$ (and it is empty otherwise). Therefore it is sufficient to show that $x \in V_i \implies x$ closed in $V_i$.

So we may consider the following situation: $U = \mathrm{Spec}(A)$ and $V = \mathrm{Spec}(B)$ are affine open neighbourhoods of $x$, for $A, B$ finitely generated $k$-algebras, where additionally $x$ is closed in $U$. Then by (6.6.2) $x$ corresponds to a maximal ideal $\mathfrak{m}$ of $A$, and a prime ideal of $\mathfrak{p}$ of $B$. As these are both stalks of $X$ we have a local $k$-algebra isomorphism $A_{\mathfrak{m}} \xrightarrow{\sim} B_{\mathfrak{p}}$, and therefore also of residue fields $k(\mathfrak{m}) \xrightarrow{\sim} k(\mathfrak{p})$. Therefore by following lemma we have $x$ is closed in $V$ as required. ∎

**Lemma 6.6.7**
*Let $A$ be a finitely-generated $k$-algebra and $\mathfrak{p}$ a prime ideal. Then the following are equivalent*

*a) $\mathfrak{p}$ is maximal*
*b) $[\mathfrak{p}]$ is a closed point of $\mathrm{Spec}(A)$.*
*c) $k(\mathfrak{p})/k$ is algebraic*
*d) $k(\mathfrak{p})/k$ is finite*

*In particular the closed points of $\mathrm{Spec}(A)$ correspond precisely to the maximal ideals of $A$.*

© David Rufino CC BY-NC-ND 4.0

*Proof.* First we observe that $k(\mathfrak{p})/k$ is always finitely generated, so $c) \iff d)$ is simply (3.19.44). Further $a) \iff b)$ is simply (6.6.2).

$d) \implies a)$ We have an inclusion $A/\mathfrak{p} \hookrightarrow k(\mathfrak{p})/k$. We deduce that $A/\mathfrak{p}$ is a finite-dimensional, integral $k$-algebra and therefore a field (3.35.56).

$a) \implies d)$ As $A$ is finitely generated, so is $k(\mathfrak{p})/k$. By Zariski's Lemma (3.35.23) it is finite. ■

© David Rufino CC BY-NC-ND 4.0

# Bibliography

[**AM69**] M. Atiyah and I.G. McDonald. *Introduction to Commutative Algebra*. Westview Press, 1969.

[**ARV10**] Jiří Ad\'amek, Jiří Rosický, and Enrico M. Vitale. *Algebraic Theories: A Categorical Introduction to General Algebra*. Cambridge University Press, 2010.

[**BS81**] Stanley Burris and H.P. Sankappanavar. *A Course in Universal Algebra*. Springer, 1981.

[**Bir35**] Garrett Birkhoff. On the Structure of Abstract Algebras. *Mathematical Proceedings of the Cambridge Philosophical Society* 31(4), 433--454, 1935.

[**Bir40**] G. Birkhoff. *Lattice Theory*. American Mathematical Society, 1940.

[**Bor94**] Francis Borceux. *Handbook of Categorical Algebra 2: Categories and Structures*. Cambridge University Press, 1994.

[**Bou89**] Nicolas Bourbaki. *Algebra: Chapters 4-7*. Springer-Verlag, 1989.

[**Bou98a**] Nicolas Bourbaki. *Algebra I: Chapters 1-3*. Springer, 1998.

[**Bou98b**] N. Bourbaki. *Commutative Algebra: Chapters 1-7*. Springer, 1998.

[**Bum98**] D. Bump. *Algebraic Geometry*. World Scientific, 1998.

[**Car71**] Henri Cartan. *Differential Calculus*. London: Kershaw Publishing Company, 1971.

[**Che51**] C. Chevalley. *Introduction to the Theory of Algebraic Functions of One Variable*. American Mathematical Society, 1951.

[**Die11**] Jean Dieudonné. *Foundations of modern analysis*. Read Books Ltd, 2011.

[**Eme10**] Matthew Emerton. Jacobson Rings, 2010.

[**For81**] Otto Forster. *Lectures on Riemann Surfaces*, 1981.

[**Fre64**] Peter Freyd. *Abelian Categories: An Introduction to the Theory of Functors*. Harper & Row, 1964.

[**Ful08**] William Fulton. Algebraic Curves. , 2008.

[**Gro57**] Alexander Grothendieck. Sur quelques points d'algèbre homologique. *Tôhoku Mathematical Journal* 9(2), 119--221, 1957.

[**Gro60**] Alexander Grothendieck. Éléments de géométrie algébrique : {I.} {Le} langage des schémas. *Publications Mathématiques de l'IHÉS* 4, 5--228, 1960.

[**Gro64**] Alexander Grothendieck. Éléments de géométrie algébrique : {IV.} {Étude} locale des schémas et des morphismes de schémas, {Première} partie. *Publications Mathématiques de l'IHÉS* 20, 5--259, 1964.

[**Hal17**] P.R. Halmos. *Naive Set Theory*. Dover Publications, 2017.

[**Har13**] Robin Hartshorne. *Algebraic Geometry*. Springer New York, 2013.

[**Hei17**] Katharina Heinrich. Some remarks on biequidimensionality of topological spaces and Noetherian schemes. *Journal of Commutative Algebra* 9(1), 49--63, 2017.

[**Hol11**] Audun Holme. *A royal road to algebraic geometry*. Springer-Verlag New York, 2011.

[**Kap74**] Irving Kaplansky. *Commutative Rings*. The University of Chicago Press, 1974.

[**Kel71**] John L Kelley. *General Topology*. Springer, 1971.

[**Lan11**] Serge Lang. *Algebra*. Springer New York, 2011.

[**Lan12**] Serge Lang. *Real and functional analysis*. Springer Science & Business Media, 2012.

[**Lan19**] Serge Lang. *Introduction to Algebraic Geometry*. Dover Publications, 2019.

[**Lan72**] Serge Lang. *Differential manifolds*. Springer, 1972.

[**Lan82**] Serge Lang. *Introduction to Algebraic and Abelian Functions*. Springer-Verlag, 1982.

[**Lan98**] Saunders Mac Lane. *Categories for the Working Mathematician*. Springer, 1998.

[**Liu06**] Qing Liu. *Algebraic Geometry and Arithmetic Curves*. Oxford University Press, 2006.

[**Mat70**] H. Matsumura. *Commutative Algebra*. Benjamin, 1970.

[**Mil14**] James Milne. Introduction to Commutative Algebra, 2014.

[**Mil17**] James Milne. Algebraic Geometry, 2017.

[**Mor12**] Patrick Morandi. *Field and Galois theory*. Springer Science & Business Media, 2012.

[**Mum99**] David Mumford. *The red book of varieties and schemes : includes the Michigan Lectures (1974) on Curves and their Jacobians*. New York: Springer, 1999.

[**Nag75**] M. Nagata. *Local Rings*. R.E. Krieger Publishing Company, 1975.

[**Neu99**] Jürgen Neukirch. *Algebraic Number Theory*. Berlin: Springer-Verlag, 1999.

[**Rey11**] Manny Reyes. Krull dimension less or equal than transcendence degree?, 2011. Math Overflow.

© David Rufino CC BY-NC-ND 4.0

[**Rie16**] Emily Riehl. *Category Theory in Context*. Dover Publications, 2016.

[**Rom05**] S. Roman. *Field Theory*. Springer New York, 2005.

[**Ser55**] Jean-Pierre Serre. Faisceaux algébriques cohérents. *Annals of Mathematics* 61(2), 197--278, 1955.

[**Ser59**] Jean-Pierre Serre. *Groupes algébriques et corps de classes*. Hermann, 1959.

[**Ser79**] Jean-Pierre Serre. *Local Fields*. New York: Springer-Verlag, 1979.

[**Sha00**] R.Y. Sharp. *Steps in Commutative Algebra*. Cambridge University Press, 2000.

[**Sha94**] Igor Shafarevich. *Basic algebraic geometry*. Springer-Verlag New York, 1994.

[**Shi55**] Goro Shimura. Reduction of Algebraic Varieties with Respect to a Discrete Valuation of the Basic Field. *American Journal of Mathematics* 77(1), 134--176, 1955.

[**Stacks**] The Stacks Project Authors. Stacks Project, 2015.

[**Sti09**] Henning Stichtenoth. *Algebraic Function Fields and Codes*. Springer Berlin Heidelberg, 2009.

[**Tat68**] John Tate. Residues of differentials on curves. *Annales scientifiques de l'École Normale Supérieure* 1(1), 149--159, 1968.

[**Wae91**] B. L. van der Waerden. *Algebra: Volume I*. Springer New York, 1991.

[**War13**] F. Warner. *Foundations of Differentiable Manifolds and Lie groups*. Springer-Verlag New York, 2013.

[**Wed16**] Torsten Wedhorn. *Manifolds, Sheaves, and Cohomology*. Springer, 2016.

[**Wil70**] Stephen Willard. *General Topology*. Dover Publications, 1970.

[**ZS75**] Oscar Zariski and Pierre Samuel. *Commutative Algebra I*. Springer New York, 1975.

[**ZS76**] Oscar Zariski and Pierre Samuel. *Commutative Algebra II*. Springer New York, 1976.

© David Rufino CC BY-NC-ND 4.0