# 3.33 Extensions of Power Series Rings

We record the Weierstrass division lemma for formal power series, used in the analysis of finite extensions of $k((t))$ below.

**Remark 3.33.1**
*[Che51] (Ch. II, §6) treats the case of (3.33.4) where the ramification index $e$ is prime to $\mathrm{char}(k)$, obtaining the sharper conclusion that $t$ itself is an $e$-th power in $k((u))$ (a literal Puiseux expansion $k\big((t^{1/e})\big)$), via Hensel's lemma applied to $W^e - v$ for the unit $v := t/u^e$. That argument is unavailable when $\mathrm{char}(k) = p \mid e$: the derivative $eW^{e-1}$ vanishes identically, so $W^e - v$ has no simple root to lift, and $t$ need not be an $e$-th power at all (an Artin-Schreier extension gives a counterexample). (3.33.4) instead only produces $t = u^e c$ for **some** unit $c$, obtained by solving $\tau G(\tau) = u^e$ by direct coefficient-stabilisation rather than root extraction; this sidesteps the characteristic obstruction entirely, at the cost of the sharper radical form, which is in any case not needed below.*

**Lemma 3.33.2 (Free Basis for a Distinguished Power Series)**
*Let $R$ be a commutative ring, $m \geq 1$ an integer, $w \in R[[X]]^\star$, and $T := X^m w \in R[[X]]$ (so the coefficients of $T$ below degree $m$ vanish, and its coefficient of $X^m$ is $w(0) \in R^\star$). Then every $p \in R[[X]]$ is uniquely*

$$p = \sum_{i=0}^{m-1} f_i(T) X^i$$

*for power series $f_0, \ldots, f_{m-1} \in R[[Y]]$ (substituted at $Y = T$, well-defined as $T$ has zero constant term (3.10.1)).*

*Proof. Drafted by Claude Sonnet 5, reviewed by David Rufino.*

**Existence.** Define $p^{(0)} := p$ and, recursively for $j \geq 0$: writing $p^{(j)} = \sum_n p^{(j)}_n X^n$, set $r_j := \sum_{i=0}^{m-1} p^{(j)}_i X^i$ (the degree-$\{<\}m$ truncation) and $\tilde{q}_j := \sum_{n \geq 0} p^{(j)}_{n+m} X^n$ (the shift), so $p^{(j)} = \tilde{q}_j X^m + r_j$; put $p^{(j+1)} := \tilde{q}_j w^{-1} \in R[[X]]$ (using $w \in R[[X]]^\star$ (3.10.2)), so that $p^{(j)} = p^{(j+1)} T + r_j$. Iterating,

$$p = p^{(J+1)} T^{J+1} + \sum_{j=0}^{J} r_j T^j \quad \text{for every } J \geq 0.$$

As $w(0) \in R^\star$, $T^{J+1} = X^{m(J+1)} w^{J+1}$ has zero coefficients below degree $m(J+1)$; so for each fixed $n$, the coefficient of $X^n$ in $p$ equals the coefficient of $X^n$ in $\sum_{j=0}^{J} r_j T^j$ once $m(J+1) > n$, and this common value no longer depends on $J$ thereafter. Writing $r_j = \sum_{i=0}^{m-1} \gamma_{j,i} X^i$, set $f_i(Y) := \sum_{j \geq 0} \gamma_{j,i} Y^j \in R[[Y]]$; the above shows $p = \sum_{i=0}^{m-1} f_i(T) X^i$, matching the coefficient of every $X^n$.

**Uniqueness.** Suppose $\sum_{i=0}^{m-1} f_i(T) X^i = 0$ with not every $f_i$ zero. Let $j_0 \geq 0$ be minimal such that some $f_i$ has a nonzero coefficient of $Y^{j_0}$, and let $i^\star$ be the least such $i$, so $f_{i^\star}(Y) = \gamma Y^{j_0} + O(Y^{j_0+1})$ for some $0 \neq \gamma \in R$, while $f_i(Y) = O(Y^{j_0+1})$ for $i < i^\star$. Then $f_i(T) = O\big(X^{m(j_0+1)}\big)$ for $i < i^\star$, and $f_{i^\star}(T) = \gamma T^{j_0} + O\big(X^{m(j_0+1)}\big) = \gamma w(0)^{j_0} X^{m j_0} + O(X^{m j_0 + 1})$; as $m j_0 + i^\star < m(j_0+1)$, the coefficient of $X^{m j_0 + i^\star}$ in $\sum_i f_i(T) X^i$ is exactly $\gamma w(0)^{j_0}$, nonzero (a nonzero element of $R$ times a unit is nonzero) – contradicting the sum being 0. ∎

**Corollary 3.33.3 (Specialisation Compatibility)**
*In the setting of (3.33.2), let $\phi : R \to S$ be a homomorphism of commutative rings, extended coefficientwise to $\phi : R[[X]] \to S[[X]]$. Then $\phi(w) \in S[[X]]^\star$, and the decomposition of $\phi(p)$ relative to $\phi(T) = X^m \phi(w)$ is $\phi(p) = \sum_{i=0}^{m-1} \phi(f_i)(\phi(T)) X^i$, where $\phi(f_i) \in S[[Y]]$ denotes $\phi$ applied coefficientwise to $f_i$.*

*Proof.* If $wv = 1$ then $\phi(w)\phi(v) = 1$, so $\phi(w) \in S[[X]]^\star$. Substitution of a zero-constant-term series into another is computed, coefficient by coefficient, by a fixed finite combination of additions and multiplications of the two series' coefficients; as $\phi$ is additive and multiplicative, it commutes with this computation, so applying $\phi$ to $p = \sum_i f_i(T) X^i$ gives $\phi(p) = \sum_i \phi(f_i)(\phi(T)) X^i$. This exhibits $\phi(p)$ in the form guaranteed unique by (3.33.2) over $S$, so it is **the** decomposition of $\phi(p)$. ∎

**Lemma 3.33.4 (Finite Separable Extensions of $k((t))$ are Power Series Fields)**
*Let $k$ be algebraically closed, $F := k((t))$, and $E/F$ a finite separable extension of degree $n$. Then there are $u \in E$, an element $\tau \in uk[[u]]$ with $\tau = u^n c$ for some $c \in k[[u]]^\star$, and a field isomorphism $\Psi : E \xrightarrow{\sim} k((u))$ with $\Psi|_F$ the substitution $t \mapsto \tau$ and $\Psi(u) = u$. In particular $1, u, \ldots, u^{n-1}$ is an $F$-basis of $E$.*

*Proof. Drafted by Claude Sonnet 5, reviewed by David Rufino.*

**Step 1: a DVR.** Let $\mathcal{O}_F := k[[t]]$ and let $C$ be the integral closure of $\mathcal{O}_F$ in $E$. By (3.24.26) $C$ is a finite $\mathcal{O}_F$-module, and by (3.32.9) it is Dedekind. By lying-over (3.24.17) there is a prime $\mathfrak{p} \subset C$ with $\mathfrak{p} \cap \mathcal{O}_F = (t)$. Put $B := C_{\mathfrak{p}}$, a DVR with fraction field $E$, and let $e := v_B(t) \geq 1$.

**Step 2: residue field and valuation.** $C/\mathfrak{p}$ is a domain, finite over $\mathcal{O}_F/(t) = k$, so $k \hookrightarrow C/\mathfrak{p}$ is injective and integral, hence $C/\mathfrak{p}$ is a field (3.24.14), finite over $k$. As $B = C_{\mathfrak{p}}$, (3.6.43) gives $B/\mathfrak{m}_B \cong \mathrm{Frac}(C/\mathfrak{p}) = C/\mathfrak{p}$ ($C/\mathfrak{p}$ already a field); as $k$ is algebraically closed, $k \to B/\mathfrak{m}_B$ is an isomorphism. Every $a \in \mathcal{O}_F^\times$ is a unit of $B$, so $v_B(x) = ev_F(x)$ for $x \in F^\times$. Let $u$ be a uniformiser of $B$, so $t = u^e w$ with $w \in B^\star$.

**Step 3: expansions.** Every $b \in B$ can be written $b = \sum_{i<e} f_i u^i$ with $f_i \in k[[t]]$. We show by induction on $N \geq 0$ that $b \equiv \sum_{i<e}\left(\sum_{j<N} c_{ij}t^j\right)u^i \pmod{t^N B}$ for some $c_{ij} \in k$. The case $N = 0$ is trivial. For the inductive step, write the current remainder as $t^N y$ with $y \in B$ (inductive hypothesis); since $B/uB = k$, repeated division by $u$ gives $y = \sum_{i<e} d_i u^i + u^e y'$ with $d_i \in k$, $y' \in B$ (there is no need to iterate past degree $e-1$: stop as soon as $e$ terms are extracted). Using $u^e = tw^{-1}$,

$$t^N y = \sum_{i<e}(t^N d_i)u^i + t^{N+1}(w^{-1}y'),$$

with $w^{-1}y' \in B$; so taking $c_{i,N} := d_i$ advances the induction, with new remainder $t^{N+1}(w^{-1}y') \in t^{N+1}B$. Put $f_i := \sum_j c_{ij}t^j \in k[[t]]$. Then $z := b - \sum_i f_i u^i$ satisfies $z \in t^N B$ for every $N$, so $v_B(z) \geq eN$ for all $N$ and $z = 0$.

**Step 4: basis and degree.** Every element of $E$ has the form $t^{-N}b$ with $b \in B$, so Step 3 gives $E = \sum_{i<e} Fu^i$. These elements are independent: if $\sum_{i<e} f_i u^i = 0$ with $f_i \in F$ not all zero, the valuations $v_B(f_i u^i) = ev_F(f_i) + i$ are distinct modulo $e$ for the nonzero $f_i$, so the minimum is attained exactly once, a contradiction. Hence $n = e$, and $1, \ldots, u^{e-1}$ is a basis. Moreover $B = \bigoplus_{i<e} k[[t]]u^i$, by the valuation estimate $\min_i(ev_F(f_i) + i) = v_B(b) \geq 0$ applied to $b = \sum_i f_i u^i$ with $f_i \in F$.

**Step 5: the minimal polynomial.** Write $w^{-1} = \sum_{i<e} g_i u^i$ with $g_i \in k[[t]]$ (Step 3). The constant $g_0(0)$ is the residue of $w^{-1}$, so $g_0(0) \in k^\star$. From $u^e = tw^{-1}$, $u$ is a root of

$$P(Y) := Y^e - t\sum_{i<e} g_i(t)Y^i \in F[Y].$$

Since $F[u] = E$ has degree $e$ over $F$, $P$ is the minimal polynomial of $u$, and $E \cong F[Y]/(P)$ with $Y \mapsto u$.

**Step 6: solving for $\tau$.** For $s \in uk[[u]]$ put $G(s) := \sum_{i<e} g_i(s)u^i \in k[[u]]$, where $g_i(s)$ is the substitution (well defined as $s$ has zero constant term). Since $G(s) \equiv g_0(0) \pmod{u}$, $G(s)$ is a unit. Define $T(s) := u^e G(s)^{-1} \in u^e k[[u]]$. For $s \equiv s' \pmod{u^N}$ we have $g_i(s) - g_i(s') \in (s-s')k[[u]] \subseteq u^N k[[u]]$, hence $G(s) \equiv G(s') \pmod{u^N}$ and $T(s) \equiv T(s') \pmod{u^{N+e}}$. Starting from $s_0 := 0$ and setting $s_{m+1} := T(s_m)$, the sequence satisfies $s_{m+1} \equiv s_m \pmod{u^{m+e}}$, so each coefficient stabilises. Let $\tau \in uk[[u]]$ be the coefficientwise limit. Then $\tau \equiv T(\tau) \pmod{u^N}$ for every $N$, so $\tau = T(\tau)$, i.e. $\tau G(\tau) = u^e$. Also $\tau = u^e c$ with $c \in k[[u]]^\star$, since $\tau G(\tau) = u^e$ and $G(\tau)$ is a unit.

**Step 7: the homomorphism.** The substitution $t \mapsto \tau$ defines a ring homomorphism $\sigma : k[[t]] \to k[[u]]$ ((3.10.11), as $\tau$ has zero constant term), and since $\tau$ is a unit times $u^e$ it extends to a field homomorphism $\sigma : F \to k((u))$. The homomorphism $F[Y] \to k((u))$, $Y \mapsto u$, extending $\sigma$, kills $P$:

$$\sigma(P)(u) = u^e - \tau\sum_{i<e} g_i(\tau)u^i = u^e - \tau G(\tau) = 0.$$

So it factors through $E \cong F[Y]/(P)$, giving a field homomorphism $\Psi : E \to k((u))$ with $\Psi|_F = \sigma$ and $\Psi(u) = u$. It is injective, as $E$ is a field.

**Step 8: surjectivity.** Since $\tau = u^e c$ with $c \in k[[u]]^\star$, the Weierstrass division lemma (3.33.2) (with $R := k$, $X := u$, $m := e$, $w := c$, so $T = \tau$) gives $k[[u]] = \bigoplus_{i<e} \sigma(k[[t]])u^i$. Multiplying by powers of $\tau$ gives $k((u)) = \sum_{i<e} \sigma(F)u^i = \sigma(F)[u]$. But $\operatorname{im}\Psi = \Psi(F[u]) = \sigma(F)[u]$, so $\Psi$ is surjective.

Hence $\Psi : E \xrightarrow{\sim} k((u))$ is an isomorphism of fields over $F$, with $\Psi(u) = u$ and $\Psi(t) = \tau = u^n c$. ∎

**Corollary 3.33.5 (Uniqueness of the Extended Valuation)**
*In the setting of (3.33.4), let $C$ be the integral closure of $k[[t]]$ in $E$. Then $C$ has exactly one prime $\mathfrak{p}$ lying over $(t)$, the DVR $C_{\mathfrak{p}}$ is the unique valuation ring of $E$ containing $k[[t]]$ whose maximal ideal contracts to $(t)$, and its ramification index is $[E : F]$.*

*Proof. Drafted by Claude Sonnet 5, reviewed by David Rufino.*

Let $\mathfrak{q}$ be any prime of $C$ over $(t)$, which exists by lying-over. Steps 2–4 of the proof of (3.33.4) apply verbatim to the DVR $B := C_{\mathfrak{q}}$ (which has residue field $k$), and give $e_{\mathfrak{q}} = v_B(t) = [E : F] = n$.

Now apply the Ramification Formula (3.32.16) to the AKLB situation $k[[t]] \subset C$ over $E/F$ (both Dedekind by (3.32.9), and $E/F$ separable), at the prime $(t)$:

$$\sum_{\mathfrak{q}\mid(t)} e_{\mathfrak{q}} f_{\mathfrak{q}} = [C/tC : k] = n.$$

Each $f_{\mathfrak{q}} = [C/\mathfrak{q} : k]$ is finite, and $C/\mathfrak{q}$ is a field algebraic over $k$, hence $f_{\mathfrak{q}} = 1$. So $\sum_{\mathfrak{q}} e_{\mathfrak{q}} = n$. By the previous paragraph each term equals $n$, so there is exactly one prime $\mathfrak{p}$ over $(t)$ and $e_{\mathfrak{p}} = n$.

For uniqueness of the valuation ring, let $V \subseteq E$ be a valuation ring with $k[[t]] \subseteq V$ and $\mathfrak{m}_V \cap k[[t]] = (t)$. By (3.25.14), $C \subseteq V$. Then $\mathfrak{m}_V \cap C$ is a prime of $C$ over $(t)$, hence equals $\mathfrak{p}$. Every element of $C \setminus \mathfrak{p}$ lies outside $\mathfrak{m}_V$, so is a unit of $V$; hence $C_{\mathfrak{p}} \subseteq V$. Since $C_{\mathfrak{p}}$ is a valuation ring of $E$ and valuation rings are maximal for domination among proper subrings of $E$ (3.25.2), $V = C_{\mathfrak{p}}$. ∎