Hurt People Hurt People

M Harper and V Knight

August 18, 2026

Abstract

Adverse childhood experiences are strongly associated with lifelong health, social, and economic outcomes, and increasing evidence suggests that their effects may propagate across generations. While empirical studies document the prevalence, transmission, and consequences of trauma, there is currently no unified framework linking these individual-level mechanisms to population-level dynamics. Here we combine a minimal analytical model with a data-informed simulation model to study the long-term evolution of trauma prevalence. The analytical model yields a closed-form threshold determining when trauma becomes self-sustaining, while the simulation model incorporates heterogeneous exposure levels and empirically grounded transmission patterns. The models show how demographic processes, intergenerational transmission, and recovery interact to produce stable endemic trauma levels or long-run decline, and establish the conditions under which interventions can eliminate endemic prevalence.

1 Introduction

Adverse childhood experiences (ACEs) are widely recognised as major determinants of lifelong health, social, and economic outcomes. The original ACE study demonstrated strong associations between early adversity and a range of adult disease risks [3]. Subsequent work has shown that exposure to multiple forms of adversity is common across populations [8]. Meta-analytic evidence further indicates that cumulative childhood adversity is strongly associated with both physical and mental health outcomes throughout the life course [6], with recent estimates suggesting that childhood maltreatment alone may account for a substantial fraction of global mental health burden [4].

Trauma may also propagate across generations. Intergenerational transmission of adversity has been documented across a range of social and developmental contexts [11, 16], and proposed mechanisms include behavioural pathways, socioeconomic constraints, and biological processes such as epigenetic regulation [22]. Developmental research further emphasises that early adversity can shape long-term stress responses and behavioural trajectories [17], while resilience and recovery processes can mitigate these effects under favourable conditions [12, 18].

Exposure to harsh or unpredictable environments has been linked to accelerated life-history strategies and earlier reproductive timing [2], and empirical studies have found associations between childhood adversity and fertility patterns [21]. Population prevalence is therefore shaped by the interplay between transmission, demographic processes, and recovery.

No theoretical framework yet links these mechanisms to the long-term population dynamics of trauma. Existing studies focus either on individual outcomes—documenting associations between ACE exposure and adult health risk [3], synthesising dose–response relationships [6], or characterising developmental consequences of early toxic stress [17]—or on statistical correlations across cohorts, such as ACE prevalence estimates [8] or intergenerational transmission rates [11]. Neither approach addresses how trauma prevalence evolves over long time horizons or responds to structural interventions. Population modelling has proven useful for linking demographic processes to long-run outcomes [1], but has not been applied to trauma dynamics.

We develop a minimal analytical model that captures the essential mechanisms required for trauma to persist or decline in a population. This model isolates the roles of differential reproductive outcomes, intergenerational transmission, and healing, and yields explicit threshold conditions for self-sustaining trauma prevalence. We complement this with a data-informed simulation model incorporating heterogeneous exposure levels and empirically grounded transmission probabilities, which we use to explore how realistic distributions of adversity shape long-run population outcomes.

Throughout this work *adversity state* refers to an individual's level of cumulative ACE exposure, not to any single event. Within a generation, individuals move between lower and higher adversity states through exposure or healing; across generations, adversity propagates via behavioural, socioeconomic, and biological pathways. These mechanisms are illustrated in Figure 1.




Figure 1: **Core model structure and simulation algorithm. (A)** Individuals occupy one of two adversity states, S (lower adversity) or T (higher adversity), shown for two successive generations. Within each generation, individuals transition from S to T with probability p (exposure) and from T to S with probability q (healing). Across generations, T-state individuals reproduce at rate αf_S relative to the baseline fertility f_S of S-state individuals ($\alpha > 1$), reflecting empirically observed associations between adversity and accelerated life-history strategies. Adversity propagates intergenerationally via behavioural, socioeconomic, and biological pathways. The product $\alpha(1-q)$ acts as a demographic analogue of R_0: endemic trauma persists whenever $\alpha(1-q) > 1$. **(B)** Flowchart of the individual-based simulation. At each annual step the three processes act sequentially on every member of the population denoted as $\mathcal{P}_t$. *Reproduction*: surviving females may give birth with probability $b(a, k)$, which combines a tempo effect, shifting the baseline schedule earlier by τ years per ACE via effective age $\tilde{a} = a + \tau \min(k, k_\tau)$, and a quantum effect, scaling the shifted probability by α for individuals with $k \geq 1$ ACEs. Each newborn is assigned a sex s' drawn from a Bernoulli with parameter p_m and an ACE count k' drawn from the empirical intergenerational conditional distribution $\Pr(k' \mid k)$. *ACE adjustment*: a surviving child ($a < 18$) acquires one additional ACE with probability p (trauma exposure); a surviving adult ($a \geq 18$) loses one ACE with probability q (healing). *Mortality*: each individual dies independently with the age- and sex-specific probability $\mu(s, a)$ taken from UN life tables; those who die are excluded from $\mathcal{P}_{t+1}$. All survivors (with age incremented by one and ACE count updated) and all newborns are collected into $\mathcal{P}_{t+1}$.

2 Methods

2.1 Analytical model

Individuals are assumed to be either traumatised or not, so the population state is $x = (x_T, x_S)$, where x_T denotes the proportion of traumatised individuals and $x_S = 1 - x_T$ the remainder.

Let f_T and f_S denote the fitness of the respective subpopulations. Consistent with evidence linking adversity to accelerated life-history strategies and elevated fertility, we set $f_T = \alpha f_S$ with $\alpha > 1$ [2, 21].

We allow both the introduction of new trauma and the possibility of recovery [18, 22], modelled through two parameters:

- p: probability that a non-traumatised individual becomes traumatised,
- q: probability that a traumatised individual recovers.

Together, α, p, and q determine the balance between transmission, recovery, and differential reproduction, and hence whether trauma prevalence grows, declines, or stabilises over time (Fig. 2A).




Figure 2: **Dynamics of trauma prevalence across mechanisms and parameters.** **A**, Time evolution of the traumatised proportion $x_T(t)$ under varying exposure (p) and healing (q) regimes. Trajectories from high and low initial conditions converge to the equilibrium level x^* (dotted line), illustrating how persistent exposure or recovery processes determine long-run prevalence. **B**, Analytical structure of the update map. Top: net growth function $\Delta(x_T) = x'_T - x_T$ for several values of the relative reproductive multiplier α, showing how equilibrium number and stability change with demographic amplification. Bottom: phase portrait for a representative parameter set, with arrows indicating the direction of population change. **C**, Parameter landscape for the invasion regime ($p = 0$). The coloured surface shows the endemic equilibrium level x^* as a function of α and healing probability q. The curve $q = 1 - \frac{1}{\alpha}$ separates parameter combinations for which trauma necessarily declines from those where it can persist endemically. **D**, Corresponding landscape with positive background exposure ($p = 0.05$). Introducing a non-zero exposure rate eliminates the trauma-free state, raising x^* across the entire parameter space and shifting the effective persistence boundary upward.

The population evolves according to the discrete replicator–mutation dynamic [7, 14],

$$x'_i = \sum_{j \in \{T,S\}} \frac{x_j f_j Q_{ji}}{\bar{f}}, \qquad i \in \{T, S\},$$

with mutation matrix

$$Q = \begin{pmatrix} 1-q & q \\ p & 1-p \end{pmatrix}.$$

The resulting update rule for the traumatised proportion is

$$x_T' = \frac{x_T f_T(1-q) + x_S f_S p}{\bar{f}}.$$

To assess persistence, consider the growth factor $\kappa_T = x_T'/x_T$. Substituting $f_T = \alpha f_S$ and $x_S = 1 - x_T$ yields

$$\kappa_T = \frac{1}{x_T} \frac{x_T\big(\alpha(1-q) - p\big) + p}{x_T(\alpha - 1) + 1}.$$

Values $\kappa_T > 1$ imply growth of the traumatised subpopulation, while $\kappa_T < 1$ imply decline, reflecting the standard growth criterion used in evolutionary dynamics [5, 14].

Interior equilibria satisfy $x_T' = x_T$, leading to the quadratic

$$(\alpha - 1)x_T^2 + (1 + p - \alpha + \alpha q)x_T - p = 0,$$

Because the constant term equals $-p < 0$ while evaluation at $x_T = 1$ gives $\alpha q \geq 0$, continuity guarantees a root in $(0, 1)$ whenever $p, q > 0$. This root defines the endemic level x_T^*, showing that any persistent introduction of trauma produces a stable long-run prevalence (Fig. 2A).

When no new trauma is introduced ($p = 0$), persistence depends on invasion from rarity. Taking the limit $x_T \to 0$ gives

$$\lim_{x_T \to 0} \kappa_T = \alpha(1 - q),$$

so trauma spreads from low prevalence precisely when

$$\alpha(1 - q) > 1 \qquad \Longleftrightarrow \qquad q < 1 - \frac{1}{\alpha}.$$

This condition acts as a demographic threshold separating extinction from endemic persistence, shown in Fig. 2C.

Long-run trauma prevalence is therefore governed by the balance between transmission, recovery, and demographic reinforcement. A sufficient reproductive advantage can sustain trauma across generations even without continual exposure, and recovery must cross a critical threshold to eliminate it. The system behaves, in this respect, like an infectious agent with a demographic amplification mechanism.

Figure 2B–C summarises the equilibrium structure. The next section embeds these mechanisms in a data-informed simulation that adds demographic stochasticity, age structure, and empirically grounded parameterisation.

3 Individual-based simulation model

To explore how stochasticity, finite-population effects, and demographic heterogeneity modify the analytical predictions, we built an individual-based simulation of the same mechanisms.

Individual state and population dynamics

Each individual is characterised by a triple (s, a, k) denoting sex $s \in \{\mathrm{M}, \mathrm{F}\}$, integer age $a \geq 0$, and ACE count $k \in \{0, 1, \ldots, 8\}$. The population at time t is the multiset $\mathcal{P}_t = \{(s_i, a_i, k_i)\}_{i=1}^{N_t}$, where N_t is the (stochastic) population size. The model evolves in discrete yearly steps; in each step three processes act sequentially on every individual: mortality, reproduction, and within-life ACE adjustment.

Mortality

Each individual (s, a, k) dies in year t with probability $\mu(s, a)$, drawn from age- and sex-specific life-table values based on UN World Population Prospects data [20]. Mortality is assumed independent of ACE count, so demographic selection operates exclusively through fertility.

Fertility and ACE transmission

Reproduction is restricted to surviving females. The baseline probability of a birth in a given year for a female of age a is $b_0(a)$, taken from UN age-specific fertility schedules [19]. Two empirically motivated mechanisms modify this baseline for ACE-exposed individuals.

Tempo effect (earlier reproductive timing). Consistent with evidence that adversity is associated with accelerated life-history strategies and earlier age at first birth [21], the fertility schedule is shifted earlier in proportion to ACE exposure. Specifically, a female with $k \geq 1$ ACEs evaluates her birth probability at an effective age

$$\tilde{a} = a + \tau \min(k, k_\tau),$$

where $\tau > 0$ is the tempo parameter (years of schedule shift per ACE). Evaluating at an older effective age brings the individual forward on the schedule, so that at real age a she faces the birth probability an unexposed individual faces at $a + \tau k$; her modal age at birth falls by τ years per ACE. Empirical estimates suggest $\tau \approx 3$ [21], which is used here. The shift is capped at $k_\tau = 3$ ACEs, because the source estimate reports its highest exposure category as 3+ and an uncapped shift would move the whole fertility window below the youngest age in the schedule. This implements a purely timing adjustment: completed fertility is unaffected in the absence of the quantum mechanism described below.

Quantum effect (differential reproductive intensity). Conditional on effective age $\tilde{a}$, the birth probability of a female with $k \geq 1$ ACEs is scaled by a multiplicative factor $\alpha > 1$:

$$b(a, k) = \begin{cases} b_0(\tilde{a}) & \text{if } k = 0, \\ \min\big(1,\, \alpha\, b_0(\tilde{a})\big) & \text{if } k \geq 1. \end{cases}$$

Under a coarse-grained binary classification (ACE-free versus ACE-exposed), this corresponds directly to the relative fitness parameter of the analytical model, in which $f_T = \alpha f_S$. The simulation therefore embeds the replicator structure of the analytical model within an age-structured, stochastic demographic framework.

Intergenerational ACE transmission. The ACE count of each newborn is drawn from an empirically estimated conditional distribution $\Pr(k_{\text{child}} \mid k_{\text{mother}})$, based on reported intergenerational ACE correlations [10]. Denoting the four maternal exposure categories as $\{0, 1, 2\text{–}3, \geq 4\}$, the conditional probabilities are summarised in Table 1. Offspring ACE counts in the range $\{2, 3\}$ and $\{4, \ldots, 8\}$ are distributed uniformly within their respective groups. This mechanism allows adversity to propagate across generations in the absence of ongoing environmental shocks, providing the simulation analogue of the transmission parameter p in the analytical model.

Table 1: Conditional probabilities $\Pr(k_{\text{child}} \mid k_{\text{mother}})$ for offspring ACE category given maternal ACE category, expressed as percentages with 95% confidence intervals. Source: [16].

	Maternal ACE category			
Child ACE category	0	1	2–3	≥ 4
0	42.6 (38.4–46.7)	37.8 (32.4–43.1)	30.4 (24.7–36.1)	25.1 (16.2–34.0)
1	25.8 (21.9–29.6)	29.7 (23.9–35.5)	28.4 (22.2–34.7)	23.2 (14.1–32.2)
2–3	25.8 (21.9–29.7)	21.5 (16.3–26.7)	21.3 (15.7–26.8)	23.8 (15.6–32.0)
≥ 4	5.8 (4.0–7.7)	11.0 (6.3–15.6)	19.9 (13.7–26.1)	27.9 (18.9–36.9)

Within-life ACE dynamics

ACE counts may change within a lifetime through two complementary processes that correspond directly to the analytical parameters p and q.

- *Trauma exposure* (p): a child aged $a < 18$ acquires one additional ACE with probability p per year, subject to the ceiling $k \leq 8$. This represents environmental and social transmission of adversity during the developmental window.
- *Healing* (q): an adult aged $a \geq 18$ reduces their ACE count by one with probability q per year, subject to the floor $k \geq 0$. This represents recovery processes including therapeutic intervention, social support, and self-forgiveness [13, 18].

The net annual change in ACE count for individual i is therefore

$$\Delta k_i = \begin{cases} +1 & \text{if } a_i < 18 \text{ and } U_i < p, \\ -1 & \text{if } a_i \geq 18 \text{ and } U_i < q, \\ 0 & \text{otherwise,} \end{cases}$$

where $U_i \sim \text{Uniform}(0, 1)$ is drawn independently each year. p and q play the same roles in the simulation as in the analytical model: p governs the rate of acquiring adversity and q the recovery rate.

Initial conditions

Simulations are initialised from a population pyramid based on UK Office for National Statistics census data [15], from which ages and sexes are sampled according to empirical proportions. Initial ACE counts are drawn from sex-stratified prevalence distributions reported in the Canadian Longitudinal Study on Aging [8].

Relationship to the analytical model

The simulation preserves the core dynamical structure—α, p, and q enter with the same meanings as in the replicator–mutation model—but replaces the mean-field assumption with explicit demographic stochasticity and age structure, and the binary ACE state with a discrete count on $\{0, \ldots, 8\}$. In the limit of large population size and binary aggregation of the ACE count, the simulation dynamics converge to the analytical predictions; departures reflect stochasticity and heterogeneity.

To identify the boundary between high- and low-adversity parameter regimes in the simulation output we applied k-means clustering ($k = 2$) to the log-transformed mean ACE surface; full details of this procedure are given in Supplementary Information .3.

Figure 3 illustrates the baseline demographic behaviour of the simulation across six parameter combinations ($q \in \{0.00, 0.01, 0.05\}$, $\alpha \in \{1.05, 1.35\}$, no background exposure): population size and mean age converge to stable levels within a few decades across all combinations (panels A–B), and the empirical survivorship curves closely match the theoretical life-table predictions irrespective of ACE parameters, confirming that mortality is ACE-independent (panel C). Population age pyramids at 50-year intervals (panels D–H) show the transition from the initial UK age structure to the long-run steady-state shape for a representative parameter set ($q = 0.05$, $\alpha = 1.35$, no background exposure).

4 Discussion

Our central result is a closed-form threshold, $q < 1 - 1/\alpha$, separating parameter regimes in which trauma necessarily declines from those in which it persists endemically. Any positive background exposure ($p > 0$) eliminates the trauma-free equilibrium entirely: the quadratic fixed-point equation always admits an interior root, so even a small rate of environmental adversity produces a stable long-run prevalence. The individual-based simulation corroborates these analytical predictions; demographic stochasticity and age structure shift the effective boundary but do not qualitatively alter the prediction.

The threshold condition $\alpha(1 - q) > 1$ is structurally analogous to the basic reproduction number R_0 in infectious-disease epidemiology [5], with the product $\alpha(1 - q)$ playing the role of R_0. Unlike standard SIS/SIR frameworks, this system contains a demographic amplification term: adversity is not merely transmitted but is reinforced by a reproductive advantage that standard epidemic models lack.

The parameter $\alpha > 1$ reflects observed associations between early adversity and accelerated life-history strategies [2, 21]. This does not imply that trauma is adaptive; it captures empirically documented tempo and quantum effects on reproductive timing and intensity. The tempo effect (earlier age at first birth) and the quantum effect (elevated birth rates conditional on effective age) make separable contributions to persistence, and the threshold condition provides a way to quantify their joint role. The empirical range of α in contemporary populations suggests that many settings may lie close to or above the persistence threshold.

The simulation broadly corroborates the analytical predictions. Without background exposure, mean ACE trajectories converge to moderate endemic levels from a range of initial conditions (Fig. 4A), consistent with the stable interior equilibrium. Positive background exposure raises the outcome substantially and widens variation across parameter combinations (Fig. 4B), confirming that even modest environmental transmission eliminates the trauma-free state. The summary panels (Fig. 4C–D) show a consistent ordering: higher α and lower q raise long-run adversity, with the effect amplified under positive background exposure. At $q = 0.05$, mean ACE counts are suppressed near zero at moderate α when $p = 0$, but only partially attenuated when $p > 0$. The heatmaps (Fig. 4E–F) show the transition is well approximated by a linear threshold $q^*(\alpha) = m\alpha + c$ ($R^2 \geq 0.99$). Fitted slopes are modest without background exposure ($m \approx 0.013$) but roughly four-fold steeper under positive background exposure ($m \approx 0.042$). The positive intercepts ($c \approx 0.025$–0.168) imply a minimum healing requirement even as $\alpha \to 1$, consistent with any $p > 0$ shifting the equilibrium landscape upward. Results for the proportion-traumatised metric are qualitatively identical (Fig. 5). The threshold separating suppression from persistence scales approximately linearly with α; background exposure raises it, so addressing transmission and healing separately is insufficient.

To drive the system below endemic persistence, q must exceed $1-1/\alpha$; any intervention that reduces α simultaneously lowers the required healing rate. Interventions combining socioeconomic support with expanded therapeutic access are therefore doubly effective, acting on both factors in $\alpha(1 - q)$. The empirical linear relationship $q^*(\alpha) = m\alpha + c$ implies that populations with higher α require proportionally more intensive recovery programmes.

Several limitations apply. Both models simplify adversity: the analytical model uses a binary state, the simulation a discrete count on $\{0, \ldots, 8\}$, and neither captures continuous severity or multi-domain ACE categorisations. Mortality is ACE-independent, though emerging evidence links high ACE burden to premature mortality, which would reduce




Figure 3: **Population-level demographic dynamics of the simulation.** Results shown for six parameter combinations ($q \in \{0.00, 0.01, 0.05\}$, $\alpha \in \{1.05, 1.35\}$, no background exposure) to demonstrate that the demographic dynamics are robust across the parameter space explored. **A**, Trajectories of total population size over 200 years (median ± IQR across replicates, one line per parameter combination). **B**, Mean age of the population over time, with the same summary. **C**, Empirical period survivorship $l(x) = N(x)/N(0)$ computed from simulation snapshots after year 100. One pair of median lines (male and female sharing the same colour and line style) is shown per parameter combination. The grey band shows the pooled inter-quartile range across all parameter combinations and replicates. Dark grey lines show the theoretical survivorship curves derived directly from the UN life-table probabilities (solid: male; dashed: female). The close agreement across all parameter combinations confirms that the simulated age structure is consistent with the embedded mortality schedule and is insensitive to the ACE transmission parameters, as expected given that mortality is modelled as ACE-independent. **D**–**H**, Population age pyramids (median counts across replicates) at years 0, 50, 100, 150, and 200 for the representative parameter set ($q = 0.05$, $\alpha = 1.35$, no background exposure), showing the transition from the initial UK census age structure to the long-run steady-state shape imposed by the empirical fertility and mortality schedules.




Figure 4: **Simulation outcomes across parameter regimes (mean ACE count).** Panels are arranged in three rows and two columns; the left column shows results without background trauma exposure ($p = 0$) and the right column with positive background exposure ($p = 0.2$). **A**–**B**, Time trajectories of mean ACEs per person for six combinations of healing probability q and reproductive multiplier α, shown as median ± IQR across simulation replicates. The dashed vertical line marks the end of the 100-year warm-up period. **C**–**D**, Mean ± SD of mean ACE count as a function of α, with one curve per healing probability. Higher α and lower q consistently raise long-run adversity levels, with the effect amplified under positive background exposure. **E**–**F**, Heatmaps of the time-averaged mean ACE count in the (q, α) parameter plane, shown on a $\log(1 + \cdot)$ scale. The dashed red contour marks the k-means threshold separating high- and low-adversity regimes; the linear fit $q^* = m\alpha + c$ (shown in each panel title) recovers a boundary consistent with the analytical condition $q > 1 - \alpha^{-1}$. Corresponding results for the proportion-traumatised metric are shown in Fig. 5.

the effective α and lower the persistence threshold. No spatial or network structure is included, so local clustering of adversity and peer transmission are not represented. Finally, α and τ were calibrated from UK and Canadian data; cross-cultural variation remains unexplored.

Extensions could include multi-domain ACE severity, spatial and network structure, and cross-national calibration of α. The threshold condition offers a direct basis for evaluating intervention cost-effectiveness. The framework gives population-level adversity science the same formal structure as epidemic theory: a minimal model with an explicit persistence threshold.

References

[1] Hal Caswell. *Matrix Population Models: Construction, Analysis, and Interpretation*. Sinauer Associates, 2001.

[2] Bruce J. Ellis, Aurelio J. Figueredo, Brad H. Brumbach, and Gabriel L. Schlomer. Fundamental dimensions of environmental risk: The impact of harsh versus unpredictable environments on the evolution and development of life history strategies. *Human Nature*, 20(2):204–268, 2009.

[3] Vincent J. Felitti, Robert F. Anda, Dale Nordenberg, David F. Williamson, Alison M. Spitz, Valerie Edwards, Mary P. Koss, and James S. Marks. Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults. *American Journal of Preventive Medicine*, 14(4):245–258, 1998.

[4] Lucinda Grummitt, Jessie R Baldwin, Johanna Lafoa'i, Katherine M Keyes, and Emma L Barrett. Burden of mental disorders and suicide attributable to childhood maltreatment. *JAMA psychiatry*, 2024.

[5] Josef Hofbauer and Karl Sigmund. *Evolutionary Games and Population Dynamics*. Cambridge University Press, 1998.

[6] Karen Hughes, Mark A. Bellis, Kate A. Hardcastle, Deepti Sethi, Alexander Butchart, Christopher Mikton, Lisa Jones, and Michael P. Dunne. The effect of multiple adverse childhood experiences on health: a systematic review and meta-analysis. *The Lancet Public Health*, 2(8):e356–e366, 2017.

[7] Lorens A. Imhof, Drew Fudenberg, and Martin A. Nowak. Evolutionary game dynamics in finite populations. *Proceedings of the National Academy of Sciences*, 102(31):10797–10800, 2005.

[8] Divya Joshi, Parminder Raina, Lil Tonmyr, Harriet L MacMillan, and Andrea Gonzalez. Prevalence of adverse childhood experiences among individuals aged 45 to 85 years: a cross-sectional analysis of the canadian longitudinal study on aging. *Canadian Medical Association Open Access Journal*, 9(1):E158–E166, 2021.

[9] Stuart P. Lloyd. Least squares quantization in PCM. *IEEE Transactions on Information Theory*, 28(2):129–137, 1982.

[10] Sheri Madigan, Chantal Cyr, Rachel Eirich, R. M. Pasco Fearon, Angela Ly, Christina Rash, Jason C. Poole, and Lenneke R. A. Alink. Testing the cycle of maltreatment hypothesis: Meta-analytic evidence of the intergenerational transmission of child maltreatment. *Development and Psychopathology*, 31(1):23–51, 2019.

[11] Sheri Madigan, Chantal Cyr, Rachel Eirich, RM Pasco Fearon, Anh Ly, Christina Rash, Julia C Poole, and Lenneke RA Alink. Testing the cycle of maltreatment hypothesis: Meta-analytic evidence of the intergenerational transmission of child maltreatment. *Development and psychopathology*, 31(1):23–51, 2019.

[12] Ann S. Masten. *Ordinary Magic: Resilience in Development*. Guilford Press, 2014.

[13] Ifigeneia Mavranezouli, Odette Megnin-Viggars, Caroline Daly, Sofia Dias, Nicky J. Welton, Sarah Stockton, Gita Bhutani, Nick Grey, Jane Leach, Neil Greenberg, Cornelius Katona, and Stephen Pilling. Research review: Psychological and psychosocial treatments for children and young people with post-traumatic stress disorder: a network meta-analysis. *Journal of Child Psychology and Psychiatry*, 61(1):18–29, 2020.

[14] Martin A. Nowak. *Evolutionary Dynamics: Exploring the Equations of Life*. Harvard University Press, Cambridge, MA, 2006.

[15] Office for National Statistics. Population estimates for the UK, England, Wales, Scotland and Northern Ireland: mid-2021. Technical report, Office for National Statistics, 2022.

[16] Adam Schickedanz, José J Escarce, Neal Halfon, Narayan Sastry, and Paul J Chung. Intergenerational associations between parents' and children's adverse childhood experience scores. *Children*, 8(9):747, 2021.

[17] Jack P. Shonkoff, Andrew S. Garner, et al. The lifelong effects of early childhood adversity and toxic stress. *Pediatrics*, 129(1):e232–e246, 2012.

[18] Victoria G Skolnick, Brian A Lynch, Latasha Smith, Magdalena Romanowicz, Gretchen Blain, and Loren Toussaint. The association between parent and child aces is buffered by forgiveness of others and self-forgiveness. *Journal of Child & Adolescent Trauma*, 16(4):995–1003, 2023.

[19] United Nations, Department of Economic and Social Affairs, Population Division. World population prospects 2022: Fertility data. Technical report, United Nations, 2022.

[20] United Nations, Department of Economic and Social Affairs, Population Division. World population prospects 2022: Mortality data. Technical report, United Nations, 2022.

[21] Kristi Williams and Brian Karl Finch. Adverse childhood experiences, early and nonmarital fertility, and women's health at midlife. *Journal of health and social behavior*, 60(3):309–325, 2019.

[22] Rachel Yehuda and Amy Lehrner. Intergenerational transmission of trauma effects: Putative role of epigenetic mechanisms. *World Psychiatry*, 15(3):243–257, 2016.

Supplementary Information

.1 Simulation outcomes: proportion-traumatised metric




Figure 5: **Simulation outcomes across parameter regimes (proportion traumatised).** Layout mirrors Fig. 4. Left column: no background trauma exposure ($p = 0$); right column: positive background exposure ($p = 0.2$). **A**–**B**, Time trajectories of the proportion of individuals with at least one ACE for the same six parameter combinations as Fig. 4, shown as median ± IQR. **C**–**D**, Mean ± SD of proportion traumatised as a function of α, one curve per healing probability. **E**–**F**, Heatmaps of the time-averaged proportion traumatised in the (q, α) parameter plane ($\log(1+\cdot)$ scale). The dashed red contour marks the k-means threshold; the linear fit $q^* = m\alpha + c$ is consistent with the mean-ACE results in Fig. 4E–F.

.2 k-means threshold across all values of p

.3 k-means threshold procedure

To identify the boundary between parameter regimes that sustain high and low levels of adversity, we applied k-means clustering with $k = 2$ [9] to the log-transformed mean ACE surface. Let $\bar{k}(q, \alpha)$ denote the mean ACE count averaged across all years and simulation replicates at healing probability q and reproductive multiplier α, and let $\ell(q, \alpha) = \log\big(1 + \bar{k}(q, \alpha)\big)$ be its $\log(1 + \cdot)$ transform, applied to compress the dynamic range of the surface and to handle zero values stably. k-means partitions the observed values of ℓ into two clusters by iteratively assigning each value to its nearest cluster centre and updating centres to cluster means until convergence (initialised at the 25th and 75th percentiles of ℓ). The scalar threshold ℓ^* is the midpoint between the two converged cluster centres, and requires




Figure 6: ***k*-means threshold contours across all values of *p*.** Each curve shows $q^*(\alpha)$ for a single value of the background exposure probability p (identified by colour, linestyle, and marker; see legend). The threshold is computed for each p separately by k-means ($k = 2$) on the $\log(1 + \cdot)$-transformed mean ACE surface and located by linear interpolation along the q-axis. Higher p shifts the threshold upward.

no user-specified cutoff. The threshold defines a level set $\{(q, \alpha) : \ell(q, \alpha) = \ell^*\}$ in the original parameter space, which we locate for each value of α by linear interpolation along the q-axis to obtain crossing points $q^*(\alpha)$. Although the threshold is identified in log-space, the resulting boundary is expressed in the untransformed coordinates (q, α), and is well described by a linear relationship

$$q^*(\alpha) = m\alpha + c,$$

recovered by ordinary least squares regression on the interpolated crossing points (fits shown in panel titles; $R^2 > 0.95$ in all cases). This linear structure implies that the critical healing probability required to suppress endemic adversity scales proportionally with the reproductive multiplier α, consistent with the analytical threshold condition $q > 1 - \alpha^{-1}$ derived in Section 2 (Fig. 4). Fig. 6 shows the resulting threshold contours for every value of p in the data.