

# VACCINATION CHECKLIST ASPLENIA & SICKLE CELL DISEASE

PROTECT EARLY – PROTECT FOR LIFE

Patient Name: \_\_\_\_\_

Date of Birth: \_\_\_\_\_ Sex: ☐ M ☐ F

Diagnosis: ☐ Asplenia ☐ Sickle Cell Disease

Planned Splenectomy Date (if any): \_\_\_\_ / \_\_\_\_ / \_\_\_\_

Allergies / Notes: \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

## AT DIAGNOSIS OR AS EARLY AS POSSIBLE

| Vaccine                        | Purpose                       | Given Date         | Due Date           | Given                    | Notes |
|--------------------------------|-------------------------------|--------------------|--------------------|--------------------------|-------|
| PCV*                           | No. of doses according to age | ____ / ____ / ____ | ____ / ____ / ____ | <input type="checkbox"/> |       |
| Hib                            | 1 dose                        | ____ / ____ / ____ | ____ / ____ / ____ | <input type="checkbox"/> |       |
| MenACWY (1 <sup>st</sup> dose) | First dose                    | ____ / ____ / ____ | ____ / ____ / ____ | <input type="checkbox"/> |       |
| MenB (1 <sup>st</sup> dose)    | First dose                    | ____ / ____ / ____ | ____ / ____ / ____ | <input type="checkbox"/> |       |

\*PCV (No. of doses according to age):

- Infants & children <2 years: Continue routine infant schedule. Give age-appropriate PCV series.
- 24 through 71 months: 2 doses of PCV20 or PCV15, or PCV13 given at least 8 weeks apart.
- From age of 71 months onward: one dose.

## AFTER 2 MONTHS (± 2 WEEKS)

| Vaccine                        | Purpose                    | Given Date         | Due Date           | Given                    | Notes |
|--------------------------------|----------------------------|--------------------|--------------------|--------------------------|-------|
| MenACWY (2 <sup>nd</sup> dose) | Second dose                | ____ / ____ / ____ | ____ / ____ / ____ | <input type="checkbox"/> |       |
| MenB (2 <sup>nd</sup> dose)    | Second dose                | ____ / ____ / ____ | ____ / ____ / ____ | <input type="checkbox"/> |       |
| PPSV23*                        | If PCV13 or PCV15 was used | ____ / ____ / ____ | ____ / ____ / ____ | <input type="checkbox"/> |       |

\*Only if PCV13 or PCV15 was used. (No PPSV23 needed if PCV20 was used)

## AFTER 4 MONTHS (≥ 6 MONTHS FROM FIRST VISIT)

| Vaccine                     | Purpose    | Given Date         | Due Date           | Given                    | Notes |
|-----------------------------|------------|--------------------|--------------------|--------------------------|-------|
| MenB (3 <sup>rd</sup> dose) | Third dose | ____ / ____ / ____ | ____ / ____ / ____ | <input type="checkbox"/> |       |

## BOOSTER DOSES

### MENACWY BOOSTERS

If the second primary dose is given:

- Before age 7 years:
  - First booster 3 years later, then every 5 years.
- At age ≥7 years:
  - First booster 5 years later, then every 5 years.



### MENB BOOSTERS

- First booster: 1 year after completion of primary series.
- Subsequent boosters: every 2–3 years while risk remains.



| Booster                 | Due Date           | Given Date         | Notes | Booster                 | Due Date           | Given Date         | Notes |
|-------------------------|--------------------|--------------------|-------|-------------------------|--------------------|--------------------|-------|
| 1 <sup>st</sup> booster | ____ / ____ / ____ | ____ / ____ / ____ |       | 1 <sup>st</sup> booster | ____ / ____ / ____ | ____ / ____ / ____ |       |
| Next booster            | ____ / ____ / ____ | ____ / ____ / ____ |       | Next booster            | ____ / ____ / ____ | ____ / ____ / ____ |       |
| Next booster            | ____ / ____ / ____ | ____ / ____ / ____ |       | Next booster            | ____ / ____ / ____ | ____ / ____ / ____ |       |
| Next booster            | ____ / ____ / ____ | ____ / ____ / ____ |       | Next booster            | ____ / ____ / ____ | ____ / ____ / ____ |       |

## PNEUMOCOCCAL (PCV) GUIDANCE



If PCV20 is used:  
No PPSV23 or booster dose is needed.



If PCV15 is used:  
Give PPSV23 (see 2nd visit).  
No further PPSV23 is recommended.

OR



If previously received PCV13 → PPSV23:  
Give PCV20 or PPSV23  
≥ 5 years after PPSV23.

1

## FIRST VISIT

(ASAP / PRE-SPLENECTOMY)



2

## SECOND VISIT

AFTER 2 MONTHS



3

## THIRD VISIT

AFTER 4 MONTHS  
(≥ 6 MONTHS FROM FIRST VISIT)



## IMPORTANT REMINDERS

- Whenever feasible, give indicated vaccines at least 14 days before planned splenectomy.
- Doses given within 14 days before surgery can still be counted as valid.
- If vaccination cannot be completed before surgery, vaccinate as soon as the patient is clinically stable after surgery.

NEXT REVIEW DATE:

\_\_\_\_ / \_\_\_\_ / \_\_\_\_