

VACCINATION CHECKLIST ASPLENIA & SICKLE CELL DISEASE

PROTECT EARLY – PROTECT FOR LIFE

Patient Name: _____

Date of Birth: _____ Sex: ☐ M ☐ F

Diagnosis: ☐ Asplenia ☐ Sickle Cell Disease

Planned Splenectomy Date (if any): ____ / ____ / ____

Allergies / Notes: _____

1

FIRST VISIT

(ASAP /

PRE-SPLENECTOMY)



AT DIAGNOSIS OR AS EARLY AS POSSIBLE

Vaccine	Purpose	Given Date	Due Date	Given	Notes
PCV*	No. of doses according to age	____ / ____ / ____	____ / ____ / ____	☐	
Hib	1 dose	____ / ____ / ____	____ / ____ / ____	☐	
MenACWY (1 st dose)	First dose	____ / ____ / ____	____ / ____ / ____	☐	
MenB (1 st dose)	First dose	____ / ____ / ____	____ / ____ / ____	☐	

*PCV (No. of doses according to age):

- Infants & children <2 years: Continue routine infant schedule. Give age-appropriate PCV series.
- 24 through 71 months: 2 doses of PCV20 or PCV15, or PCV13 given at least 8 weeks apart.
- From age of 71 months onward: one dose.

MenB / Bexsero (primary series options):

- Preferred schedule: 0, 1–2, 6 months.
- Alternative valid schedule: 0 and 6 months.
- If dose 2 is given at least 6 months after dose 1, a third primary dose is not needed.

2

SECOND VISIT

AFTER

2 MONTHS



AFTER 2 MONTHS

Vaccine	Purpose	Given Date	Due Date	Given	Notes
MenACWY (2 nd dose)	Second dose	____ / ____ / ____	____ / ____ / ____	☐	
MenB (3 rd dose)	Second dose	____ / ____ / ____	____ / ____ / ____	☐	
PPSV23*	If PCV13 or PCV15 was used	____ / ____ / ____	____ / ____ / ____	☐	

*Only if PCV13 or PCV15 was used. (No PPSV23 needed if PCV20 was used)

3

THIRD VISIT

AFTER

4 MONTHS

(≥6 MONTHS FROM FIRST VISIT)



AFTER 4 MONTHS (≥ 6 MONTHS FROM FIRST VISIT)

Vaccine	Purpose	Given Date	Due Date	Given	Notes
MenB (3 rd dose, if using 0, 1–2, 6 month schedule)	Third dose (if indicated)	____ / ____ / ____	____ / ____ / ____	☐	

BOOSTER DOSES

MENACWY BOOSTERS				MENB BOOSTERS			
If the second primary dose is given:							
• Before age 7 years: – First booster 3 years later, then every 3 years. • At age 27 years: – First booster 5 years later, then every 5 years.				• First booster: 1 year after completion of primary series. • Subsequent boosters: every 2–3 years while risk remains.			
Booster	Due Date	Given Date	Notes	Booster	Due Date	Given Date	Notes
1 st booster	____ / ____ / ____	____ / ____ / ____		1 st booster	____ / ____ / ____	____ / ____ / ____	
Next booster	____ / ____ / ____	____ / ____ / ____		Next booster	____ / ____ / ____	____ / ____ / ____	
Next booster	____ / ____ / ____	____ / ____ / ____		Next booster	____ / ____ / ____	____ / ____ / ____	
Next booster	____ / ____ / ____	____ / ____ / ____		Next booster	____ / ____ / ____	____ / ____ / ____	

PNEUMOCOCCAL (PCV) GUIDANCE



If PCV20 is used:
No PPSV23 or booster dose is needed.

OR



If PCV13 is used:
Give PPSV23 (see 2nd visit).
No further PPSV23 is recommended.

OR



If previously received PCV13 → PPSV23:
Give PCV20 or PPSV23 ≥5 years after PPSV23.



IMPORTANT REMINDERS

- Whenever feasible, give indicated vaccines at least 14 days before planned splenectomy.
- Doses given within 14 days before surgery can still be counted as valid.
- If vaccination cannot be completed before surgery, vaccinate as soon as the patient is clinically stable after surgery.

NEXT REVIEW DATE:

____ / ____ / ____