

VACCINATION CHECKLIST AFTER CHEMOTHERAPY

REBUILD PROTECTION AFTER IMMUNE RECOVERY

Patient Name: _____
Date of Birth: ____ / ____ / ____ Sex: ☐ M ☐ F
Diagnosis: _____
Last Chemotherapy Date: ____ / ____ / ____

Allergies / Notes: _____

1
FIRST VISIT
3-6 MONTHS
AFTER LAST
CHEMOTHERAPY

INACTIVATED VACCINES

Vaccine	Purpose	Given Date	Due Date	Given	Notes
PCV*	No. of doses according to age	___ / ___ / ___	___ / ___ / ___	☐	
Hexa*	Complete / continue age-appropriate series	___ / ___ / ___	___ / ___ / ___	☐	
Hepatitis A	Start standard schedule	___ / ___ / ___	___ / ___ / ___	☐	
Twinrix†	Start standard schedule	___ / ___ / ___	___ / ___ / ___	☐	
MenACWY (1st dose)	First dose	___ / ___ / ___	___ / ___ / ___	☐	
MenB (1st dose)	First dose	___ / ___ / ___	___ / ___ / ___	☐	

***PCV (No. of doses according to age):**

- **Infants & children < 2 years:** Continue routine infant schedule. Give age-appropriate PCV series.
- **24 through 71 months:** 2 doses of PCV20 or PCV15, or PCV15, at least 8 weeks apart.
- **From age 71 months onward:** one dose.

***Hexa:** complete the age-appropriate primary or catch-up series after chemotherapy if incomplete before treatment.

†Hepatitis A: 3-dose schedule (0, 6 months). **Twinrix** (from 16 years old): 3-dose schedule (0, 1, 6 months).

Check anti-HBs titer 2 months after the last hepatitis B-containing vaccine dose.

†MenB (Bexsero) primary series options:

- **Preferred schedule:** 0, 1-2, 6 months.
- **Alternative valid schedule:** 0 and 6 months.
- If dose 2 is given at least 5 months after dose 1, a third primary dose is not needed.

2
SECOND VISIT
AFTER
2 MONTHS

FOLLOW-UP

Vaccine	Purpose	Given Date	Due Date	Given	Notes
PPSV23*	If PCV13 or PCV15 was used	___ / ___ / ___	___ / ___ / ___	☐	
MenACWY (2nd dose)	Second dose	___ / ___ / ___	___ / ___ / ___	☐	
MenB (2nd dose)	Second dose	___ / ___ / ___	___ / ___ / ___	☐	
Hexa / other series	Continue if indicated	___ / ___ / ___	___ / ___ / ___	☐	

*No PPSV23 needed if PCV20 was used.

3
THIRD VISIT
AFTER
4 MONTHS
(≥ 6 MONTHS
FROM FIRST
VISIT)

IF INDICATED

Vaccine	Purpose	Given Date	Due Date	Given	Notes
MenB (3rd dose)	If using 0, 1-2, 2 month Bexsero schedule	___ / ___ / ___	___ / ___ / ___	☐	
Standard series	Continue as needed	___ / ___ / ___	___ / ___ / ___	☐	

LIVE VACCINES

START 12 MONTHS AFTER LAST CHEMOTHERAPY

Vaccine	Schedule	Given Date	Due Date	Given	Notes
MMR	1 dose, or 2 doses 1 month apart if not previously taken	___ / ___ / ___	___ / ___ / ___	☐	
Varicella	2 doses, 1-3 months apart	___ / ___ / ___	___ / ___ / ___	☐	

BOOSTER DOSES — HIGH RISK ONLY

MENACWY BOOSTERS (HIGH RISK ONLY)

- If the second primary dose is given before age 7 years: first booster 3 years later, then every 5 years.
- If given at age > 7 years: first booster 5 years later, then every 5 years.

MENB BOOSTERS (HIGH RISK ONLY)

- First booster 1 year after completion of primary series.
- Subsequent boosters every 2-3 years within risk remains.

PNEUMOCOCCAL GUIDANCE

- **IF PCV20 is used:**
No PPSV23 is needed.
- **IF PCV13 or PCV15 is used:**
Give PPSV23 after 2 months.

IMPORTANT REMINDERS

- Inactivated vaccines: 3-6 months after chemotherapy.
- Live vaccines: 12 months after chemotherapy.
- If series is incomplete before chemotherapy, complete it after therapy — do not restart.
- MenACWY / MenB boosters and extra doses for high-risk patients.
- Check anti-HBs titer 2 months after the last HepB containing vaccine dose.

NEXT REVIEW DATE: ____ / ____ / ____