



VACCINATION AFTER CHEMOTHERAPY

Restart protection after immune recovery



Vaccine talks
Egyptian edition

WHEN TO VACCINATE



Inactivated vaccines:
start **3–6 months** after
the last chemotherapy dose.




Live vaccines:
start **12 months** after
the last chemotherapy dose.



If a vaccine series was
incomplete before chemotherapy,
**complete it after therapy —
do not restart.**

VACCINATION SCHEDULE AFTER CHEMOTHERAPY




PCV: NUMBER OF DOSES ACCORDING TO AGE*

AGE	PCV RECOMMENDATION	DURATION / INTERVAL
 Infants and children < 2 years	- Continue routine infant schedule. - Give age-appropriate PCV series (PCV15, PCV15 or PCV20).	According to routine schedule.
 24 through 71 months (2 through < 6 years)	2 doses of PCV20 or PCV15 or PCV13	Given at least 8 weeks apart.
 From age 71 months onward (≥ 6 years)	One dose of PCV20 or PCV15 or PCV13	Single dose.

*Use any available PCV (PCV20, PCV15 or PCV13) according to age recommendations.

PCV: IMPORTANT POINTS

- ✓ **If PCV20 is used:**
No PPSV23 is needed. 
- ✓ **If PCV13 or PCV15 is used:**
Give PPSV23 after 2 months.  or  → 
- ✓ Complete interrupted pneumococcal series after chemotherapy. 

LIVE VACCINES (START 12 MONTHS AFTER CHEMOTHERAPY)



- ✓ **MMR:** 1 dose if one dose is needed, or 2 doses 3 month apart if not previously taken.
- ✓ **Varicella:** 2 doses, 1–3 months apart.

HEPATITIS A / TWINRIX



- ✓ **Hepatitis A:** 2-dose schedule (0, 6 months).
- ✓ **Twinrix (from 16 years old):** 3 dose schedule (0, 1, 6 months).
- ✓ **Check anti-HBs titre** 2 months after the last hepatitis B-containing vaccine dose.

MENINGOCOCCAL VACCINES

Boosters / extra doses in high-risk patients



- ✓ **MenACWY primary series:** 2 doses, 2 months apart.
- ✓ **MenB / Bexsero primary series:** preferred 0, 1–2, 6 months; alternative valid schedule 0 and 6 months. If dose 2 is given at least 6 months after dose 1, a third primary dose is not needed.
- ✓ **MenACWY boosters:** for high-risk patients only — if the second primary dose is given before age 7 years, first booster 3 years later then every 5 years, if given at age 7 years or older, first booster 3 years later then every 5 years.
- ✓ **MenB boosters:** for high-risk patients only — first booster 1 year after completion of the primary series, then every 2–3 years while risk remains.

KEY REMINDERS



Non-live vaccines:
3–6 months after
chemotherapy.



Live vaccines:
12 months after
chemotherapy.



Complete
interrupted series
after therapy.



Check anti-HBs titre
2 months after the last
HepB-containing
vaccine dose.



Document vaccines
and follow-up
dates.