

VACCINATION CHECKLIST AFTER HSCT

REBUILD PROTECTION AFTER IMMUNE RECOVERY

Patient Name: _____
 Date of Birth: ____ / ____ / ____ Sex: ☐ M ☐ F
 Diagnosis: _____
 HSCT Date: ____ / ____ / ____

Allergies / Notes: _____

INACTIVATED VACCINES

Vaccine	Purpose	Given Date	Due Date	Given	Notes
PCV (1st dose)	Start primary series	___ / ___ / ___	___ / ___ / ___	☐	
Hexa (1st dose)	Start 0, 2, 4 month series	___ / ___ / ___	___ / ___ / ___	☐	
Hepatitis A	Start standard schedule	___ / ___ / ___	___ / ___ / ___	☐	
Twinrix	Start standard schedule	___ / ___ / ___	___ / ___ / ___	☐	
MenACWY (1st dose)	First dose	___ / ___ / ___	___ / ___ / ___	☐	
MenB (1st dose)	First dose	___ / ___ / ___	___ / ___ / ___	☐	

- PCV primary series after HSCT: 3 doses, 1–2 months apart.
- If PCV20 is used: give a 4th PCV dose 6 months after dose 3.
- If PCV13 or PCV15 is used and there is no GVHD: give PPSV23 2 months after dose 3.
- If GVHD is present, give a 4th PCV dose 6 months after dose 3 instead of PPSV23.
- PPSV23 may be repeated 5 years after the last PPSV23 if PCV13 was used.
- Hexa: 3, 2, 4 months.
- Hepatitis A: 2-dose schedule (0, 6 months).
- Twinrix (from 16 years after 3-dose schedule (0, 1, 6 months).
- Check and HBs IIRs 2 months after the last HepB-containing vaccine dose.
- MenB (Bexsero) primary series options: preferred 0, 1–2, 6 months.
 Alternative valid schedule: 0 and 6 months. If dose 2 is given at least 5 months after dose 1, a 3rd primary dose is not needed.

FOLLOW-UP

Vaccine	Purpose	Given Date	Due Date	Given	Notes
PCV (2nd dose)	Continue primary series	___ / ___ / ___	___ / ___ / ___	☐	
Hexa (2nd dose)	Continue 0, 2, 4 month series	___ / ___ / ___	___ / ___ / ___	☐	
MenACWY (1st dose)	Second dose	___ / ___ / ___	___ / ___ / ___	☐	
MenB (2nd dose)	Second dose	___ / ___ / ___	___ / ___ / ___	☐	

FOLLOW-UP

Vaccine	Purpose	Given Date	Due Date	Given	Notes
PCV (2nd dose)	Complete primary PCV series	___ / ___ / ___	___ / ___ / ___	☐	
Hexa (2nd dose)	Complete 0, 2, 4 month series	___ / ___ / ___	___ / ___ / ___	☐	
Standard series	Continue as needed	___ / ___ / ___	___ / ___ / ___	☐	

PCV FOLLOW-UP / IF INDICATED

Vaccine	Purpose	Given Date	Due Date	Given	Notes
PPSV23	If PCV13 or PCV15 was used and no GVHD	___ / ___ / ___	___ / ___ / ___	☐	
PCV (4th dose)	If PCV23 was used, or if GVHD is present	___ / ___ / ___	___ / ___ / ___	☐	
Repeat PPSV23	5 years after last PPSV23 if PCV13 was used	___ / ___ / ___	___ / ___ / ___	☐	

LIVE VACCINES START 24 MONTHS AFTER HSCT

Vaccine	Purpose	Given Date	Due Date	Given	Notes
MMR	2 doses, 1 month apart	___ / ___ / ___	___ / ___ / ___	☐	
Varicella	With doctor recommendation	___ / ___ / ___	___ / ___ / ___	☐	

BOOSTER DOSES — IF RISK REMAINS

- | MENACWY BOOSTERS | MENB BOOSTERS |
|--|---|
| <ul style="list-style-type: none"> • If the second primary dose is given before age 7 years; first booster 3 years later, then every 5 years. • If given at age 7 years or older: first booster 5 years later, then every 5 years. | <ul style="list-style-type: none"> • First booster 1 year after completion of primary series. • Subsequent boosters every 2–3 years while risk remains. |

PNEUMOCOCCAL GUIDANCE

- PCV primary series: 3 doses, 1–2 months apart.
- PCV20: 4th dose 6 months after dose 3.
- PCV13 or PCV15 with no GVHD: PPSV23 after 2 months.
- GVHD: give a 4th PCV dose instead of PPSV23.
- PPSV23 may be repeated 5 years later if PCV13 was used.

IMPORTANT REMINDERS

- Inactivated vaccines: 6–12 months after HSCT.
- Live vaccines: 24 months after HSCT.
- Hexa schedule: 0, 2, 4 months.
- Check anti-HBs titre 2 months after the last HepB-containing vaccine dose.
- Varicella only with doctor recommendation.

NEXT REVIEW DATE: ____ / ____ / ____

1
FIRST VISIT
6–12 MONTHS
AFTER HSCT

2
SECOND VISIT
AFTER
2 MONTHS

3
THIRD VISIT
AFTER
4 MONTHS