

Vaccine talks

Egyptian edition



VACCINATION AFTER HSCT

Restart protection after immune reconstitution



WHEN TO VACCINATE



Inactivated vaccines:
start **6–12 months**
after HSCT.



Live vaccines:
start **24 months**
after HSCT.



PCV follow-up depends
on the PCV product used
and GVHD status.

VACCINATION SCHEDULE AFTER HSCT



FIRST VISIT
6–12 months
after HSCT



SECOND VISIT
After 2 months



THIRD VISIT
After 4 months

- ✓ PCV (1st dose)
- ✓ Hexa (1st dose)
- ✓ Hepatitis A
- ✓ Twinrix (from 15 years old)
- ✓ MenACWY (1st dose)
- ✓ MenB (1st dose)


- ✓ PCV (2nd dose)
- ✓ Hexa (2nd dose)
- ✓ MenACWY (2nd dose)
- ✓ MenB (2nd dose)

- ✓ PCV (3rd dose)
- ✓ Hexa (3rd dose)
- ✓ Continue standard schedules as needed


PCV SCHEDULE AFTER HSCT



Primary PCV series:
3 doses,
1–2 months apart.



If PCV20 is used:
give a 4th PCV20 dose
6 months after
dose 3.



**If PCV13 or PCV15 is used
and there is NO GVHD:**
give PPSV23 2 months
after dose 3.



If GVHD is present:
give a 4th PCV dose
6 months after dose 3
instead of PPSV23.




PPSV23 may be repeated
5 years after the last
PPSV23 if PCV13
was used.


PCV: IMPORTANT POINTS




PCV20:
4th dose 6 months
after dose 3.



PCV13 or PCV15 with no GVHD:
PPSV23 after 2 months.



If GVHD is present:
give a 4th PCV dose 6 months
after dose 2, PPSV23 may be
repeated 5 years later if PCV13
was used.



LIVE VACCINES (START 24 MONTHS AFTER HSCT)



- ✓ **MMR:** 2 doses, 1 month apart.
- ✓ **Varicella:** only with doctor recommendation.

HEPATITIS A / TWINRIX



- ✓ **Hepatitis A:** 2-dose schedule (0, 6 months).
- ✓ **Twinrix** (from 16 years old); 3-dose schedule (0, 1, 6 months).
- ✓ **Check anti-HBs titrs** 2 months after the last HepB-containing vaccine dose.

HEXA VACCINE



Hexa schedule: 0, 2, 4 months.

MENINGOCOCCAL VACCINES

Boosters if risk remains



- ✓ **MenACWY primary series:** 2 doses, 2 months apart.
- ✓ **MenB / Bexsere primary series:** preferred 0, 1–2, 6 months, alternative volul schedule 0 and 6 months. If dose 2 is given at least 6 months after dose 1, a 3rd primary does is not needed.
- ✓ **MenACWY boosters:** if the second primary dose is given before age 7 years, first booster 3 years later then every 5 years, if given at age 7 years of older, first booster 3 years later then every 5 years.
- ✓ **MenB boosters:** first booster 1 year after completion of the primary series, then every 2–3 years while risk remains.



KEY REMINDERS



Inactivated vaccines:
6–12 months
after HSCT.



Live vaccines:
24 months
after HSCT.



Hexa schedule:
0, 2, 4 months.



PCV follow-up
depends on PCV
product and GVHD.



Check anti-HBs titra
2 months after the last
HepB-containing
vaccine dose.



Document vaccines
and follow-up
dates.