

Reflection structure against M, $\phi_* = 5\,M_P$
$\beta = 1.2$
$\beta = 2$
$\beta = 2.9$
kinetic fraction $\pi^2/6$
parks
ρ_χ/ρ_R (dissipation)
T_J [GeV]
$\log_{10}\mathcal{R}$, first impact
elastic: no particle production
M/M_P
$M = 4 \times 10^{-31}$ (laboratory) M_P, dissipation ($\alpha = 0.08$ in row 1)
$M = 10^{-15}\,M_P$, dissipation ($\alpha = 0.08$ in row 1)
$M = 10^{-12}\,M_P$, dissipation ($\alpha = 0.08$ in row 1)
$M = 10^{-3}\,M_P$, natural mode ($\alpha = 0$)
$M = 0.5\,M_P$, natural mode ($\alpha = 0$)
$\alpha = 0.001$ (row 2)
$\alpha = 0.01$ (row 2)
$\alpha = 0.08$ (row 2)
$\alpha = 0.15$ (row 2)
$M = 10^{-9}\,M_P$, dissipation, $\alpha = 0.08$ (row 2)
$M = 10^{-6}\,M_P$, dissipation, $\alpha = 0.08$ (row 2)
$M = 10^{-3}\,M_P$, dissipation, $\alpha = 0.08$ (row 2)
$M = 0.5\,M_P$, dissipation, $\alpha = 0.08$ (row 2)
BBN, 2 MeV – 20 keV (indicative)
$\mathcal{R}$ at the first impact, main store (row 3)
$\mathcal{R}$ at the first impact, large-M diagnostic (row 3)
briefing notes' $M_{\rm crit}$, median ($1.2 \times 10^{-15}\,M_P$)
briefing notes' $M_{\rm crit}$, maximum ($8.2 \times 10^{-13}\,M_P$)