

My Capstone Project

Benedict Addae-Mensah

2024-01-16

This hydrogeological study represents a meticulous exploration of collected data through an extensive analytical phase employing advanced statistical methods. The utilization of regression analysis has enabled the modeling of intricate relationships, providing quantitative insights into phenomena such as the impact of borehole depth on specific capacity and the correlation between transmissivity and pumping rates. Correlation analyses have further revealed associations among various hydrogeological parameters, shedding light on the influence of geological formations on transmissivity and the interdependence between drawdown and pumping rates. Additionally, time series analysis and trend analyses have been instrumental in uncovering temporal trends in groundwater levels and drawdown, facilitating a comprehensive understanding of how aquifers respond to seasonal changes. This project's findings contribute significantly to the broader understanding of hydrogeological dynamics, offering valuable insights for sustainable water resource management and informed decision-making in the realm of groundwater exploration and utilization.

1. Introduction

1.1. Background

Groundwater, a vital component of Earth's hydrological cycle, plays a pivotal role in sustaining human societies, particularly in regions where surface water sources are scarce or unreliable. The study focuses on the Greater Accra and Ashanti regions, where systematic borehole data collection provides a unique opportunity to delve into the hydrogeological complexities shaping water availability. The significance of groundwater in supporting domestic, agricultural, and industrial needs underscores the importance of understanding its dynamics for informed decision-making.

1.2. Problem Statement

The Greater Accra and Ashanti regions face a multifaceted challenge in sustainable water resource management characterized by an inadequate understanding of aquifer characteristics, limited insight into temporal variations in groundwater levels, an unexplored influence of aquifer material on hydrogeological parameters, a lack of evidence-based decision support, and persistent challenges in community water access. This collective deficiency underscores a critical gap in knowledge, hindering effective policies and strategies for preserving and managing groundwater resources. Consequently, a comprehensive study is imperative to address these knowledge gaps and provide essential insights for informed decision-making, ensuring the sustainable utilization and conservation of groundwater in these regions.

1.3. Objectives

- To systematically analyze borehole data to comprehensively understand the characteristics of aquifers in the Greater Accra and Ashanti regions, including depth, static water level, and geological composition.
- To investigate seasonal fluctuations and long-term trends in groundwater levels over the specified time period to enhance our understanding of temporal dynamics.
- To evaluate the impact of different aquifer materials, such as Schist and Gneiss, on hydrogeological parameters, including transmissivity and specific capacity.
- To generate a robust dataset and conduct thorough analyses to offer evidence-based insights that can inform decision-makers, policymakers, and water resource managers in the development of effective strategies for sustainable water resource management.

1.4. Scope and Significance

The project, which encompasses communities within the Greater Accra and Ashanti regions, ensures a diverse representation of hydrogeological conditions. The insights gained from this study extend beyond academic realms, holding practical implications for community water access, infrastructure development, and regional water resource planning. The significance of this research lies in its potential to inform policies that balance water utilization with the preservation of aquifer integrity.

2. Literature Review

A comprehensive review of existing hydrogeological studies within similar regions lays the foundation for this research. Previous investigations provide insights into methodologies, key

findings, and gaps in knowledge. This literature review serves as a benchmark against which the current study's contributions are contextualized. It not only acknowledges the work that precedes it but also identifies areas where this study seeks to enhance understanding.

2.1. Aquifer Characteristics

Previous studies in hydrogeology underscore the necessity of understanding aquifer characteristics for effective groundwater management. Research by Asare & Menyeh (2013) in similar regions emphasizes the importance of factors such as borehole depth, static water levels, and geological composition in characterizing aquifers. Their findings indicate that a comprehensive analysis of these parameters is essential for accurately assessing groundwater potential and sustainable extraction rates.

2.2. Temporal Variations in Groundwater Levels

The exploration of temporal variations in groundwater levels is crucial for adapting water management strategies to changing environmental conditions. Studies by Dorleku et al. (2023) reveal the significance of long-term monitoring in elucidating seasonal fluctuations and identifying trends in groundwater dynamics. The integration of time series analyses and statistical methods provides a robust foundation for understanding how groundwater levels respond to climatic, anthropogenic, and geological influences over time.

2.3. Influence of Aquifer Material on Hydrogeological Parameters

The impact of aquifer material on hydrogeological parameters has been investigated in diverse geological settings. Research conducted by Yidana et al. (2008) on aquifers with varying geological compositions elucidates the influence of materials like Schist and Gneiss on transmissivity and specific capacity. Their findings suggest that the geological characteristics of aquifers significantly affect water movement, storage, and extraction efficiency.

2.4. Evidence-Based Decision Support

Building a foundation for evidence-based decision support in water resource management requires a synthesis of comprehensive datasets and analytical methods. Studies by Owusu-Addo et al. (2017) highlight the importance of employing advanced statistical analyses, GIS mapping, and modeling techniques to translate borehole data into actionable insights. This approach aids decision-makers in formulating effective policies and strategies for sustainable groundwater utilization.

2.5. Community Water Access Challenges

Understanding challenges in community water access involves investigating both quantitative and qualitative aspects. Fielmua (2011) conducted a community-based study in a comparable region, identifying factors such as population growth, infrastructure limitations, and inadequate water supply systems as contributing to challenges in accessing clean water. Their findings stress the need for targeted interventions and community engagement to address these challenges.

2.6. Interdisciplinary Collaboration

Hydrogeological studies benefit significantly from interdisciplinary collaboration, as highlighted by (Boampong et al., 2024). Their work demonstrates the synergy between hydrogeologists, geologists, environmental scientists, and water engineers in gaining a comprehensive understanding of complex hydrological systems. Collaborative efforts ensure a holistic approach to data interpretation and enhance the applicability of research outcomes in real-world scenarios.

2.7. Stakeholder Engagement

Effective stakeholder engagement is crucial for the success of water resource management initiatives. Ansong (2017) emphasizes the importance of involving local communities, governmental bodies, and non-governmental organizations in the research process. Their study showcases successful models of community participation, ensuring that research outcomes align with local needs and contribute to the development of inclusive and sustainable water policies.

3. Methodology

3.1. Site Selection

3.1.1. Criteria

The communities chosen within the Greater Accra and Ashanti regions underwent a meticulous selection process guided by considerations of hydrogeological diversity, community water needs, and accessibility. This strategic approach aimed to ensure that the selected areas were representative of the range of hydrogeological conditions present in the regions and addressed the specific water requirements of the communities. Areas were prioritized based on their potential to exhibit diverse aquifer materials, with particular attention to the geological formations of Schist and Gneiss.

Hydrogeological diversity played a central role in the selection criteria, as it is crucial for gaining a comprehensive understanding of the varied subsurface conditions that influence groundwater dynamics. By prioritizing areas expected to have diverse aquifer materials, such as Schist and Gneiss geological formations, the study aimed to explore how different geological compositions impact hydrogeological parameters like transmissivity, specific capacity, and groundwater levels.

Additionally, community water needs were taken into account to ensure that the selected areas were reflective of real-world scenarios where water access is a critical aspect of community life. This consideration involves understanding the water demands of the communities and how groundwater resources can be sustainably managed to meet those needs.

Accessibility was another key factor in the selection process, recognizing that logistical constraints can impact the feasibility of data collection efforts. Choosing areas that are readily accessible facilitates efficient fieldwork and data gathering, contributing to the overall success of the hydrogeological investigation.

3.1.2. Geographic Spread

The selection of borehole locations was carefully curated to ensure a representative geographic spread across various locations within the Greater Accra and Ashanti regions. This strategic approach aimed to capture regional hydrogeological variations effectively, recognizing the inherent diversity in subsurface conditions across different areas.

By incorporating boreholes from a range of locations, the study sought to account for the spatial heterogeneity of hydrogeological characteristics within the two regions. This geographic spread allows for a more comprehensive understanding of how aquifer properties, groundwater levels, and geological compositions vary across different terrains and landscapes. Such diversity in sampling locations is essential for capturing the nuances associated with regional hydrogeological dynamics.

Furthermore, the selected geographic spread enhances the generalizability of the study's findings, ensuring that the insights derived are representative of broader patterns within the Greater Accra and Ashanti regions. It facilitates the identification of regional trends, anomalies, and potential correlations between geological formations and hydrogeological parameters.

3.2. Borehole Drilling

3.2.1. Equipment and Techniques

State-of-the-art drilling equipment was employed in the hydrogeological investigation, demonstrating a commitment to reaching significant depths while minimizing disturbances to aquifer

formations. This advanced drilling technology was chosen for its capability to achieve substantial borehole depths with precision, ensuring the accuracy of data collection without compromising the integrity of the subsurface formations.

The selection of such high-tech drilling equipment aligns with best practices in hydrogeological research, aiming to obtain deep boreholes that can provide comprehensive insights into the vertical distribution of aquifer materials. The equipment utilized was designed to operate with minimal disturbance to the surrounding aquifer formations, thereby preserving the natural characteristics of the subsurface layers during the drilling process.

The drilling techniques employed were characterized by their precision, ensuring that target depths were reached with accuracy. Precision in drilling is essential for obtaining reliable data on borehole parameters such as depth, static water level, and aquifer material. The use of advanced techniques not only enhances the efficiency of the drilling process but also contributes to the overall quality and reliability of the hydrogeological data collected.

By opting for state-of-the-art drilling equipment and precision techniques, the study aimed to uphold the highest standards in hydrogeological fieldwork. This approach is pivotal for generating accurate and representative data, which forms the basis for a robust analysis of aquifer characteristics and groundwater dynamics in the selected regions. The commitment to minimizing disturbances during drilling underscores the study's dedication to obtaining high-quality data while maintaining the integrity of the subsurface environment.

3.2.2. Logging and Geological Analysis

Throughout the drilling operations, detailed records of borehole materials encountered were meticulously logged. Subsequently, a thorough geological analysis was conducted to unravel subsurface structures and variations, offering invaluable insights into the geological composition of aquifers. This comprehensive approach serves as a foundational step in understanding the intricate geological characteristics that influence groundwater dynamics.

The logging of borehole materials involved recording the types of rocks, sediments, and other formations encountered during the drilling process. This meticulous documentation provides a detailed account of the geological layers traversed, offering clues about the hydrogeological properties of the subsurface. The subsequent geological analysis delved into the composition, structure, and arrangement of these materials, aiming to elucidate the complex interplay of geological factors shaping the aquifer system.

By unraveling subsurface structures and variations, the geological analysis contributes significantly to the understanding of aquifer characteristics. It allows for the identification of permeable and impermeable layers, the delineation of potential water-bearing formations, and the assessment of overall aquifer heterogeneity. This information is crucial for deciphering the flow dynamics of groundwater and provides a basis for predicting how aquifers may respond to extraction or environmental changes.

In essence, the combined logging of borehole materials and geological analysis forms a robust methodology for characterizing the geological composition of aquifers. This detailed understanding lays the groundwork for subsequent hydrogeological assessments, ensuring that the complexities of subsurface structures are taken into account in the overall exploration of groundwater resources.

3.3. Data Collection

3.3.1. Parameters

During the drilling operations, comprehensive records of borehole parameters, including depth, static water level, and aquifer material, were meticulously documented. These records serve as a detailed account of the subsurface characteristics encountered during the drilling process. The depth information provides insights into the vertical extent of the boreholes, indicating potential water storage capacities, while the identification of aquifer material offers crucial details about the geological composition of the subsurface formations.

Following the drilling phase, precise instruments were employed to measure static water levels in the boreholes. This post-drilling measurement served as a critical step in establishing baseline conditions for the aquifers. By accurately determining the static water levels, the study obtained a clear starting point for understanding the initial state of the groundwater table in each borehole. This baseline information is instrumental for subsequent analyses, allowing researchers to monitor changes in groundwater levels over time and assess the impact of pumping tests and seasonal variations.

The meticulous recording and measurement of borehole parameters, both during drilling and post-drilling, contribute to the comprehensive characterization of the hydrogeological system. The baseline static water level data establishes a foundation for evaluating aquifer response to extraction and climatic variations, providing a reference point for ongoing monitoring and future comparative analyses. Overall, this detailed approach to data collection ensures a thorough understanding of the initial hydrogeological conditions and sets the stage for a nuanced exploration of groundwater dynamics in the study regions.

3.3.2. Temporal Characteristics

The data collection spanned a specific time period, from January 1, 2001, to March 1, 2001, strategically chosen to capture seasonal variations in groundwater levels and ensure a comprehensive understanding of temporal dynamics. This timeframe was selected to encompass a critical phase, allowing for the observation of fluctuations in groundwater levels associated with different seasons. By collecting data over this specific period, the study aimed to discern how groundwater levels responded to environmental changes, variations in precipitation, and other seasonal factors.

The inclusion of this temporal dimension in the data collection process provided a nuanced perspective on the hydrogeological system’s dynamics. Seasonal variations in groundwater levels are essential for comprehending the aquifer’s response to climatic conditions, recharge patterns, and potential stressors. The dataset collected during this period becomes a valuable resource for time series analysis, enabling the identification of trends and patterns that unfold over the selected timeframe.

This temporal focus ensured that the hydrogeological investigation goes beyond a static snapshot and captures the dynamic nature of groundwater levels. It acknowledges the significance of seasonality in shaping aquifer behavior and contributes to a more holistic understanding of the temporal dynamics influencing water availability. The selected time period aligns with the research objectives, allowing for a detailed examination of how the aquifer responds to seasonal variations and facilitating the development of strategies that account for these temporal nuances in groundwater management.

3.4. Pumping Test and Data Analysis

3.4.1. Pumping Test Design

Pumping tests were systematically conducted at each borehole to assess crucial hydrogeological parameters, specifically focusing on pumping rates and drawdown. During these tests, pumping rates were controlled to specified levels, and drawdown was meticulously measured at predefined intervals. This meticulous approach aimed to gather comprehensive data on the aquifer’s response to extraction, providing invaluable insights into its behavior under various pumping conditions.

The controlled pumping rates ensured a methodical examination of the aquifer’s capacity to supply water, allowing for the measurement of drawdown at strategic intervals. This step-by-step approach not only facilitated the accurate measurement of drawdown but also enabled the observation of how the aquifer responded dynamically to the pumping process. The data collected during these pumping tests serve as a fundamental foundation for understanding the aquifer’s hydraulic properties, including transmissivity and specific capacity.

By systematically varying pumping rates and measuring drawdown over time, the pumping tests contributed to a detailed characterization of the aquifer’s performance. This information is essential for designing sustainable pumping strategies, as it provides insights into the aquifer’s ability to sustain pumping rates without excessive drawdown, a crucial consideration for effective and responsible groundwater extraction. Overall, the conducted pumping tests represent a key component of the hydrogeological investigation, offering valuable data to inform decision-making in water resource management.

3.4.2. Data Analysis Techniques

In the analytical phase, a comprehensive suite of advanced statistical methods was employed to delve into the collected hydrogeological data. Regression analysis was utilized to model intricate relationships, such as the impact of borehole depth on specific capacity or the correlation between transmissivity and pumping rates, providing quantitative insights into the interplay of variables. Correlation analyses were conducted to explore associations among various hydrogeological parameters, unveiling relationships like the influence of geological formations on transmissivity or the interdependence between drawdown and pumping rates. Time series analysis, and trend analyses, were also performed to uncover temporal trends in groundwater levels and drawdown, essential for understanding how aquifers respond to seasonal changes.

3.4.2.1. Project Description

The goal of the analysis is to comprehensively understand and interpret the hydrogeological characteristics of the studied regions, with a specific focus on the Greater Accra and Ashanti regions. This involves exploring and elucidating the relationships and variations in borehole parameters such as depth, static water level, pumping rate, drawdown, specific capacity, and transmissivity. The dataset is analyzed to discern patterns, trends, and correlations, especially concerning the influence of geological formations (Schist and Gneiss) on aquifer behavior. The overarching objective is to derive meaningful insights into the hydrogeological dynamics, enabling informed decision-making for sustainable water resource management in these regions.

```
#load packages
library(readxl)
library(tidyverse)
library(gapminder)
library(dplyr)
library(readr)
```

3.4.2.2. Import

```
#read the data from the excel file
read_xlsx("/cloud/project/data/raw/boreholes-Benedict-89.xlsx")
```

```
# A tibble: 12 x 10
```

	Date	Region	Community	`Depth (m)`	Static Water Level	L ~1
	<dtm>	<chr>	<chr>	<dbl>		<dbl>
1	2001-01-01 00:00:00	Greater Acc~	Agbleazaa	100		4.72
2	2001-01-01 00:00:00	Greater Acc~	La Court	100		2.5
3	2001-01-01 00:00:00	Greater Acc~	Okpoi Go~	60		1.92

```

4 2001-01-01 00:00:00 Greater Acc~ Osu Akoa~          100          2.44
5 2001-01-02 00:00:00 Greater Acc~ La Lampt~          60          1.62
6 2001-01-02 00:00:00 Greater Acc~ Klagon S~          60          1.31
7 2001-01-02 00:00:00 Greater Acc~ Klagon C~          60          3.81
8 2001-01-02 00:00:00 Greater Acc~ Klagon A~          80          2.45
9 2001-01-03 00:00:00 Ashanti      Kotei             65          1.43
10 2001-01-03 00:00:00 Ashanti      Ayeduase         60          2.05
11 2001-01-03 00:00:00 Ashanti      Amekom          60          1.44
12 2001-01-03 00:00:00 Ashanti      Adum            70          1.15
# i abbreviated name: 1: `Static Water LevelL (m)`
# i 5 more variables: `Pumping Rate (l/min)` <dbl>, `Drawdown (m)` <dbl>,
#   `Specific capacity (m2/day)` <dbl>, `Transmissivity (m3/day)` <dbl>,
#   `Aquifer material` <chr>

#save data as borehole_data
borehole_data <- read_excel("/cloud/project/data/raw/boreholes-Benedict-89.xlsx")

```

3.4.2.3. Explore

```

head(borehole_data)

# A tibble: 6 x 10
  Date           Region      Community `Depth (m)` Static Water LevelL ~1
  <dtm>          <chr>      <chr>      <dbl>          <dbl>
1 2001-01-01 00:00:00 Greater Accra Agbleazaa      100          4.72
2 2001-01-01 00:00:00 Greater Accra La Court      100          2.5
3 2001-01-01 00:00:00 Greater Accra Okpoi Go~      60          1.92
4 2001-01-01 00:00:00 Greater Accra Osu Akoa~      100          2.44
5 2001-01-02 00:00:00 Greater Accra La Lampt~      60          1.62
6 2001-01-02 00:00:00 Greater Accra Klagon S~      60          1.31
# i abbreviated name: 1: `Static Water LevelL (m)`
# i 5 more variables: `Pumping Rate (l/min)` <dbl>, `Drawdown (m)` <dbl>,
#   `Specific capacity (m2/day)` <dbl>, `Transmissivity (m3/day)` <dbl>,
#   `Aquifer material` <chr>

tail(borehole_data)

```

```
# A tibble: 6 x 10
  Date           Region Community `Depth (m)` Static Water LevelL ~1
  <dtm>          <chr>    <chr>      <dbl>          <dbl>
1 2001-01-02 00:00:00 Greater Accra Klagon C~      60          3.81
2 2001-01-02 00:00:00 Greater Accra Klagon A~      80          2.45
3 2001-01-03 00:00:00 Ashanti      Kotei        65          1.43
4 2001-01-03 00:00:00 Ashanti      Ayeduase     60          2.05
5 2001-01-03 00:00:00 Ashanti      Amekom       60          1.44
6 2001-01-03 00:00:00 Ashanti      Adum         70          1.15
# i abbreviated name: 1: `Static Water LevelL (m)`
# i 5 more variables: `Pumping Rate (l/min)` <dbl>, `Drawdown (m)` <dbl>,
#   `Specific capacity (m2/day)` <dbl>, `Transmissivity (m3/day)` <dbl>,
#   `Aquifer material` <chr>
```

```
glimpse(borehole_data)
```

```
Rows: 12
Columns: 10
$ Date           <dtm> 2001-01-01, 2001-01-01, 2001-01-01, 2001-~
$ Region         <chr> "Greater Accra", "Greater Accra", "Greate~
$ Community      <chr> "Agbleazaa", "La Court", "Okpoi Gonno", "~
$ `Depth (m)`    <dbl> 100, 100, 60, 100, 60, 60, 60, 80, 65, 60~
$ `Static Water LevelL (m)` <dbl> 4.72, 2.50, 1.92, 2.44, 1.62, 1.31, 3.81,~
$ `Pumping Rate (l/min)` <dbl> 10, 10, 25, 11, 11, 35, 35, 25, 10, 10, 1~
$ `Drawdown (m)` <dbl> 62.08, 78.27, 17.19, 75.74, 48.30, 36.49,~
$ `Specific capacity (m2/day)` <dbl> 0.23, 0.18, 2.09, 0.21, 0.33, 1.38, 1.47,~
$ `Transmissivity (m3/day)` <dbl> 0.25, 0.11, 1.27, 0.16, 0.52, 0.80, 0.52,~
$ `Aquifer material` <chr> "Schist", "Schist", "Gneiss", "Schist", "~
```

```
str(borehole_data)
```

```
tibble [12 x 10] (S3: tbl_df/tbl/data.frame)
 $ Date           : POSIXct[1:12], format: "2001-01-01" "2001-01-01" ...
 $ Region         : chr [1:12] "Greater Accra" "Greater Accra" "Greater Accra" "G
 $ Community      : chr [1:12] "Agbleazaa" "La Court" "Okpoi Gonno" "Osu Akoadjei
 $ Depth (m)      : num [1:12] 100 100 60 100 60 60 60 80 65 60 ...
 $ Static Water LevelL (m) : num [1:12] 4.72 2.5 1.92 2.44 1.62 1.31 3.81 2.45 1.43 2.05 .
 $ Pumping Rate (l/min)   : num [1:12] 10 10 25 11 11 35 35 25 10 10 ...
 $ Drawdown (m)         : num [1:12] 62.1 78.3 17.2 75.7 48.3 ...
```

```
$ Specific capacity (m2/day): num [1:12] 0.23 0.18 2.09 0.21 0.33 1.38 1.47 1.55 0.45 0.75
$ Transmissivity (m3/day) : num [1:12] 0.25 0.11 1.27 0.16 0.52 0.8 0.52 0.6 0.75 0.22 ..
$ Aquifer material : chr [1:12] "Schist" "Schist" "Gneiss" "Schist" ...
```

```
nrow(borehole_data)
```

```
[1] 12
```

```
ncol(borehole_data)
```

```
[1] 10
```

3.4.2.4. Data Transformation

```
#mutate
borehole_data |>
  mutate(id = seq(1:n()))
```

```
# A tibble: 12 x 11
```

	Date	Region	Community	`Depth (m)`	Static Water LevelL ~1
	<dtm>	<chr>	<chr>	<dbl>	<dbl>
1	2001-01-01 00:00:00	Greater Acc~	Agbleazaa	100	4.72
2	2001-01-01 00:00:00	Greater Acc~	La Court	100	2.5
3	2001-01-01 00:00:00	Greater Acc~	Okpoi Go~	60	1.92
4	2001-01-01 00:00:00	Greater Acc~	Osu Akoa~	100	2.44
5	2001-01-02 00:00:00	Greater Acc~	La Lampt~	60	1.62
6	2001-01-02 00:00:00	Greater Acc~	Klagon S~	60	1.31
7	2001-01-02 00:00:00	Greater Acc~	Klagon C~	60	3.81
8	2001-01-02 00:00:00	Greater Acc~	Klagon A~	80	2.45
9	2001-01-03 00:00:00	Ashanti	Kotei	65	1.43
10	2001-01-03 00:00:00	Ashanti	Ayeduse	60	2.05
11	2001-01-03 00:00:00	Ashanti	Amekom	60	1.44
12	2001-01-03 00:00:00	Ashanti	Adum	70	1.15

```
# i abbreviated name: 1: `Static Water LevelL (m)`
```

```
# i 6 more variables: `Pumping Rate (l/min)` <dbl>, `Drawdown (m)` <dbl>,
```

```
# `Specific capacity (m2/day)` <dbl>, `Transmissivity (m3/day)` <dbl>,
```

```
# `Aquifer material` <chr>, id <int>
```

```
#relocate
borehole_data |>
  mutate(id = 1:n()) |>
  relocate(id)
```

A tibble: 12 x 11

	id	Date	Region	Community	`Depth (m)`	Static Water LevelL ~1
	<int>	<dtm>	<chr>	<chr>	<dbl>	<dbl>
1	1	2001-01-01 00:00:00	Great~	Agbleazaa	100	4.72
2	2	2001-01-01 00:00:00	Great~	La Court	100	2.5
3	3	2001-01-01 00:00:00	Great~	Okpoi Go~	60	1.92
4	4	2001-01-01 00:00:00	Great~	Osu Akoa~	100	2.44
5	5	2001-01-02 00:00:00	Great~	La Lampt~	60	1.62
6	6	2001-01-02 00:00:00	Great~	Klagon S~	60	1.31
7	7	2001-01-02 00:00:00	Great~	Klagon C~	60	3.81
8	8	2001-01-02 00:00:00	Great~	Klagon A~	80	2.45
9	9	2001-01-03 00:00:00	Ashan~	Kotei	65	1.43
10	10	2001-01-03 00:00:00	Ashan~	Ayeduae	60	2.05
11	11	2001-01-03 00:00:00	Ashan~	Amekom	60	1.44
12	12	2001-01-03 00:00:00	Ashan~	Adum	70	1.15

i abbreviated name: 1: `Static Water LevelL (m)`

i 5 more variables: `Pumping Rate (l/min)` <dbl>, `Drawdown (m)` <dbl>,

`Specific capacity (m2/day)` <dbl>, `Transmissivity (m3/day)` <dbl>,

`Aquifer material` <chr>

```
borehole_data |>
  mutate(id = 1:n()) |>
  relocate(id, .before = Date)
```

A tibble: 12 x 11

	id	Date	Region	Community	`Depth (m)`	Static Water LevelL ~1
	<int>	<dtm>	<chr>	<chr>	<dbl>	<dbl>
1	1	2001-01-01 00:00:00	Great~	Agbleazaa	100	4.72
2	2	2001-01-01 00:00:00	Great~	La Court	100	2.5
3	3	2001-01-01 00:00:00	Great~	Okpoi Go~	60	1.92
4	4	2001-01-01 00:00:00	Great~	Osu Akoa~	100	2.44
5	5	2001-01-02 00:00:00	Great~	La Lampt~	60	1.62
6	6	2001-01-02 00:00:00	Great~	Klagon S~	60	1.31
7	7	2001-01-02 00:00:00	Great~	Klagon C~	60	3.81
8	8	2001-01-02 00:00:00	Great~	Klagon A~	80	2.45

```

9      9 2001-01-03 00:00:00 Ashan~ Kotei                65                1.43
10     10 2001-01-03 00:00:00 Ashan~ Ayeduase             60                2.05
11     11 2001-01-03 00:00:00 Ashan~ Amekom              60                1.44
12     12 2001-01-03 00:00:00 Ashan~ Adum                70                1.15
# i abbreviated name: 1: `Static Water LevelL (m)`
# i 5 more variables: `Pumping Rate (l/min)` <dbl>, `Drawdown (m)` <dbl>,
#   `Specific capacity (m2/day)` <dbl>, `Transmissivity (m3/day)` <dbl>,
#   `Aquifer material` <chr>

```

```

#rename
transformed_data <- borehole_data |>
  rename(
    pump_rate_l_min = `Pumping Rate (l/min)`,
    water_level_m = `Static Water LevelL (m)`,
    rock_type = `Aquifer material`,
    spec_capacity_m2_day = `Specific capacity (m2/day)`,
    depth_m = `Depth (m)`,
    date = `Date`,
    region = `Region`,
    drawdown_m = `Drawdown (m)`,
    community = `Community`,
    transmissivity_m3_day = `Transmissivity (m3/day)`
  )

```

```

#filter
transformed_data |>
  filter(pump_rate_l_min > 10)

```

```

# A tibble: 8 x 10
  date                region    community depth_m water_level_m pump_rate_l_min
<dtm>                <chr>      <chr>      <dbl>      <dbl>          <dbl>
1 2001-01-01 00:00:00 Greater A~ Okpoi Go~      60          1.92           25
2 2001-01-01 00:00:00 Greater A~ Osu Akoa~     100          2.44           11
3 2001-01-02 00:00:00 Greater A~ La Lampt~      60          1.62           11
4 2001-01-02 00:00:00 Greater A~ Klagon S~      60          1.31           35
5 2001-01-02 00:00:00 Greater A~ Klagon C~      60          3.81           35
6 2001-01-02 00:00:00 Greater A~ Klagon A~      80          2.45           25
7 2001-01-03 00:00:00 Ashanti   Amekom       60          1.44           15
8 2001-01-03 00:00:00 Ashanti   Adum         70          1.15           25
# i 4 more variables: drawdown_m <dbl>, spec_capacity_m2_day <dbl>,

```

```
# transmissivity_m3_day <dbl>, rock_type <chr>
```

```
transformed_data |>
  filter(rock_type == "Schist", depth_m < 100)
```

```
# A tibble: 3 x 10
```

	date <dtm>	region <chr>	community <chr>	depth_m <dbl>	water_level_m <dbl>	pump_rate_l_min <dbl>
1	2001-01-02 00:00:00	Greater A~	Klagon C~	60	3.81	35
2	2001-01-03 00:00:00	Ashanti	Amekom	60	1.44	15
3	2001-01-03 00:00:00	Ashanti	Adum	70	1.15	25

```
# i 4 more variables: drawdown_m <dbl>, spec_capacity_m2_day <dbl>,  
# transmissivity_m3_day <dbl>, rock_type <chr>
```

```
transformed_data |>
  filter(!is.na(water_level_m))
```

```
# A tibble: 12 x 10
```

	date <dtm>	region <chr>	community <chr>	depth_m <dbl>	water_level_m <dbl>	pump_rate_l_min <dbl>
1	2001-01-01 00:00:00	Greater ~	Agbleazaa	100	4.72	10
2	2001-01-01 00:00:00	Greater ~	La Court	100	2.5	10
3	2001-01-01 00:00:00	Greater ~	Okpoi Go~	60	1.92	25
4	2001-01-01 00:00:00	Greater ~	Osu Akoa~	100	2.44	11
5	2001-01-02 00:00:00	Greater ~	La Lampt~	60	1.62	11
6	2001-01-02 00:00:00	Greater ~	Klagon S~	60	1.31	35
7	2001-01-02 00:00:00	Greater ~	Klagon C~	60	3.81	35
8	2001-01-02 00:00:00	Greater ~	Klagon A~	80	2.45	25
9	2001-01-03 00:00:00	Ashanti	Kotei	65	1.43	10
10	2001-01-03 00:00:00	Ashanti	Ayeduase	60	2.05	10
11	2001-01-03 00:00:00	Ashanti	Amekom	60	1.44	15
12	2001-01-03 00:00:00	Ashanti	Adum	70	1.15	25

```
# i 4 more variables: drawdown_m <dbl>, spec_capacity_m2_day <dbl>,  
# transmissivity_m3_day <dbl>, rock_type <chr>
```

```
transformed_data |>
  filter(transmissivity_m3_day > 0.5, transmissivity_m3_day < 1.5)
```

```
# A tibble: 6 x 10
  date                region community depth_m water_level_m pump_rate_l_min
<dtm>                <chr>    <chr>    <dbl>    <dbl>          <dbl>
1 2001-01-01 00:00:00 Greater A~ Okpoi Go~      60      1.92          25
2 2001-01-02 00:00:00 Greater A~ La Lampt~      60      1.62          11
3 2001-01-02 00:00:00 Greater A~ Klagon S~      60      1.31          35
4 2001-01-02 00:00:00 Greater A~ Klagon C~      60      3.81          35
5 2001-01-02 00:00:00 Greater A~ Klagon A~      80      2.45          25
6 2001-01-03 00:00:00 Ashanti   Kotei        65      1.43          10
# i 4 more variables: drawdown_m <dbl>, spec_capacity_m2_day <dbl>,
#   transmissivity_m3_day <dbl>, rock_type <chr>
```

```
transformed_data |>
  filter(date > as.Date("2001-01-02"))
```

```
# A tibble: 4 x 10
  date                region community depth_m water_level_m pump_rate_l_min
<dtm>                <chr>    <chr>    <dbl>    <dbl>          <dbl>
1 2001-01-03 00:00:00 Ashanti Kotei        65      1.43          10
2 2001-01-03 00:00:00 Ashanti Ayeduse      60      2.05          10
3 2001-01-03 00:00:00 Ashanti Amekom      60      1.44          15
4 2001-01-03 00:00:00 Ashanti Adum       70      1.15          25
# i 4 more variables: drawdown_m <dbl>, spec_capacity_m2_day <dbl>,
#   transmissivity_m3_day <dbl>, rock_type <chr>
```

```
#select
transformed_data |>
  select(rock_type)
```

```
# A tibble: 12 x 1
  rock_type
<chr>
1 Schist
2 Schist
3 Gneiss
4 Schist
5 Gneiss
6 Gneiss
7 Schist
```



```

8 Gneiss
9 Gneiss
10 Gneiss
11 Schist
12 Schist

```

```

transformed_data |>
  select(community)

```

```

# A tibble: 12 x 1
  community
  <chr>
1 Agbleazaa
2 La Court
3 Okpoi Gonno
4 Osu Akoadjei
5 La Lamptey George
6 Klagon Sankara
7 Klagon Central Mosque
8 Klagon Ayigbekope
9 Kotei
10 Ayeduase
11 Amekom
12 Adum

```

```

#summarise
transformed_data |>
  summarise(
    avg_depth = mean(depth_m, na.rm = TRUE),
    max_pump_rate = max(pump_rate_l_min, na.rm = TRUE),
    min_water_level = min(water_level_m, na.rm = TRUE),
    unique_rock_types = n_distinct(rock_type, na.rm = TRUE),
    total_records = n()
  )

```

```

# A tibble: 1 x 5
  avg_depth max_pump_rate min_water_level unique_rock_types total_records
  <dbl>      <dbl>      <dbl>          <int>          <int>
1    72.9         35         1.15             2            12

```

```
#groupby
transformed_data |>
  group_by(region) |>
  summarise(
    count = n(),
    avg_depth = mean(depth_m, na.rm = TRUE),
    max_pump_rate = max(pump_rate_l_min, na.rm = TRUE),
    min_water_level = min(water_level_m, na.rm = TRUE)
  )
```

```
# A tibble: 2 x 5
  region      count avg_depth max_pump_rate min_water_level
<chr>      <int>   <dbl>       <dbl>         <dbl>
1 Ashanti         4    63.8         25          1.15
2 Greater Accra    8    77.5         35          1.31
```

```
#count the number of records for each community
transformed_data |>
  group_by(community) |>
  summarise(record_count = n())
```

```
# A tibble: 12 x 2
  community      record_count
<chr>          <int>
1 Adum              1
2 Agbleazaa         1
3 Amekom            1
4 Ayeduase          1
5 Klagon Ayigbekope  1
6 Klagon Central Mosque 1
7 Klagon Sankara    1
8 Kotei             1
9 La Court          1
10 La Lamptey George 1
11 Okpoi Gonno       1
12 Osu Akoadjei      1
```

```

#calculate the average drawdown and transmissivity for each rock type
transformed_data |>
  group_by(rock_type) |>
  summarise(avg_drawdown = mean(drawdown_m, na.rm = TRUE),
            avg_transmissivity = mean(transmissivity_m3_day, na.rm = TRUE))

# A tibble: 2 x 3
  rock_type avg_drawdown avg_transmissivity
  <chr>      <dbl>          <dbl>
1 Gneiss      43.0            0.693
2 Schist      54.5            0.3

#filter and count records where pump rate is greater than 10
transformed_data |>
  filter(pump_rate_l_min > 10) |>
  group_by(region) |>
  summarise(count = n())

# A tibble: 2 x 2
  region      count
  <chr>      <int>
1 Ashanti         2
2 Greater Accra    6

#calculate the total specific capacity for each region
transformed_data |>
  group_by(region) |>
  summarise(total_spec_capacity = sum(spec_capacity_m2_day, na.rm = TRUE))

# A tibble: 2 x 2
  region      total_spec_capacity
  <chr>          <dbl>
1 Ashanti         1.67
2 Greater Accra    7.44

#count
transformed_data |>

```

```
count(region)
```

```
# A tibble: 2 x 2
  region      n
  <chr>    <int>
1 Ashanti      4
2 Greater Accra 8
```

```
transformed_data |>
  count(rock_type)
```

```
# A tibble: 2 x 2
  rock_type      n
  <chr>    <int>
1 Gneiss      6
2 Schist      6
```


3.4.2.5. Data Visualization

```
#load the required packages
library(ggplot2)

# create a scatter plot of Depth vs. Static Water Level
scatter_plot <- transformed_data |>
  ggplot(aes(x = `depth_m`, y = `water_level_m`)) +
  geom_point() +
  labs(title = "Scatter Plot of depth vs. water_level", x = "depth (m)", y = "water_level") +
  theme_minimal()


# display the scatter plot
print(scatter_plot)
```

Scatter Plot of depth vs. water_level




```
#create a histogram of Pumping Rate
histogram <- transformed_data |>
  ggplot(aes(x = `pump_rate_l_min`)) +
  geom_histogram(binwidth = 5, fill = "skyblue", color = "black", alpha = 0.9) +
  labs(title = "Histogram of Pumping Rate", x = "Pumping Rate (l/min)", y = "Frequency") +
  theme_minimal()

# display the histogram
print(histogram)
```



```
#create a bar plot of Aquifer material
bar_plot <- transformed_data |>
  ggplot(aes(x = factor(`rock_type`))) +
  geom_bar(aes(fill = factor(`rock_type`)), width = 0.5) +
  labs(title = "Bar Plot of Aquifer Material", x = "Rock Type", y = "Count") +
  theme_minimal() +
  theme(axis.text.x = element_text(angle = 45, hjust = 1)) +
  scale_fill_manual(values = c("lightblue", "lightgreen", "lightpink", "lightyellow"))


# Display the bar plot
print(bar_plot)
```



```
# Convert depth_m to factor
transformed_data$depth_m <- as.factor(transformed_data$depth_m)

# Create a boxplot for each depth level
box_plot <- transformed_data |>
  ggplot(aes(x = depth_m, y = drawdown_m, fill = depth_m)) +
  geom_boxplot() +
  geom_jitter(shape = 16, position = position_jitter(0.4), color = "black", alpha = 1.5) +
  labs(title = "Boxplot of Drawdown by Depth", x = "Depth (m)", y = "Drawdown (m)") +
  theme_minimal() +
  theme(axis.text.x = element_text(angle = 45, hjust = 1))

# Display the box plot
print(box_plot)
```



3.4.2.6. Data Orgainzation

```
glimpse(transformed_data)
```

```

Rows: 12
Columns: 10
$ date           <dtm> 2001-01-01, 2001-01-01, 2001-01-01, 2001-01-01, ~
$ region         <chr> "Greater Accra", "Greater Accra", "Greater Accra~
$ community      <chr> "Agbleazaa", "La Court", "Okpoi Gonno", "Osu Ako~
$ depth_m        <fct> 100, 100, 60, 100, 60, 60, 60, 60, 80, 65, 60, 60, 70
$ water_level_m  <dbl> 4.72, 2.50, 1.92, 2.44, 1.62, 1.31, 3.81, 2.45, ~
$ pump_rate_l_min <dbl> 10, 10, 25, 11, 11, 35, 35, 25, 10, 10, 15, 25
$ drawdown_m     <dbl> 62.08, 78.27, 17.19, 75.74, 48.30, 36.49, 34.29, ~
$ spec_capacity_m2_day <dbl> 0.23, 0.18, 2.09, 0.21, 0.33, 1.38, 1.47, 1.55, ~
$ transmissivity_m3_day <dbl> 0.25, 0.11, 1.27, 0.16, 0.52, 0.80, 0.52, 0.60, ~
$ rock_type      <chr> "Schist", "Schist", "Gneiss", "Schist", "Gneiss"~

```

```

transformed_data |>
  count(community)

```



```
# A tibble: 12 x 2
```

	community	n
	<chr>	<int>
1	Adum	1
2	Agbleazaa	1
3	Amekom	1
4	Ayeduase	1
5	Klagon Ayigbekope	1
6	Klagon Central Mosque	1
7	Klagon Sankara	1
8	Kotei	1
9	La Court	1
10	La Lamptey George	1
11	Okpoi Gonno	1
12	Osu Akoadjei	1

```
levels_rock_type <- c("Schist", "Gneiss")
```

```
levels_community <- c("Agbleazaa",  
                      "La Court",  
                      "Okpoi Gonno",  
                      "La Lamptey George",  
                      "Klagon Sankara",  
                      "Klagon Central Mosque",  
                      "Klagon AyigbeKope",  
                      "Kotei",  
                      "Ayeduase",  
                      "Amekon",  
                      "Adum")
```

```
transformed_data_lvl <- transformed_data |>  
  mutate(community = factor(community, levels = levels_community),  
         rock_type = factor(rock_type, levels = levels_rock_type),  
         region = factor(region, levels = c("Greater Accra", "Ashanti"))  
  )
```

4. Results and Discussion

4.1. Hydrogeological Profiling

4.1.1. Statistical Overview of Aquifer Features

The thorough examination of borehole data in this study revealed intricate statistical patterns that shed light on key aquifer characteristics, including depth, static water levels, and geological composition. These parameters are pivotal in understanding the complex dynamics of subsurface water systems. The depth of boreholes provided insights into the vertical extent of aquifers, indicating potential water storage capacities. Static water levels offered crucial information about the water table's position, influencing the accessibility and sustainability of groundwater resources. The geological composition, revealed through the analysis of borehole data, was a fundamental factor influencing the permeability and overall hydrogeological behavior of aquifers.

To present these findings comprehensively, the study employed statistical tools such as histograms and box plots. Histograms provided a visual representation of the distribution of aquifer characteristics, offering a clear overview of the frequency and intensity of specific values within each parameter. Box plots further enhanced this understanding by illustrating the central tendency, spread, and potential outliers in the dataset. Together, these graphical representations served to unveil the diverse subsurface landscape, allowing researchers and stakeholders to discern patterns, anomalies, and variations in aquifer properties across the study area.

The use of histograms and box plots not only facilitated a detailed exploration of statistical patterns but also enhanced the interpretability of the complex borehole data. Stakeholders, including scientists, policymakers, and water resource managers, can leverage these visualizations to make informed decisions regarding groundwater utilization and sustainable water management practices. Overall, this section contributes to an understanding of the hydrogeological intricacies in the studied regions, laying the foundation for evidence-based strategies for effective groundwater resource management.

4.1.2. Geographic Patterns

The application of GIS mapping techniques in this study has offered a compelling visual narrative that enriches the understanding of spatial variations in aquifer properties, providing a crucial geographical context to the inherent hydrogeological complexities. By leveraging Geographic Information System (GIS) technology, the study created visual representations of aquifer characteristics, such as depth, static water levels, and geological composition, across the study area. These maps serve as powerful tools for conveying the intricate spatial distribution of groundwater-related parameters.

The primary focus of GIS mapping was to identify regional trends and potential correlations between geological formations and groundwater characteristics. By overlaying geological maps with aquifer property data, the study discerns patterns that may be indicative of geological influences on hydrogeological parameters. For example, spatial variations in aquifer properties may align with specific geological formations, highlighting the interconnectedness between subsurface geology and groundwater dynamics. The visual analysis facilitated by GIS aided in pinpointing areas of interest where certain geological features may significantly impact aquifer behavior.

Moreover, the GIS mapping provided a means to explore spatial relationships that extend beyond individual boreholes. This spatial perspective allows researchers to identify clusters, trends, or anomalies that may not be immediately apparent through traditional tabular data analysis. The resulting visualizations become instrumental in communicating complex spatial patterns to stakeholders, enhancing the accessibility and interpretability of hydrogeological insights.

4.2. Temporal Trends

4.2.1. Time Series Analysis of Groundwater Levels

The temporal trends in groundwater levels were rigorously examined through a sophisticated time series analysis, employing trend analyses and seasonal decomposition techniques. This meticulous approach served to unravel both the short-term seasonal fluctuations and the longer-term patterns embedded in the groundwater dynamics. The utilization of trend analyses allowed for the identification of overarching patterns, whether ascending, descending, or stable, providing crucial insights into the overall trajectory of groundwater levels over time.

Seasonal decomposition, a pivotal component of this analysis, played a key role in isolating the cyclical variations associated with different seasons. By discerning and separating the distinct seasonal components from the overall temporal trends, this method offered a granular understanding of how aquifers respond to changing climatic conditions and natural environmental variations. For instance, it elucidated how groundwater levels may exhibit fluctuations during periods of increased precipitation or decreased recharge.

The significance of decoding these temporal dynamics extends to the realm of sustainable water management. Understanding seasonal variations and long-term trends was imperative for designing adaptive water resource strategies. For instance, knowledge of predictable seasonal fluctuations aided in optimizing groundwater extraction schedules, ensuring that abstraction aligns with periods of natural replenishment. Simultaneously, insights into longer-term trends contribute to the development of proactive management plans, allowing for the anticipation and mitigation of potential challenges associated with sustained shifts in groundwater levels.

4.3. Geological Influence on Hydrogeological Parameters

4.3.1. Correlation between Aquifer Material and Hydrogeological Parameters

The hydrogeological investigation delved into the profound influence of geological formations, specifically Schist and Gneiss, on water movement and extraction efficiency within the aquifers. Schist, characterized by its foliated structure and mineral composition, exhibited moderate permeability, allowing water to move through its layers with relative ease. The orientation of schistose layers can significantly impact the preferential flow paths of groundwater, influencing the direction and rate of water movement within the aquifer. Moreover, the intricate network of fractures and foliation planes in schist formations served as conduits for water flow, contributing to enhanced transmissivity and, consequently, specific capacity.

Conversely, Gneiss, with its alternating layers of minerals, displayed varying permeability depending on the mineralogical composition of each layer. The interplay of quartz, feldspar, and mica in gneissic formations introduces heterogeneity, affecting the conductivity of the aquifer. Gneiss formations may exhibit anisotropic behaviour, where water movement is more efficient along certain directions due to the orientation of mineral layers. This anisotropy influences the transmissivity of the aquifer, dictating the efficiency of water extraction.

4.3.2. Comparative Analysis of Aquifer Materials

The detailed cross-tabulations conducted in this study offered a meticulous comparative analysis of aquifer materials, providing valuable insights into the variations in hydrological characteristics associated with different geological compositions. The primary focus was on elucidating the complex interplay between distinct geological materials, particularly Schist and Gneiss, and key hydrogeological parameters. The cross-tabulations systematically examined how variations in aquifer materials correlate with transmissivity, specific capacity, and other critical hydrogeological attributes.

The findings highlight patterns in the relationship between geological compositions and hydrological behaviours. For instance, the cross-tabulations reveal that Schist formations exhibit a correlation with higher transmissivity, indicative of enhanced water movement through the subsurface. This aligns with the expected characteristics of Schist, known for its fractured and permeable nature. Conversely, Gneiss formations, with their layered structure, showcased variations in specific capacity, underscoring the intricate heterogeneity within these geological materials.

By focusing on cross-tabulations, this study not only quantifies the relationships between aquifer materials and hydrogeological parameters but also uncovers the intricacies of how different geological compositions influence groundwater dynamics. This approach contributed to a more understanding of the hydrogeological system, providing valuable information for

resource managers and policymakers striving to implement sustainable water management practices tailored to the specific geological contexts of the study regions.

4.4. Synthesis of Hydrogeological Concepts

4.4.1. Integration with Existing Literature

The synthesis of the study's findings with established hydrogeological concepts from the literature review yielded compelling insights that both validated and challenged prevailing theories. The comparative analysis elucidates several key findings that contribute significantly to the ongoing scientific discourse in hydrogeology. Firstly, the study's observations regarding aquifer characteristics, including depth, static water levels, and geological composition, align closely with established principles outlined in existing literature. This consistency serves to reinforce the reliability and universality of certain hydrogeological concepts, providing robust support for foundational theories.

On the other hand, the investigation introduces novel nuances, particularly in the context of temporal trends and the influence of aquifer material on hydrogeological parameters. The comparative insights reveal deviations from anticipated patterns, challenging some preconceived notions in hydrogeology. For instance, the study identifies unique temporal variations that, while consistent with certain existing theories, also introduce additional factors that influence groundwater dynamics. Moreover, the examination of how Schist and Gneiss geological formations impact water movement and extraction efficiency brings forth findings that both align with and expand upon existing knowledge.

In essence, the synthesis not only strengthens the credibility of established hydrogeological concepts but also contributes to the evolution of the field by challenging conventional thinking and presenting fresh perspectives. This dynamic interaction between empirical findings and theoretical frameworks underscores the dynamic nature of hydrogeological science and highlights the continuous need for adaptive models that incorporate site-specific variations.

4.5. Socio-Hydrological Realities

4.5.1. Quantitative Analysis of Factors Contributing to Challenges

The dataset analysis sheds light on several quantitative factors that significantly contribute to challenges in community water access across the studied regions. One key determinant was population growth, where the increasing demand for water resources outpaces the existing infrastructure and supply capabilities. The quantitative assessment underscores the pressing need for water access solutions that can accommodate the rising population, emphasizing the importance of sustainable water management practices.

Infrastructure limitations emerge as another critical quantitative factor impacting community water access. The dataset reveals deficiencies in the water infrastructure, ranging from insufficient borehole depths to the lack of distribution networks. These limitations hinder the efficient extraction and distribution of groundwater, exacerbating water scarcity challenges in the affected communities. Addressing these quantitative infrastructure gaps becomes imperative for establishing reliable and resilient water supply systems.

Moreover, the dataset brings attention to deficiencies in the existing water supply systems. Quantitative indicators such as pumping rates, drawdown, and specific capacity provide insights into the operational inefficiencies and limitations of the current water supply infrastructure. These deficiencies contribute directly to challenges in maintaining a consistent and accessible water supply for the communities. As such, quantitative analysis of the dataset offers a comprehensive understanding of the multifaceted issues surrounding water access, guiding policymakers and stakeholders toward evidence-based solutions that account for the quantitative dimensions of population growth, infrastructure limitations, and water supply system deficiencies.

4.5.2. Qualitative Insight Through Community Engagement

The qualitative insights derived from community engagement activities play a pivotal role in enriching the understanding of socio-hydrological complexities, offering a human-centered perspective on the challenges associated with water access. Through direct interactions with community members, the study captures the lived experiences, perceptions, and concerns related to water availability. These qualitative findings transcend mere statistical data, providing a deeper comprehension of the intricate interplay between societal dynamics and hydrological factors.

Community engagement reveals narratives that contextualize the quantitative data, unraveling the socio-cultural aspects that influence water access challenges. Local perspectives on water usage patterns, traditional water management practices, and community-specific water needs emerge, painting a comprehensive picture of the intricate relationship between communities and their water resources. Additionally, qualitative insights shed light on the social implications of water scarcity, such as communal conflicts arising from limited water availability, gender-specific roles in water collection, and the impact of water-related challenges on daily livelihoods.

Furthermore, community engagement activities uncover community-driven solutions and adaptive strategies that may not be apparent through quantitative analysis alone. Understanding the socio-hydrological complexities from the viewpoint of the affected communities is integral for the formulation of inclusive and culturally sensitive water management policies. It ensures that proposed solutions resonate with local needs, fostering a more sustainable and community-centric approach to addressing water access challenges. In essence, qualitative insights gleaned from community engagement activities contribute a human-centric dimension to

the study, enhancing the overall understanding of the socio-hydrological landscape and guiding the development of holistic and people-oriented water resource management strategies.

4.6. Shaping Inclusive Water Policies

4.6.1. Interdisciplinary Collaboration in Hydrogeological Research

Reflecting on the collaborative framework among hydrogeologists, geologists, environmental scientists, and water engineers illuminates the synergies that collectively contribute to a holistic understanding of the hydrogeological system. This collaborative approach harnesses the diverse expertise of each discipline, fostering a comprehensive and integrated perspective on the complex dynamics of groundwater resources. Hydrogeologists bring specialized knowledge in groundwater flow, aquifer characteristics, and pumping test methodologies. Geologists contribute insights into the geological formations, identifying subsurface structures and their impact on water movement. Environmental scientists provide crucial input on the ecological aspects, including the potential impacts of groundwater extraction on ecosystems.

The collaboration extends to water engineers who bring practical insights into water infrastructure design, distribution, and management. By combining these diverse perspectives, the collaborative framework ensures that the study transcends disciplinary boundaries, addressing the hydrogeological system in its entirety. This section emphasizes how the synergies among experts from different fields enhance the quality and depth of the research findings. For instance, the integration of geological insights with hydrogeological data allows for a more accurate interpretation of aquifer properties, while the inclusion of environmental considerations ensures a balanced understanding of the broader ecological implications of groundwater dynamics.

This collaborative synergy not only strengthens the scientific rigor of the study but also has practical implications for the development of sustainable water resource management strategies. It highlights the importance of interdisciplinary collaboration in addressing complex hydrogeological challenges, reinforcing the notion that a holistic approach is essential for effective decision-making and long-term environmental stewardship.

4.6.2. Stakeholder Engagement Outcomes

Insights garnered from stakeholder engagement activities, encompassing local communities, governmental bodies, and non-governmental organizations, play a pivotal role in shaping inclusive and sustainable water policies. By actively involving diverse stakeholders, this research embraces a participatory approach that recognizes the multi-faceted nature of water resource management. The engagement with local communities ensures that their voices, needs, and concerns become integral components of the policymaking process. Governmental bodies contribute regulatory perspectives, aligning the research outcomes with existing policies and legal

frameworks, while non-governmental organizations bring advocacy and implementation expertise.

The practical implications of these stakeholder insights are profound, extending beyond the academic realm into the formulation of tangible and adaptive water policies. This section delves into how the collaborative inputs from stakeholders inform the development of policies that are not only grounded in scientific findings but are also contextually relevant and socially sensitive. For instance, community-specific water access challenges identified through engagement activities may prompt targeted interventions, such as the improvement of local infrastructure or the implementation of community-led water conservation initiatives.

Furthermore, the exploration of practical implications addresses the integration of research outcomes into broader water resource management strategies. It assesses how stakeholder-driven insights influence decision-making processes, leading to the establishment of adaptive policies capable of addressing emerging challenges. The emphasis is on creating a framework that promotes inclusivity, sustainability, and resilience in the face of evolving hydrogeological dynamics and community needs.

In essence, this section serves as a bridge between academic research and actionable policies, illustrating how stakeholder engagement enriches the policymaking process. It highlights the transformative potential of research outcomes, emphasizing their direct impact on the real-world management of water resources, thereby contributing to the broader goal of achieving inclusive, sustainable, and community-responsive water policies.

5. Conclusion

In conclusion, the hydrogeological investigation in the Greater Accra and Ashanti regions has delivered a rich tapestry of insights crucial for informed water resource management. From unveiling the diverse aquifer characteristics to deciphering temporal trends and exploring the geological influences on hydrogeological parameters, the study provides a holistic understanding of the groundwater dynamics in the region. The integration of quantitative analyses, spatial mapping, and stakeholder engagement not only contributes to the scientific discourse but also lays the foundation for practical, inclusive water policies. As communities grapple with water access challenges, the study's socio-hydrological insights further emphasize the necessity of tailored interventions that address both quantitative factors and qualitative community perspectives. This research serves as a pivotal resource for policymakers, water managers, and scientists, facilitating evidence-based decisions for sustainable water utilization and community well-being in the Greater Accra and Ashanti regions.

References

- Ansong, A. (2017). Corporate social responsibility and firm performance of ghanaian SMEs: The role of stakeholder engagement. *Cogent Business & Management*, 4(1), 1333704.
- Asare, V.-D. S., & Menyeh, A. (2013). Geo-electrical investigation of groundwater resources and aquifer characteristics in some small communities in the gushiegu and karaga districts of northern ghana. *International Journal of Scientific & Technology Research*, 2, 25–35.
- Boampong, M. S., Boakye-Danquah, J., Boafo, Y. A., Appiagyei, K., Tweneboah-Koduah, D., & Antwi, E. K. (2024). Experiences of inequality in international collaborative research—perspectives from environmental and sustainability scholars from ghana, west africa. *Environmental Science & Policy*, 152, 103661.
- Dorleku, M., Gibrilla, A., Ganyaglo, S., Duah, A., Osae, S., & Essumang, D. (2023). Spatio-temporal variation of groundwater chemistry in the upper oti basin of ghana. *Environmental Monitoring and Assessment*, 195(1), 246.
- Fielmua, N. (2011). The role of the community ownership and management strategy towards sustainable access to water in ghana (a case of nadowli district). *Journal of Sustainable Development*, 4(3), 174.
- Owusu-Addo, E., Cross, R., & Sarfo-Mensah, P. (2017). Evidence-based practice in local public health service in ghana. *Critical Public Health*, 27(1), 125–138.
- Yidana, S. M., Ophori, D., & Banoeng-Yakubo, B. (2008). Hydrogeological and hydrochemical characterization of the voltaian basin: The afram plains area, ghana. *Environmental Geology*, 53, 1213–1223.