

# Slide 7–LU decomposition

## MAT2040 Linear Algebra

# Upper triangular matrix

## Definition 7.1: (upper triangular matrix)

$A = (a_{ij})_{n \times n}$  is said to be **upper triangular** if  $a_{ij} = 0$  for  $i > j$ .

A  $4 \times 4$  upper triangular matrix is given as follows:

$$\begin{bmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix}$$

\* is arbitrary number.

# Lower triangular matrix

$A = (a_{ij})_{n \times n}$  is said to be **lower triangular** if  $a_{ij} = 0$  for  $i < j$ .

A  $4 \times 4$  upper triangular matrix is given as follows:

$$\begin{bmatrix} * & 0 & 0 & 0 \\ * & * & 0 & 0 \\ * & * & * & 0 \\ * & * & * & * \end{bmatrix}$$

\* is arbitrary number

Note: the diagonal entries could be zero for upper triangular matrix and lower triangular matrix.

In lecture 2, we talked about for a square matrix with good property (here good property actually means that in every step of row reduction, the diagonal entry always be nonzero and without row exchange), a series of elementary row operation type III can be used to transform this square matrix to upper triangular form.

Recall: Illustration of the procedure for  $4 \times 4$  matrix without row exchange and the diagonal entries are all nonzero:

Using elementary row operation type III:

$$\text{Step 1: } A \triangleq \begin{bmatrix} * & * & * & * \\ * & * & * & * \\ * & * & * & * \\ * & * & * & * \end{bmatrix} \rightarrow \begin{bmatrix} \# & * & * & * \\ 0 & * & * & * \\ 0 & * & * & * \\ 0 & * & * & * \end{bmatrix}$$

$$\text{Step 2: } \begin{bmatrix} \# & * & * & * \\ 0 & * & * & * \\ 0 & * & * & * \\ 0 & * & * & * \end{bmatrix} \rightarrow \begin{bmatrix} \# & * & * & * \\ 0 & \# & * & * \\ 0 & 0 & * & * \\ 0 & 0 & * & * \end{bmatrix}$$

$$\text{Step 3: } \begin{bmatrix} \# & * & * & * \\ 0 & \# & * & * \\ 0 & 0 & * & * \\ 0 & 0 & * & * \end{bmatrix} \rightarrow \begin{bmatrix} \# & * & * & * \\ 0 & \# & * & * \\ 0 & 0 & \# & * \\ 0 & 0 & 0 & \# \end{bmatrix} \triangleq U$$

$\#$  is nonzero number,  $*$  is arbitrary number

For  $n \times n$  matrix  $A$  with good property, one can use a series of elementary row operation type III  $op_1, \dots, op_k$  (the corresponding elementary matrices are  $E_1, \dots, E_k$ ) to transform it into an upper triangular form  $U$ .

Suppose  $A \xrightarrow{op_1} A_1 \xrightarrow{op_2} A_2 \xrightarrow{op_3} \dots \xrightarrow{op_k} A_k = U$ .

By using the properties of elementary matrices, one has

$$E_1 A = A_1, E_2 A_1 = A_2, \dots, E_k A_{k-1} = A_k = U,$$

$$\text{then } E_k E_{k-1} \dots E_1 A = U$$

Thus,  $A = E_1^{-1} E_2^{-1} \dots E_k^{-1} U = LU$  where  $L = E_1^{-1} E_2^{-1} \dots E_k^{-1}$  is a lower triangular matrix.

This is because that  $E_i (i = 1, \dots, k)$  are lower triangular matrices,  $E_i^{-1} (i = 1, \dots, k)$  are also lower triangular matrices, the production  $E_1^{-1} E_2^{-1} \dots E_k^{-1}$  is also a lower triangular matrix.

$A = LU$  is called the **LU-decomposition**.

# Recall Gaussian elimination to obtain upper triangular form

**Example 7.2** Take the invertible matrix

$$A = \begin{bmatrix} 2 & 4 & 2 \\ 1 & 5 & 2 \\ 4 & -1 & 9 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 4 & 2 \\ 1 & 5 & 2 \\ 4 & -1 & 9 \end{bmatrix} \xrightarrow{\substack{R_2 \rightarrow -\frac{1}{2}R_1 + R_2 \\ R_3 \rightarrow -2R_1 + R_3}} \begin{bmatrix} 2 & 4 & 2 \\ 0 & \boxed{3} & 1 \\ 0 & -9 & 5 \end{bmatrix} \xrightarrow{R_3 \rightarrow -(-3)R_2 + R_3} \begin{bmatrix} 2 & 4 & 2 \\ 0 & \boxed{3} & 1 \\ 0 & 0 & 8 \end{bmatrix}$$

$$L_2 L_1 A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -2 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ -\frac{1}{2} & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 2 & 4 & 2 \\ 1 & 5 & 2 \\ 4 & -1 & 9 \end{bmatrix} = \begin{bmatrix} 2 & 4 & 2 \\ 0 & 3 & 1 \\ 0 & -9 & 5 \end{bmatrix}$$

$$L_3(L_2 L_1 A) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 3 & 1 \end{bmatrix} \begin{bmatrix} 2 & 4 & 2 \\ 0 & 3 & 1 \\ 0 & -9 & 5 \end{bmatrix} = \begin{bmatrix} 2 & 4 & 2 \\ 0 & 3 & 1 \\ 0 & 0 & 8 \end{bmatrix} = U$$

where  $U$  is the upper triangular matrix and  $L_3(L_2 L_1 A) = U$ .



$$\begin{aligned}
A &= L_1^{-1} L_2^{-1} L_3^{-1} \begin{bmatrix} 2 & 4 & 2 \\ 0 & 3 & 1 \\ 0 & 0 & 8 \end{bmatrix} \\
&= \begin{bmatrix} 1 & 0 & 0 \\ -\frac{1}{2} & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}^{-1} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -2 & 0 & 1 \end{bmatrix}^{-1} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 3 & 1 \end{bmatrix}^{-1} \begin{bmatrix} 2 & 4 & 2 \\ 0 & 3 & 1 \\ 0 & 0 & 8 \end{bmatrix} \\
&= \begin{bmatrix} 1 & 0 & 0 \\ \frac{1}{2} & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 2 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -3 & 1 \end{bmatrix} \begin{bmatrix} 2 & 4 & 2 \\ 0 & 3 & 1 \\ 0 & 0 & 8 \end{bmatrix} \\
&= \begin{bmatrix} 1 & 0 & 0 \\ \frac{1}{2} & 1 & 0 \\ 2 & -3 & 1 \end{bmatrix} \begin{bmatrix} 2 & 4 & 2 \\ 0 & 3 & 1 \\ 0 & 0 & 8 \end{bmatrix} \\
&= LU
\end{aligned}$$

$L$  is the lower triangular matrix.

$A = LU$  is called the **LU decomposition**.

Check the lower triangular entries of  $L$ , together with the elementary row operations. What do you observe?

$$L = \begin{bmatrix} 1 & 0 & 0 \\ \frac{1}{2} & 1 & 0 \\ 2 & -3 & 1 \end{bmatrix}$$

The entries below the diagonal of the unit lower triangular matrix  $L$  are **the multipliers** during the Gaussian Elimination process.

Do we really need to calculate  $L$  through finding the inverse of elementary matrices? **NO!**

When using the elementary row operations to transform  $A$  to an upper triangular form, we can obtain  $L$  simultaneously.

For the example 7.2

$$A = \begin{bmatrix} \boxed{2} & 4 & 2 \\ 1 & 5 & 2 \\ 4 & -1 & 9 \end{bmatrix}$$

Start with  $L = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

**Step 1:**  $\boxed{2}$  is the first pivot corresponding to elimination of first variable, now set the entries in first column of  $L$  below the number 1 equal to multipliers during the elimination in the first step. Multipliers are  $1/2$  and  $2$  for second row and third row, respectively.

Update  $L$ :  $L = \begin{bmatrix} 1 & 0 & 0 \\ \frac{1}{2} & 1 & 0 \\ 2 & 0 & 1 \end{bmatrix}$

**Step 2:** Perform elementary row operations for first column

$$\begin{bmatrix} 2 & 4 & 2 \\ 1 & 5 & 2 \\ 4 & -1 & 9 \end{bmatrix} \xrightarrow[\substack{R_2 \rightarrow -\frac{1}{2}R_1 + R_2 \\ R_3 \rightarrow -2R_1 + R_3}]{\phantom{R_2 \rightarrow -\frac{1}{2}R_1 + R_2}} \begin{bmatrix} 2 & 4 & 2 \\ 0 & \boxed{3} & 1 \\ 0 & -9 & 5 \end{bmatrix}$$

$\boxed{3}$  is the second pivot corresponding to elimination of second variable, set the entries in second column of  $L$  below the number 1 equal to the multiplier during the elimination in the second step. Multiplier is  $-3$  for the third row.

Update  $L$ :  $L = \begin{bmatrix} 1 & 0 & 0 \\ \frac{1}{2} & 1 & 0 \\ 2 & -3 & 1 \end{bmatrix}$

Now performing elementary row operations for second column to obtain the upper triangular form

$$\begin{bmatrix} 2 & 4 & 2 \\ 0 & \boxed{3} & 1 \\ 0 & -9 & 5 \end{bmatrix} \xrightarrow{R_3 \rightarrow 3R_2 + R_3} \begin{bmatrix} 2 & 4 & 2 \\ 0 & \boxed{3} & 1 \\ 0 & 0 & 8 \end{bmatrix} = U$$

One can check that

$$A = \begin{bmatrix} 1 & 0 & 0 \\ \frac{1}{2} & 1 & 0 \\ 2 & -3 & 1 \end{bmatrix} \begin{bmatrix} 2 & 4 & 2 \\ 0 & 3 & 1 \\ 0 & 0 & 8 \end{bmatrix} = LU$$

Keep tracking the multipliers during the Gaussian Elimination process, one can obtain the  $LU$  decomposition simultaneously.

# Application of LU decomposition to solve linear system

**Example 7.3** Find the solution of following system

$$A = \begin{bmatrix} 2 & 4 & 2 \\ 1 & 5 & 2 \\ 4 & -1 & 9 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} -4 \\ -7 \\ 15 \end{bmatrix}$$

$A = LU$ , thus

$$\begin{bmatrix} 1 & 0 & 0 \\ \frac{1}{2} & 1 & 0 \\ 2 & -3 & 1 \end{bmatrix} \begin{bmatrix} 2 & 4 & 2 \\ 0 & 3 & 1 \\ 0 & 0 & 8 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} -4 \\ -7 \\ 15 \end{bmatrix}$$

For  $A\mathbf{x} = LU\mathbf{x} = \mathbf{b}$ , one can first solve  $L\mathbf{y} = \mathbf{b}$  using **forward substitution**, then solve  $U\mathbf{x} = \mathbf{y}$  by **backward substitution**.

First solve the linear system  $L\mathbf{y} = \mathbf{b}$  by using **forward substitution**:

$$\begin{bmatrix} 1 & 0 & 0 \\ \frac{1}{2} & 1 & 0 \\ 2 & -3 & 1 \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = \begin{bmatrix} -4 \\ -7 \\ 15 \end{bmatrix} \Rightarrow \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = \begin{bmatrix} -4 \\ -5 \\ 8 \end{bmatrix}$$

Then solve the linear system  $U\mathbf{x} = \mathbf{y}$  by using **backward substitution**:

$$\begin{bmatrix} 2 & 4 & 2 \\ 0 & 3 & 1 \\ 0 & 0 & 8 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} \Rightarrow \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 1 \\ -2 \\ 1 \end{bmatrix}$$

In software, Matlab's command  $\mathbf{x} = A \setminus \mathbf{b}$  use LU-decomposition, along with forward and backward substitution to solve the linear system when the matrix  $A$  is nonsingular.



For general square matrices, we have the following result.

**Theorem 7.4 (LU decomposition for a square matrix)**

If  $A$  is a square matrix, then there exists a permutation matrix  $P$  such that  $PA$  has the  $LU$  decomposition, i.e.,  $PA = LU$ , where  $L$  is a unit lower triangular matrix (a lower triangular matrix whose diagonal entries are all 1's),  $U$  is an upper triangular matrix.

Proof is skipped.

**Remark:** For  $PA = LU$ , when  $U$  is an upper triangular matrix with nonzero diagonal entries, then  $A$  is nonsingular. However, if one of the diagonal entries of  $U$  is zero, then  $A$  is singular.

**Example 7.5** Take the nonsingular matrix

$$A = \begin{bmatrix} 1 & 1 & 2 \\ 2 & 2 & 3 \\ 3 & 5 & 9 \end{bmatrix}$$

$$L_2 L_1 A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -3 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 2 \\ 2 & 2 & 3 \\ 3 & 5 & 9 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 2 \\ 0 & 0 & -1 \\ 0 & 2 & 3 \end{bmatrix}$$

We find row exchange is needed at this stage! Let  $P = E_{R_2 R_3}$ , then

$$P L_2 L_1 A = \begin{bmatrix} 1 & 1 & 2 \\ 0 & 2 & 3 \\ 0 & 0 & -1 \end{bmatrix} = U$$

But this does not yields the form  $PA = LU$ .

How to get the the form  $PA = LU$ ?

Idea: do all the row exchanges first.

Starting from  $A$ , if we do the row exchange for row 2 and row 3 first, then

$$PA = \begin{bmatrix} 1 & 1 & 2 \\ 3 & 5 & 9 \\ 2 & 2 & 3 \end{bmatrix}$$

Now start with  $L = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ ,  $\boxed{1}$  is the first pivot of  $PA = \begin{bmatrix} \boxed{1} & 1 & 2 \\ 3 & 5 & 9 \\ 2 & 2 & 3 \end{bmatrix}$

Set  $L = \begin{bmatrix} 1 & 0 & 0 \\ 3 & 1 & 0 \\ 2 & 0 & 1 \end{bmatrix}$  and perform elementary row operations for  $PA$ , one can get

$$\begin{bmatrix} \boxed{1} & 1 & 2 \\ 3 & 5 & 9 \\ 2 & 2 & 3 \end{bmatrix} \xrightarrow[\substack{R_2 \rightarrow -3R_1 + R_2 \\ R_3 \rightarrow -2R_1 + R_3}]{\phantom{R_2 \rightarrow -3R_1 + R_2}} \begin{bmatrix} 1 & 1 & 2 \\ 0 & \boxed{2} & 3 \\ 0 & 0 & -1 \end{bmatrix} = U$$

Now one can check that

$$PA = L = \begin{bmatrix} 1 & 0 & 0 \\ 3 & 1 & 0 \\ 2 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 2 \\ 0 & 2 & 3 \\ 0 & 0 & -1 \end{bmatrix} = LU$$

Note that here  $A$  is nonsingular, the diagonal entries of  $U$  are nonzero,  $U$  can be further decomposed into

$$U = \begin{bmatrix} 1 & 1 & 2 \\ 0 & 2 & 3 \\ 0 & 0 & -1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & -1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 2 \\ 0 & 1 & \frac{3}{2} \\ 0 & 0 & 1 \end{bmatrix} = D\hat{U}$$

where  $D$  is a diagonal matrix with nonzero diagonal entries,  $\hat{U}$  is a unit upper triangular matrix (an upper triangular matrix whose diagonal entries are all 1's).

Thus

$$PA = LU = LD\hat{U}$$

This is the  $LDU$  decomposition.

### **Theorem 7.6 (LDU decomposition for a nonsingular matrix)**

If  $A$  is nonsingular, then there exists a permutation matrix  $P$  such that  $PA$  has the  $LDU$  decomposition, i.e.,  $PA = LDU$ , where  $L$  is a unit lower triangular matrix,  $D$  is a diagonal matrix whose diagonal entries are nonzero,  $U$  is a unit upper triangular matrix.