
Sample Midterm

DURATION OF EXAMINATION: 2 hours

Your exam sheet shall include 6 **problems**. If not, notify the instructors.

1. **(26 pt) Solving a Linear System of Equations**

Consider the linear system:

$$2x_1 - 3x_2 + x_3 + x_4 = 1 \tag{1}$$

$$-x_2 + x_3 + x_4 = 0 \tag{2}$$

$$x_1 + 2x_2 + 4x_3 - 2x_4 = -2. \tag{3}$$

- (a) (3 pt) Write out the *coefficient matrix* \mathbf{A} of this system.
- (b) (10 pt) Write out the *augmented matrix* for this system and calculate its *row-reduced echelon form*.
- (c) (7 pt) Write out the *complete set* of solutions. Note: You shall write in the form of $\mathbf{x}_p + \alpha_1 \mathbf{v}_1 + \dots \alpha_k \mathbf{v}_k$, where $\mathbf{x}_p, \mathbf{v}_1, \dots, \mathbf{v}_k$ are vectors that you need to find out.
- (d) (3 pt) What is the *rank* of the coefficient matrix \mathbf{A} ? Justify your answer.
- (e) (3 pt) Find a *basis* of the *null space* of matrix \mathbf{A} .

2. **(12 pt) Question Answering**

- (a) (3 pt) A 6th grade student Xiao Ming told you that she/he has learned from the school about how to get a specific solution of a 2×2 linear system of equations by substituting variables. Can you give an example of a 2×2 system that Xiao Ming may not be able to solve? Note: Your example should be a specific 2×2 linear system.
- (b) (4 pt) If a matrix A has full row rank, then what can we say about the solution set of $Ax = b$? If A has full column rank, what can we say about the solution set of $Ax = b$?
- (c) (3 pt) Write one 2×2 invertible matrix that is both upper triangular and symmetric.
- (d) (2 pt) What theorem makes the definition of “dimension” valid (i.e. well defined)? Choose one from the following choices.
 - (A) The column space and row space of a matrix have the same dimension.
 - (B) Any two bases of a linear space has the same number of elements.
 - (C) A square matrix has full rank iff the columns are linearly independent.
 - (D) The sum of the rank and the nullity of a matrix equals the number of columns.

3. **(16 pt) Matrix Transpose, Inverse and Multiplication**

- (a) (6 pt) Find the inverse of $\begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$ and justify what you found is indeed its inverse.
- (b) (4 pt) Let

$$A = \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix} \text{ and } B = \begin{pmatrix} 1 & 0 \\ 3 & 1 \end{pmatrix}.$$

Compute $(AB)^\top$ and $B^\top A^\top$.

- (c) (6 pt) Suppose A and B are 2×2 matrices. Suppose $A = [\mathbf{a}_1, \mathbf{a}_2] = \begin{bmatrix} \mathbf{u}_1^\top \\ \mathbf{u}_2^\top \end{bmatrix}$, $B = [\mathbf{b}_1, \mathbf{b}_2] = \begin{bmatrix} \mathbf{v}_1^\top \\ \mathbf{v}_2^\top \end{bmatrix}$. Express $B^\top A^\top$ at least in two forms using the vectors $\mathbf{a}_i, \mathbf{u}_i, \mathbf{b}_i, \mathbf{v}_i, i = 1, 2$.

Remark: Your expressions can NOT contain scalars like a_{ij} or b_{ij} . Your expressions can NOT contain matrices like A or B . Your expressions can ONLY contain vectors $\mathbf{a}_i, \mathbf{u}_i, \mathbf{b}_i, \mathbf{v}_i, i = 1, 2$.

4. (16 pt) True or False

For each question, you do NOT need to justify. (Only writing T or F is enough)

- (a) Any invertible matrix can be written as the product of elementary matrices.
- (b) If A and AB are invertible where A and B are both square matrices with the same size, then B is also invertible.
- (c) Any square matrix can be written as the product of a lower triangular matrix and an upper triangular matrix.
- (d) The time complexity of conducting Gaussian elimination for an $n \times n$ matrix is $O(n^2)$.
- (e) If the columns of a **square** matrix A are linearly independent, then the columns of A^2 are also linearly independent.
- (f) The three vectors $[0, 0, 0]$, $[1, 4, 5]$ and $[1, 2, 3]$ are linearly dependent.
- (g) Suppose $A \in \mathbb{R}^{n \times n}$ and $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$. If $A\mathbf{x} = A\mathbf{y}$, then $\mathbf{x} = \mathbf{y}$.
- (h) $\text{span}(\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}) = \text{span}(\{\mathbf{u}, \mathbf{u} - \mathbf{v} + \mathbf{w}, 2\mathbf{u} - \mathbf{v} + \mathbf{w}\})$ for any vectors $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathbb{R}^3$.

5. (15 pt) Linear Space, Column Space and Null Space

- (a) (8 pt) Suppose A is a matrix. Prove: the null space of A is a linear space.
Remark: We talked about this result in class. You cannot directly quote the theorem in class. Instead, you need to provide a proof of this result.
- (b) (7 pt) Show that if the product of two matrices $A \in \mathbb{R}^{m \times n}$ and $B \in \mathbb{R}^{n \times k}$ is a zero matrix $0 \in \mathbb{R}^{m \times k}$, i.e., $AB = 0$, then the column space of A is contained in the null space of B , i.e., $C(B) \subseteq N(A)$.

6. (15 pt) Linear Dependence and Rank

In all subproblems, I_k is the $k \times k$ identity matrix.

- (a) (5 pt) Prove: All rows of the matrix $[A, I_k, U] \in \mathbb{R}^{k \times (m+k+n)}$ are linearly independent, where $A \in \mathbb{R}^{k \times m}$ and $U \in \mathbb{R}^{k \times n}$ are two arbitrary matrices.
- (b) (5 pt) Suppose $A \in \mathbb{R}^{k \times k}$, and define the matrix $M = \begin{bmatrix} I_k & 0 \\ 0 & A \end{bmatrix} \in \mathbb{R}^{(2k) \times (2k)}$. Prove that $\text{rank}(M) = k + \text{rank}(A)$.
- (c) (5 pt) Suppose $A, B, C \in \mathbb{R}^{k \times k}$, and the matrix $Q = \begin{bmatrix} I_k & B \\ C & A \end{bmatrix} \in \mathbb{R}^{(2k) \times (2k)}$. Does $\text{rank}(Q) = k + \text{rank}(A)$ always hold? If yes, provide a proof. If not, provide an alternative expression of $\text{rank}(Q)$ in terms of k and $\text{rank}(f(A, B, C))$ where f is a certain function of A, B, C that you need to figure out, and justify the expression.

The following are some additional sample questions for the midterm.

7. Given an m by n matrix A with rank s , take s columns from A and s rows from A . Their intersection forms an $s \times s$ matrix. Show that the intersection is non-singular if these selected rows and columns are independent.
8. Let $A = CR$, where C is an m by r matrix and R is r by n . Prove that if the rank of A is r , then C has full column rank and R has full row rank
9. Let $u, v \in \mathbb{R}^n$, assume $u^T v = 0$, prove that $(I + uv^T)^{-1} = I - uv^T$.
10. Consider block elimination matrices, show that

$$\left[\begin{array}{c|ccc} I_k & & & & \\ & 1 & & & \\ & l & & I_{n-k-1} & \end{array} \right] \left[\begin{array}{c|c} I_k & u \\ & I_{n-k} \end{array} \right] = \left[\begin{array}{c|ccc} I_k & & u & & \\ & 1 & & & \\ & l & & I_{n-k-1} & \end{array} \right] \triangleq J.$$

Also, for $A = \left[\begin{array}{c|cc} u_1 & u & * \\ & \alpha & \\ & l & * \end{array} \right]$, find J of the above form s.t. $JA = \left[\begin{array}{c|cc} u_1 & 0 & * \\ & \alpha & \\ & 0 & * \end{array} \right]$

11. Compute $PA = LU$ for $A = \begin{bmatrix} 0 & 1 & 2 \\ 1 & 2 & 3 \\ 1 & 1 & 2 \end{bmatrix}$