

DSBA Linear Algebra HW5

Kirill Korolev 203-1

17th of October, 2020

1. Find the number of inversions in the following ordered line:

$$(9, 1, 3, 7, 2, 8, 5, 4, 6)$$

Inversion is such pair $(f(i), f(j))$ of permutation f for which valid $i < j$ and $f(i) > f(j)$. Basically, for one-line notation number of inversions means number of non-sorted pairs in ascending order.

Let's list these pairs for each number:

9: 1, 3, 7, 2, 8, 5, 4, 6

1: none

3: 2

7: 2, 5, 4, 6

2: none

8: 5, 4, 6

5: 4

4: none

6: none

Overall there are $8 + 1 + 4 + 3 + 1 = 17$ inversions.

2. Let $f = \begin{pmatrix} 2 & 3 & 5 & 1 & 4 \\ 4 & 1 & 2 & 3 & 5 \end{pmatrix}$ and $g = \begin{pmatrix} 1 & 5 & 3 & 4 & 2 \\ 5 & 4 & 1 & 2 & 3 \end{pmatrix}$

- (a) cycle notation for f and g

Each permutation can be represented as product of independent cycles whereas a cycle is a list of (a_1, a_2, \dots, a_k) with mapping $a_1 \rightarrow a_2, a_2 \rightarrow a_3, \dots, a_k \rightarrow a_1$.

$$f = (245)(13) \text{ and } g = (15423)$$

- (b) $f \circ g$ (using two-line notation)

$$\begin{pmatrix} 1 & 5 & 3 & 4 & 2 \\ 5 & 4 & 1 & 2 & 3 \\ \hline 5 & 4 & 1 & 2 & 3 \\ 2 & 5 & 3 & 4 & 1 \end{pmatrix} \Rightarrow f \circ g = \begin{pmatrix} 1 & 5 & 3 & 4 & 2 \\ 2 & 5 & 3 & 4 & 1 \end{pmatrix}$$

(c) $g \circ f$ (using cycle notation)

$$g \circ f = (15423)(245)(13) = (2)(4)(53)(1) \Rightarrow$$

$$g \circ f = \begin{pmatrix} 2 & 3 & 5 & 1 & 4 \\ 2 & 5 & 3 & 1 & 4 \end{pmatrix}$$

(d) $f^{-1} \circ g^{-1}$

Obtain inverse permutations by swapping rows in two-line notation:

$$f^{-1} = \begin{pmatrix} 4 & 1 & 2 & 3 & 5 \\ 2 & 3 & 5 & 1 & 4 \end{pmatrix} \quad g^{-1} = \begin{pmatrix} 5 & 4 & 1 & 2 & 3 \\ 1 & 5 & 3 & 4 & 2 \end{pmatrix}$$

$$\begin{pmatrix} 5 & 4 & 1 & 2 & 3 \\ 1 & 5 & 3 & 4 & 2 \\ 1 & 5 & 3 & 4 & 2 \\ 3 & 4 & 1 & 2 & 5 \end{pmatrix} \Rightarrow f^{-1} \circ g^{-1} = \begin{pmatrix} 5 & 4 & 1 & 2 & 3 \\ 3 & 4 & 1 & 2 & 5 \end{pmatrix}$$

(e) $sgn(f)$

Instead of calculating $sgn(f)$ by definition we can use the fact, which was proved on seminar, that $sgn(f) = (-1)^{n-d}$ where n is the number of elements in permutation and d is the number of independent cycles, therefore $sgn(f) = (-1)^{5-2} = -1$

(f) $sgn(g)$

$$sgn(g) = (-1)^{5-1} = 1$$

(g) Verify that $sgn(f) \circ sgn(g) = sgn(f \circ g)$

Remember that $f \circ g = \begin{pmatrix} 1 & 5 & 3 & 4 & 2 \\ 2 & 5 & 3 & 4 & 1 \end{pmatrix}$ and we can factorize it on independent cycles like this: $f \circ g = (12)(5)(3)(4)$. Therefore $sgn(f \circ g) = (-1)^{5-4} = -1 = -1 \cdot 1 = sgn(f) \cdot sgn(g)$

3. Let

$$f = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ 4 & 5 & 7 & 1 & 3 & 2 & 6 \end{pmatrix}$$

$$g = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 & 11 & 12 & 13 \\ 8 & 11 & 7 & 12 & 9 & 1 & 6 & 3 & 2 & 13 & 5 & 4 & 10 \end{pmatrix}$$

(a) Find f^{2020}

Split f on cycles $\Rightarrow f = (14)(25376)$, then:

$$f^{2020} = (14)^{2020}(25376)^{2020}$$

Let's use the fact that the order of a cycle of length n , in other words the minimum power to raise a permutation to obtain an identity

permutation, is also n . It is easy to see that we need to scroll a cycle n times to get the same permutation. Therefore, $(14)^{2k} = id$ and $(25376)^{5k} = id$. 2020 is divisible by 2 and 5, hence after putting everything together:

$$f^{2020} = (14)^{2020}(25376)^{2020} = id \cdot id = id$$

(b) Find g^{2077}

Split g on cycles $\Rightarrow g = (18376)(2, 11, 5, 9)(10, 13)(4, 12)$, then:

$$\begin{aligned} g^{2077} &= (18376)^{2077}(2, 11, 5, 9)^{2077}(10, 13)^{2077}(4, 12)^{2077} \\ 2077 &\equiv 2(5) \quad 2077 \equiv 1(4) \quad 2077 \equiv 1(2) \Rightarrow \\ &(18376)^{2077}(2, 11, 5, 9)^{2077}(10, 13)^{2077}(4, 12)^{2077} = (18376)^2(2, 11, 5, 9)(10, 13)(4, 12) = \\ &(13687)(2, 11, 5, 9)(10, 13)(4, 12) \\ g^{2077} &= \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 & 11 & 12 & 13 \\ 3 & 11 & 6 & 12 & 9 & 8 & 1 & 7 & 2 & 13 & 5 & 4 & 10 \end{pmatrix} \end{aligned}$$

4. Find all $f \in S_{2020}$ such that

$$f^2 = (1, 2, 3, \dots, 2019, 2020)$$

where $(1, 2, 3, \dots, 2019, 2020)$ is a long cycle of length 2020.

Let's denote this cycle as σ_{2020} and use lemma about parity of composition:

$$sgn(f \circ g) = sgn(f) \circ sgn(g)$$

We can see that there is no way to find such permutation f , because $sgn(\sigma_{2020}) = (-1)^{2020-1} = -1 \neq sgn^2(f)$