

Double Integrals in Polar Coordinates

SUGGESTED REFERENCE MATERIAL:


As you work through the problems listed below, you should reference Chapter 14.3 of the recommended textbook (or the equivalent chapter in your alternative textbook/online resource) and your lecture notes.

EXPECTED SKILLS:

- Be able to convert rectangular double integrals to polar double integrals, including converting the limits of integration, the function to be integrated, and the differential dA to $r dr d\theta$.

PRACTICE PROBLEMS:

1. Consider the region R shown below which is enclosed by $x^2 + y^2 = 1$, $x^2 + y^2 = 4$, $y = x$ and the x axis.




Fill in the missing limits of integration: $\iint_R f(x, y) dA = \int_{\square}^{\square} \int_{\square}^{\square} f(r, \theta) r dr d\theta$.

For problems 2-6, evaluate the iterated integral by converting to polar coordinates.


2. $\int_0^4 \int_0^{\sqrt{16-x^2}} \sqrt{x^2 + y^2} dy dx$
3. $\int_0^{3/\sqrt{2}} \int_x^{\sqrt{9-x^2}} (x^2 + y^2)^2 dy dx$
4. $\int_0^2 \int_0^{\sqrt{2x-x^2}} xy dy dx$

5. Evaluate $\iint_R (x - y) dA$ where $R = \{(x, y) : 4 \leq x^2 + y^2 \leq 16 \text{ and } y \leq x\}$
6. Evaluate $\iint_R e^{-(x^2+y^2)} dA$ where $R = \{(x, y) : x^2 + y^2 \leq 3 \text{ and } 0 \leq y \leq \sqrt{3}x\}$
7. Use a double integral in polar coordinates to calculate the area of the region which is inside of the cardioid $r = 2 + 2 \cos \theta$ and outside of the circle $r = 3$.
8. Use a double integral in polar coordinates to calculate the area of the region which is common to both circles $r = 3 \sin \theta$ and $r = \sqrt{3} \cos \theta$.
9. Consider the top which is bounded above by $z = \sqrt{4 - x^2 - y^2}$ and bounded below by $z = \sqrt{x^2 + y^2}$, as shown below.



Use a double integral in polar coordinates to calculate the volume of the top.

10. Consider the surfaces $x^2 + y^2 + z^2 = 16$ and $x^2 + y^2 = 4$, shown below.



Calculate the volume of the solid which is inside of $x^2 + y^2 + z^2 = 16$ but outside of $x^2 + y^2 = 4$.

11. Calculate the volume of the solid which is bounded above by $z = 9 - x^2 - y^2$, bounded below by $z = 0$, and contained within $x^2 - 3x + y^2 = 0$.

