Improper Integrals

SUGGESTED REFERENCE MATERIAL:

As you work through the problems listed below, you should reference Chapter 7.8 of the recommended textbook (or the equivalent chapter in your alternative textbook/online resource) and your lecture notes.

EXPECTED SKILLS:

- Given an improper integral, which either has an infinite interval of integration or an infinite discontinuity, be able to evaluate it using a limit.
- Know how to determine if such an integral converges (and if so, what it converges to) or diverges.

PRACTICE PROBLEMS:

For problems 1-13, evaluate each improper integral or show that it diverges.

1.
$$\int_{1}^{\infty} \frac{1}{\sqrt{x}} dx$$

2.
$$\int_{-\infty}^{3} \frac{3x}{x^2 + 1} dx$$

$$3. \int_{1}^{\infty} e^{-x} dx$$

$$4. \int_{1}^{\infty} xe^{-3x^2} dx$$

5.
$$\int_0^4 \frac{1}{x^{2/3}} dx$$

6.
$$\int_{4}^{\infty} \frac{1}{(x-2)^3} \, dx$$

7.
$$\int_{2}^{6} \frac{1}{\sqrt{x-2}} dx$$

$$8. \int_0^2 \frac{2}{\sqrt{4-x^2}} \, dx$$

9.
$$\int_0^\infty \frac{1}{x^2 + 4x + 5} dx$$
 (Hint: Complete the square)


$$10. \int_0^{\frac{\pi}{2}} \frac{\sin x}{\sqrt[3]{\cos x}} \, dx$$

11.
$$\int_0^{\frac{\pi}{2}} \sqrt{\tan x} \sec^2 x \, dx$$

12.
$$\int_0^9 \frac{1}{\sqrt[3]{(x-1)^2}} \, dx$$

$$13. \int_0^1 \frac{1}{x \ln x} \, dx$$

- 14. Find the value of the constant k so that $\int_{-\infty}^{\infty} \frac{k}{1+x^2} dx = 1.$
- 15. Compute the exact area between the graph of $y = \frac{4}{x^2 1}$ and the x-axis for $x \ge 7$.
- 16. Consider Gabriel's Horn (shown below) which is formed by revolving the curve $y = \frac{1}{x}$ on $[1, \infty)$ around the x-axis.



Show that the volume within this horn is finite.

Note: It can be shown the surface area of this horn is infinite. Thus, it appears that the horn can be filled with a finite amount of paint; but, there is not enough to paint the inside of the surface for a coating of uniform thickness. This is called **The Paradox of Gabriel's Horn**.

17. Determine the values of the constant p for which the following integral will converge and those for which it will diverge.

2

$$\int_{1}^{\infty} \frac{1}{x^{p}} dx$$

(Hint: Consider two cases – when p = 1 and when $p \neq 1$.)

18. Determine the values of the constant p for which the following integral will converge and those for which it will diverge.

$$\int_0^1 \frac{1}{x^p} \, dx$$

(Hint: Consider two cases – when p = 1 and when $p \neq 1$.)

- 19. Consider the Gamma Function: $\Gamma(\alpha) = \int_0^\infty x^{\alpha-1} e^{-x} dx$, which is defined for $\alpha > 0$.
 - (a) Compute $\Gamma(1)$.
 - (b) Compute $\Gamma(2)$.
 - (c) Compute $\Gamma(3)$.
- 20. If f(x) is a continuous for $x \ge 0$, the **Laplace Transform** of f(x) is given by:

$$\mathcal{L}\left\{f(x)\right\}(s) = \int_0^\infty f(x)e^{-sx} dx$$

and the domain of $\mathcal{L}\{f(x)\}(s)$ is the set consisting of all numbers s for which the integral converges. Laplace Transforms are useful for solving differential equations

- (a) Compute the Laplace Transform of f(x) = 1 and state its domain.
- (b) Compute the Laplace Transform of $f(x) = e^x$ and state its domain.
- (c) Compute the Laplace Transform of f(x) = x and state its domain.
- 21. **Definition:** In probability theory, the **probability density function (pdf)** of a random variable X is a function f(x) from which we can compute the probability that X lies in the interval [a, b] as follows:

$$P(a \le X \le b) = \int_a^b f(x) \, dx$$

And, in order for f(x) to be a valid pdf, it must satisfy the following:

- $f(x) \ge 0$ for all values of x
- $\bullet \int_{-\infty}^{\infty} f(x) \, dx = 1$
- (a) Verify that $f(x) = \begin{cases} 2e^{-2x} & x \ge 0 \\ 0 & x < 0 \end{cases}$ is a valid probability density function.
- (b) Using the density function in part a, compute $P(0 \le X \le 1)$.

(c) **Definition:** The **cumulative distribution function (CDF)** for a continuous random variable X is defined as:

$$F(t) = P(X \le t) = \int_{-\infty}^{t} f(x) dx$$

This CDF describes the accumulation of probability up to the real number t. Compute the CDF for the random variable X which has the density function from part a.