

# Second-Order Linear Homogeneous ODE's

---

## SUGGESTED REFERENCE MATERIAL:

As you work through the problems listed below, you should reference your lecture notes and the relevant chapters in a textbook/online resource.

## EXPECTED SKILLS:

- Show that two functions are linearly independent.
- Be able to solve second-order linear ODE's with constant coefficients by using the appropriate auxiliary equations.
- Be able to solve initial value problems and boundary value problems involving second-order linear ODE's with constant coefficients.

## PRACTICE PROBLEMS:

1. Suppose that  $y_1(x)$  and  $y_2(x)$  are solutions to the second-order ODE

$$P(x)\frac{d^2y}{dx^2} + Q(x)\frac{dy}{dx} + R(x)y = 0.$$

Show that  $y(x) = c_1y_1(x) + c_2y_2(x)$  is also a solution, where  $c_1$  and  $c_2$  are arbitrary constants.

2. Consider the functions  $y_1 = e^{m_1x}$  and  $y_2 = e^{m_2x}$ , where  $m_1$  and  $m_2$  are arbitrary real numbers with  $m_1 \neq m_2$ . Show that the functions are linearly independent.
3. Consider the functions  $y_1 = e^{\alpha x} \cos(\beta x)$  and  $y_2 = e^{\alpha x} \sin(\beta x)$ , where  $\alpha$  and  $\beta$  are arbitrary real numbers. Show that the functions are linearly independent.

**For problems 4 – 10, solve the second-order differential equation. Express your answer as an explicit function of  $x$ .**

4.  $y'' + 5y' + 4y = 0$

5.  $25y = 10\frac{dy}{dx} - \frac{d^2y}{dx^2}$

6.  $2\frac{d^2y}{dx^2} + 5\frac{dy}{dx} - 3y = 0$

7.  $y'' + 2y' + 3y = 0$

8.  $9y'' + 12y' + 4y = 0$

9.  $2\frac{d^2y}{dx^2} - 6\frac{dy}{dx} + 17y = 0$
10.  $9y'' = 4y$
11. Solve the initial value problem.
 
$$\begin{cases} y'' + 5y' + 4y = 0 \\ y(0) = 5 \\ y'(0) = 7 \end{cases}$$
12. Consider a curve  $y = y(x)$  that goes through the point  $\left(\frac{\pi}{\sqrt{2}}, e^{-\pi/\sqrt{2}}\right)$ , has a horizontal tangent line at that point, and satisfies the ODE  $y'' + 2y' + 3y = 0$ .
  - (a) Set up an initial value problem (IVP) whose solution is  $y(x)$ .
  - (b) Solve the IVP from part (a).
13. Consider a curve  $y = y(x)$  that has a  $y$ -intercept of  $(0, 1)$ , has a tangent line at the  $y$ -intercept that is perpendicular to the line  $y = -\frac{1}{2}x + 999$ , and satisfies the ODE  $9y'' = 4y$ .
  - (a) Set up an initial value problem (IVP) whose solution is  $y(x)$ .
  - (b) Solve the IVP from part (a).
14. Consider a curve  $y = y(x)$  that goes through both the origin and the point  $(\ln 2, 64)$  and also satisfies the ODE  $25y = 10\frac{dy}{dx} - \frac{d^2y}{dx^2}$ .
  - (a) A boundary value problem, or BVP, is a problem that consists of a differential equation and information about two *different* points on the solution curve.  
[Whereas an initial value problem, or IVP, consists of a differential equation and initial conditions about a *single* point.]  
Set up a BVP whose solution is  $y(x)$ .
  - (b) Solve the BVP from part (a).
15. Suppose an object of mass  $M$  is attached to a spring, and the spring is allowed to hang at its natural length. Let  $x = x(t)$  represent the position of the object when the spring is stretched (or compressed)  $x$  units from its natural length, with  $x = 0$  corresponding to the spring's natural length. If we ignore any forces acting on the object except it's

weight and the restoring force of the spring, then based on Newton's Second Law and Hooke's Law it can be shown that

$$M \frac{d^2x}{dt^2} + kx = 0$$

where  $k$  is a some positive constant, called the spring constant.

- (a) Solve the above second-order ODE.
- (b) Suppose that an object of mass 3 kg is attached to a spring with spring constant 75 whose natural length is 1 meter. If the spring is stretched to a length of 1.5 meters and is released with zero initial velocity, find the position of the object at any time  $t$ . Describe the motion of the object over time.