

Multifractal and monofractal scaling in a laboratory MHD turbulence experiment

D.A. Schaffner¹ and M.R. Brown¹

¹*Swarthmore College, Swarthmore, PA, USA*

(Dated: May 27, 2015)

Both multifractal and monofractal scaling of structure function exponents is observed in the turbulent magnetic fluctuations of the Swarthmore Spheromak Experiment (SSx) plasma. Structure function and probability distribution function (PDF) analysis exhibits multifractal scaling exponents in low frequency, inertial range fluctuations of the turbulence but monofractal scaling in higher frequency, dissipation range fluctuations. The transition from multifractal to monofractal scaling occurs rapidly suggesting a dissipation mechanism that is insensitive to turbulent structure scale size. Structure functions and PDFs are presented for both temporal and spatial measurements. Variations in the magnetic helicity in the plasma are also shown to modify multifractal scaling characteristics of the inertial range, but do not affect the monofractal scaling of the dissipation range.

INTRODUCTION

This paper presents the results of a thorough intermittency analysis of the fluctuating magnetic fields in the Swarthmore Spheromak Experiment (SSX) plasma through the use of structure functions and probability distribution functions (PDFs) of increments of both temporal and spatial measurements. The primary observation from this analysis is that temporal regions of the magnetic fluctuation data are shown to be consistent with dissipation range turbulence [1] exhibiting structure function scaling that indicates self-similarity in the turbulence structure, whereas temporal regions for inertial range turbulence do not exhibit this self-similarity. These results are discussed in the context of multifractal versus monofractal structure function scaling behavior of turbulence [2–4], where a lack of self-similarity can be attributed to the multifractal nature of the structure function scaling. Similar distinctions between inertial range and dissipation range scaling are made in recent solar wind observations [5] suggesting that physical mechanisms underlying the scaling in each regime may be universal between solar wind and laboratory based MHD turbulence. Furthermore, the relatively fast transition from multifractal to monofractal scaling suggests a dissipation mechanism which has an absolute scale, such as the generation of current sheets at the ion inertial scale length [6, 7].

Trends in scaling with magnetic helicity injection are also explored. Previous work [8] demonstrated that increased injected helicity in the SSX plasma system resulted in an increased intermittency of the raw $dB(t)/dt$ signal. Results reported here show that the $B(t)$ formed from time integration of the raw $dB(t)/dt$ signal demonstrate this same trend. The scaling behavior of the structure functions differ with helicity depending on whether they are constructed from data in the dissipation versus the inertial range. Inertial range scaling appears to vary more widely with helicity than the dissipation range. This indicates that the effect of the helicity in-

jection on intermittency has an effect only on structures generated in the inertial range, or on the physical mechanism which does the generating. Conversely, the mechanism that governs intermittency in the dissipation range is not strongly affected by the overall helicity content of the plasma.

Structure function analysis has typically been utilized to extend spectral or correlation analysis of turbulent fluctuations to higher-order statistics, particularly in scenarios where Gaussian and self-similar properties appear to break down. Examined first in hydrodynamics [2, 9], the use of the technique has been expanded to magnetohydrodynamic (MHD) fluids including the solar wind velocity fluctuations [10], solar wind magnetic fluctuations [11, 12] and magnetospheric plasmas [13, 14]. The work has led to the development of models which attempt to reconcile the observation of intermittency—non-self-similar statistics in turbulence fluctuations—with the 1941 Kolomogorov turbulence model which implies a self-similar fluctuation structure, and which can be described in terms of a monofractal scaling exponent [2, 15]. Such multifractal models relax global scale invariance to a local scale invariance. Physically, these models describe modifications of how energy is distributed to smaller scales from larger scales either through differences in the space-filling nature of the turbulence, such as the random β -model [16] or through variations in the energy transfer rate between from larger to smaller scales such as the log-normal model [17], the p -model [18] and the She-Leveque model [19, 20]. Multifractal scaling models can make predictions for dissipation scaling, in particular addressing a reduction in scaling exponents by viscosity [21, 22]. Since these models were developed with isotropic and homogeneous turbulence of conventional neutral fluids, extending intermittency models to MHD turbulence has added complications. MHD turbulence is inherently anisotropic—due to the symmetry breaking caused by embedded magnetic fields—in addition to having fluctuations, energy transfer and dissipation channels for both magnetic and velocity fields. While hydrody-

dynamic intermittency models have been directly applied to MHD turbulence [10, 23], modified models pertaining explicitly to MHD have also been developed [24–29]. Structure function analysis can be of particular use for extracting signatures of dissipation in MHD turbulence [5, 6, 29, 30].

The structure functions and PDFs are constructed by taking differences or increments at varying time scale separations. In this analysis, the increments of magnetic measurements are constructed in three ways; first, the increments of each orthogonal magnetic component ($\Delta B_r, \Delta B_\theta$, and ΔB_z) are examined separately. Then, vector magnitude is created from the vector sum of the three components at each time step and differences between the magnitudes ($\Delta|B|$) at each time step are used for the analysis. Lastly, the magnitude of the vector difference between time points is used as the increment ($|\Delta B|$). The results for these five different forms are generally similar to one another suggesting that the intermittent characteristics of the fluctuations are not strongly anisotropic. The paper also demonstrates that the trends for the fluctuations in the magnetic field are consistent with that found fluctuations in $dB(t)/dt$ which were reported in previous work [8, 31].

EXPERIMENT

Magnetohydrodynamic turbulent fluctuations are generated inside the wind tunnel configuration of the Swarthmore Spheromak Experiment (SSX) using a plasma gun source inside a flux-conserving copper boundary. This copper cylinder is 15.5cm in diameter and 86cm long and situated inside a highly-evacuated chamber ($\approx 1 \times 10^{-8}$ Torr). The details of the gun source and production of magnetic fluctuations have been previously reported [1, 32, 33]. The data analyzed in this paper is of magnetic fluctuations using an array of sixteen magnetic pickup channels. Each channel consists of a single-loop of magnetic wire, 3mm in diameter. Three loops are oriented in each cylindrical coordinate direction (B_r, B_θ, B_z) and each triplet is separated by 0.46cm spanning from about 1cm off the cylindrical axis to the edge of the cylinder boundary. The injected helicity of the plasma is scanned from 0 to $7 \times 10^{-5} Wb^2$ which corresponds to a scan of the amount of flux provided to the gun core—from 0 to $1.5 mWb$ [8].

ANALYSIS TECHNIQUES

The structure of the turbulent plasma is analyzed here through an evaluation of its intermittent character; the primary method for quantifying this characteristic is through the construction of probability distribution functions (PDFs) of increments. The increments can be

in terms of a spatial division or a temporal division. In the context of this experiment, the spatial increments are in units of separation distance between probe locations ($\Delta r_{min} = 0.46cm$), while the temporal increments are in units of sampling cadence ($\Delta \tau_{min} = 154ns$) which corresponds to the sampling frequency of 65MHz. Increments are then increased in multiples of these minimum values. Maximum separation values are limited by physical distance or data acquisition time span, as well the ability to generate enough statistics for a valid calculation.

For a given increment, Δx , the PDF of increments is constructed by computing and histogramming a list of ΔB 's,

$$\Delta B = |B(x_j + \Delta x) - B(x_j)| \quad (1)$$

which can then be normalized to the standard deviation and total count of the histogram.

A single PDF can only describe the nature of the data in terms of its relative distribution with respect to a Gaussian distribution. Large excursions of values in the tail values of the distribution are then indicative of intermittent behavior in the time series signal—i.e. large jumps in values outside of the standard deviation of the mean. Physically, these excursions can be identified with physical mechanisms in the plasma.

Further insight into the physical nature of the plasma can be gained by comparing these PDFs over a range of scales. This can be accomplished qualitatively by examining how the PDFs themselves change, but can also be quantitatively accomplished by calculating moments of these distributions, also known as structure functions. The structure function can be constructed for a given Δx by computing the average of the increments raised to some power, p , as in,

$$S^p(\Delta x) = \langle |B(x_j + \Delta x) - B(x_j)|^p \rangle \quad (2)$$

where j indicates the elements summed over.

Analysis of structure functions can provide a number of features of the fluctuations particularly when modeled as a power-law function,

$$S^p(\Delta x) = (\Delta x)^\zeta(p) \quad (3)$$

When plotted logarithmically, ζ is the slope of the structure function; this slope can indicate the relative level of intermittency compared between two signals. That is, the steeper the slope, the more intermittent (i.e. the larger the "fat tails") is the signal. A flat line is the extreme case and indicates a lack of intermittency in the signal. As one would expect, the structure function of a timeseries generated from a normal distribution has a flat structure function. Changes in slope with scale indicates a change in intermittency as a function of scale. It should be noted that this trend is not a perfect predictor of the presence of intermittency however. For example,

it can be shown that a time-series of fractional Brownian motion (fBm) can be constructed with the same structure function slope [14], but fBm produces a Gaussian probability distribution of increments. However, when intermittency is known to be present based on the PDF of increments (as is true for this SSX dataset) then the relationship between slope and degree of intermittency generally holds true.

If ζ scales as a linear function of p (i.e. $\zeta = Hp$), the system exhibits self-similarity at different scales. In this context, the self-similarity is a qualitative measure of how alike fluctuations appear at different scales. For example, a fractal from chaos theory is a construct that exhibits self-similarity. In other words, a self-similar system can be described in terms of a single fractal scaling or monofractal scaling description. The constant coefficient, H , is called the Hurst exponent as it can be shown that distributions that exhibit this self-similarity can be rescaled using the Hurst exponent as,

$$PDF(\Delta B \Delta x^{-H}) = (\Delta x)^H (PDF(\Delta B, \Delta x)) \quad (4)$$

A non-linear $\zeta(p)$, however, indicates non-self-similar behavior, which in turn can be considered to exhibit multiple fractal scaling simultaneously (i.e. multifractal behavior). These distinctions again can be used to elucidate possible physical mechanisms at play in the plasma.

In particular, it has been shown that differences in self-similarity exist between inertial range and dissipation data in the solar wind [5, 6]. Using the structure function analysis, the slope of the structure functions of the inertial range were not linear as a function of order, indicating that the inertial range turbulent fluctuations in the solar wind did not indicate self-similarity of turbulent structure, and can be considered multifractal. On the other hand, the same analysis for dissipation range structure functions exhibited linear scaling—apparently, the physical mechanism behind the dissipation in the solar wind has a self-similar, scale-invariant nature—that is, monofractal behavior.

TEMPORAL AND SPATIAL PDFS OF INCREMENTS

Representative PDFs of both temporal and spatial magnetic field magnitude ($\Delta|B|$) increments are shown in Figure 1 with temporal PDFs on the left and spatial PDFs on the right. Each PDFs is scaled to its own standard deviation and total count so that they can be cross compared. The increments chosen for the temporal PDFs are $0.092\mu s$, $0.46\mu s$, and $1.85\mu s$ which correspond to frequencies of $10.8MHz$, $2.2MHz$, and $541kHz$ respectively. These three times also approximately correspond to three different regimes of the frequency spectra, called for lack of more accurate description the dissipation, transition and inertial region. All three PDFs ex-

hibit intermittent behavior as indicated by the wings or “fat tails” for large fluctuation values which reflect larger counts compared to a true Gaussian distribution which is indicated by the dashed red curves. Note, though, that the qualitative “level” of intermittency decreases with increasing temporal increment, consistent with previous intermittency analyses on SSX [8, 31], as well as solar wind results [34].

The increments chosen for the spatial PDFs are $0.46cm$, $0.92cm$, and $1.38cm$, which corresponds to the minimum, double the minimum, and triple the minimum separation possible given the probe array in the SSX. Unlike the temporal data, the spatial data is not believed to probe as far into dissipation range scales (for reference, an ion inertial length of $0.6cm$ and an ion gyroradius of $0.1cm$ is calculated for this plasma). Thus the spatial data is likely representative of inertial range turbulence and at most transition range. The resulting PDFs support this notion as they tend to have a more Gaussian-like feature for small fluctuations and only slightly elevated tails. There does appear to be a trend toward increasing Gaussian-ness with increasing increment similar to the temporal data. The curves do not appear as smooth as the temporal data though, as the histograms have larger breaks—in particular, the $0.46cm$ increment PDF. This potentially reflects both a lower amount of increment statistics available for the spatial measurement as well as the possible variations in the PDF due to spatial variation. Asymmetry of the spatial PDFs can also be seen; however, this is likely due to influences of the boundary of which the temporal data is much less susceptible. Finer resolution of the spatial measurements would also likely improve the symmetry of the PDFs in addition to allowing observation of the dissipation scale; such improved measurements are currently being attempted.

As discussed before, the use of PDFs of increments can clearly demonstrate the lack or presence of intermittent behavior in a signal, however in a primarily qualitative way. In order to demonstrate quantitative scaling behavior, and be able to extract more physical interpretations from the data, a structure function analysis is needed and is discussed in the next section.

STRUCTURE FUNCTIONS AND SCALING WITH ORDER

To produce a more quantitative metric from the PDFs of increments, one can calculate moments of the histogram to any order p , and for each time or spatial delay Δx . This is equivalent to calculating the structure function in Equation 2 for the magnetic field increments. Integer values of p correspond to the moments of the PDF ($p = 2$ is the second moment, $p = 3$ is the third moment, etc.) though given the form of Equation 2, p is not restricted to integer values in the structure function




FIG. 1. Probability Distribution Functions (PDFs) of increments for (a-c) temporal and (d-f) spatial measurements of $\Delta|B|$. The temporal increments are (a) $0.092\mu s$, (b) $0.46\mu s$, and (c) $1.85\mu s$ while the spatial increments are (d) $0.46cm$, (e) $0.92cm$, and (f) $1.38cm$. Both temporal and spatial PDFs exhibit non-Gaussian tails indicating intermittency (Gaussian distributions are indicated with dashed red lines), but that the level of intermittency decreases with increasing increment. Spatial PDFs appear to be slightly more asymmetric which may be a result of boundary effects or low resolution compared with temporal measurements.

formalization.

Figure 2 shows the structure functions, $S^p(\Delta x)$, for temporal ($\Delta x = \Delta\tau$) data on the left and spatial ($\Delta x = \Delta r$) data on the right. The integer structure functions from $p = 1$ to $p = 6$ are shown. The structure functions vary as a function of τ or r which confirms the presence of intermittency in the signal. For the temporal results, it is clear that this variation changes as function of time scale, with steeper slopes at small values of τ transitioning to shallower slopes at large values of τ . This indicates that the relative degree of intermittency of the signal increases when analyzed at smaller scales. The transition range is also consistent with the transition from the inertial to the dissipation range of the cascade as determined by the change in spectral index of the frequency spectrum. This difference between inertial and dissipation range intermittency suggests that the physical mechanism underlying the dissipation dynamics has a stronger intermittent nature. Explanations for this behavior in

the solar wind typically focus on the presence of current sheets in the magnetic turbulence [35]. The observation of this difference between dissipation and inertial range fluctuations in this laboratory plasma suggests a similar phenomena [8].

The spatial structure functions do not show a variation with scale; however the slopes and thus the degree of intermittency is consistent with the temporal structure functions in the inertial range. Since the spatial measurements can only probe inertial range scales in the plasma, this result supports the distinction between dissipation and inertial range intermittency seen in the temporal data.

It can also be instructive to normalize the structure functions and in particular highlight the “level” of non-Gaussianness of a distribution. A normalized structure




FIG. 2. Structure functions of temporal (left) and spatial (right) measurements for integer orders $p = 1$ to $p = 6$. Fits to regions of the temporal structure functions are made to compute scaling exponents, ζ ; separate regions are selected based on the breakdown of the frequency spectrum shown in Schaffner 2014c which occurs at approximately 1MHz or $1\mu s$. Spatial measurements likely span only the inertial range and thus only one region is fit.




FIG. 3. Normalized fourth order structure functions (a.k.a flatness or kurtosis) for temporal (left) and spatial (right) measurements. The flatness of a Gaussian signal is indicated by the dashed line at $F=3$. Both figures show increasing flatness with decreasing scale indicating a rise in intermittency.

function can be constructed as,

$$S_{norm}^p(\Delta x) = \frac{S^p(\Delta x)}{(S^2(\Delta x))^{(p/2)}} \quad (5)$$

where p is the order of the moment. Using this normalized structure function, a Gaussian timeseries would have a constant value. For example, for $p = 4$, this quantity becomes the flatness or kurtosis. Figure 3 shows the flatness for both temporal and spatial data. A Gaussian timeseries would have a constant value of 3. Both Figure 3(a) and (b) show an excursion from a Gaussian distribution at smaller increments, though it is clearly

more pronounced for the temporal data.

A final characteristic, self-similarity of the turbulence, can be extracted from the structure function analysis by examining the slope of the structure functions as a function of order, $\zeta(p)$. The red lines in Figure 2 are fits to the structure functions. In the temporal data, fits are applied to two separate regions that correspond to the dissipation and inertial regions based on spectral frequency analysis, while the spatial data has only one fit for the entire region. A fit is made in each of these regions for each structure function using order p which ranges between 0.1 and 10 in steps of 0.1. The results of this scan are displayed in Figure 4 for both time and space. The five separate magnetic field measurements are shown now, with inertial range fits indicated by solid lines and dissipation range fits indicated with dashed lines. In this construction, a timeseries that exhibits self-similarity or monofractality would produce a line with a constant slope. As the increments are raised to increasingly higher powers, the value of the structure function should also increase power-law like. If the signal is self-similar, an increment between two points in the time series at one scale should have on average the same relative increment between two points at a different scale. In other words, a big fluctuation compared to a small fluctuation at one scale should appear to have the same relative ratio at a different scale. Thus, the structure function should change at a constant rate based on this ratio. If the sig-

nal is not self-similar, then the ratio is not constant and differences are increasingly accentuated by the raised exponent resulting in a non-linear scaling.

Since the original 1941 Kolmogorov turbulence theory [15] predicts a self-similar scaling, it predicts that the structure function slope should be one-third the order, or $\zeta = p/3$. The dashed purple line indicates this prediction in Figure 4. The inertial range curves for all five temporal magnetic measurements in Figure 4(a) sit relatively near the Kolmogorov theory line but none of the lines exhibiting completely linear behavior. This indicates that the inertial range fluctuations are multifractal, and not self-similar. The spatial curves in Figure 4(b) support this observation.

The dissipation range curves, in contrast, are clearly linear. This indicates that the dissipation range fluctuations are monofractal while also having a higher level of intermittency than inertial range fluctuations. The Hurst exponents for the dissipation range are consistently about 1 for the various measured B values, steeper than the K41 prediction of $H=1/3$.




FIG. 4. Scaling exponent, ζ , as a function of structure function order, p for temporal (upper) and spatial (lower) measurements. Inertial range scaling is indicated by the bold lines, while dissipation range scaling is indicated by dashed lines. Temporal scaling shows a clear distinction between multifractal (non-linear ζ vs p) in the inertial range and monofractal (linear ζ vs p) in the dissipation range. The different color curves indicate the type of magnetic structure function used (i.e. B-component, B-vector magnitude, etc.)

HELICITY SCALING

The structure function analysis is also used to examine the effect of varying the amount of magnetic helicity in the plasma on the intermittency and self-similarity of the magnetic fluctuations. Previous work has shown that the degree of intermittency in fluctuations of $\dot{B} = dB/dt$ (not B), as determined by a calculation of the flatness, increases on averaged with an increasing amount of injected magnetic helicity [8]. This work revisits that result by examining the trend in B using unnormalized structure functions and the resulting relationship between structure function slope and p .




FIG. 5. Scaling exponent versus order for different values of magnetic helicity (as set by the amount of initial flux in the plasma gun [8]). Inertial range intermittency increases with helicity as shown by increasingly steeper curves in the inertial range. Dissipation range scaling appears to be unaffected by the change in helicity.

Figure 5 shows the slope versus order for eight different helicity states for inertial range fluctuations (solid lines) and dissipation range fluctuations (dashed lines). Recalling that the steepness of the slope is indicative of relative degree of intermittency, the order of the inertial range curves for each helicity indicates increasing intermittency with increasing helicity, again consistent with the findings for an analysis of \dot{B} timeseries [8]. However, the structure function analysis here further shows that with the exception of the zero helicity state, the inertial range turbulence for any helicity is not self-similar. The dissipation range lines, however, show that the dissipation range intermittency is self-similar regardless of the amount of injected magnetic helicity in the plasma.

DISCUSSION

The observation of both multifractal and monofractal scaling of magnetic fluctuation structure functions in SSX has implications for both the dissipative nature of the turbulence and for the universality of MHD turbulence. Like that observed in Kiyani 2013, the transition between multifractal scaling of the inertial range and monofractal scaling of the dissipation range is rapid. This suggests that the dissipation mechanism is independent of the scale of the turbulence structures—in other words, the dissipation scale is absolute rather than relative. This is in contrast to the predictions and observations made for hydrodynamic turbulence where viscosity is the known dissipation mechanism. In those cases, each scaling exponent has a corresponding dissipation scale due to the viscosity. As smaller and smaller scales are approached, the viscosity “shuts-off” the scaling exponents gradually producing a scaling transition region from multifractal inertial to monofractal dissipation ranges. If the dissipation mechanism has an absolute scale, independent of each scaling exponent, than a rapid transition would be observed as this absolute scale is passed. Kiyani 2010 discusses this possibility that a dissipation scale associated with the ion inertial length may be at play. However, it is often difficult to distinguish the ion gyroscale (ρ_i) effects versus ion inertial scale (δ_i) effects in super-Alfvenic solar wind since $\beta \approx 1$. This laboratory turbulence, however, has a more resolved separation between ρ_i and δ_i) [1], and the results presented here show the multifractal to monofractal scaling transition occurring at around δ_i rather than ρ_i . Though evidence for the existence of current sheets has been demonstrated in previous SSX experiments on reconnection using stable spheromaks [36], more concrete evidence for the presence of current sheets in this more turbulent SSX plasma is still being sought; however, there is preliminary evidence from the helicity scan work [8] that these current sheets may indeed be present. These current sheets, then, could play the role of an absolute dissipation scale. It should be noted, however, that an alternative solar wind intermittency analysis [30] does not exhibit a transition in scaling from inertial to dissipation range and the authors of this work suggest the existence of a compressible inertial range cascade rather than a dissipation range.

Another interesting comparison between the results in Kiyani 2013 and those presented here is the difference in collisionality. The solar wind is a collisionless plasma with mean free path lengths on the order of AU. The SSX plasma, while still fully ionized, is a collisional plasma with ion mean free path lengths on the order of 0.1cm, approximately two orders of magnitude smaller than the ion inertial length [1]. While in MHD turbulence theory, resistivity can be thought to play a role for magnetic turbulence as viscosity would for velocity turbulence, the re-

sistivity does not appear to have the same affect for dissipation given the comparison in scaling between the collisionless solar wind and the collisional laboratory plasma. This has implications for the universality of MHD turbulence.

CONCLUSIONS

This paper has presented to results of a detailed structure function analysis of turbulent magnetic fluctuations in a laboratory plasma experiment. The structure functions indicate a distinction between multifractal scaling in inertial range regimes and monofractal scaling in dissipation range regimes. Multifractal scaling is observed in both temporal fluctuations as well as spatial fluctuations. Monofractal scaling is seen only in temporal fluctuations due to the limitations of spatial measurement resolution. The work also shows that intermittency and scaling can be modified by the magnetic helicity of the turbulent plasma; increasing helicity corresponds to increase intermittency in the inertial range and modification of the scaling exponents as a function of order. However, helicity does not appear to affect the monoscaling results of the dissipation scale. These results compare favorably to a similar analysis of inertial range and dissipation range turbulence in the solar wind [5], which suggests some amount of universality between the two MHD systems despite differences in collisionality.

ACKNOWLEDGEMENTS

The authors would like to acknowledge fruitful discussions with Peter Weck. This work has been funded by DOE OFES and NSF CMSO.

REFERENCES

-
- [1] D.A. Schaffner, M.R. Brown and V.S. Lukin. “Temporal and Spatial Turbulent Spectra of MHD Plasma and an Observation of Variance Anisotropy”. *ApJ* **790**, 126 (2014).
 - [2] Frisch, U. 1995, *Turbulence* (Cambridge: Cambridge Univ. Press).
 - [3] G. Paladin and A. Vulpiani. “Anomalous Scaling Laws in Multifractal Objects.” *Phys. Rep.* **156** 147-225 (1987).
 - [4] E. Marsch and C.-Y. Tu. “Intermittency, non-Gaussian statistics and fractal scaling of MHD fluctuations in the solar wind.” *Nonlinear Processes in Geophysics* **4**, 101-124 (1997).
 - [5] K.H. Kiyani, S.C. Chapman, F. Sahraoui, B. Hnat, O. Fauvarque, and Yu.V. Khotyainsev. Enhanced Magnetic

- Compressibility and Isotropic Scale Invariance at Sub-Ion Larmor Scales in Solar Wind Turbulence. *ApJ*. **763** 10 (2013).
- [6] K.H. Kiyani, S.C. Chapman, Yu. V. Khotyaintsev, M.W. Dunlop and F. Sahraoui. "Global Scale-Invariant Dissipation in Collisionless Plasma Turbulence." *Phys. Rev. Lett.* **103** 075006 (2009).
- [7] K. Kiyani, S. Chapman, Y. Khotyaintsev, M. Dunlop, and F. Sahraoui. "Fractal dissipation of small-scale magnetic fluctuations in solar wind turbulence as seen by CLUSTER." AIP Conf. Proc. 1216, Twelfth International Solar Wind Conference, ed. M. Maksimovic et al. (Melville, NY: AIP), **136** (2010).
- [8] D.A. Schaffner, A. Wan, and M. R. Brown, "Observation of turbulent intermittency scaling with magnetic helicity in an MHD plasma wind-tunnel", *Phys. Rev. Letters* **112**, 165001 (2014).
- [9] F. Anselmet, Y. Gagne, E.J. Hopfinger, and R.A. Antonia. "High-order velocity structure functions in turbulent shear flows." *J. Fluid Mech.* **140** 63-89 (1984).
- [10] L.F. Burlaga. "Intermittent Turbulence in the Solar Wind". *J. Geophys. Res.* **96** 5847-5851 (1991).
- [11] L.F. Burlaga. "Multifractal structure of the magnetic field and plasma in recurrent streams at 1AU." *J. Geophys. Res.* **97** 4283-4293 (1992).
- [12] C.-Y. Tu and E. Marsch. MHD Structures, Waves and Turbulence in the Solar Wind: Observations and Theories. *Space Sci. Rev.* **73** 1-210 (1995).
- [13] G. Consolini, M.F. Marcucci and M. Candidi. "Multifractal Structure of Auroral Electrojet Index Data." *Phys. Rev. Lett.* **76** 4082 (1996).
- [14] B. Hnat, S.C. Chapman, G. Rowlands, N.W. Watkins, and M.P. Freeman. "Scaling in long term data sets of geomagnetic indices and solar wind ε as seen by WIND spacecraft". *Geophys. Res. Lett.* **30** 2174 (2003).
- [15] A. N. Kolmogorov, *Dokl. Acad. Nauk. SSSR* **30**, 301, (1941).
- [16] R. Benzi, G. Paladin, G. Parisi, and A. Vulpiani. "On the multifractal nature of fully developed turbulence and chaotic systems." *J. Phys. A: Math. Gen.* **17** 3521-3531 (1984).
- [17] A.N. Kolmogorov. "A refinement of previous hypothesis concerning the local structure of turbulence in viscous incompressible fluid at high Reynolds number." *J. Fluid Mech.* **13** 82 (1962).
- [18] C. Meneveau and K.R. Sreenivasan. "Simple Multifractal Cascade Model for Fully Developed Turbulence." *Phys. Rev. Lett.* **59** 1424 (1987).
- [19] Z.-S. She and E. Leveque. "Universal Scaling Laws in Fully Developed Turbulence." *Phys. Rev. Lett.* **72** 336 (1994).
- [20] B. Dubrulle. "Intermittency in Fully Developed Turbulence: Log-Poisson Statistics and Generalized Scale Covariance." *Phys. Rev. Lett.* **73** 959 (1994).
- [21] U. Frisch and M. Vergassola. "A Prediction of the Multifractal Model: the Intermediate Dissipation Range." *Europhys. Lett.* **14** 439-444 (1991).
- [22] L. Chevillard, B. Castaing, and E. Leveque. "On the rapid increase of intermittency in the near-dissipation range of fully developed turbulence." *Eur. Phys. J. B* **45** 561-567 (2005).
- [23] C. Pagel and A. Balogh. "Intermittency in the solar wind: A comparison between solar minimum and maximum using Ulysses Data". *J. Geophys. Res.* **107** 1178 (2002).
- [24] V. Carbone. "Cascade Model for Intermittency in Fully Developed Magnetohydrodynamic Turbulence." *Phys. Rev. Lett.* **71** 1546 (1993).
- [25] D. Biskamp "Cascade Model for magnetohydrodynamic turbulence." *Phys. Rev. E* **50** 2702 (1994).
- [26] W.-C. Müller and D. Biskamp. "Scaling Properties of Three-Dimensional Magnetohydrodynamic Turbulence". *Phys. Rev. Lett.* **84** 475 (2000).
- [27] D. Biskamp and W.-C. Müller. "Scaling Properties of three-dimensional isotropic magnetohydrodynamic turbulence." *Phys. Plasmas*. **7** 4889 (2000).
- [28] S. Boldyrev. "Kolmogorov-Burgers Model for Star-Forming Turbulence." *Astrophysical J.* **569** 841-845 (2002).
- [29] J. Cho, A. Lazarian, and E.T. Vishniac. "Ordinary and Viscosity-Damped Magnetohydrodynamic Turbulence." *ApJ*. **59** 812-823 (2003).
- [30] O. Alexandrova, V. Carbone, P. Veltri and L. Sorriso-Valvo. "Small-Scale Energy Cascade of the Solar Wind Turbulence." *Astrophysical J.* **674** 1153-1157 (2008).
- [31] D.A. Schaffner *et al.* Turbulence analysis of an experimental flux rope plasma. **56** 064003 (2014).
- [32] M.R. Brown and D.A. Schaffner. "Laboratory sources of turbulent plasma: a unique MHD plasma wind tunnel". *Plasma Sources and Science Technology*. **23** 063001 (2014).
- [33] M.R. Brown and D.A. Schaffner. "SSX MHD plasma wind tunnel". *Journal of Plasma Physics*. (2015).
- [34] R. Bruno and V. Carbone. *Living Rev Solar Phys* **10** (2013).
- [35] K.T. Osman, W.H. Matthaeus, J.T. Gosling, A. Greco, S. Servidio, B. Hnat, S.C. Chapman, and T.D. Phan. "Magnetic Reconnection and Intermittent Turbulence in the Solar Wind." *Phys. Rev. Lett.* **112** 215002 (2014).
- [36] M.R. Brown, C.D. Cothran, T. Gray, C.E. Myers, and E.V. Belova. *Phys. Plasmas* **19** 080704 (2012).