

1 NEBB

The Non-Equilibrium Bounce-Back scheme [1] is a boundary condition based on the wet-node approach. Instead of the more common Half-way Bounce-Back scheme where the boundary fluid nodes are half a node away from the physical wall, the boundary fluid nodes are infinitesimally close to the physical wall in the wet-node approach. The NEBB scheme leaves the populations streaming out of the domain unchanged after streaming. The unknown populations streaming into the domain can be found using the non-equilibrium parts of the populations streaming out of the domain. Additionally, momentum corrections N_α , where α is the spatial component, are introduced [2]. This gives the following boundary condition for the unknown incoming populations i ,

$$\begin{aligned} f_i^{\text{neq},\sigma} &= f_q^{\text{neq},\sigma} - w_i c_{i\alpha} N_\alpha^\sigma, \\ f_i^\sigma &= f_q^\sigma + (f_i^{\text{eq},\sigma} - f_q^{\text{eq},\sigma}) - w_i c_{i\alpha} N_\alpha^\sigma, \end{aligned} \quad (1.1)$$

where the index q is given by $\mathbf{c}_i = -\mathbf{c}_q$. Furthermore, repeated Greek indices indicate sums. Note that the incoming populations are defined by $\mathbf{c}_i \cdot \hat{\mathbf{n}}_w > 0$, where $\hat{\mathbf{n}}_w$ is the normal vector to the boundary pointing into the domain. The density ρ and the velocity $\mathbf{u}$ are the imposed boundary density and velocity. To conserve mass, we will satisfy the condition that the amount of mass being removed from the domain is equal to the mass added to the domain at every grid point. During the streaming step, the values of the outgoing populations f_q will be overwritten by populations from the bulk, leading to mass loss. The mass gain is given by Eq. (1.1). This gives the net mass loss at the boundary due to the NEBB method,

$$\begin{aligned} \Delta m^\sigma &= \sum_i^{\text{out}} (f_i^{*,\sigma} - f_i^\sigma) + \sum_i^{\text{in}} w_i c_{i\alpha} N_\alpha^\sigma \\ &= \sum_i^{\text{out}} (f_i^{*,\sigma} - f_i^\sigma) + \frac{1}{6} N_n^\sigma, \end{aligned} \quad (1.2)$$

where outgoing populations are defined by $\mathbf{c}_i \cdot \hat{\mathbf{n}}_w < 0$. Furthermore, the index n indicates the component normal to the boundary. Note that we ignored the term $\sum_i^{\text{in}} (f_i^{\text{eq},\sigma} - f_q^{\text{eq},\sigma}) = \rho^\sigma u_n$. This is because this is not mass loss, but the expected mass flux across an open boundary. For a set velocity boundary condition, the densities can be found using the bare velocity components,

$$\begin{aligned} \rho^\sigma u_n^\sigma &= \sum_i^{\text{in}} f_i^\sigma - \sum_i^{\text{out}} f_i^\sigma + \frac{1}{2} F_n^\sigma \\ &= \rho^\sigma - f_0^\sigma - 2 \sum_i^{\text{out}} f_i^\sigma - \sum_i^{\text{par}} f_i^\sigma + \frac{1}{2} F_n^\sigma \\ \rho^\sigma &= \frac{f_0^\sigma + 2 \sum_i^{\text{out}} f_i^\sigma + \sum_i^{\text{par}} f_i^\sigma - \frac{1}{2} F_n^\sigma}{1 - u_n^\sigma}, \end{aligned} \quad (1.3)$$

where parallel populations are defined as $\mathbf{c}_i \cdot \hat{\mathbf{n}}_w = 0$ with $i \neq 0$. Furthermore, we assume that $|\hat{\mathbf{c}}_i \cdot \hat{\mathbf{n}}_w| = 1$ for $i \neq 0$. This equation can be inverted to find the normal velocity from the imposed density at the boundary. Note that the parallel velocity components are not fixed by this equation and can be chosen freely. The momentum corrections N_α^σ can be computed by plugging the incoming populations from Eq. (1.1) back into the bare velocity components,

$$\begin{aligned} \rho^\sigma u_\alpha^\sigma &= \sum_i^{\text{in}} f_i^\sigma c_{i\alpha} + \sum_i^{\text{out}} f_i^\sigma c_{i\alpha} + \sum_i^{\text{par}} f_i^\sigma c_{i\alpha} + \frac{1}{2} F_\alpha^\sigma \\ &= \sum_i^{\text{in}} \left[(f_i^{\text{eq},\sigma} - f_q^{\text{eq},\sigma}) - w_i c_{i\beta} N_\beta^\sigma \right] c_{i\alpha} + \sum_i^{\text{par}} f_i^\sigma c_{i\alpha} + \frac{1}{2} F_\alpha^\sigma. \end{aligned} \quad (1.4)$$

The normal and parallel components of the momentum corrections can be found,

$$\begin{aligned} N_\alpha^\sigma &= \frac{1}{A_\alpha} \left[\sum_i^{\text{par}} c_{i\alpha} f_i^\sigma + \sum_i^{\text{in}} c_{i\alpha} (f_i^{\text{eq},\sigma} - f_q^{\text{eq},\sigma}) - \rho^\sigma u_\alpha^\sigma + \frac{1}{2} F_\alpha^\sigma \right] \\ &= \frac{1}{A_\alpha} \left[\sum_i^{\text{par}} c_{i\alpha} (f_i^\sigma - f_i^{\text{eq},\sigma}) + \frac{1}{2} F_\alpha^\sigma \right], \end{aligned} \quad (1.5)$$

with $A_\alpha = \sum_i^{\text{in}} w_i c_{i\alpha}^2$. For the D2Q9, D3Q19 and D3Q27 lattices we have $A_\alpha = \frac{1}{6}\delta_{\alpha n} + \frac{1}{18}(1 - \delta_{\alpha n})$. To correct for the mass loss given in Eq. (1.2), we add this back to the zero-velocity component before computing the density,

$$\begin{aligned} f_0^\sigma &= f_0^{*,\sigma} + \Delta m^\sigma \\ &= f_0^{*,\sigma} + \sum_i^{\text{out}} (f_i^{*,\sigma} - f_i^\sigma) + \frac{1}{2} F_n^\sigma. \end{aligned} \quad (1.6)$$

This simplifies the computation of the densities, making it independent of forcing,

$$\rho_w^\sigma = \frac{f_0^{*,\sigma} + \sum_i^{\text{out}} (f_i^{*,\sigma} + f_i^\sigma) + \sum_i^{\text{par}} f_i^\sigma}{1 - u_n^\sigma}. \quad (1.7)$$

2 Asymptotic expansion of LBE

To test the validity of the mass corrected NEBB method, we will need to apply an asymptotic expansion in Knudsen number [3]. The Knudsen number is given as $Kn = \lambda/L$, where λ is the mean free path that decreases with relaxation time τ , and L is the characteristic length scale of the system. For simplicity, we apply the expansion to the single-component Lattice Boltzmann Equation, but now in explicitly dimensional form,

$$\begin{aligned} & \tilde{f}_i(\tilde{\mathbf{x}} + \tilde{\mathbf{c}}_i \Delta \tilde{t}, \tilde{t} + \Delta \tilde{t}) - \tilde{f}_i(\tilde{\mathbf{x}}, \tilde{t}) \\ &= -\frac{\Delta \tilde{t}}{\tilde{\tau}} \left(\tilde{f}_i(\tilde{\mathbf{x}}, \tilde{t}) - \tilde{f}_i^{\text{eq}}(\tilde{\rho}, \tilde{\mathbf{u}}) \right) + \Delta \tilde{t} \tilde{S}_i, \end{aligned} \quad (2.1)$$

where the dimensional source term is given as follows,

$$\tilde{S}_i = \left(1 - \frac{\Delta \tilde{t}}{2\tilde{\tau}} \right) w_i \left(\frac{\tilde{\mathbf{c}}_i - \tilde{\mathbf{u}}}{\tilde{c}_s^2} + \frac{\tilde{\mathbf{c}}_i \cdot \tilde{\mathbf{u}}}{\tilde{c}_s^4} \tilde{\mathbf{c}}_i \right) \cdot \tilde{\mathbf{F}}. \quad (2.2)$$

We will now nondimensionalize this equation the length scale $\tilde{L}$, the velocity scale $\tilde{c}$, the timescale $\tilde{t}_0$ and the density scale $\tilde{\rho}_0$. Note that $\tilde{c} = \Delta \tilde{x} / \Delta \tilde{t}$. Furthermore, we introduce the Strouhal number $Sh = \tilde{L} / (\tilde{t}_0 \tilde{c}) = \Delta t / \Delta x$. This gives the nondimensional version of the previous equations,

$$\begin{aligned} & f_i(\mathbf{x} + \mathbf{c}_i \Delta x, t + \Delta t) - f_i(\mathbf{x}, t) \\ &= -\frac{\Delta t}{\tau} \left(f_i(\mathbf{x}, t) - f_i^{\text{eq}}(\rho, \mathbf{u}) \right) + \Delta x S_i, \\ S_i &= \left(1 - \frac{\Delta t}{2\tau} \right) w_i \left(\frac{\mathbf{c}_i - \mathbf{u}}{c_s^2} + \frac{\mathbf{c}_i \cdot \mathbf{u}}{c_s^4} \mathbf{c}_i \right) \cdot \mathbf{F}, \end{aligned} \quad (2.3)$$

where we defined $\tilde{\mathbf{F}} = \tilde{\rho}_0 \tilde{c}^2 / \tilde{L} \mathbf{F}$. We will now introduce diffusive scaling, such that $\Delta t = \mathcal{O}(Kn^2)$ and $\Delta x = \mathcal{O}(Kn)$, implying $Sh = \mathcal{O}(Kn)$. Furthermore, if we choose a finite Reynolds number $Re = \tilde{U} \tilde{L} / \tilde{\nu} = \mathcal{O}(1)$, we get the Mach number $Ma = \tilde{U} / \tilde{c}_s = \mathcal{O}(Kn)$ from the equation $Ma \propto$

$Kn \cdot Re$. Lastly, $\tau = \mathcal{O}(Kn^2)$, which we will show is required to recover the equilibrium at leading order in Knudsen number. We will now perform a Taylor expansion on the dimensionless LBE,

$$\sum_{n=1}^{\infty} \frac{1}{n!} \left(\Delta x c_{i\alpha} \partial_{\alpha} + \Delta t \partial_t \right)^n f_i = -\frac{\Delta t}{\tau} \left(f_i - f_i^{\text{eq}} \right) + \Delta x S_i, \quad (2.4)$$

We will continue by expanding this equation in order of Knudsen number. We will expand the distribution function, the equilibrium distribution, the density, and the velocity in orders of Knudsen. Note that from the Mach number, we can state that $\mathbf{u}^{(0)} = \mathbf{0}$. Sorted by order in Knudsen, we recover the following equations,

$$\begin{aligned} f_i^{(0)} &= f_i^{\text{eq},(0)}, \\ f_i^{(1)} &= f_i^{\text{eq},(1)} - \frac{\tau}{\Delta t} \left(\Delta x c_{i\alpha} \partial_{\alpha} f_i^{(0)} - \Delta x S_i^{(0)} \right), \\ f_i^{(2)} &= f_i^{\text{eq},(2)} - \frac{\tau}{\Delta t} \left(\Delta x c_{i\alpha} \partial_{\alpha} f_i^{(1)} + \frac{1}{2} \Delta x^2 c_{i\alpha} c_{i\beta} \partial_{\alpha} \partial_{\beta} f_i^{(0)} + \Delta t \partial_t f_i^{(0)} - \Delta x S_i^{(1)} \right), \\ f_i^{(3)} &= f_i^{\text{eq},(3)} - \frac{\tau}{\Delta t} \left(\Delta x c_{i\alpha} \partial_{\alpha} f_i^{(2)} + \frac{1}{2} \Delta x^2 c_{i\alpha} c_{i\beta} \partial_{\alpha} \partial_{\beta} f_i^{(1)} + \Delta t \partial_t f_i^{(1)} + \Delta x c_{i\alpha} \partial_{\alpha} \Delta t \partial_t f_i^{(0)} \right. \\ &\quad \left. + \frac{1}{6} \Delta x^3 c_{i\alpha} c_{i\beta} c_{i\gamma} \partial_{\alpha} \partial_{\beta} \partial_{\gamma} f_i^{(0)} - \Delta x S_i^{(2)} \right). \end{aligned} \quad (2.5)$$

Here, we expanded the source term and the equations for density and velocity,

$$\begin{aligned} \rho^{(n)} &= \sum_i f^{(n)}, \\ (\rho \mathbf{u})^{(n)} &= \sum_i \mathbf{c}_i f_i^{(n)} + \frac{\Delta x}{2} \mathbf{F}^{(n-1)}. \end{aligned} \quad (2.6)$$

The macroscopic equations can be obtained by taking the zeroth and first moments of the expansion in Eq. (2.5). For simplicity, we will derive these macroscopic equations for $\rho^{(0)} = \text{const}$ and $\mathbf{F}^{(0)} = \mathbf{0}$. We can now compute the zeroth and first moments of the equation for $f^{(2)}$,

$$\begin{aligned} \rho^{(0)} \nabla \cdot \mathbf{u}^{(1)} &= 0, \\ \mathbf{0} &= -\nabla \left(c_s^2 \rho^{(1)} \right) + \mathbf{F}^{(1)}. \end{aligned} \quad (2.7)$$

We can do the same for $f^{(3)}$,

$$\begin{aligned} Sh \partial_t \rho^{(1)} + \nabla \cdot \left(\rho^{(0)} \mathbf{u}^{(2)} + \rho^{(1)} \mathbf{u}^{(1)} \right) &= 0, \\ Sh \rho^{(0)} \partial_t \mathbf{u}^{(1)} + \rho^{(0)} \nabla \cdot \left(\mathbf{u}^{(1)} \mathbf{u}^{(1)} \right) &= \\ -\nabla \left(c_s^2 \rho^{(2)} \right) + \frac{\rho^{(0)}}{Sh} \left[\nu \left(\nabla \mathbf{u}^{(1)} + \left(\nabla \mathbf{u}^{(1)} \right)^T \right) \right] &+ \mathbf{F}^{(2)}, \end{aligned} \quad (2.8)$$

with $\nu = c_s^2(\tau - 1/2)$.

3 Analysis of NEBB

To analyze the accuracy of the force-corrected and mass-corrected NEBB method, previous literature usually assumes the lack of a Knudsen layer [4]. This means that this method is used to

find out up until which order in Knudsen number the bulk equations for the distribution function fulfill the boundary condition. We start by writing Eq. (1.1) in dimensionless form,

$$\begin{aligned} f_i &= f_q + (f_i^{\text{eq}} - f_q^{\text{eq}}) - w_i c_{i\alpha} N_\alpha, \\ N_\alpha &= \frac{1}{A_\alpha} \left[\sum_i^{\text{par}} c_{i\alpha} (f_i - f_i^{\text{eq}}) + \frac{\Delta x}{2} F_\alpha \right]. \end{aligned} \quad (3.1)$$

Note that we ignored the component specifier σ for simplicity. We can expand this equation in Knudsen number. The momentum corrections are given as follows,

$$\begin{aligned} N_\alpha^{(0)} &= \frac{1}{A_\alpha} \sum_i^{\text{par}} c_{i\alpha} (f_i^{(0)} - f_i^{\text{eq},(0)}), \\ N_\alpha^{(1)} &= \frac{1}{A_\alpha} \left[\sum_i^{\text{par}} c_{i\alpha} (f_i^{(1)} - f_i^{\text{eq},(1)}) + \frac{\Delta x}{2} F_\alpha^{(0)} \right], \\ N_\alpha^{(2)} &= \frac{1}{A_\alpha} \left[\sum_i^{\text{par}} c_{i\alpha} (f_i^{(2)} - f_i^{\text{eq},(2)}) + \frac{\Delta x}{2} F_\alpha^{(1)} \right]. \end{aligned} \quad (3.2)$$

As can be seen from Eq. (2.5), the zeroth-order distribution function is equal to the zeroth-order equilibrium. This means that the momentum correction at this order is zero, and the NEBB is satisfied. The first-order distribution function gives the following momentum corrections,

$$N_\alpha^{(1)} = \frac{1}{A_\alpha} \left[-\frac{\tau}{\Delta t} \sum_i^{\text{par}} c_{i\alpha} (\Delta x c_{i\beta} \partial_\beta f_i^{\text{eq},(0)} - \Delta x S_i^{(0)}) + \frac{\Delta x}{2} F_\alpha^{(0)} \right]. \quad (3.3)$$

For the equilibrium and source term defined by previous research [5], we get the following if we assume $\mathbf{u}^{(0)} = \mathbf{0}$,

$$\begin{aligned} \sum_i^{\text{par}} c_{i\alpha} c_{i\beta} f_i^{\text{eq},(0)} &= (\delta_{\alpha n} - 1) \delta_{\alpha\beta} \frac{2}{3} p^{(0)}, \\ \sum_i^{\text{par}} c_{i\alpha} S_i^{(0)} &= (\delta_{\alpha n} - 1) \frac{2}{3} F_\alpha^{(0)}, \end{aligned} \quad (3.4)$$

where p is the pressure. This gives the following momentum correction,

$$\begin{aligned} w_i c_{i\alpha} N_\alpha^{(1)} &= \frac{w_i c_{i\alpha}}{\sum_i^{\text{in}} w_i c_{i\alpha}^2} \left[-\frac{2}{3} \frac{\tau}{\Delta t} (\delta_{\alpha n} - 1) (\partial_\alpha p^{(0)} - \Delta x F_\alpha^{(0)}) + \frac{\Delta x}{2} F_\alpha^{(0)} \right] \\ &= \sum_\alpha \left[-12 \frac{\tau}{\Delta t} (\delta_{\alpha n} - 1) w_i c_{i\alpha} (\partial_\alpha p^{(0)} - \Delta x F_\alpha^{(0)}) + 3 w_i c_{i\alpha} \Delta x F_\alpha^{(0)} \right] \end{aligned} \quad (3.5)$$

References

- [1] Q. Zou and X. He, [Physics of Fluids](#) **9**, 1591 (1997).
- [2] T. Krüger, H. Kusumaatmaja, A. Kuzmin, O. Shardt, G. Silva, and E. M. Viggien, *The Lattice Boltzmann Method: Principles and Practice*, Graduate Texts in Physics (Springer International Publishing, Cham, 2017).
- [3] T. Inamuro, M. Yoshino, and F. Ogino, [Physics of Fluids](#) **9**, 3535 (1997).
- [4] M. Junk and Z. Yang, [Journal of Statistical Physics](#) **121**, 3 (2005).
- [5] S. Saito, N. Takada, S. Baba, S. Someya, and H. Ito, en, [Physical Review E](#) **108**, 065305 (2023).