

Name



ID#

Math 150 Spring 2025 Final Exam

- Desmos is allowed
- Desmos and the only app (or website) allowed - no other apps or websites are allowed at anytime, this is **not** an open internet test.
- Must show all work to receive credit
- Clearly label all graphs and make sure both axes have a clear scale.
- Attach all scratch paper

	Question	Points
1	Integration	4 / 5
2	Riemann Sums	5 / 5
3	Volume	2 / 5
4	Limits & Continuity	5 / 5
5	Average Value	2 / 4
6	Optimization	5 / 5
7	Position, Velocity, Acceleration	0 / 4
8	Related Rates	/ 3
9	First Derivative Test	/ 3
10	Geometry	2 / 3
Total		26 / 40

1) Evaluate the integral $\int 4 \sin^5(x) \cos^3(x) dx$ using two different substitutions and discuss how the two anti-derivatives are related.

$$\cos^2 = \sin^2 + 1$$

$$4 \int \sin^5(x) \cos^3(x) dx$$

$$\frac{-\frac{8}{9}U^9}{9} + \frac{U^6}{6}$$

$$\sin^7 x \cos + 1$$

$$U = \sin$$

$$DU = \cos$$

$$\cos^2 x - 1 = -\sin^2 x$$

$$\sin^2 x - 1 = -\cos^2 x$$

$$\cos^2 x = -\sin^2 x + 1$$

$$\sin^2 x = -\cos^2 x + 1$$

ANS 1 11

$$\left(-\frac{1}{8} \sin^8 x + \frac{1}{6} \sin^6 x + C \right) \sin^5 x \cos^2 x \cos x$$

$$2) \sin^2 x \sin^2 x \cdot \sin \cos^3 x$$

$$\sin^5 x (-\sin^2 x + 1) \cos x$$

SUB

$$-(\cos^2 x + 1)^2 \sin x \cos^3 x$$

$$U = \sin x$$

$$DU = \cos x$$

AK

$$U = \cos$$

$$DU = -\sin$$

$$U^5 (-U^2 + 1)$$

$$-U^7 + U^5 \quad DU \quad \uparrow$$

$$-(U^2 + 1)^2 \cdot U^3 \quad DU$$

$$\frac{-U^{7+1}}{8} + \frac{U^{5+1}}{6}$$

$$-\frac{U^8}{8} - \frac{U^4}{4}$$

$$+U^5 - U^5$$

$$-(-U^2 + 1) U^3$$

$$(-U^5 + U^3) - U^2 + 1$$

$$-U^7 + U^3$$

$$-U^2 + 1$$

$$-\frac{U^8}{8} + \frac{U^6}{6}$$

$$= -\frac{1}{8} \sin^8 x$$

ANS 2


$$-\frac{\cos^8 x}{8} - \frac{\cos^4 x}{4} + C$$

Name _____ ID# _____

2) Estimate the signed area under the curve $y = 4 - x^2$ from 0 to 3 using:

- three (3) rectangles with the **left endpoint** approximation.
- three (3) rectangles with the **right endpoint** approximation,
- three (3) rectangles with the **midpoint** approximation,
- Find the exact signed area using FTC and compare your estimates from parts (a-c)

In each case a-c draw the graph and rectangles and use sigma notation.



LEFT $\frac{3-0}{3} = 1$

$$\sum_{N=1}^3 4 - (N-1)^2 \cdot 1 = 7$$

$4 - 2 = 2$

4×1
3
0


L	R
0, 4	3
1, 3	0
	-5

$\sum 10$

$3, -5$

LEFT $\sum_{N=1}^3 4 - (N-1)^2 \cdot 1 = 7 \checkmark$

RIGHT $\sum_{N=1}^3 4 - (N)^2 \cdot 1 = -2 \checkmark$



MID $= \sum_{N=1}^3 4 - (x - 1/2)^2 \cdot 1 = 13/4 \checkmark$

$4 \cdot 10 \sqrt{14} + 7 \cdot$


$4x - \frac{x^3}{3}$

$F(3) - F(0) = \boxed{3} \checkmark$

3) Let R be the region bounded by the curves $y = \sqrt{\ln\left(\frac{x}{2}\right)}$, $y = \sqrt{\ln\left(\frac{x^2}{4}\right)}$ and $y = 1$.

- Express the volume of R revolved around the x -axis using Washers.
- Draw the region R , a typical slice, and the washer or disk generated by the slice revolving around the x -axis.
- Express the volume of R revolved around the x -axis using Shells.
- Draw the region R , a typical slice, and the shell generated by the slice revolving around the x -axis.
- Choose one of the two integrals to calculate the exact volume.

X AXIS



$$V = \pi \int_a^b (R(y)^2 - r(y)^2) dx$$

WASHER

$$\pi \int_2^4 \left(\sqrt{\ln\left(\frac{x^2}{4}\right)}^2 - \left(\sqrt{\ln\left(\frac{x}{2}\right)} \right)^2 \right) dx$$

$$V = 10.56637$$

$$\sqrt{\ln\frac{x}{2}} = \sqrt{\ln\frac{x^2}{4}}$$

$$e \cdot \ln\frac{x}{2} = \ln\frac{x^2}{4}$$

$$\frac{x}{2} = \frac{x^2}{4}$$


$$x = 2$$

$$\sqrt{\ln\frac{x}{2}} = 1$$

$$\sqrt{\ln\frac{x}{2}} = 1 \Rightarrow \ln\frac{x}{2} = 1 \Rightarrow x = 2e$$

$$x = 2e$$

SHELL $2\pi \int_2^{2e} x \left(\sqrt{\ln\left(\frac{x^2}{4}\right)} - \sqrt{\ln\left(\frac{x}{2}\right)} \right) dx$




$$2\pi \int_2^{2e} x \left(\sqrt{\ln\left(\frac{x^2}{4}\right)} - \sqrt{\ln\left(\frac{x}{2}\right)} \right) dx$$

Name _____ ID# _____

4) Sketch the graph of a function with following limits, and classify any discontinuities.

- $\lim_{x \rightarrow -3} = 5$
- $f(-3) = 2$
- $\lim_{x \rightarrow 2^-} f(x) = -2$
- $\lim_{x \rightarrow 2^+} f(x) = 1$
- $f(2) = 3$
- $\lim_{x \rightarrow 5^+} f(x) = \infty$
- $\lim_{x \rightarrow 5^-} f(x) = -\infty$
- $\lim_{x \rightarrow \infty} f(x) = 1$




1 = INFINITY
DISCONTINUITY
2 = REMOVEABLE
DISCONTINUITY
3 = JUMP DISCONTINUITY

5) Find the exact average value of the function $f(x) = \tanh(x)$ on the interval $-1 \leq x \leq 2$.

Graph the function and indicate the approximate average value on the graph.


AVERAGE FUNCTION VALUE

$$\frac{1}{b-a} \int_a^b f(x) dx$$




$$\frac{1}{2-(-1)} \int_{-1}^2 \tanh(x) dx = 0.297$$

AVERAGE VALUE



Name _____ ID# _____

6) A sheet of paper measuring 6in by 6in is constructed into an open top box by removing four corners. Find the exact dimensions of the box with maximum volume. State the objective function and interval of interest.



MAX VALUE

$$(6-2x)(6-2x)x$$

OBJECTIVE FUNCTION $(6-2x)^2 x$

$$6/2 = 3_{\text{max}}$$



DOMAIN OF INTEREST

$$0 \leq x \leq 3$$

$$(6-2)^2$$

$$(4)^2 = 16 \text{ in}^3 \text{ MAX}$$

LOW is 0 in³ NO W



7) A payload is released at an elevation of 700 meters from a hot air balloon that is rising at a rate of 11 meters per second. Assume no wind resistance and acceleration due to gravity is 9.8 meters per second squared.

a) Find the velocity of the payload. $v(t) =$

$$v(t) = (700 + 11t) - 9.8t^2$$




$$v(t) = (700 + 11t) - 9.8t^2$$

b) Find the position of the payload. $s(t) =$

$$s(t) = 9.8t^2 + 11t - 19.6t$$

$$s(t) = 11 - 19.6t$$



c) The payload's highest point is 700 meters above the ground at time = 0 seconds.

d) Find the time when the payload strikes the ground.





$$9.03 \text{ sec}$$

Name _____ ID# _____

8) A block of ice in the shape of a perfect cube is melting. Each side of the ice cube is decreasing at a constant rate of 5 mm per minute.

a) How quickly is the surface area of the ice cube shrinking when each side is 50 mm?

b) How quickly is the volume of the ice cube shrinking when each side is 50 mm?



$$5 \text{ mm/min}$$

$$60 \text{ mm}^2/\text{SEC}$$

$$\frac{dA}{dt} =$$

$$(50-5)^2 \cdot 6(50) = 6(45)^2 \cdot 5 \text{ mm}^2/\text{min} = 6(2025) \cdot 5 = 60375 \text{ mm}^2/\text{min}$$

$$V = (50-5)^3 = 45^3 \text{ mm}^3/\text{min} = 91125 \text{ mm}^3/\text{min}$$



$$5 \times 2 = 10$$

$$6 \times (50 \times 2) = 5 \text{ mm/min}$$

$$\frac{dV}{dt} = -125 \text{ mm}^3/\text{SEC}$$

9) Given the graph of power $P(t)$ over a 24 hour period below, and that power $P(t) = E'(t)$.


Graph energy $E(t)$ and estimate the time(s) that energy is maximized and minimized.
(Hint use the first derivative test)



Name _____ ID# _____

10) Find the exact value of $\int_{-3}^3 \sqrt{9-x^2} dx$ using geometry.

$$\int_{-3}^3 \sqrt{9-x^2} dx = 14.137$$



$$9\pi \times 1/2 = 9/2\pi = V$$

Find the exact value of $\int_{-3}^3 \sqrt{9-x^2} dx$ using integration techniques.

$$\int_{-3}^3 \sqrt{9-x^2} dx = \int_{-3}^0 \sqrt{9-x^2} dx + \int_0^3 \sqrt{9-x^2} dx$$

Handwritten notes include: $(3-x)^2$, $2x = -3 dx$, and $dx = -3 dx$.



$$A^2 + B^2 = C^2$$

$$x \quad B \quad \sqrt{C^2 - x^2}$$

Hand-drawn diagram of a right triangle with hypotenuse 3 and base $\sqrt{9-x^2}$.

$$x \quad \sqrt{9-x^2} = 3 \ln(10)$$

Hand-drawn diagram of a right triangle with hypotenuse 3 and base $\sqrt{9-x^2}$.