A (Brief) Introduction to SEM

DATA 695 Research Capstone Project

David Errington

Master of Data Science & Analytics Student
Faculty of Graduate Studies

June 13, 2025



Table of Contents

- Territorial Acknowledgment
- Chapter 1: Introduction, Background, & Review
 - Review of Simple/Multiple Regression
 - Linear Regression as a Structural Equation Model
 - Limitations of Multiple Regression
- Chapter 2: Path Analysis
- Chapter 3: Confirmatory vs. Exploratory Factor Analysis
- Chapter 4: Structural Equation Models w/ Latent Variables
- References



The University of Calgary, located in the heart of Southern Alberta, both acknowledges and pays tribute to the traditional territories of the peoples of Treaty 7, which include the Blackfoot Confederacy (comprised of the Siksika, the Piikani, and the Kainai First Nations), the Tsuut'ina First Nation, and the Stoney Nakoda (including Chiniki, Bearspaw, and Goodstoney First Nations). The City of Calgary is also home to the Métis Nation of Alberta (Districts 5 and 6).



Chapter 1: Introduction, Background, & Review




What is Structural Equation Modelling?

"Structural equation modeling is a linear model framework that models both simultaneous regression equations with latent variables."

— Johnny Lin, PhD (2024)




What is Structural Equation Modelling?





What is Structural Equation Modelling?





Definitions / Terminology

- Exogenous (a.k.a. independent) variables:
 - Variables that are not expressed as a function of other variables; they exist "outside" the system of variables under study.
 - Notation:
 - *x* (single); or
 - x_i (multiple)
- Endogenous (a.k.a. dependent) variables:
 - Variables that are expressed as a function of one or more other variables; they exist "inside" the system of variables under study.
 - Notation:
 - *y* (single); or
 - y_i (multiple)



Definitions / Terminology

- Observed variable(s):
 - Variables that can be directly measured or "observed".
 - Examples: Height, weight, age, etc.
- Latent variable(s):
 - Variables that (usually) cannot be directly measured; often "inferred".
 - Example: Intelligence.
 - Notation:
 - η "eta" (single); or
 - η_i (multiple)




• Step 1: Start with linear regression equation.

$$y = \beta_0 + \beta_1 x_1 + \varepsilon$$



Step 2: Add subscript i to each variable.





Step 3: Take the intercept...

$$y_i = \beta_0 + \beta_1 x_{1i} + \varepsilon$$
Intercept




• Step 3: And replace it with α .

$$y_i = \alpha + \beta_1 x_{1i} + \varepsilon$$
Intercept




• Step 4: Take the regression coefficient(s)...






• Step 4: And replace it(them) with γ .






Step 5: Take the residual error term...





• Step 5: And replace it with ζ_i .





References

Lin, J. (2024). Introduction to Structural Equation Modeling (SEM) in R with lavaan. OARC Stats – Statistical Consulting Web Resources | UCLA. https://stats.oarc.ucla.edu/r/seminars/rsem/

