

1. Write each power series as an ordinary function.

a.)  $\sum_{n=5}^{\infty} x^n$

Notice that we can re-write this as

$$\sum_{n=5}^{\infty} x^n = \sum_{n=0}^{\infty} x^n - \sum_{n=0}^4 x^n$$

Since we know the sum starting from 0, we get

$$\sum_{n=5}^{\infty} x^n = \frac{1}{1-x} - 1 - x - x^2 - x^3 - x^4; \quad x \in (-1, 1)$$

b.)  $\sum_{n=0}^{\infty} 2^n x^n$

Notice that we can write  $2^n x^n = (2x)^n$ , then let  $y = 2x$  and we obtain

$$\sum_{n=0}^{\infty} 2^n x^n = \sum_{n=0}^{\infty} (2x)^n = \sum_{n=2(0)}^{2(\infty)} y^n = \sum_{n=0}^{\infty} y^n = \frac{1}{1-y} = \frac{1}{1-2x}$$

Note that since  $y \in (-1, 1)$  we have  $2x \in (-1, 1)$  or  $x \in (-1/2, 1/2)$ .

c.)  $\sum_{n=0}^{\infty} \frac{(-3)^{n+1} x^n}{5^{n-1}}$

We have to do some extensive algebra on this. Notice that we have

$$\frac{(-3)^{n+1} x^n}{5^{n-1}} = \frac{(-3)(-3)^n x^n}{5^{n-1}} = \frac{(5)(-3)(-3)^n x^n}{5^n} = -15 \left( \frac{(-3)^n x^n}{5^n} \right) = -15 \left( \frac{(-3x)^n}{5^n} \right) = -15 \left( \frac{-3x}{5} \right)^n$$

Thus we obtain

$$\sum_{n=0}^{\infty} \frac{(-3)^{n+1} x^n}{5^{n-1}} = \sum_{n=0}^{\infty} -15 \left( \frac{-3x}{5} \right)^n = -15 \sum_{n=0}^{\infty} \left( \frac{-3x}{5} \right)^n = -15 \left( \frac{1}{1 - \left( \frac{-3x}{5} \right)} \right) = -15 \left( \frac{1}{1 + \frac{3x}{5}} \right) = -15 \frac{5}{5 + 3x}$$

With radius of convergence  $\frac{3x}{5} \in (-1, 1)$  which gives us  $x \in (-5/3, 5/3)$ .

d.)  $\sum_{n=4}^{\infty} nx^{n-1}$

For this one we have to recognize that if this were  $\sum x^{n-1}$  it would be relatively easy to fix. Notice that

$$\frac{d}{dx} x^n = nx^{n-1}$$

Moreover, we obtain

$$\frac{d}{dx} \left( \sum_{n=4}^{\infty} x^n \right) = \frac{d}{dx} (x^4 + x^5 + x^6 + \dots) = 4x^3 + 5x^4 + 6x^5 + \dots = \sum_{n=4}^{\infty} nx^{n-1}$$

Since we know the first sum above we obtain

$$\sum_{n=4}^{\infty} nx^{n-1} = \frac{d}{dx} \left( \sum_{n=4}^{\infty} x^n \right) = \frac{d}{dx} \left( \sum_{n=0}^{\infty} x^n - \sum_{n=0}^3 x^n \right) = \frac{d}{dx} \left( \frac{1}{1-x} - 1 - x - x^2 - x^3 \right) = \frac{1}{(1-x)^2} - 1 - 2x - 3x^2$$

where  $x \in (-1, 1)$

e.)  $\sum_{n=0}^{\infty} n^2 x^{n-1}$

Just like in  $d$  we recognize that we need one derivative to get  $nx^{n-1}$ , so a second derivative should get us close to  $n^2 x^{n-1}$ . Notice that we have

$$x \frac{d}{dx} \left( \sum_{n=0}^{\infty} x^n \right) = \sum_{n=0}^{\infty} x \frac{d}{dx} x^n = \sum_{n=0}^{\infty} nx^n$$

Moreover, we have

$$\frac{d}{dx} \left( \sum_{n=0}^{\infty} nx^n \right) = \sum_{n=0}^{\infty} \frac{d}{dx} nx^n = \sum_{n=0}^{\infty} n^2 x^{n-1}$$

which gives us our target series. Thus, using geometric series, we obtain

$$\sum_{n=0}^{\infty} n^2 x^{n-1} = \frac{d}{dx} \left( \sum_{n=0}^{\infty} nx^n \right) = \frac{d}{dx} \left( x \frac{d}{dx} \left( \sum_{n=0}^{\infty} x^n \right) \right) = \frac{d}{dx} \left( x \frac{d}{dx} \left( \frac{1}{1-x} \right) \right) = \frac{d}{dx} \left( \frac{x}{(1-x)^2} \right) = \frac{(1-x)^2 - 2x(1-x)}{(1-x)^4}$$

where  $x \in (-1, 1)$  because we used a geometric series.

f.)  $\sum_{n=1}^{\infty} \frac{x^{n+3}}{n}$

Something to notice here is that

$$\frac{d}{dx} \left( \frac{x^n}{n} \right) = x^{n-1}$$

which is a geometric series that we can solve. This is backwards from what we did in  $e$  and  $d$ . Since the reverse of a derivative is an integral, we need to use an integral. Thus we obtain

$$\int \sum_{n=1}^{\infty} x^{n-1} dx = \int (1 + x + x^2 + x^3 + \dots) dx = x + \frac{x^2}{2} + \frac{x^3}{3} + \dots = \sum_{n=1}^{\infty} \frac{x^n}{n}$$

Thus we can describe the series above as

$$\sum_{n=1}^{\infty} \frac{x^{n+3}}{n} = x^3 \int \sum_{n=1}^{\infty} x^{n-1} dx = x^3 \int \frac{1}{x} \left( \frac{1}{1-x} \right) dx = x^3 (\ln(|x|) - \ln(|1-x|)) = x^3 \ln \left( \left| \frac{x}{1-x} \right| \right)$$

where we used partial fractions to solve that integral. Note we used a geometric series so  $x \in (-1, 1)$ , moreover,  $x \neq 0$  since  $\ln 0$  doesn't exist.

g.)  $\sum_{n=1}^{\infty} (-1)^n \frac{x^n}{2^n n!}$

Notice we obtain

$$\frac{x^n}{2^n n!} = \frac{\left(\frac{x}{2}\right)^n}{n!}$$

So what we can do is recognize that since

$$e^x = \sum_{n=1}^{\infty} (-1)^n \left( \frac{x^n}{n!} \right)$$

then we obtain

$$\sum_{n=1}^{\infty} (-1)^n \frac{x^n}{2^n n!} = \sum_{n=1}^{\infty} (-1)^n \left( \frac{(x/2)^n}{n!} \right) = e^{x/2}$$

Where  $x \in \mathbb{R}$  because  $e^x$  is valid everywhere.

h.)  $\sum_{n=2}^{\infty} (-1)^n \left(\frac{2}{5}\right)^n \frac{x^{2n+1}}{2n+1}$

Notice that  $\int x^{2n} dx = \frac{x^{2n+1}}{2n+1}$ , so we will need to use that at some point in this problem. Using some algebra we obtain,

$$(-1)^n \left(\frac{2}{5}\right)^n \frac{x^{2n+1}}{2n+1} = (-1)^n \left(\frac{2}{5}\right)^n \int x^{2n} dx = \int (-1)^n \left(\frac{2}{5}\right)^n x^{2n} dx = \int (-1)^n \left(\frac{2}{5}\right)^n (x^2)^n dx = \int \left(-\frac{2}{5}x^2\right)^n dx$$

Moreover, the series  $\sum \left(-\frac{2}{5}x^2\right)^n$  is relatively easy to solve using geometric series theorems. Thus we obtain

$$\begin{aligned} \sum_{n=2}^{\infty} (-1)^n \left(\frac{2}{5}\right)^n \frac{x^{2n+1}}{2n+1} &= \int \sum_{n=2}^{\infty} \left(-\frac{2}{5}x^2\right)^n dx \\ &= \int \sum_{n=0}^{\infty} \left(-\frac{2}{5}x^2\right)^n - \sum_{n=0}^1 \left(-\frac{2}{5}x^2\right)^n dx \\ &= \int \frac{1}{1 - \left(-\frac{2}{5}x^2\right)} - 1 - \left(-\frac{2}{5}x^2\right) dx \\ &= \int \frac{1}{1 + \frac{2}{5}x^2} dx - x + \frac{2}{15}x^3 \\ &= \int \frac{1}{1 + \left(\sqrt{\frac{2}{5}}x\right)^2} dx - x + \frac{2}{15}x^3 \\ &= \sqrt{\frac{5}{2}} \arctan\left(\sqrt{\frac{2}{5}}x\right) - x + \frac{2}{15}x^3 \end{aligned}$$

Notice that we used a geometric series, so we need  $\frac{2}{5}x^2 \in (-1, 1)$  or  $x \in \left(-\sqrt{\frac{5}{2}}, \sqrt{\frac{5}{2}}\right)$ .

**2. Use any method to find the exact value of each of the following convergent series.**

a.)  $\sum_{n=0}^{\infty} 3 \left(\frac{-2}{3}\right)^n$

Let  $f(x) = \sum_{n=0}^{\infty} x^n = \frac{1}{1-x}$ , the target series is given by

$$\sum_{n=0}^{\infty} 3 \left(\frac{-2}{3}\right)^n = 3f\left(\frac{-2}{3}\right) = 3 \left(\frac{1}{1 - \left(\frac{-2}{3}\right)}\right) = 3 \left(\frac{3}{5}\right) = \frac{9}{5}$$

b.)  $\sum_{n=4}^{\infty} \frac{(-1)^{n+2} n^{-3}}{2}$

c.)  $\sum_{n=1}^{\infty} n^2 \left(\frac{1}{2}\right)^n$

d.)  $\sum_{n=0}^{\infty} n(n-1) \left(\frac{3}{4}\right)^{n+1}$

e.)  $\sum_{n=0}^{\infty} \frac{(\ln 2)^n}{n!}$

f.)  $\sum_{n=2}^{\infty} (-1)^n \frac{9^n}{(2n)!}$