

Linear combinations of projections in operator algebras.

Victor Kaftal

University of Cincinnati

U. Virginia

November 18, 2014

Joint work with

Ping W. Ng (University of Louisiana)

Shuang Zhang (University of Cincinnati)

Outline

- 1 Questions
- 2 Linear combinations of projections
- 3 Positive linear combinations of projections (PCP)
- 4 Infinite sums of projections
- 5 Finite sums of projections & open questions

Questions

- For which C^* -algebras are all elements linear combinations of projections? If not all, which are?

Questions

- For which C^* -algebras are all elements linear combinations of projections? If not all, which are?
- If the elements are positive, can the coefficients be chosen to be positive? **PCP: positive combinations of projections.**

$b = \sum_1^n \lambda_j p_j$, $\lambda_j > 0$ where p_j are projections
not necessarily mutually orthogonal.

Questions

- For which C^* -algebras are all elements linear combinations of projections? If not all, which are?
- If the elements are positive, can the coefficients be chosen to be positive? **PCP: positive combinations of projections.**

$b = \sum_1^n \lambda_j p_j$, $\lambda_j > 0$ where p_j are projections
not necessarily mutually orthogonal.

The questions that started us on this project were:

- Which positive elements are infinite sums of projections?
(Converging in the strict topology in a multiplier algebra or in the strong topology in a von Neumann algebra.) This question originated in frame theory (**Dykema, Larson & all, 2004**)
It is now solved in $B(H)$.
- Which positive elements are finite sums of projections?
Still open in $B(H)$.

Which W^* -algebras are the span of their projections?

- $B(H)$:
 - Fillmore, 1967 ($n=257$),
then progress via link to commutators:
 - Percy & Topping, 1967 ($n=16$),
 - Matsumoto, 1984 ($n=10$)

Which W^* -algebras are the span of their projections?

- $B(H)$:
 - Fillmore, 1967 ($n=257$),
then progress via link to commutators:
 - Percy & Topping, 1967 ($n=16$),
 - Matsumoto, 1984 ($n=10$)
- All W^* -algebras with the exception of finite type I algebras with infinite dimensional center (e.g., $\bigoplus_1^\infty M_n(\mathbb{C})$) or algebras having such a direct summand. (Percy & Topping (1967), Fack & de La Harpe (1980), Goldstein & Paszkiewicz (1992))

Universal constants

Definition

A C^* -algebra \mathcal{A} has universal constants V, N if for every $b = b^* \in \mathcal{A}$,

$$b = \sum_1^N \alpha_j p_j \quad \text{and} \quad \sum_1^N |\alpha_j| \leq V \|b\|$$

with $\alpha_j \in \mathbb{R}$ and $p_j \in \mathcal{A}$ projections.

Universal constants for a W^* -algebra M

- M properly infinite, $N = 6$, $V = 8$
- M type II_1 , $N = 12$, $V = 14$
- M direct sum of m matrix algebras: $N = m + 4$, $V = m + 4$.

These universal constants are quite useful. E.g., Fong&Murphy's proof that positive invertible in $B(H)$ are positive combinations of projections.

Which C^* -algebras are the span of their projections?

- Purely infinite unital simple C^* -algebras.

Which C^* -algebras are the span of their projections?

- **Purely infinite unital** simple C^* -algebras.
- **Unital** simple finite **real rank zero** algebras with a unique tracial state satisfying the property of **strict comparison of projections**: $\tau(p) < \tau(q) \Rightarrow p \precsim q$ (Murray-von Neumann subequivalence);

Which C^* -algebras are the span of their projections?

- **Purely infinite unital** simple C^* -algebras.
- **Unital** simple finite **real rank zero** algebras with a unique tracial state satisfying the property of **strict comparison of projections**: $\tau(p) < \tau(q) \Rightarrow p \precsim q$ (Murray-von Neumann subequivalence);
- **Unital** simple AF-algebras, AT-algebras, or AH-algebras (with bounded dimension growth) with **real rank zero** and with $|\partial_e(T(\mathcal{A}))| < \infty$.: The collection of the tracial states $T(\mathcal{A})$ is convex and compact (in the w^* -topology), $\partial_e(T(\mathcal{A}))$ is the collection of the extreme points of $T(\mathcal{A})$.

All the above results due mainly to **Marcoux, 1998-2010** based on earlier work of **Fack and Thompsen**.

Which C^* -algebras are the span of their projections?

- **Purely infinite unital** simple C^* -algebras.
- **Unital** simple finite **real rank zero** algebras with a unique tracial state satisfying the property of **strict comparison of projections**: $\tau(p) < \tau(q) \Rightarrow p \precsim q$ (Murray-von Neumann subequivalence);
- **Unital** simple AF-algebras, AT-algebras, or AH-algebras (with bounded dimension growth) with **real rank zero** and with $|\partial_e(T(\mathcal{A}))| < \infty$.: The collection of the tracial states $T(\mathcal{A})$ is convex and compact (in the w^* -topology), $\partial_e(T(\mathcal{A}))$ is the collection of the extreme points of $T(\mathcal{A})$.

All the above results due mainly to **Marcoux, 1998-2010** based on earlier work of **Fack and Thompsen**.

More C^* -algebras that are the span of their projections

The following includes all the finite algebras studied by Marcoux:

Theorem (2014)

A unital simple separable finite C^ -algebra \mathcal{A} with real rank zero, stable rank one, strict comparison of projections is the linear span of its projections **if and only if** $|\partial_e(T(\mathcal{A}))| < \infty$.*

The key steps are

- elements in the kernel of all traces are sums of commutators (this holds also when $|\partial_e(T(\mathcal{A}))| = \infty$.)
- commutators are linear combinations of projections (by Marcoux this holds under very general conditions)
- every element can be decomposed as a linear combination of projections plus an element in the kernel of all traces (this **requires** $|\partial_e(T(\mathcal{A}))| < \infty$.)

C^* -algebras that are not the span of their projections

Already mentioned:

- W^* -algebras with finite type I direct summand with infinite dimensional center
- C^* -algebras as in previous slide but with $|\partial_e(T(\mathcal{A}))| = \infty$.

C^* -algebras that are not the span of their projections

Already mentioned:

- W^* -algebras with finite type I direct summand with infinite dimensional center
- C^* -algebras as in previous slide but with $|\partial_e(T(\mathcal{A}))| = \infty$.

What else?

- Some nonunital algebras, e.g. \mathcal{K} ($\mathcal{K} = K(H)$): linear combinations of projections must have finite rank.

Theorem (in press)

Let \mathcal{A} finite as previous slide and with $|\partial_e(T(\mathcal{A}))| < \infty$. Then $b \in \mathcal{A} \otimes \mathcal{K}$ is a linear combination of projections in $\mathcal{A} \otimes \mathcal{K}$ if and only if $\tau(R_b) < \infty \forall \tau \in T(\mathcal{A})$ (R_b range projection of b .)

Positive combinations of projections in $B(H)$

- If $b \in B(H)_+ \setminus \mathcal{K}$ then b is PCP (Fillmore, 1969).
- $\mathcal{K}_+ \ni b = \sum_1^n \lambda_j p_j$ with $\lambda_j > 0 \Rightarrow b$ has finite rank.
 $b \geq 0$ has finite rank $\Rightarrow b$ is PCP. (Both are obvious)
- If $b \in B(H)_+ \setminus \mathcal{K}$ then b is PCP. (Fong&Murphy, 1985).

In summary,

Theorem (Fong&Murphy, 1985)

$b \in B(H)_+$ is PCP if and only if either $b \in B(H)_+ \setminus \mathcal{K}$ or b has finite rank.

Positive combinations of projections in $B(H)$

- If $b \in B(H)_+ \setminus \mathcal{K}$ then b is PCP (Fillmore, 1969).
- $\mathcal{K}_+ \ni b = \sum_1^n \lambda_j p_j$ with $\lambda_j > 0 \Rightarrow b$ has finite rank.
 $b \geq 0$ has finite rank $\Rightarrow b$ is PCP. (Both are obvious)
- If $b \in B(H)_+ \setminus \mathcal{K}$ then b is PCP. (Fong&Murphy, 1985).

In summary,

Theorem (Fong&Murphy, 1985)

$b \in B(H)_+$ is PCP if and only if either $b \in B(H)_+ \setminus \mathcal{K}$ or b has finite rank.

Surprisingly, not much was known for other C^* -algebras or even W^* -algebras.

Positive combinations of projections in $B(H)$

- If $b \in B(H)_+ \setminus \mathcal{K}$ then b is PCP (Fillmore, 1969).
- $\mathcal{K}_+ \ni b = \sum_1^n \lambda_j p_j$ with $\lambda_j > 0 \Rightarrow b$ has finite rank.
 $b \geq 0$ has finite rank $\Rightarrow b$ is PCP. (Both are obvious)
- If $b \in B(H)_+ \setminus \mathcal{K}$ then b is PCP. (Fong&Murphy, 1985).

In summary,

Theorem (Fong&Murphy, 1985)

$b \in B(H)_+$ is PCP if and only if either $b \in B(H)_+ \setminus \mathcal{K}$ or b has finite rank.

Surprisingly, not much was known for other C^* -algebras or even W^* -algebras.

Yet, knowing about PCP is useful, e.g., it is the first step to obtain decomposition into sum of projections.

Positive combination of projections - the “easy” cases

Theorem (with H. Halpern, 2013)

The following positive elements in a W^ -algebra are PCP:*

- *Simple ($M_n(\mathbb{C})$, type II_1 or type III σ -finite factors): all.*
- *Type II_∞ factors: iff either b is not in the **Breuer ideal** of compact operators or the range projection R_b is finite.*
- *“Small center” (finite direct sums of factors): same as above.*
- *“Large center”: obstruction in terms of the **central essential spectrum**. (E.g., $\bigoplus_n \frac{1}{n} 1_n \in \bigoplus B(H_n)$ is not PCP.)*

Positive combination of projections - the “easy” cases

Theorem (with H. Halpern, 2013)

The following positive elements in a W^ -algebra are PCP:*

- *Simple ($M_n(\mathbb{C})$, type II_1 or type III σ -finite factors): all.*
- *Type II_∞ factors: iff either b is not in the **Breuer ideal** of compact operators or the range projection R_b is finite.*
- *“Small center” (finite direct sums of factors): same as above.*
- *“Large center”: obstruction in terms of the **central essential spectrum**. (E.g., $\bigoplus \frac{1}{n} 1_n \in \bigoplus B(H_n)$ is not PCP.)*

Theorem (2011)

Let \mathcal{A} be a purely infinite simple σ -unital C^ -algebra. Then every positive element of \mathcal{A} and of **the multiplier algebra $\mathcal{M}(\mathcal{A})$** is a PCP.*

Positive combination of projections - the “easy” cases

Theorem (with H. Halpern, 2013)

The following positive elements in a W^ -algebra are PCP:*

- *Simple ($M_n(\mathbb{C})$, type II_1 or type III σ -finite factors): all.*
- *Type II_∞ factors: iff either b is not in the **Breuer ideal** of compact operators or the range projection R_b is finite.*
- *“Small center” (finite direct sums of factors): same as above.*
- *“Large center”: obstruction in terms of the **central essential spectrum**. (E.g., $\bigoplus \frac{1}{n} 1_n \in \bigoplus B(H_n)$ is not PCP.)*

Theorem (2011)

Let \mathcal{A} be a purely infinite simple σ -unital C^ -algebra. Then every positive element of \mathcal{A} and of **the multiplier algebra $\mathcal{M}(\mathcal{A})$** is a PCP.*

More on multiplier algebras

By Marcoux, $\mathcal{M}(\mathcal{A} \otimes \mathcal{K})$ is the span of its projections. But more:

Theorem (in press)

Let \mathcal{A} be unital simple separable C^ -algebra with real rank zero, stable rank one, strict comparison of projections and with $\partial_e(T(\mathcal{A}))$ finite.*

- *If $P \in \mathcal{M}(\mathcal{A} \otimes \mathcal{K})$ is a projection and $B \in P\mathcal{M}(\mathcal{A} \otimes \mathcal{K})P$, then B is a linear combination of projections in $P\mathcal{M}(\mathcal{A} \otimes \mathcal{K})P$.*
- *If $\mathcal{M}(\mathcal{A} \otimes \mathcal{K})$ has real rank zero, then $B \in \mathcal{M}(\mathcal{A} \otimes \mathcal{K})_+$ is a PCP if and only if and only if $\tau(R_B) < \infty$ for every $\tau \in T(\mathcal{A})$ for which $B \in I_\tau := \overline{\text{span}}\{X \in \mathcal{M}(\mathcal{A} \otimes \mathcal{K})_+ \mid \tau(X) < \infty\}$.*

More on multiplier algebras

By Marcoux, $\mathcal{M}(\mathcal{A} \otimes \mathcal{K})$ is the span of its projections. But more:

Theorem (in press)

Let \mathcal{A} be unital simple separable C^ -algebra with real rank zero, stable rank one, strict comparison of projections and with $\partial_e(T(\mathcal{A}))$ finite.*

- *If $P \in \mathcal{M}(\mathcal{A} \otimes \mathcal{K})$ is a projection and $B \in P\mathcal{M}(\mathcal{A} \otimes \mathcal{K})P$, then B is a linear combination of projections in $P\mathcal{M}(\mathcal{A} \otimes \mathcal{K})P$.*
- *If $\mathcal{M}(\mathcal{A} \otimes \mathcal{K})$ has real rank zero, then $B \in \mathcal{M}(\mathcal{A} \otimes \mathcal{K})_+$ is a PCP if and only if and only if $\tau(R_B) < \infty$ for every $\tau \in T(\mathcal{A})$ for which $B \in I_\tau := \overline{\text{span}}\{X \in \mathcal{M}(\mathcal{A} \otimes \mathcal{K})_+ \mid \tau(X) < \infty\}$.*

Real rank zero needed to use Brown's interpolation theorem. We recently removed the condition that $\mathcal{M}(\mathcal{A} \otimes \mathcal{K})$ has real rank zero by proving that $\mathcal{M}(\mathcal{A} \otimes \mathcal{K})$ has strict comparison of positive elements.

Infinite sums of projections in $B(H)$

Theorem (Choi & Wu, 1994-2014)

If $b \in B(H)_+$ and $\|b\|_{\text{ess}} > 1$ then b is a sum of finitely many projections.

Theorem (Dykema, Larson & all, 2004)

If $b \in B(H)_+$ and $\|b\|_{\text{ess}} > 1$ then b is a sum of infinitely many rank-one projections.

Infinite sums of projections in $B(H)$

Theorem (Choi & Wu, 1994-2014)

If $b \in B(H)_+$ and $\|b\|_{\text{ess}} > 1$ then b is a sum of finitely many projections.

Theorem (Dykema, Larson & all, 2004)

If $b \in B(H)_+$ and $\|b\|_{\text{ess}} > 1$ then b is a sum of infinitely many rank-one projections.

Theorem (2009)

Let $b \in B(H)_+$, then b is a sum of projections iff either $\text{Tr}((b - R_b)_+) = \infty$ or $\text{Tr}((b - R_b)_+) - \text{Tr}((b - R_b)_-) \in \mathbb{N} \cup \{0\}$.

Infinite sums of projections in $B(H)$

Theorem (Choi & Wu, 1994-2014)

If $b \in B(H)_+$ and $\|b\|_{\text{ess}} > 1$ then b is a sum of finitely many projections.

Theorem (Dykema, Larson & all, 2004)

If $b \in B(H)_+$ and $\|b\|_{\text{ess}} > 1$ then b is a sum of infinitely many rank-one projections.

Theorem (2009)

Let $b \in B(H)_+$, then b is a sum of projections iff either $\text{Tr}((b - R_b)_+) = \infty$ or $\text{Tr}((b - R_b)_+) - \text{Tr}((b - R_b)_-) \in \mathbb{N} \cup \{0\}$.

Where does that integer come from?

The integer in the N&S condition

Let $b = \sum_1^\infty p_j$ with $p_j = \xi_j \otimes \xi_j$ rank-one projections.

$((\xi \otimes \eta)(\zeta) := (\zeta, \eta)\xi$ rank one operator, $\xi, \eta, \zeta \in H$).

Let $\{\chi_j\}$ be an o.n. basis,

$\theta = \sum_j^\infty \chi_j \otimes \xi_j$ frame transform, aka analysis operator

$\theta = wb^{1/2}$ polar decomposition, so $w^*w = R_b$

$\theta\theta^*$ Gram matrix of $\{\xi_j\}$

Now, an easy computation:

$$E(bw^*) = E(\theta\theta^*) = 1 \quad \text{and} \quad b = (b - R_b)_+ - (b - R_b)_- + R_b$$

The integer in the N&S condition

Let $b = \sum_1^\infty p_j$ with $p_j = \xi_j \otimes \xi_j$ rank-one projections.

$((\xi \otimes \eta)(\zeta) := (\zeta, \eta)\xi$ rank one operator, $\xi, \eta, \zeta \in H$).

Let $\{\chi_j\}$ be an o.n. basis,

$\theta = \sum_j^\infty \chi_j \otimes \xi_j$ frame transform, aka analysis operator

$\theta = wb^{1/2}$ polar decomposition, so $w^*w = R_b$

$\theta\theta^*$ Gram matrix of $\{\xi_j\}$

Now, an easy computation:

$$E(bw^*) = E(\theta\theta^*) = 1 \quad \text{and} \quad b = (b - R_b)_+ - (b - R_b)_- + R_b$$

$$E\left(w((b - R_b)_+ - (b - R_b)_- + R_b)w^*\right) = E(1) \quad \text{and} \quad wR_bw^* = ww^*$$

The integer in the N&S condition

Let $b = \sum_1^\infty p_j$ with $p_j = \xi_j \otimes \xi_j$ rank-one projections.

$((\xi \otimes \eta)(\zeta) := (\zeta, \eta)\xi$ rank one operator, $\xi, \eta, \zeta \in H$).

Let $\{\chi_j\}$ be an o.n. basis,

$\theta = \sum_j^\infty \chi_j \otimes \xi_j$ frame transform, aka analysis operator

$\theta = wb^{1/2}$ polar decomposition, so $w^*w = R_b$

$\theta\theta^*$ Gram matrix of $\{\xi_j\}$

Now, an easy computation:

$$E(bwb^*) = E(\theta\theta^*) = 1 \quad \text{and} \quad b = (b - R_b)_+ - (b - R_b)_- + R_b$$

$$E\left(w((b - R_b)_+ - (b - R_b)_- + R_b)w^*\right) = E(1) \quad \text{and} \quad wR_bw^* = ww^*$$

$$E(w((b - R_b)_+)) = E(w((b - R_b)_-)) + E(1 - ww^*)$$

The integer in the N&S condition

Let $b = \sum_1^\infty p_j$ with $p_j = \xi_j \otimes \xi_j$ rank-one projections.

$((\xi \otimes \eta)(\zeta) := (\zeta, \eta)\xi$ rank one operator, $\xi, \eta, \zeta \in H$).

Let $\{\chi_j\}$ be an o.n. basis,

$\theta = \sum_j^\infty \chi_j \otimes \xi_j$ frame transform, aka analysis operator

$\theta = wb^{1/2}$ polar decomposition, so $w^*w = R_b$

$\theta\theta^*$ Gram matrix of $\{\xi_j\}$

Now, an easy computation:

$$E(bwb^*) = E(\theta\theta^*) = 1 \quad \text{and} \quad b = (b - R_b)_+ - (b - R_b)_- + R_b$$

$$E\left(w((b - R_b)_+ - (b - R_b)_- + R_b)w^*\right) = E(1) \quad \text{and} \quad wR_bw^* = ww^*$$

$$E(w((b - R_b)_+)) = E(w((b - R_b)_-)) + E(1 - ww^*)$$

$$\text{Tr}((b - R_b)_+) - \text{Tr}((b - R_b)_-) = \text{Tr}(I - ww^*) = -\text{ind}(w),$$

The integer in the N&S condition

Let $b = \sum_1^\infty p_j$ with $p_j = \xi_j \otimes \xi_j$ rank-one projections.

$((\xi \otimes \eta)(\zeta) := (\zeta, \eta)\xi$ rank one operator, $\xi, \eta, \zeta \in H$).

Let $\{\chi_j\}$ be an o.n. basis,

$\theta = \sum_j^\infty \chi_j \otimes \xi_j$ frame transform, aka analysis operator

$\theta = wb^{1/2}$ polar decomposition, so $w^*w = R_b$

$\theta\theta^*$ Gram matrix of $\{\xi_j\}$

Now, an easy computation:

$$E(bwbw^*) = E(\theta\theta^*) = 1 \quad \text{and} \quad b = (b - R_b)_+ - (b - R_b)_- + R_b$$

$$E\left(w((b - R_b)_+ - (b - R_b)_- + R_b)w^*\right) = E(1) \quad \text{and} \quad wR_bw^* = ww^*$$

$$E(w((b - R_b)_+)) = E(w((b - R_b)_-)) + E(1 - ww^*)$$

$$\text{Tr}((b - R_b)_+) - \text{Tr}((b - R_b)_-) = \text{Tr}(I - ww^*) = -\text{ind}(w),$$

that is, the integer is the Fredholm index of the partial isometry w .

Infinite sums of projections in W^* -factors

Theorem (2009)

Let M be a W^ -factor with separable predual (e.g., H is separable), let $b \in M_+$*

- If M is type III, then b is a sum of projections iff either $\|b\| > 1$ or b is a projection.*
- If M is type II and b is a sum of projections, then*

$$\tau((b - R_b)_+) \geq \tau((b - R_b)_-).$$

The condition is also sufficient if $\tau((b - R_b)_+) = \infty$ or if b is diagonalizable (i.e., $b = \bigoplus_1^\infty \gamma_n p_n$).

Finite sums of projections in W^* -algebras: Suff condition

Theorem (2013)

Let M be a σ -finite factor. A sufficient condition for $b \in M_+$ to be a finite sum of projections is

- *If M is of type I_∞ : $\|b\|_{\text{ess}} > 1$ (usual essential norm)*
- *If M is of type II_∞ : $\|b\|_{\text{ess}} > 1$ (essential norm relative to the Breuer ideal)*
- *If M is of type III : $\|b\| > 1$ (operator norm)*
- *If M is of type II_1 and b is diagonalizable:
 $\tau((b - R_b)_+) > \tau((b - R_b)_-)$.*

Finite sums of projections in W^* -algebras: Suff condition

Theorem (2013)

Let M be a σ -finite factor. A sufficient condition for $b \in M_+$ to be a finite sum of projections is

- *If M is of type I_∞ : $\|b\|_{\text{ess}} > 1$ (usual essential norm)*
- *If M is of type II_∞ : $\|b\|_{\text{ess}} > 1$ (essential norm relative to the Breuer ideal)*
- *If M is of type III : $\|b\| > 1$ (operator norm)*
- *If M is of type II_1 and b is diagonalizable:
 $\tau((b - R_b)_+) > \tau((b - R_b)_-)$.*

But what if $\|b\|_{\text{ess}} = 1$?

Finite sums of projections in $B(H)$: Nec condition

Choi&Wu's test case, 2014

$b = 1 - k_1 \oplus 1 + k_2$ with $\text{Tr}(k_1) = \text{Tr}(k_2) < \infty$

They showed b can fail to be a finite sum of projections.

Why?

Theorem (with Halpern, 2013)

If b is a finite sum of projections, and $(I - b)_+ R_b \in K$ then $(I - b)_- R_b \in K$ and $(I - b)_+ R_b$ and $(I - b)_- R_b$ generate the same (non-closed) ideal of $B(H)$.

The same holds in W^ -algebras.*

Question

Is the condition that k_1 and k_2 as above generate the same ideals sufficient for b to be a finite sum of projections?

Finite sums of proj in C^* -algebras

Theorem (2012)

Let \mathcal{A} be purely infinite simple C^ -algebra whose K_0 is a torsion group (e.g., \mathcal{O}_n with $n < \infty$) and let $b \in \mathcal{A}_+$. Then b is a finite sum of projections if and only if either $\|b\| > 1$ or b is a projection.*

Question: Is the torsion of K_0 necessary? What about \mathcal{O}_∞ ?

Theorem (2011&2012)

Let \mathcal{A} be a purely infinite simple unital C^ -algebra and $b \in \mathcal{M}(\mathcal{A} \otimes \mathcal{K})_+ \setminus \mathcal{A} \otimes \mathcal{K}$ not a projection.*

If b is a finite or infinite sum of proj then $\|b\|_{\text{ess}} \geq 1$ and $\|b\| > 1$.

If $\|b\|_{\text{ess}} > 1$ then b is a finite sum of projections.

If $\|b\|_{\text{ess}} = 1$ and $\|b\| > 1$ then b is an infinite sum of projections.

Question: can it be a finite sum of projections?

THANK YOU