In Class Problems

Precipitation-Dissolution Equilibrium & Kinetics

CENG 340-Introduction to Environmental Engineering

Instructor: Deborah Sills September 16, 2013

1. Modified from Mihelcic and Zimmerman

Ingesting cadmium may lead to kidney damage. According to the EPA, major sources of cadmium in drinking water include corrosion of galvanized pipes; erosion of natural deposits; discharge from metal refineries; and runoff from waste batteries and paints.

One method to remove metals, such as cadmium, from water is to raise the pH and cause them to precipitate as their metal hydroxides. The precipitation-dissolution equilibrium relationship for cadmium hydroxide is described by the following equation:

where
$$pK_{sp} = 13.85$$

$$|Cd(OH)_2| \stackrel{K_{sp}}{\rightleftharpoons} |Cd^{2+}| + 2[OH^-]$$

(a) In an attempt to remove cadmium from water by precipitating cadmium hydroxide, the pH of water was raised from 6.8 to 8.0. Was the dissolved cadmium concentration reduced to below 100 mg/L at the final pH?

$$PH=8 \ [Ed^{2}] = |D| = |D|$$

(b) What pH is required to reduce cadmium concentration to below the MCL of 5 ppb?

$$pH = ? \quad \text{If I find } [OH^{-}], \text{I } cont \text{ find } pH.$$

$$[CH^{2r}] = S \text{ spb } x \text{ | } ppm = S \times p \text{ | } ppm = S \times ppm = S \times p \text{ | } ppm = S \times ppm = S$$