

Predicting Animal Space-Use with Continuous-Time Markov Chains

Devin S. Johnson^{1,*}, Michaela A. Kratofil^{2,3}, Janelle J. Gardner¹, Robin W. Baird², and
Amanda L. Bradford¹

¹Pacific Islands Fisheries Science Center, National Marine Fisheries Service, NOAA,
Honolulu, Hawaii, USA

²Cascadia Research Collective, Olympia, Washington, USA

³Marine Mammal Institute, Department of Fisheries, Wildlife, and Conservation Sciences,
Oregon State University, Newport, Oregon, USA

July 6, 2026

*Corresponding author: devin.johnson@noaa.gov

Abstract. I. Background and Problem StatementContext: Remote animal telemetry data is widely used in ecology to assess animal space-use and evaluate utilization distributions (UDs).

The Problem (Start Bias): Traditional empirical UD (like kernel density or Brownian bridge estimators) can suffer from "start bias" if tag deployments are too short for an animal to fully explore its habitat, heavily biasing results toward the initial release location.

Existing Solutions & Limitations: Predictive UD attempt to mitigate this by finding the long-run stationary distribution of a movement model. However, previous methods (like Langevin diffusions) require discrete-time approximations, are restricted to continuous covariates, and assume constant diffusion rates across habitats.

II. Proposed MethodThe Innovation: Introduction of a general class of Continuous-Time Markov Chain (CTMC) models that provide exact, closed-form predictions of the UD directly from fitted parameters.

Key Advantages: * Eliminates the need for likelihood approximations during optimization. Accommodates both continuous and discrete habitat covariates. Allows habitat factors to influence diffusion rates, not just advection rates. Software Availability: The authors developed the R package walk to fit these general CTMC models.

III. Application and Case StudyStudy System: The framework was applied to a telemetry dataset tracking five individuals from different social clusters of endangered false killer whales (*Pseudorca crassidens*) in the Hawaiian Islands.

Modeling Approach: A two-stage divide-and-conquer meta-analytic approach was used to approximate a full Bayesian hierarchical model, sharing information between individuals to improve population-level inference.

Covariates Considered: Movement was modeled relative to depth residuals, distance

from land, sea surface temperature (SST), and sea level anomaly (SLA).

IV. Results and Main Conclusions Ecological Findings: Whales exhibited preferences for shallower waters close to the main Hawaiian Islands and warmer SST conditions. Population-level coefficients generally shrank individual variations toward a common mean, except for notable variations in depth and SLA preferences.

Methodological Validation: To verify that the model’s rate truncation did not adversely affect the results, the closed-form exact truncated UDs were compared to fully numerical Eigen-decomposition methods. Bhattacharyya’s Affinity (BA) values ranged from 0.996 to 1, demonstrating that the distributions were virtually identical and validating the accuracy of the closed-form approach.

Key words: CTMC, Hidden Markov model, Limiting distribution, Satellite telemetry, Space-use, Utilization distribution

1 Introduction

Any quick literature search will reveal that remote collection and analysis of animal locations, or telemetry data, have become a ubiquitous and important tool in the advancement of animal ecology. A perennial subject within animal telemetry analysis is the assessment of space-use. That is what aspects of habitat and movement control where an individual generally occurs on the landscape. The topic of animal space-use is often used in conjunction with the terms home range, resource selection, or utilization distribution. Here we are specifically concerned with the concept of utilization distributions (UD) as defined by Powell and Mitchell (2012), a spatial probability distribution that describes an animal’s location at any random time. More specifically, what habitat characteristics influence that

distribution.

Literature on UD is generally divided into two types. The first type is what we term *empirical* UDs. Empirical UDs are constructed based on a summary of observed locations, for example, the traditional kernel density (Keating and Cherry, 2009) or Brownian bridge estimators (Horne et al., 2007). These are empirical because they place positive mass over observed locations. If an animal does not visit a particular area there will be very little if any use estimated for that area depending on how close it is to observed locations. However, absence of observed use may be a symptom of *start bias* (Whitehead and Jonsen, 2013). Start bias results from tag deployments that are not long enough for the animal to fully explore the space and habitat available to it over the course of study. Therefore, spatial use assessments based solely on the observed locations can be biased towards the release location of the animal. This is especially true in recent years where many locations can be obtained over a fine time-scale and thus produce highly autocorrelated paths. To match the increased technological sophistication of location devices, statistical methodology of empirical UD methods have increased to account for the high autocorrelation (e.g., Fleming et al. 2015; Calabrese et al. 2016). However, the most basic solution for reducing start bias is to increase data collection (i.e., tag deployment) interval length (Fieberg et al., 2010). This is easy to prescribe by analysts, but in practice may be difficult to ensure telemetry devices stay active long enough and may be out of the researchers control entirely. This aspect motivates the development of *predictive* UDs.

In order to counter start bias effects researchers can fit resource selection function (RSF) models (Manly et al., 2002) to telemetry data to investigate what space characteristics an animal might prefer relative to available habitat. A map of space preference based on resource selection is used as a UD approximation than is not subject to

start bias because it makes use of spatial habitat characteristics than can be assessed at locations away from where the animal was located during tag deployment. However, Barnett and Moorcroft (2008) describe some general results concerning the weighted distribution form Johnson et al. (2008, 2013) of a resource selection movement model in discrete time. Assuming a symmetric movement kernel, they show that the limiting distribution is proportional to the resource selection function or the square of the resource selection function depending on how large the movement kernel is relative to the spatial variation of the habitat. Although, not explicitly examined, step-selection function (SSF; Fortin et al. 2005; Avgar et al. 2016) models use this same movement kernel form. So, we conjecture similar results might hold for those types of analyses as well. Therefore, while qualitatively similar, RSFs and UD_s can be quite different depending on the scale of movement.

Predictive UD_s seek to blend the habitat association benefits of RSF analysis with movement models such that the resulting UD functions, which appear similar to RSFs, represent the actual UD. Predictive UD_s are heuristically what a kernel estimate would look like if the animal was allowed to move through its habitat for a very long (infinite) time. In essence a movement model is fitted to the telemetry data and the location of the animal is predicted from the movement model for a time far in the future. If it exists, this is the stationary or limiting distribution of the movement process. These limiting distributions are independent of the starting location, thus, removing the start bias (Whitehead and Jonsen, 2013; Wilson et al., 2018). Moorcroft et al. (1999) present an early example of a predictive UD approach modeling home range of coyotes (*Canis latrans*) with a movement model which included a response to scent marks. Wilson et al. (2018) extended their work to continuous-time Markov chains (CTMC) that included habitat based

95 movement. Although, the CTMC predicted UD was derived numerically from the fitted
 96 model components, there was no direct parameter association that could be used to relate
 97 habitat variables with the UD surface. Michelot et al. (2019) used continuous-time and
 98 space Langevin diffusions to model resource selection and movement. From this model it is
 99 possible to calculate an closed-form predictive UD with the RSF exponential form, that is,

$$\mathbf{u}(\mathbf{s}) \propto \exp\{\mathbf{x}(\mathbf{s})'\boldsymbol{\beta}\}, \quad (1)$$

100 where $\mathbf{u}(\mathbf{s})$ is the UD at location $\mathbf{s}$ and $\mathbf{x}(\mathbf{s})$ are habitat covariates at location $\mathbf{s}$. This
 101 model produces the type of analytic UD prediction that directly relates the UD to habitat
 102 variables through the coefficients $\boldsymbol{\beta}$. However. There are some drawbacks for this
 103 approach. First the likelihood for this model is not available in closed form, so a discrete
 104 time (Euler) approximation must be used. Second, this model is only applicable to
 105 continuously valued covariates because the habitat gradient surfaces, $\nabla\mathbf{x}(\mathbf{s})$, are used in
 106 the movement model (Michelot et al., 2019). Finally, the Langevin diffusion limiting
 107 distribution in (1) is only valid when the diffusion component is constant. That is the
 108 animal does not change its overall rate movement in different habitats; only the advection
 109 component is altered by habitat (See Section 3). However, Blackwell (2026) has
 110 demonstrated that this is not necessary for general diffusions.

111 Herein we consider a general class of CTMC models for which we can obtain closed
 112 form predictions of the UD from the fitted model parameters. The CTMC model has
 113 several advantages over the Langevin derived UD. First, there is no approximation
 114 necessary to calculate the likelihood for optimization. Second, continuous or discrete
 115 habitat covariates can be used for movement modeling and UD prediction. Finally, the

diffusion rate can be influenced by habitat not just the advection rate. Moreover, because there is no approximation for the likelihood that error does not propagate to the UD prediction, so it is exact. To fit these model, we also developed the R package `walk`[†] that can fit general CTMC models including the more general forms requiring numerical calculation of the UD as presented by Hanks et al. (2015) and Wilson et al. (2018).

2 Continuous-Time Markov Movement Models

Here we provide a brief description of continuous-time Markov chain (CTMC) models. More detail can be found in Norris (1998), Johnson et al. (2021), or Hanks et al. (2015). In a CTMC movement model the geographical study domain is partitioned into a cells or other discrete locations indexed by $\mathcal{C} = \{1, \dots, n\}$. Each cell, i , has a set of neighboring cells $\mathcal{N}_i = \{j \in \mathcal{C} : i \sim j\}$, where $i \sim j$ means that an animal can go from cell i to cell j in one move. For example, in a raster partition, neighboring cells might share a border (rook neighborhood) or a corner point (queen neighborhood). Here we assume that if $i \sim j$, then $j \sim i$, that is, movement is symmetric between cells. As with all CTMCs, the complete path of the animal, Y , can be decomposed into the *jump times* at which cell transitions are made, $\boldsymbol{\tau} = \{\tau_0, \tau_1, \dots, \tau_M, \tau_{M+1}\}$, and the *embedded Markov chain*, which is the sequence of discrete-space cells visited, $\mathcal{G} = \{g_0, \dots, g_M\}$. Thus, the CTMC path has the bivariate discrete-time representation $Y = \{(\tau_m, g_m) : m = 0, \dots, M\}$, where $\tau_0 = 0$ and $\tau_{M+1} = T$ is the known end of the telemetry deployment. In a slight abuse of notation, we also use g_t and $\mathcal{N}_t$ (i.e., indexed by continuous time, $t \in [0, T]$) to represent the location and neighborhood of the animal for any time, that is, $g_t = g_m$ and $\mathcal{N}_t = \mathcal{N}_{g_m}$ for $t \in [\tau_m, \tau_{m+1})$.

[†]Available from: <https://dsjohnson.r-universe.dev/walk>

It is apparent which context is being used in a particular situation.

CTMC models are defined by the conditional movement rates between cells,

$$\lambda_{ij} = \lim_{h \rightarrow 0} [g_{t+h} = j | g_t = i] / h, \quad (2)$$

where $[g_{t+h} = j | g_t = i]$ is the probability that $g_{t+h} = j$ given the animal is in cell i at time t . We use the notation $'[A|B]'$ to represent the probability density (distribution) function of A given B . The total rate of movement away from cell i is

$$\Lambda_i = \sum_{j \in \mathcal{N}_i} \lambda_{ij}. \quad (3)$$

One other useful property of CTMCs is that the residence time in a cell, $r_m = \tau_{m+1} - \tau_m$, is exponentially distributed,

$$[r_m | g_m = i] = \text{Exponential}(r_m | \Lambda_i).$$

Thus the expected residence time in cell i is Λ_i^{-1} . If we put all the rates in the $n \times n$ matrix $\mathbf{Q}$ with $(-\Lambda_1, \dots, -\Lambda_n)$ on the diagonal and λ_{ij} in the i th row, j th column. Then the transition matrix from $g_0 \in \mathcal{C}$ to $g_t \in \mathcal{C}$ is,

$$\mathbf{\Gamma}(t) = \exp(\mathbf{Q}t),$$

where “exp” is the matrix exponential function. The conditional distribution of $[g_t | g_0 = i] = \gamma_i(t)$ is the i th row of $\mathbf{\Gamma}(t)$.

If a limiting distribution exists, as $t \rightarrow \infty$, $\gamma_i(t) \rightarrow \mathbf{u}$ regardless of what cell the animal

is in currently. The conditions under which a CTMC will have a long-run limiting distribution over the study area are outlined by Wilson et al. (2018). These conditions are

1. *Homogeneity*.– The movement rate function remains constant through time.

2. *Irreducibility*.– Any cell can be reached from any other cell given sufficient time. In practice, that the study area is not separated into multiple areas by impenetrable barriers.

3. *Positive recurrence*.– After an animal leaves a cell the amount of time it takes for the animal to visit that cell again is finite.

Wilson et al. (2018) use numerical Eigen decomposition of $\mathbf{Q}$ to obtain $\mathbf{u}$, but they also present an additional, well known, CTMC result that we use. Assuming the previous conditions are met for a CTMC model, the limiting distribution can be evaluated as

$$u_i \propto \alpha_i / \Lambda_i \tag{4}$$

where $\boldsymbol{\alpha}$ is the limiting distribution of the embedded Markov chain, $\{g_0, g_1, \dots\}$. This makes intuitive sense that the long-run probability of occurrence is the long-run probability the animal is in the cell weighted by the expected amount of time it spends there.

3 Explicit Utilization Distribution CTMC Models

Now that we have a background CTMC we can turn to modeling utilization distributions (UDs) directly. Here we assume the UD has the exponential form for the i th element of $\mathbf{u}$,

$$u_i \propto \exp(v_i),$$

167 where, typically, $v_i = \mathbf{x}_i' \boldsymbol{\beta}$ and $\mathbf{x}_i$ is a vector of habitat variables for the i th cell. However,
 168 we do not necessarily need to restrict ourselves to that linear form now. We use Markov
 169 Chain Monte Carlo (MCMC) theoretical results (Michelot et al., 2018; Blackwell, 2026) to
 170 show that the limiting distribution does, indeed, have the desired exponential form.

171 The CTMC movement model we consider here has the form

$$\log(\lambda_{ij}) = \beta_0 + a_i + b_j + c_{ij}, \quad (5)$$

172 where

- 173 • β_0 is a base rate of movement,
- 174 • a_i only controls residency time when the animal is in cell i
- 175 • b_j controls movement to cell j from any cell in $\mathcal{N}_i$, and
- 176 • c_{ij} controls movement specifically from cell i to cell j .

177 Of course, each of the terms can be functions of spatial covariates. If the neighbor-specific
 178 terms are symmetric, that is, $c_{ij} = c_{ji}$, then we use the embedded chain result (4) and
 179 MCMC theory to show (Appendix A) the limiting distribution is,

$$u_i \propto \exp(-a_i + b_i),$$

180 which has the desired form. If $a_i = \mathbf{x}_{ai}' \boldsymbol{\beta}_a$ and $b_i = \mathbf{x}_{bi}' \boldsymbol{\beta}_b$, then u_i has the form

$$u_i \propto \exp(\mathbf{x}_i' \boldsymbol{\beta}),$$

181 where $\mathbf{x}'_i = (\mathbf{x}'_{ai}, \mathbf{x}'_{bi})$ and $\boldsymbol{\beta} = (-\boldsymbol{\beta}_a, \boldsymbol{\beta}_b)$. This is the discrete-space equivalent of the
182 continuous-space predictive UD given by Barnett and Moorcroft (2008). However, there is
183 another specific condition for c_{ij} that also produces a closed exponential form UD,
184 $c_{ij} = d_j - d_i$. That is if the cell pair effects are local differences of cell specific conditions
185 then the UD has the form,

$$u_i \propto \exp(-a_i + b_i + 2d_i),$$

186 (Appendix A). If $d_i = \mathbf{x}'_{di}\boldsymbol{\beta}_d$ we get the same log-linear exponential form where
187 $\mathbf{x}'_i = (\mathbf{x}'_{ai}, \mathbf{x}'_{bi}, \mathbf{x}'_{di})$ and $\boldsymbol{\beta} = (-\boldsymbol{\beta}_a, \boldsymbol{\beta}_b, 2\boldsymbol{\beta}_d)$.

188 4 CTMC parameter estimation

189 Previous estimation of CTMC movement parameters using satellite location telemetry data
190 used an imputation method (see Scharf et al. 2017) which fills in missing cell locations
191 between location acquisition times (Hanks et al., 2015; Wilson et al., 2018; Johnson et al.,
192 2021). Even though the imputation method attempts to propagate the uncertainty of the
193 missing locations, it is unclear how this affects estimation of the parameters. Therefore we
194 will use the exact, although, more computationally expensive hidden Markov model
195 (HMM) formulation of the model that is often use for CTMC analysis marine fish
196 movement from archival telemetry tags (e.g., see CH. 10 of Thorson and Kristensen 2024).

4.1 Hidden Markov model specification

4.1.1 Satellite telemetry data specification and processing

For the proposed analysis we consider satellite telemetry data for an individual of the form $\mathbf{y} = \{\mathbf{y}_1, \dots, \mathbf{y}_K\}$, where $\mathbf{y}_k = (\mathbf{s}_k, t_k, \boldsymbol{\epsilon}_k)$, $\mathbf{s}_k$ is the observed location of the individual within the study area at time t_k and $\boldsymbol{\epsilon}_k$ represents information or parameters governing location uncertainty. For example, $\boldsymbol{\epsilon}_k$ might represent the major and minor error ellipse lengths of a Kalman Filtered Argos location (McClintock et al., 2015) or a known error radius for GPS locations, e.g., 20m. From these locations we translate the continuous-space data locations and uncertainty to the discrete grid, $\mathcal{C}$, using a function, say, $p(\mathbf{y}_k, \mathcal{C}) = \mathbf{p}_k = (p_{1k}, \dots, p_{nk})$, where p_{ik} is the probability that the animal is located within cell i at time t_k . We used a general form because the function, p , will depend on how location uncertain is assessed for the specific satellite and tag hardware. However, uncertainty in the true location, $\boldsymbol{\mu}_k$, is often represented by the bivariate Normal distribution $\boldsymbol{\mu}_k \sim N(\mathbf{s}_k, \boldsymbol{\Sigma}_k)$, where $\boldsymbol{\epsilon}_k = \boldsymbol{\Sigma}_k$. Therefore p_{ik} would represent the mass of the Normal distribution contained within the i th cell. For a raster grid partitioning of the study area this can be calculated by,

$$p_{ik} = \text{CDF}(x_{i2}, y_{i2}) - \text{CDF}(x_{i2}, y_{i1}) - \text{CDF}(x_{i1}, y_{i2}) + \text{CDF}(x_{i1}, y_{i1})$$

where $[x_{i1}, x_{i2}] \times [y_{i1}, y_{i2}]$ is the bounding box for the i th cell and CDF is the bivariate normal cumulative density function with mean $\mathbf{s}_k$ and covariance $\boldsymbol{\Sigma}_k$. A Monte Carlo option for other error distributions would be to draw a sufficiently large sample from the distribution and assess the proportion of samples that falls within each cell.

4.1.2 Likelihood calculation

For an individual the likelihood is calculated with the matrix equation (Zucchini et al. 2016; Ch. 3),

$$L_{\text{CTMC}}(\mathbf{y}) = \boldsymbol{\pi}' \mathbf{P}(\mathbf{y}_1) \boldsymbol{\Gamma}(t_2 - t_1) \mathbf{P}(\mathbf{y}_2) \cdots \boldsymbol{\Gamma}(t_K - t_{K-1}) \mathbf{P}(\mathbf{y}_K) \mathbf{1} \quad (6)$$

where $\boldsymbol{\pi} = (\pi_1, \dots, \pi_n)$ is the probability at the individual is cell i at t_1 , $\mathbf{P}(\mathbf{y}_k) = \text{diag}(\mathbf{p}_k)$, $\boldsymbol{\Gamma}(t_k - t_{k-1}) = \exp\{\mathbf{Q}(t_k - t_{k-1})\}$, and $\mathbf{1}$ is a vector of 1s. The natural choice for $\boldsymbol{\pi}$ is $\mathbf{u}$, the limiting distribution defined by the CTMC movement model, but there are other choices. For example, if a telemetered individual was released at a known location after tagging, then $\boldsymbol{\pi}$ would be a vector of all 0s except for the cell containing the release site, which would be a 1. In the false killer whale example below, we use $\boldsymbol{\pi} = \mathbf{u}$.

Calculating the likelihood directly as it is presented in (6) can lead to numerical underflow and is much more computationally expensive than necessary. In order to avoid numerical underflow we evaluate the likelihood on the log scale using the alternate version presented in Section 3.2 of Zucchini et al. (2016). The computationally expensive aspect is due to the repeated evaluation of $\boldsymbol{\Gamma}(t_k - t_{k-1}) = \exp\{\mathbf{Q}(t_k - t_{k-1})\}$, which is the matrix exponential of a potentially very large dimensional $\mathbf{Q}$ matrix. However, the full evaluation of $\boldsymbol{\Gamma}(t_k - t_{k-1})$ is not necessary, we only need to evaluate quantities of the form $\mathbf{v}' \exp\{\mathbf{Q}(t_k - t_{k-1})\}$. Sherlock (2021) presents very computationally efficient ways to evaluate this product directly. Moreover, $\mathbf{Q}$ is a sparse matrix so we can also take advantage of that fact in the evaluations as well. The `walk` package implements the methods of Sherlock (2021) using sparse matrix operations for evaluating and optimizing the likelihood.

In addition to the matrix exponential evaluation numerical savings, another optional approach to save computational effort is to set a truncation point for the maximum Λ_i value. The numerical methods of Sherlock (2021) require more matrix operations for higher values of $\max(\Lambda_i)$. For some cells, the log linear λ_{ij} model may produce rates of movement that are unrealistically high for the species in question. With respect to UD's this would imply that an individual will not remain in that cell and the $u_i \approx 0$. Hewitt et al. (2023) proposed a logit rate model that restricts this behavior, but that model does not allow us to construct a CTMC with an explicit limiting distribution. So, we propose picking a truncation value, say, Λ^* and setting $\lambda_{ij}^* = (\Lambda^*/\Lambda_i)\lambda_{ij}$ if $\Lambda_i > \Lambda^*$. If Λ^* is sufficiently large, this will have negligible impact on β and $\mathbf{u}$ estimation but can dramatically help numerical likelihood evaluation. One reasonable way to choose Λ^* is to select the minimum amount of time that an animal would likely to reside in a cell, say ξ . Then set Λ^* so that the probability of a residence time $< \xi$ is, say, 0.9. That is, with a total cell movement rate of Λ^* , there is a 90% chance residency will be $< \xi$. This represents a reasonable upper bound for movement. The effect on β and $\mathbf{u}$ can always be verified by comparing the numerically calculated $\mathbf{u}$ as presented in Wilson et al. (2018) versus $u_i \propto \exp(\mathbf{x}_i' \beta)$. If there are sufficiently large discrepancies, Λ^* can be increased. See Section (5.3) for example.

Once the CTMC movement model has been fitted, recall that the parameters need be adjusted for UD inference. That is the maximum likelihood estimate of $\hat{\beta}_{\text{CTMC}} = (\hat{\beta}_a, \hat{\beta}_b, \hat{\beta}_d)$ needs to be transformed to the UD coefficients via $\hat{\beta} = \mathbf{M}\hat{\beta}_{\text{CTMC}}$, where $\mathbf{M} = \text{diag}((-1, -\mathbf{1}'_a, \mathbf{1}'_b, 2\mathbf{1}'_d)')$. Thus, the large sample covariance matrix of the UD coefficients is $\mathbf{V}(\hat{\beta}) = \mathbf{M}\mathbf{V}(\hat{\beta}_{\text{CTMC}})\mathbf{M}'$. If only the standard errors are desired, one can simply multiply the CTMC coefficient standard errors by the multipliers, but if the entire covariance matrix is desired the matrix multiplication is necessary.

5 False killer whale space-use

In this section, we will apply the modeling approach described herein to a telemetry dataset on endangered false killer whales (FKW; *Pseudorca crassidens*) in the Hawaiian Islands (Baird 2016). This particular population is resident to the insular waters of the main Hawaiian Islands (MHI; includes Kaua‘i/Ni‘ihau, Oahu, Maui Nui, and Hawai‘i Island) and is comprised of five social clusters, or groups of close family members and regular associates (Baird et al., 2008, 2019; Baird, 2016; Martien et al., 2019). These whales are known to move quickly and frequently among island areas throughout their range (Baird et al., 2010, 2012, 2021; Baird, 2016). Due to inherent challenges associated with tracking apex marine predators, this dataset is characterized by variable deployment durations and irregular location data streams. Considering these factors may drive start bias, as well as their expansive range, we deem this an appropriate dataset to empirically demonstrate the utility of predictive UD. For the purposes of this assessment, we only focus on movements of 5 individuals that were satellite tagged in 2021 and 2022. There is one representative from each social cluster in the data.

5.1 Satellite tag data

False killer whale satellite tag deployment methods have been described in detail elsewhere (see Baird et al. 2010, 2012) so they are only briefly summarized here. Individuals were tagged with Argos transmitting SPOT6 ($n = 4$) and SPLASH10 ($n = 1$) tags during dedicated small boat survey efforts throughout the main Hawaiian Islands (Baird et al. 2012, 2021; Baird 2016; see Baird et al. 2013 for details on surveys). Deployment duration ranged from 12 days to 81 days (median=40 days). Argos location data were processed

through the Douglas-Argos Filter (accessed via Movebank; Kranstauber et al. 2011; Douglas et al. 2012) to remove erroneous locations based on unrealistic travel speeds and turning angles. The final analytical dataset consisted of 5 trajectories and 4,009 filtered Argos and Fastloc GPS (SPASH10 tag) locations (Figure 1A). For the Argos locations, we used the Kalman-filter diagnostic ellipse data to construct error covariance matrices (McClintock et al., 2015). While for the Fastloc GPS locations, we used the 95% quantile error ranges from Dujon et al. (2014), based on the number of satellites acquired, as twice the standard deviation of the normal error.

5.2 Study area and habitat variables

The official management range of the MHI stock of FKW was documented in Bradford et al. (2015) and represents a minimum convex polygon surrounding a buffer of the main Hawaiian Islands (see Figure 1A). This is the study area boundary we use for the UD analysis.

For habitat characteristics we used two modeled global oceanographic variables from E.U. Copernicus Marine Service: (1) Sea Surface Temperature (SST; °C; DOI: <https://doi.org/10.48670/moi-00016>) and (2) Sea Level Anomaly (SLA; m; DOI: <https://doi.org/10.48670/moi-00148>). SST is available at 0.082° resolution and SLA is available at 0.125° degree resolution. In order to match resolution we chose to project the SLA raster to the resolution of the SST raster to avoid losing oceanographic information. Because we are using these variables to predict long term space-use, we used a single layer for each variable that was the 2 year daily average in the study area. This allowed us to model movement with respect to typical oceanographic conditions (Figure 1B-C).

In addition to the oceanographic variables, we used distance from land (m) to model

FKW’s resident nature in the MHI. Also, we also included a depth residual variables by adding data from the global, 30 arc-second ETOPO elevation model (NOAA National Centers for Environmental Information, 2022) and removing the linear trend of distance from land. The distance and depth variables were projected to the 0.082° SST resolution as well. Then all data, including telemetry data, were projected to the Hawaii Albers Equal Area projection, ESRI:102007 (Figure 1D-E)

5.3 CTMC movement model and population inference for space-use

The CTMC movement rate model we used to model the telemetry for each individual $l = 1, \dots, 5$ was

$$\log \lambda_{ij} = \beta_0 + \underbrace{\text{DEPTH}_i + \text{SST}_i + \text{SST}_i^2 + \text{SLA}_i + \text{SLA}_i^2}_{a_i \text{ portion}} + \underbrace{[\text{DIST}_j - \text{DIST}_i]}_{c_{ij} \text{ portion}}$$

In this model differences of the distance from land variable were used as a movement kernel between cells i and j . This allowed individuals to prefer movement towards (or away from) land at the same rate no matter how far away they are. In addition SST, SLA, and DIST were Z -transformed for numerical stability. The DEPTH residual was already on a nearly standard normal scale and DIST was only used as a difference, so values were not large enough to cause issue.

Estimated coefficients and predicted UD’s can be difficult to interpret for population inference and management considerations. Therefore, a Bayesian hierarchical or random

325 effects model of the form

$$[\mathbf{y}_1, \dots, \mathbf{y}_N | \boldsymbol{\beta}_1, \dots, \boldsymbol{\beta}_N] = \prod_{l=1}^N L_{\text{CTMC}}(\mathbf{y}_l | \mathbf{M}^{-1} \boldsymbol{\beta}_l)$$

$$[\boldsymbol{\beta}_1, \dots, \boldsymbol{\beta}_N | \boldsymbol{\theta}, \boldsymbol{\Sigma}] = \prod_{l=1}^N N(\boldsymbol{\beta}_l | \boldsymbol{\theta}, \boldsymbol{\Sigma})$$

326 where $\mathbf{y}_l$ is the observed data for individual $i = 1, \dots, N$ and the prior distribution for the
 327 random effect mean and variance is $[\boldsymbol{\theta}] [\boldsymbol{\Sigma}]$. Note, we have phased this model in terms of
 328 the UD coefficients, $\boldsymbol{\beta}_l$, rather than the movement coefficients, $\boldsymbol{\beta}_{\text{CTMC},l}$ because we are
 329 more interested in UD prediction here. But it is just a linear transformation between the
 330 two, so, all the results are equivalent. Even if individual inference is desired, it is useful to
 331 fit this hierarchical model because it allows us to share information between individuals.
 332 That is we can make inference to the posterior distribution $[\boldsymbol{\beta}_l | \mathbf{y}_1, \dots, \mathbf{y}_N]$. If we assume
 333 that there is some similarity in movement between individuals of the same species then we
 334 can leverage all data for better individual inference. Unfortunately, even with the
 335 computational savings, CTMC models are complex enough as to render it infeasible to fit
 336 this model jointly.

337 While we cannot fit the hierarchical model in one step, there have been several
 338 proposed 2 step divide-and-conquer methods to approximate full posterior inference of
 339 hierarchical models of this form (e.g., Lunn et al., 2013; Hooten et al., 2016; Johnson
 340 et al., 2022). Here we use the method of Johnson et al. (2022) as it allows use to use
 341 maximum likelihood estimation in the first step. So, in the first step the CTMC model was
 342 fitted separately to the data from each individual using normal location errors as described
 343 in Section 4 with maximum likelihood estimation (MLE). For the rate truncation we set
 344 $\Lambda^* = 138.16$. This resulted from selecting the minimum residence time to be 1 minute. The

second step involves fitting the multivariate linear mixed model,

$$\hat{\boldsymbol{\beta}}_l = \boldsymbol{\theta} + \mathbf{u}_l + \boldsymbol{\delta}_l; \quad l = 1, \dots, N$$

where $\hat{\boldsymbol{\beta}}_l$ is the MLE estimate of the coefficients for the l th individual, the random effect is $[\mathbf{u}_l | \boldsymbol{\Sigma}] = N(\mathbf{0}, \boldsymbol{\Sigma})$, the error term is $[\boldsymbol{\delta}_l | \mathbf{V}(\hat{\boldsymbol{\beta}}_l)] = N(\mathbf{0}, \mathbf{V}(\hat{\boldsymbol{\beta}}_l))$, and $\mathbf{V}(\hat{\boldsymbol{\beta}}_l)$ is the parameter covariance matrix from the first MLE step. For this analysis we used independent random effects, that is, $\boldsymbol{\Sigma} = \text{diag}(\sigma_1^2, \dots, \sigma_7^2)$. The `nimble` R package (de Valpine et al., 2017) was used to optimize model with respect to the parameters (i.e., maximum *a posteriori*, MAP) subject to the prior distributions,

$$[\theta_w] = N(0, \tau_w^2); \quad [\tau_w] = \text{Exponential}(1)$$

$$[\sigma_w] = \text{Exponential}(1)$$

for $w = 1, \dots, 7$. The exponential priors for the variance parameters were added to penalize the complexity of the model (i.e., regularization) to perform selection for the population level parameters (Hooten and Hobbs, 2015; Simpson et al., 2017). Following second stage model fitting we can extract the approximate MAP estimates $\bar{\boldsymbol{\theta}}$ and $\bar{\boldsymbol{\beta}}_1, \dots, \bar{\boldsymbol{\beta}}_N$, where $\bar{\boldsymbol{\beta}}_l$ is the approximate MAP of the posterior $[\boldsymbol{\beta}_l | \mathbf{y}_1, \dots, \mathbf{y}_N]$.

5.4 Results

Individual results for all individuals were generally consistent with notable deviations being individual PcTag074 did not have any effect of SLA on its UD and PcTag070 had negative effect of the linear SST term (Figure 2). All other coefficients were similar in direction, i.e., less than or greater than 0. This bolsters the idea of using the 2 stage meta-analytic

approach to share data between individuals to improve estimates. The second stage individual coefficient estimates, $\bar{\beta}_i$ illustrate that there is only a significant amount of individual variation in the DEPTH and SLA² coefficients (Figure 2). All other second stage coefficients are shrunk to the population mean estimates, $\bar{\theta}$ with nearly identical approximate credible intervals (Figure 2). There was only 1 nonsignificant population coefficient, the linear SLA term. Correspondingly, the penalized model complexity priors shrunk the SLA term to zero. All other population coefficients were different than zero. The DEPTH and DISTANCE coefficients were negative as expected because this population inhabits shallower waters close to the MHI without venturing away. The zero SLA coefficient combined with the negative SLA² term implies a maximum effect at 0, but we used A Z-transform, therefore, the maximum effect occurs at the mean original SLA value. Both SST coefficients are positive implying that the typical individual is probably located in the warmest SST conditions in the study area.

Unsurprisingly, the similarity in coefficient estimates produced similar predicted UD_s at the first stage (Figure 3). Individual PcTag074 had a more notably diffuse UD, the other four individuals had very similar looking UD_s. This may be due to the lack of SLA effect on the UD of PcTag074. After the second stage, the UD_s are even more similar as the second stage coefficients are essentially the same at the population mean coefficients (Figure 4). The UD created from the population mean coefficients is illustrated in Figure 4 (Typical). This is the UD of a so called “Typical” or median individual, not the population average UD because of the log link between the coefficients and the UD. Even after the second stage the coefficients for PcTag074 lack any SLA effect, so the UD is the only substantially different one from the typical UD.

In order to check that the rate truncation did not adversely affect the estimated UD_s we

compared the truncated numerical Eigen decomposition UD ($u_{ei,i}$; Wilson et al. 2018) with the exponential closed form version using Bhattacharyya’s Affinity (BA) (Bhattacharyya, 1943),

$$BA = \sum_{i \in \mathcal{C}} \sqrt{u_{ei,i} \cdot u_i}.$$

The values of BA range from 0 (no similarity) to 1 (identical distributions) Fieberg and Kochanny (2005). The proportion of study area cells truncated in the fitting of individual models ranged from 0–44% of the study area. However, BA ranged from 0.996–1 indicated that the truncated cells possessed very little UD mass and the closed form and exact truncated UDs were virtually identical. Therefore, we can be assured that the truncation approximation did not adversely affect parameter estimation.

6 Discussion

Acknowledgments

The findings and conclusions in the paper are those of the authors and do not necessarily represent the views of the National Marine Fisheries Service, NOAA. Reference to trade names does not imply endorsement by the National Marine Fisheries Service, NOAA.

References

- Avgar, T., Potts, J. R., Lewis, M. A., and Boyce, M. S. (2016). Integrated step selection analysis: Bridging the gap between resource selection and animal movement. *Methods in Ecology and Evolution*, 7(5):619–630.

- Baird, R. W. (2016). *The Lives of Hawai'i's Dolphins and Whales: Natural History and Conservation*. University of Hawaii press.
- Baird, R. W., Anderson, D. B., Kratofil, M. A., and Webster, D. L. (2021). Bringing the right fishermen to the table: Indices of overlap between endangered false killer whales and nearshore fisheries in Hawai'i. *Biological Conservation*, 255:108975.
- Baird, R. W., Anderson, D. B., Kratofil, M. A., Webster, D. L., and Mahaffy, S. D. (2019). Cooperative conservation and long-term management of false killer whales in Hawai'i: Geospatial analyses of fisheries and satellite tag data to understand fishery interactions. A Report to the State of Hawai'i Board of Land and Natural Resources, Contract No. 67703, Cascadia Research Collective.
- Baird, R. W., Gorgone, A. M., McSweeney, D. J., Webster, D. L., Salden, D. R., Deakos, M. H., Ligon, A. D., Schorr, G. S., Barlow, J., and Mahaffy, S. D. (2008). False killer whales (*Pseudorca crassidens*) around the main Hawaiian Islands: Long-term site fidelity, inter-island movements, and association patterns. *Marine Mammal Science*, 24(3):591–612.
- Baird, R. W., Hanson, M. B., Schorr, G. S., Webster, D. L., McSweeney, D. J., Gorgone, A. M., Mahaffy, S. D., Holzer, D. M., Oleson, E. M., and Andrews, R. D. (2012). Range and primary habitats of Hawaiian insular false killer whales: Informing determination of critical habitat. *Endangered Species Research*, 18(1):47–61.
- Baird, R. W., Schorr, G. S., Webster, D. L., McSweeney, D. J., Hanson, M. B., and Andrews, R. D. (2010). Movements and habitat use of satellite-tagged false killer whales around the main Hawaiian Islands. *Endangered Species Research*, 10:107–121.

- Baird, R. W., Webster, D. L., Aschettino, J. M., Schorr, G. S., and McSweeney, D. J. (2013). Odontocete cetaceans around the main Hawaiian Islands: Habitat use and relative abundance from small-boat sighting surveys. *Aquatic Mammals*, 39(3).
- Barnett, A. H. and Moorcroft, P. R. (2008). Analytic steady-state space use patterns and rapid computations in mechanistic home range analysis. *Journal of Mathematical Biology*, 57(1):139–159.
- Bhattacharyya, A. (1943). On a measure of divergence between two statistical populations defined by their probability distribution. *Bulletin of the Calcutta Mathematical Society*, 35:99–110.
- Blackwell, P. G. (2026). Consistent multiscale modelling of movement and habitat selection. *Journal of Mathematical Biology*, 92(6):90.
- Bradford, A. L., Oleson, E. M. E. M., Baird, R. W., Boggs, C. H., Forney, K. A., and Young, N. C. (2015). Revised stock boundaries for false killer whales (*Pseudorca crassidens*) in Hawaiian waters. <https://repository.library.noaa.gov>.
- Calabrese, J. M., Fleming, C. H., and Gurarie, E. (2016). Ctm: An R package for analyzing animal relocation data as a continuous-time stochastic process. *Methods in Ecology and Evolution*, 7(9):1124–1132.
- Chib, S. and Greenberg, E. (1995). Understanding the metropolis-hastings algorithm. *The american statistician*, 49(4):327–335.
- Citation (2015). Unknown. *Unknown*.
- de Valpine, P., Turek, D., Paciorek, C. J., Anderson-Bergman, C., Lang, D. T., and Bodik,

R. (2017). Programming with models: Writing statistical algorithms for general model structures with NIMBLE. *Journal of Computational and Graphical Statistics*, 26(2):403–413.

Douglas, D. C., Weinzierl, R., C. Davidson, S., Kays, R., Wikelski, M., and Bohrer, G. (2012). Moderating A rgos location errors in animal tracking data. *Methods in Ecology and Evolution*, 3(6):999–1007.

Dujon, A. M., Lindstrom, R. T., and Hays, G. C. (2014). The accuracy of Fastloc-GPS locations and implications for animal tracking. *Methods in Ecology and Evolution*, doi: 10.1111/2041-210X.12286.

Fieberg, J. and Kochanny, C. O. (2005). Quantifying Home-Range Overlap: The Importance of the Utilization Distribution. *The Journal of Wildlife Management*, 69(4):1346–1359.

Fieberg, J., Matthiopoulos, J., Hebblewhite, M., Boyce, M. S., and Frair, J. L. (2010). Correlation and studies of habitat selection: Problem, red herring or opportunity? *Philosophical Transactions of the Royal Society B-Biological Sciences*, 365(1550):2233–2244.

Fleming, CH., Fagan, WF., Mueller, T., Olson, KA., Leimgruber, P., and Calabrese, JM. (2015). Rigorous home range estimation with movement data: A new autocorrelated kernel density estimator. *Ecology*, 96(5):1182–1188.

Fortin, D., Beyer, H. L., Boyce, M. S., Smith, D. W., Duchesne, T., and Mao, J. S. (2005). Wolves Influence Elk Movements: Behavior Shapes a Trophic Cascade in Yellowstone National Park. *Ecology*, 86(5):1320–1330.

- Hanks, E. M., Hooten, M. B., and Alldredge, M. W. (2015). Continuous-time discrete-space models for animal movement. *The Annals of Applied Statistics*, 9:145–165.
- Hewitt, J., Gelfand, A. E., and Schick, R. S. (2023). Time-discretization approximation enriches continuous-time discrete-space models for animal movement. *The Annals of Applied Statistics*, 17(1):740–760.
- Hooten, M. B., Buderman, F. E., Brost, B. M., Hanks, E. M., and Ivan, J. S. (2016). Hierarchical animal movement models for population-level inference. *Environmetrics (London, Ont.)*, 27:322–333.
- Hooten, M. B. and Hobbs, N. T. (2015). A guide to Bayesian model selection for ecologists. *Ecological Monographs*, 85(1):3–28.
- Horne, J. S., Garton, E. O., Krone, S. M., and Lewis, J. S. (2007). Analyzing animal movements using Brownian bridges. *Ecology*, 88:2354–2363.
- Johnson, D. S., Brost, B. M., and Hooten, M. B. (2022). Greater than the sum of its parts: Computationally flexible Bayesian hierarchical modeling. *Journal of Agricultural, Biological and Environmental Statistics*, 27(2):382–400.
- Johnson, D. S., Hooten, M. B., and Kuhn, C. E. (2013). Estimating animal resource selection from telemetry data using point process models. *Journal of Animal Ecology*, 82:1155–1164.
- Johnson, D. S., Pelland, N. A., and Sterling, J. T. (2021). A continuous-time semi-Markov model for animal movement in a dynamic environment. *Annals of Applied Statistics*, 15(2):797–812.

490 Johnson, D. S., Thomas, D. L., Hoef, J. M. V., and Christ, A. (2008). A general framework
 491 for the analysis of animal resource selection from telemetry data. *Biometrics. Journal of*
 492 *the International Biometric Society*, 64(3):968–976.

493 Keating, K. A. and Cherry, S. (2009). Modeling utilization distributions in space and time.
 494 *Ecology*, 90:1971–1980.

495 Kranstauber, B., Cameron, A., Weinzerl, R., Fountain, T., Tilak, S., Wikelski, M., and
 496 Kays, R. (2011). The Movebank data model for animal tracking. *Environmental*
 497 *Modelling & Software*, 26(6):834–835.

498 Lunn, D., Barrett, J., Sweeting, M., and Thompson, S. (2013). Fully Bayesian hierarchical
 499 modelling in two stages, with application to meta-analysis. *Journal of the Royal*
 500 *Statistical Society: Series C (Applied Statistics)*, 62(4):551–572.

501 Manly, B., McDonald, L., Thomas, D., McDonald, T., and Erikson, W. (2002). *Resource*
 502 *Selection by Animals*. Kluwer Academic Publishers, 2nd edition, pp. 221 edition.

503 Martien, K. K., Taylor, B. L., Chivers, S. J., Mahaffy, S. D., Gorgone, A. M., and Baird,
 504 R. W. (2019). Fidelity to natal social groups and mating within and between social
 505 groups in an endangered false killer whale (*Pseudorca crassidens*) population.
 506 *Endangered Species Research*, 40:219–230.

507 McClintock, B. T., London, J. M., Cameron, M. F., and Boveng, P. L. (2015). Modelling
 508 animal movement using the Argos satellite telemetry location error ellipse. *Methods in*
 509 *Ecology and Evolution*, In press:doi: 10.1111/2041-210X.12311.

510 Michelot, T., Blackwell, P. G., and Matthiopoulos, J. (2018). Linking resource selection
 511 and step selection models for habitat preferences in animals. *Ecology*.

512 Michelot, T., Etienne, M.-P., and Gloaguen, P. (2019). The Langevin diffusion as a
513 continuous-time model of animal movement and habitat selection. *Methods In Ecology*
514 *and Evolution*, In press.

515 Moorcroft, P. R., Lewis, M. A., and Crabtree, R. L. (1999). Home range analysis using a
516 mechanistic home range model. *Ecology*, 80(5):1656–1665.

517 NOAA National Centers for Environmental Information (2022). ETOPO 2022 15
518 arc-second global relief model.

519 Norris, J. R. (1998). *Markov Chains*. Cambridge University Press, 2nd edition edition.

520 Powell, R. A. and Mitchell, M. S. (2012). What is a home range? *Journal of mammalogy*,
521 93(4):948–958.

522 Scharf, H., Hooten, M. B., and Johnson, D. S. (2017). Imputation Approaches for Animal
523 Movement Modeling. *Journal of Agricultural, Biological, and Environmental Statistics*,
524 22(3):335–352.

525 Sherlock, C. (2021). Direct statistical inference for finite Markov jump processes via the
526 matrix exponential. *Computational Statistics*, 36(4):2863–2887.

527 Simpson, D. P., Rue, H., Riebler, A., Martins, T. G., and Sørbye, SH. (2017). Penalising
528 model component complexity: A principled, practical approach to constructing priors.
529 *Statistical Science*, 32:1–28.

530 Thorson, J. and Kristensen, K. (2024). *Spatio-Temporal Models for Ecologists*. Chapman
531 and Hall/CRC.

- 532 Whitehead, H. and Jonsen, I. D. (2013). Inferring animal densities from tracking data
533 using markov chains. *PloS one*, 8(4):e60901.
- 534 Wilson, K., Hanks, E., and Johnson, D. (2018). Estimating animal utilization densities
535 using continuous-time Markov chain models. *Methods in Ecology and Evolution*,
536 9(5):1232–1240.
- 537 Zucchini, W., MacDonald, I. L., and Langrock, R. (2016). *Hidden Markov Models for Time*
538 *Series: An Introduction Using R*. CRC Press, 2 edition.




Figure 1: Habitat covariates




Figure 2: Parameter estimates for fitted UD surfaces. The Stage 1 values are from individually fitted models. The Stage 2 estimates are parameters estimates using the two stage method of Johnson et al. (2022) to approximate a combined random effects model. The solid vertical lines are the estimated random effect means and the dashed lines are the approximate 95% credible intervals.




Figure 3: Individual model predicted utilization distributions. These UD were predicted using only the data from each respective individual. A UD prediction for PcTag071 is absent because the resulting prediction was nonsensical given the individual movement parameter estimates. That is, large (but highly uncertain) SLA and SLA^2 coefficients resulted in only a few cells getting all the UD mass. This illustrates one of the problem when using movement models to predict space use.




Figure 4: Predicted UD after 2 stage approach.

A Proof of closed form CTMC limiting distribution

Using the Metropolis-Hastings (MH) algorithm we can show, by construction, that the limiting distribution of the CTMC we are considering has the limiting distribution stated in Section 3.1. Recall that the limiting distribution of a CTMC is $\mathbf{u}$ with i th element $u_i = \alpha_i/\Lambda_i$ where α is the limiting distribution of the embedded chain, $\mathcal{G}$. So, we can use the theoretical results of the MH algorithm to show that the discrete-time chain, $\mathcal{G}$, has the appropriate limiting distribution.

In general, the MH algorithm is designed to sample from a distribution $c^{-1}\pi(x)$ when the proportionality constant is unknown. It accomplishes this feat by constructing a Markov process $q(X_t|X_{t-1})$ to move from state X_{t-1} to X_t such that, asymptotically, $X_t \sim c^{-1}\pi(x)$ as $t \rightarrow \infty$. Full details of the MH algorithm can be found in Chib and Greenberg (1995), but the basic version proceeds as follows for desired limiting distribution, $c^{-1}\pi(x)$,

-
1. For current state $X_{t-1} = x$ propose $X_t = x'$ from probability distribution $q(x'|x)$.
 2. With probability

$$R_t = \min \left\{ \frac{\pi(x')q(x|x')}{\pi(x)q(x'|x)}, 1 \right\}$$

set $X_t = x'$ else, set $X_t = x$.

3. Repeat.
-

The acceptance step of the MH algorithm is the key feature that ensures, under certain general regularity conditions, that the distribution of $X_t \sim c^{-1}\pi(x)$ as $t \rightarrow \infty$. The Markov process $q(\cdot|\cdot)$ need not necessarily have a limiting distribution, but the acceptance step will produce a chain with limiting distribution $c^{-1}\pi(x)$. If $R_t \geq 1$ for all t , then no matter what

state is proposed by $q(\cdot|\cdot)$, it is accepted, thus, it is the same as simply drawing a realization of the $q(\cdot|\cdot)$ process. In that case, the limiting distribution of $q(\cdot|\cdot)$ is $c^{-1}\pi(x)$ by construction.

We now return to the CTMC movement model in Section 3.1 to prove the resulting limiting distribution is as stated. Recall the proposed CTMC has movement rate function of the form,

$$\lambda_{ij}(t) = \exp\{\beta_0 + a_i + b_j + c_{ij}\}.$$

The embedded Markov chain, $\mathcal{G}$, for the CTMC movement process is defined by the transition probabilities (Citation, 2015),

$$q(g_m = j | g_{m-1} = i) = \lambda_{ij} / \Lambda_i \text{ for } i \sim j,$$

where $\Lambda_i = \sum_{j \in \mathcal{N}_i} \lambda_{ij}$. All we need to do is construct an appropriate α_i , $i = 1, \dots, n$, which is only a function of cell i and no other cell j , such that

$$R_m = \frac{\alpha_j \lambda_{ji} \Lambda_i}{\alpha_i \lambda_{ij} \Lambda_j} \geq 1$$

for all $m > 0$ and all (i, j) such that i and j are neighbors. Then, α will be the limiting distribution of the embedded chain and we can use the result that the limiting distribution of the full CTMC is $u_i \propto \alpha_i / \Lambda_i$.

We begin by evaluating the MH acceptance ratio for our CTMC model. After

canceling like terms in the fraction and rearranging advantageously, we obtain,

$$\begin{aligned}
R_m &= \frac{\exp\{a_j + b_i\}\Lambda_i}{\alpha_i} \times \frac{\alpha_j}{\exp\{a_i + b_j\}\Lambda_j} \times \frac{\exp\{c_{ji}\}}{\exp\{c_{ij}\}} \\
&= \frac{\exp\{b_i - a_i\}\Lambda_i}{\alpha_i} \times \frac{\alpha_j}{\exp\{b_j - a_j\}\Lambda_j} \times \exp\{c_{ji} - c_{ij}\}
\end{aligned} \tag{7}$$

From this MH acceptance ratio, one can see that a set of sufficient conditions that satisfy

$R_m \geq 1$ for all $m = 1, 2, \dots$ and all (i, j) such that $i \sim j$ are:

1. If $c_{ij} = c_{ji}$, then $\alpha_i \propto \exp\{b_i - a_i\}\Lambda_i$, else if,
2. If $c_{ij} = d_j - d_i$ (i.e., a difference of cell indexed effects), then $c_{ji} - c_{ij} = 2d_i - 2d_j$ and $\alpha_i \propto \exp\{b_i - a_i + 2d_i\}\Lambda_i$ satisfies the condition for the limiting distribution of the embedded chain.