

Journal: Ecology

Manuscript type: Statistical Innovation

Predicting Animal Space-Use with Continuous-Time Markov Chain Movement Models

Devin S. Johnson^{1,*}, Michaela A. Kratochvil^{2,3}, Janelle J. Gardner¹, Robin W. Baird³, and Amanda L. Bradford¹

¹Pacific Islands Fisheries Science Center, National Marine Fisheries Service, NOAA, Honolulu, Hawaii, USA

²Marine Mammal Institute, Department of Fisheries, Wildlife, and Conservation Sciences, Oregon State University, Newport, Oregon, USA

³Cascadia Research Collective, Olympia, Washington, USA

*Corresponding author: devin.johnson@noaa.gov

July 27, 2026

Abstract. Remote animal telemetry data is widely used in ecology to assess animal space-use and evaluate utilization distributions (UDs). However, traditional empirical UD (like kernel density or Brownian bridge estimators) can suffer from “start bias” if tag deployments are too short for an animal to fully explore its habitat, heavily biasing results toward the initial release location. Recently, predictive UD have been proposed in an attempt to mitigate this by finding the long-run stationary distribution of a stochastic process that uses habitat covariates to describe movement. This stationary distribution is free from start bias because it uses habitat variables to predict space-use in the future based on short term movements. Predictive UD have been calculated from Langevin diffusions as well as general continuous-time Markov chain models (CTMC). While log-linear closed form UD calculations are possible for Langevin models with coefficients

directly relating habitat to space-use, CTMC UD^s have only been calculated numerically without parameters relating directly habitat to use. Here we present a general class of CTMC models and show they have a closed form log-linear predictive UD. CTMC^s have several benefits over Langevin models: (1) the exact likelihood can be used without approximation, (2) both categorical and continuous habitat variables can be used, and (3) fixed barriers to movement (e.g., land in marine settings) are automatically accounted for as part of the model. We developed the R package **walk** for fitting these models in practice and it is used to demonstrate explicit UD estimation from CTMC models with a data set of Hawaiian false killer whales (*Pseudorca crassidens*). We also illustrate a 2-stage meta-analysis method for population inference from individual UD^s.

Key words: Hidden Markov model, resource selection, Stationary distribution, Satellite telemetry, Utilization distribution

1 Introduction

Any quick literature search will reveal that remote collection and analysis of animal locations, or location telemetry data, have become a ubiquitous and important tool in the advancement of animal ecology. A perennial subject within animal telemetry analysis is the assessment of space-use. The topic of animal space-use is often used in conjunction with the terms home range (Burt, 1943) or resource selection (Johnson, 1980), but here we are specifically concerned with the concept of utilization distributions (UD; Powell and Mitchell 2012). UD^s are a spatial probability distribution that describes an animal’s location at any random time. As opposed to standard (Keating and Cherry, 2009) or correlated kernel density estimates such as Brownian bridge (Horne et al., 2007) or AKDE (Fleming et al., 2015; Calabrese et al., 2016) we consider UD models that explicitly relate

habitat variables to use through the log-linear form,

$$\mathbf{u}(\mathbf{s}) \propto \exp\{\mathbf{x}(\mathbf{s})'\boldsymbol{\beta}\}, \quad (1)$$

where $\mathbf{u}(\mathbf{s})$ is the UD at location $\mathbf{s}$ and $\mathbf{x}(\mathbf{s})$ are habitat covariates at location $\mathbf{s}$. Inference for parametric spatial UDs from temporally correlated telemetry data can be challenging (Johnson et al., 2013). Estimates that ignore the temporal aspect can suffer from start-bias (Whitehead and Jonsen, 2013; Calabrese et al., 2016) if the individual does not have a long enough deployment to fully explore its home range. Also, treating locations as independent will underestimate uncertainty in $\boldsymbol{\beta}$ inference (Alston et al., 2023).

To account for autocorrelated telemetry data predictive UDs have been proposed. Predictive UDs are derived by fitting an autocorrelated movement model to the telemetry data the stationary distribution of the movement model is treated as the UD. Predictive UDs can help eliminate start bias because they predict where the individual will be in the long-run (Whitehead and Jonsen, 2013) and the estimated parameters are estimated by explicitly modeling autocorrelation which avoids the predicament of underestimation of parameter uncertainty by treating the data as independent. Barnett and Moorcroft (2008) and Moorcroft et al. (1999) considered mechanistic stochastic differential equation (SDE) movement models but the resulting UD was time consuming to numerically assess. This same approach was also used to scale step-selection models to the resulting UD (Potts and Börger, 2022). Langevin diffusions were recently proposed as an alternative SDE model because they have a log-linear form (1) stationary distribution (Michelot et al., 2019). However, only continuously valued habitat variables could be used. Moreover, a 2-stage imputation procedure (Scharf et al., 2017) is necessary for approximate inference unless the telemetry data is sufficiently fine scale. Also, in study areas with complex barriers to movement, such as coastal marine environments, the imputation approach can become

challenging to implement.

Continuous-time Markov chain (CTMC) movement models have become another type of movement model that can easily incorporate habitat variables to describe movement. We provide a brief description here, but more technical details can be found in (Norris, 1998). In a CTMC movement model the geographical study domain is partitioned into cells or other discrete locations indexed by $\mathcal{C} = \{1, \dots, n\}$ (See Figure 1B-E). Each cell, i , has a set of neighboring cells $\mathcal{N}_i = \{j \in \mathcal{C} : i \sim j\}$, where $i \sim j$ means that an animal can go from cell i to cell j in one move. For example, in a raster partition, neighboring cells might share a border (rook neighborhood) or a corner point (queen neighborhood). CTMC models are defined by the conditional movement rates between cells, $\lambda_{ij} \geq 0$. The total rate of movement away from cell i is $\Lambda_i = \sum_{j \in \mathcal{N}_i} \lambda_{ij}$. After an individual enters cell i it will reside there an exponential, mean Λ_i^{-1} , amount of time, then transition to a neighboring cell with probability $P_{ij} = \lambda_{ij}/\Lambda_i$. Two important benefits of CTMC models over SDEs is that categorical habitat variables can be used and CTMCs can be formulated as large sparse hidden Markov models (HMMs) so that efficient evaluation of the exact likelihood is possible. Although to this point most analyses of satellite location telemetry data with CTMCs have also relied on imputation, for example, Hanks et al. (2015); Wilson et al. (2018); Johnson et al. (2021), but see Pedersen et al. (2011) for an archival tag CTMC HMM. One additional benefit of CTMCs relative to SDEs is that complex barriers can be easily accounted for simply by the definition of the cell partitions (Wilson et al., 2018), that is, for a cell j where the animal cannot travel, $\lambda_{ij} \equiv 0$ for all $i \in \mathcal{N}_j$. Or, in other words $j \notin \mathcal{C}$. If all the rates are placed in an $n \times n$ matrix $\mathbf{Q}$ with $(-\Lambda_1, \dots, -\Lambda_n)$ on the diagonal and λ_{ij} in the i th row, j th column then one can also represent a CTMC as the discrete-space SDE,

$$\frac{d}{dt}\mathbf{u}_t = \mathbf{Q}\mathbf{u}_t,$$

where $\mathbf{u}_t$ is the distribution of the individuals location at time t . The solution to this SDE is,

$$\mathbf{u}_t = \mathbf{\Gamma}(t)\mathbf{u}_0,$$

where $\mathbf{\Gamma}(t) = \exp(\mathbf{Q}t)$ is the transition probability matrix (TPM) and “exp” is the matrix exponential function. If the rates and study area are constructed such that: (1) rates are constant through time (homogeneity), (2) any cell can be reached from any other cell given sufficient time (irreducibility), and (3) once a cell is departed the probability of returning in a finite amount of time is > 0 (positive recurrence), then the CTMC will have a stationary distribution $\mathbf{u}_t \rightarrow \mathbf{u}$ as $t \rightarrow \infty$ (Whitehead and Jonsen, 2013; Wilson et al., 2018). This stationary distribution is the eigenvector of $\mathbf{Q}$ associated with the zero eigenvalue. Therefore, a general CTMC movement model does not have the interpretable log-linear form (1) with parameters that directly relate use to habitat variables as it must be numerically determined.

In order to blend the benefits of the CTMC based movement model with the closed-form, log-linear UD of the Langevin SDE model we propose a class of CTMC models for which one can obtain closed-form log-linear predictions of the UD from the fitted movement model parameters. Markov Chain Monte Carlo (MCMC) theory is used to verify that these specific models possess the desired form for their stationary distributions (e.g., Michelot et al. 2018; Blackwell 2026) and we develop the HMM likelihood for fitting the model, specifically for normally distributed location errors. To fit these models, we developed the R package `walk*` (Johnson, 2026) to estimate parameters of general CTMC models using the exact HMM likelihood (McClintock et al., 2020). Which to our knowledge has not been formulated and used for CTMC-based animal movement using noisy satellite location telemetry data. The process of fitting models and making UD inference is

*Available from: <https://dsjohnson.r-universe.dev/walk>

demonstrated with telemetry data from a false killer whale (*Pseudorca crassidens*) population in the Hawaiian Islands. In addition to individual UD inference we also adapt the 2-stage meta-analysis method of Johnson et al. (2022) for approximating population UD inference from a Bayesian random effects CTMC model.

2 Explicit Utilization Distribution CTMC Models

General CTMCs do not have closed-form stationary distributions but here we propose a class of CTMC useful for animal movement modeling for which we can directly evaluate the stationary distribution. The movement rates of this class have the form,

$$\log(\lambda_{ij}) = \mathbf{x}'_{1i}\boldsymbol{\alpha} + \mathbf{x}'_{2j}\boldsymbol{\delta} + (\mathbf{x}_{3j} - \mathbf{x}_{3i})'\boldsymbol{\gamma} + \mathbf{x}'_{4ij}\boldsymbol{\eta}, \quad (2)$$

where $\mathbf{x}_{1i}$ is a set of habitat covariates recorded in the current cell i , $\mathbf{x}_{2j}$ are other covariates recorded in cell $j \in \mathcal{N}_i$, $\mathbf{x}_{3j} - \mathbf{x}_{3i}$ are local differences of a third set of habitat covariates, and $\mathbf{x}_{4ij}$ are symmetric pair specific covariates ($\mathbf{x}_{4ij} = \mathbf{x}_{4ji}$) such as, say, distance between i and j . The covariates in $\mathbf{x}_1$ only control residence time and do not affect which cell an individual might choose upon moving. The variables in $\mathbf{x}_2$ primarily function to specify preference for neighboring cells based on the absolute value of the covariates, while the variables included via local differences, $\mathbf{x}_3$, control movement based on relative habitat differences, similar to habitat gradient effects in Langenvin diffusions. Now that we have a defined class of CTMC movement models we can evaluate the stationary utilization distribution that results from this CTMC diffusion.

2.1 Evaluating the stationary utilization distribution

To evaluate the stationary distribution and make sure one exists we first check the 3 convergence criteria. First, all rates are homogeneous through time. Second, all movement rates in (2) are positive for finite covariates and coefficients, therefore if $i \sim j$, then $j \sim i$. Finally, if the study area is not divided into separate regions for which inter-region moves are impossible, then the positive rates imply irreducibility and positive recurrence.

We have established that, outside of pathological habitat variables and study areas, a CTMC with rates defined by (2) will have a stationary utilization distribution. To assess what the actual form of the UD is we begin with another theoretical result for evaluating a CTMC stationary distribution, that is,

$$u_i \propto \pi_i / \Lambda_i, \quad (3)$$

where $\boldsymbol{\pi}$ is the stationary distribution of the embedded Markov chain of the CTMC (Norris, 1998; Wilson et al., 2018). The embedded Markov chain is the discrete-time list of cells visited by the CTMC disregarding the actual times when it moves. The embedded Markov chain is an true discrete-time Markov chain with transition probabilities, $P_{ij} = \lambda_{ij} / \Lambda_i$. Therefore, we can work in the discrete-time to find the stationary distribution of the CTMC.

We can use Markov Chain Monte Carlo (MCMC) theoretical results to show that a CTMC movement model has a stationary distribution of log-linear form (e.g., Michelot et al. 2018; Blackwell 2026). To verify the closed-form stationary distribution we need only check that the MCMC reversibility condition is satisfied for a given $\boldsymbol{\pi}$, that is,

$$\pi_i P_{ij} = \pi_j P_{ji},$$

for all i and j in $\mathcal{C}$ (Blackwell, 2026). By inserting P_{ij} and P_{ji} into the equation, one can check that it is satisfied for $\pi_i \propto \exp(-\mathbf{x}'_{1i}\alpha + \mathbf{x}'_{2i}\delta + 2\mathbf{x}'_{3i}\gamma)\Lambda_i$ (See online supplement S1). This implies, using (3), the stationary UD for the full CTMC model is,

$$\begin{aligned} u_i &\propto \exp(-\mathbf{x}'_{1i}\alpha + \mathbf{x}'_{2i}\delta + 2\mathbf{x}'_{3i}\gamma) \\ &= \exp(\mathbf{x}'_i\beta), \end{aligned} \tag{4}$$

where $\mathbf{x}_i = (\mathbf{x}'_{1i}, \mathbf{x}'_{2i}, \mathbf{x}'_{3i})'$ and $\beta = (-\alpha', \delta', 2\gamma')'$. Thus, the utilization distribution for the proposed class of CTMC movement models has the desired closed log-linear form (1).

2.2 Parameter Estimation

Previous estimation of CTMC movement parameters using satellite location telemetry data used an imputation method (see Scharf et al. 2017) which fills in missing cell locations and transition times between location acquisition times (Hanks et al., 2015; Wilson et al., 2018; Johnson et al., 2021). Even though the imputation method attempts to propagate the uncertainty of the missing locations, it is unclear how this affects estimation of CTMC parameters. Therefore we propose use the exact, although potentially more computationally expensive, hidden Markov model (HMM) formulation of the model.

2.2.1 Satellite telemetry data specification and processing

For the proposed analysis we consider satellite telemetry data for an individual of the form $\mathbf{y} = \{\mathbf{y}_1, \dots, \mathbf{y}_K\}$, where $\mathbf{y}_k = (\mathbf{s}_k, t_k, \boldsymbol{\psi}_k)$, $\mathbf{s}_k$ is the observed location of the individual within the study area at time t_k and $\boldsymbol{\psi}_k$ represents information or parameters governing location uncertainty. For example, $\boldsymbol{\psi}_k$ might represent the major and minor error ellipse lengths of a Kalman Filtered Argos location (McClintock et al., 2015) or a known error radius for GPS locations, e.g., 20m. From these locations we translate the continuous-space

163 data locations and uncertainty to the discrete space, $\mathcal{C}$, using a function, say,
 164 $p(\mathbf{y}_k, \mathcal{C}) = \mathbf{p}_k = (p_{1k}, \dots, p_{nk})$, where p_{ik} is the probability that the animal is located within
 165 cell i at time t_k . We used a general form because the function, p , will depend on how
 166 location uncertainty is assessed for the specific satellite and tag hardware. However,
 167 uncertainty in the true location, $\boldsymbol{\mu}_k$, is often represented by the bivariate Normal
 168 distribution $\boldsymbol{\mu}_k \sim N(\mathbf{s}_k, \boldsymbol{\Sigma}_k)$, where $\boldsymbol{\psi}_k = \boldsymbol{\Sigma}_k$. In that case, p_{ik} would represent the mass of
 169 the Normal distribution contained within the i th cell. For a raster grid partitioning of the
 170 study area this can be calculated by,

$$p_{ik} = \text{CDF}(x_{i2}, y_{i2}) - \text{CDF}(x_{i2}, y_{i1}) - \text{CDF}(x_{i1}, y_{i2}) + \text{CDF}(x_{i1}, y_{i1})$$

171 where $[x_{i1}, x_{i2}] \times [y_{i1}, y_{i2}]$ is the bounding box for the i th cell and CDF is the bivariate
 172 normal cumulative density function with mean $\mathbf{s}_k$ and covariance $\boldsymbol{\Sigma}_k$ (see Figure S1 for
 173 illustration of $\mathbf{p}_k$). A Monte Carlo option for very general error distributions would be to
 174 draw a sufficiently large sample from the distribution and assess the proportion of samples
 175 that falls within each cell. If the error distribution is known, this only has to be done one
 176 time.

177 2.2.2 Likelihood calculation

178 For an individual, the likelihood is calculated with the matrix equation (Zucchini et al.,
 179 2016; McClintock et al., 2020),

$$[\mathbf{y} | \boldsymbol{\alpha}, \boldsymbol{\delta}, \boldsymbol{\gamma}, \boldsymbol{\eta}] = \mathbf{u}'_1 \mathbf{P}(\mathbf{y}_1) \boldsymbol{\Gamma}(t_2 - t_1) \mathbf{P}(\mathbf{y}_2) \cdots \boldsymbol{\Gamma}(t_K - t_{K-1}) \mathbf{P}(\mathbf{y}_K) \mathbf{1} \quad (5)$$

180 where $\mathbf{u}_1$ is the location probability distribution at t_1 , $\mathbf{P}(\mathbf{y}_k) = \text{diag}(\mathbf{p}_k)$, $\boldsymbol{\Gamma}(t_k - t_{k-1})$ is the
 181 TPM evaluated over the interval $t_k - t_{k-1}$, and $\mathbf{1}$ is a vector of 1s. The natural choice for

$\mathbf{u}_1$ is $\mathbf{u}$, the stationary distribution defined by the CTMC movement model, but there are other choices. For example, if a telemetered individual was released at a known location after tagging that may not be representative of its usual movement, then $\mathbf{u}_1$ would be a vector of all 0s except for the cell containing the release site, which would be a 1. In the false killer whale example below, we use $\mathbf{u}_1 = \mathbf{u}$ because they were tagged where they were encountered in the wild. In the R package `walk` we use scaling methods to avoid numerical underflow as well as the efficient sparse matrix uniformization methods presented in Sherlock (2021) to calculate products of the form $\mathbf{v}' \exp\{\mathbf{Q}(t_k - t_{k-1})\}$.

In addition to the matrix exponential evaluation numerical savings, another optional approach to save computational effort is to set a censoring value for the maximum Λ_i value. The numerical methods of Sherlock (2021) require more matrix operations for higher values of $\max(\Lambda_i)$. For some cells, the log linear λ_{ij} model may produce rates of movement that are unrealistically high in some parts of the study area. With respect to UD this would imply that an individual will not remain in that cell and the $u_i \approx 0$. Hewitt et al. (2023) considered a logit rate model that restricts this behavior, but that model does not allow us to construct a CTMC with an explicit stationary distribution. So, we propose picking a censoring value, say, Λ^* and setting $\lambda_{ij}^* = (\Lambda^*/\Lambda_i)\lambda_{ij}$ if $\Lambda_i > \Lambda^*$. If Λ^* is sufficiently large, this will have negligible impact on $\mathbf{u}$ estimation but can dramatically help likelihood computation. One reasonable way to choose Λ^* is to select the minimum amount of time that an animal would likely reside in a cell, say r^* , then set $\Lambda^* = 1/r^*$. This implies that the expected residency time $\geq r^*$ for all cells. The effect on $\mathbf{u}$ can be verified by comparing the exact, numerically calculated $\mathbf{u}$ from the censored rate model (Wilson et al., 2018) to $u_i \propto \exp(\mathbf{x}_i' \boldsymbol{\beta})$. If discrepancies are sufficiently large, Λ^* can be increased until the discrepancy is within tolerance. In the resulting model, if $\max(\Lambda_i) \leq \Lambda^*$ the UD is identical.

Once the CTMC movement model has been fitted the maximum likelihood estimate,

208 $\hat{\boldsymbol{\beta}}_{\text{CTMC}} = (\hat{\boldsymbol{\alpha}}', \hat{\boldsymbol{\delta}}', \hat{\boldsymbol{\gamma}}')'$, needs to be transformed to the UD coefficients via $\hat{\boldsymbol{\beta}} = \mathbf{M}\hat{\boldsymbol{\beta}}_{\text{CTMC}}$ and
 209 $\mathbf{M} = \text{diag}((-1'_{\alpha}, 1'_{\delta}, 21'_{\gamma})')$ Thus, the large sample covariance matrix of the UD coefficients
 210 is $\hat{\mathbf{S}} = \mathbf{M}\hat{\mathbf{S}}_{\text{CTMC}}\mathbf{M}$, where $\hat{\mathbf{S}}_{\text{CTMC}}$ is the Hessian-derived large sample covariance matrix of
 211 $\hat{\boldsymbol{\beta}}_{\text{CTMC}}$. If only the standard errors are desired, one can simply multiply the CTMC
 212 coefficient standard errors by the multipliers, but if the entire covariance matrix is desired
 213 the matrix multiplication is necessary.

214 2.2.3 Population-level inference

215 Individual estimates of UD coefficients and predicted UD's can be difficult to interpret for
 216 population inference and management considerations. Therefore, we suggest the more
 217 appropriate model to fit is the Bayesian random effects model,

$$[\mathbf{y}_1, \dots, \mathbf{y}_N | \boldsymbol{\beta}_1, \dots, \boldsymbol{\beta}_N] = \prod_{l=1}^N [\mathbf{y}_l | \mathbf{M}^{-1}\boldsymbol{\beta}_l, \boldsymbol{\eta}_l]$$

$$[\boldsymbol{\beta}_1, \dots, \boldsymbol{\beta}_N | \boldsymbol{\theta}, \boldsymbol{\Sigma}] = \prod_{l=1}^N N(\boldsymbol{\beta}_l | \boldsymbol{\theta}, \boldsymbol{\Sigma})$$

218 where $\mathbf{y}_l$ is the observed data for individual $l = 1, \dots, N$, $\mathbf{M}^{-1}\boldsymbol{\beta}_l = (\boldsymbol{\alpha}'_l, \boldsymbol{\delta}'_l, \boldsymbol{\gamma}'_l)'$ and $\boldsymbol{\eta}_l$ are
 219 the movement parameters for individual l , and the prior distribution for the random
 220 population mean coefficient and variance is $[\boldsymbol{\theta}][\boldsymbol{\Sigma}]$. Even if individual inference is desired, it
 221 is useful to fit this hierarchical model because it allows us to share information between
 222 individuals. via the posterior distribution $[\boldsymbol{\beta}_l | \mathbf{y}_1, \dots, \mathbf{y}_N]$. If we assume that there is some
 223 similarity in movement between individuals then we can leverage all data for better
 224 individual inference. Even though the random effects model would be beneficial, however,
 225 due to the computational demands for fitting the CTMC model to one individual, a full
 226 joint random effects model is not currently feasible.

227 Although we cannot fit the full random effects model all at once we can tailor the
 228 general 2-stage meta-analysis methods of Johnson et al. (2022) to approximate inference

for the hierarchical CTMC model using maximum likelihood estimates for each individual. In the first stage individual MLEs, $\hat{\beta}_l$ and covariance matrices, $\hat{\mathbf{S}}_l$ are obtained using (5). In the second stage the MLEs are used as responses in the multivariate linear mixed model,

$$\hat{\beta}_l = \boldsymbol{\theta} + \mathbf{z}_l + \boldsymbol{\epsilon}_l; \quad l = 1, \dots, N$$

where the random effect is $[\mathbf{z}_l | \boldsymbol{\Sigma}] = N(\mathbf{0}, \boldsymbol{\Sigma})$, the error term is $[\boldsymbol{\epsilon}_l | \hat{\mathbf{S}}_l] = N(\mathbf{0}, \hat{\mathbf{S}}_l)$. REML or Laplace approximation procedures are used to fit this model subject to the prior constraint of $[\boldsymbol{\theta} | \boldsymbol{\Sigma}]$. Following second stage model fitting we can extract the approximate MAP estimates $\bar{\boldsymbol{\theta}}$ and $\bar{\beta}_l = \bar{\boldsymbol{\theta}} + \bar{\mathbf{z}}_l$, where $\bar{\beta}_l$ is the approximate MAP of the posterior $[\beta_l | \mathbf{y}_1, \dots, \mathbf{y}_N]$.

3 Illustration: False Killer Whale Space-Use

In this section, we will apply the modeling approach described in the previous sections to a telemetry dataset on endangered false killer whales (FKW; *Pseudorca crassidens*) in the Hawaiian Islands (Baird 2016). This particular population is resident to the insular waters of the main Hawaiian Islands (MHI; includes Kaua‘i/Ni‘ihau, Oahu, Maui Nui, and Hawai‘i Island) and is comprised of five social clusters, or groups of close family members and regular associates (Baird et al., 2008, 2019; Baird, 2016; Martien et al., 2019). These whales are known to move quickly and frequently among island areas throughout their range (Baird et al., 2010, 2012, 2021; Baird, 2016). Due to inherent challenges associated with tracking apex marine predators, this dataset is characterized by variable deployment durations and irregular location data streams. Considering these factors may drive start bias, as well as their expansive range, we deem this an appropriate dataset to empirically demonstrate the utility of predictive UD. For the purposes of this assessment, we only

focus on movements of 5 individuals that were satellite tagged in 2021 and 2022. There is one representative from each social cluster in the data.

3.1 Satellite tag data

False killer whale satellite tag deployment methods have been described in detail elsewhere (see Baird et al. 2010, 2012) so they are only briefly summarized here. Individuals were tagged with Argos transmitting SPOT6 ($n = 4$) and SPLASH10 ($n = 1$) tags during dedicated small boat survey efforts throughout the main Hawaiian Islands (Baird et al. 2012, 2021; Baird 2016; see Baird et al. 2013 for details on surveys). Deployment duration for these data ranged from 12 days to 81 days (median=40 days). Argos location data were processed through the Douglas-Argos Filter (accessed via Movebank; Kranstauber et al. 2011; Douglas et al. 2012) to remove erroneous locations based on unrealistic travel speeds and turning angles. The final analytical dataset consisted of 5 trajectories and 4,009 filtered Argos and Fastloc GPS (SPLASH10 tag) locations (Figure 1A). For the Argos locations, we used the Kalman-filter diagnostic ellipse data to construct error covariance matrices (McClintock et al., 2015). While for the Fastloc GPS locations, we used the 95% quantile error ranges from Dujon et al. (2014), based on the number of satellites acquired, as twice the standard deviation of the normal error.

3.2 Study area and habitat variables

The official management range of the MHI stock of FKW was documented in Bradford et al. (2015) and represents a minimum convex polygon surrounding a buffer of the main Hawaiian Islands (Figure 1A). This is the study area boundary we use for the UD analysis.

We used two modeled global oceanographic variables from E.U. Copernicus Marine Service: (1) Sea Surface Temperature (SST; °C; DOI:

<https://doi.org/10.48670/moi-00016>) and (2) Sea Level Anomaly (SLA; m; DOI: <https://doi.org/10.48670/moi-00148>) to model movement and space-use. SST is available at 0.082° resolution and SLA is available at 0.125° degree resolution. In order to match resolution we chose to project the SLA raster to the resolution of the SST raster to avoid losing oceanographic information. Because we are using these variables to predict long term space-use, we used a single layer for each variable that was the 2 year daily average in the study area. This allowed us to model movement with respect to typical oceanographic conditions (Figure 1B-C).

In addition to the oceanographic variables, we used distance from land (m) to model FKW’s resident nature in the MHI. Also, we constructed a depth residual variable by using data from the global, 30 arc-second ETOPO elevation model (NOAA National Centers for Environmental Information, 2022) and removing the linear trend of distance from land. The distance and depth variables were projected to the 0.082° SST resolution as well. Then all data, including telemetry data, were projected to the Hawaii Albers Equal Area projection, ESRI:102007 (Figure 1D-E)

3.3 CTMC model and hierarchical inference

Notationally, the CTMC movement rate model we used to model the movement of each individual $l = 1, \dots, 5$ was

$$\log \lambda_{ij} = \underbrace{(\text{Intercept}) + \text{DEPTH}_i + \text{SST}_i + \text{SST}_i^2 + \text{SLA}_i + \text{SLA}_i^2}_{\mathbf{x}_1 \text{ portion}} + \underbrace{[\text{DIST}_j - \text{DIST}_i]}_{\mathbf{x}_3 \text{ portion}}$$

In this model differences of the distance from land variable were used as a movement kernel between cells i and j . This allowed individuals to prefer movement towards (or away from) land at the same rate no matter how far away they are. In addition SST, SLA, and DIST were Z -transformed for numerical stability of the HMM likelihood. The DEPTH residual

was already on a nearly standard normal scale and DIST was only used as a difference, so values were not large enough to cause numerical issues.

In the first stage the CTMC model was fitted separately to the data from each individual using normal location errors and the HMM likelihood (5). To determine the censor value for the total movement rate we selected the minimum expected residency time to be $r^* = 1/3$ minutes. This implies a maximum total movement rate of $\Lambda^* = 60/(1/3) = 180$ because the units of time in the data are hours.

The `nimble` R package (de Valpine et al., 2017) was used to optimize the second stage multivariate linear mixed model using Laplace approximation (de Valpine et al., 2017).

The random effects and prior distributions used for the CTMC/UD coefficients are,

$$[\theta_w] = N(0, \tau_w^2); \quad [\tau_w] = \text{Exponential}(1)$$

$$\Sigma = \text{diag}(\sigma_1^2, \dots, \sigma_7^2); \quad [\sigma_w] = \text{Exponential}(1)$$

for $w = 1, \dots, 7$. The exponential hyperpriors for the variance parameters were added to penalize the complexity of the model to perform selection for the population level parameters (Hooten and Hobbs, 2015; Simpson et al., 2017).

3.4 Results

Individual results for all individuals were generally consistent with notable deviations being individual PcTag074 did not have any significant effect of SLA on its UD and PcTag070 had a negative effect of the linear SST term (Figure 2). All other coefficients were similar in direction, i.e., less than or greater than 0. This bolsters the idea of using the 2 stage meta-analytic approach to share data to improve individual estimates. The second stage individual coefficient estimates, $\bar{\beta}_i$ illustrate that there is only a significant amount of individual variation in the DEPTH and SLA² coefficients (Figure 2). All other second

stage coefficients are shrunk to the population mean estimates, $\bar{\theta}$ with nearly identical approximate credible intervals (Figure 2). There was only 1 nonsignificant population-level coefficient, the linear SLA term. Correspondingly, the penalized model complexity priors shrunk the SLA term to zero. All other population coefficients were significantly different from zero. The DEPTH and DIST coefficients were negative as expected because this population inhabits shallower waters close to the Main Hawaiian Islands without venturing too far away. The zero SLA coefficient combined with the negative SLA^2 term implies a maximum effect at 0, but we used a Z -transform, therefore, the maximum effect occurs at the mean original SLA value. Both SST coefficients are positive implying that the typical individual is probably located in the warmest SST conditions in the study area.

Unsurprisingly, the similarity in coefficient estimates produced similar predicted UD at the first stage (Figure S1, online supplement). Individual PcTag074 had a more notably diffuse UD, the other four individuals had very similar looking UD. This may be due to the lack of SLA effect on the UD of PcTag074. After the second stage, the UD are even more similar as the second stage coefficients are essentially the same at the population mean coefficients (Figure 3). The UD created from the population mean coefficients is illustrated in Figure 3 (Typical). This is the UD of a so called “Typical” or median individual, not the population average UD because of the log link between the coefficients and the UD. Even after the second stage the coefficients for PcTag074 lack any SLA effect, so the UD is the only different one from the typical UD.

In order to check that the rate truncation did not adversely affect the estimated UD we compared the truncated numerical Eigen decomposition UD ($u_{ei,i}$; Wilson et al. 2018), which is the exact UD for the censored movement models that was used in the first stage, with the exponential closed form version using Bhattacharyya’s Affinity (BA)

(Bhattacharyya, 1943),

$$BA = \sum_{i \in \mathcal{C}} \sqrt{u_{ei,i} \cdot u_i}.$$

The values of BA range from 0 (no similarity) to 1 (identical distributions) (Fieberg and Kochanny, 2005). The proportion of study area cells that are censored in the fitting of individual models ranged from 0–43% of the study area. This may, at first, seem troublesome, however, BA ranged from 0.997–1 for all of the individuals indicating that the censored cells possessed very little UD mass and the closed form target and exact censored UDs were virtually identical. Therefore, the rate censor approximation did not adversely affect parameter estimation in this case.

4 Discussion

This work introduces a continuous-time Markov chain movement process which possesses a closed form log-linear stationary distribution which is a function of spatial covariates. The coefficients in log-linear form allow direct inference of the effect of habitat variables on the utilization distribution. Moreover, because these parameters are estimated within the context of a continuous-time movement model, the temporal correlation between observations is accounted for in the standard errors of the UD coefficients. In addition to UD inference, the fitted movement parameters also provide information on resource selection and local habitat drivers of residency (Hanks et al., 2015; Johnson et al., 2021). Another benefit of the CTMC approach is that barriers to movement can be easily handled in the movement model by simple exclusion of cells on the other side of the barriers. This is especially important for analysis of marine systems such as the FKW example.

In addition to individual inference we also adapted the 2-stage meta-analytic approach of Johnson et al. (2022) to approximate inference from a Bayesian CTMC random effects model. This allows practitioners to estimate population-level space use and movement

parameters. Also, information sharing between individuals allows improved inference if movement patterns are similar between individuals. As part of this research, the R package `walk` was developed for processing continuous-space location telemetry data into a discrete-space form and fitting CTMC models using the exact HMM likelihood.

While overarching movement and space-use inference is also shared by Langevin diffusion movement models (Michelot et al., 2019), there are benefits to using the CTMC models proposed here. First, although it can be computationally intensive, one can use the exact likelihood for the movement data. For Langevin diffusion inference an Euler or Ozaki time discretization approximation is needed within a pseudo-likelihood for parameter estimation (Gloaguen et al., 2018; Michelot et al., 2019). One might make the argument that the CTMC approach just exchanges discrete space for discrete time approximation. This may be true for constructed variables such as distance from land in the FKW example. However, many habitat variables are only available as cellular rasters, e.g., SST and SLA. Therefore, the Langevin pseudo-likelihoods must also spatially approximate the gradients of these variables as well. In addition, while the Langevin model can, in theory, handle uneven gaps in telemetry data, in practice, interpolation is necessary for model fitting. If there is no location error, then straight line interpolation may be sufficient, but location uncertainty generally implies a stochastic imputation method is needed (Michelot et al., 2019; Scharf et al., 2017). The CTMC model presented here does not need data interpolation or specification of an imputation model. Finally, the Langevin models are not well suited to use of categorical habitat characteristics (Michelot et al., 2019) and a smoothing of indicator variables must be used. The CTMC model proposed here can use categorical variables just as easily as continuously valued variables.

Given the apparent positive aspects of the closed-form CTMC approach, it behooves us to look at some potential downsides. The first issue is that CTMC models are built on a defined set of cells, if it is changed, for example, the resolution of a raster grid is increased,

it is not apparent how the parameters relate to each other. If the grid is fixed based on habitat covariates, then this is probably not an issue because that is the spatial resolution that habitat information is available. However, if there is some flexibility in resolution of cells then some consideration is warranted. The second issue is that the TPM, $\Gamma(t)$, can become difficult to compute for very large study areas relative to cell resolution. Uniformization methods (Sherlock, 2021) and rate censoring as implemented in the `walk` package provide ways to mitigate this issue. But for extremely large study areas and very fast moving species (i.e., high $\max(\Lambda_i)$) it can become burdensome. Alternative uniformization methods which calculate likelihood parameter gradients (Rupp et al., 2024) may provide avenues to speed optimization.

Ultimately, bridging the gap between fine-scale movement decisions and long-term space-use remains a central challenge in spatial ecology. By offering a computationally tractable, closed-form bridge between continuous-time movement and utilization distributions, the CTMC framework provides a unified approach to movement and space-use modeling. As telemetry datasets continue to grow in resolution and volume, methods that seamlessly blend animal movement with flexible population-level inference will be essential for management and conservation. The combination of CTMC models for individual movement and meta-analysis methods for population inference provides a theoretical and practical approach for transforming raw telemetry data into scientific inference.

Acknowledgments

The findings and conclusions in the paper are those of the authors and do not necessarily represent the views of the National Marine Fisheries Service, NOAA. Reference to trade names does not imply endorsement by the National Marine Fisheries Service, NOAA.

Conflicts of interest

The authors declare that they have no competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

- Alston, J. M., Fleming, C. H., Kays, R., Streicher, J. P., Downs, C. T., Ramesh, T., Reineking, B., and Calabrese, J. M. (2023). Mitigating pseudoreplication and bias in resource selection functions with autocorrelation-informed weighting. *Methods in Ecology and Evolution*, 14(2):643–654.
- Baird, R. W. (2016). *The Lives of Hawai‘i’s Dolphins and Whales: Natural History and Conservation*. University of Hawaii press.
- Baird, R. W., Anderson, D. B., Kratofil, M. A., and Webster, D. L. (2021). Bringing the right fishermen to the table: Indices of overlap between endangered false killer whales and nearshore fisheries in Hawai‘i. *Biological Conservation*, 255:108975.
- Baird, R. W., Anderson, D. B., Kratofil, M. A., Webster, D. L., and Mahaffy, S. D. (2019). Cooperative conservation and long-term management of false killer whales in Hawai‘i: Geospatial analyses of fisheries and satellite tag data to understand fishery interactions. A Report to the State of Hawai‘i Board of Land and Natural Resources, Contract No. 67703, Cascadia Research Collective.
- Baird, R. W., Gorgone, A. M., McSweeney, D. J., Webster, D. L., Salden, D. R., Deakos, M. H., Ligon, A. D., Schorr, G. S., Barlow, J., and Mahaffy, S. D. (2008). False killer whales (*Pseudorca crassidens*) around the main Hawaiian Islands: Long-term site

fidelity, inter-island movements, and association patterns. *Marine Mammal Science*,
24(3):591–612.

Baird, R. W., Hanson, M. B., Schorr, G. S., Webster, D. L., McSweeney, D. J., Gorgone,
A. M., Mahaffy, S. D., Holzer, D. M., Oleson, E. M., and Andrews, R. D. (2012). Range
and primary habitats of Hawaiian insular false killer whales: Informing determination of
critical habitat. *Endangered Species Research*, 18(1):47–61.

Baird, R. W., Schorr, G. S., Webster, D. L., McSweeney, D. J., Hanson, M. B., and
Andrews, R. D. (2010). Movements and habitat use of satellite-tagged false killer whales
around the main Hawaiian Islands. *Endangered Species Research*, 10:107–121.

Baird, R. W., Webster, D. L., Aschettino, J. M., Schorr, G. S., and McSweeney, D. J.
(2013). Odontocete cetaceans around the main Hawaiian Islands: Habitat use and
relative abundance from small-boat sighting surveys. *Aquatic Mammals*, 39(3).

Barnett, A. H. and Moorcroft, P. R. (2008). Analytic steady-state space use patterns and
rapid computations in mechanistic home range analysis. *Journal of Mathematical
Biology*, 57(1):139–159.

Bhattacharyya, A. (1943). On a measure of divergence between two statistical populations
defined by their probability distribution. *Bulletin of the Calcutta Mathematical Society*,
35:99–110.

Blackwell, P. G. (2026). Consistent multiscale modelling of movement and habitat
selection. *Journal of Mathematical Biology*, 92(6):90.

Bradford, A. L., Oleson, E. M. E. M., Baird, R. W., Boggs, C. H., Forney, K. A., and
Young, N. C. (2015). Revised stock boundaries for false killer whales (*Pseudorca
crassidens*) in Hawaiian waters. <https://repository.library.noaa.gov>.

- Burt, W. H. (1943). Territoriality and Home Range Concepts as Applied to Mammals. *Journal of Mammalogy*, 24(3):346–352.
- Calabrese, J. M., Fleming, C. H., and Gurarie, E. (2016). Ctm: An R package for analyzing animal relocation data as a continuous-time stochastic process. *Methods in Ecology and Evolution*, 7(9):1124–1132.
- de Valpine, P., Turek, D., Paciorek, C. J., Anderson-Bergman, C., Lang, D. T., and Bodik, R. (2017). Programming with models: Writing statistical algorithms for general model structures with NIMBLE. *Journal of Computational and Graphical Statistics*, 26(2):403–413.
- Douglas, D. C., Weinzierl, R., C. Davidson, S., Kays, R., Wikelski, M., and Bohrer, G. (2012). Moderating A rgos location errors in animal tracking data. *Methods in Ecology and Evolution*, 3(6):999–1007.
- Dujon, A. M., Lindstrom, R. T., and Hays, G. C. (2014). The accuracy of Fastloc-GPS locations and implications for animal tracking. *Methods in Ecology and Evolution*, doi: 10.1111/2041-210X.12286.
- Fieberg, J. and Kochanny, C. O. (2005). Quantifying Home-Range Overlap: The Importance of the Utilization Distribution. *The Journal of Wildlife Management*, 69(4):1346–1359.
- Fleming, CH., Fagan, WF., Mueller, T., Olson, KA., Leimgruber, P., and Calabrese, JM. (2015). Rigorous home range estimation with movement data: A new autocorrelated kernel density estimator. *Ecology*, 96(5):1182–1188.
- Gloaguen, P., Etienne, M.-P., and Le Corff, S. (2018). Stochastic Differential Equation

- Based on a Multimodal Potential to Model Movement Data in Ecology. *Journal of the Royal Statistical Society Series C: Applied Statistics*, 67(3):599–619.
- Hanks, E. M., Hooten, M. B., and Alldredge, M. W. (2015). Continuous-time discrete-space models for animal movement. *The Annals of Applied Statistics*, 9:145–165.
- Hewitt, J., Gelfand, A. E., and Schick, R. S. (2023). Time-discretization approximation enriches continuous-time discrete-space models for animal movement. *The Annals of Applied Statistics*, 17(1):740–760.
- Hooten, M. B. and Hobbs, N. T. (2015). A guide to Bayesian model selection for ecologists. *Ecological Monographs*, 85(1):3–28.
- Horne, J. S., Garton, E. O., Krone, S. M., and Lewis, J. S. (2007). Analyzing animal movements using Brownian bridges. *Ecology*, 88:2354–2363.
- Johnson, D. (1980). The comparison of usage and availability measurements for evaluating resource preference. *Ecology*, 61:65–71.
- Johnson, D. S. (2026). *Walk: Model Animal Movement with Continuous-Time Markov Chains*. R package version 0.0.0.942. <https://github.com/dsjohnson/walk>.
- Johnson, D. S., Brost, B. M., and Hooten, M. B. (2022). Greater than the sum of its parts: Computationally flexible Bayesian hierarchical modeling. *Journal of Agricultural, Biological and Environmental Statistics*, 27(2):382–400.
- Johnson, D. S., Hooten, M. B., and Kuhn, C. E. (2013). Estimating animal resource selection from telemetry data using point process models. *Journal of Animal Ecology*, 82:1155–1164.
- Johnson, D. S., Pelland, N. A., and Sterling, J. T. (2021). A continuous-time semi-Markov

model for animal movement in a dynamic environment. *Annals of Applied Statistics*,
15(2):797–812.

Keating, K. A. and Cherry, S. (2009). Modeling utilization distributions in space and time.
Ecology, 90:1971–1980.

Kranstauber, B., Cameron, A., Weinzerl, R., Fountain, T., Tilak, S., Wikelski, M., and
Kays, R. (2011). The Movebank data model for animal tracking. *Environmental
Modelling & Software*, 26(6):834–835.

Martien, K. K., Taylor, B. L., Chivers, S. J., Mahaffy, S. D., Gorgone, A. M., and Baird,
R. W. (2019). Fidelity to natal social groups and mating within and between social
groups in an endangered false killer whale (*Pseudorca crassidens*) population.
Endangered Species Research, 40:219–230.

McClintock, B. T., Langrock, R., Gimenez, O., Cam, E., Borchers, D. L., Glennie, R., and
Patterson, T. A. (2020). Uncovering ecological state dynamics with hidden Markov
models. *Ecology Letters*, 23(12):1878–1903.

McClintock, B. T., London, J. M., Cameron, M. F., and Boveng, P. L. (2015). Modelling
animal movement using the Argos satellite telemetry location error ellipse. *Methods in
Ecology and Evolution*, In press:doi: 10.1111/2041-210X.12311.

Michelot, T., Blackwell, P. G., and Matthiopoulos, J. (2018). Linking resource selection
and step selection models for habitat preferences in animals. *Ecology*.

Michelot, T., Etienne, M.-P., and Gloaguen, P. (2019). The Langevin diffusion as a
continuous-time model of animal movement and habitat selection. *Methods In Ecology
and Evolution*, In press.

523 Moorcroft, P. R., Lewis, M. A., and Crabtree, R. L. (1999). Home range analysis using a
524 mechanistic home range model. *Ecology*, 80(5):1656–1665.

525 NOAA National Centers for Environmental Information (2022). ETOPO 2022 15
526 arc-second global relief model.

527 Norris, J. R. (1998). *Markov Chains*. Cambridge University Press, 2nd edition edition.

528 Pedersen, M. W., Patterson, T. A., Thygesen, U. H., and Madsen, H. (2011). Estimating
529 animal behavior and residency from movement data. *Oikos (Copenhagen, Denmark)*,
530 120(9):1281–1290.

531 Potts, J. R. and Börger, L. (2022). How to scale up from animal movement decisions to
532 spatiotemporal patterns: An approach via step selection. *Journal of Animal Ecology*.

533 Powell, R. A. and Mitchell, M. S. (2012). What is a home range? *Journal of mammalogy*,
534 93(4):948–958.

535 Rupp, K., Schill, R., Süskind, J., Georg, P., Klever, M., Lösch, A., Grasedyck, L., Wettig,
536 T., and Spang, R. (2024). Differentiated uniformization: A new method for inferring
537 Markov chains on combinatorial state spaces including stochastic epidemic models.
538 *Computational Statistics*, 39(7):3643–3663.

539 Scharf, H., Hooten, M. B., and Johnson, D. S. (2017). Imputation Approaches for Animal
540 Movement Modeling. *Journal of Agricultural, Biological, and Environmental Statistics*,
541 22(3):335–352.

542 Sherlock, C. (2021). Direct statistical inference for finite Markov jump processes via the
543 matrix exponential. *Computational Statistics*, 36(4):2863–2887.

544 Simpson, D. P., Rue, H., Riebler, A., Martins, T. G., and Sørbye, SH. (2017). Penalising

- 545 model component complexity: A principled, practical approach to constructing priors.
546 *Statistical Science*, 32:1–28.
- 547 Thorson, J. and Kristensen, K. (2024). *Spatio-Temporal Models for Ecologists*. Chapman
548 and Hall/CRC.
- 549 Whitehead, H. and Jonsen, I. D. (2013). Inferring animal densities from tracking data
550 using markov chains. *PloS one*, 8(4):e60901.
- 551 Wilson, K., Hanks, E., and Johnson, D. (2018). Estimating animal utilization densities
552 using continuous-time Markov chain models. *Methods in Ecology and Evolution*,
553 9(5):1232–1240.
- 554 Zucchini, W., MacDonald, I. L., and Langrock, R. (2016). *Hidden Markov Models for Time*
555 *Series: An Introduction Using R*. CRC Press, 2 edition.

556 **Figure 1:** Telemetry data for five false killer whales in the Main Hawaiian Island Insular
557 stock and habitat layers for CTMC modeling. The original values are presented here, but
558 the oceanographic variables (B and C) and distance from land (D) were Z -transformed in
559 model fitting.

560 **Figure 2:** Parameter estimates for fitted UD surfaces. The Stage 1 values are from
561 individually fitted models. The Stage 2 estimates are derived using the two stage method
562 of Johnson et al. (2022) to approximate inference from fitting a fully joint Bayesian random
563 effects model. The solid vertical lines are the estimated random effect means and the
564 dashed lines are the approximate 95% credible intervals.

565 **Figure 3:** Individual Predicted UD's using parameter estimates from the second stage
566 linear mixed model. The parameter estimates, $\bar{\beta}_i$ are approximate maximum *a posteriori*
567 estimates from the posterior $[\beta_i | \mathbf{y}_1, \dots, \mathbf{y}_5]$. The “Typical” UD plot is created from the
568 population mean coefficients.




Figure 1




Figure 2




Figure 3