

**Journal:** Ecology

**Manuscript type:** Statistical Innovation

# Predicting Animal Space-Use with Continuous-Time Markov Chains

Devin S. Johnson<sup>1,\*</sup>, Michaela A. Kratofil<sup>2,3</sup>, Janelle J. Gardner<sup>1</sup>, Robin W. Baird<sup>3</sup>, and  
Amanda L. Bradford<sup>1</sup>

<sup>1</sup>Pacific Islands Fisheries Science Center, National Marine Fisheries Service, NOAA,  
Honolulu, Hawaii, USA

<sup>2</sup>Marine Mammal Institute, Department of Fisheries, Wildlife, and Conservation Sciences,  
Oregon State University, Newport, Oregon, USA

<sup>3</sup>Cascadia Research Collective, Olympia, Washington, USA

\*Corresponding author: [devin.johnson@noaa.gov](mailto:devin.johnson@noaa.gov)

July 15, 2026

**Abstract.** Remote animal telemetry data is widely used in ecology to assess animal space-use and evaluate utilization distributions (UDs). However, traditional empirical UD (like kernel density or Brownian bridge estimators) can suffer from "start bias" if tag deployments are too short for an animal to fully explore its habitat, heavily biasing results toward the initial release location. Recently, predictive UD have been proposed in an attempt to mitigate this by finding the long-run stationary distribution of a stochastic process that uses habitat covariates to describe movement. This limiting distribution is free from start bias because it uses habitat variables to predict space-use in the future based on short term movements. Predictive UD have been calculated from Langevin diffusions as well as general continuous-time Markov chain models (CTMC). While log-linear closed form UD calculations are possible for Langevin models with coefficients directly relating habitat to space-use, CTMC UD have only been calculated numerically with no parameters directly relating habitat to use. Here we present a general class of CTMC models and show they have a closed form log-linear predictive UD. CTMCs have several benefits over Langevin models: (1) the exact likelihood can be used without approximation, (2) both categorical and continuous habitat variables can be used, and (3) fixed barriers to movement (e.g., land in movement of marine species) are automatically accounted for as part of the model. The R package `walk` was developed for fitting these models in practice and it is used demonstrate explicit UD estimation from CTMC models with a small data set of Hawaiian false killer whales (*Pseudorca crassidens*), including population inference from individual UD.

**Key words:** CTMC, Hidden Markov model, Limiting distribution, Satellite telemetry, Space-use, Utilization distribution

# 1 Introduction

Any quick literature search will reveal that remote collection and analysis of animal locations, or telemetry data, have become a ubiquitous and important tool in the advancement of animal ecology. A perennial subject within animal telemetry analysis is the assessment of space-use. That is what aspects of habitat and movement control where an individual generally occurs on the landscape. The topic of animal space-use is often used in conjunction with the terms home range, resource selection, or utilization distribution. Here we are specifically concerned with the concept of utilization distributions (UD) as defined by Powell and Mitchell (2012), a spatial probability distribution that describes an animal's location at any random time. More specifically, what habitat characteristics influence that distribution.

Literature on UD's is generally divided into two types. The first type is what we term *empirical* UD's. Empirical UD's are constructed based on a summary of observed locations, for example, the traditional kernel density (Keating and Cherry, 2009) or Brownian bridge estimators (Horne et al., 2007). These are empirical because they place positive mass over observed locations. If an animal does not visit a particular area there will be very little if any use estimated for that area depending on how close it is to observed locations. However, absence of observed use may be a symptom of *start bias* (Whitehead and Jonsen, 2013). Start bias results from tag deployments that are not long enough for the animal to fully explore the space and habitat available to it over the course of study. Therefore, spatial use assessments based solely on the observed locations can be biased towards the release location of the animal. This is especially true in recent years where many locations can be obtained over a fine time-scale and thus produce highly autocorrelated paths. To

match the increased technological sophistication of location devices, statistical methodology of empirical UD methods have increased to account for the high autocorrelation (e.g., Fleming et al. 2015; Calabrese et al. 2016). However, the most basic solution for reducing start bias is to increase data collection, i.e., tag deployment interval length (Fieberg et al., 2010). This is easy to prescribe by analysts, but in practice may be difficult to ensure telemetry devices stay active long enough and may be out of the researchers control. This aspect motivates the development of *predictive* UDs.

In order to counter start bias effects researchers can fit resource selection function (RSF) models (Manly et al., 2002) to telemetry data to investigate what space characteristics an animal might prefer relative to available habitat. A map of space preference based on resource selection is used as a UD approximation than is not subject to start bias because it makes use of spatial habitat characteristics than can be assessed at locations away from where the animal was located during tag deployment. However, Barnett and Moorcroft (2008) describe some general results concerning the weighted distribution from Johnson et al. (2008, 2013) of a resource selection movement model in discrete time. Assuming a symmetric movement kernel, they show that the limiting distribution is proportional to the resource selection function or the square of the resource selection function depending on how large the movement kernel is relative to the spatial variation of the habitat. Therefore, while qualitatively similar, RSFs and UDs can be quite different depending on the scale of movement relative to the variation in habitat.

Predictive UDs seek to blend the habitat association benefits of RSF analysis with movement models such that the resulting UD functions, which appear similar to RSFs, represent the actual UD. Predictive UDs are heuristically what a kernel estimate would look like if the animal was allowed to move through its habitat for a very long (infinite)

time. In essence a movement model is fitted to the telemetry data and the location of the animal is predicted from the movement model for a time far in the future. If it exists, this is the stationary or limiting distribution of the movement process. These limiting distributions are independent of the starting location, thus, removing the start bias (Whitehead and Jonsen, 2013; Wilson et al., 2018). Moorcroft et al. (1999) present an early example of a predictive UD approach modeling home range of coyotes (*Canis latrans*) with a movement model which included a response to scent marks. Wilson et al. (2018) extended their work to continuous-time Markov chains (CTMC) that included habitat based movement. Although, the CTMC predicted UD was derived numerically from the fitted model components, there was no direct parameter association that could be used to relate habitat variables with the UD surface. Michelot et al. (2019) used continuous-time and space Langevin diffusions to model resource selection and movement. From this model it is possible to calculate an closed-form predictive UD with the RSF exponential form, that is,

$$\mathbf{u}(\mathbf{s}) \propto \exp\{\mathbf{x}(\mathbf{s})'\boldsymbol{\beta}\}, \quad (1)$$

where  $\mathbf{u}(\mathbf{s})$  is the UD at location  $\mathbf{s}$  and  $\mathbf{x}(\mathbf{s})$  are habitat covariates at location  $\mathbf{s}$ . This model produces the type of closed-form UD prediction that directly relates the UD to habitat variables through the coefficients  $\boldsymbol{\beta}$ . However. There are some drawbacks for this approach. First the likelihood for this model is not available in closed form, so a discrete time (Euler) approximation must be used. Second, this model is only applicable to continuously valued covariates because the habitat gradient surfaces,  $\nabla\mathbf{x}(\mathbf{s})$ , must be used in the movement model (Michelot et al., 2019). Finally, the Langevin diffusion limiting distribution in (1) is only valid when the diffusion component is constant. That is the

animal does not change its overall rate movement in different habitats; only the advection component is altered by habitat (See Section 3). However, Blackwell (2026) has demonstrated that this is not necessary for more general diffusions.

Herein we consider a general class of CTMC models for which we can obtain closed-form predictions of the UD from the fitted model parameters. The CTMC model has several advantages over the Langevin derived UD. First, there is no approximation necessary to calculate the likelihood for optimization. Second, continuous or discrete habitat covariates can be used for movement modeling and UD prediction. Finally, the diffusion rate can be influenced by habitat not just the advection rate. Moreover, because there is no approximation for the likelihood that error does not propagate to the UD prediction, so it is exact. To fit these CTMC models, we developed the R package `walk`\* that can fit general CTMC models including the more general forms requiring numerical calculation of the UD as presented by Hanks et al. (2015) and Wilson et al. (2018). The process of fitting models and making UD inference is demonstrated with telemetry data from a false killer whale (*Pseudorca crassidens*) population in the Hawaiian Islands.

## 2 Explicit Utilization Distribution CTMC Modeling

### 2.1 A brief CTMC primer

Here we provide a brief description of continuous-time Markov chain (CTMC) models. More detail can be found in Norris (1998). In a CTMC movement model the geographical study domain is partitioned into a cells or other discrete locations indexed by  $\mathcal{C} = \{1, \dots, n\}$ . Each cell,  $i$ , has a set of neighboring cells  $\mathcal{N}_i = \{j \in \mathcal{C} : i \sim j\}$ , where  $i \sim j$

---

\*Available from: <https://dsjohnson.r-universe.dev/walk>

means that an animal can go from cell  $i$  to cell  $j$  in one move. For example, in a raster partition, neighboring cells might share a border (rook neighborhood) or a corner point (queen neighborhood). Here we assume that if  $i \sim j$ , then  $j \sim i$ , that is, it is possible to return to a cell that was just exited. As with all CTMCs, the complete path of the individual,  $\mathcal{P}$ , can be decomposed into the *jump times* at which cell transitions are made,  $\mathcal{T} = \{\tau_0, \tau_1, \dots, \tau_M\}$  and the *embedded Markov chain*, which is the sequence of cells visited,  $\mathcal{G} = \{g_0, \dots, g_M\}$ . Thus, the CTMC path has the bivariate discrete-time representation  $\{(\tau_m, g_m) : m = 0, \dots, M\}$ , where  $\tau_0 = 0$  and  $T \geq \tau_M$  is the end of the telemetry deployment. In a slight abuse of notation, we also use  $g_t$  and  $\mathcal{N}_t$  (i.e., indexed by continuous time,  $t \in [0, T]$ ) to represent the location and neighborhood of the individual for any time, that is,  $g_t = g_m$  and  $\mathcal{N}_t = \mathcal{N}_{g_m}$  for  $t \in [\tau_m, \tau_{m+1})$ .

CTMC models are defined by the conditional movement rates between cells,  $\lambda_{ij} \geq 0$ . The total rate of movement away from cell  $i$  is  $\Lambda_i = \sum_{j \in \mathcal{N}_i} \lambda_{ij}$ . One useful property of CTMCs is that the residence time in a cell,  $r_m = \tau_{m+1} - \tau_m$ , is exponentially distributed with rate  $\Lambda_i$ , therefore, the expected residence time in cell  $i$  is  $\Lambda_i^{-1}$ . To describe movement throughout the study area from time  $s$  to time  $t$  we use the  $n \times n$  transition probability matrix,  $\mathbf{\Gamma}(t - s)$ , where the  $(i, j)$  element is the conditional probability  $[g_t = j | g_s = i]$ . To calculate,  $\mathbf{\Gamma}(t - s)$  we put all the rates in the  $n \times n$  matrix  $\mathbf{Q}$  with  $(-\Lambda_1, \dots, -\Lambda_n)$  on the diagonal and  $\lambda_{ij}$  in the  $i$ th row,  $j$ th column. Then the transition matrix is  $\mathbf{\Gamma}(t - s) = \text{expm}\{\mathbf{Q}(t - s)\}$  where “expm” is the matrix exponential function.

If a limiting distribution exists, as  $t \rightarrow \infty$ ,  $\mathbf{\Gamma}(t - s) \rightarrow \mathbf{\Gamma}_\infty$  where each row is equal to the vector  $\mathbf{u}$ , the limiting distribution or in our case the predicted UD. The conditions under which a CTMC will have a long-run limiting distribution over the study area are outlined by Wilson et al. (2018). These conditions are: (1) *Homogeneity*, constant

movement rates through time, (2) *Irreducibility*, any cell can be reached from any other cell given sufficient time, and *Positive recurrence*, an individual will return to cell in before  $t = \infty$ . Wilson et al. (2018) use numerical Eigen decomposition of  $\mathbf{Q}$  to obtain  $\mathbf{u}$ , but they also present an additional result that we use for constructing CTMC models with closed form UD. Assuming the previous conditions are met for a CTMC model, the limiting distribution can be evaluated as

$$u_i \propto \alpha_i / \Lambda_i \quad (2)$$

where  $\alpha$  is the limiting distribution of the embedded Markov chain,  $\{g_0, g_1, \dots\}$ . This makes intuitive sense that the long-run probability of occurrence is the long-run probability the animal is in the cell weighted by the expected amount of time it spends there.

## 2.2 CTMC models with closed form UD

The CTMC movement model we consider for explicit UD modeling has the form,

$$\log(\lambda_{ij}) = \mathbf{x}'_{1i}\boldsymbol{\alpha} + \mathbf{x}'_{2j}\boldsymbol{\delta} + (\mathbf{x}_{3j} - \mathbf{x}_{3i})'\boldsymbol{\gamma} + \mathbf{z}'_{ij}\boldsymbol{\eta}, \quad (3)$$

where  $\mathbf{x}_{1i}$  is a set of habitat covariates recorded in the current cell  $i$ ,  $\mathbf{x}_{2j}$  are other covariates recorded in cell  $j \in \mathcal{N}_i$ ,  $\mathbf{x}_{3j} - \mathbf{x}_{3i}$  are local differences of a third set of habitat covariates, and  $\mathbf{z}_{ij}$  are symmetric pair specific covariates ( $\mathbf{z}_{ij} = \mathbf{z}_{ji}$ ) such as, say, distance between  $i$  and  $j$ . The covariates in  $\mathbf{x}_1$  only control residence time and do not affect which cell an individual might choose upon moving. Variables in  $\mathbf{x}_2$  primarily function to specify preference for neighboring cells based on the absolute value of the covariates, while the local differences control movement based on relative habitat differences. Both of these



movement effects also influence residence time as well. That is if a pull to another habitat is strong a movement to that cell is more likely in the near term.

Using Markov Chain Monte Carlo (MCMC) theoretical results (Michelot et al., 2018; Blackwell, 2026) we can show that a CTMC movement model that satisfies the previous three conditions to have a limiting distribution and has rates of the form (3), it has the limiting distribution  $u_i \propto \exp(\mathbf{x}'_i \boldsymbol{\beta})$ , where  $\mathbf{x}_i = (\mathbf{x}'_{1i}, \mathbf{x}'_{2i}, \mathbf{x}'_{3i})'$  and  $\boldsymbol{\beta} = (-\boldsymbol{\alpha}', \boldsymbol{\delta}', 2\boldsymbol{\gamma})'$  (see online supplement S1 for proof). For a given application, satifying the three general conditions is largely a function of cell partition and study area. The homogeneity requirement is satisfied by constant rates through time. For irreducibility, the study area cannot be separated in to multiple areas such that travel between them is impossible. Finally, for irreducibility, the study area cannot possess any permeable barriers such that the individual cannot return once crossed.

## 2.3 Parameter Estimation

Previous estimation of CTMC movement parameters using satellite location telemetry data used an imputation method (see Scharf et al. 2017) which fills in missing cell locations between location acquisition times (Hanks et al., 2015; Wilson et al., 2018; Johnson et al., 2021). Even though the imputation method attempts to propagate the uncertainty of the missing locations, it is unclear how this affects estimation of the parameters. Therefore we will use the exact, although potentially more computationally expensive hidden Markov model (HMM) formulation of the model that is often use for CTMC analysis marine fish movement from archival telemetry tags (e.g., see CH. 10 of Thorson and Kristensen 2024).

### 2.3.1 Satellite telemetry data specification and processing

For the proposed analysis we consider satellite telemetry data for an individual of the form  $\mathbf{y} = \{\mathbf{y}_1, \dots, \mathbf{y}_K\}$ , where  $\mathbf{y}_k = (\mathbf{s}_k, t_k, \boldsymbol{\epsilon}_k)$ ,  $\mathbf{s}_k$  is the observed location of the individual within the study area at time  $t_k$  and  $\boldsymbol{\epsilon}_k$  represents information or parameters governing location uncertainty. For example,  $\boldsymbol{\epsilon}_k$  might represent the major and minor error ellipse lengths of a Kalman Filtered Argos location (McClintock et al., 2015) or a known error radius for GPS locations, e.g., 20m. From these locations we translate the continuous-space data locations and uncertainty to the discrete grid,  $\mathcal{C}$ , using a function, say,  $p(\mathbf{y}_k, \mathcal{C}) = \mathbf{p}_k = (p_{1k}, \dots, p_{nk})$ , where  $p_{ik}$  is the probability that the animal is located within cell  $i$  at time  $t_k$ . We used a general form because the function,  $p$ , will depend on how location uncertain is assessed for the specific satellite and tag hardware. However, uncertainty in the true location,  $\boldsymbol{\mu}_k$ , is often represented by the bivariate Normal distribution  $\boldsymbol{\mu}_k \sim N(\mathbf{s}_k, \boldsymbol{\Sigma}_k)$ , where  $\boldsymbol{\epsilon}_k = \boldsymbol{\Sigma}_k$ . In that case,  $p_{ik}$  would represent the mass of the Normal distribution contained within the  $i$ th cell. For a raster grid partitioning of the study area this can be calculated by,

$$p_{ik} = \text{CDF}(x_{i2}, y_{i2}) - \text{CDF}(x_{i2}, y_{i1}) - \text{CDF}(x_{i1}, y_{i2}) + \text{CDF}(x_{i1}, y_{i1})$$

where  $[x_{i1}, x_{i2}] \times [y_{i1}, y_{i2}]$  is the bounding box for the  $i$ th cell and CDF is the bivariate normal cumulative density function with mean  $\mathbf{s}_k$  and covariance  $\boldsymbol{\Sigma}_k$ . A Monte Carlo option for other error distributions would be to draw a sufficiently large sample from the distribution and assess the proportion of samples that falls within each cell.

### 2.3.2 Likelihood calculation

For an individual the likelihood is calculated with the matrix equation (Zucchini et al. 2016; Ch. 3),

$$[\mathbf{y}|\boldsymbol{\alpha}, \boldsymbol{\delta}, \boldsymbol{\gamma}, \boldsymbol{\eta}] = \boldsymbol{\pi}' \mathbf{P}(\mathbf{y}_1) \boldsymbol{\Gamma}(t_2 - t_1) \mathbf{P}(\mathbf{y}_2) \cdots \boldsymbol{\Gamma}(t_K - t_{K-1}) \mathbf{P}(\mathbf{y}_K) \mathbf{1} \quad (4)$$

where  $\boldsymbol{\pi} = (\pi_1, \dots, \pi_n)$  is the probability at the individual is cell  $i$  at  $t_1$ ,  $\mathbf{P}(\mathbf{y}_k) = \text{diag}(\mathbf{p}_k)$ ,  $\boldsymbol{\Gamma}(t_k - t_{k-1}) = \exp\{\mathbf{Q}(t_k - t_{k-1})\}$ , and  $\mathbf{1}$  is a vector of 1s. The natural choice for  $\boldsymbol{\pi}$  is  $\mathbf{u}$ , the limiting distribution defined by the CTMC movement model, but there are other choices. For example, if a telemetered individual was released at a known location after tagging that may not be representative of its usual movement, then  $\boldsymbol{\pi}$  would be a vector of all 0s except for the cell containing the release site, which would be a 1. In the false killer whale example below, we use  $\boldsymbol{\pi} = \mathbf{u}$  because they were tagged where they were encountered in the wild. In the R package `walk` we use scaling methods to avoid numerical underflow and well as the efficient sparse matrix methods presented in Sherlock (2021) to calculate products of the form  $\mathbf{v}' \exp\{\mathbf{Q}(t_k - t_{k-1})\}$ .

In addition to the matrix exponential evaluation numerical savings, another optional approach to save computational effort is to set a censoring value for the maximum  $\Lambda_i$  value. The numerical methods of Sherlock (2021) require more matrix operations for higher values of  $\max(\Lambda_i)$ . For some cells, the log linear  $\lambda_{ij}$  model may produce rates of movement that are unrealistically high in some parts of the study area. With respect to UDs this would imply that an individual will not remain in that cell and the  $u_i \approx 0$ . Hewitt et al. (2023) considered a logit rate model that restricts this behavior, but that model does not allow us to construct a CTMC with an explicit limiting distribution. So, we propose

216 picking a censoring value, say,  $\Lambda^*$  and setting  $\lambda_{ij}^* = (\Lambda^*/\Lambda_i)\lambda_{ij}$  if  $\Lambda_i > \Lambda^*$ . If  $\Lambda^*$  is  
 217 sufficiently large, this will have negligible impact on  $\mathbf{u}$  estimation but can dramatically help  
 218 numerical likelihood evaluation. One reasonable way to choose  $\Lambda^*$  is to select the minimum  
 219 amount of time that an animal would likely to reside in a cell, say  $r^*$ . Then set  $\Lambda^*$  so that  
 220  $Pr(r < r^*|\Lambda^*)$  is near 1. This implies that cells with  $\Lambda_i > \Lambda^*$ , residence time is sufficiently  
 221 short such that we can censor the rate at  $\Lambda^*$ . The effect on  $\mathbf{u}$  can be verified by comparing  
 222 the exact, numerically calculated  $\mathbf{u}$  from the censored rate model (Wilson et al., 2018) to  
 223  $u_i \propto \exp(\mathbf{x}'_i\boldsymbol{\beta})$ . If there are sufficiently large discrepancies,  $\Lambda^*$  can be increased. When  
 224  $\Lambda^* \geq \max(\Lambda_i)$  the UD's are guaranteed to be identical.

225 Once the CTMC movement model has been fitted the maximum likelihood estimate of  
 226  $\hat{\boldsymbol{\beta}}_{\text{CTMC}} = (\boldsymbol{\alpha}', \boldsymbol{\delta}', \boldsymbol{\gamma}')'$  needs to be transformed to the UD coefficients via  $\hat{\boldsymbol{\beta}} = \mathbf{M}\hat{\boldsymbol{\beta}}_{\text{CTMC}}$  and  
 227  $\mathbf{M} = \text{diag}((-1'_{\alpha}, 1'_{\delta}, 21'_{\gamma})')$ . Thus, the large sample covariance matrix of the UD coefficients  
 228 is  $\mathbf{V}(\hat{\boldsymbol{\beta}}) = \mathbf{M}\mathbf{V}(\hat{\boldsymbol{\beta}}_{\text{CTMC}})\mathbf{M}$ . If only the standard errors are desired, one can simply multiply  
 229 the CTMC coefficient standard errors by the multipliers, but if the entire covariance matrix  
 230 is desired the matrix multiplication is necessary.

### 231 **3 False Killer Whale Space-Use**

232 In this section, we will apply the modeling approach described in the previous sections to a  
 233 telemetry dataset on endangered false killer whales (FKW; *Pseudorca crassidens*) in the  
 234 Hawaiian Islands (Baird 2016). This particular population is resident to the insular waters  
 235 of the main Hawaiian Islands (MHI; includes Kaua'i/Ni'ihau, Oahu, Maui Nui, and Hawai'i  
 236 Island) and is comprised of five social clusters, or groups of close family members and  
 237 regular associates (Baird et al., 2008, 2019; Baird, 2016; Martien et al., 2019). These

whales are known to move quickly and frequently among island areas throughout their range (Baird et al., 2010, 2012, 2021; Baird, 2016). Due to inherent challenges associated with tracking apex marine predators, this dataset is characterized by variable deployment durations and irregular location data streams. Considering these factors may drive start bias, as well as their expansive range, we deem this an appropriate dataset to empirically demonstrate the utility of predictive UD. For the purposes of this assessment, we only focus on movements of 5 individuals that were satellite tagged in 2021 and 2022. There is one representative from each social cluster in the data.

### 3.1 Satellite tag data

False killer whale satellite tag deployment methods have been described in detail elsewhere (see Baird et al. 2010, 2012) so they are only briefly summarized here. Individuals were tagged with Argos transmitting SPOT6 ( $n = 4$ ) and SPLASH10 ( $n = 1$ ) tags during dedicated small boat survey efforts throughout the main Hawaiian Islands (Baird et al. 2012, 2021; Baird 2016; see Baird et al. 2013 for details on surveys). Deployment duration ranged from 12 days to 81 days (median=40 days). Argos location data were processed through the Douglas-Argos Filter (accessed via Movebank; Kranstauber et al. 2011; Douglas et al. 2012) to remove erroneous locations based on unrealistic travel speeds and turning angles. The final analytical dataset consisted of 5 trajectories and 4,009 filtered Argos and Fastloc GPS (SPASH10 tag) locations (Figure 1A). For the Argos locations, we used the Kalman-filter diagnostic ellipse data to construct error covariance matrices (McClintock et al., 2015). While for the Fastloc GPS locations, we used the 95% quantile error ranges from Dujon et al. (2014), based on the number of satellites acquired, as twice the standard deviation of the normal error.

## 3.2 Study area and habitat variables

The official management range of the MHI stock of FKW was documented in Bradford et al. (2015) and represents a minimum convex polygon surrounding a buffer of the main Hawaiian Islands (see Figure 1A). This is the study area boundary we use for the UD analysis.

For habitat characteristics we used two modeled global oceanographic variables from E.U. Copernicus Marine Service: (1) Sea Surface Temperature (SST; °C; DOI: <https://doi.org/10.48670/moi-00016>) and (2) Sea Level Anomaly (SLA; m; DOI: <https://doi.org/10.48670/moi-00148>). SST is available at 0.082° resolution and SLA is available at 0.125° degree resolution. In order to match resolution we chose to project the SLA raster to the resolution of the SST raster to avoid losing oceanographic information. Because we are using these variables to predict long term space-use, we used a single layer for each variable that was the 2 year daily average in the study area. This allowed us to model movement with respect to typical oceanographic conditions (Figure 1B-C).

In addition to the oceanographic variables, we used distance from land (m) to model FKW's resident nature in the MHI. Also, we also included a depth residual variable by adding data from the global, 30 arc-second ETOPO elevation model (NOAA National Centers for Environmental Information, 2022) and removing the linear trend of distance from land. The distance and depth variables were projected to the 0.082° SST resolution as well. Then all data, including telemetry data, were projected to the Hawaii Albers Equal Area projection, ESRI:102007 (Figure 1D-E)

### 3.3 CTMC model and population inference

The CTMC movement rate model we used to model the telemetry for each individual

$l = 1, \dots, 5$  was

$$\log \lambda_{ij} = \beta_0 + \underbrace{\text{DEPTH}_i + \text{SST}_i + \text{SST}_i^2 + \text{SLA}_i + \text{SLA}_i^2}_{\mathbf{x}_1 \text{ portion}} + \underbrace{[\text{DIST}_j - \text{DIST}_i]}_{\mathbf{x}_3 \text{ portion}}$$

In this model differences of the distance from land variable were used as a movement kernel between cells  $i$  and  $j$ . This allowed individuals to prefer movement towards (or away from) land at the same rate no matter how far away they are. In addition SST, SLA, and DIST were  $Z$ -transformed for numerical stability. The DEPTH residual was already on a nearly standard normal scale and DIST was only used as a difference, so values were not large enough to cause issue.

Individually estimated FKW coefficients and predicted UD's can be difficult to interpret for population inference and management considerations. Therefore, we used the 2-stage meta-analysis method of Johnson et al. (2022) to approximate the Bayesian random effects model,

$$[\mathbf{y}_1, \dots, \mathbf{y}_N | \boldsymbol{\beta}_1, \dots, \boldsymbol{\beta}_N] = \prod_{l=1}^N [\mathbf{y}_l | \mathbf{M}^{-1} \boldsymbol{\beta}_l]$$

$$[\boldsymbol{\beta}_1, \dots, \boldsymbol{\beta}_N | \boldsymbol{\theta}, \boldsymbol{\Sigma}] = \prod_{l=1}^N N(\boldsymbol{\beta}_l | \boldsymbol{\theta}, \boldsymbol{\Sigma})$$

where  $\mathbf{y}_l$  is the observed data for individual  $i = 1, \dots, N$  and the prior distribution for the random effect mean and variance is  $[\boldsymbol{\theta}] [\boldsymbol{\Sigma}]$ . Even if individual inference is desired, it is useful to fit this hierarchical model because it allows us to share information between individuals. That is we can make inference to the posterior distribution  $[\boldsymbol{\beta}_l | \mathbf{y}_1, \dots, \mathbf{y}_N]$ . If we assume that there is some similarity in movement between individuals then we can

leverage all data for better individual inference.

In the first stage the CTMC model was fitted separately to the data from each individual using normal location errors. For the rate truncation we set  $\Lambda^* = 138.16$ . This resulted from selecting the minimum residence time,  $r^*$  to be 1 minute and residence probability 0.9. The R function `walk::find_clip` will calculate  $\Lambda^*$ . In the second stage we fitted the multivariate linear mixed model,

$$\hat{\beta}_l = \boldsymbol{\theta} + \mathbf{u}_l + \boldsymbol{\delta}_l; \quad l = 1, \dots, N$$

where  $\hat{\beta}_l$  is the MLE estimate of the coefficients for the  $l$ th individual, the random effect is  $[\mathbf{u}_l | \boldsymbol{\Sigma}] = N(\mathbf{0}, \boldsymbol{\Sigma})$ , the error term is  $[\boldsymbol{\delta}_l | \mathbf{V}(\hat{\beta}_l)] = N(\mathbf{0}, \mathbf{V}(\hat{\beta}_l))$ , and  $\mathbf{V}(\hat{\beta}_l)$  is the parameter covariance matrix from the first MLE step. For this analysis we used independent random effects, that is,  $\boldsymbol{\Sigma} = \text{diag}(\sigma_1^2, \dots, \sigma_7^2)$ . The `nimble` R package (de Valpine et al., 2017) was used to optimize model with respect to the parameters (i.e., maximum *a posteriori*, MAP) subject to the prior distributions,

$$[\theta_w] = N(0, \tau_w^2); \quad [\tau_w] = \text{Exponential}(1)$$

$$[\sigma_w] = \text{Exponential}(1)$$

for  $w = 1, \dots, 7$ . The exponential priors for the variance parameters were added to penalize the complexity of the model (i.e., regularization) to perform selection for the population level parameters (Hooten and Hobbs, 2015; Simpson et al., 2017). Following second stage model fitting we can extract the approximate MAP estimates  $\bar{\boldsymbol{\theta}}$  and  $\bar{\beta}_1, \dots, \bar{\beta}_N$ , where  $\bar{\beta}_l$  is the approximate MAP of the posterior  $[\beta_l | \mathbf{y}_1, \dots, \mathbf{y}_N]$ . The linear mixed model was fitted using the optimization capabilities of the R package `nimble` (Citation, 2015).



### 3.4 Results

Individual results for all individuals were generally consistent with notable deviations being individual PcTag074 did not have any effect of SLA on its UD and PcTag070 had negative effect of the linear SST term (Figure 2). All other coefficients were similar in direction, i.e., less than or greater than 0. This bolsters the idea of using the 2 stage meta-analytic approach to share data between individuals to improve estimates. The second stage individual coefficient estimates,  $\bar{\beta}_i$  illustrate that there is only a significant amount of individual variation in the DEPTH and SLA<sup>2</sup> coefficients (Figure 2). All other second stage coefficients are shrunk to the population mean estimates,  $\bar{\theta}$  with nearly identical approximate credible intervals (Figure 2). There was only 1 nonsignificant population coefficient, the linear SLA term. Correspondingly, the penalized model complexity priors shrunk the SLA term to zero. All other population coefficients were different than zero. The DEPTH and DIST coefficients were negative as expected because this population inhabits shallower waters close to the MHI without venturing away. The zero SLA coefficient combined with the negative SLA<sup>2</sup> term implies a maximum effect at 0, but we used a Z-transform, therefore, the maximum effect occurs at the mean original SLA value. Both SST coefficients are positive implying that the typical individual is probably located in the warmest SST conditions in the study area.

Unsurprisingly, the similarity in coefficient estimates produced similar predicted UDs at the first stage (Figure 3). Individual PcTag074 had a more notably diffuse UD, the other four individuals had very similar looking UDs. This may be due to the lack of SLA effect on the UD of PcTag074. After the second stage, the UDs are even more similar as the second stage coefficients are essentially the same at the population mean coefficients

(Figure 4). The UD created from the population mean coefficients is illustrated in Figure 4 (Typical). This is the UD of a so called “Typical” or median individual, not the population average UD because of the log link between the coefficients and the UD. Even after the second stage the coefficients for PcTag074 lack any SLA effect, so the UD is the only substantially different one from the typical UD.

In order to check that the rate truncation did not adversely affect the estimated UDs we compared the truncated numerical Eigen decomposition UD ( $u_{ei,i}$ ; Wilson et al. 2018), which is the exact UD for the fitted movement models, with the exponential closed form version using Bhattacharyya’s Affinity ( $BA$ ) (Bhattacharyya, 1943),

$$BA = \sum_{i \in \mathcal{C}} \sqrt{u_{ei,i} \cdot u_i}.$$

The values of  $BA$  range from 0 (no similarity) to 1 (identical distributions) (Fieberg and Kochanny, 2005). The proportion of study area cells truncated in the fitting of individual models ranged from 0–44% of the study area. However,  $BA$  ranged from 0.996–1 indicated that the truncated cells possessed very little UD mass and the closed form and exact truncated UDs were virtually identical. Therefore, we can be assured that the truncation approximation did not adversely affect parameter estimation and no adjustment is necessary for this analysis.

## 4 Discussion

## Acknowledgments

The findings and conclusions in the paper are those of the authors and do not necessarily represent the views of the National Marine Fisheries Service, NOAA. Reference to trade names does not imply endorsement by the National Marine Fisheries Service, NOAA.

## References

- Baird, R. W. (2016). *The Lives of Hawai'i's Dolphins and Whales: Natural History and Conservation*. University of Hawaii press.
- Baird, R. W., Anderson, D. B., Kratofil, M. A., and Webster, D. L. (2021). Bringing the right fishermen to the table: Indices of overlap between endangered false killer whales and nearshore fisheries in Hawai'i. *Biological Conservation*, 255:108975.
- Baird, R. W., Anderson, D. B., Kratofil, M. A., Webster, D. L., and Mahaffy, S. D. (2019). Cooperative conservation and long-term management of false killer whales in Hawai'i: Geospatial analyses of fisheries and satellite tag data to understand fishery interactions. A Report to the State of Hawai'i Board of Land and Natural Resources, Contract No. 67703, Cascadia Research Collective.
- Baird, R. W., Gorgone, A. M., McSweeney, D. J., Webster, D. L., Salden, D. R., Deakos, M. H., Ligon, A. D., Schorr, G. S., Barlow, J., and Mahaffy, S. D. (2008). False killer whales (*Pseudorca crassidens*) around the main Hawaiian Islands: Long-term site

fidelity, inter-island movements, and association patterns. *Marine Mammal Science*,  
24(3):591–612.

Baird, R. W., Hanson, M. B., Schorr, G. S., Webster, D. L., McSweeney, D. J., Gorgone,  
A. M., Mahaffy, S. D., Holzer, D. M., Oleson, E. M., and Andrews, R. D. (2012). Range  
and primary habitats of Hawaiian insular false killer whales: Informing determination of  
critical habitat. *Endangered Species Research*, 18(1):47–61.

Baird, R. W., Schorr, G. S., Webster, D. L., McSweeney, D. J., Hanson, M. B., and  
Andrews, R. D. (2010). Movements and habitat use of satellite-tagged false killer whales  
around the main Hawaiian Islands. *Endangered Species Research*, 10:107–121.

Baird, R. W., Webster, D. L., Aschettino, J. M., Schorr, G. S., and McSweeney, D. J.  
(2013). Odontocete cetaceans around the main Hawaiian Islands: Habitat use and  
relative abundance from small-boat sighting surveys. *Aquatic Mammals*, 39(3).

Barnett, A. H. and Moorcroft, P. R. (2008). Analytic steady-state space use patterns and  
rapid computations in mechanistic home range analysis. *Journal of Mathematical  
Biology*, 57(1):139–159.

Bhattacharyya, A. (1943). On a measure of divergence between two statistical populations  
defined by their probability distribution. *Bulletin of the Calcutta Mathematical Society*,  
35:99–110.

Blackwell, P. G. (2026). Consistent multiscale modelling of movement and habitat  
selection. *Journal of Mathematical Biology*, 92(6):90.

Bradford, A. L., Oleson, E. M. E. M., Baird, R. W., Boggs, C. H., Forney, K. A., and

- Young, N. C. (2015). Revised stock boundaries for false killer whales (*Pseudorca crassidens*) in Hawaiian waters. <https://repository.library.noaa.gov>.
- Calabrese, J. M., Fleming, C. H., and Gurarie, E. (2016). Ctmm: An R package for analyzing animal relocation data as a continuous-time stochastic process. *Methods in Ecology and Evolution*, 7(9):1124–1132.
- Citation (2015). Unknown. *Unknown*.
- de Valpine, P., Turek, D., Paciorek, C. J., Anderson-Bergman, C., Lang, D. T., and Bodik, R. (2017). Programming with models: Writing statistical algorithms for general model structures with NIMBLE. *Journal of Computational and Graphical Statistics*, 26(2):403–413.
- Douglas, D. C., Weinzierl, R., C. Davidson, S., Kays, R., Wikelski, M., and Bohrer, G. (2012). Moderating A rgos location errors in animal tracking data. *Methods in Ecology and Evolution*, 3(6):999–1007.
- Dujon, A. M., Lindstrom, R. T., and Hays, G. C. (2014). The accuracy of Fastloc-GPS locations and implications for animal tracking. *Methods in Ecology and Evolution*, doi: 10.1111/2041-210X.12286.
- Fieberg, J. and Kochanny, C. O. (2005). Quantifying Home-Range Overlap: The Importance of the Utilization Distribution. *The Journal of Wildlife Management*, 69(4):1346–1359.
- Fieberg, J., Matthiopoulos, J., Hebblewhite, M., Boyce, M. S., and Frair, J. L. (2010). Correlation and studies of habitat selection: Problem, red herring or opportunity?

*Philosophical Transactions of the Royal Society B-Biological Sciences*,

365(1550):2233–2244.

Fleming, CH., Fagan, WF., Mueller, T., Olson, KA., Leimgruber, P., and Calabrese, JM.

(2015). Rigorous home range estimation with movement data: A new autocorrelated

kernel density estimator. *Ecology*, 96(5):1182–1188.

Hanks, E. M., Hooten, M. B., and Alldredge, M. W. (2015). Continuous-time

discrete-space models for animal movement. *The Annals of Applied Statistics*, 9:145–165.

Hewitt, J., Gelfand, A. E., and Schick, R. S. (2023). Time-discretization approximation

enriches continuous-time discrete-space models for animal movement. *The Annals of*

*Applied Statistics*, 17(1):740–760.

Hooten, M. B. and Hobbs, N. T. (2015). A guide to Bayesian model selection for ecologists.

*Ecological Monographs*, 85(1):3–28.

Horne, J. S., Garton, E. O., Krone, S. M., and Lewis, J. S. (2007). Analyzing animal

movements using Brownian bridges. *Ecology*, 88:2354–2363.

Johnson, D. S., Brost, B. M., and Hooten, M. B. (2022). Greater than the sum of its parts:

Computationally flexible Bayesian hierarchical modeling. *Journal of Agricultural,*

*Biological and Environmental Statistics*, 27(2):382–400.

Johnson, D. S., Hooten, M. B., and Kuhn, C. E. (2013). Estimating animal resource

selection from telemetry data using point process models. *Journal of Animal Ecology*,

82:1155–1164.

Johnson, D. S., Pelland, N. A., and Sterling, J. T. (2021). A continuous-time semi-Markov

model for animal movement in a dynamic environment. *Annals of Applied Statistics*,  
15(2):797–812.

Johnson, D. S., Thomas, D. L., Hoef, J. M. V., and Christ, A. (2008). A general framework  
for the analysis of animal resource selection from telemetry data. *Biometrics. Journal of  
the International Biometric Society*, 64(3):968–976.

Keating, K. A. and Cherry, S. (2009). Modeling utilization distributions in space and time.  
*Ecology*, 90:1971–1980.

Kranstauber, B., Cameron, A., Weinzerl, R., Fountain, T., Tilak, S., Wikelski, M., and  
Kays, R. (2011). The Movebank data model for animal tracking. *Environmental  
Modelling & Software*, 26(6):834–835.

Manly, B., McDonald, L., Thomas, D., McDonald, T., and Erikson, W. (2002). *Resource  
Selection by Animals*. Kluwer Academic Publishers, 2nd edition, pp. 221 edition.

Martien, K. K., Taylor, B. L., Chivers, S. J., Mahaffy, S. D., Gorgone, A. M., and Baird,  
R. W. (2019). Fidelity to natal social groups and mating within and between social  
groups in an endangered false killer whale (*Pseudorca crassidens*) population.  
*Endangered Species Research*, 40:219–230.

McClintock, B. T., London, J. M., Cameron, M. F., and Boveng, P. L. (2015). Modelling  
animal movement using the Argos satellite telemetry location error ellipse. *Methods in  
Ecology and Evolution*, In press:doi: 10.1111/2041-210X.12311.

Michelot, T., Blackwell, P. G., and Matthiopoulos, J. (2018). Linking resource selection  
and step selection models for habitat preferences in animals. *Ecology*.

460 Michelot, T., Etienne, M.-P., and Gloaguen, P. (2019). The Langevin diffusion as a  
461 continuous-time model of animal movement and habitat selection. *Methods In Ecology*  
462 *and Evolution*, In press.

463 Moorcroft, P. R., Lewis, M. A., and Crabtree, R. L. (1999). Home range analysis using a  
464 mechanistic home range model. *Ecology*, 80(5):1656–1665.

465 NOAA National Centers for Environmental Information (2022). ETOPO 2022 15  
466 arc-second global relief model.

467 Norris, J. R. (1998). *Markov Chains*. Cambridge University Press, 2nd edition edition.

468 Powell, R. A. and Mitchell, M. S. (2012). What is a home range? *Journal of mammalogy*,  
469 93(4):948–958.

470 Scharf, H., Hooten, M. B., and Johnson, D. S. (2017). Imputation Approaches for Animal  
471 Movement Modeling. *Journal of Agricultural, Biological, and Environmental Statistics*,  
472 22(3):335–352.

473 Sherlock, C. (2021). Direct statistical inference for finite Markov jump processes via the  
474 matrix exponential. *Computational Statistics*, 36(4):2863–2887.

475 Simpson, D. P., Rue, H., Riebler, A., Martins, T. G., and Sørbye, SH. (2017). Penalising  
476 model component complexity: A principled, practical approach to constructing priors.  
477 *Statistical Science*, 32:1–28.

478 Thorson, J. and Kristensen, K. (2024). *Spatio-Temporal Models for Ecologists*. Chapman  
479 and Hall/CRC.



- 480 Whitehead, H. and Jonsen, I. D. (2013). Inferring animal densities from tracking data  
481 using markov chains. *PloS one*, 8(4):e60901.
- 482 Wilson, K., Hanks, E., and Johnson, D. (2018). Estimating animal utilization densities  
483 using continuous-time Markov chain models. *Methods in Ecology and Evolution*,  
484 9(5):1232–1240.
- 485 Zucchini, W., MacDonald, I. L., and Langrock, R. (2016). *Hidden Markov Models for Time*  
486 *Series: An Introduction Using R*. CRC Press, 2 edition.

**Figure 1:** Habitat covariates

**Figure 2:** Parameter estimates for fitted UD surfaces. The Stage 1 values are from individually fitted models. The Stage 2 estimates are parameters estimates using the two stage method of Johnson et al. (2022) to approximate a combined random effects model. The solid vertical lines are the estimated random effect means and the dashed lines are the approximate 95% credible intervals.

**Figure 3:** Individual model predicted utilization distributions. These UDs were predicted using only the data from each respective individual. A UD prediction for PcTag071 is absent because the resulting prediction was nonsensical given the individual movement parameter estimates. That is, large (but highly uncertain) SLA and  $SLA^2$  coefficients resulted in only a few cells getting all the UD mass. This illustrates one of the problem when using movement models to predict space use.

**Figure 4:** Predicted UDs after 2 stage approach.




Figure 1




Figure 2




Figure 3




Figure 4